

25:157–163, 1995.

- [63946] 2276.Wakui, K.; Nishida, T.; Masuda, J.; Itoh, T.; Katsumata, D.; Ohno,T.; Fukushima, Y.: De novo interstitial deletion of 4q[46,XX,del(4)(q27q28.2)]with intact blood group–MN locus, confining its locus to 4q28.2–4q31.1. *Jpn.J. Hum. Genet.* 36: 149–153, 1991.
- [63947] 2277.Wilson, A. F.; Elston, R. C.; Siervogel, R. M.; Weinshilboum,R.; Ward, L. J.: Linkage relationships between a major gene for catechol–O–methyltransferaseactivity and 25 polymorphic marker systems. *Am. J. Med. Genet.* 19:525–532, 1984.
- [63948] 2278.Mattera, R.; Graziano, M. P.; Yatani, A.; Zhou, Z.; Graf, R.;Codina, J.; Birnbaumer, L.; Gilman, A. G.; Brown, A. M.: Splice variantsof the alpha subunit of the G protein G(8) activate both adenylylcyclase and calcium channels. *Science* 243: 804–807, 1989.
- [63949] 2279.Mehlmann, L. M.; Jones, T. L. Z.; Jaffe, L. A.: Meiotic arrestin the mouse follicle maintained by a GS protein in the oocyte. *Science* 297:1343–1345, 2002.
- [63950] 2280.Nerlich, A.; Peschel, O.; Lohrs, U.; Parsche, F.; Betz, P.: Juvenilegigantism plus polyostotic fibrous dysplasia in the Tegernsee giant.(Letter) *Lancet* 338: 886–887, 1991.
- [63951] 2281.Patten, J. L.; Johns, D. R.; Valle, D.; Eil, C.; Gruppuso,

P.A.; Steele, G.; Smallwood, P. M.; Levine, M. A.: Mutation in the gene encoding the stimulatory G protein of adenylylate cyclase in Albright's hereditary osteodystrophy. *New Eng. J. Med.* 322: 1412–1419, 1990.

[63952] 2282. Persani, L.; Borgato, S.; Lania, A.; Filopanti, M.; Mantovani, G.; Conti, M.; Spada, A.: Relevant cAMP-specific phosphodiesterase isoforms in human pituitary: effect of Gs-alpha mutations. *J. Clin. Endocr. Metab.* 86: 3795–3800, 2001.

[63953] 2283. Peters, J.; Beechey, C. V.; Ball, S. T.; Evans, E. P.: Mapping studies of the distal imprinting region of mouse chromosome 2. *Genet. Res.* 63: 169–174, 1994.

[63954] 2284. Premawardhana, L. D. K. E.; Vora, J. P.; Mills, R.; Scanlon, M. F.: Acromegaly and its treatment in the McCune–Albright syndrome. *Clin. Endocr.* 36: 605–608, 1992.

[63955] 2285. Riminucci, M.; Fisher, L. W.; Majolagbe, A.; Corsi, A.; Lala, R.; De Sanctis, C.; Robey, P. G.; Bianco, P.: A novel GNAS1 mutation, R201G, in McCune–Albright syndrome. *J. Bone Miner. Res.* 14: 1987–1989, 1999.

[63956] 2286. Schwindinger, W. F.; Francomano, C. A.; Levine, M. A.: Identification of a mutation in the gene encoding the alpha subunit of the stimulatory G-protein of adenylyl cy-



clase in McCune–Albright syndrome. Proc.Nat. Acad. Sci. 89: 5152–5156, 1992.

[63957] 2287.Schwindinger, W. F.; Francomano, C. A.; Levine, M. A.; McKusick,V. A.: DNA light on the Tegernsee giant. (Letter) Lancet 338: 1454–1455,1991.

[63958] 2288.Shapira, H.; Mouallem, M.; Shapiro, M. S.; Weisman, Y.; Farfel,Z.: Pseudohypoparathyroidism type Ia: two new heterozygous frameshiftmutations in exons 5 and 10 of the Gs alpha gene. Human Genet. 97:73–75, 1995.

[63959] 2289.Shenker, A.; Chanson, P.; Weinstein, L. S.; Chi, P.; Spiegel,A. M.; Lomri, A.; Marie, P. J.: Osteoblastic cells derived from isolatedlesions of fibrous dysplasia contain activating somatic mutationsof the G(S)–alpha gene. Hum. Molec. Genet. 4: 1675–1676, 1995.

[63960] 2290.Shenker, A.; Weinstein, L. S.; Moran, A.; Pescovitz, O. H.; Charest,N. J.; Boney, C. M.; Van Wyk, J. J.; Merino, M. J.; Feuillan, P. P.;Spiegel, A. M.: Severe endocrine and nonendocrine manifestationsof the McCune–Albright syndrome associated with activating mutationsof stimulatory G protein Gs. J. Pediat. 123: 509–518, 1993.

[63961] 2291.Shore, E. M.; Ahn, J.; Jan de Beur, S.; Li, M.; Xu, M.; Gardiner,R. J. M.; Zasloff, M. A.; Whyte, M. P.; Levine, M. A.; Kaplan, F.S.: Paternally inherited inactivating mutations

of the GNAS1 gene in progressive osseous heteroplasia.

New Eng. J. Med. 346: 99–106, 2002.

[63962] 2292. Shore, E. M.; Kaplan, F. S.; Levine, M. A.: GNAS1 mutations and progressive osseous heteroplasia. (Letter) New Eng. J. Med. 346: 1670–1671, 2002.

[63963] 2293. Sparkes, R. S.; Cohn, V. H.; Mohandas, T.; Zollman, S.; Cire-Eversole, P.; Amatruda, T. T.; Reed, R. R.; Lochrie, M. A.; Simon, M. I.: Mapping of genes encoding the subunits of guanine nucleotide-binding protein (G-proteins) in humans. (Abstract) Cytogenet. Cell Genet. 46: 696 only, 1987.

[63964] 2294. Tinschert, S.; Gerl, H.; Gewies, A.; Jung, H.-P.; Nurnberg, P.: McCune-Albright syndrome: clinical and molecular evidence of mosaicism in an unusual giant patient. Am. J. Med. Genet. 83: 100–108, 1999.

[63965] 2295. Vallar, L.; Spada, A.; Giannattasio, G.: Altered Gs and adenylate cyclase activity in human GH-secreting pituitary adenomas. Nature 330: 566–568, 1987.

[63966] 2296. Warner, D. R.; Gejman, P. V.; Collins, R. M.; Weinstein, L. S.: A novel mutation adjacent to the switch III domain of Gs- $\alpha$  in a patient with pseudohypoparathyroidism. Molec. Endocr. 11: 1718–1727, 1997.

[63967] 2297. Warner, D. R.; Weinstein, L. S.: A mutation in the

heterotrimeric stimulatory guanine nucleotide binding protein alpha-subunit with impaired receptor-mediated activation because of elevated GTPase activity. *Proc. Nat. Acad. Sci.* 96: 4268–4272, 1999.

[63968] 2298. Warner, D. R.; Weng, G.; Yu, S.; Matalon, R.; Weinstein, L. S.: A novel mutation in the switch 3 region of Gs-alpha in a patient with Albright hereditary osteodystrophy impairs GDP binding and receptor activation. *J. Biol. Chem.* 273: 23976–23983, 1998.

[63969] 2299. Weinstein, L. S.; Gejman, P. V.; de Mazancourt, P.; American, N.; Spiegel, A. M.: A heterozygous 4-bp deletion mutation in the Gs-alpha gene (GNAS1) in a patient with Albright hereditary osteodystrophy. *Genomics* 13: 1319–1321, 1992.

[63970] 2300. Weinstein, L. S.; Gejman, P. V.; Friedman, E.; Kadowaki, T.; Collins, R. M.; Gershon, E. S.; Spiegel, A. M.: Mutations of the Gs alpha-subunit gene in Albright hereditary osteodystrophy detected by denaturing gradient gel electrophoresis. *Proc. Nat. Acad. Sci.* 87: 8287–8290, 1990.

[63971] 2301. Weinstein, L. S.; Shenker, A.; Gejman, P. V.; Merino, M. J.; Friedman, E.; Spiegel, A. M.: Activating mutations of the stimulatory G protein in the McCune–Albright syndrome. *New Eng. J. Med.* 325: 1688–1695, 1991.

- [63972] 2302.Weinstein, L. S.; Yu, S.: The role of genomic imprinting of GS-alpha in the pathogenesis of Albright hereditary osteodystrophy. *Trends Endocr. Metab.* 10: 81-85, 1999.
- [63973] 2303.Walker, M. E.; Rubinstein, P.; Allen, F. H., Jr.: Biochemical genetics of MN. *Vox Sang.* 32: 111-120, 1977.
- [63974] 2304.Weitkamp, L. R.; Adams, M. S.; Rowley, P. T.: Linkage between the MN and Hb beta loci. *Hum. Hered.* 22: 566-572, 1972.
- [63975] 2305.Weitkamp, L. R.; Lovrien, E. W.; Olaisen, B.; Fenger, K.; Gedde-Dahl, T., Jr.; Sorensen, S. A.; Conneally, P. M.; Bias, W. B.; Ott, J.: Linkage relations of the loci for the MN blood group and red cell phosphate. *Birth Defects Orig. Art. Ser.* XI(3): 276-280, 1975. Note: Alternate: *Cytogenet. Cell Genet.* 14: 446-450, 1975.....
- [63976] 2306.Neer, E. J.; Schmidt, C. J.; Nambudripad, R.; Smith, T. F.: The ancient regulatory-protein family of WD-repeat proteins. *Nature* 371:297-300, 1994.
- [63977] 2307.Byrd, P. J.; McConville, C. M.; Cooper, P.; Parkhill, J.; Stankovic, T.; McGuire, G. M.; Thick, J. A.; Taylor, A. M. R.: Mutations revealed by sequencing the 5-prime half of the gene for ataxia telangiectasia. *Hum. Molec. Genet.* 5: 145-149, 1996.
- [63978] 2308.Chong, M. J.; Murray, M. R.; Gosink, E. C.; Russell, H.

R. C.;Srinivasan, A.; Kapsetaki, M.; Korsmeyer, S. J.; McK-innon, P. J.:Atm and Bax cooperate in ionizing radiation-induced apoptosis in thecentral nervous system. Proc. Nat. Acad. Sci. 97: 889–894, 2000.

[63979] 2309.Falck, J.; Mailand, N.; Syljuasen, R. G.; Bartek, J.; Lukas, J.: The ATM–Chk2–Cdc25A checkpoint pathway guards against radioresistantDNA synthesis. Nature 410: 842–847, 2001.

[63980] 2310.Falck, J.; Petrini, J. H. J.; Williams, B. R.; Lukas, J.; Bartek,J.: The DNA damage–dependent intra–S phase checkpoint is regulatedby parallel pathways. Nature Genet. 30: 290–294, 2002.

[63981] 2311.Kanters, J. K.; Larsen, L. A.; Orholm, M.; Agner, E.; Anderson,P. S.; Vuust, J.; Christiansen, M.: Novel donor splice site mutationin the KVLQT1 gene is associated with the long QT syndrome. J. Cardiovasc.Electrophysiol. 9: 620–624, 1998.

[63982] 2312.Itoh, T.; Kikuchi, K.; Odagawa, Y.; Takata, S.; Yano, K.; Okada,S.; Haneda, N.; Ogawa, S.; Nakano, O.; Kawahara, Y.; Kasai, H.; Nakayama,T.; Fukutomi, T.; Sakurada, H.; Shimizu, A.; Yazaki, Y.; Nagai, R.;Nakamura, Y.; Tanaka, T.: Correlation of genetic etiology with respon-seto beta-adrenergic blockade among symptomatic pa-

tients with familial long-QT syndrome. J. Hum. Genet. 46: 38-40, 2001.

[63983] 2313. Keating, M.: Response to Benhorin et al. (1993). Science 260:1962 only, 1993.

[63984] 2314. Keating, M.; Atkinson, D.; Dunn, C.; Timothy, K.; Vincent, G.M.; Leppert, M.: Linkage of a cardiac arrhythmia, the long QT syndrome, and the Harvey RAS-1 gene. Science 252: 704-706, 1991.

[63985] 2315. Keating, M.; Atkinson, D.; Dunn, C.; Timothy, K.; Vincent, G.M.; Leppert, M.: Long QT syndrome is closely linked to the Harvey RAS-1 gene on chromosome 11. (Abstract) Clin. Res. 39: 317A only, 1991.

[63986] 2316. Keating, M.; Dunn, C.; Atkinson, D.; Timothy, K.; Vincent, G.M.; Leppert, M.: Consistent linkage of the long-QT syndrome to the Harvey Ras-1 locus on chromosome 11. Am. J. Hum. Genet. 49: 1335-1339, 1991.

[63987] 2317. Kerem, B.; Benhorin, J.; Kalman, Y. M.; Medina, A.; Dyer, T. D.; Blangero, J.; MacCluer, J. W.: Evidence for genetic heterogeneity in the long QT syndrome. (Abstract) Am. J. Hum. Genet. 51 (suppl.):A192 only, 1992.

[63988] 2318. Klein, H. O.; Levi, A.; Kaplinsky, E.; Di Segni, E.; David, D.: Congenital long-QT syndrome: deleterious effect of long-term high-rate ventricular pacing and defini-

tive treatment by cardiac transplantation. *Am.Heart J.* 132: 1079–1081, 1996.

[63989] 2319.Ko, Y.–L.; Chen, S.–A.; Tang, T. K.; Lin, J.–L.; Chiang, C.–E.;Chen, J.–J.; Teng, M.–S.; Chang, M.–S.; Lien, W.–P.; Wu, C.–W.: Noevidence for linkage of long QT syndrome and chromosome 11p15.5 markersin a Chinese family: evidence for genetic heterogeneity. *Hum. Genet.* 94:364–366, 1994.

[63990] 2320.Kukolich, M. K.; Telsey, A.; Ott, J.; Motulsky, A. G.: Suddeninfant death syndrome: normal QT intervals on ECGs of relatives. *Pediatrics* 60:51–54, 1977.

[63991] 2321.Larsen, L. A.; Fosdal, I.; Andersen, P. S.; Kanter, J. K.; Vuust,J.; Wettrell, G.; Christiansen, M.: Recessive Romano–Ward syndromeassociated with compound heterozygosity for two mutations in the KVLQT1gene. *Europ. J. Hum. Genet.* 7: 724–728, 1999.

[63992] 2322.Lee, M. P.; et al.; et al.: Loss of imprinting of a paternallyexpressed transcript, with antisense orientation to KVLQT1, occursfrequently in Beckwith–Wiedemann syndrome and is independent of insulin–likegrowth factor II imprinting. *Proc. Nat. Acad. Sci.* 96: 5203–5208,1999.

[63993] 2323.Lee, M. P.; Hu, R.–J.; Johnson, L. A.; Feinberg, A. P.: HumanKVLQT1 gene shows tissue–specific imprinting and

encompasses Beckwith–Wiedemann syndrome chromosomal rearrangements. *Nature Genet.* 15: 181–185, 1997.

[63994] 2324. Li, H.; Chen, Q.; Moss, A. J.; Robinson, J.; Goytia, V.; Perry, J. C.; Vincent, G. M.; Priori, S. G.; Lehmann, M. H.; Denfield, S. W.; Duff, D.; Kaine, S.; Shimizu, W.; Schwartz, P. J.; Wang, Q.; Towbin, J. A.: New mutations in the KVLQT1 potassium channel that cause long QT syndrome. *Circulation* 97: 1264–1269, 1998.

[63995] 2325. Locati, E. H.; Zareba, W.; Moss, A. J.; Schwartz, P. J.; Vincent, G. M.; Lehmann, M. H.; Towbin, J. A.; Priori, S. G.; Napolitano, C.; Robinson, J. L.; Andrews, M.; Timothy, K.; Hall, W. J.: Age- and sex-related differences in clinical manifestations in patients with congenital long-QT syndrome. *Circulation* 97: 2237–2244, 1998.

[63996] 2326. Mannens, M.; Wilde, A.: KVLQT1, the rhythm of imprinting. *Nature Genet.* 15: 113–115, 1997.

[63997] 2327. Martini, B.: Personal Communication. Thiene, Italy 11/9/1998.

[63998] 2328. Melki, J.; Kaplan, J.; Lucet, V.; Halley, L.; Clemenceau, S.; Kaplan, C.; Baule, M. S.; Frezal, J.: Long QT (Romano–Ward) syndrome: further definition of the linkage to the HLA loci. (Abstract) *Cytogenet. Cell Genet.* 46: 661 only, 1987.



- [63999] 2329.Milne, J. R.; Ward, D. E.; Spurrell, R. A. J.; Camm, A. J.: The long QT syndrome: effects of drugs and left stellate ganglion block. *Am.Heart J.* 104: 194–198, 1982.
- [64000] 2330.Mitsutake, A.; Takeshita, A.; Kuroiwa, A.; Nakamura, M.: Usefulness of the Valsalva maneuver in management of the long QT syndrome. *Circulation* 63:1029–1035, 1981.
- [64001] 2331.Moss, A. J.; McDonald, J.: Unilateral cervico–thoracic sympathetic ganglionectomy for the treatment of long Q–T interval syndrome. *NewEng. J. Med.* 285: 903–904, 1971.
- [64002] 2332.Moss, A. J.; Schwartz, P. J.: Sudden death and the idiopathic long Q–T syndrome. (Editorial) *Am. J. Med.* 66: 6–7, 1979.
- [64003] 2333.Moss, A. J.; Schwartz, P. J.; Crampton, R. S.; Tzivoni, D.; Locati, E. H.; MacCluer, J.; Hall, W. J.; Weitkamp, L.; Vincent, G. M.; Garson, A., Jr.; Robinson, J. L.; Benhorin, J.; Choi, S.: The long QT syndrome: prospective longitudinal study of 328 families. *Circulation* 84:1136–1144, 1991.
- [64004] 2334.Murray, A.; Donger, C.; Fenske, C.; Spillman, I.; Richard, P.; Dong, Y. B.; Neyroud, N.; Chevalier, P.; Denjoy, I.; Carter, N.; Syrris, P.; Afzal, A. P.; Patton, M. A.; Guicheney, P.; Jeffery, S.: Splicing mutations in KCNQ1: a mutation hot spot at codon 344 that produces in frame transcripts. *Circulation* 100: 1077–1084, 1999.

- [64005] 2335.Neyroud, N.; Denjoy, I.; Donger, C.; Gary, F.; Villain, E.; Leenhardt, A.; Benali, K.; Schwartz, K.; Coumel, P.; Guicheney, P.: Heterozygous mutation in the pore of potassium channel gene KvLQT1 causes an apparently-normal phenotype in long QT syndrome. *Europ. J. Hum. Genet.* 6: 129–133, 1998.
- [64006] 2336.Neyroud, N.; Richard, P.; Vignier, N.; Donger, C.; Denjoy, I.; Demay, L.; Shkolnikova, M.; Pesce, R.; Chevalier, P.; Hainque, B.; Coumel, P.; Schwartz, K.; Guicheney, P.: Genomic organization of the KCNQ1 K<sup>+</sup> channel gene and identification of C-terminal mutations in the long-QT syndrome. *Circ. Res.* 84: 290–297, 1999.
- [64007] 2337.Heald, R.; McLoughlin, M.; McKeon, F.: Human wee1 maintains mitotic timing by protecting the nucleus from cytoplasmically activated cdc2 kinase. *Cell* 74: 463–474, 1993.
- [64008] 2338.Igarashi, M.; Nagata, A.; Jinno, S.; Suto, K.; Okayama, H.: Wee1(+)-like gene in human cells. *Nature* 353: 80–83, 1991.
- [64009] 2339.Annunen, P.; Helaakoski, T.; Myllyharju, J.; Veijola, J.; Pihlajaniemi, T.; Kivirikko, K. I.: Cloning of the human prolyl 4-hydroxylase alpha subunit isoform alpha(II) and characterization of the type II enzyme tetramer: the alpha(I)

and alpha(II) subunits do not form a mixedal-  
pha(I)alpha(II)beta2 tetramer. J. Biol. Chem. 272:  
17342–17348,1997.

- [64010] 2340.Friedman, L.; Higgin, J. J.; Moulder, G.; Barstead, R.; Raines,R. T.; Kimble, J.: Prolyl 4–hydroxylase is required for viabilityand morphogenesis in *Caenorhabditis elegans*. Proc. Nat. Acad. Sci. 97:4736–4741, 2000.
- [64011] 2341.Helaakoski, T.; Annunen, P.; Vuori, K.; MacNeil, I. A.; Pihlajaniemi,T.; Kivirikko, K. I.: Cloning, baculovirus expression, and characterizationof a second mouse prolyl 4–hydroxylase alpha–subunit isoform: formationof an al-pha(2)beta(2) tetramer with the protein disulfide–iso-merase/betasubunit. Proc. Nat. Acad. Sci. 92: 4427–4431, 1995.
- [64012] 2342.Helaakoski, T.; Vuori, K.; Myllyla, R.; Kivirikko, K. I.; Pihlajaniemi,T.: Molecular cloning of the alpha–subunit of human prolyl 4–hydroxylase:the complete cDNA–derived amino acid sequence and evidence for alternativesplicing of RNA transcripts. Proc. Nat. Acad. Sci. 86: 4392–4396,1989.
- [64013] 2343.Pajunen, L.; Jones, T. A.; Helaakoski, T.; Pihla-janiemi, T.; Solomon,E.; Sheer, D.; Kivirikko, K. I.: Assign-ment of the gene coding forthe alpha–subunit of prolyl

4-hydroxylase to human chromosome re-  
gion10q21.3-23.1. Am. J. Hum. Genet. 45: 829-834,  
1989.

[64014] 2344.Helaakoski, T.; Veijola, J.; Vuori, K.; Rehn, M.; Chow, L. T.;Taillon-Miller, P.; Kivirikko, K. I.; Pihlajaniemi, T.: Structureand expression of the human gene for the alpha subunit of prolyl 4-hydroxylase:the two alternatively spliced types of mRNA correspond to two homologousex-  
ons the sequences of which are expressed in a variety of tissues. J.Biol. Chem. 269: 27847-27854, 1994.

[64015] 2345.Zelinski, T.; Coghlan, G.; Myal, Y.; Shiu, R. P. C.; Philipps,S.; White, L.; Lewis, M.: Genetic linkage between the Kell bloodgroup system and prolactin-inducible pro-  
tein loci: provisional assignmentof KEL to chromosome 7. Ann. Hum. Genet. 55: 137-140, 1991.

[64016] 2346.Bhalla, U. S.; Ram, P. T.; Iyengar, R.: MAP kinase phosphataseas a locus of flexibility in a mitogen-activated protein kinase signalingnetwork. Science 297: 1018-1023, 2002.

[64017] 2347.Boulton, T. G.; Nye, S. H.; Robbins, D. J.; Ip, N. Y.; Radziejewska,E.; Morgenbesser, S. D.; DePinho, R. A.; Panayotatos, N.; Cobb, M.H.; Yancopoulos, G. D.: ERKs: a family of protein-serine/threoninekinases that are acti-

vated and tyrosine phosphorylated in response to insulin and NGF. *Cell* 65: 663–675, 1991.

[64018] 2348. Cobb, M. H.; Boulton, T. G.; Robbins, D. J.: Extracellular signal-regulated kinases: ERKs in progress. *Cell Regul.* 2: 965–978, 1991.

[64019] 2349. Di Cristo, G.; Berardi, N.; Cancedda, L.; Pizzorusso, T.; Putignano, E.; Ratto, G. M.; Maffei, L.: Requirement of ERK activation for visual cortical plasticity. *Science* 292: 2337–2340, 2001.

[64020] 2350. Forcet, C.; Stein, E.; Pays, L.; Corset, V.; Llambi, F.; Tessier-Lavigne, M.; Mehlen, P.: Netrin-1-mediated axon outgrowth requires deleted in colorectal cancer-dependent MAPK activation. *Nature* 417: 443–447, 2002.

[64021] 2351. Khokhlatchev, A. V.; Canagarajah, B.; Wilsbacher, J.; Robinson, M.; Atkinson, M.; Goldsmith, E.; Cobb, M. H.: Phosphorylation of the MAP kinase ERK2 promotes its homodimerization and nuclear translocation. *Cell* 93: 605–615, 1998.

[64022] 2352. Li, L.; Wisk, M.; Gonzalez, F. A.; Davis, R. J.: Genomic loci of human mitogen-activated protein kinases. *Oncogene* 9: 647–649, 1994.

[64023] 2353. Owaki, H.; Makar, R.; Boulton, T. G.; Cobb, M. H.; Geppert, T. D.: Extracellular signal-regulated kinases in T

cells: characterization of human ERK1 and ERK2 cDNAs.

Biochem. Biophys. Res. Commun. 182:1416–1422, 1992.

[64024] 2354.Saba–El–Leil, M. K.; Malo, D.; Meloche, S.: Chromosomal localization of the mouse genes encoding the ERK1 and ERK2 isoforms of MAP kinases. Mammalian Genome 8: 141–142, 1997.

[64025] 2355.Stefanovsky, V. Y.; Pelletier, G.; Hannan, R.; Gagnon–Kugler, T.; Rothblum, L. I.; Moss, T.: An immediate response of ribosomal transcription to growth factor stimulation in mammals is mediated by ERK phosphorylation of UBF. Molec. Cell 8: 1063–1073, 2001.

[64026] 2356.Thomas, G.: MAP kinase by any other name smells just as sweet. Cell 68:3–6, 1992.

[64027] 2357.Garcia, J. I.; Zalba, G.; Detera–Wadleigh, S. D.; de Miguel, C.: Isolation of a cDNA encoding the rat MAP–kinase homolog of human p63mapk. Mammalian Genome 7: 810–814, 1996.

[64028] 2358.Gonzalez, F. A.; Raden, D. L.; Rigby, M. R.; Davis, R. J.: Heterogeneous expression of four MAP kinase isoforms in human tissues. FEBS Lett. 304:170–178, 1992.

[64029] 2359.Zhu, A. X.; Zhao, Y.; Moller, D. E.; Flier, J. S.: Cloning and characterization of p97(MAPK), a novel human homolog of rat ERK–3. Molec. Cell. Biol. 14: 8202–8211,

1994.

- [64030] 2360.Alvaro, V.; Levy, L.; Dubray, C.; Roche, A.; Peillon, F.; Querat,B.; Joubert, D.: Invasive human pituitary tumors express a point-mutatedalpha-protein kinase-C. J. Clin. Endocr. 77: 1125-1129, 1993.
- [64031] 2361.Coussens, L.; Parker, P. J.; Rhee, L.; Yang-Feng, T. L.; Chen,E.; Waterfield, M. D.; Francke, U.; Ullrich, A.: Multiple, distinctforms of bovine and human protein kinase C suggest diversity in cellularsignaling pathways. Science 233: 859-866, 1986.
- [64032] 2362.Finkenzeller, G.; Marme, D.; Hug, H.: Sequence of human proteinkinase C alpha. Nucleic Acids Res. 18: 2183 only, 1990.
- [64033] 2363.Jones, K. W.; Shapero, M. H.; Chevrette, M.; Fournier, R. E. K.: Subtractive hybridization cloning of a tissue-specific extinguisher:TSE1 encodes a regulatory subunit of protein kinase A. Cell 66:861-872, 1991.
- [64034] 2364.Latos-Bielenska, A.; Klett, C.; Just, W.; Hameister, H.: Refinementof localization of the human genes for myeloperoxidase (MPO), proteinkinase C, alpha polypeptide, PRKCA, and the DNA fragment D17S21 on chromosome 17q. Hereditas 115: 69-72, 1991.
- [64035] 2365.Linnenbach, A. J.; Huebner, K.; Reddy, E. P.; Herlyn,

M.; Parmiter, A. H.; Nowell, P. C.; Koprowski, H.: Structural alteration in the MYB protooncogene and deletion within the gene encoding alpha-type protein kinase C in human melanoma cell lines. *Proc. Nat. Acad. Sci.* 85:74–78, 1988.

[64036] 2366. Parker, P. J.; Coussens, L.; Totty, N.; Rhee, L.; Young, S.; Chen, E.; Stabel, S.; Waterfield, M. D.; Ullrich, A.: The complete primary structure of protein kinase C--the major phorbol ester receptor. *Science* 233:853–859, 1986.

[64037] 2367. Summar, M. L.; Phillips, J. A., III; Krishnamani, M. R. S.; Keefer, J.; Trofatter, J.; Schwartz, R. C.; Tsipouras, P.; Willard, H.; Ullrich, A.: Protein kinase C: a new linkage marker for growth hormone and for COL1A1. *Genomics* 5: 163–165, 1989.

[64038] 2368. Bayes, M.; Valverde, D.; Balcells, S.; Grinberg, D.; Vilageliu, L.; Benitez, J.; Ayuso, C.; Beneyto, M.; Baiget, M.; Gonzalez-Duarte, R.: Evidence against involvement of recoverin in autosomal recessive retinitis pigmentosa in 42 Spanish families. *Hum. Genet.* 96: 89–94, 1995.

[64039] 2369. Dizhoor, A. M.; Ray, S.; Kumar, S.; Niemi, G.; Spencer, M.; Brolley, D.; Walsh, K. A.; Philipov, P. P.; Hurley, J. B.; Stryer, L.: Recoverin: a calcium sensitive activator of retinal rod guanylate cyclase. *Science* 251:915–918, 1991.

[64040] 2370. Premkumar, L. S.; Ahern, G. P.: Induction of vanilloid



receptorchannel activity by protein kinase C. *Nature* 408: 985–990, 2000.

- [64041] 2371. Al-Maghteh, M.; Vithana, E. N.; Inglehearn, C. F.; Moore, T.; Bird, A. C.; Bhattacharya, S. S.: Segregation of a PRKCG mutation in two RP11 families. (Letter) *Am. J. Hum. Genet.* 62: 1248–1252, 1998.
- [64042] 2372. Dryja, T. P.; McEvoy, J.; McGee, T. L.; Berson, E. L.: No mutations in the coding region of the PRKCG gene in three families with retinitis pigmentosa linked to the RP11 locus on chromosome 19q. (Letter) *Am. J. Hum. Genet.* 65: 926–928, 1999.
- [64043] 2373. Johnson, K. J.; Jones, P. J.; Spurr, N.; Nimmo, E.; Davies, J.; Creed, H.; Weiss, M.; Williamson, R.: Linkage relationships of the protein kinase C gamma gene which exclude it as a candidate for myotonic dystrophy. *Cytogenet. Cell Genet.* 48: 13–15, 1988.
- [64044] 2374. McGee, T. L.; Devoto, M.; Ott, J.; Berson, E. L.; Dryja, T. P.: Evidence that the penetrance of mutations at the RP11 locus causing dominant retinitis pigmentosa is influenced by a gene linked to the homologous RP11 allele. *Am. J. Hum. Genet.* 61: 1059–1066, 1997.
- [64045] 2375. Saunders, A. M.; Seldin, M. F.: The syntenic relationship of proximal mouse chromosome 7 and the myotonic

dystrophy gene region on human chromosome 19q. Genomics 6: 324–332, 1990.

[64046] 2376. Vithana, E. N.; Abu-Safieh, L.; Allen, M. J.; Carey, A.; Papaioannou, M.; Chakarova, C.; Al-Maghteh, M.; Ebenezer, N. D.; Willis, C.; Moore, A. T.; Bird, A. C.; Hunt, D. M.; Bhattacharya, S. S.: A human homolog of yeast pre-mRNA splicing gene, PRP31, underlies autosomal dominant retinitis pigmentosa on chromosome 19q13.4 (RP11). Molec. Cell 8:375–381, 2001.

[64047] 2377. Gong, J.; Xu, J.; Bezanilla, M.; van Huizen, R.; Derin, R.; Li, M.: Differential stimulation of PKC phosphorylation of potassium channels by ZIP1 and ZIP2. Science 285: 1565–1569, 1999.

[64048] 2378. Hayashi, N.; Yokoyama, N.; Seki, T.; Azuma, Y.; Ohba, T.; Nishimoto, T.: RanBP1, a Ras-like nuclear G protein binding to Ran/TC4, inhibits RCC1 via Ran/TC4. Molec. Gen. Genet. 247: 661–669, 1995.

[64049] 2379. Spritz, R. A.; Strunk, K.; Surowy, C. S.; Mohrenweiser, H. W.: Human U1–70K ribonucleoprotein antigen gene: organization, nucleotide sequence, and mapping to locus 19q13.3. Genomics 8: 371–379, 1990.

[64050] 2380. Epstein, A. C. R.; Gleadle, J. M.; McNeill, L. A.; Hewitson, K. S.; O'Rourke, J.; Mole, D. R.; Mukherji, M.; Metzen,

E.; Wilson, M.I.; Dhanda, A.; Tian, Y.-M.; Masson, N.; Hamilton, D. L.; Jaakkola,P.; Barstead, R.; Hodgkin, J.; Maxwell, P. H.; Pugh, C. W.; Schofield,C. J.; Ratcliffe, P. J.: C. elegans EGL-9 and mammalian homologsdefine a family of dioxygenases that regulate HIF by prolyl hydroxylation. Cell107: 43–54, 2001.

[64051] 2381.Gatei, M.; Young, D.; Cerosaletti, K. M.; Desai-Mehta, A.; Spring,K.; Kozlov, S.; Lavin, M. F.; Gatti, R. A.; Concannon, P.; Khanna,K.: ATM-dependent phosphorylation of nibrin in response to radiationexposure. Nature Genet. 25: 115–119, 2000.

[64052] 2382.Kai, R.; Ohtsubo, M.; Sekiguchi, M.; Nishimoto, T.: Molecularcloning of a human gene that regulates chromosome condensation andis essential for cell proliferation. Molec. Cell. Biol. 6: 2027–2032,1986.

[64053] 2383.Nishimoto, T.; Seino, H.; Seki, N.; Hori, T.-A.: The human CHC1gene encoding RCC1 (regulator of chromosome condensation) (CHC1) islocalized to human chromosome 1p36.1. Genomics 23: 719–721, 1994.

[64054] 2384.Ohba, T.; Nakamura, M.; Nishitani, H.; Nishimoto, T.: Self-organizationof microtubule asters induced in Xenopus egg extracts by GTP-boundRan. Science 284: 1356–1358, 1999.

- [64055] 2385.Ohtsubo, M.; Kai, R.; Furuno, N.; Sekiguchi, T.; Sekiguchi, M.; Hayashida, H.; Kuma, K.; Miyata, T.; Fukushige, S.; Murotsu, T.; Matsubara, K.; Nishimoto, T.: Isolation and characterization of the active cDNA of the human cell cycle gene (RCC1) involved in the regulation of onset of chromosome condensation. *Genes Dev.* 1: 585–593, 1987.
- [64056] 2386.Naggert, J. K.; Mu, J.–L.: The mouse very low density lipoprotein receptor (Vldlr) gene maps to chromosome 19. *Mammalian Genome* 5:453–455, 1994.
- [64057] 2387.Meloni, A. M.; Dobbs, R. M.; Pontes, J. E.; Sandberg, A. A.: Translocation(X;1) in papillary renal cell carcinoma: a new cytogenetic subtype. *Cancer Genet. Cytogenet.* 65: 1–6, 1993.
- [64058] 2388.Zbar, B.; Tory, K.; Merino, M.; Schmidt, L.; Glenn, G.; Choyke, P.; Walther, M. M.; Lerman, M.; Linehan, W. M.: Hereditary papillary renal cell carcinoma. *J. Urol.* 151: 561–566, 1994.
- [64059] 2389.Adachi, H.; Tawaragi, Y.; Inuzuka, C.; Kubota, I.; Tsujimoto, M.; Nishihara, T.; Nakazato, H.: Primary structure of human microsomal dipeptidase deduced from molecular cloning. *J. Biol. Chem.* 265:3992–3995, 1990.
- [64060] 2390.Austruy, E.; Jeanpierre, C.; Antignac, C.; Whitmore,

S. A.; VanCong, N.; Bernheim, A.; Callen, D. F.; Junien, C.: Physical and genetic mapping of the dipeptidase gene DPEP1 to 16q24.3. *Genomics* 15: 684–687, 1993.

[64061] 2391. Campbell, B. J.; Forrester, L. J.; Zahler, W. L.; Burks, M.: Beta-lactamase activity of purified and partially characterized human renal dipeptidase. *J. Biol. Chem.* 259: 14586–14590, 1984.

[64062] 2392. Kozak, E. M.; Tate, S. S.: Glutathione-degrading enzymes of microvillus membranes. *J. Biol. Chem.* 257: 6322–6327, 1982.

[64063] 2393. Nakagawa, H.; Inazawa, J.; Inoue, K.; Misawa, S.; Kashima, K.; Adachi, H.; Nakazato, H.; Abe, T.: Assignment of the human renal dipeptidase gene (DPEP1) to band q24 of chromosome 16. (Abstract) *Cytogenet. Cell Genet.* 58: 2002 only, 1991.

[64064] 2394. Adachi, T.; Ohta, H.; Yamada, H.; Futenma, A.; Kato, K.; Hirano, K.: Quantitative analysis of extracellular-superoxide dismutase in serum and urine by ELISA with monoclonal antibody. *Clin. Chim. Acta* 212: 89–102, 1992.

[64065] 2395. Folz, R. J.; Crapo, J. D.: Extracellular superoxide dismutase (SOD3): tissue-specific expression, genomic characterization, and computer-assisted sequence analysis of the human EC SOD gene. *Genomics* 22: 162–171, 1994.

- [64066] 2396.Hendrickson, D. J.; Fisher, J. H.; Jones, C.; Ho, Y.-S.: Regional localization of human extracellular superoxide dismutase gene to 4pter-q21.Genomics 8: 736-738, 1990.
- [64067] 2397.Hjalmarsson, K.; Marklund, S. L.; Engstrom, A.; Edlund, T.: Isolation and sequence of complementary DNA encoding human extracellular superoxide dismutase. Proc. Nat. Acad. Sci. 84: 6340-6344, 1987.
- [64068] 2398.Marklund, S. L.: Extracellular superoxide dismutase in human tissues and human cell lines. J. Clin. Invest. 74: 1398-1403, 1984.
- [64069] 2399.Sandstrom, J.; Nilsson, P.; Karlsson, K.; Marklund, S. L.: 10-fold increase in human plasma extracellular superoxide dismutase content caused by a mutation in heparin-binding domain. J. Biol. Chem. 269:19163-19166, 1994.
- [64070] 2400.Yamada, H.; Yamada, Y.; Adachi, T.; Goto, H.; Ogasawara, N.; Futenma, A.; Kitano, M.; Hirano, K.; Kato, K.: Molecular analysis of extracellular-superoxide dismutase gene associated with high level in serum. Jpn. J. Hum. Genet. 40: 177-184, 1995.
- [64071] 2401.McNew, J. A.; Parlati, F.; Fukuda, R.; Johnston, R. J.; Paz, K.; Paumet, F.; Sollner, T. H.; Rothman, J. E.: Compartmental specificity of cellular membrane fusion encoded in

SNARE proteins. *Nature* 407:153–159, 2000.

[64072] 2402. Weber, T.; Zemelman, B. V.; McNew, J. A.; Westermann, B.; Gmachl, M.; Parlati, F.; Sollner, T. H.; Rothman, J. E.: SNAREpins: minimal machinery for membrane fusion. *Cell* 92: 759–772, 1998.

[64073] 2403. Rheaume, E.; Simard, J.; Morel, Y.; Mebarki, F.; Zachmann, M.; Forest, M. G.; New, M. I.; Labrie, F.: Congenital adrenal hyperplasia due to point mutations in the type II 3-beta-hydroxysteroid dehydrogenase gene. *Nature Genet.* 1: 239–245, 1992.

[64074] 2404. Amor, M.; Parker, K. L.; Globberman, H.; New, M. I.; White, P. C.: Mutation in the CYP21B gene (ile172-to-asn) causes steroid 21-hydroxylase deficiency. *Proc. Nat. Acad. Sci.* 85: 1600–1604, 1988.

[64075] 2405. Araujo, M.; Sanches, M. R.; Suzuki, L. A.; Guerra, G., Jr.; Farah, S. B.; De Mello, M. P.: Molecular analysis of CYP21 and C4 genes in Brazilian families with the classical form of steroid 21-hydroxylase deficiency. *Braz. J. Med. Biol. Res.* 29: 1–13, 1996.

[64076] 2406. Aston, C. E.; Sherman, S. L.; Morton, N. E.; Speiser, P. W.; New, M. I.: Genetic mapping of the 21-hydroxylase locus: estimation of small recombination frequencies. *Am. J. Hum. Genet.* 43: 304–310, 1988.

- [64077] 2407. Bachega, T. A. S. S.; Billerbeck, A. E. C.; Madureira, G.; Marcondes, J. A. M.; Longui, C. A.; Leite, M. V.; Arnhold, I. J. P.; Mendonca, B. B.: Molecular genotyping in Brazilian patients with the classical and nonclassical forms of 21-hydroxylase deficiency. *J. Clin. Endocr. Metab.* 83: 4416–4419, 1998.
- [64078] 2408. Aguiar-Oliveira, M. H.; Gill, M. S.; de A. Barretto, E. S.; Alcantara, M. R. S.; Miraki-Moud, F.; Menezes, C. A.; Souza, A. H. O.; Martinelli, C. E.; Pereira, F. A.; Salvatori, R.; Levine, M. A.; Shalet, S. M.; Camacho-Hubner, C.; Clayton, P. E.: Effect of severe growth hormone (GH) deficiency due to a mutation in the GH-releasing hormone receptor on insulin-like growth factors (IGFs), IGF-binding proteins, and ternary complex formation throughout life. *J. Clin. Endocr. Metab.* 84: 4118–4126, 1999.
- [64079] 2409. DeAlmeida, V. I.; Mayo, K. E.: Identification and binding domains of the growth hormone-releasing hormone receptor by analysis of mutant and chimeric receptor proteins. *Molec. Endocr.* 12: 750–765, 1998.
- [64080] 2410. Gaylinn, B. D.; Harrison, J. K.; Zysk, J. R.; Lyons, C. E.; Lynch, K. R.; Thorner, M. O.: Molecular cloning and expression of a human anterior pituitary receptor for growth hormone-releasing hormone. *Molec. Endocr.* 7: 77–84,



1993.

- [64081] 2411.Gaylinn, B. D.; von Kap-Herr, C.; Golden, W. L.; Thorner, M. O.: Assignment of the human growth hormone-releasing hormone receptor gene (GHRHR) to 7p14 by in situ hybridization. *Genomics* 19: 193–195,1994.
- [64082] 2412.Godfrey, P.; Rahal, J. O.; Beamer, W. G.; Copeland, N. G.; Jenkins, N. A.; Mayo, K. E.: GHRH receptor of little mice contains a missense mutation in the extracellular domain that disrupts receptor function. *Nature Genet.* 4: 227–232, 1993.
- [64083] 2413.Gondo, R. G.; Aguiar-Oliveira, M. H.; Hayashida, C. Y.; Toledo, S. P. A.; Abelin, N.; Levine, M. A.; Bowers, C. Y.; Souza, A. H. O.; Pereira, R. M. C.; Santos, N. L.; Salvatori, R.: Growth hormone-releasing peptide-2 stimulates GH secretion in GH-deficient patients with mutated GH-releasing hormone receptor. *J. Clin. Endocr. Metab.* 86: 3279–3283,2001.
- [64084] 2414.Lin, S.-C.; Lin, C. R.; Gukovsky, I.; Lysis, A. J.; Sawchenko, P. E.; Rosenfeld, M. G.: Molecular basis of the little mouse phenotype and implications for cell type-specific growth. *Nature* 364: 208–213,1993.
- [64085] 2415.Maheshwari, H. G.; Silverman, B. L.; Dupuis, J.; Baumann, G.: Phenotype and genetic analysis of a syndrome

caused by an inactivating mutation in the growth hormone-releasing hormone receptor: dwarfism of Sindh. J. Clin. Endocr. Metab. 83: 4065–4074, 1998.

[64086] 2416. Netchine, I.; Talon, P.; Dastot, F.; Vitaux, F.; Goossens, M.; Amselem, S.: Extensive phenotypic analysis of a family with growth hormone (GH) deficiency caused by a mutation in the GH-releasing hormone receptor gene. J. Clin. Endocr. Metab. 83: 432–436, 1998.

[64087] 2417. Petersenn, S.; Rasch, A. C.; Heyens, M.; Schulte, H. M.: Structure and regulation of the human growth hormone-releasing hormone receptor gene. Molec. Endocr. 12: 233–247, 1998.

[64088] 2418. Roelfsema, F.; Biermasz, N. R.; Veldman, R. G.; Veldhuis, J. D.; Frolich, M.; Stovkis-Brantsma, W. H.; Wit, J.-M.: Growth hormone (GH) secretion in patients with an inactivating defect of the GH-releasing hormone (GHRH) receptor is pulsatile: evidence for a role for non-GHRH inputs into the generation of GH pulses. J. Clin. Endocr. Metab. 86: 2459–2464, 2001.

[64089] 2419. Salvatori, R.; Fan, X.; Mullis, P. E.; Haile, A.; Levine, M. A.: Decreased expression of the GHRH receptor gene due to a mutation in a Pit-1 binding site. Molec. Endocr. 16: 450–458, 2002.

- [64090] 2420.Salvatori, R.; Fan, X.; Phillips, J. A., III; Espigares–Martin,R.; de Lara, I. M.; Freeman, K. L.; Plotnick, L.; Al–Ashwal, A.; Levine,M. A.: Three new mutations in the gene for the growth hormone (GH)–releasinghormone receptor in familial isolated GH deficiency type IB. J. Clin.Endocr. Metab. 86: 273–279, 2001.
- [64091] 2421.Salvatori, R.; Hayashida, C. Y.; Aguiar–Oliveira, M. H.; Phillips,J. A., III; Souza, A. H. O.; Gondo, R. G.; Toledo, S. P. A.; Conceicao,M. M.; Prince, M.; Maheshwari, H. G.; Baumann, G.; Levine, M. A.: Familial dwarfism due to a novel mutation of the growth hormone–releasinghormone receptor gene. J. Clin. Endocr. Metab. 84: 917–923, 1999.
- [64092] 2422.Vamvakopoulos, N. C.; Kunz, J.; Olberding, U.; Scherer, S. W.;Sioutopoulou, T. O.; Schneider, V.; Durkin, A. S.; Nierman, W. C.: Mapping the human growth hormone–releasing hormone receptor (GHRHR)gene to the short arm of chromosome 7 (7p13–p21) near the epidermalgrowth factor receptor (EGFR) gene. Genomics 20: 338–340, 1994.
- [64093] 2423.Wajnrajch, M. P.; Gertner, J. M.; Harbison, M. D.; Chua, S. C.,Jr.; Leibel, R. L.: Nonsense mutation in the human growth hormone–releasinghormone receptor causes growth failure analogous to the little (lit)mouse. Nature

Genet. 12: 88–90, 1996.

- [64094] 2424. Borsani, G.; Bassi, M. T.; Sperandio, M. P.; De Grandi, A.; Buoninconti, A.; Riboni, M.; Manzoni, M.; Incerti, B.; Pepe, A.; Andria, G.; Ballabio, A.; Sebastio, G.: SLC7A7, encoding a putative permease-related protein, is mutated in patients with lysinuric protein intolerance. *Nature Genet.* 21: 297–301, 1999.
- [64095] 2425. Ghizzoni, L.; Mastorakos, G.; Street, M. E.; Vottero, A.; Mazzardo, G.; Vanelli, M.; Chrousos, G. P.; Bernasconi, S.: Spontaneous thyrotropin and cortisol secretion interactions in patients with nonclassical 21-hydroxylase deficiency and control children. *J. Clin. Endocr. Metab.* 82: 3677–3683, 1997.
- [64096] 2426. Globerman, H.; Amor, H.; Parker, K. L.; New, M. I.; White, P. C.: A nonsense mutation causing steroid 21-hydroxylase deficiency. *J. Clin. Invest.* 82: 139–144, 1988.
- [64097] 2427. Gordon, M. T.; Conway, D. I.; Anderson, D. C.; Harris, R.: Genetics and biochemical variability of variants of 21 hydroxylase deficiency. *J. Med. Genet.* 22: 354–360, 1985.
- [64098] 2428. Gotoh, H.; Sagai, T.; Hata, J.-I.; Shiroishi, T.; Moriwaki, K.: Steroid 21-hydroxylase deficiency in mice. *En-*

ocrinology 123:1923–1927, 1988.

- [64099] 2429.Grosse–Wilde, H.; Weil, J.; Albert, E.; Scholz, S.; Bidlingmaier, F.; Sippel, W. G.; Knorr, D.: Genetic linkage studies between congenital adrenal hyperplasia and the HLA blood group system. *Immunogenetics* 8:41–49, 1979.
- [64100] 2430.Gueux, B.; Fiet, J.; Couillin, P.; Raux–Demay, M.–C.; Mornet, E.; Galons, H.; Villette, J.–M.; Boue, J.; Dreux, C.: Prenatal diagnosis of 21–hydroxylase deficiency congenital adrenal hyperplasia by simultaneous radioimmunoassay of 21–deoxycortisol and 17–hydroxyprogesterone in amniotic fluid. *J. Clin. Endocr. Metab.* 66: 534–537, 1988.
- [64101] 2431.Gutai, J. P.; Lee, P. A.; Johnsonbaugh, R. E.; Gareis, F.; Urban, M. D.; Migeon, C. J.: Detection of the heterozygous state in siblings of patients with congenital adrenal hyperplasia due to 21–hydroxylase deficiency. *J. Pediatr.* 94: 770–772, 1979.
- [64102] 2432.Haglund–Stengler, B.; Ritzen, E. M.; Gustafsson, J.; Luthman, H.: Haplotypes of the steroid 21–hydroxylase gene region encoding mild steroid 21–hydroxylase deficiency. *Proc. Nat. Acad. Sci.* 88:8352–8356, 1991.
- [64103] 2433.Harada, F.; Kimura, A.; Iwanaga, T.; Shimozawa, K.; Yata, J.; Sasazuki, T.: Gene conversion–like events cause steroid 21–hydroxylase deficiency in congenital adrenal

hyperplasia. *Proc. Nat. Acad. Sci.* 84:8091–8094, 1987.

[64104] 2434. Helmberg, A.; Tusie-Luna, M. T.; Tabarelli, M.; Kofler, R.; White, P. C.: R339H and P453S: CYP21 mutations associated with nonclassic steroid 21-hydroxylase deficiency that are not apparent gene conversions. *Molec. Endocr.* 6: 1318–1322, 1992.

[64105] 2435. Higashi, Y.; Tanae, A.; Inoue, H.; Fujii-Kuriyama, Y.: Evidence for frequent gene conversion in the steroid 21-hydroxylase P-450(C21) gene: implications for steroid 21-hydroxylase deficiency. *Am. J. Hum. Genet.* 42: 17–25, 1988.

[64106] 2436. Higashi, Y.; Tanae, A.; Inoue, H.; Hiromasa, T.; Fujii-Kuriyama, Y.: Aberrant splicing and missense mutations cause steroid 21-hydroxylase (P-450[C21]) deficiency in humans: possible gene conversion products. *Proc. Nat. Acad. Sci.* 85: 7486–7490, 1988.

[64107] 2437. Higashi, Y.; Yoshioka, H.; Yamane, M.; Gotoh, O.; Fujii-Kuriyama, Y.: Complete nucleotide sequence of two steroid 21-hydroxylase genes tandemly arranged in human chromosome: a pseudogene and a genuine gene. *Proc. Nat. Acad. Sci.* 83: 2841–2845, 1986.

[64108] 2438. Hirschfeld, A. J.; Fleshman, J. K.: An unusually high incidence of salt-losing congenital adrenal hyperplasia in

the Alaskan Eskimo. J.Pediat. 75: 492–494, 1969.

- [64109] 2439.Holler, W.; Scholz, S.; Knorr, D.; Bidlingmaier, F.; Keller, E.;Albert, E. D.: Genetic differences between the salt-wasting, simplevirilizing, and nonclassical types of congenital adrenal hyperplasia. J.Clin. Endocr. Metab. 60: 757–763, 1985.
- [64110] 2440.Hughes, I. A.; Dyas, J.; Riad-Fahmy, D.; Laurence, K. M.: Prenataldiagnosis of congenital adrenal hyperplasia: reliability of amnioticfluid steroid analysis. J. Med. Genet. 24: 344–347, 1987.
- [64111] 2441.Jaaskelainen, J.; Levo, A.; Voutilainen, R.; Partanen, J.: Population-widееvaluation of disease manifestation in relation to molecular genotypein steroid 21-hydroxylase (CYP21) deficiency: good correlation ina well defined population. J. Clin. Endocr. Metab. 82: 3293–3297,1997.
- [64112] 2442.Jaresch, S.; Kornely, E.; Kley, H.–K.; Schlaghecke, R.: Adrenalincidentaloma and patients with homozygous or heterozygous congenitaladrenal hyperplasia. J. Clin. Endocr. Metab. 74: 685–689, 1992.
- [64113] 2443.Jones, H. W., Jr.: Personal Communication. Baltimore, Md. 2/10/1978.
- [64114] 2444.Jospe, N.; Donohoue, P. A.; Van Dop, C.; McLean, R. H.; Bias,W. B.; Migeon, C. J.: Prevalence of polymorphic

21-hydroxylase gene(CA21HB) mutations in salt-losing congenital adrenal hyperplasia. Biochem.Biophys. Res. Commun. 142: 798-804, 1987.

- [64115] 2445.Kadair, R. G.; Block, M. B.; Katz, F. H.; Hofeldt, F. D.: 'Masked'21-hydroxylase deficiency of the adrenal presenting with gynecomastiaand bilateral testicular masses. Am. J. Med. 62: 278-282, 1977.
- [64116] 2446.Kawaguchi, H.; O'hUigin, C.; Klein, J.: Evolutionary origin ofmutations in the primate cytochrome P450c21 gene. Am. J. Hum. Genet. 50:766-780, 1992.
- [64117] 2447.Kirby-Keyser, L.; Porter, C. C.; Donohoue, P. A.: E380D: a novelpoint mutation of CYP21 in an HLA-homozygous patient with salt-losingcongenital adrenal hyperplasia due to 21-hydroxylase deficiency. Hum.Mutat. 9: 181-182, 1997.
- [64118] 2448.Kirkland, R. T.; Kirkland, J. L.; Keenan, B. S.; Bongiovanni,A. M.; Rosenberg, H. S.; Clayton, G. W.: Bilateral testicular tumorsin congenital adrenal hyperplasia. J. Clin. Endocr. Metab. 44: 369-378,1977.
- [64119] 2449.Klouda, P. T.; Harris, R.; Price, D. A.: Linkage and associationbetween HLA and 21-hydroxylase deficiency. J. Med. Genet. 17: 337-341,1980.
- [64120] 2450.Knochenhauer, E. S.; Cortet-Rudelli, C.; Cunning-



ham, R. D.; Conway-Myers, B. A.; Dewailly, D.; Azziz, R.: Carriers of 21-hydroxylase deficiency are not at increased risk for hyperandrogenism. *J. Clin. Endocr. Metab.* 82: 479–485, 1997.

[64121] 2451. Krone, N.; Braun, A.; Roscher, A. A.; Knorr, D.; Schwarz, H. P.: Predicting phenotype in steroid 21-hydroxylase deficiency? Comprehensive genotyping in 155 unrelated, well defined patients from Southern Germany. *J. Clin. Endocr. Metab.* 85: 1059–1065, 2000.

[64122] 2452. Itoh, H.; Tomita, M.; Uchino, H.; Kobayashi, T.; Kataoka, H.; Sekiya, R.; Nawa, Y.: cDNA cloning of rat pS2 peptide and expression of trefoil peptides in acetic acid-induced colitis. *Biochem. J.* 318: 939–944, 1996.

[64123] 2453. May, F. E. B.; Westley, B. R.: Close physical linkage of the genes encoding the pNR-2/pS2 protein and human spasmodic protein (hSP). *Hum. Genet.* 99: 303–307, 1997.

[64124] 2454. Seib, T.; Blin, N.; Hilgert, K.; Seifert, M.; Theisinger, B.; Engel, M.; Dooley, S.; Zang, K.-D.; Welter, C.: The three human trefoil genes TFF1, TFF2, and TFF3 are located within a region of 55 kb on chromosome 21q22.3. *Genomics* 40: 200–202, 1997.

[64125] 2455. Tomasetto, C.; Rio, M.-C.; Gautier, C.; Wolf, C.;

Hareuveni, M.;Chambon, P.; Lathe, R.: hSP, the domain-duplicated homolog of pS2protein, is co-expressed with pS2 in stomach but not in breast carcinoma. EMBOJ. 9: 407–414, 1990.

[64126] 2456.Brose, N.; Petrenko, A. G.; Sudhof, T. C.; Jahn, R.: Synaptotagmin:a calcium sensor on the synaptic vesicle surface. Science 256: 1021–1025,1992.

[64127] 2457.Fernandez–Chacon, R.; Konigstorfer, A.; Gerber, S. H.; Garcia,J.; Matos, M. F.; Stevens, C. F.; Brose, N.; Rizo, J.; Rosenmund,C.; Sudhof, T. C.: Synaptotagmin I functions as a calcium regulatorof release probability. Nature 410: 41–49, 2001.

[64128] 2458.Geppert, M.; Archer, B. T., III; Sudhof, T. C.: SynaptotagminII: a novel differentially distributed form of synaptotagmin. J.Biol. Chem. 266: 13548–13552, 1991.

[64129] 2459.Hilbush, B. S.; Morgan, J. I.: A third synaptotagmin gene, Syt3,in the mouse. Proc. Nat. Acad. Sci. 91: 8195–8199, 1994.

[64130] 2460.Jones, J. M.; Popma, S. J.; Mizuta, M.; Seino, S.; Meisler, M.H.: Synaptotagmin genes on mouse chromosomes 1, 7, and 10 and humanchromosome 19. Mammalian Genome 6: 212–213, 1995.

[64131] 2461.Kwon, O.–J.; Adamson, M. C.; Chin, H.; Kozak, C. A.:

Genetic mapping of five mouse genes encoding synaptotagmins. *Mammalian Genome* 6:880–881, 1995.

- [64132] 2462. Mackler, J. M.; Drummond, J. A.; Loewen, C. A.; Robinson, I. M.; Reist, N. E.: The C2B  $\text{Ca}^{2+}$ -binding motif of synaptotagmin is required for synaptic transmission in vivo. *Nature* 418: 340–344, 2002.
- [64133] 2463. Perin, M. S.; Fried, V. A.; Mignery, G. A.; Jahn, R.; Sudhof, T. C.: Phospholipid binding by a synaptic vesicle protein homologous to the regulatory region of protein kinase C. *Nature* 345: 260–263, 1990.
- [64134] 2464. Perin, M. S.; Johnston, P. A.; Ozcelik, T.; Jahn, R.; Francke, U.; Sudhof, T. C.: Structural and functional conservation of synaptotagmin(p65) in *Drosophila* and humans. *J. Biol. Chem.* 266: 615–622, 1991.
- [64135] 2465. Robinson, I. M.; Ranjan, R.; Schwarz, T. L.: Synaptotagmins I and IV promote transmitter release independently of  $\text{Ca}^{2+}$  binding in the C2A domain. *Nature* 418: 336–340, 2002.
- [64136] 2466. Shin, O.-H.; Rizo, J.; Sudhof, T. C.: Synaptotagmin function in dense core vesicle exocytosis studied in cracked PC12 cells. *Nature Neurosci.* 5: 649–656, 2002.
- [64137] 2467. Wang, C.-T.; Grishanin, R.; Earles, C. A.; Chang, P. Y.; Martin, T. F. J.; Chapman, E. R.; Jackson, M. B.: Synapto-

tagmin modulation of fusion pore kinetics in regulated exocytosis of dense-core vesicles. *Science* 294:1111–1115, 2001.

- [64138] 2468. Begley, C. G.; Aplan, P. D.; Davey, M. P.; Nakahara, K.; Tchorz, K.; Kurtzberg, J.; Hershfield, M. S.; Haynes, B. F.; Cohen, D. I.; Waldmann, T. A.; Kirsch, I. R.: Chromosomal translocation in a human leukemic stem-cell line disrupts the T-cell antigen receptor delta-chain diversity region and results in a previously unreported fusion transcript. *Proc. Nat. Acad. Sci.* 86: 2031–2035, 1989.
- [64139] 2469. Kikuchi, M.; Tayama, T.; Hayakawa, H.; Takahashi, I.; Hoshino, H.; Ohsaka, A.: Familial thrombocytosis. *Brit. J. Haemat.* 89: 900–902, 1995.
- [64140] 2470. Wiestner, A.; Schlemper, R. J.; van der Maas, A. P. C.; Skoda, R. C.: An activating splice donor mutation in the thrombopoietin gene causes hereditary thrombocythaemia. *Nature Genet.* 18: 49–52, 1998.
- [64141] 2471. Hruban, R. H.; Goggins, M.; Parsons, J.; Kern, S. E.: *Clin. Cancer Res.* 6: 2969–2972, 2000.
- [64142] 2472. Hruban, R. H.; van Mansfield, A. D. M.; Offerhaus, G. J. A.; van Weering, D. H. J.; Allison, D. C.; Goodman, S. N., Kensler, T. W.; Bose, K. K.; Cameron, J. L.; Bos, J. L.: *Am. J. Path.* 143: 545–554, 1993.

- [64143] 2473.Hruban, R. H.; Wilentz, R. E.; Kern, S. E. :Am. J. Pathol. 156:1821–1825, 2000.
- [64144] 2474.Ruivenkamp, C. A. L.; van Wezel, T.; Zanon, C.; Stassen, A. P.M.; Vlcek, C.; Csikos, T.; Klous, A. M.; Tripodis, N.; Perrakis, A.;Boerrigter, L.; Groot, P. C.; Lindeman, J.; Mooi, W. J.; Meijjer,G. A.; Scholten, G.; Dauwerse, H.; Paces, V.; van Zandwijk, N.; vanOmmen, G. J. B.; Demant, P.: Ptp<sup>prj</sup> is a candidate for the mouse colon-cancersusceptibility locus Scc1 and is frequently deleted in human cancers. NatureGenet. 31: 295–300, 2002.
- [64145] 2475.Watanabe, T.; Mukouyama, Y.; Rhodes, M.; Thomas, M.; Kume, T.;Oishi, M.: Chromosomal location of murine protein tyrosine phosphatase(Ptp<sup>prj</sup> and Ptp<sup>pre</sup>) genes. Genomics 29: 793–795, 1995.
- [64146] 2476.Elson, A.; Kozak, C. A.; Morton, C. C.; Weremowicz, S.; Leder,P.: The protein tyrosine phosphatase epsilon gene maps to mouse chromosome7 and human chromosome 10q26. Genomics 31: 373–375, 1996.
- [64147] 2477.Krueger, N. X.; Streuli, M.; Saito, H.: Structural diversity andevolution of human receptor-like protein tyrosine phosphatases. EMBOJ. 9: 3241–3252, 1990.
- [64148] 2478.Melhado, I. G.; Anderson, L. L.; Duncan, A. M. V.; Jirik, F. R.: The gene encoding protein-tyrosine phos-

phatase RPTP-epsilon (PTPRE) is assigned to human chromosome 10q26. *Cytogenet. Cell Genet.* 73:168-170, 1996.

- [64149] 2479. Banerjee, P.; Kleyn, P. W.; Knowles, J. A.; Lewis, C. A.; Ross, B. M.; Parano, E.; Kovats, S. G.; Lee, J. J.; Penchaszadeh, G. K.; Ott, J.; Jacobson, S. G.; Gilliam, T. C.: TULP1 mutation in two extended Dominican kindreds with autosomal recessive retinitis pigmentosa. *Nature Genet.* 18: 177-179, 1998.
- [64150] 2480. Hagstrom, S. A.; North, M. A.; Nishina, P. M.; Berson, E. L.; Dryja, T. P.: Recessive mutations in the gene encoding the tubby-like protein TULP1 in patients with retinitis pigmentosa. *Nature Genet.* 18: 174-176, 1998.
- [64151] 2481. Knowles, J. A.; Shugart, Y.; Banerjee, P.; Gilliam, T. C.; Lewis, C. A.; Jacobson, S. G.; Ott, J.: Identification of a locus, distinct from RDS-peripherin, for autosomal recessive retinitis pigmentosa on chromosome 6p. *Hum. Molec. Genet.* 3: 1401-1403, 1994.
- [64152] 2482. Iivanainen, A.; Sainio, K.; Sariola, H.; Tryggvason, K.: Primary structure and expression of a novel human laminin alpha-4 chain. *FEBS Lett.* 365: 183-188, 1995.
- [64153] 2483. Richards, A. J.; Al-Imara, L.; Carter, N. P.; Lloyd, J. C.; Leversha, M. A.; Pope, F. M.: Localization of the gene

(LAMA4) to chromosome6q21 and isolation of a partial cDNA encoding a variant laminin A chain. Genomics 22: 237–239, 1994.

- [64154] 2484. Richards, A. J.; Al-Imara, L.; Carter, N. P.; Lloyd, J. C.; Pope, F. M.: A laminin A variant gene (LAMA3) is present on chromosome6q21 (Abstract) J. Med. Genet. 31: 164, 1994.
- [64155] 2485. Dorow, D. S.; Devereux, L.; Dietzsch, E.; De Kretser, T.: Identification of a new family of human epithelial protein kinases containing two leucine/isoleucine-zipper domains. Europ. J. Biochem. 213: 701–710, 1993.
- [64156] 2486. Hanks, S. K.: Eukaryotic protein kinases. Curr. Opin. Struct. Biol. 1: 369–383, 1991.
- [64157] 2487. Balczon, R.; Bao, L.; Zimmer, W. E.: PCM-1, a 228-kD centrosome autoantigen with a distinct cell cycle distribution. J. Cell Biol. 124: 783–793, 1994.
- [64158] 2488. Graham, M.; Shutter, J. R.; Sarmiento, U.; Sarosi, I.; Stark, K. L.: Overexpression of Agt leads to obesity in transgenic mice. (Letter) Nature Genet. 17: 273–274, 1997.
- [64159] 2489. Kanetsky, P. A.; Swoyer, J.; Panossian, S.; Holmes, R.; Guerry, D.; Rebbeck, T. R.: A polymorphism in the agouti signaling protein gene is associated with human pigmen-

tation. *Am. J. Hum. Genet.* 70:770–775, 2002.

- [64160] 2490. Klebig, M. L.; Wilkinson, J. E.; Geisler, J. G.; Woychik, R. P.: Ectopic expression of the agouti gene in transgenic mice causes obesity, features of type II diabetes, and yellow fur. *Proc. Nat. Acad. Sci.* 92: 4728–4732, 1995.
- [64161] 2491. Kwon, H. Y.; Bultman, S. J.; Loffler, C.; Chen, W.-J.; Furdon, P. J.; Powell, J. G.; Usala, A.-L.; Wilkison, W.; Hansmann, I.; Woychik, R. P.: Molecular structure and chromosomal mapping of the human homolog of the agouti gene. *Proc. Nat. Acad. Sci.* 91: 9760–9764, 1994.
- [64162] 2492. Manne, J.; Argeson, A. C.; Siracusa, L. D.: Mechanisms for the pleiotropic effects of the agouti gene. *Proc. Nat. Acad. Sci.* 92: 4721–4724, 1995.
- [64163] 2493. Miltenberger, R. J.; Mynatt, R. L.; Bruce, B. D.; Wilkison, W. O.; Woychik, R. P.; Michaud, E. J.: An agouti mutation lacking the basic domain induces yellow pigmentation but not obesity in transgenic mice. *Proc. Nat. Acad. Sci.* 96: 8579–8584, 1999.
- [64164] 2494. Wilson, B. D.; Ollmann, M. M.; Kang, L.; Stoffel, M.; Bell, G. I.; Barsh, G. S.: Structure and function of ASP, the human homolog of the mouse agouti gene. *Hum. Molec. Genet.* 4: 223–230, 1995.
- [64165] 2495. Zemel, M. B.; Kim, J. H.; Woychik, R. P.; Michaud, E.



J.; Kadwell, S. H.; Patel, I. R.; Wilkison, W. O.: Agouti regulation of intracellular calcium: role in the insulin resistance of viable yellow mice. *Proc. Nat. Acad. Sci.* 92: 4733–4737, 1995.

[64166] 2496. Cardon, L. R.; Smith, S. D.; Fulker, D. W.; Kimberling, W. J.; Pennington, B. F.; DeFries, J. C.: Quantitative trait locus for reading disability on chromosome 6. *Science* 266: 276–279, 1994. Note: Erratum: *Science* 268: 1553 only, 1995.

[64167] 2497. Field, L. L.; Kaplan, B. J.: Absence of linkage of phonological coding dyslexia to chromosome 6p23–p21.3 in a large family data set. *Am. J. Hum. Genet.* 63: 1448–1456, 1998.

[64168] 2498. Fisher, S. E.; Francks, C.; Marlow, A. J.; MacPhie, I. L.; Newbury, D. F.; Cardon, L. R.; Ishikawa-Brush, Y.; Richardson, A. J.; Talcott, J. B.; Gayán, J.; Olson, R. K.; Pennington, B. F.; Smith, S. D.; DeFries, J. C.; Stein, J. F.; Monaco, A. P.: Independent genome-wide scans identify a chromosome 18 quantitative-trait locus influencing dyslexia. *Nature-Genet.* 30: 86–91, 2002.

[64169] 2499. Fisher, S. E.; Marlow, A. J.; Lamb, J.; Maestrini, E.; Williams, D. F.; Richardson, A. J.; Weeks, D. E.; Stein, J. F.; Monaco, A. P.: A quantitative-trait locus on chromosome

6p influences different aspects of developmental dyslexia.  
Am. J. Hum. Genet. 64: 146–156, 1999.

- [64170] 2500. Gayan, J.; Smith, S. D.; Cherny, S. S.; Cardon, L. R.; Fulker, D. W.; Brower, A. M.; Olson, R. K.; Pennington, B. F.; DeFries, J. C.: Quantitative-trait locus for specific language and reading deficit on chromosome 6p. Am. J. Hum. Genet. 64: 157–164, 1999.
- [64171] 2501. Kaplan, D. E.; Gayan, J.; Ahn, J.; Won, T.-W.; Pauls, D.; Olson, R. K.; DeFries, J. C.; Wood, F.; Pennington, B. F.; Page, G. P.; Smith, S. D.; Gruen, J. R.: Evidence for linkage and association with reading disability, on 6p21.3–22. Am. J. Hum. Genet. 70: 1287–1298, 2002.
- [64172] 2502. Smith, S. D.; Kimberling, W. J.; Pennington, B. F.: Screening for multiple genes influencing dyslexia. Reading Writing 3: 285–298, 1991.
- [64173] 2503. Fazioli, F.; Minichiello, L.; Matoska, V.; Castagnino, P.; Miki, T.; Wong, W. T.; Di Fiore, P. P.: Eps8, a substrate for the epidermal growth factor receptor kinase, enhances EGF-dependent mitogenic signals. EMBO J. 12: 3799–3808, 1993.
- [64174] 2504. Corvi, R.; Berger, N.; Balczon, R.; Romeo, G.: RET/PCM-1: a novel fusion gene in papillary thyroid carcinoma. Oncogene 19: 4236–4242, 2000.

[64175] 2505.Ohata, H.; Fujiwara, Y.; Koyama, K.; Nakamura, Y.: Mapping of the human autoantigen pericentriolar material 1 (PCM1) gene to chromosome 8p21.3–p22. *Genomics* 24: 404–406, 1994.

[64176] 2506.Andresen, B. S.; Christensen, E.; Corydon, T. J.; Bross, P.; Pilgaard, B.; Wanders, R. J. A.; Ruiter, J. P. N.; Simonsen, H.; Winter, V.; Knudsen, I.; Schroeder, L. D.; Gregersen, N.; Skovby, F.: Iso-branched 2-methylbutyrylglycinuria caused by short/branched-chain acyl-CoA dehydrogenase deficiency: identification of a new enzyme defect, resolution of its molecular basis, and evidence for distinct acyl-CoA dehydrogenases in isoleucine and valine metabolism. *Am. J. Hum. Genet.* 67: 1095–1103, 2000.

[64177] 2507.Takai, S.; Tanaka, M.; Sugimura, H.; Yamada, K.; Naito, Y.; Kino, I.; Matsuda, M.: Mapping of the human C3G gene coding a guanine nucleotide-releasing protein for Ras family to 9q34.3 by fluorescence in situ hybridization. *Hum. Genet.* 94: 549–550, 1994.

[64178] 2508.Tanaka, S.; Morishita, T.; Hashimoto, Y.; Hattori, S.; Nakamura, S.; Shibuya, M.; Matsuoka, K.; Takenawa, T.; Kurata, T.; Nagashima, K.; Matsuda, M.: C3G, a guanine nucleotide-releasing protein expressed ubiquitously, binds

to the Src homology 3 domains of CRK and GRB2/ASH proteins. *Proc. Nat. Acad. Sci.* 91: 3443–3447, 1994.

- [64179] 2509. Belich, M. P.; Glynn, R. J.; Senger, G.; Sheer, D.; Trowsdale, J.: Proteasome components with reciprocal expression to that of the MHC-encoded LMP proteins. *Curr. Biol.* 4: 769–776, 1994.
- [64180] 2510. McCusker, D.; Jones, T.; Sheer, D.; Trowsdale, J.: Genetic relationships of the genes encoding the human proteasome beta subunits and the proteasome PA28 complex. *Genomics* 45: 362–367, 1997.
- [64181] 2511. Kumar, S.; Tomooka, Y.; Noda, M.: Identification of a set of genes with developmentally down-regulated expression in the mouse brain. *Biochem. Biophys. Res. Commun.* 185: 1155–1161, 1992.
- [64182] 2512. Braybrooke, J. P.; Spink, K. G.; Thacker, J.; Hickson, I. D.: The RAD51 family member, RAD51L3, is a DNA-stimulated ATPase that forms a complex with XRCC2. *J. Biol. Chem.* 275: 29100–29106, 2000.
- [64183] 2513. Johnson, R. D.; Liu, N.; Jasin, M.: Mammalian XRCC2 promotes the repair of DNA double-strand breaks by homologous recombination. *Nature* 401: 397–399, 1999.
- [64184] 2514. Jones, N. J.; Cox, R.; Thacker, J.: Isolation and cross-

sensitivity of x-ray-sensitive mutants of V79-4 hamster cells. *Mutat. Res.* 183:279–286, 1987.

- [64185] 2515. Jones, N. J.; Zhao, Y.; Siciliano, M. J.; Thompson, L. H.: Assignment of the XRCC2 human DNA repair gene to chromosome 7q36 by complementation analysis. *Genomics* 26: 619–622, 1995.
- [64186] 2516. Kurumizaka, H.; Ikawa, S.; Nakada, M.; Enomoto, R.; Kagawa, W.; Kinebuchi, T.; Yamazoe, M.; Yokoyama, S.; Shibata, T.: Homologous pairing and ring and filament structure formation activities of the human Xrcc2–Rad51D complex. *J. Biol. Chem.* 277: 14315–14320, 2002.
- [64187] 2517. Liu, N.; Schild, D.; Thelen, M. P.; Thompson, L. H.: Involvement of Rad51C in two distinct protein complexes of Rad51 paralogs in human cells. *Nucleic Acids Res.* 30: 1009–1015, 2002.
- [64188] 2518. Masson, J.-Y.; Tarsounas, M. C.; Stasiak, A. Z.; Stasiak, A.; Shah, R.; McIlwraith, M. J.; Benson, F. E.; West, S. C.: Identification and purification of two distinct complexes containing the five RAD51 paralogs. *Genes Dev.* 15: 3296–3307, 2001.
- [64189] 2519. Tambini, C. E.; George, A. M.; Rommens, J. M.; Tsui, L.-C.; Scherer, S. W.; Thacker, J.: The XRCC2 DNA repair gene: identification of a positional candidate. *Genomics*

41: 84–92, 1997.

- [64190] 2520. Thacker, J.; Tambini, C. E.; Simpson, P. J.; Tsui, L.-C.; Scherer, S. W.: Localization to chromosome 7q36.1 of the human XRCC2 gene, determining sensitivity to DNA-damaging agents. *Hum. Molec. Genet.* 4:113–120, 1995.
- [64191] 2521. Kjeldsen, A. D.; Brusgaard, K.; Poulsen, L.; Kruse, T.; Rasmussen, K.; Green, A.: Mutations in the ALK-1 gene and the phenotype of hereditary hemorrhagic telangiectasia in two large Danish families. *Am. J. Med. Genet.* 98: 298–302, 2001.
- [64192] 2522. Nakagawara, A.; Liu, X.-G.; Ikegaki, N.; White, P. S.; Yamashiro, D. J.; Nycum, L. M.; Biegel, J. A.; Brodeur, G. M.: Cloning and chromosomal localization of the human TRK-B tyrosine kinase receptor gene (NTRK2). *Genomics* 25:538–546, 1995.
- [64193] 2523. Rico, B.; Xu, B.; Reichardt, L. F.: TrkB receptor signaling is required for establishment of GABAergic synapses in the cerebellum. *Nature Neurosci.* 5: 225–233, 2002.
- [64194] 2524. Slaugenhaupt, S. A.; Blumenfeld, A.; Liebert, C. B.; Mull, J.; Lucente, D. E.; Monahan, M.; Breakefield, X. O.; Maayan, C.; Parada, L.; Axelrod, F. B.; Gusella, J. F.: The human gene for neurotrophic tyrosine kinase receptor type

2 (NTRK2) is located on chromosome 9 but is not the familial dysautonomia gene. *Genomics* 25: 730–732, 1995.

[64195] 2525. Soppet, D.; Escandon, E.; Maragos, J.; Middlemas, D. S.; Reid, S. W.; Blair, J.; Burton, L. E.; Stanton, B. R.; Kaplan, D. R.; Hunter, T.; Nikolics, K.; Parada, L. F.: The neurotrophic factors brain-derived neurotrophic factor and neurotrophin-3 are ligands for the trkB tyrosine kinase receptor. *Cell* 65: 895–903, 1991.

[64196] 2526. Squinto, S. P.; Stitt, S. N.; Aldrich, T. H.; Davis, S.; Bianco, S. M.; Radziejewski, C.; Glass, D. J.; Masiakowski, P.; Furth, M. E.; Valenzuela, D. M.; DiStefano, P. S.; Yancopoulos, G. D.: trkB encodes a functional receptor for brain-derived neurotrophic factor and neurotrophin-3 but not nerve growth factor. *Cell* 65: 885–893, 1991.

[64197] 2527. Grun, F.; Hirose, Y.; Kawauchi, S.; Ogura, T.; Umesono, K.: Aldehyde dehydrogenase 6, a cytosolic retinaldehyde dehydrogenase prominently expressed in sensory neuroepithelia during development. *J. Biol. Chem.* 275: 41210–41218, 2000.

[64198] 2528. Hsu, L. C.; Chang, W.-C.; Hiraoka, L.; Hsieh, C.-L.: Molecular cloning, genomic organization, and chromosomal localization of an additional human aldehyde dehydrogenase gene, ALDH6. *Genomics* 24: 333–341, 1994.

- [64199] 2529.Kapfhamer, D.; Miller, D. E.; Lambert, S.; Bennett, V.; Glover,T. W.; Burmeister, M.: Chromosomal localization of the ankyrin-Ggene (ANK3/Ank3) to human 10q21 and mouse 10. Genomics 27: 189–191,1995.
- [64200] 2530.Bennett, M.; Reed, R.: Correspondence between a mammalian spliceosomecomponent and an essential yeast splicing factor. Science 262: 105–108,1993.
- [64201] 2531.Dresser, D. W.; Hacker, A.; Lovell-Badge, R.; Guerrier, D.: Thegenes for a spliceosome protein (SAP62) and the anti-Mullerian hormone(AMH) are contiguous. Hum. Molec. Genet. 4: 1613–1618, 1995.
- [64202] 2532.Burks, D. J.; Font de Mora, J.; Schubert, M.; Withers, D. J.; Myers,M. G.; Towery, H. H.; Altamuro, S. L.; Flint, C. L.; White, M. F.: IRS–2 pathways integrate female reproduction and energy homeostasis. Nature 407:377–382, 2000.
- [64203] 2533.Fritsche, A.; Madaus, A.; Renn, W.; Tschritter, O.; Teigeler, A.;Weisser, M.; Maerker, E.; Machicao, F.; Haring, H.; Stumvoll, M.:The prevalent Gly1057Asp polymorphism in the insulin receptor substrate–2gene is not associated with impaired insulin secretion. J. Clin.Endocr. Metab. 86: 4822–4825, 2001.
- [64204] 2534.Kubota, N.; Tobe, K.; Terauchi, Y.; Eto, K.; Yamauchi,



T.; Suzuki,R.; Tsubamoto, Y.; Komeda, K.; Nakano, R.; Miki, H.; Satoh, S.; Sekihara,H.; Sciacchitano, S.; Lesniak, M.; Aizawa, S.; Nagai, R.; Kimura,S.; Akanuma, Y.; Taylor, S. I.; Kadowaki, T.: Disruption of insulinreceptor substrate 2 causes type 2 diabetes because of liver insulinresistance and lack of compensatory beta-cell hyperplasia. Diabetes 49::1880–1889, 2000.

[64205] 2535.Sun, X. J.; Wang, L.–M.; Zhang, Y.; Yenush, L.; Myers, M. G., Jr.;Glasheen, E.; Lane, W. S.; Pierce, J. H.; White, M. F.: Role of IRS–2in insulin and cytokine signalling. Nature 377: 173–177, 1995.

[64206] 2536.Withers, D. J.; Burks, D. J.; Towery, H. H.; Altamuro, S. L.;Flint, C. L.; White, M. F.: Irs–2 coordinates Igf–1 receptor–mediatedbeta-cell development and peripheral insulin signalling. Nature Genet. 23:32–40, 1999.

[64207] 2537.Withers, D. J.; Gutierrez, J. S.; Towery, H.; Burks, D. J.; Ren,J.–M.; Previs, S.; Zhang, Y.; Bernal, D.; Pons, S.; Shulman, G. I.;Bonner–Weir, S.; White, M. F.: Disruption of IRS–2 causes type 2diabetes in mice. Nature 391: 900–902, 1998.

[64208] 2538.Eberle, F.; Dubreuil, P.; Mattei, M.–G.; Devilard, E.; Lopez, M.: The human PRR2 gene, related to the human poliovirus receptor gene(PVR), is the true homolog of the

murine Mph gene. *Gene* 159: 267–272,1995.

- [64209] 2539.Morrison, M. E.; Racaniello, V. R.: Molecular cloning and expression of a murine homolog of the human poliovirus receptor gene. *J. Virol.* 66:2807–2813, 1992.
- [64210] 2540.Wong, W. T.; Carlomagno, F.; Druck, T.; Barletta, C.; Croce, C.M.; Huebner, K.; Kraus, M. H.; Di Fiore, P. P.: Evolutionary conservation of the EPS8 gene and its mapping to human chromosome 12q23–q24. *Oncogene* 9:3057–3061, 1994.
- [64211] 2541.Avraham, K. B.; Levanon, D.; Negreanu, V.; Bernstein, Y.; Groner, Y.; Copeland, N. G.; Jenkins, N. A.: Mapping of the mouse homolog of the human runt domain gene, AML2, to the distal region of mouse chromosome 4. *Genomics* 25: 603–605, 1995.
- [64212] 2542.Bae, S.-C.; Takahashi, E.; Zhang, Y. W.; Ogawa, E.; Shigesada, K.; Namba, Y.; Satake, M.; Ito, Y.: Cloning, mapping and expression of PEBP2- $\alpha$ -C, a third gene encoding the mammalian Runt domain. *Gene* 159:245–248, 1995.
- [64213] 2543.Inoue, K.; Ozaki, S.; Shiga, T.; Ito, K.; Masuda, T.; Okado, N.; Iseda, T.; Kawaguchi, S.; Ogawa, M.; Bae, S.-C.; Yamashita, N.; Itohara, S.; Kudo, N.; Ito, Y.: Runx3 controls the axonal projection of proprioceptive dorsal root gan-

glion neurons. Nature Neurosci. 23Sept, 2002.

Note:Advance Electronic Publication.

[64214] 2544.Li, Q.-L.; Ito, K.; Sakakura, C.; Fukamachi, H.; Inoue, K.; Chi,X.-Z.; Lee, K.-Y.; Nomura, S.; Lee, C.-W.; Han, S.-B.; Kim, H.-M.;Kim, W.-J.; and 15 others: Causal relationship between the lossof RUNX3 expression and gastric cancer. Cell 109: 113–124, 2002.

[64215] 2545.Wijmenga, C.; Speck, N. A.; Dracopoli, N. C.; Hofker, M. H.; Liu,P.; Collins, F. S.: Identification of a new murine runt domain–containinggene, Cbfa3, and localization of the human homolog, CBFA3, to chromosome1p35–pter. Genomics 26: 611–614, 1995.

- [64216] 2546.Cody, J. D.; Hale, D. E.; Brkanac, Z.; Kaye, C. I.; Leach, R. J.: Growth hormone insufficiency associated with haploinsufficiency at 18q23. *Am. J. Med. Genet.* 71: 420–425, 1997.
- [64217] 2547.Habert–Ortoli, E.; Amiranoff, B.; Loquet, I.; Laburthe, M.; Mayaux,J.–F.: Molecular cloning of a functional human galanin receptor. *Proc.Nat. Acad. Sci.* 91: 9780–9783, 1994.
- [64218] 2548.Hecht, G.; Marrero, J. A.; Danilkovich, A.; Matkowskyj, K. A.;Savkovic, S. D.; Koutsouris, A.; Benya, R. V.: Pathogenic *Escherichiacoli* increase Cl– secretion from intestinal epithelia by upregulatinggalanin–1 receptor expression. *J. Clin. Invest.* 104: 253–262, 1999.
- [64219] 2549.Beckman, L.; Beckman, G.; Mi, M. P.; De Simone, J.: The humanplacental amino acid naphthylamidases: their molecular interrelationsand correlations with perinatal factors. *Hum. Hered.* 19: 249–257,1969.
- [64220] 2550.Beckman, L.; Bjorling, G.; Christodoulou, C.: Pregnancy enzymesand placental polymorphism. II. Leucine aminopeptidase. *Acta Genet.Statist. Med.* 16: 122–131, 1966.
- [64221] 2551.Horio, J.; Nomura, S.; Okada, M.; Katsumata, Y.; Nakanishi, Y.;Kumano, Y.; Takami, S.; Kinoshita, M.; Tsuji–

moto, M.; Nakazato, H.; Mizutani, S.: Structural organization of the 5-prime-end and chromosomal assignment of human placental leucine aminopeptidase/insulin-regulated membrane aminopeptidase gene. *Biochem. Biophys. Res. Commun.* 262:269–274, 1999.

- [64222] 2552. Nagasaka, T.; Nomura, S.; Okamura, M.; Tsujimoto, M.; Nakazato, H.; Oiso, Y.; Nakashima, N.; Mizutani, S.: Immunohistochemical localization of placental leucine aminopeptidase/oxytocinase in normal human placental, fetal and adult tissues. *Reprod. Fertil. Dev.* 9: 747–753, 1997.
- [64223] 2553. Røgi, T.; Tsujimoto, M.; Nakazato, H.; Mizutani, S.; Tomoda, Y.: Human placental leucine aminopeptidase/oxytocinase: a new member of type II membrane-spanning zinc metallopeptidase family. *J. Biol. Chem.* 271: 56–61, 1996.
- [64224] 2554. Scandalios, J. G.: Human serum leucine aminopeptidase: variation in pregnancy and in disease states. *J. Hered.* 58: 153–156, 1967.
- [64225] 2555. Levanon, D.; Negreanu, V.; Bernstein, Y.; Bar-Am, I.; Avivi, L.; Groner, Y.: AML1, AML2, and AML3, the human members of the runt domain gene-family: cDNA structure, expression, and chromosomal localization. *Genomics*

23:425–432, 1994.

- [64226] 2556. Bielinska, B.; Blaydes, S. M.; Buiting, K.; Yang, T.; Krajewska–Walasek, M.; Horsthemke, B.; Brannan, C. I.: De novo deletions of SNRPN exon1 in early human and mouse embryos result in a paternal to maternal imprint switch. *Nature Genet.* 25: 74–78, 2000. Note: Erratum: *Nature–Genet.* 25: 241 only, 2000.
- [64227] 2557. Raeymaekers, P.; Van Broeckhoven, C.; Backhovens, H.; Wehnert, A.; Muylle, L.; De Jonghe, P.; Gheuens, J.; Vandenberghe, A.: The Duffy blood group is linked to the alpha-spectrin locus in a large pedigree with autosomal dominant inheritance of Charcot–Marie–Tooth disease type 1. *Hum. Genet.* 78: 76–78, 1988.
- [64228] 2558. Eagle, L. R.; Yin, X.; Brothman, A. R.; Williams, B. J.; Atkin, N. B.; Prochownik, E. V.: Mutation of the MXI1 gene in prostate cancer. *Nature Genet.* 9: 249–255, 1995.
- [64229] 2559. Edwards, A.; Hammond, H. A.; Jin, L.; Caskey, C. T.; Chakraborty, R.: Genetic variation at five trimeric and tetrameric tandem repeat loci in four human population groups. *Genomics* 12: 241–253, 1992.
- [64230] 2560. Ichikawa, T.; Ichikawa, Y.; Dong, J.; Hawkins, A. L.; Griffin, C. A.; Isaacs, W. B.; Oshimura, M.; Barrett, J. C.; Isaacs, J. T.: Localization of metastasis suppressor gene(s)

for prostatic cancer to the short arm of human chromosome 11. *Cancer Res.* 52: 3486–3490, 1992.

[64231] 2561. Ichikawa, T.; Ichikawa, Y.; Isaacs, J. T.: Genetic factors and suppression of metastatic ability of prostatic cancer. *Cancer Res.* 51: 3788–3792, 1991.

[64232] 2562. Irvine, R. A.; Yu, M. C.; Ross, R. K.; Coetzee, G. A.: The CAG and GGC microsatellites of the androgen receptor gene are in linkage disequilibrium in men with prostate cancer. *Cancer Res.* 55: 1937–1940, 1995.

[64233] 2563. Peters, M. A.; Ostrander, E. A.: Prostate cancer: more than two to tango. *Nature Genet.* 27: 134–135, 2001.

[64234] 2564. Bansal, A.; Murray, D. K.; Wu, J. T.; Stephenson, R. A.; Middleton, R. G.; Meikle, A. W.: Heritability of prostate-specific antigen and relationship with zonal prostate volumes in aging twins. *J. Clin. Endocr. Metab.* 85: 1272–1276, 2000.

[64235] 2565. Cleutjens, K. B. J. M.; van der Korput, H. A. G. M.; van Eekelen, C. C. E. M.; van Rooij, H. C. J.; Faber, P. W.; Trapman, J.: An androgen response element in a far upstream enhancer region is essential for high, androgen-regulated activity of the prostate-specific antigen promoter. *Molec. Endocr.* 11: 148–161, 1997.

[64236] 2566. Nelson, R. J.; Demas, G. E.; Huang, P. L.; Fishman, M.

C.; Dawson, V. L.; Dawson, T. M.; Snyder, S. H.: Behavioural abnormalities in male mice lacking neuronal nitric oxide synthase. *Nature* 378: 383–386, 1995.

[64237] 2567. Henry, J. G.; Mitnick, M.; Dann, P. R.; Stewart, A. F.: Parathyroid hormone–related protein–(1–36) is biologically active when administered subcutaneously to humans. *J. Clin. Endocr. Metab.* 82: 900–906, 1997.

[64238] 2568. Magnaghi, P.; Agazzi, A.; Semino, O.; Ferrari, M.; Barbui, T.; D'Angelo, A.; Taramelli, R.: A recombination event in the closely linked plasminogen and apolipoprotein(a) gene loci. *Clin. Genet.* 47: 285–289, 1995.

[64239] 2569. Holick, M. F.; Ray, S.; Chen, T. C.; Tian, X.; Persons, K. S.: A parathyroid hormone antagonist stimulates epidermal proliferation and hair growth in mice. *Proc. Nat. Acad. Sci.* 91: 8014–8016, 1994.

[64240] 2570. Lanske, B.; Amling, M.; Neff, L.; Guiducci, J.; Baron, R.; Kronenberg, H. M.: Ablation of the PTHrP gene or the PTH/PTHrP receptor gene leads to distinct abnormalities in bone development. *J. Clin. Invest.* 104: 399–407, 1999.

[64241] 2571. Mangin, M.; Ikeda, K.; Dreyer, B. E.; Broadus, A. E.: Isolation and characterization of the human parathyroid hormone–like peptide gene. *Proc. Nat. Acad. Sci.* 86: 2408–2412, 1989.



- [64242] 2572. Mangin, M.; Webb, A. C.; Dreyer, B. E.; Posillico, J. T.; Ikeda, K.; Weir, E. C.; Stewart, A. F.; Bander, N. H.; Milstone, L.; Barton, D. E.; Francke, U.; Broadus, A. E.: Identification of a cDNA encoding a parathyroid hormone-like peptide from a human tumor associated with humoral hypercalcemia of malignancy. *Proc. Nat. Acad. Sci.* 85: 597–601, 1988.
- [64243] 2573. Moseley, J. M.; Kubota, M.; Diefenbach-Jagger, H.; Wettenhall, R. E. H.; Kemp, B. E.; Suva, L. J.; Rodda, C. P.; Ebeling, P. R.; Hudson, P. J.; Zajac, J. D.; Martin, T. J.: Parathyroid hormone-related protein purified from a human lung cancer cell line. *Proc. Nat. Acad. Sci.* 84: 5048–5052, 1987.
- [64244] 2574. Philbrick, W. M.; Dreyer, B. E.; Nakchbandi, I. A.; Karaplis, A. C.: Parathyroid hormone-related protein is required for tooth eruption. *Proc. Nat. Acad. Sci.* 95: 11846–11851, 1998.
- [64245] 2575. Strewler, G. J.: The physiology of parathyroid hormone-related protein. *New Eng. J. Med.* 342: 177–185, 2000.
- [64246] 2576. Suva, L. J.; Winslow, G. A.; Wettenhall, R. E. H.; Hammonds, R. G.; Moseley, J. M.; Diefenbach-Jagger, H.; Rodda, C. P.; Kemp, B. E.; Rodriguez, H.; Chen, E. Y.; Hud-

son, P. J.; Martin, T. J.; Wood, W. I.: A parathyroid hormone-related protein implicated in malignant hypercalcemia: cloning and expression. *Science* 237: 893–896, 1987.

[64247] 2577. Vortkamp, A.; Lee, K.; Lanske, B.; Segre, G. V.; Kronenberg, H. M.; Tabin, C. J.: Regulation of rate of cartilage differentiation by Indian hedgehog and PTH-related protein. *Science* 273: 613–622, 1996.

[64248] 2578. Wysolmerski, J. J.; Cormier, S.; Philbrick, W. M.; Dann, P.; Zhang, J.-P.; Roume, J.; Delezoide, A.-L.; Silve, C.: Absence of functional type 1 parathyroid hormone (PTH)/PTH-related protein receptors in humans is associated with abnormal breast development and tooth impaction. *J. Clin. Endocr. Metab.* 86: 1788–1794, 2001.

[64249] 2579. Wysolmerski, J. J.; Philbrick, W. M.; Dunbar, M. E.; Lanske, B.; Kronenberg, H.; Broadus, A. E.: Rescue of the parathyroid hormone-related protein knockout mouse demonstrates that parathyroid hormone-related protein is essential for mammary gland development. *Development* 125: 1285–1294, 1998.

[64250] 2580. Yasuda, T.; Banville, D.; Hendy, G. N.; Goltzman, D.: Characterization of the human parathyroid hormone-like peptide gene: functional and evolutionary aspects. *J. Biol.*

Chem. 264: 7720–7725, 1989.

- [64251] 2581.Mavrogiannis, L. A.; Antonopoulou, I.; Baxova, A.; Kutilek, S.;Kim, C. A.; Sugayama, S. M.; Salamanca, A.; Wall, S. A.; Morriss–Kay,G. M.; Wilkie, A. O. M.: Haploin–sufficiency of the human homeoboxgene ALX4 causes skull ossification defects. *Nature Genet.* 27: 17–18,2001.
- [64252] 2582.Qu, S.; Niswender, K. D.; Ji, Q.; van der Meer, R.; Keeney, D.;Magnuson, M. A.; Wisdom, R.: Polydactyly and ectopic ZPA formationin Alx–4 mutant mice. *Development* 124: 3999–4008, 1997.
- [64253] 2583.Wu, Y.–Q.; Badano, J. L.; McCaskill, C.; Vogel, H.; Potocki, L.;Shaffer, L. G.: Haploinsufficiency of ALX4 as a potential cause ofparietal foramina in the 11p11.2 contiguous gene–deletion syndrome. *Am.J. Hum. Genet.* 67: 1327–1332, 2000.
- [64254] 2584.Wuyts, W.; Cleiren, E.; Homfray, T.; Rasore–Quartino, A.; Vanhoenacker,F.; Van Hul, W.: The ALX4 homeobox gene is mutated in patients withossification defects of the skull (foramina parietalia permagna, OMIM168500). *J. Med. Genet.* 37: 916–920, 2000.
- [64255] 2585.Deybach, J.–C.; de Verneuil, H.; Boulechfar, S.; Grandchamp, B.;Nordmann, Y.: Point mutations in the uroporphyrinogen III synthasegene in congenital erythro–

poietic porphyria (Gunther's disease). Blood 75:1763–1765, 1990.

- [64256] 2586.Lemyre, E.; Russo, P.; Melancon, S. B.; Gagne, R.; Potier, M.; Lambert, M.: Clinical spectrum of infantile free sialic acid storage disease. Am. J. Med. Genet. 82: 385–391, 1999.
- [64257] 2587.Mancini, G. M. S.; Beerens, C. E. M. T.; Aula, P. P.; Verheijen, F. W.: Sialic acid storage diseases: a multiple lysosomal transport defect for acidic monosaccharides. J. Clin. Invest. 87: 1329–1335, 1991.
- [64258] 2588.Schleutker, J.; Laine, A.–P.; Haataja, L.; Renlund, M.; Weissenbach, J.; Aula, P.; Peltonen, L.: Linkage disequilibrium utilized to establish a refined genetic position of the Salla disease locus on 6q14–q15. Genomics 27:286–292, 1995.
- [64259] 2589.Schleutker, J.; Leppanen, P.; Mansson, J.–E.; Erikson, A.; Weissenbach, J.; Peltonen, L.; Aula, P.: Lysosomal free sialic acid storage disorders with different phenotypic presentations—infantile form sialic acid storage disease and Salla disease—represent allelic disorders on 6q14–15. Am. J. Hum. Genet. 57: 893–901, 1995.
- [64260] 2590.Tondeur, M.; Libert, J.; Vamos, E.; Van Hoof, F.; Thomas, G. H.; Strecker, G.: Infantile form of sialic acid

storage disorder: clinical, ultrastructural, and biochemical studies in two siblings. *Europ. J. Pediatr.* 139: 142–147, 1982.

- [64261] 2591. Verheijen, F. W.; Verbeek, E.; Aula, N.; Beerens, C. E. M. T.; Havelaar, A. C.; Joosse, M.; Peltonen, L.; Aula, P.; Galjaard, H.; van der Spek, P. J.; Mancini, G. M. S.: A new gene, encoding an anion transporter, is mutated in sialic acid storage diseases. *Nature Genet.* 23:462–465, 1999.
- [64262] 2592. Krasnewich, D. M.; Tietze, F.; Krause, W.; Pretzlaff, R.; Wenger, D. A.; Diwadkar, V.; Gahl, W. A.: Clinical and biochemical studies in an American child with sialuria. *Biochem. Med. Metab. Biol.* 49:90–96, 1993.
- [64263] 2593. Leroy, J. G.; Seppala, R.; Huizing, M.; Dacremont, G.; De Simpel, H.; Van Coster, R. N.; Orvisky, E.; Krasnewich, D. M.; Gahl, W. A.: Dominant inheritance of sialuria, an inborn error of feedback inhibition. *Am. J. Hum. Genet.* 68: 1419–1427, 2001.
- [64264] 2594. Seppala, R.; Lehto, V.-P.; Gahl, W. A.: Mutations in the human UDP-N-acetylglucosamine 2-epimerase gene define the disease sialuria and the allosteric site of the enzyme. *Am. J. Hum. Genet.* 64: 1563–1569, 1999.
- [64265] 2595. Murrell, J. R.; Spillanti, M. G.; Zolo, P.; Guazzelli, M.; Smith, M. J.; Hasegawa, M.; Redi, F.; Crowther, R. A.;

Pietrini, P.; Ghetti, B.; Goedert, M.: Tau gene mutation G389R causes a tauopathy with abundant Pick body-like inclusions and axonal deposits. *J. Neuropath. Exp. Neurol.* 58: 1207–1226, 1999.

[64266] 2596. Nagase, T.; Ishikawa, K.; Kikuno, R.; Hirose, M.; Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XV. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.* 6: 337–345, 1999.

[64267] 2597. Neumann, M.; Schulz-Schaeffer, W.; Crowther, R. A.; Smith, M. J.; Spillantini, M. G.; Goedert, M.; Kretzschmar, H. A.: Pick's disease associated with the novel tau gene mutation K369I. *Ann. Neurol.* 50:503–513, 2001.

[64268] 2598. Neve, R. L.; Harris, P.; Kosik, K. S.; Kurnit, D. M.; Donlon, T. A.: Identification of cDNA clones for the human microtubule-associated protein tau and chromosomal localization of the genes for tau and microtubule-associated protein 2. *Molec. Brain Res.* 1: 271–280, 1986.

[64269] 2599. Pickering-Brown, S.; Baker, M.; Yen, S.-H.; Liu, W.-K.; Hasegawa, M.; Cairns, N.; Lantos, P. L.; Rossor, M.; Iwatsubo, T.; Davies, Y.; Allsop, D.; Furlong, R.; Owen, F.; Hardy, J.; Mann, D.; Hutton, M.: Pick's disease is associated with mutations in the tau gene. *Ann. Neurol.* 48:

859–867, 2000.

- [64270] 2600.Poorkaj, P.: Personal Communication. Seattle, Wash. 11/10/1998.
- [64271] 2601.Poorkaj, P.; Bird, T. D.; Wijsman, E.; Nemens, E.; Garruto, R.M.; Anderson, L.; Andreadis, A.; Wiederholt, W. C.; Raskind, M.; Schellenberg, G. D.: Tau is a candidate gene for chromosome 17 frontotemporal dementia. *Ann.Neurol.* 43: 815–825, 1998.
- [64272] 2602.Poorkaj, P.; Kas, A.; D'Souza, I.; Zhou, Y.; Pham, Q.; Stone, M.; Olson, M. V.; Schellenberg, G. D.: A genomic sequence analysis of the mouse and human microtubule-associated protein tau. *MammalianGenome* 12: 700–712, 2001.
- [64273] 2603.Rapoport, M.; Dawson, H. N.; Binder, L. I.; Vitek, M. P.; Ferreira, A.: Tau is essential to beta-amyloid-induced neurotoxicity. *Proc.Nat. Acad. Sci.* 99: 6364–6369, 2002.
- [64274] 2604.Rizzu, P.; Van Swieten, J. C.; Joosse, M.; Hasegawa, M.; Stevens, M.; Tibben, A.; Niermeijer, M. F.; Hillebrand, M.; Ravid, R.; Oostra, B. A.; Goedert, M.; van Duijn, C. M.; Heutink, P.: High prevalence of mutations in the microtubule-associated protein tau in a population study of frontotemporal dementia in the Netherlands. *Am. J. Hum.Genet.* 64: 414–421, 1999.

- [64275] 2605.Spillantini, M. G.; Murrell, J. R.; Goedert, M.; Farlow, M. R.;Klug, A.; Ghetti, B.: Mutation in the tau gene in familial multiplesystem tauopathy with presenile dementia. Proc. Nat. Acad. Sci. 95:7737–7741, 1998.
- [64276] 2606.Spillantini, M. G.; Yoshida, H.; Rizzini, C.; Lantos, P. L.; Khan,N.; Rossor, M. N.; Goedert, M.; Brown, J.: A novel tau mutation (N296N)in familial dementia with swollen achromatic neurons and corticobasal inclusion bodies. Ann. Neurol. 48: 939–943, 2000.
- [64277] 2607.Stambolic, V.; Ruel, L.; Woodgett, J. R.: Lithium inhibits glycogensynthase kinase–3 activity and mimics Wingless signalling in intactcells. Curr. Biol. 6: 1664–1668, 1996. Note: Erratum: Curr. Biol.7: 196 only, 1997.
- [64278] 2608.Stamer, K.; Vogel, R.; Thies, E.; Mandelkow, E.; Mandelkow, E.–M.: Tau blocks traffic of organelles, neurofilaments, and APP vesicles in neurons and enhances oxidative stress. J. Cell Biol. 156: 1051–1063,2002.
- [64279] 2609.Varani, L.; Hasegawa, M.; Spillantini, M. G.; Smith, M. J.; Murrell,J. R.; Ghetti, B.; Klug, A.; Goedert, M.; Varani, G.: Structure of tau exon 10 splicing regulatory element RNA and destabilization by mutations of frontotemporal dementia and parkinsonism linked to chromosome17. Proc.



Nat. Acad. Sci. 96: 8229–8234, 1999.

- [64280] 2610. Verpillat, P.; Camuzat, A.; Hannequin, D.; Thomas-Anterion, C.; Puel, M.; Belliard, S.; Dubois, B.; Didic, M.; Michel, B.-F.; Lacomblez, L.; Moreaud, O.; Sella, F.; Golfier, V.; Campion, D.; Clerget-Darpoux, F.; Brice, A.: Association between the extended tau haplotype and frontotemporal dementia. Arch. Neurol. 59: 935–939, 2002.
- [64281] 2611. Wilhelmsen, K. C.; Lynch, T.; Pavlou, E.; Higgins, M.; Hygaard, T. G.: Localization of disinhibition-dementia-parkinsonism-amyotrophy complex to 17q21–22. Am. J. Hum. Genet. 55: 1159–1165, 1994.
- [64282] 2612. Yasuda, M.; Yokoyama, K.; Nakayasu, T.; Nishimura, Y.; Matsui, M.; Yokoyama, T.; Miyoshi, K.; Tanaka, C.: A Japanese patient with frontotemporal dementia and parkinsonism by a tau P301S mutation. Neurology 55: 1224–1227, 2000.
- [64283] 2613. Zhukareva, V.; Mann, D.; Pickering-Brown, S.; Uryu, K.; Shuck, T.; Shah, K.; Grossman, M.; Miller, B. L.; Hulette, C. M.; Feinstein, S. C.; Trojanowski, J. Q.; Lee, V. M.-Y.: Sporadic Pick's disease: a tauopathy characterized by a spectrum of pathological tau isoforms in gray and white matter. Ann. Neurol. 51: 730–739, 2002.
- [64284] 2614. Weintraub, H.; Davis, R.; Tapscott, S.; Thayer, M.;

Krause, M.;Benezra, R.; Blackwell, T. K.; Turner, D.; Rupp, R.; Hollenberg, S.;Zhuang, Y.; Lassar, A.: The myoD gene family: nodal point during specification of the muscle cell lineage. *Science* 251: 761–766,1991.

[64285] 2615.Amack, J. D.; Mahadevan, M. S.: The myotonic dystrophy expandedCUG repeat tract is necessary but not sufficient to disrupt C2C12myoblast differentiation. *Hum. Molec. Genet.* 10: 1879–1887, 2001.

[64286] 2616.Chen, S. L.; Dowhan, D. H.; Hosking, B. M.; Muscat, G. E. O.:The steroid receptor coactivator, GRIP-1, is necessary for MEF-2C-dependentgene expression and skeletal muscle differentiation. *Genes Dev.* 14:1209–1228, 2000.

[64287] 2617.Hasty, P.; Bradley, A.; Morris, J. H.; Edmondson, D. G.; Venuti,J. M.; Olson, E. N.; Klein, W. H.: Muscle deficiency and neonataldeath in mice with a targeted mutation in the myogenin gene. *Nature* 364:501–506, 1993.

[64288] 2618.Nabeshima, Y.; Hanaoka, K.; Hayasaka, M.; Esumi, E.; Li, S.; Nonaka,I.; Nabeshima, Y.–I.: Myogenin gene disruption results in perinatallethality because of severe muscle defect. *Nature* 364: 532–535,1993.

[64289] 2619.Braun, T.; Bober, E.; Winter, B.; Rosenthal, N.; Arnold, H. H.: Myf-6, a new member of the human gene

family of myogenic determination factors: evidence for a gene cluster on chromosome 12. *EMBO J.* 9:821–831, 1990.

[64290] 2620. Braun, T.; Grzeschik, K.-H.; Bober, E.; Arnold, H.-H.: The MYF genes, a group of human muscle determining factors, are localized on different human chromosomes. (Abstract) *Cytogenet. Cell Genet.* 51:969, 1989.

[64291] 2621. Olson, E.; Edmondson, D.; Wright, W. E.; Lin, V. K.; Guenet, J.-L.; Simon-Chazottes, D.; Thompson, L. H.; Stallings, R. L.; Schroeder, W. T.; Duvic, M.; Brock, D.; Heilin, D.; Siciliano, M. J.: Myogenin is in an evolutionarily conserved linkage group on human chromosome 1q31–q41 and unlinked to other mapped muscle regulatory factor genes. *Genomics* 8:427–434, 1990.

[64292] 2622. Cupelli, L.; Renault, B.; Leblanc-Straceski, J.; Banks, A.; Ward, D.; Kucherlapati, R. S.; Krauter, K.: Assignment of the human myogenic factors 5 and 6 (MYF5, MYF6) gene cluster to 12q21 by in situ hybridization and physical mapping of the locus between D12S350 and D12S106. *Cytogenet. Cell Genet.* 72: 250–251, 1996.

[64293] 2623. Ott, M.-O.; Bober, E.; Lyons, G.; Arnold, H.-H.; Buckingham, M.: Early expression of the myogenic regulatory gene, Myf-5, in precursor cells of skeletal muscle in

the mouse embryo. *Development* 111:1097–1107, 1991.

[64294] 2624.Rudnicki, M. A.; Schneegelsberg, P. N. J.; Stead, R. H.; Braun,T.; Arnold, H.–H.; Jaenisch, R.: MyoD or Myf–5 is required for theformation of skeletal muscle. *Cell* 75: 1351–1359, 1993.

[64295] 2625.Akaboshi, E.: Cloning of the human myoglobin gene. *Gene* 33:241–249, 1985.

[64296] 2626.Boulton, F. E.; Huntsman, R. G.; Lehmann, H.; Lorkin, P. A.; Romero–Herrera,A. E.: Myoglobin variants. (Abstract) *Biochem. J.* 118: 39P only,1970.

[64297] 2627.Boulton, F. E.; Huntsman, R. G.; Lorkin, P. A.; Lehmann, H.: Abnormalhuman myoglobin: 53(D4) glutamic acid lysine. *Nature* 223: 832–833,1969.

[64298] 2628.Boulton, F. E.; Huntsman, R. G.; Yawson, G. I.; Romero–Herrera,A. E.; Lorkin, P. A.: The second variant of human myoglobin: 138(H16)arginine to glutamine. *Brit. J. Haemat.* 20A: 69–74, 1971.

[64299] 2629.Boyer, S. H.: Similar incidence and non–randomness among humanmyoglobin and hemoglobin mutants in general populations: implicationsfor the study of myoglobin in muscle disease.In: *Pathogenesis of HumanMuscular Dystrophies. Proc. Vth Int. Cong. of Muscular Dystrophy Assoc.,Durango, Colo., June 21–25, 1976. Amster–*

dam: Excerpta Medica (pub.) 1977.

- [64300] 2630. Boyer, S. H.; Fainer, D. C.; Naughton, M. A.: Myoglobin inherited structural variation in man. *Science* 140: 1228–1231, 1963.
- [64301] 2631. Dozier, C.; Walbaum, S.; Leprince, D.; Stehelin, D.: EcoRI RFLP linked to the human myb gene. *Nucleic Acids Res.* 14: 1928 only, 1986.
- [64302] 2632. Garry, D. J.; Ordway, G. A.; Lorenz, J. N.; Radford, N. B.; Chin, E. R.; Grange, R. W.; Bassel-Duby, R.; Williams, R. S.: Mice without myoglobin. *Nature* 395: 905–908, 1998.
- [64303] 2633. Godecke, A.; Flogel, U.; Zanger, K.; Ding, Z.; Hirchenhain, J.; Decking, U. K. M.; Schrader, J.: Disruption of myoglobin in mice induces multiple compensatory mechanisms. *Proc. Nat. Acad. Sci.* 96: 10495–10500, 1999.
- [64304] 2634. Jeffreys, A.; Wilson, V.; Thein, S.: Hypervariable 'minisatellite' regions in human DNA. *Science* 314: 67–73, 1985.
- [64305] 2635. Jeffreys, A. J.; Wilson, V.; Blanchetot, A.; Weller, P.; Geurtsvan Kessel, A.; Spurr, N.; Solomon, E.; Goodfellow, P.: The human myoglobin gene: a third dispersed globin locus in the human genome. *Nucleic Acids Res.* 12: 3235–3243, 1984.

- [64306] 2636. McKeithan, T. W.; Rowley, J. D.; Shows, T. B.; Diaz, M. O.: Cloning of the chromosome translocation breakpoint junction of the t(14;19) in chronic lymphocytic leukemia. *Proc. Nat. Acad. Sci.* 84: 9257–9260, 1987.
- [64307] 2637. de Groot, N. G.; Otting, N.; Doxiadis, G. G. M.; Balla-Jhagjhoorsingh, S. S.; Heeney, J. L.; van Rood, J. J.; Gagneux, P.; Bontrop, R. E.: Evidence for an ancient selective sweep in the MHC class I gene repertoire of chimpanzees. *Proc. Nat. Acad. Sci.* 99: 11748–11753, 2002.
- [64308] 2638. Del Pozzo, G.; Guardiola, J.: Mummy DNA fragment identified. (Letter) *Nature* 339: 431–432, 1989.
- [64309] 2639. Dorf, M. E.; Balner, H.; DeGroot, M. L.; Benacerraf, B.: Histocompatibility-linked immune-response genes in the Rhesus monkey. *Transplant. Proc.* 6: 119–124, 1974.
- [64310] 2640. Edwards, J. H.; Allen, F. H.; Glenn, K. P.; Lamm, L. U.; Robson, E. B.: The linkage relationships of HL-A. In: *Histocompatibility Testing*. Baltimore: Williams and Wilkins (pub.) 1973.
- [64311] 2641. Engelfriet, C. P.; Britten, A.: The cytotoxic test for leucocyte antibodies. A simple and reliable technique. *Vox Sang.* 10: 660–674, 1965.
- [64312] 2642. Erlich, H.; Lee, J. S.; Petersen, J. W.; Bugawan, T.; Demars, R.: Molecular analysis of HLA class I and class II anti-

gen loss mutants reveals a homozygous deletion of the DR, DQ, and part of the DP region: implications for class II gene order. Hum. Immun. 16: 205–219, 1986.

- [64313] 2643. Erlich, H. A.; Stetler, D.; Sheng–Dong, R.; Ness, D.; Grumet, C.: Segregation and mapping analysis of polymorphic HLA class I restriction fragments: detection of a novel fragment. Science 222: 72–74, 1983.
- [64314] 2644. Fellous, M.; Dausset, J.: Probable haploid expression of HL–A antigens on human spermatozoan. Nature 225: 191–193, 1970.
- [64315] 2645. Figueroa, F.; Gunther, E.; Klein, J.: MHC polymorphism pre–dates speciation. (Letter) Nature 335: 265–267, 1988.
- [64316] 2646. Francke, U.; Pellegrino, M. A.: Assignment of the major histocompatibility complex to a region of the short arm of human chromosome 6. Proc. Nat. Acad. Sci. 74: 5776 only, 1977.
- [64317] 2647. Francke, U.; Pellegrino, M. A.: Assignment of the major histocompatibility complex to a region of the short arm of human chromosome 6. Proc. Nat. Acad. Sci. 74: 1147–1151, 1977.
- [64318] 2648. Gill, T. J., III; Cramer, D. V.; Kunz, H. W.: The major histocompatibility complex—comparison in the mouse,

man, and the rat: a review. *Am.J. Path.* 90: 735–777, 1978.

[64319] 2649.Gladstone, P.; Furesz, L.; Pious, D.: Gene dosage and gene expression in the HLA region: evidence from deletion variants. *Proc. Nat. Acad.Sci.* 79: 1235–1239, 1982.

[64320] 2650.Gluecksohn-Waelsch, S.; Erickson, R. P.: The T-locus of the mouse: implications for mechanisms of development. *Curr. Top. Dev. Biol.* 5:281–316, 1970.

[64321] 2651.Goodfellow, P. N.; Jones, E.; Van Heyningen, V.; Solomon, E.; Kennett, R.; Bobrow, M.; Bodmer, W. F.: Linkage relationships of the HL-A system and beta-2-microglobulin. *Birth Defects Orig. Art.Ser.* 11(3): 162–167, 1975. Note: *Alternate: Cytogenet. Cell Genet.* 14: 332–337, 1975.

[64322] 2652.Gruen, J. R.; Nalabolu, S. R.; Chu, T. W.; Bowlus, C.; Fan, W.F.; Goei, V. L.; Wei, H.; Sivakamasundari, R.; Liu, Y.-C.; Xu, H.X.; Parimoo, S.; Nallur, G.; Ajioka, R.; Shukla, H.; Bray-Ward, P.; Pan, J.; Weissman, S. M.: A transcription map of the major histocompatibility complex (MHC) class I region. *Genomics* 36: 70–85, 1996.

[64323] 2653.Hedrick, S. M.: Dawn of the hunt for nonclassical MHC function. *Cell* 70:177–180, 1992.



- [64324] 2654.Helenius, A.; Morein, B.; Fries, E.; Simons, K.; Robinson, P.;Schirrmacher, V.; Terhorst, C.; Strominger, J. L.: Human (HLA-A andHLA-B) and murine (H-2K and H-2D) histocompatibility antigens arecell surface receptors for Semliki Forest virus. Proc. Nat. Acad.Sci. 75: 3846–3850, 1978.
- [64325] 2655.Hirai, M.; Takahashi, E.; Ishida, T.; Hori, T.: Chromosomal localizationof the major histocompatibility complex (MHC) in the rhesus monkeyand chimpanzee by fluorescence in situ hybridization. Cytogenet.Cell Genet. 57: 204–205, 1991.
- [64326] 2656.Hood, L.; Steinmetz, M.; Goodenow, R.: Genes of the major histocompatibilitycomplex. Cell 28: 685–687, 1982.
- [64327] 2657.Jordan, B. R.; Bregegere, F.; Kourilsky, P.: Human HLA gene segmentisolated by hybridization with mouse H-2 cDNA probes. Nature 290:521–523, 1981.
- [64328] 2658.Kissmeyer-Nielsen, F.; Jorgensen, F.; Lamm, L. U.: The HL-A systemin clinical medicine. Johns Hopkins Med. J. 131: 385–400, 1972.
- [64329] 2659.Kissmeyer-Nielsen, F.; Svejgaard, A.; Ahrons, S.; Nielsen, L.S.: Crossing-over within the HL-A system. Nature 224: 75–76, 1969.
- [64330] 2660.Kissmeyer-Nielsen, F.; Svejgaard, A.; Hauge, M.: Ge-

netics of the human HL-A transplantation system. *Nature* 219: 1116–1119, 1968.

[64331] 2661. Kissmeyer-Nielsen, F.; Thorsby, E.: Human transplantation antigens. *Transplant. Rev.* 4: 1–176, 1970.

[64332] 2662. Klein, J.: The major histocompatibility complex of the mouse. *Science* 203: 516–521, 1979.

[64333] 2663. Klein, J.; Figueroa, F.: Evolution of the major histocompatibility complex. *CRC Crit. Rev. Immun.* 6: 295–386, 1986.

[64334] 2664. Kompf, J.; Bissbort, S.; Gohler, F.; Schunter, F.; Wernet, P.: Mapping of the linkage group GLO-Bf-HLA-B, C, A-PGM. I. Recombination frequencies. *Hum. Genet.* 44: 313–319, 1978.

[64335] 2665. Lamm, L. U.; Friedrich, U.; Petersen, G. B.; Jorgensen, J.; Nielsen, J.; Therkelsen, A. J.; Kissmeyer-Nielsen, F.: Assignment of the major histocompatibility complex to chromosome no. 6 in a family with a pericentric inversion. *Hum. Hered.* 24: 273–284, 1974.

[64336] 2666. Lamm, L. U.; Kissmeyer-Nielsen, F.; Svejgaard, A.; Petersen, G. B.; Thorsby, E.; Mayr, W.; Hogman, C.: On the orientation of the HL-A region and the PGM(3) locus in the chromosome. *Tissue Antigens* 2: 205–214, 1972.

[64337] 2667. Lamm, L. U.; Svejgaard, A.; Kissmeyer-Nielsen, F.:

Further evidence for PGM(3): HL-A is another linkage in man. *Nature* N.B. 231: 109–110, 1971.

[64338] 2668. Lamm, L. U.; Thorsen, I.-L.; Petersen, G. B.; Jorgensen, J.; Henningsen, K.; Bech, B.; Kissmeyer-Nielsen, F.: Data on the HL-A linkage group. *Ann. Hum. Genet.* 38: 383–390, 1975.

[64339] 2669. Lawlor, D. A.; Dickel, C. D.; Hauswirth, W. W.; Parham, P.: Ancient HLA genes from 7,500-year-old archaeological remains. *Nature* 349: 785–788, 1991.

[64340] 2670. Lawlor, D. A.; Ward, F. E.; Ennis, P. D.; Jackson, A. P.; Parham, P.: HLA-A and B polymorphisms predate the divergence of humans and chimpanzees. (Letter) *Nature* 335: 268–271, 1988.

[64341] 2671. Lawrance, S. K.; Smith, C. L.; Srivastava, R.; Cantor, C. R.; Weissman, S. M.: Megabase-scale mapping of the HLA gene complex by pulsed field gel electrophoresis. *Science* 235: 1387–1390, 1987.

[64342] 2672. Levine, F.; Erlich, H.; Mach, B.; Leach, R.; White, R.; Pious, D.: Deletion mapping of HLA and chromosome 6p genes. *Proc. Nat. Acad. Sci.* 82: 3741–3745, 1985.

[64343] 2673. Slate, D. L.; Ruddle, F. H.: Genetics of the interferon system. *Pharm. Therap.* 4: 221–230, 1979.

[64344] 2674. Stock, A. D.; Hsu, T. C.: Evolutionary conservatism in

arrangement of genetic material. A comparative analysis of chromosome banding between the Rhesus macaque ( $2n=42$ , 84 arms) and the African green monkey ( $2n=60$ , 120 arms). *Chromosoma* 43: 211–224, 1973.

[64345] 2675. Streuli, M.; Nagata, S.; Weissmann, C.: At least three human type alpha interferons: structure of alpha-2. *Science* 209: 1343–1347, 1980.

[64346] 2676. Szabo, S. J.; Sullivan, B. M.; Stemmann, C.; Satoskar, A. R.; Sleckman, B. P.; Glimcher, L. H.: Distinct effects of T-bet in T(H)1 lineage commitment and IFN-gamma production in CD4 and CD8 T cells. *Science* 295: 338–342, 2002.

[64347] 2677. Takayanagi, H.; Ogasawara, K.; Hida, S.; Chiba, T.; Murata, S.; Sato, K.; Takaoka, A.; Yokochi, T.; Oda, H.; Tanaka, K.; Nakamura, K.; Taniguchi, T.: T-cell-mediated regulation of osteoclastogenesis by signalling cross-talk between RANKL and IFN-gamma. *Nature* 408: 600–605, 2000.

[64348] 2678. Tan, Y. H.; Creagan, R. P.; Ruddle, F. H.: Assignment of the genes of the human interferon system to chromosomes 2 and 5. *Cytogenet. Cell Genet.* 13: 155–157, 1974.

[64349] 2679. Tan, Y. H.; Ke, Y. H.; Armstrong, J. A.; Ho, M.: The regulation of cellular interferon production: enhancement by antimetabolites. *Proc. Nat. Acad. Sci.* 67: 464–471,

1970.

- [64350] 2680. Trent, J. M.; Olson, S.; Lawn, R. M.: Chromosomal localization of human leukocyte, fibroblast, and immune interferon genes by means of in situ hybridization. *Proc. Nat. Acad. Sci.* 79: 7809–7813, 1982.
- [64351] 2681. Tzoneva, M.; Ganey, V.; Galabov, A.; Georgieva, K.: Selective immunodeficiency with defect in interferon-gamma induction in two sibs with recurrent infections. *Clin. Genet.* 33: 454–456, 1988.
- [64352] 2682. White, A. C., Jr.; Robinson, P.; Okhuysen, P. C.; Lewis, D. E.; Shahab, I.; Lahoti, S.; DuPont, H. L.; Chappell, C. L.: Interferon-gamma expression in jejunal biopsies in experimental human cryptosporidiosis correlates with prior sensitization and control of oocyst excretion. *J. Infect. Dis.* 181: 701–709, 2000.
- [64353] 2683. Yip, Y. K.; Barrowclough, B. S.; Urban, C.; Vilcek, J.: Purification of two subspecies of human gamma (immune) interferon. *Proc. Nat. Acad. Sci.* 79: 1820–1824, 1982.
- [64354] 2684. Zimonjic, D. B.; Rezanka, L. J.; Evans, C. H.; Polymeropoulos, M. H.; Trent, J. M.; Popescu, N. C.: Mapping of the immune interferon gamma gene (IFNG) to chromosome band 12q14 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 71: 247–248, 1995.

- [64355] 2685.Zohlhofer, D.; Richter, T.; Neumann, F.-J.; Nuhrenberg, T.; Wessely,R.; Brandl, R.; Murr, A.; Klein, C. A.; Baeuerle, P. A.: Transcriptomeanalysis reveals a role of interferon-gamma in human neointoma formation. Molec.Cell 7: 1059-1069, 2001.
- [64356] 2686.Zoon, K. C.; Smith, M. E.; Bridgen, P. J.; Anfinsen, C. B.; Hunkapiller,M. W.; Hood, L. E.: Amino terminal sequence of the major componentof human lymphoblastoid interferon. Science 207: 527-528, 1980.
- [64357] 2687.Clauss, I. M.; Wathelet, M. G.; Szpirer, J.; Content, J.; Islam,M. Q.; Levan, G.; Szpirer, C.; Huez, G. A.: Chromosomal localizationof two human genes inducible by interferons, double-stranded RNA,and viruses. Cytogenet. Cell Genet. 53: 166-168, 1990.
- [64358] 2688.Itzhaki, J. E.; Barnett, M. A.; MacCarthy, A. B.; Buckle, V. J.;Brown, W. R. A.; Porter, A. C. G.: Targeted breakage of a human chromosomemediated by cloned human telomeric DNA. Nature Genet. 2: 283-287,1992.
- [64359] 2689.Itzhaki, J. E.; Porter, A. C. G.: Targeted disruption of a humaninterferon-inducible gene detected by secretion of human growth hormone. NucleicAcids Res. 19: 3835-3842, 1991.
- [64360] 2690.Kelly, J. M.; Porter, A. C. G.; Chernajovsky, Y.;

Gilbert, C. S.; Stark, G. R.; Kerr, I. M.: Characterization of a human gene inducible by alpha- and beta-interferons and its expression in mouse cells. EMBO J. 5: 1601–1606, 1986.

[64361] 2691. Porter, A. C. G.; Chernajovsky, Y.; Dale, T. C.; Gilbert, C. S.; Stark, G. R.; Kerr, I. M.: Interferon response element of the human gene 6–16. EMBO J. 7: 85–92, 1988.

[64362] 2692. Fu, X.-Y.; Schindler, C.; Imbrota, T.; Aebersold, R.; Darnell, J. E., Jr.: The proteins of ISGF-3, the interferon alpha-induced transcriptional activator, define a gene family involved in signal transduction. Proc. Nat. Acad. Sci. 89: 7840–7843, 1992.

[64363] 2693. Levy, D. E.; Kessler, D. S.; Pine, R.; Darnell, J. E., Jr.: Cytoplasmic activation of ISGF3, the positive regulator of interferon-alpha-stimulated transcription, reconstituted in vitro. Genes Dev. 3: 1362–1371, 1989.

[64364] 2694. McCusker, D.; Wilson, M.; Trowsdale, J.: Organization of the genes encoding the human proteasome activators PA28-alpha and beta. Immunogenetics 49: 438–445, 1999.

[64365] 2695. Schindler, C.; Fu, X.-Y.; Imbrota, T.; Aebersold, R.; Darnell, J. E., Jr.: Proteins of transcription factor ISGF-3: one gene encodes the 91- and 84-kDa ISGF-3 proteins

that are activated by interferon alpha. Proc. Nat. Acad. Sci. 89: 7836–7839, 1992.

- [64366] 2696. Suhara, W.; Yoneyama, M.; Yonekawa, H.; Fujita, T.: Structure of mouse interferon stimulated gene factor 3 gamma (ISGF3 gamma/p48) cDNA and chromosomal localization of the gene. J. Biochem. 119:231–234, 1996.
- [64367] 2697. Veals, S. A.; Schindler, C.; Leonard, D.; Fu, X.-Y.; Aebersold, R.; Darnell, J. E., Jr.; Levy, D. E.: Subunit of an alpha-interferon-responsive transcription factor is related to interferon regulatory factor and myb families of DNA-binding proteins. Molec. Cell. Biol. 12: 3315–3324, 1992.
- [64368] 2698. Eason, D. D.; Shepherd, A. T.; Blanck, G.: Interferon regulatory factor 1 tryptophan 11 to arginine point mutation abolishes DNA binding. Biochim. Biophys. Acta 1446: 140–144, 1999.
- [64369] 2699. Savtchenko, E. S.; Tomic, M.; Ivker, R.; Blumenberg, M.: Three parallel linkage groups of human acidic keratin genes. Genomics 7:394–407, 1990.
- [64370] 2700. Bowden, P. E.; Haley, J. L.; Kinsky, A.; Rothnagel, J. A.; Jones, D. O.; Turner, R. J.: Mutation of a type II keratin gene (K6a) in pachyonychia congenita. Nature Genet. 10: 363–365, 1995.
- [64371] 2701. Rosenberg, M.; Fuchs, E.; Le Beau, M. M.; Eddy, R. L.;



Shows, T.B.: Three epidermal and one simple epithelial type II keratin genes map to human chromosome 12. Cytogenet. Cell Genet. 57: 33–38, 1991.

- [64372] 2702. Takahashi, K.; Paladini, R. D.; Coulombe, P. A.: Cloning and characterization of multiple human genes and cDNAs encoding highly related type II keratin 6 isoforms. J. Biol. Chem. 270: 18581–18592, 1995.
- [64373] 2703. Tyner, A.; Eichman, M.; Fuchs, E.: The sequence of a type II keratin gene expressed in human skin: conservation of structure among all intermediate filament genes. Proc. Nat. Acad. Sci. 82: 4683–4687, 1985.
- [64374] 2704. Wang, N.; Perkins, K. L.: Involvement of band 3p14 in t(3;8) hereditary renal carcinoma. Cancer Genet. Cytogenet. 11: 479–481, 1984.
- [64375] 2705. Schoenmakers, E. F. P. M.; Huysmans, C.; Van de Ven, W. J. M.: Allelic knockout of novel splice variants of human recombination repair gene RAD51B in t(12;14) uterine leiomyomas. Cancer Res. 59: 19–23, 1999.
- [64376] 2706. Funk, C. D.; Funk, L. B.; FitzGerald, G. A.; Samuelsson, B.: Characterization of human 12-lipoxygenase genes. Proc. Nat. Acad. Sci. 89: 3962–3966, 1992.
- [64377] 2707. Yoshida, M. C.; Sasaki, M.; Mise, K.; Semba, K.; Nishizawa, M.; Yamamoto, T.; Toyoshima, K.: Regional

mapping of the human proto-oncogene c-yes-1 to chromosome 18 at band q21.3. Jpn. J. Cancer Res. 76:559–562, 1985.

- [64378] 2708. Goossens, M.; Brauner, R.; Czernichow, P.; Duquesnoy, P.; Rappaport, R.: Isolated growth hormone (GH) deficiency type 1A associated with a double deletion in the human GH gene cluster. J. Clin. Endocr. Metab. 62: 712–716, 1986.
- [64379] 2709. Harper, M. E.; Barrera-Saldana, H. A.; Saunders, G. F.: Chromosomal localization of the human placental lactogen-growth hormone gene cluster to 17q22–24. Am. J. Hum. Genet. 34: 227–234, 1982.
- [64380] 2710. Harmer, D.; Evans, D. A. P.; Eze, L. C.; Jolly, M.; Whibley, E. J.: The relationship between the acetylator and the sparteine hydroxylation polymorphisms. J. Med. Genet. 23: 155–156, 1986.
- [64381] 2711. Hendershot, L. M.; Valentine, V. A.; Lee, A. S.; Morris, S. W.; Shapiro, D. N.: Localization of the gene encoding human BiP/GRP78, the endoplasmic reticulum cognate of the HSP70 family, to chromosome 9q34. Genomics 20: 281–284, 1994.
- [64382] 2712. Law, M. L.; Seeliger, M. B.; Lee, A. S.; Kao, F. T.: Genetic mapping of the structural gene coding for a glucose–

regulated protein (GRP78) of 78k-dalton to the long arm of human chromosome 9. (Abstract) Cytogenet. Cell Genet. 37: 518–519, 1984.

- [64383] 2713. Lee, A. S.; Deleage, A.; Scharff, D.: Highly conserved glucose-regulated protein in hamster and chicken cells: preliminary characterization of its cDNA clone. Proc. Nat. Acad. Sci. 78: 4922–4925, 1981.
- [64384] 2714. Lee, A. S.; Deleage, A. M.; Baker, V.; Chow, P. C.: Transcriptional regulation of two genes specifically induced by glucose starvation in hamster mutant fibroblast cell line. J. Biol. Chem. 258: 597–603, 1983.
- [64385] 2715. Muresan, Z.; Arvan, P.: Enhanced binding to the molecular chaperone BiP slows thyroglobulin export from the endoplasmic reticulum. Molec. Endocr. 12: 458–467, 1998.
- [64386] 2716. McCombie, R. R.; Dolphin, C. T.; Povey, S.; Phillips, I. R.; Shephard, E. A.: Localization of human flavin-containing monooxygenase genes FMO2 and FMO5 to chromosome 1q. Genomics 34: 426–429, 1996.
- [64387] 2717. Bevilacqua, M. P.; Nelson, R. M.: Selectins. J. Clin. Invest. 91: 379–387, 1993.
- [64388] 2718. Florian, V.; Schluter, T.; Bohnsack, R.: A new member of the sorting nexin family interacts with the C-

terminus of P-selectin. Biochem.Biophys. Res. Commun. 281: 1045–1050, 2001.

[64389] 2719.Herrmann, S.-M.; Ricard, S.; Nicaud, V.; Mallet, C.; Evans, A.;Ruidavets, J.-B.; Arveiler, D.; Luc, G.; Cambien, F.: The P-selectingene is highly polymorphic: reduced frequency of the pro715 allelecarriers in patients with myocardial infarction. Hum. Molec. Genet. 7:1277–1284, 1998.

[64390] 2720.Johnston, G. I.; Bliss, G. A.; Newman, P. J.; McEver, R. P.: Structureof the human gene encoding granule membrane protein-140, a memberof the selectin family of adhesion receptors for leukocytes. J. Biol.Chem. 265: 21381–21385, 1990.

[64391] 2721.Johnston, G. I.; Cook, R. G.; McEver, R. P.: Cloning of GMP-140,a granule membrane protein of platelets and endothelium: sequencessimilarity to proteins involved in cell adhesion and inflammation. Cell 56:1033–1044, 1989.

[64392] 2722.Johnston, G. I.; Le Beau, M. M.; Lemons, R. S.; McEver, R. P.:Cloning of GMP-140: chromosomal localization, molecular heterogeneityand identification of cDNAs predicting both membrane bound and solubleproteins. (Abstract) Blood 72 (suppl.): 327a only, 1988.

- [64393] 2723.Lages, B.; Shattil, S. J.; Bainton, D. F.; Weiss, H. J.: Decreased content and surface expression of alpha-granule membrane protein GMP-140 in one of two types of platelet alpha-delta storage pool deficiency. *J.Clin. Invest.* 87: 919-929, 1991.
- [64394] 2724.Mayadas, T. N.; Johnson, R. C.; Rayburn, H.; Hynes, R. O.; Wagner, D. D.: Leukocyte rolling and extravasation are severely compromised in P selectin-deficient mice. *Cell* 74: 541-554, 1993.
- [64395] 2725.Mazurov, A. V.; Vinogradov, D. V.; Khaspekova, S. G.; Krushinsky, A. V.; Gerdeva, L. V.; Vasiliev, S. A.: Deficiency of P-selectin in a patient with grey platelet syndrome. *Europ. J. Haemat.* 57:38-41, 1996.
- [64396] 2726.Souttou, B.; Juhl, H.; Hackenbruck, J.; Rockseisen, M.; Klomp, H. J.; Raulais, D.; Vigny, M.; Wellstein, A.: Relationship between serum concentrations of the growth factor pleiotrophin and pleiotrophin-positive tumors. *J. Nat. Cancer Inst.* 90: 1468-1473, 1998.
- [64397] 2727.Weber, D.; Klomp, H.-J.; Czubyko, F.; Wellstein, A.; Juhl, H.: Pleiotrophin can be rate-limiting for pancreatic cancer cell growth. *Cancer Res.* 60: 5284-5288, 2000.
- [64398] 2728.Zhang, N.; Yeh, H.-J.; Zhong, R.; Li, Y.-S.; Deuel, T. F.: A dominant-negative pleiotrophin mutant introduced

by homologous recombination leads to germ-cell apoptosis in male mice. *Proc. Nat. Acad. Sci.* 96: 6734–6738, 1999.

[64399] 2729. Eddy, R. L.; Kretschmer, P. J.; Fairhurst, J. L.; Shows, T. B.; Bohlen, P.; O'Hara, B.; Kovesdi, I.: A human gene family of neurite outgrowth-promoting proteins: heparin-binding neurite outgrowth promoting factor maps to 11p11–11p13. (Abstract) *Cytogenet. Cell Genet.* 58:1958 only, 1991.

[64400] 2730. Kaname, T.; Kuwano, A.; Murano, I.; Uehara, K.; Muramatsu, T.; Kajii, T.: Midkine gene (MDK), a gene for pre-natal differentiation and neuroregulation, maps to band 11p11.2 by fluorescence in situ hybridization. *Genomics* 17: 514–515, 1993.

[64401] 2731. Simon-Chazottes, D.; Matsubara, S.; Miyauchi, T.; Muramatsu, T.; Guenet, J.-L.: Chromosomal localization of two cell surface-associated molecules of potential importance in development: midkine (Mdk) and basigin (Bsg). *Mammalian Genome* 2: 269–271, 1992.

[64402] 2732. Cool, D. R.; Normant, E.; Shen, F.; Chen, H.-C.; Pannell, L.; Zhang, Y.; Loh, Y. P.: Carboxypeptidase E is a regulated secretory pathway sorting receptor: genetic obliteration leads to endocrine disorders in Cpe(fat) mice. *Cell*

88: 73–83, 1997.

- [64403] 2733.Beg, A. A.; Sha, W. C.; Bronson, R. T.; Ghosh, S.; Baltimore, D.: Embryonic lethality and liver degeneration in mice lacking the RelA component of NF-kappa B. *Nature* 376: 167–170, 1995.
- [64404] 2734.Chen, L.; Fischle, W.; Verdin, E.; Greene, W. C.: Duration of nuclear NF-kappa-B action regulated by reversible acetylation. *Science* 293:1653–1657, 2001.
- [64405] 2735.Deloukas, P.; Dauwerse, J. G.; van Ommen, G. J. B.; van Loon, A.P. G. M.: The human NFKB3 gene encoding the p65 subunit of transcription factor NF-kappa-B is located on chromosome 11q12. *Genomics* 19: 592–594, 1994.
- [64406] 2736.Deloukas, P.; van Loon, A. P. G. M.: Genomic organization of the gene encoding the p65 subunit of NF-kappa-B: multiple variants of the p65 protein may be generated by alternative splicing. *Hum. Molec.Genet.* 2: 1895–1900, 1993.
- [64407] 2737.Neurath, M. F.; Pettersson, S.; Myer zum Buschenfelde, K.-H.; Strober, W.: Local administration of antisense phosphorothioate oligonucleotides to the p65 subunit of NF-kappa-B abrogates established experimental colitis in mice. *Nature Med.* 2: 998–1004, 1996.

- [64408] 2738.Zhong, H.; Voll, R. E.; Ghosh, S.: Phosphorylation of NF-kappaB by PKA stimulates transcriptional activity by promoting a novelbivalent interaction with the coactivator CBP/p300. *Molec. Cell* 1:661–671, 1998.
- [64409] 2739.Biamonti, G.; Buvoli, M.; Bassi, M. T.; Morandi, C.; Cobianchi,F.; Riva, S.: Isolation of an active gene encoding human hnRNP proteinA1: evidence for alternative splicing. *J. Molec. Biol.* 207: 491–503,1989.
- [64410] 2740.Buvoli, M.; Biamonti, G.; Tsoulfas, P.; Bassi, M. T.; Ghetti, A.;Riva, S.; Morandi, C.: cDNA cloning of human hnRNP protein A1 revealsthe existence of multiple mRNA isoforms. *Nucleic Acids Res.* 16:3751–3770, 1988.
- [64411] 2741.Michael, W. M.; Choi, M.; Dreyfuss, G.: A nuclear export signalin hnRNP A1: a signal-mediated, temperature-dependent nuclear proteinexport pathway. *Cell* 83: 415–422, 1995.
- [64412] 2742.Saccone, S.; Biamonti, G.; Maugeri, S.; Bassi, M. T.; Bunone, G.;Riva, S.; Della Valle, G.: Assignment of the human heterogeneousnuclear ribonucleoprotein A1 gene (HNRPA1) to chromosome 12q13.1 bycDNA competitive in situ hybridization. *Genomics* 12: 171–174, 1992.
- [64413] 2743.Pollard, A. J.; Sparey, C.; Robson, S. C.; Krainer, A. R.; Europe-Finner,G. N.: Spatio-temporal expression of



the trans-acting splicing factors SF2/ASF and heterogeneous ribonuclear proteins A1/A1B in the myometrium of the pregnant human uterus: a molecular mechanism for regulating regional protein isoform expression in vivo. *J. Clin. Endocr. Metab.* 85:1928–1936, 2000.

- [64414] 2744. Nicoloso, M.; Caizergues-Ferrer, M.; Michot, B.; Azum, M.-C.; Bachellerie, J.-P.: U20, a novel small nucleolar RNA, is encoded in an intron of the nucleolin gene in mammals. *Molec. Cell. Biol.* 14: 5766–5776, 1994.
- [64415] 2745. Srivastava, M.; Fleming, P. J.; Pollard, H. B.; Burns, A. L.: Cloning and sequencing of the human nucleolin cDNA. *FEBS Lett.* 250:99–105, 1989.
- [64416] 2746. Srivastava, M.; McBride, O. W.; Fleming, P. J.; Pollard, H. B.; Burns, A. L.: Genomic organization and chromosomal localization of the human nucleolin gene. *J. Biol. Chem.* 265: 14922–14931, 1990.
- [64417] 2747. Turc-Carel, C.; Pietrzak, E.; Kakati, S.; Kinniburgh, A. J.; Sandberg, A. A.: INT1 maps to 12q12–12q13. (Abstract) *Cytogenet. Cell Genet.* 46:706 only, 1987.
- [64418] 2748. van't Veer, L. J.; Geurts van Kessel, A. H. M.; van Heerikhuizen, H.; van Ooyen, A.; Nusse, R.: Molecular cloning and chromosomal assignment of the human homolog of int-1, a mouse gene implicated in mammary tu-

morigenesis. *Molec. Cell. Biol.* 4: 2532–2534, 1984.

[64419] 2749. van Ooyen, A.; Kwee, V.; Nusse, R.: The nucleotide sequence of the human int-1 mammary oncogene; evolutionary conservation of coding and non-coding sequences. *EMBO J.* 4: 2905–2909, 1985.

[64420] 2750. Wolda, S. L.; Moon, R. T.: Cloning and developmental expression in *Xenopus laevis* of seven additional members of the Wnt family. *Oncogene* 7:1941–1947, 1992.

[64421] 2751. Satijn, D. P. E.; Gunster, M. J.; van der Vlag, J.; Hamer, K. M.; Schul, W.; Alkema, M. J.; Saurin, A. J.; Freemont, P. S.; van Driel, R.; Otte, A. P.: RING1 is associated with the polycomb group protein complex and acts as a transcriptional repressor. *Molec. Cell. Biol.* 17:4105–4113, 1997.

[64422] 2752. Armstrong, B. C.; Krystal, G. W.: Isolation and characterization of complementary DNA for N-cym, a gene encoded by the DNA strand opposite to N-myc. *Cell Growth Differ.* 3: 385–390, 1992.

[64423] 2753. Brodeur, G. M.; Seeger, R. C.: Gene amplification in human neuroblastomas: basic mechanisms and clinical implications. *Cancer Genet. Cytogenet.* 19:101–111, 1986.

[64424] 2754. Brodeur, G. M.; Seeger, R. C.; Schwab, M.; Varmus, H. E.; Bishop, J. M.: Amplification of N-myc in untreated

human neuroblastomas correlates with advanced disease stage. *Science* 224: 1121–1124, 1984.

[64425] 2755. Campbell, G. R.; Zimmerman, K.; Blank, R. D.; Alt, F. W.; D'Eustachio, P.: Chromosomal location of N-myc and L-myc genes in the mouse. *Oncogene Res.* 4: 47–54, 1989.

[64426] 2756. Corvi, R.; Amler, L. C.; Savelyeva, L.; Gehring, M.; Schwab, M.: MYCN is retained in single copy at chromosome 2 band p23–24 during amplification in human neuroblastoma cells. *Proc. Nat. Acad. Sci.* 91:5523–5527, 1994.

[64427] 2757. Emanuel, B. S.; Balaban, G.; Boyd, J. P.; Grossman, A.; Negishi, M.; Parmiter, A.; Glick, M. C.: N-myc amplification in multiple homogeneously staining regions in two human neuroblastomas. *Proc. Nat. Acad. Sci.* 82:3736–3740, 1985.

[64428] 2758. Garson, J. A.; van den Berghe, J. A.; Kemshead, J. T.: Novel non-isotopic in situ hybridization technique detects small (1 kb) unique sequences in routinely G-banded human chromosomes: fine mapping of N-myc and beta-NGF genes. *Nucleic Acids Res.* 15: 4761–4770, 1987.

[64429] 2759. Guo, C.; White, P. S.; Weiss, M. J.; Hogarty, M. D.; Thompson, P. M.; Stram, D. O.; Gerbing, R.; Matthay, K. K.;

Seeger, R. C.; Brodeur, G. M.; Maris, J. M.: Allelic deletion at 11q23 is common in MYCN single copy neuroblastomas. *Oncogene* 18: 4948–4957, 1999.

[64430] 2760. Kanda, N.; Schreck, R.; Alt, F.; Bruns, G.; Baltimore, D.; Latt, S.: Isolation of amplified DNA sequences from IMR-32 human neuroblastoma cells: facilitation by fluorescence-activated flow sorting of metaphase chromosomes. *Proc. Nat. Acad. Sci.* 80: 4069–4073, 1983.

[64431] 2761. Kohl, N. E.; Gee, C. E.; Alt, F. W.: Activated expression of the N-myc gene in human neuroblastomas and related tumors. *Science* 226: 1335–1337, 1984.

[64432] 2762. Kohl, N. E.; Kanda, N.; Schreck, R. R.; Bruns, G.; Latt, S. A.; Gilbert, F.; Alt, F. W.: Transposition and amplification of oncogene-related sequences in human neuroblastomas. *Cell* 35: 359–367, 1983.

[64433] 2763. Kohl, N. E.; Legouy, E.; DePinho, R. A.; Nisen, P. D.; Smith, R. K.; Gee, C. E.; Alt, F. W.: Human N-myc is closely related in organization and nucleotide sequence to c-myc. *Nature* 319: 73–77, 1986.

[64434] 2764. Lee, W.-H.; Murphree, A. L.; Benedict, W. F.: Expression and amplification of the N-myc gene in primary retinoblastoma. *Nature* 309: 458–460, 1984.

[64435] 2765. Michitsch, R. W.; Melera, P. W.: Nucleotide sequence

of the 3-prime exon of the human N-myc gene. *Nucleic Acids Res.* 13: 2545–2558, 1985.

- [64436] 2766. Ramsay, G.; Stanton, L.; Schwab, M.; Bishop, J. M.: Human proto-oncogene N-myc encodes nuclear proteins that bind DNA. *Molec. Cell. Biol.* 6:4450–4457, 1986.
- [64437] 2767. Reiter, J. L.; Brodeur, G. M.: High-resolution mapping of a 130-kb core region of the MYCN amplicon in neuroblastomas. *Genomics* 32:97–103, 1996.
- [64438] 2768. Schwab, M.: Amplification of N-myc in human neuroblastomas. *Trends Genet.* 1: 271–275, 1985.
- [64439] 2769. Schwab, M.; Alitalo, K.; Klempnauer, K.-H.; Varmus, H. E.; Bishop, J. M.; Gilbert, F.; Brodeur, G.; Goldstein, M.; Trent, J.: Amplified DNA with limited homology to myc cellular oncogene is shared by human neuroblastoma cell lines and a neuroblastoma tumour. *Nature* 305:245–248, 1983.
- [64440] 2770. Schwab, M.; Ellison, J.; Busch, M.; Rosenau, W.; Varmus, H. E.; Bishop, J. M.: Enhanced expression of the human gene N-myc consequent to amplification of DNA may contribute to malignant progression of neuroblastoma. *Proc. Nat. Acad. Sci.* 81: 4940–4944, 1984.
- [64441] 2771. Schwab, M.; Varmus, H. E.; Bishop, J. M.; Grzeschik, K.-H.; Naylor, S. L.; Sakaguchi, A. Y.; Brodeur, G.; Trent, J.:

Chromosome localization in normal human cells and neuroblastomas of a gene related to c-myc. *Nature* 308:288–291, 1984.

[64442] 2772. Seeger, R. C.; Brodeur, G. M.; Sather, H.; Dalton, A.; Siegel, S. E.; Wong, K. Y.; Hammond, D.: Association of multiple copies of the N-myc oncogene with rapid progression of neuroblastomas. *New Eng. J. Med.* 313: 1111–1116, 1985.

[64443] 2773. Shiloh, Y.; Shipley, J.; Brodeur, G. M.; Bruns, G.; Korf, B.; Donlon, T.; Schreck, R. R.; Seeger, R.; Sakai, K.; Latt, S. A.: Differential amplification, assembly, and relocation of multiple DNA sequences in human neuroblastomas and neuroblastoma cell lines. *Proc. Nat. Acad. Sci.* 82: 3761–3765, 1985.

[64444] 2774. Bahou, W. F.; Nierman, W. C.; Durkin, A. S.; Potter, C. L.; Demetrick, D. J.: Chromosomal assignment of the human thrombin receptor gene: localization to region q13 of chromosome 5. *Blood* 82: 1532–1537, 1993.

[64445] 2775. Coughlin, S. R.; Vu, T.-K. H.; Hung, D. T.; Wheaton, V. I.: Characterization of a functional thrombin receptor: issues and opportunities. *J. Clin. Invest.* 89: 351–355, 1992.

[64446] 2776. Griffin, C. T.; Srinivasan, Y.; Zheng, Y.-W.; Huang,

W.; Coughlin, S. R.: A role for thrombin receptor signaling in endothelial cells during embryonic development. *Science* 293: 1666–1670, 2001.

- [64447] 2777. Poirier, C.; O'Brien, E. P.; Bueno Brunialti, A. L.; Chambard, J.-C.; Swank, R. T.; Guenet, J.-L.: The gene encoding the thrombin receptor (Cf2r) maps to mouse chromosome 13. *Mammalian Genome* 7:322, 1996.
- [64448] 2778. Schmidt, V. A.; Nierman, W. C.; Feldblyum, T. V.; Maglott, D. R.; Bahou, W. F.: The human thrombin receptor and proteinase activated receptor-2 genes are tightly linked on chromosome 5q13. *Brit. J. Haemat.* 97: 523–529, 1997.
- [64449] 2779. Schmidt, V. A.; Vitale, E.; Bahou, W. F.: Genomic cloning and characterization of the human thrombin receptor gene: structural similarity to the proteinase activated receptor-2 gene. *J. Biol. Chem.* 271:9307–9312, 1996.
- [64450] 2780. Vu, T.-K. H.; Hung, D. T.; Wheaton, V. I.; Coughlin, S. R.: Molecular cloning of a functional thrombin receptor reveals a novel proteolytic mechanism of receptor activation. *Cell* 64: 1057–1068, 1991.
- [64451] 2781. Dean, G.: *The Porphyrrias. A Story of Inheritance and Environment.* Philadelphia: J. B. Lippincott (pub.) (2nd ed.):

1972.

- [64452] 2782. Bennett, M. K.; Calakos, N.; Scheller, R. H.: Syntaxin: a synaptic protein implicated in docking of synaptic vesicles at presynaptic active zones. *Science* 257: 255–259, 1992.
- [64453] 2783. Bennett, M. K.; Garcia-Arreaga, J. E.; Elferink, L. A.; Peterson, K.; Fleming, A. M.; Hazuka, C. D.; Scheller, R. H.: The syntaxin family of vesicular transport receptors. *Cell* 74: 863–873, 1993.
- [64454] 2784. Fernandez, I.; Ubach, J.; Dulubova, I.; Zhang, X.; Sudof, T. C.; Rizo, J.: Three-dimensional structure of an evolutionarily conserved N-terminal domain of syntaxin 1A. *Cell* 94: 841–849, 1998.
- [64455] 2785. Fisher, R. J.; Pevsner, J.; Burgoyne, R. D.: Control of fusion pore dynamics during exocytosis by Munc18. *Science* 291: 875–878, 2001.
- [64456] 2786. Martin-Martin, B.; Nabokina, S. M.; Lazo, P. A.; Mollinedo, F.: Co-expression of several human syntaxin genes in neutrophils and differentiating HL-60 cells: various isoforms and detection of syntaxin 1. *J. Leuko. Biol.* 65: 397–406, 1999.
- [64457] 2787. Nakayama, T.; Fujiwara, T.; Miyazawa, A.; Asakawa, S.; Shimizu, N.; Shimizu, Y.; Mikoshiba, K.; Akagawa, K.:



Mapping of the humanHPC-1/syntaxin 1A gene (STX1A) to chromosome 7 band q11.2. Genomics 42:173–176, 1997.

- [64458] 2788.Naren, A. P.; Di, A.; Cormet-Boyaka, E.; Boyaka, P. N.; McGhee,J. R.; Zhou, W.; Akagawa, K.; Fujiwara, T.; Thome, U.; Engelhardt,J. F.; Nelson, D. J.; Kirk, K. L.: Syntaxin 1A is expressed in airwayepithelial cells, where it modulates CFTR C1- currents. J. Clin.Invest. 105: 377–386, 2000.
- [64459] 2789.Richmond, J. E.; Weimer, R. M.; Jorgensen, E. M.: An open formof syntaxin bypasses the requirement for UNC-13 in vesicle priming. Nature 412:338–341, 2001.
- [64460] 2790.Zhang, R.; Maksymowych, A. B.; Simpson, L. L.: Cloning and sequenceanalysis of a cDNA encoding human syntaxin 1A, a polypeptide essentialfor exocytosis. Gene 159: 293–294, 1995.
- [64461] 2791.Shepley, M. P.; Sherry, B.; Weiner, H. L.: Monoclonal antibodyidentification of a 100-kDa membrane protein in HeLa cells and humanspinal cord involved in poliovirus attachment. Proc. Nat. Acad. Sci. 85:7743–7747, 1988.
- [64462] 2792.Siddique, T.; Bartlett, R. J.; McKinney, R.; Hung, W.-Y.; Bruns,G.; Wilfert, C.; Roses, A. D.: The poliovirus sensitivity (PVS) ison chromosome 19q13-qter. (Abstract)

Cytogenet. Cell Genet. 40:745 only, 1985.

- [64463] 2793.Siddique, T.; McKinney, R.; Hung, W.-Y.; Bartlett, R. J.; Bruns,G.; Mohandas, T. K.; Ropers, H.-H.; Wilfert, C.; Roses, A. D.: Thepoliovirus sensitivity (PVS) gene is on chromosome 19q12-q13.2. Genomics 3:156-160, 1988.
- [64464] 2794.Solecki, D. J.; Gromeier, M.; Mueller, S.; Bernhardt, G.; Wimmer,E.: Expression of the human poliovirus receptor/CD155 gene is activatedby Sonic hedgehog. J. Biol. Chem. 277: 25697-25702, 2002.
- [64465] 2795.Morris, C. M.; Bodger, M. P.: Localization of the human poly(A)-bindingprotein gene (PAB1) to chromosomal regions 3q22-q25, 12q13-q14, and13q12-q13 by in situ hybridization. Genomics 15: 209-211, 1993.
- [64466] 2796.Alkhatib, H. M.; Chen, D.; Cherney, B.; Bhatia, K.; Notario, V.;Giri, C.; Stein, G.; Slattey, E.; Roeder, R. G.; Smulson, M. E.:Cloning and expression of cDNA for human poly(ADP-ribose) polymerase. Proc.Nat. Acad. Sci. 84: 1224-1228, 1987.
- [64467] 2797.Auer, B.; Nagl, U.; Herzog, H.; Schneider, R.; Schweiger, M.:Human nuclear NAD+ ADP-riboseyltransferase(polymerizing): organizationof the gene. DNA 8: 575-580, 1989.
- [64468] 2798.Baumgartner, M.; Schneider, R.; Auer, B.; Herzog, H.;

Schweiger, M.; Hirsch-Kauffmann, M.: Fluorescence in situ mapping of the human nuclear NAD<sup>+</sup> ADP-ribosyltransferase gene (ADPRT) and two secondary sites to human chromosomal bands 1q42, 13q34, and 14q24. *Cytogenet. Cell Genet.* 61: 172–174, 1992.

[64469] 2799. Bhatia, K. G.; Cherney, B. W.; Huppi, K.; Magrath, I. T.; Cossman, J.; Sausville, E.; Barriga, F.; Johnson, B.; Gause, B.; Bonney, G.; Neequayi, J.; DeBernardi, M.; Smulson, M.: A deletion linked to a poly(ADP-ribose) polymerase gene on chromosome 3q33-qter occurs frequently in the normal black population as well as in multiple tumor DNA. *Cancer Res.* 50: 5406–5413, 1990.

[64470] 2800. Cherney, B. W.; McBride, O. W.; Chen, D.; Alkhatib, H.; Bhatia, K.; Hensley, P.; Smulson, M. E.: cDNA sequence, protein structure, and chromosomal location of the human gene for poly(ADP-ribose) polymerase. *Proc. Nat. Acad. Sci.* 84: 8370–8374, 1987.

[64471] 2801. d'Adda di Fagagna, F.; Hande, M. P.; Tong, W.-M.; Lansdorp, P. M.; Wang, Z.-Q.; Jackson, S. P.: Functions of poly(ADP-ribose) polymerase in controlling telomere length and chromosomal stability. *Nature Genet.* 23: 76–80, 1999.

[64472] 2802. Doll, J. A.; Suarez, B. K.; Donis-Keller, H.: Associa-

tion between prostate cancer in black Americans and an allele of the PADPRP pseudogene locus on chromosome 13. (Letter) Am. J. Hum. Genet. 58: 425–428, 1996.

[64473] 2803. Flick, K.; Schneider, R.; Auer, B.; Hirsch-Kauffmann, M.; Schweiger, M.: No abnormalities in the NAD(+) ADP-ribosyltransferase (polymerizing) gene of transformed cells from a Fanconi's anemia patient. (Letter) Hum. Genet. 89: 690–691, 1992.

[64474] 2804. Grube, K.; Burkle, A.: Poly(ADP-ribose) polymerase activity in mononuclear leukocytes of 13 mammalian species correlates with species-specific life span. Proc. Nat. Acad. Sci. 89: 11759–11763, 1992.

[64475] 2805. Herzog, H.; Zabel, B.; Schneider, R.; Auer, B.; Hirsch-Kauffmann, M.; Schweiger, M.: Human NAD(+):protein ADP ribosyltransferase (ADPRT): synthesis of active enzyme in E. coli, sequence of its cDNA from HeLa cells and chromosomal localization. Biol. Chem. Hoppe-Seyler 369: 836–837, 1988.

[64476] 2806. Herzog, H.; Zabel, B. U.; Schneider, R.; Auer, B.; Hirsch-Kauffmann, M.; Schweiger, M.: Human nuclear NAD(+) ADP-ribosyltransferase: localization of the gene on chromosome 1q41–q42 and expression of an active human enzyme in Escherichia coli. Proc. Nat. Acad. Sci. 86:

3514–3518,1989.

- [64477] 2807.Keijzer, W.; Stefanini, M.; Bootsma, D.; Verkerk, A.; Geurts vanKessel, A. H. M.; Jongkind, J. F.; Westerveld, A.: Localization of a gene involved in complementation of the defect in xeroderma pigmentosum group A cells on human chromosome 1. *Exp. Cell Res.* 169: 490–501,1987.
- [64478] 2808.Kurosaki, T.; Ushiro, H.; Mitsuuchi, Y.; Suzuki, S.; Matsuda, M.; Matsuda, Y.; Katunuma, N.; Kangawa, K.; Matsuo, H.; Hirose, T.; Inayama, S.; Shizuta, Y.: Primary structure of human poly(ADP–ribose)synthetase as deduced from cDNA sequence. *J. Biol. Chem.* 262: 15990–15997,1987.
- [64479] 2809.Loetscher, P.; Alvarez–Gonzalez, R.; Althaus, F. R.: Poly(ADP–ribose) may signal changing metabolic conditions to the chromatin of mammalian cells. *Proc. Nat. Acad. Sci.* 84: 1286–1289, 1987.
- [64480] 2810.Lyn, D.; Cherney, B. W.; Lalande, M.; Berenson, J. R.; Lichtenstein, A.; Lupold, S.; Bhatia, K. G.; Smulson, M.: A duplicated region is responsible for the poly(ADP–ribose) polymerase polymorphism, on chromosome 13, associated with a predisposition to cancer. *Am. J. Hum. Genet.* 52:124–134, 1993.
- [64481] 2811.McBride, O. W.; Cherney, B.; Takourdin, C.; Smulson,

M. E.: Humanpoly(ADP-ribose) polymerase sequences are located on chromosomes 1,13, and 14. (Abstract) Cytogenet. Cell Genet. 46: 659-660, 1987.

[64482] 2812.Pieper, A. A.; Brat, D. J.; Krug, D. K.; Watkins, C. C.; Gupta,A.; Blackshaw, S.; Verma, A.; Wang, Z.-Q.; Snyder, S. H.: Poly(ADP-ribose)polymerase-deficient mice are protected from streptozotocin-induceddiabetes. Proc. Nat. Acad. Sci. 96: 3059-3064, 1999.

[64483] 2813.Schweiger, M.; Auer, B.; Burtscher, H. J.; Hirsch-Kauffmann, M.;Klocker, H.; Schneider, R.: DNA repair in human cells: biochemistryof the hereditary diseases Fanconi's anaemia and Cockayne syndrome. Europ.J. Biochem. 165: 235-242, 1987.

[64484] 2814.Simbulan-Rosenthal, C. M.; Haddad, B. R.; Rosenthal, D. S.; Weaver,Z.; Coleman, A.; Luo, R.; Young, H. M.; Wang, Z.-Q.; Ried, T.; Smulson,M. E.: Chromosomal aberrations in PARP -/- mice: genome stabilizationin immortalized cells by reintroduction of poly(ADP-ribose) polymerasecDNA. Proc. Nat. Acad. Sci. 96: 13191-13196, 1999.

[64485] 2815.Smithies, O.; Gregg, R. G.; Boggs, S. S.; Koralewski, M. A.; Kucherlapati,R. S. :Nature (London) 317: 230-234, 1985.

- [64486] 2816.Bravo, R.: Synthesis of the nuclear protein cyclin (PCNA) and its relationship with DNA replication. *Exp. Cell Res.* 163: 287–293, 1986.
- [64487] 2817.Hasan, S.; Hassa, P. O.; Imhof, R.; Hottiger, M. O.: Transcription coactivator p300 binds PCNA and may have a role in DNA repair synthesis. *Nature* 410:387–391, 2001.
- [64488] 2818.Hoege, C.; Pfander, B.; Moldovan, G.–L.; Pyrowolakis, G.; Jentsch, S.: RAD6–dependent DNA repair is linked to modification of PCNA by ubiquitin and SUMO. *Nature* 419: 135–141, 2002.
- [64489] 2819.Ku, D.–H.; Travali, S.; Calabretta, B.; Huebner, K.; Baserga, R.: Human gene for proliferating cell nuclear antigen has pseudogenes and localizes to chromosome 20. *Somat. Cell Molec. Genet.* 15: 297–307, 1989.
- [64490] 2820.Mann, M. J.; Gibbons, G. H.; Kernoff, R. S.; Diet, F. P.; Tsao, P. S.; Cooke, J. P.; Kaneda, Y.; Dza, V. J.: Genetic engineering of vein grafts resistant to atherosclerosis. *Proc. Nat. Acad. Sci.* 92:4502–4506, 1995.
- [64491] 2821.Rao, V. V. N. G.; Schnittger, S.; Hansmann, I.: Chromosomal localization of the human proliferating cell nuclear antigen (PCNA) gene to or close to 20p12 by in situ hybridization. *Cytogenet. Cell Genet.* 56:169–170, 1991.
- [64492] 2822.Suzuka, I.; Daidoji, H.; Matsuoka, M.; Kadowaki, K.;

Takasaki,Y.; Nakane, P. K.; Moriuchi, T.: Gene for proliferating-cell nuclearantigen (DNA polymerase delta auxiliary protein) is present in bothmammalian and higher plant genomes. Proc. Nat. Acad. Sci. 86: 3189–3193,1989.

[64493] 2823.Taniguchi, Y.; Katsumata, Y.; Koido, S.; Suemizu, H.; Yoshimura,S.; Moriuchi, T.; Okumura, K.; Kagotani, K.; Taguchi, H.; Imanishi,T.; Gojobori, T.; Inoko, H.: Cloning, sequencing, and chromosomallocalization of two tandemly arranged human pseudogenes for the proliferatingcell nuclear antigen (PCNA). Mammalian Genome 7: 906–908, 1996.

[64494] 2824.Travali, S.; Ku, D.–H.; Rizzo, M. G.; Ottavio, L.; Baserga, R.;Calabretta, B.: Structure of the human gene for the proliferatingcell nuclear antigen. J. Biol. Chem. 264: 7466–7472, 1989.

[64495] 2825.Webb, G.; Parsons, P.; Chenevix–Trench, G.: Localization of thegene for human proliferating nuclear antigen/cyclin by in situ hybridization. Hum.Genet. 86: 84–86, 1990.

[64496] 2826.Fonatsch, C.; Duchrow, M.; Rieder, H.; Schluter, C.; Gerdes, J.: Assignment of the human Ki–67 gene (MKI67) to 10q25–qter. Genomics 11:476–477, 1991.

[64497] 2827.Schluter, C.; Duchrow, M.; Wohlenberg, C.; Becker,



M. H. G; Key,G.; Flad, H.-D.; Gerdes, J.: The cell proliferation-associated antigen of antibody Ki-67: a very large, ubiquitous nuclear protein with numerous repeated elements, representing a new kind of cell cycle-maintaining proteins. J. Cell. Biol. 123: 513-522, 1993.

[64498] 2828.Schonk, D. M.; Kuijpers, H. J. H.; van Drunen, E.; van Dalen, C.H.; Geurts van Kessel, A. H. M.; Verheijen, R.; Ramaekers, F. C. S.: Assignment of the gene(s) involved in the expression of the proliferation-related Ki-67 antigen to human chromosome 10. Hum. Genet. 83: 297-299, 1989.

[64499] 2829.Traut, W.; Scholzen, T.; Winking, H.; Kubbutat, M. H. G.; Gerdes,J.: Assignment of the murine Ki-67 gene (Mki67) to chromosome band 7F3-F5 by in situ hybridization. Cytogenet. Cell Genet. 83: 12-13,1998.

[64500] 2830.Cooke, N. E.; Baxter, J. D.: Structural analysis of the prolactin gene suggests a separate origin for its 5-prime end. Nature 297:603-606, 1982.

[64501] 2831.Cooke, N. E.; Coit, D.; Shine, J.; Baxter, J. D.; Martial, J. A.: Human prolactin: cDNA structural analysis and evolutionary comparisons. J.Biol. Chem. 256: 4007-4016, 1981.

[64502] 2832.D'Alessandro, E.; Santiemma, V.; Lo Re, M. L.; Ligas,

C.; Del Porto, G.: 6p23 deletion mosaicism in a woman with recurrent abortions and idiopathic hypoprolactinemia. *Am. J. Med. Genet.* 44: 220–222, 1992.

[64503] 2833. DiMattia, G. E.: Personal Communication. London, Ontario, Canada 3/23/1998.

[64504] 2834. DiMattia, G. E.; Gellersen, B.; Duckworth, M. L.; Friesen, H. G.: Human prolactin gene expression: the use of an alternative noncoding exon in decidua and the IM-9-P3 lymphoblast cell line. *J. Biol. Chem.* 265:16412–16421, 1990.

[64505] 2835. Evans, A. M.; Petersen, J. W.; Sekhon, G. S.; DeMars, R. I.: Use of human lymphoblastoid deletion mutants to map the prolactin gene on human chromosome 6p. (Abstract) *Am. J. Hum. Genet.* 43: A143 only, 1988.

[64506] 2836. Farid, N. R.; Noel, E. P.; Sampson, L.; Russell, N. A.: Prolactin-secreting adenomata are possibly associated with HLA-B8. *Tissue Antigens* 15:333–335, 1980.

[64507] 2837. Gellersen, B.; Kempf, R.; Telgmann, R.; DiMattia, G. E.: Nonpituitary human prolactin gene transcription is independent of Pit-1 and differentially controlled in lymphocytes and in endometrial stroma. *Molec. Endocr.* 8:356–373, 1994.

[64508] 2838. Erdile, L. F.; Heyer, W.-D.; Kolodner, R.; Kelly, T. J.:

Characterization of a cDNA encoding the 70-kDa single-stranded DNA-binding subunit of human replication protein A and the role of the protein in DNA replication. *J. Biol. Chem.* 266: 12090–12098, 1991.

[64509] 2839. Gomes, X. V.; Wold, M. S.: Functional domains of the 70-kilodalton subunit of human replication protein A. *Biochemistry* 35: 10558–10568, 1996.

[64510] 2840. Shen, L. X.; Basilion, J. P.; Stanton, V. P., Jr.: Single-nucleotide polymorphisms can cause different structural folds of mRNA. *Proc. Nat. Acad. Sci.* 96: 7871–7876, 1999.

[64511] 2841. Umbricht, C. B.; Erdile, L. F.; Jabs, E. W.; Kelly, T. J.: Cloning, overexpression, and genomic mapping of the 14-kDa subunit of human replication protein A. *J. Biol. Chem.* 268: 6131–6138, 1993.

[64512] 2842. Umbricht, C. B.; Griffin, C. A.; Hawkins, A. L.; Grzeschik, K. H.; O'Connell, P.; Leach, R.; Green, E. D.; Kelly, T. J.: High-resolution genomic mapping of the three human replication protein A genes (RPA1, RPA2, and RPA3). *Genomics* 20: 249–257, 1994.

[64513] 2843. Bilbe, G.; Delabie, J.; Bruggen, J.; Richener, H.; Asselbergs, F. A. M.; Cerletti, N.; Sorg, C.; Odink, K.; Tarcsay, L.; Wiesendanger, W.; DeWolf-Peeters, C.; Shipman, R.: Restin: a novel intermediate filament-associated protein

highly expressed in the Reed–Sternberg cells of Hodgkin's disease. *EMBO J.* 11: 2103–2113, 1992.

[64514] 2844. Delabie, J.; Shipman, R.; Bruggen, J.; De Strooper, B.; van Leuven, F.; Tarcsay, L.; Cerletti, N.; Odink, K.; Diehl, V.; Bilbe, G.; DeWolf–Peeters, C.: Expression of the novel intermediate filament–associated protein restin in Hodgkin's disease and anaplastic large–cell lymphoma. *Blood* 80:2891–2896, 1992.

[64515] 2845. Fukata, M.; Watanabe, T.; Noritake, J.; Nakagawa, M.; Yamaga, M.; Kuroda, S.; Matsuura, Y.; Iwamatsu, A.; Perez, F.; Kaibuchi, K.: Rac1 and Cdc42 capture microtubules through IQGAP1 and CLIP–170. *Cell* 109:873–885, 2002.

[64516] 2846. Griparic, L.; Keller, T. C. S., III: Identification and expression of two novel CLIP–170/restin isoforms expressed predominantly in muscle. *Biochim. Biophys. Acta* 1405: 35–46, 1998.

[64517] 2847. Hilliker, C.; Delabie, J.; Speleman, F.; Bilbe, G.; Bruggen, J.; Van Leuven, F.; Van Den Berghe, H.: Localization of the gene (RSN) coding for restin, a marker for Reed–Sternberg cells in Hodgkin's disease, to human chromosome band 12q24.3 and YAC cloning of the locus. *Cytogenet. Cell Genet.* 65: 172–176, 1994.

[64518] 2848. Durbin, H.; Rodrigues, N.; Bodmer, W. F.: Further

characterization, isolation and identification of the epithelial cell-surface antigen defined by monoclonal antibody AUAI. *Int. J. Cancer* 45: 562-565, 1990.

- [64519] 2849. Meyaard, L.; van der Vuurst de Vries, A.-R.; de Ruiter, T.; Lanier, L. L.; Phillips, J. H.; Clevers, H.: The epithelial cellular adhesion molecule (Ep-CAM) is a ligand for the leukocyte-associated immunoglobulin-like receptor (LAIR). *J. Exp. Med.* 194: 107-112, 2001.
- [64520] 2850. Perez, M. S.; Walker, L. E.: Isolation and characterization of a cDNA encoding the KS1/4 epithelial carcinoma marker. *J. Immun.* 142:3662-3667, 1989.
- [64521] 2851. Spurr, N. K.; Durbin, H.; Sheer, D.; Parkar, M.; Bobrow, L.; Bodmer, W. F.: Characterization and chromosomal assignment of a human cell surface antigen defined by the monoclonal antibody AUAI. *Int. J. Cancer* 38: 631-636, 1986.
- [64522] 2852. Strnad, J.; Hamilton, A. E.; Beavers, L. S.; Gamboa, G. C.; Apelgren, L. D.; Taber, L. D.; Sportsman, J. R.; Bumol, T. F.; Sharp, J. D.; Gadski, R. A.: Molecular cloning and characterization of a human adenocarcinoma/epithelial cell surface antigen complementary DNA. *Cancer Res.* 49: 314-317, 1989.
- [64523] 2853. Szala, S.; Froehlich, M.; Scollon, M.; Kasai, Y.;

Steplewski, Z.;Koprowski, H.; Linnenbach, A. J.: Molecular cloning of cDNA for thecarcinoma-associated antigen GA733-2. Proc. Nat. Acad. Sci. 87:3542-3546, 1990.

[64524] 2854.Hitchins, M. P.; Monk, D.; Bell, G. M.; Ali, Z.; Preece, M. A.;Stanier, P.; Moore, G. E.: Maternal repression of the human GRB10gene in the developing central nervous system; evaluation of the rolefor GRB10 in Silver-Russell syndrome. Europ. J. Hum. Genet. 9: 82-90,2001.

[64525] 2855.McCann, J. A.; Zheng, H.; Islam, A.; Goodyer, C. G.; Polychronakos,C.: Evidence against GRB10 as the gene responsible for Silver-Russellsyndrome. Biochem. Biophys. Res. Commun. 286: 943-948, 2001.

[64526] 2856.Miyoshi, N.; Kuroiwa, Y.; Kohda, T.; Shitara, H.; Yonekawa, H.;Kawabe, T.; Hasegawa, H.; Barton, S. C.; Surani, M. A.; Kaneko-Ishino,T.; Ishino, F.: Identification of the Meg1/Grb10 imprinted gene onmouse proximal chromosome 11, a candidate for the Silver-Russell syndrome gene. Proc. Nat. Acad. Sci. 95: 1102-1107, 1998.

[64527] 2857.Yoshihashi, H.; Maeyama, K.; Kosaki, R.; Ogata, T.; Tsukahara,M.; Goto, Y.; Hata, J.; Matsuo, N.; Smith, R. J.; Kosaki, K.: Imprintingof human GRB10 and its mutations in two patients with Russell-Silversyndrome. Am. J. Hum. Genet. 67: 476-482, 2000.

- [64528] 2858.Grob, P. M.; Ross, A. H.; Koprowski, H.; Bothwell, M.: Characterization of the human melanoma nerve growth factor receptor. *J. Biol. Chem.* 260:8044–8049, 1985.
- [64529] 2859.Stanton, L. W.; Schwab, M.; Bishop, J. M.: Nucleotide sequence of the human N-myc gene. *Proc. Nat. Acad. Sci.* 83: 1772–1776, 1986.
- [64530] 2860.Brown, M. A.; Nicolai, H.; Xu, C.-F.; Griffiths, B. L.; Jones, K. A.; Solomon, E.; Hosking, L.; Trowsdale, J.; Black, D. M.; McFarlane, R.: Regulation of BRCA1. (Letter) *Nature* 372: 733 only, 1994.
- [64531] 2861.Brown, M. A.; Xu, C.-F.; Nicolai, H.; Griffiths, B.; Chambers, J. A.; Black, D.; Solomon, E.: The 5-prime end of the BRCA1 gene lies within a duplicated region of human chromosome 17q21. *Oncogene* 12:2507–2513, 1996.
- [64532] 2862.Campbell, I. G.; Nicolai, H. M.; Foulkes, W. D.; Senger, G.; Stamp, G. W.; Allan, G.; Boyer, C.; Jones, K.; Bast, R. C., Jr.; Solomon, E.; Trowsdale, J.; Black, D. M.: A novel gene encoding a B-box protein within the BRCA1 region at 17q21.1. *Hum. Molec. Genet.* 3: 589–594, 1994.
- [64533] 2863.Kawashima, K.; Shikama, H.; Imoto, K.; Izawa, M.; Naruke, T.; Okabayashi, K.; Nishimura, S.: Close correlation between restriction fragment length polymorphism of the

L-MYC gene and metastasis of human lungcancer to the lymph nodes and other organs. Proc. Nat. Acad. Sci. 85:2353–2356, 1988.

[64534] 2864.Kaye, F.; Battey, J.; Nau, M.; Brooks, B.; Seifter, E.; De Greve,J.; Birrer, M.; Sausville, E.; Minna, J.: Structure and expressionof the human L-myc gene reveal a complex pattern of alternative mRNAprocessing. Molec. Cell. Biol. 8: 186–195, 1988.

[64535] 2865.McBride, O. W.; Kirsch, I.; Hollis, G.; Nau, M.; Battey, J.; Minna,J.: Human L-myc (MYCL) proto-oncogene is on chromosome 1p32.(Abstract) Cytogenet.Cell Genet. 40: 694 only, 1985.

[64536] 2866.Nau, M. M.; Brooks, B. J.; Battey, J.; Sausville, E.; Gazdar, A.F.; Kirsch, I. R.; McBride, O. W.; Bertness, V.; Hollis, G. F.; Minna,J. D.: L-myc, a new myc-related gene amplified and expressed in humansmall cell lung cancer. Nature 318: 69–73, 1985.

[64537] 2867.Rouleau, G. A.; Bazanowski, A.; Gusella, J. F.; Haines, J. L.:A genetic map of chromosome 1: comparison of different data sets andlinkage programs. Genomics 7: 313–318, 1990.

[64538] 2868.Speleman, F.; Van Camp, G.; Van Roy, N.: Reassignment of MYCL1to human chromosome 1p34.3 by fluores–



cence in situ hybridization. *Cytogenet. Cell Genet.* 72: 189–190, 1996.

- [64539] 2869. Van Roy, N.; Cheng, N. C.; Laureys, G.; Opdenakker, G.; Versteeg, R.; Speleman, F.: Molecular cytogenetic analysis of 1;17 translocations in neuroblastoma. *Europ. J. Cancer* 31A: 530–535, 1995.
- [64540] 2870. Dean, M.; Park, M.; Le Beau, M. M.; Robins, T. S.; Diaz, M. O.; Rowley, J. D.; Blair, D. G.; Vande Woude, G. F.: The human met oncogene is related to the tyrosine kinase oncogenes. *Nature* 318: 385–388, 1985.
- [64541] 2871. Akiyama, T.; Sudo, C.; Ogawara, H.; Toyoshima, K.; Yamamoto, T.: The product of the human c-erbB-2 gene: a 185-kilodalton glycoprotein with tyrosine kinase activity. *Science* 232: 1644–1646, 1986.
- [64542] 2872. Ameyaw, M.-M.; Tayeb, M.; Thornton, N.; Folayan, G.; Tariq, M.; Mobarek, A.; Evans, D. A. P.; Ofori-Adjei, D.; McLeod, H. L.: Ethnic variation in the HER-2 codon 655 genetic polymorphism previously associated with breast cancer. *J. Hum. Genet.* 47: 172–175, 2002.
- [64543] 2873. Chan, J. Y. C.; Lerman, M. I.; Prabhakar, B. S.; Isozaki, O.; Santisteban, P.; Koppers, R. C.; Oates, E. L.; Notkins, A. L.; Kohn, L. D.: Cloning and characterization of a cDNA that encodes a 70-kDa novel human thyroid au-

toantigen. J. Biol. Chem. 264: 3651–3654, 1989.

- [64544] 2874. Goedecke, W.; Eijpe, M.; Offenberg, H. H.; van Aalderen, M.; Heyting, C.: Mre11 and Ku70 interact in somatic cells, but are differentially expressed in early meiosis. Nature Genet. 23: 194–198, 1999.
- [64545] 2875. Hartley, K. O.; Gell, D.; Smith, G. C. M.; Zhang, H.; Divecha, N.; Connelly, M. A.; Admon, A.; Lees–Miller, S. P.; Anderson, C. W.; Jackson, S. P.: DNA–dependent protein kinase catalytic subunit: a relative of phosphatidylinositol 3–kinase and the ataxia telangiectasia gene product. Cell 82: 849–856, 1995.
- [64546] 2876. Koike, M.; Matsuda, Y.; Mimori, T.; Harada, Y.–N.; Shiomi, N.; Shiomi, T.: Chromosomal localization of the mouse and rat DNA double–strand break repair genes Ku p70 and Ku p80/XRCC5 and their mRNA expression in various mouse tissues. Genomics 38: 38–44, 1996.
- [64547] 2877. Kusano, K.; Johnson–Schlitz, D. M.; Engels, W. R.: Sterility of *Drosophila* with mutations in the Bloom syndrome gene—complementation by Ku70. Science 291: 2600–2602, 2001.
- [64548] 2878. Li, G. C.; Ouyang, H.; Li, X.; Nagasawa, H.; Little, J. B.; Chen, D. J.; Ling, C. C.; Fuks, Z.; Cordon–Cardo, C.: Ku70: a candidate tumor suppressor gene for murine T cell

lymphoma. *Molec. Cell* 2:1–8, 1998.

- [64549] 2879. McBride, O. W.; Chan, J. Y. C.; Notkins, A. L.; Kohn, L. D.; Lerman, M.: The TSH receptor gene is located on human chromosome 22 and homologous sequences are present on chromosomes 1q, 8, 10, and Xq. (Abstract) *Am. J. Hum. Genet.* 41: A177, 1987.
- [64550] 2880. Mitchell, A. L.; Bale, A. E.; Chan, J.; Kohn, L.; Gonzalez, F.; McBride, O. W.: Localization of TSHR gene and cytochrome p450 IID subfamily on chromosome 22 by linkage analysis. (Abstract) *Cytogenet. Cell Genet.* 51: 1045, 1989.
- [64551] 2881. Reeves, W. H.; Stoecker, Z. M.: Molecular cloning of cDNA encoding the p70 (Ku) lupus autoantigen. *J. Biol. Chem.* 264: 5047–5052, 1989.
- [64552] 2882. Takata, M.; Sasaki, M. S.; Sonoda, E.; Morrison, C.; Hashimoto, M.; Utsumi, H.; Yamaguchi-Iwai, Y.; Shinohara, A.; Takeda, S.: Homologous recombination and non-homologous end-joining pathways of DNA double-strand break repair have overlapping roles in the maintenance of chromosomal integrity in vertebrate cells. *EMBO J.* 17: 5497–5508, 1998.
- [64553] 2883. Takiguchi, Y.; Kurimasa, A.; Chen, F.; Pardington, P. E.; Kuriyama, T.; Okinaka, R. T.; Moyzis, R.; Chen, D. J.:

Genomic structure and chromosomal assignment of the mouse Ku70 gene. *Genomics* 35: 129–135, 1996.

[64554] 2884. Tuteja, N.; Tuteja, R.; Ochem, A.; Taneja, P.; Huang, N. W.; Simoncsits, A.; Susic, S.; Rahman, K.; Marusic, L.; Chen, J.; Zhang, J.; Wang, S.; Pongor, S.; Falaschi, A.: Human DNA helicase II: a novel DNA unwinding enzyme identified as the Ku autoantigen. *EMBO J.* 13: 4991–5001, 1994.

[64555] 2885. Walker, J. R.; Corpina, R. A.; Goldberg, J.: Structure of the Ku heterodimer bound to DNA and its implications for double-strand break repair. *Nature* 412: 607–614, 2001.

[64556] 2886. Baer, A. N.; Woosley, R. L.; Pincus, T.: Further evidence for the lack of association between acetylator phenotype and systemic lupus erythematosus. *Arthritis Rheum.* 29: 508–514, 1986.

[64557] 2887. Esteller, M.; Garcia-Foncillas, J.; Andion, E.; Goodman, S. N.; Hidalgo, O. F.; Vanaclocha, V.; Baylin, S. B.; Herman, J. G.: Inactivation of the DNA-repair gene MGMT and the clinical response of gliomas to alkylating agents. *New Eng. J. Med.* 343: 1350–1354, 2000.

[64558] 2888. Gardner, E.; Rydberg, B.; Karran, P.; Ponder, B. A. J.: Localization of the human O(6)-methylguanine-DNA

methyltransferase gene to chromosome10q24.33-qter.

Genomics 11: 475–476, 1991.

[64559] 2889.Natarajan, A. T.; Vermeulen, S.; Darroudi, F.; Valentine, M. B.;Brent, T. P.; Mitra, S.; Tano, K.: Chromosomal localization of humanO(6)–methylguanine–DNA methyltransferase (MGMT) gene by in situ hybridization. Mutagenesis 7:83–85, 1992.

[64560] 2890.Rydberg, B.; Spurr, N.; Karran, P.: cDNA cloning and chromosomalassignment of the human O(6)–methylguanine–DNA methyltransferase:cDNA expression in Escherichia coli and gene expression in human cells. J.Biol. Chem. 265: 9563–9569, 1990.

[64561] 2891.Tano, K.; Shiota, S.; Collier, J.; Foote, R. S.; Mitra, S.: Isolationand structural characterization of a cDNA clone encoding the humanDNA repair protein for O(6)–alkylguanine. Proc. Nat. Acad. Sci. 87:686–690, 1990.

[64562] 2892.Zunino, A.; Arena, G.; Rossi, O.; Archidiacono, N.; Rocchi, M.;Romeo, G.; Abbondandolo, A.: Chromosomal assignment of human O6–methylguanine–DNA–methyltransferasegene by hamster–human somatic cell hybrids. Mutagenesis 6: 395–397,1991.

- [64563] 2893. Van Someren, H.; Westerveld, A.; Hagemeijer, A.; Mees, J. R.; Meera Khan, P.; Zaalberg, O. B.: Human antigen and enzyme markers in man–Chinese hamster somatic cell hybrids: evidence for synteny between the HL–A, PGM–3, ME–1, and IPO–B loci. *Proc. Nat. Acad. Sci.* 71:962–965, 1974.
- [64564] 2894. Yoshida, H.; Mitarai, T.; Kawamura, T.; Kitajima, T.; Miyazaki, Y.; Nagasawa, R.; Kawaguchi, Y.; Kubo, H.; Ichikawa, I.; Sakai, O.: Role of the deletion polymorphism of the angiotensin converting enzyme gene in the progression and therapeutic responsiveness of IgA nephropathy. *J. Clin. Invest.* 96: 2162–2169, 1995.
- [64565] 2895. Zheng, F.; Kundu, G. C.; Zhang, Z.; Ward, J.; DeMayo, F.; Mukherjee, A. B.: Uteroglobin is essential in preventing immunoglobulin A nephropathy in mice. *Nature Med.* 5: 1018–1025, 1999.
- [64566] 2896. Anisowicz, A.; Zajchowski, D.; Stenman, G.; Sager, R.: Functional diversity of GRO gene expression in human fibroblasts and mammary epithelial cells. *Proc. Nat. Acad. Sci.* 85: 9645–9649, 1988.
- [64567] 2897. Horuk, R.; Yansura, D. G.; Reilly, D.; Spencer, S.; Bourell, J.; Henzel, W.; Rice, G.; Unemori, E.: Purification, receptor binding analysis, and biological characterization

of human melanoma growthstimulating activity (MGSA):  
evidence for a novel MGSA receptor. J.Biol. Chem. 268:  
541–546, 1993.

[64568] 2898.Richmond, A.; Balentien, E.; Thomas, H. G.; Flaggs,  
G.; Barton,D. E.; Spiess, J.; Bordoni, R.; Francke, U.;  
Derynck, R.: Molecularcharacterization and chromosomal  
mapping of melanoma growth stimulatoryactivity, a  
growth factor structurally related to beta-  
thromboglobulin. EMBOJ. 7: 2025–2033, 1988.

[64569] 2899.Sakaguchi, A. Y.; Lalley, P. A.; Ghosh Choudhury, G.;  
Martinez,L.; Han, E. S.; Killary, A. M.; Naylor, S. L.; Wang,  
L.–M.: Mousemelanoma growth stimulatory activity gene  
(Mgsa) is polymorphic andsyntenic with the W, Patch,  
Rumpwhite, and recessive spotting locion chromosome 5.  
Genomics 5: 629–632, 1989.

[64570] 2900.Chen, S.–H.; Anderson, J. E.; Giblett, E. R.: Human  
red cell 2,3–diphosphoglyceratemutase and monophos-  
phoglycerate mutase: genetic evidence for two separate-  
loci. Am. J. Hum. Genet. 29: 405–407, 1977.

[64571] 2901.Daikh, D. I.; Douglass, J. O.; Adelman, J. P.: Structure  
and expressionof the human motilin gene. DNA 8:  
615–621, 1989.

[64572] 2902.Fan, Y.–S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.;

Henry, W.M.; Yano, H.; Seino, Y.; Fujita, J.; Yamada, Y.; Inagaki, N.; Takeda, J.; Imura, H.; Bell, G. I.; Shows, T. B.: Localization of the human motilin gene (MLN) to chromosome 6p21.2. (Abstract) Cytogenet. Cell Genet. 51: 977 only, 1989.

- [64573] 2903. Feighner, S. D.; Tan, C. P.; McKee, K. K.; Palyha, O. C.; Hreniuk, D. L.; Pong, S.-S.; Austin, C. P.; Figueroa, D.; MacNeil, D.; Cascieri, M. A.; Nargund, R.; Bakshi, R.; Abramovitz, M.; Stocco, R.; Kargman, S.; O'Neill, G.; Van Der Ploeg, L. H. T.; Evans, J.; Patchett, A. A.; Smith, R. G.; Howard, A. D.: Receptor for motilin identified in the human gastrointestinal system. Science 284: 2184–2188, 1999.
- [64574] 2904. Gasparini, P.; Grifa, A.; Savasta, S.; Merlo, I.; Bisceglia, L.; Totaro, A.; Zelante, L.: The motilin gene: subregional localisation, tissue expression, DNA polymorphisms and exclusion as a candidate gene for the HLA-associated immotile cilia syndrome. Hum. Genet. 94: 671–674, 1994.
- [64575] 2905. Strissel, P. L.; Strick, R.; Tomek, R. J.; Roe, B. A.; Rowley, J. D.; Zeleznik-Le, N. J.: DNA structural properties of AF9 are similar to MLL and could act as recombination hot spots resulting in MLL/AF9 translocations and leukemogenesis. Hum. Molec. Genet. 9: 1671–1679, 2000.
- [64576] 2906. Isnard, P.; Depetris, D.; Mattei, M.-G.; Ferrier, P.;



Djabali,M.: cDNA cloning, expression and chromosomal localization of themurine AF-4 gene involved in human leukemia. *Mammalian Genome* 9:1065–1068, 1998.

[64577] 2907.Lovett, B. D.; Lo Nigro, L.; Rappaport, E. F.; Blair, I. A.; Osheroff,N.; Zheng, N.; Megonigal, M. D.; Williams, W. R.; Nowell, P. C.; Felix,C. A.: Near-precise interchromosomal recombination and functionalDNA topoisomerase II cleavage sites at MLL and AF-4 genomic breakpointsin treatment-related acute lymphoblastic leukemia with t(4;11) translocation. *Proc.Nat. Acad. Sci.* 98: 9802–9807, 2001.

[64578] 2908.Uckun, F. M.; Herman–Hatten, K.; Crotty, M.–L.; Sensel, M. G.;Sather, H. N.; Tuel–Ahlgren, L.; Sarquis, M. B.; Bostrom, B.; Nachman,J. B.; Steinherz, P. G.; Gaynon, P. S.; Heerema, N.: Clinical significanceof MLL–AF4 fusion transcript expression in the absence of a cytogenetically–detectable t(4;11)(q21;q23) chromosomal translocation. *Blood* 92:810–821, 1998.

[64579] 2909.Prasad, R.; Gu, Y.; Alder, H.; Nakamura, T.; Canaani, O.; Saito,H.; Huebner, K.; Gale, R. P.; Nowell, P. C.; Kuriyama, K.; Miyazaki,Y.; Croce, C. M.; Canaani, E.: Cloning of the ALL-1 fusion partner,the AF-6 gene, involved in acute myeloid leukemias with the

t(6;11)chromosome translocation. Cancer Res. 53:  
5624–5628, 1993.

- [64580] 2910.Saha, V.; Lillington, D. M.; Shelling, A. N.; Chaplin, T.; Yaspo,M.–L.; Ganesan, T. S.; Young, B. D.: AF6 gene on chromosome band6q27 maps distal to the minimal region of deletion in epithelial ovariancancer. Genes Chromo–somes Cancer 14: 220–222, 1995.
- [64581] 2911.Gerling, I. C.; Karlsen, A. E.; Chapman, H. D.; Ander–sen, H. U.;Boel, E.; Cunningham, J. M.; Nerup, J.; Leiter, E. H.: The inducible nitric oxide synthase gene, Nos2, maps to mouse chromosome 11. MammalianGenome 5: 318–320, 1994.
- [64582] 2912.Van Cong, N.; Moullec, J.: Linkage probable entre les groupes de phosphatase acide des globules rouges et le systeme Lewis. Ann.Genet. 14: 121–125, 1971.
- [64583] 2913.Wakita, Y.; Narahara, K.; Takahashi, Y.; Kikkawa, K.; Kimura,S.; Oda, M.; Kimoto, H.: Duplication of 2p25: confirmation of the assignment of soluble acid phosphatase (ACP1) locus to 2p25. Hum.Genet. 71: 259–260, 1985.
- [64584] 2914.Weitkamp, L. R.; Janzen, M. K.; Guttormsen, S. A.; Gershowitz,H.: Inherited pericentric inversion of chromosome number two: a linkage study. Ann. Hum. Genet. 33: 53–59, 1969.

- [64585] 2915.Weitkamp, L. R.; Lovrien, E. W.; Olaisen, B.; Fenger, K.; Gedde-Dahl,T., Jr.; Sorensen, S. A.; Conneally, P. M.; Bias, W. B.; Ott, J.:Linkage relations of the loci for the MN blood group and red cellphosphate. Birth Defects Orig. Art. Ser. 11(3): 276–280, 1975. Note:Alternate: Cytogenet. Cell Genet. 14: 446–450, 1975...
- [64586] 2916.Wo, Y.–Y. P.; McCormack, A. L.; Shabanowitz, J.; Hunt, D. F.;Davis, J. P.; Mitchell, G. L.; Van Etten, R. L.: Sequencing, cloning,and expression of human red cell-type acid phosphatase, a cytoplasmicphosphotyrosyl protein phosphatase. J. Biol. Chem. 267: 10856–10865,1992.
- [64587] 2917.Yoshihara, C. M.; Mohrenweiser, H. W.: Characterization of ACP1(TIC–1),an electrophoretic variant of erythrocyte acid phosphatase restrictedto the Ticuna Indians of Central Amazonas. Am. J. Hum. Genet. 32:898–907, 1980.
- [64588] 2918.Griffiths, L. R.: Authors response to 'Chromosome 17 and the inducible nitric oxide synthase gene in human essential hypertension' by Rutherford et al., Human Genetics, published on-line September 2001. (Letter) Hum.Genet. 110: 100–103, 2002.
- [64589] 2919.Jenkins, N. A.; Rothe, H.; Gilbert, D. J.; Copeland, N. G.; Kolb,H.: Mapping of the gene for inducible nitric oxide

(NO) synthase of mouse macrophages to chromosome 11, close to Evi-2, nu, and Idd-4. *Genomics* 19:402-404, 1994.

[64590] 2920. Johannesen, J.; Pie, A.; Pociot, F.; Kristiansen, O. P.; Karlsen, A. E.; Nerup, J. The Danish Study Group of Diabetes in Childhood; The Danish Insulin-Dependent Diabetes Mellitus Epidemiology and Genetics Group: Linkage of the human inducible nitric oxide synthase gene to type 1 diabetes. *J. Clin. Endocr. Metab.* 86: 2792-2796, 2001.

[64591] 2921. Lowenstein, C. J.; Glatt, C. S.; Bredt, D. S.; Snyder, S. H.: Cloned and expressed macrophage nitric oxide synthase contrasts with the brain enzyme. *Proc. Nat. Acad. Sci.* 89: 6711-6715, 1992.

[64592] 2922. Marsden, P. A.; Heng, H. H. Q.; Duff, C. L.; Shi, X.-M.; Tsui, L.-C.; Hall, A. V.: Localization of the human gene for inducible nitric oxide synthase (NOS2) to chromosome 17q11.2-q12. *Genomics* 19:183-185, 1994.

[64593] 2923. Mehrabian, M.; Xia, Y.-R.; Wen, P.-Z.; Warden, C. H.; Herschman, H. R.; Luscis, A. J.: Localization of murine macrophage inducible nitric oxide synthase to mouse chromosome 11. *Genomics* 22: 646-647, 1994.

[64594] 2924. Morris, B. J.: Critique of 'Chromosome 17 and the inducible nitric oxide synthase gene in human essential

hypertension' by Rutherford et al., Human Genetics, published online September 2001. (Letter) Hum.Genet. 110: 98–99, 2002.

- [64595] 2925. Nathan, C.; Shiloh, M. U.: Reactive oxygen and nitrogen intermediates in the relationship between mammalian hosts and microbial pathogens. Proc.Nat. Acad. Sci. 97: 8841–8848, 2000.
- [64596] 2926. Nicholson, S.; Bonecini-Almeida Mda, G; Lapa e Silva, J. R.; Nathan, C.; Xie, Q. W.; Mumford, R.; Weidner, J. R.; Calaycay, J.; Geng, J.; Boechat, N.; et al.; et al.: Inducible nitric oxide synthase in pulmonary alveolar macrophages from patients with tuberculosis. J. Exp. Med. 183:2293–2302, 1996.
- [64597] 2927. Nozaki, Y.; Hasegawa, Y.; Ichiyama, S.; Nakashima, I.; Shimokata, K.: Mechanism of nitric oxide-dependent killing of Mycobacterium bovis BCG in human alveolar macrophages. Infect. Immun. 65: 3644–3647, 1997.
- [64598] 2928. Vouldoukis, I.; Riveros-Moreno, V.; Dugas, B.; Ouaz, F.; Becherel, P.; Debre, P.; Moncada, S.; Mossalayi, M. D.: The killing of Leishmania major by human macrophages is mediated by nitric oxide induced after ligation of the Fc-epsilon-RII/CD23 surface antigen. Proc. Nat. Acad. Sci. 92: 7804–7808, 1995.

- [64599] 2929.Xu, W.; Charles, I. G.; Liu, L.; Moncada, S.; Emson, P.: Molecularcloning and structural organization of the human inducible nitricoxide synthase gene (NOS2). *Biochem. Biophys. Res. Commun.* 219:784–788, 1996.
- [64600] 2930.Xu, W.; Charles, I. G.; Moncada, S.; Gorman, P.; Sheer, D.; Liu,L.; Emson, P.: Mapping of the genes encoding human inducible andendothelial nitric oxide synthase (NOS2 and NOS3) to the pericentricregion of chromosome 17 and to chromosome 7, respectively. *Genomics* 21:419–422, 1994.
- [64601] 2931.Bredt, D. S.; Hwang, P. M.; Glatt, C. E.; Lowenstein, C.; Reed,R. R.; Snyder, S. H.: Cloned and expressed nitric oxide synthasestructurally resembles cytochrome P–450 reductase. *Nature* 351: 714–718,1991.
- [64602] 2932.Brenman, J. E.; Chao, D. S.; Xia, H.; Aldape, K.; Bredt, D. S.: Nitric oxide synthase complexed with dystrophin and absent fromskeletal muscle sarcolemma in Duchenne muscular dystrophy. *Cell* 82:743–752, 1995.
- [64603] 2933.Burnett, A. L.; Lowenstein, C. J.; Bredt, D. S.; Chang, T. S. K.;Snyder, S. H.: Nitric oxide: a physiologic mediator of penile erection. *Science* 257:401–403, 1992.
- [64604] 2934.Day, B. J.; Patel, M.; Calavetta, L.; Chang, L.–Y.; Stamler, J.S.: A mechanism of paraquat toxicity involving

nitric oxide synthase. Proc.Nat. Acad. Sci. 96:  
12760–12765, 1999.

- [64605] 2935.Deans, Z.; Dawson, S. J.; Xie, J.; Young, A. P.; Wallace, D.; Latchman,D. S.: Differential regulation of the two neuronal nitric-oxide synthasegene promoters by the Oct-2 transcription factor. J. Biol. Chem. 271:32153–32158, 1996.
- [64606] 2936.Gu, Z.; Kaul, M.; Yan, B.; Kridel, S. J.; Cui, J.; Strongin, A.;Smith, J. W.; Liddington, R. C.; Lipton, S. A.: S-nitrosylation ofmatrix metalloproteinases: signaling pathway to neuronal cell death. Science 297: 1186–1190, 2002.
- [64607] 2937.Kharazia, V. N.; Schmidt, H. H. H. W.; Weinberg, R. J.: Type I nitric oxide synthase fully accounts for NADPH-diaphorase in rat striatum, but not cortex. Neuroscience 62: 983–987, 1994.
- [64608] 2938.Kishimoto, J.; Spurr, N.; Liao, M.; Lizhi, L.; Emson, P.; Xu, W.: Localization of brain nitric oxide synthase (NOS) to human chromosome12. Genomics 14: 802–804, 1992.
- [64609] 2939.Kuo, R. C.; Baxter, G. T.; Thompson, S. H.; Stricker, S. A.; Patton,C.; Bonaventura, J.; Epel, D.: NO is necessary and sufficient foregg activation at fertilization. Nature 406: 633–636, 2000.

- [64610] 2940.Lee, C. G. L.; Gregg, A. R.; O'Brien, W. E.: Localization of the neuronal form of nitric oxide synthase to mouse chromosome 5. *Mammalian Genome* 6: 56–57, 1995.
- [64611] 2941.Magee, T.; Fuentes, A. M.; Garban, H.; Rajavashisth, T.; Marquez, D.; Rodriguez, J. A.; Rajfer, J.; Gonzalez-Cadavid, N. F.: Cloning of a novel neuronal nitric oxide synthase expressed in penis and lower urinary tract. *Biochem. Biophys. Res. Commun.* 226: 145–151, 1996.
- [64612] 2942.Devlin, R.; Henderson, H.; Monsalve, V.; Brunzell, J.; Deeb, S.; Hayden, M. R.: The molecular biology of hypertriglyceridemia: characterization of mutations in patients with lipoprotein lipase deficiency. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A4, 1989.
- [64613] 2943.Craig, W. J.: Leigh disease with deficiency of lipoamide dehydrogenase: treatment failure with dichloroacetate. *Pediat. Neurol.* 14: 69–71, 1996.
- [64614] 2944.Chynn, E. W.; Walton, D. S.; Hahn, L. B.; Dryja, T. P.: Norrie disease: diagnosis of a simplex case by DNA analysis. *Arch. Ophthalmol.* 114: 1136–1138, 1996.
- [64615] 2945.Thomas, S. A.; Palmiter, R. D.: Impaired maternal behavior in mice lacking norepinephrine and epinephrine. *Cell* 91: 583–592, 1997.



- [64616] 2946.Thomas, S. A.; Palmiter, R. D.: Thermoregulatory and metabolicphenotypes of mice lacking noradrenaline and adrenaline. *Nature* 387:94–97, 1997.
- [64617] 2947.Weinshilboum, R. M.: Catecholamine biochemical genetics in humanpopulations.In: Breakefield, X. O.: Neu-rogenetics: Genetic Approachesto the Nervous System. New York: Elsevier/North Holland (pub.) 1979. Pp. 257–282.
- [64618] 2948.Weinshilboum, R. M.; Schrott, H. G.; Raymond, F. A.; Weidman,W. H.; Elveback, L. R.: Inheritance of very low serum dopamine–beta–hydroxylaseactivity. *Am. J. Hum. Genet.* 27: 573–585, 1975.
- [64619] 2949.Wilson, A. F.; Elston, R. C.; Siervogel, R. M.; Tran, L. D.:Linkage of a gene regulating dopamine–beta–hydroxylase activity andthe ABO blood group locus. *Am. J. Hum. Genet.* 42: 160–166, 1988.
- [64620] 2950.Wilson, A. F.; Elston, R. C.; Siervogel, R. M.; Tran, L. D.:Linkage of a gene regulating dopamine–beta–hydroxylase activity andthe ABO blood group locus. (Abstract) *Am. J. Hum. Genet.* 41: A191only, 1987.
- [64621] 2951.Zabetian, C. P.; Anderson, G. M.; Buxbaum, S. G.; Elston, R. C.;Ichinose, H.; Nagatsu, T.; Kim, K.–S.; Kim, C.–H.; Malison, R. T.;Gelernter, J.; Cubells, J. F.: A quanti–

tative-trait analysis of human plasma-dopamine beta-hydroxylase activity: evidence for a major functional polymorphism at the DBH locus. *Am. J. Hum. Genet.* 68: 515-522, 2001.

[64622] 2952. Monros, E.; Molto, M. D.; Martinez, F.; Canizares, J.; Blanca, J.; Vilchez, J. J.; Prieto, F.; de Frutos, R.; Palau, F.: Phenotype correlation and intergenerational dynamics of the Friedreich ataxia GAA trinucleotide repeat. *Am. J. Hum. Genet.* 61: 101-110, 1997.

[64623] 2953. Montermini, L.; Andermann, E.; Labuda, M.; Richter, A.; Pandolfo, M.; Cavalcanti, F.; Pianese, L.; Iodice, L.; Farina, G.; Monticelli, A.; Turano, M.; Filla, A.; De Michele, G.; Coccozza, S.: The Friedreich ataxia GAA triplet repeat: pre-mutation and normal alleles. *Hum. Molec. Genet.* 6: 1261-1266, 1997.

[64624] 2954. Ohshima, K.; Montermini, L.; Wells, R. D.; Pandolfo, M.: Inhibitory effects of expanded GAA-TTC triplet repeats from intron I of the Friedreich ataxia gene on transcription and replication in vivo. *J. Biol. Chem.* 273:14588-14595, 1998.

[64625] 2955. Pianese, L.; Cavalcanti, F.; De Michele, G.; Filla, A.; Campanella, G.; Calabrese, O.; Castaldo, I.; Monticelli, A.; Coccozza, S.: The effect of parental gender on the GAA dy-

namic mutation in the FRDA gene. (Letter) Am. J. Hum. Genet. 60: 460–463, 1997.

[64626] 2956. Puccio, H.; Koenig, M.: Recent advances in the molecular pathogenesis of Friedreich ataxia. Hum. Molec. Genet. 9: 887–892, 2000.

[64627] 2957. Robinson, B. H.; Sherwood, W. G.; Kahler, S.; O'Flynn, M. E.; Nadler, H.: Lipoamide dehydrogenase deficiency. (Letter) New Eng. J. Med. 304: 53–54, 1981.

[64628] 2958. Rodius, F.; Duclos, F.; Wrogemann, K.; Le Paslier, D.; Ougen, P.; Billault, A.; Belal, S.; Musenger, C.; Brice, A.; Durr, A.; Mignard, C.; Sirugo, G.; Weissenbach, J.; Cohen, D.; Hentati, F.; Ben Hamida, M.; Mandel, J.-L.; Koenig, M.: Recombinations in individuals homozygous by descent localize the Friedreich ataxia locus in a cloned 450-kb interval. Am. J. Hum. Genet. 54: 1050–1059, 1994.

[64629] 2959. Macdonald, D.; Aguiar, R. C.; Mason, P. J.; Goldman, J. M.; Cross, N. C.: A new myeloproliferative disorder associated with chromosomal translocations involving 8p11: a review. Leukemia 9: 1628–1630, 1995.

[64630] 2960. Popovici, C.; Adelaide, J.; Ollendorff, V.; Chaffanet, M.; Guasch, G.; Jacrot, M.; Leroux, D.; Birnbaum, D.; Pebusque, M.-J.: Fibroblast growth factor receptor 1 is fused to FIM in stem-cell myeloproliferative disorder with

t(8;13)(p12;q12). Proc. Nat. Acad. Sci. 95:  
5712–5717,1998.

- [64631] 2961. Popovici, C.; Zhang, B.; Gregoire, M.-J.; Jonveaux, P.; Lafage-Pochitaloff, M.; Birnbaum, D.; Pebusque, M.-J.: The t(6;8)(q27;p11) translocation in a stem cell myeloproliferative disorder fuses a novel gene, FOP, to fibroblast growth factor receptor 1. Blood 93: 1381–1389, 1999.
- [64632] 2962. Thomas, W.; Rubenstein, M.; Goto, M.; Drayna, D.: A genetic analysis of the Werner syndrome region on human chromosome 8p. Genomics 16:685–690, 1993.
- [64633] 2963. Wood, S.; Schertzer, M.; Yaremko, M. L.: Sequence identity locates CEBPD and FGFR1 to mapped human loci within proximal 8p. Cytogenet. Cell Genet. 70: 188–191, 1995.
- [64634] 2964. Xiao, S.; Nalabolu, S. R.; Aster, J. C.; Ma, J.; Abruzzo, L.; Jaffe, E. S.; Stone, R.; Weissman, S. M.; Hudson, T. J.; Fletcher, J. A.: FGFR1 is fused with a novel zinc-finger gene, ZNF198, in the t(8;13) leukaemia/lymphoma syndrome. Nature Genet. 18: 84–87, 1998.
- [64635] 2965. Andersson, S. M.; Pispas, J. P.: Purification and properties of human liver tyrosine aminotransferase. Clin. Chim. Acta 125: 117–123, 1982.
- [64636] 2966. Barton, D. E.; Yang-Feng, T. L.; Francke, U.: The hu-

man tyrosineaminotransferase gene mapped to the long arm of chromosome 16 (region16q22–q24) by somatic cell hybrid analysis and in situ hybridization.Hum. Genet. 72: 221–224, 1986.

- [64637] 2967.Chen, S.–H.; Giblett, E. R.: Genetic variation of soluble glutamic–oxaloacetictransaminase in man. Am. J. Hum. Genet. 23: 419–424, 1971.
- [64638] 2968.Craig, I. W.; Tolley, E.; Bobrow, M.; van Heyningen, V.: Assignmentof a gene necessary for the expression of mitochondrial glutamic–oxaloacetictransaminase in human–mouse hybrid cells. Cytogenet. Cell Genet. 22:190–194, 1978.
- [64639] 2969.Davidson, R. G.; Cortner, J. A.; Rattazzi, M. C.; Ruddle, F. H.;Lubs, H. A.: Genetic polymorphisms of human mitochondrial glutamicoxaloacetic transaminase. Science 169: 391–392, 1970.
- [64640] 2970.DeLorenzo, R. J.; Ruddle, F. H.: Glutamate transaminase (GOT)genetics in mus musculus: linkage, polymorphism, and phenotypes ofthe GOT–2 and GOT–1 loci. Biochem. Genet. 4: 259–273, 1970.
- [64641] 2971.Francke, U.; Weitkamp, L. R.: Report of the committee on the geneticconstitution of chromosome 6. Cytogenet. Cell Genet. 25: 32–38,1979.

- [64642] 1972.Jeremiah, S. J.; Povey, S.; Burley, M. W.; Kielty, C.; Lee, M.;Spowart, G.; Corney, G.; Cook, P. J. L.: Mapping studies on humanmitochondrial glutamate oxaloacetate transaminase. *Ann. Hum. Genet.* 46:145–152, 1982.
- [64643] 1973.Pol, S.; Bousquet–Lemerrier, B.; Pave–Preux, M.; Bulle, F.; Passage,E.; Hanoune, J.; Mattei, M. G.; Barouki, R.: Chromosomal localizationof human aspartate amino–transferase genes by in situ hybridization.*Hum. Genet.* 83: 159–164, 1989.
- [64644] 1974.Pol, S.; Bousquet–Lemerrier, B.; Pave–Preux, M.; Pawlak, A.; Nalpas,B.; Berthelot, P.; Hanoune, J.; Barouki, R.: Nucleotide sequenceand tissue distribution of the human mitochondrial aspartate aminotransferasemRNA. *Biochem. Biophys. Res. Commun.* 157: 1309–1315, 1988.
- [64645] 1975.Tolley, E.; van Heyningen, V.; Brown, R.; Bobrow, M.; Craig, I.W.: Assignment to chromosome 16 of a gene necessary for the expressionof human mitochondrial glutamate oxaloacetate transaminase (aspartateaminotransferase) (E.C. 2.6.1.1). *Biochem. Genet.* 18: 947–954,1980.
- [64646] 1976.Toyomasu, T.; Sakakibara, S.; Kagamiyama, H.; Matsumoto, H.:Genetic polymorphism of mitochondrial glutamate–oxaloacetate transaminasein Japanese. *Hum. Genet.*

66: 90–91, 1984.

- [64647] 2977. Tsuji, S.; Kobayashi, H.; Uchida, Y.; Ihara, Y.; Miyatake, T.: Molecular cloning of human growth inhibitory factor cDNA and its down-regulation in Alzheimer's disease. *EMBO J.* 11: 4843–4850, 1992.
- [64648] 2978. Uchida, Y.; Takio, K.; Titani, K.; Ihara, Y.; Tomonaga, M.: The growth inhibitory factor that is deficient in the Alzheimer's disease brain is a 68 amino acid metallothionein-like protein. *Neuron* 7: 337–347, 1991.
- [64649] 2979. Hoshino, S.; Miyazawa, H.; Enomoto, T.; Hanaoka, F.; Kikuchi, Y.; Kikuchi, A.; Ui, M.: A human homologue of the yeast GST1 gene codes for a GTP-binding protein and is expressed in a proliferation-dependent manner in mammalian cells. *EMBO J.* 8: 3807–3814, 1989.
- [64650] 2980. Kikuchi, Y.; Shimatake, H.; Kikuchi, A.: A yeast gene required for the G1-to-S transition encodes a protein containing an A-kinase target site and GTPase domain. *EMBO J.* 7: 1175–1182, 1988.
- [64651] 2981. Ozawa, K.; Murakami, Y.; Eki, T.; Yokoyama, K.; Soeda, E.; Hoshino, S.; Ui, M.; Hanaoka, F.: Mapping of the human GSPT1 gene, a human homologue of the yeast GST1 gene, to chromosomal band 16p13.1. *Somat. Cell Molec. Genet.* 18: 189–194, 1992.

- [64652] 2982.Harris, H.; Hopkinson, D. A.; Robson, E. B.: The incidence of rare alleles determining electrophoretic variants: data on 43 enzyme loci in man. *Ann. Hum. Genet.* 37: 237–253, 1974.
- [64653] 2983.Beutler, E.; Gelbart, T.; Kuhl, W.: Human red cell glucose-6-phosphate dehydrogenase: all active enzyme has sequence predicted by the X chromosome-encoded cDNA. *Cell* 62: 7–9, 1990.
- [64654] 2984.Henikoff, S.; Smith, J. M.: The human mRNA that provides the N-terminus of chimeric G6PD encodes GMP reductase. *Cell* 58: 1021–1022, 1989.
- [64655] 2985.Kanno, H.; Huang, I.-Y.; Kan, Y. W.; Yoshida, A.: Two structural genes on different chromosomes are required for encoding the major subunit of human red cell glucose-6-phosphate dehydrogenase. *Cell* 58: 595–606, 1989.
- [64656] 2986.Kondoh, T.; Kanno, H.; Chang, L.; Yoshida, A.: Genomic structure and expression of human guanosine monophosphate reductase. *Hum. Genet.* 88: 219–224, 1991.
- [64657] 2987.Kondoh, T.; Kanno, H.; Chang, L.; Yoshida, A.: Identification of common variant alleles of the human guanosine monophosphate reductase gene. *Hum. Genet.* 88: 225–227, 1991.



- [64658] 1988.Mason, P. J.; Bautista, J. M.; Vulliamy, T. J.; Turner, N.; Luzzatto,L.: Human red cell glucose-6-phosphate dehydrogenase is encoded onlyon the X chromosome. *Cell* 62: 9-10, 1990.
- [64659] 1989.Murano, I.; Tsukahara, M.; Kajii, T.; Yoshida, A.: Mapping ofthe human guanosine monophosphate reductase gene (GMPR) to chromosome6p23 by fluorescence in situ hybridization. *Genomics* 19: 179-180,1994.
- [64660] 1990.Yoshida, A.; Kan, Y. W.: Origin of 'fused' glucose-6-phosphatedehydrogenase. *Cell* 62: 11-12, 1990.
- [64661] 1991.Brady, W. A.; Kokoris, M. S.; Fitzgibbon, M.; Black, M. E.: Cloning,characterization, and modeling of mouse and human guanylate kinases. *J.Biol. Chem.* 271: 16734-16740, 1996.
- [64662] 1992.Dallapiccola, B.; Lungarotti, M. S.; Falorni, A.; Magnani, M.;Dacha, M.: Evidence for the assignment of GUK1 gene locus to 1q32-q43segment from gene dosage effect. *Ann. Genet.* 23: 83-85, 1980.
- [64663] 1993.Kajiwara, K.; Berson, E. L.; Dryja, T. P.: Digenic retinitispigmentosa due to mutations at the unlinked peripherin/RDS and ROM1loci. *Science* 264: 1604-1608, 1994.
- [64664] 1994.Kajiwara, K.; Hahn, L. B.; Mukai, S.; Travis, G. H.;

Berson, E.L.; Dryja, T. P.: Mutations in the human retinal degeneration slow gene in autosomal dominant retinitis pigmentosa. *Nature* 354: 480–483, 1991.

[64665] 1995. Kajiwar, K.; Sandberg, M. A.; Berson, E. L.; Dryja, T. P.: A null mutation in the human RDS/peripherin gene in a family with autosomal dominant retinitis punctata albescens. (Abstract) *Am. J. Hum. Genet.* 51(suppl.): A6 only, 1992.

[64666] 1996. Kajiwar, K.; Sandberg, M. A.; Berson, E. L.; Dryja, T. P.: A null mutation in the human peripherin/RDS gene in a family with autosomal dominant retinitis punctata albescens. *Nature Genet.* 3: 208–212, 1993.

[64667] 1997. Kedzierski, W.; Nusinowitz, S.; Birch, D.; Clarke, G.; McInnes, R. R.; Bok, D.; Travis, G. H.: Deficiency of rds/peripherin causes photoreceptor death in mouse models of digenic and dominant retinitis pigmentosa. *Proc. Nat. Acad. Sci.* 98: 7718–7723, 2001.

[64668] 1998. Keen, T. J.; Inglehearn, C. F.: Mutations and polymorphisms in the human peripherin-RDS gene and their involvement in inherited retinal degeneration. *Hum. Mutat.* 8: 297–303, 1996.

[64669] 1999. Kikawa, E.; Nakazawa, M.; Chida, Y.; Shiono, T.; Tamai, M.: A novel mutation (asn244-to-lys) in the periph-

erin/RDS gene causing autosomal dominant retinitis pigmentosa associated with bull's-eye maculopathy detected by nonradioisotopic SSCP. *Genomics* 20: 137–139, 1994.

[64670] 3000. Kim, R. Y.; Dollfus, H.; Keen, T. J.; Fitzke, F. W.; Arden, G. B.; Bhattacharya, S. S.; Bird, A. C.: Autosomal dominant pattern dystrophy of the retina associated with a 4-base pair insertion at codon 140 in the peripherin/RDS gene. *Arch. Ophthalmol.* 113: 451–455, 1995.

[64671] 3001. Kohl, S.; Christ-Adler, M.; Apfelstedt-Sylla, E.; Kellner, U.; Eckstein, A.; Zrenner, E.; Wissinger, B.: RDS/peripherin gene mutations are frequent causes of central retinal dystrophies. *J. Med. Genet.* 34: 620–626, 1997.

[64672] 3002. Ma, J.; Norton, J. C.; Allen, A. C.; Burns, J. B.; Hasel, K. W.; Burns, J. L.; Sutcliffe, J. G.; Travis, G. H.: Retinal degeneration slow (rds) in mouse results from simple insertion of a t haplotype-specific element into protein-coding exon II. *Genomics* 28: 212–219, 1995.

[64673] 3003. Nichols, B. E.; Drack, A. V.; Vandenberg, K.; Kimura, A. E.; Sheffield, V. C.; Stone, E. M.: A 2 base pair deletion in the RDS gene associated with butterfly-shaped pigment dystrophy of the fovea. *Hum. Molec. Genet.* 2: 601–603, 1993.

[64674] 3004. Nichols, B. E.; Sheffield, V. C.; Vandenberg, K.;

Drack, A. V.; Kimura, A. E.; Stone, E. M.: Butterfly-shaped pigment dystrophy of the fovea caused by a point mutation in codon 167 of the RDS gene. *Nature Genet.* 3: 202–207, 1993.

[64675] 3005. Payne, A. M.; Downes, S. M.; Bessant, D. A. R.; Bird, A. C.; Bhattacharya, S. S.: Founder effect, seen in the British population, of the 172 peripherin/RDS mutation – and further refinement of genetic positioning of the peripherin/RDS gene. (Letter) *Am. J. Hum. Genet.* 192–195, 1998.

[64676] 3006. Pendleton, J. W.; Violette, S. M.; Hunihan, L. W.; Greene, L. A.; Ruddle, F. H.: The peripherin gene maps to mouse chromosome 15. *Genomics* 9:369–372, 1991.

[64677] 3007. Reig, C.; Serra, A.; Gean, E.; Vidal, M.; Arumi, J.; De la Calzada, M. D.; Antich, J.; Carballo, M.: A point mutation in the RDS-peripherin gene in a Spanish family with central areolar choroidal dystrophy. *Ophthalmol. Genet.* 16: 35–44, 1995.

[64678] 3008. Sarra, G.-M.; Stephens, C.; de Alwis, M.; Bainbridge, J. W. B.; Smith, A. J.; Thrasher, A. J.; Ali, R. R.: Gene replacement therapy in the retinal degeneration slow (rds) mouse: the effect on retinal degeneration following partial transduction of the retina. *Hum. Molec. Genet.* 10:

2353–2361, 2001.

- [64679] 3009.Travis, G. H.; Brennan, M. B.; Danielson, P. E.; Kozak, C. A.;Sutcliffe, J. G.: Identification of a photoreceptor-specific mRNAencoded by the gene responsible for retinal degeneration slow (rds). *Nature* 338:70–73, 1989.
- [64680] 3010.Travis, G. H.; Christerson, L.; Danielson, P. E.; Klisak, I.;Sparkes, R. S.; Hahn, L. B.; Dryja, T. P.; Sutcliffe, J. G.: Thehuman retinal degeneration slow (RDS) gene: chromosome assignmentand structure of the mRNA. *Genomics* 10: 733–739, 1991.
- [64681] 3011.Travis, G. H.; Hepler, J. E.: A medley of retinal dystrophies. *NatureGenet.* 3: 191–192, 1993.
- [64682] 3012.Cramer, P.; Bushnell, D. A.; Fu, J.; Gnatt, A. L.; Maier–Davis,B.; Thompson, N. E.; Burgess, R. R.; Edwards, A. M.; David, P. R.;Kornberg, R. D.: Architecture of RNA polymerase II and implicationsfor the transcription mechanism. *Science* 288: 640–648, 2000.
- [64683] 3013.Yudkovsky, N.; Ranish, J. A.; Hahn, S.: A transcription reinitiationintermediate that is stabilized by activator. *Nature* 408: 225–229,2000.
- [64684] 3014.Dammann, R.; Pfeifer, G. P.: Cloning and characterization of thehuman RNA polymerase I subunit hRPA40. *Biochim. Biophys. Acta* 1396:153–157, 1998.

- [64685] 3015.Fanciulli, M.; Bruno, T.; Di Padova, M.; De Angelis, R.; Lovari,S.; Floridi, A.; Passananti, C.: The interacting RNA polymerase II subunits, hRPB11 and hRPB3, are coordinately expressed in adult human tissues and down-regulated by doxorubicin. FEBS Lett. 427: 236–240,1998.
- [64686] 3016.Pati, U. K.; Weissman, S. M.: The amino acid sequence of the human RNA polymerase II 33–kDa subunit hRPB 33 is highly conserved among eukaryotes. J. Biol. Chem. 265: 8400–8403, 1990.
- [64687] 3017.Becker, D. M.; Fikes, J. D.; Guarente, L.: A cDNA encoding a human CCAAT-binding protein cloned by functional complementation in yeast. Proc.Nat. Acad. Sci. 88: 1968–1972, 1991.
- [64688] 3018.Henrion, A. A.; Martinez, A.; Mattei, M.–G.; Kahn, A.; Raymondjean,M.: Structure, sequence, and chromosomal location of the gene for USF2 transcription factors in mouse. Genomics 25: 36–43, 1995.
- [64689] 3019.Gharani, N.; Waterworth, D. M.; Batty, S.; White, D.; Gilling–Smith,C.; Conway, G. S.; McCarthy, M.; Franks, S.; Williamson, R.: Association of the steroid synthesis gene CYP11a with polycystic ovary syndrome and hyperandrogenism. Hum. Molec. Genet. 6: 397–402, 1997.
- [64690] 3020.Jakimiuk, A. J.; Weitsman, S. R.; Magoffin, D. A.:

5- $\alpha$ -reductase activity in women with polycystic ovary syndrome. *J. Clin. Endocr. Metab.* 84: 2414–2418, 1999.

[64691] 3021. Kuttann, F.; Couillin, P.; Girard, F.; Billaud, L.; Vincens, M.; Boucekkine, C.; Thalabard, J.-C.; Maudelonde, T.; Spritzer, P.; Mowszowicz, I.; Boue, A.; Mauvais-Jarvis, P.: Late-onset adrenal hyperplasia in hirsutism. *New Eng. J. Med.* 313: 224–231, 1985.

[64692] 3022. Kuttann, F.; Mowszowicz, I.; Schaison, G.; Mauvais-Jarvis, P.: Androgen production and skin metabolism in hirsutism. *J. Endocr.* 75: 83–91, 1977.

[64693] 3023. Kurtz, T. W.: The ACE of hearts. *Nature* 359: 588–589, 1992.

[64694] 3024. Lindpaintner, K.; Lee, M.; Larson, M. G.; Rao, V. S.; Pfeffer, M. A.; Ordovas, J. M.; Schaefer, E. J.; Wilson, A. F.; Wilson, P. W. F.; Vasan, R. S.; Myers, R. H.; Levy, D.: Absence of association or genetic linkage between the angiotensin-converting-enzyme gene and left ventricular mass. *New Eng. J. Med.* 334: 1023–1028, 1996.

[64695] 3025. Lindpaintner, K.; Pfeffer, M. A.; Kreutz, R.; Stampfer, M. J.; Grodstein, F.; LaMotte, F.; Buring, J.; Hennekens, C. H.: A prospective evaluation of an angiotensin-converting-enzyme gene polymorphism and the risk of ischemic heart disease. *New Eng. J. Med.* 332: 706–711, 1995.

- [64696] 3026.Marre, M.; Bernadet, P.; Gallois, Y.; Savagner, F.; Guyene, T.–T.;Hallab, M.; Cambien, F.; Passa, P.; Alhenc–Gelas, F.: Relationships between angiotensin I converting enzyme gene polymorphism, plasma levels, and diabetic retinal and renal complications. *Diabetes* 43:384–388, 1994.
- [64697] 3027.Marre, M.; Jeunemaitre, X.; Gallois, Y.; Rodier, M.; Chatellier, G.; Sert, C.; Dusselier, L.; Kahal, Z.; Chaillous, L.; Halimi, S.; Muller, A.; Sackmann, H.; Bauduceau, B.; Bled, F.; Passa, P.; Alhenc–Gelas, F.: Contribution of genetic polymorphism in the renin–angiotensin system to the development of renal complications in insulin–dependent diabetes. *J. Clin. Invest.* 99: 1585–1595, 1997.
- [64698] 3028.Mattei, M.–G.; Hubert, C.; Alhenc–Gelas, F.; Roeckel, N.; Corvol, P.; Soubrier, F.: Angiotensin–I converting enzyme gene is on chromosome 17. (Abstract) *Cytogenet. Cell Genet.* 51: 1041, 1989.
- [64699] 3029.Montgomery, H. E.; Marshall, R.; Hemingway, H.; Myerson, S.; Clarkson, P.; Dollery, C.; Hayward, M.; Holliman, D. E.; Jubb, M.; World, M.; Thomas, E. L.; Brynes, A. E.; Saeed, N.; Barnard, M.; Bell, J. D.; Prasad, K.; Rayson, M.; Talmud, P. J.; Humphries, S. E.: Human gene for physical performance. (Letter) *Nature* 393: 221–222, 1998.



- [64700] 3030.Ohishi, M.; Fujii, K.; Minamino, T.; Hagaki, J.; Kamitani, A.;Rakugi, H.; Zhao, Y.; Mikami, H.; Miki, T.; Ogihara, T.: A potent genetic risk factor for restenosis. (Letter) Nature Genet. 5: 324–325,1993.
- [64701] 3031.Oike, Y.; Hata, A.; Ogata, Y.; Numata, Y.; Shido, K.; Kondo, K.: Angiotensin converting enzyme as a genetic risk factor for coronaryartery spasm: implication in the pathogenesis of myocardial infarction. J.Clin. Invest. 96: 2975–2979, 1995.
- [64702] 3032.Okabe, T.; Fumisawa, M.; Yotsumoto, M.; Takaru, F.; Lanzillo,J. J.; Fanburg, B. L.: Familial elevation of serum angiotensin–convertingenzyme. Quart. J. Med. 216: 55–61, 1985.
- [64703] 3033.Montgomery, H. E.; Clarkson, P.; Dollery, C. M.; Prasad, K.; Losi,M.–A.; Hemingway, H.; Statters, D.; Jubb, M.; Girvain, M.; Varnava,A.; World, M.; Deanfield, J.; Talmud, P.; McEwan, J. R.; McKenna,W. J.; Humphries, S.: Association of angiotensin–converting enzymegene I/D polymorphism with change in left ventricular mass in responseto physical training. Circulation 96: 741–747, 1997.
- [64704] 3034.Pfeffer, M. A.; Braunwald, E.; Moye, L. A.; Basta, L.; Brown,E. J., Jr.; Cuddy, T. E.; Davis, B. R.; Geltman, E. M.;

Goldman, S.; Flaker, G. C.; Klein, M.; Lamas, G. A.; Packer, M.; Rouleau, J.; Rouleau, J. L.; Rutherford, J.; Wertheimer, J. H.; Hawkins, C. M.: Effect of captopril on mortality and morbidity in patients with left ventricular dysfunction after myocardial infarction: results of the survival and ventricular enlargement trial. *New Eng. J. Med.* 327: 669–677, 1992.

[64705] 3035. Ramaraj, P.; Kessler, S. P.; Colmenares, C.; Sen, G. C.: Selective restoration of male fertility in mice lacking angiotensin-converting enzymes by sperm-specific expression of the testicular isozyme. *J. Clin. Invest.* 102: 371–378, 1998.

[64706] 3036. Rigat, B.; Hubert, C.; Alhenc-Gelas, F.; Cambien, F.; Corvol, P.; Soubrier, F.: An insertion/deletion polymorphism in the angiotensin-converting enzyme gene accounting for half the variance of serum enzyme levels. *J. Clin. Invest.* 86: 1343–1346, 1990.

[64707] 3037. Rigat, B.; Hubert, C.; Corvol, P.; Soubrier, F.: PCR detection of the insertion/deletion polymorphism of the human angiotensin converting enzyme gene (DCP1) (dipeptidylcarboxypeptidase 1). *Nucleic Acids Res.* 20: 1433, 1992.

[64708] 3038. Ruiz, J.; Blanche, H.; Cohen, N.; Velho, G.; Cambien,

F.; Cohen,D.; Passa, P.; Froguel, P.: Insertion/deletion polymorphism of theangiotensin–converting enzyme gene is strongly associated with coronaryheart disease in non–insulin–dependent diabetes mellitus. Proc. Nat.Acad. Sci. 91: 3662–3665, 1994.

[64709] 3039.Schachter, F.; Faure–Delanef, L.; Guenot, F.; Rouger, H.; Froguel,P.; Lesueur–Ginot, L.; Cohen, D.: Genetic associations with humanlongevity at the APOE and ACE loci. Nature Genet. 6: 29–32, 1994.

[64710] 3040.Schmidt, S.; Schone, N.; Ritz, E.; Giesel, R.; Bergis, K.; Strojek,K.; Greszczak, W.; Schroter, W.; Willms, B. H. L.; Petzold, R.; Henrichs,H. R.; Rambausek, M.; Schwarzbeck, A.; Kohr, B.; Schneider, P.; Bosch,A.: Association of ACE gene polymorphism and diabetic nephropathy? KidneyInt. 47: 1176–1181, 1995.

[64711] 3041.Schunkert, H.; Hense, H.–W.; Holmer, S. R.; Stender, M.; Perz,S.; Keil, U.; Lorell, B. H.; Riegger, G. A. J.: Association betweena deletion polymorphism of the angiotensin–converting–enzyme geneand left ventricular hypertrophy. New Eng. J. Med. 330: 1634–1638,1994.

[64712] 3042.Singer, D. R. J.; Missouris, C. G.; Jeffery, S.: Angiotensin–convertingenzyme gene polymorphism: what to do about all the confusion? (Editorial) Circulation

94:236–239, 1996.

[64713] 3043.Soubrier, F.; Alhenc-Gelas, F.; Hubert, C.; Allegrini, J.; John, M.; Tregear, G.; Corvol, P.: Two putative active centers in human angiotensin I-converting enzyme revealed by molecular cloning. *Proc. Nat. Acad. Sci.* 85: 9386–9390, 1988.

[64714] 3044.Tarnow, L.; Cambien, F.; Rossing, P.; Nielsen, F. S.; Hansen, B. V.; Lecerf, L.; Poirier, O.; Danilov, S.; Parving, H.-H.: Lack of relationship between an insertion/deletion polymorphism in the angiotensin I-converting enzyme gene and diabetic nephropathy and proliferative retinopathy in IDDM patients. *Diabetes* 44: 489–494, 1995.

[64715] 3045.Tiret, L.; Rigat, B.; Visvikis, S.; Breda, C.; Corvol, P.; Cambien, F.; Soubrier, F.: Evidence, from combined segregation and linkage analysis, that a variant of the angiotensin I-converting enzyme (ACE) gene controls plasma ACE levels. *Am. J. Hum. Genet.* 51: 197–205, 1992.

[64716] 3046.Vleming, L. J.; van der Pijl, J. W.; Lemkes, H. H. P. J.; Westendorp, R. G. J.; Maassen, J. A.; Daha, M. R.; van Es, L. A.; van Kooten, C.: The DD genotype of the ACE gene polymorphism is associated with progression of diabetic nephropathy to end stage renal failure in IDDM. *Clin. Nephrol.* 51: 133–140, 1999.

- [64717] 3047.Williams, A. G.; Rayson, M. P.; Jubb, M.; World, M.; Woods, D.R.; Hayward, M.; Martin, J.; Humphries, S. E.; Montgomery, H. E.:The ACE gene and muscle performance. *Nature* 403: 614 only, 2000.
- [64718] 3048.Reddi, A. H.: BMP-1: resurrection as procollagen C-proteinase. *Science* 271:5-6, 1996.
- [64719] 3049.Takahara, K.; Lee, S.; Wood, S.; Greenspan, D. S.: Structuralorganization and genetic localization of the human bone morphogeneticprotein 1/mammalian tolloid gene. *Genomics* 29: 9-15, 1995.
- [64720] 3050.Takahara, K.; Lyons, G. E.; Greenspan, D. S.: Bone morphogeneticprotein-1 and a mammalian tolloid homologue (mTld) are encoded byalternatively spliced transcripts which are differentially expressedin some tissues. *J. Biol. Chem.* 269: 32572-32578, 1994.
- [64721] 3051.Scott, I. C.; Blitz, I. L.; Pappano, W. N.; Imamura, Y.; Clark,T. G.; Steiglit, B. M.; Thomas, C. L.; Maas, S. A.; Takahara, K.;Cho, K. W. Y.; Greenspan, D. S.: Mammalian BMP-1/Tolloid-relatedmetalloproteinases, including novel family member mammalian Tolloid-like2, have differential enzymatic activities and distributions of expressionrelevant to patterning and skeletogenesis. *Dev. Biol.* 213: 283-300,1999.

- [64722] 3052. Yoshiura, K.; Tamura, T.; Hong, H.-S.; Ohta, T.; Soejima, H.; Kishino, T.; Jinno, Y.; Niikawa, N.: Mapping of the bone morphogenetic protein 1 gene (BMP1) to 8p21: removal of BMP1 from candidacy for the bone disorder in Langer-Giedion syndrome. *Cytogenet. Cell Genet.* 64:208–209, 1993.
- [64723] 3053. Hahn, G. V.; Cohen, R. B.; Wozney, J. M.; Levitz, C. L.; Shore, E. M.; Zasloff, M. A.; Kaplan, F. S.: A bone morphogenetic protein subfamily: chromosomal localization of human genes for BMP5, BMP6, and BMP7. *Genomics* 14: 759–762, 1992.
- [64724] 3054. Lyons, K.; Graycar, J. L.; Lee, A.; Hashmi, S.; Lindquist, P. B.; Chen, E. Y.; Hogan, B. L. M.; Derynck, R.: Vgr-1, a mammalian gene related to *Xenopus* Vg-1, is a member of the transforming growth factor beta gene superfamily. *Proc. Nat. Acad. Sci.* 86: 4554–4558, 1989.
- [64725] 3055. Rickard, D. J.; Hofbauer, L. C.; Bonde, S. K.; Gori, F.; Spelsberg, T. C.; Riggs, B. L.: Bone morphogenetic protein-6 production in human osteoblastic cell lines: selective regulation by estrogen. *J. Clin. Invest.* 101: 413–422, 1998.
- [64726] 3056. Marker, P. C.; King, J. A.; Copeland, N. G.; Jenkins, N. A.; Kingsley, D. M.: Chromosomal localization, embryonic

expression, and imprinting tests for Bmp7 on distal mouse chromosome 2. *Genomics* 28: 576–580, 1995.

[64727] 3057. Ozkaynak, E.; Rueger, D. C.; Drier, E. A.; Corbett, C.; Ridge, R. J.; Sampath, T. K.; Oppermann, H.: OP-1 cDNA encodes an osteogenic protein in the TGF- $\beta$  family. *EMBO J.* 9: 2085–2093, 1990.

[64728] 3058. Solursh, M.; Langille, R. M.; Wood, J.; Sampath, T. K.: Osteogenic protein-1 is required for mammalian eye development. *Biochem. Biophys. Res. Commun.* 218: 438–443, 1996.

[64729] 3059. Modi, W. S.; Dean, M.; Pollock, D. D.; Seuanez, H. N.; Christakos, S.: Chromosomal localization of the calbindin gene. (Abstract) *Cytogenet. Cell Genet.* 58: 1930 only, 1991.

[64730] 3060. Parmentier, M.; De Vijlder, J. J. M.; Muir, E.; Szpirer, C.; Islam, M. Q.; Geurts van Kessel, A.; Lawson, D. E. M.; Vassart, G.: The human calbindin 27 kDa gene: structural organization of the 5-prime and 3-prime regions, chromosomal assignment and restriction fragment length polymorphism. *Genomics* 4: 309–319, 1989.

[64731] 3061. Parmentier, M.; Lawson, D. E. M.; Vassart, G.: Human 27-kDa calbindin complementary DNA sequence: evolutionary and functional implications. *Europ. J. Biochem.*

170: 207–215, 1987.

- [64732] 3062.Parmentier, M.; Passage, E.; Vassart, G.; Mattei, M.–G.: The humancalbindin D28k (CALB1) and calretinin (CALB2) genes are located at8q21.3–q22.1 and 16q22–q23, respectively, suggesting a common duplicationwith the carbonic anhydrase isozyme loci. Cytogenet. Cell Genet. 57:41–43, 1991.
- [64733] 3063.Parmentier, M.; Vassart, G.: HindIII RFLP on chromosome 8 detectedwith a calbindin 27 kDa cDNA probe, HBSC21. Nucleic Acids Res. 16:9373 only, 1988.
- [64734] 3064.Seto–Ohshima, A.; Emson, P. C.; Lawson, E.; Mountjoy, C. Q.; Carrasco,L. H.: Loss of matrix calcium–binding protein–containing neuronsin Huntington's disease. Lancet I: 1252–1254, 1988.
- [64735] 3065.Parmentier, M.; Szpirer, J.; Levan, G.; Vassart, G.: The humangenes for calbindin 27 and 29 kDa proteins are located on chromosomes8 and 16, respectively. Cytogenet. Cell Genet. 52: 85–87, 1989.
- [64736] 3066.Schiffmann, S. N.; Cheron, G.; Lohof, A.; d'Alcantara, P.; Meyer,M.; Parmentier, M.; Schurmans, S.: Impaired motor coordination andPurkinje cell excitability in mice lacking calretinin. Proc. Nat.Acad. Sci. 96: 5257–5262, 1999.
- [64737] 3067.Davies, A. A.; Moss, S. E.; Crompton, M. R.; Jones, T.



A.; Spurr, N. K.; Sheer, D.; Kozak, C.; Crumpton, M. J.: The gene coding for the p68 calcium-binding protein is localized to bands q32-q34 of human chromosome 5, and to mouse chromosome 11. *Hum. Genet.* 82: 234-238, 1989.

[64738] 3068. Smith, P. D.; Davies, A.; Crumpton, M. J.; Moss, S. E.: Structure of the human annexin VI gene. *Proc. Nat. Acad. Sci.* 91: 2713-2717, 1994.

[64739] 3069. Sudhof, T. C.; Slaughter, C. A.; Leznicki, I.; Barjon, P.; Reynolds, G. A.: Human 67-kDa calelectrin contains a duplication of four repeats found in 35-kDa lipocortins. *Proc. Nat. Acad. Sci.* 85: 664-668, 1988.

[64740] 3070. Blow, J. J.; Laskey, R. A.: A role for the nuclear envelope in controlling DNA replication within the cell cycle. *Nature* 332: 546-548, 1988.

[64741] 3071. Burkhardt, R.; Schulte, D.; Hu, D.; Musahl, C.; Gohring, F.; Knippers, R.: Interactions of human nuclear proteins P1Mcm3 and P1Cdc46. *Europ. J. Biochem.* 228: 431-438, 1995.

[64742] 3072. Chong, J. P.; Mahbubani, H. M.; Khoo, C. Y.; Blow, J. J.: Purification of an MCM-containing complex as a component of the DNA replication licensing system. *Nature* 375: 418-421, 1995.

[64743] 3073. Hu, B.; Burkhardt, R.; Schulte, D.; Musahl, C.; Knip-

pers, R.: TheP1 family: a new class of nuclear mammalian proteins related to theyeast Mcm replication proteins. Nu-  
cleic Acids Res. 21: 5289–5293,1993.

- [64744] 3074.Kubota, Y.; Mimura, S.; Nishimoto, S.; Takisawa, H.;  
Nojima, H.: Identification of the yeast MCM3–related pro-  
tein as a componentof Xenopus DNA replication licensing  
factor. Cell 81: 601–609, 1995.
- [64745] 3075.Labib, K.; Tercero, J. A.; Diffley, J. F. X.: DNA replica-  
tionfork progression requires uninterrupted MCM2–7  
function. Science 288:1643–1647, 2000.
- [64746] 3076.Madine, M. A.; Khoo, C. Y.; Mills, A. D.; Laskey, R. A.:  
MCM3complex required for cell cycle regulation of DNA  
replication in vertebratecells. Nature 375: 421–424, 1995.
- [64747] 3077.Mincheva, A.; Todorov, I.; Werner, D.; Fink, T. M.;  
Lichter, P.: The human gene for nuclear protein BM28  
(CDCL1), a new member ofthe early S–phase family of  
proteins, maps to chromosome band 3q21. Cytogenet.Cell  
Genet. 65: 276–277, 1994.
- [64748] 3078.Tsuruga, H.; Yabuta, N.; Hashizume, K.; Ikeda, M.;  
Endo, Y.; Nojima,H.: Expression, nuclear localization and  
interactions of human MCM/P1proteins. Biochem. Bio-  
phys. Res. Commun. 236: 118–125, 1997.
- [64749] 3079.Adams, M. D.; Dubnick, M.; Kerlavage, A. R.;

Moreno, R.; Kelley, J. M.; Utterback, T. R.; Nagle, J. W.; Fields, C.; Venter, J. C.: Sequence identification of 2,375 human brain genes. *Nature* 355:632–634, 1992.

[64750] 3080. Tugendreich, S.; Boguski, M. S.; Seldin, M. S.; Hieter, P.: Linking yeast genetics to mammalian genomes: identification and mapping of the human homolog of CDC27 via the expressed sequence tag (EST) database. *Proc. Nat. Acad. Sci.* 90: 10031–10035, 1993.

[64751] 3081. Demetrick, D. J.; Beach, D. H.: Chromosome mapping of human CDC25A and CDC25B phosphatases. *Genomics* 18: 144–147, 1993.

[64752] 3082. Fauman, E. B.; Cogswell, J. P.; Lovejoy, B.; Rocque, W. J.; Holmes, W.; Montana, V. G.; Piwnicka-Worms, H.; Rink, M. J.; Saper, M. A.: Crystal structure of the catalytic domain of the human cell cycle control phosphatase, Cdc25A. *Cell* 93: 617–625, 1998.

[64753] 3083. Galaktionov, K.; Lee, A. K.; Eckstein, J.; Draetta, G.; Meckler, J.; Loda, M.; Beach, D.: CDC25 phosphatases as potential human oncogenes. *Science* 269:1575–1577, 1995.

[64754] 3084. Mailand, N.; Falck, J.; Lukas, C.; Syljuasen, R. G.; Welcker, M.; Bartek, J.; Lukas, J.: Rapid destruction of human Cdc25A in response to DNA damage. *Science* 288:

1425–1429, 2000.

- [64755] 3085.Lane, S. A.; Baker, E.; Sutherland, G. R.; Tonks, I.; Hayward,N.; Ellem, K.: The human cell cycle gene CDC25B is located at 20p13. *Genomics* 15:693–694, 1993.
- [64756] 3086.Lincoln, A. J.; Wickramasinghe, D.; Stein, P.; Schultz, R. M.;Palko, M. E.; De Miguel, M. P.; Tessarollo, L.; Donovan, P. J.: Cdc25bphosphatase is required for resumption of meiosis during oocyte maturation. *NatureGenet.* 30: 446–449, 2002.
- [64757] 3087.Dullaart, R. P.; Hoogenberg, K.; Riemens, S. C.; Groener, J. E.;van Tol, A. Sluiter, W. J.; Stulp, B. K.: Cholesteryl ester transferprotein gene polymorphism is a determinant of HDL cholesterol andof the lipoprotein response to a lipid–lowering diet in type 1 diabetes. *Diabetes* 46:2082–2087, 1997.
- [64758] 3088.Durlach, A.; Clavel, C.; Girard–Globa, A.; Durlach, V.: Sex–dependentassociation of a genetic polymorphism of cholesteryl ester transferprotein with high–density lipoprotein cholesterol and macrovascularpathology in type II diabetic patients. *J. Clin. Endocr. Metab.* 84:3656–3659, 1999.
- [64759] 3089.Freeman, D. J.; Griffin, B. A.; Holmes, A. P.; Lindsay, G. M.;Gaffney, D.; Packard, C. J.; Shepherd, J.: Regulation

of plasma HDLcholesterol and subfraction distribution by genetic and environmental factors: associations between the TaqI B RFLP in the CETP gene and smoking and obesity. *Arterioscler. Thromb.* 14: 336–344, 1994.

[64760] 3090. Hannuksela, M. L.; Liinamaa, M. J.; Kesaniemi, Y. A.; Savolainen, M. J.: Relation of polymorphisms in the cholesteryl ester transfer protein gene to transfer protein activity and plasma lipoprotein levels in alcohol drinkers. *Atherosclerosis* 110: 35–44, 1994.

[64761] 3091. Herrera, V. L. M.; Makrides, S. C.; Xie, H. X.; Adari, H.; Krauss, R. M.; Ryan, U. S.; Ruiz-Opazo, N.: Spontaneous combined hyperlipidemia, coronary heart disease and decreased survival in Dahl salt-sensitive hypertensive rats transgenic for human cholesteryl ester transfer protein. *Nature Med.* 5: 1383–1389, 1999.

[64762] 3092. Fumeron, F.; Betoulle, D.; Luc, G.; Behague, I.; Ricard, S.; Poirier, O.; Jemaa, R.; Evans, A.; Arveiler, D.; Marques-Vidal, P.; Bard, J.-M.; Fruchart, J.-C.; Ducimetiere, P.; Apfelbaum, M.; Cambien, F.: Alcohol intake modulates the effect of a polymorphism of the cholesteryl ester transfer protein gene on plasma high density lipoprotein and the risk of myocardial infarction. *J. Clin. Invest.* 96: 1664–1671, 1995.

- [64763] 3093.Inazu, A.; Jiang, X.-C.; Haraki, T.; Yagi, K.; Kamon, N.; Koizumi,J.; Mabuchi, H.; Takeda, R.; Takata, K.; Moriyama, Y.; Doi, M.; Tall,A.: Genetic cholesteryl ester transfer protein deficiency causedby two prevalent mutations as a major determinant of increased levelsof high density lipoprotein cholesterol. *J. Clin. Invest.* 94: 1872–1882,1994.
- [64764] 3094.Kondo, I.; Berg, K.; Drayna, D.; Lawn, R.: DNA polymorphism atthe locus for human cholesteryl ester transfer protein (CETP) is associatedwith high density lipoprotein cholesterol and apolipoprotein levels. *Clin.Genet.* 35: 49–56, 1989.
- [64765] 3095.Kuivenhoven, J. A.; de Knijff, P.; Boer, J. M. A.; Smalheer, H.A.; Botma, G. J.; Seidell, J. C.; Kastelein, J. J.; Pritchard, P.H.: Heterogeneity at the CETP gene locus: influence on plasma CETPconcentrations and HDL cholesterol levels. *Arterioscler. Thromb.Vasc. Biol.* 17: 560–568, 1997.
- [64766] 3096.Kuivenhoven, J. A.; Jukema, J. W.; Zwinderman, A. H.; de Knijff,P.; McPherson, R.; Bruschke, A. V. G.; Lie, K. I.; Kastelein, J. J.P.: The role of a common variant of the cholesteryl ester transferprotein gene in the progression of coronary atherosclerosis. *NewEng. J. Med.* 338: 86–93,

1998.

- [64767] 3097.Lusis, A. J.; Zollman, S.; Sparkes, R. S.; Klisak, I.; Mohandas, T.; Drayna, D.; Lawn, R. M.: Assignment of the human gene for cholesterylester transfer protein to chromosome 16q12–16q21. *Genomics* 1: 232–242, 1987.
- [64768] 3098.Bentley, K. L.; Ferguson–Smith, A. C.; Miki, T.; Kidd, K. K.; Ruddle, F. H.: Physical linkage of Hox 2.1 and nerve growth factor receptor.(Abstract) *Cytogenet. Cell Genet.* 51: 961 only, 1989.
- [64769] 3099.Bibel, M.; Barde, Y.–A.: Neurotrophins: key regulators of cell fate and cell shape in the vertebrate nervous system. *Genes Dev.* 14:2919–2937, 2000.
- [64770] 3100.Bothwell, M.: p75(NTR): a receptor after all. *Science* 272: 506–507, 1996.
- [64771] 3101.Carter, B. D.; Kaltschmidt, C.; Kaltschmidt, B.; Offenhauser, N.; Bohm–Matthaei, R.; Baeuerle, P. A.; Barde, Y.–A.: Selective activation of NF–kappa–B by nerve growth factor through the neurotrophin receptor p75. *Science* 272: 542–545, 1996.
- [64772] 3102.Carter, B. D.; Lewin, G. R.: Neurotrophins live or let die: does p75(NTR) decide? *Neuron* 18: 187–190, 1997.
- [64773] 3103.Chao, M. V.; Bothwell, M. A.; Ross, A. H.; Koprowski, H.; Lanahan, A. A.; Buck, C. R.; Sehgal, A.: Gene transfer

and molecular cloning of the human NGF receptor. *Science* 232: 518–521, 1986.

[64774] 3104. Dobrowsky, R. T.; Werner, M. H.; Castellino, A. M.; Chao, M. V.; Hannun, Y. A.: Activation of the sphingomyelin cycle through the low-affinity neurotrophin receptor. *Science* 265: 1596–1599, 1994.

[64775] 3105. Colonna, M.; Bresnahan, M.; Bahram, S.; Strominger, J. L.; Spies, T.: Allelic variants of the human putative peptide transporter involved in antigen processing. *Proc. Nat. Acad. Sci.* 89: 3932–3936, 1992.

[64776] 3106. Huh, G. S.; Boulanger, L. M.; Du, H.; Riquelme, P. A.; Brotz, T. M.; Shatz, C. J.: Functional requirement for class I MHC in CNS development and plasticity. *Science* 290: 2155–2159, 2000.

[64777] 3107. Karttunen, J. T.; Lehner, P. J.; Gupta, S. S.; Hewitt, E. W.; Cresswell, P.: Distinct functions and cooperative interaction of the subunits of the transporter associated with antigen processing (TAP). *Proc. Nat. Acad. Sci.* 98: 7431–7436, 2001.

[64778] 3108. Cullen, M.; Erlich, H.; Klitz, W.; Carrington, M.: Molecular mapping of a recombination hotspot located in the second intron of the human TAP2 locus. *Am. J. Hum. Genet.* 56: 1350–1358, 1995.



- [64779] 3109.de la Salle, H.; Donato, L.; Zimmer, J; Plebani, A.; Hanau, D.;Bonnevill, M.; Tongio, M.–M.: HLA class I deficiencies.In: Ochs,H. D.; Smith, C. I. E.; Puck, J. M. (eds.): Primary ImmunodeficiencyDiseases: A Molecular and Genetic Approach. New York: Oxford UniversityPress 1999. Pp. 181–188.
- [64780] 3110.de la Salle, H.; Hanau, D.; Fricker, D.; Urlacher, A.; Kelly, A.;Salamero, J.; Powis, S. H.; Donato, L.; Bausinger, H.; Laforet, M.;Jeras, M.; Spehner, D.; Bieber, T.; Falkenrodt, A.; Cazenave, J.–P.;Trowsdale, J.; Tongio, M.–M.: Homozygous human TAP peptide transporter mutation in HLA class I deficiency. Science 265: 237–241, 1994.
- [64781] 3111.Jeffreys, A. J.; Ritchie, A.; Neumann, R.: High resolution analysis of haplotype diversity and meiotic crossover in the human TAP2 recombination hotspot. Hum. Molec. Genet. 9: 725–733, 2000.
- [64782] 3112.Powis, S. H.; Mockridge, I.; Kelly, A.; Kerr, L.–A.; Glynn, R.;Gileadi, U.; Beck, S.; Trowsdale, J.: Polymorphism in a second ABC transporter gene located within the class II region of the human major histocompatibility complex. Proc. Nat. Acad. Sci. 89: 1463–1467, 1992.
- [64783] 3113.Glauder, J.; Ragg, H.; Rauch, J.; Engels, J. W.: Human peptidylglycine  $\alpha$ -amidating monooxygenase: cDNA,

cloning and functional expression of a truncated form in COS cells. *Biochem. Biophys. Res. Commun.* 169:551–558, 1990.

- [64784] 3114. Lossie, A. C.; Eipper, B. A.; Hand, T. A.; Camper, S. A.: Localization of the peptidylglycine alpha-amidating monooxygenase gene (Pam) introduces a region of homology between human chromosome 5q and mouse chromosome 1. *Mammalian Genome* 5: 738–739, 1994.
- [64785] 3115. Ouafik, L'H.; Mattei, M. G.; Giraud, P.; Oliver, C.; Eipper, B.A.; Mains, R. E.: Localization of the gene encoding peptidylglycine alpha-amidating monooxygenase (PAM) to human chromosome 5q14–5q21. *Genomics* 18: 319–321, 1993.
- [64786] 3116. Jackson, R. S.; Creemers, J. W. M.; Ohagi, S.; Raffin-Sanson, M.-L.; Sanders, L.; Montague, C. T.; Hutton, J. C.; O'Rahilly, S.: Obesity and impaired prohormone processing associated with mutations in the human prohormone convertase 1 gene. *Nature Genet.* 16: 303–306, 1997.
- [64787] 3117. Naggert, J. K.; Fricker, L. D.; Varlamov, O.; Nishina, P. M.; Rouille, Y.; Steiner, D. F.; Carroll, R. J.; Paigen, B. J.; Leiter, E. H.: Hyperproinsulinaemia in obese fat/fat mice associated with a carboxypeptidase E mutation which reduces enzyme activity. *Nature Genet.* 10: 135–142, 1995.

- [64788] 3118.O'Rahilly, S.; Gray, H.; Humphreys, P. J.; Krook, A.; Polonsky, K. S.; White, A.; Gibson, S.; Taylor, K.; Carr, C.: Brief report: impaired processing of prohormones associated with abnormalities of glucose homeostasis and adrenal function. *New Eng. J. Med.* 333:1386–1390, 1995.
- [64789] 3119.Ohagi, S.; Sakaguchi, H.; Sanke, T.; Tatsuta, H.; Hanabusa, T.; Nanjo, K.: Human prohormone convertase 3 gene: exon–intron organization and molecular scanning for mutations in Japanese subjects with NIDDM. *Diabetes* 45:897–901, 1996.
- [64790] 3120.Furuta, M.; Carroll, R.; Martin, S.; Swift, H.; Ravazzola, M.; Orci, L.; Steiner, D.: Incomplete processing of proinsulin to insulin accompanied by elevation of Des-31,32 proinsulin intermediates in islets of mice lacking active PC2. *J. Biol. Chem.* 273: 3431–3437, 1998.
- [64791] 3121.Furuta, M.; Yano, H.; Zhou, A.; Rouille, Y.; Holst, J.; Carroll, R.; Ravazzola, M.; Orci, L.; Furuta, H.; Steiner, D.: Defective prohormone processing and altered pancreatic islet morphology in mice lacking active SPC2. *Proc. Nat. Acad. Sci.* 94: 6646–6651, 1997.
- [64792] 3122.Furuta, M.; Zhou, A.; Webb, G.; Carroll, R.; Ravazzola, M.; Orci, L.; Steiner, D. F.: Severe defect in proglucagon processing in islet A-cells of prohormone

convertase 2 null mice. J. Biol. Chem. 276:27197–27202, 2001.

[64793] 3123. Gabreels, B. A. T. F.; Swaab, D. F.; de Kleijn, D. P. V.; Seidah, N. G.; Van de Loo, J.-W.; Van de Ven, W. J. M.; Martens, G. J. M. and van Leeuwen, F. W.: Attenuation of the polypeptide 7B2, prohormone convertase PC2, and vasopressin in the hypothalamus of some Prader-Willi patients: indications for a processing defect. J. Clin. Endocr. Metab. 83:591–599, 1998.

[64794] 3124. Maglott, D. R.; Feldblyum, T. V.; Durkin, A. S.; Nierman, W. C.: Radiation hybrid mapping of SNAP, PCSK2, and THBD (human chromosome 20p). Mammalian Genome 7: 400–401, 1996.

[64795] 3125. Ohagi, S.; LaMendola, J.; LeBeau, M. M.; Espinosa, R., III; Takeda, J.; Smeekens, S. P.; Chan, S. J.; Steiner, D. F.: Identification and analysis of the gene encoding human PC2, a prohormone convertase expressed in neuroendocrine tissues. Proc. Nat. Acad. Sci. 89: 4977–4981, 1992.

[64796] 3126. Taylor, N. A.; Shennan, K. I. J.; Cutler, D. F.; Docherty, K.: Mutations within the propeptide, the primary cleavage site or the catalytic site, or deletion of C-terminal sequences, prevents secretion of proPC2 from transfected

COS-7 cells. *Biochem. J.* 321: 367-373, 1997.

- [64797] 3127. Chan, W.-Y.; Liu, Q.-R.; Borjigin, J.; Busch, H.; Renert, O. M.; Tease, L. A.; Chan, P.-K.: Characterization of the cDNA encoding human nucleophosmin and studies of its role in normal and abnormal growth. *Biochemistry* 28: 1033-1039, 1989.
- [64798] 3128. Cheng, G.-X.; Zhu, X.-H.; Men, X.-Q.; Wang, L.; Huang, Q.-H.; Jin, X. L.; Xiong, S.-M.; Zhu, J.; Guo, W.-M.; Chen, J.-Q.; Xu, S.-F.; So, E.; Chan, L.-C.; Waxman, S.; Zelen, A.; Chen, G.-Q.; Dong, S.; Liu, J.-X.; Chen, S.-J.: Distinct leukemia phenotypes in transgenic mice and different corepressor interactions generated by promyelocytic leukemia variant fusion genes PLZF-RAR-alpha and NPM-RAR-alpha. *Proc. Nat. Acad. Sci.* 96: 6318-6323, 1999.
- [64799] 3129. Dutta, S.; Akey, I. V.; Dingwall, C.; Hartman, K. L.; Laue, T.; Nolte, R. T.; Head, J. F.; Akey, C. W.: The crystal structure of nucleoplasmin-core: implications for histone binding and nucleosome assembly. *Molec. Cell* 8: 841-853, 2001.
- [64800] 3130. Okuda, M.; Horn, H. F.; Tarapore, P.; Tokuyama, Y.; Smulian, A. G.; Chan, P.-K.; Knudsen, E. S.; Hofmann, I. A.; Snyder, J. D.; Bove, K. E.; Fukasawa, K.: Nucleophosmin/B23 is a target of CDK2/cyclin E in centrosome duplication.

Cell 103: 127–140, 2000.

- [64801] 3131.Redner, R. L.; Rush, E. A.; Faas, S.; Rudert, W. A.; Corey, S.J.: The t(5;17) variant of acute promyelocytic leukemia expresses a nucleophosmin–retinoic acid receptor fusion. *Blood* 87: 882–886, 1996.
- [64802] 3132.Moore, M. D.; Cooper, N. R.; Tack, B. F.; Nemerow, G. R.: Molecular cloning of the cDNA encoding the Epstein–Barr virus/C3d receptor (complement receptor type 2) of human B lymphocytes. *Proc. Nat. Acad. Sci.* 84:9194–9198, 1987.
- [64803] 3133.Prota, A. E.; Sage, D. R.; Stehle, T.; Fingerhuth, J. D.: The crystal structure of human CD21: implications for Epstein–Barr virus and C3d binding. *Proc. Nat. Acad. Sci.* 99: 10641–10646, 2002.
- [64804] 3134.Rodriguez de Cordoba, S.; Rubinstein, P.: Quantitative variations of the C3b/C4b receptor (CR1) in human erythrocyte are controlled by genes within the regulator of complement activation (RCA) gene cluster. *J. Exp. Med.* 164: 1274–1283, 1986.
- [64805] 3135.Szakonyi, G.; Guthridge, J. M.; Li, D.; Young, K.; Holers, V. M.; Chen, X. S.: Structure of complement receptor 2 in complex with its C3d ligand. *Science* 292: 1725–1728, 2001.

- [64806] 3136. Weis, J. J.; Tedder, T. F.; Fearon, D. T.: Identification of a 145,000 M(r) membrane protein as the C3d receptor (CR2) of human B lymphocytes. *Proc. Nat. Acad. Sci.* 81: 881–885, 1984.
- [64807] 3137. Yefenof, E.; Klein, G.; Jondal, M.; Oldstone, M. B. A.: Surface markers on human B- and T-lymphocytes. IX. Two color immunofluorescence studies on the association between EBV receptors and complement receptors on the surface of lymphoid cell lines. *Int. J. Cancer* 17: 693–700, 1976.
- [64808] 3138. Potter, E.; Behan, D. P.; Fischer, W. H.; Linton, E. A.; Lowry, P. J.; Vale, W. W.: Cloning and characterization of the cDNAs for human and rat corticotropin releasing factor-binding proteins. *Nature* 349: 423–426, 1991.
- [64809] 3139. Vamvakopoulos, N. C.; Sioutopoulou, T. O.; Durkin, S. A.; Nierman, W. C.; Wasmuth, J. J.; McPherson, J. D.: Mapping the human corticotropin releasing hormone binding protein gene (CRHBP) to the long arm of chromosome 5 (5q11.2–q13.3). *Genomics* 25: 325–327, 1995.
- [64810] 3140. Arbiser, J. L.; Morton, C. C.; Bruns, G. A. P.; Majzoub, J. A.: Human corticotropin releasing hormone gene is located on the long arm of chromosome 8. *Cytogenet. Cell Genet.* 47: 113–116, 1988.

- [64811] 3141.Behan, D. P.; Heinrichs, S. C.; Troncoso, J. C.; Liu, X.-J.; Kawas,C. H.; Ling, N.; De Souza, E. B.: Displacement of corticotropin releasingfactor from its binding protein as a possible treatment for Alzheimer'sdisease. *Nature* 378: 284–287, 1995.
- [64812] 3142.Cheng, Y.-H.; Nicholson, R. C.; King, B.; Chan, E.-C.; Fitter,J. T.; Smith, R.: Corticotropin–releasing hormone gene expressionin primary placental cells is modulated by cyclic adenosine 3–prime,5–prime–monophosphate. *J.Clin. Endocr. Metab.* 85: 1239–1244, 2000.
- [64813] 3143.Habib, K. E.; Weld, K. P.; Rice, K. C.; Pushkas, J.; Champoux,M.; Listwak, S.; Webster, E. L.; Atkinson, A. J.; Schulkin, J.; Contoreggi,C.; Chrousos, G. P.; McCann, S. M.; Suomi, S. J.; Higley, J. D.; Gold,P. W.: Oral administra–tion of a corticotropin–releasing hormone receptorantag–onist significantly attenuates behavioral, neuroendocrine, andautonomic responses to stress in primates. *Proc. Nat. Acad. Sci.* 97:6079–6084, 2000.
- [64814] 3144.Inder, W. J.; Prickett, T. C. R.; Ellis, M. J.; Hull, L.; Reid,R.; Benny, P. S.; Livesey, J. H.; Donald, R. A.: The util–ity of plasmaCRH as a predictor of preterm delivery. *J. Clin. Endocr. Metab.* 86:5706–5710, 2001.
- [64815] 3145.Kellogg, J.; Luty, J. A.; Thompson, R.; Luo, X. Y.; Ma–



genis, R.E.; Litt, M.: Corticotropin releasing hormone (CRH) maps to human chromosome 8 and identifies a TaqI RFLP. *Cytogenet. Cell Genet.* 51:1022, 1989.

[64816] 3146. Knapp, L. T.; Keegan, C. E.; Seasholtz, A. F.; Camper, S. A.: Corticotropin-releasing hormone (Crh) maps to mouse chromosome 3. *Mammalian Genome* 4: 615–617, 1993.

[64817] 3147. Kylo, J. H.; Collins, M. M.; Vetter, K. L.; Cuttler, L.; Rosenfield, R. L.; Donohue, P. A.: Linkage of congenital isolated adrenocorticotrophic hormone deficiency to the corticotropin releasing hormone locus using simple sequence repeat polymorphisms. *Am. J. Med. Genet.* 62: 262–267, 1996.

[64818] 3148. Majzoub, J. A.: Personal Communication. Boston, Mass. 3/3/1995.

[64819] 3149. Mandel, H.; Berant, M.; Gotfried, E.; Hochberg, Z.: Autosomal recessive hypothalamic corticotropin deficiency: a new entity and its metabolic consequences. (Abstract) *Am. J. Hum. Genet.* 47 (suppl.):A66, 1990.

[64820] 3150. McLean, M.; Bisits, A.; Davies, J.; Woods, R.; Lowry, P.; Smith, R.: A placental clock controlling the length of human pregnancy. *Nature Med.* 1: 460–463, 1995.

[64821] 3151. Muglia, L.; Jacobson, L.; Dikkes, P.; Majzoub, J. A.:

Corticotropin-releasing hormone deficiency reveals major fetal but not adult glucocorticoid need. *Nature* 373: 427–432, 1995.

- [64822] 3152. Robinson, B. G.; Emanuel, R. L.; Frim, D. M.; Majzoub, J. A.: Glucocorticoid stimulates expression of corticotropin-releasing hormone gene in human placenta. *Proc. Nat. Acad. Sci.* 85: 5244–5248, 1988.
- [64823] 3153. Scatena, C. D.; Adler, S.: Trans-acting factors dictate the species-specific placental expression of corticotropin-releasing factor genes in choriocarcinoma cell lines. *Endocrinology* 137: 3000–3008, 1996.
- [64824] 3154. Scatena, C. D.; Adler, S.: Characterization of a human-specific regulator of placental corticotropin-releasing hormone. *Molec. Endocr.* 12: 1228–1240, 1998.
- [64825] 3155. Shibahara, S.; Morimoto, Y.; Furutani, Y.; Notake, M.; Takahashi, H.; Shimizu, S.; Horikawa, S.; Numa, S.: Isolation and sequence analysis of the human corticotropin-releasing factor precursor gene. *EMBO J.* 2: 775–779, 1983.
- [64826] 3156. Stratakis, C. A.; Chrousos, G. P.: Neuroendocrinology and pathophysiology of the stress system. *Ann. N.Y. Acad. Sci.* 771: 1–18, 1995.
- [64827] 3157. Stratakis, C. A.; Sarlis, N. J.; Berrettini, W. H.; Badner, J. A.; Chrousos, G. P.; Gershon, E. S.; Detera-Wadleigh, S.

D.: Lack of linkage between the corticotropin-releasing hormone (CRH) gene and bipolar affective disorder. *Molec. Psychiat.* 2: 483–485, 1997.

[64828] 3158. Xu, B.; Sano, T.; Yamada, S.; Li, C. C.; Hirokawa, M.: Expression of corticotropin-releasing hormone messenger ribonucleic acid in human pituitary corticotroph adenomas associated with proliferative potential. *J. Clin. Endocr. Metab.* 85: 1220–1225, 2000.

[64829] 3159. Zouboulis, C. C.; Seltmann, H.; Hiroi, N.; Chen, W.; Young, M.; Oeff, M.; Scherbaum, W. A.; Orfanos, C. E.; McCann, S. M.; Bornstein, S. R.: Corticotropin-releasing hormone: an autocrine hormone that promotes lipogenesis in human sebocytes. *Proc. Nat. Acad. Sci.* 99: 7148–7153, 2002.

[64830] 3160. Chen, R.; Lewis, K. A.; Perrin, M. H.; Vale, W. W.: Expression cloning of a human corticotropin-releasing-factor receptor. *Proc. Nat. Acad. Sci.* 90: 8967–8971, 1993.

[64831] 3161. Dieterich, K. D.; Gundelfinger, E. D.; Ludecke, D. K.; Lehnert, H.: Mutation and expression analysis of corticotropin-releasing factor 1 receptor in adrenocorticotropin-secreting pituitary adenomas. *J. Clin. Endocr. Metab.* 83: 3327–3331, 1998.

[64832] 3162. Grammatopoulos, D.; Dai, Y.; Chen, J.; Karteris, E.;

Papadopoulou,N.; Easton, A. J.; Hillhouse, E. W.: Human corticotropin–releasing hormone receptor: differences in subtype expression between pregnant and nonpregnant myometria. *J. Clin. Endocr. Metab.* 83: 2539–2544, 1998.

[64833] 3163. Leproult, R.; Colecchia, E. F.; L'Hermite–Baleriaux, M.; Van Cauter, E.: Transition from dim to bright light in the morning induces an immediate elevation of cortisol levels. *J. Clin. Endocr. Metab.* 86:151–157, 2001.

[64834] 3164. Liaw, C. W.; Grigoriadis, D. E.; Lovenberg, T. W.; De Souza, E. B.; Maki, R. A.: Localization of ligand–binding domains of human corticotropin–releasing factor receptor: a chimeric receptor approach. *Molec. Endocr.* 11: 980–985, 1997.

[64835] 3165. Lublin, D. M.; Mallinson, G.; Poole, J.; Reid, M. E.; Thompson, E. S.; Ferdman, B. R.; Telen, M. J.; Anstee, D. J.; Tanner, M. J. A.: Molecular basis of reduced or absent expression of decay–accelerating factor in Cromer blood group phenotypes. *Blood* 84: 1276–1282, 1994.

[64836] 3166. Medof, M. E.; Lublin, D. M.; Holers, V. M.; Ayers, D. J.; Getty, R. R.; Leykam, J. F.; Atkinson, J. P.; Tykocinski, M. L.: Cloning and characterization of cDNAs encoding the complete sequence of decay–accelerating factor of human complement. *Proc. Nat. Acad. Sci.* 84: 2007–2011, 1987.

- [64837] 3167. Reid, M. E.; Chandrasekaran, V.; Sausais, L.; Pierre, J.; Bullock, R.: Disappearance of antibodies to Cromer blood group system antigens during mid pregnancy. *Vox Sang.* 71: 48–50, 1996.
- [64838] 3168. Wang, L.; Uchikawa, M.; Tsuneyama, H.; Tokunaga, K.; Tadokoro, K.; Juji, T.: Molecular cloning and characterization of decay-accelerating factor deficiency in Cromer blood group Inab phenotype. *Blood* 91:680–684, 1998.
- [64839] 3169. Danielson, K. G.; Fazzio, A.; Cohen, I.; Cannizzaro, L. A.; Eichstetter, I.; Iozzo, R. V.: The human decorin gene: intron–exon organization, discovery of two alternatively spliced exons in the 5–prime untranslated region, and mapping of the gene to chromosome 12q23. *Genomics* 15:146–160, 1993.
- [64840] 3170. Dyne, K. M.; Valli, M.; Forlino, A.; Mottes, M.; Kresse, H.; Cetta, G.: Deficient expression of the small proteoglycan decorin in a case of severe/lethal osteogenesis imperfecta. *Am. J. Med. Genet.* 63:161–166, 1996.
- [64841] 3171. Ion, A.; Crosby, A. H.; Kremer, H.; Kenmochi, N.; Van Reen, M.; Fenske, C.; Van Der Burgt, I.; Brunner, H. G.; Montgomery, K.: Detailed mapping, mutation analysis, and intragenic polymorphism identification in candidate Noonan syndrome genes MYL2, DCN, EPS8, and RPL6. *J. Med.*

Genet. 37: 884–886, 2000.

- [64842] 3172. McBride, O. W.; Fisher, L. W.; Young, M. F.: Localization of PGI(biglycan, BGN) and PGII (decorin, DCN, PG-40) genes on human chromosomes Xq13–qter and 12q, respectively. *Genomics* 6: 219–255, 1990.
- [64843] 3173. Moscatello, D. K.; Santra, M.; Mann, D. M.; McQuillan, D. J.; Wong, A. J.; Iozzo, R. V.: Decorin suppresses tumor cell growth by activating the epidermal growth factor receptor. *J. Clin. Invest.* 101: 406–412, 1998.
- [64844] 3174. Pulkkinen, L.; Alitalo, T.; Krusius, T.; Peltonen, L.: Expression of decorin in human tissues and cell lines and defined chromosomal assignment of the gene locus (DCN). *Cytogenet. Cell Genet.* 60: 107–111, 1992.
- [64845] 3175. Pulkkinen, L.; Kainulainen, K.; Krusius, T.; Makinen, P.; Schollin, J.; Gustavsson, K.–H.; Peltonen, L.: Deficient expression of the gene coding for decorin in a lethal form of Marfan syndrome. *J. Biol. Chem.* 265: 17780–17785, 1990.
- [64846] 3176. Schollin, J.; Bjarke, B.; Gustavson, K.–H.: Probable homozygotic form of the Marfan syndrome in a newborn child. *Acta Paediat. Scand.* 77: 452–456, 1988.
- [64847] 3177. Scholzen, T.; Solursh, M.; Suzuki, S.; Reiter, R.; Morgan, J. L.; Buchberg, A. M.; Siracusa, L. D.; Iozzo, R. V.: The

murine decorin:complete cDNA cloning, genomic organization, chromosomal assignment,and expression during organogenesis and tissue differentiation. J.Biol. Chem. 269: 28270–28281, 1994.

[64848] 3178.Vogel, K. G.; Clark, P. E.: Small proteoglycan synthesis by skinfibroblasts cultured from elderly donors and patients with defineddefects in types I and III collagen metabolism. Europ. J. Cell Biol. 49:236–243, 1989.

[64849] 3179.Comer, K. A.; Falany, J. L.; Falany, C. N.: Cloning and expressionof human liver dehydroepiandrosterone sulphotransferase. Biochem.J. 289: 233–240, 1993.

[64850] 3180.Durocher, F.; Morissette, J.; Dufort, I.; Simard, J.; Luu-The,V.: Genetic linkage mapping of the dehydroepiandrosterone sulfotransferase(STD) gene on the chromosome 19q13.3 region. Genomics 29: 781–783,1995.

[64851] 3181.Kong, A.-N. T.; Yang, L.; Ma, M.; Tao, D.; Bjornsson, T. D.: Molecularcloning of the alcohol/hydroxysteroid form (hSTa) of sulfotransferasefrom human liver. Biochem. Biophys. Res. Commun. 187: 448–454, 1992.

[64852] 3182.Otterness, D.; Mohrenweiser, H. W.; Brandriff, B. F.; Weinshilboum,R. M.: Dehydroepiandrosterone sulfotransferase gene (STD): localizationto human chromosome

19q13.3. Cytogenet. Cell Genet. 70: 45–47, 1995.

- [64853] 3183.Otterness, D. M.; Her, C.; Aksoy, S.; Kimura, S.; Wieben, E. D.;Weinshilboum, R. M.: Human dehydroepiandrosterone sulfotransferasegene: molecular cloning and structural characterization. DNA CellBiol. 14: 331–341, 1995.
- [64854] 3184.Otterness, D. M.; Wieben, E. D.; Wood, T. C.; Watson, W. G.; Madden,B. J.; McCormick, D. J.; Weinshilboum, R. M.: Human liver dehydroepiandrosteronesulfotransferase: molecular cloning and expression of cDNA. Molec.Pharm. 41: 865–872, 1992.
- [64855] 3185.von Lindern, M.; Fornerod, M.; van Baal, S.; Jaegle, M.; de Wit,T.; Buijs, A.; Grosveld, G.: The translocation (6;9), associatedwith a specific subtype of acute myeloid leukemia, results in thefusion of two genes, dek and can, and the expression of a chimeric,leukemia-specific dek-can mRNA. Molec. Cell. Biol. 12: 1687–1697,1992.
- [64856] 3186.Schulte-Korne, G.; Grimm, T.; Nothen, M. M.; Muller-Myhsok, B.;Cichon, S.; Vogt, I. R.; Propping, P.; Remschmidt, H.: Evidence forlinkage of spelling disability to chromosome 15. (Letter) Am. J.Hum. Genet. 63: 279–282, 1998.
- [64857] 3187.Yoon, J.-W.; Yoon, C.-S.; Lim, H.-W.; Huang, Q. Q.;



Kang, Y.;Pyun, K. H.; Hirasawa, K.; Sherwin, R. S.; Jun, H.-S.: Control of autoimmune diabetes in NOD mice by GAD expression or suppression in beta cells. *Science* 284: 1183–1187, 1999.

[64858] 3188.Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hirose, M.; Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XII. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.* 5: 355–364, 1998.

[64859] 3189.Clancy, K. P.; Berger, R.; Cox, M.; Bleskan, J.; Walton, K. A.; Hart, I.; Patterson, D.: Localization of the L-glutamine synthetase gene to chromosome 1q23. *Genomics* 38: 418–420, 1996.

[64860] 3190.Gibbs, C. S.; Campbell, K. E.; Wilson, R. H.: Sequence of a human glutamine synthetase cDNA. *Nucleic Acids Res.* 15: 6293 only, 1987.

[64861] 3191.Gunnersen, D.; Haley, B.: Detection of glutamine synthetase in the cerebrospinal fluid of Alzheimer diseased patients: a potential diagnostic biochemical marker. *Proc. Nat. Acad. Sci.* 89: 11949–11953, 1992.

[64862] 3192.Helou, K.; Das, A. T.; Lamers, W. H.; Hoovers, J. M. N.; Szpirer, C.; Szpirer, J.; Klinga-Levan, K.; Levan, G.: FISH

mapping of three ammonia metabolism genes (Glul, Cps1, Glud1) in rat, and the chromosomal localization of GLUL in human and Cps1 in mouse. *Mammalian Genome* 8:362–364, 1997.

- [64863] 3193. Pesole, G.; Bozzetti, M. P.; Lanave, C.; Preparata, G.; Saccone, C.: Glutamine synthetase gene evolution: a good molecular clock. *Proc. Nat. Acad. Sci.* 88: 522–526, 1991.
- [64864] 3194. Wang, Y.; Kudoh, J.; Kubota, R.; Asakawa, S.; Minoshima, S.; Shimizu, N.: Chromosomal mapping of a family of human glutamine synthetase genes: functional gene (GLUL) on 1q25, pseudogene (GLULP) on 9p13, and three related genes (GLULL1, GLULL2, GLULL3) on 5q33, 11p15, and 11q24. *Genomics* 37: 195–199, 1996.
- [64865] 3195. Smith, M. D.; Dawson, S. J.; Latchman, D. S.: Inhibition of neuronal process outgrowth and neuronal specific gene activation by the Brn-3b transcription factor. *J. Biol. Chem.* 272: 1382–1388, 1997.
- [64866] 3196. Horisberger, M. A.; Wathélet, M.; Szpirer, J.; Szpirer, C.; Islam, Q.; Levan, G.; Huez, G.; Content, J.: cDNA cloning and assignment to chromosome 21 of IFI-78K gene, the human equivalent of murine Mx gene. *Somat. Cell Molec. Genet.* 14: 123–131, 1988.
- [64867] 3197. Huber, P.; Aebi, M.; Grob, R.; Pravtcheva, D.; Ruddle,

F.; Haller,O.: Chromosomal localization of two human Mx genes. (Abstract) *Experientia* 44:A84 only, 1988.

[64868] 3198.Li, Y.; Youssoufian, H.: MxA overexpression reveals a common genetic link in four Fanconi anemia complementation groups. *J. Clin. Invest.* 100:2873–2880, 1997.

[64869] 3199.Winqvist, R.; Lundstrom, K.; Salminen, M.; Laatikainen, M.; Ulmanen,I.: Mapping of human catechol-O-methyltransferase gene to 22q11.2 and detection of a frequent RFLP with BglI. (Abstract) *Cytogenet.Cell Genet.* 58: 2051 only, 1991.

[64870] 3200.Bu, L.; Jin, Y.; Shi, Y.; Chu, R.; Ban, A.; Eiberg, H.; Andres,L.; Jiang, H.; Zheng, G.; Qian, M.; Cui, B.; Xia, Y.; Liu, J.; Hu,L.; Zhao, G.; Hayden, M. R.; Kong, X.: Mutant DNA-binding domain of HSF4 is associated with autosomal dominant lamellar and Marner cataract. *Nature Genet.* 31: 276–278, 2002.

[64871] 3201.Eiberg, H.; Marner, E.; Rosenberg, T.; Mohr, J.: Marner's cataract(CAM) assigned to chromosome 16: linkage to haptoglobin. *Clin. Genet.* 34:272–275, 1988.

[64872] 3202.Marner, E.: A family with eight generations of hereditary cataract. *Acta Ophthal.* 27: 537–551, 1949.

[64873] 3203.Pokutta, S.; Weis, W. I.: Structure of the dimerization and beta-catenin-binding region of alpha-catenin. *Molec.*

Cell 5: 533–543, 2000.

[64874] 3204. Bailey, A.; Norris, A. L.; Leek, J. P.; Clissold, P. M.; Carr, I. M.; Ogilvie, D. J.; Morrison, J. F. J.; Meredith, D. M.; Markham, A. F.: Yeast artificial chromosome cloning of the beta-catenin locus on human chromosome 3p21–22. Chromosome Res. 3: 201–203, 1995.

[64875] 3205. Blaker, H.; Hofmann, W. J.; Rieker, R. J.; Penzel, R.; Graf, M.; Otto, H. F.: Beta-catenin accumulation and mutation of the CTNNB1 gene in hepatoblastoma. Genes Chromosomes Cancer 25: 399–402, 1999.

[64876] 3206. Chan, T. A.; Wang, Z.; Dang, L. H.; Vogelstein, B.; Kinzler, K. W.: Targeted inactivation of CTNNB1 reveals unexpected effects of beta-catenin mutation. Proc. Nat. Acad. Sci. 99: 8265–8270, 2002.

[64877] 3207. Chenn, A.; Walsh, C. A.: Regulation of cerebral cortical size by control of cell cycle exit in neural precursors. Science 297: 365–369, 2002.

[64878] 3208. Eastman, Q.; Grosschedl, R.: Regulation of LEF-1/TCF transcription factors by Wnt and other signals. Curr. Opin. Cell Biol. 11: 233–240, 1999.

[64879] 3209. Gat, U.; DasGupta, R.; Degenstein, L.; Fuchs, E.: De novo hair follicle morphogenesis and hair tumors in mice expressing a truncated beta-catenin in skin. Cell 95:

605–614, 1998.

- [64880] 3210.Asakura, H.; Zwain, I. H.; Yen, S. S. C.: Expression of genes encoding corticotropin–releasing factor (CRF), type 1 CRF receptor, and CRF–binding protein and localization of the gene products in the human ovary. *J. Clin. Endocr. Metab.* 82: 2720–2725, 1997.
- [64881] 3211.Lynch, E. D.; Lee, M. K.; Morrow, J. E.; Welcsh, P. L.; Leon, P.E.; King, M.–C.: Nonsyndromic deafness DFNA1 associated with mutation of the human homolog of the *Drosophila* gene *diaphanous*. *Science* 278:1315–1318, 1997.
- [64882] 3212.Hamann, J.; Vogel, B.; van Schijndel, G. M.; van Lier, R. A.: The seven–span transmembrane receptor CD97 has a cellular ligand (CD55, DAF). *J. Exp. Med.* 184: 1185–1189, 1996.
- [64883] 3213.Hourcade, D.; Garcia, A. D.; Post, T. W.; Taillon–Miller, P.; Holers, V. M.; Wagner, L. M.; Bora, N. S.; Atkinson, J. P.: Analysis of the human regulators of complement activation (RCA) gene cluster with yeast artificial chromosomes (YACs). *Genomics* 12: 289–300, 1992.
- [64884] 3214.Lemons, R. S.; Le Beau, M. M.; Lublin, D. M.; Holers, V. M.; Tykocinski, M. L.; Medof, M. E.; Atkinson, J. P.: The gene encoding decay–accelerating factor (DAF) is located

in the complement regulatory locus on the long arm of chromosome 1. (Abstract) *Cytogenet. Cell Genet.* 46:646–647, 1987.

- [64885] 3215. Lublin, D. M.; Lemons, R. S.; Le Beau, M. M.; Holers, V. M.; Tykocinski, M. L.; Medof, M. E.; Atkinson, J. P.: The gene encoding decay-accelerating factor (DAF) is located in the complement regulatory locus on the long arm of chromosome 1. (Abstract) *Clin. Res.* 35: 460A only, 1987.
- [64886] 3216. Swanson, J. M.; Flodman, P.; Kennedy, J.; Spence, M. A.; Moyzis, R.; Schuck, S.; Murias, M.; Moriarity, J.; Barr, C.; Smith, M.; Posner, M.: Dopamine genes and ADHD. *Neurosci. Behav. Rev.* 24: 21–25, 2000.
- [64887] 3217. Swanson, J. M.; Sunohara, G. A.; Kennedy, J. L.; Regino, R.; Fineberg, E.; Wigal, T.; Lerner, M.; Williams, L.; LaHoste, G. J.; Wigal, S.: Association of the dopamine receptor D4 (DRD4) gene with a refined phenotype of attention deficit hyperactivity disorder (ADHD): a family-based approach. *Molec. Psychiat.* 3: 38–41, 1998.
- [64888] 3218. Tomitaka, M.; Tomitaka, S.; Otuka, Y.; Kim, K.; Matuki, H.; Sakamoto, K.; Tanaka, A.: Association between novelty seeking and dopamine receptor D4 (DRD4) exon III polymorphism in Japanese subjects. *Am. J. Med. Genet. (Neuropsychiat. Genet.)* 88: 469–471, 1999.

- [64889] 3219. Van Tol, H. H. M.; Bunzow, J. R.; Guan, H. C.; Sunahara, R. K.; Seeman, P.; Niznik, H. B.; Civelli, O.: Cloning of the gene for a human dopamine D4 receptor with high affinity for the antipsychotic clozapine. *Nature* 350: 610–614, 1991.
- [64890] 3220. Van Tol, H. H. M.; Wu, C. M.; Guan, H.-C.; Ohara, K.; Bunzow, J. R.; Civelli, O.; Kennedy, J.; Seeman, P.; Niznik, H. B.; Jovanovic, V.: Multiple dopamine D4 receptor variants in the human population. *Nature* 358:149–152, 1992.
- [64891] 3221. Wiesbeck, G. A.; Mauerer, C.; Thome, J.; Jakob, F.; Boening, J.: Neuroendocrine support for a relationship between 'novelty seeking' and dopaminergic function in alcohol-dependent men. *Psychoneuroendocrinology* 20:755–761, 1995.
- [64892] 3222. Beischlag, T. V.; Marchese, A.; Meador-Woodruff, J. H.; Damask, S. P.; O'Dowd, B. F.; Tyndale, R. F.; Van Tol, H. H. M.; Seeman, P.; Niznik, H. B.: The human dopamine D5 receptor gene: cloning and characterization of the 5-prime flanking and promoter region. *Biochemistry* 34: 5960–5970, 1995.
- [64893] 3223. Eubanks, J. H.; Altherr, M.; Wagner-McPherson, C.; McPherson, J. D.; Wasmuth, J. J.; Evans, G. A.: Localization

of the D5 dopaminereceptor gene to human chromosome 4p15.1–p15.3, centromeric to theHuntington's disease locus. *Genomics* 12: 510–516, 1992.

[64894] 3224.Grandy, D. K.; Allen, L. J.; Zhang, Y.; Magenis, R. E.; Civelli,O.: Chromosomal localization of three human D5 dopamine receptorgenes. *Genomics* 13: 968–973, 1992.

[64895] 3225.Grandy, D. K.; Zhang, Y.; Bouvier, C.; Zhou, Q.–Y.; Johnson, R.A.; Allen, L.; Buck, K.; Bunzow, J. R.; Salon, J.; Civelli, O.: Multiplehuman D5 dopamine receptor genes: a functional receptor and two pseudogenes. *Proc.Nat. Acad. Sci.* 88: 9175–9179, 1991.

[64896] 3226.Grosson, C. L. S.; MacDonald, M. E.; Duyao, M. P.; Ambrose, C.M.; Roffler–Tarlov, S.; Gusella, J. F.: Syntenic conservation of theHuntington's disease gene and surrounding loci on mouse chromosome5. *Mammalian Genome* 5: 424–428, 1994.

[64897] 3227.Liu, F.; Wan, Q.; Pristupa, Z. B.; Yu, X.–M.; Wang, Y. T.; Niznik,H. B.: Direct protein–protein coupling enables cross–talk between dopamine D5 and gamma–aminobutyric acid A receptors. *Nature* 403:274–280, 2000.

[64898] 3228.Misbahuddin, A.; Placzek, M. R.; Chaudhuri, K. R.; Wood, N. W.;Bhatia, K. P.; Warner, T. T.: A polymorphism



in the dopamine receptor DRD5 is associated with blepharospasm. *Neurology* 58: 124–126, 2002.

- [64899] 3229. Placzek, M. R.; Misbahuddin, A.; Chaudhuri, K. R.; Wood, N. W.; Bhatia, K. P.; Warner, T. T.: Cervical dystonia is associated with a polymorphism in the dopamine (D5) receptor gene. *J. Neurol. Neurosurg. Psychiat.* 71: 262–264, 2001.
- [64900] 3230. Polymeropoulos, M. H.; Xiao, H.; Merrill, C. R.: The human D5 dopamine receptor (DRD5) maps on chromosome 4. *Genomics* 11: 777–778, 1991.
- [64901] 3231. Sherrington, R.; Mankoo, B.; Attwood, J.; Kalsi, G.; Curtis, D.; Buetow, K.; Povey, S.; Gurling, H.: Cloning of the human dopamine D5 receptor gene and identification of a highly polymorphic microsatellite for the DRD5 locus that shows tight linkage to the chromosome 4p reference marker RAF1P1. *Genomics* 18: 423–425, 1993.
- [64902] 3232. Tiberi, M.; Jarvie, K. R.; Silvia, C.; Falardeau, P.; Gingrich, J. A.; Godinot, N.; Bertrand, L.; Yang-Feng, T. L.; Fremeau, R. T., Jr.; Caron, M. G.: Cloning, molecular characterization, and chromosomal assignment of a gene encoding a second D-1 dopamine receptor subtype: differential expression pattern in rat brain compared with the D-1A receptor. *Proc. Nat. Acad. Sci.* 88:

7491–7495, 1991.

- [64903] 3233. Weinshank, R. L.; Adham, N.; Macchi, M.; Olsen, M. A.; Branchek, T. A.; Hartig, P. R.: Molecular cloning and characterization of a high affinity dopamine receptor (D-1-beta) and its pseudogene. *J. Biol. Chem.* 266: 22427–22435, 1991.
- [64904] 3234. Byerley, W.; Hoff, M.; Holik, J.; Caron, M. G.; Giros, B.: VNTR polymorphism for the human dopamine transporter gene (DAT1). *Hum. Molec. Genet.* 2: 335, 1993.
- [64905] 3235. Cook, E. H., Jr.; Stein, M. A.; Krasowski, M. D.; Cox, N. J.; Olkon, D. M.; Kieffer, J. E.; Leventhal, B. L.: Association of attention-deficit disorder and the dopamine transporter gene. *Am. J. Hum. Genet.* 56:993–998, 1995.
- [64906] 3236. Doucette-Stamm, L.; Blakely, D. J.; Tian, J.; Mockus, S.; Mao, J.: Population genetic study of the human dopamine transporter gene (DAT1). *Genet. Epidemiol.* 12: 303–308, 1995.
- [64907] 3237. Gainetdinov, R. R.; Wetsel, W. C.; Jones, S. R.; Levin, E. D.; Jaber, M.; Caron, M. G.: Role of serotonin in the paradoxical calming effect of psychostimulants on hyperactivity. *Science* 283: 397–401, 1999.
- [64908] 3238. Gelernter, J.; Vandenbergh, D.; Kruger, S. D.; Pauls, D. L.; Kurlan, R.; Pakstis, A. J.; Kidd, K. K.; Uhl, G.: The

dopamine transporter protein gene (SLC6A3): primary linkage mapping and linkage studies in Tourette syndrome. *Genomics* 30: 459–463, 1995.

- [64909] 3239. Gill, M.; Daly, G.; Heron, S.; Hawi, Z.; Fitzgerald, M.: Confirmation of association between attention deficit hyperactivity disorder and a dopamine transporter polymorphism. *Molec. Psychiat.* 2: 311–313, 1997.
- [64910] 3240. Giros, B.; El Mestikawy, S.; Godinot, N.; Zheng, K.; Han, H.; Yang-Feng, T.; Caron, M. G.: Cloning, pharmacological characterization, and chromosome assignment of the human dopamine transporter. *Molec. Pharm.* 42: 383–390, 1992.
- [64911] 3241. Giros, B.; Jaber, M.; Jones, S. R.; Wightman, R. M.; Caron, M. G.: Hyperlocomotion and indifference to cocaine and amphetamine in mice lacking the dopamine receptor. *Nature* 370: 606–612, 1996.
- [64912] 3242. Goldman, D.: Dopamine transporter, alcoholism and other diseases. *Nature Med.* 1: 624–625, 1995.
- [64913] 3243. Korn, W. T.; Schatzki, S. C.; DiSciullo, A. J.; Scully, R. E.: Papillary cystadenoma of the broad ligament in von Hippel–Lindau disease. *Am. J. Obstet. Gynec.* 163: 596–598, 1990.
- [64914] 3244. Isnard, R.; Kalotka, H.; Durr, A.; Cossee, M.; Schmitt,

M.; Pousset, F.; Thomas, D.; Brice, A.; Koenig, M.; Komajda, M.: Correlation between left ventricular hypertrophy and GAA trinucleotide repeat length in Friedreich's ataxia. *Circulation* 95: 2247–2249, 1997.

[64915] 3245. Koutnikova, H.; Campuzano, V.; Foury, F.; Dolle, P.; Cazzalini, O.; Koenig, M.: Studies of human, mouse and yeast homologues indicate a mitochondrial function for frataxin. *Nature Genet.* 16: 345–351, 1997.

[64916] 3246. Kromberg, J. G. R.; Jenkins, T.: Albinism in the South African Negro. III. Genetic counselling issues. *J. Biosoc. Sci.* 16: 99–108, 1984.

[64917] 3247. Kugelman, T. P.; Van Scott, E. J.: Tyrosinase activity in melanocytes of human albinos. *J. Invest. Derm.* 37: 73–76, 1961.

[64918] 3248. Lee, S.-T.; Nicholls, R. D.; Bunday, S.; Laxova, R.; Musarella, M.; Spritz, R. A.: Mutations of the P gene in oculocutaneous albinism, ocular albinism, and Prader-Willi syndrome plus albinism. *New Eng. J. Med.* 330: 529–534, 1994.

[64919] 3249. Lee, S.-T.; Nicholls, R. D.; Jong, M. T. C.; Fukai, K.; Spritz, R. A.: Organization and sequence of the human P gene and identification of a new family of transport proteins. *Genomics* 26: 354–363, 1995.

- [64920] 3250.Lee, S.-T.; Nicholls, R. D.; Schnur, R. E.; Guida, L. C.; Lu-Kuo,J.; Spinner, N. B.; Zackai, E. H.; Spritz, R. A.: Diverse mutations of the P gene among African-Americans with type II (tyrosinase-positive) oculocutaneous albinism (OCA2). *Hum. Molec. Genet.* 3: 2047-2051,1994.
- [64921] 3251.Ludowese, C. J.; Thompson, K. J.; Sekhon, G. S.; Pauli, R. M.: Absence of predictable phenotypic expression in proximal 15q duplications. *Clin.Genet.* 40: 194-201, 1991.
- [64922] 3252.Manga, P.: Identification and molecular characterisation of the genes for brown and rufous oculocutaneous albinism in southern Africa. Ph.D.Thesis: University of the Witwatersrand, Johannesburg , 1997.
- [64923] 3253.Manga, P.; Kromberg, J. G. R.; Turner, A.; Jenkins, T.; Ramsay,M.: In southern Africa, brown oculocutaneous albinism (BOCA) mapsto the OCA2 locus on chromosome 15q: P-gene mutations identified. *Am.J. Hum. Genet.* 68: 782-787, 2001.
- [64924] 3254.Baker, E.; Chen, L. Z.; Smith, C. A.; Callen, D. F.; Goodwin, R.;Sutherland, G. R.: Chromosomal location of the human tumor necrosisfactor receptor genes. *Cytogenet. Cell Genet.* 57: 117-118, 1991.
- [64925] 3255.Chan, F. K.-M.; Chun, H. J.; Zheng, L.; Siegel, R. M.;

Bui, K.L.; Lenardo, M. J.: A domain in TNF receptors that mediates ligand-independent receptor assembly and signaling. *Science* 288: 2351–2354, 2000.

[64926] 3256. Atuk, N. O.; McDonald, T.; Wood, T.; Carpenter, J. T.; Walzak, M. P.; Donaldson, M.; Gillenwater, J. Y.; Turner, S. M.; Westfall, V.: Familial pheochromocytoma, hypercalcemia, and von Hippel–Lindau disease: a ten-year study of a large family. *Medicine* 58: 209–218, 1979.

[64927] 3257. Bender, B. U.; Altehofer, C.; Januszewicz, A.; Gartner, R.; Schmidt, H.; Hoffmann, M. M.; Heidemann, P. H.; Neumann, H. P. H.: Functioning thoracic paraganglioma: association with Von Hippel–Lindau syndrome. *J. Clin. Endocr. Metab.* 82: 3356–3360, 1997.

[64928] 3258. Bender, B. U.; Eng, C.; Olschewski, M.; Berger, D. P.; Laubenberger, J.; Altehofer, C.; Kirste, G.; Orszagh, M.; van Velthoven, V.; Miosczka, H.; Schmidt, D.; Neumann, H. P. H.: VHL c.505 T–C mutation confers a high age related penetrance but no increased overall mortality. *J. Med. Genet.* 38: 508–514, 2001.

[64929] 3259. Bender, B. U.; Gutsche, M.; Glasker, S.; Muller, B.; Kirste, G.; Eng, C.; Neumann, H. P. H.: Differential genetic alterations in von Hippel–Lindau syndrome-associated and sporadic pheochromocytomas. *J. Clin. Endocr. Metab.* 85:

4568–4574, 2000.

- [64930] 3260. Bonnet, P.; Dechaume, J.; Blanc, E.: L'anevrisme cirsoïde de l'artère lenticulaire l'anevrisme racémeux, ses relations avec l'anevrisme cirsoïde de la face et l'anevrisme cirsoïde du cerveau. *Bull. Soc. Ophtal. Franc.* 51: 521–524, 1938.
- [64931] 3261. Bradley, J. F.; Collins, D. L.; Schimke, R. N.; Parrott, H. N.; Rothberg, P. G.: Two distinct phenotypes caused by two different missense mutations in the same codon of the VHL gene. *Am. J. Med. Genet.* 87: 163–167, 1999.
- [64932] 3262. Brauch, H.; Kishida, T.; Glavac, D.; Chen, F.; Pausch, F.; Hofler, H.; Latif, F.; Lerman, M. I.; Zbar, B.; Neumann, H. P. H.: Von Hippel–Lindau (VHL) disease with pheochromocytoma in the Black Forest region of Germany: evidence for a founder effect. *Hum. Genet.* 95: 551–556, 1995.
- [64933] 3263. Brown, D. G.; Hilal, S. K.; Tenner, M. S.: Wyburn–Mason syndrome. *Arch. Neurol.* 28: 67–68, 1973.
- [64934] 3264. Cahill, G. F.; Melicow, M. M.; Guerry, D.: The renal lesions in von Hippel–Lindau's disease. *Trans. Am. Assoc. Genitourinary Surg.* 35: 271–281, 1942.
- [64935] 3265. Cameron, S. J.; Doig, A.: Cerebellar tumors presented with clinical features of pheochromocytoma. *Lancet* I: 492–494, 1970.
- [64936] 3266. Chapman, R. C.; Diaz–Perez, R.: Pheochromocytoma

associated with cerebellar hemangioblastoma. J.A.M.A. 182: 1014–1017, 1962.

- [64937] 3267. Chen, F.; Slife, L.; Kishida, T.; Mulvihill, J.; Tisherman, S.E.; Zbar, B.: Genotype–phenotype correlation in von Hippel–Lindau disease: identification of a mutation associated with VHL type 2A. J. Med. Genet. 33: 716–717, 1996.
- [64938] 3268. Christoferson, L. A.; Gustafson, M. B.; Petersen, A. G.: Von Hippel–Lindau's disease. J.A.M.A. 178: 280–282, 1961.
- [64939] 3269. Chen, F.; Kishida, T.; Yao, M.; Hustad, T.; Glavac, D.; Dean, M.; Gnarr, J. R.; Orcutt, M. L.; Duh, F. M.; Glenn, G.; Green, J.; Hsia, Y. E.; Lamiell, J.; Li, H.; Wei, M. H.; Schmidt, L.; Tory, K.; Kuzman, I.; Stackhouse, T.; Latif, F.; Linehan, W. M.; Lerman, M.; Zbar, B.: Germline mutations in the von Hippel–Lindau disease tumor suppressor gene: correlations with phenotype. Hum. Mutat. 5: 66–75, 1995.
- [64940] 3270. Collins, E. T.: Intra-ocular growths (two cases, brother and sister, with peculiar vascular new growth, probably primarily retinal, affecting both eyes). Trans. Ophthal. Soc. U.K. 14: 141–149, 1894.
- [64941] 3271. Crossey, P. A.; Foster, K.; Richards, F. M.; Phipps, M. E.; Latif, F.; Tory, K.; Jones, M. H.; Bentley, E.; Kumar, R.;



Lerman, M. I.; Zbar, B.; Affara, N. A.; Ferguson-Smith, M. A.; Maher, E. R.: Molecular genetic investigations of the mechanism of tumorigenesis in von Hippel-Lindau disease: analysis of allele loss in VHL tumours. *Hum. Genet.* 93:53-58, 1994.

[64942] 3272. Crossey, P. A.; Richards, F. M.; Foster, K.; Green, J. S.; Prowse, A.; Latif, F.; Lerman, M. I.; Zbar, B.; Affara, N. A.; Ferguson-Smith, M. A.; Maher, E. R.: Identification of intragenic mutations in the von Hippel-Lindau disease tumour suppressor gene and correlation with disease phenotype. *Hum. Molec. Genet.* 3: 1303-1308, 1994.

[64943] 3273. Cushing, H.; Bailey, P.: Hemangiomas of cerebellum and retina (Lindau's disease), with the report of a case. *Arch. Ophthalmol.* 57:447-463, 1928.

[64944] 3274. Davies, D. R.; Norman, A. M.; Whitehouse, R. W.; Evans, D. G. R.: Non-expression of von Hippel-Lindau phenotype in an obligate gene carrier. *Clin. Genet.* 45: 104-106, 1994.

[64945] 3275. Decker, H.-J. H.; Neumann, H. P. H.; Walter, T. A.; Sandberg, A. A.: 3p involvement in a renal cell carcinoma in von Hippel-Lindau syndrome: region of tumor breakpoint clustering on 3p? *Cancer Genet. Cytogenet.* 33: 59-65, 1988.

- [64946] 3276.Duan, D. R.; Humphrey, J. S.; Chen, D. Y. T.; Weng, Y.; Sukegawa,J.; Lee, S.; Gnarra, J. R.; Linehan, W. M.; Klausner, R. D.: Characterizationof the VHL tumor suppressor gene product: localization, complex formation,and the effect of natural inactivating mutations. Proc. Nat. Acad.Sci. 92: 6459–6463, 1995.
- [64947] 3277.Duan, D. R.; Pause, A.; Burgess, W. H.; Aso, T.; Chen, D. Y. T.;Garrett, K. P.; Conaway, R. C.; Conaway, J. W.; Linehan, W. M.; Klausner,R. D.: Inhibition of transcription elongation by the VHL tumor suppressorprotein. Science 269: 1402–1406, 1995.
- [64948] 3278.Schoof, E.; Girstl, M.; Frobenius, W.; Kirschbaum, M.; Dorr, H.G.; Rascher, W.; Dotsch, J.: Decreased gene expression of 11–beta–hydroxysteroiddehydrogenase type 2 and 15–hydroxyprostaglandin dehydrogenase inhuman placenta of patients with preeclampsia. J. Clin. Endocr. Metab. 86:1313–1317, 2001.
- [64949] 3279.Hurvitz, J. R.; Suwairi, W. M.; Van Hul, W.; El–Shanti, H.; Superti–Furga,A.; Roudier, J.; Holderbaum, D.; Pauli, R. M.; Herd, J. K.; Van Hul,E.; Rezai–Delui, H.; Legius, E.; Le Merrer, M.; Al–Alami, J.; Bahabri,S. A.; Warman, M. L.: Mutations in the CCN gene family member WISP3cause progressive pseudorheumatoid dysplasia. Nature Genet.

23:94–98, 1999.

- [64950] 3280.Spranger, J.; Albert, C.; Schilling, F.; Bartsocas, C.; Stoss,H.: Progressive pseudorheumatoid arthritis of childhood (PPAC): ahereditary disorder simulating rheumatoid arthritis. *Europ. J. Pediat.* 140:34–40, 1983.
- [64951] 3281.Bahabri, S. A.; Suwairi, W. M.; Laxer, R. M.; Polinkovsky, A.;Dalaan, A. A.; Warman, M. L.: The campto-dactyly–arthropathy–coxavara–pericarditis syndrome: clinical features and genetic mappingto human chromosome 1. *Arthritis Rheum.* 41: 730–735, 1998.
- [64952] 3282.Marcelino, J.; Carpten, J. D.; Suwairi, W. M.; Gutierrez, O. M.;Schwartz, S.; Robbins, C.; Sood, R.; Makalowska, I.; Baxevanis, A.;Johnstone, B.; Laxer, R. M.; Zemel, L.; and 13 others: CACP, encodinga secreted proteoglycan, is mutated in camptodactyly–arthropathy–coxavara–pericarditis syndrome. *Nature Genet.* 23: 319–322, 1999.
- [64953] 3283.Tannin, G. M.; Agarwal, A. K.; Monder, C.; New, M. I.; White,P. C.: The human gene for 11–beta–hydroxysteroid dehydrogenase: structure,tissue distribution, and chromosomal localization. *J. Biol. Chem.* 266:16653–16658, 1991.
- [64954] 3284.Andrechek, E. R.; Hardy, W. R.; Girgis–Gabardo, A. A.; Perry, R.L. S.; Butler, R.; Graham, F. L.; Kahn, R. C.;

Rudnicki, M. A.; Muller, W. J.: ErbB2 is required for muscle spindle and myoblast cell survival. *Molec. Cell. Biol.* 22: 4714–4722, 2002.

[64955] 3285. Bargmann, C. I.; Hung, M.-C.; Weinberg, R. A.: The NEU oncogene encodes an epidermal growth factor receptor-related protein. *Nature* 319:226–230, 1986.

[64956] 3286. Chan, R.; Hardy, W. R.; Laing, M. A.; Hardy, S. E.; Muller, W. J.: The catalytic activity of the ErbB-2 receptor tyrosine kinase is essential for embryonic development. *Molec. Cell. Biol.* 22: 1073–1078, 2002.

[64957] 3287. Coussens, L.; Yang-Feng, T. L.; Liao, Y.-C.; Chen, E.; Gray, A.; McGrath, J.; Seeburg, P. H.; Libermann, T. A.; Schlessinger, J.; Francke, U.; Levinson, A.; Ullrich, A.: Tyrosine kinase receptor with extensive homology to EGF receptor shares chromosomal location with NEU oncogene. *Science* 230:1132–1139, 1985.

[64958] 3288. Crone, S. A.; Zhao, Y.-Y.; Fan, L.; Gu, Y.; Minamisawa, S.; Liu, Y.; Peterson, K. L.; Chen, J.; Kahn, R.; Condorelli, G.; Ross, J., Jr.; Chien, K. R.; Lee, K.-F.: ErbB2 is essential in the prevention of dilated cardiomyopathy. *Nature Med.* 8: 459–465, 2002.

[64959] 3289. Dankort, D.; Maslikowski, B.; Warner, N.; Kanno, N.; Kim, H.; Wang, Z.; Moran, M. F.; Oshima, R. G.; Cardiff, R.

D.; Muller, W. J.: Grb2 and Shc adapter proteins play distinct roles in Neu (ErbB-2) induced mammary tumorigenesis: implications for human breast cancer. *Molec. Cell. Biol.* 21: 1540-1551, 2001.

[64960] 3290. De Boer, J. G.: A new mutator phenotype in breast cancer? (Commentary) *Proc. Nat. Acad. Sci.* 99: 3368-3369, 2002.

[64961] 3291. De Placido, S.; Carlomagno, C.; De Laurentiis, M.; Bianco, A.R.: C-erbB2 expression predicts tamoxifen efficacy in breast cancer patients. *Breast Cancer Res. Treat.* 52: 55-64, 1998.

[64962] 3292. Di Fiore, P. P.; Pierce, J. H.; Kraus, M. H.; Segatto, O.; King, C. R.; Aaronson, S. A.: erbB-2 is a potent oncogene when overexpressed in NIH/3T3 cells. *Science* 237: 178-182, 1987.

[64963] 3293. Doherty, J. K.; Bond, C.; Jardim, A.; Adelman, J. P.; Clinton, G. M.: The HER-2/neu receptor tyrosine kinase gene encodes a secreted autoinhibitor. *Proc. Nat. Acad. Sci.* 96: 10869-10874, 1999.

[64964] 3294. Ehsani, A.; Low, J.; Wallace, R. B.; Wu, A. M.: Characterization of a new allele of the human ERBB2 gene by allele-specific competition hybridization. *Genomics* 15: 426-429, 1993.

- [64965] 3295.Francke, U.: Personal Communication. New Haven, Connecticut 4/1988.
- [64966] 3296.Fukushige, S.-I.; Matsubara, K.-I.; Yoshida, M.; Sasaki, M.; Suzuki,T.; Semba, K.; Toyoshima, K.; Yamamoto, T.: Localization of a novelv-erbB-related gene, c-erbB-2, on human chromosome 17 and its amplification in a gastric cancer cell line. *Molec. Cell. Biol.* 6: 955-958, 1986.
- [64967] 3297.Kaneko, Y.; Homma, C.; Maseki, N.; Sakurai, M.; Toyoshima, K.;Yamamoto, T.: Human c-erbB-2 remains on chromosome 17 in band q21 in the 15;17 translocation associated with acute promyelocytic leukemia. *Jpn.J. Cancer Res.* 78: 16-19, 1987.
- [64968] 3298.Lin, W.; Sanchez, H. B.; Deerinck, T.; Morris, J. K.; Ellisman,M.; Lee, K. F.: Aberrant development of motor axons and neuromuscularsynapses in erbB2-deficient mice. *Proc. Nat. Acad. Sci.* 97: 1299-1304,2000.
- [64969] 3299.Liu, S.; Liu, W.; Jakubczak, J. L.; Erexson, G. L.; Tindall, K.R.; Chan, R.; Muller, W. J.; Adhya, S.; Garges, S.; Merlino, G.:Genetic instability favoring transversions associated with ErbB2-inducedmammary tumorigenesis. *Proc. Nat. Acad. Sci.* 99: 3770-3775, 2002.
- [64970] 3300.Mehta, R. R.; McDermott, J. H.; Hieken, T. J.; Marler,

K. C.;Patel, M. K.; Wild, L. D.; Das Gupta, T. K.: Plasma c-erbB2 levels in breast cancer patients: prognostic significance in predicting response to chemotherapy. J. Clin. Oncol. 16: 2409–2416, 1998.

[64971] 3301.Morris, J. K.; Lin, W.; Hauser, C.; Marchuk, Y.; Getman, D.; Lee,K.-F.: Rescue of the cardiac defect in ErbB2 mutant mice reveals essential roles of ErbB2 in peripheral nervous system development. Neuron 23:273–283, 1999.

[64972] 3302.Muleris, M.; Almeida, A.; Malfoy, B.; Dutrillaux, B.: Assignment of v-erb-b2 avian erythroblastic leukemia viral oncogene homolog 2 (ERBB2) to human chromosome band 17q21.1 by in situ hybridization. Cytogenet. Cell Genet. 76: 34–35, 1997.

[64973] 3303.Ozcelik, C.; Erdmann, B.; Pilz, B.; Wettschureck, N.; Britsch, S.; Hubner, N.; Chien, K. R.; Birchmeier, C.; Garratt, A. N.: Conditional mutation of the ErbB2 (HER2) receptor in cardiomyocytes leads to dilated cardiomyopathy. Proc. Nat. Acad. Sci. 99: 8880–8885, 2002.

[64974] 3304.Papewalis, J.; Nikitin, A. Y.; Rajewsky, M. F.: G to A polymorphism at amino acid codon 655 of the human erbB-2/HER2 gene. Nucleic Acids Res. 19: 5452 only, 1991.

[64975] 3305.Pegram, M. D.; Finn, R. S.; Arzoo, K.; Beryt, M.;

Pietras, R.J.; Slamon, D. J.: The effect of HER-2/neu over-expression on chemotherapeutic drug sensitivity in human breast and ovarian cancer cells. *Oncogene* 15:537–547, 1997.

[64976] 3306. Pietras, R. J.; Pegram, M. D.; Finn, R. S.; Maneval, D. A.; Slamon, D. J.: Remission of human breast cancer xenografts on therapy with humanized monoclonal antibody to HER-2 receptor and DNA-reactive drugs. *Oncogene* 17:2235–2249, 1998.

[64977] 3307. Popescu, N. C.; King, C. R.; Kraus, M. H.: Localization of the human erbB-2 gene on normal and rearranged chromosomes 17 to bands q12–21.32. *Genomics* 4: 362–366, 1989.

[64978] 3308. Qiu, Y.; Ravi, L.; Kung, H.-J.: Requirement of ErbB2 for signalling by interleukin-6 in prostate carcinoma cells. *Nature* 393: 83–85, 1998.

[64979] 3309. Semba, K.; Kamata, N.; Toyoshima, K.; Yamamoto, T.: A v-erbB-related protooncogene, c-erbB-2, is distinct from the c-erbB-1/epidermal growth factor-receptor gene and is amplified in a human salivary gland adenocarcinoma. *Proc. Nat. Acad. Sci.* 82: 6497–6501, 1985.

[64980] 3310. Slamon, D. J.; Godolphin, W.; Jones, L. A.; Holt, J. A.; Wong, S. G.; Keith, D. E.; Levin, W. J.; Stuart, S. G.; Udove,



J.; Ullrich,A.; Press, M. F.: Studies of the HER-2/neu proto-oncogene in humanbreast and ovarian cancer. Science 244: 707–712, 1989.

[64981] 3311.Slamon, D. J.; Leyland–Jones, B.; Shak, S.; Fuchs, H.; Paton,V.; Bajamonde, A.; Fleming, T.; Eiermann, W.; Wolter, J.; Pegram,M.; Baselga, J.; Norton, L.: Use of chemotherapy plus a monoclonalantibody against HER2 for metastatic breast cancer that overexpressesHER2. New Eng. J. Med. 344: 783–792, 2001.

[64982] 3312.Beutler, E.; Kuhl, W.; Gelbart, T.: 6–Phosphogluconolactonasedeficiency, a hereditary erythrocyte enzyme deficiency: possible interactionwith glucose–6–phosphate dehydrogenase deficiency. Proc. Nat. Acad.Sci. 82: 3876–3878, 1985.

[64983] 3313.Benkmann, H.–G.; Paik, Y. K.; Chen, L. Z.; Goedde, H. W.: Polymorphismof 6–PGD in South Korea: a new genetic variant 6–PGD Korea. Hum.Genet. 74: 204–205, 1986.

[64984] 3314.Blake, N. M.; Kirk, R. L.: New genetic variant of 6–phosphogluconatedehydrogenase in Australian aborigines. Nature 221: 278 only, 1969.

[64985] 3315.Bowman, J. E.; Carson, P. E.; Frischer, H.; De Garay, A. L.: Geneticsof starch–gel electrophoretic variants of hu–

man 6-phosphogluconicdehydrogenase: population and family studies in the United Statesand in Mexico. *Nature* 210: 811–812, 1966.

[64986] 3316.Brewer, G. J.; Dern, R. J.: A new inherited enzymatic deficiencyof human erythrocytes: 6-phosphogluconate dehydrogenase deficiency. *Am.J. Hum. Genet.* 16: 472–476, 1964.

[64987] 3317.Burgerhout, W.; Van Someren, H.; Bootsma, D.: Cytological mappingof the genes assigned to the human A1 chromosome by use of radiation–inducedchromosome breakage in a human–Chinese hamster hybrid cell line. *Humangenetik* 20:159–162, 1973.

[64988] 3318.Davidson, R. G.: Electrophoretic variants of human 6-phosphogluconatedehydrogenase: population and family studies and description of anew variant. *Ann. Hum. Genet.* 30: 355–362, 1967.

[64989] 3319.Dern, R. J.; Brewer, G. J.; Tashian, R. E.; Shows, T. B.: Hereditaryvariation of erythrocytic 6-phosphogluconate dehydrogenase. *J. Lab.Clin. Med.* 67: 255–264, 1966.

[64990] 3320.Douglas, G. R.; McAlpine, P. J.; Hamerton, J. L.: Regional localizationof loci for human PGM–1 and 6PGD on human chromosome 1 by use of hybridsof Chinese hamster–human somatic cells. *Proc. Nat. Acad. Sci.*

70:2737–2740, 1973.

- [64991] 3321.Fildes, R. A.; Parr, C. W.: Human red–cell phosphogluconate dehydrogenases. *Nature* 200:890–891, 1963.
- [64992] 3322.Nelson, M. S.: Biochemical and genetic characterization of theLowell variant: a new phenotype of 6–phosphogluconate dehydrogenase. *Hum.Genet.* 62: 333–336, 1982.
- [64993] 3323.Nevo, S.: A new rare PGD variant, PGD Mediteranean. *Hum. Genet.* 81:199 only, 1989.
- [64994] 3324.Parr, C. W.: Erythrocyte phosphogluconate dehydrogenase polymorphism. *Nature* 210:487–489, 1966.
- [64995] 3325.Parr, C. W.; Fitch, L. I.: Inherited quantitative variations of human phosphogluconate dehydrogenase. *Ann. Hum. Genet.* 30: 339–353, 1967.
- [64996] 3326.Ritter, H.; Toriverdiau, G.; Wendt, G. G.; Zilch, I.: Genetic and linkage analysis on 6–PGD. *Humangenetik* 14: 73–75, 1971.
- [64997] 3327.Tariverdian, G.; Ropers, H.–H.; Op't Hof, J.; Ritter, H.: ZurGenetik der 6–Phosphogluconatdehydrogenase (EC: 1.1.1.44): Eine neueVariante F (Freiburg). *Humangenetik* 10: 355–357, 1970.
- [64998] 3328.Weitkamp, L. R.: Genetic linkage relationships of the ADA and6–PGD loci in 'Humangenetik.' (Letter) *Human–*

genetik 15: 359–360,1972.

- [64999] 3329.Weitkamp, L. R.; Guttormsen, S. A.; Greendyke, R. M.: Genetic linkage between a locus for 6-PGD and the Rh locus: evaluation of possible heterogeneity in the recombination fraction between sexes and among families. Am. J. Hum. Genet. 23: 462–470, 1971.
- [65000] 3330.Weitkamp, L. R.; Guttormsen, S. A.; Shreffler, D. C.; Sing, C.F.; Napier, J. A.: Genetic linkage relations of the loci for 6-phosphogluconate dehydrogenase and adenosine deaminase in man. Am. J. Hum. Genet. 22:216–220, 1970.
- [65001] 3331.Westerveld, A.; Meera Khan, P.: Evidence for linkage between human loci for 6-phosphogluconate dehydrogenase and phosphoglucomutase(1) in man–Chinese hamster somatic cell hybrids. Nature 236: 30–32,1972.
- [65002] 3332.McEver, R. P.; Beckstead, J. H.; Moore, K. L.; Marshall–Carlson, L.; Bainton, D. F.: GMP–140, a platelet alpha-granule membrane protein, is also synthesized by vascular endothelial cells and is localized in Weibel–Palade bodies. J. Clin. Invest. 84: 92–99, 1989.
- [65003] 3333.Thomas, K. R.; Capecchi, M. R. :Cell 51: 503–512, 1987.
- [65004] 3334.Scita, G.; Nordstrom, J.; Carbone, R.; Tenca, P.; Giar-

dina, G.;Gutkind, S.; Bjarnegard, M.; Betsholtz, C.; Di Fiore, P. P.: EPS8and E3B1 transduce signals from Ras to Rac. Nature 401: 290–293,1999.

[65005] 3335.Remmers, E. F.; Goldmuntz, E. A.; Cash, J. M.; Crofford, L. J.;Misiewicz–Poltorak, B.; Zha, H.; Wilder, R. L.: Genetic map of ninepolymorphic loci comprising a single linkage group on rat chromosome10: evidence for linkage conservation with human chromosome 17 andmouse chromosome 11. Genomics 14: 618–623, 1992.

[65006] 3336.Sarraf, P.; Mueller, E.; Smith, W. M.; Wright, H. M.; Kum, J.B.; Aaltonen, L. A.; de la Chapelle, A.; Spiegelman, B. M.; Eng, C.: Loss–of–function mutations in PPAR–gamma associated with human coloncancer. Molec. Cell 3: 799–804, 1999.

[65007] 3337.Scheidler, S.; Fredericks, W. J.; Rauscher, F. J., III; Barr,F. G.; Vogt, P. K.: The hybrid PAX3–FKHR fusion protein of alveolarrhabdomyosarcoma transforms fibroblasts in culture. Proc. Nat. Acad.Sci. 93: 9805–9809, 1996.

[65008] 3338.Sublett, J. E.; Jeon, I–S.; Shapiro, D. N.: The alveolar rhabdomyosarcomaPAX3/FKHR fusion protein is a transcriptional activator. Oncogene 11:545–552, 1995.

[65009] 3339.Whang–Peng, J.; Knutsen, T.; Theil, K.; Horowitz, M. E.; Triche,T.: Cytogenetic studies in subgroups of rhab–

domyosarcoma. *GenesChromosomes Cancer* 5: 299–310, 1992.

[65010] 3340. Richard, I.; Broux, O.; Chiannikulchai, N.; Fougere-  
ousse, F.; Allamand, V.; Bourg, N.; Brenguier, L.; Devaud,  
C.; Pasturaud, P.; Roudaut, C.; Lorenzo, F.; Sebastiani-Ka-  
batchis, C.; Schultz, R. A.; Polymeropoulos, M. H.; Gyapay,  
G.; Auffray, C.; Beckmann, J. S.: Regional localization of  
human chromosome 15 loci. *Genomics* 23: 619–627,  
1994.

[65011] 3341. Zuniga, A.; Haramis, A.-P. G.; McMahon, A. P.;  
Zeller, R.: Signal relay by BMP antagonism controls the  
SHH/FGF4 feedback loop in vertebrate limb buds. *Nature*  
401: 598–602, 1999.

[65012] 3342. Bao, L.; Gerard, N. P.; Eddy, R. L., Jr.; Shows, T. B.;  
Gerard, C.: Mapping of genes for the human C5a receptor  
(C5AR), human FMLPreceptor (FPR), and two FMLP receptor  
homologue orphan receptors (FPRH1, FPRH2) to chromo-  
some 19. *Genomics* 13: 437–440, 1992.

[65013] 3343. Cohen-Haguenauer, O.; Van Cong, N.; Prud'homme,  
J. F.; Jegou-Foubert, C.; Gross, M. S.; De Tand, M. F.; Mil-  
grom, E.; Frezal, J.: A gene expressed in human breast  
cancer and regulated by estrogen in MCF-7 cells is located  
on chromosome 21. (Abstract) *Cytogenet. Cell Genet.*

40:606 only, 1985.

- [65014] 3344.Jakowlew, S. B.; Breathnach, R.; Jeltsch, J.-M.; Masiakowski, P.; Chambon, P.: Sequence of the pS2 mRNA induced by estrogen in the human breast cancer cell line MCF-7. *Nucleic Acids Res.* 12: 2861-2878, 1984.
- [65015] 3345.Lefebvre, O.; Chenard, M.-P.; Masson, R.; Linares, J.; Dierich, A.; LeMeur, M.; Wendling, C.; Tomasetto, C.; Chambon, P.; Rio, M.-C.: Gastric mucosa abnormalities and tumorigenesis in mice lacking the pS2 trefoil protein. *Science* 274: 259-262, 1996.
- [65016] 3346.Moisan, J.-P.; Mattei, M.-G.; Mandel, J.-L.: Chromosome localization and polymorphism of an oestrogen-inducible gene specifically expressed in some breast cancers. *Hum. Genet.* 79: 168-171, 1988.
- [65017] 3347.Moisan, J. P.; Mattei, M. G.; Baeteman-Volkel, M. A.; Mattei, J.F.; Brown, A. M. C.; Garnier, J. M.; Jeltsch, J. M.; Masiakowsky, P.; Roberts, M.; Mandel, J. L.: A gene expressed in human mammary tumor cells under estrogen control (BCEI) is located in 21q22.3 and defines an RFLP. (Abstract) *Cytogenet. Cell Genet.* 40: 701-702, 1985.
- [65018] 3348.Roberts, M.; Wallace, J.; Jeltsch, J.-M.; Berry, M.: The 5'-prime flanking region of the human pS2 gene mediates its transcriptional activation by estrogen in MCF-7 cells.

Biochem. Biophys. Res. Commun. 151:306–313, 1988.

- [65019] 3349.Seib, T.; Blin, N.; Hilgert, K.; Seifert, M.; Theisinger, B.;Engel, M.; Dooley, S.; Zang, K.–D.; Welter, C.: The three human trefoilgenes TFF1, TFF2, and TFF3 are located within a region of 55 kb on chromosome 21q22.3. Genomics 40: 200–202, 1997.
- [65020] 3350.Watkins, P. C.; Tanzi, R. E.; Roy, J.; Stuart, N.; Stanislovitis,P.; Gusella, J. F.: A cosmid genetic linkage map of chromosome 21and localization of the breast cancer estrogen–inducible (BCEI) gene.(Abstract) Am. J. Hum. Genet. 41: A189 only, 1987.
- [65021] 3351.Ponnambalam, S.; Jackson, A. P.; LeBeau, M. M.; Pravtcheva, D.;Ruddle, F. H.; Alibert, C.; Parham, P.: Chromosomal location andsome structural features of human clathrin light–chain genes (CLTAand CLTB). Genomics 24: 440–444, 1994.
- [65022] 3352.Andersen, T. I.; Wooster, R.; Laake, K.; Collins, N.; Warren, W.;Skrede, M.; Eeles, R.; Tveit, K. M.; Johnston, S. R. D.; Dowsett,M.; Olsen, A. O.; Moller, P.; Stratton, M. R.; Borresen–Dale, A.–L.: Screening for ESR mutations in breast and ovarian cancer patients. Hum.Mutat. 9: 531–536, 1997.
- [65023] 3353.Auboeuf, D.; Honig, A.; Berget, S. M.; O'Malley, B.



W.: Coordinateregulation of transcription and splicing by steroid receptor coregulators. *Science* 298:416–419, 2002.

- [65024] 3354.Balleine, R. L.; Hunt, S. M. N.; Clarke, C. L.: Coexpression ofalternatively spliced estrogen and progesterone receptor transcripts in human breast cancer. *J. Clin. Endocr. Metab.* 84: 1370–1377, 1999.
- [65025] 3355.Becherini, L.; Gennari, L.; Masi, L.; Mansani, R.; Marsart, F.;Morelli, A.; Falchetti, A.; Gonnelli, S.; Fiorelli, G.; Tanini, A.;Brandi, M. L.: Evidence of a linkage disequilibrium between polymorphisms in the human estrogen receptor–alpha gene and their relationship to bone mass variation in postmenopausal Italian women. *Hum. Molec.Genet.* 9: 2043–2050, 2000.
- [65026] 3356.Bord, S.; Horner, A.; Beavan, S.; Compston, J.: Estrogen receptorsalpha and beta are differentially expressed in developing human bone. *J.Clin. Endocr. Metab.* 86: 2309–2314, 2001.
- [65027] 3357.Castagnoli, A.; Maestri, I.; Bernardi, F.; Del Senno, L.: PvuII RFLP inside the human estrogen receptor gene. *Nucleic Acids Res.* 15:866 only, 1987.
- [65028] 3358.Chaidarun, S. S.; Alexander, J. M.: A tumor-specific truncatedestrogen receptor splice variant enhances estrogen

gen-stimulated geneexpression. *Molec. Endocr.* 12:  
1355–1366, 1998.

[65029] 3359.Chiang, C.–H.; Cheng, K. W.; Igarashi, S.; Nathwani, P. S.; Leung,P. C. K.: Hormonal regulation of estrogen receptor alpha and betagene expression in human granulosa–luteal cells in vitro. *J. Clin.Endocr. Metab.* 85:  
3828–3839, 2000.

[65030] 3360.Clark, G. M.; McGuire, W. L.: Steroid receptors and other prognosticfactors in primary breast cancer. *Semin. Oncol.* 15 (suppl. 1): 20–25,1988.

[65031] 3361.Couse, J. F.; Hewitt, S. C.; Bunch, D. O.; Sar, M.; Walker, V.R.; Davis, B. J.; Korach, K. S.: Postnatal sex reversal of the ovariesin mice lacking estrogen receptors alpha and beta. *Science* 286:2328–2331, 1999.

[65032] 3362.Davis, V. L.; Chan, C.–C.; Schoen, T. J.; Couse, J. F.; Chader,G. J.; Korach, K. S.: An estrogen receptor repressor induces cataractformation in transgenic mice. *Proc. Nat. Acad. Sci.* 99: 9427–9432,2002.

[65033] 3363.Esmaeli, B.; Harvey, J. T.; Hewlett, B.: Immunohistochemical evidence for estrogen receptors in meibomian glands. *Ophthalmology* 107:180–184, 2000.

[65034] 3364.Fan, S.; Wang, J.–A.; Yuan, R.; Ma, Y.; Meng, Q.; Erdos, M. R.;Pestell, R. G.; Yuan, F.; Auborn, K. J.; Goldberg,

I. D.; Rosen, E.M.: BRCA1 inhibition of estrogen receptor signaling in transfected cells. *Science* 284: 1354–1356, 1999.

[65035] 3365. Fuqua, S. A. W.; Chamness, G. C.; McGuire, W. L.: Estrogen receptor mutations in breast cancer. *J. Cell. Biochem.* 51: 135–139, 1993.

[65036] 3366. Gosden, J. R.; Middleton, P. G.; Rout, D.: Localization of the human oestrogen receptor gene to chromosome 6q24–q27 by in situ hybridization. *Cytogenet. Cell Genet.* 43: 218–220, 1986.

[65037] 3367. Green, S.; Walter, P.; Kumar, V.; Krust, A.; Bornert, J.-M.; Argos, P.; Chambon, P.: Human oestrogen receptor cDNA: sequence, expression and homology to v-erb-A. *Nature* 320: 134–139, 1986.

[65038] 3368. Greene, G. L.; Gilna, P.; Waterfield, M.; Baker, A.; Hort, Y.; Shine, J.: Sequence and expression of human estrogen receptor complementary DNA. *Science* 231: 1150–1154, 1986.

[65039] 3369. Heine, P. A.; Taylor, J. A.; Iwamoto, G. A.; Lubahn, D. B.; Cooke, P. S.: Increased adipose tissue in male and female estrogen receptor- $\alpha$  knockout mice. *Proc. Nat. Acad. Sci.* 97: 12729–12734, 2000.

[65040] 3370. Herrington, D. M.; Howard, T. D.; Hawkins, G. A.;

Reboussin, D.M.; Xu, J.; Zheng, S. L.; Brosnihan, K. B.; Meyers, D. A.; Bleecker, E. R.: Estrogen-receptor polymorphisms and effects of estrogen replacement on high-density lipoprotein cholesterol in women with coronary disease. *New Eng. J. Med.* 346: 967–974, 2002.

[65041] 3371. Issa, J.-P. J.; Ottaviano, Y. L.; Celano, P.; Hamilton, S. R.; Davidson, N. E.; Baylin, S. B.: Methylation of the oestrogen receptor CpG island links ageing and neoplasia in human colon. *Nature Genet.* 7:536–540, 1994.

[65042] 3372. Jeltsch, J. M.; Roberts, M.; Schatz, C.; Garnier, J. M.; Brown, A. M. C.; Chambon, P.: Structure of the human oestrogen-responsive gene pS2. *Nucleic Acids Res.* 15: 1401–1414, 1987.

[65043] 3373. Bing, D. H.; Almeda, S.; Isliker, H.; Lahav, J.; Hynes, R. O.: Fibronectin binds to the C1q component of complement. *Proc. Nat. Acad. Sci.* 79: 4198–4201, 1982.

[65044] 3374. Bittner, M.; Meltzer, P.; Chen, Y.; Jiang, Y.; Seftor, E.; Hendrix, M.; Radmacher, M.; Simon, R.; Yakhini, Z.; Bendor, A.; Sampas, N.; Dougherty, E.; and 16 others: Molecular classification of cutaneous malignant melanoma by gene expression profiling. *Nature* 406: 536–540, 2000.

[65045] 3375. Clark, E. A.; Golub, T. R.; Lander, E. S.; Hynes, R. O.: Genomic analysis of metastasis reveals an essential role for

RhoC. Nature 406:532–535, 2000.

- [65046] 3376.Clemmensen, I.: Fibronectin and its role in connective tissue diseases. (Editorial) Europ. J. Clin. Invest. 11: 145–146, 1981.
- [65047] 3377.Croce, C. M.: Personal Communication. Philadelphia, Pa. 1/12/1983.
- [65048] 3378.Eun, C. K.; Klinger, H. P.: Human chromosome 11 affects the expression of fibronectin fibers in human–mouse cell hybrids. Cytogenet.Cell Genet. 27: 57–65, 1980.
- [65049] 3379.Gutman, A.; Kornblihtt, A. R.: Identification of a third region of cell–specific alternative splicing in human fibronectin mRNA. Proc.Nat. Acad. Sci. 84: 7179–7182, 1987.
- [65050] 3380.Henry, I.; Jeanpierre, M.; Weil, D.; Grzeschik, K. H.; Ramirez,F.; Junien, C.: The structural gene for fibronectin (FN) maps to 2q323–qter. (Abstract) Cytogenet. Cell Genet. 40: 650 only, 1985.
- [65051] 3381.Hirano, H.; Yamada, Y.; Sullivan, M.; de Crombrughe, B.; Pastan,I.; Yamada, K. M.: Isolation of genomic DNA clones spanning the entire fibronectin gene. Proc. Nat. Acad. Sci. 80: 46–50, 1983.
- [65052] 3382.Jhanwar, S. C.; Jensen, J. T.; Kaelbling, M.; Chaganti,

R. S.K.; Klinger, H. P.: In situ localization of human fibronectin (FN) genes to chromosome regions 2p14–p16, 2q34–q36, and 11q12.1–q13.5 in germ line cells, but to chromosome 2 sites only in somatic cells. *Cytogenet. Cell Genet.* 41: 47–53, 1986.

[65053] 3383. Koch, G. A.; Schoen, R. C.; Klebe, R. J.; Shows, T. B.: Assignment of a fibronectin gene to human chromosome 2 using monoclonal antibodies. *Exp. Cell Res.* 141: 293–302, 1982.

[65054] 3384. Kornblihtt, A. R.; Umezawa, K.; Vibe-Pedersen, K.; Baralle, F.E.: Primary structure of human fibronectin: differential splicing may generate at least 10 polypeptides from a single gene. *EMBO J.* 4:1755–1759, 1985.

[65055] 3385. Kornblihtt, A. R.; Vibe-Pedersen, K.; Baralle, F. E.: Isolation and characterization of cDNA clones for human and bovine fibronectins. *Proc. Nat. Acad. Sci.* 80: 3218–3222, 1983.

[65056] 3386. Kornblihtt, A. R.; Vibe-Pedersen, K.; Baralle, F. E.: Human fibronectin: molecular cloning evidence for two mRNA species differing by an internal segment coding for a structural domain. *EMBO J.* 3: 221–226, 1984.

[65057] 3387. Kurkinen, M.; Vartio, T.; Vaheri, A.: Polypeptides of human plasma fibronectin are similar but not identical.

Biochim. Biophys. Acta 624:490–498, 1980.

- [65058] 3388.Matsuura, H.; Takio, K.; Titani, K.; Greene, T.; Levery, S. B.;Salyan, M. E. K.; Hakomori, S.: The oncofetal structure of humanfibronectin defined by monoclonal antibody FDC-6. J. Biol. Chem. 263:3314–3322, 1988.
- [65059] 3389.McDonagh, J.: Fibronectin: a molecular glue. Arch. Path. Lab.Med. 105: 393–396, 1981.
- [65060] 3390.Mosesson, M. W.; Amrani, D. L.: The structure and biologic activitiesof plasma fibronectin. Blood 56: 145–158, 1980.
- [65061] 3391.Odermatt, E.; Tamkun, J. W.; Hynes, R. O.: Repeating modularstructure of the fibronectin gene: relationship to protein structureand subunit variation. Proc. Nat. Acad. Sci. 82: 6571–6575, 1985.
- [65062] 3392.Owerbach, D.; Doyle, D.; Shows, T. B.: Genetics of the large,external, transformation–sensitive (LETs) protein: assignment of a gene coding for expression of LETs to human chromosome 8. Proc. Nat.Acad. Sci. 75: 5640–5644, 1978.
- [65063] 3393.Prowse, K.; Tricoli, J.; Klebe, R.; Shows, T.: Chromosome 2 assignmentof the structural gene for fibronectin (FN) using a cloned probe.(Abstract) Cytogenet. Cell Genet. 40: 724 only, 1985.

- [65064] 3394.Prowse, K. R.; Tricoli, J. V.; Klebe, R. J.; Shows, T. B.: Assignmentof the human fibronectin structural gene to chromosome 2. *Cytogenet.Cell Genet.* 41: 42–46, 1986.
- [65065] 3395.Rennard, S. I.; Church, R. L.; Rohrbach, D. H.; Shupp, D. E.;Abe, S.; Hewitt, A. T.; Murray, J. C.; Martin, G. R.: Localizationof the human fibronectin (FN) gene on chromosome 8 by a specific enzymeimmunoassay. *Biochem. Genet.* 19: 551–566, 1981.
- [65066] 3396.Ridley, A.: Molecular switches in metastasis. *Nature* 406: 466–467,2000.
- [65067] 3397.Li, Q. J.; Ashraf, M. F.; Shen, D.; Green, W. R.; Stark, W. J.;Chan, C.–C; O'Brien, T. P.: The role of apoptosis in the pathogenesisof Fuchs endothelial dystrophy of the cornea. *Arch. Ophthal.* 119:1597–1604, 2001.
- [65068] 3398.McCurley, R. S.; Recinos, A., III; Olsen, A. S.; Gingrich, J. C.;Szczepaniak, D.; Cameron, H. S.; Krauss, R.; Weston, B. W.: Physicalmaps of human alpha(1,3)fucosyltransferase genes FUT3–FUT6 on chromosomes19p13.3 and 11q21. *Genomics* 26: 142–146, 1995.
- [65069] 3399.Weston, B. W.; Nair, R. P.; Larsen, R. D.; Lowe, J. B.: Isolationof a novel human alpha(1,3)fucosyltransferase gene and molecular comparisonto the human Lewis blood group alpha(1,3/1,4)fucosyltransferase gene:syntenic, ho–



mologous, nonallelic genes encoding enzymes with distinct acceptor substrate specificities. *J. Biol. Chem.* 267: 4152–4160, 1992.

[65070] 3400. Weston, B. W.; Smith, P. L.; Kelly, R. J.; Lowe, J. B.: Molecular cloning of a fourth member of a human  $\alpha$ -(1,3)-fucosyltransferase gene family: multiple homologous sequences that determine expression of the Lewis x, sialyl Lewis x, and difucosyl sialyl Lewis x epitopes. *J. Biol. Chem.* 267: 24575–24584, 1992.

[65071] 3401. Brinkman–Van der Linden, E. C. M.; Mollicone, R.; Oriol, R.; Larson, G.; Van den Eijnden, D. H.; Van Dijk, W.: A missense mutation in the FUT6 gene results in total absence of  $\alpha$ -3-fucosylation of human  $\alpha$ -1-acid glycoprotein. *J. Biol. Chem.* 271: 14492–14495, 1996.

[65072] 3402. Billardon, C.; Van Cong, N.; Picard, J. Y.; Dekaouel, C.; Rebourcet, R.; Weil, D.; Feingold, J.; Frezal, J.: Linkage studies of enzyme markers in man–mouse somatic cell hybrids. *Ann. Hum. Genet.* 36: 273–284, 1973.

[65073] 3403. Cook, P. J. L.; Hamerton, J. L.: Report of the committee on the genetic constitution of chromosome 1. *Cytogenet. Cell Genet.* 25: 9–20, 1979.

[65074] 3404. Zhou, X.–P.; Smith, W. M.; Gimm, O.; Mueller, E.; Gao, X.; Sarraf, P.; Prior, T. W.; Plass, C.; van Deimling, A.;

Black, P. M.; Yates, A. J.; Eng, C.: Over-representation of PPAR-gamma sequence variants in sporadic cases of glioblastoma multiforme: preliminary evidence for common low penetrance modifiers for brain tumour risk in the general population. *J. Med. Genet.* 37: 410-414, 2000.

[65075] 3405. Bingham, C.; Bulman, M. P.; Ellard, S.; Allen, L. I. S.; Lipkin, G. W.; van't Hoff, W. G.; Woolf, A. S.; Rizzoni, G.; Novelli, G.; Nicholls, A. J.; Hattersley, A. T.: Mutations in the hepatocyte nuclear factor-1-beta gene are associated with familial hypoplastic glomerulocystic kidney disease. *Am. J. Hum. Genet.* 68: 219-224, 2001.

[65076] 3406. Kaplan, B. S.; Gordon, I.; Pincott, J.; Barratt, T. M.: Familial hypoplastic glomerulocystic kidney disease: a definite entity with dominant inheritance. *Am. J. Med. Genet.* 34: 569-573, 1989.

[65077] 3407. Rizzoni, G.; Loirat, C.; Levy, M.; Milanesi, C.; Zachello, G.; Mathieu, H.: Familial hypoplastic glomerulocystic kidney: a new entity? *Clin. Nephrol.* 18: 263-268, 1982.

[65078] 3408. Karasawa, M.; Zwacka, R. M.; Reuter, A.; Fink, T.; Hsieh, C. L.; Lichter, P.; Francke, U.; Weiher, H.: The human homolog of the glomerulosclerosis gene Mpv17: structure and genomic organization. *Hum. Molec. Genet.*

2:1829–1834, 1993.

- [65079] 3409.Fong, H. K. W.; Yoshimoto, K. K.; Eversole–Cire, P.; Simon, M.I.: Identification of a GTP–binding protein alpha subunit that lacks an apparent ADP–ribosylation site for pertussis toxin. *Proc. Nat.Acad. Sci.* 85: 3066–3070, 1988.
- [65080] 3410.Matsuoka, M.; Itoh, H.; Kozasa, T.; Kaziro, Y.: Sequence analysis of cDNA and genomic DNA for a putative pertussis toxin–insensitive guanine nucleotide–binding regulatory protein alpha subunit. *Proc.Nat. Acad. Sci.* 85: 5384–5388, 1988.
- [65081] 3411.Wilkie, T. M.; Gilbert, D. J.; Olsen, A. S.; Chen, X.–N.; Amatruda, T. T.; Korenberg, J. R.; Trask, B. J.; de Jong, P.; Reed, R. R.; Simon, M. I.; Jenkins, N. A.; Copeland, N. G.: Evolution of the mammalian G protein alpha subunit multi–gene family. *Nature Genet.* 1: 85–91, 1992.
- [65082] 3412.Magovcevic, I.; Ang, S.–L.; Seidman, J. G.; Tolman, C. J.; Neer, E. J.; Morton, C. C.: Regional localization of the human G protein alpha(i2) (GNAI2) gene: assignment to 3p21 and a related sequence (GNAI2L) to 12p12–p13. *Genomics* 12: 125–129, 1992.
- [65083] 3413.Neer, E. J.; Michel, T.; Eddy, R.; Shows, T.; Seidman, J. G.: Genes for two homologous G–protein alpha subunits

map to different human chromosomes. Hum. Genet. 77: 259–262, 1987.

- [65084] 3414. Colombo, M. P.; Martinotti, A.; Howard, T. A.; Schneider, C.; D'Eustachio, P.; Seldin, M. F.: Localization of growth arrest-specific genes on mouse chromosomes 1, 7, 8, 11, 13, and 16. Mammalian Genome 2: 130–134, 1992.
- [65085] 3415. Del Sal, G.; Collavin, L.; Ruaro, M. E.; Edomi, P.; Saccone, S.; Della Valle, G.; Schneider, C.: Structure, function, and chromosomal mapping of the growth-suppressing human homologue of the murine *gas1* gene. Proc. Nat. Acad. Sci. 91: 1848–1852, 1994.
- [65086] 3416. Del Sal, G.; Ruaro, M. E.; Philipson, L.; Schneider, C.: The growth arrest-specific gene, *gas1*, is involved in growth suppression. Cell 70: 595–607, 1992.
- [65087] 3417. Evdokiou, A.; Webb, G. C.; Peters, G. B.; Dobrovic, A.; O'Keefe, D. S.; Forbes, I. J.; Cowled, P. A.: Localization of the human growth arrest-specific gene (*GAS1*) to chromosome bands 9q21.3–q22, a region frequently deleted in myeloid malignancies. Genomics 18: 731–733, 1993.
- [65088] 3418. Schneider, C.; King, R. M.; Philipson, L.: Genes specifically expressed at growth arrest of mammalian cells. Cell 54: 787–793, 1988.
- [65089] 3419. Webb, G. C.; Cowled, P. A.; Evdokiou, A.; Ford, J. H.;

Forbes, I. J.: Assignment, by in situ hybridization, of the growth arrest-specific gene, Gas-1, to mouse chromosome 13, bands B3-C2. *Genomics* 14: 548-549, 1992.

[65090] 3420. Lebo, R. V.; Cheung, M.-C.; Bruce, B. D.: Rapid gene mapping by dual laser chromosome sorting and spot blot DNA analysis. (Abstract) *Am. J. Hum. Genet.* 36: 101S only, 1984.

[65091] 3421. Kudo, J.; Chao, L.-Y.; Narni, F.; Saunders, G. F.: Structure of the human gene encoding the invariant gamma-chain of class II histocompatibility antigens. *Nucleic Acids Res.* 13: 8827-8841, 1985.

[65092] 3422. O'Sullivan, D. M.; Larhammar, D.; Wilson, M. C.; Peterson, P. A.; Quaranta, V.: Structure of the human Ia-associated invariant (gamma)-chain gene: identification of 5-prime sequences shared with major histocompatibility-complex class II genes. *Proc. Nat. Acad. Sci.* 83: 4484-4488, 1986.

[65093] 3423. Riese, R. J.; Shi, G.-P.; Villadangos, J.; Stetson, D.; Driessen, C.; Lennon-Dumenil, A.-M.; Chu, C.-L.; Naumov, Y.; Behar, S. M.; Ploegh, H.; Locksley, R.; Chapman, H. A.: Regulation of CD1 function and NK1.1+ T cell selection and maturation by cathepsin S. *Immunity* 15: 909-919, 2001.

- [65094] 3424.Topilski, I.; Harmelin, A.; Flavell, R. A.; Levo, Y.; Shachar, I.: Preferential Th1 immune response in invariant chain-deficient mice. *J. Immun.* 168: 1610–1617, 2002.
- [65095] 3425.Abderrahim, H.; Sambucy, J.-L.; Iris, F.; Ougen, P.; Billault, A.; Chumakov, I. M.; Dausset, J.; Cohen, D.; Le Paslier, D.: Cloning the human major histocompatibility complex in YACs. *Genomics* 23:520–527, 1994.
- [65096] 3426.Adman, R.; Pious, D. A.: Isoantigenic variants: isolation from human diploid cells in culture. *Science* 168: 370–372, 1970.
- [65097] 3427.Arnaiz-Villena, A.; Festenstein, H.: HLA genotyping by using spermatozoa: evidence for haploid gene expression. *Lancet* II: 707–709, 1976.
- [65098] 3428.Auffray, C.; Strominger, J. L.: Molecular genetics of the human major histocompatibility complex. *Adv. Hum. Genet.* 15: 197–247, 1986.
- [65099] 3429.Awdeh, Z. L.; Raum, D.; Yunis, E. J.; Alper, C. A.: Extended HLA/complement allele haplotypes: evidence for T/t-like complex in man. *Proc. Nat. Acad. Sci.* 80: 259–263, 1983.
- [65100] 3430.Bach, F. H.; Amos, D. B.: Hu-1 major histocompatibility locus in man. *Science* 156: 1506–1508, 1967.
- [65101] 3431.Bach, M. L.; Bach, F. H.: The genetics of histocom-

patibility. Hosp.Practice 5(8): 33–44, 1970.

[65102] 3432.Balner, H.: The major histocompatibility complex of primates:evolutionary aspects and comparative histogenetics. Phil. Trans.Roy. Soc. London B 292: 109–119, 1981.

[65103] 3433.Balner, H.; D'Amaro, J.; Visser, T. P.: Tissue typing of chimpanzees:I. Evidence for two allelic series of leukocyte antigens. Transplant.Proc. 6: 141–149, 1974.

[65104] 3434.Berger, R.; Bernheim, A.; Sasportes, M.; Hauptmann, G.; Hors,J.; Legrand, L.; Fellous, M.: Regional mapping of the HLA on the short arm of chromosome 6. Clin. Genet. 15: 245–251, 1979.

[65105] 3435.Bernard, J.: La decouverte du systeme principal d'histocompatibilite de l'homme. (Editorial) Presse Med. 75: 2369 only, 1967.

[65106] 3436.Bodmer, W. F.: Personal Communication. London, England 9/26/1986.

[65107] 3437.Bodmer, W. F.; Bodmer, J. G.; Adler, S.; Payne, R.; Bialek, J.: Genetics of '4' and 'LA' human leukocyte groups. Ann. N.Y. Acad.Sci. 129: 473–489, 1966.

[65108] 3438.Bodmer, W. F.; Bodmer, J. G.; Tripp, M.: Recombination between the LA and 4 loci of the HL–A system.In: Histocompatibility Testing 1970. Copenhagen: Munksgaard

(pub.) 1970. Pp. 187–191.

- [65109] 3439. Bodmer, W. F. (ed.): Evolutionary significance of the HL-A system. *Nature* 237:139–145, 1972.
- [65110] 3440. Bodmer, W. F. (ed.): The HLA System (17 papers). *Brit. Med. Bull.* 34(3): 213–316, 1978.
- [65111] 3441. Borgaonkar, D. S.; Bias, W. B.: HL-A loci and chromosome 6. *Birth Defects Orig. Art. Ser.* X(3): 67–68, 1974.
- [65112] 3442. Borgaonkar, D. S.; Bias, W. B.; Chase, G. A.; Sadasi-  
van, G.; Herr, H. M.; Golomb, H. M.; Bahr, G. F.; Kunkel, L.  
M.: Identification of C6–G21 translocation chromosome by  
the Q–M and Giemsa banding techniques in a patient with  
Down's syndrome, with possible assignment of Gm locus.  
*Clin. Genet.* 4: 53–57, 1973.
- [65113] 3443. Breuning, M. H.; van den Berg–Loonen, E. M.;  
Bernini, L. F.; Bijlsma, J. B.; van Loghem, E.; Meera Khan, P.;  
Nijenhuis, L. E.: Localization of HLA on the short arm of  
chromosome 6. *Hum. Genet.* 37: 131–139, 1977.
- [65114] 3444. Bull, R. W.; Benson, J. W.; Pearson, G.; Mann, J. D.:  
HLA substantiation of a trisomic human chromosome 6.  
*Transplant. Proc.* 10: 747–748, 1978.
- [65115] 3445. Campbell, R. D.; Trowsdale, J.: Map of the human  
MHC. *Immun. Today* 14: 349–352, 1993.
- [65116] 3446. Cardoso, C. S.; Alves, H.; Mascarenhas, M.;



Goncalves, R.; Oliveira,P.; Rodrigues, P.; Cruz, E.; de Sousa, M.; Porto, G.: Co-selection of the H63D mutation and the HLA-A29 allele: a new paradigm of linkage disequilibrium? Immunogenetics 53: 1002–1008, 2002.

[65117] 3447.Carrington, M.; Nelson, G. W.; Martin, M. P.; Kissner, T.; Vlahov,D.; Goedert, J. J.; Kaslow, R.; Buchbinder, S.; Hoots, K.; O'Brien,S. J.: HLA and HIV-1: heterozygote advantage and B\*35-Cw\*04 disadvantage. Science 283:1748–1752, 1999.

[65118] 3448.Ceppellini, R.; Van Rood, J. J.: The HL-A system. I. Genetics and molecular biology. Seminars Hemat. 11: 233–252, 1974.

[65119] 3449.Cohen, D.; Paul, P.; Font, M.-P.; Cohen, O.; Sayagh, B.; Marcadet,A.; Busson, M.; Mahouy, G.; Cann, H. M.; Dausset, J.: Analysis of HLA class I genes with restriction endonuclease fragments: implications for polymorphism of the human major histocompatibility complex. Proc.Nat. Acad. Sci. 80: 6289–6292, 1983.

[65120] 3450.Colombani, J.: Changing the name of the major histocompatibility complex. Res. Immun. 143: 411–417, 1992.

[65121] 3451.Dausset, J.: The major histocompatibility complex in man: past,present, and future concepts. Science 213:

1469–1474, 1981.

- [65122] 3452.Dausset, J.: Personal Communication. Paris, France 1/12/1983.
- [65123] 3453.Dausset, J.: Similarities between the HL-A system and other immunogeneticsystems. (Editorial) Vox Sang. 23: 153–164, 1972.
- [65124] 3454.Ion, R.; Telvi, L.; Chaussain, J.–L.; Barbet, J. P.; Nunes, M.;Safar, A.; Rethore, M.–O.; Fellous, M.; McElreavey, K.: Failure of testicular development associated with a rearrangement of 9p24.1 proximal to the SNF2 gene. Hum. Genet. 102: 151–156, 1998.
- [65125] 3455.Matsushime, H.; Jinno, A.; Takagi, N.; Shibuya, M.: A novel mammalian protein kinase gene (mak) is highly expressed in testicular germ cells at and after meiosis. Molec. Cell. Biol. 10: 2261–2268, 1990.
- [65126] 3456.Taketo, M.; Jinno, A.; Yamaguchi, S.; Matsushime, H.; Shibuya, M.;Seldin, M. F.: Mouse Mak gene for male germ cell-associated kinase maps to chromosome 13. Genomics 19: 397–398, 1994.
- [65127] 3457.Cohen, P. T. W.; Omenn, G. S.: Genetic variation of the cytoplasmic and mitochondrial malic enzymes in the monkey: *Macaca nemestrina*. Biochem.Genet. 7: 289–301, 1972.

- [65128] 3458. Gonzalez-Manchon, C.; Butta, N.; Ferrer, M.; Ayuso, M. S.; Parrilla, R.: Molecular cloning and functional characterization of the humancytosolic malic enzyme promoter: thyroid hormone responsiveness. *DNACell Biol.* 16: 533-544, 1997.
- [65129] 3459. Loeber, G.; Dworkin, M. B.; Infante, A.; Ahorn, H.: Characterization of cytosolic malic enzyme in human tumor cells. *FEBS Lett.* 344:181-186, 1994.
- [65130] 3460. Meera Khan, P.; Hagemeijer, A.; Wijnen, L. M. M.; van der Goes, R. G. M.: PGM3 and ME1 are probably in the 6pter-q12 region. (Abstract) *Cytogenet. Cell Genet.* 37: 537 only, 1984.
- [65131] 3461. Nass, S. J.; Olowson, M.; Miyashita, N.; Moriwaki, K.; Balling, R.; Imai, K.: Mapping of the Mod-1 locus on mouse chromosome 9. *Mammalian Genome* 4: 333-337, 1993.
- [65132] 3462. Povey, S.; Wilson, D. E., Jr.; Harris, H.; Gormley, I. P.; Perry, P.; Buckton, K. E.: Sub-unit structure of soluble and mitochondrial malic enzyme: demonstration of human mitochondrial enzyme in human-mouse hybrids. *Ann. Hum. Genet.* 39: 203-212, 1975.
- [65133] 3463. Itoh, T.; Tanaka, T.; Nagai, R.; Kamiya, T.; Sawayama, T.; Nakayama, T.; Tomoike, H.; Sakurada, H.; Yazaki, Y.; Nakamura, Y.: Genomic organization and muta-

tional analysis of HERG, a gene responsible for familial long QT syndrome. Hum. Genet. 102: 435–439, 1998.

- [65134] 3464. Jiang, C.; Atkinson, D.; Towbin, J. A.; Splawski, I.; Lehmann, M. H.; Li, H.; Timothy, K.; Taggart, R. T.; Schwartz, P. J.; Vincent, G. M.; Moss, A. J.; Keating, M. T.: Two long QT syndrome loci map to chromosomes 3 and 7 with evidence for further heterogeneity. Nature Genet. 8: 141–147, 1994.
- [65135] 3465. Jongbloed, R. J. E.; Wilde, A. A. M.; Geelen, J. L. M. C.; Doevendans, P.; Schaap, C.; Van Langen, I.; van Tintelen, J. P.; Cobben, J. M.; Beaufort-Krol, G. C. M.; Geraedts, J. P. M.; Smeets, H. J. M.: Novel KCNQ1 and HERG missense mutations in Dutch long-QT families. Hum. Mutat. 13: 301–310, 1999.
- [65136] 3466. Kagan, A.; Yu, Z.; Fishman, G. I.; McDonald, T. V.: The dominant negative LQT2 mutation A561V reduces wild-type HERG expression. J. Biol. Chem. 275: 11241–11248, 2000.
- [65137] 3467. Kuperschmidt, S.; Yang, T.; Chanthaphaychith, S.; Wang, Z.; Towbin, J. A.; Roden, D. M.: Defective human ether- $\alpha$ -go-go-related gene trafficking linked to an endoplasmic reticulum retention signal in the C terminus. J. Biol. Chem. 277: 27442–27448, 2002.

- [65138] 3468. Larsen, L. A.; Svendsen, I. H.; Jensen, A. M.; Kanter, J. K.; Andersen, P. S.; Moller, M.; Sorensen, S. A.; Sandoe, E.; Jacobsen, J. R.; Vuust, J.; Christiansen, M.: Long QT syndrome with a high mortality rate caused by a novel G572R missense mutation in KCNH2. *Clin. Genet.* 57: 125–130, 2000.
- [65139] 3469. Li, X.; Xu, J.; Li, M.: The human delta-1261 mutation of the HERG potassium channel results in a truncated protein that contains a subunit interaction domain and decreases the channel expression. *J. Biol. Chem.* 272: 705–708, 1997.
- [65140] 3470. Miller, C.: The inconstancy of the human heart. *Nature* 379: 767–768, 1996.
- [65141] 3471. Moss, A. J.; Zareba, W.; Kaufman, E. S.; Gartman, E.; Peterson, D. R.; Benhorin, J.; Towbin, J. A.; Keating, M. T.; Priori, S. G.; Schwartz, P. J.; Vincent, G. M.; Robinson, J. L.; Andrews, M. L.; Feng, C.; Hall, W. J.; Medina, A.; Zhang, L.; Wang, Z.: Increased risk of arrhythmic events in long-QT syndrome with mutations in the pore region of the human ether-a-go-go-related gene potassium channel. *Circulation* 105: 794–799, 2002.
- [65142] 3472. Nakajima, T.; Kurabayashi, M.; Ohyama, Y.; Kaneko, Y.; Furukawa, T.; Itoh, T.; Taniguchi, Y.; Tanaka, T.; Naka-

mura, Y.; Hiraoka, M.; Nagai, R.: Characterization of S818L mutation in HERG C-terminus in LQT2: modification of activation-deactivation gating properties. FEBS Lett. 481: 197–203, 2000.

[65143] 3473. Priori, S. G.; Barhanin, J.; Hauer, R. N. W.; Haverkamp, W.; Jongsma, H. J.; Kleber, A. G.; McKenna, W. J.; Roden, D. M.; Rudy, Y.; Schwartz, K.; Schwartz, P. J.; Towbin, J. A.; Wilde, A. M.: Genetic and molecular basis of cardiac arrhythmias: impact on clinical management. Parts I and II. Circulation 99: 518–528, 1999.

[65144] 3474. Priori, S. G.; Barhanin, J.; Hauer, R. N. W.; Haverkamp, W.; Jongsma, H. J.; Kleber, A. G.; McKenna, W. J.; Roden, D. M.; Rudy, Y.; Schwartz, K.; Schwartz, P. J.; Towbin, J. A.; Wilde, A. M.: Genetic and molecular basis of cardiac arrhythmias: impact on clinical management. Part III. Circulation 99: 674–681, 1999.

[65145] 3475. Priori, S. G.; Napolitano, C.; Schwartz, P. J.: Low penetrance in the long QT syndrome: clinical impact. Circulation 99: 529–533, 1999.

[65146] 3476. Rajamani, S.; Anderson, C. L.; Anson, B. D.; January, C. T.: Pharmacological rescue of human K<sup>+</sup> channel long-QT2 mutations. Circulation 105: 2830–2835, 2002.

[65147] 3477. Sanguinetti, M. C.; Jiang, C.; Curran, M. E.; Keating,

M. T.:A mechanistic link between an inherited and an acquired cardiac arrhythmia:HERG encodes the I(Kr) potassium channel. *Cell* 81: 299–307, 1995.

[65148] 3478.Satler, C. A.; Vesely, M. R.; Duggal, P.; Ginsburg, G. S.; Beggs,A. H.: Multiple different missense mutations in the pore region ofHERG in patients with long QT syndrome. *Hum. Genet.* 102: 265–272,1998.

[65149] 3479.Satler, C. A.; Walsh, E. P.; Vesely, M. R.; Plummer, M. H.; Ginsburg,G. S.; Jacob, H. J.: Novel missense mutation in the cyclic nucleotide–bindingdomain of HERG causes long QT syndrome. *Am. J. Med. Genet.* 65: 27–35,1996.

[65150] 3480.Smith, P. L.; Baukrowtiz, T.; Yellen, G.: The inward rectificationmechanism of the HERG cardiac potassium channel. *Nature* 379: 833–835,1996.

[65151] 3481.Splawski, I.; Shen, J.; Timothy, K. W.; Vincent, G. M.; Lehmann,M. H.; Keating, M. T.: Genomic structure of three long QT syndromegenes: KVLQT1, HERG, and KCNE1. *Genomics* 51: 86–97, 1998.

[65152] 3482.Tanaka, T.; Nagai, R.; Tomoike, H.; Takata, S.; Yano, K.; Yabuta,K.; Haneda, N. Nakano, O.; Shibata, A.; Sawayama, T.; Kasai, H.; Yazaki,Y.; Nakamura, Y.: Four novel KVLQT1 and four novel HERG mutationsin familial long–QT syndrome. *Circulation* 95: 565–567, 1997.

- [65153] 3483.Thomas, P. J.; Qu, B.-H.; Pedersen, P. L.: Defective proteinfolding as a basis of human disease. Trends Biochem. Sci. 20: 456–459,1995.
- [65154] 3484.Trudeau, M. C.; Warmke, J. W.; Ganetzky, B.; Robertson, G. A.: HERG, a human inward rectifier in the voltage-gated potassium channelfamily. Science 269: 92–95, 1995.
- [65155] 3485.Warmke, J. W.; Ganetzky, B.: A family of potassium channel genesrelated to eag in Drosophila and mammals. Proc. Nat. Acad. Sci. 91:3438–3442, 1994.
- [65156] 3486.Yoshida, H.; Horie, M.; Otani, H.; Kawashima, T.; Onishi, Y.;Sasayama, S.: Bradycardia-induced long QT syndrome caused by a denovo missense mutation in the S2–S3 inner loop of HERG. Am. J. Med.Genet. 98: 348–352, 2001.
- [65157] 3487.Zareba, W.; Moss, A. J.; Schwartz, P. J.; Vincent, G. M.; Robinson,J. L.; Priori, S. G.; Benhorin, J.; Locati, E. H.; Towbin, J. A.;Keating, M. T.; Lehmann, M. H.; Hall, W. J.; International Long-QTSyndrome Registry Research Group: Influence of the genotype on theclinical course of the long-QT syndrome. New Eng. J. Med. 339: 960–965,1998.
- [65158] 3488.Zhou, Z.; Gong, Q.; Epstein, M. L.; January, C. T.: HERG channel dysfunction in human long QT syndrome:



intracellular transport and functional defects. J. Biol. Chem. 273: 21061–21066, 1998.

- [65159] 3489. Rosengard, A. M.; Krutzsch, H. C.; Shearn, A.; Biggs, J. R.; Barker, E.; Margulies, I. M. K.; King, C. R.; Liotta, L. A.; Steeg, P. S.: Reduced nm23/awd protein in tumour metastasis and aberrant Drosophila development. Nature 342: 177–180, 1989.
- [65160] 3490. Steeg, P. S.; Bevilacqua, G.; Kopper, L.; Thorgeirsson, U. P.; Talmadge, J. E.; Liotta, L. A.; Sobel, M. E.: Evidence for a novel gene associated with a low tumor metastatic potential. J. Nat. Cancer Inst. 80: 200–204, 1988.
- [65161] 3491. Subramanian, C.; Cotter, M. A., II; Robertson, E. S.: Epstein–Barr virus nuclear protein EBNA–3C interacts with the human metastasis suppressor Nm23–H1: a molecular link to cancer metastasis. Nature Med. 7: 350–355, 2001.
- [65162] 3492. Varesco, L.; Caligo, M. A.; Simi, P.; Black, D. M.; Nardini, V.; Casarino, L.; Rocchi, M.; Ferrara, G.; Solomon, E.; Bevilacqua, G.: The NM23 gene maps to human chromosome band 17q22 and shows a restriction fragment length polymorphism with BglII. Genes Chromosomes Cancer 4: 84–88, 1992.
- [65163] 3493. Backer, J. M.; Mendola, C. E.; Kovesdi, I.; Fairhurst, J. L.; O'Hara, B.; Eddy, R. L., Jr.; Shows, T. B.; Mathew, S.;

Murty, V. V. V. S.; Chaganti, R. S. K.: Chromosomal localization and nucleoside diphosphate kinase activity of human metastasis-suppressor genes NM23-1 and NM23-2. *Oncogene* 8:497-502, 1993.

[65164] 3494. Chandrasekharappa, S. C.; Gross, L. A.; King, S. E.; Collins, F.S.: The human NME2 gene lies within 18kb of NME1 in chromosome 17. *Genes Chromosomes Cancer* 6: 245-248, 1993.

[65165] 3495. Gilles, A. M.; Presecan, E.; Vonica, A.; Lascu, I.: Nucleoside diphosphate kinase from human erythrocytes: structural characterization of the two polypeptide chains responsible for heterogeneity of the hexameric enzyme. *J. Biol. Chem.* 266: 8784-8789, 1991.

[65166] 3496. Kelsell, D. P.; Black, D. M.; Solomon, E.; Spurr, N. K.: Localization of a second NM23 gene, NME2, to chromosome 17q21-q22. *Genomics* 17:522-524, 1993.

[65167] 3497. Stahl, J. A.; Leone, A.; Rosengard, A. M.; Porter, L.; King, C.R.; Steeg, P. S.: Identification of a second human nm23 gene, nm23-H2. *Cancer Res.* 51: 445-449, 1991.

[65168] 3498. Boyes, J.; Bird, A.: DNA methylation inhibits transcription indirectly via a methyl-CpG binding protein. *Cell* 64: 1123-1134, 1991.

[65169] 3499. Cross, S. H.; Meehan, R. R.; Nan, X.; Bird, A.: A com-

ponent of the transcriptional repressor MeCP1 shares a motif with DNA methyltransferase and HRX proteins. *Nature Genet.* 16: 256–259, 1997.

- [65170] 3500. Hendrich, B.; Abbott, C.; McQueen, H.; Chambers, D.; Cross, S.; Bird, A.: Genomic structure and chromosomal mapping of the murine and human Mbd1, Mbd2, Mbd3, and Mbd4 genes. *Mammalian Genome* 10:906–912, 1999.
- [65171] 3501. Hempstead, B. L.; Martin-Zanca, D.; Kaplan, D. R.; Parada, L.F.; Chao, M. V.: High-affinity NGF binding requires coexpression of the trk proto-oncogene and the low-affinity NGF receptor. *Nature* 350:678–683, 1991.
- [65172] 3502. Huebner, K.; Isobe, M.; Chao, M.; Bothwell, M.; Ross, A. H.; Finan, J.; Hoxie, J. A.; Sehgal, A.; Buck, C. R.; Lanhahn, A.; Nowell, P.C.; Koprowski, H.; Croce, C. M.: The nerve growth factor receptor gene is at human chromosome region 17q12–17q22, distal to the chromosome 17 breakpoint in acute leukemias. *Proc. Nat. Acad. Sci.* 83: 1403–1407, 1986.
- [65173] 3503. Ip, N. Y.; Stitt, T. N.; Tapley, P.; Klein, R.; Glass, D. J.; Fandl, J.; Greene, L. A.; Barbacid, M.; Yancopoulos, G. D.: Similarities and differences in the way neurotrophins interact with the Trk receptors in neuronal and nonneuronal cells. *Neuron* 10: 137–149, 1993.

- [65174] 3504. Johnson, D.; Lanahan, A.; Buck, C. R.; Sehgal, A.; Morgan, C.; Mercer, E.; Bothwell, M.; Chao, M.: Expression and structure of the human NGF receptor. *Cell* 47: 545–554, 1986.
- [65175] 3505. Lee, K. F.; Li, E.; Huber, J.; Landis, S. C.; Sharpe, A. H.; Chao, M. V.; Jaenisch, R.: Targeted mutation of the gene encoding the low affinity NGF receptor p75 leads to deficits in the peripheral sensory nervous system. *Cell* 69: 737–749, 1992.
- [65176] 3506. Mischel, P. S.; Smith, S. G.; Vining, E. R.; Valletta, J. S.; Mobley, W. C.; Reichardt, L. F.: The extracellular domain of p75(NTR) is necessary to inhibit neurotrophin–3 signaling through TrkA. *J. Biol. Chem.* 276: 11294–11301, 2001.
- [65177] 3507. Rettig, W. J.; Thomson, T. M.; Spengler, B. A.; Biedler, J. L.; Old, L. J.: Assignment of human nerve growth factor receptor gene to chromosome 17 and regulation of receptor expression in somatic cell hybrids. *Somat. Cell Molec. Genet.* 12: 441–447, 1986.
- [65178] 3508. Tuffereau, C.; Benejean, J.; Blondel, D.; Kieffer, B.; Flamand, A.: Low-affinity nerve-growth factor receptor (p75NTR) can serve as a receptor for rabies virus. *EMBO J.* 17: 7250–7259, 1998.
- [65179] 3509. Welcher, A. A.; Bitler, C. M.; Radeke, M. J.; Shooter,

E. M.:Nerve growth factor binding domain of the nerve growth factor receptor. *Proc.Nat. Acad. Sci.* 88: 159–163, 1991.

- [65180] 3510.Breakefield, X. O.; Orloff, G.; Castiglione, C.; Coussens, L.;Axelrod, F. B.; Ullrich, A.: Structural gene for beta-nerve growthfactor not defective in familial dysautonomia. *Proc. Nat. Acad. Sci.* 81:4213–4216, 1984.
- [65181] 3511.Carrier, A.; Rosier, M.–F.; Guillemot, F.; Goguel, A.–F.; Pulcini,F.; Bernheim, A.; Auffray, C.; Devignes, M.–D.: Integrated physical,genetic, and genic map covering 3 Mb around the human NGF gene (NGFB)at 1p13. *Genomics* 31: 80–89, 1996.
- [65182] 3512.Chuang, H.; Prescott, E. D.; Kong, H.; Shields, S.; Jordt, S.–E.;Basbaum, A. I.; Chao, M. V.; Julius, D.: Bradykinin and nerve growthfactor release the capsaicin receptor from PtdIns(4,5)P2–mediatedinhibition. *Nature* 411: 957–962, 2001.
- [65183] 3513.Darby, J. K.; Feder, J.; Selby, M.; Riccardi, V.; Ferrell, R.;Siao, D.; Goslin, K.; Rutter, W.; Shooter, E. M.; Cavalli–Sforza,L. L.: A discordant sibship analysis between beta–NGF and neurofibromatosis. *Am.J. Hum. Genet.* 37: 52–59, 1985.
- [65184] 3514.Darby, J. K.; Kidd, J. R.; Pakstis, A. J.; Sparkes, R. S.;

Cann, H. M.; Ferrell, R. E.; Gerhard, D. G.; Riccardi, V.; Ege-land, J. A.; Shooter, E. M.; Cavalli-Sforza, L. L.; Kidd, K. K.: Linkage relationships of the gene for the beta subunit of nerve growth factor (NGFB) with other chromosome 1 marker loci. *Cytogenet. Cell Genet.* 39: 158–160, 1985.

[65185] 3515. Dracopoli, N. C.: Personal Communication. Cambridge, Mass. 4/29/1988.

[65186] 3516. Dracopoli, N. C.; Rose, E.; Whitfield, G. K.; Guidon, P. T.; Bale, S. J.; Chance, P. A.; Kourides, I. A.; Housman, D. E.: Two thyroid hormone regulated genes, the beta-subunits of nerve growth factor (NGFB) and thyroid stimulating hormone (TSHB), are located less than 310 kb apart in both human and mouse genomes. *Genomics* 3: 161–167, 1988.

[65187] 3517. Francke, U.; de Martinville, B.; Coussens, L.; Ullrich, A.: The human gene for the beta subunit of nerve growth factor is located on the proximal short arm of chromosome 1. *Science* 222: 1248–1251, 1983.

[65188] 3518. Garson, J. A.; van den Berghe, J. A.; Kemshead, J. T.: Novel non-isotopic in situ hybridization technique detects small (1 kb) unique sequences in routinely G-banded human chromosomes: fine mapping of N-myc and beta-NGF genes. *Nucleic Acids Res.* 15: 4761–4770, 1987.

- [65189] 3519.Lee, R.; Kermani, P.; Teng, K. K.; Hempstead, B. L.: Regulation of cell survival by secreted proneurotrophins. Science 294: 1945–1948, 2001.
- [65190] 3520.Levi-Montalcini, R.: The nerve growth factor thirty-five years later. Science 237: 1154–1162, 1987.
- [65191] 3521.MacInnis, B. L.; Campenot, R. B.: Retrograde support of neuronal survival without retrograde transport of nerve growth factor. Science 295:1536–1539, 2002.
- [65192] 3522.Middleton-Price, H.; van den Berghe, J.; Harding, A.; Scott, J.; Malcolm, S.: Analysis of markers on chromosome 1. (Abstract) Cytogenet. Cell Genet. 46: 662, 1987.
- [65193] 3523.Mitchell, E. L. D.; Jones, D.; White, G. R. M.; Varley, J. M.; Santibanez Koref, M. F.: Determination of the gene order of the three loci CD2, NGFB, and NRAS at human chromosome band 1p13 and refinement of their localisation at the subband level by fluorescence in situ hybridization. Cytogenet. Cell Genet. 70: 183–185, 1995.
- [65194] 3524.Munke, M.; Lindgren, V.; de Martinville, B.; Francke, U.: Comparative analysis of mouse–human hybrids with rearranged chromosomes 1 by in situ hybridization and Southern blotting: high-resolution mapping of NRAS, NGFB, and AMY on human chromosome 1. Somat. Cell Molec. Genet. 10: 589–599, 1984.

- [65195] 3525.Ramer, M. S.; Priestley, J. V.; McMahon, S. B.: Functional regeneration of sensory axons into the adult spinal cord. *Nature* 403: 312–316, 2000.
- [65196] 3526.Sanico, A. M.; Stanisz, A. M.; Gleeson, T. D.; Bora, S.; Proud, D.; Bienenstock, J.; Koliatsos, V. E.; Togias, A.: Nerve growth factor expression and release in allergic inflammatory disease of the upper airways. *Am. J. Resp. Crit. Care Med.* 161: 1631–1635, 2000.
- [65197] 3527.Ullrich, A.; Gray, A.; Berman, C.; Dull, T. J.: Human beta-nerve growth factor gene sequence highly homologous to that of mouse. *Nature* 303: 821–825, 1983.
- [65198] 3528.Beauchemin, N.; Draber, P.; Dveksler, G.; Gold, P.; Gray-Owen, S.; Grunert, F.; Hammarstrom, S.; Holmes, K. V.; Karlsson, A.; Kuroki, M.; Lin, S.-H.; Lucka, L.; and 13 others: Redefined nomenclature for members of the carcinoembryonic antigen family. *Exp. Cell Res.* 252: 243–249, 1999.
- [65199] 3529.Inazawa, J.; Abe, T.; Inoue, K.; Misawa, S.; Oikawa, S.; Nakazato, H.; Yoshida, M. C.: Regional assignment of nonspecific cross-reacting antigen (NCA) of the CEA gene family to chromosome 19 at band q13.2. *Cytogenet. Cell Genet.* 52: 28–31, 1989.
- [65200] 3530.Willcocks, T. C.; Craig, S. P.; Craig, I. W.: Assignment



of the coding sequence for carcinoembryonic antigen (CEA) and normal cross-reacting antigen (NCA) to human chromosome 19q13. *Ann. Hum. Genet.* 53: 141–148, 1989.

[65201] 3531. Qian, F.; Kruse, U.; Lichter, P.; Sippel, A. E.: Chromosomal localization of the four genes (NFIA, B, C, and X) for the human transcription factor nuclear factor I by FISH.

*Genomics* 28: 66–73, 1995.

[65202] 3532. Santoro, C.; Mermod, N.; Andrews, P. C.; Tjian, R.: A family of human CCAAT-box-binding proteins active in transcription and DNA replication: cloning and expression of multiple cDNAs. *Nature* 334: 218–224, 1988.

[65203] 3533. Scherthan, H.; Seisenberger, C.; Greulich, K.; Winnacker, E.-L.: Mapping of the murine nuclear factor I/X gene (Nfix) to mouse chromosome 8 C1–2 by FISH. *Genomics* 22: 247–249, 1994.

[65204] 3534. Seisenberger, C.; Winnacker, E.-L.; Scherthan, H.: Localisation of the human nuclear factor I/X (NFI/X) gene to chromosome 19p13 and detection of five other related loci at 1p21–22, 1q42–43, 5q15, 11p13 and 20q13 by FISH. *Hum. Genet.* 91: 535–537, 1993.

[65205] 3535. Auphan, N.; DiDonato, J. A.; Rosette, C.; Helmberg, A.; Karin, M.: Immunosuppression by glucocorticoids: inhibition of NF- $\kappa$ B activity through induction of I-

kappa-B synthesis. Science 270: 286–290,1995.

[65206] 3536.Baeuerle, P. A.: I-kappa-B--NF-kappa-B structures: at the interface of inflammation control. Cell 95: 729–731, 1998.

[65207] 3537.Haskill, S.; Beg, A. A.; Tompkins, S. M.; Morris, J. S.; Yurochko,A. D.; Sampson-Johannes, A.; Mondal, K.; Ralph, P.; Baldwin, A. S.,Jr.: Characterization of an immediate-early gene induced in adherent monocytes that encodes I-kappa-B-like activity. Cell 65: 1281–1289,1991.

[65208] 3538.Huxford, T.; Huang, D.-B.; Malek, S.; Ghosh, G.: The crystal structure of the I-kappa-B-alpha/NF-kappa-B complex reveals mechanisms of NF-kappa-Binactivation. Cell 95: 759–770, 1998.

[65209] 3539.Ito, C. Y.; Adey, N.; Bautch, V. L.; Baldwin, A. S., Jr.: Structure and evolution of the human IKBA gene. Genomics 29: 490–495, 1995.

[65210] 3540.Jacobs, M. D.; Harrison, S. C.: Structure of an I-kappa-B-alpha/NF-kappa-B complex. Cell 95: 749–758, 1998.

[65211] 3541.Jung, M.; Zhang, Y.; Lee, S.; Ditschilo, A.: Correction of radiation sensitivity in ataxia telangiectasia cells by a truncated I-kappa-B-alpha. Science 268:1619–1621, 1995.

- [65212] 3542. Le Beau, M. M.; Ito, C.; Cogswell, P.; Espinosa, R., III; Fernald, A. A.; Baldwin, A. S., Jr.: Chromosomal localization of the genes encoding the p50/p105 subunits of NF- $\kappa$ -B (NFKB2) and the I- $\kappa$ -B/MAD-3 (NFKBI) inhibitor of NF- $\kappa$ -B to 4q24 and 14q13, respectively. *Genomics* 14:529–531, 1992.
- [65213] 3543. Neish, A. S.; Gewirtz, A. T.; Zeng, H.; Young, A. N.; Hobert, M. E.; Karmali, V.; Rao, A. S.; Madara, J. L.: Prokaryotic regulation of epithelial responses by inhibition of I- $\kappa$ -B- $\alpha$  ubiquitination. *Science* 289:1560–1563, 2000.
- [65214] 3544. Rupec, R. A.; Poujol, D.; Grosgeorge, J.; Carle, G. F.; Livolsi, A.; Peyron, J.-F.; Schmid, R. M.; Baeuerle, P. A.; Messer, G.: Structural analysis, expression, and chromosomal localization of the mouse *ikb* gene. *Immunogenetics* 49: 395–403, 1999.
- [65215] 3545. Scheinman, R. I.; Cogswell, P. C.; Lofquist, A. K.; Baldwin, A. S., Jr.: Role of transcriptional activation of I- $\kappa$ -B- $\alpha$  in mediation of immunosuppression by glucocorticoids. *Science* 270:283–286, 1995.
- [65216] 3546. Compton, D. A.; Szilak, I.; Cleveland, D. W.: Primary structure of NuMA, an intranuclear protein that defines a novel pathway for segregation of proteins at mitosis. *J. Cell*

Biol. 116: 1395–1408,1992.

- [65217] 3547.Lydersen, B. K.; Pettijohn, D. E.: Human-specific nuclear protein that associates with the polar region of the mitotic apparatus: distribution in a human/hamster hybrid cell. Cell 22: 489–499, 1980.
- [65218] 3548.Merdes, A.; Ramyar, K.; Vechio, J. D.; Cleveland, D. W.: A complex of NuMA and cytoplasmic dynein is essential for mitotic spindle assembly. Cell 87:447–458, 1996.
- [65219] 3549.Sparks, C. A.; Bangs, P. L.; McNeil, G. P.; Lawrence, J. B.; Fey, E. G.: Assignment of the nuclear mitotic apparatus protein NuMA gene to human chromosome 11q13. Genomics 17: 222–224, 1993.
- [65220] 3550.Wells, R. A.; Catzavelos, C.; Kamel-Reid, S.: Fusion of retinoic acid receptor alpha to NuMA, the nuclear mitotic apparatus protein, by a variant translocation in acute promyelocytic leukaemia. Nature Genet. 17: 109–113, 1997.
- [65221] 3551.Wiese, C.; Wilde, A.; Moore, M. S.; Adam, S. A.; Merdes, A.; Zheng, Y.: Role of importin-beta in coupling Ran to downstream targets in microtubule assembly. Science 291: 653–656, 2001.
- [65222] 3552.Yang, C. H.; Lambie, E. J.; Snyder, M.: NuMA: an unusually long coiled-coil related protein in the mammalian

nucleus. J. Cell Biol. 116:1303–1317, 1992.

[65223] 3553.Halila, R.; Apostolou, S.; Winqvist, R.; Callen, D.; Prockop, D.J.; Peltonen, L.: Isolation and genomic assignment of a candidatecDNA clone for type III procollagen N–proteinase. (Abstract) Am.J. Hum. Genet. 51 (suppl.): A128 only, 1992.

[65224] 3554.Halila, R.; Peltonen, L.; Prockop, D. J.: Isolation of a candidatecDNA clone for type III procollagen N–proteinase from human placentalcDNA library. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A192 only,1989.

[65225] 3555.Nomura, N.; Nagase, T.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Sato,S.; Seki, N.; Kawarabayasi, Y.; Ishikawa, K.; Tabata, S.: Predictionof the coding sequences of unidentified human genes. II. The codingsequences of 40 new genes (KIAA0041–KIAA0080) deduced by analysisof cDNA clones from human cell line KG–1. DNA Res. 1: 223–229, 1994.

[65226] 3556.Freedman, D. A.; Epstein, C. B.; Roth, J. C.; Levine, A. J.: Agenetic approach to mapping the p53 binding site in the MDM2 protein. Molec.Med. 3: 248–259, 1997.

[65227] 3557.Fuchs, S. Y.; Adler, V.; Buschmann, T.; Yin, Z.; Wu, X.; Jones,S. N.; Ronai, Z.: JNK targets p53 ubiquitination and degradatin in nonstressed cells. Genes Dev. 12:

2658–2663, 1998.

- [65228] 3558.Mayo, L. D.; Donner, D. B.: A phosphatidylinositol 3–kinase/Akt pathway promotes translocation of Mdm2 from the cytoplasm to the nucleus. *Proc.Nat. Acad. Sci.* 98: 11598–11603, 2001.
- [65229] 3559.Mitchell, E. L. D.; White, G. R. M.; Santibanez–Koref, M. F.;Varley, J. M.; Heighway, J.: Mapping of gene loci in the q13–q15 region of chromosome 12. *Chromosome Res.* 3: 261–262, 1995.
- [65230] 3560.Momand, J.; Zambetti, G. P.; Olson, D. C.; George, D. L.; Levine,A. J.: The mdm–2 oncogene product forms a complex with the p53 protein and inhibits p53–mediated transactivation. *Cell* 69: 1237–1245, 1992.
- [65231] 3561.Oliner, J. D.; Kinzler, K. W.; Meltzer, P. S.; George, D. L.;Vogelstein, B.: Amplification of a gene encoding a p53–associated protein in human sarcomas. *Nature* 358: 80–83, 1992.
- [65232] 3562.Ries, S.; Biederer, C.; Woods, D.; Shifman, O.; Shirasawa, S.;Sasazuki, T.; McMahon, M.; Oren, M.; McCormick, F.: Opposing effects of Ras on p53: transcriptional activation of mdm2 and induction of p19(ARF). *Cell* 103: 321–330, 2000.
- [65233] 3563.Shenoy, S. K.; McDonald, P. H.; Kohout, T. A.;

Lefkowitz, R. J.: Regulation of receptor fate by ubiquitination of activated beta-2-adrenergic receptor and beta-arrestin. *Science* 294: 1307–1313, 2001.

[65234] 3564. Shieh, S.-Y.; Ikeda, M.; Taya, Y.; Prives, C.: DNA damage-induced phosphorylation of p53 alleviates inhibition by MDM2. *Cell* 91: 325–334, 1997.

- [65235] 3565. Testa, J. R.; Bellacosa, A.: AKT plays a central role in tumorigenesis. *Proc. Nat. Acad. Sci.* 98: 10983–10985, 2001.
- [65236] 3566. Xiao, Z.-X.; Chen, J.; Levine, A. J.; Modjtahedi, N.; Xing, J.; Sellers, W. R.; Livingston, D. M.: Interaction between the retinoblastoma protein and the oncoprotein MDM2. *Nature* 375: 694–698, 1995.
- [65237] 3567. Zhang, Y.; Xiong, Y.: A p53 amino-terminal nuclear export signal inhibited by DNA damage-induced phosphorylation. *Science* 292: 1910–1915, 2001.
- [65238] 3568. Balazs, I.: Personal Communication. New York, N. Y. 8/23/1983.
- [65239] 3569. Bezieau, S.; Devilder, M.-C.; Avet-Loiseau, H.; Mellerin, M.-P.; Puthier, D.; Pennarun, E.; Rapp, M.-J.; Harousseau, J.-L.; Moisan, J.-P.; Bataille, R.: High incidence of N and K-Ras activating mutations in multiple myeloma and primary plasma cell leukemia at diagnosis. *Hum. Mutat.* 18: 212–224, 2001.
- [65240] 3570. Bos, J. L.; Toksoz, D.; Marshall, C. J.; Verlaan-de Vries, M.; Veeneman, G. H.; van der Eb, A. J.; van Boom, J. H.; Janssen, J. W. G.; Steenvoorden, A. C. M.: Amino-acid substitutions at codon 13 of the N-ras oncogene in human acute myeloid leukaemia. *Nature* 315: 726–730, 1985.



- [65241] 3571.Davis, M.; Malcolm, S.; Hall, A.: The N-ras oncogene is located on the short arm of chromosome 1. (Abstract) Cytogenet. Cell Genet. 37:448–449, 1984.
- [65242] 3572.Davis, M.; Malcolm, S.; Hall, A.; Marshall, C. J.: Localisation of the human N-ras oncogene to chromosome 1cen–p21 by in situ hybridisation. EMBOJ. 2: 2281–2283, 1983.
- [65243] 3573.Deka, R.; Majumder, P. P.; Warren, A. C.; Surti, U.; Hoffner, L.; Hauselman, E.; Antonarakis, S. E.; Ferrell, R. E.; Chakravarti, A.: Gene-centromere mapping using ovarian teratomas: results from chromosomes 1p, 13q and 21q. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A137 only, 1989.
- [65244] 3574.de Martinville, B.; Cunningham, J. M.; Murray, M. J.; Francke, U.: The N-ras oncogene assigned to chromosome 1 (p31–cen) by somatic cell hybrid analysis. (Abstract) Cytogenet. Cell Genet. 37: 531 only, 1984.
- [65245] 3575.Hall, A.; Brown, R.: Human N-ras: cDNA cloning and gene structure. Nucleic Acids Res. 13: 5255–5268, 1985.
- [65246] 3576.Hall, A.; Marshall, C. J.; Spurr, N. K.; Weiss, R. A.: Identification of transforming gene in two human sarcoma cell lines as a new member of the ras gene family located on chromosome 1. Nature 303: 396–400, 1983.

- [65247] 3577.Linder, D.; McCaw, B. F.; Hecht, F.: Parthenogenic origin of benign ovarian teratomas. *New Eng. J. Med.* 292: 63–66, 1975.
- [65248] 3578.Nitta, N.; Ochiai, M.; Nagao, M.; Sugimura, T.: Amino-acid substitution at codon 13 of the N-ras oncogene in rectal cancer in a Japanese patient. *Jpn.J. Cancer Res.* 78: 21–26, 1987.
- [65249] 3579.Nobori, T.; Hexdall, L. E.; Carson, D. A.: A polymorphic region defined by pCN2 (the 3-prime nontranslated region of N-ras) maps to chromosome 9cen-p12. *Hum. Genet.* 87: 433–437, 1991.
- [65250] 3580.Parrington, J. M.; West, L. F.; Povey, S.: The origin of ovarian teratomas. *J. Med. Genet.* 21: 4–12, 1984.
- [65251] 3581.Popescu, N. C.; Amsbaugh, S. C.; DiPaolo, J. A.; Tronick, S. R.; Aaronson, S. A.; Swan, D. C.: Chromosomal localization of three human ras genes by in situ molecular hybridization. *Somat. Cell Molec. Genet.* 11: 149–155, 1985.
- [65252] 3582.Povey, S.; Morton, N. E.; Sherman, S. L.: Report of the committee on the genetic constitution of chromosomes 1 and 2 (HGM8). *Cytogenet. Cell Genet.* 40: 67–106, 1985.
- [65253] 3583.Rabin, M.; Watson, M.; Barker, P.; Ryan, J.; Breg, W. R.; Ruddle, F. H.: Chromosomal assignment of human c-

fos and N-ras oncogenes.(Abstract) Am. J. Hum. Genet.  
35: 148A only, 1983.

- [65254] 3584.Rabin, M.; Watson, M.; Barker, P. E.; Ryan, J.; Breg, W. R.; Ruddle,F. H.: NRAS transforming gene maps to region p11–p13 on chromosome1 by in situ hybridization. Cytogenet. Cell Genet. 38: 70–72, 1984.
- [65255] 3585.Ryan, J.; Barker, P. E.; Shimizu, K.; Wigler, M.; Ruddle, F. H.: Chromosomal assignment of a family of human oncogenes. Proc. Nat.Acad. Sci. 80: 4460–4463, 1983.
- [65256] 3586.Ashkar, S.; Weber, G. F.; Panoutsakopoulou, V.; Sanchirico, M.E.; Jansson, M.; Zawaideh, S.; Rittling, S. R.; Denhardt, D. T.; Glimchar,M. J.; Cantor, H.: Eta–1 (osteopontin): an early component of type–1(cell–mediated) immunity. Science 287: 860–864, 2000.
- [65257] 3587.Baccarini–Contri, M.; Taparelli, F.; Pasquali–Ronchetti, I.: Osteopontinis a constitutive component of normal elastic fibers in human skinand aorta. Matrix Biol. 14: 553–560, 1994.
- [65258] 3588.Beck, G. R., Jr.; Zerler, B.; Moran, E.: Phosphate is a specificsignal for induction of osteopontin gene expression. Proc. Nat. Acad.Sci. 97: 8352–8357, 2000.
- [65259] 3589.Crosby, A. H.; Lyu, M. S.; Lin, K.; McBride, O. W.; Kerr, J. M.;Aplin, H. M.; Fisher, L. W.; Young, M. F.; Kozak,

C. A.; Dixon, M.J.: Mapping of the human and mouse bone sialoprotein and osteopontin loci. *Mammalian Genome* 7: 149–151, 1996.

[65260] 3590. Fisher, L. W.; McBride, O. W.; Termine, J. D.; Young, M. F.: Human bone sialoprotein: deduced protein sequence and chromosomal localization. *J. Biol. Chem.* 265: 2347–2351, 1990.

[65261] 3591. Graf, K.; Do, Y. S.; Ashizawa, N.; Meehan, W. P.; Giachelli, C.M.; Marboe, C. C.; Fleck, E.; Hsueh, W. A.: Myocardial osteopontin expression is associated with left ventricular hypertrophy. *Circulation* 96:3063–3071, 1997.

[65262] 3592. Kiefer, M. C.; Bauer, D. M.; Barr, P. J.: The cDNA and derived amino acid sequence for human osteopontin. *Nucleic Acids Res.* 17:3306, 1989.

[65263] 3593. Kim, J.-H.; Skates, S. J.; Ueda, T.; Wong, K.; Schorge, J. O.; Feltmate, C. M.; Berkowitz, R. S.; Cramer, D. W.; Mok, S. C.: Osteopontin as a potential diagnostic biomarker for ovarian cancer. *J.A.M.A.* 287:1671–1679, 2002.

[65264] 3594. Kohri, K.; Nomura, S.; Kitamura, Y.; Nagata, T.; Yoshioka, K.; Iguchi, M.; Yamate, T.; Umekawa, T.; Suzuki, Y.; Sinohara, H.; Kurita, T.: Structure and expression of the mRNA encoding urinary stone protein (osteopontin). *J. Biol. Chem.* 268: 15180–15184, 1993.

- [65265] 3595.Kohri, K.; Suzuki, Y.; Yoshida, K.; Yamamoto, K.; Amasaki, N.;Yamate, T.; Umekawa, T.; Iguchi, M.; Sino-hara, H.; Kurita, T.: Molecularcloning and sequencing of cDNA encoding urinary stone protein, whichis identical to osteopontin. *Biochem. Biophys. Res. Commun.* 184:859–864, 1992.
- [65266] 3596.Liaw, L.; Birk, D. E.; Ballas, C. B.; Whitsitt, J. S.; Davidson,J. M.; Hogan, B. L. M.: Altered wound healing in mice lacking a functionalosteopontin gene (spp1). *J. Clin. Invest.* 101: 1468–1478, 1998.
- [65267] 3597.Morimoto, I.; Sasaki, Y.; Ishida, S.; Imai, K.; Tokino, T.: Identificationof the osteopontin gene as a direct target of TP53. *Genes ChromosomesCancer* 33: 270–278, 2002.
- [65268] 3598.Reinholt, F. P.; Hultenby, K.; Oldberg, A.; Heinegard, D.: Osteopontin--apossible anchor of osteoclasts to bone. *Proc. Nat. Acad. Sci.* 87:4473–4475, 1990.
- [65269] 3599.Singh, K.; Balligand, J.–L.; Fischer, T. A.; Smith, T. W.; Kelly,R. A.: Glucocorticoids increase osteopontin expression in cardiacmyocytes and microvascular endothelial cells: role in regulation ofinducible nitric oxide synthase. *J. Biol. Chem.* 270: 28471–28478,1995.
- [65270] 3600.Weber, G. F.; Ashkar, S.; Glimcher, M. J.; Cantor, H.: Receptor–ligandinteraction between CD44 and osteopon–

tin (Eta-1). Science 271: 509-512,1996.

- [65271] 3601. Staal, A.; van Wijnen, A. J.; Birkenhager, J. C.; Pols, H. A.P.; Prah, J.; DeLuca, H.; Gaub, M.-P.; Lian, J. B.; Stein, G. S.; van Leeuwen, J. P. T. M.; Stein, J. L.: Distinct conformations of vitamin D receptor/retinoid X receptor-alpha heterodimers are specified by dinucleotide differences in the vitamin D-responsive elements of the osteocalcin and osteopontin genes. Molec. Endocr. 10: 1444-1456,1996.
- [65272] 3602. Young, M. F.; Kerr, J. M.; Termine, J. D.; Wewer, U. M.; Wang, M. G.; McBride, O. W.; Fisher, L. W.: cDNA cloning, mRNA distribution and heterogeneity, chromosomal location and RFLP analysis of human osteopontin. Genomics 7: 491-502, 1990.
- [65273] 3603. Vogelstein, B.; Fearon, E. R.; Hamilton, S. R.; Kern, S. E.; Preisinger, A. C.; Leppert, M.; Nakamura, Y.; White, R.; Smits, A. M. M.; Bos, J. L.: Genetic alterations during colorectal-tumor development. New Eng. J. Med. 319: 525-532, 1988.
- [65274] 3604. Zhao, C.; Takita, J.; Tanaka, Y.; Setou, M.; Nakagawa, T.; Takeda, S.; Yang, H. W.; Terada, S.; Nakata, T.; Takei, Y.; Saito, M.; Tsuji, S.; Hayashi, Y.; Hirokawa, N.: Charcot-Marie-Tooth disease type 2A caused by mutation in a microtubule motor KIF1B-beta. Cell 105:587-597, 2001.

- [65275] 3605.Maemura, K.; Kurihara, H.; Kurihara, Y.; Oda, H.; Ishikawa, T.;Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Yazaki, Y.: Sequenceanalysis, chromosomal location, and developmental expression of themouse preproendothelin-1 gene. *Genomics* 31: 177–184, 1996.
- [65276] 3606.Maggi, M.; Barni, T.; Fantoni, G.; Mancina, R.; Pupilli, C.; Luconi,M.; Crescioli, C.; Serio, M.; Vannelli, G. B.: Expression and biologicaleffects of endothelin-1 in human gonadotropin-releasing hormone-secretingneurons. *J. Clin. Endocr. Metab.* 85: 1658–1665, 2000.
- [65277] 3607.Napolitano, M.; Miceli, F.; Calce, A.; Vacca, A.; Gulino, A.;Apa, R.; Lanzone, A.: Expression and relationship between endothelin-1messenger ribonucleic acid (mRNA) and inducible/endothelial nitricoxide synthase mRNA isoforms from normal and preeclamptic placentas. *J.Clin. Endocr. Metab.* 85: 2318–2323, 2000.
- [65278] 3608.Okafor, M. C.; Delamere, N. A.: The inhibitory influence of endothelinon active sodium-potassium transport in porcine lens. *Invest. Ophthal.Vis. Sci.* 42: 1018–1023, 2001.
- [65279] 3609.Pache, M.; Kaiser, H. J.; Haufschild, T.; Lubeck, P.; Flammer,J.: Increased endothelin-1 plasma levels in giant cell arteritis:a report on four patients. *Am. J. Ophthal.*

133: 160–162, 2002.

- [65280] 3610.Pages, J.-C.; Drieu, C.; Blanche, H.; Beckmann, J.; Cann, H. M.: A short tandem repeat polymorphism at the endothelin 1 (EDN1) locus. *Hum.Molec. Genet.* 2: 90, 1993.
- [65281] 3611.Pezzetti, F.; Scapoli, L.; Martinelli, M.; Carinci, F.; Brunelli,G.; Carls, F. P.; Palomba, F.; Gombos, F.; Carinci, P.; Tognon, M.: Linkage analysis of candidate endothelin pathway genes in nonsyndromicfamilial orofacial cleft. *Ann. Hum. Genet.* 64: 341–347, 2000.
- [65282] 3612.Yanagisawa, H.; Hammer, R. E.; Richardson, J. A.; Williams, S.C.; Clouthier, D. E.; Yanagisawa, M.: Role of endothelin–1/endothelin–Areceptor–mediated signaling pathway in the aortic arch patterningin mice. *J. Clin. Invest.* 102: 22–33, 1998.
- [65283] 3613.Yanagisawa, H.; Yanagisawa, M.; Kapur, R. P.; Richardson, J. A.;Williams, S. C.; Clouthier, D. E.; de Wit, D.; Emoto, N.; Hammer,R. E.: Dual genetic pathways of endothelin–mediated intercellularsignaling revealed by targeted disruption of endothelin convertingenzyme–1 gene. *Development* 125: 825–836, 1998.
- [65284] 3614.Zeidel, M. L.; Brady, H. R.; Kone, B. C.; Gullans, S. R.; Brenner,B. M.: Endothelin, a peptide inhibitor of



Na(+)-K(+)-ATPase in intact renal tubular epithelial cells.  
Am. J. Physiol. 257: C1101-C1107, 1989.

- [65285] 3615. Bloch, K. D.; Hong, C. C.; Eddy, R. L.; Shows, T. B.; Quertermous, T.: cDNA cloning and chromosomal assignment of the endothelin 2 gene: vasoactive intestinal contractor peptide is rat endothelin 2. Genomics 10:236-242, 1991.
- [65286] 3616. Deng, A. Y.; Dene, H.; Pravenec, M.; Rapp, J. P.: Genetic mapping of two new blood pressure quantitative trait loci in the rat by genotyping endothelin system genes. J. Clin. Invest. 93: 2701-2709, 1994.
- [65287] 3617. Ohkubo, S.; Ogi, K.; Hosoya, M.; Matsumoto, H.; Suzuki, N.; Kimura, C.; Onda, H.; Fujino, M.: Specific expression of human endothelin-2 (ET-2) gene in a renal adenocarcinoma cell line: molecular cloning of cDNA encoding the precursor of ET-2 and its characterization. FEBS Lett. 274: 136-140, 1990.
- [65288] 3618. Baynash, A. G.; Hosoda, K.; Giaid, A.; Richardson, J. A.; Emoto, N.; Hammer, R. E.; Yanagisawa, M.: Interaction of endothelin-3 with endothelin-B receptor is essential for development of epidermal melanocytes and enteric neurons. Cell 79: 1277-1285, 1994.
- [65289] 3619. Gros, P.; Croop, J.; Housman, D.: Mammalian mul-

tidrug resistance gene: complete cDNA sequence indicates strong homology to bacterial transport proteins. *Cell* 47: 371–380, 1986.

- [65290] 3620. Gros, P.; Neriah, Y. B.; Croop, J. M.; Housman, D. E.: Isolation and expression of a complementary DNA that confers multidrug resistance. *Nature* 323:728–731, 1986.
- [65291] 3621. Lerman, C.; Caporaso, N. E.; Audrain, J.; Main, D.; Bowman, E. D.; Lockshin, B.; Boyd, N. R.; Shields, P. G.: Evidence suggesting the role of specific genetic factors in cigarette smoking. *Health Psych.* 18: 14–20, 1999.
- [65292] 3622. Lossie, A. C.; Vandenberg, D. J.; Uhl, G. R.; Camper, S. A.: Localization of the dopamine transporter gene, *Dat1*, on mouse chromosome
- [65293] 3623. *Mammalian Genome* 5: 117–118, 1994. 13. Sabol, S. Z.; Nelson, M. L.; Fisher, C.; Gunzerath, L.; Brody, C. L.; Hu, S.; Sirota, L. A.; Marcus, S. E.; Greenberg, B. D.; Lucas, F. R., IV; Benjamin, J.; Murphy, D. L.; Hamer, D. H.: A genetic association for cigarette smoking behavior. *Health Psych.* 18: 7–13, 1999.
- [65294] 3624. Tiihonen, J.; Kuikka, J.; Bergstrom, K.; Hakola, P.; Karhu, J.; Ryyanen, O.-P.; Fohr, J.: Altered striatal dopamine re-uptake site densities in habitually violent and non-violent alcoholics. *Nature Med.* 1: 654–657, 1995.

- [65295] 3625.Vandenbergh, D. J.; Persico, A. M.; Hawkins, A. L.; Griffin, C.A.; Li, X.; Jabs, E. W.; Uhl, G. R.: Human dopamine transporter gene(DAT1) maps to chromosome 5p15.3 and displays a VNTR. *Genomics* 14:1104–1106, 1992.
- [65296] 3626.Vandenbergh, D. J.; Persico, A. M.; Uhl, G. R.: A human dopaminetransporter cDNA predicts reduced glycosylation, displays a novelrepetitive element and provides racially–dimorphic TaqI RFLPs. *Molec.Brain Res.* 15: 161–166, 1992.
- [65297] 3627.Waldman, I. D.; Rowe, D. C.; Abramowitz, A.; Kozel, S. T.; Mohr,J. H.; Sherman, S. L.; Cleveland, H. H.; Sanders, M. L.; Gard, J.M. C.; Stever, C.: Association and linkage of the dopamine transportergene and attention–deficit hyperactivity disorder in children: heterogeneityowing to diagnostic subtype and severity. *Am. J. Hum. Genet.* 63:1767–1776, 1998.
- [65298] 3628.Huang, H.; Mahler–Araujo, B. M.; Sankila, A.; Chimelli, L.; Yonekawa,Y.; Kleihues, P.; Ohgaki, H.: APC mutations in sporadic medulloblastomas. *Am.J. Path.* 156: 433–437, 2000.
- [65299] 3629.Behan, D. P.; Potter, E.; Lewis, K. A.; Jenkins, N. A.; Copeland,N.; Lowry, P. J.; Vale, W. W.: Cloning and struc–

ture of the humancorticotropin releasing factor-binding protein gene (CRHBP). *Genomics* 16:63–68, 1993.

- [65300] 3630.Lublin, D. M.; Lemons, R. S.; Le Beau, M. M.; Holers, V. M.; Tykocinski, M. L.; Medof, M. E.; Atkinson, J. P.: The gene encoding decay-accelerating factor (DAF) is located in the complement-regulatory locus on the long arm of chromosome 1. *J. Exp. Med.* 165: 1731–1736, 1987.
- [65301] 3631.von Boehmer, H.; Sarukhan, A.: GAD, a single autoantigen for diabetes. *Science* 284: 1135–1136, 1999.
- [65302] 3632.Williamson, E. A.; Ince, P. G.; Harrison, D.; Kendall-Taylor, P.; Harris, P. E.: G-protein mutations in human pituitary adrenocorticotrophic hormone-secreting adenomas. *Europ. J. Clin. Invest.* 25: 128–131, 1995.
- [65303] 3633.Marsden, V. S.; O'Connor, L.; O'Reilly, L. A.; Silke, J.; Metcalf, D.; Ekert, P. G.; Huang, D. C. S.; Cecconi, F.; Kuida, K.; Tomaselli, K. J.; Roy, S.; Nicholson, D. W.; Vaux, D. L.; Bouillet, P.; Adams, J. M.; Strasser, A.: Apoptosis initiated by Bcl-2-regulated caspase activation independently of the cytochrome c/Apaf-1/caspase-9 apoptosome. *Nature* 419:634–637, 2002.
- [65304] 3634.Huelsken, J.; Vogel, R.; Erdmann, B.; Cotsarelis, G.; Birchmeier, W.: Beta-catenin controls hair follicle morphogenesis and stem cell differentiation in the skin. *Cell* 105:

533–545, 2001.

- [65305] 3635. Ilyas, M.; Tomlinson, I. P. M.; Rowan, A.; Pignatelli, M.; Bodmer, W. F.: Beta-catenin mutations in cell lines established from human colorectal cancers. *Proc. Nat. Acad. Sci.* 94: 10330–10334, 1997.
- [65306] 3636. Iwao, K.; Nakamori, S.; Kameyama, M.; Imaoka, S.; Kinoshita, M.; Fukui, T.; Ishiguro, S.; Nakamura, Y.; Miyoshi, Y.: Activation of the beta-catenin gene by interstitial deletions involving exon 3 in primary colorectal carcinomas without adenomatous polyposis coli mutations. *Cancer Res.* 58: 1021–1026, 1998.
- [65307] 3637. Kawasaki, Y.; Senda, T.; Ishidata, T.; Koyama, R.; Morishita, T.; Iwayama, Y.; Higuchi, O.; Akiyama, T.: Asef, a link between the tumor suppressor APC and G-protein signaling. *Science* 289: 1194–1197, 2000.
- [65308] 3638. Koch, A.; Denkhaus, D.; Albrecht, S.; Leuschner, I.; von Schweinitz, D.; Pietsch, T.: Childhood hepatoblastomas frequently carry a mutated degradation targeting box of the beta-catenin gene. *Cancer Res.* 59: 269–273, 1999.
- [65309] 3639. Korinek, V.; Barker, N.; Morin, P. J.; van Wichen, D.; de Weger, R.; Kinzler, K. W.; Vogelstein, B.; Clevers, H.: Constitutive transcriptional activation by a beta-catenin–Tcf complex in APC–/– colon carcinoma. *Science*

275:1784–1787, 1997.

[65310] 3640.Kraus, C.; Liehr, T.; Hulsken, J.; Behrens, J.; Birchmeier, W.;Grzeschik, K.–H.; Ballhausen, W. G.: Localization of the human beta–catenin gene (CTNNB1) to 3p21: a region implicated in tumor development. *Genomics* 23:272–274, 1994.

[65311] 3641.Legoix, P.; Bluteau, O.; Bayer, J.; Perret, C.; Balabaud, C.;Belghiti, J.; Franco, D.; Thomas, G.; Laurent–Puig, P.; Zucman–Rossi,J.: Beta–catenin mutations in hepatocellular carcinoma correlatewith a low rate of loss of heterozygosity. *Oncogene* 18: 4044–4046,1999.

[65312] 3642.Lin, S.–Y.; Xia, W.; Wang, J. C.; Kwong, K. Y.; Spohn, B.; Wen,Y.; Pestell, R. G.; Hung, M.–C.: Beta–catenin, a novel prognosticmarker for breast cancer: its roles in cyclin D1 expression and cancerprogression. *Proc. Nat. Acad. Sci.* 97: 4262–4266, 2000.

[65313] 3643.McCrea, P. D.; Turck, C. W.; Gumbiner, B.: A homolog of the armadilloprotein in *Drosophila* (plakoglobin) associated with E–cadherin. *Science* 254:1359–1361, 1991.

[65314] 3644.Morin, P. J.; Sparks, A. B.; Korinek, V.; Barker, N.; Clevers,H.; Vogelstein, B.; Kinzler, K. W.: Activation of beta–catenin–Tcfsignaling in colon cancer by mutations in

beta-catenin or APC. Science 275:1787–1790, 1997.

- [65315] 3645. Murase, S.; Mosser, E.; Schuman, E. M.: Depolarization drives beta-catenin into neuronal spines promoting changes in synaptic structure and function. Neuron 35: 91–105, 2002.
- [65316] 3646. Nollet, F.; Berx, G.; Molemans, F.; van Roy, F.: Genomic organization of the human beta-catenin gene (CTNNB1). Genomics 32: 413–424, 1996.
- [65317] 3647. Peifer, M.: Cancer, catenins, and cuticle pattern: a complex connection. Science 262: 1667–1668, 1993.
- [65318] 3648. Rodova, M.; Islam, M. R.; Maser, R. L.; Calvet, J. P.: The polycystic kidney disease–1 promoter is a target of the beta-catenin/T-cell factor pathway. J. Biol. Chem. 277: 29577–29583, 2002.
- [65319] 3649. Roose, J.; Huls, G.; van Beest, M.; Moerer, P.; van der Horn, K.; Goldschmeding, R.; Logtenberg, T.; Clevers, H.: Synergy between tumor suppressor APC and the beta-catenin–Tcf4 target Tcf1. Science 285: 1923–1926, 1999.
- [65320] 3650. Rubinfeld, B.; Robbins, P.; El-Gamil, M.; Albert, I.; Porfiri, E.; Polakis, P.: Stabilization of beta-catenin by genetic defects in melanoma cell lines. Science 275: 1790–1792, 1997.
- [65321] 3651. Saadi-Kheddouchi, S.; Berrebi, D.; Romagnolo, B.;

Cluzeaud, F.;Peuchmaur, M.; Kahn, A.; Vandewalle, A.; Perret, C.: Early developmentof polycystic kidney disease in transgenic mice expressing an activatedmutant of the beta-catenin gene. *Oncogene* 20: 5972–5981, 2001.

[65322] 3652.Sagae, S.; Kobayashi, K.; Nishioka, Y.; Sugimura, M.; Ishioka,S.; Nagata, M.; Terasawa, K.; Tokino, T.; Kudo, R.: Mutational analysisof beta-catenin gene in Japanese ovar-ian carcinomas: frequent mutationsin endometrioid carci-nomas. *Jpn. J. Cancer Res.* 90: 510–515, 1999.

[65323] 3653.Trent, J. M.; Wiltshire, R.; Su, L.-K.; Nicolaides, N. C.; Vogelstein,B.; Kinzler, K. W.: The gene for the APC-binding protein beta-catenin(CTNNB1) maps to chromo-some 3p22, a region frequently altered in humanmalig-nancies. *Cytogenet. Cell Genet.* 71: 343–344, 1995.

[65324] 3654.van Hengel, J.; Nollet, F.; Berx, G.; van Roy, N.; Speleman, F.;van Roy, F.: Assignment of the human beta-catenin gene (CTNNB1) to3p22–p21.3 by fluorescence in situ hybridization. *Cytogenet. CellGenet.* 70: 68–70, 1995.

[65325] 3655.Chan, S. J.; San Segundo, B.; McCormick, M. B.; Steiner, D. F.: Nucleotide and predicted amino acid se-quences of cloned human andmouse preprocathepsin B cDNAs. *Proc. Nat. Acad. Sci.* 83: 7721–7725,1986.

[65326] 3656.Deussing, J.; Roth, W.; Rommerskirch, W.; Wiederan-



ders, B.; von Figura, K.; Peters, C.: The genes of the lysosomal cysteine proteinases cathepsin B, H, L, and S map to different mouse chromosomes. *Mammalian Genome* 8: 241–245, 1997.

[65327] 3657. Shigemitsu, K.; Sekido, Y.; Usami, N.; Mori, S.; Sato, M.; Horio, Y.; Hasegawa, Y.; Bader, S. A.; Gazdar, A. F.; Minna, J. D.; Hida, T.; Yoshioka, H.; Imaizumi, M.; Ueda, Y.; Takahashi, M.; Shimokata, K.: Genetic alteration of the beta-catenin gene (CTNNB1) in human lung cancer and malignant mesothelioma and identification of a new 3p21.3 homozygous deletion. *Oncogene* 20: 4249–4257, 2001.

[65328] 3658. Geck, P.; Sonnenschein, C.; Soto, A. M.: The D13S171 marker, misannotated to BRCA2, links the AS3 gene to various cancers. (Letter) *Am. J. Hum. Genet.* 69: 461–463, 2001.

[65329] 3659. Geck, P.; Szelei, J.; Jimenez, J.; Sonnenschein, C.; Soto, A. M.: Early gene expression during androgen-induced inhibition of proliferation of prostate cancer cells: a new suppressor candidate on chromosome 13, in the BRCA2–Rb1 locus. *J. Steroid Biochem. Molec. Biol.* 68:41–50, 1999.

[65330] 3660. Kuske, M. D. A.; Johnson, J. P.: Assignment of the

human PHLDA1 gene to chromosome 12q15 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 89: 1 only, 2000.

- [65331] 3661. Reddy, P. H.; Stockburger, E.; Gillevet, P.; Tagle, D. A.: Mapping and characterization of novel (CAG)<sub>n</sub> repeat cDNAs from adult human brain derived by the oligo capture method. *Genomics* 46: 174–182, 1997.
- [65332] 3662. Bontekoe, C. J. M.; McIlwain, K. L.; Nieuwenhuizen, I. M.; Yuva-Paylor, L. A.; Nellis, A.; Willemsen, R.; Fang, Z.; Kirkpatrick, L.; Bakker, C. E.; McAninch, R.; Cheng, N. C.; Merriweather, M.; Hoogeveen, A. T.; Nelson, D.; Paylor, R.; Oostra, B. A.: Knockout mouse model for Fxr2: a model for mental retardation. *Hum. Molec. Genet.* 11: 487–498, 2002.
- [65333] 3663. Tamanini, F.; Willemsen, R.; van Unen, L.; Bontekoe, C.; Galjaard, H.; Oostra, B. A.; Hoogeveen, A. T.: Differential expression of FMR1, FXR1 and FXR2 proteins in human brain and testis. *Hum. Molec. Genet.* 6: 1315–1322, 1997.
- [65334] 3664. Zhang, Y.; O'Connor, J. P.; Siomi, M. C.; Srinivasan, S.; Dutra, A.; Nussbaum, R. L.; Dreyfuss, G.: The fragile X mental retardation syndrome protein interacts with novel homologs FXR1 and FXR2. *EMBO J.* 14: 5358–5366, 1995.
- [65335] 3665. Lee, J. K.; Bhakta, S.; Rosen, S. D.; Hemmerich, S.: Cloning and characterization of a mammalian N–

acetylglucosamine-6-sulfotransferase that is highly restricted to intestinal tissue. *Biochem. Biophys. Res. Commun.* 263: 543-549, 1999.

- [65336] 3666. Raymond, C. S.; Parker, E. D.; Kettlewell, J. R.; Brown, L. G.; Page, D. C.; Kusz, K.; Jaruzelska, J.; Reinberg, Y.; Fletjer, W. L.; Bardwell, V. J.; Hirsch, B.; Zarkower, D.: A region of human chromosome 9p required for testis development contains two genes related to known sexual regulators. *Hum. Molec. Genet.* 8: 989-996, 1999.
- [65337] 3667. Colonna, M.; Samaridis, J.: Cloning of immunoglobulin-superfamily members associated with HLA-C and HLA-B recognition by human natural killer cells. *Science* 268: 405-408, 1995.
- [65338] 3668. Lanier, L. L.: Turning on natural killer cells. *J. Exp. Med.* 191: 1259-1262, 2000.
- [65339] 3669. Lanier, L. L.: Natural killer cells: from no receptors to too many. *Immunity* 6: 371-378, 1997.
- [65340] 3670. Lanier, L. L.: Follow the leader: NK cell receptors for classical and nonclassical MHC class I. *Cell* 92: 705-707, 1998.
- [65341] 3671. Ljunggren, H. G.; Karre, K.: In search of the 'missing self': MHC molecules and NK cell recognition. *Immun. Today* 11: 237-244, 1990.

- [65342] 3672.Rajalingam, R.; Krausa, P.; Shilling, H. G.; Stein, J. B.; Balamurugan,A.; McGinnis, M. D.; Cheng, N. W.; Mehra, N. K.; Parham, P.: DistinctiveKIR and HLA diversity in a panel of north Indian Hindus. *Immunogenetics* 53:1009–1019, 2002.
- [65343] 3673.Wagtmann, N.; Biassoni, R.; Cantoni, C.; Verdiani, S.; Malnati,M. S.; Vitale, M.; Bottino, C.; Moretta, L.; Moretta, A.; Long, E.O.: Molecular clones of the p58 NK cell receptor reveal immunoglobulin–relatedmolecules with diversity in both the extra– and intracellular domains. *Immunity* 2:439–449, 1995.
- [65344] 3674.Wagtmann, N.; Rajagopalan, S.; Winter, C. C.; Peruzzi, M.; Long,E. O.: Killer cell inhibitory receptors specific for HLA–C and HLA–Bidentified by direct binding and by functional transfer. *Immunity* 3:801–809, 1995.
- [65345] 3675.Wende, H.; Colonna, M.; Ziegler, A.; Volz, A.: Organization ofthe leukocyte receptor cluster (LRC) on human chromosome 19q13.4. *MammalianGenome* 10: 154–160, 1999.
- [65346] 3676.Dohring, C.; Samaridis, J.; Colonna, M.: Alternatively splicedforms of human killer inhibitory receptors. *Immunogenetics* 44: 227–230,1996.
- [65347] 3677.Matsumoto, N.; Laub, F.; Aldabe, R.; Zhang, W.;

Ramirez, F.; Yoshida,T.; Terada, M.: Cloning the cDNA for a new human zinc finger protein defines a group of closely related Kruppel-like transcription factors. *J.Biol. Chem.* 273: 28229–28237, 1998.

- [65348] 3678.Okazaki, I.; Kinoshita, K.; Muramatsu, M.; Yoshikawa, K.; Honjo,T.: The AID enzyme induces class switch recombination in fibroblasts. *Nature* 416:340–345, 2002.
- [65349] 3679.Petersen–Mahrt, S. K.; Harris, R. S.; Neuberger, M. S.: AID mutates *E. coli* suggesting a DNA deamination mechanism for antibody diversification. *Nature* 418:99–104, 2002.
- [65350] 3680.Revy, P.; Muto, T.; Levy, Y.; Geissmann, F.; Plebani, A.; Sanal,O.; Catalan, N.; Forveille, M.; Dufourcq–Lagelouse, R.; Gennery, A.;Tezcan, I.; Ersoy, F.; and 9 others: Activation–induced cytidine deaminase (AID) deficiency causes the autosomal recessive form of the hyper–IgM syndrome (HIGM2). *Cell* 102: 565–575, 2000.
- [65351] 3681.Yoshikawa, K.; Okazaki, I.; Eto, T.; Kinoshita, K.; Muramatsu,M.; Nagaoka, H.; Honjo, T.: AID enzyme–induced hypermutation in an actively transcribed gene in fibroblasts. *Science* 296: 2033–2036,2002.
- [65352] 3682.Geiszt, M.; Kopp, J. B.; Varnai, P.; Leto, T. L.: Identi–

fication of Renox, an NAD(P)H oxidase in kidney. *Proc. Nat. Acad. Sci.* 97:8010–8014, 2000.

[65353] 3683. Shiose, A.; Kuroda, J.; Tsuruya, K.; Hirai, M.; Hirakata, H.; Naito, S.; Hattori, M.; Sakaki, Y.; Sumimoto, H.: A novel superoxide-producing NAD(P)H oxidase in kidney. *J. Biol. Chem.* 276: 1417–1423, 2001.

[65354] 3684. Kokame, K.; Kato, H.; Miyata, T.: Homocysteine-responsive genes in vascular endothelial cells identified by differential display analysis: GRP78/BiP and novel genes. *J. Biol. Chem.* 271: 29659–29665, 1996.

[65355] 3685. Park, H.; Adams, M. A.; Lachat, P.; Bosman, F.; Pang, S. C.; Graham, C. H.: Hypoxia induces the expression of a 43-kDa protein (PROXY-1) in normal and malignant cells. *Biochem. Biophys. Res. Commun.* 276:321–328, 2000.

[65356] 3686. Shimono, A.; Okuda, T.; Kondoh, H.: N-myc-dependent repression of Ndr1, a gene identified by direct subtraction of whole mouse embryonic DNAs between wild type and N-myc mutant. *Mech. Dev.* 83: 39–52, 1999.

[65357] 3687. van Belzen, N.; Dinjens, W. N. M.; Diesveld, M. P. G.; Groen, N. A.; van der Made, A. C. J.; Nozawa, Y.; Vliestra, R.; Trapman, J.; Bosman, F. T.: A novel gene which is up-regulated during colon epithelial cell differentiation and down-regulated in colorectal neoplasms. *Lab. Invest.* 77:

85–92, 1997.

- [65358] 3688.Zhou, D.; Salnikow, K.; Costa, M.: Cap43, a novel gene specifically induced by Ni<sup>2+</sup> compounds. *Cancer Res.* 58: 2182–2189, 1998.
- [65359] 3689.Janz, R.; Sudhof, T. C.: Cellugyrin, a novel ubiquitous form of synaptogyrin that is phosphorylated by pp60(c-src). *J. Biol. Chem.* 273:2851–2857, 1998.
- [65360] 3690.Ludwig, J.; Kerscher, S.; Brandt, U.; Pfeiffer, K.; Getlawi, F.; Apps, D. K.; Schagger, H.: Identification and characterization of a novel 9.2-kDa membrane sector-associated protein of vacuolar proton-ATPase from chromaffin granules. *J. Biol. Chem.* 273: 10939–10947, 1998.
- [65361] 3691.Nanji, M.; Coronado, V. A.; Cox, D. W.: ATP6H, a subunit of vacuolar ATPase involved in metal transport: evaluation in canine copper toxicosis. *Mammalian Genome* 12: 617–621, 2001.
- [65362] 3692.Arama, E.; Yanai, A.; Kilfin, G.; Bernstein, A.; Motro, B.: Murine NIMA-related kinases are expressed in patterns suggesting distinct functions in gametogenesis and a role in the nervous system. *Oncogene* 16:1813–1823, 1998.
- [65363] 3693.Fry, A. M.; Meraldi, P.; Nigg, E. A.: A centrosomal function for the human Nek2 protein kinase, a member of the NIMA family of cell cycle regulators. *EMBO J.* 17:

470–481, 1998.

- [65364] 3694.Schultz, S. J.; Fry, A. M.; Sutterlin, C.; Ried, T.; Nigg, E. A.: Cell cycle–dependent expression of Nek2, a novel human protein kinase related to the NIMA mitotic regulator of *Aspergillus nidulans*. *Cell Growth Diff.* 5: 625–635, 1994.
- [65365] 3695.Schultz, S. J.; Nigg, E. A.: Identification of 21 novel human protein kinases, including 3 members of a family related to the cell cycle regulator nimA of *Aspergillus nidulans*. *Cell Growth Diff.* 4:821–830, 1993.
- [65366] 3696.Tamari, M.; Daigo, Y.; Nakamura, Y.: Isolation and characterization of a novel serine threonine kinase gene on chromosome 3q22–21.3. *J.Hum. Genet.* 44: 116–120, 1999.
- [65367] 3697.Tamari, M.; Daigo, Y.; Ishikawa, S.; Nakamura, Y.: Genomic structure of a novel human gene (XYLB) on chromosome 3p22–p21.3 encoding a xylulokinase–like protein. *Cytogenet. Cell Genet.* 82: 101–104, 1998.
- [65368] 3698.Migone, T.–S.; Zhang, J.; Luo, X.; Zhuang, L.; Chen, C.; Hu, B.; Hong, J. S.; Perry, J. W.; Chen, S.–F.; Zhou, J. X. H.; Cho, Y. H.; Ullrich, S.; and 14 others: TL1A is a TNF–like ligand for DR3 and TR6/DcR3 and functions as a T cell costimulator. *Immunity* 16: 479–492, 2002.



- [65369] 3699.Zhai, Y.; Ni, J.; Jiang, G.-W.; Lu, J.; Xing, L.; Lincoln, C.;Carter, K. C.; Janat, F.; Kozak, D.; Xu, S.; Rojas, L.; Aggarwal,B. B.; Ruben, S.; Li, L.-Y.; Gentz, R.; Yu, G.-L.: VEGI, a novelcytokine of the tumor necrosis factor family, is an angiogenesis inhibitorthat suppresses the growth of colon carcinomas in vivo. *FASEB J.* 13:181–189, 1999.
- [65370] 3700.Yue, T.-L.; Ni, J.; Romanic, A. M.; Gu, J.-L.; Keller, P.; Wang,C.; Kumar, S.; Yu, G.; Hart, T. K.; Wang, X.; Xia, Z.; DeWolf, W.E., Jr.; Feuerstein, G. Z.: TL1, a novel tumor necrosis factor–likecytokine, induces apoptosis in endothelial cells: involvement of activationof stress protein kinases (stress–activated protein kinase and p38mitogen–activated protein kinase) and caspase–3–like protease. *J.Biol. Chem.* 274: 1479–1486, 1999.
- [65371] 3701.Hertzel, A. V.; Bernlohr, D. A.: Cloning and chromosomal locationof the murine keratinocyte lipid–binding protein gene. *Gene* 221:235–243, 1998.
- [65372] 3702.Madsen, P.; Rasmussen, H. H.; Leffers, H.; Honore, B.; Celis, J.E.: Molecular cloning and expression of a novel keratinocyte protein(psoriasis–associated fatty acid–binding protein [PA–FABP]) that ishighly up–regulated in psoriatic skin and that shares similarity tofatty acid–binding proteins. *J. Invest. Derm.* 99: 299–305, 1992.

- [65373] 3703.Siegenthaler, G.; Hotz, R.; Chatellard–Gruaz, D.; Didierjean, L.;Hellman, U.; Saurat, J. H.: Purification and characterization ofthe human epidermal fatty acid–binding protein: localization duringepidermal cell differentiation in vivo and in vitro. *Biochem J.* 302:363–371, 1994.
- [65374] 3704.Gu, Z.; Flemington, C.; Chittenden, T.; Zambetti, G. P.: Ei24,a p53 response gene involved in growth suppression and apoptosis. *Molec.Cell. Biol.* 20: 233–241, 2000.
- [65375] 3705.Gu, Z.; Gilbert, D. J.; Valentine, V. A.; Jenkins, N. A.; Copeland,N. G.; Zambetti, G. P.: The p53–inducible gene EI24/PIG8 localizesto human chromosome 11q23 and the proximal region of mouse chromosome9. *Cytogenet. Cell Genet.* 89: 230–233, 2000.
- [65376] 3706.Polyak, K.; Xia, Y.; Zweier, J. L.; Kinzler, K. W.; Vogelstein,B.: A model for p53–induced apoptosis. *Nature* 389: 300–305, 1997.
- [65377] 3707.Contente, A.; Dittmer, A.; Koch, M. C.; Roth, J.; Dobbelstein,M.: A polymorphic microsatellite that mediates induction of PIG3by p53. *Nature Genet.* 30: 315–320, 2002.
- [65378] 3708.Hernandez, M.–C.; Andres–Barquin, P. J.; Holt, I.; Israel, M. A.: Cloning of human ENC–1 and evaluation of its

expression and regulation in nervous system tumors. *Exp. Cell Res.* 242: 470–477, 1998.

- [65379] 3709. Hernandez, M.-C.; Andres-Barquin, P. J.; Israel, M. A.: Assignment of the ectodermal-neural cortex 1 gene (Enc1) to mouse chromosome band 13D1 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 89: 158–159, 2000.
- [65380] 3710. Hernandez, M.-C.; Andres-Barquin, P. J.; Kuo, W. L.; Israel, M. A.: Assignment of the ectodermal-neural cortex 1 gene (ENC1) to human chromosome band 5q13 by in situ hybridization. *Cytogenet. Cell Genet.* 87: 89–90, 1999.
- [65381] 3711. Kim, T.-A.; Lim, J.; Ota, S.; Raja, S.; Rogers, R.; Rivnay, B.; Avraham, H.; Avraham, S.: NRP/B, a novel nuclear matrix protein, associates with p110(RB) and is involved in neuronal differentiation. *J. Cell Biol.* 141: 553–566, 1998.
- [65382] 3712. Rosenfeld, M. G.; Mermod, J.-J.; Amara, S. G.; Swanson, L. W.; Sawchenko, P. E.; Rivier, J.; Vale, W. W.; Evans, R. M.: Production of a novel neuropeptide encoded by the calcitonin gene via tissue-specific RNA processing. *Nature* 304: 129–135, 1983.
- [65383] 3713. Saller, B.; Feldmann, G.; Haupt, K.; Broecker, M.; Janssen, O. E.; Roggendorf, M.; Mann, K.; Lu, M.: RT-PCR-based detection of circulating calcitonin-producing

cells in patients with advanced medullary thyroidcancer. J. Clin. Endocr. Metab. 87: 292–296, 2002.

- [65384] 3714.Simpson, N. E.; Goodfellow, P. J.; Riddell, D. C.; Hamerton, J.L.; Holden, J. J. A.; White, B. N.: Assignment of the calcitoningene to chromosome 11 and probable exclusion of linkage between thegene and the locus for multiple endocrine neoplasia type 2. (Abstract) Am.J. Hum. Genet. 36: 153S, 1984.
- [65385] 3715.Struthers, A. D.; Brown, M. J.; Macdonald, D. W. R.; Beacham,J. L.; Stevenson, J. C.; Morris, H. R.; MacIntyre, I.: Human calcitoningene related peptide: a potent endogenous vasodilator in man. Clin.Sci. 70: 389–393, 1986.
- [65386] 3716.Testa, J. R.: Personal Communication. Baltimore, Md. 3/1984.
- [65387] 3717.Tiller–Borcich, J. K.; Capili, H.; Gordan, G. S.: Human braincalcitonin gene–related peptide (CGRP) is concentrated in the locuscaeruleus. Neuropeptides 11: 55–61, 1988.
- [65388] 3718.Tippins, J. R.: CGRP: a novel neuropeptide from the calcitoningene is the most potent vasodilator known. J. Hypertension 4 (suppl.5): S102–S105, 1986.
- [65389] 3719.Tschopp, F. A.; Henke, H.; Petermann, J. B.; Tobler, P. H.; Janzer,R.; Hokfelt, T.; Lundberg, J. M.; Cuello, C.;

Fischer, J. A.: Calcitonin-related peptide and its binding sites in the human central nervous system and pituitary. *Proc. Nat. Acad. Sci.* 82: 248–252, 1985.

[65390] 3720.Christopoulos, G.; Perry, K. J.; Morfis, M.;

Tilakaratne, N.; Gao, Y.; Fraser, N. J.; Main, M. J.; Foord, S. M.; Sexton, P. M.: Multiple amylin receptors arise from receptor activity-modifying protein interaction with the calcitonin receptor gene product. *Molec. Pharm.* 56: 235–242, 1999.

[65391] 3721.de Vernejoul, M.-C.: Personal Communication. Paris, France 1/19/1999.

[65392] 3722.Gorn, A. H.; Lin, H. Y.; Yamin, M.; Auron, P. E.; Flannery, M.R.; Tapp, D. R.; Manning, C. A.; Lodish, H. F.; Krane, S. M.; Goldring, S. R.: Cloning, characterization, and expression of a human calcitonin receptor from an ovarian carcinoma cell line. *J. Clin. Invest.* 90:1726–1735, 1992.

[65393] 3723.Chung, C. D.; Liao, J.; Liu, B.; Rao, X.; Jay, P.; Berta, P.; Shuai, K.: Specific inhibition of Stat3 signal transduction by PIAS3. *Science* 278:1803–1805, 1997.

[65394] 3724.Hagen, F. S.; Grant, F. J.; Kuijper, J. L.; Slaughter, C. A.; Moomaw, C. R.; Orth, K.; O'Hara, P. J.; Munford, R. S.: Expression and characterization of recombinant human acylglycerol hydrolase, a leukocyte enzyme that deacylates

bacterial lipopolysaccharides. *Biochemistry* 30:  
8415–8423, 1991.

- [65395] 3725. Whitmore, T. E.; Mathewes, S. L.; O'Hara, P. J.; Durnam, D. M.: Chromosomal localization of the acyloxyacyl hydrolase (AOAH) gene to 7p14–p12 using fluorescence in situ hybridization. *Genomics* 21:457–458, 1994.
- [65396] 3726. Greenberg, F.; Faucett, A.; Rose, E.; Bancalari, L.; Kardon, N.B.; Mizejewski, G.; Haddow, J. E.; Alpert, E.: Congenital deficiency of alpha-fetoprotein. *Am. J. Obstet. Gynec.* 167: 509–511, 1992.
- [65397] 3727. Greenberg, F.; Rose, E.; Alpert, E.: Hereditary persistence of alpha-fetoprotein. *Gastroenterology* 98: 1083–1085, 1990.
- [65398] 3728. Hammer, R. E.; Krumlauf, R.; Camper, S. A.; Brinster, R. L.; Tilghman, S. M.: Diversity of alpha-fetoprotein gene expression in mice is generated by a combination of separate enhancer elements. *Science* 235:53–58, 1986.
- [65399] 3729. Ingram, R. S.; Scott, R. W.; Tilghman, S. M.: Alpha-fetoprotein and albumin genes are in tandem in the mouse genome. *Proc. Nat. Acad. Sci.* 78: 4694–4698, 1981.
- [65400] 3730. Jagodzinski, L. L.; Sargent, T. D.; Yang, M.; Glackin, C.; Bonner, J.: Sequence homology between RNAs encoding rat alpha-fetoprotein and rat serum albumin. *Proc. Nat.*

Acad. Sci. 78: 3521–3525, 1981.

[65401] 3731.Magenis, R. E.; Luo, X. Y.; Dugaiczky, A.; Ryan, S. C.; Oosterhuis,J. E.: Chromosomal localization of the albumin and alpha-fetoproteingenes in the orangutan (*Pongo pygmaeus*) and gorilla (*Gorilla gorilla*). (Abstract) Cytogenet. Cell Genet. 51: 1037, 1989.

[65402] 3732.McVey, J. H.; Michaelides, K.; Hansen, L. P.; Ferguson-Smith,M.; Tilghman, S.; Krumlauf, R.; Tuddenham, E. G. D.: A G-to-A substitution in an HNF I binding site in the human alpha-fetoprotein gene is associated with hereditary persistence of alpha-fetoprotein (HPAFP). Hum. Molec.Genet. 2: 379–384, 1993.

[65403] 3733.Morinaga, T.; Sakai, M.; Wegmann, T. G.; Tamaoki, T.: Primary structures of human alpha-fetoprotein and its mRNA. Proc. Nat. Acad.Sci. 80: 4604–4608, 1983.

[65404] 3734.Motulsky, A. G.: Personal Communication. Seattle, Wash. 1983.

[65405] 3735.Olsson, M.; Lindahl, G.; Ruoslahti, E.: Genetic control of alpha-fetoproteinsynthesis in the mouse. J. Exp. Med. 145: 819–827, 1977.

[65406] 3736.Rose, E.; Greenberg, F.; Alpert, E.: Hereditary persistence of alpha fetoprotein. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A61, 1989.

- [65407] 3737. Ruoslahti, E.; Terry, W. D.: Alpha fetoprotein and serum albumin show sequence homology. *Nature* 260: 804–805, 1976.
- [65408] 3738. Sakai, M.; Morinaga, T.; Urano, Y.; Watanabe, K.; Wegmann, T.G.; Tamaoki, T.: The human alpha-fetoprotein gene: sequence organization and the 5-prime flanking region. *J. Biol. Chem.* 260: 5055–5060, 1985.
- [65409] 3739. Staples, J.: Alpha fetoprotein, cancer, and benign conditions. (Letter) *Lancet* II: 1277, 1986.
- [65410] 3740. Szpirer, J.; Levan, G.; Thorn, M.; Szpirer, C.: Gene mapping in the rat by mouse–rat somatic cell hybridization: synteny of the albumin and alpha-fetoprotein genes and assignment to chromosome 14. *Cytogenet. Cell Genet.* 38: 142–149, 1984.
- [65411] 3741. Tilghman, S. M.: Personal Communication. Princeton, N. J. 8/12/1992.
- [65412] 3742. Tilghman, S. M.; Belayew, A.: Transcriptional control of the murine albumin/alpha-fetoprotein locus during development. *Proc. Nat. Acad. Sci.* 79: 5254–5257, 1982.
- [65413] 3743. Urano, Y.; Sakai, M.; Watanabe, K.; Tamaoki, T.: Tandem arrangement of the albumin and alpha-fetoprotein genes in the human genome. *Gene* 32: 255–261, 1984.



- [65414] 3744.Vogt, T. F.; Solter, D.; Tilghman, S. M.: Raf, a trans-acting locus, regulates the alpha-fetoprotein gene in a cell-autonomous manner. *Science* 236:301–303, 1987.
- [65415] 3745.Voigtlander, T.; Vogel, F.: Low alpha-fetoprotein and serum albumin levels in Morbus Down may point to a common regulatory mechanism. *Hum.Genet.* 71: 276–277, 1985.
- [65416] 3746.Watanabe, K.; Saito, A.; Tamaoki, T.: Cell-specific enhancer activity in a far upstream region of the human alpha-fetoprotein gene. *J.Biol. Chem.* 262: 4812–4818, 1987.
- [65417] 3747.Abbott, C. A.; Baker, E.; Sutherland, G. R.; McCaughan, G. W.: Genomic organization, exact localization, and tissue expression of the human CD26 (dipeptidyl peptidase IV) gene. *Immunogenetics* 40:331–338, 1994.
- [65418] 3748.Callebaut, C.; Krust, B.; Jacotot, E.; Hovanessian, A. G.: T cell activation antigen, CD26, as a cofactor for entry of HIV in CD4+ cells. *Science* 262:2045–2050, 1993.
- [65419] 3749.Darmoul, D.; Lacasa, M.; Chantret, I.; Swallow, D. M.; Trugnan, G.: Isolation of a cDNA probe for the human intestinal dipeptidylpeptidase IV and assignment of the gene locus DPP4 to chromosome 2. *Ann. Hum.Genet.* 54: 191–197, 1990.

- [65420] 3750.Herbschleb-Voogt, E.; Grzeschik, K.-H.; Pearson, P. L.; Meera Khan,P.: Assignment of adenosine deaminase complexing protein (ADCP) gene(s)to human chromosome 2 in rodent-human somatic cell hybrids. Hum.Genet. 59: 317-323, 1981.
- [65421] 3751.Kameoka, J.; Tanaka, T.; Nojima, Y.; Schlossman, S. F.; Morimoto,C.: Direct association of adenosine deaminase with a T cell activationantigen, CD26. Science 261: 466-469, 1993.
- [65422] 3752.Koch, G. A.; Shows, T. B.: Genes on human chromosomes 2 and 6are required for expression of the adenosine deaminase complexingprotein (ADCP) in human-mouse somatic cell hybrids. (Abstract) Cytogenet.Cell Genet. 25: 174, 1979.
- [65423] 3753.Marguet, D.; Baggio, L.; Kobayashi, T.; Bernard, A.-M.; Pierres,M.; Nielsen, P. F.; Ribel, U.; Watanabe, T.; Drucker, D. J.; Wagtmann,N.: Enhanced insulin secretion and improved glucose tolerance in mice lacking CD26. Proc. Nat. Acad. Sci. 97: 6874-6879, 2000.
- [65424] 3754.Mathew, S.; Morrison, M. E.; Murty, V. V. V. S.; Houghton, A. N.;Chaganti, R. S. K.: Assignment of the DPP4 gene encoding adenosinedeaminase binding protein (CD26/dipeptidylpeptidase IV) to 2q23. Genomics

22:211–212, 1994.

- [65425] 3755.Misumi, Y.; Hayashi, Y.; Arakawa, F.; Ikehara, Y.: Molecularcloning and sequence analysis of human dipeptidyl peptidase IV, aserine proteinase on the cell surface. Biochim. Biophys. Acta 1131:333–336, 1992.
- [65426] 3756.Morrison, M. E.; Vijayasaradhi, S.; Engelstein, D.; Albino, A.P.; Houghton, A. N.: A marker for neoplastic progression of humanmelanocytes is a cell surface ectopeptidase. J. Exp. Med. 117: 1135–1143,1993.
- [65427] 3757.Van Cong, N.; Weil, D.; Gross, M.–S.; Foubert, C.; Jami, J.; Frezal,J.: Controle genetique et epigenetique de l'expression de l'adenosinedeaminase. Analyse des cellules humaines et hybrides homme–rongeur. Ann.Genet. 24: 141–147, 1981.
- [65428] 3758.Boison, D.; Scheurer, L.; Zumsteg, V.; Rulicke, T.; Litynski, P.;Fowler, B.; Brandner, S.; Mohler, H.: Neonatal hepatic steatosisby disruption of the adenosine kinase gene. Proc. Nat. Acad. Sci. 99:6985–6990, 2002.
- [65429] 3759.Chan, T.–S.; Cregan, R. P.; Reardon, M. P.: Adenosine kinase asa new selective marker in somatic cell genetics: isolation of adenosinekinase–deficient mouse cell lines and human–mouse hybrid cell linescontaining adenosine kinase. Somat. Cell Genet. 4: 1–12, 1978.

- [65430] 3760.Francke, U.; Thompson, L.: Regional mapping, by exclusion, of adenosine kinase (ADK) on human chromosome 10 using the gene dosage approach. (Abstract) Cytogenet. Cell Genet. 25: 156, 1979.
- [65431] 3761.Bundey, S.: Recent views on genetic factors in retinoblastoma.(Abstract) J. Med. Genet. 17: 386–387, 1980.
- [65432] 3762.Bundey, S.; Morten, J. E. N.: An unusual pedigree with retinoblastoma.Does it shed light on the delayed mutation and host resistance theories? Hum.Genet. 59: 434–436, 1981.
- [65433] 3763.Cance, W. G.; Brennan, M. F.; Dudas, M. E.; Huang, C.-M.; Cordon-Cardo,C.: Altered expression of the retinoblastoma gene product in human sarcomas. New Eng. J. Med. 323: 1457–1462, 1990.
- [65434] 3764.Canning, S.; Dryja, T. P.: Short, direct repeats at the breakpointsof deletions of the retinoblastoma gene. Proc. Nat. Acad. Sci. 86:5044–5048, 1989.
- [65435] 3765.Carlson, E. A.; Desnick, R. J.: Mutational mosaicism and genetic counseling in retinoblastoma. Am. J. Med. Genet. 4: 365–381, 1979.
- [65436] 3766.Cavenee, W. K.: The genetic basis of neoplasia: the retinoblastoma paradigm. Trends Genet. 2: 299–300,

1986.

- [65437] 3767.Cavenee, W. K.; Dryja, T. P.; Phillips, R. A.; Benedict, W. F.; Godbout, R.; Gallie, B. L.; Murphree, A. L.; Strong, L. C.; White, R. L.: Expression of recessive alleles by chromosomal mechanisms in retinoblastoma. *Nature* 305: 779–784, 1983.
- [65438] 3768.Cavenee, W. K.; Hansen, M. F.; Nordenskjold, M.; Kock, E.; Maumenee, I.; Squire, J. A.; Phillips, R. A.; Gallie, B. L.: Genetic origin of mutations predisposing to retinoblastoma. *Science* 228: 501–503, 1985.
- [65439] 3769.Cavenee, W. K.; Murphree, A. L.; Shull, M. M.; Benedict, W. F.; Sparkes, R. S.; Kock, E.; Nordenskjold, M.: Prediction of familial predisposition to retinoblastoma. *New Eng. J. Med.* 314: 1201–1207, 1986.
- [65440] 3770.Chano, T.; Ikegawa, S.; Kontani, K.; Okabe, H.; Baldini, N.; Saeki, Y.: Identification of RB1CC1, a novel human gene that can induce RB1 in various human cells. *Oncogene* 21: 1295–1298, 2002.
- [65441] 3771.Chauveinc, L.; Mosseri, V.; Quintana, E.; Desjardins, L.; Schlienger, P.; Doz, F.; Dutrillaux, B.: Osteosarcoma following retinoblastoma: age at onset and latency period. *Ophthalmic Genet.* 22: 77–88, 2001.
- [65442] 3772.Chen, P.-L.; Scully, P.; Shew, J.-Y.; Wang, J. Y. J.; Lee,

W.-H.: Phosphorylation of the retinoblastoma gene product is modulated during the cell cycle and cellular differentiation. *Cell* 58: 1193–1198, 1989.

[65443] 3773. Connolly, M. J.; Payne, R. H.; Johnson, G.; Gallie, B. L.; Alderdice, P. W.; Marshall, W. H.; Lawton, R. D.: Familial, EsD-linked, retinoblastoma with reduced penetrance and variable expressivity. *Hum. Genet.* 65:122–124, 1983.

[65444] 3774. Cowell, J. K.; Bia, B.: A novel missense mutation in patients from a retinoblastoma pedigree showing only mild expression of the tumor phenotype. *Oncogene* 16: 3211–3213, 1998.

[65445] 3775. Cowell, J. K.; Rutland, P.; Hungerford, J.; Jay, M.: Deletion of chromosome region 13q14 is transmissible and does not always predispose to retinoblastoma. *Hum. Genet.* 80: 43–45, 1988.

[65446] 3776. Cowell, J. K.; Rutland, P.; Jay, M.; Hungerford, J.: Deletion of the esterase D locus from a survey of 200 retinoblastoma patients. *Hum. Genet.* 72: 164–167, 1986.

[65447] 3777. Cowell, J. K.; Smith, T.; Bia, B.: Frequent constitutional C to T mutations in CGA-arginine codons in the RB1 gene produce premature stop codons in patients with bilateral (hereditary) retinoblastoma. *Europ. J. Hum. Genet.* 2: 281–290, 1994.

- [65448] 3778.Dahiya, A.; Wong, S.; Gonzalo, S.; Gavin, M.; Dean, D. C.: Linking the Rb and Polycomb pathways. *Molec. Cell* 8: 557–568, 2001.
- [65449] 3779.Davison, E. V.; Gibbons, B.; Aherne, G. E. S.; Roberts, D. F.: Chromosomes in retinoblastoma patients. *Clin. Genet.* 15: 505–508, 1979.
- [65450] 3780.DeCaprio, J. A.; Ludlow, J. W.; Lynch, D.; Furukawa, Y.; Griffin, J.; Piwnica-Worms, H.; Huang, C.-M.; Livingston, D. M.: The product of the retinoblastoma susceptibility gene has properties of a cell cycle regulatory element. *Cell* 58: 1085–1095, 1989.
- [65451] 3781.de Grouchy, J.; Turleau, C.; Cabanis, M. O.; Richardet, J. M.: Retinoblastome et deletion intercalaire du chromosome 13. *Arch. Franc. Pediat.* 37: 531–535, 1980.
- [65452] 3782.Dryja, T.; Cavenee, W.; Epstein, J.; Rapaport, J.; Goorin, A.; Koufos, A.: Chromosome 13 homozygosity in osteogenic sarcoma without retinoblastoma. (Abstract) *Am. J. Hum. Genet.* 36: 28S, 1984.
- [65453] 3783.Dryja, T. P.; Bruns, G. A. P.; Gallie, B.; Petersen, R.; Green, W.; Rapaport, J. M.; Albert, D. M.; Gerald, P. S.: Low incidence of deletion of the esterase D locus in retinoblastoma patients. *Hum. Genet.* 64: 151–155, 1983.
- [65454] 3784.Dryja, T. P.; Cavenee, W.; White, R.; Rapaport, J. M.;

Petersen,R.; Albert, D. M.; Bruns, G. A. P.: Homozygosity of chromosome 13in retinoblastoma. New Eng. J. Med. 310: 550–553, 1984.

[65455] 3785.Dryja, T. P.; Friend, S.; Weinberg, R. A.: Isolation of a cDNAfragment derived from human retina mRNA which detects a locus within13q14 often deleted in retinoblastomas. (Abstract) Am. J. Hum. Genet. 39:A29, 1986.

[65456] 3786.Dryja, T. P.; Mukai, S.; Petersen, R.; Rapaport, J. M.; Walton,D.; Yandell, D. W.: Parental origin of mutations of the retinoblastomagene. Nature 339: 556–558, 1989.

[65457] 3787.Dryja, T. P.; Mukai, S.; Rapaport, J. M.; Yandell, D. W.: Parentalorigin of mutations of the retinoblastoma gene. (Abstract) Am. J.Hum. Genet. 45 (suppl.): A19, 1989.

[65458] 3788.Messer Peters, P. G.; Kamarck, M. E.; Hemler, M. E.; Strominger,J. L.; Ruddle, F. H.: Genetic and biochemical characterization ofa human surface determinant on somatic cell hybrids: the 4F2 antigen. Somat.Cell Genet. 8: 825–834, 1982.

[65459] 3789.Markert, C. L.; Silvers, W. K.: The effects of genotype and cellenvironment on melanoblast differentiation in the house mouse. Genetics 41:429–450, 1956.

[65460] 3790.Van Cong, N.; Moreau–Gachelin, F.; Ray, D.; Gross,



M. S.; de Tand, M. F.; Tavitian, A.; Frezal, J.: Assignment of SPI1 oncogene to chromosome 11 (somatic cell hybrid analysis), region p11.22 (in situ hybridization). (Abstract) Cytogenet. Cell Genet. 51: 1097 only, 1989.

[65461] 3791. Bell, L. R.; Maine, E. M.; Schedl, P.; Cline, T. W.: Sex-lethal, a *Drosophila* sex determination switch gene, exhibits sex-specific RNA splicing and sequence similarity to RNA binding proteins. *Cell* 55: 1037–1046, 1988.

[65462] 3792. Fletcher, C. F.; Okano, H. J.; Gilbert, D. J.; Yang, Y.; Yang, C.; Copeland, N. G.; Jenkins, N. A.; Darnell, R. B.: Mouse chromosomal locations of nine genes encoding homologs of human paraneoplastic neurologic disorder antigens. *Genomics* 45: 313–319, 1997.

[65463] 3793. King, P. H.; Levine, T. D.; Fremeau, R. T., Jr.; Keene, J. D.: Mammalian homologs of *Drosophila* elav localized to a neuronal subset can bind in vitro to the 3-prime UTR of mRNA encoding the *Id* transcriptional repressor. *J. Neurosci.* 14: 1943–1952, 1994.

[65464] 3794. Levine, T. D.; Gao, F.; King, P. H.; Andrews, L. G.; Keene, J. D.: He1-N1: an autoimmune RNA-binding protein with specificity for 3-prime uridylate-rich untranslated regions of growth factor mRNAs. *Molec. Cell. Biol.* 13: 3494–3504, 1993.

- [65465] 3795.Muresu, R.; Baldini, A.; Gress, T.; Posner, J. B.; Furneaux, H.M.; Siniscalco, M.: Mapping of the gene coding for a paraneoplasticencephalomyelitis antigen (HuD) to human chromosome site 1p34. Cytogenet.Cell Genet. 65: 177–178, 1994.
- [65466] 3796.Robinow, S.; Campos, A. R.; Yao, K.–M.; White, K.: The elav gene product of Drosophila, required in neurons, has three RNP consensus motifs. Science 242: 1570–1572, 1988.
- [65467] 3797.Szabo, A.; Dalmau, J.; Manley, G.; Rosenfeld, M.; Wong, E.; Henson, J.; Posner, J. B.; Furneaux, H. M.: HuD, a paraneoplastic encephalomyelitis antigen, contains RNA-binding domains and is homologous to elav and sex-lethal. Cell 67: 325–333, 1991.
- [65468] 3798.Clinton, M.; Frangou–Lazaridis, M.; Panneerselvam, C.; Horecker, B. L.: The sequence of human parathymosin deduced from a cloned human kidney cDNA. Biochem. Biophys. Res. Commun. 158: 855–862, 1989.
- [65469] 3799.Szabo, P.; Clinton, M.; Macera, M.; Horecker, B. L.: Localization of the gene coding for parathymosin to chromosome 17 in humans. Cytogenet.Cell Genet. 50: 91–92, 1989.
- [65470] 3800.Arnold, A.; Kim, H. G.; Gaz, R. D.; Eddy, R. L.;

Fukushima, Y.; Byers, M. G.; Shows, T. B.; Kronenberg, H. M.: Molecular cloning and chromosomal mapping of DNA rearranged with the parathyroid hormone gene in a parathyroid adenoma. *J. Clin. Invest.* 83: 2034–2040, 1989.

[65471] 3801. Mehta, A. D.; Rock, R. S.; Rief, M.; Spudich, J. A.; Mooseker, M. S.; Cheney, R. E.: Myosin-V is a processive actin-based motor. *Nature* 400:590–593, 1999.

[65472] 3802. Mercer, J. A.; Seperack, P. K.; Strobel, M. C.; Copeland, N. G.; Jenkins, N. A.: Novel myosin heavy chain encoded by murine dilute coat colour locus. *Nature* 349: 709–713, 1991.

[65473] 3803. Moore, K. J.; Testa, J. R.; Francke, U.; Milatovich, A.; Copeland, N. G.; Jenkins, N. A.: Cloning and regional assignment of the human myosin heavy chain 12 (MYH12) gene to chromosome band 15q21. *Cytogenet. Cell Genet.* 69: 53–58, 1995.

[65474] 3804. Pastural, E.; Barrat, F. J.; Dufourcq-Lagelouse, R.; Certain, S.; Sanal, O.; Jabado, N.; Seger, R.; Griscelli, C.; Fischer, A.; de Saint Basile, G.: Griscelli disease maps to chromosome 15q21 and is associated with mutations in the myosin-Va gene. *Nature Genet.* 16:289–292, 1997.  
Note: Erratum: *Nature Genet.* 23: 373 only, 1999.

- [65475] 3805.Rief, M.; Rock, R. S.; Mehta, A. D.; Mooseker, M. S.; Cheney,R. E.; Spudich, J. A.: Myosin-V stepping kinetics: a molecular model for processivity. *Proc. Nat. Acad. Sci.* 97: 9482–9486, 2000.
- [65476] 3806.Russell, E. S.: A quantitative histological study of the pigment found in the coat-color mutants of the house mouse. IV. The nature of the effects of genic substitution in five major allelic series. *Genetics* 34:146–166, 1949.
- [65477] 3807.Musahl, C.; Schulte, D.; Burkhart, R.; Knippers, R.: A human homologue of the yeast replication protein Cdc21: interactions with other Mcm proteins. *Europ. J. Biochem.* 230: 1096–1101, 1995.
- [65478] 3808.Kim, N.-S.; Kato, T.; Abe, N.; Kato, S.: Nucleotide sequence of human cDNA encoding eukaryotic initiation factor 4A1. *Nucleic Acids Res.* 21: 2012 only, 1993.
- [65479] 3809.Kukimoto, I.; Watanabe, S.; Taniguchi, K.; Ogata, T.; Yoshiike, K.; Kanda, T.: Characterization of the cloned promoter of the human initiation factor 4A1 gene. *Biochem. Biophys. Res. Commun.* 233:844–847, 1997.
- [65480] 3810.Nielsen, P. J.; McMaster, G. K.; Trachsel, H.: Cloning of eukaryotic protein synthesis initiation factor genes: isolation and characterization of cDNA clones encoding factor eIF-4A. *Nucleic Acids Res.* 13: 6867–6880, 1985.

- [65481] 3811. Anderson, D. M.; Maraskovsky, E.; Billingsley, W. L.; Dougall, W. C.; Tometsko, M. E.; Roux, E. R.; Teepe, M. C.; DuBose, R. F.; Cosman, D.; Galibert, L.: A homologue of the TNF receptor and its ligand enhance T-cell growth and dendritic-cell function. *Nature* 390:175–179, 1997.
- [65482] 3812. Croucher, P. I.; Shipman, C. M.; Lippitt, J.; Perry, M.; Asosingh, K.; Hijzen, A.; Brabbs, A. C.; van Beek, E. J. R.; Holen, I.; Skerry, T. M.; Dunstan, C. R.; Russell, G. R.; Van Camp, B.; Vanderkerken, K.: Osteoprotegerin inhibits the development of osteolytic bone disease in multiple myeloma. *Blood* 98: 3534–3540, 2001.
- [65483] 3813. Fata, J. E.; Kong, Y.-Y.; Li, J.; Sasaki, T.; Irie-Sasaki, J.; Moorehead, R. A.; Elliott, R.; Scully, S.; Voura, E. B.; Lacey, D. L.; Boyle, W. J.; Khokha, R.; Penninger, J. M.: The osteoclast differentiation factor osteoprotegerin–ligand is essential for mammary gland development. *Cell* 103:41–50, 2000.
- [65484] 3814. Kim, N.; Odgren, P. R.; Kim, D.-K.; Marks, S. C., Jr.; Choi, Y.: Diverse roles of the tumor necrosis factor family member TRANCE in skeletal physiology revealed by TRANCE deficiency and partial rescue by a lymphocyte-expressed TRANCE transgene. *Proc. Nat. Acad. Sci.* 97:10905–10910, 2000.

- [65485] 3815.Kong, Y.-Y.; Feige, U.; Sarosi, I.; Bolon, B.; Tafuri, A.; Morony,S.; Capparelli, C.; Li, J.; Elliott, R.; McCabe, S.; Wong, T.; Campagnuolo,G.; and 9 others: Activated T cells regulate bone loss and jointdestruction in adjuvant arthri-  
tis through osteoprotegerin ligand. *Nature* 402:304–309, 1999.
- [65486] 3816.Lacey, D. L.; Timms, E.; Tan, H.-L.; Kelley, M. J.; Dunstan, C.R.; Burgess, T.; Elliott, R.; Colombero, A.; Eli-  
liott, G.; Scully,S.; Hsu, H.; Sullivan, J.; Hawkins, N.; Davy, E.; Capparelli, C.;Eli, A.; Qian, Y.-X.; Kaufman, S.; Sarosi, I.; Shalhoub, V.; Senaldi,G.; Guo, J.; Delaney, J.; Boyle, W. J.: Osteoprotegerin ligand isa cytokine that regulates os-  
teoclast differentiation and activation. *Cell* 93:165–176, 1998.
- [65487] 3817.Pearse, R. N.; Sordillo, E. M.; Yaccoby, S.; Wong, B. R.; Liau,D. F.; Colman, N.; Michaeli, J.; Epstein, J.; Choi, Y.: Multiplemyeloma disrupts the TRANCE/osteoprotegerin  
cytokine axis to triggerbone destruction and promote tu-  
mor progression. *Proc. Nat. Acad.Sci.* 98: 11581–11586, 2001.
- [65488] 3818.Wong, B. R.; Rho, J.; Arron, J.; Robinson, E.; Orlinick, J.; Chao,M.; Kalachikov, S.; Cayani, E.; Bartlett, F. S., III; Frankel, W.N.; Lee, S. Y.; Choi, Y.: TRANCE is a novel lig-

and of the tumor necrosis factor receptor family that activates c-Jun N-terminal kinase in T cells. *J. Biol. Chem.* 272: 25190–25194, 1997.

- [65489] 3819. Udell, C. M.; Lee, S. K.; Davey, S.: HRAD1 and MRAD1 encode mammalian homologues of the fission yeast rad1+ cell cycle checkpoint control gene. *Nucleic Acids Res.* 26: 3971–3976, 1998.
- [65490] 3820. Carvajal, J. J.; Pook, M. A.; dos Santos, M.; Doudney, K.; Hillermann, R.; Minogue, S.; Williamson, R.; Hsuan, J. J.; Chamberlain, S.: The Friedreich's ataxia gene encodes a novel phosphatidylinositol-4-phosphate 5-kinase. *Nature Genet.* 14: 157–162, 1996.
- [65491] 3821. Carvajal, J. J.; Pook, M. A.; Doudney, K.; Hillermann, R.; Wilkes, D.; Al-Mahdawi, S.; Williamson, R.; Chamberlain, S.: Friedreich's ataxia: a defect in signal transduction? *Hum. Molec. Genet.* 4: 1411–1419, 1995.
- [65492] 3822. Pook, M. A.; Carvajal, J. J.; Doudney, K.; Hillermann, R.; Chamberlain, S.: Exon-intron structure of a 2.7-kb transcript of the STM7 gene with phosphatidylinositol-4-phosphate 5-kinase activity. *Genomics* 42: 170–172, 1997.
- [65493] 3823. Carfi, A.; Willis, S. H.; Whitbeck, J. C.; Krummenacher, C.; Cohen, G. H.; Eisenberg, R. J.; Wiley, D. C.: Her-

pes simplex virus glycoproteinD bound to the human receptor HveA. *Molec. Cell* 8: 169–179, 2001.

[65494] 3824.Hsu, H.; Solovyev, I.; Colombero, A.; Elliott, R.; Kelley, M.;Boyle, W. J.: ATAR, a novel tumor necrosis factor receptor familymember, signals through TRAF2 and TRAF5. *J. Biol. Chem.* 272: 13471–13474,1997.

[65495] 3825.Marsters, S. A.; Ayres, T. M.; Skubatch, M.; Gray, C. L.; Rothe,M.; Ashkenazi, A.: Herpesvirus entry mediator, a member of the tumornecrosis factor receptor (TNFR) family, interacts with members ofthe TNFR-associated factor family and activates the transcriptionfactors NF-kappa-B and AP-1. *J. Biol. Chem.* 272: 14029–14032, 1997.

[65496] 3826.Montgomery, R. I.; Warner, M. S.; Lum, B. J.; Spear, P. G.: Herpes simplex virus-1 entry into cells mediated by a novel member of theTNF/NGF receptor family. *Cell* 87: 427–436, 1996.

[65497] 3827.Kwon, B. S.; Tan, K. B.; Ni, J.; Oh, K.-O.; Lee, Z. H.; Kim, K.K.; Kim, Y.-J.; Wang, S.; Gentz, R.; Yu, G.-L.; Harrop, J.; Lyn, S.D.; Silverman, C.; Porter, T. G.; Truneh, A.; Young, P. R.: A newlyidentified member of the tumor necrosis factor receptor superfamilywith a wide tissue distribution and involvement in lymphocyte activation. *J.Biol. Chem.* 272: 14272–14276, 1997.



- [65498] 3828. Morel, Y.; Schiano de Colella, J.-M.; Harrop, J.; Deen, K. C.; Holmes, S. D.; Wattam, T. A.; Khandekar, S. S.; Truneh, A.; Sweet, R. W.; Gastaut, J.-A.; Olive, D.; Costello, R. T.: Reciprocal expression of the TNF family receptor herpes virus entry mediator and its ligand LIGHT on activated T cells: LIGHT down-regulates its own receptor. *J. Immun.* 165: 4397-4404, 2000.
- [65499] 3829. Guan, K.-L.; Butch, E.: Isolation and characterization of a novel dual specific phosphatase, HVH2, which selectively dephosphorylates the mitogen-activated protein kinase. *J. Biol. Chem.* 270: 7197-7203, 1995.
- [65500] 3830. Smith, A.; Price, C.; Cullen, M.; Muda, M.; King, A.; Ozanne, B.; Arkinstall, S.; Ashworth, A.: Chromosomal localization of three human dual specificity phosphatase genes (DUSP4, DUSP6, and DUSP7). *Genomics* 42: 524-527, 1997.
- [65501] 3831. Furukawa, T.; Yatsuoka, T.; Youssef, E. M.; Abe, T.; Yokoyama, T.; Fukushige, S.; Soeda, E.; Hoshi, M.; Hayashi, Y.; Sunamura, M.; Kobari, M.; Horii, A.: Genomic analysis of DUSP6, a dual specificity MAP kinase phosphatase, in pancreatic cancer. *Cytogenet. Cell Genet.* 82: 156-159, 1998.
- [65502] 3832. Groom, L. A.; Sneddon, A. A.; Alessi, D. R.; Dowd, S.; Keyse, S. M.: Differential regulation of the MAP, SAP and

RK/p38 kinases by Pyst1, a novel cytosolic dual-specificity phosphatase. *EMBO J.* 15:3621–3632, 1996.

[65503] 3833. Muda, M.; Boschert, U.; Dickinson, R.; Martinou, J.-C.; Martinou, I.; Camps, M.; Schlegel, W.; Arkinstall, S.: MKP-3, a novel cytosolic protein-tyrosine phosphatase that exemplifies a new class of mitogen-activated protein kinase phosphatase. *J. Biol. Chem.* 271: 4319–4326, 1996.

[65504] 3834. Smith, A.; Price, C.; Cullen, M.; Muda, M.; King, A.; Ozanne, B.; Arkinstall, S.; Ahsworth, A.: Chromosomal localization of three human dual specificity phosphatase genes (DUSP4, DUSP6, and DUSP7). *Genomics* 42:524–527, 1997.

[65505] 3835. Nishihira, J.; Fujinaga, M.; Kuriyama, T.; Suzuki, M.; Sugimoto, H.; Nakagawa, A.; Tanaka, I.; Sakai, M.: Molecular cloning of human D-dopachrome tautomerase cDNA: N-terminal proline is essential for enzyme activation. *Biochem. Biophys. Res. Commun.* 243: 538–544, 1998.

[65506] 3836. Blouin, J.-L.; Sail, G. D.; Guipponi, M.; Rossier, C.; Pappasavas, M.-P.; Antonarakis, S. E.: Isolation of the human BACH1 transcription regulator gene, which maps to chromosome 21q22.1. *Hum. Genet.* 102:282–288, 1998.

[65507] 3837. Conley, Y. P.; Erturk, D.; Keverline, A.; Mah, T. S.;

Keravala,A.; Barnes, L. R.; Bruchis, A.; Hess, J. F.; FitzGerald, P. G.; Weeks,D. E.; Ferrell, R. E.; Gorin, M. B.: A juvenile-onset, progressive cataract locus on chromosome 3q21-q22 is associated with a missense mutation in the beaded filament structural protein-2. *Am. J. Hum. Genet.* 66: 1426-1431, 2000.

[65508] 3838. Strehl, S.; Glatt, K.; Liu, Q. M.; Glatt, H.; Lalande, M.: Characterization of two novel protocadherins (PCDH8 and PCDH9) localized on human chromosome 13 and mouse chromosome 14. *Genomics* 53: 81-89, 1998.

[65509] 3839. Ohbayashi, N.; Hoshikawa, M.; Kimura, S.; Yamasaki, M.; Fukui, S.; Itoh, N.: Structure and expression of the mRNA encoding a novel fibroblast growth factor, FGF-18. *J. Biol. Chem.* 273: 18161-18164, 1998.

[65510] 3840. Loeffen, J.; Smeets, R.; Smeitink, J.; Triepels, R.; Sengers, R.; Trijbels, F.; van den Heuvel, L.: The human NADH:ubiquinone oxidoreductase NDUF5 (15 kDa) subunit: cDNA cloning, chromosomal localization, tissue distribution and the absence of mutations in isolated complex I-deficient patients. *J. Inherit. Metab. Dis.* 22: 19-28, 1999.

[65511] 3841. Jackson, A.; Panayiotidis, P.; Foroni, L.: The human homologue of the *Drosophila* tailless gene (TLX): charac-

terization and mapping to a region of common deletion in human lymphoid leukemia on chromosome 6q21. *Genomics* 50: 34–43, 1998.

- [65512] 3842. Monaghan, A. P.; Bock, D.; Gass, P.; Schwager, A.; Wolfer, D. P.; Lipp, H.-P.; Schutz, G.: Defective limbic system in mice lacking the *tailless* gene. *Nature* 390: 515–517, 1997.
- [65513] 3843. Yu, R. T.; McKeown, M.; Evans, R. M.; Umesono, K.: Relationship between *Drosophila* gap gene *tailless* and a vertebrate nuclear receptor *Tlx*. *Nature* 370: 375–379, 1994.
- [65514] 3844. Plougastel, B.; Trowsdale, J.: Cloning of NKG2–F, a new member of the NKG2 family of human natural killer cell receptor genes. *Europ. J. Immun.* 27: 2835–2839, 1997.
- [65515] 3845. Sutherland, C. L.; Chalupny, N. J.; Schooley, K.; Van den Bos, T.; Kubin, M.; Cosman, D.: UL16–binding proteins, novel MHC class I–related proteins, bind to NKG2D and activate multiple signaling pathways in primary NK cells. *J. Immun.* 168: 671–679, 2002.
- [65516] 3846. Wu, J.; Song, Y.; Bakker, A. B. H.; Bauer, S.; Spies, T.; Lanier, L. L.; Phillips, J. H.: An activating immunoreceptor complex formed by NKG2D and DAP10. *Science* 285:

730–732, 1999.

- [65517] 3847.Chang, C.; Rodriguez, A.; Carretero, M.; Lopez–Botet, M.; Phillips,J. H.; Lanier, L. L.: Molecular characteri–zation of human CD94: atype II membrane glycoprotein related to the C–type lectin superfamily. *Europ.J. Immun.* 25: 2433–2437, 1995.
- [65518] 3848.Lazetic, S.; Chang, C.; Houchins, J. P.; Lanier, L. L.; Phillips,J. H.: Human natural killer cell receptors involved in MHC classI recognition are disulfide–linked het–erodimers of CD94 and NKG2 subunits. *J.Immun.* 157: 4741–4745, 1996.
- [65519] 3849.Rodriguez, A.; Carretero, M.; Glienke, J.; Bellon, T.; Ramirez,A.; Lehrach, H.; Francis, F.; Lopez–Botet, M.: Structure of the humanCD94 C–type lectin gene. *Immuno–genetics* 47: 305–309, 1998.
- [65520] 3850.Volkmer, E.; Karnitz, L. M.: Human homologs of *Schizosaccharomycespombe* Rad1, Hus1, and Rad9 form a DNA damage–responsive protein complex. *J.Biol. Chem.* 274: 567–570, 1999.
- [65521] 3851.Brandner, J. M.; Reidenbach, S.; Franke, W. W.: Evi–dence that'pinin,' reportedly a differentiation–specific desmosomal protein,is actually a widespread nuclear pro–tein. *Differentiation* 62: 119–127,1997.

- [65522] 3852.Brandner, J. M.; Reidenbach, S.; Kuhn, C.; Franke, W. W.: Identification and characterization of a novel kind of nuclear protein occurring free in nucleoplasm and in ribonucleoprotein structures of the 'speckle' type. *Europ. J. Cell Biol.* 75: 295–308, 1998.
- [65523] 3853.Degen, W. G. J.; Agterbos, M. A.; Muylers, J. P. P.; Bloemers, H. P. J.; Swart, G. W. M.: memA/DRS, a putative mediator of multiprotein complexes, is overexpressed in the metastasizing human melanoma cell lines BLM and MV3. *Biochim. Biophys. Acta* 1444: 384–394, 1999.
- [65524] 3854.Ouyang, P.; Sugrue, S. P.: Characterization of pinin, a novel protein associated with the desmosome–intermediate filament complex. *J. Cell Biol.* 135: 1027–1042, 1996.
- [65525] 3855.Puente, X. S.; Lopez–Otin, C.: Cloning and expression analysis of a novel human serine hydrolase with sequence similarity to prokaryotic enzymes involved in the degradation of aromatic compounds. *J. Biol. Chem.* 270: 12926–12932, 1995.
- [65526] 3856.Puente, X. S.; Pendas, A. M.; Lopez–Otin, C.: Structural characterization and chromosomal localization of the gene encoding human biphenyl hydrolase–related protein (BPHL). *Genomics* 51: 459–462, 1998.

- [65527] 3857.Janssen, J. W. G.; Schleithoff, L.; Bartram, C. R.; Schulz, A.S.: An oncogenic fusion product of the phosphatidylinositol 3-kinasep85-beta subunit and HUMORF8, a putative deubiquitinating enzyme. *Oncogene* 16:1767-1772, 1998.
- [65528] 3858.Chadwick, B. P.; Williamson, J.; Sheer, D.; Frischauf, A.-M.:cDNA cloning and chromosomal mapping of a mouse gene with homologyto NTPases. *Mammalian Genome* 9: 162-164, 1998.
- [65529] 3859.Yeung, G.; Mulero, J. J.; McGowan, D. W.; Bajwa, S. S.; Ford, J.E.: CD39L2, a gene encoding a human nucleoside diphosphatase, predominantlyexpressed in the heart. *Biochemistry* 39: 12916-12923, 2000.
- [65530] 3860.Ji, H.; Liu, Y. E.; Jia, T.; Wang, M.; Liu, J.; Xiao, G.; Joseph,B. K.; Rosen, C.; Shi, Y. E.: Identification of a breast cancer-specificgene, BCSG1, by direct differential cDNA sequencing. *Cancer Res.* 57:759-764, 1997.
- [65531] 3861.Lavedan, C.; Leroy, E.; Dehejia, A.; Buchholtz, S.; Dutra, A.;Nussbaum, R. L.; Polymeropoulos, M. H.: Identification, localizationand characterization of the human gamma-synuclein gene. *Hum. Genet.* 103:106-112, 1998.
- [65532] 3862.Ninkina, N. N.; Alimova-Kost, M. V.; Paterson, J. W. E.; Delaney,L.; Cohen, B. B.; Imreh, S.; Gnuchev, N. V.;

Davies, A. M.; Buchman, V. L.: Organization, expression and polymorphism of the human persyngene. Hum. Molec. Genet. 7: 1417–1424, 1998.

[65533] 3863. Spillantini, M. G.; Crowther, R. A.; Jakes, R.; Hasegawa, M.; Goedert, M.: alpha-synuclein in filamentous inclusions of Lewy bodies from Parkinson's disease and dementia with Lewy bodies. Proc. Nat. Acad. Sci. 95: 6469–6473, 1998.

[65534] 3864. Seroussi, E.; Pan, H.-Q.; Kedra, D.; Roe, B. A.; Duman, J. P.: Characterization of the human NIPSNAP1 gene from 22q12: a member of a novel gene family. Gene 212: 13–20, 1998.

[65535] 3865. Wang, X.-Y.; Smith, D. I.; Lui, W.; James, C. D.: GBAS, a novel gene encoding a protein with tyrosine phosphorylation sites and a transmembrane domain, is co-amplified with EGFR. Genomics 49: 448–451, 1998.

[65536] 3866. Kurima, K.; Warman, M. L.; Krishnan, S.; Domowicz, M.; Krueger, R. C., Jr.; Deyrup, A.; Schwartz, N. B.: A member of a family of sulfate-activating enzymes causes murine brachymorphism. Proc. Nat. Acad. Sci. 95: 8681–8685, 1998.

[65537] 3867. Li, H.; Deyrup, A.; Mensch, J. R., Jr.; Domowicz, M.; Konstantinidis, A. K.; Schwartz, N. B.: The isolation and



characterization of cDNA encoding the mouse bifunctional ATP sulfurylase-adenosine 5-prime-phosphosulfate kinase. *J. Biol. Chem.* 270: 29453–29459, 1995.

[65538] 3868. Hess, J. F.; Casselman, J. T.; FitzGerald, P. G.: Chromosomal locations of the genes for the beaded filament proteins CP 115 and CP 47. *Curr. Eye Res.* 14: 11–18, 1995.

[65539] 3869. Hess, J. F.; Casselman, J. T.; FitzGerald, P. G.: Gene structure and cDNA sequence identify the beaded filament protein CP49 as a highly divergent type I intermediate filament protein. *J. Biol. Chem.* 271: 6729–6735, 1996.

[65540] 3870. Jakobs, P. M.; Hess, J. F.; FitzGerald, P. G.; Kramer, P.; Weleber, R. G.; Litt, M.: Autosomal-dominant congenital cataract associated with a deletion mutation in the human beaded filament protein gene BFSP2. *Am. J. Hum. Genet.* 66: 1432–1436, 2000.

[65541] 3871. Holzinger, A.; Kammerer, S.; Roscher, A. A.: Primary structure of human PMP69, a putative peroxisomal ABC-transporter. *Biochem. Biophys. Res. Commun.* 237: 152–157, 1997.

[65542] 3872. Holzinger, A.; Roscher, A. A.; Landgraf, P.; Lichtner, P.; Kammerer, S.: Genomic organization and chromosomal

localization of the human peroxisomal membrane protein-1-like protein (PXMP1-L) gene encoding a peroxisomal ABC transporter. FEBS Lett. 426: 238–242, 1998.

- [65543] 3873. Shani, N.; Jimenez-Sanchez, G.; Steel, G.; Dean, M.; Valle, D.: Identification of a fourth half ABC transporter in the human peroxisomal membrane. Hum. Molec. Genet. 6: 1925–1931, 1997.
- [65544] 3874. Fink, M.; Duprat, F.; Lesage, F.; Reyes, R.; Romey, G.; Heurteaux, C.; Lazdunski, M.: Cloning, functional expression and brain localization of a novel unconventional outward rectifier K<sup>+</sup> channel. EMBO J. 15: 6854–6862, 1996.
- [65545] 3875. Lesage, F.; Lazdunski, M.: Mapping of human potassium channel genes TREK-1 (KCNK2) and TASK (KCNK3) to chromosomes 1q41 and 2p23. Genomics 51: 478–479, 1998.
- [65546] 3876. Duprat, F.; Lesage, F.; Fink, M.; Reyes, R.; Heurteaux, C.; Lazdunski, M.: TASK, a human background K<sup>+</sup> channel to sense external pH variations near physiological pH. EMBO J. 16: 5464–5471, 1997.
- [65547] 3877. Manjunath, N. A.; Bray-Ward, P.; Goldstein, S. A. N.; Gallagher, P. G.: Assignment of the 2P domain, acid-sensitive potassium channel OAT1 gene KCNK3 to human chromosome bands 2p24.1–p23.3 and murine 5B by in

situ hybridization. Cytogenet. Cell Genet. 86:  
242–243,1999.

- [65548] 3878.Hacker, B. M.; Tomlinson, J. E.; Wayman, G. A.; Sul-  
tana, R.; Chan,G.; Villacres, E.; Disteché, C.; Storm, D. R.:  
Cloning, chromosomal mapping, and regulatory properties  
of the human type 9 adenylyl cyclase(ADCY9). Genomics  
50: 97–104, 1998.
- [65549] 3879.Paterson, J. M.; Smith, S. M.; Harmar, A. J.; Antoni, F.  
A.: Control of a novel adenylyl cyclase by calcineurin.  
Biochem. Biophys. Res. Commun. 214: 1000–1008, 1995.
- [65550] 3880.Premont, R. T.; Matsuoka, I.; Mattei, M. G.; Pouille,  
Y.; Defer,N.; Hanoune, J.: Identification and characteriza-  
tion of a widely expressed form of adenylyl cyclase. J. Biol.  
Chem. 271: 13900–13907,1996.
- [65551] 3881.Toyota, T.; Hattori, E.; Meerabux, J.; Yamada, K.;  
Saito, K.; Shibuya,H.; Nankai, M.; Yoshikawa, T.: Molecular  
analysis, mutation screening, and association study of  
adenylate cyclase type 9 gene (ADCY9) in mood disorders.  
Am. J. Med. Genet. (Neuropsychiat. Genet.) 114:84–92,  
2002.
- [65552] 3882.Chi, N.-W.; Lodish, H. F.: Tankyrase is a Golgi-  
associated mitogen-activated protein kinase substrate that  
interacts with IRAP in GLUT4 vesicles. J. Biol. Chem. 275:

38437–38444, 2000.

- [65553] 3883.Seimiya, H.; Smith, S.: The telomeric poly(ADP-ribose) polymerase,tankyrase 1, contains multiple binding sites for telomeric repeatbinding factor 1 (TRF1) and a novel acceptor, 182-kDa tankyrase-bindingprotein (TAB182). J. Biol. Chem. 277: 14116–14126, 2002.
- [65554] 3884.Smith, S.; Gariat, I.; Schmitt, A.; de Lange, T.: Tankyrase, apoly(ADP-ribose) polymerase at human telomeres. Science 282: 1484–1487,1998.
- [65555] 3885.Zhu, L.; Smith, S.; de Lange, T.; Seldin, M. F.: Chromosomal mappingof the tankyrase gene in human and mouse. Genomics 57: 320–321,1999.
- [65556] 3886.Occhiodoro, T.; Bernheim, L.; Liu, J.-H.; Bijlenga, P.; Sinnreich,M.; Bader, C. R.; Fischer-Lougheed, J.: Cloning of a human ether-a-go-gopotassium channel expressed in myoblasts at the onset of fusion. FEBSLett. 434: 177–182, 1998.
- [65557] 3887.Hidai, H.; Bardales, R.; Goodwin, R.; Quertermous, T.; Quertermous,E. E.: Cloning of capsulin, a basic helix-loop-helix factor expressedin progenitor cells of the pericardium and the coronary arteries. Mech.Dev. 73: 33–43, 1998.

- [65558] 3888.Lu, J.; Chang, P.; Richardson, J. A.; Gan, L.; Weiler, H.; Olson, E. N.: The basic helix-loop-helix transcription factor capsulin controls spleen organogenesis. *Proc. Nat. Acad. Sci.* 97: 9525–9530, 2000.
- [65559] 3889.Lu, J.; Richardson, J. A.; Olson, E. N.: Capsulin: a novel bHLH transcription factor expressed in epicardial progenitors and mesenchyme of visceral organs. *Mech. Dev.* 73: 23–32, 1998.
- [65560] 3890.Quaggin, S. E.; Vanden Heuvel, G. B.; Igarashi, P.: Pod-1, a mesoderm-specific basic-helix-loop-helix protein expressed in mesenchymal and glomerular epithelial cells in the developing kidney. *Mech. Dev.* 71: 37–48, 1998.
- [65561] 3891.Robb, L.; Mifsud, L.; Hartley, L.; Biben, C.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Harvey, R. P.: Epicardin: a novel basic helix-loop-helix transcription factor gene expressed in epicardium, branchial arch myoblasts, and mesenchyme of developing lung, gut, kidney, and gonads. *Dev. Dyn.* 213: 105–113, 1998.
- [65562] 3892.Stacey, P.; Rulten, S.; Dapling, A.; Phillips, S. C.: Molecular cloning and expression of human cGMP-binding cGMP-specific phosphodiesterase (PDE5). *Biochem. Biophys. Res. Commun.* 247: 249–254, 1998.

- [65563] 3893. Yanaka, N.; Kotera, J.; Ohtsuka, A.; Akatsuka, H.; Imai, Y.; Michibata, H.; Fujishige, K.; Kawai, E.; Takebayashi, S.-I.; Okumura, K.; Omori, K.: Expression, structure and chromosomal localization of the human cGMP-binding cGMP-specific phosphodiesterase PDE5A gene. *Eur. J. Biochem.* 255: 391–399, 1998.
- [65564] 3894. Hess, G. F.; Drong, R. F.; Weiland, K. L.; Slightom, J. L.; Sclafani, R. A.; Hollingsworth, R. E.: A human homolog of the yeast CDC7 gene is overexpressed in some tumors and transformed cell lines. *Gene* 211: 133–140, 1998.
- [65565] 3895. Jiang, W.; Hunter, T.: Identification and characterization of a human protein kinase related to budding yeast Cdc7p. *Proc. Nat. Acad. Sci.* 94: 14320–14325, 1997.
- [65566] 3896. Engelender, S.; Kaminsky, Z.; Guo, X.; Sharp, A. H.; Amaravi, R. K.; Kleiderlein, J. J.; Margolis, R. L.; Troncoso, J. C.; Lanahan, A. A.; Worley, P. F.; Dawson, V. L.; Dawson, T. M.; Ross, C. A.: Synphilin-1 associates with alpha-synuclein and promotes the formation of cytosolic inclusions. *Nature Genet.* 22: 110–114, 1999.
- [65567] 3897. Jakes, R.; Spillantini, M. G.; Goedert, M.: Identification of two distinct synucleins from human brain. *FEBS Lett.* 345: 27–32, 1994.
- [65568] 3898. Tory, K.; Latif, F.; Modi, W.; Schmidt, L.; Wei, M. H.;

Li, H.;Cobler, P.; Orcutt, M. L.; Delisio, J.; Geil, L.; Zbar, B.; Lerman,M. I.: A genetic linkage map of 96 loci on the short arm of human chromosome 3. *Genomics* 13: 275–286, 1992.

[65569] 3899.Amiel, J.; Salomon, R.; Attie, T.; Pelet, A.; Trang, H.; Mokhtari,M.; Gaultier, C.; Munnich, A.; Lyonnet, S.: Mutations of the RET–GDNFsignaling pathway in Ondine's curse. (Letter) *Am. J. Hum. Genet.* 62:715–717, 1998.

[65570] 3900.Angrist, M.; Bolk, S.; Thiel, B.; Puffenberger, E. G.; Hofstra,R. M.; Buys, C. H. C. M.; Cass, D. T.; Chakravarti, A.: Mutationanalysis of the RET receptor tyrosine kinase in Hirschsprung disease. *Hum.Molec. Genet.* 4: 821–830, 1995.

[65571] 3901.Antinolo, G.; Marcos, I.; Fernandez, R. M.; Romero, M.; Borrego,S.: A novel germline point mutation, c.2304G(T, in codon 768 of theRET proto–oncogene in a patient with medullary thyroid carcinoma.(Letter) *Am. J. Med. Genet.* 110: 85–87, 2002.

[65572] 3902.Attie, T.; Pelet, A.; Edery, P.; Eng, C.; Mulligan, L. M.; Amiel,J.; Boutrand, L.; Beldjord, C.; Nihoul–Fekete, C.; Munnich, A.; Ponder,B. A. J.; Lyonnet, S.: Diversity of RET proto–oncogene mutationsin familial and sporadic Hirschsprung disease. *Hum. Molec. Genet.* 4:1381–1386,

1995.

- [65573] 3903. Attie-Bitach, T.; Abitbol, M.; Gerard, M.; Delezoide, A.-L.; Auge, J.; Pelet, A.; Amiel, J.; Pachnis, V.; Munnich, A.; Lyonnet, S.; Vekemans, M.: Expression of the RET proto-oncogene in human embryos. *Am. J. Med. Genet.* 80: 481–486, 1998.
- [65574] 3904. Auricchio, A.; Griseri, P.; Carpentieri, M. L.; Betsos, N.; Staiano, A.; Tozzi, A.; Priolo, M.; Thompson, H.; Boccia-  
rdi, R.; Romeo, G.; Ballabio, A.; Ceccherini, I.: Double heterozygosity for a RET substitution interfering with splicing and an EDNRB missense mutation in Hirschsprung disease. (Letter) *Am. J. Hum. Genet.* 64: 1216–1221, 1999.
- [65575] 3905. Batourina, E.; Choi, C.; Paragas, N.; Bello, N.; Hensle, T.; Costantini, F. D.; Schuchardt, A.; Bacallao, R. L.; Mendelsohn, C. L.: Distal ureter morphogenesis depends on epithelial cell remodeling mediated by vitamin A and Ret. *Nature Genet.* 32: 109–115, 2002. Note: Erratum: *Nature Genet.* 32: 331 only, 2002.
- [65576] 3906. Batourina, E.; et al; et al: Vitamin A controls epithelial/mesenchymal interactions through Ret expression. *Nature Genet.* 27: 74–78, 2001.
- [65577] 3907. Berndt, I.; Reuter, M.; Saller, B.; Frank-Raue, K.; Groth, P.; Grubendorf, M.; Raue, F.; Ritter, M. M.; Hoppner,



W.: A new hot spot for mutations in the ret protooncogene causing familial medullary thyroid carcinoma and multiple endocrine neoplasia type 2A. *J. Clin. Endocr. Metab.* 83: 770–774, 1998.

[65578] 3908. Boccia, L. M.; Green, J. S.; Joyce, C.; Eng, C.; Taylor, S. A. M.; Mulligan, L. M.: Mutation of RET codon 768 is associated with the FMTC phenotype. *Clin. Genet.* 51: 81–85, 1997.

[65579] 3909. Bolino, A.; Schuffenecker, I.; Luo, Y.; Seri, M.; Silengo, M.; Tocco, T.; Chabrier, G.; Houdent, C.; Murat, A.; Schlumberger, M.; Tourniaire, J.; Lenoir, G. M.; Romeo, G.: RET mutations in exons 13 and 14 of FMTC patients. *Oncogene* 10: 2415–2419, 1995.

[65580] 3910. Bolk, S.; Angrist, M.; Schwartz, S.; Silvestri, J. M.; Weese-Mayer, D. E.; Chakravarti, A.: Congenital central hypoventilation syndrome: mutation analysis of the receptor tyrosine kinase RET. *Am. J. Med. Genet.* 63: 603–609, 1996.

[65581] 3911. Bolk Gabriel, S.; Salomon, R.; Pelet, A.; Angrist, M.; Amiel, J.; Fornage, M.; Attie-Bitach, T.; Olson, J. M.; Hofstra, R.; Buys, C.; Steffann, J.; Munnich, A.; Lyonnet, S.; Chakravarti, A.: Segregation at three loci explains familial and population risk in Hirschsprung disease. *Nature Genet.*

31: 89–93, 2002.

- [65582] 3912.Ceccherini, I.; Hofstra, R. M.; Luo, Y.; Stulp, R. P.; Barone,V.; Stelwagen, T.; Bocciardi, R.; Nijveen, H.; Bolino, A.; Seri, M.;Ronchetto, P.; Pasini, B.; Bozzano, M.; Buys, C. H. C. M.; Romeo,G.: DNA polymorphisms and conditions for SSCP analysis of the 20exons of the Ret proto-oncogene. *Oncogene* 9: 3025–3029, 1994.
- [65583] 3913.Azen, E. A.; Goodman, P. A.; Lalley, P. A.: Human salivary proline-richprotein genes on chromosome 12. *Am. J. Hum. Genet.* 37: 418–424,1985.
- [65584] 3914.Mamula, P. W.; Heerema, N. A.; Palmer, C. G.; Lyons, K. M.; Karn,R. C.: Localization of the human salivary protein complex (SPC) tochromosome band 12p13.2. *Cytogenet. Cell Genet.* 39: 279–284, 1985.
- [65585] 3915.Azen, E. A.: Genetic protein polymorphism in human saliva: aninterpretive review. *Biochem. Genet.* 16: 79–99, 1978.
- [65586] 3916.Azen, E. A.; Denniston, C.: Genetic polymorphism of PIF (parotidisolectric focusing variant) proteins with linkage to the PPP (parotidproline-rich protein) gene complex. *Biochem. Genet.* 19: 475–485,1981.
- [65587] 3917.Azen, E. A.; Denniston, C. L.: Genetic polymorphism of human salivaryproline-rich proteins: further genetic

analysis. *Biochem. Genet.* 12:109–120, 1974.

- [65588] 3918.Azen, E. A.; Kim, H.-S.; Goodman, P.; Flynn, S.; Maeda, N.: Alleles at the PRH1 locus coding for the human salivary–acidic proline–rich proteins Pa, Db, and PIF. *Am. J. Hum. Genet.* 41: 1035–1047, 1987.
- [65589] 3919.Azen, E. A.; Lush, I. E.; Taylor, B. A.: Close linkage of mouse genes for salivary proline–rich proteins (PRPs) and taste. *Trends Genet.* 2: 199–200, 1986.
- [65590] 3920.Azen, E. A.; Maeda, N.: Molecular genetics of human salivary proteins and their polymorphisms. *Adv. Hum. Genet.* 17: 141–199, 1988.
- [65591] 3921.Bernstein, F.: Ergebnisse einer biostatistischen zusammenfassenden Betrachtung ueber die erblichen Blutstrukturen des Menschen. *Klin. Wschr.* 3: 1495–1497, 1924.
- [65592] 3922.Bernstein, F.: Zusammenfassende Betrachtungen ueber die erblichen Blutstrukturen des Menschen. *Z. indukt. Abstamm. u. Vererb. Lehre* 37:237–270, 1925. Note: Translation: Blood Transfusion Division, U.S. Army Medical Research Laboratory, Fort Knox, Kentucky 40121.
- [65593] 3923.Friedman, R. D.; Merritt, A. D.: Partial purification and characterization of a polymorphic protein (Pa) in human parotid saliva. *Am. J. Hum. Genet.* 27: 304–314, 1975.

- [65594] 3924.Friedman, R. D.; Merritt, A. D.; Rivas, M. L.: Genetic studies of human acidic salivary protein (Pa). *Am. J. Hum. Genet.* 27: 292–303, 1975.
- [65595] 3925.Ikemoto, S.; Minaguchi, K.; Hinohara, H.: Genetic polymorphisms of human parotid salivary proteins (Pa, Pb, Pr, Db and Pm) and salivary amylase isozyme in Japanese population. *Hum. Hered.* 27: 328–331, 1977.
- [65596] 3926.Kim, H.-S.; Maeda, N.: Structures of two HaeIII-type genes in the human salivary proline-rich protein multigene family. *J. Biol.Chem.* 261: 6712–6718, 1986.
- [65597] 3927.Maeda, N.: Inheritance of the human salivary proline-rich proteins: a reinterpretation in terms of six loci forming two subfamilies. *Biochem.Genet.* 23: 455–464, 1985.
- [65598] 3928.Maeda, N.; Kim, H.-S.; Azen, E. A.; Smithies, O.: Differential RNA splicing and post-translational cleavages in the human salivary proline-rich protein gene system. *J. Biol. Chem.* 260: 11123–11130, 1985.
- [65599] 3929.O'Connell, P.; Lathrop, G. M.; Law, M.; Leppert, M.; Nakamura, Y.; Hoff, M.; Kumlin, E.; Thomas, W.; Elsner, T.; Ballard, L.; Goodman, P.; Azen, E.; Sadler, J. E.; Cai, G. Y.; Lalouel, J.-M.; White, R.: A primary genetic linkage map for human chromosome 12. *Genomics* 1:93–102, 1987.

- [65600] 3930.O'Hanlon, K.; Weissbecker, K.; Cortessis, V.; Spence, M. A.; Azen,E. A.: Genes for salivary proline-rich proteins and taste for phenylthioureaare not closely linked in humans. *Cytogenet. Cell Genet.* 49: 315-317,1988.
- [65601] 3931.Stern, C.: *Principles of Human Genetics*. San Francisco: W.H. Freeman (pub.) (3rd ed.): 1973. Pp. 256-262.
- [65602] 3932.von Dungern, E.; Hirschfeld, L.: Ueber Vererbung gruppenspezifischerStrukturen des Blutes. II. *Z. Immunforsch.* 6: 284-292, 1910. Note:Translation: G. P. Pohlmann: *Transfusion* 2: 70-74, 1962.
- [65603] 3933.Yu, P. L.; Karn, R. C.; Merritt, A. D.: Multipoint mapping ofthe human parotid salivary proteins: Pr, Pa, Db. (Abstract) *Am. J.Hum. Genet.* 30: 129A only, 1978.
- [65604] 3934.Yu, P. L.; Schwartz, R. C.; Merritt, A. D.; Azen, E. A.; Rivas,M. L.; Karn, R. C.; Craft, M. A.: Linkage relationships of the proline-richsalivary proteins (Pr, Pa, Db). *Cytogenet. Cell Genet.* 22: 655-658,1978.
- [65605] 3935.Furlong, C. E.; Richter, R. J.; Chapline, C.; Crabb, J. W.: Purificationof rabbit and human serum paraoxonase. *Biochemistry* 30: 10133-10140,1991.
- [65606] 3936.Furlong, C. E.; Richter, R. J.; Seidel, S. L.; Motulsky, A. G.: Role of genetic polymorphism of human plasma paraoxonase/arylesterasein hydrolysis of the insecticide

metabolites chlorpyrifos oxon and paraoxon. Am. J. Hum. Genet. 43: 230–238, 1988.

[65607] 3937. Garin, M.-C. B.; James, R. W.; Dussoix, P.; Blanche, H.; Passa, P.; Froguel, P.; Ruiz, J.: Paraoxonase polymorphism met-leu54 is associated with modified serum concentrations of the enzyme: a possible link between the paraoxonase gene and increased risk of cardiovascular disease in diabetes. J. Clin. Invest. 99: 62–66, 1997.

[65608] 3938. Geldmacher-von Mallinckrodt, M.: Polymorphism of human serum paraoxonase. Hum. Genet. 45 (suppl. 1): 65–68, 1978.

[65609] 3939. Geldmacher-von Mallinckrodt, M.; Lindorft, H. H.; Petenyi, M.; Flugel, M.; Fischer, T.; Hiller, T.: Genetisch determinierter Polymorphismus des menschlichen Serum-Paroxonase (E.C.3.1.1.2). Humangenetik 17:331–335, 1973.

[65610] 3940. Hassett, C.; Richter, R. J.; Humbert, R.; Chapline, C.; Crabb, J. W.; Omiecinski, C. J.; Furlong, C. E.: Characterization of cDNA clones encoding rabbit and human serum paraoxonase: the mature protein retains its signal sequence. Biochemistry 30: 10141–10149, 1991.

[65611] 3941. Humbert, R.; Adler, D. A.; Disteche, C. M.; Hassett, C.; Omiecinski, C. J.; Furlong, C. E.: The molecular basis of

the human serum paraoxonase activity polymorphism. *Nature Genet.* 3: 73–76, 1993.

- [65612] 3942. Ito, T.; Yasue, H.; Yoshimura, M.; Nakamura, S.; Nakayama, M.; Shimasaki, Y.; Harada, E.; Mizuno, Y.; Kawano, H.; Ogawa, H.: Paraoxonase gene gln192-to-arg (Q192R) polymorphism is associated with coronary artery spasm. *Hum. Genet.* 110: 89–94, 2002.
- [65613] 3943. Kao, Y.-L.; Donaghue, K.; Chan, A.; Knight, J.; Silink, M.: A variant of paraoxonase (PON1) gene is associated with diabetic retinopathy in IDDM. *J. Clin. Endocr. Metab.* 83: 2589–2592, 1998.
- [65614] 3944. La Du, B. N.: The human serum paraoxonase/arylesterase polymorphism. (Editorial) *Am. J. Hum. Genet.* 43: 227–229, 1988.
- [65615] 3945. Li, W.-F.; Furlong, C. E.; Costa, L. G.: Paraoxonase protects against chlorpyrifos toxicity in mice. *Toxic. Lett.* 76: 219–226, 1995.
- [65616] 3946. Li, W.-F.; Matthews, C.; Disteché, C. M.; Costa, L. G.; Furlong, C. E.: Paraoxonase (Pon1) gene in mice: sequencing, chromosomal localization and developmental expression. *Pharmacogenetics* 7: 137–144, 1997.
- [65617] 3947. Mackness, B.; Mackness, M. I.; Arrol, S.; Turkie, W.; Durrington, P. N.: Effect of the human serum paraoxonase

55 and 192 genetic polymorphisms on the protection by high density lipoprotein against low density lipoprotein oxidative modification. FEBS Lett. 423: 57–60, 1998.

- [65618] 3948. Mochizuki, H.; Scherer, S. W.; Xi, T.; Nickle, D. C.; Majer, M.; Huizenga, J. J.; Tsui, L.-C.; Prochazka, M.: Human PON2 gene at 7q21.3: cloning, multiple mRNA forms, and missense polymorphisms in the coding sequence. Gene 213: 149–157, 1998.
- [65619] 3949. Mueller, R. F.; Hornung, S.; Furlong, C. E.; Anderson, J.; Giblett, E. R.; Motulsky, A. G.: Plasma paraoxonase polymorphism: a new enzyme assay, population, family, biochemical, and linkage studies. Am. J. Hum. Genet. 35: 393–408, 1983.
- [65620] 3950. Navab, M.; Hama-Levy, S.; Van Lenten, B. J.; Fonarow, G. C.; Cardinez, C. J.; Castellani, L. W.; Brennan, M.-L.; Lusis, A. J.; Fogelman, A. M.: Mildly oxidized LDL induces an increased apolipoprotein J/paraoxonase ratio. J. Clin. Invest. 99: 2005–2019, 1997.
- [65621] 3951. Neel, J. V.; Tanis, R. J.; Migliazza, E. C.; Spielman, R. S.; Salzano, F. M.; Oliver, W. J.; Morrow, M.; Bachofer, S.: Genetic studies of the Macushi and Wapishana Indians. I. Rare genetic variants and a 'private polymorphism' of esterase A. Hum. Genet. 36: 81–108, 1977.



- [65622] 3952.Nielsen, A.; Eiberg, H.; Mohr, J.: Number of loci responsible for the inheritance of high and low activity of paraoxonase. Clin.Genet. 29: 216–221, 1986.
- [65623] 3953.Odawara, M.; Tachi, Y.; Yamashita, K.: Paraoxonase polymorphism Gln192–Arg is associated with coronary heart disease in Japanese noninsulin–dependent diabetes mellitus. J. Clin. Endocr. Metab. 82: 2257–2260, 1997.
- [65624] 3954.Ortigoza–Ferado, J.; Richter, R. J.; Hornung, S. K.; Motulsky, A. G.; Furlong, C. E.: Paraoxon hydrolysis in human serum mediated by a genetically variable arylesterase and albumin. Am. J. Hum. Genet. 36:295–305, 1984.
- [65625] 3955.Paolisso, G.; Manzella, D.; Tagliamonte, M. R.; Barberi, M.; Marfella, R.; Zito, G.; Bonafe, M.; Giugliano, D.; Franceschi, C.; Varricchio, M.: The BB–paraoxonase genotype is associated with impaired brachial reactivity after acute hypertriglyceridemia in healthy subjects. J.Clin. Endocr. Metab. 86: 1078–1082, 2001.
- [65626] 3956.Playfer, J. R.; Eze, L. C.; Bullen, M. F.; Evans, D. A. P.: Genetic polymorphism and interethnic variability of plasma paraoxonase activity. J.Med. Genet. 13: 337–342, 1976.
- [65627] 3957.Schmiegelow, K.; Eiberg, H.; Tsui, L.–C.; Buchwald, M.; Phelan, P. D.; Williamson, R.; Warwick, W.; Niebuhr, E.; Mohr, J.; Schwartz, M.; Koch, C.: Linkage between the loci

for cystic fibrosis and paraoxonase. Clin.Genet. 29: 374–377, 1986.

- [65628] 3958.Serrato, M.; Marian, A. J.: A variant of human paraoxonase/arylesterase(HUMPONA) gene is a risk factor for coronary artery disease. J. Clin.Invest. 96: 3005–3008, 1995.
- [65629] 3959.Shih, D. M.; Gu, L.; Xia, Y.–R.; Navab, M.; Li, W.–F.; Hama, S.;Castellani, L. W.; Furlong, C. E.; Costa, L. G.; Fogelman, A. M.;Lusis, A. J.: Mice lacking serum paraoxonase are susceptible to organophosphatetoxicity and atherosclerosis. Nature 394: 284–287, 1998.
- [65630] 3960.Simpson, N. E.: Serum arylesterase levels of activity in twinsand their parents. Am. J. Hum. Genet. 23: 375–382, 1971.
- [65631] 3961.Sorenson, R. C.; Primo–Parmo, S. L.; Camper, S.; La Du, B. N.: The genetic mapping and gene structure of mouse paraoxonase/arylesterase. Genomics 30:431–438, 1995.
- [65632] 3962.Tashian, R. E.: Genetic variation and evolution of the carboxylicesterases and carbonic anhydrases of primate erythrocytes. Am. J.Hum. Genet. 17: 257–272, 1965.
- [65633] 3963.Tashian, R. E.; Shaw, M. W.: Inheritance of an erythrocyte acetylesterasevariant of man. Am. J. Hum. Genet.

14: 295–300, 1962.

- [65634] 3964. Tsui, L.-C.; Buchwald, M.; Barker, D.; Braman, J. C.; Knowlton, R.; Schumm, J. W.; Eiberg, H.; Mohr, J.; Kennedy, D.; Plavsic, N.; Zsiga, M.; Markiewicz, D.; Akots, G.; Brown, V.; Helms, C.; Gravius, T.; Parker, C.; Rediker, K.; Donis-Keller, H.: Cystic fibrosis locus defined by a genetically linked polymorphic DNA marker. *Science* 230:1054–1057, 1985.
- [65635] 3965. Gogolin, K. J.; Wray, L. K.; Slaughter, C. A.; Harris, H.: A monoclonal antibody that reacts with nonallelic enzyme glycoproteins. *Science* 216:59–61, 1982.
- [65636] 3966. Goldstein, D. J.; Rogers, C. E.; Harris, H.: Expression of alkaline phosphatase loci in mammalian tissues. *Proc. Nat. Acad. Sci.* 77:2857–2860, 1980.
- [65637] 3967. Gould, B. S.: Studies on the source of serum phosphatase: the nature of the increased serum phosphatase in rats after fat feeding. *Arch. Biochem.* 4: 175–181, 1944.
- [65638] 3968. Griffin, C. A.; Smith, M.; Henthorn, P. S.; Harris, H.; Weiss, M. J.; Raducha, M.; Emanuel, B. S.: Human placental and intestinal alkaline phosphatase genes map to 2q34–q37. *Am. J. Hum. Genet.* 41:1025–1034, 1987.
- [65639] 3969. Henthorn, P. S.; Raducha, M.; Edwards, Y. H.; Weiss, M. J.; Slaughter, C.; Lafferty, M. A.; Harris, H.: Nucleotide

and amino acid sequences of human intestinal alkaline phosphatase: close homology to placental alkaline phosphatase. *Proc. Nat. Acad. Sci.* 84: 1234–1238, 1987.

[65640] 3970. Henthorn, P. S.; Raducha, M.; Kadesch, T.; Weiss, M. J.; Harris, H.: Sequence and characterization of the human intestinal alkaline phosphatase gene. *J. Biol. Chem.* 263: 12011–12019, 1988.

[65641] 3971. Langman, M. J. S.; Leuthold, E.; Robson, E. B.; Harris, J.; Luffman, J. E.; Harris, H.: Influence of diet on the 'intestinal' component of serum alkaline phosphatase in people of different ABO blood groups and secretor status. *Nature* 212: 41–43, 1966.

[65642] 3972. Lehmann, F.-G.: Human alkaline phosphatases: evidence of three isoenzymes (placental, intestinal and liver–bone–kidney–type) by lectin–binding affinity and immunological specificity. *Biochim. Biophys. Acta* 616: 41–59, 1980.

[65643] 3973. Rajput, B.; Degen, S. F.; Reich, E.; Waller, E. K.; Axelrod, J.; Eddy, R. L.; Shows, T. B.: Chromosomal locations of human tissue plasminogen activator and urokinase genes. *Science* 230: 672–674, 1985.

[65644] 3974. Chen, I.-T.; Roufa, D. J.: The transcriptionally active human ribosomal protein S17 gene. *Gene* 70: 107–116,

1988.

- [65645] 3975.Lang, F.; Klingel, K.; Wagner, C. A.; Stegen, C.; Warntges, S.;Friedrich, B.; Lanzendorfer, M.; Melzig, J.; Moschen, I.; Steuer,S.; Waldegger, S.; Sauter, M.; and 9 others: Deranged transcriptionalregulation of cell-vol-ume-sensitive kinase hSGK in diabetic nephropathy. Proc.Nat. Acad. Sci. 97: 8157–8162, 2000.
- [65646] 3976.Ebner, R.; Chen, R.–H.; Shum, L.; Lawler, S.; Zioncheck, T. F.;Lee, A.; Lopez, A. R.; Derynck, R.: Cloning of a type I TGF–betareceptor and its effect on TGF–beta binding to the type II receptor. Science 260:1344–1348, 1993.
- [65647] 3977.Franzen, P.; ten Dijke, P.; Ichijo, H.; Yamashita, H.; Schulz,P.; Heldin, C.–H.; Miyazono, K.: Cloning of a TGF–beta type I receptorthat forms a heteromeric complex with the TGF–beta type II receptor. Cell 75:681–692, 1993.
- [65648] 3978.Inman, G. J.; Nicolas, F. J.; Hill, C. S.: Nucleocyto-plasmic shuttlingof Smads 2, 3, and 4 permits sensing of TGF–beta receptor activity. Molec.Cell 10: 283–294, 2002.
- [65649] 3979.Johnson, D. W.; Qumsiyeh, M.; Benkhalifa, M.; Marchuk, D. A.:Assignment of human transforming growth factor–beta type I and typelll receptor genes (TGFB1 and TGFB3) to 9q33–q34 and 1p32–p33, re–

spectively. *Genomics* 28:356–357, 1995.

[65650] 3980.Kuan, J.; Kono, D. H.: *Tgfbr1* maps to chromosome 4. *MammalianGenome* 9: 95–96, 1998.

[65651] 3981.Pasche, B.; Luo, Y.; Rao, P. H.; Nimer, S. D.; Dmitro-  
vsky, E.;Caron, P.; Luzzatto, L.; Offit, K.; Cordon-Cardo,  
C.; Renault, B.;Satagopan, J. M.; Murty, V. V.; Massague, J.:  
Type I transforminggrowth factor beta receptor maps to  
9q22 and exhibits a polymorphismand a rare variant  
within a polyalanine tract. *Cancer Res.* 58:  
2727–2732,1998.

[65652] 3982.D'Eustachio, P.; Meyuhas, O.; Ruddle, F.; Perry, R. P.:  
Chromosomal distribution of ribosomal protein genes in  
the mouse. *Cell* 24: 307–312,1981.

[65653] 3983.Filipenko, M. L.; Iantsen, E. I.; Muravlev, A. I.;  
Kopantsev, E.P.; Karpova, G. G.; Mertvetsov, N. P.: Map-  
ping the genes for ribosomal proteins S14 and S17 on hu-  
man chromosomes using cDNA from a panel of hybrid  
cells. *Bioorg. Khim.* 21: 349–353, 1995. Note: Article in  
Russian.

[65654] 3984.Woolford, J. L., Jr.; Hereford, L. M.; Rosbash, M.: Iso-  
lation of cloned DNA sequences containing ribosomal pro-  
tein genes from *Saccharomyces cerevisiae*. *Cell* 18:  
1247–1259, 1979.

- [65655] 3985.Nucifora, G.; Begy, C. R.; Erickson, R.; Drabkin, H. A.; Rowley, J. D.: The 3;21 translocation in myelodysplasia results in a fusion transcript between the AML1 gene and the gene for EAP, a highly conserved protein associated with the Epstein-Barr virus small RNA EBER 1. *Proc. Nat. Acad. Sci.* 90: 7784-7788, 1993.
- [65656] 3986.Chang, A.; Schalkwijk, J.; Happle, R.; van de Kerkhof, P. C. M.: Elastase-inhibiting activity in scaling skin disorders. *Acta Derm.Venerol.* 70: 147-151, 1990.
- [65657] 3987.Kuijpers, A. L. A.; Pfundt, R.; Zeeuwen, P. L. J. M.; Molhuizen, H. O. F.; Mariman, E. C. M.; van de Kerkhof, P. C. M.; Schalkwijk, J.: SKALP/elafin gene polymorphisms are not associated with pustular forms of psoriasis. *Clin. Genet.* 54: 96-101, 1998.
- [65658] 3988.Molhuizen, H. O. F.; Alkemade, H. A. C.; Zeeuwen, P. L. J. M.; de Jongh, G. J.; Wieringa, B.; Schalkwijk, J.: SKALP/elafin: an elastase inhibitor from cultured human keratinocytes: purification, cDNA sequence, and evidence for transglutaminase cross-linking. *J. Biol. Chem.* 268:12028-12032, 1993.
- [65659] 3989.Molhuizen, H. O. F.; Zeeuwen, P. L. J. M.; Olde Weghuis, D.; Geurts van Kessel, A.; Schalkwijk, J.: Assignment of the human gene encoding the epidermal serine

proteinase inhibitor SKALP (PI3) to chromosomeregion 20q12–q13. Cytogenet. Cell Genet. 66: 129–131, 1994.

[65660] 1990.Sallenave, J.–M.; Ryle, A. P.: Purification and characterization of elastase–specific inhibitor: sequence homology with mucus proteinaseinhibitor. Biol. Chem. Hoppe–Seyler 372: 13–21, 1991.

[65661] 1991.Schalkwijk, J.; Chang, A.; Janssen, P.; de Jongh, G. J.; Mier, P. D.: Skin–derived antileucoproteases (SKALPs): characterization of two new elastase inhibitors from psoriatic epidermis. Brit. J.Derm. 122: 631–641, 1990.

[65662] 1992.Schalkwijk, J.; de Roo, C.; de Jongh, G. J.: Skin–derived antileukoproteinase(SKALP), an elastase inhibitor from human keratinocytes: purification and biochemical properties. Biochim. Biophys. Acta 1096: 148–154, 1991.

[65663] 1993.Wiedow, O.; Schroeder, J.–M.; Gregory, H.; Young, J. A.; Christophers, E.: Elafin: an elastase–specific inhibitor of human skin: purification, characterization, and complete amino acid sequence. J. Biol. Chem. 265:14791–14795, 1990.

[65664] 1994.Begley, C. G.; Visvader, J.; Green, A. R.; Aplan, P. D.; Metcalf, D.; Kirsch, I. R.; Gough, N. M.: Molecular cloning and chromosomal localization of the murine homolog of the human helix–loop–helix gene SCL. Proc. Nat. Acad. Sci.



88: 869–873, 1991.

- [65665] 3995.Finger, L. R.; Kagan, J.; Christopher, G.; Kurtzberg, J.; Hershfield, M. S.; Nowell, P. C.; Croce, C. M.: Involvement of the TCL5 gene on human chromosome 1 in T-cell leukemia and melanoma. *Proc. Nat. Acad. Sci.* 86: 5039–5043, 1989.
- [65666] 3996.Gottgens, B.; Barton, L. M.; Chapman, M. A.; Sinclair, A. M.; Knudsen, B.; Grafham, D.; Gilbert, J. G. R.; Rogers, J.; Bentley, D. R.; Green, A. R.: Transcriptional regulation of the stem cell leukemia gene (SCL)—comparative analysis of five vertebrate SCL loci. *Genome Res.* 12:749–759, 2002.
- [65667] 3997.Gottgens, B.; Gilbert, J. G.; Barton, L. M.; Grafham, D.; Rogers, J.; Bentley, D. R.; Green, A. R.: Long-range comparison of human and mouse SCL loci: localized regions of sensitivity to restriction endonucleases correspond precisely with peaks of conserved noncoding sequences. *Genome Res.* 11: 87–97, 2001.
- [65668] 3998.Kocher, O.; Cheresch, P.; Lee, S. W.: Identification and partial characterization of a novel membrane-associated protein (MAP17) up-regulated in human carcinomas and modulating cell replication and tumor growth. *Am. J. Path.* 149: 493–500, 1996.

- [65669] 3999.Kozak, M.: An analysis of 5-prime-noncoding sequences from 699vertebrate messenger RNAs. *Nucleic Acids Res.* 15: 8125–8148, 1987.
- [65670] 4000.Kurtzberg, J.; Bigner, S. H.; Hershfield, M. S.: Establishment of the DU.528 human lymphohemopoietic stem cell line. *J. Exp. Med.* 162:1561–1578, 1985.
- [65671] 4001.Robb, L.; Lyons, I.; Li, R.; Hartley, L.; Kontgen, F.; Harvey, R. P.; Metcalf, D.; Begley, C. G.: Absence of yolk sac hematopoiesis from mice with a targeted disruption of the *scl* gene. *Proc. Nat. Acad. Sci.* 92: 7075–7079, 1995.
- [65672] 4002.Shivdasani, R. A.; Mayer, E. L.; Orkin, S. H.: Absence of blood formation in mice lacking the T-cell leukaemia oncoprotein tal-1/SCL. *Nature* 373:432–434, 1995.
- [65673] 4003.Sinclair, A. M.; Bench, A. J.; Bloor, A. J. C.; Li, J.; Gottgens, B.; Stanley, M. L.; Miller, J.; Piltz, S.; Hunter, S.; Nacheva, E. P.; Sanchez, M.-J.; Green, A. R.: Rescue of the lethal *scl*<sup>-/-</sup> phenotype by the human SCL locus. *Blood* 99: 3931–3938, 2002.
- [65674] 4004.Williams, M.; Lyu, M. S.; Yang, Y. L.; Lin, E. P.; Dunbrack, R.; Birren, B.; Cunningham, J.; Hunter, K.: *ler5*, a novel member of the slow-kinetics immediate-early genes. *Genomics* 44: 327–334, 1999.
- [65675] 4005.Xia, Y.; Brown, L.; Yang, C. Y.-C.; Tsou Tsan, J.; Si-

ciliano, M. J.; Espinosa, R., III; Le Beau, M. M.; Baer, R. J.:  
TAL2, a helix-loop-helix gene activated by the  
(7;9)(q34;q32) translocation in human T-cell leukemia.  
Proc. Nat. Acad. Sci. 88: 11416–11420, 1991.

[65676] 4006. Durick, K.; Gill, G. N.; Taylor, S. S.: Shc and Enigma  
are both required for mitogenic signaling by Ret/ptc2.  
Molec. Cell. Biol. 18:2298–2308, 1998.

[65677] 4007. Saadat, M.; Nomoto, K.; Mizuno, Y.; Kikuchi, K.;  
Yoshida, M. C.: Assignment of the gene encoding type  
1-gamma protein phosphatase catalytic subunit (PPP1CC)  
on human, rat, and mouse chromosomes. Jpn. J. Hum.  
Genet. 41: 159–165, 1996.

[65678] 4008. Cohen, P.; Cohen, P. T. W.: Protein phosphatases  
come of age. J. Biol. Chem. 264: 21435–21438, 1989.

[65679] 4009. Groves, M. R.; Hanlon, N.; Turowski, P.; Hemmings,  
B. A.; Barford, D.: The structure of the protein phosphatase  
2A PR65/A subunit reveals the conformation of its 15  
tandemly repeated HEAT motifs. Cell 96:99–110, 1999.

[65680] 4010. Jones, T. A.; Barker, H. M.; da Cruz e Silva, E. F.;  
Mayer-Jaekel, R. E.; Hemmings, B. A.; Spurr, N. K.; Sheer,  
D.; Cohen, P. T. W.: Localization of the genes encoding the  
catalytic subunits of protein phosphatase 2A to human  
chromosome bands 5q23–q31 and

8p12–p11.2, respectively. Cytogenet. Cell Genet. 63: 35–41, 1993.

[65681] 4011.Stone, S. R.; Mayer, R.; Wernet, W.; Maurer, F.; Hofsteenge, J.; Hemmings, B. A.: The nucleotide sequence of the cDNA encoding the human lung protein phosphatase 2A alpha catalytic subunit. Nucleic Acids Res. 16: 11365 only, 1988.

[65682] 4012.Hemmings, B. A.; Wernet, W.; Mayer, R.; Maurer, F.; Hofsteenge, J.; Stone, S. R.: The nucleotide sequence of the cDNA encoding the human lung protein phosphatase 2A–beta catalytic subunit. Nucleic Acids Res. 16: 11366 only, 1988.

[65683] 4013.Imbert, A.; Chaffanet, M.; Essioux, L.; Noguchi, T.; Adelaide, J.; Kerangueven, F.; Le Paslier, D.; Bonaiti–Pellie, C.; Sobol, H.; Birnbaum, D.; Pebusque, M.–J.: Integrated map of the chromosome 8p12–p21 region, a region involved in human cancers and Werner syndrome. Genomics 32:29–38, 1996.

[65684] 4014.Adamkiewicz, T. V.; McSherry, C.; Bach, F. H.; Houchins, J. P.: Natural killer lectin–like receptors have divergent carboxy–termini, distinct from C–type lectins. Immunogenetics 39: 218 only, 1994.

[65685] 4015.Houchins, J. P.; Yabe, T.; McSherry, C.; Bach, F. H.:

DNA sequence analysis of NKG2, a family of related cDNA clones encoding type II integral membrane proteins on human natural killer cells. *J. Exp. Med.* 173: 1017–1020, 1991.

[65686] 4016. Plougastel, B.; Jones, T.; Trowsdale, J.: Genomic structure, chromosomal location, and alternative splicing of the human NKG2A gene. *Immunogenetics* 44:286–291, 1996.

[65687] 4017. Plougastel, B.; Trowsdale, J.: Sequence analysis of a 62-kb region overlapping the human KLRC cluster of genes. *Genomics* 49: 193–199, 1998.

[65688] 4018. Renedo, M.; Arce, I.; Rodriguez, A.; Carretero, M.; Lanier, L.L.; Lopez-Botet, M.; Fernandez-Ruiz, E.: The human natural killer gene complex is located on chromosome 12p12–p13. *Immunogenetics* 46:307–311, 1997.

[65689] 4019. Yabe, T.; McSherry, C.; Bach, F. H.; Fisch, P.; Schall, R. P.; Sondel, P. M.; Houchins, J. P.: A multigene family on human chromosome 12 encodes natural killer-cell lectins. *Immunogenetics* 37: 455–460, 1993.

[65690] 4020. Cooper, A. M.; Kipnis, A.; Turner, J.; Magram, J.; Ferrante, J.; Orme, I. M.: Mice lacking bioactive IL-12 can generate protective, antigen-specific cellular responses to mycobacterial infection only if the IL-12 p40 subunit is

present. *J. Immun.* 168: 1322–1327, 2002.

[65691] 4021. Gubler, U.; Chua, A. O.; Schoenhaut, D. S.; Dwyer, C. M.; McComas, W.; Motyka, R.; Nabavi, N.; Wolitzky, A. G.; Quinn, P. M.; Familletti, P. C.; Gately, M. K.: Coexpression of two distinct genes is required to generate secreted bioactive cytotoxic lymphocyte maturation factor.

*Proc. Nat. Acad. Sci.* 88: 4143–4147, 1991.

[65692] 4022. Hall, S. S.: IL-12 at the crossroads. *Science* 268: 1432–1434, 1995.

[65693] 4023. Merberg, D. M.; Wolfe, S. F.; Clark, S. C.: Sequence similarity between NKSF and the IL6/GCSF family. (Letter) *Immun. Today* 13: 77–78, 1992.

[65694] 4024. Schwarz, A.; Stender, S.; Berneburg, M.; Bshmi, M.; Kulms, D.; van Steeg, H.; Grosse-Heitmeyer, K.; Krutmann, J.; Schwarz, T.: Interleukin-12 suppresses ultraviolet radiation-induced apoptosis by inducing DNA repair. *Nature Cell Biol.* 4: 26–31, 2002.

[65695] 4025. Schweitzer, P. A.; Noben-Trauth, N.; Pelsue, S. C.; Johnson, K. R.; Wolf, S. F.; Shultz, L. D.: Genetic mapping of the IL-12 alpha chain gene (Il12a) on mouse chromosome 3. *Mammalian Genome* 7: 394–395, 1996.

[65696] 4026. Sieburth, D.; Jabs, E. W.; Warrington, J. A.; Li, X.; Lasota, J.; LaForgia, S.; Kelleher, K.; Huebner, K.; Wasmuth, J.

J.; Wolf, S.F.: Assignment of genes encoding a unique cytokine (IL12) composed of two unrelated subunits to chromosomes 3 and 5. *Genomics* 14: 59–62, 1992.

[65697] 4027. Wolf, S. F.; Sieburth, D.; Sypek, J.: Interleukin 12: a key modulator of immune function. *Stem Cells* 12: 154–168, 1994.

[65698] 4028. Wolf, S. F.; Temple, P. A.; Kobayashi, M.; Young, D.; Dicig, M.; Lowe, L.; Dzialo, R.; Fitz, L.; Ferenz, C.; Hewick, R. M.; Kelleher, K.; Herrmann, S. H.; Clark, S. C.; Azzoni, L.; Chan, S. H.; Trinchieri, G.; Perussia, B.: Cloning of cDNA for natural killer cell stimulatory factor, a heterodimeric cytokine with multiple biologic effects on T and natural killer cells. *J. Immun.* 146: 3074–3081, 1991.

[65699] 4029. Mizuno, T.; Kaibuchi, K.; Yamamoto, T.; Kawamura, M.; Sakoda, T.; Fujioka, H.; Matsuura, Y.; Takai, Y.: A stimulatory GDP/GTP exchange protein for smg p21 is active on the post-translationally processed form of c-Ki-ras p21 and rhoA p21. *Proc. Nat. Acad. Sci.* 88: 6442–6446, 1991.

[65700] 4030. Lindner, T. H.; Njolstad, P. R.; Horikawa, Y.; Bostad, L.; Bell, G. I.; Sovik, O.: A novel syndrome of diabetes mellitus, renal dysfunction and genital malformation associated with a partial deletion of the pseudo-POU domain of hepatocyte nuclear factor-1-beta. *Hum. Molec. Genet.* 8:

2001–2008, 1999.

- [65701] 4031. Anderson, L. A.; Hall, J. M.; Lee, M. K.; Lebo, R. V.; King, M.C.: Polymorphisms near chromosome 1cen. (Abstract) Cytogenet. CellGenet. 51: 951, 1989.
- [65702] 4032. Gendler, S. J.; Lancaster, C. A.; Taylor–Padimitriou, J.; Duhig, T.; Peat, N.; Burchell, J.; Pemberton, L.; Lalani, E.–N.; Wilson, D.: Molecular cloning and expression of human tumor–associated polymorphic epithelial mucin. J. Biol. Chem. 265: 15286–15293, 1990.
- [65703] 4033. Karlsson, S.; Swallow, D. M.; Griffiths, B.; Corney, G.; Hopkinson, D. A.; Dawnay, A.; Cartron, J. P.: A genetic polymorphism of a human urinary mucin. Ann. Hum. Genet. 47: 263–269, 1983.
- [65704] 4034. Kingsmore, S. F.; Spicer, A. P.; Gendler, S. J.; Seldin, M. F.: Genetic mapping of the tumor–associated mucin 1 gene on mouse chromosome 3. Mammalian Genome 6: 378, 1995.
- [65705] 4035. Lan, M. S.; Batra, S. K.; Qi, W.–N.; Metzgar, R. S.; Hollingsworth, M. A.: Cloning and sequencing of a human pancreatic tumor mucin cDNA. J. Biol. Chem. 265: 15294–15299, 1990.
- [65706] 4036. Ligtenberg, M. J.; Gennissen, A. M.; Vos, H. L.; Hilkens, J.: A single nucleotide polymorphism in an exon



dictates allele dependent differential splicing of episialin mRNA. *Nucleic Acids Res.* 19:297–301, 1991.

- [65707] 4037. Ligtenberg, M. J. L.; Vos, H. L.; Gennissen, A. M. C.; Hilkens, J.: Episialin, a carcinoma-associated mucin, is generated by a polymorphic gene encoding splice variants with alternative amino termini. *J. Biol. Chem.* 265: 5573–5578, 1990.
- [65708] 4038. Middleton-Price, H.; Gendler, S.; Malcolm, S.: Close linkage of PUM and SPTA within chromosome band 1q21. *Ann. Hum. Genet.* 52: 273–278, 1988.
- [65709] 4039. Pratt, W. S.; Islam, I.; Swallow, D. M.: Two additional polymorphisms within the hypervariable MUC1 gene: association of alleles either side of the VNTR region. *Ann. Hum. Genet.* 60: 21–28, 1996.
- [65710] 4040. Silva, F.; Carvalho, F.; Peixoto, A.; Seixas, M.; Almeida, R.; Carneiro, F.; Mequita, P.; Figueiredo, C.; Nogueira, C.; Swallow, D. M.; Amorim, A.; David, L.: MUC1 gene polymorphism in the gastric carcinogenesis pathway. *Europ. J. Hum. Genet.* 9: 548–552, 2001.
- [65711] 4041. Swallow, D. M.; Gendler, S.; Griffiths, B.; Corney, G.; Taylor-Papadimitriou, J.; Bramwell, M. E.: The human tumour-associated epithelial mucins are coded by an expressed hypervariable gene locus PUM. *Nature* 328:82–84,

1987.

- [65712] 4042. Swallow, D. M.; Gendler, S.; Griffiths, B.; Kearney, A.; Povey, S.; Sheer, D.; Palmer, R. W.; Taylor-Papadimitriou, J.: The hypervariable gene locus PUM, which codes for the tumour associated epithelial mucins, is located on chromosome 1, within the region 1q21-24. (Abstract) Cytogenet. Cell Genet. 46: 701, 1987.
- [65713] 4043. Swallow, D. M.; Gendler, S.; Griffiths, B.; Kearney, A.; Povey, S.; Sheer, D.; Palmer, R. W.; Taylor-Papadimitriou, J.: The hypervariable gene locus PUM, which codes for the tumour associated epithelial mucins, is located on chromosome 1, within the region 1q21-q24. Ann. Hum. Genet. 51: 289-294, 1987.
- [65714] 4044. Swallow, D. M.; Griffiths, B.; Noades, J.; Corney, G.: Linkage between the expressed hypervariable gene locus PUM and the gene coding for the Duffy blood group FY. Ann. Hum. Genet. 52: 269-271, 1988.
- [65715] 4045. Cole, S. P. C.; Bhardwaj, G.; Gerlach, J. H.; Mackie, J. E.; Grant, C. E.; Almquist, K. C.; Stewart, A. J.; Kurz, E. U.; Duncan, A. M. V.; Deeley, R. G.: Overexpression of a transporter gene in a multidrug-resistant human lung cancer cell line. Science 258: 1650-1654, 1992.
- [65716] 4046. Conrad, S.; Kauffmann, H.-M.; Ito, K.; Deeley, R. G.;

Cole, S.P. C.; Schrenk, D.: Identification of human multidrug resistance protein 1 (MRP1) mutations and characterization of a G671V substitution. *J.Hum. Genet.* 46: 656–663, 2001.

[65717] 4047. Grant, C. E.; Kurz, E. U.; Cole, S. P. C.; Deeley, R. G.: Analysis of the intron–exon organization of the human multidrug–resistance protein gene (MRP) and alternative splicing of its mRNA. *Genomics* 45:368–378, 1997.

[65718] 4048. Lorico, A.; Bertola, A.; Baum, C.; Fodstad, O.; Rappa, G.: Role of multidrug resistance protein 1 in protection from heavy metal oxyanions: investigations in vitro and in Mrp1–deficient mice. *Biochem. Biophys. Res. Commun.* 291: 617–622, 2002.

[65719] 4049. Robbiani, D. F.; Finch, R. A.; Jager, D.; Muller, W. A.; Sartorelli, A. C.; Randolph, G. J.: The leukotriene C4 transporter MRP1 regulates CCL19 (MIP–3–beta, ELC)–dependent mobilization of dendritic cells to lymph nodes. *Cell* 103: 757–768, 2000.

[65720] 4050. Schultz, M. J.; Wijnholds, J.; Peppelenbosch, M. P.; Vervoordeldonk, M. J. B. M.; Speelman, P.; van Deventer, S. J. H.; Borst, P.; van der Poll, T.: Mice lacking the multidrug resistance protein 1 are resistant to *Streptococcus pneumoniae*–induced pneumonia. *J. Immun.* 166:4059–4064,

2001.

- [65721] 4051.Zaman, G. J. R.; Flens, M. J.; van Leusden, M. R.; de Haas, M.; Mulder, H. S.; Lankelma, J.; Pinedo, H. M.; Scheper, R. J.; Baas, F.; Broxterman, H. J.; Borst, P.: The human multidrug resistance-associated protein MRP is a plasma membrane drug-efflux pump. *Proc. Nat. Acad. Sci.* 91: 8822–8826, 1994.
- [65722] 4052.Polymeropoulos, M. H.; Xiao, H.; Glodek, A.; Gorski, M.; Adams, M. D.; Moreno, R. F.; Fitzgerald, M. G.; Venter, J. C.; Merrill, C. R.: Chromosomal assignment of 46 brain cDNAs. *Genomics* 12: 492–496, 1992.
- [65723] 4053.Riess, O.; Epplen, C.; Siedlaczek, I.; Epplen, J. T.: Chromosomal assignment of the human smg GDP dissociation stimulator gene to human chromosome 4q21–q25. *Hum. Genet.* 92: 629–630, 1993.
- [65724] 4054.Doria, A.; Caldwell, J. S.; Ji, L.; Reynet, C.; Rich, S. S.; Weremowicz, S.; Morton, C. C.; Warram, J. H.; Kahn, C. R.; Krolewski, A. S.: Trinucleotide repeats at the rad locus: allele distributions in NIDDM and mapping to a 3-cM region on chromosome 16q. *Diabetes* 44: 243–247, 1995.
- [65725] 4055.Orho, M.; Carlsson, M.; Kanninen, T.; Groop, L. C.: Polymorphism at the rad gene is not associated with NIDDM in Finns. *Diabetes* 45: 429–433, 1996.

- [65726] 4056.Reynet, C.; Kahn, C. R.: Rad: a member of the Ras family overexpressed in muscle of type II diabetic humans. *Science* 262: 1441–1444, 1993.
- [65727] 4057.le Gallic, L.; Fort, P.: Structure of the human ARHG locus encoding the Rho/Rac-like RhoG GTPase. *Genomics* 42: 157–160, 1997.
- [65728] 4058.Taviaux, S. A.; Vincent, S.; Fort, P.; Demaille, J. G.: Localization of ARHG, a member of the RAS homolog gene family, to 11p15.5–11p15.4 by fluorescence in situ hybridization. *Genomics* 16: 788–790, 1993.
- [65729] 4059.Vincent, S.; Jeanteur, P.; Fort, P.: Growth-regulated expression of rhoG, a new member of the ras homolog gene family. *Molec. Cell Biol.* 12: 3138–3148, 1992.
- [65730] 4060.Allan, B. B.; Moyer, B. D.; Balch, W. E.: Rab1 recruitment of p115 into a cis-SNARE complex: programming budding COPII vesicles for fusion. *Science* 289: 444–448, 2000.
- [65731] 4061.Wedemeyer, N.; Lengeling, A.; Ronsiek, M.; Korthaus, D.; Baer, K.; Wuttke, M.; Jockusch, H.: YAC contigs of the Rab1 and wobbler (wr) spinal muscular atrophy gene region on proximal mouse chromosome 11 and of the homologous region on human chromosome 2p. *Genomics* 32:447–454, 1996.

- [65732] 4062.Rousseau–Merck, M. F.; Zahraoui, A.; Touchot, N.; Tavitian, A.;Berger, R.: Chromosome assignment of four RAS–related RAB genes. *Hum.Genet.* 86: 350–354, 1991.
- [65733] 4063.Barbosa, M. D. F. S.; Johnson, S. A.; Achey, K.; Gutierrez, M.;Wakeland, E. K.; Zerial, M.; Kingsmore, S. F.: The Rab protein family:genetic mapping of six Rab genes in the mouse. *Genomics* 30: 439–444,1995.
- [65734] 4064.Rousseau–Merck, M.–F.; Zahraoui, A.; Touchot, N.; Tavitian, A.;Berger, R.: Chromosome assignment of four RAS–related RAB genes. *Hum.Genet.* 86: 350–354, 1991.
- [65735] 4065.Bucci, C.; Parton, R. G.; Mather, I. H.; Stunnenberg, H.; Simons,K.; Hoflack, B.; Zerial, M.: The small GTPase rab5 functions as aregulatory factor in the early endocytic pathway. *Cell* 70: 715–728,1992.
- [65736] 4066.Stenmark, H.; Vitale, G.; Ullrich, O.; Zerial, M.: Rabaptin–5is a direct effector of the small GTPase Rab5 in endocytic membranefusion. *Cell* 83: 423–432, 1995.
- [65737] 4067.Xiao, G.–H.; Shoarinejad, F.; Jin, F.; Golemis, E. A.; Yeung, R.S.: The tuberous sclerosis 2 gene product, tuberlin, functions asa Rab5 GTPase activating protein (GAP) in modulating endocytosis. *J.Biol. Chem.* 272: 6097–6100, 1997.
- [65738] 4068.Korenberg, J. R.; Chen, X.–N.; Adams, M. D.; Venter,

J. C.: Towarda cDNA map of the human genome. *Genomics* 29: 364–370, 1995.

[65739] 4069. Wilson, D. B.; Wilson, M. P.: Identification and sub-cellular localization of human rab5b, a new member of the ras-related superfamily of GTPases. *J. Clin. Invest.* 89: 996–1005, 1992.

[65740] 4070. Asha, H.; de Ruiter, N. D.; Wang, M.-G.; Hariharan, I. K.: The Rap1 GTPase functions as a regulator of morphogenesis in vivo. *EMBO J.* 18: 605–615, 1999.

[65741] 4071. Boussiotis, V. A.; Freeman, G. J.; Berezovskaya, A.; Barber, D. L.; Nadler, L. M.: Maintenance of human T cell anergy: blocking of IL-2 gene transcription by activated Rap1. *Science* 278: 124–128, 1997.

[65742] 4072. Kitayama, H.; Sugimoto, Y.; Matsuzaki, T.; Ikawa, Y.; Noda, M.: A ras-related gene with transformation suppressor activity. *Cell* 56:77–84, 1989.

[65743] 4073. Knox, A. L.; Brown, N. H.: Rap1 GTPase regulation of adherens junction positioning and cell adhesion. *Science* 295: 1285–1288, 2002.

[65744] 4074. Muller-Ladner, U.; Judex, M.; Ballhorn, W.; Kullmann, F.; Distler, O.; Schlottmann, K.; Gay, R. E.; Scholmerich, J.; Gay, S.: Activation of the IL-4 STAT pathway in rheumatoid synovium. *J. Immun.* 164:3894–3901, 2000.

- [65745] 4075.Aita, N.; Ishii, K.; Akamatsu, Y.; Ogasawara, Y.; Tanabe, S.:Cloning and expression of human liver rhodanese cDNA. *Biochem. Biophys.Res. Commun.* 231: 56–60, 1997.
- [65746] 4076.Cagianut, B.; Rhyner, K.; Furrer, W.; Schnebli, H. P.: Thiosulphate–sulphurtransferase (rhodanese) deficiency in Leber's hereditary optic atrophy.(Letter) *Lancet* II: 981–982, 1981.
- [65747] 4077.Nikoskelainen, E.: New aspects of the genetic, etiology, andclinical puzzle of Leber's disease. *Neurology* 34: 1482–1484, 1984.
- [65748] 4078.Scott, E. M.; Wright, R. C.: Genetic polymorphism of rhodanese from human erythrocytes. *Am. J. Hum. Genet.* 32: 112–114, 1980.
- [65749] 4079.Weng, L.; Heinrikson, R. L.; Westley, J.: Active site cysteinyland arginyl residues of rhodanese. *J. Biol. Chem.* 253: 8109–8119,1978.
- [65750] 4080.Whitehouse, D. B.; Pilz, A. J.; Porta, G.; Hopkinson, D. A.: Rhodanese isozymes in human tissues. *Ann. Hum. Genet.* 52: 1–10, 1988.
- [65751] 4081.Whitehouse, D. B.; Poole, C. J. M.; Kind, P. R. N.; Hopkinson,D. A.: Rhodanese isozymes in three subjects with Leber's optic neuropathy. *J.Med. Genet.* 26: 113–115,



1989.

- [65752] 4082.Alloway, P. G.; Howard, L.; Dolph, P. J.: The formation of stablerhodopsin-arrestin complexes induces apoptosis and photoreceptor celldegeneration. *Neuron* 28: 129–138, 2000.
- [65753] 4083.Byrne, J. A.; Smith, P. J.: The 11p15.5 ribonucleotide reductaseM1 subunit locus is not imprinted in Wilms' tumour and hepatoblastoma. *Hum.Genet.* 91: 275–277, 1993.
- [65754] 4084.Bouchard, B.; Del Marmol, V.; Jackson, I. J.; Cherif, D.; Dubertret,L.: Molecular characterization of a human tyrosinase-related-protein-2cDNA: patterns of expression in melanocytic cells. *Europ. J. Biochem.* 219:127–134, 1994.
- [65755] 4085.Budd, P. S.; Jackson, I. J.: Structure of the mouse tyrosinase-relatedprotein-2/dopachrome tautomerase (Tyrp2/Dct) gene and sequence oftwo novel slaty alleles. *Genomics* 29: 35–43, 1995.
- [65756] 4086.Cassady, J. L.; Sturm, R. A.: Sequence of the human dopachrometautomerase-encoding TRP-2 cDNA. *Gene* 143: 295–298, 1994.
- [65757] 4087.Jackson, I. J.; Chambers, D. M.; Tsukamoto, K.; Copeland, N. G.;Gilbert, D. J.; Jenkins, N. A.; Hearing, V.: A

second tyrosinase-related protein, TRP-2, maps to and is mutated at the mouse slaty locus. EMBOJ. 11: 527-535, 1992.

[65758] 4088. Khong, H. T.; Rosenberg, S. A.: Pre-existing immunity to tyrosinase-related protein (TRP)-2, a new TRP-2 isoform, and the NY-ESO-1 melanoma antigen in a patient with a dramatic response to immunotherapy. J. Immun. 168:951-956, 2002.

[65759] 4089. Kwon, B. S.: Pigmentation genes: the tyrosinase gene family and the pmel 17 gene family. J. Invest. Derm. 100: 134S-140S, 1993.

[65760] 4090. Sturm, R. A.; Baker, E.; Sutherland, G. R.: Assignment of the tyrosinase-related protein-2 gene (TYRP2) to human chromosome 13q31-q32 by fluorescence in situ hybridization: extended synteny with mouse chromosome 14. Genomics 21: 293-296, 1994.

[65761] 4091. Sturm, R. A.; O'Sullivan B. J.; Box, N. F.; Smith, A. G.; Smit, S. E.; Puttick, E. R. J.; Parsons, P. G.; Dunn, I. S.: Chromosomal structure of the human TYRP1 and TYRP2 loci and comparison of the tyrosinase-related protein gene family. Genomics 29: 24-34, 1995.

[65762] 4092. Ouafik, L'H.; Stoffers, D. A.; Campbell, T. A.; Johnson, R. C.; Bloomquist, B. T.; Mains, R. E.; Eipper, B. A.: The

multifunctionalpeptidylglycine alpha-amidating monooxygenase gene: exon/intron organizationof catalytic, processing, and routing domains. *Molec. Endocr.* 6:1571–1584, 1992.

- [65763] 4093.King, C. R.; Schimke, R. N.; Arthur, T.; Davoren, B.; Collins,D.: Proximal 3p deletion in renal cell carcinoma cells from a patientwith von Hippel–Lindau disease. *Cancer Genet. Cytogenet.* 27: 345–348,1987.
- [65764] 4094.Li, F. P.; Decker, H.–J. H.; Zbar, B.; Stanton, V. P., Jr.; Kovacs,G.; Seizinger, B. R.; Aburatani, H.; Sandberg, A. A.; Berg, S.; Hosoe,S.; Brown, R. S.: Clinical and genetic studies of renal cell carcinomasin a family with a constitutional chromosome 3;8 translocation: geneticsof familial renal carcinoma. *Ann. Intern. Med.* 118: 106–111, 1993.
- [65765] 4095.Neel, B. G.; Jhanwar, S. C.; Chaganti, R. S. K.; Hayward, W. S.: Two human c–onc genes are located on the long arm of chromosome8. *Proc. Nat. Acad. Sci.* 79: 7842–7846, 1982.
- [65766] 4096.Ohta, M.; Inoue, H.; Cotticelli, M. G.; Kastury, K.; Baffa, R.;Palazzo, J.; Siprashvili, Z.; Mori, M.; McCue, P.; Druck, T.; Croce,C. M.; Huebner, K.: The FHIT gene, spanning the chromosome 3p14.2fragile site and renal carcinoma–associated t(3;8) breakpoint, isabnormal in diges–

tive tract cancers. *Cell* 84: 587–597, 1996.

[65767] 4097. Roses, A. D.: From genes to mechanisms to therapies: lessons to be learned from neurological disorders. *Nature Med.* 2: 267–269, 1996.

[65768] 4098. Wood, J. D.; Yuan, J.; Margolis, R. L.; Colomer, V.; Duan, K.; Kushi, J.; Kaminsky, Z.; Kleiderlein, J. J., Jr.; Sharp, A. H.; Ross, C. A.: Atrophin-1, the DRPLA gene product, interacts with two families of WW domain-containing proteins. *Molec. Cell. Neurosci.* 11: 149–160, 1998.

[65769] 4099. Aksoy, M.; Erdem, S.: Combination of hereditary elliptocytosis and heterozygous beta-thalassemia: a family study. *J. Med. Genet.* 5: 298–301, 1968.

[65770] 4100. Alloisio, N.; Dorleac, E.; Delaunay, J.; Girot, R.; Galand, C.; Boivin, P.: A shortened variant of red cell membrane protein 4.1. *Blood* 60: 265–267, 1982.

[65771] 4101. Alloisio, N.; Dorleac, E.; Girot, R.; Delaunay, J.: Analysis of red cell membrane in a family with hereditary elliptocytosis: total or partial absence of protein 4.1. *Hum. Genet.* 59: 68–71, 1981.

[65772] 4102. Alloisio, N.; Morle, L.; Dorleac, E.; Gentilhomme, O.; Bachir, D.; Guetarni, D.; Colonna, P.; Bost, M.; Zouaoui, Z.; Roda, L.; Roussel, D.; Delaunay, J.: The heterozygous form

of 4.1(-) hereditary elliptocytosis[the 4.1(-) trait]. Blood 65: 46-51, 1985.

[65773] 4103.Bahary, N.; Zorich, G.; Pachter, J. E.; Leibel, R. L.; Friedman,J. M.: Molecular genetic linkage maps of mouse chromosomes 4 and

[65774] 4104.Genomics 11: 33-47, 1991.6. Baklouti, F.; Huang, S.-C.; Vulliamy, T. J.; Delaunay, J.; Benz,E. J., Jr.: Organization of the human protein 4.1 genomic locus:new insights into the tissue-specific alternative splicing of thepre-mRNA. Genomics 39: 289-302, 1997.

[65775] 4105.Bannerman, R. M.; Renwick, J. H.: The hereditary elliptocytoses:clinical and linkage data. Ann. Hum. Genet. 26: 23-38, 1962.

[65776] 4106.Clarke, C. A.; Donohoe, W. T. A.; Finn, R.; McConnell, R. B.; Sheppard,P. M.; Nicol, D. S. H.: Data on linkage in man: ovalocytosis, sicklingand the Rhesus blood group complex. Ann. Hum. Genet. 24: 283-287,1960.

[65777] 4107.Conboy, J.; Kan, Y. W.; Shohet, S. B.; Mohandas, N.: Molecularcloning of protein 4.1, a major structural element of the human erythrocytemembrane skeleton. Proc. Nat. Acad. Sci. 83: 9512-9516, 1986.

[65778] 4108.Conboy, J.; Marchesi, S.; Kim, R.; Agre, P.; Kan, Y.

W.; Mohandas, N.: Molecular analysis of insertion/deletion mutations in protein 4.1 in elliptocytosis. II. Determination of molecular genetic origin of rearrangements. *J. Clin. Invest.* 86: 524–530, 1990.

[65779] 4109. Conboy, J.; Mohandas, N.; Tchernia, G.; Kan, Y. W.: Molecular basis of hereditary elliptocytosis due to protein 4.1 deficiency. *New Eng. J. Med.* 315: 680–685, 1986.

[65780] 4110. Conboy, J. G.: Structure, function, and molecular genetics of erythroid membrane skeletal protein 4.1 in normal and abnormal red blood cells. *Seminars Hemat.* 30: 58–73, 1993.

[65781] 4111. Conboy, J. G.; Chan, J.; Mohandas, N.; Kan, Y. W.: Multiple protein 4.1 isoforms produced by alternative splicing in human erythroid cells. *Proc. Nat. Acad. Sci.* 85: 9062–9065, 1988.

[65782] 4112. Conboy, J. G.; Chasis, J. A.; Winardi, R.; Tchernia, G.; Kan, Y. W.; Mohandas, N.: An isoform-specific mutation in the protein 4.1 gene results in hereditary elliptocytosis and complete deficiency of protein 4.1 in erythrocytes but not in nonerythroid cells. *J. Clin. Invest.* 91: 77–82, 1993.

[65783] 4113. Conboy, J. G.; Mohandas, N.; Wang, C.; Tchernia, G.; Shohet, S. B.; Kan, Y. W.: Molecular cloning and characterization of the gene coding for red cell membrane skeletal

protein 4.1. (Abstract) Blood 66(suppl. 1): 31A, 1985.

- [65784] 4114.Cook, P. J. L.; Noades, J. E.; Newton, M. S.; de Mey, R.: On the orientation of the Rh:E1-1 linkage group. Ann. Hum. Genet. 41:157-162, 1977.
- [65785] 4115.Lane, P. W.: Association of megacolon with two recessive spotting genes in the mouse. J. Hered. 57: 29-31, 1966.
- [65786] 4116.Matsushima, Y.; Shinkai, Y.; Kobayashi, Y.; Sakamoto, M.; Kunieda, T.; Tachibana, M.: A mouse model of Waardenburg syndrome type 4 with a new spontaneous mutation of the endothelin-B receptor gene. Mammalian Genome 13: 30-35, 2002.
- [65787] 4117.Metallinos, D. L.; Bowling, A. T.; Rine, J.: A missense mutation in the endothelin-B receptor gene is associated with lethal whitefoal syndrome: an equine version of Hirschsprung disease. Mammalian Genome 9: 426-431, 1998.
- [65788] 4118.Nakamuta, M.; Takayanagi, R.; Sakai, Y.; Sakamoto, S.; Hagiwara, H.; Mizuno, T.; Saito, Y.; Hirose, S.; Yamamoto, M.; Nawata, H.: Cloning and sequence analysis of a cDNA encoding human non-selective type of endothelin receptor. Biochem. Biophys. Res. Commun. 177:34-39, 1991.

- [65789] 4119.Ogawa, Y.; Nakao, K.; Arai, H.; Nakagawa, O.; Hosoda, K.; Suga,S.; Nakanishi, S.; Imura, H.: Molecular cloning of a non–isopeptide–selectivehuman endothelin receptor. *Biochem. Biophys. Res. Commun.* 178: 248–255,1991.
- [65790] 4120.Pao, M. M.; Tsutsumi, M.; Liang, G.; Uzvolgyi, E.; Gonzales, F.A.; Jones, P. A.: The endothelin receptor B (EDNRB) promoter displaysheterogeneous, site specific methylation patterns in normal and tumorcells. *Hum. Molec. Genet.* 10: 903–910, 2001.
- [65791] 4121.Puffenberger, E. G.; Hosoda, K.; Washington, S. S.; Nakao, K.;deWit, D.; Yanagisawa, M.; Chakravarti, A.: A missense mutation ofthe endothelin–B receptor gene in multigenic Hirschsprung's disease. *Cell* 79:1257–1266, 1994.
- [65792] 4122.Santschi, E. M.; Purdy, A. K.; Valberg, S. J.; Vrotsos, P. D.;Kaese, H.; Mickelson, J. R.: Endothelin receptor B polymorphism associatedwith lethal white foal syndrome in horses. *Mammalian Genome* 9: 306–309,1998.
- [65793] 4123.Shin, M. K.; Levorse, J. M.; Ingram, R. S.; Tilghman, S. M.:The temporal requirement for endothelin receptor–B signalling duringneural crest development. *Nature* 402: 496–501, 1999.



- [65794] 4124. Shin, M. K.; Russell, L. B.; Tilghman, S. M.: Molecular characterization of four induced alleles at the Ednrb locus. Proc. Nat. Acad. Sci. 94:13105–13110, 1997.
- [65795] 4125. Svensson, P.-J.; Anvret, M.; Molander, M.-L.; Nordenskjold, A.: Phenotypic variation in a family with mutations in two Hirschsprung-related genes (RET and endothelin receptor B). Hum. Genet. 103: 145–148, 1998.
- [65796] 4126. Syrris, P.; Carter, N. D.; Patton, M. A.: Novel non-sense mutation of the endothelin-B receptor gene in a family with Waardenburg-Hirschsprung disease. Am. J. Med. Genet. 87: 69–71, 1999.
- [65797] 4127. Takayanagi, R.; Ohnaka, K.; Takasaki, C.; Ohashi, M.; Nawata, H.: Multiple subtypes of endothelin receptors in porcine tissues: characterization by ligand binding, affinity labeling and regional distribution. Regul. Pept. 32: 23–37, 1991.
- [65798] 4128. Tanaka, H.; Moroi, K.; Iwai, J.; Takahashi, H.; Ohnuma, N.; Hori, S.; Takimoto, M.; Nishiyama, M.; Masaki, T.; Yanagisawa, M.; Sekiya, S.; Kimura, S.: Novel mutations of the endothelin B receptor gene in patients with Hirschsprung's disease and their characterization. J. Biol. Chem. 273: 11378–11383, 1998.
- [65799] 4129. Vane, J. R.: Endothelins come home to roost. Nature

348: 673,1990.

- [65800] 4130.Verheij, J. B. G. M.; Kunze, J.; Osinga, J.; van Essen, A. J.;Hofstra, R. M. W.: ABCD syndrome is caused by a homozygous mutation in the EDNRB gene. *Am. J. Med. Genet.* 108: 223–225, 2002.
- [65801] 4131.Yang, G. C.; Croaker, D.; Zhang, A. L.; Manglick, P.; Cartmill,T.; Cass, D.: A dinucleotide mutation in the endothelin-B receptorgene is associated with lethal white foal syndrome (LWFS); a horsevariant of Hirschsprung disease (HSCR). *Hum. Molec. Genet.* 7: 1047–1052,1998.
- [65802] 4132.Hanks, M.; Wurst, W.; Anson–Cartwright, L.; Auerbach, A. B.; Joyner,A. L.: Rescue of the En–1 mutant phenotype by replacement of En–1with En–2. *Science* 269: 679–682, 1995.
- [65803] 4133.Johnson, R. L.; Tabin, C. J.: Molecular models for vertebratelimb development. *Cell* 90: 979–990, 1997.
- [65804] 4134.Kohler, A.; Logan, C.; Joyner, A. L.; Muenke, M.: Regional assignmentof the human homeobox–containing gene EN1 to chromosome 2q13–q21. *Genomics* 15:233–235, 1993.
- [65805] 4135.Logan, C.; Hanks, M. C.; Noble–Topham, S.; Nalainathan, D.; Provart,N. J.; Joyner, A. L.: Cloning and sequence comparison of the mouse,human, and chicken en–

grailed genes reveal potential functional domains and regulatory regions. *Dev. Genet.* 13: 345–358, 1992.

[65806] 4136. Logan, C.; Willard, H. F.; Rommens, J. M.; Joyner, A. L.: Chromosomal localization of the human homeo box-containing genes, EN1 and EN2. *Genomics* 4:206–209, 1989.

[65807] 4137. Loomis, C. A.; Harris, E.; Michaud, J.; Wurst, W.; Hanks, M.; Joyner, A. L.: The mouse engrailed-1 gene and ventral limb patterning. *Nature* 382:360–363, 1996.

[65808] 4138. Martin, G. R.; Richman, M.; Reinsch, S.; Nadeau, J. H.; Joyner, A.: Mapping of the two mouse engrailed-like genes: close linkage of En-1 to dominant hemimelia (Dh) on chromosome 1 and of En-2 to hemimelic extra-toes (Hx) on chromosome 5. *Genomics* 6: 302–308, 1990.

[65809] 4139. Matsui, T.; Hirai, M.; Hirano, M.; Kurosawa, Y.: The HOX complex neighbored by the EVX gene, as well as two other homeobox-containing genes, the GBX-class and the EN class, are located on the same chromosomes 2 and 7 in humans. *FEBS Lett.* 336: 107–110, 1993.

[65810] 4140. Wurst, W.; Auerbach, A. B.; Joyner, A. L.: Multiple developmental defects in Engrailed-1 mutant mice: an early mid-hindbrain deletion and patterning defects in forelimbs and sternum. *Development* 120:2065–2075,

1994.

- [65811] 4141.Kohl, S.; Baumann, B.; Rosenberg, T.; Kellner, U.; Lorenz, B.;Vadala, M.; Jacobson, S. G.; Wissinger, B.: Mutations in the cone photoreceptor G-protein alpha-subunit gene GNAT2 in patients with achromatopsia. *Am. J. Hum. Genet.* 71: 422–425, 2002.
- [65812] 4142.Morris, T. A.; Fong, S.-L.: Characterization of the gene encoding human cone transducin alpha-subunit (GNAT2). *Genomics* 17: 442–448, 1993.
- [65813] 4143.Sparkes, R. S.; Cohn, V. H.; Cire-Eversole, P.; Blatt, C.; Amatruda, T. T.; Weiner, L. P.; Nesbitt, M.; Reed, R. R.; Lochrie, M. A.; Fournier, R. E. K.; Simon, M. I.: Mapping of genes encoding the subunits of guanine nucleotide-binding proteins (G-proteins) in the mouse. (Abstract) *Cytogenet. Cell Genet.* 46: 696 only, 1987.
- [65814] 4144.Chipev, C. C.; Korge, B. P.; Markova, N.; Bale, S. J.; DiGiovanna, J. J.; Compton, J. G.; Steinert, P. M.: A leucine-to-proline mutation in the H1 subdomain of keratin 1 causes epidermolytic hyperkeratosis. *Cell* 70:821–828, 1992.
- [65815] 4145.Compton, J. G.: Epidermal disease: faulty keratin filaments take their toll. *Nature Genet.* 6: 6–7, 1994.
- [65816] 4146.Compton, J. G.; DiGiovanna, J. J.; Santucci, S. K.;

Kearns, K.S.; Amos, C. I.; Abangan, D. L.; Korge, B. P.; McBride, O. W.; Steinert, P. M.; Bale, S. J.: Linkage of epidermolytic hyperkeratosis to the type II keratin gene cluster on chromosome 12q. *Nature Genet.* 1:301–305, 1992.

[65817] 4147. Fraser, R. D. B.; MacRae, T. P.; Suzuki, E.: Structure of the  $\alpha$ -keratin microfibril. *J. Molec. Biol.* 108: 435–452, 1976.

[65818] 4148. Kimonis, V.; DiGiovanna, J. J.; Yang, J.-M.; Doyle, S. Z.; Bale, S. J.; Compton, J. G.: A mutation in the V1 end domain of keratin 1 in non-epidermolytic palmar-plantar keratoderma. *J. Invest. Derm.* 103:764–769, 1994.

[65819] 4149. Lee, L. D.; Ludwig, K.; Baden, H. P.: Matrix proteins of human hair as a tool for identification of individuals. *Forensic Sci.* 11:115–121, 1978.

[65820] 4150. Lessin, S. R.; Huebner, K.; Isobe, M.; Croce, C. M.; Steinert, P. M.: Chromosomal mapping of human keratin genes: evidence of non-linkage. *J. Invest. Derm.* 91: 572–578, 1988.

[65821] 4151. Nadeau, J. H.: Personal Communication. Bar Harbor, Me. 7/29/1987.

[65822] 4152. Le Marchand-Brustel, Y.; Gremeaux, T.; Ballotti, R.; van Obberghen, E.: Insulin receptor tyrosine kinase is defective in skeletal muscle of insulin-resistant obese mice.

Nature 315: 676–679, 1985.

[65823] 4153.Leme, C. E.; Wajchenberg, B. L.; Lerario, A. C.; Goldman, J.;Borges, J. L. C.: Acanthosis nigricans, hirsutism, insulin resistanceand insulin receptor defect. Clin. Endocr. 17: 43–49, 1982.

[65824] 4154.Longo, N.; Langley, S. D.; Griffin, L. D.; Elsas, L. J.: Activationof glucose transport by a natural mutation in the human insulin receptor. Proc.Nat. Acad. Sci. 90: 60–64, 1993.

[65825] 4155.Longo, N.; Langley, S. D.; Griffin, L. D.; Elsas, L. J., II:Reduced mRNA and a nonsense mutation in the insulin-receptor geneproduce heritable severe insulin resistance. Am. J. Hum. Genet. 50:998–1007, 1992.

[65826] 4156.Maassen, J. A.; Klinkhamer, M. P.; Odink, R. J. H.; Sips, H.;van der Zon, G. C. M.; Wieringa, T.; Krans, H. M. J.; Moller, W.:Improper expression of insulin receptors on fibroblasts from a leprechaunpatient. Europ. J. Biochem. 172: 725–729, 1988.

[65827] 4157.Mariani, S.; Pedone, A.; Meschi, F.; Di Natale, B.; Caputo, R.;Broggi, U.; Chiumello, G.: Insulin resistance in a child with acanthosisnigricans type A. Acta Paediat. Scand. 71: 667–670, 1982.

[65828] 4158.Michael, M. D.; Kulkarni, R. N.; Postic, C.; Previs, S.

F.; Shulman, G. I.; Magnuson, M. A.; Kahn, C. R.: Loss of insulin signaling in hepatocytes leads to severe insulin resistance and progressive hepatic dysfunction. *Molec. Cell* 6: 87–97, 2000.

[65829] 4159. Moller, D. E.; Cohen, O.; Yamaguchi, Y.; Assiz, R.; Grigorescu, F.; Eberle, A.; Morrow, L. A.; Moses, A. C.; Flier, J. S.: Prevalence of mutations in the insulin receptor gene in subjects with features of the type A syndrome of insulin resistance. *Diabetes* 43: 247–255, 1994.

[65830] 4160. Moller, D. E.; Flier, J. S.: Insulin resistance—mechanisms, syndromes, and implications. *New Eng. J. Med.* 325: 938–948, 1991.

[65831] 4161. Moller, D. E.; Flier, J. S.: Detection of an alteration in the insulin-receptor gene in a patient with insulin resistance, acanthosis nigricans, and the polycystic ovary syndrome (type A insulin resistance). *New Eng. J. Med.* 319: 1526–1529, 1988.

[65832] 4162. Moller, D. E.; Yokota, A.; Ginsberg-Fellner, F.; Flier, J. S.: Functional properties of a naturally occurring trp(1200)-to-ser(1200) mutation of the insulin receptor. *Molec. Endocr.* 4: 1183–1191, 1990.

[65833] 4163. Moller, D. E.; Yokota, A.; Pazianos, A.; Flier, J. S.: A missense mutation in one allele of the tyrosine kinase do-

main of the insulinreceptor gene is associated with dominantly inherited insulin resistance.(Abstract) Clin. Res. 38: 435A only, 1990.

- [65834] 4164.Moller, D. E.; Yokota, A.; White, M. F.; Pazianos, A. G.; Flier,J. S.: A naturally occurring mutation of insulin receptor alanine1134 impairs tyrosine kinase function and is associated with dominantlyinherited insulin resistance. J. Biol. Chem. 265: 14979–14985, 1990.
- [65835] 4165.Moncada, V. Y.; Hedo, J. A.; Serrano–Rios, M.; Taylor, S. I.:Insulin–receptor biosynthesis in cultured lymphocytes from an insulin–resistantpatient (Rabson–Mendenhall syndrome): evidence for defect before insertionof receptor into plasma membrane. Diabetes 35: 802–807, 1986.
- [65836] 4166.Norton, K. I.; Glicklich, M.; Kupchik, G.; Gray, C. E.; Ludman,M.: Leprechaunism: a case report with radiographic features. Dysmorph.Clin. Genet. 4: 57–62, 1990.
- [65837] 4167.Odawara, M.; Kadowaki, T.; Yamamoto, R.; Shibasaki, Y.; Tobe,K.; Accili, D.; Bevins, C.; Mikami, Y.; Matsuura, N.; Akanuma, Y.;Takaku, F.; Taylor, S. I.; Kasuga, M.: Human diabetes associatedwith a mutation in the tyrosine kinase domain of the insulin receptor. Science 245:66–68, 1989.
- [65838] 4168.Ojamaa, K.; Hedo, J. A.; Roberts, C. T., Jr.; Moncada,



V. Y.;Gorden, P.; Ullrich, A.; Taylor, S. I.: Defects in human insulinreceptor gene expression. *Molec. Endocr.* 2: 242–247, 1988.

[65839] 4169.Prince, M. J.; Smith, F. E.; Peters, E. J.; Stuart, C. A.: Functionalcharacteristics of decreased insulin receptors on fibroblasts obtainedfrom a subject with severe insulin resistance and acanthosis nigricans. *Diabetes* 35:148–154, 1986.

[65840] 4170.Quin, J. D.; Fisher, B. M.; Paterson, K. R.; Inoue, A.; Beastall,G. H.; MacCuish, A. C.: Acute response to recombinant insulin–likegrowth factor–I in a patient with Mendenhall's syndrome. *New Eng.J. Med.* 323: 1425–1426, 1991.

[65841] 4171.Quon, M. J.; Guerre–Millo, M.; Zarnowski, M. J.; Butte, A. J.;Em, M.; Cushman, S. W.; Taylor, S. I.: Tyrosine kinase–deficientmutant human insulin receptors (met1153–to–ile) overexpressed in transfectedrat adipose cells fail to mediate translocation of epitope–taggedGLUT4. *Proc. Nat. Acad. Sci.* 91: 5587–5591, 1994.

[65842] 4172.Rabson, S. M.; Mendenhall, E. N.: Familial hypertrophy of pinealbody, hyperplasia of adrenal cortex and diabetes mellitus. *Am. J.Clin. Path.* 26: 283–290, 1956.

[65843] 4173.Rajala, R. V. S.; Anderson, R. E.: Interaction of the

insulinreceptor beta-subunit with phosphatidylinositol  
3-kinase in bovineROS. Invest. Ophthal. Vis. Sci. 42:  
3110-3117, 2001.

- [65844] 4174.Roth, R. A.; Cassell, D. J.: Insulin receptor: evidence  
thatit is a protein kinase. Science 219: 299-301, 1983.
- [65845] 4175.Rubin, C. S.: Personal Communication. Bronx, N. Y.  
12/8/1984.
- [65846] 4176.Rudiger, H. W.; Ahrens, P.; Dreyer, M.; Frorath, B.;  
Loffel, C.;Schmidt-Preuss, U.: Impaired insulin-induced  
RNA synthesis secondaryto a genetically defective insulin  
receptor. Hum. Genet. 69: 76-78,1985.
- [65847] 4177.Rudiger, H. W.; Dreyer, M.; Kuhnau, J.; Bartelheimer,  
H.: Familialinsulin-resistant diabetes secondary to an  
affinity defect of theinsulin receptor. Hum. Genet. 64:  
407-411, 1983.
- [65848] 4178.Salmeen, A.; Andersen, J. N.; Myers, M. P.; Tonks, N.  
K.; Barford,D.: Molecular basis for the dephosphorylation  
of the activation segmentof the insulin receptor by protein  
tyrosine phosphatase 1B. Molec.Cell 6: 1401-1412, 2000.
- [65849] 4179.Scarlett, J. A.; Kolterman, O. G.; Moore, P.; Saekow,  
M.; Insel,J.; Griffin, J.; Mako, M.; Rubenstein, A. H.; Olef-  
sky, J. M.: Insulinresistance and diabetes due to a genetic  
defect in insulin receptors. J.Clin. Endocr. Metab. 55:

123–132, 1982.

- [65850] 4180.Seino, S.; Seino, M.; Nishi, S.; Bell, G. I.: Structure of the human insulin receptor gene and characterization of its promoter. *Proc.Nat. Acad. Sci.* 86: 114–118, 1989.
- [65851] 4181.Polymeropoulos, M. H.; Torres, R.; Yanovski, J. A.; Chandrasekharappa,S. C.; Ledbetter, D. H.: The human corticotropin–releasing factor receptor (CRHR) gene maps to chromosome 17q12–q22. *Genomics* 28:123–124, 1995.
- [65852] 4182.Kusafuka, T.; Wang, Y.; Puri, P.: Novel mutations of the endothelin–Receptor gene in isolated patients with Hirschsprung's disease. *Hum.Molec. Genet.* 5: 347–349, 1996.
- [65853] 4183.Comings, D. E.: Evidence for ancient tetraploidy and conservation of linkage groups in mammalian chromosomes. *Nature* 238: 455–467,1972.
- [65854] 4184.Deol, M. S.: Genetical studies on the skeleton of the mouse. XXVIII.Tail–short. *Proc. Roy. Soc. Ser. B.* 155: 78–95, 1961.
- [65855] 4185.Hart, C. P.; Awgulewitsch, A.; Fainsod, A.; McGinnis, W.; Ruddle,F. H.: Homeo box gene complex on mouse chromosome 11: molecular cloning,expression in embryogenesis, and homology to a human homeo box locus.

Cell 43:9–18, 1985.

[65856] 4186. Hauser, C. A.; Joyner, A. L.; Klein, R. D.; Learned, T. K.; Martin, G. R.; Tjian, R.: Expression of homologous homeo-box-containing genes in differentiated human teratocarcinoma cells and mouse embryos. Cell 43:19–28, 1985.

[65857] 4187. Levine, M.; Rubin, G. M.; Tjian, R.: Human DNA sequences homologous to a protein coding region conserved between homeotic genes of Drosophila. Cell 38:667–673, 1984.

[65858] 4188. Manley, J. L.; Levine, M. S.: The homeo box and mammalian development. Cell 43:1–2, 1985.

[65859] 4189. Meijlink, F.; de Laaf, R.; Verrijzer, P.; Destree, O.; Kroezen, V.; Hilkens, J.; Deschamps, J.: A mouse homeobox containing gene on chromosome 11: sequence and tissue-specific expression. Nucleic Acids Res. 15: 6773–6786, 1987.

[65860] 4190. Miki, T.; Murphy, P. D.; Pletcher, B. A.; Kidd, J. R.; Ferguson-Smith, A. C.; Ruddle, F. H.; Kidd, K. K.: HOX2 maps to 17q near PPY and NGFR. (Abstract) Cytogenet. Cell Genet. 46: 662 only, 1987.

[65861] 4191. Munke, M.; Cox, D. R.; Jackson, I. J.; Hogan, B. L. M.; Francke, U.: The murine Hox-2 cluster of homeo box con-

taining genes maps distal on chromosome 11 near the tail-short (Ts) locus. *Cytogenet. CellGenet.* 42: 236–240, 1986.

[65862] 4192. Ruddle, F. H.: Personal Communication. New Haven, Conn. 1987.

[65863] 4193. Ruddle, F. H.: Personal Communication. New Haven, Conn. 1985.

[65864] 4194. Schughart, K.; Utset, M. F.; Awgulewitsch, A.; Ruddle, F. H.: Structure and expression of Hox-2.2, a murine homeobox-containing gene. *Proc. Nat. Acad. Sci.* 85: 5582–5586, 1988.

[65865] 4195. Trainor, P. A.; Ariza-McNaughton, L.; Krumlauf, R.: Role of the isthmus and FGFs in resolving the paradox of neural crest plasticity and prepattern. *Science* 295: 1288–1291, 2002.

[65866] 4196. Kaur, S.; Singh, G.; Stock, J. L.; Schreiner, C. M.; Kier, A. B.; Yager, K. L.; Mucenski, M. L.; Scott, W. J., Jr.; Potter, S. S.: Dominant mutation of the murine Hox-2.2 gene results in developmental abnormalities. *J. Exp. Zool.* 264: 323–336, 1992.

[65867] 4197. Lill, M. C.; Fuller, J. F.; Herzig, R.; Crooks, G. M.; Gasson, J. C.: The role of the homeobox gene, HOX B7, in human myelomonocytic differentiation. *Blood* 85:

692–697, 1995.

- [65868] 4198.Simeone, A.; Mavilio, F.; Acampora, D.; Giampaolo, A.; Faiella, A.; Zappavigna, V.; D'Esposito, M.; Pannese, M.; Russo, G.; Boncinelli, E.; Peschle, C.: Two human homeobox genes, c1 and c8: structure analysis and expression in embryonic development. *Proc. Nat. Acad. Sci.* 84:4914–4918, 1987.
- [65869] 4199.Yaron, Y.; McAdara, J. K.; Lynch, M.; Hughes, E.; Gasson, J. C.: Identification of novel functional regions important for the activity of HOXB7 in mammalian cells. *J. Immun.* 166: 5058–5067, 2001.
- [65870] 4200.Sakai, K.; Yamada, M.; Horiba, N.; Wakui, M.; Demura, H.; Suda, T.: The genomic organization of the human corticotropin-releasing factor type-1 receptor. *Gene* 219: 125–130, 1998.
- [65871] 4201.Sillaber, I.; Rammes, G.; Zimmermann, S.; Mahal, B.; Ziegler, W.; Wurst, W.; Holsboer, F.; Spanagel, R.: Enhanced and delayed stress-induced alcohol drinking in mice lacking functional CRH1 receptors. *Science* 296:931–933, 2002.
- [65872] 4202.Smith, G. W.; Aubry, J.-M.; Dellu, F.; Contarino, A.; Bilezikian, L. M.; Gold, L. H.; Chen, R.; Marchuk, Y.; Hauser, C.; Bentley, C.A.; Sawchenko, P. E.; Koob, G. F.;

Vale, W.; Lee, K.-F.: Corticotropin-releasing factor receptor 1-deficient mice display decreased anxiety, impaired stress response, and aberrant neuroendocrine development. *Neuron* 20:1093–1102, 1998.

[65873] 4203. Timpl, P.; Spanagel, R.; Sillaber, I.; Kresse, A.; Reul, J. M. H. M.; Stalla, G. K.; Blanquet, V.; Steckler, T.; Holboer, F.; Wurst, W.: Impaired stress response and reduced anxiety in mice lacking a functional corticotropin-releasing hormone receptor 1. *Nature Genet.* 19:162–166, 1998.

[65874] 4204. Azim, A. C.; Knoll, J. H. M.; Beggs, A. H.; Chisti, A. H.: Isoform cloning, actin binding, and chromosomal localization of human erythrocyte dematin, a member of the villin superfamily. *J. Biol. Chem.* 270:17407–17413, 1995.

[65875] 4205. Azim, A. C.; Marfatia, S. M.; Korsgren, C.; Dotimas, E.; Cohen, C. M.; Chishti, A. H.: Human erythrocyte dematin and protein 4.2 (pallidin) are ATP binding proteins. *Biochemistry* 35: 3001–3006, 1996.

[65876] 4206. Chishti, A. H.; Faquin, W.; Wu, C.-C.; Branton, D.: Purification of erythrocyte dematin (protein 4.9) reveals an endogenous protein kinase that modulates actin-bundling activity. *J. Biol. Chem.* 264:8985–8991, 1989.

[65877] 4207. Gilligan, D. M.; Bennett, V.: The junctional complex of the membrane skeleton. *Seminars Hemat.* 30: 74–83,

1993.

- [65878] 4208.Khanna, R.; Chang, S. H.; Andrabi, S.; Azam, M.; Kim, A.; Rivera,A.; Brugnara, C.; Low, P. S.; Liu, S.-C.; Chishti, A. H.: Headpiecedomain of dematin is required for the stability of the erythrocytemembrane. *Proc. Nat. Acad. Sci.* 99: 6637–6642, 2002.
- [65879] 4209.Peters, L. L.; Eicher, E. M.; Azim, A. C.; Chishti, A. H.: Thegene encoding the erythrocyte membrane skeleton protein dematin (Epb4.9)maps to mouse chromosome 14. *Genomics* 26: 634–635, 1995.
- [65880] 4210.Rana, A. P.; Ruff, P.; Maalouf, G. J.; Speicher, D. W.; Chishti,A. H.: Cloning of human erythroid dematin reveals another memberof the villin family. *Proc. Nat. Acad. Sci.* 90: 6651–6655, 1993.
- [65881] 4211.Correas, I.; Speicher, D. W.; Marchesi, V. T.: Struc-  
ture of thespectrin–actin binding site of erythrocyte pro-  
tein 4.1. *J. Biol.Chem.* 261: 13362–13366, 1986.
- [65882] 4212.Albig, W.; Drabent, B.; Kunz, J.; Kalff–Suske, M.; Grzeschik, K.–H.;Doenecke, D.: All known human H1 his-  
tone genes except the H1(0) geneare clustered on chro-  
mosome 6. *Genomics* 16: 649–654, 1993.
- [65883] 4213.Tanguay, R. M.; Berube, D.; Gagne, R.: Localization  
of histonegenes to chromosomes 6, 12, and 1 by in situ



hybridization. (Abstract) Cytogenet.Cell Genet. 46: 702 only, 1987.

- [65884] 4214.Cheng, J.; Baumhueter, S.; Cacalano, G.; Carver-Moore, K.; Thibodeaux,H.; Thomas, R.; Broxmeyer, H. E.; Cooper, S.; Hague, N.; Moore, M.;Lasky, L. A.: Hematopoietic defects in mice lacking the sialomucinCD34. Blood 87: 479–490, 1996.
- [65885] 4215.He, X.–Y.; Antao, V. P.; Basila, D.; Marx, J. C.; Davis, B. R.: Isolation and molecular characterization of the human CD34 gene. Blood 79:2296–2302, 1992.
- [65886] 4216.Howell, S. M.; Molgaard, H. V.; Greaves, M. F.; Spurr, N. K.:Localisation of the gene coding for the haemopoietic stem cell antigenCD34 to chromosome 1q32. Hum. Genet. 87: 625–627, 1991.
- [65887] 4217.Okuno, Y.; Iwasaki, H.; Huettner, C. S.; Radomska, H. S.; Gonzalez,D. A.; Tenen, D. G.; Akashi, K.: Differential regulation of the humanand murine CD34 genes in hematopoietic stem cells. Proc. Nat. Acad.Sci. 99: 6246–6251, 2002.
- [65888] 4218.Satterthwaite, A. B.; Burn, T. C.; Le Beau, M. M.; Tenen, D. G.: Structure of the gene encoding CD34, a human hematopoietic stemcell antigen. Genomics 12: 788–794, 1992.

- [65889] 4219.Simmons, D. L.; Satterthwaite, A. B.; Tenen, D. G.; Seed, B.:Molecular cloning of a cDNA encoding CD34, a sialomucin of human hematopoieticstem cells. J. Immun. 148: 267–271, 1992.
- [65890] 4220.Sutherland, D. R.; Stewart, A. K.; Keating, A.: CD34 antigen:molecular features and potential clinical applications. Stem Cells 11(suppl. 3): 50–57, 1993.
- [65891] 4221.Sutherland, D. R.; Watt, S. M.; Dowden, G.; Karhi, K.; Baker, M.A.; Greaves, M. F.; Smart, J. E.: Structural and partial amino acidsequence analysis of the human hemopoietic progenitor cell antigenCD34. Leukemia 2: 793–803, 1988.
- [65892] 4222.Tenen, D. G.; Satterthwaite, A. B.; Borson, R.; Simmons, D.; Eddy,R. L.; Shows, T. B.: Chromosome 1 localization of the gene for CD34,a surface antigen of human stem cells. Cytogenet. Cell Genet. 53:55–57, 1990.
- [65893] 4223.Clegg, J. B.: Can the product of the theta gene be a real globin? Nature 329:465–466, 1987.
- [65894] 4224.Fei, Y. J.; Fujita, S.; Huisman, T. H. J.: Two different thetaglobin gene deletions observed among black newborn babies. Brit.J. Haemat. 68: 249–253, 1988.
- [65895] 4225.Hsu, S.–L.; Marks, J.; Shaw, J.–P.; Tam, M.; Higgs, D. R.; Shen,C. C.; Shen, C.–K. J.: Structure and expression of

the human theta-1globin gene. *Nature* 331: 94–96, 1988.

[65896] 4226. Leung, S.; Proudfoot, N. J.; Whitelaw, E.: The gene for theta-globin is transcribed in human fetal erythroid tissues. *Nature* 329: 551–554, 1987.

[65897] 4227. Marks, J.; Shaw, J.-P.; Perez-Stabile, C.; Hu, W.-S.; Ayres, T.M.; Shen, C.; Shen, C.-K. J.: The primate alpha-globin gene family: a paradigm of the fluid genome. *Cold Spring Harbor Symp. Quant. Biol.* 51:499–508, 1986.

[65898] 4228. Marks, J.; Shaw, J.-P.; Shen, C.-K. J.: Sequence organization and genomic complexity of primary theta-1 globin gene, a novel alpha-globin-like gene. *Nature* 321: 785–788, 1986.

[65899] 4229. Peschle, C.; Mavilio, F.; Care, A.; Migliaccio, G.; Migliaccio, A. R.; Salvo, G.; Samoggia, P.; Petti, S.; Guerriero, R.; Marinucci, M.; Lazzaro, D.; Russo, G.; Mastroberardino, G.: Haemoglobin switching in human embryos: asynchrony of zeta-to-alpha- and epsilon-to-gamma-globin switches in primitive and definitive erythropoietic lineage. *Nature* 313:235–238, 1985.

[65900] 4230. Utsch, B.; Albers, N.; Dame, C.; Bartmann, P.; Lentze, M. J.; Ludwig, M.: Homozygous alpha-thalassemia associated with hypospadias: SEA-type deletion does not affect expression of the -14 gene and loss of the theta-1-globin

gene on 16p13.3 is compensated by its duplicate theta-  
2on chromosome 10. (Letter) Am. J. Med. Genet. 101:  
286–287, 2001.

[65901] 4231.Feo, S.; Davies, B.; Fried, M.: The mapping of seven  
intron-containingribosomal protein genes shows they are  
unlinked in the human genome. Genomics 13:201–207,  
1992.

[65902] 4232.Barton, D. E.; Crimando, C.; Hortsch, M.; Francke, U.:  
The genesfor ribophorins I and II are on human chromo-  
somes 3q and 20 and mousechromosomes 6 and 12, re-  
spectively. (Abstract) Cytogenet. Cell Genet. 46:577 only,  
1987.

[65903] 4233.Crimando, C.; Hortsch, M.; Gausepohl, H.; Meyer, D.  
I.: Humanribophorins I and II: the primary structure and  
membrane topologyof two highly conserved round endo-  
plasmic reticulum-specific glycoproteins. EMBOJ. 6:  
75–82, 1987.

[65904] 4234.Elsasser, S.; Gali, R. R.; Schwickart, M.; Larsen, C. N.;  
Leggett,D. S.; Muller, B.; Feng, M. T.; Tubing, F.; Dittmar,  
G. A. G.; Finley,D.: Proteasome subunit Rpn1 binds ubiq-  
uitin-like protein domains. NatureCell Biol. 4: 725–730,  
2002.

[65905] 4235.Kelleher, D. J.; Kreibich, G.; Gilmore, R.: Oligosac-

charyltransferase activity is associated with a protein complex composed of ribophorins I and II and a 48 kd protein. Cell 69: 55–65, 1992.

- [65906] 4236. Horbach, J. M. L. M.; Brenninkmeyer, S. J.; van de Velde, C. J. H.; Nieuwenhuizen Kruseman, A. C.: A forme fruste of von Hippel–Lindau disease—a combination of adrenal pheochromocytoma and ipsilateral renal cell carcinoma: a case report. Surgery 105: 436–441, 1989.
- [65907] 4237. Horton, W. A.; Wong, V.; Eldridge, R.: Von Hippel–Lindau disease—clinical and pathological manifestations in 9 families with 50 affected members. Arch. Intern. Med. 136: 769–777, 1976.
- [65908] 4238. Hull, M. T.; Roth, L. M.; Glover, J. L.; Walker, P. D.: Metastatic carotid body paraganglioma in von Hippel–Lindau disease: an electron microscopic study. Arch. Path. Lab. Med. 106: 235–239, 1982.
- [65909] 4239. Hosoe, S.; Brauch, H.; Latif, F.; Glenn, G.; Daniel, L.; Bale, S.; Choyke, P.; Gorin, M.; Oldfield, E.; Berman, A.; Goodman, J.; Orcutt, M. L.; Hampsch, K.; Delisio, J.; Modi, W.; McBride, W.; Anglard, P.; Weiss, G.; Walther, M. M.; Linehan, W. M.; Lerman, M. I.; Zbar, B.: Localization of the von Hippel–Lindau disease gene to a small region of chromosome 3. Genomics 8: 634–640, 1990.

- [65910] 4240.Huson, S. M.; Harper, P. S.; Hourihan, M. D.; Cole, G.; Weeks, R. D.; Compston, D. A. S.: Cerebellar haemangioblastoma and von Hippel–Lindau disease. *Brain* 109: 1297–1310, 1986.
- [65911] 4241.Iliopoulos, O.; Kibel, A.; Gray, S.; Kaelin, W. G., Jr.: Tumour suppression by the human von Hippel–Lindau gene product. *Nature Med.* 1:822–826, 1995.
- [65912] 4242.Iliopoulos, O.; Levy, A. P.; Jiang, C.; Kaelin, W. G., Jr.; Goldberg, M. A.: Negative regulation of hypoxia-inducible genes by the vonHippel–Lindau protein. *Proc. Nat. Acad. Sci.* 93: 10595–10599, 1996.
- [65913] 4243.Iliopoulos, O.; Ohh, M.; Kaelin, W. G., Jr.: pVHL(19) is a biologically active product of the von Hippel–Lindau gene arising from internal translation initiation. *Proc. Nat. Acad. Sci.* 95: 11661–11666, 1998.
- [65914] 4244.Ivan, M.; Kondo, K.; Yang, H.; Kim, W.; Valiando, J.; Ohh, M.; Salic, A.; Asara, J. M.; Lane, W. S.; Kaelin, W. G., Jr.: HIF- $\alpha$  targeted for VHL-mediated destruction by proline hydroxylation: implications for O(2) sensing. *Science* 292: 464–468, 2001.
- [65915] 4245.Ivanov, S. V.; Kuzmin, I.; Wei, M.–H.; Pack, S.; Geil, L.; Johnson, B. E.; Stanbridge, E. J.; Lerman, M. I.: Down-regulation of transmembrane carbonic anhydrases in renal

cell carcinoma cell lines by wild-type von Hippel–Lindau transgenes. *Proc. Nat. Acad. Sci.* 95: 12596–12601, 1998.

[65916] 4246. Jaakkola, P.; Mole, D. R.; Tian, Y.-M.; Wilson, M. I.; Gielbert, J.; Gaskell, S. J.; von Kriegsheim, A.; Hebestreit, H. F.; Mukherji, M.; Schofield, C. J.; Maxwell, P. H.; Pugh, C. W.; Ratcliffe, P. J.: Targeting of HIF- $\alpha$  to the von Hippel–Lindau ubiquitylation complex by O(2)-regulated prolyl hydroxylation. *Science* 292: 468–472, 2001.

[65917] 4247. James, G. P.: Personal Communication. Springfield, Ohio 8/15/1998.

[65918] 4248. Jennings, A. M.; Smith, C.; Cole, D. R.; Jennings, C.; Shortland, J. R.; Williams, J. L.; Brown, C. B.: Von Hippel–Lindau disease in a large British family: clinicopathological features and recommendations for screening and follow-up. *Quart. J. Med.* 66: 233–249, 1988.

[65919] 4249. Kanno, H.; Kondo, K.; Ito, S.; Yamamoto, I.; Fujii, S.; Torigoe, S.; Sakai, N.; Hosaka, M.; Shuin, T.; Yao, M.: Somatic mutations of the von Hippel–Lindau tumor suppressor gene in sporadic central nervous system human glioblastomas. *Cancer Res.* 54: 4845–4847, 1994.

[65920] 4250. Kanno, H.; Saljooque, F.; Yamamoto, I.; Hattori, S.; Yao, M.; Shuin, T.; U, H.-S.: Role of the von Hippel–Lindau tumor suppressor protein during neuronal differentiation.

Cancer Res. 60: 2820–2824,2000.

[65921] 4251.Kaplan, C.; Sayre, G. P.; Greene, L. F.: Bilateral nephrogeniccarcinomas in Lindau–von Hippel disease. J. Urol. 86: 36–42, 1961.

[65922] 4252.Karsdorp, N.; Elderson, A.; Wittebol–Post, D.; Hene, R. J.; Vos,J.; Feldberg, M. A. M.; van Gils, A. P. G.; Jansen–Schillhorn vanVeen, J. M.; Vroom, T. M.; Hoppener, J. W. M.; Lips, C. J. M.: VonHippel–Lindau disease: new strategies in early detection and treatment. Am.J. Med. 97: 158–168, 1994.

[65923] 4253.Keeler, L. L., III; Klauber, G. T.: von Hippel–Lindau diseaseand renal cell carcinoma in a 16–year–old boy. J. Urol. 147: 1588–1591,1992.

[65924] 4254.Kenck, C.; Wilhelm, M.; Bugert, P.; Staehler, G.; Kovacs, G.:Mutation of the VHL gene is associated exclusively with the developmentof non–papillary renal cell carcinomas. J. Path. 179: 157–161, 1996.

[65925] 4255.Kerr, D. J.; Scheithauer, B. W.; Miller, G. M.; Ebersold, M. J.;McPhee, T. J.: Hemangioblastoma of the optic nerve: case report. Neurosurgery 36:573–581, 1995.

[65926] 4256.Kibel, A.; Iliopoulos, O.; DeCaprio, J. A.; Kaelin, W. G., Jr.: Binding of the von Hippel–Lindau tumor suppressor protein to elonginB and C. Science 269: 1444–1446,



1995.

- [65927] 4257.Kiechle-Schwarz, M.; Neumann, H. P. H.; Decker, H.-J. H.; Dietrich,C.; Wullich, B.; Schempp, W.: Cytogenetic studies on three pheochromocytomasderived from patients with von Hippel-Lindau syndrome. Hum. Genet. 82:127-130, 1989.
- [65928] 4258.Larsen, N.; Samuelsson, T.; Swieb, C.: The Signal RecognitionParticle Database (SRPDB). Nucleic Acids Res. 26: 177-178, 1998.
- [65929] 4259.Pool, M. R.; Stumm, J.; Fulga, T. A.; Sinning, I.; Dobberstein,B.: Distinct modes of signal recognition particle interaction withthe ribosome. Science 297: 1345-1348, 2002.
- [65930] 4260.Yaffe, M. B.; Farr, G. W.; Miklos, D.; Horwich, A. L.; Sternlicht,M. L.; Sternlicht, H.: TCP1 complex is a molecular chaperone in tubulinbiogenesis. Nature 358: 245-248, 1992.
- [65931] 4261.Brown, M. H.; Gorman, P. A.; Sewell, W. A.; Spurr, N. K.; Sheer,D.; Crumpton, M. J.: The gene coding for the human T-lymphocyte CD2antigen is located on chromosome 1p. Hum. Genet. 76: 191-195, 1987.
- [65932] 4262.Clayton, L. K.; Ramachandran, H.; Pravtcheva, D.; Chen, Y.-F.;Diamond, D. J.; Ruddie, F. H.; Reinherz, E. L.:

The gene for T11(CD2) maps to chromosome 1 in humans and to chromosome 3 in mice. *J.Immun.* 140: 3617–3621, 1988.

[65933] 4263.Diamond, D. J.; Clayton, L. K.; Sayre, P. H.; Reinherz, E. L.:Exon–intron organization and sequence comparison of human and murineT11 (CD2) genes. *Proc. Nat. Acad. Sci.* 85: 1615–1619, 1988.

[65934] 4264.Festenstein, R.; Tolaini, M.; Corbella, P.; Mamalaki, C.; Parrington,J.; Fox, M.; Miliou, A.; Jones, M.; Kioussis, D.: Locus control regionfunction and heterochromatin–induced position effect variegation. *Science* 271:1123–1125, 1996.

[65935] 4265.Kingsmore, S. F.; Watson, M. L.; Moseley, W. S.; Seldin, M. F.: Physical linkage of genes encoding the lymphocyte adhesion moleculesCD2 and its ligand LFA–3. *Immunogenetics* 30: 123–125, 1989.

[65936] 4266.Lang, G.; Wotton, D.; Owen, M. J.; Sewell, W. A.; Brown, M. H.;Mason, D. Y.; Crumpton, M. J.; Kioussis, D.: The structure of thehuman CD2 gene and its expression in transgenic mice. *EMBO J.* 7:1675–1682, 1988.

[65937] 4267.Richardson, N. E.; Chang, H.–C.; Brown, N. R.; Hussey, R. E.; Sayre,P. H.; Reinherz, E. L.: Adhesion domain of human T11 (CD2) is encodedby a single exon.

Proc. Nat. Acad. Sci. 85: 5176–5180, 1988.

- [65938] 4268.Seed, B.; Aruffo, A.: Molecular cloning of the CD2 antigen, the T-cell erythrocyte receptor, by a rapid immunoselection procedure. Proc.Nat. Acad. Sci. 84: 3365–3369, 1987.
- [65939] 4269.Sewell, W. A.; Brown, M. H.; Dunne, J.; Owen, M. J.; Crumpton, M. J.: Molecular cloning of the human T-lymphocyte surface CD2 (T11) antigen. Proc. Nat. Acad. Sci. 83: 8718–8722, 1986.
- [65940] 4270.Wang, J.; Smolyar, A.; Tan, K.; Liu, J.; Kim, M.; Sun, Z. J.; Wagner, G.; Reinherz, E. L.: Structure of a heterophilic adhesion complex between the human CD2 and CD58 (LFA-3) counterreceptors. Cell 97:791–803, 1999.
- [65941] 4271.Brissenden, J. E.; Derynck, R.; Francke, U.: Transforming growth factor alpha gene (TGFA) maps to human chromosome 2 close to the breakpoint of the t(2;8) variant translocation in Burkitt lymphoma. (Abstract) Cytogenet. Cell Genet. 40: 589 only, 1985.
- [65942] 4272.Collin, G. B.; Marshall, J. D.; Naggert, J. K.; Nishina, P. M.: TGFA: exon–intron structure and evaluation as a candidate gene for Alstrom syndrome. (Letter) Clin. Genet. 55: 61–62, 1999.
- [65943] 4273.Ellis, D. L.; Kafka, S. P.; Chow, J. C.; Nanney, L. B.;

Inman,W. H.; McCadden, M. E.; King, L. E., Jr.: Melanoma, growth factors,acanthosis nigricans, the sign of Leser-Trelat, and multiple acrochordons:a possible role for alpha-transforming growth factor in cutaneousparaneoplastic syndromes. New Eng. J. Med. 317: 1582–1587, 1987.

[65944] 4274.Fernandez-Larrea, J.; Merlos-Suarez, A.; Urena, J. M.; Baselga,J.; Arribas, J.: A role for a PDZ protein in the early secretorypathway for the targeting of proTGF-alpha to the cell surface. Molec.Cell 3: 423–433, 1999.

[65945] 4275.Fowler, K. J.; Mann, G. B.; Dunn, A. R.: Linkage of the murinetransforming growth factor-alpha gene with Igk, Ly-2, and Fabp1 onchromosome 6. Genomics 16: 782–784, 1993.

[65946] 4276.Tam, J. P.; Scheikh, M. A.; Solomon, D. S.; Ossowski, L.: Efficientsynthesis of human type alpha transforming growth factor: its physicaland biological characterization. Proc. Nat. Acad. Sci. 83: 8082–8086,1986.

[65947] 4277.Tricoli, J. V.; Nakai, H.; Byers, M. G.; Rall, L. B.; Bell, G.I.; Shows, T. B.: Assignment of the gene coding for human TGF-alphato chromosome 2p13. (Abstract) Cyto-genet. Cell Genet. 40: 762 only,1985.

[65948] 4278.Tricoli, J. V.; Nakai, H.; Byers, M. G.; Rall, L. B.; Bell,

G.I.; Shows, T. B.: The gene for human transforming growth factor  $\alpha$  is on the short arm of chromosome 2. Cytogenet. Cell Genet. 42:94–98, 1986.

[65949] 4279.Dong, C.; Zhu, S.; Wang, T.; Yoon, W.; Li, Z.; Alvarez, R. J.;ten Dijke, P.; White, B.; Wigley, F. M.; Goldschmidt–Clermont, P.J.: Deficient Smad7 expression: a putative molecular defect in scleroderma. Proc. Nat. Acad. Sci. 99: 3908–3913, 2002.

[65950] 4280.Dickinson, M. E.; Kobrin, M. S.; Silan, C. M.; Kingsley, D. M.;Justice, M. J.; Miller, D. A.; Ceci, J. D.; Lock, L. F.; Lee, A.;Buchberg, A. M.; Siracusa, L. D.; Lyons, K. M.; Derynck, R.; Hogan,B. L. M.; Copeland, N. G.; Jenkins, N. A.: Chromosomal localizationof seven members of the murine TGF–beta superfamily suggests closelinkage to several morphogenetic mutant loci. Genomics 6: 505–520,1990.

[65951] 4281.Heldin, C.–H.; Miyazono, K.; ten Dijke, P.: TGF–beta signallingfrom cell membrane to nucleus through SMAD proteins. Nature 390:465–471, 1997.

[65952] 4282.Thomas, S. A.; Matsumoto, A. M.; Palmiter, R. D.: Noradrenalineis essential for mouse fetal development. Nature 374: 643–646, 1995.

[65953] 4283.Drebin, J. A.; Hartzell, S. W.; Griffin, C.; Campbell, M.

J.; Niederhuber, J. E.: Molecular cloning and chromosomal localization of the human homologue of a B-lymphocyte specific protein tyrosine kinase (blk). *Oncogene* 10:477–486, 1995.

[65954] 4284. Dymecki, S.; Niederhuber, J.; Desiderio, S.: Specific expression of a novel tyrosine kinase gene, Blk, in B lymphoid cells. *Science* 247:332–336, 1990.

[65955] 4285. Islam, K. B.; Rabbani, H.; Larsson, C.; Sanders, R.; Smith, C. I. E.: Molecular cloning, characterization, and chromosomal localization of a human lymphoid tyrosine kinase related to murine Blk. *J. Immun.* 154:1265–1272, 1995.

[65956] 4286. Kozak, C. A.; Dymecki, S. M.; Niederhuber, J. E.; Desiderio, S. V.: Genetic mapping of the gene for a novel tyrosine kinase, Blk, to mouse chromosome 14. *Genomics* 9: 762–764, 1991.

[65957] 4287. Kikuchi, S.; Hata, M.; Fukumoto, K.; Yamane, Y.; Matsui, T.; Tamura, A.; Yonemura, S.; Yamagishi, H.; Keppler, D.; Tsukita, S.; Tsukita, S.: Radixin deficiency causes conjugated hyperbilirubinemia with loss of Mrp2 from bile canalicular membranes. *Nature Genet.* 31:320–325, 2002.

[65958] 4288. Wilgenbus, K. K.; Milatovich, A.; Francke, U.; Furthmayr, H.: Molecular cloning, cDNA sequence, and chromo-

somal assignment of the human radixin gene and two dispersed pseudogenes. *Genomics* 16: 199–206, 1993.

- [65959] 4289. Geppert, M.; Goda, Y.; Stevens, C. F.; Sudhof, T. C.: The small GTP-binding protein Rab3A regulates a late step in synaptic vesicle fusion. *Nature* 387: 810–814, 1997.
- [65960] 4290. Kapfhamer, D.; Valladares, O.; Sun, Y.; Nolan, P. M.; Rux, J. J.; Arnold, S. E.; Veasey, S. C.; Bucan, M.: Mutations in Rab3a alter circadian period and homeostatic response to sleep loss in the mouse. *Nature Genet.* 32: 290–295, 2002.
- [65961] 4291. Rousseau-Merck, M. F.; Zahraoui, A.; Bernheim, A.; Touchot, N.; Migliarina, R.; Tavitian, A.; Berger, R.: Chromosome mapping of the human RAS related RAB3A and RAB3B genes to chromosomes 19p13.2 and 1p31–p32, respectively. (Abstract) *Cytogenet. Cell Genet.* 51: 1070 only, 1989.
- [65962] 4292. Rousseau-Merck, M. F.; Zahraoui, A.; Bernheim, A.; Touchot, N.; Migliarina, R.; Tavitian, A.; Berger, R.: Chromosome mapping of the human RAS-related RAB3A gene to 19p13.2. *Genomics* 5: 694–698, 1989.
- [65963] 4293. Zahraoui, A.; Touchot, N.; Chardin, P.; Tavitian, A.: The human rab genes encode a family of GTP-binding proteins related to yeast YPT1 and SEC4 products involved

in secretion. J. Biol. Chem. 264:12394–12401, 1989.

[65964] 4294.Durkin, A. S.; Maglott, D. R.; Nierman, W. C.: Chromosomal assignment of 38 human brain expressed sequence tags (ESTs) by analyzing fluorescently labeled PCR products from hybrid cell panels. Genomics 14: 808–810, 1992.

[65965] 4295.Hussey, D. J.; Nicola, M.; Moore, S.; Peters, G. B.; Dobrovic, A.: The (4;11)(q21;p15) translocation fuses the NUP98 and RAP1GDS1 genes and is recurrent in T-cell acute lymphocytic leukemia. Blood 94:2072–2079, 1999.

[65966] 4296.Germain, P.; Iyer, J.; Zechel, C.; Gronemeyer, H.: Coregulator recruitment and the mechanism of retinoic acid receptor synergy. Nature 415:187–192, 2002.

[65967] 4297.McNamara, P.; Seo, S.; Rudic, R. D.; Sehgal, A.; Chakravarti, D.; FitzGerald, G. A.: Regulation of CLOCK and MOP4 by nuclear hormone receptors in the vasculature: a humoral mechanism to reset a peripheral clock. Cell 105: 877–889, 2001.

[65968] 4298.Almasan, A.; Mangelsdorf, D. J.; Ong, E. S.; Wahl, G. M.; Evans, R. M.: Chromosomal localization of the human retinoid X receptors. Genomics 20:397–403, 1994.

[65969] 4299.Claudel, T.; Leibowitz, M. D.; Fievet, C.; Tailleux, A.; Wagner, B.; Repa, J. J.; Torpier, G.; Lobaccaro, J.-M.; Pater-



niti, J. R.; Mangelsdorf, D. J.; Heyman, R. A.; Auwerx, J.: Reduction of atherosclerosis in apolipoprotein E knockout mice by activation of the retinoid X receptor. *Proc. Nat. Acad. Sci.* 98: 2610–2615, 2001.

[65970] 4300. de Urquiza, A. M.; Liu, S.; Sjöberg, M.; Zetterstrom, R. H.; Griffiths, W.; Sjövall, J.; Perlmann, T.: Docosahexaenoic acid, a ligand for the retinoid X receptor in mouse brain. *Science* 290: 2140–2144, 2000.

[65971] 4301. Thomas, G. D.; Sander, M.; Lau, K. S.; Huang, P. L.; Stull, J. T.; Victor, R. G.: Impaired metabolic modulation of alpha-adrenergic vasoconstriction in dystrophin-deficient skeletal muscle. *Proc. Nat. Acad. Sci.* 95: 15090–15095, 1998.

[65972] 4302. Xie, J.; Roddy, P.; Rife, T. K.; Murad, F.; Young, A. P.: Two closely linked but separable promoters for human neuronal nitric oxide synthase gene transcription. *Proc. Nat. Acad. Sci.* 92: 1242–1246, 1995.

[65973] 4303. Xu, W.; Gorman, P.; Sheer, D.; Bates, G.; Kishimoto, J.; Lizhi, L.; Emson, P.: Regional localization of the gene coding for human brain nitric oxide synthase (NOS1) to 12q24.2–24.31 by fluorescent in situ hybridization. *Cytogenet. Cell Genet.* 64: 62–63, 1993.

[65974] 4304. Taniyama, T.; Takai, S.; Miyazaki, E.; Fukumura, R.;

Sato, J.;Kobayashi, Y.; Hirakawa, T.; Moore, K. W.; Yamada, K.: The humaninterleukin-10 receptor gene maps to chromosome 11q23.3. Hum. Genet. 95:99-101, 1995.

[65975] 4305.Bono, M. R.; Alcaide-Loridan, C.; Couillin, P.; Letouze, B.; Grisard,M. C.; Jouin, H.; Fellous, M.: Human chromosome 16 encodes a factorinvolved in induction of class II major histocompatibility antigensby interferon gamma. Proc. Nat. Acad. Sci. 88: 6077-6081, 1991.

[65976] 4306.Bono, R.; Hatat, D.; Couillin, P.; Grisard, M. C.; Van Cong, N.;Fisher, D.; Fellous, M.: Receptor for human gamma interferon is specifiedby human chromosomes 6 and 21. (Abstract) Cytogenet. Cell Genet. 46:584, 1987.

[65977] 4307.Dorman, S. E.; Holland, S. M.: Mutation in the signal-transducingchain of the interferon-gamma receptor and susceptibility to mycobacterialinfection. J. Clin. Invest. 101: 2364-2369, 1998.

[65978] 4308.Jung, V.; Rashidbaigi, A.; Jones, C.; Tischfield, J. A.; Shows,T. B.; Pestka, S.: Human chromosomes 6 and 21 are required for sensitivityto human interferon gamma. Proc. Nat. Acad. Sci. 84: 4151-4155,1987.

[65979] 4309.Langer, J. A.; Rashidbaigi, A.; Lai, L.-W.; Patterson, D.; Jones,C.: Sublocalization on chromosome 21 of human interferon-alpha receptorgene and the gene for an inter-

feron-gamma response protein. *Somat.Cell Molec. Genet.* 16: 231-240, 1990.

[65980] 4310.Mariano, T. M.; Muthukumaran, G.; Donnelly, R. J.; Wang, N.; Adamson,M. C.; Pestka, S.; Kozak, C. A.: Genetic mapping of the gene forthe mouse interferon-gamma receptor signaling subunit to the distalend of chromosome 16. *Mammalian Genome* 7: 321-322, 1996.

[65981] 4311.Rhee, S.; Ebensperger, C.; Dembic, Z.; Pestka, S.: The structureof the gene for the second chain of the human interferon-gamma receptor. *J.Biol. Chem.* 271: 28947-28952, 1996.

[65982] 4312.Soh, J.; Donnelly, R. J.; Mariano, T. M.; Cook, J. R.; Schwartz,B.; Pestka, S.: Identification of a yeast artificial chromosome cloneencoding an accessory factor for the human interferon gamma receptor:evidence for multiple accessory factors. *Proc. Nat. Acad. Sci.* 90:8737-8741, 1993.

[65983] 4313.Badovinac, V. P.; Tvinnereim, A. R.; Harty, J. T.: Regulationof antigen-specific CD8(+) T cell homeostasis by perforin and interferon-gamma. *Science* 290:1354-1357, 2000.

[65984] 4314.Ben-Asouli, Y.; Banai, Y.; Pel-Or, Y.; Shir, A.; Kaempfer, R.:Human interferon-gamma mRNA autoregu-

lates its translation through a pseudoknot that activates the interferon-inducible protein kinase PKR. *Cell* 108: 221–232, 2002.

- [65985] 4315. Binder, G. K.; Griffin, D. E.: Interferon- $\gamma$ -mediated site-specific clearance of alphavirus from CNS neurons. *Science* 293: 303–306, 2001.
- [65986] 4316. Blalock, J. E.; Smith, E. M.: Human leukocyte interferon: structural and biological relatedness to adrenocorticotrophic hormone and endorphins. *Proc. Nat. Acad. Sci.* 77: 5972–5974, 1980.
- [65987] 4317. Bureau, J. F.; Bihl, F.; Brahic, M.; Le Paslier, D.: The gene coding for interferon- $\gamma$  is linked to the D12S335 and D12S313 microsatellites and to the MDM2 gene. *Genomics* 28: 109–112, 1995.
- [65988] 4318. Burke, D. C.: The status of interferon. *Sci. Am.* 236(4): 42–50, 1977.
- [65989] 4319. Cassingena, R.; Chany, C.; Vignal, M.; Suarez, H.; Estrade, S.; Lazar, P.: Use of monkey-mouse hybrid cells for the study of the cellular regulation of interferon production and action. *Proc. Nat. Acad. Sci.* 68: 580–584, 1971.
- [65990] 4320. Creagan, R. P.; Tan, Y. H.; Chen, S.-H.; Ruddle, F. H.: Somatic cell genetic analysis of the interferon system. *Fed.*

Proc. 34: 2222–2226,1975.

- [65991] 4321.Devos, R.; Cheroutre, H.; Taya, Y.; Degrave, W.; Van Heuverswyn,H.; Fiers, W.: Molecular cloning of human immune interferon cDNAand its expression in eukaryotic cells. Nucleic Acids Res. 10: 2487–2502,1982.
- [65992] 4322.Diaz, M. O.; Bohlander, S.; Allen, G.: Nomenclature of the humaninterferon genes. J. Interferon Res. 13: 443–444, 1993.
- [65993] 4323.Diefenbach, A.; Schindler, H.; Rollinghoff, M.; Yokoyama, W. M.;Bogdan, C.: Requirement for type 2 NO synthase for IL–12 signalingin innate immunity. Science 284: 951–955, 1999.
- [65994] 4324.Gray, P. W.; Goeddel, D. V.: Structure of the human immune interferongene. Nature 298: 859–863, 1982.
- [65995] 4325.Isaacs, A.; Lindenmann, J.: Virus interference. I. The interferon. Proc.Roy. Soc. London 147B: 258–267, 1957.
- [65996] 4326.Isaacs, A.; Lindenmann, J.; Valentine, R. C.: Virus interference.II. Some properties of interferon. Proc. Roy. Soc. London 147B:268–273, 1957.
- [65997] 4327.Knight, E., Jr.: Human fibroblast interferon: amino acid analysisand amino terminal amino acid sequence. Science 207: 525–526, 1980.
- [65998] 4328.Lipinski, M.; Virelizier, J. L.; Tursz, T.; Griscelli, C.:

Naturalkiller and killer cell activities in patients with primary immunodeficiencies or defects in immune interferon production. *Europ. J. Immun.* 10:246–249, 1980.

[65999] 4329. Maeda, S.; McCandliss, R.; Gross, M.; Sloma, A.; Familletti, P.C.; Tabor, J. M.; Evinger, M.; Levy, W. P.; Pestka, S.: Construction and identification of bacterial plasmids containing nucleotide sequence for human leukocyte interferon. *Proc. Nat. Acad. Sci.* 77: 7010–7013, 1980.

[66000] 4330. Mantei, N.; Schwarzstein, M.; Streuli, M.; Panem, S.; Nagata, S.; Weissmann, C.: The nucleotide sequence of a cloned human leukocyte interferon cDNA. *Gene* 10: 1–10, 1980.

[66001] 4331. Nagata, S.; Mantei, N.; Weissmann, C.: The structure of one of the eight or more distinct chromosomal genes for human interferon- $\alpha$ . *Nature* 287:401–408, 1980.

[66002] 4332. Nagata, S.; Taira, H.; Hall, A.; Johnstrud, L.; Streuli, M.; Escodi, J.; Boll, W.; Cantell, K.; Weissmann, C.: Synthesis in *E. coli* of a polypeptide with human leukocyte interferon activity. *Nature* 284:316–320, 1980.

[66003] 4333. Nathan, C. F.; Murray, H. W.; Wiebe, M. E.; Rubin, B. Y.: Identification of interferon- $\gamma$  as the lymphokine that activates human macrophage oxidative metabolism

and antimicrobial activity. J. Exp. Med. 158:670–689, 1983.

[66004] 4334.Naylor, S. L.; Sakaguchi, A. Y.; Shows, T. B.; Law, M. L.; Goeddel, D. V.; Gray, P. W.: Human immune interferon gene is located on chromosome 12. J. Exp. Med. 57: 1020–1027, 1983.

[66005] 4335.Shimizu, A.; Sakai, Y.; Ohno, K.; Masaki, S.; Kuwano, R.; Takahashi, Y.; Miyashita, N.; Watanabe, T.: A molecular genetic linkage map of mouse chromosome 10, including the Myb, S100b, Pah, Sl, and Ifg genes. Biochem. Genet. 30: 529–535, 1992.

[66006] 4336.Blumenberg, M.; Savtchenko, E. S.: Linkage of human keratin genes. Cytogenet. Cell Genet. 42: 65–71, 1986.

[66007] 4337.Fuchs, E.; Coppock, S.; Green, H.; Cleveland, D.: Two distinct classes of keratin genes and their evolutionary significance. Cell 27:75–84, 1981.

[66008] 4338.Fuchs, E.; Green, H.: Changes in keratin gene expression during terminal differentiation of the keratinocyte. Cell 19: 1033–1042, 1980.

[66009] 4339.Hanukoglu, I.; Fuchs, E.: The cDNA sequence of a human epidermal keratin: divergence of sequence but conservation of structure among intermediate filament

proteins. Cell 31: 243–252, 1982.

[66010] 4340.Lee, L. D.; Baden, H. P.: Organisation of the polypeptide chains in mammalian keratin. Nature 264: 377–379, 1976.

[66011] 4341.Marchuk, D.; McCrohon, S.; Fuchs, E.: Complete sequence of a gene encoding a human type I keratin: sequences homologous to enhancer elements in the regulatory region of the gene. Proc. Nat. Acad. Sci. 82:1609–1613, 1985.

[66012] 4342.Nadeau, J. H.; Berger, F. G.; Cox, D. R.; Crosby, J. L.; Davisson, M. T.; Ferrara, D.; Fuchs, E.; Hart, C.; Hunihan, L.; Lalley, P. A.; Langley, S. H.; Martin, G. R.; Nichols, L.; Phillips, S. J.; Roderick, T. H.; Roop, D. R.; Ruddle, F. H.; Skow, L. C.; Compton, J. G.: A family of type I keratin genes and the homeobox-2 gene complex are closely linked to the rex locus on mouse chromosome 11. Genomics 5:454–462, 1989.

[66013] 4343.Ray Chaudhury, A.; Marchuk, D.; Lindhurst, M.; Fuchs, E.: Three tightly linked genes encoding human type I keratins: conservation of sequence in the 5-prime-untranslated leader and 5-prime-upstream regions of coexpressed keratin genes. Molec. Cell. Biol. 6: 539–548, 1986.



- [66014] 4344. Romano, V.; Bosco, P.; Raimondi, E.; Feo, S.; Leube, R.; Franke, W.; Ceratto, N.: Chromosomal mapping and physical linkage analysis of human acidic cytokeratin genes. (Abstract) *Cytogenet. Cell Genet.* 58:2009–2010, 1991.
- [66015] 4345. Heim, S.; Nilbert, M.; Vanni, R.; Floderus, U.-M.; Mandahl, N.; Liedgren, S.; Lecca, U.; Mitelman, F.: A specific translocation, t(12;14)(q14–15;q23–24), characterizes a subgroup of uterine leiomyomas. *Cancer Genet. Cytogenet.* 32: 13–17, 1988.
- [66016] 4346. Nucifora, G.; Begy, C. R.; Kobayashi, H.; Roulston, D.; Claxton, D.; Pedersen-Bjergaard, J.; Parganas, E.; Ihle, J. N.; Rowley, J. D.: Consistent intergenic splicing and production of multiple transcripts between AML1 at 21q22 and unrelated genes at 3q26 in (3;21)(q26;q22) translocations. *Proc. Nat. Acad. Sci.* 91: 4004–4008, 1994.
- [66017] 4347. Nucifora, G.; Rowley, J. D.: AML1 and the 8;21 and 3;21 translocations in acute and chronic myeloid leukemia. *Blood* 86: 1–14, 1995.
- [66018] 4348. Tan, M.; Jing, T.; Lan, K.-H.; Neal, C. L.; Li, P.; Lee, S.; Fang, D.; Nagata, Y.; Liu, J.; Arlinghaus, R.; Hung, M.-C.; Yu, D.: Phosphorylation on tyrosine-15 of p34(Cdc2) by

ErbB2 inhibits p34(Cdc2) activation and is involved in resistance to taxol-induced apoptosis. *Molec.Cell* 9: 993–1004, 2002.

- [66019] 4349. Gromeier, M.; Solecki, D.; Patel, D. D.; Wimmer, E.: Expression of the human poliovirus receptor/CD155 gene during development of the central nervous system: implications for the pathogenesis of poliomyelitis. *Virology* 273:248–257, 2000.
- [66020] 4350. He, Y.; Bowman, V. D.; Mueller, S.; Bator, C. M.; Bella, J.; Peng, X.; Baker, T. S.; Wimmer, E.; Kuhn, R. J.; Rossmann, M. G.: Interaction of the poliovirus receptor with poliovirus. *Proc. Nat. Acad. Sci.* 97:79–84, 2000.
- [66021] 4351. Koike, S.; Horie, H.; Ise, I.; Okitsu, A.; Yoshida, M.; Iizuka, N.; Takeuchi, K.; Takegami, T.; Nomoto, A.: The poliovirus receptor protein is produced both as membrane-bound and secreted forms. *EMBO J.* 9: 3217–3224, 1990.
- [66022] 4352. Koike, S.; Taya, C.; Kurata, T.; Abe, S.; Ise, I.; Yonekawa, H.; Nomoto, A.: Transgenic mice susceptible to poliovirus. *Proc. Nat. Acad. Sci.* 88: 951–955, 1991.
- [66023] 4353. Mendelsohn, C.; Johnson, B.; Lionetti, K. A.; Nobis, P.; Wimmer, E.; Racaniello, V. R.: Transformation of a human poliovirus receptor gene into mouse cells. *Proc. Nat.*

Acad. Sci. 83: 7845–7849, 1986.

- [66024] 4354. Mendelsohn, C. L.; Wimmer, E.; Racaniello, V. R.: Cellular receptor for poliovirus: molecular cloning, nucleotide sequence, and expression of a new member of the immunoglobulin superfamily. *Cell* 56: 855–865, 1989.
- [66025] 4355. Miller, D. A.; Miller, O. J.; Dev, V. G.; Hashmi, S.; Tantravahi, R. R.; Medrano, L.; Green, H.: Human chromosome 19 carries a poliovirus receptor gene. *Cell* 1: 167–174, 1974.
- [66026] 4356. Larrea, F.; Escorza, A.; Granados, J.; Valencia, X.; Valero, A.; Cravioto, M. C.; Perez-Palacios, G.: Familial occurrence of big-big prolactin as the predominant immunoreactive human prolactin species in blood. *Fertil. Steril.* 47: 956–963, 1987.
- [66027] 4357. Carter, R. E.; Cerosaletti, K. M.; Burkin, D. J.; Fournier, R. E. K.; Jones, C.; Greenberg, B. D.; Citron, B. A.; Festoff, B. W.: The gene for the serpin thrombin inhibitor (P17), protease nexin 1, is located on human chromosome 2q33–q35 and on syntenic regions in the mouse and sheep genomes. *Genomics* 27: 196–199, 1995.
- [66028] 4358. Gloor, S.; Odink, K.; Guenther, J.; Nick, H.; Monard, D.: A glia-derived neurite promoting factor with protease inhibitory activity belongs to the protease nexins. *Cell* 47:

687–693, 1986.

- [66029] 4359.Hall, C. L.; Yang, B.; Yang, X.; Zhang, S.; Turley, M.; Samuel,S.; Lange, L. A.; Wang, C.; Curpen, G. D.; Savani, R. C.; Greenberg,A. H.; Turley, E. A.: Overexpression of the hyaluronan receptor RHAMM is transforming and is also required for H-ras transformation. *Cell* 82:19–26, 1995.
- [66030] 4360.Itano, N.; Kimata, K.: Expression cloning and molecular characterization of HAS protein, a eukaryotic hyaluronan synthase. *J. Biol. Chem.* 271:9875–9878, 1996.
- [66031] 4361.Itano, N.; Kimata, K.: Molecular cloning of human hyaluronan synthase. *Biochem.Biophys. Res. Commun.* 222: 816–820, 1996.
- [66032] 4362.Shyjan, A. M.; Heldin, P.; Bucher, E. C.; Yoshino, T.; Briskin,M. J.: Functional cloning of the cDNA for a human hyaluronan synthase. *J.Biol. Chem.* 271: 23395–23399, 1996.
- [66033] 4363.Spicer, A. P.; Seldin, M. F.; Olsen, A. S.; Brown, N.; Wells, D.E.; Doggett, N. A.; Itano, N.; Kimata, K.; Inazawa, J.; McDonald,J. A.: Chromosomal localization of the human and mouse hyaluronan synthase genes. *Genomics* 41: 493–497, 1997.
- [66034] 4364.Watanabe, K.; Yamaguchi, Y.: Molecular identification of a putative human hyaluronan synthase. *J. Biol.*

Chem. 271: 22945–22948, 1996.

[66035] 4365.Johansson, M.; Karlsson, A.: Cloning and expression of human deoxyguanosinekinase cDNA. Proc. Nat. Acad. Sci. 93: 7258–7262, 1996.

[66036] 4366.Taanman, J.–W.; Kateeb, I.; Muntau, A. C.; Jaksch, M.; Cohen, N.;Mandel, H.: A novel mutation in the deoxyguanosine kinase gene causingdepletion of mitochondrial DNA. Ann. Neurol. 52: 237–239, 2002.

[66037] 4367.Cahill, D. P.; da Costa, L. T.; Carson–Walter, E. B.; Kinzler,K. W.; Vogelstein, B.; Lengauer, C.: Characterization of MAD2B andother mitotic spindle checkpoint genes. Genomics 58: 181–187, 1999.

[66038] 4368.Chen, R.–H.; Waters, J. C.; Salmon, E. D.; Murray, A. W.: Associationof spindle assembly checkpoint component X MAD2 with unattached kinetochores. Science 274:242–245, 1996.

[66039] 4369.Dobles, M.; Liberal, V.; Scott, M. L.; Benezra, R.; Sorger, P.K.: Chromosome missegregation and apoptosis in mice lacking the mitoticcheckpoint protein Mad2. Cell 101: 635–645, 2000.

[66040] 4370.Krishnan, R.; Goodman, B.; Jin, D.–Y.; Jeang, K.–T.; Collins, C.;Stetten, G.; Spencer, F.: Map location and gene structure of theHomo sapiens mitotic arrest deficient 2

(MAD2L1) gene at 4q27. Genomics 49:475–478, 1998.

- [66041] 4371.Li, X.; Nicklas, R. B.: Mitotic forces control a cell-cycle checkpoint. Nature 373:630–632, 1995.
- [66042] 4372.Li, Y.; Benezra, R.: Identification of a human mitotic checkpointgene: hsMAD2. Science 274: 246–248, 1996.
- [66043] 4373.Luo, X.; Tang, Z.; Rizo, J.; Yu, H.: The Mad2 spindle checkpointprotein undergoes similar major conformational changes upon bindingto either Mad1 or Cdc20. Molec. Cell 9: 59–71, 2002.
- [66044] 4374.Michel, L. S.; Liberal, V.; Chatterjee, A.; Kirchwegger, R.; Pasche,B.; Gerald, W.; Dobles, M.; Sorger, P. K.; Murty, V. V. V. S.; Benezra,R.: MAD2 haplo-insufficiency causes premature anaphase and chromosomeinstability in mammalian cells. Nature 409: 355–359, 2001.
- [66045] 4375.Nelson, K. K.; Schlondorff, J.; Blobel, C. P.: Evidence for aninteraction of the metalloprotease–disintegrin tumour necrosis factoralpha convertase (TACE) with mitotic arrest deficient 2 (MAD2), andof the metalloprotease–disintegrin MDC9 with a novel MAD2–relatedprotein, MAD2–beta. Biochem. J. 343: 673–680, 1999.
- [66046] 4376.Shonn, M. A.; McCarroll, R.; Murray, A. W.: Requirement of thespindle checkpoint for proper chromosome segregation in budding yeastmeiosis. Science 289:

300–303, 2000.

- [66047] 4377. Xu, L.; Deng, H. X.; Yang, Y.; Xia, J. H.; Hung, W. Y.; Siddique, T.: Assignment of mitotic arrest deficient protein 2 (MAD2L1) to human chromosome band 5q23.3 by in situ hybridization. *Cytogenet. Cell Genet.* 78: 63–64, 1997.
- [66048] 4378. De La Rosa, J.; Ostrowski, J.; Hryniewicz, M. M.; Kredich, N. M.; Kotb, M.; LeGros, H. L., Jr.; Valentine, M.; Geller, A. M.: Chromosomal localization and catalytic properties of the recombinant alpha subunit of human lymphocyte methionine adenosyltransferase. *J. Biol. Chem.* 270: 21860–21868, 1995.
- [66049] 4379. Wang, D.; Stravopodis, D.; Teglund, S.; Kitazawa, J.; Ihle, J. N.: Naturally occurring dominant negative variants of Stat5. *Molec. Cell Biol.* 16: 6141–6148, 1996.
- [66050] 4380. Hou, J.; Schindler, U.; Henzel, W. J.; Ho, T. C.; Brasseur, M.; McKnight, S. L.: An interleukin-4-induced transcription factor: IL-4Stat. *Science* 265: 1701–1706, 1994.
- [66051] 4381. Leek, J. P.; Hamlin, P. J.; Bell, S. M.; Lench, N. J.: Assignment of the STAT6 gene (STAT6) to human chromosome band 12q13 by in situ hybridization. *Cytogenet. Cell Genet.* 79: 208–209, 1997.
- [66052] 4382. Patel, B. K. R.; Keck, C. L.; O'Leary, R. S.; Popescu, N.

C.;LaRochelle, W. J.: Localization of the human Stat6 gene to chromosome12q13.3–q14.1, a region implicated in multiple solid tumors. *Genomics* 52:192–200, 1998.

[66053] 4383.Patel, B. K. R.; Pierce, J. H.; LaRochelle, W. J.: Regulation of interleukin 4–mediated signaling by naturally occurring dominantnegative and attenuated forms of human Stat6. *Proc. Nat. Acad. Sci.* 95:172–177, 1998.

[66054] 4384.Quelle, F. W.; Shimoda, K.; Thierfelder, W.; Fischer, C.; Kim,A.; Ruben, S. M.; Cleveland, J. L.; Pierce, J. H.; Keegan, A. D.;Nelms, K.; Paul, W. E.; Ihle, J. N.: Cloning of murine Stat6 andhuman Stat6, Stat proteins that are tyrosine phosphorylated in response to IL–4 and IL–4 but are not required for mitogenesis. *Molec. Cell.Biol.* 15: 3336–3343, 1995.

[66055] 4385.Liu, Y.; Chiu, I.–M.: Assignment of FGF12, the human FGF homologousfactor 1 gene, to chromosome 3q29–3qter by fluorescence in situ hybridization. *Cytogenet.Cell Genet.* 78: 48–49, 1997.

[66056] 4386.Caslini, C.; Spinelli, O.; Cazzaniga, G.; Golay, J.; De Gioia,L.; Pedretti, A.; Breviario, F.; Amaru, R.; Barbui, T.; Biondi, A.;Introna, M.; Rambaldi, A.: Identification of two novel isoforms ofthe ZNF162 gene: a growing family of signal transduction and activator of RNA proteins. *Ge–*



nomics 42: 268–277, 1997.

[66057] 4387.Kramer, A.; Quentin, M.; Mulhauser, F.: Diverse modes of alternativesplicing of human splicing factor SF1 deduced from the exon–intronstructure of the gene. *Gene* 211: 29–37, 1998.

[66058] 4388.Reddy, U. R.; Phatak, S.; Allen, C.; Nycum, L. M.; Sulman, E. P.;White, P. S.; Biegel, J. A.: Localization of the human Ror1 gene(NTRKR1) to chromosome 1p31–p32 by fluorescence in situ hybridizationand somatic cell hybrid analysis. *Genomics* 41: 283–285, 1997.

[66059] 4389.Reddy, U. R.; Phatak, S.; Pleasure, D.: Human neural tissues expressa truncated Ror1 receptor tyrosine kinase, lacking both extracellularand transmembrane domains. *Oncogene* 13: 1555–1559, 1996.

[66060] 4390.Gong, Y.; Chitayat, D.; Kerr, B.; Chen, T.; Babul–Hirji, R.; Pal,A.; Reiss, M.; Warman, M. L.: Brachydactyly type B: clinical description,genetic mapping to chromosome 9q, and evidence for a shared ancestralmutation. *Am. J. Hum. Genet.* 64: 578–585, 1999.

[66061] 4391.Innis, J. W.; Mortlock, D. P.: Limb development: molecular dysmorphologyis at hand! *Clin. Genet.* 53: 337–348, 1998.

[66062] 4392.Manouvrier–Hanu, S.; Holder–Espinasse, M.; Lyon–

net, S.: Geneticsof limb anomalies in humans. Trends Genet. 15: 409–417, 1999.

[66063] 4393.Oishi, I.; Takeuchi, S.; Hashimoto, R.; Nagabukuro, A.; Ueda, T.;Liu, Z.-J.; Hatta, T.; Akira, S.; Matsuda, Y.; Yamamura, H.; Otani,H.; Minami, Y.: Spatio-temporally regulated expression of receptortyrosine kinases, mRor1, mRor2, during mouse development: implicationsin development and function of the nervous system. Genes Cells 4:41–56, 1999.

[66064] 4394.Schwabe, G. C.; Tinschert, S.; Buschow, C.; Meinecke, P.; Wolff,G.; Gillessen-Kaesbach, G.; Oldridge, M.; Wilkie, A. O. M.; Komec,R.; Mundlos, S.: Distinct mutations in the receptor tyrosine kinasegene ROR2 cause brachydactyly type B. Am. J. Hum. Genet. 67: 822–831,2000.

[66065] 4395.Takeuchi, S.; Takeda, K.; Oishi, I.; Nomi, M.; Ikeya, M.; Itoh,K.; Tamura, S.; Ueda, T.; Hatta, T.; Otani, H.; Terashima, T.; Takada,S.; Yamamura, H.; Akira, S.; Minami, Y.: Mouse Ror2 receptor tyrosinekinase is required for the heart development and limb formation. GenesCells 5: 71–78, 2000.

[66066] 4396.van Bokhoven, H.; Brunner, H. G.: Splitting p63. Am. J. Hum.Genet. 71: 1–13, 2002.

[66067] 4397.Jaffrey, S. R.; Snyder, S. H.: PIN: an associated pro-

tein inhibitor of neuronal nitric oxide synthase. *Science* 274: 774–777, 1996.

[66068] 4398. Garkavtsev, I.; Demetrick, D.; Riabowol, K.: Cellular localization and chromosome mapping of a novel candidate tumor suppressor gene (ING1). *Cytogenet. Cell Genet.* 76: 176–178, 1997.

[66069] 4399. Garkavtsev, I.; Kazarov, A.; Gudkov, A.; Riabowol, K.: Suppression of the novel growth inhibitor p33(ING1) promotes neoplastic transformation. *Nature Genet.* 14: 415–420, 1996. Note: Erratum: *Nature Genet.* 23: 373 only, 1999.

[66070] 4400. Saito, A.; Furukawa, T.; Fukushige, S.; Koyama, S.; Hoshi, M.; Hayashi, Y.; Horii, A.: p24/ING1-ALT1 and p47/ING1-ALT2, distinct alternative transcripts of p33/ING1. *J. Hum. Genet.* 45: 177–181, 2000.

[66071] 4401. Zeremski, M.; Horrigan, S. K.; Grigorian, I. A.; Westbrook, C. A.; Gudkov, A. V.: Localization of the candidate tumor suppressor gene ING1 to human chromosome 13q34. *Somat. Cell Molec. Genet.* 23: 233–236, 1997.

[66072] 4402. Gilligan, D. M.; Lozovatsky, L.; Gwynn, B.; Brugnara, C.; Mohandas, N.; Peters, L. L.: Targeted disruption of the beta adducin gene (Add2) causes red blood cell spherocytosis in mice. *Proc. Nat. Acad. Sci.* 96: 10717–10722,

1999.

- [66073] 4403. Adamson, M. C.; Dennis, C.; Delaney, S.; Christiansen, J.; Monkley, S.; Kozak, C. A.; Wainwright, B.: Isolation and genetic mapping of two novel members of the murine Wnt gene family, Wnt11 and Wnt12, and the mapping of Wnt5a and Wnt7a. *Genomics* 24: 9–13, 1994.
- [66074] 4404. Bui, T. D.; Lako, M.; Lejeune, S.; Curtis, A. R. J.; Strachan, T.; Lindsay, S.; Harris, A. L.: Isolation of a full-length human WNT7A gene implicated in limb development and cell transformation, and mapping to chromosome 3p25. *Gene* 189: 25–29, 1997.
- [66075] 4405. Ikegawa, S.; Kumano, Y.; Okui, K.; Fujiwara, T.; Takahashi, E.; Nakamura, Y.: Isolation, characterization and chromosomal assignment of the human WNT7A gene. *Cytogenet. Cell Genet.* 74: 149–152, 1996.
- [66076] 4406. Li, S.; Chiang, T.-C.; Davis, G. R.; Williams, R. M.; Wilson, V. P.; McLachlan, J. A.: Decreased expression of Wnt7a mRNA is inversely associated with the expression of estrogen receptor- $\alpha$  in human uterine leiomyoma. *J. Clin. Endocr. Metab.* 86: 454–457, 2001.
- [66077] 4407. Pichaud, F.; Delage-Mourroux, R.; Pidoux, E.; Julienne, A.; Rousseau-Merck, M.-F.: Chromosomal localization of the type-I 15-PGDH gene to 4q34–q35.

Hum.Genet. 99: 279–281, 1997.

[66078] 4408.Walder, K.; Norman, R. A.; Hanson, R. L.; Schrauwen, P.; Neverova,M.; Jenkinson, C. P.; Easlick, J.; Warden, C. H.; Pecqueur, C.; Raimbault,S.; Ricquier, D.; Harper, M.; Silver, K.; Shuldiner, A. R.; Solanes,G.; Lowell, B. B.; Chung, W. K.; Leibel, R. L.; Pratley, R.; Ravussin,E.: Association between uncoupling protein polymorphisms (UCP2–UCP3)and energy metabolism/obesity in Pima Indians. Hum. Molec. Genet. 7:1431–1435, 1998.

[66079] 4409.Lindsay, E. A.; Rizzu, P.; Antonacci, R.; Jurecic, V.; Delmas–Mata,J.; Lee, C.–C.; Kim, U.–J.; Scambler, P. J.; Baldini, A.: A transcriptionmap in the CATCH22 critical region: identification, mapping, and orderingof four novel transcripts expressed in heart. Genomics 32: 104–112,1996.

[66080] 4410.Takai, S.; Hinoda, Y.; Adachi, T.; Imai, K.; Oshima, M.: A humanUPD (sic)–GalNAc: polypeptide, N–acetylgalactosaminyltransferase type1 gene is located at the chromosomal region 18q12.1. Hum. Genet. 99:293–294, 1997.

[66081] 4411.White, T.; Bennett, E. P.; Takio, K.; Sorensen, T.; Bonding, N.;Clausen, H.: Purification and cDNA cloning of a human UDP–

N-  
acetyl-al-  
pha-  
D-galac-  
tosamine:polypeptideN-acetylgalactosaminyltransferase. J.  
Biol. Chem. 270: 24156–24165,1995.

- [66082] 4412.Candau, R.; Moore, P. A.; Wang, L.; Barlev, N.; Ying, C. Y.; Rosen,C. A.; Berger, S. L.: Identification of human proteins functionally conserved with the yeast putative adaptors ADA2 and GCN5. Molec.Cell. Biol. 16: 593–602, 1996.
- [66083] 4413.Carter, K. C.; Wang, L.; Shell, B. K.; Zamir, I.; Berger, S. L.;Moore, P. A.: The human transcriptional adaptor genes TADA2L andGCN5L2 colocalize to chromosome 17q12–q21 and display a similar tissueexpression pattern. Genomics 40: 497–500, 1997.
- [66084] 4414.Ogryzko, V. V.; Kotani, T.; Zhang, X.; Schiltz, R. L.; Howard,T.; Yang, X.–J.; Howard, B. H.; Qin, J.; Nakatani, Y.: Histone–likeTAFs within the PCAF histone acetylase complex. Cell 94: 35–44,1998.
- [66085] 4415.Struhl, K.; Moqtaderi, Z.: The TAFs in the HAT. Cell 94: 1–4,1998.
- [66086] 4416.Imhof, M. O.; McDonnell, D. P.: Yeast RSP5 and its

human homolog hRPF1 potentiate hormone-dependent activation of transcription by human progesterone and glucocorticoid receptors. *Molec. Cell. Biol.* 16:2594–2605, 1996.

[66087] 4417. Kumar, S.; Harvey, K. F.; Kinoshita, M.; Copeland, N. G.; Noda, M.; Jenkins, N. A.: cDNA cloning, expression analysis, and mapping of the mouse Nedd4 gene. *Genomics* 40: 435–443, 1997.

[66088] 4418. Cabin, D. E.; Gardiner, K.; Reeves, R. H.: Molecular genetic characterization and comparative mapping of the human PCP4 gene. *Somat. Cell Molec. Genet.* 22: 167–175, 1996.

[66089] 4419. Chen, H.; Bouras, C.; Antonarakis, S. E.: Cloning of the cDNA for a human homolog of the rat PEP-19 gene and mapping to chromosome 21q22.2–q22.3. *Hum. Genet.* 98: 672–677, 1996.

[66090] 4420. Hubert, R. S.; Korenberg, J. R.: PCP4 maps between D21S345 and P31P10SP6 on chromosome 21q22.2–q22.3. *Cytogenet. Cell Genet.* 78:44–45, 1997.

[66091] 4421. Collum, R. G.; Fisher, P. E.; Datta, M.; Mellis, S.; Thiele, C.; Huebner, K.; Croce, C. M.; Israel, M. A.; Theil, T.; Moroy, T.; DePinho, R.; Alt, F. W.: A novel POU homeodomain gene specifically expressed in cells of the devel-

oping mammalian nervous system. *Nucleic Acids Res.* 20: 4919–4925, 1992.

[66092] 4422. Gerrero, M. R.; McEvilly, R. J.; Turner, E.; Lin, C. R.; O'Connell, S.; Jenne, K. J.; Hobbs, M. V.; Rosenfeld, M. G.: Brn-3.0: a POU-domain protein expressed in the sensory immune and endocrine systems that functions on elements distinct from known octamer motifs. *Proc. Nat. Acad. Sci.* 90: 10841–10845, 1993.

[66093] 4423. Still, I. H.; Cowell, J.: The Brn-3a transcription factor gene (POU4F1) maps close to the locus for the variant late infantile form of neuronal ceroid-lipofuscinosis. *Cytogenet. Cell Genet.* 74: 225–226, 1996.

[66094] 4424. Block, M. R.; Glick, B. S.; Wilcox, C. A.; Wieland, F. T.; Rothman, J. E.: Purification of an N-ethylmaleimide-sensitive protein catalyzing vesicular transport. *Proc. Nat. Acad. Sci.* 85: 7852–7856, 1988.

[66095] 4425. Glick, B. S.; Rothman, J. E.: Possible role for fatty acyl-coenzyme A in intracellular protein transport. *Nature* 326: 309–312, 1987.

[66096] 4426. Hoyle, J.; Phelan, J. P.; Bermingham, N.; Fisher, E. M. C.: Localization of human and mouse N-ethylmaleimide-sensitive factor (NSF) gene: a two-domain member of the AAA family that is involved in membrane



fusion. *Mammalian Genome* 7: 850–852, 1996.

[66097] 4427. Rothman, J. E.: Mechanisms of intracellular protein transport. *Nature* 372:55–63, 1994.

[66098] 4428. Wilson, D. W.; Wilcox, C. A.; Flynn, G. C.; Chen, E.; Kuang, W.-J.; Henzel, W. J.; Block, M. R.; Ullrich, A.; Rothman, J. E.: A fusion protein required for vesicle-mediated transport in both mammalian cells and yeast. *Nature* 339: 355–359, 1989.

[66099] 4429. Dikstein, R.; Zhou, S.; Tjian, R.: Human TAF(II)105 is a cell type-specific TFIID subunit related to hTAF(II)130. *Cell* 87: 137–146, 1996.

[66100] 4430. Freiman, R. N.; Albright, S. R.; Zheng, S.; Sha, W. C.; Hammer, R. E.; Tjian, R.: Requirement of tissue-selective TBP-associated factor TAFII105 in ovarian development. *Science* 293: 2084–2087, 2001.

[66101] 4431. Allikmets, R.; Gerrard, B.; Hutchinson, A.; Dean, M.: Characterization of the human ABC superfamily: isolation and mapping of 21 new genes using the expressed sequence tags database. *Hum. Molec. Genet.* 5:1649–1655, 1996.

[66102] 4432. Rizzu, P.; Lindsay, E. A.; Taylor, C.; O'Donnell, H.; Levy, A.; Scambler, P.; Baldini, A.: Cloning and comparative mapping of a gene from the commonly deleted region of

DiGeorge and velocardiofacial syndromes conserved in *C. elegans*. *Mammalian Genome* 7: 639–643, 1996.

- [66103] 4433. Bennett, E. P.; Hassan, H.; Clausen, H.: cDNA cloning and expression of a novel human UDP-N-acetyl- $\alpha$ -D-galactosamine. *J. Biol. Chem.* 271:17006–17012, 1996.
- [66104] 4434. Bennett, E. P.; Weghuis, D. O.; Merks, G.; Geurts van Kessel, A.; Eiberg, H.; Clausen, H.: Genomic organization and chromosomal localization of three members of the UDP-N-acetylgalactosamine:polypeptide N-acetylgalactosaminyltransferase family. *Glycobiology* 8: 547–555, 1998.
- [66105] 4435. Zara, J.; Hagen, F. K.; Ten Hagen, K. G.; Van Wuyckhuyse, B. C.; Tabak, L. A.: Cloning and expression of mouse UDP-GalNAc:polypeptide N-acetylgalactosaminyltransferase-T3. *Biochem. Biophys. Res. Commun.* 228:38–44, 1996.
- [66106] 4436. Zhao, Y.-Y.; Liu, Y.; Stan, R.-V.; Fan, L.; Gu, Y.; Dalton, N.; Chu, P.-H.; Peterson, K.; Ross, J., Jr.; Chien, K. R.: Defects in caveolin-1 cause dilated cardiomyopathy and pulmonary hypertension in knockout mice. *Proc. Nat. Acad. Sci.* 99: 11375–11380, 2002.
- [66107] 4437. Mok, S. C.; Wong, K.-K.; Chan, R. K. W.; Lau, C. C.;

Tsao, S.-W.; Knapp, R. C.; Berkowitz, R. S.: Molecular cloning of differentially expressed genes in human epithelial ovarian cancer. *Gynecol. Oncol.* 52:247–252, 1994.

[66108] 4438. Xu, X.-X.; Yang, W.; Jackowski, S.; Rock, C. O.: Cloning of a novel phosphoprotein regulated by colony-stimulating factor 1 shares a domain with the *Drosophila* disabled gene product. *J. Biol. Chem.* 270:14184–14191, 1995.

[66109] 4439. Galbiati, F.; Volonte, D.; Gil, O.; Zanazzi, G.; Salzer, J. L.; Sargiacomo, M.; Scherer, P. E.; Engelman, J. A.; Schlegel, A.; Parenti, M.; Okamoto, T.; Lisanti, M. P.: Expression of caveolin-1 and -2 in differentiating PC12 cells and dorsal root ganglion neurons: caveolin-2 is up-regulated in response to cell injury. *Proc. Nat. Acad. Sci.* 95:10257–10262, 1998.

[66110] 4440. Scherer, P. E.; Okamoto, T.; Chun, M.; Nishimoto, I.; Lodish, H. F.; Lisanti, M. P.: Identification, sequence, and expression of caveolin-2 defines a caveolin gene family. *Proc. Nat. Acad. Sci.* 93: 131–135, 1996.

[66111] 4441. Campbell, H. D.; Webb, G. C.; Fountain, S.; Young, I. G.: The human PIN1 peptidyl-prolyl cis/trans isomerase gene maps to human chromosome 19p13 and the closely related PIN1L gene to 1p31. *Genomics* 44:157–162, 1997.

- [66112] 4442.Liou, Y.-C.; Ryo, A.; Huang, H.-K.; Lu, P.-J.; Bronson, R.; Fujimori, F.; Uchida, T.; Hunter, T.; Lu, K. P.: Loss of Pin1 function in the mouse causes phenotypes resembling cyclin D1-null phenotypes. *Proc. Nat. Acad. Sci.* 99: 1335–1340, 2002.
- [66113] 4443.Lu, K. P.; Hanes, S. D.; Hunter, T.: A human peptidyl-prolyl isomerase essential for regulation of mitosis. *Nature* 380: 544–547, 1996.
- [66114] 4444.Lu, P.-J.; Wulf, G.; Zhou, X. Z.; Davies, P.; Lu, K. P.: The prolyl isomerase Pin1 restores the function of Alzheimer-associated phosphorylated tau protein. *Nature* 399: 784–788, 1999.
- [66115] 4445.Maleszka, R.; Hanes, S. D.; Hackett, R. L.; de Couet, H. G.; Gabor Miklos, G. L.: The *Drosophila melanogaster* dodo (dod) gene, conserved in humans, is functionally interchangeable with the ESS1 cell division gene of *Saccharomyces cerevisiae*. *Proc. Nat. Acad. Sci.* 93: 447–451, 1996.
- [66116] 4446.Winkler, K. E.; Swanson, K. I.; Kornbluth, S.; Means, A. R.: Requirement of the prolyl isomerase Pin1 for the replication checkpoint. *Science* 287:1644–1647, 2000.
- [66117] 4447.Takahashi, T.; Fournier, A.; Nakamura, F.; Wang, L.-H.; Murakami, Y.; Kalb, R. G.; Fujisawa, H.; Strittmatter,

S. M.: Plexin–neuropilin–1 complexes form functional semaphorin–3A receptors. *Cell* 99: 59–69, 1999.

[66118] 4448. Santagata, S.; Boggon, T. J.; Baird, C. L.; Gomez, C. A.; Zhao, J.; Shan, W. S.; Myszka, D. G.; Shapiro, L.: G–protein signaling through tubby proteins. *Science* 292: 2041–2050, 2001.

[66119] 4449. Sasaki, T.; Irie–Sasaki, J.; Horie, Y.; Bachmaier, K.; Fata, J. E.; Li, M.; Suzuki, A.; Bouchard, D.; Ho, A.; Redston, M.; Gallinger, S.; Khokha, R.; Mak, T. W.; Hawkins, P. T.; Stephens, L.; Scherer, S. W.; Tsao, M.; Penninger, J. M.: Colorectal carcinomas in mice lacking the catalytic subunit of PI(3)K–gamma. *Nature* 406: 897–902, 2000.

[66120] 4450. Sasaki, T.; Irie–Sasaki, J.; Jones, R. G.; Oliveira–dos–Santos, A. J.; Stanford, W. L.; Bolon, B.; Wakeham, A.; Itie, A.; Bouchard, D.; Kozieradzki, I.; Joza, N.; Mak, T. W.; Ohashi, P. S.; Suzuki, A.; Penninger, J. M.: Function of PI3K–gamma in thymocyte development, T cell activation, and neutrophil migration. *Science* 287: 1040–1046, 2000.

[66121] 4451. Stoyanov, B.; Volinia, S.; Hanck, T.; Rubio, I.; Loubtchenkov, M.; Malek, D.; Stoyanova, S.; Vanhaesebroeck, B.; Dhand, R.; Nurnberg, B.; Gierschik, P.; Seedorf, K.; Hsuan, J. J.; Waterfield, M. D.; Wetzker, R.: Cloning and characterization of a G protein–activated human phos–

phoinositide-3kinase. Science 269: 690-693, 1995.

- [66122] 4452.Hotten, G.; Neidhardt, H.; Schneider, C.; Pohl, J.: Cloning of a new member of the TGF-beta family: a putative new activin betaC chain. Biochem. Biophys. Res. Commun. 206: 608-613, 1995.
- [66123] 4453.Ling, N.; Ying, S. Y.; Ueno, N.; Shimasaki, S.; Esch, F.; Hotta, M.; Guillemin, R.: Pituitary FSH is released by a heterodimer of the beta-subunits from the two forms of inhibin. Nature 321: 779-782, 1986.
- [66124] 4454.Schmitt, J.; Hotten, G.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Pohl, J.; Schrewe, H.: Structure, chromosomal localization, and expression analysis of the mouse inhibin/activin betaC (Inhbc) gene. Genomics 32: 358-366, 1996.
- [66125] 4455.Savitsky, K.; Ziv, Y.; Bar-Shira, A.; Gilad, S.; Tagle, D. A.; Smith, S.; Uziel, T.; Sfez, S.; Nahmias, J.; Sartiel, A.; Eddy, R.L.; Shows, T. B.; Collins, F. S.; Shiloh, Y.; Rotman, G.: A human gene (DDX10) encoding a putative DEAD-box RNA helicase at 11q22-q23. Genomics 33:199-206, 1996.
- [66126] 4456.Albertsen, H. M.; Smith, S. A.; Melis, R.; Williams, B.; Holik, P.; Stevens, J.; White, R.: Sequence, genomic structure, and chromosomal assignment of human DOC-2. Ge-

nomics 33: 207–213, 1996.

[66127] 4457.Mok, S. C.; Chan, W. Y.; Wong, K. K.; Cheung, K. K.; Lau, C. C.; Ng, S. W.; Baldini, A.; Colitti, C. V.; Rock, C. O.; Berkowitz, R.S.: DOC-2, a candidate tumor suppressor gene in human epithelial ovarian cancer. *Oncogene* 16: 2381–2387, 1998.

[66128] 4458.Kimura, T.; Arakawa, Y.; Inoue, S.; Fukushima, Y.; Kondo, I.; Koyama, K.; Hosoi, T.; Orimo, A.; Muramatsu, M.; Nakamura, Y.; Abe, T.; Inazawa, J.: The brain finger protein gene (ZNF179), a member of the RING finger family, maps within the Smith–Magenis syndrome region at 17p11.2. *Am.J. Med. Genet.* 69: 320–324, 1997.

[66129] 4459.Matsuda, Y.; Inue, S.; Seki, N.; Hosoi, T.; Orimo, A.; Muramatsu, M.; Hori, T.: Chromosome mapping of human (ZNF179), mouse, and rat genes for brain finger protein (bfp), a member of the RING finger family. *Genomics* 33: 325–327, 1996.

[66130] 4460.Ambrose, H. J.; Blake, D. J.; Nawrotzki, R. A.; Davies, K. E.: Genomic organization of the mouse dystrobrevin gene: comparative analysis with the dystrophin gene. *Genomics* 39: 359–369, 1997.

[66131] 4461.Blake, D. J.; Nawrotzki, R.; Loh, N. Y.; Gorecki, D. C.; Davies, K. E.: Beta–dystrobrevin, a member of the dys–

trophin-related protein family. *Proc. Nat. Acad. Sci.* 95: 241–246, 1998.

- [66132] 4462. Ichida, F.; Tsubata, S.; Bowles, K. R.; Haneda, N.; Uese, K.; Miyawaki, T.; Dreyer, W. J.; Messina, J.; Li, H.; Bowles, N. E.; Towbin, J. A.: Novel gene mutations in patients with left ventricular noncompaction or Barth syndrome. *Circulation* 103: 1256–1263, 2001.
- [66133] 4463. England, S. K.; Uebele, V. N.; Shear, H.; Kodali, J.; Bennett, P. B.; Tamkun, M. M.: Characterization of a voltage-gated K<sup>+</sup> channel beta subunit expressed in human heart. *Proc. Nat. Acad. Sci.* 92: 6309–6313, 1995.
- [66134] 4464. Jones, J. M.; Bentley, E.; Meisler, M. H.; Darling, S. M.: Genetic mapping of the voltage-gated shaker potassium channel beta subunit *Kcnab1* to mouse chromosome 3. *Mammalian Genome* 9: 260 only, 1998.
- [66135] 4465. Leicher, T.; Bähring, R.; Isbrandt, D.; Pongs, O.: Co-expression of the *KCNA3B* gene product with Kv1.5 leads to a novel A-type potassium channel. *J. Biol. Chem.* 273: 35095–35101, 1998.
- [66136] 4466. Leicher, T.; Roeper, J.; Weber, K.; Wang, X.; Pongs, O.: Structural and functional characterization of human potassium channel subunit beta-1 (*KCNA1B*). *Neuropharmacology* 35: 787–795, 1996.



- [66137] 4467. McCormack, K.; McCormack, T.; Tanouye, M.; Rudy, B.; Stuhmer, W.: Alternative splicing of the human Shaker K<sup>+</sup> channel beta-1 gene and functional expression of the beta-2 gene product. *FEBS Lett.* 370:32–36, 1995.
- [66138] 4468. Schultz, D.; Litt, M.; Smith, L.; Thayer, M.; McCormack, K.: Localization of two potassium channel beta subunit genes, KCNA1B and KCNA2B. *Genomics* 31:389–391, 1996.
- [66139] 4469. Collin, G. B.; Nishina, P. M.; Marshall, J. D.; Naggert, J. K.: Human DCTN1: genomic structure and evaluation as a candidate for Alstrom syndrome. *Genomics* 53: 359–364, 1998.
- [66140] 4470. Holzbaur, E. L. F.; Tokito, M. K.: Localization of the DCTN1 gene encoding p150(Glued) to human chromosome 2p13 by fluorescence in situ hybridization. *Genomics* 31: 398–399, 1996.
- [66141] 4471. Holzbaur, E. L. F.; Vallee, R. B.: Dyneins: molecular structure and cellular function. *Ann. Rev. Cell Biol.* 10: 339–372, 1994.
- [66142] 4472. Jang, W.; Weber, J. S.; Tokito, M. K.; Holzbaur, E. L. F.; Meisler, M. H.: Mouse p150(Glued) (dynactin 1) cDNA sequence and evaluation as a candidate for the neuromuscular disease mutation mnd2. *Biochem. Biophys. Res.*

Commun. 231: 344–347, 1997.

- [66143] 4473.Korthaus, D.; Wedemeyer, N.; Lengeling, A.; Ronsiek, M.; Jockusch,H.; Schmitt-John, T.: Integrated radiation hybrid map of human chromosome2p13: possible involvement of dynactin in neuromuscular diseases. Genomics 43:242–244, 1997.
- [66144] 4474.Pushkin, A.; Abuladze, N.; Newman, D.; Tatishchev, S.; Kurtz, I.: Genomic organization of the DCTN1–SLC4A5 locus encoding both NBC4and p150(Glued). Cytogenet. Cell Genet. 95: 163–168, 2001.
- [66145] 4475.Miyata, A.; Hara, S.; Yokoyama, C.; Inoue, H.; Ullrich, V.; Tanabe,T.: Molecular cloning and expression of human prostacyclin synthase. Biochem.Biophys. Res. Commun. 200: 1728–1734, 1994.
- [66146] 4476.Inoue, H.; Ishii, H.; Alder, H.; Snyder, E.; Druck, T.; Huebner,K.; Croce, C. M.: Sequence of the FRA3B common fragile region: implicationsfor the mechanism of FHIT deletion. Proc. Nat. Acad. Sci. 94: 14584–14589,1997.
- [66147] 4477.Morikawa, H.; Nakagawa, Y.; Hashimoto, K.; Niki, M.; Egashira,Y.; Hirata, I.; Katsu, K.; Akao, Y.: Frequent altered expressionof fragile histidine triad protein in human colorectal adenomas. Biochem.Biophys. Res. Commun. 278: 205–210, 2000.

- [66148] 4478. Nelson, D. R.; Koymans, L.; Kamataki, T.; Stegeman, J. J.; Feyereisen, R.; Waxman, D. J.; Waterman, M. R.; Gotoh, O.; Coon, M. J.; Estabrook, R. W.; Gunsalus, I. C.; Nebert, D. W.: P450 superfamily: update on new sequences, gene mapping, accession numbers and nomenclature. *Pharmacogenetics* 6:1–42, 1996.
- [66149] 4479. Wang, L.-H.; Chen, L.: Organization of the gene encoding human prostacyclin synthase. *Biochem. Biophys. Res. Commun.* 226: 631–637, 1996.
- [66150] 4480. Yokoyama, C.; Yabuki, T.; Inoue, H.; Tone, Y.; Hara, S.; Hatae, T.; Nagata, M.; Takahashi, E.-I.; Tanabe, T.: Human gene encoding prostacyclin synthase (PTGIS): genomic organization, chromosomal localization, and promoter activity. *Genomics* 36: 296–304, 1996.
- [66151] 4481. Brindle, N. P. J.; Holt, M. R.; Davies, J. E.; Price, C. J.; Critchley, D. R.: The focal-adhesion vasodilator-stimulated phosphoprotein (VASP) binds to the proline-rich domain in vinculin. *Biochem. J.* 318: 753–757, 1996.
- [66152] 4482. Haffner, C.; Jarchau, T.; Reinhard, M.; Hoppe, J.; Lohmann, S. M.; Walter, V.: Molecular cloning, structural analysis and functional expression of the proline-rich focal adhesion and microfilament-associated protein VASP. *EMBO J.* 14: 19–27, 1995.

- [66153] 4483.Zimmer, M.; Fink, T.; Fischer, L.; Hauser, W.; Scherer, K.; Lichter,P.; Walter, U.: Cloning of the VASP (vasodilator–stimulated phosphoprotein)genes in human and mouse: structure, sequence, and chromosomal localization. *Genomics* 36:227–233, 1996.
- [66154] 4484.Brezin, A. P.; Adam, M. F.; Belmouden, A.; Lureau, M.–A.; Chaventre,A.; Copin, B.; Gomez, L.; Dupont de Dinechin, S.; Berkani, M.; Valtot,F.; Rouland, J.–F.; Dascotte, J.–C.; Bach, J.–F.; Garchon, H.–J.:Founder effect in GLC1A–linked familial open–angle glaucoma in northern–France. *Am. J. Med. Genet.* 76: 438–445, 1998.
- [66155] 4485.Colomb, E.; Nguyen, T. D.; Bechetoille, A.; Dascotte, J.–C.; Valtot,F.; Brezin, A. P.; Berkani, M.; Copin, B.; Gomez, L.; Polansky, J.R.; Garchon, H.–J.: Association of a single nucleotide polymorphismin the TIGR/MYOCILIN gene promoter with the severity of primary open–angleglaucoma. *Clin. Genet.* 60: 220–225, 2001.
- [66156] 4486.Escribano, J.; Ortego, J.; Coca–Prados, M.: Isolation and characterizationof cell–specific cDNA clones from a subtractive library of the ocularciliary body of a single normal human donor: transcription and synthesisof plasma proteins. *J. Biochem.* 118: 921–931, 1995.
- [66157] 4487.Fingert, J. H.; Ying, L.; Swiderski, R. E.; Nystuen, A.

M.; Arbour, N. C.; Alward, W. L. M.; Sheffield, V. C.; Stone, E. M.: Characterization and comparison of the human and mouse GLC1A glaucoma genes. *Genome Res.* 8: 377–384, 1998.

[66158] 4488. Fleck, B. W.; Cullen, J. F.: Autosomal dominant juvenile onset glaucoma affecting six generations in an Edinburgh family. (Letter) *Brit. J. Ophthalmol.* 70: 715 only, 1986.

[66159] 4489. Kubota, R.; Noda, S.; Wang, Y.; Minoshima, S.; Asakawa, S.; Kudoh, J.; Mashima, Y.; Oguchi, Y.; Shimizu, N.: A novel myosin-like protein (myocilin) expressed in the connecting cilium of the photoreceptor: molecular cloning, tissue expression, and chromosomal mapping. *Genomics* 41:360–369, 1997.

[66160] 4490. Fingert, J. H.; Heon, E.; Liebmann, J. M.; Yamamoto, T.; Craig, J. E.; Rait, J.; Kawase, K.; Hoh, S.-T.; Buys, Y. M.; Dickinson, J.; Hockey, R. R.; Williams-Lyn, D.; Trope, G.; Kitazawa, Y.; Ritch, R.; Mackey, D. A.; Alward, W. L. M.; Sheffield, V. C.; Stone, E. M.: Analysis of myocilin mutations in 1703 glaucoma patients from five different populations. *Hum. Molec. Genet.* 8: 899–905, 1999.

[66161] 4491. Mansergh, F. C.; Kenna, P. F.; Ayuso, C.; Kiang, A.-S.; Humphries, P.; Farrar, G. J.: Novel mutations in the TIGR gene in early and late onset open angle glaucoma.

Hum. Mutat. 11: 244–251, 1998.

[66162] 4492. Michels–Rautenstrauss, K. G.; Mardin, C. Y.; Budde, W. M.; Liehr, T.; Polansky, J.; Nguyen, T.; Timmerman, V.; Van Broeckhoven, C.; Naumann, G. O. H.; Pfeiffer, R. A.; Rautenstrauss, B. W.: Juvenile open angle glaucoma: fine mapping of the TIGR gene to 1q24.3–q25.2 and mutation analysis. Hum. Genet. 102: 103–106, 1998.

[66163] 4493. Morissette, J.; Clepet, C.; Moisan, S.; Dubois, S.; Winstall, E.; Vermeeren, D.; Nguyen, T. D.; Polansky, J. R.; Cote, G.; Anctil, J.–L.; Amyot, M.; Plante, M.; Falardeau, P.; Raymond, V.: Homozygotes carrying an autosomal dominant TIGR mutation do not manifest glaucoma. (Letter) Nature Genet. 19: 319–321, 1998.

[66164] 4494. Shepard, A. R.; Jacobson, N.; Fingert, J. H.; Stone, E. M.; Sheffield, V. C.; Clark, A. F.: Delayed secondary glucocorticoid responsiveness of MYOC in human trabecular meshwork cells. Invest. Ophthalmol. Vis. Sci. 42: 3173–3181, 2001.

[66165] 4495. Stoilova, D.; Child, A.; Brice, G.; Crick, R. P.; Fleck, B. W.; Sarfarazi, M.: Identification of a new 'TIGR' mutation in a family with juvenile-onset primary open angle glaucoma. Ophthalmic Genet. 18: 109–118, 1997.

[66166] 4496. Stone, E. M.: Personal Communication. Iowa City,

Iowa 6/16/1999.

[66167] 4497.Stone, E. M.: Personal Communication. Iowa City, Iowa 11/20/1997.

[66168] 4498.Suzuki, Y.; Shirato, S.; Taniguchi, F.; Ohara, K.; Nishimaki,K.; Ohta, S.: Mutations in the TIGR gene in familial primary open-angleglaucoma in Japan. (Letter) Am. J. Hum. Genet. 1202–1204, 1997.

[66169] 4499.Vasconcellos, J. P. C.; Melo, M. B.; Costa, V. P.; Tsukumo, D.M. L.; Basseres, D. S.; Bordin, S.; Saad, S. T. O.; Costa, F. F.:Novel mutation in the MYOC gene in primary open angle glaucoma patients. J.Med. Genet. 37: 301–303, 2000.

[66170] 4500.Wiggs, J. L.; Allingham, R. R.; Vollrath, D.; Jones, K. H.; DeLa Paz, M.; Kern, J.; Patterson, K.; Babb, V. L.; Del Bono, E. A.;Broomer, B. W.; Pericak–Vance, M. A.; Haines, J. L.: Prevalence ofmutations in TIGR/myocilin in patients with adult and juvenile primaryopen-angle glaucoma. (Letter) Am. J. Hum. Genet. 63: 1549–1552,1998.

[66171] 4501.Zhou, Z.; Vollrath, D.: A cellular assay distinguishes normaland mutant TIGR/myocilin protein. Hum. Molec. Genet. 8: 2221–2228,1999.

[66172] 4502.Hackam, A. S.; Yassa, A. S.; Singaraja, R.; Metzler, M.; Gutekunst,C.–A.; Gan, L.; Warby, S.; Wellington, C. L.;

Vaillancourt, J.; Chen, N.; Gervais, F. G.; Raymond, L.; Nicholson, D. W.; Hayden, M. R.: Huntingtin interacting protein 1 induces apoptosis via a novel caspase-dependent death effector domain. *J. Biol. Chem.* 275: 41299–41308, 2000.

[66173] 4503. Himmelbauer, H.; Wedemeyer, N.; Haaf, T.; Wanker, E. E.; Schalkwyk, L. C.; Lehrach, H.: IRS-PCR-based genetic mapping of the huntingtin interacting protein gene (HIP1) on mouse chromosome 5. *Mammalian Genome* 9: 26–31, 1998.

[66174] 4504. Manabe, T.; Noda, Y.; Mamiya, T.; Katagiri, H.; Houtani, T.; Nishi, M.; Noda, T.; Takahashi, T.; Sugimoto, T.; Nabeshima, T.; Takeshima, H.: Facilitation of long-term potentiation and memory in mice lacking nociceptin receptors. *Nature* 394: 577–581, 1998.

[66175] 4505. Mollereau, C.; Simons, M.-J.; Soularue, P.; Liners, F.; Vassart, G.; Meunier, J.-C.; Parmentier, M.: Structure, tissue distribution, and chromosomal localization of the prepronociceptin gene. *Proc. Nat. Acad. Sci.* 93: 8666–8700, 1996.

[66176] 4506. Nishi, M.; Houtani, T.; Noda, Y.; Mamiya, T.; Sato, K.; Doi, T.; Kuno, J.; Takeshima, H.; Nukada, T.; Nabeshima, T.; Yamashita, T.; Noda, T.; Sugimoto, T.: Unrestrained no-



ciceptive response and disregulation of hearing ability in mice lacking the nociceptin/orphanin FQ receptor. *EMBO J.* 16: 1858–1864, 1997.

- [66177] 4507. Okuda–Ashitaka, E.; Minami, T.; Tachibana, S.; Yoshihara, Y.; Nishiuchi, Y.; Kimura, T.; Ito, S.: Nocistatin, a peptide that blocks nociceptin action in pain transmission. *Nature* 392: 286–289, 1998.
- [66178] 4508. Pan, Z. Z.; Hirakawa, N.; Fields, H. L.: A cellular mechanism for the bidirectional pain–modulating actions of orphanin FQ/nociceptin. *Neuron* 26: 515–522, 2000.
- [66179] 4509. Serhan, C. N.; Fierro, I. M.; Chiang, N.; Pouliot, M.: Cutting edge: nociceptin stimulates neutrophil chemotaxis and recruitment: inhibition by aspirin–triggered–15–epi–lipoxin A(4). *J. Immun.* 166: 3650–3654, 2001.
- [66180] 4510. Kanai, N.; Lu, R.; Satriano, J. A.; Bao, Y.; Wolkoff, A. W.; Schuster, V. L.: Identification and characterization of a prostaglandin transporter. *Science* 268: 866–869, 1995.
- [66181] 4511. Lu, R.; Kanai, N.; Bao, Y.; Schuster, V. L.: Cloning, in vitro expression, and tissue distribution of a human prostaglandin transporter cDNA (hPGT). *J. Clin. Invest.* 98: 1142–1149, 1996.
- [66182] 4512. Lu, R.; Schuster, V. L.: Molecular cloning of the gene

for the human prostaglandin transporter hPGT: gene organization, promoter activity, and chromosomal localization. *Biochem. Biophys. Res. Commun.* 246:805–812, 1998.

[66183] 4513. Frucht, D. M.; Aringer, M.; Galon, J.; Danning, C.; Brown, M.; Fan, S.; Centola, M.; Wu, C.-Y.; Yamada, N.; Galaway, H. E.; O'Shea, J. J.: Stat4 is expressed in activated peripheral blood monocytes, dendritic cells, and macrophages at sites of Th1-mediated inflammation. *J. Immun.* 164: 4659–4664, 2000.

[66184] 4514. Loughna, S.; Sato, T. N.: A combinatorial role of angiopoietin-1 and orphan receptor TIE1 pathways in establishing vascular polarity during angiogenesis. *Molec. Cell* 7: 233–239, 2001.

[66185] 4515. Sato, T. N.; Tozawa, Y.; Deutsch, U.; Wolburg-Buchholz, K.; Fujiwara, Y.; Gendron-Maguire, M.; Gridley, T.; Wolburg, H.; Risau, W.; Qin, Y.: Distinct roles of the receptor tyrosine kinases Tie-1 and Tie-2 in blood vessel formation. *Nature* 376: 70–74, 1995.

[66186] 4516. Tafuri, A.; Shahinian, A.; Bladt, F.; Yoshinaga, S. K.; Jordana, M.; Wakeham, A.; Boucher, L.-M.; Bouchard, D.; Chan, V. S. F.; Duncan, G.; Odermatt, B.; Ho, A.; Itie, A.; Horan, T.; Whoriskey, J. S.; Pawson, T.; Penninger, J. M.; Ohashi, P. S.; Mak, T. W.: ICOS is essential for effective T-

helper-cell responses. *Nature* 409: 105–109, 2001.

- [66187] 4517. Yoshinaga, S. K.; Whoriskey, J. S.; Khare, S. D.; Sarmiento, U.; Guo, J.; Horan, T.; Shih, G.; Zhang, M.; Coccia, M. A.; Kohno, T.; Tafuri-Bladt, A.; Brankow, D.; and 14 others: T-cell co-stimulation through B7RP-1 and ICOS. *Nature* 402: 827–832, 1999.
- [66188] 4518. Paige, A. J. W.; Taylor, K. J.; Taylor, C.; Hillier, S. G.; Farrington, S.; Scott, D.; Porteous, D. J.; Smyth, J. F.; Gabra, H.; Watson, J. E. V.: WWOX: a candidate tumor suppressor gene involved in multiple tumor types. *Proc. Nat. Acad. Sci.* 98: 11417–11422, 2001.
- [66189] 4519. Ried, K.; Finnis, M.; Hobson, L.; Mangelsdorf, M.; Dayan, S.; Nancarrow, J. K.; Woollatt, E.; Kremmidiotis, G.; Gardner, A.; Venter, D.; Baker, E.; Richards, R. I.: Common chromosomal fragile site FRA16D sequence: identification of the FOR gene spanning FRA16D and homozygous deletions and translocation breakpoints in cancer cells. *Hum. Molec. Genet.* 9: 1651–1663, 2000.
- [66190] 4520. Clark, S. W.; Staub, O.; Clark, I. B.; Holzbaur, E. L. F.; Paschal, B. M.; Vallee, R. B.; Meyer, D. I.: Beta-centractin: characterization and distribution of a new member of the contractin family of actin-related proteins. *Molec. Biol. Cell* 5: 1301–1310, 1994.

- [66191] 4521.Elsea, S. H.; Clark, I. B.; Juyal, R. C.; Meyer, D. J.; Meyer,D. I.; Patel, P. I.: Assignment of beta-centractin (CTRN2) to human chromosome 2 bands q11.1–q11.2 with somatic cell hybrids and in situ hybridization. *Cytogenet. Cell Genet.* 84: 48–49, 1999.
- [66192] 4522.Ho, A. M.; Johnson, M. D.; Kingsley, D. M.: Role of the mouse ank gene in control of tissue calcification and arthritis. *Science* 289:265–270, 2000.
- [66193] 4523.Yu, H.; Peters, J.–M.; King, R. W.; Page, A. M.; Hieter, P.; Kirschner,M. W.: Identification of a cullin homology region in a subunit of the anaphase–promoting complex. *Science* 279: 1219–1222, 1998.
- [66194] 4524.Zhao, N.; Lai, F.; Fernald, A. A.; Eisenbart, J. D.; Espinosa,R., III.; Wang, P. W.; Le Beau, M. M.: Human CDC23: cDNA cloning,mapping to 5q31, genomic structure, and evaluation as a candidate tumor suppressor gene in myeloid leukemias. *Genomics* 53: 184–190,1998.
- [66195] 4525.Brambilla, R.; Draetta, G.: Molecular cloning of PISSLRE, a novel putative member of the cdk family of protein serine/threonine kinases. *Oncogene* 9:3037–3041, 1994.
- [66196] 4526.Grana, X.; Claudio, P. P.; De Luca, A.; Sang, N.; Giordano, A.: PISSLRE, a human novel CDC2–related protein

kinase. *Oncogene* 9:2097–2103, 1994.

- [66197] 4527.Ma, W.-J.; Cheng, S.; Campbell, C.; Wright, A.; Furneaux, H.:Cloning and characterization of HuR, a ubiquitously expressed Elav-likeprotein. *J. Biol. Chem.* 271: 8144–8151, 1996.
- [66198] 4528.Ma, W.-J.; Furneaux, H.: Localization of the human HuR gene tochromosome 19p13.2. *Hum. Genet.* 99: 32–33, 1997.
- [66199] 4529.Beaudet, A. L.; O'Brien, W. E.; Bock, H.-G. O.; Freytag, S. O.;Su, T.-S.: The human argininosuccinate synthetase locus and citrullinemia. *Adv.Hum. Genet.* 15: 161–196, 1986.
- [66200] 4530.Beaudet, A. L.; Su, T.-S.; Bock, H.-G.; D'Eustachio, P.; Ruddle,F. H.; O'Brien, W. E.: Use of a cloned cDNA to study human argininosuccinatesynthetase. (Abstract) *Am. J. Hum. Genet.* 33: 36A only, 1981.
- [66201] 4531.Beaudet, A. L.; Su, T.-S.; O'Brien, W. E.; D'Eustachio, P.; Barker,P. E.; Ruddle, F. H.: Dispersion of argininosuccinate-synthetase-likehuman genes to multiple autosomes and the X chromosome. *Cell* 30:287–293, 1982.
- [66202] 4532.Daiger, S. P.; Wildin, R. S.; Su, T.-S.: Sequences on the humanY chromosome homologous to the autosomal gene for argininosuccinatesynthetase. *Nature* 298:

682–684, 1982.

[66203] 4533. Underwood, K. W.; Song, C.; Kriz, R. W.; Chang, X. J.; Knopf, J. L.; Lin, L.-L.: A novel calcium-independent phospholipase A(2), cPLA(2)-gamma, that is prenylated and contains homology to cPLA(2). *J. Biol. Chem.* 273:21926–21932, 1998.

[66204] 4534. Deak, M.; Clifton, A. D.; Lucocq, J. M.; Alessi, D. R.: Mitogen- and stress-activated protein kinase-1 (MSK1) is directly activated by MAPK and SAPK2/p38, and may mediate activation of CREB. *EMBO J.* 17:4426–4441, 1998.

[66205] 4535. Jiang, C.; Yu, L.; Tu, Q.; Zhao, Y.; Zhang, H.; Zhao, S.: Assignment of a member of the ribosomal protein S6 kinase family, RPS6KA5, to human chromosome 14q31–q32.1 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 87: 261–261, 1999.

[66206] 4536. Ackerman, S. L.; Knowles, B. B.: Cloning and mapping of the UNC5C gene to human chromosome 4q21–q23. *Genomics* 52: 205–208, 1998.

[66207] 4537. Ackerman, S. L.; Kozak, L. P.; Przyborski, S. A.; Rund, L. A.; Boyer, B. B.; Knowles, B. B.: The mouse rostral cerebellar malformation gene encodes an UNC-5-like protein. *Nature* 386: 838–842, 1997.

[66208] 4538. Lane, P. W.; Bronson, R. T.; Spencer, C. A.: Rostral

cerebellar malformation, (rcm): a new recessive mutation on chromosome 3 of the mouse. *J. Hered.* 83: 315–318, 1992.

[66209] 4539. Leonardo, E. D.; Hinck, L.; Masu, M.; Keino-Masu, K.; Ackerman, S. L.; Tessier-Lavigne, M.: Vertebrate homologues of *C. elegans* UNC-5 are candidate netrin receptors. *Nature* 386: 833–838, 1997.

[66210] 4540. Przyborski, S. A.; Knowles, B. B.; Ackerman, S. L.: Embryonic phenotype of *Unc5h3* mutant mice suggests chemorepulsion during the formation of the rostral cerebellar boundary. *Development* 125: 41–50, 1998.

[66211] 4541. Marsters, S. A.; Sheridan, J. P.; Pitti, R. M.; Huang, A.; Skubatch, M.; Baldwin, D.; Yuan, J.; Gurney, A.; Goddard, A. D.; Godowski, P.; Ashkenazi, A.: A novel receptor for Apo2L/TRAIL contains a truncated death domain. *Curr. Biol.* 7: 1003–1006, 1997.

[66212] 4542. Pan, G.; O'Rourke, K.; Chinnaiyan, A. M.; Gentz, R.; Ebner, R.; Ni, J.; Dixit, V. M.: The receptor for the cytotoxic ligand TRAIL. *Science* 276: 111–113, 1997.

[66213] 4543. MacFarlane, M.; Ahmad, M.; Srinivasula, S. M.; Fernandes-Alnemri, T.; Cohen, G. M.; Alnemri, E. S.: Identification and molecular cloning of two novel receptors for the cytotoxic ligand TRAIL. *J. Biol. Chem.* 272: 25417–25420,

1997.

- [66214] 4544.Pai, S. I.; Wu, G. S.; Ozoren, N.; Wu, L.; Jen, J.; Sidransky,D.; El-Deiry, W. S.: Rare loss-of-function mutation of a death receptorgene in head and neck cancer. Cancer Res. 58: 3513–3518, 1998.
- [66215] 4545.Pan, G.; Ni, J.; Wei, Y.–F.; Yu, G.; Gentz, R.; Dixit, V. M.:An antagonist decoy receptor and a death domain-containing receptorfor TRAIL. Science 277: 815–818, 1997.
- [66216] 4546.Schneider, P.; Bodmer, J.–L.; Thome, M.; Hofmann, K.; Holler, N.;Tschopp, J.: Characterization of two receptors for TRAIL. FEBS Lett. 416:329–334, 1997.
- [66217] 4547.Screaton, G. R.; Mongkolsapaya, J.; Xu, X.–N.; Cowper, A. E.; McMichael,A. J.; Bell, J. I.: TRICK2, a new alternatively spliced receptorthat transduces the cytotoxic signal from TRAIL. Curr. Biol. 7:693–696, 1997.
- [66218] 4548.Sheridan, J. P.; Marsters, S. A.; Pitti, R. M.; Gurney, A.; Skubatch,M.; Baldwin, D.; Ramakrishnan, L.; Gray, C. L.; Baker, K.; Wood, W.I.; Goddard, A. D.; Godowski, P.; Ashkenazi, A.: Control of TRAIL-inducedapoptosis by a family of signaling and decoy receptors. Science 277:818–821, 1997.
- [66219] 4549.Boeglin, W. E.; Kim, R. B.; Brash, A. R.: A



12R-lipoxygenase in human skin: mechanistic evidence, molecular cloning, and expression. *Proc. Nat. Acad. Sci.* 95: 6744–6749, 1998.

[66220] 4550. Sun, D.; McDonnell, M.; Chen, X.-S.; Lakkis, M. M.; Li, H.; Isaacs, S. N.; Elsea, S. H.; Patel, P. I.; Funk, C. D.: Human 12(R)-lipoxygenase and the mouse ortholog: molecular cloning, expression, and gene chromosomal assignment. *J. Biol. Chem.* 273: 33540–33547, 1998.

[66221] 4551. Itoh, A.; Miyabayashi, T.; Ohno, M.; Sakano, S.: Cloning and expression of three mammalian homologues of *Drosophila* slit suggest possible roles for Slit in the formation and maintenance of the nervous system. *Molec. Brain Res.* 62: 175–186, 1998.

[66222] 4552. Nakayama, M.; Nakajima, D.; Nagase, T.; Nomura, N.; Seki, N.; Ohara, O.: Identification of high molecular weight proteins with multiple EGF-like motifs by motif-trap screening. *Genomics* 51: 27–34, 1998.

[66223] 4553. Duchateau, P. N.; Pullinger, C. R.; Cho, M. H.; Eng, C.; Kane, J. P.: Apolipoprotein L gene family: tissue-specific expression, splicing, promoter regions; discovery of a new gene. *J. Lipid Res.* 42: 620–630, 2001.

[66224] 4554. Duchateau, P. N.; Pullinger, C. R.; Orellana, R. E.; Kunitake, S. T.; Naya-Vigne, J.; O'Connor, P. M.; Malloy, M.

J.; Kane, J. P.: Apolipoprotein L, a new human high density lipoprotein apolipoprotein expressed by the pancreas: identification, cloning, characterization, and plasma distribution of apolipoprotein L. *J. Biol. Chem.* 272:25576–25582, 1997.

- [66225] 4555. Monajemi, H.; Fontijn, R. D.; Pannekoek, H.; Horrevoets, A. J. G.: The apolipoprotein L gene cluster has emerged recently in evolution and is expressed in human vascular tissue. *Genomics* 79: 539–546, 2002.
- [66226] 4556. Page, N. M.; Butlin, D. J.; Lomthaisong, K.; Lowry, P. J.: The human apolipoprotein L gene cluster: identification, classification, and sites of distribution. *Genomics* 74: 71–78, 2001.
- [66227] 4557. Lin, C.-Y.; Huang, P.-H.; Liao, W.-L.; Cheng, H.-J.; Huang, C.-F.; Kuo, J.-C.; Patton, W. A.; Massenburg, D.; Moss, J.; Lee, F.-J. S.: ARL4, an ARF-like protein that is developmentally regulated and localized to nuclei and nucleoli. *J. Biol. Chem.* 275: 37815–37823, 2000.
- [66228] 4558. Schurmann, A.; Breiner, M.; Becker, W.; Huppertz, C.; Kainulainen, H.; Kentrup, H.; Joost, H.-G.: Cloning of two novel ADP-ribosylation factor-like proteins and characterization of their differential expression in 3T3-L1 cells. *J. Biol. Chem.* 269: 15683–15688, 1994.

- [66229] 4559.Schurmann, A.; Koling, S.; Jacobs, S.; Saftig, P.; Kraub, S.; Wennemuth,G.; Kluge, R.; Joost, H.–G.: Reduced sperm count and normal fertility in male mice with targeted disruption of the ADP–ribosylation factor–like4 (Arl4) gene. *Molec. Cell. Biol.* 22: 2761–2768, 2002.
- [66230] 4560.Lehnert, K.; Ni, J.; Leung, E.; Gough, S. M.; Weaver, A.; Yao,W.–P.; Liu, D.; Wang, S.–X.; Morris, C. M.; Krisansen, G. W.: Cloning,sequence analysis, and chromosomal localization of the novel humanintegrin alpha–11 subunit (ITGA11). *Genomics* 60: 179–187, 1999.
- [66231] 4561.Velling, T.; Kusche–Gullberg, M.; Sejersen, T.; Gullberg, D.:cDNA cloning and chromosomal localization of human alpha–11 integrin:a collagen–binding, I domain–containing, beta–1–associated integrinalpha–chain present in muscle tissues. *J. Biol. Chem.* 274: 25735–25742,1999.
- [66232] 4562.Mital, R.; Kobayashi, R.; Hernandez, N.: RNA polymerase III transcriptionfrom the human U6 and adenovirus type 2 VAI promoters has differentrequirements for human BRF, a subunit of human TFIIIB. *Molec. Cell.Biol.* 16: 7031–7042, 1996.
- [66233] 4563.Wang, Z.; Roeder, R. G.: Structure and function of a human transcriptionfactor TFIIIB subunit that is evolution–

arily conserved and contains both TFIIB- and high-mobility-group protein 2-related domains. *Proc. Nat. Acad. Sci.* 92: 7026–7030, 1995.

- [66234] 4564. Comai, L.; Zomerdijk, J. C. B. M.; Beckmann, H.; Zhou, S.; Admon, A.; Tjian, R.: Reconstitution of transcription factor SL1: exclusive binding of TBP by SL1 or TFIID subunits. *Science* 266: 1966–1972, 1994.
- [66235] 4565. Di Pietro, C.; Rapisarda, A.; Amico, V.; Bonaiuto, C.; Viola, A.; Scalia, M.; Motta, S.; Amato, A.; Engel, H.; Messina, A.; Sichel, G.; Grzeschik, K.-H.; Purrello, M.: Genomic localization of the human genes TAF1A, TAF1B and TAF1C, encoding TAFI48, TAFI63 and TAFI110 subunits of class I general transcription initiation factor SL1. *Cytogenet. Cell Genet.* 89: 133–136, 2000.
- [66236] 4566. Nakamura, T.; Takeuchi, K.; Muraoka, S.; Takezoe, H.; Takahashi, N.; Mori, N.: A neurally enriched coronin-like protein, ClipnC, is a novel candidate for an actin cytoskeleton-cortical membrane-linking protein. *J. Biol. Chem.* 274: 13322–13327, 1999.
- [66237] 4567. Kim, K. I.; Baek, S. H.; Jeon, Y.-J.; Nishimori, S.; Suzuki, T.; Uchida, S.; Shimbara, N.; Saitoh, H.; Tanaka, K.; Chung, C. H.: A new SUMO-1-specific protease, SUSP1, that is highly expressed in reproductive organs. *J. Biol.*

Chem. 275: 14102–14106, 2000.

[66238] 4568.Ali, A.; Hoeflich, K. P.; Woodgett, J. R. Glycogen synthase kinase-3: properties, functions, and regulation.

Chem. Rev. 101: 2527–2540, 2001.

[66239] 4569.Dajani, R.; Fraser, E.; Roe, S. M.; Young, N.; Good, V.; Dale, T. C.; Pearl, L. H.: Crystal structure of glycogen synthase kinase3-beta: structural basis for phosphate-primed substrate specificity and autoinhibition. Cell 105: 721–732, 2001.

[66240] 4570.Frame, S.; Cohen, P.; Biondi, R. M.: A common phosphate binding site explains the unique substrate specificity of GSK3 and its inactivation by phosphorylation. Molec. Cell 7: 1321–1327, 2001.

[66241] 4571.Hoeflich, K. P.; Luo, J.; Rubie, E. A.; Tsao, M.-S.; Jin, O.; Woodgett, J. R.: Requirement for glycogen synthase kinase-3-beta in cell survival and NF-kappa-B activation. Nature 406: 86–90, 2000.

[66242] 4572.Lau, K. F.; Miller, C. C.; Anderton, B. H.; Shaw, P. C.: Molecular cloning and characterization of the human glycogen synthase kinase-3-beta promoter. Genomics 60: 121–128, 1999.

[66243] 4573.Lau, K. F.; Miller, C. C.; Anderton, B. H.; Shaw, P. C.: Expression analysis of glycogen synthase kinase-3 in hu-

man tissues. J. PeptideRes. 54: 85–91, 1999.

[66244] 4574.Lucas, J. J.; Hernandez, F.; Gomez–Ramos, P.; Moran, M. A.; Hen,R.; Avila, J.: Decreased nuclear beta–catenin, tau hyperphosphorylationand neurodegeneration in GSK–3–beta conditional transgenic mice. EMBOJ. 20: 27–39, 2001.

[66245] 4575.Plyte, S. E.; Hughes, K.; Nikolakaki, E.; Pulverer, B. J.; Woodgett,J. R.: Glycogen synthase kinase–3: functions in oncogenesis and development. Biochim.Biophys. Acta 1114: 147–162, 1992.

[66246] 4576.Shaw, P. C.; Davies, A. F.; Lau, K. F.; Garcia–Barcelo, M.; Waye,M. M.; Lovestone, S.; Miller, C. C.; Anderton, B. H.: Isolation andchromosomal mapping of human glyco–gen synthase kinase–3 alpha and3 beta encoding genes. Genome 41: 720–727, 1998.

[66247] 4577.Stambolic, V.; Woodgett, J. R.: Mitogen inactivation of glycogensynthase kinase–3 beta in intact cells via ser–ine 9 phosphorylation. Biochem.J. 303: 701–704, 1994.

[66248] 4578.Wang, Q. M.; Fiol, C. J.; DePaoli–Roach, A. A.; Roach, P. J.:Glycogen synthase kinase–3 beta is a dual specificity kinase differentiallyregulated by tyrosine and serine/ threonine phosphorylation. J. Biol.Chem. 269: 14566–14574, 1994.

- [66249] 4579.Saitoh, T.; Moriwaki, J.; Koike, J.; Takagi, A.; Miwa, T.; Shiokawa,K.; Katoh, M.: Molecular cloning and characterization of FRAT2, encoding a positive regulator of the WNT signaling pathway. *Biochem. Biophys.Res. Commun.* 281: 815–820, 2001.
- [66250] 4580.Abbaszade, I.; Liu, R.–Q.; Yang, F.; Rosenfeld, S. A.; Ross, O.H.; Link, J. R.; Ellis, D. M.; Tortorella, M. D.; Pratta, M. A.; Hollis,J. M.; Wynn, R.; Duke, J. L.; and 15 others: Cloning and characterization of ADAMTS11, an aggrecanase from the ADAMTS family. *J. Biol. Chem.* 274:23443–23450, 1999.
- [66251] 4581.Duan, H.; Wang, Y.; Aviram, M.; Swaroop, M.; Loo, J. A.; Bian,J.; Tian, Y.; Mueller, T.; Bisgaier, C. L.; Sun, Y.: SAG, a novel zinc RING finger protein that protects cells from apoptosis induced by redox agents. *Molec. Cell. Biol.* 19: 3145–3155, 1999.
- [66252] 4582.Murakami, A.; Yajima, T.; Sakuma, H.; McLaren, M. J.; Inana, G.: X-arrestin: a new retinal arrestin mapping to the X chromosome. *FEBS Lett.* 334: 203–209, 1993.
- [66253] 4583.Sakuma, H.; Murakami, A.; Fujimaki, T.; Inana, G.: Isolation and characterization of the human X-arrestin gene. *Gene* 224: 87–95, 1998.

- [66254] 4584.Ellison, J.; Passage, M.; Yu, L.-C.; Yen, P.; Mohandas, T. K.;Shapiro, L.: Directed isolation of human genes that escape X inactivation. *Somat.Cell Molec. Genet.* 18: 259–268, 1992.
- [66255] 4585.Ellison, J. W.; Ramos, C.; Yen, P. H.; Shapiro, L. J.: Structureand expression of the human pseudoautosomal gene XE7. *Hum. Molec.Genet.* 1: 691–696, 1992.
- [66256] 4586.Amar, L. C.; Dandolo, L.; Hanauer, A.; Cook, A. R.; Arnaud, D.;Mandel, J. L.; Avner, P.: Conservation and reorganization of locion the mammalian X chromosome: a molecular framework for the identificationof homologous subchromosomal regions in man and mouse. *Genomics* 2:220–230, 1988.
- [66257] 4587.Chin, L.-S.; Li, L.; Ferreira, A.; Kosik, K. S.; Greengard, P.: Impairment of axonal development and of synaptogenesis in hippocampalneurons of synapsin I-deficient mice. *Proc. Nat. Acad. Sci.* 92:9230–9234, 1995.
- [66258] 4588.Klagges, B. R. E.; Heimbeck, G.; Godenschwege, T. A.; Hofbauer,A.; Pflugfelder, G. O.; Reifegerste, R.; Reisch, D.; Schaupp, M.;Buchner, S.; Buchner, E.: Invertebrate synapsins: a single gene codesfor several isoforms in *Drosophila*. *J. Neurosci.* 16: 3154–3165,1996.
- [66259] 4589.Li, L.; Chin, L.-S.; Shupliakov, O.; Brodin, L.; Sihra, T.



S.; Hvalby, O.; Jensen, V.; Zheng, D.; McNamara, J. O.; Greengard, P.; Andersen, P.: Impairment of synaptic vesicle clustering and of synaptic transmission, and increased seizure propensity, in synapsin I-deficient mice. *Proc. Nat. Acad. Sci.* 92: 9235–9239, 1995.

[66260] 4590. Sudhof, T. C.: The structure of the human synapsin I gene and protein. *J. Biol. Chem.* 265: 7849–7852, 1990.

[66261] 4591. Banting, G. S.; Pym, B.; Goodfellow, P. N.: Biochemical analysis of an antigen produced by both human sex chromosomes. *EMBO J.* 4: 1967–1972, 1985.

[66262] 4592. Hagiwara, T.; Tanaka, K.; Takai, S.; Maeno-Hikichi, Y.; Mukainaka, Y.; Wada, K.: Genomic organization, promoter analysis, and chromosomal localization of the gene for the mouse glial high-affinity glutamate transporter Slc1a3. *Genomics* 33: 508–515, 1996.

[66263] 4593. Harada, T.; Harada, C.; Watanabe, M.; Inoue, Y.; Sakagawa, T.; Nakayama, N.; Sasaki, S.; Okuyama, S.; Watase, K.; Wada, K.; Tanaka, K.: Functions of the two glutamate transporters GLAST and GLT-1 in the retina. *Proc. Nat. Acad. Sci.* 95: 4663–4666, 1998.

[66264] 4594. Keppen, L. D.; Gollin, S. M.; Edwards, D.; Sawyer, J.; Wilson, W.; Overhauser, J.: Clinical phenotype and molecular analysis of a three-generation family with an interstitial

deletion of the shortarm of chromosome 5. *Am. J. Med. Genet.* 44: 356–360, 1992.

- [66265] 4595. Kirschner, M. A.; Arriza, J. L.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Magenis, E.; Amara, S. G.: The mouse and human excitatory amino acid transporter gene (EAAT1) maps to mouse chromosome 15 and a region of syntenic homology on human chromosome 5. *Genomics* 22:631–633, 1994.
- [66266] 4596. Shashidharan, P.; Huntley, G. W.; Meyer, T.; Morrison, J. H.; Plaitakis, A.: Neuron-specific human glutamate transporter: molecular cloning, characterization and expression in human brain. *Brain Res.* 662:245–250, 1994.
- [66267] 4597. Stoffel, W.; Sasse, J.; Duker, M.; Muller, R.; Hofmann, K.; Fink, T.; Lichter, P.: Human high affinity, Na(+)-dependent L-glutamate/L-aspartate transporter GLAST-1 (EAAT-1): gene structure and localization to chromosome 5p11–p12. *FEBS Lett.* 386: 189–193, 1996.
- [66268] 4598. Takai, S.; Yamada, K.; Kawakami, H.; Tanaka, K.; Nakamura, S.: Localization of the gene (SLC1A3) encoding human glutamate transporter (GluT-1) to 5p13 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 69: 209–210, 1995.
- [66269] 4599. Agulnik, A. I.; Mitchell, M. J.; Lerner, J. L.; Woods, D.

R.; Bishop, C. E.: A mouse Y chromosome gene encoded by a region essential for spermatogenesis and expression of male-specific minor histocompatibility antigens. *Hum. Molec. Genet.* 3: 873–878, 1994.

[66270] 4600. Agulnik, A. I.; Mitchell, M. J.; Mattei, M.-G.; Borsani, G.; Avner, P. A.; Lerner, J. L.; Bishop, C. E.: A novel X gene with a widely transcribed Y-linked homologue escapes X-inactivation in mouse and human. *Hum. Molec. Genet.* 3: 879–884, 1994.

[66271] 4601. Lingenfelter, P. A.; Adler, D. A.; Poslinski, D.; Thomas, S.; Elliott, R. W.; Chapman, V. M.; Disteché, C. M.: Escape from X inactivation of *Smcx* is preceded by silencing during mouse development. *Nature Genet.* 18: 212–213, 1998.

[66272] 4602. Wu, J.; Ellison, J.; Salido, E.; Yen, P.; Mohandas, T.; Shapiro, L. J.: Isolation and characterization of XE169, a novel human gene that escapes X-inactivation. *Hum. Molec. Genet.* 3: 153–160, 1994.

[66273] 4603. Wu, J.; Salido, E. C.; Yen, P. H.; Mohandas, T. K.; Heng, H. H. Q.; Tsui, L.-C.; Park, J.; Chapman, V. M.; Shapiro, L. J.: The murine *Xe169* gene escapes X-inactivation like its human homologue. *Nature Genet.* 7: 491–496, 1994.

- [66274] 4604. Evans, H. J.; Buckton, K. E.; Spowart, G.; Carothers, A. D.: Heteromorphic X chromosomes in 46,XX males: evidence for the involvement of X-Y interchange. *Hum. Genet.* 49: 11–31, 1979.
- [66275] 4605. Hu, X.; Burghes, A. H. M.; Ray, P. N.; Thompson, M. W.; Murphy, E. G.; Worton, R. G.: Partial gene duplication in Duchenne and Becker muscular dystrophies. *J. Med. Genet.* 25: 369–376, 1988.
- [66276] 4606. Berger, W.; van de Pol, D.; Warburg, M.; Gal, A.; Bleeker-Wagemakers, L.; de Silva, H.; Meindl, A.; Meitinger, T.; Cremers, F.; Ropers, H.-H.: Mutations in the candidate gene for Norrie disease. *Hum. Molec. Genet.* 1: 461–465, 1992.
- [66277] 4607. Bleeker-Wagemakers, L. M.; Friedrich, U.; Gal, A.; Wienker, T. F.; Warburg, M.; Ropers, H.-H.: Close linkage between Norrie disease, a cloned DNA sequence from the proximal short arm, and the centromere of the X chromosome. *Hum. Genet.* 71: 211–214, 1985.
- [66278] 4608. Chen, Z.-Y.; Battinelli, E. M.; Hendriks, R. W.; Powell, J. F.; Middleton-Price, H.; Sims, K. B.; Breakefield, X. O.; Craig, I. W.: Norrie disease gene: characterization of deletions and possible function. *Genomics* 16: 533–535, 1993.
- [66279] 4609. Chen, Z.-Y.; Battinelli, E. M.; Woodruff, G.; Young, I.;

Breakefield,X. O.; Craig, I. W.: Characterization of a mutation within the NDPgene in a family with a manifesting female carrier. Hum. Molec. Genet. 2:1727–1729, 1993.

[66280] 4610.Chen, Z.–Y.; Hendriks, R. W.; Jobling, M. A.; Powell, J. F.; Breakefield,X. O.; Sims, K. B.; Craig, I. W.: Isolation and characterization of a candidate gene for Norrie disease. Nature Genet. 1: 204–208,1992.

[66281] 4611.Chen, Z.–Y.; Sims, K. B.; Coleman, M.; Donnai, D.; Monaco, A.;Breakefield, X. O.; Davies, K. E.; Craig, I. W.: Characterization of a YAC containing part or all of the Norrie disease locus. Hum.Molec. Genet. 1: 161–164, 1992.

[66282] 4612.Clarke, E.: 'Pseudo–glioma' in both eyes. Trans. Ophthal. Soc.U.K. 18: 136–138, 1898.

[66283] 4613.Dahlberg–Parrow, R.: Congenital sex–linked pseudoglioma and gravenmental deficiency. Acta Ophthal. 34: 250–254, 1956.

[66284] 4614.Diergaarde, P. J.; Wieringa, B.; Bleeker–Wagemakers, E. M.; Sims,K. B.; Breakefield, X. O.; Ropers, H.–H.: Physical fine–mapping of a deletion spanning the Norrie gene. Hum. Genet. 84: 22–26, 1989.

[66285] 4615.Donnai, D.; Mountford, R. C.; Read, A. P.: Norrie disease resulting from a gene deletion: clinical features and DNA studies. J. Med.Genet. 25: 73–78, 1988.

- [66286] 4616.Duke–Elder, J. R.: Pseudoglioma in children: aspects of clinical and pathological diagnosis. *Sth. Med. J.* 51: 754–759, 1958.
- [66287] 4617.Forssman, H.: Mental deficiency and pseudoglioma, a syndrome inherited as an X–linked recessive. *Am. J. Ment. Defic.* 64: 984–987, 1960.
- [66288] 4618.Fuchs, S.; Kellner, U.; Wedemann, H.; Gal, A.: Missense mutation (Arg121Trp) in the Norrie disease gene associated with X–linked exudative vitreoretinopathy. *Hum. Mutat.* 6: 257–259, 1995.
- [66289] 4619.Fuchs, S.; Xu, S. Y.; Caballero, M.; Salcedo, M.; La O, A.; Wedemann, H.; Gal, A.: A missense point mutation (Leu13Arg) of the Norrie disease gene in a large Cuban kindred with Norrie disease. *Hum. Molec. Genet.* 3: 655–656, 1994.
- [66290] 4620.Collins, F. A.; Murphy, D. L.; Reiss, A. L.; Sims, K. B.; Lewis, J. G.; Freund, L.; Karoum, F.; Zhu, D.; Maumenee, I. H.; Antonarakis, S. E.: Clinical, biochemical, and neuropsychiatric evaluation of a patient with a contiguous gene syndrome due to a microdeletion Xp11.3 including the Norrie disease locus and monoamine oxidase (MAOA and–MAOB) genes. *Am. J. Med. Genet.* 42: 127–134, 1992.
- [66291] 4621.Fullwood, P.; Jones, J.; Bunday, S.; Dudgeon, J.;

Fielder, A.R.; Kilpatrick, M. W.: X linked exudative vitreo-retinopathy: clinical features and genetic linkage analysis. Brit. J. Ophthal. 77: 168–170, 1993.

[66292] 4622. Gal, A.; Bleeker-Wagemakers, L.; Wienker, T. F.; Warburg, M.; Ropers, H.-H.: Localization of the gene for Norrie disease by linkage to the DXS7 locus. (Abstract) Cytogenet. Cell Genet. 40: 633, 1985.

[66293] 4623. Gal, A.; Stolzenberger, C.; Wienker, T.; Wieacker, P.; Ropers, H.-H.; Friedrich, U.; Bleeker-Wagemakers, L.; Pearson, P.; Warburg, M.: Norrie's disease: close linkage with genetic markers from the proximal short arm of the X chromosome. Clin. Genet. 27: 282–283, 1985.

[66294] 4624. Gal, A.; Uhlhaas, S.; Glaser, D.; Grimm, T.: Prenatal exclusion of Norrie disease with flanking DNA markers. Am. J. Med. Genet. 31: 449–453, 1988.

[66295] 4625. Harendra de Silva, D. G.; de Silva, D. B. K.: Norrie's disease in an Asian family. Brit. J. Ophthal. 72: 62–64, 1988.

[66296] 4626. Bermingham, N. A.; Martin, J. E.; Fisher, E. M. C.: The mouse lysosomal membrane protein 1 gene as a candidate for the motor neuron degeneration (mnd) locus. Genomics 32: 266–271, 1996.

[66297] 4627. Howe, C. L.; Granger, B. L.; Hull, M.; Green, S. A.;

Gabel, C.A.; Helenius, A.; Mellman, I.: Derived protein sequence, oligosaccharides, and membrane insertion of the 120-kDa lysosomal membrane glycoprotein (lgp120): identification of a highly conserved family of lysosomal membrane glycoproteins. *Proc. Nat. Acad. Sci.* 85: 7577–7581, 1988.

[66298] 4628. Mattei, M.-G.; Matterson, J.; Chen, J. W.; Williams, M. A.; Fukuda, M.: Two human lysosomal membrane glycoproteins, h-lamp-1 and h-lamp-2, are encoded by genes localized to chromosome 13q34 and chromosome Xq24–25, respectively. *J. Biol. Chem.* 265: 7548–7551, 1990.

[66299] 4629. Schleutker, J.; Haataja, L.; Renlund, M.; Puhakka, L.; Viitala, J.; Peltonen, L.; Aula, P.: Confirmation of the chromosomal localization of human lamp genes and their exclusion as candidate genes for Salla disease. *Hum. Genet.* 88: 95–97, 1991.

[66300] 4630. Viitala, J.; Carlsson, S. R.; Siebert, P. D.; Fukuda, M.: Molecular cloning of cDNAs encoding lamp A, a human lysosomal membrane glycoprotein with apparent M(r) about 120,000. *Proc. Nat. Acad. Sci.* 85: 3743–3747, 1988.

[66301] 4631. Andreae, S.; Piras, F.; Burdin, N.; Triebel, F.: Matura-



tion and activation of dendritic cells induced by lymphocyte activation gene-3 (CD223). *J. Immun.* 168: 3874–3880, 2002.

[66302] 4632. Triebel, F.; Jitsukawa, S.; Baixeras, E.; Roman-Roman, S.; Genevee, C.; Viegas-Pequignot, E.; Hercend, T.: LAG-3, a novel lymphocyte activation gene closely related to CD4. *J. Exp. Med.* 171: 1393–1405, 1990.

[66303] 4633. LeClair, K. P.; Rabin, M.; Nesbitt, M. N.; Pravtcheva, D.; Ruddle, F. H.; Palfree, R. G. E.; Bothwell, A.: Murine Ly-6 multigene family is located on chromosome 15. *Proc. Nat. Acad. Sci.* 84: 1638–1642, 1987.

[66304] 4634. Ravetch, J. V.; Lanier, L. L.: Immune inhibitory receptors. *Science* 290:84–89, 2000.

[66305] 4635. Fischer, A.; Durandy, A.; Sterkers, G.; Griscelli, C.: Role of the LFA-1 molecule in cellular interactions required for antibody production in humans. *J. Immun.* 136: 3198–3203, 1986.

[66306] 4636. Marlin, S. D.; Morton, C. C.; Anderson, D. C.; Springer, T. A.: LFA-1 immunodeficiency disease: definition of the genetic defect and chromosomal mapping of alpha and beta subunits of the lymphocyte function-associated antigen 1 (LFA-1) by complementation in hybrid cells. *J. Exp. Med.* 164: 855–867, 1986.

- [66307] 4637.Futscher, B. W.; Oshiro, M. M.; Wozniak, R. J.; Holtan, N.; Hanigan, C. L.; Duan, H.; Domann, F. E.: Role for DNA methylation in the control of cell type-specific maspin expression. *Nature Genet.* 31: 175–179, 2002.
- [66308] 4638.Ngamkitidechakul, C.; Burke, J. M.; O'Brien, W. J.; Twining, S.S.: Maspin: synthesis by human cornea and regulation of in vitro stromal cell adhesion to extracellular matrix. *Invest. Ophthalmol. Vis.Sci.* 42: 3135–3141, 2001.
- [66309] 4639.Zou, Z.; Anisowicz, A.; Hendrix, M. J. C.; Thor, A.; Neveu, M.; Sheng, S.; Rafidi, K.; Seftor, E.; Sager, R.: Maspin, a serpin with tumor-suppressing activity in human mammary epithelial cells. *Science* 263:526–529, 1994.
- [66310] 4640.Blackwood, E.; Eisenman, R. N.: Max: a helix-loop-helix zipper protein that forms a sequence-specific DNA-binding complex with Myc. *Science* 251:1211–1217, 1991.
- [66311] 4641.Eisenman, R. N.: Personal Communication. Seattle, Wash. 7/27/1994.
- [66312] 4642.Gilladoga, A. D.; Edelhoff, S.; Blackwood, E. M.; Eisenman, R.N.; Distech, C. M.: Mapping of MAX to human chromosome 14 and mouse chromosome 12 by in situ hybridization. *Oncogene* 7: 1249–1251, 1992.
- [66313] 4643.Loeffen, J. L. C. M.; Triepels, R. H.; van den Heuvel,

L. P.; Schuelke, M.; Buskens, C. A. F.; Smeets, R. J. P.; Trij-  
bels, J. M. F.; Smeitink, J. A. M.: cDNA of eight nuclear en-  
coded subunits of NADH:ubiquinone oxidoreductase: hu-  
man complex I cDNA characterization completed.

Biochem. Biophys. Res. Commun. 253: 415–422, 1998.

[66314] 4644. Bulavin, D. V.; Higashimoto, Y.; Popoff, I. J.; Gaarde,  
W. A.; Basrur, V.; Potapova, O.; Appella, E.; Fornace, A. J.,  
Jr.: Initiation of a G2/M checkpoint after ultraviolet radia-  
tion requires p38 kinase. Nature 411: 102–107, 2001.

[66315] 4645. Gould, K. L.; Moreno, S.; Tonks, N. K.; Nurse, P.:  
Complementation of the mitotic activator, p80(cdc25), by a  
human protein-tyrosine phosphatase. Science 250:  
1573–1576, 1990.

[66316] 4646. Moreno, S.; Nurse, P.: Clues to action of cdc25 pro-  
tein. (Letter) Nature 351: 194 only, 1991.

[66317] 4647. Raj, K.; Ogston, P.; Beard, P.: Virus-mediated killing  
of cells that lack p53 activity. Nature 412: 914–917, 2001.  
Note: Addendum: Nature 416: 202 only, 2002.

[66318] 4648. Sadhu, K.; Reed, S. I.; Richardson, H.; Russell, P.:  
Human homolog of fission yeast cdc25 mitotic inducer is  
predominantly expressed in G(2). Proc. Nat. Acad. Sci. 87:  
5139–5143, 1990.

[66319] 4649. Strausfeld, U.; Labbe, J. C.; Fesquet, D.; Cavadore, J.

C.; Picard,A.; Sadhu, K.; Russell, P.; Doree, M.: Dephos-  
phorylation and activationof a p34(cdc2)/cyclin B complex  
in vitro by human CDC25 protein. Nature 351:242–245,  
1991.

[66320] 4650.Taviaux, S. A.; Demaille, J. G.: Localization of human  
cell cycleregulatory genes CDC25C to 5q31 and WEE1 to  
11p15.3–11p.15.1 by fluorescencein situ hybridization.  
Genomics 15: 194–196, 1993.

[66321] 4651.Willman, C. L.; Sever, C. E.; Pallavicini, M. G.; Harada,  
H.; Tanaka,N.; Slovak, M. L.; Yamamoto, H.; Harada, K.;  
Meeker, T. C.; List,A. F.; Taniguchi, T.: Deletion of IRF–1,  
mapping to chromosome 5q31.1,in human leukemia and  
preleukemic myelodysplasia. Science 259: 968–971,1993.

[66322] 4652.Dear, T. N.; Campbell, K.; Rabbitts, T. H.: Molecular  
cloningof putative odorant–binding and odorant–  
metabolizing proteins. Biochemistry 30:10376–10382,  
1991.

[66323] 4653.Lacazette, E.; Gachon, A. M.; Pitiot, G.: A novel hu-  
man odorant–bindingprotein gene family resulting from  
genomic duplicons at 9q34: differentialexpression in the  
oral and genital spheres. Hum. Molec. Genet. 9:289–301,  
2000.

[66324] 4654.Pevsner, J.; Reed, R. R.; Feinstein, P. G.; Snyder, S.

H.: Molecular cloning of odorant-binding protein: member of a ligand carrier family. *Science* 241:336–339, 1988.

[66325] 4655. Raming, K.; Krieger, J.; Strotmann, J.; Boekhoff, I.; Kubick, S.; Baumstark, C.; Breer, H.: Cloning and expression of odorant receptors. *Nature* 361:353–356, 1993.

[66326] 4656. Unterman, R. D.; Lynch, K. R.; Nakhasi, H. L.; Dolan, K. P.; Hamilton, J. W.; Cohn, D. V.; Feigelson, P.: Cloning and sequence of several  $\alpha$ -2-microglobulin cDNAs. *Proc. Nat. Acad. Sci.* 78: 3478–3482, 1981.

[66327] 4657. Miles, S. A.; Martinez-Maza, O.; Rezai, A.; Magpantay, L.; Kishimoto, T.; Nakamura, S.; Radka, S. F.; Linsley, P. S.: Oncostatin M as a potent mitogen for AIDS-Kaposi's sarcoma-derived cells. *Science* 255:1432–1434, 1992.

[66328] 4658. Modur, V.; Feldhaus, M. J.; Weyrich, A. S.; Jicha, D. L.; Prescott, S. M.; Zimmerman, G. A.; McIntyre, T. M.: Oncostatin M is a proinflammatory mediator: in vivo effects correlate with endothelial cell expression of inflammatory cytokines and adhesion molecules. *J. Clin. Invest.* 100:158–168, 1997.

[66329] 4659. Nair, B. C.; DeVico, A. L.; Nakamura, S.; Copeland, T. D.; Chen, Y.; Patel, A.; O'Neil, T.; Oroszlan, S.; Gallo, R. C.; Sarngadharan, M. G.: Identification of a major growth factor for AIDS-Kaposi's sarcoma cells as oncostatin M. *Sci-*

ence 255: 1430–1432, 1992.

- [66330] 4660. Rose, T. M.; Bruce, A. G.: Oncostatin M is a member of a cytokine family that includes leukemia-inhibitory factor, granulocyte colony-stimulating factor, and interleukin 6. *Proc. Nat. Acad. Sci.* 88: 8641–8645, 1991.
- [66331] 4661. Rose, T. M.; Lagrou, M. J.; Fransson, I.; Werelius, B.; Delattre, O.; Thomas, G.; de Jong, P. J.; Todaro, G. J.; Dumaniski, J. P.: The genes for oncostatin M (OSM) and leukemia inhibitory factor (LIF) are tightly linked on human chromosome 22. *Genomics* 17: 136–140, 1993.
- [66332] 4662. Zarling, J. M.; Shoyab, M.; Marquardt, H.; Hanson, M. B.; Lioubin, M. N.; Todaro, G. J.: Oncostatin M: a growth regulator produced by differentiated histiocytic lymphoma cells. *Proc. Nat. Acad. Sci.* 83: 9739–9743, 1986.
- [66333] 4663. Harder, K. W.; Parsons, L. M.; Armes, J.; Evans, N.; Kountouri, N.; Clark, R.; Quillici, C.; Grail, D.; Hodgson, G. S.; Dunn, A. R.; Hibbs, M. L.: Gain- and loss-of-function Lyn mutant mice define a critical inhibitory role for Lyn in the myeloid lineage. *Immunity* 15: 603–615, 2001.
- [66334] 4664. Fujimoto, M.; Kantaputra, P. N.; Ikegawa, S.; Fukushima, Y.; Sonta, S.; Matsuo, M.; Ishida, T.; Matsumoto, T.; Kondo, S.; Tomita, H.; Deng, H.-X.; D'urso, M.; Rinaldi, M. M.; Ventruto, V.; Takagi, T.; Nakamura, Y.; Ni-

ikawa, N.: The gene for mesomelic dysplasia Kantaputratype is mapped to chromosome 2q24–q32. *J. Hum. Genet.* 43: 32–36, 1998.

[66335] 4665. Ventruto, V.; Pisciotto, R.; Renda, S.; Fosta, B.; Rinaldi, M.M.; Stabile, M.; Cavaliere, M. L.; Esposito, M.: Multiple skeletal familial abnormalities associated with balanced reciprocal translocation 2;8(q32;p13). *Am. J. Med. Genet.* 16: 589–594, 1983.

[66336] 4666. Joyner, A. L.; Herrup, K.; Auerbach, B. A.; Davis, C. A.; Rossant, J.: Subtle cerebellar phenotype in mice homozygous for a targeted deletion of the *En-2* homeobox. *Science* 251: 1239–1243, 1991.

[66337] 4667. Joyner, A. L.; Skarnes, W. C.; Rossant, J.: Production of a mutation in mouse *En-2* gene by homologous recombination in embryonic stem cells. *Nature* 338: 153–156, 1989.

[66338] 4668. Poole, S. J.; Law, M. L.; Kao, F.-T.; Lau, Y.-F.: Isolation and chromosomal localization of the human *En-2* gene. *Genomics* 4: 225–231, 1989.

[66339] 4669. Kiss, C.; Li, J.; Szeles, A.; Gizatullin, R. Z.; Kashuba, V. I.; Lushnikova, T.; Protopopov, A. I.; Kelve, M.; Kiss, H.; Kholodnyuk, I. D.; Imreh, S.; Klein, G.; Zabarovsky, E. R.: Assignment of the *ARHA* and *GPX1* genes to human chro-

mosome bands 3p21.3 by in situ hybridization and with somatic cell hybrids. *Cytogenet. Cell Genet.* 79: 228–230, 1997.

[66340] 4670. Ma, J. J.; Nishimura, M.; Mine, H.; Kuroki, S.; Nukina, M.; Ohta, M.; Saji, H.; Obayashi, H.; Kawakami, H.; Saida, T.; Uchiyama, T.: Genetic contribution of the tumor necrosis factor region in Guillain–Barre syndrome. *Ann. Neurol.* 44: 815–818, 1998.

[66341] 4671. Giuli, G.; Roechel, N.; Scholl, U.; Mattei, M.–G.; Guellaen, G.: Colocalization of the genes coding for the alpha-3 and beta-3 subunits of soluble guanylyl cyclase to human chromosome 4 at q31.3–q33. *Hum. Genet.* 91: 257–260, 1993.

[66342] 4672. Cameron, H. S.; Szczepaniak, D.; Weston, B. W.: Expression of human chromosome 19p alpha-(1,3)-fucosyltransferase genes in normal tissues: alternative splicing, polyadenylation, and isoforms. *J. Biol. Chem.* 270: 20112–20122, 1995.

[66343] 4673. Koszdin, K. L.; Bowen, B. R.: The cloning and expression of a human alpha-1,3 fucosyltransferase capable of forming the E-selectin ligand. *Biochem. Biophys. Res. Commun.* 187: 152–157, 1992.

[66344] 4674. Mollicone, R.; Reguigne, I.; Fletcher, A.; Aziz, A.;



Rustam, M.; Weston, B. W.; Kelly, R. J.; Lowe, J. B.; Oriol, R.: Molecular basis for plasma alpha(1,3)-fucosyltransferase gene deficiency (FUT6). *J. Biol. Chem.* 269: 12662–12671, 1994.

[66345] 4675. Pang, H.; Koda, Y.; Soejima, M.; Schlaphoff, T.; Du Toit, E. D.; Kimura, H.: Allelic diversity of the human plasma alpha(1,3)fucosyltransferase gene (FUT6). *Ann. Hum. Genet.* 63: 277–284, 1999.

[66346] 4676. Schnyder–Candrian, S.; Borsig, L.; Moser, R.; Berger, E. G.: Localization of alpha–1,3–fucosyltransferase VI in Weibel–Palade bodies of human endothelial cells. *Proc. Nat. Acad. Sci.* 97: 8369–8374, 2000.

[66347] 4677. Doonan, S.; Barra, D.; Bossa, F.: Structural and genetic relationships between cytosolic and mitochondrial isoenzymes. *Int. J. Biochem.* 16: 1193–1199, 1984.

[66348] 4678. Allen, S. J.; O'Donnell, A.; Alexander, N. D. E.; Alpers, M. P.; Peto, T. E. A.; Clegg, J. B.; Weatherall, D. J.: Alpha(+)-thalassaemia protects children against disease caused by other infections as well as malaria. *Proc. Nat. Acad. Sci.* 94: 14736–14741, 1997.

[66349] 4679. Keats, B. J. B.; Morton, N. E.; Rao, D. C.: Possible linkages (lod score over 1.5) and a tentative map of the Jk–Km linkage group. *Cytogenet. Cell Genet.* 22: 304–308,

1978.

- [66350] 4680.Keats, B. J. B.; Morton, N. E.; Rao, D. C.: Likely linkage: In Vwith Jk. Hum. Genet. 39: 157–159, 1977.
- [66351] 4681.Davenne, M.; Maconochie, M. K.; Neun, R.; Pattyn, A.; Chambon,P.; Krumlauf, R.; Rijli, F. M.: Hoxa2 and Hoxb2 control dorsoventral patterns of neuronal development in the rostral hindbrain. Neuron 22:677–691, 1999.
- [66352] 4682.Pollock, R. A.; Sreenath, T.; Ngo, L.; Bieberich, C. J.: Gain of function mutations for paralogous Hox genes: implications for the evolution of Hox gene function. Proc. Nat. Acad. Sci. 92: 4492–4496, 1995.
- [66353] 4683.Vieille–Grosjean, I.; Huber, P.: Transcription factor GATA–1 regulates human HOXB2 gene expression in erythroid cells. J. Biol. Chem. 270:4544–4550, 1995.
- [66354] 4684.Cannizzaro, L. A.; Croce, C. M.; Griffin, C. A.; Simone, A.; Boncinelli, E.; Huebner, K.: Human homeo box–containing genes located at chromosomal regions 2q31–2q37 and 12q12–12q13. Am. J. Hum. Genet. 41: 1–15, 1987.
- [66355] 4685.Chariot, A.; Castronovo, V.; Le, P.; Gillet, C.; Sobel, M. E.; Gielen, J.: Cloning and expression of a new HOXC6 transcript encoding a repressing protein. Biochem. J. 319: 91–97, 1996.

- [66356] 4686. Gibbs, S.; Fijneman, R.; Wiegant, J.; Geurts van Kessel, A.; vande Putte, P.; Backendorf, C.: Molecular characterization and evolution of the SPRR family of keratinocyte differentiation markers encoding small proline-rich proteins. *Genomics* 16: 630–637, 1993.
- [66357] 4687. Marvin, K. W.; George, M. D.; Fujimoto, W.; Saunders, N. A.; Bernacki, S. H.; Jetten, A. M.: Cornifin, a cross-linked envelope precursor in keratinocytes that is down-regulated by retinoids. *Proc. Nat. Acad. Sci.* 89: 11026–11030, 1992.
- [66358] 4688. Bressler, J.; Tsai, T.-F.; Wu, M.-Y.; Tsai, S.-F.; Ramirez, M. A.; Armstrong, D.; Beaudet, A. L.: The SNRPN promoter is not required for genomic imprinting of the Prader-Willi/Angelman domain in mice. *Nature Genet.* 28: 232–240, 2001.
- [66359] 4689. Buiting, K.; Saitoh, S.; Gross, S.; Dittrich, B.; Schwartz, S.; Nicholls, R. D.; Horsthemke, B.: Inherited microdeletions in the Angelman and Prader-Willi syndromes define an imprinting centre on human chromosome 15. *Nature Genet.* 9: 395–400, 1995.
- [66360] 4690. Dittrich, B.; Buiting, K.; Korn, B.; Rickard, S.; Buxton, J.; Saitoh, S.; Nicholls, R. D.; Poustka, A.; Winterpacht, A.; Zabel, B.; Horsthemke, B.: Imprint switching on human

chromosome 15 may involve alternativetranscripts of the SNRPN gene. *Nature Genet.* 14: 163–170, 1996.

[66361] 4691.Gallagher, R. C.; Pils, B.; Albalwi, M.; Francke, U.: Evidencefor the role of PWCR1/HBII–85 C/D box small nucleolar RNAs in Prader–Willisyndrome. *Am. J. Hum. Genet.* 71: 669–678, 2002.

[66362] 4692.Glenn, C. C.; Porter, K. A.; Jong, M. T. C.; Nicholls, R. D.; Driscoll,D. J.: Functional imprinting and epigenetic modification of the humanSNRPN gene. *Hum. Molec. Genet.* 2: 2001–2005, 1993.

[66363] 4693.Gray, T. A.; Saitoh, S.; Nicholls, R. D.: An imprinted, mammalianbicistronic transcript encodes two independent proteins. *Proc. Nat.Acad. Sci.* 96: 5616–5621, 1999.

[66364] 4694.Kuslich, C. D.; Kobori, J. A.; Mohapatra, G.; Gregorio–King, C.;Donlon, T. A.: Prader–Willi syndrome is caused by disruption of theSNRPN gene. *Am. J. Hum. Genet.* 64: 70–76, 1999.

[66365] 4695.Leff, S. E.; Brannan, C. I.; Reed, M. L.; Ozcelik, T.; Francke,U.; Copeland, N. G.; Jenkins, N. A.: Maternal imprinting of the mouseSnrpn gene and conserved linkage homology with the human Prader–Willisyndrome region. *Nature Genet.* 2: 259–264, 1992.

[66366] 4696.Li, S.; Klein, E. S.; Russo, A. F.; Simmons, D. M.;

Rosenfeld, M. G.: Isolation of cDNA clones encoding small nuclear ribonucleoparticle-associated proteins with different tissue specificities. *Proc. Nat. Acad. Sci.* 86:9778–9782, 1989.

[66367] 4697. Lyko, F.; Buiting, K.; Horsthemke, B.; Paro, R.: Identification of a silencing element in the human 15q11–q13 imprinting center by using transgenic *Drosophila*. *Proc. Nat. Acad. Sci.* 95: 1698–1702, 1998.

[66368] 4698. McAllister, G.; Amara, S. G.; Lerner, M. R.: Tissue-specific expression and cDNA cloning of small nuclear ribonucleoprotein-associated polypeptide N. *Proc. Nat. Acad. Sci.* 85: 5296–5300, 1988.

[66369] 4699. Mutirangura, A.; Jayakumar, A.; Sutcliffe, J. S.; Nakao, M.; McKinney, M. J.; Buiting, K.; Horsthemke, B.; Beaudet, A. L.; Chinault, A. C.; Ledbetter, D. H.: A complete YAC contig of the Prader–Willi/Angelman chromosome region (15q11–q13) and refined localization of the SNRPN-gene. *Genomics* 18: 546–552, 1993.

[66370] 4700. Ozcelik, T.; Leff, S.; Robinson, W.; Donlon, T.; Lalande, M.; Sanjines, E.; Schinzel, A.; Francke, U.: Small nuclear ribonucleoprotein polypeptide N (SNRPN), an expressed gene in the Prader–Willi syndrome critical region. *Nature Genet.* 2: 265–269, 1992.

- [66371] 4701. Reed, M. L.; Leff, S. E.: Maternal imprinting of human SNRPN, a gene deleted in Prader-Willi syndrome. *Nature Genet.* 6: 163-167, 1994.
- [66372] 4702. Saitoh, S.; Wada, T.: Parent-of-origin specific histone acetylation and reactivation of a key imprinted gene locus in Prader-Willi syndrome. *Am. J. Hum. Genet.* 66: 1958-1962, 2000.
- [66373] 4703. Kantaputra, P. N.; Gorlin, R. J.; Langer, L. O., Jr.: Dominant mesomelic dysplasia, ankle, carpal, and tarsal synostosis type: a new autosomal dominant bone disorder. *Am. J. Med. Genet.* 44: 730-737, 1992.
- [66374] 4704. Karin, M.; Eddy, R. L.; Henry, W. M.; Haley, L. L.; Byers, M. G.; Shows, T. B.: Human metallothionein genes are clustered on chromosome 16. *Proc. Nat. Acad. Sci.* 81: 5494-5498, 1984.
- [66375] 4705. West, A. K.; Stallings, R.; Hildebrand, C. E.; Chiu, R.; Karin, M.; Richards, R. I.: Human metallothionein genes: structure of the functional locus at 16q13. *Genomics* 8: 513-518, 1990.
- [66376] 4706. Wen, L.; Huang, J.-K.; Johnson, B. H.; Reeck, G. R.: A human placental cDNA clone that encodes nonhistone chromosomal protein HMG-1. *Nucleic Acids Res.* 17: 1197-1214, 1989.

- [66377] 4707.Laing, N. G.; Wilton, S. D.; Akkari, P. A.; Dorosz, S.; Boundy,K.; Kneebone, C.; Blumbergs, P.; White, S.; Watkins, H.; Love, D.R.; Haan, E.: A mutation in the alpha tropomyosin gene TPM3 associatedwith autosomal dominant nemaline myopathy. *Nature Genet.* 9: 75–79,1995.
- [66378] 4708.Shirakawa, H.; Yoshida, M.: Structure of a gene coding for humanHMG2 protein. *J. Biol. Chem.* 267: 6641–6645, 1992.
- [66379] 4709.Wanschura, S.; Schoenmakers, E. F. P. M.; Huysmans, C.; Bartnitzke,S.; Van de Ven, W. J. M.; Bullerdiek, J.: Mapping of the human HMG2gene to 4q31. *Genomics* 31: 264–265, 1996.
- [66380] 4710.Bustin, M.: Regulation of DNA-dependent activities by the functionalmotifs of the high-mobility-group chromosomal proteins. *Molec. Cell.Biol.* 19: 5237–5246, 1999.
- [66381] 4711.Landsman, D.; Bustin, M.: Chromosomal proteins HMG–14 and HMG–17:distinct multigene families coding for similar types of transcripts. *J.Biol. Chem.* 261: 16087–16091, 1986.
- [66382] 4712.Landsman, D.; Soares, N.; Gonzalez, F. J.; Bustin, M.: Chromosomalprotein HMG–17: complete human cDNA sequence and evidence for a multigenefamily. *J. Biol. Chem.* 261: 7479–7484, 1986.

- [66383] 4713.Mitchell, A.; McBride, W.; Landsman, D.; Bustin, M.: Chromosomal mapping of HMG-17 gene to human chromosome 1p. (Abstract) Am. J.Hum. Genet. 43: A152 only, 1988.
- [66384] 4714.Mitchell, A. L.; Bale, A. E.; Bustin, M.; Landsman, D.; Popescu,N.; McBride, O. W.: Localization of HMG17 gene to chromosome 1p35-36.1.(Abstract) Cytogenet. Cell Genet. 51: 1045 only, 1989.
- [66385] 4715.Popescu, N.; Landsman, D.; Bustin, M.: Mapping the human genecoding for chromosomal protein HMG-17. Hum. Genet. 85: 376-378,1990.
- [66386] 4716.Porkka, K.; Laakkonen, P.; Hoffman, J. A.; Bernasconi, M.; Ruoslahti,E.: A fragment of the HMGN2 protein homes to the nuclei of tumorcells and tumor endothelial cells in vivo. Proc. Nat. Acad. Sci. 99:7444-7449, 2002.
- [66387] 4717.Srikantha, J.; Landsman, D.; Bustin, M.: Retrosequences forhuman chromosomal protein HMG-17. J. Molec. Biol. 197: 405-413,1987.
- [66388] 4718.Ding, H.-F.; Rimsky, S.; Batson, S. C.; Bustin, M.; Hansen, U.: Stimulation of RNA polymerase II elongation by chromosomal proteinHMG-14. Science 265: 796-799, 1994.



- [66389] 4719.Landsman, D.; McBride, O. W.; Soares, N.; Crippa, M. P.; Srikantha,T.; Bustin, M.: Chromosomal protein HMG-14: identification, characterization,and chromosome localization of a functional gene from the large human-multigene family. J. Biol. Chem. 264: 3421-3427, 1989.
- [66390] 4720.Pash, J.; Popescu, N.; Matocha, M.; Rapoport, S.; Bustin, M.:Chromosomal protein HMG-14 gene maps to the Down syndrome region ofhuman chromosome 21 and is overexpressed in mouse trisomy 16. Proc.Nat. Acad. Sci. 87: 3836-3840, 1990.
- [66391] 4721.Petersen, M. B.; Economou, E. P.; Slaugenhaupt, S. A.; Chakravarti,A.; Antonarakis, S. E.: Linkage analysis of the human HMG14 geneon chromosome 21 using a GT dinucleotide repeat as polymorphic marker. Genomics 7:136-138, 1990.
- [66392] 4722.Bajalica, S.; Allander, S. V.; Ehrenborg, E.; Brondum-Nielsen,K.; Luthman, H.; Larsson, C.: Localization of the human insulin-likegrowth-factor-binding protein 4 gene to chromosomal region 17q12-21.1. Hum.Genet. 89: 234-236, 1992.
- [66393] 4723.Kiefer, M. C.; Schmid, C.; Waldvogel, M.; Schlapfer, I.; Futo,E.; Masiarz, F. R.; Green, K.; Barr, P. J.; Zapf, J.: Characterizationof recombinant human insulin-like

growth factor binding proteins 4,5, and 6 produced in yeast. J. Biol. Chem. 267: 12692–12699, 1992.

[66394] 4724. Shimasaki, S.; Uchiyama, F.; Shimonaka, M.; Ling, N.: Molecular cloning of the cDNAs encoding a novel insulin-like growth factor-binding protein from rat and human. Molec. Endocr. 4: 1451–1458, 1990.

[66395] 4725. Tonin, P.; Ehrenborg, E.; Lenoir, G.; Feunteun, J.; Lynch, H.; Morgan, K.; Zazzi, H.; Vivier, A.; Pollak, M.; Huynh, H.; Luthman, H.; Larsson, C.; Narod, S.: The human insulin-like growth factor-binding protein 4 gene maps to chromosome region 17q12–q21.1 and is close to the gene for hereditary breast–ovarian cancer. Genomics 18: 414–417, 1993.

[66396] 4726. Zazzi, H.; Nikoshkov, A.; Hall, K.; Luthman, H.: Structure and transcription regulation of the human insulin-like growth factor binding protein 4 gene (IGFBP4). Genomics 49: 401–410, 1998.

[66397] 4727. Kou, K.; James, P. L.; Clemmons, D. R.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Rotwein, P.: Identification of two clusters of mouse insulin-like growth factor binding protein genes on chromosomes 1 and 11. Genomics 21: 653–655, 1994.

[66398] 4728. Brink, P. A.; Steyn, L. T.; Coetzee, G. A.; van der

Westhuyzen,D. R.: Familial hypercholesterolemia in South African Afrikaners:PvuII and StuI DNA polymorphisms in the LDL-receptor gene consistentwith a predominating founder gene effect. Hum. Genet. 77: 32-35,1987.

[66399] 4729.Brown, M. S.; Goldstein, J. L.: Analysis of a mutant strain ofhuman fibroblasts with a defect in the internal-ization of receptor-boundlow density lipoproteins. Cell 9: 663-674, 1976.

[66400] 4730.Brown, M. S.; Goldstein, J. L.: Familial hypercholesterolemia:defective binding of lipoproteins to cultured fibroblasts associatedwith impaired regulation of 3-hydroxy-3-methylglutaryl coenzyme atreductase activity. Proc. Nat. Acad. Sci. 71: 788-792, 1974.

[66401] 4731.Davis, C. G.; Lehrman, M. A.; Russell, D. W.; Anderson, R. G.W.; Brown, M. S.; Goldstein, J. L.: The J.D. mutation in familialhypercholesterolemia: amino acid substitution in cytoplasmic domainimpedes internalization of LDL receptors. Cell 45: 15-24, 1986.

[66402] 4732.De Braekeleer, M.: Hereditary disorders in Saguenay-Lac-St-Jean(Quebec, Canada). Hum. Hered. 41: 141-146, 1991.

[66403] 4733.Defesche, J. C.; Kastelein, J. J. P.: Molecular epidemiologyof familial hypercholesterolaemia. (Letter) Lancet

352: 1643–1644,1998.

- [66404] 4734.Durst, R.; Colombo, R.; Shpitzen, S.; Ben Avi, L.; Friedlander,Y.; Wexler, R.; Raal, F. J.; Marais, D. A.; Defesche, J. C.; Mandelshtam,M. Y.; Kotze, M. J.; Leitersdorf, E.; Meiner, V.: Recent origin andspread of a common Lithuanian mutation, G197del LDLR, causing familialhypercholesterolemia: positive selection is not always necessary toaccount for disease incidence among Ashkenazi Jews. *Am. J. Hum. Genet.* 68:1172–1188, 2001.
- [66405] 4735.Ekstrom, U.; Abrahamson, M.; Floren, C.–H.; Tollig, H.; Wettrell,G.; Nilsson, G.; Sun, X.–M.; Soutar, A. K.; Nilsson–Ehle, P.: Anindividual with a healthy phenotype in spite of a pathogenic LDL receptormutation (C240F). *Clin. Genet.* 55: 332–339, 1999.
- [66406] 4736.Elston, R. C.; Namboodiri, K. K.; Go, R. C. P.; Siervogel, R.M.; Glueck, C. J.: Probable linkage between essential familial hypercholesterolemiaand third complement component (C3). *Cytogenet. Cell Genet.* 16:294–297, 1976.
- [66407] 4737.Feussner, G.; Dobmeyer, J.; Nissen, H.; Hansen, T. S.: UnusualXanthomas in a young patient with heterozygous familial hypercholesterolemiaand type III hyperlipoproteinemia. *Am. J. Med. Genet.* 65: 149–154,1996.
- [66408] 4738.Francke, U.; Brown, M. S.; Goldstein, J. L.: Assign–

ment of the human gene for the low density lipoprotein receptor to chromosome 19: synteny of a receptor, a ligand, and a genetic disease. *Proc. Nat. Acad. Sci.* 81: 2826–2830, 1984.

[66409] 4739. Frank, S. L.; Taylor, B. A.; Lusis, A. J.: Linkage of the mouse LDL receptor gene on chromosome 9. *Genomics* 5: 646–648, 1989.

[66410] 4740. Copeland, N. G.; Silan, C. M.; Kingsley, D. M.; Jenkins, N. A.; Cannizzaro, L. A.; Croce, C. M.; Huebner, K.; Sims, J. E.: Chromosomal location of murine and human IL-1 receptor genes. *Genomics* 9: 44–50, 1991.

[66411] 4741. Dale, M.; Nicklin, M. J.: Interleukin-1 receptor cluster: gene organization of IL1R2, IL1R1, IL1RL2 (IL-1Rrp2), IL1RL1 (T1/ST2), and IL18R1 (IL-1Rrp) on human chromosome 2q. *Genomics* 57: 177–179, 1999.

[66412] 4742. Dower, S. K.; Kronheim, S. R.; Hopp, T. P.; Cantrell, M.; Deeley, M.; Gillis, S.; Henney, C. S.; Urdal, D. L.: The cell surface receptors for interleukin-1(alpha) and interleukin-1(beta) are identical. *Nature* 324: 266–268, 1986.

[66413] 4743. Sims, J. E.; Acres, R. B.; Grubin, C. E.; McMahan, C. J.; Wignall, J. M.; March, C. J.; Dower, S. K.: Cloning the interleukin 1 receptor from human T cells. *Proc. Nat. Acad. Sci.* 86: 8946–8950, 1989.

- [66414] 4744.Habets, G. G. M.; van der Kammen, R. A.; Willemsen, V.; Balemans, M.; Wiegant, J.; Collard, J. G.: Sublocalization of an invasion-inducing locus and other genes on human chromosome 7. *Cytogenet. Cell Genet.* 60:200–205, 1992.
- [66415] 4745.Ballantyne, C. M.; Kozak, C. A.; O'Brien, W. E.; Beaudet, A. L.: Assignment of the gene for intercellular adhesion molecule-1 (Icam-1) to proximal mouse chromosome 9. *Genomics* 9: 547–550, 1991.
- [66416] 4746.Bella, J.; Kolatkar, P. R.; Marlbor, C. W.; Greve, J. M.; Rossmann, M. G.: The structure of the two amino-terminal domains of human ICAM-1 suggests how it functions as a rhinovirus receptor and as an LFA-1 integrin ligand. *Proc. Nat. Acad. Sci.* 95: 4140–4145, 1998.
- [66417] 4747.Bellamy, R.; Kwiatkowski, D.; and Hill, A. V. S.: Absence of an association between intercellular adhesion molecule 1, complement receptor 1 and interleukin 1 receptor antagonist gene polymorphisms and severe malaria in a West African population. *Trans. R. Soc. Trop. Med. Hyg.* 92: 312–316, 1998.
- [66418] 4748.Craig, A.; Fernandez-Reyes, D.; Mesri, M.; McDowall, A.; Altieri, D. C.; Hogg, N.; Newbold, C.: A functional analysis of a natural variant of intercellular adhesion molecule-1 (ICAM-1-Kilifi). *Hum. Molec. Genet.* 9: 525–530, 2000.

- [66419] 4749.Fernandez-Reyes, D.; Craig, A. G.; Kyes, S. A.; Peshu, N.; Snow,R. W.; Berendt, A. R.; Marsh, K.; Newbold, C. I.: A high frequencyAfrican coding polymorphism in the N-terminal domain of ICAM-1 predisposingto cerebral malaria in Kenya. *Hum. Molec. Genet.* 6: 1357-1360, 1997.
- [66420] 4750.Greve, J. M.; Davis, G.; Meyer, A. M.; Forte, C. P.; Yost, S. C.;Marlor, C. W.; Kamarck, M. E.; McClelland, A.: The major human rhinovirusreceptor is ICAM-1. *Cell* 56: 839-847, 1989.
- [66421] 4751.Hill, A. V. S.: The immunogenetics of resistance to malaria. *Proc.Assoc. Am. Phys.* 111: 272-277, 1999.
- [66422] 4752.Katz, F. E.; Parkar, M.; Stanley, K.; Murray, L. J.; Clark, E.A.; Greaves, M. F.: Chromosome mapping of cell membrane antigensexpressed on activated B cells. *Europ. J. Immun.* 15: 103-106, 1985.
- [66423] 4753.Le Beau, M. M.; Ryan, D., Jr.; Pericak-Vance, M. A.: Report ofthe committee on the genetic constitution of chromosomes 18 and 19. *Cytogenet.Cell Genet.* 51: 338-357, 1989.
- [66424] 4754.Lu, T. T.; Cyster, J. G.: Integrin-mediated long-term B cellretention in the splenic marginal zone. *Science* 297: 409-412, 2002.

- [66425] 4755. Prieto, J.; Takei, F.; Gendelman, R.; Christenson, B.; Biberfeld, P.; Patarroyo, M.: MALA-2, mouse homologue of human adhesion molecule ICAM-1 (CD54). *Europ. J. Immun.* 19: 1551-1557, 1989.
- [66426] 4756. Simmons, D.; Makgoba, M. W.; Seed, B.: ICAM, an adhesion ligand of LFA-1, is homologous to the neural cell adhesion molecule NCAM. *Nature* 331: 624-627, 1988.
- [66427] 4757. Gedde-Dahl, T., Jr.; Dupuy, B. M.; Jonassen, R.; Winberg, J.-O.; Anton-Lamprecht, I.; Olaisen, B.: Junctional epidermolysis bullosa inversa (locus EBR2A) assigned to 1q31 by linkage and association of LAMC1. *Hum. Molec. Genet.* 3: 1387-1391, 1994.
- [66428] 4758. Kallunki, T.; Ikonen, J.; Chow, L. T.; Kallunki, P.; Tryggvason, K.: Structure of the human laminin B2 chain gene reveals extensive divergence from the laminin B1 chain gene. *J. Biol. Chem.* 266: 221-228, 1991.
- [66429] 4759. Kallunki, T.; Pikkarainen, T.; Tryggvason, K.; Savolainen, E.-R.: A Pst I polymorphism in the human laminin B2 chain gene on 1q25-q31. *Nucleic Acids Res.* 17: 4423 only, 1989.
- [66430] 4760. Mattei, M.-G.; Weil, D.; Passage, E.; Van Cong, N.; Pribula-Conway, D.; Timpl, R.; Chu, M.-L.: Human gene for laminin B2 chain (LAMB2) maps to the long arm of chro-



mosome 1. (Abstract) Cytogenet. CellGenet. 46: 659 only, 1987.

[66431] 4761. Mattei, M.-G.; Weil, D.; Pribula-Conway, D.; Bernard, M. P.; Passage, E.; Van Cong, N.; Timpl, R.; Chu, M.-L.: cDNA cloning, expression and mapping of human laminin B2 gene to chromosome 1q31. Hum. Genet. 79:235-241, 1988.

[66432] 4762. Miner, J. H.; Patton, B. L.; Lentz, S. I.; Gilbert, D. J.; Jenkins, N. A.; Copeland, N. G.; Sanes, J. R.: The laminin alpha chains: expression, developmental transitions, and chromosomal locations of alpha1-5, identification of heterodimeric laminins 8-11, and cloning of a novel alpha3 isoform. J. Cell Biol. 137: 685-701, 1997.

[66433] 4763. Crozat, A.; Aman, P.; Mandahl, N.; Ron, D.: Fusion of CHOP to a novel RNA-binding protein in human myxoid liposarcoma. Nature 363:640-644, 1993.

[66434] 4764. Mrozek, K.; Karakousis, C. P.; Bloomfield, C. D.: Chromosome 12 breakpoints are cytogenetically different in benign and malignant lipogenic tumors: localization of breakpoints in lipoma to 12q15 and in myxoid liposarcoma to 12q13.3. Cancer Res. 53: 1670-1675, 1993.

[66435] 4765. Panagopoulos, I.; Aman, P.; Mertens, F.; Mandahl, N.; Rydholm, A.; Bauer, H. F. C.; Mitelman, F.: Genomic PCR

detects tumor cells in peripheral blood from patients with myxoid liposarcoma. *Genes Chromosomes Cancer* 17: 102–107, 1996.

- [66436] 4766. Park, J. S.; Luethy, J. D.; Wang, M. G.; Fagnoli, J.; Fornace, A. J., Jr.; McBride, O. W.; Holbrook, N. J.: Isolation, characterization and chromosomal localization of the human GADD153 gene. *Gene* 116:259–267, 1992.
- [66437] 4767. Rabbitts, T. H.; Forster, A.; Larson, R.; Nathan, P.: Fusion of the dominant negative transcription regulator CHOP with a novel gene FUS by translocation t(12;16) in malignant liposarcoma. *Nature Genet.* 4:175–180, 1993.
- [66438] 4768. Ron, D.; Habener, J. F.: CHOP, a novel developmentally regulated nuclear protein that dimerizes with transcription factors C/EBP and LAP and functions as a dominant-negative inhibitor of gene transcription. *Genes Dev.* 6: 439–453, 1992.
- [66439] 4769. Kaneda, Y.; Yoshida, M. C.; Kohno, K.; Uchida, T.; Okada, Y.: Chromosomal assignment of the gene for human elongation factor-2. *Proc. Nat. Acad. Sci.* 81: 3158–3162, 1984.
- [66440] 4770. Rapp, G.; Klaudiny, J.; Hagendorff, G.; Luck, M. R.; Scheit, K. H.: Complete sequence of the coding region of human elongation factor 2 (EF-2) by enzymatic amplifica-

tion of cDNA from human ovarian granulosa cells. *Biol. Chem. Hoppe-Seyler* 370: 1071–1075, 1989.

- [66441] 4771. Chen, I.-T.; Dixit, A.; Rhoads, D. D.; Roufa, D. J.: Homologous ribosomal proteins in bacteria, yeast, and humans. *Proc. Nat. Acad. Sci.* 83: 6907–6911, 1986.
- [66442] 4772. Kenmochi, N.; Kawaguchi, T.; Rozen, S.; Davis, E.; Goodman, N.; Hudson, T. J.; Tanaka, T.; Page, D. C.: A map of 75 human ribosomal protein genes. *Genome Res.* 8: 509–523, 1998.
- [66443] 4773. Nakamichi, N. N.; Kao, F.-T.; Wasmuth, J.; Roufa, D. J.: Ribosomal protein gene sequences map to human chromosomes 5, 8, and 17. *Somat. Cell Molec. Genet.* 12: 225–236, 1986.
- [66444] 4774. Justice, M. J.; Siracusa, L. D.; Gilbert, D. J.; Heisterkamp, N.; Groffen, J.; Chada, K.; Silan, C. M.; Copeland, N. G.; Jenkins, N. A.: A genetic linkage map of mouse chromosome 10: localization of eighteen molecular markers using a single interspecific backcross. *Genetics* 125: 855–866, 1990.
- [66445] 4775. King, M. C.: Personal Communication. Berkeley, Calif. 5/1996.
- [66446] 4776. Korach, K. S.: Insights from the study of animals lacking functional estrogen receptor. *Science* 266:

1524–1527, 1994.

- [66447] 4777.Kos, M.; Reid, G.; Denger, S.; Gannon, F.: Minireview: genomic organization of the human ER- $\alpha$  gene promoter region. *Molec. Endocr.* 15:2057–2063, 2001.
- [66448] 4778.Kumar, R.; Wang, R.-A.; Mazumdar, A.; Talukder, A. H.; Mandal, M.; Yang, Z.; Bagheri-Yarmand, R.; Sahin, A.; Hortobagyi, G.; Adam, L.; Barnes, C. J.; Vadlamudi, R. K.: A naturally occurring MTA1 variant sequesters oestrogen receptor- $\alpha$  in the cytoplasm. *Nature* 418:654–657, 2002.
- [66449] 4779.Lawson, J. S.; Field, A. S.; Champion, S.; Tran, D.; Ishikura, H.; Trichopoulos, D.: Low oestrogen receptor  $\alpha$  expression in normal breast tissue underlies low breast cancer incidence in Japan. (Letter) *Lancet* 354: 1787–1788, 1999.
- [66450] 4780.Lonard, D. M.; Nawaz, Z.; Smith, C. L.; O'Malley, B. W.: The 26S proteasome is required for estrogen receptor- $\alpha$  and coactivator turnover and for efficient estrogen receptor- $\alpha$  transactivation. *Molec. Cell* 5: 939–948, 2000.
- [66451] 4781.Lorentzon, M.; Lorentzon, R.; Backstrom, T.; Nordstrom, P.: Estrogen receptor gene polymorphism, but not estradiol levels, is related to bone density in healthy ado-

lescent boys: a cross-sectional and longitudinal study. *J. Clin. Endocr. Metab.* 84: 4597–4601, 1999.

- [66452] 4782. Mader, S.; Kumar, V.; de Verneuil, H.; Chambon, P.: Three amino acids of the oestrogen receptor are essential to its ability to distinguish oestrogen from a glucocorticoid-responsive element. *Nature* 338:271–274, 1989.
- [66453] 4783. McGuire, W. L.; Chamness, G. C.; Fuqua, S. A. W.: Estrogen receptor variants in clinical breast cancer. *Molec. Endocr.* 5: 1571–1577, 1991.
- [66454] 4784. McGuire, W. L.; Chamness, G. C.; Fuqua, S. A. W.: Abnormal estrogen receptor in clinical breast cancer. *J. Steroid Biochem. Molec. Biol.* 43:243–247, 1992.
- [66455] 4785. McInerney, E. M.; Ince, B. A.; Shapiro, D. J.; Katzenellenbogen, B. S.: A transcriptionally active estrogen receptor mutant is a novel type of dominant negative inhibitor of estrogen action. *Molec. Endocr.* 10:1519–1526, 1996.
- [66456] 4786. Menasce, L. P.; White, G. R. M.; Harrison, C. J.; Boyle, J. M.: Localization of the estrogen receptor locus (ESR) to chromosome 6q25.1 by FISH and a simple post-FISH banding technique. *Genomics* 17:263–265, 1993.
- [66457] 4787. Metzger, D.; White, J. H.; Chambon, P.: The human oestrogen receptor functions in yeast. *Nature* 334: 31–36,

1988.

- [66458] 4788. Murphy, L. C.; Wang, M.; Coutt, A.; Dotzlaw, H.: Novel mutations in the estrogen receptor messenger RNA in human breast cancers. *J. Clin. Endocr. Metab.* 81: 1420–1427, 1996.
- [66459] 4789. Pelletier, G.; El-Alfy, M.: Immunocytochemical localization of estrogen receptors alpha and beta in the human reproductive organs. *J. Clin. Endocr. Metab.* 85: 4835–4840, 2000.
- [66460] 4790. Ponglikitmongkol, M.; Green, S.; Chambon, P.: Genomic organization of the human oestrogen receptor gene. *EMBO J.* 7: 3385–3388, 1988.
- [66461] 4791. Reese, J. C.; Katzenellenbogen, B. S.: Characterization of a temperature-sensitive mutation in the hormone binding domain of the human estrogen receptor: studies in cell extracts and intact cells and their implications for hormone-dependent transcriptional activation. *J. Biol. Chem.* 267: 9868–9873, 1992.
- [66462] 4792. Reese, J. C.; Katzenellenbogen, B. S.: Mutagenesis of cysteines in the hormone binding domain of the human estrogen receptor: alterations in binding and transcriptional activation by covalently and reversibly attaching ligands. *J. Biol. Chem.* 266: 10880–10887, 1991.

- [66463] 4793.Scott, G. K.; Kushner, P.; Vigne, J.-L.; Benz, C. C.: Truncatedforms of DNA-binding estrogen receptors in human breast cancer. *J.Clin. Invest.* 88: 700–706, 1991.
- [66464] 4794.Shang, Y.; Brown, M.: Molecular determinants for the tissue specificityof SERMs. *Science* 295: 2465–2468, 2002.
- [66465] 4795.Shiau, A. K.; Barstad, D.; Loria, P. M.; Cheng, L.; Kushner, P.J.; Agard, D. A.; Greene, G. L.: The structural basis of estrogenreceptor/coactivator recognition and the antagonism of this interactionby tamoxifen. *Cell* 95: 927–937, 1998.
- [66466] 4796.Shupnik, M. A.; Pitt, L. K.; Soh, A. Y.; Anderson, A.; Lopes,M. B.; Laws, E. R., Jr.: Selective expression of estrogen receptoralpha and beta isoforms in human pituitary tumors. *J. Clin. Endocr.Metab.* 83: 3965–3972, 1998.
- [66467] 4797.Simoncini, T.; Hafezi-Moghadam, A.; Brazil, D. P.; Ley, K.; Chin,W. W.; Liao, J. K.: Interaction of oestrogen receptor with the regulatorysubunit of phosphatidylinositol-3-OH kinase. *Nature* 407: 538–541,2000.
- [66468] 4798.Sluyser, M.: Mutations in the estrogen receptor gene. *Hum. Mutat.* 6:97–103, 1995.
- [66469] 4799.Sluyser, M.; Mester, J.: Oncogenes homologous to steroid receptors?(Letter) *Nature* 315: 546 only, 1985.

- [66470] 4800.Smith, E. P.; Boyd, J.; Frank, G. R.; Takahashi, H.; Cohen, R.M.; Specker, B.; Williams, T. C.; Lubahn, D. B.; Korach, K. S.: Estrogenresistance caused by a mutation in the estrogen–receptor gene in aman. New Eng. J. Med. 331: 1056–1061, 1994.
- [66471] 4801.Sudhir, K.; Chou, T. M.; Chatterjee, K.; Smith, E. P.; Williams,T. C.; Kane, J. P.; Malloy, M. J.; Korach, K. S.; Rubanyi, G. M.:Premature coronary artery disease associated with a disruptive mutationin the estrogen receptor gene in a man. Circulation 96: 3774–3777,1997.
- [66472] 4802.Takeyama, J.; Suzuki, T.; Inoue, S.; Kaneko, C.; Nagura, H.; Harada,N.; Sasano, H.: Expression and cellular localization of estrogenreceptors alpha and beta in the human fetus. J. Clin. Endocr. Metab. 86:2258–2262, 2001.
- [66473] 4803.Walter, P.; Green, S.; Greene, G.; Krust, A.; Bornert, J.–M.;Jeltsch, J.–M.; Staub, A.; Jensen, E.; Scrace, G.; Waterfield, M.;Chambon, P.: Cloning of the human estrogen receptor cDNA. Proc.Nat. Acad. Sci. 82: 7889–7893, 1985.
- [66474] 4804.Weel, A. E. A. M.; Uitterlinden, A. G.; Westendorp, I. C. D.;Burger, H.; Schuit, S. C. E.; Hofman, A.; Helmerhorst, T. J. M.; vanLeeuwen, J. P. T. M.; Pols, H. A. P.: Estrogen receptor polymorphismpredicts the onset of natural and surgical menopause. J. Clin. Endocr.Metab. 84:



3146–3150, 1999.

[66475] 4805.Weis, K. E.; Ekena, K.; Thomas, J. A.; Lazennec, G.; Katzenellenbogen, B. S.: Constitutively active human estrogen receptors containing amino acid substitutions for tyrosine 537 in the receptor protein. *Molec.Endocr.* 10: 1388–1398, 1996.

[66476] 4806.Liu, B. P.; Fournier, A.; GrandPre, T.; Strittmatter, S. M.: Myelin-associated glycoprotein as a functional ligand for the Nogo-66 receptor. *Science* 297:1190–1193, 2002.

[66477] 4807.Costa, R. M.; Federov, N. B.; Kogan, J. H.; Murphy, G. G.; Stern, J.; Ohno, M.; Kucherlapati, R.; Jacks, T.; Silva, A. J.: Mechanism for the learning deficits in a mouse model of neurofibromatosis type 1. *Nature* 415: 526–530, 2002.

[66478] 4808.van de Vijver, M. J.; Peterse, J. L.; Mooi, W. J.; Wisman, P.; Lomans, J.; Dalesio, O.; Nusse, R.: NEU-protein overexpression in breast cancer: association with comedo-type ductal carcinoma in situ and limited prognostic value in stage II breast cancer. *New Eng.J. Med.* 319: 1239–1245, 1988.

[66479] 4809.Xie, D.; Shu, X. O.; Deng, Z.; Wen, W.-Q.; Creek, K. E.; Dai, Q.; Gao, Y.-T.; Jin, F.; Zheng, W.: Population-based, case-control study of HER2 genetic polymorphism and breast cancer risk. *J. Nat.Cancer Inst.* 92: 412–417, 2000.

- [66480] 4810. Yamamoto, T.; Ikawa, S.; Akiyama, T.; Semba, K.; Nomura, N.; Miyajima, N.; Saito, T.; Toyoshima, K.: Similarity of protein encoded by the human c-erb-B-2 gene to epidermal growth factor receptor. *Nature* 319:230–234, 1986.
- [66481] 4811. Yang–Feng, T. L.; Schechter, A. L.; Weinberg, R. A.; Francke, U.: Oncogene from rat neuro/glioblastomas (human gene symbol NGL) is located on the proximal long arm of human chromosome 17 and EGFR is confirmed at 7p13–q11.2. (Abstract) *Cytogenet. Cell Genet.* 40:784 only, 1985.
- [66482] 4812. Yu, D.; Jing, T.; Liu, B.; Yao, J.; Tan, M.; McDonnell, T. J.; Hung, M.–C.: Overexpression of ErbB2 blocks Taxol–induced apoptosis by upregulation of p21Cip1, which inhibits p34Cdc2 kinase. *Molec. Cell* 2: 581–591, 1998.
- [66483] 4813. Klappacher, G. W.; Lunyak, V. V.; Sykes, D. B.; Sawka–Verhelle, D.; Sage, J.; Brard, G.; Ngo, S. D.; Gangadharan, D.; Jacks, T.; Kamps, M. P.; Rose, D. W.; Rosenfeld, M. G.: An induced Ets repressor complex regulates growth arrest during terminal macrophage differentiation. *Cell* 109:169–180, 2002.
- [66484] 4814. Klemsz, M.; Hromas, R.; Raskind, W.; Bruno, E.; Hoffman, R.: PE–1, a novel ETS oncogene family member,

localizes to chromosome 1q21–q23. *Genomics* 20:291–294, 1994.

- [66485] 4815.Kastury, K.; Li, J.; Druck, T.; Su, H.; Vogt, P. K.; Croce, C.M.; Huebner, K.: The human homologue of the retroviral oncogene qin maps to chromosome 14q13. *Proc. Nat. Acad. Sci.* 91: 3616–3618, 1994.
- [66486] 4816.Li, J.; Chang, H. W.; Lai, E.; Parker, E. J.; Vogt, P. K.: The oncogene qin codes for a transcriptional repressor. *Cancer Res.* 55:5540–5544, 1995.
- [66487] 4817.Li, J.; Vogt, P. K.: The retroviral oncogene qin belongs to the transcription factor family that includes the homeotic gene fork head. *Proc. Nat. Acad. Sci.* 90: 4490–4494, 1993.
- [66488] 4818.Murphy, D. B.; Wiese, S.; Burfeind, P.; Schmundt, D.; Mattei, M.–G.; Schulz–Schaeffer, W.; Thies, U.: Human brain factor 1, a new member of the fork head gene family. *Genomics* 21: 551–557, 1994.
- [66489] 4819.Wiese, S.; Murphy, D. B.; Schlung, A.; Burfeind, P.; Schmundt, D.; Schnulle, V.; Mattei, M.–G.; Thies, U.: The genes for human brainfactor 1 and 2, members of the fork head gene family, are clustered on chromosome 14q. *Biochim. Biophys. Acta* 1262: 105–112, 1995.
- [66490] 4820.Bustelo, X. R.: Regulatory and signaling properties of

the Vavfamily. *Molec. Cell. Biol.* 20: 1461–1477, 2000.

- [66491] 4821. Bustelo, X. R.; Barbacid, M.: Tyrosine phosphorylation of the VAV proto-oncogene product in activated B cells. *Science* 256: 1196–1199, 1992.
- [66492] 4822. Denkiner, D. J.; Borges, C. R.; Butler, C. L.; Cushman, A. M.; Kawahara, R. S.: Genomic organization and regulation of the vav proto-oncogene. *Biochim. Biophys. Acta* 1491: 253–262, 2000.
- [66493] 4823. Fackler, O. T.; Luo, W.; Geyer, M.; Alberts, A. S.; Peterlin, B. M.: Activation of Vav by Nef induces cytoskeletal rearrangements and downstream effector functions. *Molec. Cell* 3: 729–739, 1999.
- [66494] 4824. Fischer, K.-D.; Zmuidzinas, A.; Gardner, S.; Barbacid, M.; Bernstein, A.; Guidos, C.: Defective T-cell receptor signalling and positive selection of Vav-deficient CD4(+) CD8(+) thymocytes. *Nature* 374: 474–477, 1995.
- [66495] 4825. Katzav, S.; Martin-Zanca, D.; Barbacid, M.: VAV, a novel human oncogene derived from a locus ubiquitously expressed in hematopoietic cells. *EMBO J.* 8: 2283–2290, 1989.
- [66496] 4826. Martinerie, C.; Cannizzaro, L. A.; Croce, C. M.; Huebner, K.; Katzav, S.; Barbacid, M.: The human VAV proto-oncogene maps to chromosome region

19p12–19p13.2. Hum. Genet. 86: 65–68, 1990.

[66497] 4827. Moores, S. L.; Selfors, L. M.; Fredericks, J.; Breit, T.; Fujikawa, K.; Alt, F. W.; Brugge, J. S.; Swat, W.: Vav family proteins couple to diverse cell surface receptors. Molec. Cell. Biol. 20: 6364–6373, 2000.

[66498] 4828. Tarakhovsky, A.; Turner, M.; Schaal, S.; Mee, P. J.; Duddy, L. P.; Rajewsky, K.; Tybulewicz, V. L. J.: Defective antigen receptor-mediated proliferation of B and T cells in the absence of Vav. Nature 374: 467–470, 1995.

[66499] 4829. Zhang, R.; Alt, F. W.; Davidson, L.; Orkin, S. H.; Swat, W.: Defective signalling through the T- and B-cell antigen receptors in lymphoid cells lacking the vav proto-oncogene. Nature 374: 470–473, 1995.

[66500] 4830. Fukuhara, S.; Rowley, J. D.; Variakojis, D.; Sweet, D. L.: Chromosome abnormalities in poorly differentiated lymphocytic lymphoma. Cancer Res. 39: 3119–3128, 1979.

[66501] 4831. Ohno, H.; Fukuhara, S.; Takahashi, R.; Mihara, K.; Sugiyama, T.; Doi, S.; Uchino, H.; Toyoshima, K.: c-myc and bcl-2 genes located on 18q21.3 in a follicular lymphoma cell line carrying a t(14;18) chromosomal translocation. Int. J. Cancer 39: 785–788, 1987.

[66502] 4832. Semba, K.; Nishizawa, M.; Satoh, H.; Fukushige, S.;

Yoshida, M.C.; Sasaki, M.; Matsubara, K.; Yamamoto, T.; Toyoshima, K.: Nucleotide sequence and chromosomal mapping of the human c-yes-2 gene. Jpn.J. Cancer Res. 79: 710–717, 1988.

[66503] 4833.Semba, K.; Yamanashi, Y.; Nishizawa, M.; Sukegawa, J.; Yoshida, M.; Sasaki, M.; Yamamoto, T.; Toyoshima, K.: Location of the c-yes gene on the human chromosome and its expression in various tissues. Science 227:1038–1040, 1985.

[66504] 4834.Silverman, G. A.; Kuo, W.-L.; Taillon-Miller, P.; Gray, J. W.: Chromosomal reassignment: YACs containing both YES1 and thymidylate synthase map to the short arm of chromosome 18. Genomics 15: 442–445, 1993.

[66505] 4835.Sukegawa, J.; Semba, K.; Yamanashi, Y.; Nishizawa, M.; Miyajima, N.; Yamamoto, T.; Toyoshima, K.: Characterization of cDNA clones for the human c-yes gene. Molec. Cell. Biol. 7: 41–47, 1987.

[66506] 4836.Alves, S. E.; Lopez, V.; McEwen, B. S.; Weiland, N. G.: Differential colocalization of estrogen receptor beta (ER-beta) with oxytocin and vasopressin in the paraventricular and supraoptic nuclei of the female rat brain: an immunocytochemical study. Proc. Nat. Acad. Sci. 95:3281–3286, 1998.

- [66507] 4837. Inoue, T.; Kimura, T.; Azuma, C.; Inazawa, J.; Take-mura, M.; Kikuchi, T.; Kubota, Y.; Ogita, K.; Saji, F.: Struc-tural organization of the human oxytocin receptor gene. *J. Biol. Chem.* 269: 32451–32456, 1994.
- [66508] 4838. Kimura, T.; Tanizawa, O.; Mori, K.; Brownstein, M. J.; Okayama, H.: Structure and expression of a human oxy-tocin receptor. *Nature* 356: 526–529, 1992.
- [66509] 4839. Simmons, C. F., Jr.; Clancy, T. E.; Quan, R.; Knoll, J. H. M.: The oxytocin receptor gene (OXTR) localizes to hu-man chromosome 3p25 by fluorescence in situ hybridiza-tion and PCR analysis of somatic cell hybrids. *Genomics* 26: 623–625, 1995.
- [66510] 4840. Yang, M.; Wang, W.; Zhong, M.; Philippi, A.; Lichtarge, O.; Sanborn, B. M.: Lysine 270 in the third intra-cellular domain of the oxytocin receptor is an important determinant for G- $\alpha$ -q coupling specificity. *Molec. Endocr.* 16: 814–823, 2002.
- [66511] 4841. Owerbach, D.; Rutter, W. J.; Cooke, N. E.; Martial, J. A.; Shows, T. B.: The prolactin gene is located on chromo-some 6 in humans. *Science* 212: 815–816, 1981.
- [66512] 4842. Shome, B.; Parlow, A. F.: Human pituitary prolactin (hPRL): the entire linear amino acid sequence. *J. Clin. En-docr. Metab.* 45: 1112–1115, 1977.

- [66513] 4843.Sun, Z.; Lee, M. S.; Rhee, H. K.; Arrandale, J. M.; Dannies, P.S.: Inefficient secretion of human H27A–prolactin, a mutant that does not bind Zn(2+). *Molec. Endocr.* 11: 1544–1551, 1997.
- [66514] 4844.Taggart, R. T.; Mohandas, T. K.; Bell, G. I.: Assignment of the human preprogastricsin (PGC) to chromosome 6 and regional localization of PGC (6pter–p21.1), prolactin PRL (6pter–p21.1). (Abstract) *Cytogenet.Cell Genet.* 46: 701–702, 1987.
- [66515] 4845.Truong, A. T.; Duez, C.; Belayew, A.; Renard, A.; Pictet, R.; Bell, G. I.; Martial, J. A.: Isolation and characterization of the human prolactin gene. *EMBO J.* 3: 429–437, 1984.
- [66516] 4846.Slate, D. L.; Ruddle, F. H.: Fibroblast interferon in man is coded by two loci on separate chromosomes. *Cell* 16: 171–180, 1979.
- [66517] 4847.Rosenberg, M.; RayChaudhury, A.; Shows, T. B.; Le Beau, M. M.; Fuchs, E.: A group of type I keratin genes on human chromosome 17: characterization and expression. *Molec. Cell. Biol.* 8: 722–736, 1988.
- [66518] 4848.Gross, J. A.; Johnston, J.; Mudri, S.; Enselman, R.; Dillon, S.R.; Madden, K.; Xu, W.; Parrish–Novak, J.; Foster, D.; Lofton–Day, C.; Moore, M.; Littau, A.; Grossman, A.;



Haugen, H.; Foley, K.; Blumberg, H.; Harrison, K.; Kindsvo-  
gel, W.; Clegg, C. H.: TACI and BCMA are receptors for a  
TNF homologue implicated in B-cell autoimmune disease.  
Nature 404:995–999, 2000.

[66519] 4849. de Wit, T. P. M.; Morton, H. C.; Capel, P. J. A.; van de  
Winkel, J. G. J.: Structure of the gene for the human  
myeloid IgA Fc receptor (CD89). J. Immun. 155:  
1203–1209, 1995.

[66520] 4850. Kremer, E. J.; Kalatzis, V.; Baker, E.; Callen, D. F.;  
Sutherland, G. R.; Maliszewski, C. R.: The gene for the hu-  
man IgA Fc receptor maps to 19q13.4. Hum. Genet. 89:  
107–108, 1992.

[66521] 4851. Maliszewski, C. R.; March, C. J.; Schoenborn, M. A.;  
Gimpel, S.; Shen, L.: Expression cloning of a human Fc re-  
ceptor for IgA. J. Exp. Med. 172: 1665–1672, 1990.

[66522] 4852. Monteiro, R. C.; Hostoffer, R. W.; Cooper, M. D.;  
Bonner, J. R.; Gartland, G. L.; Kubagawa, H.: Definition of  
immunoglobulin A receptor on eosinophils and their en-  
hanced expression in allergic individuals. J. Clin. Invest.  
92: 1681–1685, 1993.

[66523] 4853. Narita, I.; Goto, S.; Saito, N.; Sakatsume, M.; Jin, S.;  
Omori, K.; Gejyo, F.: Genetic polymorphisms in the pro-  
moter and 5-prime UTR region of the Fc alpha receptor

(CD89) are not associated with a risk of IgA nephropathy. J. Hum. Genet. 46: 694–698, 2001.

[66524] 4854. Pleass, R. J.; Andrews, P. D.; Kerr, M. A.; Woof, J. M.: Alternative splicing of the human IgA Fc receptor CD89 in neutrophils and eosinophils. Biochem. J. 318: 771–777, 1996.

[66525] 4855. Shimokawa, T.; Tsuge, T.; Okumura, K.; Ra, C.: Identification and characterization of the promoter for the gene encoding the human myeloid IgA Fc receptor (Fc-alpha-R, CD89). Immunogenetics 51: 945–954, 2000.

[66526] 4856. Tsuge, T.; Shimokawa, T.; Horikoshi, S.; Tomino, Y.; Ra, C.: Polymorphism in promoter region of Fc-alpha receptor gene in patients with IgA nephropathy. Hum. Genet. 108: 128–133, 2001.

[66527] 4857. Lalley, P. A.; Sakaguchi, A. Y.; Eddy, R. L.; Honey, N. H.; Bell, G. I.; Shen, L.-P.; Rutter, W. J.; Jacobs, J. W.; Heinrich, G.; Chin, W. W.; Naylor, S. L.: Mapping polypeptide hormone genes in the mouse: somatostatin, glucagon, calcitonin, and parathyroid hormone. Cytogenet. Cell Genet. 44: 92–97, 1987.

[66528] 4858. Dillon, J. S.; Tanizawa, Y.; Wheeler, M. B.; Leng, X.-H.; Ligon, B. B.; Rabin, D. U.; Yoo-Warren, H.; Permutt, M. A.; Boyd, A. E., III: Cloning and functional expression of

the human glucagon-like peptide-1 (GLP-1) receptor. *Endocrinology* 133: 1907-1910, 1993.

[66529] 4859. Kershaw, E. E.; Chua, S. C., Jr.; Leibel, R. L.: Localization of a (CA)<sub>n</sub> repeat in glucagon-like peptide-1 receptor gene (Glp1r) to proximal mouse chromosome 17 and its linkage to other markers. *Mammalian Genome* 6: 301-303, 1995.

[66530] 4860. Stoffel, M.; Espinosa, R., III; Le Beau, M. M.; Bell, G. I.: Human glucagon-like peptide-1 receptor gene: localization to chromosome band 6p21 by fluorescence in situ hybridization and linkage of a highly polymorphic simple tandem repeat DNA polymorphism to other markers on chromosome 6. *Diabetes* 42: 1215-1218, 1993.

[66531] 4861. Thorens, B.: Expression cloning of the pancreatic beta cell receptor for the gluco-incretin hormone glucagon-like peptide 1. *Proc. Nat. Acad. Sci.* 89: 8641-8645, 1992.

[66532] 4862. Cohen, P.; Miyazaki, M.; Socci, N. D.; Hagge-Greenberg, A.; Liedtke, W.; Soukas, A. A.; Sharma, R.; Hudgins, L. C.; Ntambi, J. M.; Friedman, J. M.: Role for stearoyl-CoA desaturase-1 in leptin-mediated weight loss. *Science* 297: 240-243, 2002.

[66533] 4863. Comuzzie, A. G.; Hixson, J. E.; Almasy, L.; Mitchell,

B. D.; Mahaney, M. C.; Dyer, T. D.; Stern, M. P.; MacCluer, J. W.; Blangero, J.: A major quantitative trait locus determining serum leptin levels and fat mass is located on human chromosome 2. *Nature Genet.* 15: 273–276, 1997.

[66534] 4864. Dahms, N. M.; Lobel, P.; Breitmeyer, J.; Chirgwin, J. M.; Kornfeld, S.: 46 kd mannose 6-phosphate receptor: cloning, expression, and homology to the 215 kd mannose 6-phosphate receptor. *Cell* 50: 181–192, 1987.

[66535] 4865. Ludwig, T.; Ruther, U.; Metzger, R.; Copeland, N. G.; Jenkins, N. A.; Lobel, P.; Hoflack, B.: Gene and pseudogene of the mouse cation-dependent mannose 6-phosphate receptor: genomic organization, expression, and chromosomal localization. *J. Biol. Chem.* 267: 12211–12219, 1992.

[66536] 4866. Pohlmann, R.; Boeker, M. W. C.; von Figura, K.: The two mannose 6-phosphate receptors transport distinct complements of lysosomal proteins. *J. Biol. Chem.* 270: 27311–27318, 1995.

[66537] 4867. Pohlmann, R.; Nagel, G.; Schmidt, B.; Stein, M.; Lorkowski, G.; Krentler, C.; Cully, J.; Meyer, H. E.; Grzeschik, K.-H.; Mersmann, G.; Hasilik, A.; von Figura, K.: Cloning of a cDNA encoding the human cation-dependent mannose 6-phosphate-specific receptor. *Proc. Nat. Acad. Sci.* 84: 5575–5579, 1987.

- [66538] 4868.Roberts, D. L.; Weix, D. J.; Dahms, N. M.; Kim, J.-J. P.: Molecularbasis of lysosomal enzyme recognition: three-dimensional structureof the cation-dependent mannose 6-phosphate receptor. *Cell* 93: 639–648,1998.
- [66539] 4869.Watanabe, H.; Grubb, J. H.; Sly, W. S.: The overex-pressed human46-kDa mannose 6-phosphate receptor mediates endocytosis and sortingof beta-glucuronidase. *Proc. Nat. Acad. Sci.* 87: 8036–8040, 1990.
- [66540] 4870.Vigon, I.; Mornon, J.-P.; Cocault, L.; Mitjavila, M.-T.; Tambourin,P.; Gisselbrecht, S.; Souyri, M.: Molecular cloning and characterizationof MPL, the human homolog of the v-mpl oncogene: identification ofa member of the hematopoietic growth factor receptor superfamily. *Proc.Nat. Acad. Sci.* 89: 5640–5644, 1992.
- [66541] 4871.Bucan, M.; Gatalica, B.; Nolan, P.; Chung, A.; Leroux, A.; Grossman,M. H.; Nadeau, J. H.; Emanuel, B. S.; Budarf, M.: Comparative mappingof 9 human chromosome 22q loci in the laboratory mouse. *Hum. Molec.Genet.* 2: 1245–1252, 1993.
- [66542] 4872.Griffiths, D. F. R.; Williams, G. T.; Williams, E. D.: Duodenalcarcinoid tumours, phaeochromocytoma and neurofibromatosis: isletcell tumour, phaeochromocytoma and the von Hippel–Lindau complex:two distinctive neu-

roendocrine syndromes. Quart. J. Med. 245:  
769–782,1987.

- [66543] 4873.Mackman, N.; Morrissey, J. H.; Fowler, B.; Edgington, T. S.: Completesequene of the human tissue factor gene, a highly regulated cellularreceptor that initiates the coagulation protease cascade. Biochemistry 28:1755–1762, 1989.
- [66544] 4874.Popescu, N. C.; Bowden, P. E.; DiPaolo, J. A.: Two type II keratingenes are localized on human chromosome 12. Hum. Genet. 82: 109–112,1989.
- [66545] 4875.Pulkkinen, L.; Christiano, A. M.; Knowlton, R. G.; Uitto, J.:Epidermolytic hyperkeratosis (bullous congenital ichthyosiform erythroderma):genetic linkage to chromosome 12q in the region of the type II keratingene cluster. J. Clin. Invest. 91: 357–361, 1993.
- [66546] 4876.Rothnagel, J. A.; Dominey, A. M.; Dempsey, L. D.; Longley, M.A.; Greenhalgh, D. A.; Gagne, T. A.; Huber, M.; Frenk, E.; Hohl, D.;Roop, D. R.: Mutations in the rod domains of keratins 1 and 10 ineptidermolytic hyperkeratosis. Science 257: 1128–1130, 1992.
- [66547] 4877.Schimkat, M.; Baur, M. P.; Henke, J.: Inheritance of some electrophoreticphenotypes of human hair. Hum. Genet. 85: 311–314, 1990.

- [66548] 4878.Sybert, V. P.; Francis, J. S.; Corden, L. D.; Smith, L. T.; Weaver,M.; Stephens, K.; McLean, W. H. I.: Cyclic ichthyosis with epidermolytichyperkeratosis: a phenotype conferred by mutations in the 2B domainof keratin K1. Am. J. Hum. Genet. 64: 732–738, 1999.
- [66549] 4879.Syder, A. J.; Yu, Q.–C.; Paller, A. S.; Giudice, G.; Pearson,R.; Fuchs, E.: Genetic mutations in the K1 and K10 genes of patientswith epidermolytic hyperkeratosis: correlation between location anddisease severity. J. Clin. Invest. 93: 1533–1542, 1994.
- [66550] 4880.Yoon, S.–J.; LeBlanc–Straceski, J.; Ward, D.; Krauter, K.; Kucherlapati,R.: Organization of the human keratin type II gene cluster at 12q13. Genomics 24:502–508, 1994.
- [66551] 4881.Hermouet, S.; Merendino, J. J., Jr.; Gutkind, J. S.; Spiegel, A.M.: Activating and inactivating mutations of the alpha subunit ofG(i2) protein have opposite effects on proliferation of NIH 3T3 cells. Proc.Nat. Acad. Sci. 88: 10455–10459, 1991.
- [66552] 4882.Lerman, B. B.; Dong, B.; Stein, K. M.; Markowitz, S. M.; Linden,J.; Catanzaro, D. F.: Right ventricular outflow tract tachycardiadue to a somatic cell mutation in G protein subunit–alpha–i2. J.Clin. Invest. 101: 2862–2868,

1998.

- [66553] 4883. Lyons, J.; Landis, C. A.; Harsh, G.; Vallar, L.; Grunewald, K.; Feichtinger, H.; Duh, Q.-Y.; Clark, O. H.; Kawasaki, E.; Bourne, H.R.; McCormick, F.: Two G protein oncogenes in human endocrine tumors. *Science* 249:655–659, 1990.
- [66554] 4884. Rudolph, U.; Finegold, M. J.; Rich, S. S.; Harriman, G. R.; Srinivasan, Y.; Brabet, P.; Boulay, G.; Bradley, A.; Birnbaumer, L.: Ulcerative colitis and adenocarcinoma of the colon in G  $\alpha(i2)$ -deficient mice. *Nature Genet.* 10: 143–150, 1995.
- [66555] 4885. Baron, B.; Fernandez, M. A.; Toledo, F.; Le Roscouet, D.; Mayau, V.; Martin, N.; Buttin, G.; Debatisse, M.: The highly conserved Chinese hamster GNAI3 gene maps less than 60 kb from the AMPD2 gene and lacks the intronic U6 snRNA present in its human counterpart. *Genomics* 24:288–294, 1994.
- [66556] 4886. Goodman, F. R.: Limb malformations and the human HOX genes. *Am. J. Med. Genet.* 112: 256–265, 2002.
- [66557] 4887. Magli, M. C.; Barba, P.; Celetti, A.; De Vita, G.; Cillo, C.; Boncinelli, E.: Coordinate regulation of HOX genes in human hematopoietic cells. *Proc. Nat. Acad. Sci.* 88: 6348–6352, 1991.



- [66558] 4888.Pravtcheva, D.; Newman, M.; Hunihan, L.; Lonai, P.; Ruddle, F.H.: Chromosome assignment of the murine Hox-4.1 gene. *Genomics* 5:541-545, 1989.
- [66559] 4889.Taniguchi, Y.; Komatsu, N.; Moriuchi, T.: Overexpression of theHOX4A (HOXD3) homeobox gene in human erythroleukemia HEL cells results in altered adhesive properties. *Blood* 85: 2786-2794, 1995.
- [66560] 4890.Zakany, J.; Duboule, D.: Hox genes and the making of sphincters. *Nature* 401:761-762, 1999.
- [66561] 4891.Mavilio, F.; Simeone, A.; Giampaolo, A.; Faiella, A.; Zappavigna,V.; Acampora, D.; Poiana, G.; Russo, G.; Peschle, C.; Boncinelli,E.: Differential and stage-related expression in embryonic tissues of a new human homeobox gene. *Nature* 324: 664-668, 1986.
- [66562] 4892.Oliver, G.; Sidell, N.; Fiske, W.; Heinzmann, C.; Mohandas, T.;Sparkes, R. S.; De Robertis, E. M.: Complementary homeo protein gradients in developing limb buds. *Genes Dev.* 3: 641-650, 1989.
- [66563] 4893.Shaw, D. J.; Meredith, A. L.; Brook, J. D.; Sarfarazi, M.; Harley,H. G.; Huson, S. M.; Bell, G. I.; Harper, P. S.: Linkage relationships of the insulin receptor gene with the complement component 3, LDLreceptor, apolipoprotein C2 and myotonic dystrophy loci on chromosome19. *Hum.*

Genet. 74: 267–269, 1986.

[66564] 4894. Blangy, A.; Lane, H. A.; d'Herin, P.; Harper, M.; Kress, M.; Nigg, E. A.: Phosphorylation by p34(cdc2) regulates spindle association of human Eg5, a kinesin-related motor essential for bipolar spindle formation in vivo. *Cell* 83: 1159–1169, 1995.

[66565] 4895. Shimada, F.; Suzuki, Y.; Taira, M.; Hashimoto, N.; Nozaki, O.; Makino, H.; Yoshida, S.: Abnormal messenger ribonucleic acid (mRNA) transcribed from a mutant insulin receptor gene in a patient with type A insulin resistance. *Diabetologia* 35: 639–644, 1992.

[66566] 4896. Shimada, F.; Taira, M.; Suzuki, Y.; Hashimoto, N.; Nozaki, O.; Taira, M.; Tatibana, M.; Ebina, Y.; Tawata, M.; Onaya, T.; Makino, H.; Yoshida, S.: Insulin-resistant diabetes associated with partial deletion of insulin-receptor gene. *Lancet* 335: 1179–1181, 1990.

[66567] 4897. Taira, M.; Taira, M.; Hashimoto, N.; Shimada, F.; Suzuki, Y.; Kanatsuka, A.; Nakamura, F.; Ebina, Y.; Tatibana, M.; Makino, H.; Yoshida, S.: Human diabetes associated with a deletion of the tyrosine kinase domain of the insulin receptor. *Science* 245: 63–66, 1989.

[66568] 4898. Takahashi, Y.; Kadowaki, H.; Ando, A.; Quin, J. D.; MacCuish, A. C.; Yazaki, Y.; Akanuma, Y.; Kadowaki, T.:

Two aberrant splicings caused by mutations in the insulin receptor gene in cultured lymphocytes from a patient with Rabson–Mendenhall's syndrome. *J. Clin. Invest.* 101:588–594, 1998.

[66569] 4899. Taylor, S. I.; Cama, A.; Accili, D.; Barbetti, F.; Imano, E.; Kadowaki, H.; Kadowaki, T.: Genetic basis of endocrine disease 1: molecular genetics of insulin resistant diabetes mellitus. *J. Clin. Endocr. Metab.* 73: 1158–1163, 1991.

[66570] 4900. Taylor, S. I.; Marcus–Samuels, B.; Ryan–Young, J.; Leventhal, S.; Elders, M. J.: Genetics of the insulin receptor defect in a patient with extreme insulin resistance. *J. Clin. Endocr. Metab.* 62: 1130–1135, 1986.

[66571] 4901. Taylor, S. I.; Underhill, L. H.; Hedo, J. A.; Roth, J.; Serrano Rios, M.; Blizzard, R. M.: Decreased insulin binding to cultured cells from a patient with the Rabson–Mendenhall syndrome: dichotomy between studies with cultured lymphocytes and cultured fibroblasts. *J. Clin. Endocr. Metab.* 56: 856–861, 1983.

[66572] 4902. Tevaarwerk, G. J. M.; Strickland, K. P.; Lin, C. H.; Hudson, A. J.: Studies on insulin resistance and insulin receptor binding in myotonia dystrophica. *J. Clin. Endocr. Metab.* 49: 216–222, 1979.

[66573] 4903. Ullrich, A.; Bell, J. R.; Chen, E. Y.; Herrera, R.;

Petruzzelli, L. M.; Dull, T. J.; Gray, A.; Coussens, L.; Liao, Y.-C.; Tsubokawa, M.; Mason, A.; Seeburg, P. H.; Grunfeld, C.; Rosen, O. M.; Ramachandran, J.: Human insulin receptor and its relationship to the tyrosine kinase family of oncogenes. *Nature* 313: 756–761, 1985.

[66574] 4904. van der Vorm, E. R.; Kuipers, A.; Kielkopf–Renner, S.; Krans, H. M. J.; Moller, W.; Maassen, J. A.: A mutation in the insulin receptor that impairs proreceptor processing but not insulin binding. *J. Biol. Chem.* 269: 14297–14302, 1994.

[66575] 4905. van der Vorm, E. R.; van der Zon, G. C. M.; Moller, W.; Krans, H. M. J.; Lindhout, D.; Maassen, J. A.: An arg for gly substitution at position 31 in the insulin receptor, linked to insulin resistance, inhibits receptor processing and transport. *J. Biol. Chem.* 267: 66–71, 1992.

[66576] 4906. Ward, G. M.; Harrison, L. C.: Structure of the human erythrocyte insulin receptor. *Diabetes* 35: 101–105, 1986.

[66577] 4907. Wertheimer, E.; Lu, S.-P.; Backeljauw, P. F.; Davenport, M. L.; Taylor, S. I.: Homozygous deletion of the human insulin receptor gene results in leprechaunism. *Nature Genet.* 5: 71–73, 1993.

[66578] 4908. Williams, D. L.; Look, A. T.; Melvin, S. L.; Roberson, P. K.; Dahl, G.; Flake, T.; Stass, S.: New chromosomal

translocations correlate with specific immunophenotypes of childhood acute lymphoblastic leukemia. *Cell* 36:101–109, 1984.

[66579] 4909. Williams, J. F.; McClain, D. A.; Dull, T. J.; Ullrich, A.; Olefsky, J. M.: Characterization of an insulin receptor mutant lacking the subunit processing site. *J. Biol. Chem.* 265: 8463–8469, 1990.

[66580] 4910. Yamamoto–Honda, R.; Koshio, O.; Tobe, K.; Shibasaki, Y.; Momomura, K.; Odawara, M.; Kadowaki, T.; Takaku, F.; Akanuma, Y.; Kasuga, M.: Phosphorylation state and biological function of a mutant human insulin receptor val(996). *J. Biol. Chem.* 265: 14777–14783, 1990.

[66581] 4911. Yang–Feng, T. L.; Francke, U.; Ullrich, A.: Gene for human insulin receptor: localization to site on chromosome 19 involved in pre–B–cell leukemia. *Science* 228: 728–731, 1985.

[66582] 4912. Yoshimasa, Y.; Seino, S.; Whittaker, J.; Kakehi, T.; Kosaki, A.; Kuzuya, H.; Imura, H.; Bell, G. I.; Steiner, D. F.: Insulin–resistant diabetes due to a point mutation that prevents insulin proreceptor processing. *Science* 240: 784–787, 1988.

[66583] 4913. Kurachi, H.; Jobo, K.; Ohta, M.; Kawasaki, T.; Itoh,

N.: A newmember of the insulin receptor family, insulin receptor-related receptor, is expressed preferentially in the kidney. *Biochem. Biophys. Res. Commun.* 187: 934–939, 1992.

[66584] 4914. Shier, P.; Watt, V. M.: Primary structure of a putative receptor for a ligand of the insulin family. *J. Biol. Chem.* 264: 14605–14608, 1989.

[66585] 4915. Shier, P.; Willard, H. F.; Watt, V. M.: Localization of the insulin receptor-related receptor gene to human chromosome 1. *Cytogenet. Cell Genet.* 54: 80–81, 1990.

[66586] 4916. Whitmore, T. E.; Maurer, M. F.; Day, H. L.; Jelmberg, A. C.; Dasovich, M. M.; Sundborg, L. M.; Burkhead, S. K.; Heipel, M. D.; Madden, K. L.; Kramer, J. M.; Kuijper, J. L.; Xu, W. F.; Jaspers, S. R.; Holly, R. D.; Lok, S.: The assignment of the human insulin receptor-related receptor gene (INSRR) to chromosome 1q21–q23 by the use of radiation hybrid mapping. *Cytogenet. Cell Genet.* 87: 93–94, 1999.

[66587] 4917. Alnemri, E. S.; Livingston, D. J.; Nicholson, D. W.; Salvesen, G.; Thornberry, N. A.; Wong, W. W.; Yuan, J.: Human ICE/CED-3 protease nomenclature. (Letter) *Cell* 87: 171 only, 1996.

[66588] 4918. Casano, F. J.; Rolando, A. M.; Mudgett, J. S.; Molin–

eaux, S. M.: The structure and complete nucleotide sequence of the murine gene encoding interleukin-1-beta converting enzyme (ICE). *Genomics* 20:474-481, 1994.

[66589] 4919. Cerretti, D. P.; Hollingsworth, L. T.; Kozlosky, C. J.; Valentine, M. B.; Shapiro, D. N.; Morris, S. W.; Nelson, N.: Molecular characterization of the gene for human interleukin-1-beta converting enzyme (IL1BC). *Genomics* 20:468-473, 1994.

[66590] 4920. Cerretti, D. P.; Kozlosky, C. J.; Mosley, B.; Nelson, N.; Van Ness, K.; Greenstreet, T. A.; March, C. J.; Kronheim, S. R.; Druck, T.; Cannizzaro, L. A.; Huebner, K.; Black, R. A.: Molecular cloning of the interleukin-1-beta converting enzyme. *Science* 256: 97-100, 1992.

[66591] 4921. Humke, E. W.; Ni, J.; Dixit, V. M.: ERICE, a novel FLICE-activatable caspase. *J. Biol. Chem.* 273: 15702-15707, 1998.

[66592] 4922. Koenig, U.; Eckhart, L.; Tschachler, E.: Evidence that caspase-13 is not a human but a bovine gene. *Biochem. Biophys. Res. Commun.* 285:1150-1154, 2001.

[66593] 4923. Baker, E.; Hort, Y. J.; Ball, H.; Sutherland, G. R.; Shine, J.; Herzog, H.: Assignment of the human neuropeptide Y gene to chromosome 7p15.1 by nonisotopic in situ hybridization. *Genomics* 26: 163-164, 1995.

- [66594] 4924.Carr, L. G.; Foroud, T.; Bice, P.; Gobbett, T.; Ivashina, J.; Edenberg,H.; Lumeng, L.; Li, T. K.: A quantitative trait locus for alcoholconsumption in selectively bred rat lines. Alcohol Clin. Exp. Res. 22:884–887, 1998.
- [66595] 4925.Dockray, G. J.: Neuropeptide Y: in search of a function. Neurochem.Int. 8: 9–11, 1986.
- [66596] 4926.Erickson, J. C.; Clegg, K. E.; Palmiter, R. D.: Sensitivity toleptin and susceptibility to seizures of mice lacking neuropeptideY. Nature 381: 415–421, 1996.
- [66597] 4927.Erickson, J. C.; Hollopeter, G.; Palmiter, R. D.: Attenuationof the obesity syndrome of ob/ob mice by the loss of neuropeptideY. Science 274: 1704–1706, 1996.
- [66598] 4928.Hansel, D. E.; Eipper, B. A.; Ronnett, G. V.: Neuropeptide Y functionsas a neuroproliferative factor. Nature 410: 940–943, 2001.
- [66599] 4929.Karvonen, M. K.; Koulu, M.; Pesonen, U.; Uusitupa, M. I. J.; Tammi,A.; Viikari, J.; Simell, O.; Ronnema, T.: Leucine 7 to proline 7polymorphism in the preproneuropeptide Y is associated with birthweight and serum triglyceride concentration in preschool-aged children. J.Clin. Endocr. Metab. 85: 1455–1460, 2000.
- [66600] 4930.Karvonen, M. K.; Pesonen, U.; Koulu, M.; Niskanen, L.; Laakso,M.; Rissanen, A.; Dekker, J. M.; 't Hart, L. M.;



Valve, R.; Uusitupa, M. I.: Association of a leucine(7)-to-proline(7) polymorphism in the signal peptide of neuropeptide Y with high serum cholesterol and LDL cholesterol levels. *Nature Med.* 4: 1434–1437, 1998.

[66601] 4931. Kauhanen, J.; Karvonen, M. K.; Pesonen, U.; Koulu, M.; Tuomainen, T.-P.; Uusitupa, M. I. J.; Salonen, J. T.: Neuropeptide Y polymorphism and alcohol consumption in middle-aged men. *Am. J. Med. Genet.* 93:117–121, 2000.

[66602] 4932. Maccarrone, C.; Jarrott, B.: Neuropeptide Y: a putative neurotransmitter. *Neurochem. Int.* 8: 13–22, 1986.

[66603] 4933. Meisler, M. H.; Spence, J. E.; Dixon, J. E.; Caldwell, R. M.; Minth, C. D.; Beaudet, A. L.: Exclusion of close linkage between the loci for cystic fibrosis and neuropeptide Y on human chromosome 7. *Cytogenet. Cell Genet.* 44: 175–176, 1987.

[66604] 4934. Minth, C. D.; Andrews, P. C.; Dixon, J. E.: Characterization, sequence, and expression of the cloned human neuropeptide Y gene. *J. Biol. Chem.* 261: 11974–11979, 1986.

[66605] 4935. Minth, C. D.; Bloom, S. R.; Polak, J. M.; Dixon, J. E.: Cloning, characterization, and DNA sequence of a human cDNA encoding neuropeptide tyrosine. *Proc. Nat. Acad. Sci.* 81: 4577–4581, 1984.

- [66606] 4936. Takeuchi, T.; Gumucio, D.; Eddy, R.; Meisler, M.; Minth, C.; Dixon, J.; Yamada, T.; Shows, T.: Assignment of the related pancreatic polypeptide (PPY) and neuropeptide Y (NPY) genes to regions on human chromosomes 17 and 7. (Abstract) Cytogenet. Cell Genet. 40: 759 only, 1985.
- [66607] 4937. Takeuchi, T.; Gumucio, D. L.; Yamada, T.; Meisler, M. H.; Minth, C. D.; Dixon, J. E.; Eddy, R. E.; Shows, T. B.: Genes encoding pancreatic polypeptide and neuropeptide Y are on human chromosomes 17 and 7. J. Clin. Invest. 77: 1038–1041, 1986.
- [66608] 4938. Terenghi, G.; Polak, J. M.; Hamid, Q.; O'Brien, E.; Denny, P.; Legon, S.; Dixon, J.; Minth, C. D.; Palay, S. L.; Yasargil, G.; Chan-Palay, V.: Localization of neuropeptide Y mRNA in neurons of human cerebral cortex by means of in situ hybridization with a complementary RNA probe. Proc. Nat. Acad. Sci. 84: 7315–7318, 1987.
- [66609] 4939. Thiele, T. E.; Marsh, D. J.; Ste. Marie, L.; Bernstein, I. L.; Palmiter, R. D.: Ethanol consumption and resistance are inversely related to neuropeptide Y levels. Nature 396: 366–369, 1998.
- [66610] 4940. Niskanen, L.; Karvonen, M. K.; Valve, R.; Koulu, M.; Pesonen, U.; Mercuri, M.; Rauramaa, R.; Toyry, J.; Laakso, M.; Uusitupa, M. I. J.: Leucine 7 to proline 7 polymorphism

in the neuropeptide Y gene is associated with enhanced carotid atherosclerosis in elderly patients with type 2 diabetes and control subjects. *J. Clin. Endocr. Metab.* 85: 2266–2269, 2000.

[66611] 4941. Uusitupa, M. I. J.; Karvonen, M. K.; Pesonen, U.; Koulu, M.: Neuropeptide Y: a novel link between the neuroendocrine system and cholesterol metabolism. *Ann. Med.* 30: 508–510, 1998.

[66612] 4942. Herzog, H.; Baumgartner, M.; Vivero, C.; Selbie, L. A.; Auer, B.; Shine, J.: Genomic organization, localization, and allelic differences in the gene for the human neuropeptide Y Y1 receptor. *J. Biol. Chem.* 268:6703–6707, 1993.

[66613] 4943. Herzog, H.; Darby, K.; Ball, H.; Hort, Y.; Beck-Sickinger, A.; Shine, J.: Overlapping gene structure of the human neuropeptide Y receptor subtypes Y1 and Y5 suggests coordinate transcriptional regulation. *Genomics* 41:315–319, 1997.

[66614] 4944. Herzog, H.; Hort, Y. J.; Ball, H. J.; Hayes, G.; Shine, J.; Selbie, L. A.: Cloned human neuropeptide Y receptor couples to two different second messenger systems. *Proc. Nat. Acad. Sci.* 89: 5794–5798, 1992.

[66615] 4945. Larhammar, D.; Blomqvist, A. G.; Yee, F.; Jazin, E.;

Yoo, H.; Wahlestedt, C.: Cloning and functional expression of a human neuropeptide Y/peptide YY receptor of the Y1 type. *J. Biol. Chem.* 267: 10935–10938, 1992.

[66616] 4946. Lutz, C. M.; Frankel, W. N.; Richards, J. E.; Thompson, D. A.: Neuropeptide Y receptor genes on human chromosome 4q31–q32 map to conserved linkage groups on mouse chromosomes 3 and 8. *Genomics* 41:498–500, 1997.

[66617] 4947. Naveilhan, P.; Hassani, H.; Lucas, G.; Blakeman, K. H.; Hao, J.-X.; Xu, X.-J.; Wiesenfeld-Hallin, Z.; Thoren, P.; Ernfors, P.: Reduced antinociception and plasma extravasation in mice lacking a neuropeptide Y receptor. *Nature* 409: 513–517, 2001.

[66618] 4948. Ammar, D. A.; Eadie, D. M.; Wong, D. J.; Ma, Y.-Y.; Kolakowski, L. F., Jr.; Yang-Feng, T. L.; Thompson, D. A.: Characterization of the human type 2 neuropeptide Y receptor gene (NPY2R) and localization to the chromosome 4q region containing the type 1 neuropeptide Y receptor gene. *Genomics* 38: 392–398, 1996.

[66619] 4949. Gerald, C.; Walker, M. W.; Vaysse, P. J.-J.; He, C.; Branchek, T. A.; Weinshank, R. L.: Expression cloning and pharmacological characterization of a human hippocampal neuropeptide Y/peptide YY Y2 receptor subtype. *J. Biol.*

Chem. 270: 26758–26761, 1995.

- [66620] 4950.Weydert, A.; Daubas, P.; Lazaridis, I.; Barton, P.; Garner, I.;Leader, D. P.; Bonhomme, F.; Catalan, J.; Simon, D.; Guenet, J. L.;Gros, F.; Buckingham, M. E.: Genes for skeletal muscle myosin heavychains are clustered and are not located on the same mouse chromosomeas a cardiac myosin heavy chain gene. Proc. Nat. Acad. Sci. 82:7183–7187, 1985.
- [66621] 4951.Jandreski, M. A.; Sole, M. J.; Liew, C.–C.: Two different formsof beta myosin heavy chain are expressed in human striated muscle. Hum.Genet. 77: 127–131, 1987.
- [66622] 4952.Leinwand, L. A.; Fournier, R. E. K.; Nadal–Ginard, B.; Shows, T.B.: Multigene family for sarcomeric myosin heavy chain in mouse andhuman DNA: localization on a single chromosome. Science 221: 766–769,1983.
- [66623] 4953.Leinwand, L. A.; Fournier, R. E. K.; Nadal–Ginard, B.; Shows, T.B.: Assignment of the sarcomeric myosin heavy chain multigene familyto chromosome 17 in humans and chromosome 11 in the mouse. (Abstract) Cytogenet.Cell Genet. 37: 521–522, 1984.
- [66624] 4954.Leinwand, L. A.; Saez, L.; McNally, E.; Bernardo, N.–G.: Isolationand characterization of human myosin heavy chain genes. Proc. Nat.Acad. Sci. 80: 3716–3720,

1983.

- [66625] 4955.Soussi–Yanicostas, N.; Whalen, R. G.; Petit, C.: Five skeletalmyosin heavy chain genes are organized as a multigene complex in thehuman genome. *Hum. Molec. Genet.* 2: 563–569, 1993.
- [66626] 4956.Weiss, A.; McDonough, D.; Wertman, B.; Acakpo–Satchivi, L.; Montgomery,K.; Kucherlapati, R.; Leinwand, L.; Krauter, K.: Organization ofhuman and mouse skeletal myosin heavy chain gene clusters is highlyconserved. *Proc. Nat. Acad. Sci.* 96: 2958–2963, 1999.
- [66627] 4957.Yoon, S.–J.; Seiler, S. H.; Kucherlapati, R.; Leinwand, L.: Organizationof the human skeletal myosin heavy chain gene cluster. *Proc. Nat.Acad. Sci.* 89: 12078–12082, 1992.
- [66628] 4958.Karsch–Mizrachi, I.; Feghali, R.; Shows, T. B.; Leinwand, L. A.: Generation of a full–length human perinatal myosin heavy chain cDNA. *Gene* 89:289–294, 1990.
- [66629] 4959.Weiss, A.; Schiaffino, S.; Leinwand, L. A.: Comparative sequenceanalysis of the complete human sarcomeric myosin heavy chain family:implications for functional diversity. *J. Molec. Biol.* 290: 61–75,1999.
- [66630] 4960.Castilla, L. H.; Garrett, L.; Adya, N.; Orlic, D.; Dutra, A.; Anderson,S.; Owens, J.; Eckhaus, M.; Bodine, D.; Liu, P.

P.: The fusion gene Cbfb-MYH11 blocks myeloid differentiation and predisposes mice to acute myelomonocytic leukaemia. (Letter) *Nature Genet.* 23: 144–146, 1999.

[66631] 4961. Deng, Z.; Liu, P.; Marlton, P.; Claxton, D. F.; Lane, S.; Callen, D. F.; Collins, F. S.; Siciliano, M. J.: Smooth muscle myosin heavy chain locus (MYH11) maps to 16p13.13–p13.12 and establishes a new region of conserved synteny between human 16p and mouse 16. *Genomics* 18:156–159, 1993.

[66632] 4962. Stewart, R. J.; Pesavento, P. A.; Woerpel, D. N.; Goldstein, L. S. B.: Identification and partial characterization of six members of the kinesin superfamily in *Drosophila*. *Proc. Nat. Acad. Sci.* 88:8470–8474, 1991.

[66633] 4963. Tihiy, F.; Kress, M.; Harper, M.; Dutrillaux, B.; Lemieux, N.: Localization of the human kinesin-related gene to band 10q24 by fluorescence in situ hybridization. *Genomics* 13: 1371–1372, 1992.

[66634] 4964. Shimura, H.; Schlossmacher, M. G.; Hattori, N.; Frosch, M. P.; Trockenbacher, A.; Schneider, R.; Mizuno, Y.; Kosik, K. S.; Selkoe, D. J.: Ubiquitination of a new form of alpha-synuclein by parkin from human brain: implications for Parkinson's disease. *Science* 293:263–269, 2001.

[66635] 4965. Spillantini, M. G.; Divane, A.; Goedert, M.: Assign-

ment of human alpha-synuclein (SNCA) and beta-synuclein (SNCB) genes to chromosomes 4q21 and 5q35. *Genomics* 27: 379–381, 1995.

- [66636] 4966. Spillantini, M. G.; Schmidt, M. L.; Lee, V. M.-Y.; Trojanowski, J. Q.; Jakes, R.; Goedert, M.: Alpha-synuclein in Lewy bodies. *Nature* 388:839–840, 1997.
- [66637] 4967. Calogero, S.; Grassi, F.; Aguzzi, A.; Voigtlander, T.; Ferrier, P.; Ferrari, S.; Bianchi, M. E.: The lack of chromosomal protein Hmg1 does not disrupt cell growth but causes lethal hypoglycaemia in newborn mice. *Nature Genet.* 22: 276–280, 1999.
- [66638] 4968. Ferrari, S.; Finelli, P.; Rocchi, M.; Bianchi, M. E.: The active gene that encodes human high mobility group 1 protein (HMG1) contains introns and maps to chromosome 13. *Genomics* 35: 367–371, 1996.
- [66639] 4969. Scaffidi, P.; Misteli, T.; Bianchi, M. E.: Release of chromatin protein HMGB1 by necrotic cells triggers inflammation. *Nature* 418:191–195, 2002.
- [66640] 4970. Taguchi, A.; Blood, D. C.; del Toro, G.; Canet, A.; Lee, D. C.; Qu, W.; Tanji, N.; Lu, Y.; Lalla, E.; Fu, C.; Hofmann, M. A.; Kislinger, T.; Ingram, M.; Lu, A.; Tanaka, H.; Hori, O.; Ogawa, S.; Stern, D. M.; Schmidt, A. M.: Blockade of RAGE–amphoterin signalling suppresses tumour growth



and metastases. *Nature* 405: 354–360, 2000.

- [66641] 4971.Wang, H.; Bloom, O.; Zhang, M.; Vishnubhakat, J. M.; Ombrellino,M.; Che, J.; Frazier, A.; Yang, H.; Ivanova, S.; Borovikova, L.; Manogue,K. R.; Faist, E.; Abraham, E.; Andersson, J.; Andersson, U.; Molina,P. E.; Abumrad, N. N.; Sama, A.; Tracey, K. J.: HMG–1 as a late mediatorof endotoxin lethality in mice. *Science* 285: 248–251, 1999.
- [66642] 4972.Cetta, F.; Chiappetta, G.; Melillo, R. M.; Petracci, M.; Montalto,G.; Santoro, M.; Fusco, A.: The ret/ptc1 oncogene is activated infamilial adenomatous polyposis–associated thyroid papillary carcinomas. *J.Clin. Endocr. Metab.* 83: 1003–1006, 1998.
- [66643] 4973.Decker, R. A.; Peacock, M. L.; Watson, P.: Hirschsprung diseasein MEN 2A: increased spectrum of RET exon 10 genotypes and stronggenotype–phenotype correlation. *Hum. Molec. Genet.* 7: 129–134, 1998.
- [66644] 4974.Donis–Keller, H.; Dou, S.; Chi, D.; Carlson, K. M.; Toshima, K.;Lairmore, T. C.; Howe, J. R.; Moley, J. F.; Goodfellow, P.; Wells,S. A., Jr.: Mutations in the RET proto-oncogene are associated withMEN 2A and FMTC. *Hum. Molec. Genet.* 2: 851–856, 1993.
- [66645] 4975.Doray, B.; Salomon, R.; Amiel, J.; Pelet, A.; Touraine, R.; Billaud,M.; Attie, T.; Bachy, B.; Munnich, A.; Lyonnet, S.:

Mutation of the RET ligand, neurturin, supports multigenic inheritance in Hirschsprung disease. *Hum. Molec. Genet.* 7: 1449–1452, 1998.

[66646] 4976. Edery, P.; Lyonnet, S.; Mulligan, L. M.; Pelet, A.; Dow, E.; Abel, L.; Holder, S.; Nihoul-Fekete, C.; Ponder, B. A. J.; Munnich, A.: Mutations of the RET proto-oncogene in Hirschsprung's disease. *Nature* 367:378–380, 1994.

[66647] 4977. Eng, C.: The RET proto-oncogene in multiple endocrine neoplasia type 2 and Hirschsprung's disease. *New Eng. J. Med.* 335: 943–951, 1996.

[66648] 4978. Eng, C.; Crossey, P. A.; Mulligan, L. M.; Healey, C. S.; Houghton, C.; Prowse, A.; Chew, S. L.; Dahia, P. L. M.; O'Riordan, J. L. H.; Toledo, S. P. A.; Smith, D. P.; Maher, E. R.; Ponder, B. A. J.: Mutations in the RET proto-oncogene and the von Hippel–Lindau disease tumour suppressor gene in sporadic and syndromic pheochromocytomas. *J. Clin. Genet.* 32: 934–937, 1995.

[66649] 4979. Eng, C.; Mulligan, L. M.: Mutations of the RET proto-oncogene in the multiple endocrine neoplasia type 2 syndromes, related sporadic tumours, and Hirschsprung disease. *Hum. Mutat.* 9: 97–109, 1997.

[66650] 4980. Eng, C.; Mulligan, L. M.; Smith, D. P.; Healey, C. S.; Frilling, A.; Raue, F.; Neumann, H. P. H.; Pfragner, R.;

Behmel, A.; Lorenzo, M. J.; Stonehouse, T. J.; Ponder, M. A.; Ponder, B. A. J.: Mutation of the RET protooncogene in sporadic medullary thyroid carcinoma. *Genes Chromosomes Cancer* 12: 209–212, 1995.

[66651] 4981. Eng, C.; Smith, D. P.; Mulligan, L. M.; Healey, C. S.; Zvelebil, M. J.; Stonehouse, T. J.; Ponder, M. A.; Jackson, C. E.; Waterfield, M. D.; Ponder, B. A. J.: A novel point mutation in the tyrosine kinase domain of the RET protooncogene in sporadic medullary thyroid carcinoma and in a family with FMTC. *Oncogene* 10: 509–513, 1995.

[66652] 4982. Eng, C.; Smith, D. P.; Mulligan, L. M.; Nagai, M. A.; Healey, C. S.; Ponder, M. A.; Gardner, E.; Scheumann, G. F. W.; Jackson, C. E.; Tunnacliffe, A.; Ponder, B. A. J.: Point mutation within the tyrosine kinase domain of the RET protooncogene in multiple endocrine neoplasia type 2B and related sporadic tumors. *Hum. Molec. Genet.* 3: 237–241, 1994.

[66653] 4983. Fearon, E. R.: Human cancer syndromes: clues to the origin and nature of cancer. *Science* 278: 1043–1050, 1997.

[66654] 4984. Fitze, G.; Schreiber, M.; Kuhlisch, E.; Schackert, H. K.; Roesner, D.: Association of RET protooncogene codon 45 polymorphism with Hirschsprung disease. (Letter) *Am.*

J. Hum. Genet. 65: 1469–1473, 1999.

[66655] 4985.Gardner, E.; Mulligan, L. M.; Eng, C.; Healey, C. S.; Kwok, J.B. J.; Ponder, M. A.; Ponder, B. A. J.: Haplotype analysis of MEN2 mutations. Hum. Molec. Genet. 3: 1771–1774, 1994.

[66656] 4986.Grieco, M.; Santoro, M.; Berlingieri, M. T.; Melillo, R. M.; Donghi, R.; Bongarzone, I.; Pierotti, M. A.; Della Porta, G.; Fusco, A.; Vecchio, G.: PTC is a novel rearranged form of the ret proto-oncogene and is frequently detected in vivo in human thyroid papillary carcinomas. Cell 60:557–563, 1990.

[66657] 4987.Hofstra, R. M. W.; Landsvater, R. M.; Ceccherini, I.; Stulp, R.P.; Stelwagen, T.; Luo, Y.; Pasini, B.; Hoppener, J. W. M.; Ploosvan Amstel, H. K.; Romeo, G.; Lips, C. J. M.; Buys, C. H. C. M.: A mutation in the RET proto-oncogene associated with multiple endocrine neoplasia type 2B and sporadic medullary thyroid carcinoma. Nature 367:375–376, 1994.

[66658] 4988.Hoppener, J. W. M.; Lips, C. J. M.: RET receptor tyrosine kinase gene mutations: molecular biological, physiological and clinical aspects. Europ. J. Clin. Invest. 26: 613–624, 1996.

[66659] 4989.Hoppner, W.; Ritter, M. M.: A duplication of 12 bp in

the critical cysteine rich domain of the RET proto-oncogene results in a distinct phenotype of multiple endocrine neoplasia type 2A. Hum. Molec. Genet. 6:587–590, 1997.

[66660] 1990. Ikeda, I.; Ishizaka, Y.; Tahira, T.; Suzuki, T.; Onda, M.; Sugimura, T.; Nagao, M.: Specific expression of the ret proto-oncogene in human neuroblastoma cell lines. Oncogene 5: 1291–1296, 1990.

[66661] 1991. Ishizaka, Y.; Itoh, F.; Tahira, T.; Ikeda, I.; Sugimura, T.; Tucker, J.; Fertitta, A.; Carrano, A. V.; Nagao, M.: Human ret proto-oncogene mapped to chromosome 10q11.2. Oncogene 4: 1519–1521, 1989.

[66662] 1992. Japon, M. A.; Urbano, A. G.; Saez, C.; Segura, D. I.; Cerro, A. L.; Dieguez, C.; Alvarez, C. V.: Glial-derived neurotrophic factor and RET gene expression in normal human anterior pituitary cell types and in pituitary tumors. J. Clin. Endocr. Metab. 87: 1879–1884, 2002.

[66663] 1993. Julies, M. G.; Moore, S. W.; Kotze, M. J.; du Plessis, L.: Novel RET mutations in Hirschsprung's disease patients from the diverse South African population. Europ. J. Hum. Genet. 9: 419–423, 2001.

[66664] 1994. Klugbauer, S.; Demidchik, E. P.; Lengfelder, E.; Rabes, H. M.: Detection of a novel type of RET rearrange-

ment (PTC5) in thyroidcarcinomas after Chernobyl and analysis of the involved RET–fusedgene RFG5. Cancer Res. 58: 198–203, 1998.

- [66665] 4995.de Oca Luna, R. M.; Tabor, A. D.; Eberspaecher, H.; Hulboy, D.L.; Worth, L. L.; Colman, M. S.; Finlay, C. A.; Lozano, G.: The organizationand expression of the mdm2 gene. Genomics 33: 352–357, 1996.
- [66666] 4996.Watson, C. E.; Draganov, D. I.; Billecke, S. S.; Bisgaier, C.L.; La Du, B. N.: Rabbits possess a serum paraoxonase polymorphismsimilar to the human Q192R. Pharmacogenetics 11: 123–134, 2001.
- [66667] 4997.Rajput, B.; Marshall, A.; Killary, A. M.; Lalley, P. A.; Naylor,S. L.; Belin, D.; Rickles, R. J.; Strickland, S.: Chromosomal assignmentsof genes for tissue plasminogen activator and urokinase in mouse. Somat.Cell Molec. Genet. 13: 581–586, 1987.
- [66668] 4998.Yamasaki, Y.; Sakamoto, K.; Watada, H.; Kajimoto, Y.; Hori, M.: The arg–192 isoform of paraoxonase with low sarin–hydrolyzing activityis dominant in the Japanese. Hum. Genet. 101: 67–68, 1997.
- [66669] 4999.Fong, L.; Brockstedt, D.; Benike, C.; Breen, J. K.; Strang, G.;Ruegg, C. L.; Engleman, E. G.: Dendritic cell–based xenoantigen vaccinationfor prostate cancer im–

munotherapy. J. Immun. 167: 7150–7156, 2001.

[66670] 5000.Li, S. S.–L.; Sharief, F. S.: The prostatic acid phosphatase (ACPP) gene is localized to human chromosome 3q21–q23. Genomics 17: 765–766, 1993.

[66671] 5001.Sharief, F. S.; Lee, H.; Leuderman, M. M.; Lundwall, A.; Deaven, L. L.; Lee, C.; Li, S. S.–L.: Human prostatic acid phosphatase: cDNA cloning, gene mapping and protein sequence homology with lysosomal acid phosphatase. Biochem. Biophys. Res. Commun. 160: 79–86, 1989.

[66672] 5002.Sharief, F. S.; Li, S. S.–L.: Nucleotide sequence of human prostatic acid phosphatase ACPP gene, including 7 Alu repeats. Biochem. Molec. Biol. Int. 33: 561–565, 1994.

[66673] 5003.Sharief, F. S.; Li, S. S.–L.: Structure of human prostatic acid phosphatase gene. Biochem. Biophys. Res. Commun. 184: 1468–1476, 1992.

[66674] 5004.Tailor, P. G.; Govindan, M. V.; Patel, P. C.: Nucleotide sequence of human prostatic acid phosphatase determined from a full-length cDNA clone. Nucleic Acids Res. 18: 4928 only, 1990.

[66675] 5005.Winqvist, R.; Virkkunen, P.; Grzeschik, K.–H.; Vihko, P.: Chromosomal localization to 3q21–qter and two TaqI RFLPs of the human prostate-specific acid phosphatase gene (ACPP). Cytogenet. Cell Genet. 52: 68–71, 1989.

- [66676] 5006. Winqvist, R.; Virkkunen, P.; Grzeschik, K.-H.; Vihko, P.: The gene for prostate-specific acid phosphatase is located in segment q21-qter of human chromosome 3 and shows two RFLPs with TaqI. (Abstract) *Cytogenet. Cell Genet.* 51: 1108 only, 1989.
- [66677] 5007. Yeh, L.-C. C.; Lee, A. J.; Lee, N. E.; Lam, K.-W.; Lee, J. C.: Molecular cloning of cDNA for human prostatic acid phosphatase. *Gene* 60:191-196, 1987.
- [66678] 5008. Badger, K. S.; Sussman, H. H.: Structural evidence that human liver and placental alkaline phosphatase isoenzymes are coded by different genes. *Proc. Nat. Acad. Sci.* 73: 2201-2205, 1976.
- [66679] 5009. Beckman, L.; Beckman, G.; Christodoulou, C.; Ifekwunigwe, A.: Variations in human placental alkaline phosphatase. *Acta Genet. Statist. Med.* 17: 406-412, 1967.
- [66680] 5010. Beckman, L.; Bjorling, G.; Christodoulou, C.: Pregnancy enzymes and placental polymorphism: alkaline phosphatase. *Acta Genet. Statist. Med.* 16: 59-73, 1966.
- [66681] 5011. Boyer, S. H.: Alkaline phosphatase in human sera and placenta. *Science* 134:1002-1004, 1961.
- [66682] 5012. Antalis, T. M.; Clark, M. A.; Barnes, T.; Lehrbach, P. R.; Devine, P. L.; Schevzov, G.; Goss, N. H.; Stephens, R. W.; Tolstoshev, P.: Cloning and expression of a cDNA coding



for a human monocyte-derived plasminogen activator inhibitor. *Proc. Nat. Acad. Sci.* 85: 985–989, 1988.

- [66683] 5013. Bartuski, A. J.; Kamachi, Y.; Schick, C.; Overhauser, J.; Silverman, G. A.: Cytoplasmic antiproteinase 2 (PI8) and bomapin (PI10) map to the serpin cluster at 18q21.3. *Genomics* 43: 321–328, 1997.
- [66684] 5014. Dickinson, J. L.; Bates, E. J.; Ferrante, A.; Antalis, T. M.: Plasminogen activator inhibitor type 2 inhibits tumor necrosis factor  $\alpha$ -induced apoptosis: evidence for an alternate biological function. *J. Biol. Chem.* 270: 27894–27904, 1995.
- [66685] 5015. Dougherty, K. M.; Pearson, J. M.; Yang, A. Y.; Westrick, R. J.; Baker, M. S.; Ginsburg, D.: The plasminogen activator inhibitor-2 gene is not required for normal murine development or survival. *Proc. Nat. Acad. Sci.* 96: 686–691, 1999.
- [66686] 5016. Kruithof, E. K. O.; Baker, M. S.; Bunn, C. L.: Biological and clinical aspects of plasminogen activator inhibitor type 2. *Blood* 86: 4007–4024, 1995.
- [66687] 5017. Oldenburg, M.; Wijnen, J. T.; van den Berg, E. A.; le Clercq, E.; Kooistra, T.; Meera Khan, P.: Assignment of plasminogen activator-inhibitor type 2 (PAI2) to chromosome 18. (Abstract) *Cytogenet. Cell Genet.* 51: 1055,

1989.

- [66688] 5018.Samia, J. A.; Alexander, S. J.; Horton, K. W.; Auron, P. E.; Byers, M. G.; Shows, T. B.; Webb, A. C.: Chromosomal organization and localization of the human urokinase inhibitor gene: perfect structural conservation with ovalbumin. *Genomics* 6: 159–167, 1990.
- [66689] 5019.Silverman, G. A.; Jockel, J. I.; Domer, P. H.; Mohr, R. M.; Taillon-Miller, P.; Korsmeyer, S. J.: Yeast artificial chromosome cloning of a two-megabase-size contig within chromosomal band 18q21 establishes physical linkage between BCL2 and plasminogen activator inhibitor type-2. *Genomics* 9:219–228, 1991.
- [66690] 5020.Webb, A. C.; Alexander, S. J.; Samia, J. A.; Auron, P. E.; Byers, M. G.; Shows, T. B.: Localization of the urokinase-type plasminogen activator inhibitor (PLANH2) gene to the long arm of chromosome 18 at 18q21.2–q22.(Abstract) *Cytogenet. Cell Genet.* 51: 1103, 1989.
- [66691] 5021.Webb, A. C.; Collins, K. L.; Snyder, S. E.; Alexander, S. J.; Rosenwasser, L. J.; Eddy, R. L.; Shows, T. B.; Auron, P. E.: Human monocyte arg-serpin cDNA: sequence, chromosomal assignment, and homology to plasminogen activator-inhibitor. *J. Exp. Med.* 166: 77–94, 1987.

- [66692] 5022.Borglum, A.; Anette, B.; Roldan, A. L.; Blasi, F.; Bolund, L.; Kruse, T. A.: Assignment of the gene for urokinase-type plasminogenactivator receptor to chromosome 19q13. (Abstract) Cytogenet. CellGenet. 58: 2016–2017, 1991.
- [66693] 5023.Borglum, A. D.; Byskov, A.; Ragno, P.; Roldan, A. L.; Tripputi, P.; Cassani, G.; Dano, K.; Blasi, F.; Bolund, L.; Kruse, T. A.: Assignment of the urokinase-type plasminogen activator receptor gene (PLAUR) to chromosome 19q13.1–q13.2. Am. J. Hum. Genet. 50: 492–497, 1992.
- [66694] 5024.Dewerchin, M.; Van Nuffelen, A.; Wallays, G.; Bouche, A.; Moons, L.; Carmeliet, P.; Mulligan, R. C.; Collen, D.: Generation and characterization of urokinase receptor-deficient mice. J. Clin. Invest. 97: 870–878, 1996.
- [66695] 5025.Foca, C.; Moses, E. K.; Quinn, M. A.; Rice, G. E.: Differential mRNA expression of urokinase-type plasminogen activator, plasminogenactivator receptor and plasminogen activator inhibitor type-2 in normal human endometria and endometrial carcinomas. Gynec. Oncol. 79: 244–250, 2000.
- [66696] 5026.Memarzadeh, S.; Kozak, K. R.; Chang, L.; Natarajan, S.; Shintaku, P.; Reddy, S. T.; Farias-Eisner, R.: Urokinase plasminogen activator receptor: prognostic biomarker for

endometrial cancer. Proc. Nat.Acad. Sci. 99:  
10647–10652, 2002. Note: Erratum: Proc. Nat. Acad.Sci.  
99: 12501 only, 2002.

- [66697] 5027.Min, H. Y.; Semnani, R.; Mizukami, I. F.; Watt, K.;  
Todd, R. F.,III; Liu, D. Y.: cDNA for Mo3, a monocyte acti-  
vation antigen, encodes the human receptor for urokinase  
plasminogen activator. J. Immun. 148:3636–3642, 1992.
- [66698] 5028.Rijneveld, A. W.; Levi, M.; Florquin, S.; Speelman, P.;  
Carmeliet,P.; van der Poll, T.: Urokinase receptor is neces-  
sary for adequate host defense against pneumococcal  
pneumonia. J. Immun. 168: 3507–3511,2002.
- [66699] 5029.Amagai, M.; Nishikawa, T.; Nousari, H. C.; Anhalt, G.  
J.; Hashimoto,T.: Antibodies against desmoglein 3  
(pemphigus vulgaris antigen)are present in sera from pa-  
tients with paraneoplastic pemphigus and cause acanthol-  
ysis in vivo in neonatal mice. J. Clin. Invest. 102:775–782,  
1998.
- [66700] 5030.Arnemann, J.; Spurr, N. K.; Buxton, R. S.: The human  
gene (DSG3) coding for the pemphigus vulgaris antigen is,  
like the genes coding for the other two known  
desmogleins, assigned to chromosome 18. Hum.Genet.  
89: 347–350, 1992.
- [66701] 5031.Ishikawa, H.; Silos, S. A.; Tamai, K.; Copeland, N. G.;

Gilbert,D. J.; Jenkins, N. A.; Uitto, J.: cDNA cloning and chromosomal assignmentof the mouse gene for desmoglein 3 (Dsg3), the pemphigus vulgarisantigen. Mammalian Genome 5: 803–804, 1994.

[66702] 5032.Silos, S. A.; Tamai, K.; Li, K.; Kivirikko, S.; Kouba, D.; Christiano,A. M.; Uitto, J.: Cloning of the gene for human pemphigus vulgarisantigen (desmoglein 3), a desmosomal cadherin. J. Biol. Chem. 271:17504–17511, 1996.

[66703] 5033.Allikmets, R.; Seddon, J. M.; Bernstein, P. S.; Hutchinson, A.;Atkinson, A.; Sharma, S.; Gerrard, B.; Li, W.; Metzker, M. L.; Wadelius,C.; Caskey, C. T.; Dean, M.; Petrukhin, K.: Evaluation of the Bestdisease gene in patients with age–related macular degeneration andother maculopathies. Hum. Genet. 104: 449–453, 1999.

[66704] 5034.Bascom, R. A.; Liu, L.; Chen, J.; Duncan, A.; Kimberling, W. J.;Moller, C. G.; Humphries, P.; Nathans, J.; McInnes, R. R.: ROM1:a candidate gene for autosomal dominant retinitis pigmentosa (ADRP),Usher syndrome type 1, and Best vitelliform macular dystrophy. (Abstract) Am.J. Hum. Genet. 51 (suppl.): A6, 1992.

[66705] 5035.Best, F.: Ueber eine hereditaere Maculaaffektion. Z. Augenheilk. 13:199–212, 1905.

[66706] 5036.Braley, A. E.: Dystrophy of the macula. Am. J. Oph–

thal. 61:1–24, 1966.

- [66707] 5037.Braley, A. E.; Spivey, B. E.: Hereditary vitelline macular degeneration:a clinical and functional evaluation of a new pedigree with variableexpressivity and dominant inheritance. Arch. Ophthal. 72: 743–762,1964.
- [66708] 5038.Brecher, R.; Bird, A. C.: Adult vitelliform macular dystrophy. Eye 4:210–215, 1990.
- [66709] 5039.Davis, C. T.; Hollenhorst, R. W.: Hereditary degeneration of the macula: occurring in five generations. Am. J. Ophthal. 39: 637–643,1955.
- [66710] 5040.Deutman, A. F.: Electro-oculography in families with vitelliformdystrophy of the fovea: detection of the carrier state. Arch. Ophthal. 81:305–316, 1969.
- [66711] 5041.Falls, H. F.: Hereditary congenital macular degeneration. Am.J. Hum. Genet. 1: 96–104, 1949.
- [66712] 5042.Forsman, K.; Graff, C.; Nordstrom, S.; Johansson, K.; Westermarck,E.; Lundgren, E.; Gustavson, K.–H.; Wadelius, C.; Holmgren, G.: Thegene for Best's macular dystrophy is located at 11q13 in a Swedishfamily. Clin. Genet. 42: 156–159, 1992.
- [66713] 5043.Francois, J.: Vitelliform degeneration of the macula. Bull.N.Y. Acad. Med. 44: 18–27, 1968.
- [66714] 5044.Friedenwald, J. S.; Maumenee, A. E.: Peculiar macular

lesions with unaccountably good vision. Arch. Ophthalmol. 45: 567–570, 1951.

[66715] 5045. Goodstadt, L.; Ponting, C. P.: Sequence variation and disease in the wake of the draft human genome. Hum. Molec. Genet. 10: 2209–2214, 2001.

[66716] 5046. Graff, C.; Eriksson, A.; Forsman, K.; Sandgren, O.; Holmgren, G.; Wadelius, C.: Refined genetic localization of the Best disease gene in 11q13 and physical mapping of linked markers on radiation hybrids. Hum. Genet. 101: 263–270, 1997.

[66717] 5047. Graff, C.; Forsman, K.; Larsson, C.; Nordstrom, S.; Lind, L.; Johansson, K.; Sandgren, O.; Weissenbach, J.; Holmgren, G.; Gustavson, K.-H.; Wadelius, C.: Fine mapping of Best's macular dystrophy localizes the gene in close proximity to but distinct from the D11S480/ROM1 loci. Genomics 24: 425–434, 1994.

[66718] 5048. Hagemeijer, A.; Hoovers, J.; Smit, E. M. E.; Bootsma, D.: Replication pattern of the X chromosomes in three X/autosomal translocations. Cytogenet. Cell Genet. 18: 333–348, 1977.

[66719] 5049. Hou, Y.-C.; Richards, J. E.; Bingham, E. L.; Pawar, H.; Scott, K.; Segal, M.; Lunetta, K. L.; Boehnke, M.; Sieving, P. A.: Linkage study of Best's vitelliform macular dystrophy

(VMD2) in a large NorthAmerican family. Hum. Hered. 46: 211–220, 1996.

[66720] 5050.Jung, E. E.: Ueber eine Sippe mit angeborener Macu-  
ladegeneration. Giessen: Seibert (pub.) 1936.

[66721] 5051.Cullen, T. S.: Embryology, Anatomy, and Diseases of  
the UmbilicusTogether with Diseases of the Urachus.  
Philadelphia: W. B. Saunders(pub.) 1916.

[66722] 5052.Friedman, J. M.: Umbilical dysmorphology: the im-  
portance of contemplatingthe belly button. Clin. Genet.  
28: 343–347, 1985.

[66723] 5053.Legius, E.; de Die–Smulders, C. E. M.; Verbraak, F.;  
Habex, H.;Decorte, R.; Marynen, P.; Fryns, J. P.; Cassiman,  
J. J.: Geneticheterogeneity in Rieger eye malformation. J.  
Med. Genet. 31: 340–341,1994.

[66724] 5054.Phillips, J. C.; Del Bono, E. A.; Haines, J. L.; Pralea, A.  
M.;Cohen, J. S.; Greff, L. J.; Wiggs, J. L.: A second locus for  
Riegersyndrome maps to chromosome 13q14. Am. J.  
Hum. Genet. 59: 613–619,1996.

[66725] 5055.Chu, X.; Thompson, D.; Yee, L. J.; Sung, L. A.: Ge-  
nomic organizationof mouse and human erythrocyte  
tropomodulin genes encoding the pointedend capping  
protein for the actin filaments. Gene 256: 271–281,2000.

[66726] 5056.Fowler, V. M.; Sussmann, M. A.; Miller, P. G.; Flucher,



B. E.;Daniels, M. P.: Tropomodulin is associated with the free (pointed)ends of the thin filaments in rat skeletal muscle. *J. Cell Biol.* 120:411–420, 1993.

[66727] 5057.Gilligan, D. M.; Bennett, V.: The junctional complex of the membraneskeleton. *Semin. Hemat.* 30: 74–83, 1993.

[66728] 5058.Lench, N. J.; Telford, E. A.; Andersen, S. E.; Moynihan, T. P.;Robinson, P. A.; Markham, A. F.: An EST and STS–based YAC contigmap of human chromosome 9q22.3 *Genomics* 38: 199–205, 1996.

[66729] 5059.Sung, L. A.; Fan, Y.–S.; Lin, C. C.: Gene assignment, expression,and homology of human tropomodulin. *Genomics* 34: 92–96, 1996.

[66730] 5060.Sung, L. A.; Fan, Y. S.; Lambert, K.; Fowler, V.; Chien, S.; Lin,C.: Assignment of human erythrocyte tropomodulin gene to q22 of chromosome9. (Abstract) *Cytogenet. Cell Genet.* 58: 1944, 1991.

[66731] 5061.Sung, L. A.; Fowler, V. M.; Lambert, K.; Chien, S.: Molecularcloning of human erythroid tropomodulin. (Abstract) *FASEB J.* 5:A1625, 1991.

[66732] 5062.Sung, L. A.; Fowler, V. M.; Lambert, K.; Sussman, M. A.; Karr,D.; Chien, S.: Molecular cloning and characterization of human fetalliver tropomodulin: a tropomyosin–bind–

ing protein. J. Biol. Chem. 267:2616–2621, 1992.

[66733] 5063.White, R. A.; Dowler, L. L.; Woo, M.; Adkison, L. R.; Pal, S.; Gershon, D.; Fowler, V. M.: The tropomodulin (Tmod) gene maps to chromosome 4, closely linked to Mup1. Mammalian Genome 6: 332–333, 1995.

[66734] 5064.Helander, H. M.; Koivuranta, K. T.; Horelli-Kuitunen, N.; Palvimo, J. J.; Palotie, A.; Hiltunen, J. K.: Molecular cloning and characterization of the human mitochondrial 2,4-dienoyl-CoA reductase gene (DECR). Genomics 46:112–119, 1997.

[66735] 5065.Roe, C. R.; Millington, D. S.; Norwood, D. L.; Kodo, N.; Sprecher, H.; Mohammed, B. S.; Nada, M.; Schulz, H.; McVie, R.: 2,4-Dienoyl-coenzyme A reductase deficiency: a possible new disorder of fatty acid oxidation. J. Clin. Invest. 85: 1703–1707, 1990.

[66736] 5066.Tauchi, H.; Matsuura, S.; Isomura, M.; Kinjo, T.; Nakamura, A.; Sakamoto, S.; Kondo, N.; Endo, S.; Komatsu, K.; Nakamura, Y.: Sequence analysis of an 800-kb genomic DNA region on chromosome 8q21 that contains the Nijmegen breakage syndrome gene, NBS1. Genomics 55: 242–247, 1999.

[66737] 5067.Hamajima, N.; Matsuda, K.; Sakata, S.; Tamaki, N.; Sasaki, M.; Nonaka, M.: A novel gene family defined by hu-

man dihydropyrimidinase and three related proteins with differential tissue distribution. *Gene* 180:157–163, 1996.

[66738] 5068. Wheeler, J. M. D.; Warren, B. F.; Mortensen, N. J. M.; Kim, H.C.; Biddolph, S. C.; Elia, G.; Beck, N. E.; Williams, G. T.; Shepherd, N. A.; Bateman, A. C.; Bodmer, W. F.: An insight into the genetic pathway of adenocarcinoma of the small intestine. *Gut* 50: 218–223, 2002.

[66739] 5069. Bardwick, P. A.; Zvaifler, N. J.; Gill, G. N.; Newman, D.; Greenway, G. D.; Resnick, D. L.: Plasma cell dyscrasia with polyneuropathy, organomegaly, endocrinopathy, M protein, and skin changes: the POEMS syndrome. Report on two cases and a review of the literature. *Medicine* 59:311–322, 1980.

[66740] 5070. Alhonen–Hongisto, L.; Leinonen, P.; Sinervirta, R.; Laine, R.; Winqvist, R.; Alitalo, K.; Janne, O. A.; Janne, J.: Mouse and human ornithine decarboxylase genes: methylation polymorphism and amplification. *Biochem.J.* 242: 205–210, 1987.

[66741] 5071. Brabant, M.; McConlogue, L.; van Daalen Wetters, T.; Coffino, P.: Mouse ornithine decarboxylase gene: cloning, structure, and expression. *Proc. Nat. Acad. Sci.* 85: 2200–2204, 1988.

[66742] 5072. Cox, D. R.; Trouillot, T.; Ashley, P. L.; Brabant, M.;

Coffino,P.: A functional mouse ornithine decarboxylase gene (Odc) maps to chromosome 12: further evidence of homology between mouse chromosome 12 and the short arm of human chromosome 2. *Cytogenet. Cell Genet.* 48:92–94, 1988.

[66743] 5073. Bandmann, O.; Davis, M. B.; Marsden, C. D.; Wood, N. W.: The human homologue of the weaver mouse gene in familial and sporadic Parkinson's disease. *Neuroscience* 72: 877–879, 1996.

[66744] 5074. Dohner, P. H.; Fakhrazadeh, S. S.; Chen, C.-S.; Jockel, J.; Johansen, L.; Silverman, G. A.; Kersey, J. H.; Korsmeyer, S. J.: Acute mixed-lineage leukemia t(4;11)(q21;q23) generates an MLL–AF4 fusion product. *Proc. Nat. Acad. Sci.* 90: 7884–7888, 1993.

[66745] 5075. Gu, Y.; Nakamura, T.; Alder, H.; Prasad, R.; Canaani, O.; Cimino, G.; Croce, C. M.; Canaani, E.: The t(4;11) chromosome translocation of human acute leukemias fuses the ALL-1 gene, related to Drosophila trithorax, to the AF-4 gene. *Cell* 71: 701–708, 1992.

[66746] 5076. Nakamura, T.; Alder, H.; Gu, Y.; Prasad, R.; Canaani, O.; Kamada, N.; Gale, R. P.; Lange, B.; Crist, W. M.; Nowell, P. C.; Croce, C. M.; Canaani, E.: Genes on chromosomes 4, 9, and 19 involved in 11q23 abnormalities in acute

leukemia share sequence homology and/or common motifs. Proc. Nat. Acad. Sci. 90: 4631–4635, 1993.

[66747] 5077. Raffini, L. J.; Slater, D. J.; Rappaport, E. F.; Lo Nigro, L.; Cheung, N.-K. V.; Biegel, J. A.; Nowell, P. C.; Lange, B. J.; Felix, C. A.: Panhandle and reverse-panhandle PCR enable cloning of der(11) and der(other) genomic breakpoint junctions of MLL translocations and identify complex translocation of MLL, AF-4, and CDK6. Proc. Nat. Acad. Sci. 99: 4568–4573, 2002.

[66748] 5078. Hickok, N. J.; Seppanen, P. J.; Gunsalus, G. L.; Janne, O. A.: Complete amino acid sequence of human ornithine decarboxylase deduced from complementary DNA. DNA 6: 179–187, 1987.

[66749] 5079. Michalak, M.; Quackenbush, E. J.; Letarte, M.: Inhibition of  $\text{Na}^+/\text{Ca}^{2+}$  exchanger activity in cardiac and skeletal muscle sarcolemmal vesicles by monoclonal antibody 44D7. J. Biol. Chem. 261: 92–95, 1986.

[66750] 5080. Posillico, J. T.; Srikanta, S.; Brown, E. M.; Eisenbarth, G. S.: The 4F2 cell surface protein modulates intracellular calcium. (Abstract) Clin. Res. 33: 385A only, 1985.

[66751] 5081. Posillico, J. T.; Srikanta, S.; Eisenbarth, G.; Quaranta, V.; Kajiji, S.; Brown, E. M.: Binding of monoclonal antibody (4F2) to its cell surface antigen on dispersed adenomatous

parathyroid cells raises cytosolic calcium and inhibits parathyroid hormone secretion. *J. Clin. Endocr. Metab.* 64: 43–50, 1987.

- [66752] 5082. Quackenbush, E.; Clabby, M.; Gottesdiener, K. M.; Barbosa, J.; Jones, N. H.; Strominger, J. L.; Speck, S.; Leiden, J. M.: Molecular cloning of complementary DNAs encoding the heavy chain of the human 4F2 cell-surface antigen: a type II membrane glycoprotein involved in normal and neoplastic cell growth. *Proc. Nat. Acad. Sci.* 84: 6526–6530, 1987. Note: Correction: *Proc. Nat. Acad. Sci.* 84: 6526–6530, 1987.
- [66753] 5083. Aljada, A.; Ghanim, H.; Saadeh, R.; Dandona, P.: Insulin inhibits NF- $\kappa$ B and MCP-1 expression in human aortic endothelial cells. *J. Clin. Endocr. Metab.* 86: 450–453, 2001.
- [66754] 5084. Harroch, S.; Furtado, G. C.; Brueck, W.; Rosenbluth, J.; Lafaille, J.; Chao, M.; Buxbaum, J. D.; Schlessinger, J.: A critical role for the protein tyrosine phosphatase receptor type Z in functional recovery from demyelinating lesions. *Nature Genet.* 30 Sept, 2002. Note: Advance Electronic Publication.
- [66755] 5085. Marsden, P. A.; Heng, H. H. Q.; Scherer, S. W.; Stewart, R. J.; Hall, A. V.; Shi, X.-M.; Tsui, L.-C.; Schappert, K.

T.: Structure and chromosomal localization of the human constitutive endothelial nitric oxide synthase gene. *J. Biol. Chem.* 268: 17478–17488, 1993.

[66756] 5086. Miyamoto, Y.; Saito, Y.; Nakayama, M.; Shimasaki, Y.; Yoshimura, T.; Yoshimura, M.; Harada, M.; Kajiyama, N.; Kishimoto, I.; Kuwahara, K.; Hino, J.; Ogawa, E.; Hamanaka, I.; Kamitani, S.; Takahashi, N.; Kawakami, R.; Kangawa, K.; Yasue, H.; Nakao, K.: Replication protein A1 reduces transcription of the endothelial nitric oxide synthase gene containing a –786T–C mutation associated with coronary spastic angina. *Hum. Molec. Genet.* 9: 2629–2637, 2000.

[66757] 5087. Nakayama, M.; Yasue, H.; Yoshimura, M.; Shimasaki, Y.; Kugiyama, K.; Ogawa, H.; Motoyama, T.; Saito, Y.; Ogawa, Y.; Miyamoto, Y.; Nakao, K.: T(–786)–C mutation in the 5–prime–flanking region of the endothelial nitric oxide synthase gene is associated with coronary spasm. *Circulation* 99: 2864–2870, 1999.

[66758] 5088. Bloch, K. D.; Wolfram, J. R.; Brown, D. M.; Roberts, J. D., Jr.; Zapol, D. G.; Lepore, J. J.; Filippov, G.; Thomas, J. E.; Jacob, H. J.; Bloch, D. B.: Three members of the nitric oxide synthase II gene family (NOS2A, NOS2B, and NOS2C) colocalize to human chromosome 17. *Genomics* 27: 526–530, 1995.

- [66759] 5089.Chartrain, N. A.; Geller, D. A.; Koty, P. P.; Sitrin, N. F.; Nussler,A. K.; Hoffman, E. P.; Billiar, T. R.; Hutchinson, N. I.; Mudgett,J. S.: Molecular cloning, structure, and chromosomal localizationof the human inducible nitric oxide synthase gene. *J. Biol. Chem.* 269:6765–6772, 1994.
- [66760] 5090.Choi, H.–S.; Rai, P. R.; Chu, H. W.; Cool, C.; Chan, E. D.: Analysisof nitric oxide synthase and nitrotyrosine expression in human pulmonarytuberculosis. *Am. J. Resp. Crit. Care Med.* 166: 178–186, 2002.
- [66761] 5091.Facchetti, F.; Vermi, W.; Fiorentini, S.; Chilosi, M.; Caruso,A.; Duse, M.; Notarangelo, L. D.; Badolato, R.: Expression of inducible nitric oxide synthase in human granulomas and histiocytic reactions. *Am.J. Path.* 154: 145–152, 1999.
- [66762] 5092.Pyronnet, S.; Pradayrol, L.; Sonenberg, N.: A cell cycle–dependentinternal ribosome entry site. *Molec. Cell* 5: 607–616, 2000.
- [66763] 5093.Radford, D. M.; Nakai, H.; Byers, M. G.; Eddy, R. L.; Haley, L.L.; Henry, W. M.; Shows, T. B.: Mapping the ornithine decarboxylase gene (ODC1 and ODC2) to 2p25 and 7q31–qter, respectively. (Abstract) *Cytogenet.Cell Genet.* 46: 678 only, 1987.
- [66764] 5094.Villani, V.; Coffino, P.; D'Eustachio, P.: Linkage ge–



netics of mouse ornithine decarboxylase (Odc). *Genomics* 5: 636–638, 1989.

- [66765] 5095. Winqvist, R.; Makela, T. P.; Seppanen, P.; Janne, O. A.; Alhonen-Hongisto, L.; Janne, J.; Grzeschik, K.-H.; Ali-talo, K.: Human ornithine decarboxylase sequences map to chromosome regions 2pter–p23 and 7cen–qter but are not coamplified with the NMYC oncogene. *Cytogenet. Cell Genet.* 42:133–140, 1986.
- [66766] 5096. Yang-Feng, T. L.; Barton, D. E.; Thelander, L.; Lewis, W. H.; Srinivasan, P. R.; Francke, U.: Ribonucleotide reductase M2 subunit sequences mapped to four different chromosomal sites in humans and mice: functional locus identified by its amplification in hydroxyurea-resistant cell lines. *Genomics* 1: 77–86, 1987.
- [66767] 5097. Clark, C. C.; Cohen, I.; Eichstetter, I.; Cannizzaro, L. A.; McPherson, J. D.; Wasmuth, J. J.; Iozzo, R. V.: Molecular cloning of the human proto-oncogene Wnt-5A and mapping of the gene (WNT5A) to chromosome 3p14–p21. *Genomics* 18: 249–260, 1993.
- [66768] 5098. He, X.; Saint-Jeannet, J.-P.; Wang, Y.; Nathans, J.; Dawid, I.; Varmus, H.: A member of the frizzled protein family mediating axis induction by Wnt-5A. *Science* 275: 1652–1654, 1997.

- [66769] 5099. Adelaide, J.; Mattei, M.-G.; Marics, I.; Raybaud, F.; Planche, J.; De Lapeyriere, O.; Birnbaum, D.: Chromosomal localization of the hst oncogene and its co-amplification with the int.2 oncogene in a human melanoma. *Oncogene* 2: 413–416, 1988.
- [66770] 5100. Dudley, A. T.; Ros, M. A.; Tabin, C. J.: A re-examination of proximodistal patterning during vertebrate limb development. *Nature* 418: 539–544, 2002.
- [66771] 5101. Feldman, B.; Poueymirou, W.; Papaionnou, V. E.; DeChiara, T. M.; Goldfarb, M.: Requirement of FGF-4 for postimplantation mouse development. *Science* 267: 246–249, 1995.
- [66772] 5102. Huebner, K.; Ferrari, A. C.; Delli Bovi, P.; Croce, C. M.; Basilico, C.: The FGF-related oncogene, K-FGF, maps to human chromosome region 11q13, possibly near int-2. *Oncogene Res.* 3: 263–270, 1988.
- [66773] 5103. Sakamoto, H.; Mori, M.; Taira, M.; Yoshida, T.; Matsukawa, S.; Shimizu, K.; Sekiguchi, M.; Terada, M.; Sugimura, T.: Transforming gene from human stomach cancers and a noncancerous portion of stomach mucosa. *Proc. Nat. Acad. Sci.* 83: 3997–4001, 1986.
- [66774] 5104. Sun, X.; Lewandoski, M.; Meyers, E. N.; Liu, Y.-H.; Maxson, R.E., Jr.; Martin, G. R.: Conditional inactivation of

Fgf4 reveals complexity of signalling during limb bud development. *Nature Genet.* 25:83–86, 2000.

[66775] 5105.Sun, X.; Mariani, F. V.; Martin, G. R.: Functions of FGF signalling from the apical ectodermal ridge in limb development. *Nature* 418:501–508, 2002.

[66776] 5106.Lesch, K.-P.; Bengel, D.; Heils, A.; Sabol, S. Z.; Greenberg, B.D.; Petri, S.; Benjamin, J.; Muller, C. R.; Hamer, D. H.; Murphy, D. L.: Association of anxiety-related traits with a polymorphism in the serotonin transporter gene regulatory region. *Science* 274:1527–1530, 1996.

[66777] 5107.Sharp, D.; Blinderman, L.; Combs, K. A.; Kienzle, B.; Ricci, B.; Wager-Smith, K.; Gil, C. M.; Turck, C. W.; Bouma, M.-E.; Rader, D.J.; Aggerbeck, L. P.; Gregg, R. E.; Gordon, D. A.; Wetterau, J. R.: Cloning and gene defects in microsomal triglyceride transfer protein associated with abetalipoproteinaemia. *Nature* 365: 65–69, 1993.

[66778] 5108.Julier, C.; Lathrop, M.; Lalouel, J. M.; Kaplan, J. C.: Use of multilocus tests of gene order: example for chromosome 22. (Abstract) *Cytogenet.Cell Genet.* 40: 663–664, 1985.

[66779] 5109.Julier, C.; Lathrop, M.; Lalouel, J. M.; Reghis, A.; Szajnert, M. F.; Kaplan, J. C.: New restriction fragment length polymorphism on human chromosome 22 at loci SIS, MB

and IGLV. (Abstract) Cytogenet. Cell Genet. 40: 664 only, 1985.

[66780] 5110. Julier, C.; Reghis, A.; Szajnert, M. F.; Kaplan, J. C.; Lathrop, G. M.; Lalouel, J. M.: A preliminary linkage map of human chromosome 22. (Abstract) Cytogenet. Cell Genet. 40: 665 only, 1985.

[66781] 5111. Nakamura, Y.; Leppert, M.; O'Connell, P.; Wolff, R.; Holm, T.; Culver, M.; Martin, C.; Fujimoto, E.; Hoff, M.; Kumlin, E.; White, R.: Variable number of tandem repeat (VNTR) markers for human genemapping. Science 235: 1616–1622, 1987.

[66782] 5112. Romero-Herrera, A. E.; Lehmann, H.: The amino acid sequence of human myoglobin and its minor fractions. Proc. Roy. Soc. London 186B: 249–279, 1974.

[66783] 5113. Romero-Herrera, A. E.; Lehmann, H.: Primary structure of human myoglobin. Nature N.B. 232: 149–152, 1971.

[66784] 5114. Weller, P.; Jeffreys, A. J.; Wilson, V.; Blanchetot, A.: Organization of the human myoglobin gene. EMBO J. 3: 439–446, 1984.

[66785] 5115. Wyman, A. R.; White, R.: A highly polymorphic locus in human DNA. Proc. Nat. Acad. Sci. 77: 6754–6758, 1980.

[66786] 5116. Almarri, A.; Batchelor, J. R.: HLA and hepatitis B in–

fection. Lancet 344:1194–1195, 1994.

- [66787] 5117. Ayala, F. J.; Escalante, A.; O'hUigin, C.; Klein, J.: Molecular genetics of speciation and human origins. Proc. Nat. Acad. Sci. 91:6787–6794, 1994.
- [66788] 5118. Bell, J. I.; Estess, P.; St. John, T.; Saiki, R.; Watling, D. L.; Erlich, H. A.; McDevitt, H. O.: DNA sequence and characterization of human class II major histocompatibility complex beta chains from the DR1 haplotype. Proc. Nat. Acad. Sci. 82: 3405–3409, 1985.
- [66789] 5119. Bodmer, J. G.; Marsh, S. G. E.; Albert, E.: Nomenclature for factors of the HLA system, 1989. Immun. Today 11: 3–10, 1990.
- [66790] 5120. Bodmer, J. G.; Marsh, S. G. E.; Albert, E. D.; Bodmer, W. F.; Bontrop, R. E.; Charron, D.; Dupont, B.; Erlich, H. A.; Fauchet, R.; Mach, B.; Mayr, W. R.; Parham, P.; Sasazuki, T.; Schreuder, G. M. T.; Strominger, J. L.; Svejgaard, A.; Terasaki, P. I.: Nomenclature for factors of the HLA system, 1996. Europ. J. Immunogenet. 24: 105–151, 1997.
- [66791] 5121. Bodmer, J. G.; Marsh, S. G. E.; Albert, E. D.; Bodmer, W. F.; Dupont, B.; Erlich, H. A.; Mach, B.; Mayr, W. R.; Parham, P.; Sasazuki, T.; Schreuder, G. M. T.; Strominger, J. L.; Svejgaard, A.; Terasaki, P. I.: Nomenclature for factors of the HLA system, 1994. Tissue Antigens 44:1–18, 1994.

- [66792] 5122. Bodmer, W. F.; Bodmer, J. G.; Batchelor, J. R.; Festenstein, H.; Morris, P. J.: Histocompatibility Testing 1977. Copenhagen: Munksgaard(pub.) 1978.
- [66793] 5123. Boss, J. M.; Strominger, J. L.: Cloning and sequence analysis of the human major histocompatibility complex gene DC-3-beta. Proc. Nat. Acad. Sci. 81: 5199-5203, 1984.
- [66794] 5124. Charron, D. J.; McDevitt, H. O.: Analysis of HLA-D region-associated molecules with monoclonal antibody. Proc. Nat. Acad. Sci. 76: 6567-6571, 1979.
- [66795] 5125. Chelladurai, M.; Honn, K. V.; Walz, D. A.: HLA-DR is a procoagulant. Biochem. Biophys. Res. Commun. 178: 467-473, 1991.
- [66796] 5126. Corte, G.; Damiani, G.; Calabi, F.; Fabbi, M.; Bargellesi, A.: Analysis of HLA-DR polymorphism by two-dimensional peptide mapping. Proc. Nat. Acad. Sci. 78: 534-538, 1981.
- [66797] 5127. Das, H. K.; Lawrance, S. K.; Weissman, S. M.: Structure and nucleotide sequence of the heavy chain gene of HLA-DR. Proc. Nat. Acad. Sci. 80: 3543-3547, 1983.
- [66798] 5128. Delovitch, T. L.; Falk, J. A.: Evidence for structural homology between murine and human Ia antigens. Immunogenetics 8: 405-418, 1979.

- [66799] 5129.Dunham, I.; Sargent, C. A.; Dawkins, R. L.; Campbell, R. D.:An analysis of variation in the long-range genomic organization ofthe human major histocompatibility complex class II region by pulsed-fieldgel electrophoresis. *Genomics* 5: 787–796, 1989.
- [66800] 5130.Erlich, H. A.; Stetler, D.; Saiki, R.; Gladstone, P.; Pious, D.: Mapping of the genes encoding the HLA-DR alpha chain and the HLA-relatedantigens to a chromosome 6 deletion by using genomic blotting. *Proc.Nat. Acad. Sci.* 80: 2300–2304, 1983.
- [66801] 5131.Ferber, K. M.; Keller, E.; Albert, E. D.; Ziegler, A.–G.: Predictivevalue of human leukocyte antigen class II typing for the developmentof islet autoantibodies and insulin-dependent diabetes postpartumin women with gestational diabetes. *J. Clin. Endocr. Metab.* 84:2342–2348, 1999.
- [66802] 5132.Fuller, T. C.; Einarson, M.; Pinto, C.; Ahern, A.; Yunis, E. J.: Genetic evidence that HLA-DR (Ia) specifications include multipleHLA-D determinants on a single haplotype. *Transplant. Proc.* 10:781–784, 1978.
- [66803] 5133.Hardy, D. A.; Bell, J. I.; Long, E. O.; Lindsten, T.; McDevitt,H. O.: Mapping of the class II region of the human major histocompatibilitycomplex by pulsed-field gel electrophoresis. *Nature* 323: 453–455,1986.

- [66804] 5134.Hui, K.; Festenstein, H.; de Klein, A.; Grosveld, G.; Grosveld,F.: HLA-DR genotyping by restriction fragment length polymorphismanalyses. Immunogenetics 22: 231-239, 1985.
- [66805] 5135.Kaufman, J. F.; Strominger, J. L.: HLA-DR light chain has a polymorphicN-terminal region and a conserved immunoglobulin-like C-terminal region. Nature 297:694-697, 1982.
- [66806] 5136.Korman, A. J.; Auffray, C.; Schamboeck, A.; Strominger, J. L.: The amino acid sequence and gene organization of the heavy chainof the HLA-DR antigen: homology to immunoglobulins. Proc. Nat. Acad.Sci. 79: 6013-6017, 1982.
- [66807] 5137.Korman, A. J.; Knudsen, P. J.; Kaufman, J. F.; Strominger, J.L.: cDNA clones for the heavy chain of HLA-DR antigens obtained afterimmunopurification of polysomes by monoclonal antibody. Proc. Nat.Acad. Sci. 79: 1844-1848, 1982.
- [66808] 5138.Kratzin, H.; Yang, C.; Gotz, H.; Pauly, E.; Kolbel, S.; Egert,G.; Thinnes, F. P.; Wernet, P.; Altevogt, P.; Hilschmann, N.: Primärstrukturmenschlicher Histokompatibilitätsantigene der Klasse II. 1. Mitteilung:aminosauresequenz der N-terminalen 198 Reste



der beta-Kette des HLA-Dw2,2;DR2,2-Alloantigens.

HoppeSeylers Z. Physiol. Chem. 362: 1665–1669, 1981.

[66809] 5139.Lamm, L. U.: Another segregant series, DR, in HLA. Cytogenet.Cell Genet. 22: 309–312, 1978.

[66810] 5140.Larhammar, D.; Schenning, L.; Gustafsson, K.; Wiman, K.; Claesson,L.; Rask, L.; Peterson, P. A.: Complete amino acid sequence of anHLA-DR antigen-like beta chain as predicted from the nucleotide sequence:similarities with immunoglobulins and HLA-A, -B, and -C antigens. Proc.Nat. Acad. Sci. 79: 3687–3691, 1982.

[66811] 5141.Lee, J. S.; Trowsdale, J.; Bodmer, W. F.: cDNA clones codingfor the heavy chain of human HLA-DR antigen. Proc. Nat. Acad. Sci. 79:545–549, 1982.

[66812] 5142.Levine, F.; Mach, B.; Long, E.; Erlich, H.; Pious, D.: Mappingin the HLA-D region with deletion variants and cloned genes. (Abstract) Cytogenet.Cell Genet. 37: 523 only, 1984.

[66813] 5143.Mann, D. L.; Abelson, L.; Harris, S. D.; Amos, D. B.: Secondgenetic locus in the HLA region for human B-cell alloantigens. Nature 259:145–146, 1976.

[66814] 5144.Mann, D. L.; Abelson, L.; Henkart, P.; Harris, S. D.; Amos, D.B.: Specific B-lymphocyte alloantigens linked to HLA. Proc. Nat.Acad. Sci. 72: 5103–5106, 1975.

- [66815] 5145. Markert, M. L.; Cresswell, P.: Polymorphism of human B-cell alloantigens: evidence for three loci within the HLA system. *Proc. Nat. Acad. Sci.* 77:6101–6104, 1980.
- [66816] 5146. Mayer, W. E.; O'hUigin, C.; Klein, J.: Resolution of the HLA-DRB6 puzzle: a case of grafting a de novo-generated exon on an existing gene. *Proc. Nat. Acad. Sci.* 90: 10720–10724, 1993.
- [66817] 5147. McDevitt, H. O.; Bodmer, W. F.: HLA, immune-response genes, and disease. *Lancet* I: 1269–1275, 1974.
- [66818] 5148. Disteche, C. M.; Adler, D. A.; Tedder, T. F.; Saito, H.: Mapping of the genes for LYAM1, a new lymphocyte adhesion molecule, and for LAR, a new receptor-linked protein tyrosine phosphatase, to human chromosome 1. (Abstract) *Cytogenet. Cell Genet.* 51: 990 only, 1989.
- [66819] 5149. Wijmenga, C.; Hansen, R. S.; Gimelli, G.; Bjorck, E. J.; Davies, E. G.; Valentine, D.; Belohradsky, B. H.; van Dongen, J. J.; Smeets, D. F. C. M.; van den Heuvel, L. P. W. J.; Luyten, J. A. F. M.; Strengman, E.; Weemaes, C.; Pearson, P. L.: Genetic variation in ICF syndrome: evidence for genetic heterogeneity. *Hum. Mutat.* 16: 509–517, 2000.
- [66820] 5150. Ehrlich, M.; Buchanan, K. L.; Tsien, F.; Jiang, G.; Sun, B.; Uicker, W.; Weemaes, C. M. R.; Smeets, D.; Sperling, K.; Belohradsky, B. H.; Tommerup, N.; Miesek, D. E.; Rouillard,

J.-M.; Kuick, R.; Hanash, S.M.: DNA methyltransferase 3B mutations linked to the ICF syndrome cause dysregulation of lymphogenesis genes. *Hum. Molec. Genet.* 10:2917–2931, 2001.

[66821] 5151. Fredrickson, D. S.; Levy, R. I.: Familial hyperlipoproteinemia. In: Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: *The Metabolic Basis of Inherited Disease*. New York: McGraw-Hill (pub.) (3rd ed.): 1972. Pp. 545–614.

[66822] 5152. Ameis, D.; Kobayashi, J.; Davis, R. C.; Ben-Zeev, O.; Malloy, M. J.; Kane, J. P.; Lee, G.; Wong, H.; Havel, R. J.; Schotz, M. C.: Familial chylomicronemia (type I hyperlipoproteinemia) due to a single missense mutation in the lipoprotein lipase gene. *J. Clin. Invest.* 87:1165–1170, 1991.

[66823] 5153. Auwerx, J. H.; Babirak, S. P.; Fujimoto, W. Y.; Iverius, P. H.; Brunzell, J. D.: Defective enzyme protein in lipoprotein lipase deficiency. *Europ. J. Clin. Invest.* 19: 433–437, 1989.

[66824] 5154. Beg, O. U.; Meng, M. S.; Skarlatos, S. I.; Previato, L.; Brunzell, J. D.; Brewer, H. B., Jr.; Fojo, S. S.: Lipoprotein lipase (Bethesda): a single amino acid substitution (ala176-to-thr) leads to abnormal heparin binding and loss of enzymic activity. *Proc. Nat. Acad. Sci.*

87:3474–3478, 1990.

- [66825] 5155. Benlian, P.; De Gennes, J. L.; Foubert, L.; Zhang, H.; Gagne, S.E.; Hayden, M.: Premature atherosclerosis in patients with familial chylomicronemia caused by mutations in the lipoprotein lipase gene. *New Eng. J. Med.* 335: 848–854, 1996.
- [66826] 5156. Berger, G. M. B.: An incomplete form of familial lipoprotein lipase deficiency presenting with type I hyperlipoproteinemia. *Am. J. Clin. Path.* 88: 369–373, 1987.
- [66827] 5157. Berger, H.; Richter, A.; Gilardi, A.; Wagner, H.: Essential familial hyperlipaemia in a 2-year-old child. *Ann. Paediat.* 199: 445–466, 1962.
- [66828] 5158. Bergeron, J.; Normand, T.; Bharucha, A.; Ven Murthy, M. R.; Julien, P.; Gagne, C.; Dionne, C.; De Braekeleer, M.; Brun, D.; Hayden, M.R.; Lupien, P. J.: Prevalence, geographical distribution and genealogical investigations of mutation 188 of lipoprotein lipase gene in the French-Canadian population of Quebec. *Clin. Genet.* 41: 206–210, 1992.
- [66829] 5159. Bertolini, S.; Simone, M. L.; Pes, G. M.; Ghisellini, M.; Rolleri, M.; Bellocchio, A.; Elicio, N.; Masturzo, P.; Calandra, S.: Pseudodominance of lipoprotein lipase (LPL) deficiency due to a nonsense mutation (tyr302-to-term) in exon 6 of

LPL gene in an Italian family from Sardinia(LPL–Olbia).

Clin. Genet. 57: 140–147, 2000.

- [66830] 5160.Boer, J. M. A.; Kuivenhoven, J. A.; Feskens, E. J. M.; Schouten,E. G.; Havekes, L. M.; Seidell, J. C.; Kastelein, J. J. P.; Kromhout,D.: Physical activity modulates the effect of a lipoprotein lipasemutation (D9N) on plasma lipids and lipoproteins. Clin. Genet. 56:158–163, 1999.
- [66831] 5161.Boggs, J. D.; Hsia, D. Y.–Y.; Mais, R. F.; Bigler, J. A.: Thegenetic mechanism of idiopathic hyperlipemia. New Eng. J. Med. 257:1101–1108, 1957.
- [66832] 5162.Breckenridge, W. C.; Little, A. C.; Steiner, G.; Chow, A.; Poapst,M.: Hypertriglyceridemia associated with defi–ciency of C–II apoprotein in plasma lipoproteins. New Eng. J. Med. 298: 1265–1273, 1978.
- [66833] 5163.Brunzell, J. D.; Chait, A.; Nikkila, E. A.; Ehnholm, C.; Huttunen,J. K.; Steiner, G.: Heterogeneity of primary lipoprotein lipase deficiency. Metabolism 29:624–629, 1980.
- [66834] 5164.Busca, R.; Martinez, M.; Vilella, E.; Pognonec, P.; Deeb, S.;Auwerx, J.; Reina, M.; Vilaro, S.: The mutation gly142–to–glu inhuman lipoprotein lipase produces a missorted protein that is divertedto lysosomes. J. Biol. Chem. 271: 2139–2146, 1996.

- [66835] 5165. Cantin, B.; Boudriau, S.; Bertrand, M.; Brun, L.-D.; Gagne, C.; Rogers, P. A.; Ven Murthy, M. R.; Lupien, P.-J.; Julien, P.: Hemolysis in primary lipoprotein lipase deficiency. *Metabolism* 44: 652–658, 1995.
- [66836] 5166. Chimienti, G.; Capurso, A.; Resta, F.; Pepe, G.: A G-to-C change at the donor splice site of intron 1 causes lipoprotein lipase deficiency in a Southern-Italian family. *Biochem. Biophys. Res. Commun.* 187: 620–627, 1992.
- [66837] 5167. Clark, A. G.; Weiss, K. M.; Nickerson, D. A.; Taylor, S. L.; Buchanan, A.; Stengard, J.; Salomaa, V.; Vartiainen, E.; Perola, M.; Boerwinkle, E.; Sing, C. F.: Haplotype structure and population genetic inferences from nucleotide-sequence variation in human lipoprotein lipase. *Am. J. Hum. Genet.* 63: 595–612, 1998.
- [66838] 5168. Clee, S. M.; Loubser, O.; Collins, J.; Kastelein, J. J. P.; Hayden, M. R.: The LPL S447X cSNP is associated with decreased blood pressure and plasma triglycerides, and reduced risk of coronary artery disease. *Clin. Genet.* 60: 293–300, 2001.
- [66839] 5169. De Braekeleer, M.; Dionne, C.; Gagne, C.; Julien, P.; Brun, D.; Ven Murthy, M. R.; Lupien, P.-J.: Founder effect in familial hyperchylomicronemia among French Canadians of Quebec. *Hum. Hered.* 41: 168–173, 1991.

- [66840] 5170. De Bruin, T. W. A.; Maily, F.; Van Barlingen, H. H. J. J.; Fisher, R.; Castro Cabezas, M.; Talmud, P.; Dallinga-Thie, G. M.; Humphries, S. E.: Lipoprotein lipase gene mutations D9N and N291S in four pedigrees with familial combined hyperlipidaemia. *Europ. J. Clin. Invest.* 26:631–639, 1996.
- [66841] 5171. Deeb, S. S.; Peng, R.: Structure of the human lipoprotein lipase gene. *Biochemistry* 28: 4131–4135, 1989.
- [66842] 5172. Henderson, H. E.; Hassan, F.; Berger, G. M. B.; Hayden, M. R.: The lipoprotein lipase gly188-to-glu mutation in South Africans of Indian descent: evidence suggesting common origins and an increased frequency. *J. Med. Genet.* 29: 119–122, 1992.
- [66843] 5173. Henderson, H. E.; Hassan, F.; Marais, D.; Hayden, M. R.: A new mutation destroying disulphide bridging in the C-terminal domain of lipoprotein lipase. *Biochem. Biophys. Res. Commun.* 227: 189–194, 1996.
- [66844] 5174. Henderson, H. E.; Ma, Y.; Hassan, M. F.; Monsalve, M. V.; Marais, A. D.; Winkler, F.; Gubernator, K.; Peterson, J.; Brunzell, J. D.; Hayden, M. R.: Amino acid substitution (ile194-to-thr) in exon 5 of the lipoprotein lipase gene causes lipoprotein lipase deficiency in three unrelated

probands: support for a multicentric origin. J.Clin. Invest. 87: 2005–2011, 1991.

- [66845] 5175.Sekiguchi, K.; Klos, A. M.; Kurachi, K.; Yoshitake, S.; Hakomori,S.: Human liver fibronectin complementary DNAs: identification oftwo different messenger RNAs possibly encoding the alpha and betasubunits of plasma fibronectin. Biochemistry 25: 4936–4941, 1986.
- [66846] 5176.Shirakami, A.; Shigekiyo, T.; Hirai, Y.; Takeichi, T.; Kawauchi,S.; Saito, S.; Miyoshi, K.: Plasma fibronectin deficiency in eightmembers of one family. Lancet I: 473–474, 1986.
- [66847] 5177.Shows, T. B.: Personal Communication. Buffalo, N. Y. 1982.
- [66848] 5178.Skow, L. C.; Adkison, L.; Womack, J. E.; Beamer, W. G.; Taylor,B. A.: Mapping of the mouse fibronectin gene (Fn-1) to chromosome1: conservation of the Idh-1--Cryg--Fn-1 synteny group in mammals. Genomics 1:283–286, 1987.
- [66849] 5179.Smith, M.; Gold, L. I.; Pearlstein, E.; Krinsky, A.: Expressionof mouse and human fibronectin in hybrid cells. (Abstract) Cytogenet.Cell Genet. 25: 205 only, 1979.
- [66850] 5180.Smith, M.; Krinsky, A. M.; Arredondo-Vega, F. X.; Pearlstein,E.: Production of soluble fibronectin by RAG x



human fibroblast hybrids.(Abstract) Cytogenet. Cell Genet. 32: 318 only, 1982.

- [66851] 5181.Wu, B.-L.; Milunsky, A.; Wyandt, H.; Hoth, C.; Baldwin, C.; Skare,J.: In situ hybridization applied to Waardenburg syndrome. Cytogenet.Cell Genet. 63: 29–32, 1993.
- [66852] 5182.Zardi, L.; Cianfriglia, M.; Balza, E.; Carnemolla, B.; Siri, A.;Croce, C. M.: Species-specific monoclonal antibodies in the assignmentof the gene for human fibronectin to chromosome 2. EMBO J. 1: 929–933,1982.
- [66853] 5183.Zardi, L.; Siri, A.; Carnemolla, B.; Santi, L.; Gardner, W. D.;Hoch, S. O.: Fibronectin: a chromatin-associated protein? Cell 18:649–657, 1979.
- [66854] 5184.Adkison, L. R.; White, R. A.; Haney, D. M.; Lee, J. C.; Pusey,K. T.; Gardner, J.: The fibronectin receptor, alpha subunit (Itga5)maps to murine chromosome 15, distal to D15Mit16. Mammalian Genome 5:456–457, 1994.
- [66855] 5185.Argraves, W. S.; Pytela, R.; Suzuki, S.; Millan, J. L.; Pierschbacher,M. D.; Ruoslahti, E.: cDNA sequences from the alpha subunit of thefibronectin receptor predict a transmembrane domain and a short cytoplasmicpeptide. J. Biol. Chem. 261: 12922–12924, 1986.
- [66856] 5186.Argraves, W. S.; Suzuki, S.; Arai, H.; Thompson, K.; Pierschbacher,M. D.; Ruoslahti, E.: Amino acid sequence of

the human fibronectin receptor. *J. Cell Biol.* 105: 1183–1190, 1987.

- [66857] 5187. Fitzgerald, L. A.; Poncz, M.; Steiner, B.; Rall, S. C., Jr.; Bennett, J. S.; Phillips, D. R.: Comparison of cDNA-derived protein sequences of the human fibronectin and vitronectin receptor alpha-subunits and platelet glycoprotein IIb. *Biochemistry* 26: 8158–8165, 1987.
- [66858] 5188. Krissansen, G. W.; Yuan, Q.; Jenkins, D.; Jiang, W.-M.; Rooke, L.; Spurr, N. K.; Eccles, M.; Leung, E.; Watson, J. D.: Chromosomal locations of the genes coding for the integrin beta-6 and beta-7 subunits. *Immunogenetics* 35: 58–61, 1992.
- [66859] 5189. Sosnoski, D.; Emanuel, B. S.; Hawkins, A. L.; van Tuinen, P.; Ledbetter, D. H.; Nussbaum, R. L.; Kaos, F.-T.; Schwartz, E.; Phillips, D.; Bennett, J. S.; Fitzgerald, L. A.; Poncz, M.: Chromosomal localization of the genes for the vitronectin and fibronectin receptors alpha-subunits and for platelet glycoproteins IIb and IIIa. *J. Clin. Invest.* 81: 1993–1998, 1988.
- [66860] 5190. Spurr, N. K.; Rooke, L.: Confirmation of the assignment of the vitronectin (VNRA) and fibronectin (FNRA) receptor alpha-subunits. *Ann. Hum. Genet.* 55: 217–223, 1991.

- [66861] 5191.Akula, S. M.; Pramod, N. P.; Wang, F.-Z.; Chandran, B.: Integrinalpha-3/beta-1 (CD 49c/29) is a cellular receptor for Kaposi's sarcoma-associatedherpesvirus (KSHV/HHV-8) entry into the target cells. *Cell* 108:407-419, 2002.
- [66862] 5192.Arregui, C.; Pathre, P.; Lilien, J.; Balsamo, J.: The nonreceptortyrosine kinase Fer mediates cross-talk between N-cadherin and beta-1-integrins. *J.Cell Biol.* 149: 1263-1273, 2000.
- [66863] 5193.Giuffra, L. A.; Lichter, P.; Wu, J.; Kennedy, J. L.; Pakstis, A.J.; Rogers, J.; Kidd, J. R.; Harley, H.; Jenkins, T.; Ward, D. C.;Kidd, K. K.: Genetic and physical mapping and population studiesof a fibronectin receptor beta-subunit-like sequence on human chromosome19. *Genomics* 8: 340-346, 1990.
- [66864] 5194.Giuffra, L. A.; Wu, J.; Lichter, P.; Kennedy, J. L.; Castiglione,C.; Pakstis, A. J.; Ward, D.; Kidd, K. K.: Mapping of a fibronectinreceptor beta subunit-like sequence to chromosome 19. (Abstract) *Am.J. Hum. Genet.* 45 (suppl.): A141 only, 1989.
- [66865] 5195.Goodfellow, P. J.; Nevanlinna, H. A.; Gorman, P.; Sheer, D.; Lam,G.; Goodfellow, P. N.: Assignment of the gene encoding the beta-subunitof the human fibronectin

receptor (beta-FNR) to chromosome 10p11.2. *Ann.Hum. Genet.* 53: 15–22, 1989.

- [66866] 5196.Graus-Porta, D.; Blaess, S.; Senften, M.; Littlewood-Evans, A.;Damsky, C.; Huang, Z.; Orban, P.; Klein, R.; Schittny, J. C.; Muller,U.: Beta-1-class integrins regulate the development of laminae andfolia in the cerebral and cerebellar cortex. *Neuron* 31: 367–379,2001.
- [66867] 5197.Hynes, R. O.: Integrins: a family of cell surface receptors. *Cell* 48:549–554, 1987.
- [66868] 5198.Johansson, S.; Forsberg, E.; Lundgren, B.: Comparison of fibronectinreceptors from rat hepatocytes and fibroblasts. *J. Biol. Chem.* 262:7819–7824, 1987.
- [66869] 5199.Lu, T. T.; Cyster, J. G.: Integrin-mediated long-term B cell retentionin the splenic marginal zone. *Science* 297: 409–412, 2002.
- [66870] 5200.Messer Peters, P.; Kamarck, M. E.; Hemler, M. E.; Strominger,J. L.; Ruddle, F. H.: Genetic and biochemical characterization ofhuman lymphocyte cell surface antigens: the A-1A5 and A-3A4 determinants. *J.Exp. Med.* 159: 1441–1454, 1984.
- [66871] 5201.Pytela, R.; Pierschbacher, M. D.; Ginsberg, M. H.; Plow, E. F.;Ruoslahti, E.: Platelet membrane glycoprotein IIb/IIIa: member ofa family of arg-gly-asp-specific adhe-

sion receptors. Science 231:1159–1162, 1986.

- [66872] 5202.Woods, V. L., Jr.; Pischel, K. D.; Avery, E. D.; Bluestein, H.G.: Antigenic polymorphism of human very late activation protein-2(platelet glycoprotein Ia-IIa): platelet alloantigen Hc(a). J. Clin.Invest. 83: 978–985, 1989.
- [66873] 5203.Wu, J. S.; Giuffra, L. A.; Goodfellow, P. J.; Myers, S.; Carson, N. L.; Anderson, L.; Hoyle, L. S.; Simpson, N. E.; Kidd, K. K.: The beta subunit locus of the human fibronectin receptor: DNA restriction fragment length polymorphism and linkage mapping studies. Hum. Genet. 83:383–390, 1989.
- [66874] 5204.Wilcox, A. S.; Warrington, J. A.; Gardiner, K.; Berger, R.; Whiting, P.; Altherr, M. R.; Wasmuth, J. J.; Patterson, D.; Sikela, J. M.: Human chromosomal localization of genes encoding the gamma-1 and gamma-2 subunits of the gamma-aminobutyric acid receptor indicates members of this gene family are often clustered in the genome. Proc. Nat. Acad. Sci. 89: 5857–5861, 1992.
- [66875] 5205.Huang, F.; Shi, L. J.; Heng, H. H. Q.; Fei, J.; Guo, L.-H.: Assignment of the human GABA transporter gene (GABATHG) locus to chromosome 3p24–p25. Genomics 29:302–304, 1995.

- [66876] 5206.Lam, D. M.-K.; Fei, J.; Zhang, X.-Y.; Tam, A. C. W.; Zhu, L.-H.;Huang, F.; King, S. C.; Guo, L.-H.: Molecular cloning and structureof the human (GABATHG) GABA transporter gene. *Molec. Brain Res.* 19:227–232, 1993.
- [66877] 5207.Nelson, H.; Mandiyan, S.; Nelson, N.: Cloning of the human brainGABA transporter. *FEBS Lett.* 269: 181–184, 1990.
- [66878] 5208.Figlewicz, D. A.; Delattre, O.; Guellaen, G.; Krizus, A.; Thomas,G.; Zucman, J.; Rouleau, G. A.: Mapping of human gamma–glutamyl transpeptidasegenes on chromosome 22 and other human autosomes. *Genomics* 17: 299–305,1993.
- [66879] 5209.Pawlak, A.; Wu, S.-J.; Bulle, F.; Suzuki, A.; Chikhi, N.; Ferry,N.; Baik, J.-H.; Siegrist, S.; Guellaen, G.: Different gamma–glutamyltranspeptidase mRNAs are expressed in human liver and kidney. *Biochem.Biophys. Res. Commun.* 164: 912–918, 1989.
- [66880] 5210.Rajpert–De Meyts, E.; Heisterkamp, N.; Groffen, J.: Cloning andnucleotide sequence of human gamma–glutamyl transpeptidase. *Proc.Nat. Acad. Sci.* 85: 8840–8844, 1988.
- [66881] 5211.Lopez de Castro, J. A.; Strominger, J. L.; Strong, D. M.; Orr,H. T.: Structure of crossreactive human histocom–

patibility antigens HLA-A28 and HLA-A2: possible implications for the generation of HLA polymorphism. *Proc. Nat. Acad. Sci.* 79: 3813–3817, 1982.

[66882] 5212. Malissen, M.; Malissen, B.; Jordan, B. R.: Exon/intron organization and complete nucleotide sequence of an HLA gene. *Proc. Nat. Acad. Sci.* 79: 893–897, 1982.

[66883] 5213. Mann, D. L.; Rogentine, G. N., Jr.; Fahey, J. L.; Nathenson, S. G.: Molecular heterogeneity of human lymphoid (HL-A) alloantigens. *Science* 163:1460–1462, 1969.

[66884] 5214. MacDonald, K. S.; Fowke, K. R.; Kimani, J.; Dunand, V. A.; Nagelkerke, N. J. D.; Ball, T. B.; Oyugi, J.; Njagi, E.; Gaur, L. K.; Brunham, R. C.; Wade, J.; Luscher, M. A.; Krausa, P.; Rowland-Jones, S.; Ngugi, E.; Bwayo, J. J.; Plummer, F. A.: Influence of HLA supertypes on susceptibility and resistance to human immunodeficiency virus type 1 infection. *J. Infect. Dis.* 181: 1581–1589, 2000.

[66885] 5215. Mayr, W. R.; Mayr, D.: Analysis of the linkage between the HL-A loci and the genes of other markers. *Humangenetik* 24: 129–133, 1974.

[66886] 5216. Messer, G.; Zemmour, J.; Orr, H. T.; Parham, P.; Weiss, E. H.; Girdlestone, J.: HLA-J, a second inactivated class I HLA gene related to HLA-G and HLA-A. *J. Immun.* 148: 4043–4053, 1992.

- [66887] 5217.Morton, C. C.; Kirsch, I. R.; Nance, W. E.; Evans, G. A.; Korman,A. J.; Strominger, J. L.: Orientation of loci within the human majorhistocompatibility complex by chromosomal in situ hybridization. Proc.Nat. Acad. Sci. 81: 2816–2820, 1984.
- [66888] 5218.Morton, N. E.; Rao, D. C.; Lindsten, J.; Hulten, M.; Yee, S.:A chiasma map of man. Hum. Hered. 27: 38–51, 1977.
- [66889] 5219.Mulley, J. C.; Hay, J.; Sheffield, L. J.; Sutherland, G. R.:Regional localization for HLA by recombination with a fragile siteat 6p23. Am. J. Hum. Genet. 35: 1284–1288, 1983.
- [66890] 5220.Newell, W. R.; Trowsdale, J.; Beck, S.: MHCDB–database of thehuman MHC. Immunogenetics 40: 109–115, 1994.
- [66891] 5221.Orr, H. T.; Bach, F. H.; Ploegh, H. L.; Strominger, J. L.; Kavathas,P.; DeMars, R.: Use of HLA loss mutants to analyse the structureof the human major histocompatibil–ity complex. Nature 296: 454–456,1982.
- [66892] 5222.Paabo, S.: Molecular cloning of ancient Egyptian mummy DNA. Nature 314:644–645, 1985.
- [66893] 5223.Patel, R.; Mickey, M. R.; Terasaki, P. I.: Serotyping for homotransplantationof kidneys from unrelated donors.



New Eng. J. Med. 279: 501–506,1968.

- [66894] 5224. Payne, R.; Tripp, M.; Weigle, J.; Bodmer, W. F.; Bodmer, J. G.: A new leukocyte isoantigen system in man. Cold Spring Harbor Symp. Quant. Biol. 29: 285–295, 1964.
- [66895] 5225. Polacek, L. A.; Phillips, R. B.; Hackbarth, S. A.; Duquesnoy, R. J.: A linkage study of the HLA region using C-band heteromorphisms. Clin. Genet. 23: 177–185, 1983.
- [66896] 5226. Ragoussis, J.; Bloemer, K.; Pohla, H.; Messer, G.; Weiss, E. H.; Ziegler, A.: A physical map including a new class I gene (cda12) of the human major histocompatibility complex (A2/B13 haplotype) derived from a monosomy 6 mutant cell line. Genomics 4: 301–308, 1989.
- [66897] 5227. Ragoussis, J.; van der Bliek, A.; Trowsdale, J.; Ziegler, A.: Mapping of HLA genes using pulsed field gradient electrophoreses. FEBS Lett. 204: 1–4, 1986.
- [66898] 5228. Salter, R. D.; Norment, A. M.; Chen, B. P.; Clayberger, C.; Krensky, A. M.; Littman, D. R.; Parham, P.: Polymorphism in the alpha(3) domain of HLA-A molecules affects binding to CD8. Nature 338: 345–347, 1989.
- [66899] 5229. Schunter, F.; Wernet, P.; Kompf, J.; Bissbort, S.; Gohler, F.: Mapping of the linkage group GLO-Bf-HLA-B, C, A-PGM. II. Segregation analysis. Hum. Genet. 44:

321–331, 1978.

- [66900] 5230. Shivdasani, R. A.; Haluska, F. G.; Dock, N. L.; Dover, J. S.; Kineke, E. J.; Anderson, K. C.: Graft-versus-host disease associated with transfusion of blood from unrelated HLA-homozygous donors. *New Eng. J. Med.* 328: 766–770, 1993.
- [66901] 5231. Shukla, H.; Gillespie, G. A.; Srivastava, R.; Collins, F.; Chorney, M. J.: A class I jumping clone places the HLA-G gene approximately 100 kilobases from HLA-H within the HLA-A subregion of the human MHC. *Genomics* 10:905–914, 1991.
- [66902] 5232. Snell, G. D.: Studies in histocompatibility. *Science* 213: 172–178, 1981.
- [66903] 5233. Solheim, B. G.; Bratlie, A.; Sandberg, L.; Staub-Nielsen, L.; Thorsby, E.: Further evidence of a third HLA-A locus. *Tissue Antigens* 3:439–453, 1973.
- [66904] 5234. Strachan, T.: Molecular genetics and polymorphism of class I HLA antigens. *Brit. Med. Bull.* 43: 1–14, 1987.
- [66905] 5235. Szpak, Y.; Vieville, J.-C.; Tabary, T.; Naud, M.-C.; Chopin, M.; Edelson, C.; Cohen, J. H. M.; Dausset, J.; de Kozak, Y.; Pla, M.: Spontaneous retinopathy in HLA-A29 transgenic mice. *Proc. Nat. Acad. Sci.* 98: 2572–2576, 2001.

- [66906] 5236.Teshima, T.; Ordemann, R.; Reddy, P.; Gagin, S.; Liu, C.; Cooke,K. R.; Ferrara, J. L. M.: Acute graft-versus-host disease does not require alloantigen expression on host epithelium. *Nature Med.* 8:575–581, 2002.
- [66907] 5237.Thorsby, E.; Sandberg, L.; Lindholm, A.; Kissmeyer-Nielsen, F.: The HL-A system: evidence of a third sub-locus. *Scand. J. Haemat.* 7:195–200, 1970.
- [66908] 5238.Tragardh, L.; Rask, L.; Wiman, K.; Fohlman, J.; Peterson, P. A.: Amino acid sequence of an immunoglobulin HLA antigen heavy chain domain. *Proc. Nat. Acad. Sci.* 76: 5839–5842, 1979.
- [66909] 5239.Trowsdale, J.: Personal Communication. London, England 1/12/1983.
- [66910] 5240.Trowsdale, J.; Ragoussis, J.; Campbell, R. D.: Map of the human MHC. *Immun. Today* 12: 443–446, 1991.
- [66911] 5241.Van Leeuwen, A.; Eernisse, J. G.; Van Rood, J. J.: A new leucocyte group with two alleles: leucocyte group five. *Vox Sang.* 9: 431–446, 1964.
- [66912] 5242.Van Rood, J. J.: Leucocyte grouping and organ transplantation. *Brit.J. Haemat.* 16: 211–220, 1969.100. Van Rood, J. J.: Tissue typing and organ transplantation. *Lancet* I:1142–1146, 1969.101. Van Rood, J. J.; Amos, D. B.: In memoriam: Ruggero Ceppellini, 1917–1988. *Hum.*

Immun. 23: 1–3, 1988.102. Van Rood, J. J.; Van Leeuwen, A.: Leukocyte grouping: a method and its application. J. Clin. Invest. 42: 1382–1390, 1963.103. Van Someren, H.; Westerveld, A.; Hagemeyer, A.; Mees, J. R.; Meera Khan, P.; Zaalberg, O. B.: Human antigen and enzyme markers in man–Chinese hamster somatic cell hybrids: evidence for synteny between the HL–A, PGM–3, ME–1, and IPO–B loci. Proc. Nat. Acad. Sci. 71:962–965, 1974.104. Walford, R. L.; Finkelstein, S.; Hanna, C.; Collins, Z.: Third sublocus in the HL–A human transplantation system. Nature 224: 74–75, 1969.105. Weitkamp, L. R.; Van Rood, J. J.; Thorsby, E.; Bias, W. B.; Fotino, M.; Lawler, S. D.; Dausset, J.; Mayr, W. R.; Bodmer, J.; Ward, F. S.; Signalet, J.; Payne, R.; Kissmeyer–Nielsen, F.; Gatti, R. A.; Sachs, J. A.; Lamm, L. U.: The relation of parental sex and age to recombination in the HL–A system. Hum. Hered. 23: 197–205, 1973.106. Wolski, K. P.; Schmid, F. R.; Mittal, K. K.: Genetic linkage between the HL–A system and a deficit of the second component (C2) of complement. Science 188: 1020–1022, 1975.107. Ziegler, A.; Ragoussis, J.; Fonatsch, C.; Weiss, E.: A physical map of the human major histocompatibility (HLA) complex derived from a monosomy 6 mutant cell line. (Abstract) Cytogenet. Cell Genet.

51:1116 only, 1989.

- [66913] 5243. Wildhage, I.; Trusheim, H.; Goke, B.; Lankat-Buttgereit, B.: Geneexpression of the human glucagon-like peptide-1 receptor is regulated by Sp1 and Sp3. *Endocrinology* 140: 624-631, 1999.
- [66914] 5244. Chambers, S. M.; Morris, B. J.: Glucagon receptor gene mutation in essential hypertension. (Letter) *Nature Genet.* 12: 122, 1996.
- [66915] 5245. Hager, J.; Hansen, L.; Vaisse, C.; Vionnet, N.; Philippi, A.; Poller, W.; Velho, G.; Carcassi, C.; Contu, L.; Julier, C.; Cambien, F.; Passa, P.; Lathrop, M.; Kindsvogel, W.; Demenais, F.; Nishimura, E.; Froguel, P.: A missense mutation in the glucagon receptor gene is associated with non-insulin-dependent diabetes mellitus. *Nature Genet.* 9: 299-304, 1995.
- [66916] 5246. Lok, S.; Kuijper, J. L.; Jelinek, L. J.; Kramer, J. M.; Whitmore, T. E.; Sprecher, C. A.; Mathewes, S.; Grant, F. J.; Biggs, S. H.; Rosenberg, G. B.; Sheppard, P. O.; O'Hara, P. J.; Foster, D. C.; Kindsvogel, W.: The human glucagon receptor encoding gene: structure, cDNA sequence and chromosomal localization. *Gene* 140: 203-209, 1994.
- [66917] 5247. Menzel, S.; Stoffel, M.; Espinosa, R., III; Fernald, A. A.; LeBeau, M. M.; Bell, G. I.: Localization of the glucagon

receptor gene to human chromosome band 17q25. *Genomics* 20: 327–328, 1994.

[66918] 5248. Weitkamp, L. R.: Concerning the linkage relationships of the Gc and MNSs loci. *Hum. Genet.* 43: 215–220, 1978.

[66919] 5249. Boguszewski, C. L.; Svensson, P.-A.; Jansson, T.; Clark, R.; Carlsson, L. M. S.; Carlsson, B.: Cloning of two novel growth hormone transcripts expressed in human placenta. *J. Clin. Endocr. Metab.* 83: 2878–2885, 1998.

[66920] 5250. Chen, E. Y.; Liao, Y.-C.; Smith, D. H.; Barrera-Saldana, H. A.; Gelinas, R. E.; Seeburg, P. H.: The human growth hormone locus: nucleotide sequence, biology, and evolution. *Genomics* 4: 479–497, 1989.

[66921] 5251. Frankenne, F.; Rentier-Delrue, F.; Scippo, M.-L.; Martial, J.; Hennen, G.: Expression of the growth hormone variant gene in human placenta. *J. Clin. Endocr. Metab.* 64: 635–637, 1987.

[66922] 5252. Lewis, U. J.; Dunn, J. T.; Bonewald, L. F.; Seavey, B. K.; VanderLaan, W. P.: A naturally occurring structural variant of human growth hormone. *J. Biol. Chem.* 253: 2679–2687, 1978.

[66923] 5253. Liebhaver, S. A.; Urbanek, M.; Ray, J.; Tuan, R. S.; Cooke, N. E.: Characterization and histologic localization

of human growthhormone-variant gene expression in the placenta. *J. Clin. Invest.* 83:1985-1991, 1989.

[66924] 5254. MacLeod, J. N.; Liebhaber, S. A.; MacGillivray, M. H.; Cooke, N.E.: Identification of a splice-site mutation in the human growthhormone-variant gene. *Am. J. Hum. Genet.* 48: 1168-1174, 1991.

[66925] 5255. MacLeod, J. N.; Worsley, I.; Ray, J.; Friesen, H. G.; Liebhaber, S. A.; Cooke, N. E.: Human growth hormone variant is a biologically active somatogen and lactogen. *Endocrinology* 128: 1298-1302, 1991.

[66926] 5256. Owerbach, D.; Rutter, W. J.; Martial, J. A.; Baxter, J. D.; Shows, T. B.: Genes for growth hormone, chorionic somatomammotropin and growth hormone-like genes on chromosome 17 in humans. *Science* 209:289-292, 1980.

[66927] 5257. Harada, H.; Fujita, T.; Miyamoto, M.; Kimura, Y.; Maruyama, M.; Furia, A.; Miyata, T.; Taniguchi, T.: Structurally similar but functionally distinct factors, IRF-1 and IRF-2, bind to the same regulatory elements of IFN and IFN-inducible genes. *Cell* 58: 729-739, 1989.

[66928] 5258. Wong, P.; Colucci-Guyon, E.; Takahashi, K.; Gu, C.; Babinet, C.; Coulombe, P. A.: Introducing a null mutation in the mouse K6-alpha and K6-beta genes reveals their essential structural role in the oral mucosa. *J. Cell Biol.* 150:

921–928, 2000.

- [66929] 5259. Nakashima, K.; Yanagisawa, M.; Arakawa, H.; Kimura, N.; Hisatsune, T.; Kawabata, M.; Miyazono, K.; Taga, T.: Synergistic signaling in fetal brain by STAT3–Smad1 complex bridged by p300. *Science* 284:479–482, 1999.
- [66930] 5260. Bae, J.; Leo, C. P.; Hsu, S. Y.; Hsueh, A. J. W.: MCL-1S, a splicing variant of the antiapoptotic BCL-2 family member MCL-1, encodes a proapoptotic protein possessing only the BH3 domain. *J. Biol. Chem.* 275:25255–25261, 2000.
- [66931] 5261. Craig, R. W.; Jabs, E. W.; Zhou, P.; Kozopas, K. M.; Hawkins, A. L.; Rochelle, J. M.; Seldin, M. F.; Griffin, C. A.: Human and mouse chromosomal mapping of the myeloid cell leukemia-1 gene: MCL1 maps to human chromosome 1q21, a region that is frequently altered in preneoplastic and neoplastic disease. *Genomics* 23: 457–463, 1994.
- [66932] 5262. Kozopas, K. M.; Yang, T.; Buchan, H. L.; Zhou, P.; Craig, R. W.: MCL1, a gene expressed in programmed myeloid cell differentiation, has sequence similarity to BCL2. *Proc. Nat. Acad. Sci.* 90: 3516–3520, 1993.
- [66933] 5263. Rinkenberger, J. L.; Horning, S.; Klocke, B.; Roth, K.; Korsmeyer, S. J.: Mcl-1 deficiency results in peri-



implantation embryonic lethality. *GenesDev.* 14: 23–27, 2000.

[66934] 5264.Hizawa, N.; Yamaguchi, E.; Furuya, K.; Ohnuma, N.; Kodama, N.;Kojima, J.; Ohe, M.; Kawakami, Y.: Association between high serumtotal IgE levels and D11S97 on chromosome 11q13 in Japanese subjects. *J.Med. Genet.* 32: 363–369, 1995.

[66935] 5265.Sandford, A. J.; Shirakawa, T.; Moffatt, M. F.; Daniels, S. E.;Ra, C.; Faux, J. A.; Young, R. P.; Nakamura, Y.; Lathrop, G. M.; Cookson,W. O. C. M.; Hopkin, J. M.: Localisation of atopy and beta subunitof high-affinity IgE receptor (FCER1) on chromosome 11q. *Lancet* 341:332–334, 1993.

[66936] 5266.Harada, H.; Kondo, T.; Ogawa, S.; Tamura, T.; Kitagawa, M.; Tanaka,N.; Lamphier, M. S.; Hirai, H.; Taniguchi, T.: Accelerated exon skippingof IRF–1 mRNA in human myelodysplasia/leukemia: a possible mechanismof tumor suppressor inactivation *Oncogene* 9: 3313–3320, 1994.

[66937] 5267.Harada, H.; Willison, K.; Sakakibara, J.; Miyamoto, M.; Fujita,T.; Taniguchi, T.: Absence of the type I IFN system in EC cells:transcriptional activator (IRF–1) and repressor (IRF–2) genes aredevelopmentally regulated. *Cell* 63: 303–312, 1990.

[66938] 5268.Itoh, S.; Harada, H.; Nakamura, Y.; White, R.;

Taniguchi, T.: Assignment of the human interferon regulatory factor-1 (IRF1) gene to chromosome 5q23-q31. *Genomics* 10: 1097-1099, 1991.

[66939] 5269. Ko, J.; Gendron-Fitzpatrick, A.; Splitter, G. A.: Susceptibility of IFN regulatory factor-1 and IFN consensus sequence binding protein-deficient mice to brucellosis. *J. Immun.* 168: 2433-2440, 2002.

[66940] 5270. Miyamoto, M.; Fujita, T.; Kimura, Y.; Maruyama, M.; Harada, H.; Sudo, Y.; Miyata, T.; Taniguchi, T.: Regulated expression of a gene encoding a nuclear factor, IRF-1, that specifically binds to IFN-beta gene regulatory elements. *Cell* 54: 903-913, 1988.

[66941] 5271. Nozawa, H.; Oda, E.; Ueda, S.; Tamura, G.; Maesawa, C.; Muto, T.; Taniguchi, T.; Tanaka, N.: Functionally inactivating point mutation in the tumor-suppressor IRF-1 gene identified in human gastric cancer. *Int. J. Cancer* 77: 522-527, 1998.

[66942] 5272. Tamura, G.; Sakata, K.; Nishizuka, S.; Maesawa, C.; Suzuki, Y.; Terashima, M.; Eda, Y.; Satodate, R.: Allelotype of adenoma and differentiated adenocarcinoma of the stomach. *J. Path.* 180: 371-377, 1996.

[66943] 5273. Willman, C. L.; Sever, C. E.; Pallavicini, M. G.; Harada, H.; Tanaka, N.; Slovak, M. L.; Yamamoto, H.; Harada, K.;

Meeker, T. C.;List, A. F.; Taniguchi, T.: Deletion of IRF-1, mapping to chromosome5q31.1, in human leukemia and preleukemic myelodysplasia. *Science* 259:968-971, 1993.

[66944] 5274.Yamada, G.; Ogawa, M.; Akagi, K.; Miyamoto, H.; Nakano, N.; Itoh,S.; Miyazaki, J.; Nishikawa, S.; Yamamura, K.; Taniguchi, T.: Specificdepletion of the B-cell population induced by aberrant expressionof human interferon regulatory factor 1 gene in transgenic mice. *Proc.Nat. Acad. Sci.* 88: 532-536, 1991.

[66945] 5275.Harada, H.; Kitagawa, M.; Tanaka, N.; Yamamoto, H.; Harada, K.;Ishihara, M.; Taniguchi, T.: Anti-oncogenic and oncogenic potentialsof interferon regulatory factors-1 and -2. *Science* 259: 971-974,1993.

[66946] 5276.Harada, H.; Takahashi, E.-I.; Itoh, S.; Harada, K.; Hori, T.-A.;Taniguchi, T.: Structure and regulation of the human interferon regulatoryfactor 1 (IRF-1) and IRF-2 genes: implications for a gene networkin the interferon system. *Molec. Cell. Biol.* 14: 1500-1509, 1994.

[66947] 5277.Hida, S.; Ogasawara, K.; Sato, K.; Abe, M.; Takayanagi, H.; Yokochi,T.; Sato, T.; Hirose, S.; Shirai, T.; Taki, S.; Taniguchi, T.: CD8+T cell-mediated skin disease in mice lacking IRF-2, the transcriptionalattenuator of interferon-alpha/beta signaling. *Immunity* 13:

643–655,2000.

- [66948] 5278.Nishio, Y.; Noguchi, E.; Ito, S.; Ichikawa, E.; Umebayashi, Y.;Otsuka, F.; Arinami, T.: Mutation and association analysis of theinterferon regulatory factor 2 gene (IRF2) with atopic dermatitis. *J.Hum. Genet.* 46: 664–667, 2001.
- [66949] 5279.Guo, B.; Yu, Y.; Leibold, E. A.: Iron regulates cytoplasmic levelsof a novel iron-responsive element-binding protein without aconitaseactivity. *J. Biol. Chem.* 269: 24252–24260, 1994.
- [66950] 5280.Henderson, B. R.; Seiser, C.; Kuhn, L. C.: Characterization ofa second RNA-binding protein in rodents with specificity for iron-responsiveelements. *J. Biol. Chem.* 268: 27327–27334, 1993.
- [66951] 5281.Hentze, M. W.; Kuhn, L. C.: Molecular control of vertebrate ironmetabolism: mRNA-based regulatory circuits operated by iron, nitricoxide, and oxidative stress. *Proc. Nat. Acad. Sci.* 93: 8175–8182,1996.
- [66952] 5282.Iwai, K.; Klausner, R. D.; Rouault, T. A.: Requirements for iron-regulateddegradation of the RNA binding protein, iron regulatory protein 2. *EMBOJ.* 14: 5350–5357, 1995.
- [66953] 5283.Rouault, T. A.; Tang, C. K.; Kaptain, S.; Burgess, W.

H.; Haile, D. J.; Samaniego, F.; McBride, O. W.; Harford, J. B.; Klausner, R.D.: Cloning of the cDNA encoding an RNA regulatory protein: the human iron-responsive element-binding protein. *Proc. Nat. Acad. Sci.* 87:7958–7962, 1990.

[66954] 5284. LaVaute, T.; Smith, S.; Cooperman, S.; Iwai, K.; Land, W.; Meyron-Holtz, E.; Drake, S. K.; Miller, G.; Abu-Asab, M.; Tsokos, M.; Switzer, R., III; Grinberg, A.; Love, P.; Tresser, N.; Rouault, T. A.: Targeted deletion of the gene encoding iron regulatory protein-2 causes misregulation of iron metabolism and neurodegenerative disease in mice. *Nature Genet.* 27: 209–214, 2001.

[66955] 5285. Samaniego, F.; Chin, J.; Iwai, K.; Rouault, T. A.; Klausner, R.D.: Molecular characterization of a second iron-responsive element binding protein, iron regulatory protein 2: structure, function, and post-translational regulation. *J. Biol. Chem.* 269: 30904–30910, 1994.

[66956] 5286. Smith, F. J. D.; Jonkman, M. F.; van Goor, H.; Coleman, C. M.; Covello, S. P.; Uitto, J.; McLean, W. H. I.: A mutation in human keratin K6b produces a phenocopy of the K17 disorder pachyonychia congenita type 2. *Hum. Molec. Genet.* 7: 1143–1148, 1998.

[66957] 5287. Moll, R.; Franke, W. W.; Schiller, D. L.; Geiger, B.;

Krepler,R.: The catalog of human cytokeratins: patterns of expression innormal epithelia, tumors and cultured cells. Cell 31: 11–24, 1982.

[66958] 5288.Casanova, M. L.; Bravo, A.; Ramirez, A.; Morreale de Escobar, G.;Were, F.; Merlino, G.; Vidal, M.; Jorcano, J. L.: Exocrine pancreaticdisorders in transsgenic (sic) mice expressing human keratin 8. J.Clin. Invest. 103: 1587–1595, 1999.

[66959] 5289.Jackson, B. W.; Grund, C.; Schmid, E.; Burke, K.; Franke, W.; Illmensee,K.: Formation of cytoskeletal elements during mouse embryogenesis:intermediate filaments of the cytokeratin type and desmosomes in preimplantationembryos. Differentiation 17: 161–179, 1980.

[66960] 5290.Ku, N.–O.; Gish, R.; Wright, T. L.; Omary, M. B.: Keratin 8 mutationsin patients with cryptogenic liver disease. New Eng. J. Med. 344:1580–1587, 2001.

[66961] 5291.Ku, N.–O.; Wright, T. L.; Terrault, N. A.; Gish, R.; Omary, M.B.: Mutation of human keratin 18 in association with cryptogeniccirrhosis. J. Clin. Invest. 99: 19–23, 1997.

[66962] 5292.Waseem, A.; Alexander, C. M.; Steel, J. B.; Lane, E. B.: Embryonicsimple epithelial keratins 8 and 18: chromosomal location emphasizesdifference from other keratin pairs. New Biologist 2: 464–478, 1990.

- [66963] 5293.Yamamoto, R.; Kao, L.-C.; McKnight, C. E.; Strauss, J. F., III: Cloning and sequence of cDNA for human placental cytokeratin 8:regulation of the mRNA in trophoblastic cells by cAMP. *Molec. Endocr.* 4:370–374, 1990.
- [66964] 5294.Fan, Q. R.; Long, E. O.; Wiley, D. C.: Crystal structure of the human natural killer cell inhibitory receptor KIR2DL1–HLA–Cw4 complex. *Nature Immun.* 2: 452–460, 2001.
- [66965] 5295.Yen, J.-H.; Moore, B. E.; Nakajima, T.; Scholl, D.; Schaid, D.J.; Weyand, C. M.; Goronzy, J. J.: Major histocompatibility complex class I–recognizing receptors are disease risk genes in rheumatoid arthritis. *J. Exp. Med.* 193: 1159–1167, 2001.
- [66966] 5296.Kogan, S. C.; Lagasse, E.; Atwater, S.; Bae, S.; Weissman, I.; Ito, Y.; Bishop, J. M.: The PEBP2–beta–MYH11 fusion created by inv(16)(p13;q22) in myeloid leukemia impairs neutrophil maturation and contributes to granulocytic dysplasia. *Proc. Nat. Acad. Sci.* 95: 11863–11868, 1998.
- [66967] 5297.Liu, P.; Tarle, S. A.; Hajra, A.; Claxton, D. F.; Marlton, P.; Freedman, M.; Siciliano, M. J.; Collins, F. S.: Fusion between transcription factor CBF–beta/PEBP2–beta and a myosin heavy chain in acute myeloid leukemia. *Science*

261: 1041–1044, 1993.

- [66968] 5298.Liu, P. P.; Hajra, A.; Wijmenga, C.; Collins, F. S.: Molecularpathogenesis of the chromosome 16 inversion in the M4Eo subtype ofacute myeloid leukemia. *Blood* 85: 2289–2302, 1995.
- [66969] 5299.Lutterbach, B.; Hou, Y.; Durst, K. L.; Hiebert, S. W.: The inv(16)encodes an acute myeloid leukemia 1 transcriptional corepressor. *Proc.Nat. Acad. Sci.* 96: 12822–12827, 1999.
- [66970] 5300.O'Reilly, J.; Chipper, L.; Springall, F.; Herrmann, R.: A uniquestructural abnormality of chromosome 16 resulting in a CBF–beta–MYH11fusion transcript in a patient with acute myeloid leukemia, FAB M4. *CancerGenet. Cytogenet.* 121: 52–55, 2000.
- [66971] 5301.Ogawa, E.; Inuzuka, M.; Maruyama, M.; Satake, M.; Naito–Fujimoto,M.; Ito, Y.; Shigesada, K.: Molecular cloning and characterizationof PEBP2–beta, the heterodimeric partner of a novel *Drosophila* runt–relatedDNA binding protein PEBP2–alpha. *Virology* 194: 314–331, 1993.
- [66972] 5302.Wang, S.; Wang, Q.; Crute, B. E.; Melnikova, I. N.; Keller, S.R.; Speck, N. A.: Cloning and characterization of subunits of theT–cell receptor and murine leukemia virus



enhancer core-binding factor. *Molec.Cell. Biol.* 13: 3324–3339, 1993.

- [66973] 5303. Pellegata, N. S.; Dieguez-Lucena, J. L.; Joensuu, T.; Lau, S.; Montgomery, K. T.; Krahe, R.; Kivela, T.; Kucherlapati, R.; Forsius, H.; de la Chapelle, A.: Mutations in KERA, encoding keratocan, cause cornea plana. *Nature Genet.* 25: 91–95, 2000.
- [66974] 5304. Fabrizi, G. M.; Rizzuto, R.; Nakase, H.; Mita, S.; Lomax, M. I.; Grossman, L. I.; Schon, E. A.: Sequence of a cDNA specifying subunit VIIa of human cytochrome c oxidase. *Nucleic Acids Res.* 17: 7107 only, 1989.
- [66975] 5305. Wolz, W.; Kress, W.; Mueller, C. R.: Genomic sequence structure and organization of the human gene for cytochrome c oxidase subunit (COX7A1) VIIa-M. *Genomics* 45: 438–442, 1997.
- [66976] 5306. Brooks, B. A.; McBride, O. W.; Dolphin, C. T.; Farrall, M.; Scambler, P. J.; Gonzalez, F. J.; Idle, J. R.: The gene CYP3 encoding P450PCN1 (nifedipine oxidase) is tightly linked to the gene COL1A2 encoding collagen type 1 alpha on 7q21–q22.1. *Am. J. Hum. Genet.* 43: 280–284, 1988.
- [66977] 5307. Chen, H.; Sandler, D. P.; Taylor, J. A.; Shore, D. L.; Liu, E.; Bloomfield, C. D.; Bell, D. A.: Increased risk for myelodysplastic syndromes in individuals with glutathione

transferase theta 1 (GSTT1) gene defect. Lancet 347: 295–297, 1996.

- [66978] 5308. Daly, A. K.; Salh, B. S.; Bilton, D.; Allen, J.; Knight, A. D.; Webb, A. K.; Braganza, J. M.; Idle, J. R.: Deficient nifedipine oxidation: a rare inherited trait associated with cystic fibrosis kindreds. Pharmacogenetics 2:19–24, 1992.
- [66979] 5309. Elshourbagy, N. A.; Guzelian, P. S.: Separation, purification, and characterization of a novel form of hepatic cytochrome P-450 from rats treated with pregnenolone-16- $\alpha$ -carbonitrile. J. Biol. Chem. 255:1279–1285, 1980.
- [66980] 5310. Felix, C. A.; Walker, A. H.; Lange, B. J.; Williams, T. M.; Winick, N. J.; Cheung, N.-K. V.; Lovett, B. D.; Nowell, P. C.; Blair, I. A.; Rebbeck, T. R.: Association of CYP3A4 genotype with treatment-related leukemia. Proc. Nat. Acad. Sci. 95: 13176–13181, 1998.
- [66981] 5311. Forrester, L. M.; Neal, G. E.; Judah, D. J.; Glancey, M. J.; Wolf, C. R.: Evidence for involvement of multiple forms of cytochrome P-450 in aflatoxin B(1) metabolism in human liver. Proc. Nat. Acad. Sci. 87:8306–8310, 1990.
- [66982] 5312. Gonzalez, F. J.; Schmid, B. J.; Umeno, M.; McBride, O. W.; Hardwick, J. P.; Meyer, U. A.; Gelboin, H. V.; Idle, J. R.: Human P450PCN1: sequence, chromosome localization,

and direct evidence through cDNA expression that P450PCN1 is nifedipine oxidase. DNA 7: 79–86, 1988.

- [66983] 5313. Hoyo–Vadillo, C.; Castaneda–Hernandez, G.; Herrera, J. E.; Vidal–Garate, J.; Moreno–Ramos, A.; Chavez, F.; Hong, E.: Pharmacokinetics of nifedipine slow release tablet in Mexican subjects: further evidence for an oxidation polymorphism. J. Clin. Pharm. 29: 816–820, 1989.
- [66984] 5314. Inoue, K.; Inazawa, J.; Nakagawa, H.; Shimada, T.; Yamazaki, H.; Guengerich, F. P.; Abe, T.: Assignment of the human cytochrome P–450 nifedipine oxidase gene (CYP3A4) to chromosome 7 at band q22.1 by fluorescence in situ hybridization. Jpn. J. Hum. Genet. 37: 133–138, 1992.
- [66985] 5315. Kittles, R. A.; Chen, W.; Panguluri, R. K.; Ahaghotu, C.; Jackson, A.; Adebamowo, C. A.; Griffin, R.; Williams, T.; Ukoli, F.; Adams–Campbell, L.; Kwagyan, J.; Isaacs, W.; Freeman, V.; Dunston, G. M.: CYP3A4–V and prostate cancer in African Americans: causal or confounding association because of population stratification? Hum. Genet. 110: 553–560, 2002.
- [66986] 5316. Kleinbloesem, C. H.; van Brummelen, P.; Faber, H.; Danhof, M.; Vermeulen, N. P. E.; Breimer, D. D.: Variability in nifedipine pharmacokinetics and dynamics: a new ox-

ation polymorphism in man. *Biochem. Pharm.*  
33:3721–3724, 1984.

- [66987] 5317. Lehmann, J. M.; McKee, D. D.; Watson, M. A.; Willson, T. M.; Moore, J. T.; Kliewer, S. A.: The human orphan nuclear receptor PXR is activated by compounds that regulate CYP3A4 gene expression and cause drug interactions. *J. Clin. Invest.* 102: 1016–1023, 1998.
- [66988] 5318. Lown, K. S.; Bailey, D. G.; Fontana, R. J.; Janardan, S. K.; Adair, C. H.; Fortlage, L. A.; Brown, M. B.; Guo, W.; Watkins, P. B.: Grapefruit juice increases felodipine oral availability in humans by decreasing intestinal CYP3A protein expression. *J. Clin. Invest.* 99: 2545–2553, 1997.
- [66989] 5319. Molowa, D. T.; Schuetz, E. G.; Wrighton, S. A.; Watkins, P. B.; Kremers, P.; Mendez-Picon, G.; Parker, G. A.; Guzelian, P. S.: Complete cDNA sequence of a cytochrome P-450 inducible by glucocorticoids in human liver. *Proc. Nat. Acad. Sci.* 83: 5311–5315, 1986.
- [66990] 5320. Paris, P. L.; Kupelian, P. A.; Hall, J. M.; Williams, T. L.; Levin, H.; Klein, E. A.; Casey, G.; Witte, J. S.: Association between a CYP3A4 genetic variant and clinical presentation in African-American prostate cancer patients. *Cancer Epidemiol. Biomarkers Prev.* 8: 901–905, 1999.
- [66991] 5321. Rebbeck, T. R.; Jaffe, J. M.; Walker, A. H.; Wein, A. J.;

Malkowicz, S. B.: Modification of clinical presentation of prostate tumors by a novel genetic variant in CYP3A4. *J. Nat. Cancer Inst.* 90: 1225–1229, 1998.

- [66992] 5322. Renwick, A. G.; Robertson, D. R. C.; Macklin, B.; Challenor, V.; Waller, D. G.; George, C. F.: The pharmacokinetics of oral nifedipine—a population study. *Brit. J. Clin. Pharm.* 25: 701–708, 1988.
- [66993] 5323. Nebert, D. W.; Adesnik, M.; Coon, M. J.; Estabrook, R. W.; Gonzalez, F. J.; Guengerich, F. P.; Gunsalus, I. C.; Johnson, E. F.; Kemper, B.; Levin, W.; Phillips, I. R.; Sato, R.; Waterman, M. R.: The P450 gene superfamily: recommended nomenclature. *DNA* 6: 1–11, 1987.
- [66994] 5324. Spielberg, S. P.: Personal Communication. Toronto, Ontario, Canada 2/26/1988.
- [66995] 5325. Thum, T.; Borlak, J.: Gene expression in distinct regions of the heart. *Lancet* 355: 979–983, 2000.
- [66996] 5326. Chen, Y.; Faraco, J.; Yin, W.; Germiller, J.; Francke, U.; Bonadio, J.: Structure, chromosomal localization, and expression pattern of the murine Magp gene. *J. Biol. Chem.* 268: 27381–27389, 1993.
- [66997] 5327. Faraco, J.; Bashir, M.; Rosenbloom, J.; Francke, U.: Characterization of the human gene for microfibril-associated glycoprotein (MFAP2), assignment to chromosome

1p36.1–p35, and linkage to D1S170. *Genomics* 25:630–637, 1995.

[66998] 5328. Gibson, M. A.; Hughes, J. L.; Fanning, J. C.; Cleary, E. G.: The major antigen of elastin-associated microfibrils is a 31-kDa glycoprotein. *J. Biol. Chem.* 261: 11429–11436, 1986.

[66999] 5329. Low, F. N.: Microfibrils: fine filamentous components of the tissuespace. *Anat. Rec.* 142: 131–137, 1962.

[67000] 5330. Aberdam, E.; Bertolotto, C.; Sviderskaya, E. V.; de Thillot, V.; Hemesath, T. J.; Fisher, D. E.; Bennett, D. C.; Ortonne, J.-P.; Ballotti, R.: Involvement of microphthalmia in the inhibition of melanocyte lineage differentiation and of melanogenesis by agouti signal protein. *J. Biol. Chem.* 273: 19560–19565, 1998.

[67001] 5331. Kelavkar, U. P.; Badr, K. F.: Effects of mutant p53 expression on human 15-lipoxygenase–promoter activity and murine 12/15-lipoxygenase gene expression: evidence that 15-lipoxygenase is a mutator gene. *Proc. Nat. Acad. Sci.* 96: 4378–4383, 1999.

[67002] 5332. Sigal, E.; Craik, C. S.; Highland, E.; Grunberger, D.; Costello, L. L.; Dixon, R. A. F.; Nadel, J. A.: Molecular cloning and primary structure of human 15-lipoxygenase. *Biochem. Biophys. Res. Commun.* 157:457–464, 1988.

- [67003] 5333.Yoshimoto, T.; Suzuki, H.; Yamamoto, S.; Takai, T.; Yokoyama, C.;Tanabe, T.: Cloning and sequence analysis of the cDNA for arachidonate12-lipoxygenase of porcine leukocytes. *Proc. Nat. Acad. Sci.* 87:2142–2146, 1990.
- [67004] 5334.Minoshima, S.; Fukuyama, R.; Yamamoto, T.; Shimizu, N.: Mappingof human long-chain acyl-CoA synthetase to chromosome 4. (Abstract) *Cytogenet.Cell Genet.* 58: 1888 only, 1991.
- [67005] 5335.Cantu, E. S.; Sprinkle, T. J.; Ghosh, B.; Singh, I.: The humanpalmitoyl-CoA ligase (FACL2) gene maps to the chromosome 4q34–q35region by fluorescence in situ hybridization (FISH) and somatic cellhybrid panels. *Genomics* 28: 600–602, 1995.
- [67006] 5336.Benson, D. W.; MacRae, C. A.; Vesely, M. R.; Walsh, E. P.; Seidman,J. G.; Seidman, C. E.; Satler, C. A.: Missense mutation in the poreregion of HERG causes familial long QT syndrome. *Circulation* 93:1791–1795, 1996.
- [67007] 5337.Brook, J. D.; Shaw, D. J.; Meredith, A. L.; Bruns, G. A. P.; Harper,P. S.: Localisation of genetic markers and orientation of the linkagegroup on chromosome 19. *Hum. Genet.* 68: 282–285, 1984.
- [67008] 5338.Bruns, G. A. P.; Regina, V. M.; Gerald, P. S.: Lysosomal DNaseand chromosome 19.(Abstract) *J. Cell Biol.* 83:

444a only, 1979.

- [67009] 5339.Kawane, K.; Fukuyama, H.; Kondoh, G.; Takeda, J.; Ohsawa, Y.; Uchiyama,Y.; Nagata, S.: Requirement of DNase II for definitive erythropoiesisin the mouse fetal liver. *Science* 292: 1546–1549, 2001.
- [67010] 5340.Yasuda, T.; Nadano, D.; Sawazaki, K.; Kishi, K.: Genetic polymorphismof human deoxyribonuclease II (DNase II): low activity levels in urineand leukocytes are due to an autosomal recessive allele. *Ann. Hum.Genet.* 56: 1–10, 1992.
- [67011] 5341.Yasuda, T.; Takeshita, H.; Iida, R.; Nakajima, T.; Hosomi, O.;Nakashima, Y.; Kishi, K.: Molecular cloning of the cDNA encodinghuman deoxyribonuclease II. *J. Biol. Chem.* 273: 2610–2616, 1998.
- [67012] 5342.Yasuda, T.; Takeshita, H.; Iida, R.; Nakajima, T.; Hosomi, O.;Nakashima, Y.; Mogi, K.; Kishi, K.: Chromosomal localization of ahuman deoxyribonuclease II gene (DNASE2) to 19p13.2–p13.1 using boththe polymerase chain reaction and fluorescence in situ hybridizationanalysis. *Biochem. Biophys. Res. Commun.* 244: 815–818, 1998.
- [67013] 5343.Baylin, S. B.: Tying it all together: epigenetics, genetics, cellcycle, and cancer. *Science* 277: 1948–1949,



1997.

- [67014] 5344. Bestor, T.; Laudano, A.; Mattaliano, R.; Ingram, V.: Cloning and sequencing of a cDNA encoding DNA methyltransferase of mouse cells: the carboxyl-terminal domain of the mammalian enzymes is related to bacterial restriction methyltransferases. *J. Molec. Biol.* 203: 971–983, 1988.
- [67015] 5345. Bestor, T. H.: The DNA methyltransferases of mammals. *Hum. Molec. Genet.* 9: 2395–2402, 2000.
- [67016] 5346. Bestor, T. H.: DNA methylation: evolution of a bacterial immune function into a regulator of gene expression and genome structure in higher eukaryotes. *Phil. Trans. Roy. Soc. London B* 326: 179–187, 1990.
- [67017] 5347. Chuang, L. S.-H.; Ian, H.-I.; Koh, T.-W.; Ng, H.-H.; Xu, G.; Li, B. F. L.: Human DNA-(cytosine-5) methyltransferase-PCNA complex as a target for p21(WAF1). *Science* 277: 1996–2000, 1997.
- [67018] 5348. El-Deiry, W. S.; Nelkin, B. D.; Celano, P.; Chiu Yen, R.-W.; Falco, J. P.; Hamilton, S. R.; Baylin, S. B.: High expression of the DNA methyltransferase gene characterizes human neoplastic cells and progression stages of colon cancer. *Proc. Nat. Acad. Sci.* 88: 3470–3474, 1991.
- [67019] 5349. Fuks, F.; Burgers, W. A.; Brehm, A.; Hughes-Davies,

L.; Kouzarides,T.: DNA methyltransferase Dnmt1 associates with histone deacetylase activity. *Nature Genet.* 24: 88–91, 2000.

[67020] 5350.Howell, C. Y.; Bestor, T. H.; Ding, F.; Latham, K. E.; Mertineit,C.; Trasler, J. M.; Chaillet, J. R.: Genomic imprinting disrupted by a maternal effect mutation in the Dnmt1 gene. *Cell* 104: 829–838,2001.

[67021] 5351.Hsu, D.–W.; Lin, M.–J.; Lee, T.–L.; Wen, S.–C.; Chen, X.; Shen,C.–K. J.: Two major forms of DNA (cytosine–5) methyltransferase in human somatic tissues. *Proc. Nat. Acad. Sci.* 96: 9751–9756, 1999.

[67022] 5352.Lee, P. P.; Fitzpatrick, D. R.; Beard, C.; Jessup, H. K.; Lehar,S.; Makar, K. W.; Perez–Melgosa, M.; Sweetser, M. T.; Schlissel, M.S.; Nguyen, S.; Cherry, S. R.; Tsai, J. H.; Tucker, S. M.; Weaver,W. M.; Kelso, A.; Jaenisch, R.; Wilson, C. B.: A critical role for Dnmt1 and DNA methylation in T cell development, function, and survival. *Immunity* 15:763–774, 2001.

[67023] 5353.Li, E.; Bestor, T. H.; Jaenisch, R.: Targeted mutation of the DNA methyltransferase gene results in embryonic lethality. *Cell* 69:915–926, 1992.

[67024] 5354.Lyko, F.; Ramsahoye, B. H.; Kashevsky, H.; Tudor, M.; Mastrangelo,M.–A.; Orr–Weaver, T. L.; Jaenisch, R.:

Mammalian (cytosine-5) methyltransferases cause genomic DNA methylation and lethality in *Drosophila*. *Nature Genet.* 23: 363–366, 1999.

[67025] 5355. Rhee, I.; Bachman, K. E.; Park, B. H.; Jair, K.-W.; Yen, R.-W. C.; Schuebel, K. E.; Cui, H.; Feinberg, A. P.; Lengauer, C.; Kinzler, K. W.; Baylin, S. B.; Vogelstein, B.: DNMT1 and DNMT3b cooperate to silence genes in human cancer cells. *Nature* 416: 552–556, 2002.

[67026] 5356. Rhee, I.; Jair, K.-W.; Yen, R.-W. C.; Lengauer, C.; Herman, J. G.; Kinzler, K. W.; Vogelstein, B.; Baylin, S. B.; Schuebel, K. E.: CpG methylation is maintained in human cancer cells lacking DNMT1. *Nature* 404: 1003–1007, 2000.

[67027] 5357. Robertson, K. D.; Ait-Si-Ali, S.; Yokochi, T.; Wade, P. A.; Jones, P. L.; Wolffe, A. P.: DNMT1 forms a complex with Rb, E2F1 and HDAC1 and represses transcription from E2F-responsive promoters. *Nature Genet.* 25: 338–342, 2000.

[67028] 5358. Rouleau, J.; Tanigawa, G.; Szyf, M.: The mouse DNA methyltransferase 5-prime region: a unique housekeeping gene promoter. *J. Biol. Chem.* 267: 7368–7377, 1992.

[67029] 5359. Rountree, M. R.; Bachman, K. E.; Baylin, S. B.: DNMT1 binds HDAC2 and a new co-repressor, DMAP1, to form a

complex at replication foci. *NatureGenet.* 25: 269–277, 2000.

[67030] 5360. Tucker, K. L.; Talbot, D.; Lee, M. A.; Leonhardt, H.; Jaenisch, R.: Complementation of methylation deficiency in embryonic stem cells by a DNA methyltransferase mini-gene. *Proc. Nat. Acad. Sci.* 93: 12920–12925, 1996.

[67031] 5361. Yen, R.-W. C.; Vertino, P. M.; Nelkin, B. D.; Yu, J. J.; El-Deiry, W.; Cumaraswamy, A.; Lennon, G. G.; Trask, B. J.; Celano, P.; Baylin, S. B.: Isolation and characterization of the cDNA encoding human DNA methyltransferase. *Nucleic Acids Res.* 20: 2287–2291, 1992.

[67032] 5362. Ye, Q.; Chung, L. W. K.; Li, S.; Zhau, H. E.: Identification of a novel FAS/ER- $\alpha$  fusion transcript expressed in human cancer cells. *Biochim. Biophys. Acta* 1493: 373–377, 2000.

[67033] 5363. Zuppan, P.; Hall, J. M.; Lee, M. K.; Ponglikitmongkol, M.; King, M.-C.: Possible linkage of the estrogen receptor gene to breast cancer in a family with late-onset disease. *Am. J. Hum. Genet.* 48: 1065–1068, 1991.

[67034] 5364. Zuppan, P. J.; Hall, J. M.; Ponglikitmongkol, M.; Spielman, R.; King, M. C.: Polymorphisms at the estrogen receptor (ESR) locus and linkage relationships on chromosome 6q. (Abstract) *Cytogenet. Cell Genet.* 51: 1116 only,

1989.

- [67035] 5365. Calabi, F.; Cilli, V.: CBFA2T1, a gene rearranged in human leukemia, is a member of a multigene family. *Genomics* 52: 332–341, 1998.
- [67036] 5366. Erickson, P.; Gao, J.; Chang, K.-S.; Look, T.; Whisenant, E.; Raimondo, S.; Lasher, R.; Trujillo, J.; Rowley, J. D.; Drabkin, H. A.: Identification of breakpoints in t(8;21) acute myelogenous leukemia and isolation of a fusion transcript, AML1/ETO, with similarity to *Drosophila* segmentation gene, runt. *Blood* 80: 1825–1831, 1992.
- [67037] 5367. Miyamoto, T.; Weissman, I. L.; Akashi, K.: AML1/ETO-expressing nonleukemic stem cells in acute myelogenous leukemia with 8;21 chromosomal translocation. *Proc. Nat. Acad. Sci.* 97: 7521–7526, 2000.
- [67038] 5368. Miyoshi, H.; Kozu, T.; Shimizu, K.; Enomoto, K.; Maseki, N.; Kaneko, Y.; Kamada, N.; Ohki, M.: The t(8;21) translocation in acute myeloid leukemia results in production of an AML1–MTG8 fusion transcript. *EMBOJ.* 12: 2715–2721, 1993.
- [67039] 5369. Miyoshi, H.; Shimizu, K.; Kozu, T.; Maseki, N.; Kaneko, Y.; Ohki, M.: t(8;21) breakpoints on chromosome 21 in acute myeloid leukemia are clustered within a limited region of a single gene, AML1. *Proc. Nat. Acad. Sci.* 88:

10431–10434, 1991.

- [67040] 5370.Linggi, B.; Muller–Tidow, C.; van de Locht, L.; Hu, M.; Nip, J.; Serve, H.; Berdel, W. E.; van der Reijden, B.; Quelle, D. E.; Rowley, J. D.; Cleveland, J.; Jansen, J. H.; Pandolfi, P. P.; Hiebert, S.W.: The t(8;21) fusion protein, AML1–ETO, specifically represses the transcription of the p14(ARF) tumor suppressor in acute myeloid leukemia. *Nature Med.* 8: 743–750, 2002.
- [67041] 5371.Minucci, S.; Maccarana, M.; Cioce, M.; De Luca, P.; Gelmetti, V.; Segalla, S.; Di Croce, L.; Giavara, S.; Matteucci, C.; Gobbi, A.; Bianchini, A.; Colombo, E.; Schiavoni, I.; Badaracco, G.; Hu, X.; Lazar, M. A.; Landsberger, N.; Nervi, C.; Pelicci, P. G.: Oligomerization of RAR and AML1 transcription factors as a novel mechanism of oncogenic activation. *Molec. Cell* 5: 811–820, 2000.
- [67042] 5372.Niwa–Kawakita, M.; Miyoshi, H.; Gotoh, O.; Matsushima, Y.; Nishimura, M.; Shisa, H.; Ohki, M.: Cloning and gene mapping of the mouse homologue of the CBFA2T1 gene associated with human acute myeloid leukemia. *Genomics* 29:755–759, 1995.
- [67043] 5373.Nucifora, G.; Rowley, J. D.: The AML1 and ETO genes in acute myeloid leukemia with a t(8;21). *Leukemia Lymphoma* 14: 353–362, 1994.

- [67044] 5374.Schoch, C.; Kohlmann, A.; Schnittger, S.; Brors, B.; Dugas, M.;Mergenthaler, S.; Kern, W.; Hiddemann, S.; Eils, R.; Haferlach, T.: Acute myeloid leukemias with reciprocal rearrangements can be distinguishedby specific gene expression profiles. *Proc. Nat. Acad. Sci.* 99:10008–10013, 2002.
- [67045] 5375.Wolford, J. K.; Bogardus, C.; Prochazka, M.: Polymorphism inthe 3–prime untranslated region of MTG8 is associated with obesityin Pima Indian males. *Biochem. Biophys. Res. Commun.* 246: 624–626,1998.
- [67046] 5376.Wolford, J. K.; Prochazka, M.: Structure and expression of thehuman MTG8/ETO gene. *Gene* 212: 103–109, 1998.
- [67047] 5377.Yergeau, D. A.; Hetherington, C. J.; Wang, Q.; Zhang, P.; Sharpe,A. H.; Binder, M.; Marin–Padilla, M.; Tenen, D. G.; Speck, N. A.;Zhang, D.–E.: Embryonic lethality and impairment of haematopoiesisin mice heterozygous for an AML1–ETO fusion gene. *Nature Genet.* 15:303–306, 1997.
- [67048] 5378.Dorfman, J.; Lazaris–Karatzas, A.; Malo, D.; Sonenberg, N.; Gros,P.: Chromosomal assignment of one of the mammalian translation initiationfactor eIF–4E genes. *Genomics* 9: 785–788, 1991.
- [67049] 5379.Jones, R. M.; Branda, J.; Johnston, K. A.; Polymenis,

M.; Gadd, M.; Rustgi, A.; Callanan, L.; Schmidt, E. V.: An essential E box in the promoter of the gene encoding the mRNA cap-binding protein (eukaryotic initiation factor 4E) is a target for activation by c-myc. *Molec. Cell. Biol.* 16: 4754–4764, 1996.

[67050] 5380. Jones, R. M.; MacDonald, M. E.; Branda, J.; Altherr, M. R.; Louis, D. N.; Schmidt, E. V.: Assignment of the human gene encoding eukaryotic initiation factor 4E (EIF4E) to the region q21–25 on chromosome 4. *Somat. Cell Molec. Genet.* 23: 221–223, 1997.

[67051] 5381. Pause, A.; Belsham, G. J.; Gingras, A.-C.; Donze, O.; Lin, T.-A.; Lawrence, J. C., Jr.; Sonenberg, N.: Insulin-dependent stimulation of protein synthesis by phosphorylation of a regulator of 5-prime-cap function. *Nature* 371: 762–767, 1994.

[67052] 5382. Pelletier, J.; Brook, J. D.; Housman, D. E.: Assignment of two of the translation initiation factor-4E (EIF4EL1 and EIF4EL2) genes to human chromosomes 4 and 20. *Genomics* 10: 1079–1082, 1991.

[67053] 5383. Pyronnet, S.; Imataka, H.; Gingras, A.-C.; Fukunaga, R.; Hunter, T.; Sonenberg, N.: Human eukaryotic translation initiation factor 4G (eIF4G) recruits Mnk1 to phosphorylate eIF4E. *EMBO J.* 18: 270–279, 1999.



- [67054] 5384. Rychlik, W.; Domier, L. L.; Gardner, P. R.; Hellmann, G. M.; Rhoads, R. E.: Amino acid sequence of the mRNA cap-binding protein from human tissues. *Proc. Nat. Acad. Sci.* 84: 945–949, 1987.
- [67055] 5385. Waskiewicz, A. J.; Flynn, A.; Proud, C. G.; Cooper, J. A.: Mitogen-activated protein kinases activate the serine/threonine kinases Mnk1 and Mnk2. *EMBO J.* 16: 1909–1920, 1997.
- [67056] 5386. Aman, P.; Panagopoulos, I.; Lassen, C.; Fioretos, T.; Mencinger, M.; Toresson, H.; Hoglund, M.; Forster, A.; Rabbitts, T. H.; Ron, D.; Mandahl, N.; Mitelman, F.: Expression patterns of the human sarcoma-associated genes FUS and EWS and the genomic structure of FUS. *Genomics* 37:1–8, 1996.
- [67057] 5387. Aurias, A.; Rimbaut, C.; Buffe, D.; Zucker, J.-M.; Mazabraud, A.: Translocation involving chromosome 22 in Ewing's sarcoma: a cytogenetic study of four fresh tumors. *Cancer Genet. Cytogenet.* 12: 21–25, 1984.
- [67058] 5388. Ben-David, Y.; Giddens, E. B.; Letwin, K.; Bernstein, A.: Erythroleukemia induction by Friend murine leukemia virus: insertional activation of a new member of the ets gene family, Fli-1, closely linked to c-ets-1. *Genes Dev.* 5: 908–918, 1991.

- [67059] 5389. Budarf, M.; Sellinger, B.; Griffin, C.; Emanuel, B. S.: Comparativemapping of the constitutional and tumor-associated 11;22 translocations. *Am.J. Hum. Genet.* 45: 128–139, 1989.
- [67060] 5390. Zhang, Y.; Saison, M.; Spaepen, M.; De Strooper, B.; Van Leuven, F.; David, G.; Van den Berghe, H.; Cassiman, J.-J.: Mapping of human fibronectin receptor beta subunit gene to chromosome 10. *Somat. Cell Molec. Genet.* 14: 99–104, 1988.
- [67061] 5391. Argraves, W. S.; Dickerson, K.; Burgess, W. H.; Ruoslahti, E.: Fibulin, a novel protein that interacts with the fibronectin receptor–beta subunit cytoplasmic domain. *Cell* 58: 623–629, 1989.
- [67062] 5392. Korenberg, J. R.; Chen, X.-N.; Tran, H.; Argraves, W. S.: Localization of the human gene for fibulin–1 (FBLN1) to chromosome band 22q13.3. *Cytogenet. Cell Genet.* 68: 192–193, 1995.
- [67063] 5393. Mattei, M.-G.; Pan, T.-C.; Zhang, R.-Z.; Timpl, R.; Chu, M.-L.: The fibulin–1 gene (FBLN1) is located on human chromosome 22 and on mouse chromosome 15. *Genomics* 22: 437–438, 1994.
- [67064] 5394. Collod, G.; Chu, M.-L.; Sasaki, T.; Coulon, M.; Timpl, R.; Renkart, L.; Weissenbach, J.; Jondeau, G.; Bourdarias, J.

P.; Junien, C.; Boileau, C.: Fibulin-2: genetic mapping and exclusion as a candidate gene in Marfan syndrome type 2. *Europ. J. Hum. Genet.* 4: 292–295, 1996.

[67065] 5395. Kuivaniemi, H.; Marshall, A.; Ganguly, A.; Chu, M.-L.; Abbott, W. M.; Tromp, G.: Fibulin-2 exhibits high degree of variability, but no structural changes concordant with abdominal aortic aneurysms. *Europ. J. Hum. Genet.* 6: 642–646, 1998.

[67066] 5396. Pavlakis, G. N.; Hizuka, N.; Gorden, P.; Seeburg, P. H.; Hamber, D. H.: Expression of two human growth hormone genes in monkey cells infected by simian virus 40 recombinants. *Proc. Nat. Acad. Sci.* 78: 7398–7402, 1981.

[67067] 5397. Leone, A.; McBride, O. W.; Weston, A.; Wang, M. G.; Anglard, P.; Cropp, C. S.; Goepel, J. R.; Lidereau, R.; Callahan, R.; Marston Linehan, W.; Rees, R. C.; Harris, C. C.; Liotta, L. A.; Steeg, P. S.: Somatic allelic deletion of nm23 in human cancer. *Cancer Res.* 51: 2490–2493, 1991.

[67068] 5398. Broide, D. H.; Hoffman, H.; Sriramaraio, P.: Genes that regulate eosinophilic inflammation. *Am. J. Hum. Genet.* 65: 302–307, 1999.

[67069] 5399. Campbell, H. D.; Tucker, W. Q. J.; Hort, Y.; Martinson, M. E.; Mayo, G.; Clutterbuck, E. J.; Sanderson, C. J.; Young, I. G.: Molecular cloning, nucleotide sequence, and

expression of the gene encoding human eosinophil differentiation factor (interleukin-5). *Proc. Nat. Acad. Sci.* 84: 6629-6633, 1987.

- [67070] 5400. Chandrasekharappa, S. C.; Rebelsky, M. S.; Firak, T. A.; Le Beau, M. M.; Westbrook, C. A.: A long-range restriction map of the interleukin-4 and interleukin-5 linkage group on chromosome 5. *Genomics* 6: 94-99, 1990.
- [67071] 5401. Coffman, R. L.; Seymour, B. W. P.; Hudak, S.; Jackson, J.; Rennick, D.: Antibody to interleukin-5 inhibits helminth-induced eosinophilia in mice. *Science* 245: 308-310, 1989.
- [67072] 5402. Cogan, E.; Schandene, L.; Crusiaux, A.; Cochaux, P.; Velu, T.; Goldman, M.: Clonal proliferation of type 2 helper T cells in a man with the hypereosinophilic syndrome. *New Eng. J. Med.* 330: 535-538, 1994.
- [67073] 5403. Kozak, C.: Personal Communication. Bethesda, Md. 6/9/1988.
- [67074] 5404. Pereira, E.; Goldblatt, J.; Rye, P.; Sanderson, C.; Le Souef, P.: Mutation analysis of interleukin-5 in an asthmatic cohort. *Hum. Mutat.* 11: 51-54, 1998.
- [67075] 5405. Rodrigues, V., Jr.; Abel, L.; Piper, K.; Dessein, A. J.: Segregation analysis indicates a major gene in the control of interleukin-5 production in humans infected with Schis-

tosoma mansonii. Am. J. Hum. Genet. 59:453–461, 1996.

[67076] 5406.Simon, H. U.; Plotz, S. G.; Dummer, R.; Blaser, K.: Abnormal clones of T cells producing interleukin-5 in idiopathic eosinophilia. NewEng. J. Med 341: 1112–1120, 1999.

[67077] 5407.Sutherland, G. R.; Baker, E.; Callen, D. F.; Campbell, H. D.;Young, Y. G.; Sanderson, C. J.; Garson, O. M.; Lopez, A. F.; Vadas,M. A.: Interleukin-5 is at 5q31 and is deleted in the 5q- syndrome. Blood 71:1150–1152, 1988.

[67078] 5408.Tanabe, T.; Konishi, M.; Mizuta, T.; Noma, T.; Honjo, T.: Molecular cloning and structure of the human interleukin-5 gene. J. Biol. Chem. 262:16580–16584, 1987.

[67079] 5409.Yokota, T.; Coffman, R. L.; Hagiwara, H.; Rennick, D. M.; Takebe,Y.; Yokota, K.; Gemmell, L.; Shrader, B.; Yang, G.; Meyerson, P.;Luh, J.; Hoy, P.; Pene, J.; Briere, F.; Spits, H.; Banchereau, J.;de Vries, J.; Lee, F. D.; Arai, N.; Arai, K.: Isolation and characterization of lymphokine cDNA clones encoding mouse and human IgA-enhancing factor and eosinophil colony-stimulating factor activities: relationship to interleukin 5. Proc. Nat. Acad. Sci. 84: 7388–7392, 1987.

[67080] 5410.Geijsen, N.; Uings, I. J.; Pals, C.; Armstrong, J.; McKinnon, M.;Raaijmakers, J. A. M.; Lammers, J.-W. J.; Koen-

derman, L.; Coffey, P. J.: Cytokine-specific transcriptional regulation through an IL-5R- $\alpha$ -interacting protein. *Science* 293: 1136–1138, 2001.

[67081] 5411. Gough, N. M.; Rakar, S.: Localization of the IL-5 receptor gene to the distal half of murine chromosome 6 using recombinant inbred strains of mice. *Genomics* 12: 855–856, 1992.

[67082] 5412. Isobe, M.; Kumura, Y.; Murata, Y.; Takaki, S.; Tomimaga, A.; Takatsu, K.; Ogita, Z.: Localization of the gene encoding the  $\alpha$  subunit of human interleukin-5 receptor (IL5RA) to chromosome region 3p24–3p26. *Genomics* 14: 755–758, 1992.

[67083] 5413. Jacob, C. O.; Myktyyn, K.; Varcony, T.; Drabkin, H. A.: Mapping of the interleukin 5 receptor gene to human chromosome 3p25–p26 and to mouse chromosome 6 close to the Raf-1 locus with polymorphic tandem repeat sequences. *Mammalian Genome* 4: 435–439, 1993.

[67084] 5414. Sun, Z.; Yergeau, D. A.; Tuypens, T.; Tavernier, J.; Paul, C. C.; Baumann, M. A.; Tenen, D. G.; Ackerman, S. J.: Identification and characterization of a functional promoter region in the human eosinophil IL-5 receptor  $\alpha$  subunit gene. *J. Biol. Chem.* 270: 1462–1471, 1995.

[67085] 5415. Tavernier, J.; Tuypens, T.; Plaetinck, G.; Verhee, A.;

Fiers, W.;Devos, R.: Molecular basis of the membrane–anchored and two solubleisoforms of the human interleukin 5 receptor alpha subunit. Proc.Nat. Acad. Sci. 89: 7041–7045, 1992.

[67086] 5416.Zhang, J.; Kuvelkar, R.; Cheewatrakoolpong, B.; Williams, S.; Egan,R. W.; Billah, M. M.: Evidence for multiple promoters of the humanIL–5 receptor alpha subunit gene: a novel 6–base pair element determinescell–specific promoter function. J. Immun. 15: 5412–5421, 1997.

[67087] 5417.Povey, S.; Morton, N. E.; Sherman, S. L.: Report of the committeeon the genetic constitution of chromosomes 1 and 2. Cytogenet. CellGenet. 40: 67–106, 1985.

[67088] 5418.Bird, A.; Taggart, M.; Frommer, M.; Miller, O. J.; Macleod, D.: A fraction of the mouse genome that is derived from islands of nonmethylated,CpG–rich DNA. Cell 40: 91–99, 1985.

[67089] 5419.Pikkarainen, T.; Kallunki, T.; Tryggvason, K.: Human laminin B2chain: comparison of the complete amino acid sequence with the B1chain reveals variability in sequence homology between different structuraldomains. J. Biol. Chem. 263: 6751–6758, 1988.

[67090] 5420.Sasaki, M.; Yamada, Y.: The laminin B2 chain has a multidomainstructure homologous to the B1 chain. J. Biol.

Chem. 262: 17111–17117,1987.

- [67091] 5421.Smyth, N.; Vatansever, H. S.; Murray, P.; Meyer, M.; Frie, C.;Paulsson, M.; Edgar, D.: Absence of basement membranes after targetingthe LAMC1 gene results in embryonic lethality due to failure of endodermdifferentiation. J. Cell Biol. 144: 151–160, 1999.
- [67092] 5422.Benlian, P.; Foubert, L.; Gagne, E.; Bernard, L.; De Gennes, J.L.; Langlois, S.; Robinson, W.; Hayden, M.: Complete paternal isodisomyfor chromosome 8 unmasked by lipoprotein lipase deficiency. Am. J.Hum. Genet. 59: 431–436, 1996.
- [67093] 5423.Schoenmakers, E. P. P. M.; Wanschura, S.; Mols, R.; Bullerdiek,J.; Van den Berghe, H.; Van de Ven, W. J. M.: Recurrent rearrangementsin the high mobility group protein gene, HMGI-C, in benign mesenchymaltumours. Nature Genet. 10: 436–444, 1995.
- [67094] 5424.Turc-Carel, C.; Pietrzak, E.; Kakati, S.; Kinniburgh, A. J.; Sandberg,A. A.: The human int-1 gene is located at chromosome region 12q12–12q13and is not rearranged in myxoid liposarcoma with t(12;16)(q13;p11). Onco-geneRes. 1: 397–405, 1987.
- [67095] 5425.Beck, Y.; Oren, R.; Amit, B.; Levanon, A.; Gorecki, M.; Hartman,J. R.: Human Mn superoxide dismutase cDNA se–



quence. *Nucleic Acids Res.* 15: 9076, 1987.

[67096] 5426.Church, S. L.; Grant, J. W.; Meese, E. U.; Trent, J. M.: Sublocalization of the gene encoding manganese superoxide dismutase (MnSOD/SOD2) to 6q25 by fluorescence in situ hybridization and somatic cell hybrid mapping. *Genomics* 14: 823–825, 1992.

[67097] 5427.Creagan, R.; Tischfield, J.; Ricciuti, F.; Ruddle, F. H.: Chromosome assignments of genes in man using mouse-human somatic cell hybrids: mitochondrial superoxide dismutase (indophenol oxidase-B, tetrameric) to chromosome 6. *Humangenetik* 20: 203–209, 1973.

[67098] 5428.Figueroa, F.; Vincek, V.; Kasahara, M.; Bell, G. I.; Klein, J.: Mapping of the Sod-2 locus into the t complex on mouse chromosome 17. *Immunogenetics* 28: 260–264, 1988.

[67099] 5429.Heckl, K.: Isolation of cDNAs encoding human manganese superoxide dismutase. *Nucleic Acids Res.* 16: 6224, 1988.

[67100] 5430.Li, Y.; Huang, T.-T.; Carlson, E. J.; Melov, S.; Ursell, P. C.; Olson, J. L.; Noble, L. J.; Yoshimura, M. P.; Berger, C.; Chan, P.H.; Wallace, D. C.; Epstein, C. J.: Dilated cardiomyopathy and neonatal lethality in mutant mice lacking manganese superoxide dismutase. *Nature Genet.* 11: 376–381,

1995.

- [67101] 5431.Melov, S.; Coskun, P.; Patel, M.; Tuinstra, R.; Cottrell, B.; Jun,A. S.; Zastawny, T. H.; Dizdaroglu, M.; Goodman, S. I.; Huang, T.–T.;Miziorko, H.; Epstein, C. J.; Wallace, D. C.: Mitochondrial diseasein superoxide dismutase 2 mutant mice. *Proc. Nat. Acad. Sci.* 96:846–851, 1999.
- [67102] 5432.Melov, S.; Schneider, J. A.; Day, B. J.; Hinerfeld, D.; Coskun,P.; Mirra, S. S.; Crapo, J. D.; Wallace, D. C.: A novel neurologicalphenotype in mice lacking mitochondrial manganese superoxide dismutase. *NatureGenet.* 18: 159–163, 1998.
- [67103] 5433.Michelson, A. M.; McCord, J. M.; Fridovich, I.: Super-oxide andSuperoxide Dismutases. New York: Academic Press , 1977.
- [67104] 5434.Hassett, C.; Robinson, K. B.; Beck, N. B.; Omiecinski, C. J.:The human microsomal epoxide hydrolase gene (EPHX1): complete nucleotidesequence and structural characterization. *Genomics* 23: 433–442,1994.
- [67105] 5435.Jackson, M. R.; Craft, J. A.; Burchell, B.: Nucleotide and deducedamino acid sequence of human liver micro-somal epoxide hydrolase. *NucleicAcids Res.* 15: 7188 only, 1987.
- [67106] 5436.Kutt, H.; Brennan, R.; Dehejia, H.; Verebely, K.:

Diphenylhydantoinintoxication: a complication of isoni-  
azid therapy. *Am. Rev. Resp.Dis.* 101: 377–383, 1970.

[67107] 5437.Kutt, H.; Verebely, K.; McDowell, F.: Inhibition of  
diphenylhydantoinmetabolism in rat and in rat liver mi-  
crosome by antitubercular drugs. *Neurology* 18:706–710,  
1968.

[67108] 5438.Kutt, H.; Wolk, M.; Scherman, R.; McDowell, F.: In-  
sufficientparahydroxylation as a cause of diphenylhydan-  
toin toxicity. *Neurology* 14:542–548, 1964.

[67109] 5439.McGlynn, K. A.; Rosvold, E. A.; Lustbader, E. D.; Hu,  
Y.; Clapper,M. L.; Zhou, T.; Wild, C. P.; Xia, X.–L.; Baffoe-  
Bonnie, A.; Ofori-Adjei,D.; Chen, G.–C.; London, W. T.;  
Shen, F.–M.; Buetow, K. H.: Susceptibilityto hepatocellular  
carcinoma is associated with genetic variation inthe enzy-  
matic detoxification of aflatoxin B1. *Proc. Nat. Acad. Sci.*  
92:2384–2387, 1995.

[67110] 5440.Nadeau, J. H.: Personal Communication. Bar Harbor,  
Me. 6/22/1988.

[67111] 5441.Phelan, M. C.; Pellock, J. M.; Nance, W. E.: Discordant  
expressionof fetal hydantoin syndrome in a pair of dizy-  
gotic twins with differentfathers. (Abstract) *Am. J. Hum.*  
*Genet.* 33: 67A only, 1981.

[67112] 5442.Sabry, M. A.; Farag, T. I.: Hand anomalies in fetal–

hydantoin syndrome: from nail/phalangeal hypoplasia to unilateral acheiria. (Letter) *Am. J. Med. Genet.* 62: 410–412, 1996.

- [67113] 5443. Skoda, R. C.; Demierre, A.; McBride, O. W.; Gonzalez, F. J.; Meyer, U. A.: Human microsomal xenobiotic epoxide hydrolase: complementary DNA sequence, complementary DNA-directed expression in COS-1 cells, and chromosomal localization. *J. Biol. Chem.* 263: 1549–1554, 1988.
- [67114] 5444. Spielberg, S. P.; Gordon, G. B.; Blake, D. A.; Goldstein, D. A.; Herlong, H. F.: Predisposition to phenytoin hepatotoxicity assessed in vitro. *New Eng. J. Med.* 305: 722–727, 1981.
- [67115] 5445. Strickler, S. M.; Dansky, L. V.; Miller, M. A.; Seni, M. H.; Andermann, E.; Spielberg, S. P.: Genetic predisposition to phenytoin-induced birth defects. *Lancet* II: 746–749, 1985.
- [67116] 5446. Vasko, M. R.; Bell, R. D.; Daly, D. D.: Inheritance of diphenylhydantoin hypometabolism: a pharmacokinetic study of one family. (Abstract) *Clin. Pharm. Therap.* 21: 120 only, 1977.
- [67117] 5447. Vasko, M. R.; Bell, R. D.; Daly, D. D.; Pippenger, C. E.: Inheritance of phenytoin hypometabolism: a kinetic study of one family. *Clin. Pharm. Therap.* 27: 96–103,

1980.

- [67118] 5448. Vermeij, P.; Ferrari, M. D.; Buruma, O. J. S.; Veenema, H.; deWolff, F. A.: Inheritance of poor phenytoin parahydroxylation capacity in a Dutch family. *Clin. Pharm. Therap.* 44: 588–593, 1988.
- [67119] 5449. Vesell, E. S.: Pharmacogenetics: multiple interactions between genes and environment as determinants of drug response. (Editorial) *Am. J. Med.* 66: 183–187, 1979.
- [67120] 5450. Zusterzeel, P. L. M.; Peters, W. H. M.; Visser, W.; Hermsen, K. J. M.; Roelofs, H. M. J.; Steegers, E. A. P.: A polymorphism in the gene for microsomal epoxide hydrolase is associated with pre-eclampsia. *J. Med. Genet.* 38: 234–237, 2001.
- [67121] 5451. Grant, D. F.; Spearow, J. L.; Storms, D. H.; Edelhoff, S.; Adler, D. A.; Disteché, C. M.; Taylor, B. A.; Hammock, B. D.: Chromosomal mapping and expression levels of a mouse soluble epoxide hydrolase gene. *Pharmacogenetics* 4: 64–72, 1994.
- [67122] 5452. Larsson, C.; White, I.; Johansson, C.; Stark, A.; Meijer, J.: Localization of the human soluble epoxide hydrolase gene (EPHX2) to chromosomal region 8p21–p12. *Hum. Genet.* 95: 356–358, 1995.
- [67123] 5453. Norris, K. K.; DeAngelo, T. M.; Vesell, E. S.: Genetic

and environmental factors that regulate cytosolic epoxide hydrolase activity in normal human lymphocytes. *J. Clin. Invest.* 84: 1749–1756, 1989.

[67124] 5454. Vesell, E. S.: Genetic factors that regulate cytosolic epoxide hydrolase activity in normal human lymphocytes. *Ann. Genet.* 34:167–172, 1991.

[67125] 5455. Fonatsch, C.; Latza, U.; Durkop, H.; Rieder, H.; Stein, H.: Assignment of the human CD30 (Ki-1) gene to 1p36. *Genomics* 14: 825–826, 1992.

[67126] 5456. Pauling, L.: The discovery of the superoxide radical. *Trends Biochem. Sci.* 4(11): 270–271, 1979.

[67127] 5457. Ma, T.; Yang, B.; Umenishi, F.; Verkman, A. S.: Closely spaced tandem arrangement of AQP2, AQP5, and AQP6 genes in a 27-kilobase segment at chromosome locus 12q13. *Genomics* 43: 387–389, 1997.

[67128] 5458. Benn, P.; Chern, C. J.; Bruns, G.; Craig, I. W.; Croce, C. M.: Assignment of the genes for human beta-glucuronidase and mitochondrial malate dehydrogenase to the region pter–q22 of chromosome 7. *Cytogenet. Cell Genet.* 19: 273–280, 1977.

[67129] 5459. Blake, N. M.: Malate dehydrogenase types in the Asian–Pacific area, and a description of new phenotypes. *Hum. Genet.* 43: 69–80, 1978.

- [67130] 5460. Davidson, R. G.; Cortner, J. A.: Mitochondrial malate dehydrogenase: a new genetic polymorphism in man. *Science* 157: 1569–1571, 1967.
- [67131] 5461. Shimizu, N.; Shimizu, Y.; Ruddle, F. H.: Assignment of the human mitochondrial NAD-linked malate dehydrogenase gene to the p22–qter region of chromosome 7. *Cytogenet. Cell Genet.* 22: 441–445, 1978.
- [67132] 5462. Van Heyningen, V.; Bobrow, M.; Bodmer, W. F.; Gardiner, S. E.; Povey, S.; Hopkinson, D. A.: Chromosome assignment of some human enzyme loci: mitochondrial malate dehydrogenase to 7, mannosephosphate isomerase and pyruvate kinase to 15 and probably, esterase D to 13. *Ann. Hum. Genet.* 38: 295–303, 1975.
- [67133] 5463. Larson, L. M.; Bruce, A. W.; Saumur, J. H.; Wasdahl, W. A.: Further evidence by gene dosage for the regional assignment of erythrocyte acid phosphatase (ACP1) and malate dehydrogenase (MDH1) loci on chromosome 2p. *Clin. Genet.* 22: 220–225, 1982.
- [67134] 5464. D'Andrea, R.; Harrison-Findik, D.; Butcher, C. M.; Finnie, J.; Blumbergs, P.; Bartley, P.; McCormack, M.; Jones, K.; Rowland, R.; Gonda, T. J.; Vadas, M. A.: Dysregulated hematopoiesis and a progressive neurological disorder induced by expression of an activated form of the human

common beta chain in transgenic mice. *J. Clin. Invest.* 102:1951–1960, 1998.

- [67135] 5465. Rosenblum, J. S.; Gilula, N. B.; Lerner, R. A.: On signal sequence polymorphisms and diseases of distribution. *Proc. Nat. Acad. Sci.* 93:4471–4473, 1996.
- [67136] 5466. Yoshimitsu, K.; Nichi, Y.; Kobayashi, Y.; Yoshimura, O.; Ohama, K.; Oguma, N.; Usui, T.: Decreased superoxide dismutase-2 activity in a patient with ring chromosome 6. *Am. J. Med. Genet.* 28: 211–214, 1987.
- [67137] 5467. Klugbauer, S.; Rabes, H. M.: The transcription coactivator HTIF1 and a related protein are fused to the RET receptor tyrosine kinase in childhood papillary thyroid carcinomas. *Oncogene* 18: 4388–4393, 1999.
- [67138] 5468. Lairmore, T. C.; Dou, S.; Howe, J. R.; Chi, D.; Carlson, K.; Veile, R.; Mishra, S. K.; Wells, S. A., Jr.; Donis-Keller, H.: A 1.5-megabase yeast artificial chromosome contig from human chromosome 10q11.2 connecting three genetic loci (RET, D10S94, and D10S102) closely linked to the MEN2A locus. *Proc. Nat. Acad. Sci.* 90: 492–496, 1993.
- [67139] 5469. Lombardo, F.; Baudin, E.; Chiefari, E.; Arturi, F.; Bardet, S.; Caillou, B.; Conte, C.; Dallapiccola, B.; Giuffrida, D.; Bidart, J.-M.; Schlumberger, M.; Filetti, S.: Familial



medullary thyroid carcinoma:clinical variability and low aggressiveness associated with RET mutationat codon 804. J. Clin. Endocr. Metab. 87: 1674–1680, 2002.

[67140] 5470.Lore, F.; Di Cairano, G.; Talidis, F.: Unilateral renal agenesisin a family with medullary thyroid carcinoma. (Letter) New Eng. J.Med. 342: 1218–1219, 2000.

[67141] 5471.Machens, A.; Gimm, O.; Hinze, R.; Hoppner, W.; Boehm, B. O.; Dralle,H.: Genotype–phenotype correlations in hereditary medullary thyroidcarcinoma: oncological features and biochemical properties. J. Clin.Endocr. Metab. 86: 1104–1109, 2001.

[67142] 5472.Mendelsohn, C.; et al; et al: Function of the retinoic acid receptors(RARs) during development (II). Multiple abnormalities at variousstages of organogenesis in RAR double mutants. Development 120:2749–2771, 1994.

[67143] 5473.Menko, F. H.; van der Luijt, R. B.; de Valk, I. A. J.; Toorians,A. W. F. T.; Sepers, J. M.; van Diest, P. J.; Lips, C. J. M.: AtypicalMEN type 2B associated with two germline RET mutations on the sameallele not involving codon 918. J. Clin. Endocr. Metab. 87: 393–397,2002.

[67144] 5474.Mulligan, L. M.; Kwok, J. B. J.; Healey, C. S.; Elsdon, M. J.;Eng, C.; Gardner, E.; Love, D. R.; Mole, S. E.; Moore, J. K.; Papi,L.; Ponder, M. A.; Telenius, H.; Tunnacliffe, A.;

Ponder, B. A. J.: Germ-line mutations of the RET proto-oncogene in multiple endocrineneoplasia type 2A. *Nature* 363: 458–460, 1993.

[67145] 5475. Munnes, M.; Fanaei, S.; Schmitz, B.; Muiznieks, I.; Holschneider, A. M.; Doerfler, W.: Familial form of Hirschsprung disease: nucleotide sequence studies reveal point mutations in the RET proto-oncogene in two of six families but not in other candidate genes. *Am. J. Med. Genet.* 94: 19–27, 2000.

[67146] 5476. Nakata, T.; Kitamura, Y.; Shimizu, K.; Tanaka, S.; Fujimori, M.; Yokoyama, S.; Ito, K.; Emi, M.: Fusion of a novel gene, ELKS, to RET due to translocation t(10;12)(q11;p13) in a papillary thyroid carcinoma. *Genes Chromosomes Cancer* 25: 97–103, 1999.

[67147] 5477. Niccoli-Sire, P.; Murat, A.; Rohmer, V.; Franc, S.; Chabrier, G.; Baldet, L.; Maes, B.; Savagner, F.; Giraud, S.; Bezieau, S.; Kottler, M.-L.; Morange, S.; Conte-Devolx, B.: The French Calcitonin Tumors Study Group (GETC): Familial medullary thyroid carcinoma with noncysteine RET mutations: phenotype-genotype relationship in a large series of patients. *J. Clin. Endocr. Metab.* 86: 3746–3753, 2001.

[67148] 5478. Pachnis, V.; Mankoo, B.; Costantini, F.: Expression of the c-ret proto-oncogene during mouse embryogenesis.

Development 119: 1005–1017,1993.

- [67149] 5479.Pasini, B.; Hofstra, R. M. W.; Yin, L.; Bocciardi, R.; Santamaria,G.; Grootsholten, P. M.; Ceccherini, I.; Patrone, G.; Priolo, M.;Buys, C. H. C. M.; Romeo, G.: The physical map of the human RET proto-oncogene. *Oncogene* 11:1737–1743, 1995.
- [67150] 5480.Pelet, A.; Geneste, O.; Edery, P.; Pasini, A.; Chappuis, S.; Attie,T.; Munnich, A.; Lenoir, G.; Lyonnet, S.; Billaud, M.: Various mechanismscause RET-mediated signaling defects in Hirschsprung's disease. *J.Clin. Invest.* 101: 1415–1423, 1998.
- [67151] 5481.Pigny, P.; Bauters, C.; Wemeau, J.–L.; Houcke, M. L.; Crepin,M.; Caron, P.; Giraud, S.; Calender, A.; Buisine, M.–P.; Kerckaert,J.–P.; Porchet, N.: A novel 9–base pair duplication in RET exon 8in familial medullary thyroid carcinoma. *J. Clin. Endocr. Metab.* 84:1700–1704, 1999.
- [67152] 5482.Pierotti, M. A.; Santoro, M.; Jenkins, R. B.; Sozzi, G.; Bongarzone,I.; Grieco, M.; Monzini, N.; Miozzo, M.; Hermann, M. A.; Fusco, A.;Hay, I. D.; Della Porta, G.; Vecchio, G.: Characterization of aninversion on the long arm of chromosome 10 juxtaposing D10S170 andRET and creating the oncogenic sequence RET/PTC. *Proc. Nat. Acad.Sci.* 89: 1616–1620, 1992.

- [67153] 5483.Rodrigues, G. A.; Park, M.: Dimerization mediated through a leucinezipper activates the oncogenic potential of the met receptor tyrosinekinase. *Molec. Cell. Biol.* 13: 6711–6722, 1993.
- [67154] 5484.Romeo, G.; Ronchetto, P.; Luo, Y.; Barone, V.; Seri, M.; Ceccherini,I.; Pasini, B.; Bocciardi, R.; Lerone, M.; Kaariainen, H.; Martucciello,G.: Point mutations affecting the tyrosine kinase domain of the RETproto–oncogene in Hirschsprung's disease. *Nature* 367: 377–378, 1994.
- [67155] 5485.Salvatore, D.; Barone, M. V.; Salvatore, G.; Melillo, R. M.; Chiappetta,G.; Mineo, A.; Fenzi, G.; Vecchio, G.; Fusco, A.; Santoro, M.: Tyrosines1015 and 1062 are in vivo autophosphorylation sites in Ret and Ret–derivedoncoproteins. *J. Clin. Endocr. Metab.* 85: 3898–3907, 2000.
- [67156] 5486.Robinson, M. F.; Cote, G. J.; Nunziata, V.; Brandi, M. L.; Ferrer,J. P.; Martins Bugalho, M. J. G.; Almeida Ruas, M. M.; Chik, C.; Colantuoni,V.; Gagel, R. F.: Mutation of a specific codon of the RET proto–oncogenein the multiple endocrine neoplasia type 2A/cutaneous lichen amyloidosis syndrome. (Abstract) Fifth International Workshop on Multiple EndocrineNeoplasia, Stockholm, Archipelago , 1994.

- [67157] 5487.Boyer, S. H.: Human organ alkaline phosphatases: discrimination by several means including starch gel electrophoresis of antienzyme-enzymesupernatant fluids. Ann. N.Y. Acad. Sci. 103: 938-950, 1963.
- [67158] 5488.Deutman, A. F.; van Blommestein, J. D. A.; Henkes, H. E.; Waardenburg, P. J.; Solleveld-van Driest, E.: Butterfly-shaped pigment dystrophy of the fovea. Arch. Ophthalmol. 83: 558-569, 1970.
- [67159] 5489.Okuda, T.; Cleveland, J. L.; Downing, J. R.: PCTAIRE-1 and PCTAIRE-3, two members of a novel cdc2/CDC28-related protein kinase gene family. Oncogene 7:2249-2258, 1992.
- [67160] 5490.Okuda, T.; Valentine, V. A.; Shapiro, D. N.; Downing, J. R.: Cloning of genomic loci and chromosomal localization of the human PCTAIRE-1 and -3 protein kinase genes. Genomics 21: 217-221, 1994.
- [67161] 5491.Donald, L. J.: The genetics of placental alkaline phosphatase: a possible 'null' allele. Ann. Hum. Genet. 38: 7-18, 1974.
- [67162] 5492.Donald, L. J.; Robson, E. B.: Rare variants of placental alkaline phosphatase. Ann. Hum. Genet. 37: 303-313, 1974.
- [67163] 5493.Edwards, J. H.; Wingham, J.: Data on linkage between

the locus determining placental alkaline phosphatase and other markers. *Ann. Hum. Genet.* 30: 233–237, 1967.

[67164] 5494. Garattini, E.; Margolis, J.; Heimer, E.; Felix, A.; Udenfriend, S.: Human placental alkaline phosphatase in liver and intestine. *Proc. Nat. Acad. Sci.* 82: 6080–6084, 1985.

[67165] 5495. Gogolin, K. J.; Slaughter, C. A.; Harris, H.: Electrophoresis of enzyme–monoclonal antibody complexes: studies of human placental alkaline phosphatase polymorphism. *Proc. Nat. Acad. Sci.* 78: 5061–5065, 1981.

[67166] 5496. Henthorn, P. S.; Knoll, B. J.; Raducha, M.; Rothblum, K. N.; Slaughter, C.; Weiss, M.; Lafferty, M. A.; Fischer, T.; Harris, H.: Products of two common alleles at the locus for human placental alkaline phosphatase differ by seven amino acids. *Proc. Nat. Acad. Sci.* 83: 5597–5601, 1986.

[67167] 5497. Kam, W.; Clauser, E.; Kim, Y. S.; Kan, Y. W.; Rutter, W. J.: Cloning, sequencing, and chromosomal localization of human term placental alkaline phosphatase cDNA. *Proc. Nat. Acad. Sci.* 82: 8715–8719, 1985.

[67168] 5498. Knoll, B. J.; Rothblum, K. N.; Longley, M.: Nucleotide sequence of the human placental alkaline phosphatase gene: evolution of the 5–prime flanking region by deletion/substitution. *J. Biol. Chem.* 263: 12020–12027, 1988.

[67169] 5499. Lucarelli, P.; Scacchi, R.; Corbo, R. M.; Benincasa, A.;

Palmarino,R.: Human placental alkaline phosphatase electrophoretic alleles:quantitative studies. Am. J. Hum. Genet. 34: 331–336, 1982.

[67170] 5500.Martin, D.; Spurr, N. K.; Trowsdale, J.: RFLP of the human placentalalkaline phosphatase gene (PLAP). Nucleic Acids Res. 15: 9104 only,1987.

[67171] 5501.Martin, D.; Tucker, D. F.; Gorman, P.; Sheer, D.; Spurr, N. K.;Trowsdale, J.: The human placental alkaline phosphatase gene andrelated sequences map to chromosome 2 band q37. Ann. Hum. Genet. 51:145–152, 1987.

[67172] 5502.Millan, J. L.; Beckman, G.; Jeppsson, A.; Stigbrand, T.: Geneticvariants of placental alkaline phosphatase as detected by a monoclonalantibody. Hum. Genet. 60: 145–149, 1982.

[67173] 5503.Millan, J. L.; Stigbrand, T.: Antigenic determinants of humanplacental and testicular placental–like alkaline phosphatases as mappedby monoclonal antibodies. Europ. J. Biochem. 136: 1–7, 1983.

[67174] 5504.Palmarino, R.; Corbo, R. M.; Lucarelli, P.: Human placental alkalinephosphatase: analysis of genetically determined rare variants. Hum.Biol. 51: 341–352, 1979.

[67175] 5505.Raimondi, E.; Talarico, D.; Moro, L.; Rutter, W. J.; Della Valle,G.; De Carli, L.: Regional mapping of the hu–

man placental alkalinephosphatase gene (ALPP) to 2q37 by in situ hybridization. *Cytogenet.Cell Genet.* 47: 98–99, 1988.

[67176] 5506.Robinson, J. C.; Goldsmith, L. A.: Genetically determined variants of serum alkaline phosphatase: a review. *Vox Sang.* 13: 289–307, 1967.

[67177] 5507.Robson, E. B.; Harris, H.: Genetics of the alkaline phosphatase polymorphism of the human placenta. *Nature* 207: 1257–1259, 1965.

[67178] 5508.Slaughter, C. A.; Gogolin, K. J.; Coseo, M. C.; Meyer, L. J.; Lesko, J.; Harris, H.: Discrimination of human placental alkalinephosphatase allelic variants by monoclonal antibodies. *Am. J. Hum.Genet.* 35: 1–20, 1983.

[67179] 5509.Almind, K.; Delahaye, L.; Hansen, T.; Van Obberghen, E.; Pedersen, O.; Kahn, C. R.: Characterization of the Met326Ile variant of phosphatidylinositol3–kinase p85–alpha. *Proc. Nat. Acad. Sci.* 99: 2124–2128, 2002.

[67180] 5510.Cannizzaro, L. A.; Skolnik, E. Y.; Margolis, B.; Croce, C. M.; Schlesinger, J.; Huebner, K.: The human gene encoding phosphatidylinositol3–kinase associated p85–alpha is at chromosome region 5q12–13. *CancerRes.* 51: 3818–3820, 1991.

[67181] 5511.Vagnarelli, P.; Raimondi, E.; Mazzieri, R.; De Carli, L.;



Mignatti,P.: Assignment of the human urokinase receptor gene (PLAUR) to 19q13. Cytogenet.Cell Genet. 60: 197–199, 1992.

[67182] 5512.Chase, P. B.; Yang, J.–M.; Thompson, F. H.; Halonen, M.; Regan,J. W.: Regional mapping of the human platelet–activating factor receptorgene (PTAFR) to 1p35–p34.3 by fluorescence in situ hybridization. Cytogenet.Cell Genet. 72: 205–207, 1996.

[67183] 5513.Prescott, S. M.; Zimmerman, G. A.; McIntyre, T. M.: Platelet–activatingfactor. J. Biol. Chem. 265: 17381–17384, 1990.

[67184] 5514.Seyfried, C. E.; Schweickart, V. L.; Godiska, R.; Gray, P. W.:The human platelet–activating factor receptor gene (PTAFR) containsno introns and maps to chromosome 1. Genomics 13: 832–834, 1992.

[67185] 5515.Abe, A.; Emi, N.; Tanimoto, M.; Terasaki, H.; Marunouchi, T.; Saito,H.: Fusion of the platelet–derived growth factor receptor beta to a novel gene CEV14 in acute myelogenous leukemia after clonal evolution. Blood 90:4271–4277, 1997.

[67186] 5516.Buchberg, A. M.; Jenkins, N. A.; Copeland, N. G.: Localizationof the murine macrophage colony–stimulating factor gene to chromosome3 using interspecific backcross

analysis. Genomics 5: 363–367, 1989.

- [67187] 5517.Claesson–Welsh, L.; Eriksson, A.; Moren, A.; Severinsson, L.; Ek,B.; Ostman, A.; Betsholtz, C.; Heldin, C.–H.: cDNA cloning and expression of a human platelet–derived growth factor (PDGF) receptor specific for B–chain–containing PDGF molecules. Molec. Cell. Biol. 8: 3476–3486,1988.
- [67188] 5518.Escobedo, J. A.; Fried, V. A.; Daniel, T. O.; Williams, L. T.:Primary structure of the platelet–derived growth factor. (Abstract) Clin.Res. 34: 544A, 1986.
- [67189] 5519.Gronwald, R. G. K.; Grant, F. J.; Haldeman, B. A.; Hart, C. E.;O'Hara, P. J.; Hagen, F. S.; Ross, R.; Bowen–Pope, D. F.; Murray,M. J.: Cloning and expression of a cDNA coding for the human platelet–derived growth factor receptor: evidence for more than one receptor class. Proc.Nat. Acad. Sci. 85: 3435–3439, 1988.
- [67190] 5520.Klinghoffer, R. A.; Mueting–Nelsen, P. F.; Faerman, A.; Shani,M.; Soriano, P.: The two PDGF receptors maintain conserved signaling in vivo despite divergent embryological functions. Molec. Cell 7:343–354, 2001.
- [67191] 5521.Kulkarni, S.; Heath, C.; Parker, S.; Chase, A.; Iqbal, S.; Pocock,C. F.; Kaeda, J.; Cwynarski, K.; Goldman, J. M.; Cross, N. C. P.:Fusion of H4/D10S170 to the platelet–de–

rived growth factor receptorbeta in BCR-ABL-negative myeloproliferative disorders with a t(5;10)(q33;q21). CancerRes. 60: 3592-3598, 2000.

- [67192] 5522. Leal, F.; Williams, L. T.; Robbins, K. C.; Aaronson, S. A.: Evidence that the v-sis gene product transforms by interaction with the receptor for platelet-derived growth factor. Science 230: 327-330, 1985.
- [67193] 5523. Magnusson, M. K.; Meade, K. E.; Brown, K. E., Arthur, D. C.; Krueger, L. A.; Barrett, A. J.; Dunbar, C. E.: Rabaptin-5 is a novel fusion partner to platelet-derived growth factor beta receptor in chronic myelomonocytic leukemia. Blood 98: 2518-2525, 2001.
- [67194] 5524. Matsui, T.; Heidaran, M.; Miki, T.; Popescu, N.; La Rochelle, W.; Kraus, M.; Pierce, J.; Aaronson, S.: Isolation of a novel receptor cDNA establishes the existence of two PDGF receptor genes. Science 243:800-804, 1989.
- [67195] 5525. Ross, T. S.; Bernard, O. A.; Berger, R.; Gilliland, D. G.: Fusion of Huntingtin interacting protein 1 to platelet-derived growth factor-beta receptor (PDGF-beta-R) in chronic myelomonocytic leukemia with t(5;7)(q33;q11.2). Blood 91:4419-4426, 1998.
- [67196] 5526. Steer, E. J.; Cross, N. C. P.: Myeloproliferative disorders with translocations of chromosome 5q31-35: role of

the platelet-derived growth factor receptor beta. *Acta Haemat.* 107: 113–122, 2002.

[67197] 5527. Treacher Collins Syndrome Collaborative Group: Positional cloning of a gene involved in the pathogenesis of Treacher Collins syndrome. *Nature Genet.* 12: 130–136, 1996.

[67198] 5528. Yarden, Y.; Escobedo, J. A.; Kuang, W.-J.; Yang-Feng, T. L.; Daniel, T. O.; Tremble, P. M.; Chen, E. Y.; Ando, M. E.; Harkins, R. N.; Francke, U.; Fried, V. A.; Ullrich, A.; Williams, L. T.: Structure of the receptor for platelet-derived growth factor helps define a family of closely related growth factor receptors. *Nature* 323: 226–232, 1986.

[67199] 5529. Berthet, M.; Denjoy, I.; Donger, C.; Demay, L.; Hamoude, H.; Klug, D.; Schulze-Bahr, E.; Richard, P.; Funke, H.; Schwartz, K.; Coumel, P.; Hainque, B.; Guicheney, P.: C-terminal HERG mutations: the role of hypokalemia and a KCNQ1-associated mutation in cardiac event occurrence. *Circulation* 99:1464–1470, 1999.

[67200] 5530. Allcock, R. J. N.; Williams, J. H.; Price, P.: The central MHC gene, BAT1, may encode a protein that down-regulates cytokine production. *Genes Cells* 6: 487–494, 2001.

[67201] 5531. Nunes, M.; Peelman, L.; Vaiman, M.; Bourgeaux, N.;

Chardon, P.: Characterization of six new loci within the swine major histocompatibility complex class III region. *Mammalian Genome* 5: 616–622, 1994.

[67202] 5532. Peelman, L. J.; Chardon, P.; Nunes, M.; Renard, C.; Geffrotin, C.; Vaiman, M.; Van Zeveren, A.; Coppieters, W.; van de Weghe, A.; Bouquet, Y.; Choy, W. W.; Strominger, J. L.; Spies, T.: The BAT1 gene in the MHC encodes an evolutionarily conserved putative nuclear RNA helicase of the DEAD family. *Genomics* 26: 210–218, 1995.

[67203] 5533. Spies, T.; Blanck, G.; Bresnahan, M.; Sands, J.; Strominger, J. L.: A new cluster of genes within the human major histocompatibility complex. *Science* 243: 214–217, 1989.

[67204] 5534. Spies, T.; Bresnahan, M.; Strominger, J. L.: Human major histocompatibility complex contains a minimum of 19 genes between the complement cluster and HLA-B. *Proc. Nat. Acad. Sci.* 86: 8955–8958, 1989.

[67205] 5535. Colosimo, A.; Calabrese, G.; Gennarelli, M.; Ruzzo, A. M.; Sangiuolo, F.; Magnani, M.; Palka, G.; Novelli, G.; Dallapiccola, B.: Assignment of the hexokinase type 3 gene (HK3) to human chromosome band 5q35.3 by somatic cell hybrids and in situ hybridization. *Cytogenet. Cell Genet.* 74: 187–188, 1996.

- [67206] 5536.Furuta, H.; Nishi, S.; Le Beau, M. M.; Fernald, A. A.; Yano, H.;Bell, G. I.: Sequence of human hexokinase III cDNA and assignmentof the human hexokinase III gene (HK3) to chromosome band 5q35.2 byfluorescence in situ hybridization. *Genomics* 36: 206–209, 1996.
- [67207] 5537.Harris, H.; Hopkinson, D. A.: Average heterozygosity per locusin man: an estimate based on the incidence of enzyme polymorphism. *Ann.Hum. Genet.* 36: 9–20, 1972.
- [67208] 5538.Hopkinson, D. A.; Edwards, Y. H.; Harris, H.: The distributionof subunit numbers and subunit sizes of enzymes: a study of the productsof 100 gene loci. *Ann. Hum. Genet.* 39: 383–411, 1976.
- [67209] 5539.Povey, S.; Corney, G.; Harris, H.: Genetically determined polymorphismin a form of hexokinase, HK III, found in human leukocytes. *Ann.Hum. Genet.* 38: 407–415, 1975.
- [67210] 5540.Buhr, A.; Bianchi, M. T.; Baur, R.; Courtet, P.; Pignay, V.; Boulenger,J. P.; Gallati, S.; Hinkle, D. J.; Macdonald, R. L.; Sigel, E.: Functionalcharacterization of the new human GABA(A) receptor mutation beta–3(R192H). *Hum.Genet.* 111: 154–160, 2002.
- [67211] 5541.Buxbaum, J. D.; Silverman, J. M.; Smith, C. J.; Greenberg, D. A.;Kilifarski, M.; Reichert, J.; Cook, E. H., Jr.; Fang,

Y.; Song, C.-Y.; Vitale, R.: Association between a GABRB3 polymorphism and autism. *Molec. Psychiat.* 7: 311–316, 2002.

[67212] 5542. Holopainen, I. E.; Metsähonkala, E.-L.; Kokkonen, H.; Parkkola, R. K.; Manner, T. E.; Nagren, K.; Korpi, E. R.: Decreased binding of [11C]flumazenil in Angelman syndrome patients with GABA-A receptor beta-3 subunit deletions. *Ann. Neurol.* 49: 110–113, 2001.

[67213] 5543. Homanics, G. E.; DeLorey, T. M.; Firestone, L. L.; Quinlan, J. J.; Handforth, A.; Harrison, N. L.; Krasowski, M. D.; Rick, C. E. M.; Korpi, E. R.; Makela, R.; Brilliant, M. H.; Hagiwara, N.; Ferguson, C.; Snyder, K.; Olsen, R. W.: Mice devoid of gamma-aminobutyrate type A receptor beta3 subunit have epilepsy, cleft palate, and hypersensitive behavior. *Proc. Nat. Acad. Sci.* 94: 4143–4148, 1997.

[67214] 5544. Knoll, J. H. M.; Cheng, S.-D.; Lalande, M.: Allele specificity of DNA replication timing in the Angelman/Prader-Willi syndrome imprinted chromosomal region. *Nature Genet.* 6: 41–46, 1994.

[67215] 5545. Laposky, A. D.; Homanics, G. E.; Baile, A.; Mendelson, W. B.: Deletion of the GABA(A) receptor beta 3 subunit eliminates the hypnotic actions of oleamide in mice. *Neuroreport* 12: 4143–4147, 2001.

- [67216] 5546. Meguro, M.; Mitsuya, K.; Sui, H.; Shigenami, K.; Kugoh, H.; Nakao, M.; Oshimura, M.: Evidence for uniparental, paternal expression of the human GABA-A receptor subunit genes, using microcell-mediated chromosome transfer. *Hum. Molec. Genet.* 6: 2127–2133, 1997.
- [67217] 5547. Saitoh, S.; Kubota, T.; Ohta, T.; Jinno, Y.; Niikawa, N.; Sugimoto, T.; Wagstaff, J.; Lalande, M.: Familial Angelman syndrome caused by imprinted submicroscopic deletion encompassing GABA(A) receptor beta(3)-subunit gene. (Letter) *Lancet* 339: 366–367, 1992.
- [67218] 5548. Scapoli, L.; Martinelli, M.; Pezzetti, F.; Carinci, F.; Bodo, M.; Tognon, M.; Carinci, P.: Linkage disequilibrium between GABRB3 gene and nonsyndromic familial cleft lip with or without cleft palate. *Hum. Genet.* 110: 15–20, 2002.
- [67219] 5549. Sinnott, D.; Wagstaff, J.; Glatt, K.; Woolf, E.; Kirkness, E.J.; Lalande, M.: High-resolution mapping of the gamma-aminobutyric acid receptor subunit beta-3 and alpha-5 gene cluster on chromosome 15q11–q13, and localization of breakpoints in two Angelman syndrome patients. *Am. J. Hum. Genet.* 52: 1216–1229, 1993.
- [67220] 5550. Tanabe, A.; Taketani, S.; Endo-Ichikawa, Y.; Toku-



naga, R.; Ogawa,Y.; Hiramoto, M.: Analysis of the candidate genes responsible for non-syndromic cleft lip and palate in Japanese people. Clin. Sci. 99:105–111, 2000.

[67221] 5551.Wagstaff, J.; Chaillet, J. R.; Lalande, M.: The GABA(A) receptor beta-3 subunit gene: characterization of a human cDNA from chromosome 15q11q13 and mapping to a region of conserved synteny on mouse chromosome 7. Genomics 11: 1071–1078, 1991.

[67222] 5552.Wagstaff, J.; Knoll, J. H. M.; Fleming, J.; Kirkness, E. F.; Martin-Gallardo,A.; Greenberg, F.; Graham, J. M., Jr.; Menninger, J.; Ward, D.; Venter,J. C.; Lalande, M.: Localization of the gene encoding the GABA(A)receptor beta-3 subunit to the Angelman/Prader-Willi region of human chromosome 15. Am. J. Hum. Genet. 49: 330–337, 1991.

[67223] 5553.Seeburg, P. H.: The human growth hormone gene family: nucleotide sequences show recent divergence and predict a new polypeptide hormone. DNA 1:239–249, 1982.

[67224] 5554.Schneider, P. M.; Witzel-Schlomp, K.; Rittner, C.; Zhang, L.:The endogenous retroviral insertion in the human complement C4 gene modulates the expression of homologous genes by antisense inhibition. Immunogenet–

ics 53:1–9, 2001.

- [67225] 5555.Suto, Y.; Tokunaga, K.; Watanabe, Y.; Hirai, M.: Visual demonstration of the organization of the human complement C4 and 21-hydroxylase genes by high-resolution fluorescence in situ hybridization. *Genomics* 33:321–324, 1996.
- [67226] 5556.Teisberg, P.; Akesson, I.; Olaisen, B.; Gedde-Dahl, T., Jr.; Thorsby, E.: Genetic polymorphism of C4 in man and localization of a structural C4 locus to the HLA gene complex of chromosome 6. *Nature* 264: 253–254, 1976.
- [67227] 5557.Boerkoel, C. F.; Takashima, H.; Stankiewicz, P.; Garcia, C. A.; Leber, S. M.; Rhee-Morris, L.; Lupski, J. R.: Periaxin mutations cause recessive Dejerine-Sottas neuropathy. *Am. J. Hum. Genet.* 68:325–333, 2001. Note: Erratum: *Am. J. Hum. Genet.* 68: 557 only, 2001.
- [67228] 5558.Pischel, K. D.; Marlin, S. D.; Springer, T. A.; Woods, V. L., Jr.; Bluestein, H. G.: Polymorphism of lymphocyte function-associated antigen-1 demonstrated by a lupus patient's alloantiserum. *J. Clin. Invest.* 79: 1607–1614, 1987.
- [67229] 5559.Akey, J. M.; Wang, H.; Xiong, M.; Wu, H.; Liu, W.; Shriver, M.D.; Jin, L.: Interaction between the melanocortin-1 receptor and P genes contributes to inter-

individual variation in skin pigmentation phenotypes in a Tibetan population. Hum. Genet. 108: 516–520, 2001.

- [67230] 5560. Chottiner, E. G.; Shewach, D. S.; Datta, N. S.; Ashcraft, E.; Gribbin, D.; Ginsburg, D.; Fox, I. H.; Mitchell, B. S.: Cloning and expression of human deoxycytidine kinase cDNA. Proc. Nat. Acad. Sci. 88: 1531–1535, 1991.
- [67231] 5561. Huang, S.-H.; Tomich, J. M.; Wu, H.; Jong, A.; Holcenberg, J.: Human deoxycytidine kinase: sequence of cDNA clones and analysis of expression in cell lines with and without enzyme activity. J. Biol. Chem. 264: 14762–14768, 1989. Note: Correction: J. Biol. Chem. 266:5353 only, 1991.
- [67232] 5562. Johansson, M.; Brismar, S.; Karlsson, A.: Human deoxycytidine kinase is located in the cell nucleus. Proc. Nat. Acad. Sci. 94:11941–11945, 1997.
- [67233] 5563. Song, J. J.; Walker, S.; Chen, E.; Johnson, E. E., II; Spychala, J.; Gribbin, T.; Mitchell, B. S.: Genomic structure and chromosomal localization of the human deoxycytidine kinase gene. Proc. Nat. Acad. Sci. 90: 431–434, 1993.
- [67234] 5564. Nurnberg, P.; Thiele, H.; Chandler, D.; Hohne, W.; Cunningham, M. L.; Ritter, H.; Leschik, G.; Uhlmann, K.; Mischung, C.; Harrop, K.; Goldblatt, J.; Borochowitz, Z. U.; Kotzot, D.; Westermann, F.; Mundlos, S.; Braun, H.-S.;

Laing, N.; Tinschert, S.: Heterozygous mutations in ANKH, the human ortholog of the mouse progressive ankylosis-gene, result in craniometaphyseal dysplasia. *Nature Genet.* 28: 37–41, 2001.

[67235] 5565. Nurnberg, P.; Tinschert, S.; Mrug, M.; Hampe, J.; Muller, C. R.; Fuhrmann, E.; Braun, H.-S.; Reis, A.: The gene for autosomal dominant craniometaphyseal dysplasia maps to chromosome 5q and is distinct from the growth hormone-receptor gene. *Am. J. Hum. Genet.* 61: 918–923, 1997.

[67236] 5566. Soriano, P.; Montgomery, C.; Geske, R.; Bradley, A.: Targeted disruption of the c-src proto-oncogene leads to osteopetrosis in mice. *Cell* 64:693–702, 1991.

[67237] 5567. Stegmann, A. P. A.; Honders, M. W.; Bolk, M. W. J.; Wessels, J.; Willemze, R.; Landegent, J. E.: Assignment of the human deoxycytidine kinase (DCK) gene to chromosome 4 band q13.3–q21.1. *Genomics* 17:528–529, 1993.

[67238] 5568. Sanchez-Madrid, F.; Nagy, J.; Robbins, E.; Simon, P.; Springer, T. A.: A human leukocyte differentiation antigen family with distinct alpha subunits and a common beta subunit: the lymphocyte function-associated antigen (LFA-1), the C3bi complement receptor (OKM1/Mac-1), and the p150,95 molecule. *J. Exp. Med.* 158: 1785–1803,

1983.

- [67239] 5569.Saint-Ruf, C.; Panigada, M.; Azogui, O.; Debey, P.; von Boehmer, H.; Grassi, F.: Different initiation of pre-TCR and gamma-delta-TCR signalling. *Nature* 406: 524–527, 2000.
- [67240] 5570.Delague, V.; Bareil, C.; Tuffery, S.; Bouvagnet, P.; Chouery, E.; Koussa, S.; Maisonobe, T.; Loiselet, J.; Megarbane, A.; Claustres, M.: Mapping of a new locus for autosomal recessive demyelinating Charcot-Marie-Tooth disease to 19q13.1–13.3 in a large consanguineous Lebanese family: exclusion of MAG as a candidate gene. *Am. J. Hum. Genet.* 67: 236–243, 2000.
- [67241] 5571.Guilbot, A.; Williams, A.; Ravise, N.; Verny, C.; Brice, A.; Sherman, D. L.; Brophy, P. J.; LeGuern, E.; Delague, V.; Bareil, C.; Megarbane, A.; Claustres, M.: A mutation in periaxin is responsible for CMT4F, an autosomal recessive form of Charcot-Marie-Tooth disease. *Hum. Molec. Genet.* 10: 415–421, 2001.
- [67242] 5572.Sugino, N.; Suzuki, T.; Kashida, S.; Karube, A.; Takiguchi, S.; Kato, H.: Expression of Bcl-2 and Bax in the human corpus luteum during the menstrual cycle in early pregnancy: regulation by human chorionic gonadotropin. *J. Clin. Endocr. Metab.* 85: 4379–4386, 2000.

- [67243] 5573.Vaskivuo, T. E.; Anttonen, M.; Herva, R.; Billig, H.; Dorland,M.; Te Velde, E. R.; Stenback, F.; Heikinheimo, M.; Tapanainen, J.S.: Survival of human ovarian follicles from fetal to adult life:apoptosis, apoptosis-related proteins, and transcription factor GATA-4. J.Clin. Endocr. Metab. 86: 3421-3429, 2001.
- [67244] 5574.Cleary, M. L.; Mellentin, J. D.; Spies, J.; Smith, S. D.: Chromosomaltranslocation involving the beta T cell receptor gene in acute leukemia.J. Exp. Med. 167: 682-687, 1988.
- [67245] 5575.Kuo, S. S.; Mellentin, J. D.; Copeland, N. G.; Gilbert, D. J.;Jenkins, N. A.; Cleary, M. L.: Structure, chromosome mapping, andexpression of the mouse Lyl-1 gene. Onco-gene 6: 961-968, 1991.
- [67246] 5576.Inoguchi, K.; Yoshioka, H.; Khaleduzzaman, M.; Ni-nomiya, Y.: ThemRNA for alpha-1(XIX) collagen chain, a new member of FACITs, containsa long unusual 3-prime untranslated region and displays many uniquesplicing variants. J. Biochem. 117: 137-146, 1995.
- [67247] 5577.Khaleduzzaman, M.; Sumiyoshi, H.; Ueki, Y.; Inoguchi, K.; Ninomiya,Y.; Yoshioka, H.: Structure of the hu-man type XIX collagen (COL19A1)gene, which suggests it has arisen from an ancestor gene of the FACITfamily. Ge-

nomics 45: 304–312, 1997.

[67248] 5578. Myers, J. C.; Sun, M. J.; D'Ippolito, J. A.; Jabs, E. W.; Neilson, E. G.; Dion, A. S.: Human cDNA clones transcribed from an unusually high molecular weight RNA encode a new collagen chain. *Gene* 123:211–217, 1993.

[67249] 5579. Yoshioka, H.; Zhang, H.; Ramirez, F.; Mattei, M.-G.; Moradi-Ameli, M.; van der Rest, M.; Gordon, M. K.: Synteny between the loci for a novel FACIT-like collagen (D6S228E) and alpha 1(IX) collagen (COL9A1) on 6q12–q14 in humans. *Genomics* 13: 884–886, 1992.

[67250] 5580. Gerecke, D. R.; Olson, P. F.; Koch, M.; Knoll, J. H. M.; Taylor, R.; Hudson, D. L.; Champlaud, M.-F.; Olsen, B. R.; Burgeson, R. E.: Complete primary structure of two splice variants of collagen XII, and assignment of alpha-1(XII) collagen (COL12A1), alpha-1(IX) collagen (COL9A1), and alpha-1(XIX) collagen (COL19A1) to human chromosome 6q12–q13. *Genomics* 41: 236–242, 1997.

[67251] 5581. Copeland, N. G.; Jenkins, N. A.; Gilbert, D. J.; Eppig, J. T.; Maltais, L. J.; Miller, J. C.; Dietrich, W. F.; Weaver, A.; Lincoln, S. E.; Steen, R. G.; Stein, L. D.; Nadeau, J. H.; Lander, E. S.: A genetic linkage map of the mouse: current applications and future prospects. *Science* 262: 57–66, 1993.

[67252] 5582. Tong, Q.; Dalgin, G.; Xu, H.; Ting, C.-N.; Leiden, J.

M.; Hotamisligil, G. S.: Function of GATA transcription factors in preadipocyte–adipocyte transition. *Science* 290: 134–138, 2000.

[67253] 5583. Litt, M.; Buroker, N. E.; Kondoleon, S.; Douglass, J.; Liston, D.; Sheehy, R.; Magenis, R. E.: Chromosomal localization of the human proenkephalin and prodynorphin genes. *Am. J. Hum. Genet.* 42: 327–334, 1988.

[67254] 5584. Litt, M.; Buroker, N. E.; Kondoleon, S. K.; Liston, D.; Douglass, J.; Sheehy, R.; Magenis, R. E.: Chromosomal localization of the human proenkephalin and prodynorphin genes. (Abstract) *Cytogenet. Cell Genet.* 46: 651 only, 1987.

[67255] 5585. Berkovic, S. F.; McIntosh, A.; Howell, R. A.; Mitchell, A.; Sheffield, L. J.; Hopper, J. L.: Familial temporal lobe epilepsy: a common disorder identified in twins. *Ann. Neurol.* 40: 227–235, 1996.

[67256] 5586. Cendes, F.; Lopes-Cendes, I.; Andermann, E.; Andermann, F.: Familial temporal lobe epilepsy: a clinically heterogeneous syndrome. *Neurology* 50: 554–557, 1998.

[67257] 5587. Horikawa, S.; Takai, T.; Toyosato, M.; Takahashi, H.; Noda, M.; Kakidani, H.; Kubo, T.; Hirose, T.; Inayama, S.; Hayashida, H.; Miyata, T.; Numa, S.: Isolation and structural organization of the human preproenkephalin B gene.



Nature 306: 611–614, 1983.

[67258] 5588.Grigorenko, E. L.; Wood, F. B.; Meyer, M. S.; Hart, L. A.; Speed, W. C.; Shuster, A.; Pauls, D. L.: Susceptibility loci for distinct components of developmental dyslexia on chromosomes 6 and 15. Am.J. Hum. Genet. 60: 27–39, 1997.

[67259] 5589.Giuli, G.; Scholl, U.; Bulle, F.; Guellaen, G.: Molecular cloning of the cDNAs coding for the two subunits of soluble guanylyl cyclase from human brain. FEBS Lett. 304: 83–88, 1992.

[67260] 5590.Feder, J. N.; Li, L.; Jan, L. Y.; Jan, Y. N.: Genomic cloning and chromosomal localization of HRY, the human homolog of the Drosophila segmentation gene, hairy. Genomics 20: 56–61, 1994.

[67261] 5591.Votruba, M.; Payne, A.; Moore, A. T.; Bhattacharya, S. S.: Dominant optic atrophy: exclusion and fine genetic mapping of the candidate gene, HRY. Mammalian Genome 9: 784–787, 1998.

[67262] 5592.Yan, B.; Heus, J.; Lu, N.; Nichols, R. C.; Raben, N.; Plotz, P.H.: Transcriptional regulation of the human acid alpha-glucosidase gene: identification of a repressor element and its transcription factors Hes-1 and YY1. J. Biol. Chem. 276: 1789–1793, 2001.

- [67263] 5593. Yan, B.; Raben, N.; Plotz, P. H.: Hes-1, a known transcriptional repressor, acts as a transcriptional activator for the human acid alpha-glucosidase gene in human fibroblast cells. *Biochem. Biophys. Res. Commun.* 291: 582-587, 2002.
- [67264] 5594. Blanchette, F.; Day, R.; Dong, W.; Laprise, M.-H.; Dubois, C. M.: TGF-beta-1 regulates gene expression of its own converting enzyme furin. *J. Clin. Invest.* 99: 1974-1983, 1997.
- [67265] 5595. Copeland, N. G.; Gilbert, D. J.; Chretien, M.; Seidah, N. G.; Jenkins, N. A.: Regional localization of three convertases, PC1 (Nec-1), PC2 (Nec-2), and furin (Fur), on mouse chromosomes. *Genomics* 13: 1356-1358, 1992.
- [67266] 5596. Dubois, C. M.; Laprise, M.-H.; Blanchette, F.; Gentry, L. E.; Leduc, R.: Processing of transforming growth factor beta-1 precursor by human furin convertase. *J. Biol. Chem.* 270: 10618-10624, 1995.
- [67267] 5597. Hendy, G. N.; Bennett, H. P. J.; Gibbs, B. F.; Lazure, C.; Day, R.; Seidah, N. G.: Parathyroid hormone is preferentially cleaved to parathyroid hormone by the pro-hormone convertase furin: a mass spectrometric study. *J. Biol. Chem.* 270: 9517-9525, 1995.
- [67268] 5598. Mbikay, M.; Seidah, N. G.; Chretien, M.; Simpson, E.

M.: Chromosomal assignment of the genes for proprotein convertases PC4, PC5, and PACE4 in mouse and human. *Genomics* 26: 123–129, 1995.

- [67269] 5599. Roebroek, A. J. M.; Schalken, J. A.; Leunissen, J. A. M.; Onnekink, C.; Bloemers, H. P. J.; Van de Ven, W. J. M.: Evolutionary conserved close linkage of the c-fes/fps proto-oncogene and genetic sequences encoding a receptor-like protein. *EMBO J.* 5: 2197–2202, 1986.
- [67270] 5600. Schalken, J. A.; Roebroek, A. J. M.; Oomen, P. P. C. A.; Wagenaar, S. S.; Debruyne, F. M. J.; Bloemers, H. P. J.; Van de Ven, W. J. M.: FUR gene expression as a discriminating marker for small cell and non-small cell lung carcinomas. *J. Clin. Invest.* 80: 1545–1549, 1987.
- [67271] 5601. Seidah, N. G.; Mattei, M. G.; Gaspar, L.; Benjannet, S.; Mbikay, M.; Chretien, M.: Chromosomal assignments of the genes for neuroendocrine convertase PC1 (NEC1) to human 5q15–21, neuroendocrine convertase PC2 (NEC2) to human 20p11.1–11.2, and furin (mouse 7[D1–E2] region). *Genomics* 11:103–107, 1991.
- [67272] 5602. Curtis, E. J.; Fraser, F. C.; Warburton, D.: Congenital cleft lip and palate. *Am. J. Dis. Child.* 102: 853–857, 1961.

- [67273] 5603.Awdeh, Z. L.; Alper, C. A.: Inherited structural polymorphism of the fourth component of human complement. *Proc. Nat. Acad. Sci.* 77:3576–3580, 1980.
- [67274] 5604.Awdeh, Z. L.; Ochs, H. D.; Alper, C. A.: Genetic analysis of C4 deficiency. *J. Clin. Invest.* 67: 260–263, 1981.
- [67275] 5605.Brade, V.; Hall, R. E.; Colten, H. R.: Biosynthesis of pro-C3, a precursor of the third component of complement. *J. Exp. Med.* 146:759–765, 1977.
- [67276] 5606.Carroll, M. C.; Campbell, R. D.; Bentley, D. R.; Porter, R. R.: A molecular map of the human major histocompatibility complex class III region linking complement genes C4, C2 and factor B. *Nature* 307:237–241, 1984.
- [67277] 5607.Carroll, M. C.; Porter, R. R.: Cloning of a human complement component C4 gene. *Proc. Nat. Acad. Sci.* 80: 264–267, 1983.
- [67278] 5608.Chan, A. C.; Mitchell, K. R.; Munns, T. W.; Karp, D. R.; Atkinson, J. P.: Identification and partial characterization of the secreted form of the fourth component of human complement: evidence that it is different from major plasma form. *Proc. Nat. Acad. Sci.* 80: 268–272, 1983.
- [67279] 5609.Cream, J. J.; Olaisen, B.; Teisberg, P.; Soler, A. V.; Thompson, R. A.: Genetic basis of acquired C4 deficiency. *Clin. Genet.* 16:297–300, 1979.

- [67280] 5610.Cunningham–Rundles, C.; Dupont, B.; Jersild, C.; Tegoli, C.; Whitsett,C.; Good, R. A.: Are HLA and Chido related antigenic groups? Transplant.Proc. 9: 33–38, 1977.
- [67281] 5611.Cunningham–Rundles, C.; Tegoli, J.; Dupont, B.; Whitsett, C.; Good,R. A.: Chemical studies on the Chido antigen. Transplant. Proc. 9:647–652, 1977.
- [67282] 5612.Curman, B.; Ostberg, L.; Sandberg, L.; Malmheden–Eriksson, I.;Stalenheim, G.; Rask, L.; Peterson, P. A.: H–2 linked Ss protein is C–4 component of complement. Nature 258: 243–245, 1975.
- [67283] 5613.Ellman, L.; Green, I.; Frank, M.: Genetically controlled total deficiency of the fourth component of complement in the guinea pig. Science 170:74–75, 1970.
- [67284] 5614.Fontaine, M.; Daveau, M.; Lebreton, J. P.: A common antigenic determinant on human C4b and C3b. Molec. Immun. 17: 1075–1078, 1980.
- [67285] 5615.Giles, C. M.: A new genetic variant for Chido. Vox Sang. 46:149–156, 1984.
- [67286] 5616.Hall, R. E.; Colten, H. R.: Genetic defect in biosynthesis of the precursor form of the fourth component of complement. Science 199:69–70, 1978.
- [67287] 5617.Hall, R. E.; Colten, H. R.: Cell–free synthesis of the fourth component of guinea pig complement (C4): identification

cation of a precursor of serum C4 (pro-C4). *Proc. Nat. Acad. Sci.* 74: 1707–1710, 1977.

[67288] 5618. Harris, J. P.; Tegoli, J.; Swanson, J.; Fisher, N.; Gavin, J.; Noades, J.: A nebulous antibody responsible for cross-matching difficulties (Chido). *Vox Sang.* 12: 140–142, 1967.

[67289] 5619. Hobart, M. J.; Lachmann, P. J.: Allotypes of complement components in man. *Transplant. Rev.* 32: 26–42, 1976.

[67290] 5620. Kramer, J.; Fulop, T.; Rajczy, K.; Ahn Tuan, N.; Fust, G.: A marked drop in the incidence of the null allele of the B gene of the fourth component of complement (C4B\*Q0) in elderly subjects: C4B\*Q0 as a probable negative selection factor for survival. *Hum. Genet.* 86: 595–598, 1991.

[67291] 5621. Kronke, M.; Geezy, A. F.; Hadding, U.; Bitter-Suermann, D.: Linkage of C4 and C4 deficiency to Bf and G2LA. *Immunogenetics* 5: 461–466, 1977.

[67292] 5622. Lachmann, P. J.; Grennan, D.; Martin, A.; Demant, P.: Identification of Ss protein as murine C4. *Nature* 258: 242–243, 1975.

[67293] 5623. Lhotta, K.; Schlogl, A.; Uring-Lambert, B.; Kronenberg, F.; König, P.: Complement C4 phenotypes in patients with end-stage renal disease. *Nephron* 72: 442–446,

1996.

- [67294] 5624. Mascart-Lemone, F.; Hauptmann, G.; Goetz, J.; Duchateau, J.; Delespesse, G.; Vray, B.; Dab, I.: Genetic deficiency of C4 presenting with recurrent infections and a SLE-like disease: genetic and immunologic studies. *Am.J. Med.* 75: 295–304, 1983.
- [67295] 5625. Meo, T.; Krasteff, T.; Shreffler, D. C.: Immunochemical characterization of murine H-2 controlled Ss (serum substance) protein through identification of its human homologue as the fourth component of complement. *Proc.Nat. Acad. Sci.* 72: 4536–4540, 1975.
- [67296] 5626. Middleton, J.; Crookston, M. C.: Chido-substance in plasma. *VoxSang.* 23: 256–261, 1972.
- [67297] 5627. Middleton, J.; Crookston, M. C.; Falk, J. A.; Robson, E. B.; Cook, P. J. L.; Batchelor, J. R.; Bodmer, J.; Ferrara, G. B.; Festenstein, J.; Harris, H.; Kissmeyer-Nielsen, F.; Lawler, S. D.; Sachs, J. A.; Wolf, E.: Linkage of Chido and HL-A. *Tissue Antigens* 4: 366–373, 1974.
- [67298] 5628. O'Neill, G. J.: The genetic control of Chido and Rodgers bloodgroup substances. *Seminars Hemat.* 18: 32–38, 1981.
- [67299] 5629. O'Neill, G. J.; Yang, S. Y.; Dupont, B.: Two HLA-linked loci controlling the fourth component of human

complement. Proc. Nat.Acad. Sci. 75: 5165–5169, 1978.

[67300] 5630.O'Neill, G. J.; Yang, S. Y.; Dupont, B.: Chido and Rodgers bloodgroups: relationships to C4 and HLA. Transplant. Proc. 10: 749–751,1978.

[67301] 5631.O'Neill, G. J.; Yang, S. Y.; Tegoli, J.; Berger, R.; Dupont, B.: Chido and Rodgers blood groups are distinct antigenic components of human C4. Nature 273: 668–670, 1978.

[67302] 5632.Ochs, H. D.; Rosenfeld, S. I.; Thomas, E. D.; Giblett, E. R.;Alper, C. A.; Dupont, B.; Schaller, J. G.; Gilliland, B. C.; Hansen,J. A.; Wedgwood, R. J.: Linkage between the gene (or genes) controlling synthesis of the fourth component of complement and the major histocompatibility complex. New Eng. J. Med. 296: 470–475, 1977.

[67303] 5633.Olaisen, B.; Teisberg, P.; Nordhagen, R.; Michaelsen, T.; Gedde-Dahl,T., Jr.: Human complement C4 locus is duplicated on some chromosomes. Nature 279:736–737, 1979.

[67304] 5634.Partanen, J.; Kere, J.; Wessberg, S.; Koskimies, S.: Determination of deletion sizes in the MHC-linked complement C4 and steroid 21-hydroxylase genes by pulsed-field gel electrophoresis. Genomics 5: 345–349,1989.

[67305] 5635.Partanen, J.; Koskimies, S.; Johansson, E.: C4 null



phenotypes among lupus erythematosus patients are predominantly the result of deletions covering C4 and closely linked 21-hydroxylase A genes. *J. Med. Genet.* 25: 387–391, 1988.

[67306] 5636. Petersen, G. B.; Sorensen, I. J.; Buskjaer, L.; Lamm, L. U.: Genetic studies of complement C4 in man. *Hum. Genet.* 53: 31–36, 1979.

[67307] 5637. Kanai, Y.; Hediger, M. A.: Primary structure and functional characterization of a high-affinity glutamate transporter. *Nature* 360: 467–471, 1992.

[67308] 5638. Lin, C. G.; Orlov, I.; Ruggiero, A. M.; Dykes-Hoberg, M.; Lee, A.; Jackson, M.; Rothstein, J. D.: Modulation of the neuronal glutamate transporter EAAC1 by the interacting protein GTRAP3-18. *Nature* 410: 84–88, 2001.

[67309] 5639. Smith, C. P.; Weremowicz, S.; Kanai, Y.; Stelzner, M.; Morton, C. C.; Hediger, M. A.: Assignment of the gene coding for the human high-affinity glutamate transporter EAAC1 to 9p24: potential role in dicarboxylic aminoaciduria and neurodegenerative disorders. *Genomics* 20: 335–336, 1994.

[67310] 5640. Hall, C. R.; Cole, W. G.; Haynes, R.; Hecht, J. T.: Reevaluation of a genetic model for the development of exostosis in hereditary multiple exostosis. *Am. J. Med.*

Genet. 112: 1–5, 2002.

[67311] 5641.Kirby, M. L.; Waldo, K. L.: Neural crest and cardiovascular patterning. Circ.Res. 77: 211–215, 1995.

[67312] 5642.Kurihara, Y.; Kurihara, H.; Oda, H.; Maemura, K.; Nagai, R.; Ishikawa,T.; Yazaki, Y.: Aortic arch malformations and ventricular septaldefect in mice deficient in endothelin-1. J. Clin. Invest. 96: 293–300,1995.

[67313] 5643.Lacey, S. W.; Sanders, J. M.; Rothberg, K. G.; Anderson, R. G.W.; Kamen, B. A.: Complementary DNA for the folate binding proteinincorrectly predicts anchoring to the membrane by glycosyl–phosphatidylinositol. J.Clin. Invest. 84: 715–720, 1989.

[67314] 5644.Istvan, E. S.; Deisenhofer, J.: Structural mechanism for statininhibition of HMG–CoA reductase. Science 292: 1160–1164, 2001.

[67315] 5645.Piedrahita, J. A.; Oetama, B.; Bennett, G. D.; van Waes, J.; Kamen,B. A.; Richardson, J.; Lacey, S. W.; Anderson, R. G. W.; Finnell,R. H.: Mice lacking the folic acid–binding protein Folbp1 are defectivein early embryonic development. Nature Genet. 23: 228–232, 1999.

[67316] 5646.Shaw, G. M.; Jensvold, N. G.; Wasserman, C. R.; Lammer, E. J.: Epidemiologic characteristics of phenotypically distinct neuraltube defects among 0.7 million California

births, 1983–1987. *Teratology* 49:143–149, 1994.

- [67317] 5647. Werler, M. M.; Shapiro, S.; Mitchell, A. A.: Periconceptional folic acid exposure and risk of occurrent neural tube defects. *J.A.M.A.* 269:1257–1261, 1993.
- [67318] 5648. Aittomäki, K.: The genetics of XX gonadal dysgenesis. *Am. J. Hum. Genet.* 54: 844–851, 1994.
- [67319] 5649. Aittomäki, K.; Herva, R.; Stenman, U.–H.; Juntunen, K.; Ylostalo, P.; Hovatta, O.; de la Chapelle, A.: Clinical features of primary ovarian failure caused by a point mutation in the follicle-stimulating hormone receptor gene. *J. Clin. Endocr. Metab.* 81: 3722–3726, 1996.
- [67320] 5650. Aittomäki, K.; Lucena, J. L. D.; Pakarinen, P.; Sistonen, P.; Tapanainen, J.; Gromoll, J.; Kaskikari, R.; Sankila, E.–M.; Lehvaslaiho, H.; Engel, A. R.; Nieschlag, E.; Huh-taniemi, I.; de la Chapelle, A.: Mutation in the follicle-stimulating hormone receptor gene causes hereditary hypergonadotropic ovarian failure. *Cell* 82: 959–968, 1995.
- [67321] 5651. Al-Hendy, A.; Moshynska, O.; Saxena, A.; Feyles, V.: Association between mutations of the follicle-stimulating-hormone receptor and repeated twinning. *Lancet* 356: 914 only, 2000.
- [67322] 5652. Arey, B. J.; Stevis, P. E.; Deecher, D. C.; Shen, E. S.; Frail, D. E.; Negro-Vilar, A.; Lopez, F. J.: Induction of

promiscuous Gprotein coupling of the follicle-stimulating hormone (FSH) receptor:a novel mechanism for transducing pleiotropic actions of FSH isoforms. *Molec.Endocr.* 11: 517-526, 1997.

- [67323] 5653.Beau, I.; Touraine, P.; Meduri, G.; Gougeon, A.; Desroches, A.;Matuchansky, C.; Milgrom, E.; Kuttenn, F.; Misrahi, M.: A novel phenotyperelated to partial loss of function mutations of the follicle stimulatinghormone receptor. *J. Clin. Invest.* 102: 1352-1359, 1998.
- [67324] 5654.Doherty, E.; Pakarinen, P.; Tiitinen, A.; Kiilavuori, A.; Huhtaniemi,I.; Forrest, S.; Aittomaki, K.: A novel mutation in the FSH receptorinhibiting signal transduction and causing primary ovarian failure. *J.Clin. Endocr. Metab.* 87: 1151-1155, 2002.
- [67325] 5655.Gromoll, J.; Pekel, E.; Nieschlag, E.: The structure and organizationof the human follicle-stimulating hormone receptor (FSHR) gene. *Genomics* 35:308-311, 1996.
- [67326] 5656.Gromoll, J.; Ried, T.; Holtgreve-Grez, H.; Nieschlag, E.; Gudermann,T.: Localization of the human FSH receptor to chromosome 2p21 usinga genomic probe comprising exon 10. *J. Molec. Endocr.* 12: 265-271,1994.
- [67327] 5657.Gromoll, J.; Simoni, M.: Follicle-stimulating-hormone receptorand twinning. (Letter) *Lancet* 357:

230 only, 2001.

- [67328] 5658.Heckert, L. L.; Daley, I. J.; Griswold, M. D.: Structural organization of the follicle-stimulating hormone receptor gene. *Molec. Endocr.* 6:70–80, 1992.
- [67329] 5659.Kelton, C. A.; Cheng, S. V. Y.; Nugent, N. P.; Schweickhardt, R. L.; Rosenthal, J. L.; Overton, S. A.; Wands, G. D.; Kuzeja, J.B.; Luchette, C. A.; Chappel, S. C.: The cloning of the human folliclestimulating hormone receptor and its expression in COS-7, CHO, and Y-1 cells. *Molec. Cell. Endocr.* 89: 141–151, 1992.
- [67330] 5660.Kotlar, T. J.; Young, R. H.; Albanese, C.; Crowley, W. F., Jr.; Scully, R. E.; Jameson, J. L.: A mutation in the follicle-stimulating hormone receptor occurs frequently in human ovarian sex cord tumors. *J.Clin. Endocr. Metab.* 82: 1020–1026, 1997.
- [67331] 5661.Minegish, T.; Nakamura, K.; Takakura, Y.; Ibuki, Y.; Igarashi, M.: Cloning and sequencing of human FSH receptor cDNA. *Biochem.Biophys. Res. Commun.* 175: 1125–1130, 1991.
- [67332] 5662.Jiang, M.; Aittomaki, K.; Nilsson, C.; Pakarinen, P.; Litia, A.; Torresani, T.; Simonsen, H.; Goh, V.; Pettersson, K.; de la Chapelle, A.; Huhtaniemi, I.: The frequency of an inactivating point mutation(566C–T) of the human follicle–

stimulating hormone receptor gene in four populations using allele-specific hybridization and time-resolved fluorometry. *J. Clin. Endocr. Metab.* 83: 4338–4343, 1998.

[67333] 5663. Oktay, K.; Briggs, D.; Gosden, R. G.: Ontogeny of follicle-stimulating hormone receptor gene expression in isolated human ovarian follicles. *J. Clin. Endocr. Metab.* 82: 3748–3751, 1997.

[67334] 5664. Rousseau-Merck, M. F.; Atger, M.; Loosfelt, H.; Milgrom, E.; Berger, R.: The chromosomal localization of the human follicle-stimulating hormone receptor gene (FSHR) on 2p21–p16 is similar to that of the luteinizing hormone receptor gene. *Genomics* 15: 222–224, 1993.

[67335] 5665. Simoni, M.; Gromoll, J.; Hoppner, W.; Kamischke, A.; Krafft, T.; Stahle, D.; Nieschlag, E.: Mutational analysis of the follicle-stimulating hormone (FSH) receptor in normal and infertile men: identification and characterization of two discrete FSH receptor isoforms. *J. Clin. Endocr. Metab.* 84: 751–755, 1999.

[67336] 5666. Sipila, K.; Aula, P.: Database for the mutations of the Finnish disease heritage. *Hum. Mutat.* 19: 16–22, 2002.

[67337] 5667. Tapanainen, J. S.; Aittomaki, K.; Min, J.; Vaskivuo, T.; Huhtaniemi, I. T.: Men homozygous for an inactivating

mutation of the follicle-stimulating hormone (FSH) receptor gene present variable suppression of spermatogenesis and fertility. *Nature Genet.* 15: 205–206, 1997.

[67338] 5668. Tortoriello, D. V.; Sidis, Y.; Holtzman, D. A.; Holmes, W. E.; Schneyer, A. L.: Human follistatin-related protein: a structural homologue of follistatin with nuclear localization. *Endocrinology* 142:3426–3434, 2001.

[67339] 5669. Lee, H.-K.; Barbarosie, M.; Kameyama, K.; Bear, M. F.; Huganir, R. L.: Regulation of distinct AMPA receptor phosphorylation sites during bidirectional synaptic plasticity. *Nature* 405: 955–959, 2000.

[67340] 5670. Mack, V.; Burnashev, N.; Kaiser, K. M. M.; Rozov, A.; Jensen, V.; Hvalby, O.; Seeburg, P. H.; Sakmann, B.; Sprengel, R.: Conditional restoration of hippocampal synaptic potentiation in GluR-A-deficient mice. *Science* 292: 2501–2504, 2001.

[67341] 5671. Puckett, C.; Gomez, C. M.; Korenberg, J. R.; Tung, H.; Meier, T. J.; Chen, X. N.; Hood, L.: Molecular cloning and chromosomal localization of one of the human glutamate receptor genes. *Proc. Nat. Acad. Sci.* 88:7557–7561, 1991.

[67342] 5672. Shi, S.-H.; Hayashi, Y.; Petralia, R. S.; Zaman, S. H.; Wenthold, R. J.; Svoboda, K.; Malinow, R.: Rapid spine de-

livery and redistribution of AMPA receptors after synaptic NMDA receptor activation. *Science* 284:1811–1816, 1999.

- [67343] 5673. Zamanillo, D.; Sprengel, R.; Hvalby, O.; Jensen, V.; Burnashev, N.; Rozov, A.; Kaiser, K. M. M.; Koster, H. J.; Borchardt, T.; Worley, P.; Lubke, J.; Frotscher, M.; Kelly, P. H.; Sommer, B.; Andersen, P.; Seeburg, P. H.; Sakmann, B.: Importance of AMPA receptors for hippocampal synaptic plasticity but not for spatial learning. *Science* 284:1805–1811, 1999.
- [67344] 5674. Hardingham, G. E.; Fukunaga, Y.; Bading, H.: Extrasynaptic NMDARs oppose synaptic NMDARs by triggering CREB shut-off and cell death pathways. *Nature Neurosci.* 5: 405–414, 2002.
- [67345] 5675. Takano, H.; Onodera, O.; Tanaka, H.; Mori, H.; Sakimura, K.; Hori, T.; Kobayashi, H.; Mishina, M.; Tsuji, S.: Chromosomal localization of the epsilon-1, epsilon-3, and zeta-1 subunit genes of the human NMDA receptor channel. *Biochem. Biophys. Res. Commun.* 197: 922–926, 1993.
- [67346] 5676. Meera Khan, P.; Doppert, B. A.; Hagemeijer, A.; Westerveld, A.: The human loci for phosphopyruvate hydratase and guanylate kinase are syntenic with the PGD-PGM1 linkage group in man–Chinese hamster somatic cell



hybrids. *Cytogenet. Cell Genet.* 13: 130–131, 1974.

- [67347] 5677. Berkovitz, G. D.; Guerami, A.; Brown, T. R.; MacDonald, P. C.; Migeon, C. J.: Familial gynecomastia with increased extraglandular aromatization of plasma carbon(19)–steroids. *J. Clin. Invest.* 75:1763–1769, 1985.
- [67348] 5678. Bloch, D. B.; Bloch, K. D.; Iannuzzi, M.; Collins, F. S.; Neer, E. J.; Seidman, J. G.; Morton, C. C.: The gene for the  $\alpha$ -i-1 subunit of human guanine nucleotide binding protein maps near the cystic fibrosis locus. *Am. J. Hum. Genet.* 42: 884–888, 1988.
- [67349] 5679. Bray, P.; Carter, A.; Guo, V.; Puckett, C.; Kamholz, J.; Spiegel, A.; Nirenberg, M.: Human cDNA clones for an  $\alpha$  subunit of G(i) signal–transduction protein. *Proc. Nat. Acad. Sci.* 84: 5115–5119, 1987.
- [67350] 5680. Itoh, H.; Toyama, R.; Kozasa, T.; Tsukamoto, T.; Matsuoka, M.; Kaziro, Y.: Presence of three distinct molecular species of G(i) protein  $\alpha$  subunit: structure of rat cDNAs and human genomic DNAs. *J. Biol. Chem.* 263: 6656–6664, 1988.
- [67351] 5681. Suki, W. N.; Abramowitz, J.; Mattera, R.; Codina, J.; Birnbaumer, L.: The human genome encodes at least three non-allelic G proteins with  $\alpha$ -i-type subunits. *FEBS Lett.* 220: 187–192, 1987.

- [67352] 5682.Sullivan, K. A.; Liao, Y.-C.; Alborzi, A.; Beiderman, B.; Chang,F.-H.; Masters, S. B.; Levinson, A. D.; Bourne, H. R.: Inhibitoryand stimulatory G proteins of adenylate cy- clase: cDNA and amino acidsequences of the alpha chains. Proc. Nat. Acad. Sci. 83: 6687-6691,1986.
- [67353] 5683.Jiang, M.; Gold, M. S.; Boulay, G.; Spicher, K.; Peyton, M.; Brabet,P.; Srinivasan, Y.; Rudolph, U.; Ellison, G.; Birn- baumer, L.: Multipleneurological abnormalities in mice deficient in the G protein G(o). Proc.Nat. Acad. Sci. 95: 3269-3274, 1998.
- [67354] 5684.Kroll, S. D.; Chen, J.; De Vivo, M.; Carty, D. J.; Buku, A.; Premont,R. T.; Iyengar, R.: The Q205LGo-alpha subunit expressed in NIH-3T3cells induces transformation. J. Biol. Chem. 267: 23183-23188, 1992.
- [67355] 5685.Murtagh, J. J., Jr.; Eddy, R.; Shows, T. B.; Moss, J.; Vaughan,M.: Different forms of Go alpha mRNA arise by alternative splicingof transcripts from a single gene on human chromosome 16. Molec.Cell. Biol. 11: 1146-1155, 1991.
- [67356] 5686.Ram, P. T.; Horvath, C. M.; Iyengar, R.: Stat3-mediated transformationof NIH-3T3 cells by the constitutively active Q205L G-alpha(O) protein. Science 287:142-144, 2000.

- [67357] 5687. Strathmann, M.; Wilkie, T. M.; Simon, M. I.: Alternative splicing produces transcripts encoding two forms of the alpha subunit of GTP-binding protein G(o). *Proc. Nat. Acad. Sci.* 87: 6477–6481, 1990.
- [67358] 5688. Tsukamoto, T.; Toyama, R.; Itoh, H.; Kozasa, T.; Matsuoka, M.; Kaziro, Y.: Structure of the human gene and two rat cDNAs encoding the alpha chain of GTP-binding regulatory protein G(o): two different mRNAs are generated by alternative splicing. *Proc. Nat. Acad. Sci.* 88:2974–2978, 1991.
- [67359] 5689. Valenzuela, D.; Han, X.; Mende, U.; Fankhauser, C.; Mashimo, H.; Huang, P.; Pfeiffer, J.; Neer, E. J.; Fishman, M. C.: G-alpha-o is necessary for muscarinic regulation of Ca(2+) channels in mouse heart. *Proc. Nat. Acad. Sci.* 94: 1727–2732, 1997.
- [67360] 5690. Belluscio, L.; Gold, G. H.; Nemes, A.; Axel, R.: Mice deficient in G(olf) are anosmic. *Neuron* 20: 69–81, 1998.
- [67361] 5691. Buck, L. B.: Information coding in the mammalian olfactory system. *Cold Spring Harbor Symp. Quant. Biol.* 61: 147–155, 1996.
- [67362] 5692. Jones, D. T.; Reed, R. R.: Golf: an olfactory neuron specific G protein involved in odorant signal transduction. *Science* 244:790–795, 1989.

- [67363] 5693.Lindgren, V.; Luskey, K. L.; Russell, D. W.; Francke, U.: Humangenes involved in cholesterol metabolism: chromosomal mapping of the loci for the low density lipoprotein receptor and 3-hydroxy-3-methylglutaryl-coenzyme A reductase with cDNA probes. *Proc. Nat. Acad. Sci.* 82: 8567–8571, 1985.
- [67364] 5694.Luskey, K. L.: Conservation of promoter sequence but not complex intron splicing pattern in human and hamster genes for 3-hydroxy-3-methylglutarylcoenzyme A reductase. *Molec. Cell. Biol.* 7: 1881–1893, 1987.
- [67365] 5695.Mohandas, T.; Heinzmann, C.; Sparkes, R. S.; Wasmuth, J.; Edwards, P.; Lusk, A. J.: Assignment of human 3-hydroxy-3-methylglutarylcoenzyme A reductase gene to q13–q23 region of chromosome 5. *Somat.Cell Molec. Genet.* 12: 89–94, 1986.
- [67366] 5696.Osborne, T. F.; Goldstein, J. L.; Brown, M. S.: 5-prime end of HMG CoA reductase gene contains sequences responsible for cholesterol-mediated inhibition of transcription. *Cell* 42: 203–212, 1985.
- [67367] 5697.Van Doren, M.; Broihier, H. T.; Moore, L. A.; Lehmann, R.: HMG-CoA reductase guides migrating primordial germ cells. *Nature* 396: 466–469, 1998.
- [67368] 5698.Hou, J.; Parrish, J.; Ludecke, H.-J.; Sapru, M.; Wang,

Y.; Chen,W.; Hill, A.; Siegel–Bartelt, J.; Northrup, H.; Elder, F. F. B.; Chinault,C.; Horsthemke, B.; Wagner, M. J.; Wells, D. E.: A 4–megabase YACcontig that spans the Langer–Giedion syndrome region on human chromosome8q24.1: use in refining the location of the trichorhinophalangeal–syndrome and multiple exostoses genes (TRPS1 and EXT1). *Genomics* 29:87–97, 1995.

[67369] 5699.Ludecke, H.–J.; Wagner, M. J.; Nardmann, J.; La Pillo, B.; Parrish,J. E.; Willems, P. J.; Haan, E. A.; Frydman, M.; Hamers, G. J. H.;Wells, D. E.; Horsthemke, B.: Molecular dissection of a contiguousgene syndrome: localization of the genes involved in the Langer–Giedionsyndrome. *Hum. Molec. Genet.* 4: 31–36, 1995.

[67370] 5700.Koizumi, T.; Hendel, E.; Lalley, P. A.; Tchetgen, M.–B. N.; Nadeau,J. H.: Homologs of genes and anonymous loci on human chromosome 13map to mouse chromosomes 8 and 14. *Mammalian Genome* 6: 263–268,1995.

[67371] 5701.Devriendt, K.; Zhang, J.; van Leuven, F.; van den Berghe, H.; Cassiman,J. J.; Marynen, P.: A cluster of alpha 2–macroglobulin–related genes(alpha 2 M) on human chromosome 12p: cloning of the pregnancy–zoneprotein gene and an alpha 2M pseudogene. *Gene* 81: 325–334, 1989.

- [67372] 5702.Dow, D. J.; Lindsey, N.; Cairns, N. J.; Brayne, C.; Robinson, D.;Huppert, F. A.; Paykel, E. S.; Xuereb, J.; Wilcock, G.; Whittaker,J. L.; Rubinsztein, D. C.: Alpha-2 macroglobulin polymorphism andAlzheimer disease risk in the UK. (Letter) *Nature Genet.* 22: 16-17,1999.
- [67373] 5703.Fukushima, Y.; Bell, G. I.; Shows, T. B.: The polymorphic humanalpha-2-macroglobulin gene (A2M) is located in chromosome region 12p12.3-p13.3. *Cytogenet.Cell Genet.* 48: 58-59, 1988.
- [67374] 5704.Gallango, M. L.; Castillo, O.: Alpha-2-macroglobulin polymorphism:a new genetic system detected by immuno-electrophoresis. *J. Immunogenet.* 1:147-151, 1974.
- [67375] 5705.Hilliker, C.; Overbergh, L.; Petit, P.; Van Leuven, F.; Van denBerghe, H.: Assignment of mouse alpha-2-macroglobulin gene to chromosome6 band F1-G3. *Mammalian Genome* 3: 469-471, 1992.
- [67376] 5706.Kan, C.-C.; Solomon, E.; Belt, K. T.; Chain, A. C.; Hiorns, L.R.; Fey, G.: Nucleotide sequence of cDNA encoding human alpha-2-macroglobulinand assignment of the chromosomal locus. *Proc. Nat. Acad. Sci.* 82:2282-2286, 1985.
- [67377] 5707.Leikola, J.; Fudenberg, H. H.; Kasukawa, R.; Milgrom,

F.: A new genetic polymorphism of human serum: alpha(2) macroglobulin (AL-M). *Am.J. Hum. Genet.* 24: 134-144, 1972.

[67378] 5708.Liao, A.; Nitsch, R. M.; Greenberg, S. M.; Finckh, U.; Blacker,D.; Albert, M.; Rebeck, G. W.; Gomez-Isla, T.; Clatworthy, A.; Binetti,G.; Hock, C.; Mueller-Thomsen, T.; Mann, U.; Zuchowski, K.; Beisiegel,U.; Staehelin, H.; Growdon, J. H.; Tanzi, R. E.; Hyman, B. T.: Genetic association of an alpha-2-macroglobulin (val1000ile) polymorphism and Alzheimer's disease. *Hum. Molec. Genet.* 7: 1953-1956, 1998.

[67379] 5709.Marynen, P.; Bell, G. I.; Cavalli-Sforza, L. L.: Three RFLPs associated with the human alpha-2-macroglobulin gene (A2M). *Nucleic Acids Res.* 13: 8287 only, 1985.

[67380] 5710.Marynen, P.; Zhang, J.; Devriendt, K.; Cassiman, J.-J.: Alpha-2-macroglobulin, pregnancy zone protein and an alpha-2-macroglobulin pseudogene map to chromosome 12p12.2-p13. (Abstract) *Cytogenet. Cell Genet.* 51:1040 only, 1989.

[67381] 5711.Matthijs, G.; Devriendt, K.; Cassiman, J.-J.; van den Berghe,H.; Marynen, P.: Structure of the human alpha-2 macroglobulin gene and its promoter (sic). *Biochem. Biophys. Res. Commun.* 184: 596-603, 1992.

- [67382] 5712. Mattheijs, G.; Marynen, P.: A deletion polymorphism in the human  $\alpha$ -2-macroglobulin (A2M) gene. *Nucleic Acids Res.* 19: 5102 only, 1991.
- [67383] 5713. Mattheijs, G.; Marynen, P. A.: A deletion polymorphism in the human  $\alpha$ -2-macroglobulin (A2M) gene. *Nucleic Acids Res.* 19: 5102 only, 1991.
- [67384] 5714. Poller, W.; Barth, J.; Voss, B.: Detection of an alteration of the  $\alpha$ -2-macroglobulin gene in a patient with chronic lung disease and serum  $\alpha$ -2-macroglobulin deficiency. *Hum. Genet.* 83: 93–96, 1989.
- [67385] 5715. Poller, W.; Faber, J.-P.; Klobeck, G.; Olek, K.: Cloning of the human  $\alpha$ -2-macroglobulin gene and detection of mutations in two functional domains: the bait region and the thiolester site. *Hum. Genet.* 88: 313–319, 1992.
- [67386] 5716. Rogaeva, E. A.; Premkumar, S.; Grubber, J.; Serneels, L.; Scott, W. K.; Kawarai, T.; Song, Y.; Hill, D. M.; Abou-Donia, S. M.; Martin, E. R.; Vance, J. J.; Yu, G.; and 18 others: An  $\alpha$ -2-macroglobulin insertion-deletion polymorphism in Alzheimer disease. (Letter) *Nature Genet.* 22: 19–21, 1999.
- [67387] 5717. Rudrasingham, V.; Wavrant-De Vrieze, F.; Lambert, J.-C.; Chakraverty, S.; Kehoe, P.; Crook, R.; Amouyel, P.; Wu, W.; Rice, F.; Perez-Tur, J.; Frigard, B.; Morris, J. C.; and



11 others: Alpha-2 macroglobulin gene and Alzheimer disease. (Letter) *Nature Genet.* 22: 17–19, 1999.

- [67388] 5718. Sottrup-Jensen, L.; Stepanik, T. M.; Kristensen, T.; Lonblad, P. B.; Jones, C. M.; Wierzbicki, D. M.; Magnusson, S.; Domdey, H.; Wetsel, R. A.; Lundwall, A.; Tack, B. F.; Fey, G. H.: Common evolutionary origin of alpha-2-macroglobulin and complement components C3 and C4. *Proc. Nat. Acad. Sci.* 82: 9–13, 1985.
- [67389] 5719. Umans, L.; Serneels, L.; Hilliker, C.; Stas, L.; Overbergh, L.; De Strooper, B.; Van Leuven, F.; Van den Berghe, H.: Molecular cloning of the mouse gene coding for alpha-2-macroglobulin and targeting of the gene in embryonic stem cells. *Genomics* 22: 519–529, 1994.
- [67390] 5720. Buckler, A. J.; Chang, D. D.; Graw, S. L.; Brook, J. D.; Haber, D. A.; Sharp, P. A.; Housman, D. E.: Exon amplification: a strategy to isolate mammalian genes based on RNA splicing. *Proc. Nat. Acad. Sci.* 88: 4005–4009, 1991.
- [67391] 5721. Brennan, R. W.; Dehejia, H.; Kutt, H.; Verebely, K.; McDowell, F.: Diphenylhydantoin intoxication attendant to slow inactivation of isoniazid. *Neurology* 20: 687–693, 1970.
- [67392] 5722. Brown, S.; Chalmers, D. E.: Microsomal epoxide hydrolase activity in human x mouse hybrid cells. *Biochem.*

Biophys. Res. Commun. 137:775–780, 1986.

- [67393] 5723. Buehler, B. A.; Delimont, D.; van Waes, M.; Finnell, R. H.: Prenatal prediction of risk of the fetal hydantoin syndrome. *New Eng. J. Med.* 322:1567–1571, 1990.
- [67394] 5724. Chodirker, B. N.; Chudley, A. E.; Reed, M. H.; Persaud, T. V. N.: Possible prenatal hydantoin effect in a child born to a nonepileptic mother. *Am. J. Med. Genet.* 27: 373–378, 1987.
- [67395] 5725. De Smet, L.; Debeer, P.: Fetal hydantoin syndrome with unilateral atypical cleft hand: additional evidence for vascular disruption. *Genet. Counsel.* 13: 157–161, 2002.
- [67396] 5726. de Wolff, F. A.; Vermeij, P.; Ferrari, M. D.; Buruma, O. J. S.; Breimer, D. D.: Impairment of phenytoin parahydroxylation as a cause of severe intoxication. *Ther. Drug Monit.* 5: 213–215, 1983.
- [67397] 5727. Gennis, M. A.; Vemuri, R.; Burns, E. A.; Hill, J. V.; Miller, M. A.; Spielberg, S. P.: Familial occurrence of hypersensitivity to phenytoin. *Am. J. Med.* 91: 631–634, 1991.
- [67398] 5728. Goldman, A. S.; Van Dyke, D. C.; Gupta, C.; Katsumata, M.: Elevated glucocorticoid receptor levels in lymphocytes of children with the fetal hydantoin syndrome (FHS). *Am. J. Med. Genet.* 28: 607–618, 1987.
- [67399] 5729. Hanson, J. W.; Myrianthopoulos, N. C.; Sedgwick, M.

H. A.; Smith,D. W.: Risks to the offspring of women treated with hydantoin anticonvulsants,with emphasis on the fetal hydantoin syndrome. J. Pediat. 89: 662–668,1976.

[67400] 5730.Hartsfield, J. K., Jr.; Sutcliffe, M. J.; Everett, E. T.; Hassett,C.; Omiecinski, C. J.; Saari, J. A.: Assignment of microsomal epoxidehydrolase (EPHX1) to human chromosome 1q42.1 by in situ hybridization. Cytogenet.Cell Genet. 83: 44–45, 1998.

[67401] 5731.Dorner, M. H.; Salfeld, J.; Will, H.; Leibold, E. A.; Vass, J.K.; Munro, H. N.: Structure of human ferritin light subunit messengerRNA: comparison with heavy subunit message and functional implications. Proc.Nat. Acad. Sci. 82: 3139–3143, 1985.

[67402] 5732.Filie, J. D.; Buckler, C. E.; Kozak, C. A.: Genetic mapping ofthe mouse ferritin light chain gene and 11 pseudogenes on 11 mousechromosomes. Mammalian Genome 9: 111–113, 1998.

[67403] 5733.Girelli, D.; Bozzini, C.; Zecchina, G.; Tinazzi, E.; Bosio, S.;Piperno, A.; Ramenghi, U.; Peters, J.; Levi, S.; Camaschella, C.;Corrocher, R.: Clinical, biochemical and molecular findings in a series of families with hereditary hyperferritinaemia–cataract syndrome. Brit.J. Haemat.

115: 334–340, 2001.

- [67404] 5734. Girelli, D.; Corrocher, R.; Bisceglia, L.; Olivieri, O.; De Franceschi, L.; Zelante, L.; Gasparini, P.: Molecular basis for the recently described hereditary hyperferritinemia–cataract syndrome: A mutation in the iron–responsive element of ferritin L–subunit gene (the 'Verona mutation'). *Blood* 86: 4050–453, 1995.
- [67405] 5735. Girelli, D.; Corrocher, R.; Bisceglia, L.; Olivieri, O.; Zelante, L.; Panozzo, G.; Gasparini, P.: Hereditary hyperferritinemia–cataract syndrome caused by a 29–base pair deletion in the iron responsive element of ferritin L–subunit gene. *Blood* 90: 2084–2088, 1997.
- [67406] 5736. Girelli, D.; Olivieri, O.; De Franceschi, L.; Corrocher, R.; Bergamaschi, G.; Cazzola, M.: A linkage between hereditary hyperferritinaemia not related to iron overload and autosomal dominant congenital cataract. *Brit. J. Haemat.* 90: 931–934, 1995.
- [67407] 5737. Lebo, R. V.; Kan, Y. W.; Cheung, M.–C.; Jain, S. K.; Drysdale, J.: Human ferritin light chain gene sequences mapped to several sorted chromosomes. *Hum. Genet.* 71: 325–328, 1985.
- [67408] 5738. Martin, M. E.; Fargion, S.; Brissot, P.; Pellat, B.; Beaumont, C.: A point mutation in the bulge of the iron–

responsive element of the L ferritin gene in two families with the hereditary hyperferritinemia–cataract syndrome. Blood 91: 319–323, 1998.

- [67409] 5739. McGill, J. R.; Boyd, D.; Barrett, K. J.; Drysdale, J. W.; Moore, C. M.: Localization of human ferritin H (heavy) and L (light) subunits by in situ hybridization. (Abstract) Am. J. Hum. Genet. 36: 146S, 1984.
- [67410] 5740. McLeod, J. L.; Craig, J.; Gumley, S.; Roberts, S.; Kirkland, M. A.: Mutation spectrum in Australian pedigrees with hereditary hyperferritinaemia–cataract syndrome reveals novel and de novo mutations. Brit. J. Haemat. 118: 1179–1182, 2002.
- [67411] 5741. Mumford, A. D.; Vulliamy, T.; Lindsay, J.; Watson, A.: Hereditary hyperferritinemia–cataract syndrome: two novel mutations in the L–ferritin iron–responsive element. (Letter) Blood 91: 367–368, 1998.
- [67412] 5742. Munro, H. N.; Aziz, N.; Leibold, E. A.; Murray, M.; Rogers, J.; Vass, J. K.; White, K.: The ferritin genes: structure, expression, and regulation. Ann. N.Y. Acad. Sci. 526: 113–123, 1988.
- [67413] 5743. Santoro, C.; Marone, M.; Ferrone, M.; Costanzo, F.; Colombo, M.; Minganti, C.; Cortese, R.; Silengo, L.: Cloning of the gene coding for human L apoferritin. Nucleic Acids

Res. 14: 2863–2876, 1986.

[67414] 5744.Watanabe, N.; Drysdale, J. W.: Evidence for distinct mRNAs for ferritin subunits. *Biochem. Biophys. Res. Commun.* 98: 507–511, 1981.

[67415] 5745.Usala, A.-L.; Madigan, T.; Burguera, B.; Sinha, M. K.; Caro, J.F.; Cunningham, P.; Powell, J. G.; Butler, P. C.: Treatment of insulin-resistant diabetic ketoacidosis with insulin-like growth factor I in an adolescent with insulin-dependent diabetes. *New Eng. J. Med.* 327: 853–857, 1992.

[67416] 5746.Vaessen, N.; Janssen, J. A.; Heutink, P.; Hofman, A.; Lamberts, S. W. J.; Oostra, B. A.; Pols, H. A. P.; van Duijn, C. M.: Association between genetic variation in the gene for insulin-like growth factor-1 and low birthweight. *Lancet* 359: 1036–1037, 2002.

[67417] 5747.Van Wyk, J. J.; Svoboda, M. E.; Underwood, L. E.: Evidence from radioligand assays that somatomedin-C and insulin-like growth factor-I are similar to each other and different from other somatomedins. *J. Clin. Endocr. Metab.* 50: 206–208, 1980.

[67418] 5748.Hochwald, G. M.; Thorbecke, G. J.: Abnormal metabolism or reduced transport of CSF gamma-trace microprotein in hereditary cerebral hemorrhage with amyloi-

dosis. (Letter) New Eng. J. Med. 312: 1127–1128, 1985.

- [67419] 5749.Jensson, O.; Arnason, A.; Thorsteinsson, L.; Petursdottir, I.;Gudmundsson, G.; Blondal, H.; Grubb, A.; Lofberg, H.; Luyendijk, W.;Bots, G. T. A. M.; Frangione, B.: Cystatin C (gamma-trace) amyloidosis.In:Turk, V.: Cysteine Proteinases and their Inhibitors. New York:Walter de Gruyter and Co. (pub.) 1986.
- [67420] 5750.Jensson, O.; Gudmundsson, G.; Arnason, A.; Blondal, H.; Petursdottir,I.; Thorsteinsson, L.; Grubb, A.; Lofberg, H.; Cohen, D.; Frangione,B.: Hereditary cystatin C (gamma-trace) amyloid angiopathy of theCNS causing cerebral hemorrhage. Acta Neurol. Scand. 76: 102–114,1987.
- [67421] 5751.Jensson, O.; Palsdottir, A.; Thorsteinsson, L.; Arnason, A.:The saga of cystatin C gene mutation causing amyloid angiopathy andbrain hemorrhage--clinical genetics in Iceland. Clin. Genet. 36:368–377, 1989.
- [67422] 5752.Lofberg, H.; Grubb, A. O.; Nilsson, E. K.; Jensson, O.; Gudmundsson,G.; Blondal, H.; Arnason, A.; Thorsteinsson, L.: Immunohistochemicalcharacterization of the amyloid deposits and quantitation of pertinentcerebrospinal fluid proteins in hereditary cerebral hemorrhage withamyloidosis. Stroke 18: 431–440, 1987.

- [67423] 5753.Palsdottir, A.; Abrahamson, M.; Thorsteinsson, L.; Arnason, A.;Olafsson, I.; Grubb, A.; Jensson, O.: Mutation in cystatin C gene causes hereditary brain haemorrhage. Lancet II: 603–604, 1988.
- [67424] 5754.Zalin, A. M.; Jones, S.; Fitch, N. J. S.; Ramsden, D. B.: Familial nephropathic non-neuropathic amyloidosis: clinical features, immunohistochemistry and chemistry. Quart. J. Med. 81: 945–956, 1991.
- [67425] 5755.Ceccherini, I.; Romei, C.; Barone, V.; Pacini, F.; Martino, E.; Loviselli, A.; Pinchera, A.; Romeo, G.: Identification of the cys634-to-tyr mutation of the RET proto-oncogene in a pedigree with multiple endocrine neoplasia type 2A and localized cutaneous lichen amyloidosis. J.Endocr. Invest. 17: 201–204, 1994.
- [67426] 5756.Winkelmann, B. R.; Nauck, M.; Klein, B.; Russ, A. P.; Bohm, B.O.; Siekmeier, R.; Ihnken, K.; Verho, M.; Gross, W.; Marz, W.: Deletion polymorphism of the angiotensin I-converting enzyme gene is associated with increased plasma angiotensin-converting enzyme activity but not with increased risk for myocardial infarction and coronary artery disease. Ann. Intern. Med. 125: 19–25, 1996.
- [67427] 5757.Woods, D.; Onambele, G.; Woledge, R.; Skelton, D.; Bruce, S.; Humphries, S. E.; Montgomery, H.: Angiotensin-I



converting enzyme genotype-dependent benefit from hormone replacement therapy in isometric muscle strength and bone mineral density. *J. Clin. Endocr. Metab.* 86:2200–2204, 2001.

- [67428] 5758. Yoshioka, T.; Xu, Y.; Yoshida, H.; Shiraga, H.; Muraki, T.; Ito, K.: Deletion polymorphism of the angiotensin converting enzyme gene predicts persistent proteinuria in Henoch–Schönlein purpura nephritis. *Arch. Dis. Child.* 79: 394–399, 1998.
- [67429] 5759. Zhu, X.; Bouzekri, N.; Southam, L.; Cooper, R. S.; Adeyemo, A.; McKenzie, C. A.; Luke, A.; Chen, G.; Elston, R. C.; Ward, R.: Linkage and association analysis of angiotensin I-converting enzyme (ACE) gene polymorphisms with ACE concentration and blood pressure. *Am. J. Hum. Genet.* 68: 1139–1148, 2001.
- [67430] 5760. Zhu, X.; McKenzie, C. A.; Forrester, T.; Nickerson, D. A.; Broeckel, U.; Schunkert, H.; Doering, A.; Jacob, H. J.; Cooper, R. S.; Rieder, M. J.: Localization of a small genomic region associated with elevated ACE. *Am. J. Hum. Genet.* 67: 1144–1153, 2000. Note: Erratum: *Am. J. Hum. Genet.* 67: 1365 only, 2000.
- [67431] 5761. Axton, R.; Hanson, I.; Danes, S.; Sellar, G.; van Heyningen, V.; Prosser, J.: The incidence of PAX6 mutation

in patients with simple aniridia: an evaluation of mutation detection in 12 cases. *J. Med. Genet.* 34: 279–286, 1997.

[67432] 5762. Beauchamp, G. R.: Anterior segment dysgenesis keratolenticular adhesion and aniridia. *J. Pediatr. Ophthalm. Strabismus* 17: 55–58, 1978.

[67433] 5763. Crolla, J. A.; Cross, I.; Atkey, N.; Wright, M.; Oley, C. A.: FISH studies in a patient with sporadic aniridia and t(7;11)(q31.2;p13). *J. Med. Genet.* 33: 66–68, 1996.

[67434] 5764. Fantes, J.; Redeker, B.; Breen, M.; Boyle, S.; Brown, J.; Fletcher, J.; Jones, S.; Bickmore, W.; Fukushima, Y.; Mannens, M.; Danes, S.; van Heyningen, V.; Hanson, I.: Aniridia-associated cytogenetic rearrangements suggest that a position effect may cause the mutant phenotype. *Hum. Molec. Genet.* 4: 415–422, 1995.

[67435] 5765. Sarkar, F. H.; Gupta, S. L.: Receptors for human gamma interferon: binding and crosslinking of 125-I-labeled recombinant human gamma interferon to receptors on WISH cells. *Proc. Nat. Acad. Sci.* 81: 5160–5164, 1984.

[67436] 5766. Slate, D. L.; Ruddle, F. H.: Antibodies to chromosome 21 coded cell surface components can block response to human interferon. *Cytogenet. Cell Genet.* 22: 265–269, 1978.

- [67437] 5767.Slate, D. L.; Shulman, L.; Lawrence, J. B.; Revel, M.; Ruddle, F. H.: Presence of human chromosome 21 alone is sufficient for hybridcell sensitivity to human interferon. *J. Virol.* 25: 319–325, 1978.
- [67438] 5768.Takaoka, A.; Mitani, Y.; Suemori, H.; Sato, M.; Yokochi, T.; Noguchi, S.; Tanaka, N.; Taniguchi, T.: Cross talk between interferon- $\gamma$  and  $\alpha/\beta$  signaling components in caveolar membrane domains. *Science* 288:2357–2360, 2000.
- [67439] 5769.Tan, Y. H.: Chromosome 21 and the cell growth inhibitory effect of human interferon preparations. *Nature* 260: 141–143, 1976.
- [67440] 5770.Tan, Y. H.; Schneider, E. L.; Tischfield, J.; Epstein, C. J.; Ruddle, F. H.: Human chromosome 21 dosage: effect on the expression of the interferon induced antiviral state. *Science* 186: 61–63, 1974.
- [67441] 5771.Tan, Y. H.; Tischfield, J.; Ruddle, F. H.: The linkage of genes for the human interferon-induced antiviral protein and indophenoloxidase to chromosome G-21. *J. Exp. Med.* 37: 317–330, 1973.
- [67442] 5772.Weil, J.; Tucker, G.; Epstein, L. B.; Epstein, C. J.: Interferon induction of (2-prime–5-prime) oligoisoadenylate synthetase in diploid and trisomy 21 human fibroblasts:

relation to dosage of the interferonreceptor gene (IFRC).  
Hum. Genet. 65: 108–111, 1983.

[67443] 5773.Wiranowska–Stewart, M.; Stewart, W. E., II: The role of humanchromosome 21 in sensitivity to interferons. J. Gen. Virol. 37:629–633, 1977.

[67444] 5774.Aguet, M.; Dembic, Z.; Merlin, G.: Molecular cloning and expressionof the human interferon–gamma receptor. Cell 55: 273–280, 1988.

[67445] 5775.Alcaide–Loridan, C.; Le Coniat, M.; Bono, R.; Benech, P.; Couillin,P.; Van Cong, N.; Fisher, D. N.; Berger, R.; Fel– lous, M.: Mappingof the human interferon gamma re– sponse. (Abstract) Cytogenet. CellGenet. 51: 949 only, 1989.

[67446] 5776.Branca, A. A.; Baglioni, C.: Evidence that types I and II interferonshave different receptors. Nature 294: 768–770, 1981.

[67447] 5777.Casanova, J.–L.; Jouanguy, E.; Lamhamedi, S.; Blanche, S.; Fischer,A.: Immunological conditions of chil– dren with BCG disseminated infection.(Letter) Lancet 346: 581 only, 1995.

[67448] 5778.Celada, A.; Allen, R.; Esparza, I.; Gray, P. W.; Schreiber, R.D.: Demonstration and partial characteriza– tion of the interferon–gammareceptor on human

mononuclear phagocytes. J. Clin. Invest. 76:  
2196–2205, 1985.

[67449] 5779. Lee, W. M.; Galbraith, R. M.: The extracellular actin-scavenger system and actin toxicity. New Eng. J. Med. 326: 1335–1341, 1992.

[67450] 5780. Pilz, A.; Moseley, H.; Peters, J.; Abbott, C.: Comparative mapping of mouse chromosome 2 and human chromosome 9q: the genes for gelsolin and dopamine beta-hydroxylase map to mouse chromosome 2. Genomics 12: 715–719, 1992.

[67451] 5781. Brahe, C.; Servidei, S.; Zappata, S.; Ricci, E.; Tonali, P.; Neri, G.: Genetic homogeneity between childhood-onset and adult-onset autosomal recessive spinal muscular atrophy. Lancet 346: 741–742, 1995.

[67452] 5782. Eglen, R. M.; Whiting, R. L.: Muscarinic receptor subtypes: a critique of the current classification and a proposal for a working nomenclature. J. Auton. Pharm. 6: 323–346, 1986.

[67453] 5783. Goyal, R. K.: Muscarinic receptor subtypes: physiology and clinical implications. New Eng. J. Med. 321: 1022–1029, 1989.

[67454] 5784. Liao, C. F.; Themmen, A. P.; Joho, R.; Barberis, C.; Birnbaumer, M.; Birnbaumer, L.: Molecular cloning and ex-

pression of a fifth muscarinicacetylcholine receptor. J. Biol. Chem. 264: 7328–7337, 1989.

[67455] 5785.Peralta, E. G.; Ashkenazi, A.; Winslow, J. W.; Smith, D. H.; Ramachandran,J.; Capon, D. J.: Distinct primary structures, ligand–binding propertiesand tissue–specific expression of four human muscarinic acetylcholinereceptors. EMBO J. 6: 3923–3929, 1987.

[67456] 5786.Burchill, S. A.; Wheeldon, J.; Cullinane, C.; Lewis, I. J.: EWS–FLI1fusion transcripts identified in patients with typical neuroblastoma. Europ.J. Cancer 33: 239–243, 1997.

[67457] 5787.Pan, T.–C.; Sasaki, T.; Zhang, R.–Z.; Fassler, R.; Timpl, R.; Chu,M.–L.: Structure and expression of fibulin–2, a novel extracellularmatrix protein with multiple EGF–like repeats and consensus motifsfor calcium binding. J. Cell Biol. 123: 1269–1277, 1993.

[67458] 5788.Fling, S. P.; Arp, B.; Pious, D.: HLA–DMA and –DMB genes are bothrequired for MHC class II/peptide complex formation in antigen–presentingcells. Nature 368: 554–558, 1994.

[67459] 5789.Morris, P.; Shaman, J.; Attaya, M.; Amaya, M.; Goodman, S.; Bergman,C.; Monaco, J. J.; Mellins, E.: An essential role for HLA–DM in antigenpresentation by class II major histocompatibility molecules. Nature 368:551–554, 1994.

- [67460] 5790.Hiroi, S.; Harada, H.; Nishi, H.; Satoh, M.; Nagai, R.; Kimura,A.: Polymorphisms in the SOD2 and HLA-DRB1 genes are associated withnonfamilial idiopathic dilated cardiomyopathy in Japanese. *Biochem.Biophys. Res. Commun.* 261: 332-339, 1999.
- [67461] 5791.Nishi, H.; Koga, Y.; Koyanagi, T.; Harada, H.; Imaizumi, T.; Toshima,H.; Sasazuki, T.; Kimura, A.: DNA typing of HLA class II genes inJapanese patients with dilated cardiomyopathy. *J. Molec. Cell. Cardiol.* 27:2385-2392, 1995.
- [67462] 5792.Delattre, O.; Zucman, J.; Plougastel, B.; Desmaze, C.; Melot, T.;Peter, M.; Kovar, H.; Joubert, I.; de Jong, P.; Rouleau, G.; Aurias,A.; Thomas, G.: Gene fusion with an ETS DNA-binding domain causedby chromosome translocation in human tumours. *Nature* 359: 162-165,1992.
- [67463] 5793.Douglass, E. C.; Valentine, M.; Green, A. A.; Hayes, F. A.; Thompson,E. I.: t(11;22) and other chromosomal rearrangements in Ewing's sarcoma. *J.Nat. Cancer Inst.* 77: 1211-1213, 1986.
- [67464] 5794.Ewing, J.: Diffuse endothelioma of bone. *Proc. N. Y. Path. Soc.* 21:17-24, 1921.
- [67465] 5795.Gerald, W. L.; Rosai, J.; Ladanyi, M.: Characterization of thegenomic breakpoint and chimeric transcripts in the

EWS-WT1 gene fusion of desmoplastic small round cell tumor. *Proc. Nat. Acad. Sci.* 92:1028–1032, 1995.

- [67466] 5796. Geurts van Kessel, A.; Turc-Carel, C.; de Klein, A.; Grosveld, G.; Lenoir, G.; Bootsma, D.: Translocation of oncogene c-sis from chromosome 22 to chromosome 11 in a Ewing sarcoma-derived cell line. *Molec. Cell. Biol.* 5: 427–429, 1985.
- [67467] 5797. Gill, S.; McManus, A. P.; Crew, A. J.; Benjamin, H.; Sheer, D.; Gusterson, B. A.; Pinkerton, C. R.; Patel, K.; Cooper, C. S.; Shipley, J. M.: Fusion of the EWS gene to a DNA segment from 9q22–31 in a human myxoid chondrosarcoma. *Genes Chromosomes Cancer* 12: 307–310, 1995.
- [67468] 5798. Ginsberg, J. P.; de Alava, E.; Ladanyi, M.; Wexler, L. H.; Kovar, H.; Paulussen, M.; Zoubek, A.; Dockhorn-Dworniczak, B.; Juergens, H.; Wunder, J. S.; Andrulis, I. L.; Malik, R.; Sorensen, P. H. B.; Womer, R. B.; Barr, F. G.: EWS-FLI1 and EWS-ERG gene fusions are associated with similar clinical phenotypes in Ewing's sarcoma. *J. Clin. Oncol.* 17: 1809–1814, 1999.
- [67469] 5799. Griffin, C. A.; McKeon, C.; Israel, M. A.; Geggion, A.; Ghysdael, J.; Stehelin, D.; Douglass, E. C.; Green, A. A.; Emanuel, B. S.: Comparison of constitutional and tumor-



associated 11;22 translocations:nonidentical breakpoints on chromosomes 11 and 22. Proc. Nat. Acad.Sci. 83: 6122–6126, 1986.

- [67470] 5800.Jeon, I.-S.; Davis, J. N.; Braun, B. S.; Sublett, J. E.; Roussel,M. F.; Denny, C. T.; Shapiro, D. N.: A variant Ewing's sarcoma translocation(7;22) fuses the EWS gene to the ETS gene ETV1. Oncogene 10: 1229–1234,1995.
- [67471] 5801.Maletz, N.; McMorrow, L. E.; Greco, M. A.; Wolman, S. R.: Ewing'ssarcoma: pathology, tissue culture, and cytogenetics. Cancer 58:252–257, 1986.
- [67472] 5802.Mastrangelo, T.; Modena, P.; Tornielli, S.; Bullrich, F.; Testi,M. A.; Mezzelani, A.; Radice, P.; Azzarelli, A.; Pilotti, S.; Croce,C. M.; Pierotti, M. A.; Sozzi, G.: A novel zinc finger gene is fusedto EWS in small round cell tumor. Oncogene 19: 3799–3804, 2000.
- [67473] 5803.May, W. A.; Gishizky, M. L.; Lessnick, S. L.; Lunsford, L. B.;Lewis, B. C.; Delattre, O.; Zucman, J.; Thomas, G.; Denny, C. T.:Ewing sarcoma 11;22 translocation produces a chimeric transcriptionfactor that requires the DNA-binding domain encoded by FLI1 for transformation. Proc.Nat. Acad. Sci. 90: 5752–5756, 1993.
- [67474] 5804.Morohoshi, F.; Ootsuka, Y.; Arai, K.; Ichikawa, H.; Mitani, S.;Munakata, N.; Ohki, M.: Genomic structure of

the human RBP56/hTAF(II)68 and FUS/TLS genes. *Gene* 221: 191–198, 1998.

- [67475] 5805. Mugneret, F.; Aurias, A.; Lizard, S.; Turc-Carel, C.: Der(16)t(1;16)(q11;q11.1) is a consistent secondary chromosome change in Ewing's sarcoma. (Abstract) *Cytogenet. Cell Genet.* 46: 665 only, 1987.
- [67476] 5806. Mugneret, F.; Lizard, S.; Aurias, A.; Turc-Carel, C.: Chromosomes in Ewing's sarcoma. II. Nonrandom additional changes, trisomy 8 and der(16)t(1;16). *Cancer Genet. Cytogenet.* 32: 239–245, 1988.
- [67477] 5807. Ohno, T.; Ouchida, M.; Lee, L.; Gatalica, Z.; Rao, V. N.; Reddy, E. S. P.: The EWS gene, involved in Ewing family of tumors, malignant melanoma of soft parts and desmoplastic small round cell tumors, codes for an RNA binding protein with novel regulatory domains. *Oncogene* 9: 3087–3097, 1994.
- [67478] 5808. Peter, M.; Couturier, J.; Pacquement, H.; Michon, J.; Thomas, G.; Magdelenat, H.; Delattre, O.: A new member of the ETS family fused to EWS in Ewing tumors. *Oncogene* 14: 1159–1164, 1997.
- [67479] 5809. Plougastel, B.; Mattei, M.-G.; Thomas, G.; Delattre, O.: Cloning and chromosome localization of the mouse Ews gene. *Genomics* 23: 278–281, 1994.

- [67480] 5810.Plougastel, B.; Zucman, J.; Peter, M.; Thomas, G.; Delattre, O.: Genomic structure of the EWS gene and its relationship to EWSR1,a site of tumor-associated chromosome translocation. *Genomics* 18:609–615, 1993.
- [67481] 5811.Selleri, L.; Hermanson, G. G.; Eubanks, J. H.; Lewis, K. A.; Evans,G. A.: Molecular localization of the t(11;22)(q24;q12) translocationof Ewing sarcoma by chromosomal in situ suppression hybridization. *Proc.Nat. Acad. Sci.* 88: 887–891, 1991.
- [67482] 5812.Tanaka, K.; Iwakuma, T.; Harimaya, K.; Sato, H.; Iwamoto, Y.:EWS–Fli1 antisense oligodeoxynucleotide inhibits proliferation ofhuman Ewing's sarcoma and primitive neuroectodermal tumor cells. *J.Clin. Invest.* 99: 239–247, 1997.
- [67483] 5813.Tilly, H.; Bastard, C.; Chevallier, B.; Halkin, E.; Monconduit,M.: Chromosomal abnormalities in secondary Ewing's sarcoma. (Letter) *Lancet* II:812 only, 1984.
- [67484] 5814.Triche, T. J.; Askin, F. B.: Neuroblastoma and the differentialdiagnosis of small-, round-, blue-cell tumors. *Hum. Path.* 14: 569–595,1983.
- [67485] 5815.Turc–Carel, C.; Aurias, A.; Mugneret, F.; Lizard, S.; Sidaner,I.; Philip, I.; Philip, T.; Lenoir, G. M.; Mazabraud, A.: Outstandingconsistency of the chromosomal break–

points 11q24 and 22q12 in Ewing'ssarcoma. (Abstract)  
Cytogenet. Cell Genet. 46: 706 only, 1987.

[67486] 5816.Turc-Carel, C.; Aurias, A.; Mugneret, F.; Lizard, S.;  
Sidaner,I.; Volk, C.; Thiery, J. P.; Olschwang, S.; Philip, I.;  
Berger, M.P.; Philip, T.; Lenoir, G. M.; Mazabraud, A.:  
Chromosomes in Ewing'ssarcoma. I. An evaluation of 85  
cases and remarkable consistency oft(11;22)(q24;q12).  
Cancer Genet. Cytogenet. 32: 229-238, 1988.

[67487] 5817.Turc-Carel, C.; Philip, I.; Berger, M.-P.; Philip, T.;  
Lenoir,G. M.: Chromosomal translocations in Ewing's sar-  
coma. New Eng. J.Med. 309: 497-498, 1983.

[67488] 5818.Turc-Carel, C.; Philip, I.; Berger, M.-P.; Philip, T.;  
Lenoir,G. M.: Chromosome study of Ewing's sarcoma (ES)  
cell lines: consistencyof a reciprocal translocation  
t(11;22)(q24;q12). Cancer Genet. Cytogenet. 12:1-19,  
1984.

[67489] 5819.Whang-Peng, J.; Freter, C. E.; Knutsen, T.; Nanfro, J.  
J.; Gazdar,A.: Translocation t(11;22) in esthesioneuroblas-  
toma. Cancer Genet.Cytogenet. 29: 155-157, 1987.

[67490] 5820.Whang-Peng, J.; Triche, T. J.; Knutsen, T.; Miser, J.;  
Douglass,E. C.; Israel, M. A.: Chromosome translocation in  
peripheral neuroepithelioma. NewEng. J. Med. 311:  
584-585, 1984.

- [67491] 5821.Zhang, R.-Z.; Pan, T.-C.; Zhang, Z.-Y.; Mattei, M.-G.; Timpl, R.;Chu, M.-L.: Fibulin-2 (FBLN2): human cDNA sequence, mRNA expression,and mapping of the gene on human and mouse chromosomes. *Genomics* 22:425-430, 1994.
- [67492] 5822.Baden, H. P.; Roth, S. I.; Goldsmith, L. A.; Lee, L. D.: Keratohyalinprotein in disorders of keratinization. *J. Invest. Derm.* 62: 411-414,1974.
- [67493] 5823.Gan, S.-Q.; McBride, O. W.; Idler, W. W.; Markova, N.; Steinert,P. M.: Organization, structure, and polymorphisms of the human profilaggringene. *Biochemistry* 29: 9432-9440, 1990.
- [67494] 5824.Holbrook, K. A.; Dale, B. A.; Brown, K. S.: Abnormal epidermalkeratinization in the repeated epilation mutant mouse. *J. Cell Biol.* 92:387-397, 1982.
- [67495] 5825.McKinley-Grant, L. J.; Idler, W. W.; Bernstein, I. A.; Parry, D.A. D.; Cannizzaro, L.; Croce, C. M.; Huebner, K.; Lessin, S. R.; Steinert,P. M.: Characterization of a cDNA clone encoding human filaggrinand localization of the gene to chromosome region 1q21. *Proc. Nat.Acad. Sci.* 86: 4848-4852, 1989.
- [67496] 5826.Rothnagel, J. A.; Longley, M. A.; Bundman, D. S.; Naylor, S. L.;Lalley, P. A.; Jenkins, N. A.; Gilbert, D. J.;

Copeland, N. G.; Roop, D. R.: Characterization of the mouse loricrin gene: linkage with profilaggrin and the flaky tail and soft coat mutant loci on chromosome 3. *Genomics* 23: 450–456, 1994.

[67497] 5827. Sybert, V. P.; Dale, B. A.; Holbrook, K. A.: Ichthyosis vulgaris: identification of a defect in synthesis of filaggrin correlated with an absence of keratohyaline granules. *J. Invest. Derm.* 84: 191–194, 1985.

[67498] 5828. Volz, A.; Korge, B. P.; Compton, J. G.; Ziegler, A.; Steinert, P. M.; Mischke, D.: Physical mapping of a functional cluster of epidermal differentiation genes on chromosome 1q21. *Genomics* 18: 92–99, 1993.

[67499] 5829. Barbosa, C. A. A.; Koury, W. H.; Krieger, H.: Linkage data on MN and the Hb beta locus. *Am. J. Hum. Genet.* 27: 797–801, 1975.

[67500] 5830. Riddell, D. C.; Wang, H.; Umbenhauer, D. R.; Beaume, P. H.; Guengerich, F. P.; Hamerton, J. L.: Regional assignment for the genes encoding human P450III A3 (CYP3) and P450II C9 (CYP2C). (Abstract) *Cytogenet. Cell Genet.* 46: 682, 1987.

[67501] 5831. Yoder, J. A.; Yen, R.-W. C.; Vertino, P. M.; Bestor, T. H.; Baylin, S. B.: New 5-prime regions of the murine and human genes for DNA (cytosine-5)-methyltransferase. *J.*

Biol. Chem. 271: 31092–31097,1996.

[67502] 5832.Hofmann, S.; Lichtner, P.; Schuffenhauer, S.; Gerbitz, K.-D.; Meitinger,T.: Assignment of the human genes coding for cytochrome c oxidasesubunits Va (COX5A), VIc (COX6C) and VIIc (COX7C) to chromosome bands15q25, 8q22–q23 and 5q14 and of three pseudogenes (COX5AP1, COX6CP1,COX7CP1) to 14q22, 16p12 and 13q14–q21 by FISH and radiation hybridmapping. Cytogenet. Cell Genet. 83: 226–227, 1998.

[67503] 5833.Martin–Gallardo, A.; McCombie, W. R.; Gocayne, J. D.; FitzGerald,M. G.; Wallace, S.; Lee, B. M. B.; Lamerdin, J.; Trapp, S.; Kelley,J. M.; Liu, L.–I.; Dubnick, M.; Johnston–Dow, L. A.; Kerlavage, A.R.; de Jong, P.; Carrano, A.; Fields, C.; Venter, J. C.: AutomatedDNA sequencing and analysis of 106 kilobases from human chromosome19q13.3. Nature Genet. 1: 34–39, 1992.

[67504] 5834.Schellens, J. H. M.; Soons, P. A.; Breimer, D. D.: Lack of bimodalityin nifedipine plasma kinetics in a large population of healthy subjects. Biochem.Pharm. 37: 2507–2510, 1988.

[67505] 5835.Shet, M. S.; Fisher, C. W.; Holmans, P. L.; Estabrook, R. W.:Human cytochrome P450 3A4: enzymatic properties of a purified recombinantfusion protein containing

NADPH-P450 reductase. Proc. Nat. Acad.Sci. 90:  
11748-11752, 1993.

[67506] 5836.Shimada, T.; Guengerich, F. P.: Evidence for cytochrome P-450(NF),the nifedipine oxidase, being the principal enzyme involved in thebioactivation of aflatoxins in human liver. Proc. Nat. Acad. Sci. 86:462-465, 1989.

[67507] 5837.Spurr, N. K.; Gough, A. C.; Stevenson, K.; Wolf, C. R.: The humancytochrome P450 CYP3 locus: assignment to chromosome 7q22-qter. Hum.Genet. 81: 171-174, 1989.

[67508] 5838.Walker, A. H.; Jaffe, J. M.; Gunasegaram, S.; Cummings, S. A.;Huang, C.-S.; Chern, H.-D.; Olopade, O. I.; Weber, B. L.; Rebbeck,T. R.: Characterization of an allelic variant in the nifedipine-specific element of CYP3A4: ethnic distribution and implications for prostatecancer risk. Hum. Mutat. 12: 289-293, 1998.

[67509] 5839.Watkins, P. B.; Wrighton, S. A.; Maurel, P.; Schuetz, E. G.; Mendez-Picon,G.; Parker, G. A.; Guzelian, P. S.: Identification of an inducible form of cytochrome P-450 in human liver. Proc. Nat. Acad. Sci. 82:6310-6314, 1985.

[67510] 5840.Wolf, C. R.; Smith, C. A. D.; Gough, A. C.; Moss, J. E.; Vallis,K. A.; Howard, G.; Carey, F. J.; Mills, K.; McNee, W.; Carmichael,J.; Spurr, N. K.: Relationship between the debrisoquine hydroxylase polymorphism and cancer suscep-



tibility. *Carcinogenesis* 13: 1035–1038,1992.

- [67511] 5841.Wrighton, S. A.; Stevens, J. C.: The human hepatic cytochromesP450 involved in drug metabolism. *Crit. Rev. Toxicol.* 22: 1–21,1992.
- [67512] 5842.Wrighton, S. A.; Vandenbranden, M.: Isolation and characterizationof human fetal liver cytochrome P450HLp2: a third member of the P450IIIgene family. *Arch. Biochem. Biophys.* 268: 144–151, 1989.
- [67513] 5843.Xie, W.; Barwick, J. L.; Downes, M.; Blumberg, B.; Simon, C. M.;Nelson, M. C.; Neuschwander–Tetri, B. A.; Brunt, E. M.; Guzelian,P. S.; Evans, R. M.: Humanized xenobiotic response in mice expressingnuclear receptor SXR. *Nature* 406: 435–439, 2000.
- [67514] 5844.Gray, I. C.; Nobile, C.; Muresu, R.; Ford, S.; Spurr, N. K.: A2.4–megabase physical map spanning the CYP2C gene cluster on chromosome10q24. *Genomics* 28: 328–332, 1995.
- [67515] 5845.Guengerich, F. P.; Distlerath, L. M.; Reilly, P. E. B.; Wolff,T.; Shimada, T.; Umbenhauer, D. R.; Martin, M. V.: Human–liver cytochromesP–450 involved in polymorphisms of drug oxidation. *Xenobiotica* 16:367–378, 1986.
- [67516] 5846.Lenka, N.; Vijayasathy, C.; Mullick, J.; Avadhani, N. G.: Structuralorganization and transcription regulation of

nuclear genes encoding the mammalian cytochrome c oxidase complex. *Prog. Nucleic Acid Res. Molec. Biol.* 61: 309–344, 1998.

- [67517] 5847. Otsuka, M.; Mizuno, Y.; Yoshida, M.; Kagawa, Y.; Ohta, S.: Nucleotide sequence of cDNA encoding human cytochrome c oxidase subunit VIc. *Nucleic Acids Res.* 16: 10916 only, 1988.
- [67518] 5848. Akbari, O.; Freeman, G. J.; Meyer, E. H.; Greenfield, E. A.; Chang, T. T.; Sharpe, A. H.; Berry, G.; DeKruyff, R. H.; Umetsu, D. T.: Antigen-specific regulatory T cells develop via the ICOS-ICOS-ligand pathway and inhibit allergen-induced airway hyperreactivity. *Nature Med.* 8: 1024–1032, 2002.
- [67519] 5849. Cox, D. R.; Burmeister, M.; Price, E. R.; Kim, S.; Myers, R. M.: Radiation hybrid mapping: a somatic cell genetic method for constructing high-resolution maps of mammalian chromosomes. *Science* 250: 245–250, 1990.
- [67520] 5850. Crawley, J. B.; Williams, L. M.; Mander, T.; Brennan, F. M.; Foxwell, B. M.: Interleukin-10 stimulation of phosphatidylinositol 3-kinase and p70 S6 kinase is required for the proliferative but not the antiinflammatory effects of the cytokine. *J. Biol. Chem.* 271: 16357–16362, 1996.
- [67521] 5851. Eskdale, J.; Gallagher, G.; Verweij, C. L.; Keijsers, V.;

Westendorp, R. G. J.; Huizinga, T. W. J.: Interleukin 10 secretion in relation to human IL-10 locus haplotypes. *Proc. Nat. Acad. Sci.* 95: 9465–9470, 1998.

[67522] 5852. Eskdale, J.; Kube, D.; Tesch, H.; Gallagher, G.: Mapping of the human IL10 gene and further characterization of the 5-prime flanking sequence. *Immunogenetics* 46: 120–128, 1997.

[67523] 5853. Farmer, M. A.; Sundberg, J. P.; Bristol, I. J.; Churchill, G. A.; Li, R.; Elson, C. O.; Leiter, E. H.: A major quantitative trait locus on chromosome 3 controls colitis severity in IL-10-deficient mice. *Proc. Nat. Acad. Sci.* 98: 13820–13825, 2001.

[67524] 5854. Franchimont, D.; Martens, H.; Hagelstein, M.-T.; Louis, E.; Dewe, W.; Chrousos, G. P.; Belaiche, J.; Geenen, V.: Tumor necrosis factor  $\alpha$  decreases, and interleukin-10 increases, the sensitivity of human monocytes to dexamethasone: potential regulation of the glucocorticoid receptor. *J. Clin. Endocr. Metab.* 84: 2834–2839, 1999.

[67525] 5855. Gesser, B.; Leffers, H.; Jinqun, T.; Vestergaard, C.; Kirstein, N.; Sindet-Pedersen, S.; Jensen, S. L.; Thstrup-Pedersen, K.; Larsen, C. G.: Identification of functional domains on human interleukin 10. *Proc. Nat. Acad. Sci.* 94:

14620–14625, 1997.

- [67526] 5856. Gibson, A. W.; Edberg, J. C.; Wu, J.; Westendorp, R. G. J.; Huizinga, T. W. J.; Kimberly, R. P.: Novel single nucleotide polymorphisms in the distal IL-10 promoter affect IL-10 production and enhance the risk of systemic lupus erythematosus. *J. Immun.* 166: 3915–3922, 2001.
- [67527] 5857. Goudy, K.; Song, S.; Wasserfall, C.; Zhang, Y. C.; Kapturczak, M.; Muir, A.; Powers, M.; Scott-Jorgensen, M.; Campbell-Thompson, M.; Crawford, J. M.; Ellis, T. M.; Flotte, T. R.; Atkinson, M. A.: Adeno-associated virus vector-mediated IL-10 gene delivery prevents type 1 diabetes in NOD mice. *Proc. Nat. Acad. Sci.* 98: 13913–13918, 2001.
- [67528] 5858. Grove, J.; Daly, A. K.; Bassendine, M. F.; Gilvarry, E.; Day, C. P.: Interleukin 10 promoter region polymorphisms and susceptibility to advanced alcoholic liver disease. *Gut* 46: 540–545, 2000.
- [67529] 5859. Kim, J. M.; Brannan, C. I.; Copeland, N. G.; Jenkins, N. A.; Khan, T. A.; Moore, K. W.: Structure of the mouse IL-10 gene and chromosomal localization of the mouse and human genes. *J. Immun.* 148: 3618–3623, 1992.
- [67530] 5860. Kitagawa, N.; Goto, M.; Kurozumi, K.; Maruo, S.; Fukayama, M.; Naoe, T.; Yasukawa, M.; Hino, K.; Suzuki, T.;

Todo, S.; Takada, K.: Epstein–Barr virus–encoded poly(A)–RNA supports Burkitt's lymphomagrowth through interleukin–10 induction. EMBO J. 19: 6742–6750,2000.

[67531] 5861.Kuhn, R.; Lohler, J.; Rennick, D.; Rajewsky, K.;

Muller, K.:Interleukin–10–deficient mice develop chronic enterocolitis. Cell 75:263–274, 1993.

[67532] 5862.Lee, T.–S.; Chau, L.–Y.: Heme oxygenase–1 mediates the anti–inflammatoryeffect of interleukin–10 in mice. Nature Med. 8: 240–246, 2002.

[67533] 5863.Meng, X.; Sawamura, D.; Tamai, K.; Hanada, K.;

Ishida, H.; Hashimoto,I.: Keratinocyte gene therapy for systemic diseases: circulatinginterleukin 10 released from gene–transferred keratinocytes inhibitscontact hypersensitivity at distant areas of the skin. J. Clin. Invest. 101:1462–1467, 1998.

[67534] 5864.Pinderski Oslund, L. J.; Hedrick, C. C.; Olvera, T.;

Hagenbaugh,A.; Territo, M.; Berliner, J. A.; Fyfe, A. I.: Interleukin–10 blocksatherosclerotic events in vitro and in vivo. Arterioscler. Thromb.Vasc. Biol. 19: 2847–2853, 1999.

[67535] 5865.Rosenwasser, L. J.; Borish, L.: Genetics of atopy and asthma:the rationale behind promoter–based candidate gene studies (IL–4 andIL–10). Am. J. Resp. Crit. Care Med.

156: S152–S155, 1997.

- [67536] 5866. Shin, H. D.; Winkler, C.; Stephens, J. C.; Bream, J.; Young, H.; Goedert, J. J.; O'Brien, T. R.; Vlahov, D.; Buchbinder, S.; Giorgi, J.; Rinaldo, C.; Donfield, S.; Willoughby, A.; O'Brien, S. J.; Smith, M. W.: Genetic restriction of HIV-1 pathogenesis to AIDS by promoter alleles of IL10. *Proc. Nat. Acad. Sci.* 97: 14467–14472, 2000.
- [67537] 5867. Terkeltaub, R. A.: IL-10: an 'immunologic scalpel' for atherosclerosis? (Editorial) *Atheroscler. Thromb. Vasc. Biol.* 19: 2823–2825, 1999.
- [67538] 5868. Turner, D. M.; Williams, D. M.; Sankaran, D.; Lazarus, M.; Sinnott, P. J.; Hutchinson, I. V.: An investigation of polymorphism in the interleukin-10 gene promoter. *Europ. J. Immunogenet.* 24: 1–8, 1997.
- [67539] 5869. Vieira, P.; de Waal-Malefyt, R.; Dang, M.-N.; Johnson, K. E.; Kastelein, R.; Fiorentino, D. F.; de Vries, J. E.; Roncarolo, M.-G.; Mosmann, T. R.; Moore, K. W.: Isolation and expression of human cytokine synthesis inhibitory factor cDNA clones: homology to Epstein-Barr virus open reading frame BCRF1. *Proc. Nat. Acad. Sci.* 88: 1172–1176, 1991.
- [67540] 5870. Westendorp, R. G. J.; Langermans, J. A. M.; Huizinga, T. W. G.; Elouali, A. H.; Boomsma, D. I.; Verweij, C. L.; Van-

denbroucke, J.P.: Genetic influence on cytokine production and fatal meningococcal disease. *Lancet* 349: 170–173, 1997.

[67541] 5871. Armstrong, E.; Cannizzaro, L.; Bergman, M.; Huebner, K.; Alitalo, K.: The c-src tyrosine kinase (CSK) gene, a potential antioncogene, localizes to human chromosome region 15q23–q25. *Cytogenet. Cell Genet.* 60:119–120, 1992.

[67542] 5872. Cloutier, J.-F.; Veillette, A.: Association of inhibitory tyrosine protein kinase p50(csk) with protein tyrosine phosphatase PEP in T cells and other hemopoietic cells. *EMBO J.* 15: 4909–4918, 1996.

[67543] 5873. Partanen, J.; Armstrong, E.; Bergman, M.; Makela, T. P.; Hirvonen, H.; Huebner, K.; Alitalo, K.: Cyl encodes a putative cytoplasmic tyrosine kinase lacking the conserved tyrosine autophosphorylation site (Y416-src). *Oncogene* 6: 2013–2018, 1991.

[67544] 5874. Choubey, D.; Snoddy, J.; Chaturvedi, V.; Toniato, E.; Opdenakker, G.; Thakur, A.; Samanta, H.; Engel, D. A.; Lengyel, P.: Interferon  $\alpha$  gene activators: indications for repeated gene duplication during the evolution of a cluster of interferon-activatable genes on murine chromosome 1. *J. Biol. Chem.* 264: 17182–17189, 1989.

- [67545] 5875.Ahuja, H. G.; Felix, C. A.; Aplan, P. D.: The t(11;20)(p15;q11)chromosomal translocation associated with therapy-related myelodysplasticsyndrome results in an NUP98-TOP1 fusion. *Blood* 94: 3258-3261, 1999.
- [67546] 5876.Zhang, F. R.; Delattre, O.; Rouleau, G.; Couturier, J.; Lefrancois,D.; Thomas, G.; Aurias, A.: The neuroepithe- lioma breakpoint on chromosome22 is proximal to the meningioma locus. *Genomics* 6: 174-177, 1990.
- [67547] 5877.Siciliano, M. J.; Bachinski, L.; Dolf, G.; Carrano, A. V.; Thompson,L. H.: Chromosomal assignments of human DNA repair genes that complementChinese hamster ovary (CHO) cell mutants. (Abstract) *Cytogenet. CellGenet.* 46: 691-692, 1987.
- [67548] 5878.Thompson, L. H.; Carrano, A. V.; Sato, K.; Salazar, E. P.; White,B. F.; Stewart, S. A.; Minkler, J. L.; Siciliano, M. J.: Identificationof nucleotide-excision-repair genes on hu- man chromosomes 2 and 13by functional complementa- tion in hamster-human hybrids. *Somat. CellMolec. Genet.* 13: 539-551, 1987.
- [67549] 5879.Sul, H. S.; Wise, L. S.; Brown, M. L.; Rubin, C. S.: Cloning ofcDNA sequences for murine ATP-citrate lyase: construction of recombinantplasmids using an immunop- urified mRNA template and evidence for thenutritional



regulation of ATP-citrate lyase mRNA content in mouse-liver. J. Biol. Chem. 259: 1201–1205, 1984.

- [67550] 5880.Brandt, P.; Ibrahim, E.; Bruns, G. A. P.; Neve, R. L.: Determination of the nucleotide sequence and chromosomal localization of the ATP2B2 gene encoding human Ca(2+)-pumping ATPase isoform PMCA2. Genomics 14:484–487, 1992.
- [67551] 5881.Kozel, P. J.; Friedman, R. A.; Erway, L. C.; Yamoah, E. N.; Liu, L. H.; Riddle, T.; Duffy, J. J.; Doetschman, T.; Miller, M. L.; Cardell, E. L.; Shull, G. E.: Balance and hearing deficits in mice with a null mutation in the gene encoding plasma membrane Ca(2+)-ATPase isoform 2. J. Biol. Chem. 273: 18693–18696, 1998.
- [67552] 5882.Richards, F. M.; Phipps, M. E.; Latif, F.; Yao, M.; Crossey, P. A.; Foster, K.; Linehan, W. M.; Affara, N. A.; Lerman, M. I.; Zbar, B.; Ferguson-Smith, M. A.; Maher, E. R.: Mapping the von Hippel-Lindau disease tumour suppressor gene: identification of germline deletions by pulsed field gel electrophoresis. Hum. Molec. Genet. 2: 879–882, 1993.
- [67553] 5883.Street, V. A.; McKee-Johnson, J. W.; Fonseca, R. C.; Tempel, B. L.; Noben-Trauth, K.: Mutations in a plasma membrane Ca(2+)-ATPase gene cause deafness in deaf-

waddler mice. *Nature Genet.* 19: 390–394,1998.

[67554] 5884.Wang, M. G.; Yi, H.; Hilfiker, H.; Carafoli, E.; Strehler, E. E.;McBride, O. W.: Localization of two genes encoding plasma membraneCa(2+)-ATPases isoforms 2 (ATP2B2) and 3 (ATP2B3) to human chromosomes3p26-p25 and Xq28, respectively. *Cytogenet. Cell Genet.* 67:

41–45,1994.

[67555] 5885.Shi, G.-P.; Munger, J. S.; Meara, J. P.; Rich, D. H.; Chapman,H. A.: Molecular cloning and expression of human alveolar macrophagecathepsin S, an elastinolytic cysteine protease. *J. Biol. Chem.* 267:7258–7262, 1992.

[67556] 5886.Shi, G.-P.; Webb, A. C.; Foster, K. E.; Knoll, J. H. M.; Lemere,C. A.; Munger, J. S.; Chapman, H. A.: Human cathepsin S: chromosomallocalization, gene structure, and tissue distribution. *J. Biol. Chem.* 269:11530–11536, 1994.

[67557] 5887.Grundmann, U.; Nerlich, C.; Rein, T.; Zettlmeissl, G.: CompletecDNA sequence encoding human beta-galactoside alpha-2,6-sialyltransferase. *NucleicAcids Res.* 18: 667 only, 1990.

[67558] 5888.Wang, X.; Vertino, A.; Eddy, R. L.; Byers, M. G.; Jani-Sait, S.N.; Shows, T. B.; Lau, J. T. Y.: Chromosome mapping and organizationof the human beta-galactoside al-

pha-2,6-sialyltransferase gene: differential and cell-type specific usage of upstream exon sequences in B-lymphoblastoid cells. *J. Biol. Chem.* 268: 4355–4361, 1993.

- [67559] 5889. Xiang, M.; Zhou, L.; Macke, J. P.; Yoshioka, T.; Hendry, S. H. C.; Eddy, R. L.; Shows, T. B.; Nathans, J.: The Brn-3 family of POU-domain factors: primary structure, binding specificity, and expression in subsets of retinal ganglion cells and somatosensory neurons. *J. Neurosci.* 15:4762–4785, 1995.
- [67560] 5890. Bouillaud, F.; Villarroya, F.; Hentz, E.; Raimbault, S.; Cassard, A.-M.; Ricquier, D.: Detection of brown adipose tissue uncoupling protein mRNA in adult patients by a human genomic probe. *Clin. Sci.* 75:21–27, 1988.
- [67561] 5891. Bouillaud, F.; Weissenbach, J.; Ricquier, D.: Complete cDNA-derived amino acid sequence of rat brown fat uncoupling protein. *J. Biol. Chem.* 261: 1487–1491, 1986.
- [67562] 5892. Cassard, A. M.; Bouillaud, F.; Mattei, M. G.; Hentz, E.; Raimbault, S.; Thomas, M.; Ricquier, D.: Human uncoupling protein gene: structure, comparison with rat gene, and assignment to the long arm of chromosome
- [67563] 5893. Echtay, K. S.; Roussel, D.; St-Pierre, J.; Jekabsons, M. B.; Cadenas, S.; Stuart, J. A.; Harper, J. A.; Roebuck, S. J.;

Morrison, A.; Pickering, S.; Clapham, J. C.; Brand, M. D.:  
Superoxide activates mitochondrial uncoupling proteins.  
Nature 415: 96–99, 2002.

[67564] 5894. Echtay, K. S.; Winkler, E.; Klingenberg, M.: Coenzyme  
Q is an obligatory cofactor for uncoupling protein function.  
Nature 408: 609–613, 2000.

[67565] 5895. J. Cell. Biochem. 43: 255–264, 1990. 4. Clement, K.;  
Ruiz, J.; Cassard-Doulcier, A. M.; Bouillaud, F.; Ricquier, D.;  
Basdevant, A.; Guy-Grand, B.; Froguel, P.: Additive effect of  
A-G (–3826) variant of the uncoupling protein gene and  
the trp64arg mutation of the beta 3-adrenergic receptor  
gene on weight gain in morbid obesity. Int. J. Obes. Relat.  
Metab. Disord. 20: 1062–1066, 1996.

[67566] 5896. Enerback, S.; Jacobsson, A.; Simpson, E. M.; Guerra,  
C.; Yamashita, H.; Harper, M.-E.; Kozak, L. P.: Mice lacking  
mitochondrial uncoupling protein are cold-sensitive but  
not obese. Nature 387: 90–93, 1997.

[67567] 5897. Fletcher, C.; Norman, D. J.; Germond, E.; Heintz, N.:  
A multilocus linkage map of mouse chromosome 8. Ge-  
nomics 9: 737–741, 1991.

[67568] 5898. Jacobson, A.; Stadler, U.; Glotzer, M. A.; Kozak, L. P.:  
Mitochondrial uncoupling protein from mouse brown fat:  
molecular cloning, genetic mapping and mRNA expression.

J. Biol. Chem. 260: 16250–16254, 1985.

- [67569] 5899. Lowell, B. B.; S-Susulic, V.; Hamann, A.; Lawitts, J. A.; Himms-Hagen, J.; Boyer, B. B.; Kozak, L. P.; Flier, J. S.: Development of obesity in transgenic mice after genetic ablation of brown adipose tissue. *Nature* 366:740–742, 1993.
- [67570] 5900. Oppert, J. M.; Vohl, M. C.; Chagnon, M.; Dionne, F. T.; Cassard-Doucier, A. M.; Ricquier, D.; Perusse, L.; Bouchard, C.: DNA polymorphism in the uncoupling protein (UCP) gene and human body fat. *Int. J. Obes. Relat. Metab. Disord.* 18: 526–531, 1994.
- [67571] 5901. Ridley, R. G.; Patel, H. V.; Gerber, G. E.; Morton, R. C.; Freeman, K. B.: Complete nucleotide and derived amino acid sequence of cDNA encoding the mitochondrial uncoupling protein of rat brown adipose tissue: lack of mitochondrial targeting presequence. *Nucleic Acids Res.* 14: 4025–4035, 1986.
- [67572] 5902. Urhammer, S. A.; Fridberg, M.; Sorensen, T. I.; Echwald, S. M.; Andersen, T.; Tybjaerg-Hansen, A.; Clausen, J. O.; Pedersen, O.: Studies of genetic variability of the uncoupling protein 1 gene in Caucasian subjects with juvenile-onset obesity. *J. Clin. Endocr. Metab.* 82: 4069–4074, 1997.

- [67573] 5903.Rodriguez de Cordoba, S.; Rubinstein, P.; Ferreira, A.: Highresolution isoelectric focusing of immunoprecipitated proteins underdenaturing conditions: a simple analytical method applied to the studyof complement component polymorphisms. *J. Immun. Methods* 69: 165–172,1984.
- [67574] 5904.Muller, U.; Kupke, K. G.: The genetics of primary torsion dystonia. *Hum.Genet.* 84: 107–115, 1990.
- [67575] 5905.Di Marzo, V.; Goparaju, S. K.; Wang, L.; Liu, J.; Batkai, S.; Jarai,Z.; Fezza, F.; Miura, G. I.; Palmiter, R. D.; Sugiura, T.; Kunos,G.: Leptin–regulated endocannabinoids are involved in maintainingfood intake. *Nature* 410: 822–825, 2001.
- [67576] 5906.Gerard, C. M.; Mollereau, C.; Vassart, G.; Parmentier, M.: Molecularcloning of a human cannabinoid receptor which is also expressed intestis. *Biochem. J.* 279: 129–134, 1991.
- [67577] 5907.Hoehe, M. R.; Caenazzo, L.; Martinez, M. M.; Hsieh, W.–T.; Modi,W. S.; Gershon, E. S.; Bonner, T. I.: Genetic and physical mappingof the human cannabinoid receptor gene to chromosome 6q14–q15. *NewBiologist* 3: 880–885, 1991.
- [67578] 5908.Ledent, C.; Valverde, O.; Cossu, G.; Petitet, F.;

Aubert, J.-F.; Beslot, F.; Bohme, G. A.; Imperato, A.; Pedrazzini, T.; Roques, B.P.; Vassart, G.; Fratta, W.; Parmentier, M.: Unresponsiveness to cannabinoids and reduced addictive effects of opiates in CB(1) receptor knockout mice. *Science* 283 401–404, 1999.

[67579] 5909. Marsicano, G.; Wotjak, C. T.; Azad, S. C.; Bisogno, T.; Rammes, G.; Cascio, M. G.; Hermann, H.; Tang, J.; Hofmann, C.; Zieglgansberger, W.; Di Marzo, V.; Lutz, B.: The endogenous cannabinoid system controls extinction of aversive memories. *Nature* 418: 530–534, 2002.

[67580] 5910. Matsuda, L. A.; Lolait, S. J.; Brownstein, M. J.; Young, A. C.; Bonner, T. I.: Structure of a cannabinoid receptor and functional expression of the cloned cDNA. *Nature* 346: 561–564, 1990.

[67581] 5911. Modi, W. S.; Bonner, T. I.: Localization of the cannabinoid (sic) receptor locus using non-isotopic in situ hybridization. (Abstract) *Cytogenet. Cell Genet.* 58: 1915 only, 1991.

[67582] 5912. Panikashvili, D.; Simeonidou, C.; Ben-Shabat, S.; Hanus, L.; Breuer, A.; Mechoulam, R.; Shohami, E.: An endogenous cannabinoid (2-AG) is neuroprotective after brain injury. *Nature* 413: 527–531, 2001.

[67583] 5913. Beechey, C.; Tweedie, S.; Spurr, N.; Ball, S.; Peters, J.;

Edwards,Y.: Mapping of mouse carbonic anhydrase-3, Car-3: another locus in the homologous region of mouse chromosome 3 and human chromosome 8. *Genomics* 6:692-696, 1990.

[67584] 5914.Carter, N.; Jeffery, S.; Shiels, A.; Edwards, Y.; Tipler, T.; Hopkinson,D. A.: Characterization of human carbonic anhydrase III from skeletal muscle. *Biochem. Genet.* 17: 837-854, 1979.

[67585] 5915.Edwards, Y. H.; Lloyd, J.; Parkar, M.; Povey, S.: Human muscle specific carbonic anhydrase, CA3, is on chromosome 8. (Abstract) *Cytogenet.Cell Genet.* 40: 621 only, 1985.

[67586] 5916.Edwards, Y. H.; Lloyd, J. C.; Parkar, M.; Povey, S.: The gene for human muscle specific carbonic anhydrase (CAIII) is assigned to chromosome 8. *Ann. Hum. Genet.* 50: 41-47, 1986.

[67587] 5917.Heath, R.; Carter, N. D.; Jeffery, S.; Edwards, R. J.; Watts, D.C.; Watts, R. L.: Evaluation of carrier detection of Duchenne muscular dystrophy using carbonic anhydrase III and creatine kinase. *Am. J.Med. Genet.* 21: 291-296, 1985.

[67588] 5918.Lloyd, J.; Brownson, C.; Tweedie, S.; Charlton, J.; Edwards, Y.H.: Human muscle carbonic anhydrase: gene



structure and DNA methylation patterns in fetal and adult tissues. *Genes Dev.* 1: 594–602, 1987.

[67589] 5919. Lloyd, J.; McMillan, S.; Hopkinson, D.; Edwards, Y. H.: Nucleotide sequence and derived amino acid sequence of a cDNA encoding human muscle carbonic anhydrase. *Gene* 41: 233–239, 1986.

[67590] 5920. Lloyd, J. C.; Isenberg, H.; Hopkinson, D. A.; Edwards, Y. H.: Isolation of a cDNA clone for the human muscle specific carbonic anhydrase, CA III. *Ann. Hum. Genet.* 49: 241–251, 1985.

[67591] 5921. Wade, R.; Gunning, P.; Eddy, R.; Shows, T.; Kedes, L.: Nucleotide sequence, tissue-specific expression, and chromosome location of human carbonic anhydrase III: the human CAIII gene is located on the same chromosome as the closely linked CAI and CAII genes. *Proc. Nat. Acad. Sci.* 83: 9571–9575, 1986.

[67592] 5922. Hall, C.; Monfries, C.; Smith, P.; Lim, H. H.; Kozma, R.; Ahmed, S.; Vanniasingham, V.; Leung, T.; Lim, L.: Novel human brain cDNA encoding a 34,000 M(r) protein n-chimaerin, related to both the regulatory domain of protein kinase C and BCR, the product of the breakpoint cluster region gene. *J. Molec. Biol.* 211: 11–16, 1990.

[67593] 5923. Hall, C.; Sin, W. C.; Teo, M.; Michael, G. J.; Smith, P.;

Dong,J. M.; Lim, H. H.; Manser, E.; Spurr, N. K.; Jones, T. A.; Lim, L.: Alpha-2-chimerin, an SH2-containing GTPase-activating protein for the ras-related protein p21-rac derived by alternate splicing of the human  $\alpha$ -chimerin gene, is selectively expressed in brain regions and testes. *Molec. Cell. Biol.* 13: 4986–4998, 1993.

[67594] 5924. Brewer, C.; Holloway, S.; Zawalnyski, P.; Schinzel, A.; FitzPatrick, D.: A chromosomal deletion map of human malformations. *Am. J. Hum. Genet.* 63: 1153–1159, 1998.

[67595] 5925. Brewer, C. M.; Leek, J. P.; Green, A. J.; Holloway, S.; Bonthron, D. T.; Markham, A. F.; FitzPatrick, D. R.: A locus for isolated cleft palate, located on human chromosome 2q32. *Am. J. Hum. Genet.* 65:387–396, 1999.

[67596] 5926. Carter, C. O.; Evans, K.; Coffey, R.; Roberts, J. A. F.; Buck, A.; Roberts, M. F.: A family study of isolated cleft palate. *J. Med. Genet.* 19: 329–331, 1982.

[67597] 5927. Christensen, K.; Holm, N. V.; Olsen, J.; Kock, K.; Fogh-Andersen, P.: Selection bias in genetic-epidemiological studies of cleft lip and palate. *Am. J. Hum. Genet.* 51: 654–659, 1992.

[67598] 5928. Christensen, K.; Mitchell, L. E.: Familial recurrence-pattern analysis of nonsyndromic isolated cleft palate: a Danish registry study. *Am. J. Hum. Genet.* 58: 182–190,

1996.

- [67599] 5929.Jenkins, M.; Stady, C.: Dominant inheritance of cleft of the softpalate. Hum. Genet. 53: 341–342, 1980.
- [67600] 5930.Shields, E. D.; Bixler, D.; Fogh–Andersen, P.: Cleft palate:a genetic and epidemiologic investigation. Clin. Genet. 20: 13–24,1981.
- [67601] 5931.Hwang, S. J.; Beaty, T. H.; Panny, S. R.; Street, N. A.; Joseph,J. M.; Gordon, S.; McIntosh, I.; Francomano, C. A.: Association studyof transforming growth factor alpha (TGF–alpha) TaqI polymorphismand oral clefts: indication of gene–environment interaction in a population–basedsample of infants with birth defects. Am. J. Epi–demiol. 141: 629–636,1995.
- [67602] 5932.Van Dyke, D. C.; Goldman, A. S.; Spielman, R. S.; Zmijewski, C.M.: Segregation of HLA in families with oral clefts: evidence againstlinkage between isolated cleft palate and HLA. Am. J. Med. Genet. 15:85–88, 1983.
- [67603] 5933.Donald, L. J.; Wang, H. S.; Hamerton, J. L.: Are there additionalCKBB loci?(Abstract) Cytogenet. Cell Genet. 32: 267–268, 1982.
- [67604] 5934.McLellan, R. A.; Oscarson, M.; Alexandrie, A.–K.; Seidegard, J.;Evans, D. A. P.; Rannug, A.; Ingelman–Sundberg, M.: Characterizationof a human glutathione S–transferase

mu cluster containing a duplicated GSTM1 gene that causes ultrarapid enzyme activity. *Molec. Pharm.* 52:958–965, 1997.

- [67605] 5935. Pearson, W. R.; Vorachek, W. R.; Xu, S.; Berger, R.; Hart, I.; Vannais, D.; Patterson, D.: Identification of class-mu glutathionetransferase genes GSTM1–GSTM5 on human chromosome 1p13. *Am. J. Hum. Genet.* 53: 220–233, 1993.
- [67606] 5936. Sanchez–Corral, P.; Pardo–Manuel de Villena, F.; Rey–Campos, J.; Rodriguez de Cordoba, S.: C4BPAL1, a member of the human regulator of complement activation (RCA) gene cluster that resulted from the duplication of the gene coding for the alpha–chain of C4b–binding protein. *Genomics* 17: 185–193, 1993.
- [67607] 5937. Hillarp, A.; Dahlback, B.: Novel subunit in C4b–binding protein required for protein S binding. *J. Biol. Chem.* 263: 12759–12764, 1988.
- [67608] 5938. Hillarp, A.; Dahlback, B.: Cloning of cDNA coding for the beta–chain of human complement component C4b–binding protein: sequence homology with the alpha chain. *Proc. Nat. Acad. Sci.* 87: 1183–1187, 1990.
- [67609] 5939. Hillarp, A.; Pardo–Manuel, F.; Ramos Ruiz, R.; Rodriguez de Cordoba, R.; Dahlback, B.: The human

C4b-binding protein beta-chain gene. J.Biol. Chem. 268: 15017–15023, 1993.

[67610] 5940.Pardo-Manuel, F.; Rey-Campos, J.; Hillarp, A.; Dahlback, B.; Rodriguez de Cordoba, S.: Human genes for the alpha and beta chains of complement C4b-binding protein are closely linked in a head-to-tail arrangement. Proc.Nat. Acad. Sci. 87: 4529–4532, 1990.

[67611] 5941.Rodriguez de Cordoba, S.; Perez-Blas, M.; Ramos-Ruiz, R.; Sanchez-Corral,P.; Pardo-Manuel de Villena, F.; Rey-Campos, J.: The gene coding for the beta-chain of C4b-binding protein (C4BPB) has become a pseudogene in the mouse. Genomics 21: 501–509, 1994.

[67612] 5942.Collaborative Study on the Genetics of Asthma: A genome-wide search for asthma susceptibility loci in ethnically diverse populations. Nature Genet. 15: 389–392, 1997.

[67613] 5943.Gavett, S. H.; O'Hearn, D. J.; Li, X.; Huang, S. K.; Finkelman, F. D.; Wills-Karp, M.: Interleukin 12 inhibits antigen-induced airway hyperresponsiveness, inflammation, and Th2 cytokine expression in mice. J. Exp. Med. 182: 1527–1536, 1995.

[67614] 5944.Karp, C. L.; Grupe, A.; Schadt, E.; Ewart, S. L.; Keane-Moore, M.; Cuomo, P. J.; Kohl, J.; Wahl, L.; Kuperman, D.;

Germer, S.; Aud,D.; Peltz, G.; Wills–Karp, M.: Identification of complement factor5 as a susceptibility locus for experimental allergic asthma. *NatureImmun.* 1: 221–226, 2000.

- [67615] 5945.Ober, C.; Cox, N. J.; Abney, M.; Di Rienzo, A.; Lander, E. S.;Changyaleket, B.; Gidley, H.; Kurtz, B.; Lee, J.; Nance, M.; Pettersson,A.; Prescott, J.; Richardson, A.; Schlenker, E.; Summerhill, E.; Willadsen,S.; Parry, R.; Collaborative Study on the Genetics of Asthma: Genome–wide search for asthma susceptibility loci in a founder population. *Hum.Molec. Genet.* 7: 1393–1398, 1998.
- [67616] 5946.Gennarelli, M.; Novelli, G.; Cobo, A.; Baiget, M.; Dal–lapiccola,B.: 3–Prime creatine kinase (M–type) polymorphisms linked to myotonicdystrophy in Italian and Spanish populations. *Hum. Genet.* 87: 654–656,1991.
- [67617] 5947.Nigro, J. M.; Schweinfest, C. W.; Rajkovic, A.; Pavlovic, J.; Jamal,S.; Dottin, R. P.; Hart, J. T.; Kamarck, M. E.; Rae, P. M. M.; Carty,M. D.; Martin–DeLeon, P.: cDNA cloning and mapping of the human creatinekinase M gene to 19q13. *Am. J. Hum. Genet.* 40: 115–125, 1987.
- [67618] 5948.Perryman, M. B.; Kerner, S. A.; Bohlmeier, T. J.; Roberts, R.:Isolation and sequence analysis of a full–length cDNA for human Mcreatine kinase. *Biochem. Biophys. Res. Commun.* 140: 981–989, 1986.

- [67619] 5949.Povey, S.; Inwood, M.; Tanyar, A.; Bobrow, M.: The expressionof the BB isozyme of creatine kinase. (Abstract) Cytogenet. CellGenet. 25: 198 only, 1979.
- [67620] 5950.Povey, S.; Inwood, M.; Tanyar, A.; Bobrow, M.: The expressionof creatine kinase isozymes in human cultured cells. Ann. Hum. Genet. 43:15–26, 1979.
- [67621] 5951.Roman, D.; Billadello, J.; Gordon, J.; Grace, A.; Sobel, B.; Strauss,A.: Complete nucleotide sequence of dog heart creatine kinase mRNA:conservation of amino acid se– quence within and among species. Proc.Nat. Acad. Sci. 82: 8394–8398, 1985.
- [67622] 5952.Rosenberg, U. B.; Kunz, G.; Frischauf, A.; Lehrach, H.; Mahr,R.; Eppenberger, H. M.; Perriard, J.–C.: Molecular cloning and expressionduring myogenesis of sequences coding for M–creatine kinase. Proc.Nat. Acad. Sci. 79: 6589–6592, 1982.
- [67623] 5953.Schweinfest, C. W.; Nigro, J. M.; Rajkovic, A.; Dottin, R. P.;Hart, J. M.; Karmack, M. E.; Rae, P. M. M.: Localization of the humancreatine kinase–M gene to chromosome 19. (Abstract) Cytogenet. CellGenet. 40: 740–741, 1985.
- [67624] 5954.Smeets, H.; Bachinski, L.; Coerwinkel, M.; Schepens, J.; Hoeijmakers,J.; van Duin, M.; Grzeschik, K.–H.; Weber, C. A.; de Jong, P.; Siciliano,M. J.; Wieringa, B.: A long–

range restriction map of the human chromosome19q13 region: close physical linkage between CKMM and the ERCC1 andERCC2 genes. Am. J. Hum. Genet. 46: 492–501, 1990.

[67625] 5955.Stallings, R. L.; Olson, E.; Strauss, A. W.; Thompson, L. H.;Bachinski, L. L.; Siciliano, M. J.: Human creatine kinase genes onchromosomes 15 and 19, and proximity of the gene for the muscle formto the genes for apolipoprotein C2 and excision repair. Am. J. Hum.Genet. 43: 144–151, 1988.

[67626] 5956.Steeghs, K.; Benders, A.; Oerlemans, F.; de Haan, A.; Heerschap,A.; Ruitenbeek, W.; Jost, C.; van Deursen, J.; Perryman, B.; Pette,D.; Bruckwilder, M.; Koudijs, J.; Jap, P.; Veerkamp, J.; Wieringa,B.: Altered Ca(2+) responses in muscles with combined mitochondrialand cytosolic creatine kinase deficiencies. Cell 89: 93–103, 1997.

[67627] 5957.Watts, D. C.: Creatine kinase (adenosine 5–prime–triphosphate–creatinephosphotransferase).In: Boyer, P. D.: The Enzymes. New York: AcademicPress (pub.) (3rd ed.) 8: 1973. Pp. 384–455.

[67628] 5958.Weil, D.; Van Cong, N.; Gross, M.–S.; Foubert, C.; Frezal, J.: Localisation du gene de la creatine kinase BB sur le chromosome14 par l'analyse des hybrides homme–



rongeur. Ann. Genet. 23: 150–154,1980.

[67629] 5959.Kishi, K.; Yasuda, T.; Awazu, S.; Mizuta, K.: Genetic polymorphism of human urine deoxyribonuclease I. Hum. Genet. 81: 295–297, 1989.

[67630] 5960.Kishi, K.; Yasuda, T.; Ikehara, Y.; Sawazaki, K.; Sato, W.; Iida, R.: Human serum deoxyribonuclease I (DNase I) polymorphism: pattern similarities among isozymes from serum, urine, kidney, liver, and pancreas. Am. J. Hum. Genet. 47: 121–126, 1990.

[67631] 5961.Napirei, M.; Karsunky, H.; Zevnik, B.; Stephan, H.; Mannherz, H.G.; Moroy, T.: Features of systemic lupus erythematosus in Dnase1-deficient mice. Nature Genet. 25: 177–181, 2000.

[67632] 5962.Yasuda, T.; Awazu, S.; Sato, W.; Iida, R.; Tanaka, Y.; Kishi, K.: Human genetically polymorphic deoxyribonuclease: purification, characterization, and multiplicity of urine deoxyribonuclease I. J. Biochem. 108:393–398, 1990.

[67633] 5963.Yasuda, T.; Kishi, K.; Yanagawa, Y.; Yoshida, A.: Structure of the human deoxyribonuclease I (DNase I) gene: identification of the nucleotide substitution that generates its classical genetic polymorphism. Ann. Hum. Genet. 59: 1–15, 1995.

[67634] 5964.Yasuda, T.; Nadano, D.; Iida, R.; Takeshita, H.; Lane,

S. A.; Callen, D. F.; Kishi, K.: Chromosomal assignment of the human deoxyribonuclease I gene, DNASE1 (DNL1), to band 16p13.3 using the polymerase chain reaction. *Cytogenet. Cell Genet.* 70: 221–223, 1995.

[67635] 5965. Yasutomo, K.; Horiuchi, T.; Kagami, S.; Tsukamoto, H.; Hashimura, C.; Urushihara, M.; Kuroda, Y.: Mutation of DNASE1 in people with systemic lupus erythematosus. *Nature Genet.* 28: 313–314, 2001.

[67636] 5966. Muller, U.; Steinberger, D.; Nemeth, A. H.: Clinical and molecular genetics of primary dystonias. *Neurogenetics* 1: 165–177, 1998.

[67637] 5967. Chambers, I.; Frampton, J.; Goldfarb, P.; Affara, N.; McBain, W.; Harrison, P. R.: The structure of the mouse glutathione peroxidase gene: the selenocysteine in the active site is encoded by the 'termination' codon, TGA. *EMBO J.* 5: 1221–1227, 1986.

[67638] 5968. Chu, F.-F.; Esworthy, R. S.; Doroshov, J. H.; Doan, K.; Liu, X.-F.: Expression of glutathione peroxidase in human liver in addition to kidney, heart, lung, and breast in humans and rodents. *Blood* 79:3233–3238, 1992.

[67639] 5969. Takahashi, K.; Akasaka, M.; Yamamoto, Y.; Kobayashi, C.; Mizoguchi, J.; Koyama, J.: Primary structure of human plasma glutathione peroxidase deduced from

cDNA sequences. J. Biochem. 108: 145–148, 1990.

[67640] 5970.Kelner, M. J.; Montoya, M. A.: Structural organization of the human selenium–dependent phospholipid hydroperoxide glutathione peroxidase gene (GPX4): chromosomal localization to 19p13.3. Biochem. Biophys. Res. Commun. 249: 53–55, 1998.

[67641] 5971.Roveri, A.; Casasco, A.; Maiorino, M.; Dalan, P.; Caligaro, A.; Ursini, F.: Phospholipid hydroperoxide glutathione peroxidase of rat testis. J. Biol. Chem. 267: 6142–6146, 1992.

[67642] 5972.Comstock, K. E.; Johnson, K. J.; Rifken, D.; Henner, W. D.: Isolation and analysis of the gene and cDNA for a human mu class glutathione S–transferase, GSTM4. J. Biol. Chem. 268: 16958–16965, 1993.

[67643] 5973.Ross, V. L.; Board, P. G.: Molecular cloning and heterologous expression of an alternatively spliced human mu class glutathione S–transferase transcript. Biochem. J. 294: 373–380, 1993.

[67644] 5974.Ross, V. L.; Board, P. G.; Webb, G. C.: Chromosomal mapping of the human mu class glutathione S–transferases to 1p13. Genomics 18:87–91, 1993.

[67645] 5975.Taylor, J. B.; Oliver, J.; Sherrington, R.; Pemble, S. E.: Structure of human glutathione S–transferase class mu

genes. *Biochem. J.* 274:587–593, 1991.

[67646] 5976.Board, P.; Coggan, M.; Johnston, P.; Ross, V.; Suzuki, T.; Webb,G.: Genetic heterogeneity of the human glutathione transferases:a complex of gene families. *Pharm. Therap.* 48: 357–369, 1990.

[67647] 5977.Chen, C.–L.; Liu, Q.; Relling, M. V.: Simultaneous characterizationof glutathione S–transferase M1 and T1 polymorphisms by polymerasechain reaction in American whites and blacks. *Pharmacogenetics* 6:187–191, 1996.

[67648] 5978.DeJong, J. L.; Chang, C.–M.; Whang–Peng, J.; Knut–sen, T.; Tu, C.–P.D.: The human liver glutathione S–transferase gene superfamily: expressionand chromosome mapping of an H(b) subunit cDNA. *Nucleic Acids Res.* 16:8541–8554, 1988.

[67649] 5979.DeJong, J. L.; Mohandas, T.; Tu, C.–P. D.: The human H(b) (mu)class glutathione S–transferases are encoded by a dispersed gene family. *Biochem.Biophys. Res. Commun.* 180: 15–22, 1991.

[67650] 5980.Godschalk, R. W. L.; Dallinga, J. W.; Wikman, H.; Risch, A.; Kleinjans,J. C. S.; Bartsch, H.; Van Schooten, F.–J.: Modulation of DNA andprotein adducts in smokers by genetic polymorphisms in GSTM1, GSTT1,NAT1 and NAT2. *Pharmacogenetics* 11: 389–398, 2001.

- [67651] 5981.Harada, S.; Abei, M.; Tanaka, N.; Agarwal, D. P.; Goedde, H. W.: Liver glutathione S-transferase polymorphism in Japanese and its pharmacogenetic importance. Hum. Genet. 75: 322–325, 1987.
- [67652] 5982.Lee, K. A.; Kim, S. H.; Woo, H. Y.; Hong, Y. J.; Cho, H. C.: Increased frequencies of glutathione S-transferase (GSTM1 and GSTT1) gene deletions in Korean patients with acquired aplastic anemia. Blood 98: 3483–3485, 2001.
- [67653] 5983.Mannervik, B.: The isozymes of glutathione transferase. Adv.Enzym. Relat. Areas Molec. Biol. 57: 357–417, 1985.
- [67654] 5984.Mannervik, B.; Awasthi, Y. C.; Board, P. G.; Hayes, J. D.; Dillio, C.; Ketterer, B.; Listowsky, I.; Morgenstern, R.; Muramatsu, M.; Pearson, W. R.; Pickett, C. B.; Sato, K.; Widersten, M.; Wolf, R. C.: Nomenclature for human glutathione transferases. (Letter) Biochem.J. 282: 305–306, 1992.
- [67655] 5985.Arce, M. A.; Thompson, E. S.; Wagner, S.; Coyne, K. E.; Ferdman, B. A.; Lublin, D. M.: Molecular cloning of RhD cDNA derived from a gene present in RhD-positive, but not RhD-negative individuals. Blood 82:651–655, 1993.
- [67656] 5986.Bennett, P. R.; Le Van Kim, C.; Colin, Y.; Warwick, R. M.; Cherif-Zahar, B.; Fisk, N. M.; Cartron, J.-P.: Prenatal

determination of fetalRhD type by DNA amplification. New Eng. J. Med. 329: 607–610, 1993.

- [67657] 5987.Bowman, J. M.: RhD hemolytic disease of the new-born. (Editorial) NewEng. J. Med. 339: 1775–1777, 1998.
- [67658] 5988.Cartron, J.–P.: Defining the Rh blood group antigens: biochemistryand molecular genetics. Blood Rev. 8: 199–212, 1994.
- [67659] 5989.Colin, Y.; Cherif–Zahar, B.; Le Van Kim, C.; Raynal, V.; Van Huffel,V.; Cartron, J.–P.: Genetic basis of the RhD–positive and RhD–negativeblood group polymorphism as determined by Southern analysis. Blood 78:2747–2752, 1991.
- [67660] 5990.Diamond, L. K.; Blackfan, K. D.; Baty, J. M.: Erythrob–lastosisfetalis and its association with universal edema of the fetus, icterusgravis neonatorum and anemia of the newborn. J. Pediat. 1: 269–309,1932.
- [67661] 5991.du Toit, E. D.; Martell, R. W.; Botha, I.; Kriel, C. J.: Anti–Dantibodies in the Rh–positive mothers. S. Afr. Med. J. 75: 452,1989.
- [67662] 5992.Garratty, G.: Severe reactions associated with trans–fusion ofpatients with sickle cell disease. Transfusion 37: 357–361, 1997.
- [67663] 5993.Huang, C.–H.: Personal Communication. New York

City, N. Y. 10/11/1996.

- [67664] 5994.Huang, C.-H.; Chen, Y.; Reid, M.; Ghosh, S.: Genetic recombinationat the human RH locus: a family study of the red-cell Evans phenotypereveals a transfer of exons 2-6 from the RHD to the RHCE gene. Am.J. Hum. Genet. 59: 825-833, 1996.
- [67665] 5995.Huang, C.-H.; Reid, M. E.; Chen, Y.; Coghlan, G.; Okubo, Y.:Molecular definition of red cell Rh haplotypes by tightly linked SphIRFLPs. Am. J. Hum. Genet. 58: 133-142, 1996.
- [67666] 5996.Hyland, C. A.; Wolter, L. C.; Liew, Y. W.; Saul, A.: A Southernanalysis of Rh blood group genes: association between restrictionfragment length polymorphism patterns and Rh serotypes. Blood 83:566-572, 1994.
- [67667] 5997.Issitt, P. D.: The Rh blood group system, 1988: eight new antigensin nine years and some observations on the biochemistry and genetics of the system. Transfusion Med. Rev. 3: 1-12, 1989.
- [67668] 5998.Kemp, T. J.; Poulter, M.; Carritt, B.: A recombination hot spotin the Rh genes revealed by analysis of unrelated donors with therare D-- phenotype. Am. J. Hum. Genet. 59: 1066-1073, 1996.
- [67669] 5999.Legler, T. J.; Eber, S. W.; Lakomek, M.; Lynen, R.;

Maas, J. H.;Pekrun, A.; Repas–Humpe, M.; Schroter, W.; Kohler, M.: Applicationof RHD and RHCE genotyping for correct blood group determination inchronically trans–fused patients. Transfusion 39: 852–855, 1999.

[67670] 6000.Le Van Kim, C.; Cherif–Zahar, B.; Raynal, V.; Mouro, I.; Lopez,M.; Cartron, J. P.; Colin, Y.: Multiple Rh messen–ger RNA isoformsare produced by alternative splicing. Blood 80: 1074–1078, 1992.

[67671] 6001.Le Van Kim, C.; Mouro, I.; Cherif–Zahar, B.; Raynal, V.; Cherrier,C.; Cartron, J.–P.; Colin, Y.: Molecular cloning and primary structureof the human blood group RhD polypeptide. Proc. Nat. Acad. Sci. 89:10925–10929, 1992.

[67672] 6002.Levine, P.; Katzin, E. M.; Burnham, L.: Isoimmuniza–tion in pregnancy:its possible bearing on the etiology of erythroblastosis foetalis. J.A.M.A. 116:825–827, 1941.

[67673] 6003.Lo, Y. M. D.; Hjelm, N. M.; Fidler, C.; Sargent, I. L.; Murphy,M. F.; Chamberlain, P. F.; Poon, P. M. K.; Redman, C. W. G.; Wainscoat,J. S.: Prenatal diagnosis of fetal RhD status by molecular analysisof maternal plasma. New Eng. J. Med. 339: 1734–1738, 1998.

[67674] 6004.Miyoshi, O.; Yabe, R.; Wakui, K.; Fukushima, Y.; Koizumi, S.;Uchikawa, M.; Kajii, T.; Numakura, C.; Taka–hashi, S.; Hayasaka, K.;Niikawa, N.: Two cases of mosaic



RhD blood-group phenotypes and paternalisodisomy for chromosome 1. *Am. J. Med. Genet.* 104: 250–256, 2001.

- [67675] 6005.Okuda, H.; Kawano, M.; Iwamoto, S.; Tanaka, M.; Seno, T.; Okubo,Y.; Kajii, E.: The RHD gene is highly detectable in RhD-negativeJapanese donors. *J. Clin. Invest.* 100: 373–379, 1997.
- [67676] 6006.Race, R. R.: An 'incomplete' antibody in human serum. (Letter) *Nature* 153:771–772, 1944.
- [67677] 6007.Race, R. R.; Sanger, R.: *Blood Groups in Man*. Oxford: Blackwell(pub.) (6th ed.): 1975.
- [67678] 6008.Rouillac, C.; Le Van Kim, C.; Beolet, M.; Cartron, J.-P.; Colin,Y.: Leu110-to-pro substitution in the RhD polypeptide is responsiblefor the D(VII) category blood group phenotype. *Am. J. Hemat.* 49:87–88, 1995.
- [67679] 6009.Smythe, J. S.; Avent, N. D.; Judson, P. A.; Parsons, S. F.; Martin,P. G.; Anstee, D. J.: Expression of RHD and RHCE gene products usingretroviral transduction of K562 cells establishes the molecular basisof Rh blood group antigens. *Blood* 87: 2968–2973, 1996.
- [67680] 6010.Spanos, T.; Karageorga, M.; Ladis, V.; Peristeri, J.; Hatziliami,A.; Kattamis, C.: Red cell alloantibodies in patients with thalassemia. *VoxSang.* 58: 50–55, 1990.
- [67681] 6011.Wagner, F. F.; Flegel, W. A.: RHD gene deletion oc-

curred in the Rhesus box. Blood 95: 3662–3668, 2000.

[67682] 6012. Wagner, F. F.; Gassner, C.; Muller, T. H.; Schonitzer, D.; Schunter, F.; Flegel, W. A.: Molecular basis of weak D phenotypes. Blood 93:385–393, 1999.

[67683] 6013. Wagner, F. F.; Ladewig, B.; Angert, K. S.; Heymann, G. A.; Eicher, N. I.; Flegel, W. A.: The DAU allele cluster of the RHD gene. Blood 100:306–311, 2002.

[67684] 6014. Wiener, A. S.: The Rh series of allelic genes. Science 100:595–597, 1944.

[67685] 6015. Wang, X.; Chan, S. J.; Eddy, R. L.; Byers, M. G.; Fukushima, Y.; Henry, W. M.; Haley, L. L.; Steiner, D. F.; Shows, T. B.: Chromosome assignment of cathepsin B (CTSB) to 8p22 and cathepsin H (CTSH) to 15q24–q25. (Abstract) Cytogenet. Cell Genet. 46: 710–711, 1987.

[67686] 6016. Tynnela, J.; Sohar, I.; Sleat, D. E.; Gin, R. M.; Donnelly, R. J.; Baumann, M.; Haltia, M.; Lobel, P.: A mutation in the ovine cathepsin D gene causes a congenital lysosomal storage disease with profound neurodegeneration. EMBO J. 19: 2786–2792, 2000.

[67687] 6017. Gelb, B. D.; Shi, G.-P.; Heller, M.; Weremowicz, S.; Morton, C.; Desnick, R. J.; Chapman, H. A.: Structure and chromosomal assignment of the human cathepsin K gene. Genomics 41: 258–262, 1997.

- [67688] 6018. Shiang, R.; Lidral, A. C.; Ardinger, H. H.; Buetow, K. H.; Romitti, P. A.; Munger, R. G.; Murray, J. C.: Association of transforming growth-factor alpha gene polymorphisms with nonsyndromic cleft palate only (CPO). *Am. J. Hum. Genet.* 53: 836–843, 1993.
- [67689] 6019. Shields, E. D.; Bixler, D.; Fogh-Andersen, P.: Facial clefts in Danish twins. *Cleft Palate J.* 16: 1–6, 1979.
- [67690] 6020. Pollack, M. S.; Ochs, H. D.; Dupont, B.: HLA typing of cultured amniotic cells for the prenatal diagnosis of complement C4 deficiency. *Clin. Genet.* 18: 197–200, 1980.
- [67691] 6021. Porter, R. R.; Reid, K. B. M.: The biochemistry of complement. *Nature* 275: 699–704, 1978.
- [67692] 6022. Rittner, C.; Bertrams, J.: On the significance of C2, C4, and factor B polymorphisms in disease. *Hum. Genet.* 56: 235–247, 1981.
- [67693] 6023. Rittner, C.; Hauptmann, G.; Grosse-Wilde, H.; Grosshans, E.; Tongio, M. M.; Mayer, S.: Linkage between HL-A (major histocompatibility complex) and genes controlling the fourth component of complement. In: *Histocompatibility Testing 1975*. Copenhagen: Munksgaard (pub.) 1976. Pp. 945–953.
- [67694] 6024. Roos, M. H.; Mollenhauer, E.; Demant, P.; Rittner, C.: A molecular basis for the two locus model of human com-

plement component C4. *Nature* 298:854–856, 1982.

[67695] 6025. Rosenfeld, S. I.; Ruddy, S.; Austen, K. F.: Structural polymorphism of the fourth component of human complement. *J. Clin. Invest.* 48:2283–2292, 1969.

[67696] 6026. Roychoudhury, A. K.; Nei, M.: *Human Polymorphic Genes: World Distribution*. New York: Oxford Univ. Press (pub.) 1988.

[67697] 6027. Schaller, J. G.; Gilliland, B. G.; Ochs, H. D.; Leddy, J. P.; Agodoa, L. C. Y.; Rosenfeld, S. I.: Severe systemic lupus erythematosus with nephritis in a boy with deficiency of the fourth component of complement. *Arthritis Rheum.* 20: 1519–1525, 1977.

[67698] 6028. Shevach, E. M.; Frank, M. M.; Green, I.: Linkage of gene controlling the synthesis of the fourth component of complement to the major histocompatibility complex of the guinea pig. *Immunogenetics* 3: 595–602, 1976.

[67699] 6029. Shreffler, D. C.: The S region of the mouse major histocompatibility complex (H-2): genetic variation and functional role in complement system. *Transplant. Rev.* 32: 140–167, 1976.

[67700] 6030. Torisu, M.; Sonozaki, H.; Inai, S.; Arata, M.: Deficiency of the fourth component of complement in man. *J. Immunogenet.* 104:728–737, 1970.

- [67701] 6031.Wank, R.; Schendel, D. J.; O'Neill, G. J.; Riethmuller, G.; Held,E.; Feucht, H. E.: Rare variant of complement C4 is seen in highfrequency in patients with primary glomerulonephritis. *Lancet* I:872–874, 1984.
- [67702] 6032.Welch, T. R.; Beischel, L.: C4 uremic variant: an acquired C4allotype. *Immunogenetics* 22: 553–562, 1985.
- [67703] 6033.Welch, T. R.; Beischel, L. S.; Choi, E.; Balakrishnan, K.; Bishof,N. A.: Uniparental isodisomy 6 associated with deficiency of thefourth component of complement. *J. Clin. Invest.* 86: 675–678, 1990.
- [67704] 6034.Whitehead, A. S.; Colten, H. R.; Chang, C. C.; Demars, R.: Localizationof the human MHC–linked complement genes between HLA–B and HLA–DRby using HLA mutant cell lines. *J. Immun.* 134: 641–643, 1985.
- [67705] 6035.Wilson, W. A.; Perez, M. C.: Complete C4B deficiency in blackAmericans with systemic lupus erythematosus. *J. Rheum.* 15: 1855–1858,1988.
- [67706] 6036.Andersson, A.; Dahlback, B.; Hanson, C.; Hillarp, A.; Levan, G.;Szpirer, J.; Szpirer, C.: Genes for C4b–binding protein alpha– andbeta–chains (C4BPA and C4BPB) are located on chromosome 1, band 1q32,in humans and on chromosome 13 in rats. *Somat. Cell Molec. Genet.* 16:493–500, 1990.

- [67707] 6037.Aso, T.; Okamura, S.; Matsuguchi, T.; Sakamoto, N.; Sata, T.; Niho,Y.: Genomic organization of the alpha chain of the human C4b-bindingprotein gene. *Biochem. Biophys. Res. Commun.* 174: 222–227, 1991.
- [67708] 6038.Barnum, S. R.; Kristensen, T.; Chaplin, D. D.; Seldin, M. F.; Tack,B. F.: Molecular analysis of the murine C4b-binding protein gene:chromosome assignment and partial gene organization. *Biochemistry* 28:8312–8317, 1989.
- [67709] 6039.Gigli, I.; Fujita, T.; Nussenzweig, V.: Modulation of the classicalpathway C3 convertase by plasma proteins C4 binding protein and C3binactivator. *Proc. Nat. Acad. Sci.* 76: 6596–6600, 1979.
- [67710] 6040.Hatch, J. A.; Atkinson, J. P.; Suarez, B. K.; Dykman, T. R.: Evaluationof linkage of the human C3b/C4b receptor to HLA. *J. Immun.* 132:2168–2169, 1984.
- [67711] 6041.Hing, S.; Day, A. J.; Linton, S. J.; Ripoche, J.; Sim, R. B.; Reid,K. B. M.; Solomon, E.: Assignment of complement components C4 bindingprotein (C4BP) and factor H (FH) to human chromosome 1q, using cDNAprobes. *Ann. Hum. Genet.* 52: 117–122, 1988.
- [67712] 6042.Kaidoh, T.; Natsuume–Sakai, S.; Takahashi, M.: Murine bindingprotein of the fourth component of com–

plement: structural polymorphism and its linkage to the major histocompatibility complex. *Proc. Nat. Acad. Sci.* 78: 3794–3798, 1981.

[67713] 6043. Matsuguchi, T.; Okamura, S.; Aso, T.; Sata, T.; Niho, Y.: Molecular cloning of the cDNA coding for proline-rich protein (PRP): identity of PRP as C4b-binding protein.

*Biochem. Biophys. Res. Commun.* 165:138–144, 1989.

[67714] 6044. Nagasawa, S.; Stroud, R. M.: Purification and characterization of a macromolecular weight cofactor for C3b-inactivator, C4bC3bINA-cofactor, of human plasma.

*Molec. Immun.* 17: 1365–1372, 1980.

[67715] 6045. Rey-Campos, J.; Rubinstein, P.; Rodriguez de Cordoba, S.: A physical map of the human regulator of complement activation gene cluster linking the complement

genes CR1, CR2, DAF, and C4BP. *J. Exp. Med.*

167:664–669, 1988.

[67716] 6046. Rodriguez de Cordoba, S.; Dykman, T. R.; Ginsberg-Fellner, F.; Ercilla, G.; Aqua, M.; Atkinson, J. P.; Rubinstein, P.: Evidence for linkage between the loci coding for the

binding protein for the fourth component of human complement (C4BP) and for the C3b/C4b receptor. *Proc. Nat.*

*Acad. Sci.* 81: 7890–7892, 1984.

[67717] 6047. Rodriguez de Cordoba, S.; Ferreira, A.; Nussen-

zweig, V.; Rubinstein, P.: Genetic polymorphism of human C4-binding protein. *J. Immun.* 131:1565–1569, 1983.

[67718] 6048. Rodriguez de Cordoba, S.; Lublin, D. M.; Rubinstein, P.; Atkinson, J. P.: Human genes for three complement components that regulate the activation of C3 are tightly linked. *J. Exp. Med.* 161: 1189–1195, 1985.

[67719] 6049. Sullivan, F. X.; Kumar, R.; Kriz, R.; Stahl, M.; Xu, G.-Y.; Rouse, J.; Chang, X.; Boodhoo, A.; Potvin, B.; Cumming, D. A.: Molecular cloning of human GDP-mannose 4,6-dehydratase and reconstitution of GDP-fucose biosynthesis in vitro. *J. Biol. Chem.* 273: 8193–8202, 1998.

[67720] 6050. Justice, M. J.; Siracusa, L. D.; Gilbert, D. J.; Heisterkamp, N.; Groffen, J.; Chada, K.; Silan, C. M.; Copeland, N. G.; Jenkins, N. A.: A genetic linkage map of mouse chromosome 10: localization of eighteen molecular markers using a single interspecific backcross. *Genetics* 125:855–866, 1990.

[67721] 6051. Mouillet-Richard, S.; Ermonval, M.; Chebassier, C.; Laplanche, J. L.; Lehmann, S.; Launay, J. M.; Kellermann, O.: Signal transduction through prion protein. *Science* 289: 1925–1928, 2000.

[67722] 6052. Parravicini, V.; Gadina, M.; Kovarova, M.; Odom, S.; Gonzalez-Espinosa, C.; Furumoto, Y.; Saitoh, S.; Samelson,



L. E.; O'Shea, J. J.; Rivera, J.: Fyn kinase initiates complementary signals required for IgE-dependent mast cell degranulation. *Nature Immun.* 3: 741–748, 2002.

[67723] 6053. Casari, G.; Barlassina, C.; Cusi, D.; Zagato, L.; Muirhead, R.; Righetti, M.; Nembri, P.; Amar, K.; Gatti, M.; Macciardi, F.; Binelli, G.; Bianchi, G.: Association of the alpha-adducin locus with essential hypertension. *Hypertension* 25: 320–326, 1995.

[67724] 6054. Crosby, A. H.; Edwards, S. J.; Murray, J. C.; Dixon, M. J.: Genomic organization of the human osteopontin gene: exclusion of the locus from a causative role in the pathogenesis of dentinogenesis imperfecta type II. *Genomics* 27: 155–160, 1995.

[67725] 6055. Abernethy, T. J.; Avery, O. T.: The occurrence during acute infection of a protein not normally present in the blood. I. Distribution of the reactive protein in patients' sera and the effect of calcium on the flocculation reaction with C polysaccharide of pneumococcus. *J. Exp. Med.* 73: 173–182, 1941.

[67726] 6056. Floyd-Smith, G. A.; Whitehead, A. S.; Colten, H. R.; Francke, U.: The human C-reactive protein gene (CRP) and serum amyloid P component gene (APCS) are located on the proximal long arm of chromosome 1. *Immunogenetics*

24:171–176, 1986.

- [67727] 6057. Tillett, W. S.; Francis, T., Jr.: Serological reactions in pneumoniawith a nonprotein somatic fraction of pneumococcus. *J. Exp. Med.* 52:561–585, 1930.
- [67728] 6058. Benger, J. C.; Teshima, I.; Walter, M. A.; Brubacher, M. G.; Daouk, G. H.; Cox, D. W.: Localization and genetic linkage of the human immunoglobulin heavy chain genes and the creatine kinase brain (CKB) gene: identification of a hot spot for recombination. *Genomics* 9:614–622, 1991.
- [67729] 6059. Brubacher, M. G.; Benger, J. C.; Billingsley, G. D.; Hofker, M. H.; Nakamura, Y.; White, R.; Cox, D. W.: A genetic linkage map of the distal region of human chromosome 14. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A133 only, 1989.
- [67730] 6060. Hoo, J. J.; Goedde, H. W.: Determination of brain type creatine kinase for diagnosis of perinatal asphyxia—choice of method. (Letter) *Pediat. Res.* 16: 806 only, 1982.
- [67731] 6061. Kaye, F. J.; McBride, O. W.; Battey, J. F.; Gazdar, A. F.; Sausville, E. A.: Human creatine kinase–B complementary DNA: nucleotide sequence, gene expression in lung cancer, and chromosomal assignment to two distinct loci. *J. Clin. Invest.* 79: 1412–1420, 1987.

- [67732] 6062.Ma, T. S.; Ifegwu, J.; Watts, L.; Siciliano, M. J.; Roberts, R.;Perryman, M. B.: Serial Alu sequence transposition interrupting a human B creatine kinase pseudogene. *Genomics* 10: 390–399, 1991.
- [67733] 6063.Mariman, E. C. M.; Schepens, J. T. G.; Wieringa, B.: Complete nucleotide sequence of the human creatine kinase B gene. *Nucleic Acids Res.* 17: 6385 only, 1989.
- [67734] 6064.Pfeiffer, F. E.; Homburger, H. A.; Yanagihara, T.: Creatine kinase BB isoenzyme in CSF in neurologic diseases: measurement by radioimmunoassay. *Arch. Neurol.* 40: 169–172, 1983.
- [67735] 6065.Stallings, R. L.; Olson, E.; Strauss, A. W.; Thompson, L. H.; Bachinski, L. L.; Siciliano, M. J.: Human creatine kinase genes on chromosomes 15 and 19, and proximity of the gene for the muscle form to the genes for apolipoprotein C2 and excision repair. *Am. J. Hum. Genet.* 43:144–151, 1988.
- [67736] 6066.Villarreal-Levy, G.; Ma, T. S.; Kerner, S. A.; Roberts, R.; Perryman, M. B.: Human creatine kinase: isolation and sequence analysis of cDNA clones for the B subunit, development of subunit specific probes, and determination of gene copy number. *Biochem. Biophys. Res. Commun.* 144:1116–1127, 1987.

- [67737] 6067.Bark, C. J.: Mitochondrial creatine kinase: a poor prognostic sign. *J.A.M.A.* 243: 2058–2060, 1980.
- [67738] 6068.Steeghs, K.; Merckx, G.; Wieringa, B.: The ubiquitous mitochondrial creatine kinase gene maps to a conserved region on human chromosome 15q15 and mouse chromosome 2 bands F1–F3. *Genomics* 24: 193–195, 1994.
- [67739] 6069.Klein, S. C.; Haas, R. C.; Perryman, M. B.; Billadello, J. J.; Strauss, A. W.: Regulatory element analysis and structural characterization of the human sarcomeric mitochondrial creatine kinase gene. *J. Biol.Chem.* 266: 18058–18065, 1991.
- [67740] 6070.Richard, I.; Devaud, C.; Cherif, D.; Cohen, D.; Beckmann, J. S.: The gene for creatine kinase, mitochondrial 2 (sarcomeric; CKMT2), maps to chromosome 5q13.3. *Genomics* 18: 134–136, 1993.
- [67741] 6071.Bailly, J.; MacKenzie, A. E.; Leblond, S.; Korneluk, R. G.: Assessment of a creatine kinase isoform M defect as a cause of myotonic dystrophy and the characterization of two novel CKMM polymorphisms. *Hum. Genet.* 86:457–462, 1991.
- [67742] 6072.Bohner, J.; Stein, W.; Kuhlmann, E.; Eggstein, M.: Serum creatine kinase BB linked to immunoglobulin G. *Clin. Chim. Acta* 97: 83–88, 1979.

- [67743] 6073.Dawson, D. M.; Eppenberger, H. M.; Eppenberger, M. E.: Multiplemolecular forms of creatine kinases. Ann. N.Y. Acad. Sci. 151: 616–626,1968.
- [67744] 6074.Schonk, D.; van Dijk, P.; Riegmann, P.; Trapman, J.; Holm, C.;Willcocks, T. C.; Sillekens, P.; van Venrooij, W.; Wimmer, E.; Geurtsvan Kessel, A.; Ropers, H.–H.; Wieringa, B.: Assignment of sevengegenes to distinct intervals on the midportion of human chromosome19q surrounding the myotonic dystrophy gene region. Cytogenet. CellGenet. 54: 15–19, 1990.
- [67745] 6075.Coles, L. S.; Diamond, P.; Occhiodoro, F.; Vadas, M. A.; Shannon,M. F.: Cold shock domain proteins repress transcription from theGM–CSF promoter. Nucleic Acids Res. 24: 2311–2317, 1996.
- [67746] 6076.Didier, D. K.; Schiffenbauer, J.; Woulfe, S. L.; Zacheis, M.; Schwartz,B. D.: Characterization of the cDNA encoding a protein binding tothe major histocompatibility complex class II Y box. Proc. Nat. Acad.Sci. 85: 7322–7326, 1988.
- [67747] 6077.Kudo, S.; Mattei, M.–G.; Fukuda, M.: Characterization of the gene for dbpA, a family member of the nucleic–acid–binding proteins containinga cold–shock domain. Europ. J. Biochem. 231: 72–82, 1995.
- [67748] 6078.Makino, Y.; Ohga, T.; Toh, S.; Koike, K.; Okumura,

K.; Wada, M.; Kuwano, M.; Kohno, K.: Structural and functional analysis of the human Y-box binding protein (YB-1) gene promoter. *Nucleic Acid Res.* 24:1873–1878, 1996.

[67749] 6079. Sakura, H.; Maekawa, T.; Imamoto, F.; Yasuda, K.; Ishii, S.: Two human genes isolated by a novel method encode DNA-binding proteins containing a common region of homology. *Gene* 73: 499–507, 1988.

[67750] 6080. Spitkovsky, D. D.; Royer-Pokora, B.; Delius, H.; Kisseljov, F.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Royer, H.-D.: Tissue restricted expression and chromosomal localization of the YB-1 gene encoding a 42kD nuclear CCAAT binding protein. *Nucleic Acids Res.* 20:797–803, 1992.

[67751] 6081. Levi-Strauss, M.; Carroll, M. C.; Steinmetz, M.; Meo, T.: A previously undetected MHC gene with an unusual periodic structure. *Science* 240:201–204, 1988.

[67752] 6082. Speiser, P. W.; White, P. C.: Structure of the human RD gene: a highly conserved gene in the class III region of the major histocompatibility complex. *DNA* 8: 745–751, 1989.

[67753] 6083. Surowy, C. S.; Gosink, J. J.; Spritz, R. A.: cDNA cloning of the human 'RD' protein, a possible novel RNA-binding protein. (Abstract) *Am. J. Hum. Genet.* 43: A203

only, 1988.

- [67754] 6084.Yamaguchi, Y.; Takagi, T.; Wada, T.; Yano, K.; Furuya, A.; Sugimoto,S.; Hasegawa, J.; Handa, H.: NELF, a multisubunit complex containingRD, cooperates with DSIF to repress RNA polymerase II elongation. *Cell* 97:41–51, 1999.
- [67755] 6085.Gross, M.–S.; Guyonnet–Duperat, V.; Porchet, N.; Bernheim, A.;Aubert, J. P.; Nguyen, V. C.: Mucin 4 (MUC4) gene: regional assignment(3q29) and RFLP analysis. *Ann. Genet.* 35: 21–26, 1992.
- [67756] 6086.Porchet, N.; Van Cong, N.; Dufosse, J.; Audie, J. P.; Guyonnet–Duperat,V.; Gross, M. S.; Denis, C.; Degand, P.; Bernheim, A.; Aubert, J.P.: Molecular cloning and chromosomal localization of a novel humantracheo–bronchial mucin cDNA containing tandemly repeated sequencesof 48 base pairs. *Biochem. Biophys. Res. Commun.* 175: 414–422,1991.
- [67757] 6087.Van Cong, N.; Aubert, J.–P.; Gross, M.–S.; Porchet, N.; Degand,P.; Bernheim, A.: Tracheobronchial mucin 4 (MUC4) gene: assignmentto 3q29 and polymorphism of VNTR type. (Abstract) *Cytogenet. CellGenet.* 58: 1879–1880, 1991.
- [67758] 6088.Pigny, P.; Pratt, W. S.; Laine, A.; Leclercq, A.; Swal–

low, D. M.; Nguyen, V. C.; Aubert, J. P.; Porchet, N.: The MUC5AC gene: RFLP analysis with the Jer58 probe. Hum. Genet. 96: 367–368, 1995.

[67759] 6089. Garcia, J. V.; Jones, C.; Miller, A. D.: Localization of the amphotropic murine leukemia virus receptor gene to the pericentromeric region of human chromosome 8. J. Virol. 65: 6316–6319, 1991.

[67760] 6090. van Zeijl, M.; Johann, S. V.; Closs, E.; Cunningham, J.; Eddy, R.; Shows, T. B.; O'Hara, B.: A human amphotropic retrovirus receptor is a second member of the gibbon ape leukemia virus receptor family. Proc. Nat. Acad. Sci. 91: 1168–1172, 1994.

[67761] 6091. van Zeijl, M.; Johann, S. V.; Eddy, R. L.; Shows, T. B.; O'Hara, B.: Assignment of GLVR2, a receptor for murine amphotropic virus to human chromosome 8. (Abstract) Human Genome Mapping Workshop 93 18 only, 1993.

[67762] 6092. Cawthon, R. M.; O'Connell, P.; Buchberg, A. M.; Viskochil, D.; Weiss, R. B.; Culver, M.; Stevens, J.; Jenkins, N. A.; Copeland, N. G.; White, R.: Identification and characterization of transcripts from the neurofibromatosis 1 region: the sequence and genomic structure of EVI2 and mapping of other transcripts. Genomics 7: 555–565, 1990.



[67763] 6093.Collins, F. S.: Personal Communication. Bethesda, Md. 11/17/1993.

[67764] 6094.O'Connell, P.; Buchberg, A.; Cawthon, R. M.; Culver, M.; Stevens,J.; Viskochil, D.; Carey, J. C.; Fountain, J.; Wallace, M.; Jenkins,N.; Copeland, N.; Collins, F. S.; White, R.: The human homologueof the murine Evi-2 oncogene lies between von Recklinghausen NF translocationbreakpoints. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A210 only,1989.

[67765] 6095.O'Connell, P.; Viskochil, D.; Buchberg, A. M.; Fountain, J.; Cawthon,R. M.; Culver, M.; Stevens, J.; Rich, D. C.; Ledbetter, D. H.; Wallace,M.; Carey, J. C.; Jenkins, N. A.; Copeland, N. G.; Collins, F. S.;White, R.: The human homolog of murine Evi-2 lies between two vonRecklinghausen neurofibromatosis translocations. Genomics 7: 547-554,1990.

[67766] 6096.Blanck, O.; Perrin, C.; Mziaut, H.; Darbon, H.; Mattei, M. G.;Miquelis, R.: Molecular cloning, cDNA analysis, and localizationof a monomer of the N-acetylglucosamine-specific receptor of the thyroid,NAGR1, to chromosome 19p13.3-13.2. Genomics 21: 18-26, 1994. Note:Erratum: Genomics 27: 561 only, 1995.

- [67767] 6097.Datar, K. V.; Dreyfuss, G.; Swanson, M. S.: The human hnRNP M proteins: identification of a methionine/arginine-rich repeat motif in ribonucleoproteins. *Nucleic Acids Res.* 21: 439–446, 1993.
- [67768] 6098.Hull, E.; Sarkar, M.; Spruijt, M. P. N.; Hoppener, J. W. M.; Dunn, R.; Schachter, H.: Organization and localization to chromosome 5 of the human UDP-N-acetylglucosamine:alpha-3-D-mannoside beta-1,2-N-acetylglucosaminyltransferase 1 gene. *Biochem. Biophys. Res. Commun.* 176: 608–615, 1991.
- [67769] 6099.Ioffe, E.; Stanley, P.: Mice lacking N-acetylglucosaminyltransferase I activity die at mid-gestation, revealing an essential role for complex or hybrid N-linked carbohydrates. *Proc. Nat. Acad. Sci.* 91: 728–732, 1994.
- [67770] 6100.Kumar, R.; Stanley, P.: Transfection of a human gene that corrects the Lec1 glycosylation defect: evidence for transfer of the structural gene for N-acetylglucosaminyltransferase I. *Molec. Cell. Biol.* 9:5713–5717, 1989. Note: Erratum: *Molec. Cell. Biol.* 10: 3857 only, 1990.
- [67771] 6101.Kumar, R.; Yang, J.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Stanley, P.: Cloning and expression of the murine

gene and chromosomal location of the human gene encoding N-acetylglucosaminyltransferase I. *Glycobiology* 2:383–393, 1992.

- [67772] 6102. Kumar, R.; Yang, J.; Larsen, R. D.; Stanley, P.: Cloning and expression of N-acetylglucosaminyltransferase I, the medial Golgi transferase that initiates complex N-linked carbohydrate formation. *Proc. Nat. Acad. Sci.* 87: 9948–9952, 1990.
- [67773] 6103. Metzler, M.; Gertz, A.; Sarkar, M.; Schachter, H.; Schrader, J.W.; Marth, J. D.: Complex asparagine-linked oligosaccharides are required for morphogenic events during post-implantation development. *EMBO J.* 13: 2056–2065, 1994.
- [67774] 6104. Pownall, S.; Kozak, C. A.; Schappert, K.; Sarkar, M.; Hull, E.; Schachter, H.; Marth, J. D.: Molecular cloning and characterization of the mouse UDP-N-acetylglucosamine:alpha-3-D-mannoside beta-1,2-N-acetylglucosaminyltransferase I gene. *Genomics* 12: 699–704, 1992.
- [67775] 6105. Puthalakath, H.; Burke, J.; Gleeson, P. A.: Glycosylation defect in lec1 Chinese hamster ovary mutant is due to a point mutation in N-acetylglucosaminyltransferase 1 gene. *J. Biol. Chem.* 271: 27818–27822, 1996.

- [67776] 6106.Shows, T. B.: Personal Communication. Buffalo, N.Y. 4/13/1999.
- [67777] 6107.Yip, B.; Chen, S.-H.; Mulder, H.; Hoppener, J. W. M.; Schachter,H.: Organization of the human beta-1,2-N-acetylglucosaminyltransferase gene (MGAT1), which controls complex and hybrid N-glycan synthesis. *Biochem.J.* 321: 465-474, 1997.
- [67778] 6108.Jaiswal, A. K.; Bell, D. W.; Radjendirane, V.; Testa, J. R.: Localization of human NQO1 gene to chromosome 16q22 and NQO2-6p25 and associated polymorphisms. *Pharmacogenetics* 9: 413-418, 1999.
- [67779] 6109.Jaiswal, A. K.; Burnett, P.; Adesnik, M.; McBride, O. W.: Nucleotide and deduced amino acid sequence of a human cDNA (NQO2) corresponding to a second member of the NAD(P)H:quinone oxidoreductase gene family: extensive polymorphism at the NQO2 gene locus on chromosome 6. (Abstract) *Biochemistry* 29:1899-1906, 1990.
- [67780] 6110.Tan, J.; D'Agostaro, G. A. F.; Bendiak, B.; Reck, F.; Sarkar,M.; Squire, J. A.; Leong, P.; Schachter, H.: The human UDP-N-acetylglucosamine:alpha-6-D-mannoside-beta-1,2-N-acetylgluco

saminyltransferase II gene (MGAT2): cloning of genomic DNA, localization to chromosome14q21, expression in insect cells and purification of the recombinantprotein. *Europ. J. Biochem.* 231: 317–328, 1995.

[67781] 6111.Kondo, M.; Scherer, D. C.; Miyamoto, T.; King, A. G.; Akashi,K.; Sugamura, K.; Weissman, I. L.: Cell–fate conversion of lymphoid–committedprogenitors by instructive actions of cytokines. *Nature* 407: 383–386,2000.

[67782] 6112.Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Larson, R. A.;Rowley, J. D.; Gasson, J. C.; Golde, D. W.; Sherr, C. J.: Evidencefor the involvement of GM–CSF and FMS in the deletion (5q) in myeloiddisorders. *Science* 231: 984–987, 1986.

[67783] 6113.Morgan, R.; Hecht, B. K.; Sandberg, A. A.; Hecht, F.; Smith, S.D.: Chromosome 5q35 breakpoint in malignant histiocytosis. (Letter) *NewEng. J. Med.* 314: 1322 only, 1986.

[67784] 6114.Ridge, S. A.; Worwood, M.; Oscier, D.; Jacobs, A.; Padua, R. A.: FMS mutations in myelodysplastic, leukemic, and normal subjects. *Proc.Nat. Acad. Sci.* 87: 1377–1380, 1990.

[67785] 6115.Roberts, W. M.; Look, A. T.; Roussel, M. F.; Sherr, C. J.: Tandemlinkage of human CSF–1 receptor (c–fms) and

PDGF receptor genes. *Cell* 55:655–661, 1988.

- [67786] 6116.Sapi, E.; Flick, M. B.; Kacinski, B. M.: The first intron of human c-fms proto-oncogene contains a processed pseudogene (RPL7P) for ribosomal protein L7. *Genomics* 22: 641–645, 1994.
- [67787] 6117.Verbeek, J. S.; Roebroek, A. J. M.; van den Ouweland, A. M. W.; Bloemers, H. P. J.; Van de Ven, W. J. M.: Human c-fms proto-oncogene: comparative analysis with an abnormal allele. *Molec. Cell. Biol.* 5:422–426, 1985.
- [67788] 6118.Verbeek, J. S.; van Heerikhuizen, H.; de Pauw, B. E.; Haanen, C.; Bloemers, H. P. J.; Van de Ven, W. J. M.: A hereditary abnormal c-fms proto-oncogene in a patient with acute lymphocytic leukaemia and congenital hypothyroidism. *Brit. J. Haemat.* 61: 135–138, 1985.
- [67789] 6119.Yarden, Y.; Ullrich, A.: Growth factor receptor tyrosine kinases. *Ann.Rev. Biochem.* 57: 443–478, 1988.
- [67790] 6120.Brown, J. R.; Ye, H.; Bronson, R. T.; Dikkes, P.; Greenberg, M.E.: A defect in nurturing in mice lacking the immediate early gene fosB. *Cell* 86: 297–309, 1996.
- [67791] 6121.Forsdyke, D.: Personal Communication. Kingston, Ontario, Canada 6/8/1992.
- [67792] 6122.Lazo, P. S.; Dorfman, K.; Noguchi, T.; Mattei, M. G.; Bravo, R.: Structure and mapping of the fosB gene: FosB

downregulates the activity of the fosB promoter. *Nucleic Acids Res.* 20: 343–350, 1992.

[67793] 6123. Nakabeppu, Y.; Nathans, D.: A naturally occurring truncated form of FosB that inhibits Fos/Jun transcriptional activity. *Cell* 64:751–759, 1991.

[67794] 6124. Siderovski, D. P.; Blum, S.; Forsdyke, R. E.; Forsdyke, D. R.: A set of human putative lymphocyte G0/G1 switch genes includes genes homologous to rodent cytokine and zinc finger protein-encoding genes. *Dev Cell Biol.* 9: 579–587, 1990.

[67795] 6125. Kelz, M. B.; Chen, J.; Carlezon, W. A., Jr.; Whisler, K.; Gilden, L.; Beckmann, A. M.; Steffen, C.; Zhang, Y.-J.; Marotti, L.; Self, D. W.; Tkatch, T.; Baranauskas, G.; Surmeier, D. J.; Neve, R. L.; Duman, R. S.; Picciotto, M. R.; Nestler, E. J.: Expression of the transcription factor delta-FosB in the brain controls sensitivity to cocaine. *Nature* 401: 272–276, 1999.

[67796] 6126. Boyd, M. T.; Vlatkovic, N.; Haines, D. S.: A novel cellular protein (MTBP) binds to MDM2 and induces a G1 arrest that is suppressed by MDM2. *J. Biol. Chem.* 275: 31883–31890, 2000.

[67797] 6127. Bueso-Ramos, C. E.; Yang, Y.; deLeon, E.; McCown, P.; Stass, S. A.; Albitar, M.: The human MDM-2 oncogene is

overexpressed in leukemias. *Blood* 82:2617–2623, 1993.

[67798] 6128. Buschmann, T.; Fuchs, S. Y.; Lee, C.-G.; Pan, Z.-Q.; Ronai, Z.: SUMO-1 modification of Mdm2 prevents its self-ubiquitination and increases Mdm2 ability to ubiquitinate p53. *Cell* 101: 753–762, 2000. Note: Erratum: *Cell* 107: 549 only, 2001.

[67799] 6129. Hoffmeyer, S.; Burk, O.; von Richter, O.; Arnold, H. P.; Brockmoller, J.; John, A.; Cascorbi, I.; Gerloff, T.; Roots, I.; Eichelbaum, M.; Brinkmann, U.: Functional polymorphisms of the human multidrug-resistance gene: multiple sequence variations and correlation of one allele with P-glycoprotein expression and activity in vivo. *Proc. Nat. Acad. Sci.* 97: 3473–3478, 2000.

[67800] 6130. Fruman, D. A.; Snapper, S. B.; Yballe, C. M.; Davidson, L.; Yu, J. Y.; Alt, F. W.; Cantley, L. C.: Impaired B cell development and proliferation in absence of phosphoinositide 3-kinase p85- $\alpha$ . *Science* 283:393–397, 1999.

[67801] 6131. Fukao, T.; Tanabe, M.; Terauchi, Y.; Ota, T.; Matsuda, S.; Asano, T.; Kadowaki, T.; Takeuchi, T.; Koyasu, S.: PI3K-mediated negative feedback regulation of IL-12 production in DCs. *Nature Immun.* 3:875–881, 2002.

[67802] 6132. Fukao, T.; Yamada, T.; Tanabe, M.; Terauchi, Y.; Ota,



T.; Takayama, T.; Asano, T.; Takeuchi, T.; Kadowaki, T.; Hata, J.; Koyasu, S.: Selective loss of gastrointestinal mast cells and impaired immunity in PI3K-deficient mice. *Nature Immun.* 3: 295–304, 2002.

[67803] 6133. Hoyle, J.; Yulug, I. G.; Egan, S. E.; Fisher, E. M. C.: The gene that encodes the phosphatidylinositol-3 kinase regulatory subunit (p85- $\alpha$ ) maps to chromosome 13 in the mouse. *Genomics* 24: 400–402, 1994.

[67804] 6134. Otsu, M.; Hiles, I.; Gout, I.; Fry, M. J.; Ruiz-Larrea, F.; Panayotou, G.; Thompson, A.; Dhand, R.; Hsuan, J.; Totty, N.; Smith, A. D.; Morgan, S. J.; Courtneidge, S. A.; Parker, P. J.; Waterfield, M. D.: Characterization of two 85 kd proteins that associate with receptor tyrosine kinases, middle-T/pp60(c-src) complexes, and PI3-kinase. *Cell* 65: 91–104, 1991.

[67805] 6135. Skolnik, E. Y.; Margolis, B.; Mohammadi, M.; Lowenstein, E.; Fischer, R.; Drepps, A.; Ullrich, A.; Schlessinger, J.: Cloning of PI3-kinase-associated p85 utilizing a novel method for expression/cloning of target proteins for receptor tyrosine kinases. *Cell* 65: 83–90, 1991.

[67806] 6136. Suzuki, H.; Terauchi, Y.; Fujiwara, M.; Aizawa, S.; Yazaki, Y.; Kadowaki, T.; Koyasu, S.: Xid-like immunodeficiency in mice with disruption of the p85- $\alpha$  subunit of

phosphoinositide 3-kinase. *Science* 283:390–392, 1999.

[67807] 6137. Terauchi, Y.; Tsuji, Y.; Satoh, S.; Minoura, H.; Murakami, K.; Okuno, A.; Inukai, K.; Asano, T.; Kaburagi, Y.; Ueki, K.; Nakajima, H.; Hanafusa, T.; and 18 others: Increased insulin sensitivity and hypoglycaemia in mice lacking the p85- $\alpha$  subunit of phosphoinositide 3-kinase. *Nature Genet.* 21: 230–235, 1999.

[67808] 6138. Volinia, S.; Patracchini, P.; Otsu, M.; Hiles, I.; Gout, I.; Calzolari, E.; Bernardi, F.; Rooke, L.; Waterfield, M. D.: Chromosomal localization of human p85- $\alpha$ , a subunit of phosphatidylinositol 3-kinase, and its homologue p85- $\beta$ . *Oncogene* 7: 789–793, 1992.

[67809] 6139. Bonne, G.; Di Barletta, M. R.; Varnous, S.; Becane, H.-M.; Hammouda, E.-H.; Merlini, L.; Muntoni, F.; Greenberg, C. R.; Gary, F.; Urtizberea, J.-A.; Duboc, D.; Fardeau, M.; Toniolo, D.; Schwartz, K.: Mutations in the gene encoding lamin A/C cause autosomal dominant Emery-Dreifuss muscular dystrophy. *Nature Genet.* 21: 285–288, 1999.

[67810] 6140. Budarf, M. L.; Konkle, B. A.; Ludlow, L. B.; Michaud, D.; Li, M.; Yamashiro, D. J.; McDonald-McGinn, D.; Zackai, E. H.; Driscoll, D. A.: Identification of a patient with Bernard-Soulier syndrome and a deletion in the DiGeorge/

velo-cardio-facial chromosomal region in 22q11.2. Hum. Molec. Genet. 4: 763–766, 1995.

[67811] 6141. Kelly, M. D.; Essex, D. W.; Shapiro, S. S.; Meloni, F. J.; Druck, T.; Huebner, K.; Konkle, B. A.: Complementary DNA cloning of the alternatively expressed endothelial cell glycoprotein Ib-beta (GPIb-beta) and localization of the GPIb-beta gene to chromosome 22. J. Clin. Invest. 93: 2417–2424, 1994.

[67812] 6142. Kunishima, S.; Lopez, J. A.; Kobayashi, S.; Imai, N.; Kamiya, T.; Saito, H.; Naoe, T.: Missense mutations of the glycoprotein (GP)Ib-beta gene impairing the GPIb alpha/beta disulfide linkage in a family with giant platelet disorder. Blood 89: 2404–2412, 1997.

[67813] 6143. Lopez, J. A.; Chung, D. W.; Fujikawa, K.; Hagen, F. S.; Davie, E. W.; Roth, G. J.: The alpha and beta chains of human platelet glycoproteinIb are both transmembrane proteins containing a leucine-rich amino acid sequence. Proc. Nat. Acad. Sci. 85: 2135–2139, 1988.

[67814] 6144. Lopez, J. A.; Chung, D. W.; Fujikawa, K.; Hagen, F. S.; Papayannopoulou, T.; Roth, G. J.: Cloning of the alpha chain of human platelet glycoproteinIb: a transmembrane protein with homology to leucine-rich alpha 2-glycoprotein. Proc. Nat. Acad. Sci. 84: 5615–5619,

1987.

- [67815] 6145. Roth, G. J.: The wanderings of a platelet gene: what is 'neo'Ib-beta telling us? (Editorial) J. Clin. Invest. 93: 2301-2302, 1994.
- [67816] 6146. Yagi, M.; Edelhoff, S.; Disteche, C. M.; Roth, G. J.: Structural characterization and chromosomal location of the gene encoding human platelet glycoprotein Ib-beta. J. Biol. Chem. 269: 17424-17427, 1994.
- [67817] 6147. Zieger, B.; Hashimoto, Y.; Ware, J.: Alternative expression of platelet glycoprotein Ib-beta mRNA from an adjacent 5-prime gene with an imperfect polyadenylation signal sequence. J. Clin. Invest. 99: 520-525, 1997.
- [67818] 6148. Bakker, E.; Pearson, P. L.; Meera Khan, P.; Schreuder, G. M. T.; Madan, K.: Orientation of major histocompatibility (MHC) genes relative to the centromere of human chromosome 6. Clin. Genet. 15: 198-202, 1979.
- [67819] 6149. Bender, K.; Grzeschik, K. H.: Assignment of the genes for human glyoxalase I to chromosome 6 and for human esterase D to chromosome 13. Cytogenet. Cell Genet. 16: 93-96, 1976.
- [67820] 6150. Beretta, M.; Schiliro, G.; Russo, A.; Barbujani, G.; Mazzetti, P.; Russo, G.; Barrai, I.: A new rare variant of the glyoxalase I system of the red cell: GLO-Sicily. Am. J. Hum.

Genet. 35: 1042–1047,1983.

- [67821] 6151. Blanche, H.; Zoghbi, H. Y.; Jabs, E. W.; de Gouyon, B.; Zunec, R.; Dausset, J.; Cann, H. M.: A centromere-based genetic map of the short arm of human chromosome 6. Genomics 9: 420–428, 1991.
- [67822] 6152. Carter, N. D.; West, C. M.; Bernard, J. M.; Farid, N. R.; Larsen, B.; Marshall, W. H.: Linkage of glyoxalase I and HLA in two Newfoundland communities. Hum. Hered. 28: 397–400, 1978.
- [67823] 6153. Giblett, E. R.; Lewis, M.: Gene linkage studies on glyoxalase I. Cytogenet. Cell Genet. 16: 313 only, 1976.
- [67824] 6154. Goldman, D.; O'Brien, S. J.; Lucas-Derse, S.; Dean, M.: Linkage mapping of human polymorphic proteins identified by two-dimensional electrophoresis. Genomics 11: 875–884, 1991.
- [67825] 6155. Hansen, H. E.; Eriksen, B.: HLA-GLO linkage analysis in 57 informative families. Hum. Hered. 29: 355–360, 1979.
- [67826] 6156. Karlsson, S.; Arnason, A.; Jensson, O.: GLO polymorphism in Iceland. Hum. Hered. 30: 383–385, 1980.
- [67827] 6157. Kavathas, P.; DeMars, R.: A new variant glyoxalase I allele that is readily detectable in stimulated lymphocytes and lymphoblastoid cell lines but not in circulating lym-

phocytes or erythrocytes. Am.J. Hum. Genet. 33: 935–945, 1981.

[67828] 6158.Kompe, J.; Bissbort, S.; Gussmann, S.; Ritter, H.: Polymorphism of red cell glyoxalase I (E.C.4.4.1.5), a new genetic marker in man: investigation of 169 mother–child combinations. Humangenetik 27:141–143, 1975.

[67829] 6159.Kompe, J.; Bissbort, S.; Ritter, H.: Red cell glyoxalase I (E.C.4.4.1.5): formal genetics and linkage relations. Humangenetik 28: 249–251, 1975.

[67830] 6160.Kompe, J.; Siebert, G.; Ritter, H.; Heilbronner, H.; Schunter, F.; Wernet, P.; Gupta, D.; Moeller, H.: Data on linkage relations between GLO and 21–hydroxylase. Hum. Genet. 54: 419–420, 1980.

[67831] 6161.Meo, T.; Douglas, T.; Rijnbeek, A.–M.: Glyoxalase I polymorphism in the mouse: a new genetic marker linked to H-2. Science 198:311–313, 1977.

[67832] 6162.Olaisen, B.; Gedde–Dahl, T., Jr.; Thorsby, E.: Localization of the human GLO gene locus. Hum. Genet. 32: 301–304, 1976.

[67833] 6163.Parr, C. W.; Bagster, I. A.; Welch, S. G.: Human red cell glyoxalase I polymorphism. Biochem. Genet. 15: 109–114, 1977.

[67834] 6164.Reinsmoen, N. L.; Friend, P. S.; Miller, W. V.;

Burgdorf, A.; Giblett, E. R.; Yunis, E. J.: Inheritance of recombinant HLA-GLO haplotype suggesting the gene sequence. *Nature* 267: 276–278, 1977.

[67835] 6165. Rubinstein, P.; Suciú-Foca, N.: Glyoxalase 1: a possible 'null' allele. *Hum. Hered.* 29: 217–220, 1979.

[67836] 6166. Schimandle, C. M.; Vander Jagt, D. L.: Isolation and kinetic analysis of the multiple forms of glyoxalase-1 from human erythrocytes. *Arch. Biochem. Biophys.* 195: 261–268, 1979.

[67837] 6167. Sparkes, R. S.; Sparkes, M. C.; Crist, M.; Anderson, C. E.: Glyoxalase 1 'null' allele in a new family: identification by abnormal segregation pattern and quantitative assay. *Hum. Genet.* 64: 146–147, 1983.

[67838] 6168. Teng, Y. S.; Tan, S. G.; Lopez, C. G.: Red cell glyoxalase and placental soluble aconitase polymorphisms in the three major ethnic groups of Malaysia. *Jpn. J. Hum. Genet.* 23: 211–215, 1978.

[67839] 6169. Kartner, N.; Evernden-Porelle, D.; Bradley, G.; Ling, V.: Detection of P-glycoprotein in multidrug-resistant cell lines by monoclonal antibodies. *Nature* 316: 820–823, 1985.

[67840] 6170. Kim, R. B.; Fromm, M. F.; Wandel, C.; Leake, B.; Wood, A. J. J.; Roden, D. M.; Wilkinson, G. R.: The drug

transporter P-glycoprotein limits oral absorption and brain entry of HIV-1 protease inhibitors. *J.Clin. Invest.* 101: 289-294, 1998.

[67841] 6171. Lankas, G. R.; Wise, L. D.; Cartwright, M. E.; Pippert, T.; Umbenhauer, D. R.: Placental P-glycoprotein deficiency enhances susceptibility to chemically induced birth defects in mice. *Reprod. Toxicol.* 12:457-463, 1998.

[67842] 6172. Martinsson, T.; Levan, G.: Localization of the multidrug resistance-associated 170 kDa P-glycoprotein gene to mouse chromosome 5 and to homogeneously staining regions in multidrug-resistant mouse cells by in situ hybridization. *Cytogenet. Cell Genet.* 45: 99-101, 1987.

[67843] 6173. Mealey, K. L.; Bentjen, S. A.; Gay, J. M.; Cantor, G. H.: Ivermectin sensitivity in collies is associated with a deletion mutation of the *mdr1* gene. *Pharmacogenetics* 11: 727-733, 2001.

[67844] 6174. Mickley, L. A.; Spengler, B. A.; Knutsen, T. A.; Biedler, J. L.; Fojo, T.: Gene rearrangement: a novel mechanism for MDR-1 gene activation. *J.Clin. Invest.* 99: 1947-1957, 1997.

[67845] 6175. Pastan, I.; Gottesman, M.: Multiple-drug resistance in human cancer. *New Eng. J. Med.* 316: 1388-1393, 1987.

[67846] 6176. Pulliam, J. D.; Seward, R. L.; Henry, R. T.; Steinberg,



S. A.: Investigating ivermectin toxicity in collies. *Vet. Med.* 80: 33–40, 1985.

- [67847] 6177. Randolph, G. J.; Beaulieu, S.; Pope, M.; Sugawara, I.; Hoffman, L.; Steinman, R. M.; Muller, W. A.: A physiologic function for p-glycoprotein (MDR-1) during the migration of dendritic cells from skin via afferent lymphatic vessels. *Proc. Nat. Acad. Sci.* 95: 6924–6929, 1998.
- [67848] 6178. Riordan, J. R.; Deuchars, K.; Kartner, N.; Alon, N.; Trent, J.; Ling, V.: Amplification of P-glycoprotein genes in multidrug-resistant mammalian cell lines. *Nature* 316: 817–819, 1985.
- [67849] 6179. Roninson, I. B.; Chin, J. E.; Choi, K.; Gros, P.; Housman, D. E.; Fojo, A.; Shen, D.; Gottesman, M. M.; Pastan, I.: Isolation of human *mdr* DNA sequences amplified in multidrug-resistant KB carcinoma cells. *Proc. Nat. Acad. Sci.* 83: 4538–4542, 1986.
- [67850] 6180. Ruiz, J. C.; Choi, K.; Von Hoff, D. D.; Roninson, I. B.; Wahl, G. M.: Autonomously replicating episomes contain *MDR1* genes in a multidrug-resistant human cell line. *Molec. Cell. Biol.* 9: 109–115, 1989.
- [67851] 6181. Safa, A. R.; Stern, R. K.; Choi, K.; Agresti, M.; Tamai, I.; Mehta, N. D.; Roninson, I. B.: Molecular basis of preferential resistance to colchicine in multidrug-resistant hu-

man cells conferred by gly185-to-val185 substitution in P-glycoprotein. Proc. Nat. Acad. Sci. 87: 7225–7229, 1990.

[67852] 6182. Schinkel, A. H.; Smit, J. J. M.; van Tellingen, O.; Beijnen, J. H.; Wagenaar, E.; van Deemter, L.; Mol, C. A. A. M.; van der Valk, M. A.; Robanus-Maandag, E. C.; te Riele, H. P. J.; Berns, A. J. M.; Borst, P.: Disruption of the mouse *mdr1a* P-glycoprotein gene leads to a deficiency in the blood-brain barrier and to increased sensitivity to drugs. Cell 77: 491–502, 1994.

[67853] 6183. Shen, D.-W.; Fojo, A.; Chin, J. E.; Roninson, I. B.; Richert, N.; Pastan, I.; Gottesman, M. M.: Human multidrug-resistant cell lines: increased *mdr1* expression can precede gene amplification. Science 232: 643–645, 1986.

[67854] 6184. Slovak, M. L.; Hoeltge, G. A.; Trent, J. M.: Cytogenetic alterations associated with the acquisition of doxorubicin resistance: possible significance of chromosome 7 alterations. Cancer Res. 47: 6646–6652, 1987.

[67855] 6185. Smit, J. W.; Huisman, M. T.; van Tellingen, O.; Wiltshire, H. R.; Schinkel, A. H.: Absence or pharmacological blocking of placental P-glycoprotein profoundly increases fetal drug exposure. J. Clin. Invest. 104: 1441–1447, 1999.

[67856] 6186. Synold, T. W.; Dussault, I.; Forman, B. M.: The orphan nuclear receptor SXR coordinately regulates drug

metabolism and efflux. *NatureMed.* 7: 584–590, 2001.

- [67857] 6187. Taylor, H. R.; Pacque, M.; Munoz, B.; Greene, B. M.: Impact of mass treatment of onchocerciasis with ivermectin on the transmission of infection. *Science* 250: 116–118, 1990.
- [67858] 6188. Trent, J.; Bell, D.; Willard, H.; Ling, V.: Chromosomal localization in normal human cells and CHO cells of a sequence derived from P-glycoprotein (PGY1). (Abstract) *Cytogenet. Cell Genet.* 40: 761–762, 1985.
- [67859] 6189. Trent, J. M.; Witkowski, C. M.: Clarification of the chromosomal assignment of the human P-glycoprotein/MDR1 gene: possible coincidence with the cystic fibrosis and c-met oncogene. *Cancer Genet. Cytogenet.* 26: 187–190, 1987.
- [67860] 6190. Trezise, A. E. O.; Romano, P. R.; Gill, D. R.; Hyde, S. C.; Sepulveda, F. V.; Buchwald, M.; Higgins, C. F.: The multidrug resistance and cystic fibrosis genes have complementary patterns of epithelial expression. *EMBO J.* 11: 4291–4303, 1992.
- [67861] 6191. Ueda, K.; Cardarelli, C.; Gottesman, M. M.; Pastan, I.: Expression of a full-length cDNA for the human 'MDR1' gene confers resistance to colchicine, doxorubicin, and vinblastine. *Proc. Nat. Acad. Sci.* 84: 3004–3008, 1987.

- [67862] 6192.Ueda, K.; Clark, D. P.; Chen, C.; Roninson, I. B.; Gottesman, M. M.; Pastan, I.: The human multidrug resistance (mdr1) gene: cDNA cloning and transcription initiation. *J. Biol. Chem.* 262: 505–508, 1987.
- [67863] 6193.Ueda, K.; Cornwell, M. M.; Gottesman, M. M.; Pastan, I.; Roninson, I. B.; Ling, V.; Riordan, J. R.: The mdr1 gene, responsible for multidrug-resistance, codes for P-glycoprotein. *Biochem. Biophys. Res. Commun.* 141: 956–962, 1986.
- [67864] 6194.Ueda, K.; Pastan, I.; Gottesman, M. M.: Isolation and sequence of the promoter region of the human multidrug-resistance (P-glycoprotein) gene. *J. Biol. Chem.* 262: 17432–17436, 1987.
- [67865] 6195.Umbenhauer, D. R.; Lankas, G. R.; Pippert, T. R.; Wise, L. D.; Cartwright, M. E.; Hall, S. J.; Beare, C. M.: Identification of a P-glycoprotein-deficient subpopulation in the CF-1 mouse strain using a restriction fragment length polymorphism. *Toxicol. Appl. Pharm.* 146:88–94, 1997.
- [67866] 6196.van Helvoort, A.; Smith, A. J.; Sprong, H.; Fritzsche, I.; Schinkel, A. H.; Borst, P.; van Meer, G.: MDR1 P-glycoprotein is a lipid translocase of broad specificity, while MDR3 P-glycoprotein specifically translocates phosphatidylcholine. *Cell* 87: 507–517, 1996.

- [67867] 6197.Woods, K. A.; Camacho–Hubner, C.; Bergman, R. N.; Barter, D.;Clark, A. J. L.; Savage, M. O.: Effects of insulin–like growth factorI (IGF–I) therapy on body composition and insulin resistance in IGF–Igene deletion. J. Clin. Endocr. Metab. 85: 1407–1411, 2000.
- [67868] 6198.Woods, K. A.; Camacho–Hubner, C.; Savage, M. O.; Clark, A. J.L.: Intrauterine growth retardation and postnatal growth failureassociated with deletion of the insulin–like growth factor I gene. NewEng. J. Med. 335: 1363–1367, 1996.
- [67869] 6199.Yang–Feng, T. L.; Brissenden, J. E.; Ullrich, A.; Francke, U.: Sub–regional localization of human genes for insulin–like growthfactors I (IGF1) and II (IGF2) by in situ hybridization. (Abstract) Cytogenet.Cell Genet. 40: 782 only, 1985.
- [67870] 6200.Yanovski, J. A.; Sovik, K. N.; Nguyen, T. T.; Sebring, N. G.:Insulin–like growth factors and bone mineral density in African Americanand white girls. J. Pediat. 137: 826–832, 2000.
- [67871] 6201.Zhu, J.; Kahn, C. R.: Analysis of a peptide hormone–receptorinteraction in the yeast two–hybrid system. Proc. Nat. Acad. Sci. 94:13063–13068, 1997.
- [67872] 6202.Kobayashi, M.; Takamatsu, K.; Saitoh, S.; Miura, M.;

Noguchi, T.: Molecular cloning of hippocalcin, a novel calcium-binding protein of the recoverin family exclusively expressed in hippocampus. *Biochem. Biophys. Res. Commun.* 189: 511–517, 1992.

[67873] 6203. Takamatsu, K.; Kobayashi, M.; Saitoh, S.; Fujishiro, M.; Noguchi, T.: Molecular cloning of human hippocalcin cDNA and chromosomal mapping of its gene. *Biochem. Biophys. Res. Commun.* 200: 606–611, 1994.

[67874] 6204. Borrego, S.; Ruiz, A.; Saez, M. E.; Gimm, O.; Gao, X.; Lopez-Alonso, M.; Hernandez, A.; Wright, F. A.; Antinolo, G.; Eng, C.: RET genotypes comprising specific haplotypes of polymorphic variants predispose to isolated Hirschsprung disease. *J. Med. Genet.* 37: 572–578, 2000.

[67875] 6205. Borrego, S.; Saez, M. E.; Ruiz, A.; Gimm, O.; Lopez-Alonso, M.; Antinolo, G.; Eng, C.: Specific polymorphisms in the RET proto-oncogene are over-represented in patients with Hirschsprung disease and may represent loci modifying phenotypic expression. *J. Med. Genet.* 36: 771–774, 1999.

[67876] 6206. Goldstein, J. L.; Brown, M. S.; Stone, N. J.: Genetics of the LDL receptor: evidence that the mutations affecting binding and internalization are allelic. *Cell* 12: 629–641, 1977.

- [67877] 6207. Goldstein, J. L.; Dana, S. E.; Brunschede, G. Y.; Brown, M. S.: Genetic heterogeneity in familial hypercholesterolemia: evidence for two different mutations affecting functions of low-density lipoprotein receptor. *Proc. Nat. Acad. Sci.* 72: 1092–1096, 1975.
- [67878] 6208. Hobbs, H. H.; Brown, M. S.; Goldstein, J. L.: Molecular genetics of the LDL receptor gene in familial hypercholesterolemia. *Hum. Mutat.* 1:445–466, 1992.
- [67879] 6209. Hobbs, H. H.; Brown, M. S.; Goldstein, J. L.; Russell, D. W.: Deletion of exon encoding cysteine-rich repeat of low density lipoprotein receptor alters its binding specificity in a subject with familial hypercholesterolemia. *J. Biol. Chem.* 261: 13114–13120, 1986.
- [67880] 6210. Hobbs, H. H.; Russell, D. W.; Brown, M. S.; Goldstein, J. L.: The LDL receptor locus in familial hypercholesterolemia: mutational analysis of a membrane protein. *Annu. Rev. Genet.* 24: 133–170, 1990.
- [67881] 6211. Horsthemke, B.; Beisiegel, U.; Dunning, A.; Havinga, J. R.; Williamson, R.; Humphries, S.: Unequal crossing-over between two Alu-repetitive DNA sequences in the low-density-lipoprotein-receptor gene: a possible mechanism for the defect in a patient with familial hypercholesterolaemia. *Europ. J. Biochem.* 164: 77–81, 1987.

- [67882] 6212.Khachadurian, A. K.: The inheritance of essential familial hypercholesterolemia. *Am.J. Med.* 37: 402–407, 1964.
- [67883] 6213.Kingsley, D. M.; Krieger, M.: Receptor-mediated endocytosis of low density lipoprotein: somatic cell mutants define multiple genes required for expression of surface-receptor activity. *Proc. Nat.Acad. Sci.* 81: 5454–5458, 1984.
- [67884] 6214.Knight, B. L.; Gavigan, S. J. P.; Soutar, A. K.; Patel, D. D.: Defective processing and binding of low-density lipoprotein receptors in fibroblasts from a familial hypercholesterolaemic subject. *Europ.J. Biochem.* 179: 693–698, 1989.
- [67885] 6215.Knoblauch, H.; Muller-Myhsok, B.; Busjahn, A.; Ben Avi, L.; Bähring, S.; Baron, H.; Heath, S. C.; Uhlmann, R.; Faulhaber, H.-D.; Shpitzen, S.; Aydin, A.; Reshef, A.; and 11 others: A cholesterol-lowering gene maps to chromosome 13q. *Am. J. Hum. Genet.* 66: 157–166, 2000.
- [67886] 6216.Komuro, I.; Kato, H.; Nakagawa, T.; Takahashi, K.; Mimori, A.; Takeuchi, F.; Nishida, Y.; Miyamoto, T.: Case report: the longest-lived patient with homozygous familial hypercholesterolemia secondary to a defect in internalization of the LDL receptor. *Am. J. Med. Sci.* 294:341–345,



1987.

- [67887] 6217.Kotze, M. J.; Langenhoven, E.; Retief, A. E.; Steyn, K.; Marais, M. P.; Grobbelaar, J. J.; Oosthuizen, C. J. J.; Weich, H. F. H.; Benade, A. J. S.: Haplotype associations of three DNA polymorphisms at the human low density lipoprotein receptor gene locus in familial hypercholesterolemia. *J. Med. Genet.* 24: 750–755, 1987.
- [67888] 6218.Grundy, H. O.; Peltz, G.; Moore, K. W.; Golbus, M. S.; Jackson, L. G.; Lebo, R. V.: The polymorphic Fc-gamma receptor II gene maps to human chromosome 1q. *Immunogenetics* 29: 331–339, 1989.
- [67889] 6219.Kudo, A.; Melchers, F.: A second gene, Vpre-B in the lambda 5 locus of the mouse, which appears to be selectively expressed in pre-B lymphocytes. *EMBO J.* 6: 2267–2272, 1987.
- [67890] 6220.Mattei, M.-G.; Fumoux, F.; Roeckel, N.; Fougereau, M.; Schiff, C.: The human pre-B-specific lambda-like cluster is located in the 22q11.2–22q12.3 region, distal to the IgC-lambda locus. *Genomics* 9:544–546, 1991.
- [67891] 6221.Minegishi, Y.; Coustan-Smith, E.; Wang, Y.-H.; Cooper, M. D.; Campana, D.; Conley, M. E.: Mutations in the human lambda-5/14.1 gene result in B cell deficiency and agammaglobulinemia. *J. Exp. Med.* 187: 71–77, 1998.

- [67892] 6222.Pillai, S.; Baltimore, D.: The omega and iota surrogate immunoglobulin light chains. *Curr. Top. Microbiol. Immun.* 137: 136–139, 1988.
- [67893] 6223.Sakaguchi, N.; Melchers, F.: Lambda 5, a new light-chain-related locus selectively expressed in pre-B lymphocytes. *Nature* 324: 579–582, 1986.
- [67894] 6224.Schiff, C.; Milili, M.; Fougereau, M.: Isolation of early immunoglobulin lambda-like gene transcripts in human fetal liver. *Europ. J. Immun.* 19:1873–1878, 1989.
- [67895] 6225.Lin, P.-F.; Slate, D. L.; Lawyer, F. C.; Ruddle, F. H.: Assignment of the murine interferon sensitivity and cytoplasmic superoxide dismutase genes to chromosome 16. *Science* 209: 285–287, 1980.
- [67896] 6226.Holmes, W. E.; Lee, J.; Kuang, W.-J.; Rice, G. C.; Wood, W. I.: Structure and functional expression of a human interleukin-8 receptor. *Science* 253:1278–1280, 1991.
- [67897] 6227.Sligh, J. E., Jr.; Ballantyne, C. M.; Rich, S. S.; Hawkins, H.K.; Smith, C. W.; Bradley, A.; Beaudet, A. L.: Inflammatory and immune responses are impaired in mice deficient in intercellular adhesion molecule 1. *Proc. Nat. Acad. Sci.* 90: 8529–8533, 1993.
- [67898] 6228.Nishimura, D.; Buetow, K. H.; Yamada, Y.; Murray, J.

C.: RFLPs and linkage relationships of the human laminin B2 gene. *Genomics* 3:393–395, 1988.

- [67899] 6229. Karaplis, A. C.; Luz, A.; Glowacki, J.; Bronson, R. T.; Tybulewicz, V. L. J.; Kronenberg, H. M.; Mulligan, R. C.: Lethal skeletal dysplasia from targeted disruption of the parathyroid hormone-related peptide gene. *Genes Dev.* 8: 277–289, 1994.
- [67900] 6230. Bevilacqua, G.; Sobel, M. E.; Liotta, L. A.; Steeg, P. S.: Association of low nm23 RNA levels in human primary infiltrating ductal breast carcinomas with lymph node involvement and other histopathological indicators of high metastatic potential. *Cancer Res.* 49: 5185–5190, 1989.
- [67901] 6231. Chang, C. L.; Zhu, X.; Thoraval, D. H.; Ungar, D.; Rawwas, J.; Hora, N.; Strahler, J. R.; Hanash, S. M.; Radany, E.: nm23-H1 mutation in neuroblastoma. (Letter) *Nature* 370: 335–336, 1994.
- [67902] 6232. Dooley, S.; Seib, T.; Engel, M.; Theisinger, B.; Janz, H.; Piontek, K.; Zang, K.-D.; Welter, C.: Isolation and characterization of the human genomic locus coding for the putative metastasis control gene nm23-H1. *Hum. Genet.* 93: 63–66, 1994.
- [67903] 6233. Eddy, R. L.; Mendola, C. E.; Fairhurst, J. L.; Shows, T. B.; O'Hara, B.; Kovesdi, I.; Backer, J. M.: Metastasis sup-

pressor gene is mapped to human chromosome 17p11-qter. (Abstract) Cytogenet. Cell Genet. 58:2005 only, 1991.

- [67904] 6234. Gilles, A.-M.; Presecan, E.; Vonica, A.; Lascu, I.: Nucleosidediphosphate kinase from human erythrocytes: structural characterization of the two polypeptide chains responsible for heterogeneity of the hexameric enzyme. J. Biol. Chem. 266: 8784-8789, 1991.
- [67905] 6235. Keim, D.; Hailat, N.; Melhem, R.; Zhu, X. X.; Lascu, I.; Veron, M.; Strahler, J.; Hanash, S. M.: Proliferation-related expression of p19/nm23 nucleoside diphosphate kinase. J. Clin. Invest. 89:919-924, 1992.
- [67906] 6236. Drayna, D.; Fielding, C.; McLean, J.; Baer, B.; Castro, G.; Chen, E.; Comstock, L.; Henzel, W.; Kohr, W.; Rhee, L.; Wion, K.; Lawn, R.: Cloning and expression of human apolipoprotein D cDNA. J. Biol. Chem. 261: 16535-16539, 1986.
- [67907] 6237. Drayna, D.; Scott, J. D.; Lawn, R.: Multiple RFLPs at the human apolipoprotein D (APOD) locus. Nucleic Acids Res. 15: 9617 only, 1987.
- [67908] 6238. Whittington, J. E.; Keats, B. J. B.; Jackson, J. F.; Currier, R. D.; Terasaki, P. I.: Linkage studies on glyoxalase I (GLO), pepsinogen (PG), spinocerebellar ataxia (SCA1), and

HLA. Cytogenet. Cell Genet. 28:145–150, 1980.

- [67909] 6239.Ziegler, A.; Fonatsch, C.; Kompf, J.: Mapping of the locus for glyoxalase 1 (GLO1) on human chromosome 6 using mutant cell lines.(Abstract) Cytogenet. Cell Genet. 40: 787 only, 1985.
- [67910] 6240.Agar, N. S.; Board, P. G.; Bell, K.: Studies of erythrocyte glyoxalase II in various domestic species: discovery of glyoxalase II deficiency in the horse. Animal Blood Groups Biochem. Genet. 15: 67–70, 1984.
- [67911] 6241.Ball, J. C.; Vander Jagt, D. L.: Purification of S-2-hydroxyacylglutathione hydrolase (glyoxalase II) from rat erythrocytes. Anal. Biochem. 98:472–477, 1979.
- [67912] 6242.Board, P. G.: Genetic polymorphism of human erythrocyte glyoxalase II. Am. J. Hum. Genet. 32: 690–694, 1980.
- [67913] 6243.Board, P. G.; Gibbs, C. J., Jr.; Gajdusek, D. C.: Polymorphism of erythrocyte glyoxalase II in anthropoid primates. Folia Primatol. 36:138–143, 1981.
- [67914] 6244.Honey, N. K.; Shows, T. B.: Assignment of the glyoxalase II gene (HAGH) to human chromosome 16. Hum. Genet. 58: 358–361, 1981.
- [67915] 6245.Mulley, J. C.; Barton, N.; Callen, D. F.: Localisation of human PGP and HAGH genes to 16p13.3. Cytogenet. Cell

Genet. 53: 175–176,1990.

[67916] 6246.Mulley, J. C.; Callen, D. F.: New regional localisations for HAGHand PGP on human chromosome 16. Hum. Genet. 74: 423–424, 1986.

[67917] 6247.Ridderstrom, M.; Saccucci, F.; Hellman, U.; Bergman, T.; Principato,G.; Mannervik, B.: Molecular cloning, heterologous expression, and characterization of human glyoxalase II. J. Biol. Chem. 271: 319–323,1996.

[67918] 6248.Valentine, W. N.; Paglia, D. E.; Neerhout, R. C.; Konrad, P. N.: Erythrocyte glyoxalase II deficiency with coincidental hereditaryelliptocytosis. Blood 36: 797–808, 1970.

[67919] 6249.Beranova, M.; Oliveira, L. M. B.; Bedecarrats, G. Y.; Schipani,E.; Vallejo, M.; Ammini, A. C.; Quintos, J. B.; Hall, J. E.; Martin,K. A.; Hayes, F. J.; Pitteloud, N.; Kaiser, U. B.; Crowley, W. F.,Jr.; Seminara, S. B.: Prevalence, phenotypic spectrum, and modes of inheritance of gonadotropin-releasing hormone receptor mutations in idiopathic hypogonadotropic hypogonadism. J. Clin. Endocr. Metab. 86:1580–1588, 2001.

[67920] 6250.Caron, P.; Chauvin, S.; Christin–Maitre, S.; Bennet, A.; Lahlou,N.; Counis, R.; Bouchard, P.; Kottler, M.–L.: Resistance of hypogonadicpatients with mutated GnRH receptor genes to pulsatile GnRH administration. J.Clin. En–

docr. Metab. 84: 990–996, 1999.

- [67921] 6251.Costa, E. M. F.; Bedecarrats, G. Y.; Mendonca, B. B.; Arnhold, I. J. P.; Kaiser, U. B.; Latronico, A. C.: Two novel mutations in the gonadotropin-releasing hormone receptor gene in Brazilian patients with hypogonadotropic hypogonadism and normal olfaction. J. Clin. Endocr. Metab. 86: 2680–2686, 2001.
- [67922] 6252.de Roux, N.; Young, J.; Brailly-Tabard, S.; Misrahi, M.; Milgrom, E.; Schaison, G.: The same molecular defects of the gonadotropin-releasing hormone receptor determine a variable degree of hypogonadism in affected kindred. J. Clin. Endocr. Metab. 84: 567–572, 1999.
- [67923] 6253.de Roux, N.; Young, J.; Misrahi, M.; Genet, R.; Chanson, P.; Schaison, G.; Milgrom, E.: A family with hypogonadotropic hypogonadism and mutations in the gonadotropin-releasing hormone receptor. New Eng. J. Med. 337: 1597–1602, 1997.
- [67924] 6254.Fan, N. C.; Jeung, E.-B.; Peng, C.; Olofsson, J. I.; Krisinger, J.; Leung, P. C. K.: The human gonadotropin-releasing hormone (GnRH) receptor gene: cloning, genomic organization and chromosomal assignment. Molec. Cell. Endocr. 103: R1–R6, 1994.
- [67925] 6255.Iwashita, T.; Murakami, H.; Asai, N.; Takahashi, M.:

Mechanism of Ret dysfunction by Hirschsprung mutations affecting its extracellular domain. *Hum. Molec. Genet.* 5: 1577–1580, 1996.

- [67926] 6256. Lehrman, M. A.; Goldstein, J. L.; Brown, M. S.; Russell, D. W.; Schneider, W. J.: Internalization-defective LDL receptors produced by genes with nonsense and frameshift mutations that truncate the cytoplasmic domain. *Cell* 41: 735–743, 1985.
- [67927] 6257. Bird, A. P.: CpG islands as gene markers in the vertebrate nucleus. *Trends Genet.* 3: 342–347, 1987.
- [67928] 6258. Maartmann-Moe, K.; Wang, H. S.; Donald, L. J.; Hamerton, J. L.; Berg, K.: Data from hybrid cell lines raise the possibility that factors controlling the low density lipoprotein receptor activity may reside on human chromosome 21, 5 or both. (Abstract) *Cytogenet. Cell Genet.* 32: 295–296, 1982.
- [67929] 6259. Miyake, Y.; Tajima, S.; Funahashi, T.; Yamamoto, A.: Analysis of a recycling-impaired mutant of low density lipoprotein receptor in familial hypercholesterolemia. *J. Biol. Chem.* 264: 16584–16590, 1989.
- [67930] 6260. Seftel, H. C.; Baker, S. G.; Sandler, M. P.; Forman, M. B.; Joffe, B. I.; Mendelsohn, D.; Jenkins, T.; Mieny, C. J.: A host of hypercholesterolaemic homozygotes in South



Africa. Brit. Med. J. 281: 633–636, 1980.

- [67931] 6261.Grundy, H. O.; Peltz, G.; Barsh, G.; Moore, K.; Golbus, M. S.;Lebo, R. V.: Immunoglobulin G Fc receptor II and Fc receptor IIIgenes map to chromosome 1 by spot-blot chromosome analysis. (Abstract) Am.J. Hum. Genet. 43: A145, 1988.
- [67932] 6262.Hibbs, M. L.; Bonadonna, L.; Scott, B. M.; McKenzie, I. F. C.;Hogarth, P. M.: Molecular cloning of a human immunoglobulin G Fcreceptor. Proc. Nat. Acad. Sci. 85: 2240–2244, 1988.
- [67933] 6263.Hibbs, M. L.; Hogarth, P. M.; McKenzie, I. F. C.: The mouse Ly–17locus identifies a polymorphism of the Fc receptor. Immunogenetics 22:335–348, 1985.
- [67934] 6264.Lebo, R. V.; Lynch, E. D.; Wiegant, J.; Moore, K.; Trounstein,M.; van der Ploeg, M.: Multicolor fluorescence in situ hybridizationand pulsed field electrophoresis dissect CMT1B gene region. Hum.Genet. 88: 13–20, 1991.
- [67935] 6265.Moser, K. L.; Neas, B. R.; Salmon, J. E.; Yu, H.; Gray-McGuire,C.; Asundi, N.; Bruner, G. R.; Fox, J.; Kelly, J.; Henshall, S.; Bacino,D.; Dietz, M.; Hogue, R.; Koelsch, G.; Nightingale, L.; Shaver, T.;Abdou, N. I.; Albert, D. A.; Carson, C.; Petri, M.; Treadwell, E.L.; James, J. A.; Harley, J. B.: Genome scan of human systemic lupuserythematosus: ev–

idence for linkage on chromosome 1q in African-American pedigrees. *Proc. Nat. Acad. Sci.* 95: 14869–14874, 1998.

- [67936] 6266. Oakey, R. J.; Watson, M. L.; Seldin, M. F.: Construction of a physical map on mouse and human chromosome 1: comparison of 13 Mb of mouse and 11 Mb of human DNA. *Hum. Molec. Genet.* 1: 613–620, 1992.
- [67937] 6267. Peltz, G. A.; Grundy, H. O.; Lebo, R. V.; Yssel, H.; Barsh, G. S.; Moore, K. W.: Human Fc-gamma-RIII: cloning, expression, and identification of the chromosomal locus of two Fc receptors for IgG. *Proc. Nat. Acad. Sci.* 86: 1013–1017, 1989.
- [67938] 6268. Qiu, W. Q.; de Bruin, D.; Brownstein, B. H.; Pearse, R.; Ravetch, J. V.: Organization of the human and mouse low-affinity Fc-gamma-R genes: duplication and recombination. *Science* 248: 732–735, 1990.
- [67939] 6269. Salmon, J. E.; Millard, S.; Schachter, L. A.; Arnett, F. C.; Ginzler, E. M.; Gourley, M. F.; Ramsey-Goldman, R.; Peterson, M. G. E.; Kimberly, R. P.: Fc-gamma-RIIA alleles are heritable risk factors for lupus nephritis in African Americans. *J. Clin. Invest.* 97: 1348–1354, 1996.
- [67940] 6270. Sammartino, L.; Webber, L. M.; Hogarth, P. M.; McKenzie, I. F. C.; Garson, O. M.: Assignment of the gene

coding for human FcRII(CD32) to bands q23q24 on chromosome 1. Immunogenetics 28: 380–381,1988.

[67941] 6271.Stein, M.-P.; Edberg, J. C.; Kimberly, R. P.; Mangan, E. K.; Bharadwaj,D.; Mold, C.; Du Clos, T. W.: C-reactive protein binding to Fc-gamma-RIIa on human monocytes and neutrophils is allele-specific. J. Clin. Invest.

105:369–376, 2000.

[67942] 6272.Unkeless, J. C.: Function and heterogeneity of human Fc receptors for immunoglobulin G. J. Clin. Invest. 83: 355–361, 1989.

[67943] 6273.Chang-Yeh, A.; Jabs, E. W.; Li, X.; Dracopoli, N. C.; Huang, R.C. C.: The IPP gene is assigned to human chromosome 1p32–1p22. Genomics 15:239–241, 1993.

[67944] 6274.Chang-Yeh, A.; Mold, D. E.; Brilliant, M. H.; Huang, R. C. C.:The mouse intracisternal A particle-promoted placental gene retrotransposition is mouse-strain-specific. Proc. Nat. Acad. Sci. 90: 292–296, 1993.

[67945] 6275.Chang-Yeh, A.; Mold, D. E.; Huang, R. C. C.: Identification of a novel murine IAP-promoted placenta-expressed gene. Nucleic Acids Res. 19: 3667–3672, 1991.

[67946] 6276.Mold, D. E.; Chang-Yeh, A.; Huang, R. C. C.: Cell lineage-specific expression of the MIPP gene. Biochem. Biophys. Res. Commun. 177:1062–1067, 1991.

- [67947] 6277.Cama, A.; de la Luz Sierra, M.; Quon, M. J.; Ottini, L.; Gorden,P.; Taylor, S. I.: Substitution of glutamic acid for alanine 1135in the putative 'catalytic loop' of the tyrosine kinase domain ofthe human insulin receptor: a mutation that impairs proteolytic processinginto subunits and inhibits receptor tyrosine kinase activity. J.Biol. Chem. 268: 8060–8069, 1993.
- [67948] 6278.Cama, A.; Quon, M. J.; de la Luz Sierra, M.; Taylor, S. I.: Substitutionof isoleucine for methionine at position 1153 in the beta–subunitof the human insulin receptor. J. Biol. Chem. 267: 8383–8389, 1992.
- [67949] 6279.Caro, J. F.; Raju, S. M.; Sinha, M. K.; Goldfine, I. D.; Dohm,G. L.: Heterogeneity of human liver, muscle, and adipose tissue insulinreceptor. Biochem. Biophys. Res. Commun. 151: 123–129, 1988.
- [67950] 6280.Christiansen, K.; Tranum–Jensen, J.; Carlsen, J.; Vinten, J.:A model for the quaternary structure of human placental insulin receptordeduced from electron microscopy. Proc. Nat. Acad. Sci. 88: 249–252,1991.
- [67951] 6281.Cocozza, S.; Porcellini, A.; Riccardi, G.; Monticelli, A.; Condorelli,G.; Ferrara, A.; Pianese, L.; Miele, C.; Capaldo, B.; Beguinot, F.;Varrone, S.: NIDDM associated with mutation in tyrosine kinase domainof insulin receptor

gene. Diabetes 41: 521–526, 1992.

[67952] 6282. Due, C.; Simonsen, M.; Olsson, L.: The major histocompatibility complex class I heavy chain as a structural subunit of the human cell membrane insulin receptor: implications for the range of biological functions of histocompatibility antigens. Proc. Nat. Acad. Sci.

83:6007–6011, 1986.

[67953] 6283. Ebina, Y.; Ellis, L.; Jarnagin, K.; Edery, M.; Graf, L.; Clauser, E.; Ou, J.-H.; Masiarz, F.; Kan, Y. W.; Goldfine, I. D.; Roth, R. A.; Rutter, W. J.: The human insulin receptor cDNA: the structural basis for hormone activated transmembrane signalling. Cell 40: 747–758, 1985.

[67954] 6284. Elbein, S. C.; Corsetti, L.; Ullrich, A.; Permutt, M. A.: Multiple restriction fragment length polymorphisms at the insulin receptor locus: a highly informative marker for linkage analysis. Proc. Nat. Acad. Sci. 83: 5223–5227, 1986.

[67955] 6285. Elbein, S. C.; Sorensen, L. K.; Schumacher, M. C.: Methionine for valine substitution in exon 17 of the insulin receptor gene in a pedigree with familial NIDDM. Diabetes 42: 429–434, 1993.

[67956] 6286. Ferrannini, E.; Muggeo, M.; Navalesi, R.; Pilo, A.: Impaired insulin degradation in a patient with insulin resis-

tance and acanthosisnigricans. Am. J. Med. 73: 148–154, 1982.

[67957] 6287.Grigorescu, F.; Flier, J. S.; Kahn, C. R.: Characterization of binding and phosphorylation defects of erythrocyte insulin receptors in the type A syndrome of insulin resistance. Diabetes 35: 127–138, 1986.

[67958] 6288.Grigorescu, F.; Flier, J. S.; Kahn, C. R.: Defect in insulin receptor phosphorylation in erythrocytes and fibroblasts associated with severe insulin resistance. J. Biol. Chem. 259: 15003–15006, 1984.

[67959] 6289.Grunberger, G.; Zick, Y.; Gordon, G.: Defect in phosphorylation of insulin receptors in cells from an insulin-resistant patient with normal insulin binding. Science 223: 932–934, 1984.

[67960] 6290.Hone, J.; Accili, D.; Al-Gazali, L. I.; Lestringant, G.; Orban, T.; Taylor, S. I.: Homozygosity for a new mutation (ile119-to-met) in the insulin receptor gene in five sibs with familial insulin resistance. J. Med. Genet. 31: 715–716, 1994.

[67961] 6291.Kadowaki, T.; Bevins, C. L.; Cama, A.; Ojamaa, K.; Marcus-Samuels, B.; Kadowaki, H.; Beitz, L.; McKeon, C.; Taylor, S. I.: Two mutant alleles of the insulin receptor gene in a patient with extreme insulin resistance. Science

240: 787–790, 1988.

- [67962] 6292.Kadowaki, T.; Kadowaki, H.; Accili, D.; Taylor, S. I.: Substitution of lysine for asparagine at position 15 in the alpha-subunit of the human insulin receptor: a mutation that impairs transport of receptors to the cell surface and decreases the affinity of insulin binding. *J.Biol. Chem.* 265: 19143–19150, 1990.
- [67963] 6293.Kadowaki, T.; Kadowaki, H.; Accili, D.; Yazaki, Y.; Taylor, S.I.: Substitution of arginine for histidine at position 209 in the alpha-subunit of the human insulin receptor: a mutation that impairs receptor dimerization and transport of receptors to the cell surface. *J.Biol. Chem.* 266: 21224–21231, 1991.
- [67964] 6294.Kadowaki, T.; Kadowaki, H.; Rechler, M. M.; Serrano-Rios, M.; Roth, J.; Gorden, P.; Taylor, S. I.: Five mutant alleles of the insulin receptor gene in patients with genetic forms of insulin resistance. *J.Clin. Invest.* 86: 254–264, 1990.
- [67965] 6295.Kadowaki, T.; Kadowaki, H.; Taylor, S. I.: A nonsense mutation causing decreased levels of insulin receptor mRNA: detection by a simplified technique for direct sequencing of genomic DNA amplified by the polymerase chain reaction. *Proc. Nat. Acad. Sci.* 87: 658–662, 1990.

- [67966] 6296.Kahn, C. R.; Flier, J. S.; Bar, R. S.; Archer, J. A.; Gordon, P.; Martin, M. M.; Roth, J.: The syndromes of insulin resistance and acanthosis nigricans: insulin receptor disorders in man. *New Eng. J. Med.* 294: 739–745, 1976.
- [67967] 6297.Kahn, C. R.; Goldstein, B. J.: Molecular defects in insulin action. *Science* 245:13 only, 1989.
- [67968] 6298.Kahn, C. R.; White, M. F.: The insulin receptor and the molecular mechanism of insulin action. *J. Clin. Invest.* 82: 1151–1156, 1988.
- [67969] 6299.Kakehi, T.; Hisatomi, A.; Kuzuya, H.; Yoshimasa, Y.; Okamoto, M.; Yamada, K.; Nishimura, H.; Kosaki, A.; Nawata, H.; Umeda, F.; Ibayashi, H.; Imura, H.: Defective processing of insulin–receptor precursor in cultured lymphocytes from a patient with extreme insulin resistance. *J. Clin. Invest.* 81: 2020–2022, 1988.
- [67970] 6300.Kittur, D.; Shimizu, Y.; DeMars, R.; Edidin, M.: Insulin binding to human B lymphoblasts is a function of HLA haplotype. *Proc. Nat. Acad. Sci.* 84: 1351–1355, 1987.
- [67971] 6301.Klinkhamer, M. P.; Groen, N. A.; van der Zon, G. C. M.; Lindhout, D.; Sandkuyl, L. A.; Krans, H. M. J.; Moller, W.; Maassen, J. A.: A leucine–to–proline mutation in the insulin receptor in a family with insulin resistance. *EMBO J.* 8: 2503–2507, 1989.



- [67972] 6302.Krook, A.; Brueton, L.; O'Rahilly, S.: Homozygous nonsense mutation in the insulin receptor gene in infant with leprechaunism. *Lancet* 342:277–278, 1993.
- [67973] 6303.Kulkarni, R. N.; Bruning, J. C.; Winnay, J. N.; Postic, C.; Magnuson, M. A.; Kahn, C. R.: Tissue-specific knockout of the insulin receptor in pancreatic beta cells creates an insulin secretory defect similar to that in type 2 diabetes. *Cell* 96: 329–339, 1999.
- [67974] 6304.Chan, A. M.–L.; Hilkens, J.; Kroezen, V.; Mitchell, P. J.; Scambler, P.; Wainwright, B. J.; Williamson, R.; Cooper, C. S.: Molecular cloning and localization to chromosome 6 of mouse INT1L1 gene. *Somat. Cell Molec. Genet.* 15: 555–562, 1989.
- [67975] 6305.Huguet, E. L.; McMahon, J. A.; McMahon, A. P.; Bicknell, R.; Harris, A. L.: Differential expression of human Wnt genes 2, 3, 4, and 7 in human breast cell lines and normal and disease states of human breast tissue. *Cancer Res.* 54: 2615–2621, 1994.
- [67976] 6306.McCoy, P. A.; Shao, Y.; Wolpert, C. M.; Donnelly, S. L.; Ashley-Koch, A.; Abel, H. L.; Ravan, S. A.; Abramson, R. K.; Wright, H. H.; DeLong, G. R.; Cuccaro, M. L.; Gilbert, J. R.; Pericak-Vance, M. A.: No association between the WNT2 gene and autistic disorder. *Am. J. Med. Genet.*

(Neuropsychiat.Genet.) 114: 106–109, 2002.

- [67977] 6307.Nusse, R.; Brown, A.; Papkoff, J.; Scambler, P.; Shackleford, G.;McMahon, A.; Moon, R.; Varmus, H.: A new nomenclature for int-1 andrelated genes: the Wnt gene family. Cell 64: 231–232, 1991.
- [67978] 6308.Wainwright, B. J.; Scambler, P. J.; Stanier, P.; Watson, E. K.;Bell, G.; Wicking, C.; Estivill, X.; Courtney, M.; Bour, A.; Pedersen,P. S.; Williamson, R.; Farrall, M.: Isolation of a human gene withprotein sequence similarity to human and murine int-1 and the Drosophilasegment polarity mutant wingless. EMBO J. 7: 1743–1748, 1988.
- [67979] 6309.Wassink, T. H.; Piven, J.; Vieland, V. J.; Huang, J.; Swiderski,R. E.; Pietila, J.; Braun, T.; Beck, G.; Folstein, S. E.; Haines,J. L.; Sheffield, V. C.: Evidence supporting WNT2 as an autism susceptibilitygene. Am. J. Med. Genet. 105: 406–413, 2001.
- [67980] 6310.Kluck, P. M. C.; Wiegant, J.; Jansen, R. P. M.; Bolk, M. W. J.;Raap, A. K.; Willemze, R.; Landegent, J. E.: The human interleukin-6receptor alpha-chain gene is localized on chromosome 1 band q21. Hum.Genet. 90: 542–544, 1993.
- [67981] 6311.Szpirer, J.; Szpirer, C.; Riviere, M.; Houart, C.; Bau-mann, M.;Fey, G. H.; Poli, V.; Cortese, R.; Islam, M. Q.;

Levan, G.: The interleukin-6-dependent DNA-binding protein gene (transcription factor 5: TCF5) maps to human chromosome 20 and rat chromosome 3, the IL6 receptor locus (IL6R) to human chromosome 1 and rat chromosome 2, and the rat IL6 gene to rat chromosome 4. *Genomics* 10: 539–546, 1991.

[67982] 6312. Yamasaki, K.; Taga, T.; Hirata, Y.; Yawata, H.; Kawanishi, Y.; Seed, B.; Taniguchi, T.; Hirano, T.; Kishimoto, T.: Cloning and expression of the human interleukin-6 (BSF-2/IFN-beta-2) receptor. *Science* 241:825–828, 1988.

[67983] 6313. Alli, C.; Consalez, G. G.: Linkage mapping of *Csrp* to proximal mouse chromosome 3. *Mammalian Genome* 9: 172 only, 1998.

[67984] 6314. Erdel, M.; Weiskirchen, R.: Assignment of CSRP1 encoding the LIM domain protein CRP1, to human chromosome 1q32 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 83: 10–11, 1998.

[67985] 6315. Liebhaber, S. A.; Emery, J. G.; Urbanek, M.; Wang, X.; Cooke, N.E.: Characterization of a human cDNA encoding a widely expressed and highly conserved cysteine-rich protein with an unusual zinc-finger motif. *Nucleic Acids Res.* 18: 3871–3879, 1990.

- [67986] 6316. Weiskirchen, R.; Pino, J. D.; Macalima, T.; Bister, K.; Beckerle, M. C.: The cysteine-rich protein family of highly related LIM domain proteins. *J. Biol. Chem.* 270: 28946–28954, 1995.
- [67987] 6317. Yamauchi, M.; Yamauchi, N.; Phear, G.; Spurr, N. K.; Martinsson, T.; Weith, A.; Meuth, M.: Genomic organization and chromosomal localization of the human CTP synthetase gene (CTPS). *Genomics* 11: 1088–1096, 1991.
- [67988] 6318. Wang, X.; Lee, G.; Liebhaber, S. A.; Cooke, N. E.: Human cysteine-rich protein: a member of the LIM/double-finger family displaying coordinate serum induction with c-myc. *J. Biol. Chem.* 267: 9176–9184, 1992.
- [67989] 6319. Wang, X.; Ray, K.; Szpirer, J.; Levan, G.; Liebhaber, S. A.; Cooke, N. E.: Analysis of the human cysteine-rich protein gene (CSRP), assignment to chromosome 1q24–1q32, and identification of an associated MspI polymorphism. *Genomics* 14: 391–397, 1992.
- [67990] 6320. Courseaux, A.; Grosgeorge, J.; Gaudray, P.; Pannett, A. A. J.; Forbes, S. A.; Williamson, C.; Bassett, D.; Thakker, R. V.; Teh, B. T.; Farnebo, F.; Shepherd, J.; Skogseid, B.; Larsson, C.; Giraud, S.; Zhang, C. X.; Salandre, J.; Calender, A.: Definition of the minimal MEN1 candidate area based on a 5-Mb integrated map of proximal 11q13. *Genomics*

37:354–365, 1996.

- [67991] 6321. Bullock, S.; Hayward, C.; Manson, J.; Brock, D. J. H.; Raeburn, J. A.: Quantitative immunoassays for diagnosis and carrier detection in cystic fibrosis. *Clin. Genet.* 21: 336–341, 1982.
- [67992] 6322. Dorin, J. R.; Emslie, E.; van Heyningen, V.: Related calcium-binding proteins map to the same subregion of chromosome 1q and to an extended region of synteny on mouse chromosome 3. *Genomics* 8: 420–426, 1990.
- [67993] 6323. Dorin, J. R.; Novak, M.; Hill, R. E.; Brock, D. J. H.; Secher, D. S.; van Heyningen, V.: A clue to the basic defect in cystic fibrosis from cloning the CF antigen gene. *Nature* 326: 614–617, 1987.
- [67994] 6324. Frizzell, R. A.; Rechkemmer, G.; Shoemaker, R. L.: Altered regulation of airway epithelial cell chloride channels in cystic fibrosis. *Science* 233:558–560, 1986.
- [67995] 6325. Manson, J. C.; Brock, D. J. H.: Development of a quantitative immunoassay for the cystic fibrosis gene. *Lancet* I: 330–331, 1980.
- [67996] 6326. Herzig, K. H.; Schon, I.; Tatemoto, K.; Ohe, Y.; Li, Y.; Folsch, U. R.; Owyang, C.: Diazepam binding inhibitor is a potent cholecystokinin-releasing peptide in the intestine. *Proc. Nat. Acad. Sci.* 93: 7927–7932, 1996.

- [67997] 6327.Li, Y.; Hao, Y.; Owyang, C.: Diazepam-binding inhibitor mediates feedback regulation of pancreatic secretion and postprandial release of cholecystokinin. *J. Clin. Invest.* 105: 351–359, 2000.
- [67998] 6328.Rose, T. M.; Schultz, E. R.; Todaro, G. J.: Molecular cloning of the gene for the yeast homolog (ACB) of diazepam binding inhibitor/endozepine/acyl-CoA-binding protein. *Proc. Nat. Acad. Sci.* 89: 11287–11291, 1992.
- [67999] 6329.Spannagel, A. W.; Green, G. M.; Guan, D.; Liddle, R. A.; Faull, K.; Reeve, J. R.: Purification and characterization of a luminal cholecystokinin-releasing factor from rat intestinal secretion. *Proc. Nat. Acad. Sci.* 93: 4415–4420, 1996.
- [68000] 6330.Todd, S.; Naylor, S. L.: New chromosomal mapping assignments for argininosuccinate synthetase pseudogene 1, interferon-beta-3 gene, and the diazepam binding inhibitor gene. *Somat. Cell Molec. Genet.* 18:381–385, 1992.
- [68001] 6331.Webb, N. R.; Rose, T. M.; Malik, N.; Marquardt, H.; Shoyab, M.; Todaro, G. J.; Lee, D. C.: Bovine and human cDNA sequences encoding a putative benzodiazepine receptor ligand. *DNA* 6: 71–79, 1987.

- [68002] 6332. Anagnou, N. P.; Antonarakis, S. E.; O'Brien, S. J.; Modi, W. S.; Nienhuis, A. W.: Chromosomal localization and racial distribution of the polymorphic human dihydrofolate reductase pseudogene (DHFRPI). *Am. J. Hum. Genet.* 42: 345–352, 1988.
- [68003] 6333. Anagnou, N. P.; Antonarakis, S. E.; O'Brien, S. J.; Nienhuis, A. W.: Chromosomal localization and racial distribution of the polymorphic hDHFR-psi-1 pseudogene. (Abstract) *Clin. Res.* 33: 328A only, 1985.
- [68004] 6334. Anagnou, N. P.; Antonarakis, S. E.; O'Brien, S. J.; Nienhuis, A. W.: A novel form of human polymorphism involving the hDHFR-psi-1 pseudogene identifies three RFLPs. *Nucleic Acids Res.* 15: 5501 only, 1987.
- [68005] 6335. Anagnou, N. P.; O'Brien, S. J.; Shimada, T.; Nash, W. G.; Chen, M.-J.; Nienhuis, A. W.: Chromosomal organization of the human dihydrofolate reductase genes: dispersion, selective amplification and a novel form of polymorphism. *Proc. Nat. Acad. Sci.* 81: 5170–5174, 1984.
- [68006] 6336. Blakley, R. L.; Sorrentino, B. P.: In vitro mutations in dihydrofolate reductase that confer resistance to methotrexate: potential for clinical application. *Hum. Mutat.* 11: 259–263, 1998.
- [68007] 6337. Chen, M.-J.; Shimada, T.; Moulton, A. D.; Harrison,

M.; Nienhuis, A. W.: Intronless human dihydrofolate reductase genes are derived from processed RNA molecules.

Proc. Nat. Acad. Sci. 79: 7435–7439, 1982.

[68008] 6338. Craik, C. S.; Rutter, W. J.; Fletterick, R.: Splice junctions: association with variation in protein structure. Science 220: 1125–1129, 1983.

[68009] 6339. Erbe, R. W.: Inborn errors of folate metabolism. New Eng. J. Med. 293: 753–757 and 807–812, 1975.

[68010] 6340. Funanage, V. L.; Myoda, T. T.; Moses, P. A.; Cowell, H. R.: Assignment of the human dihydrofolate reductase gene to the q11–q22 region of chromosome 5. Molec. Cell. Biol. 4: 2010–2016, 1984.

[68011] 6341. Hoffbrand, A. V.; Tripp, E.; Jackson, B. F. A.; Luck, W. E.; Frater-Schroder, M.: Hereditary abnormal transcobalamin II previously diagnosed as congenital dihydrofolate reductase deficiency. (Letter) New Eng. J. Med. 310: 789–790, 1984.

[68012] 6342. Killary, A. M.; Leach, R. J.; Moran, R. G.; Fournier, R. E. K.: Assignment of genes encoding dihydrofolate reductase and hexosaminidase B to Mus musculus chromosome 13. Somat. Cell Molec. Genet. 12: 641–648, 1986.

[68013] 6343. Maurer, B.; Barker, P. E.; Masters, J. N.; D'Eustachio, P.; Ruddle, F. H.; Attardi, G.: Chromosomal location of the



normal human DHFR gene and of its amplified copies in methotrexate resistant cell variants. (Abstract) Cytogenet. Cell Genet. 37: 534 only, 1984.

- [68014] 6344. Maurer, B. J.; Barker, P. E.; Masters, J. N.; Ruddle, F. H.; Attardi, G.: Human dihydrofolate reductase gene is located in chromosome 5 and is unlinked to the related pseudogenes. Proc. Nat. Acad. Sci. 81:1484–1488, 1984.
- [68015] 6345. Maurer, B. J.; Carlock, L.; Wasmuth, J.; Attardi, G.: Assignment of human dihydrofolate reductase gene to band q23 of chromosome 5 and of related pseudogene psiHD1 to chromosome 3. Somat. Cell Molec. Genet. 11: 79–85, 1985.
- [68016] 6346. Myoda, T. T.; Funanage, V. L.: Personal Communication. Wilmington, Del. 10/7/1983.
- [68017] 6347. Singer, M. J.; Mesner, L. D.; Friedman, C. L.; Trask, B. J.; Hamlin, J. L.: Amplification of the human dihydrofolate reductase gene via double minutes is initiated by chromosome breaks. Proc. Nat. Acad. Sci. 97: 7921–7926, 2000.
- [68018] 6348. Tauro, G. P.; Danks, D. M.; Rowe, P. B.; Van der Weyden, M. B.; Schwarz, M. A.; Collins, V. L.; Neal, B. W.: Dihydrofolate reductase deficiency causing megaloblastic anemia in two families. New Eng. J. Med. 294: 466–470, 1976.
- [68019] 6349. Walters, T. R.: Congenital megaloblastic anemia re-

sponsive to N(5)-formyltetrahydrofolic acid administration.

J. Pediatr. 70: 686–687, 1967.

[68020] 6350. Ali, G.; Wasco, W.; Cai, X.; Szabo, P.; Sheu, K.-F. R.; Cooper, A. J. L.; Gaston, S. M.; Gusella, J. F.; Tanzi, R. E.; Blass, J. P.: Isolation, characterization, and mapping of gene encoding dihydrolipoylsuccinyltransferase (E2k) of human alpha-ketoglutarate dehydrogenase complex. Somat. Cell Molec. Genet. 20: 99–105, 1994.

[68021] 6351. Nakano, K.; Matuda, S.; Sakamoto, T.; Takase, C.; Nakagawa, S.; Ohta, S.; Ariyama, T.; Inazawa, J.; Abe, T.; Miyata, T.: Human dihydrolipoamidesuccinyltransferase: cDNA cloning and localization on chromosome 14q24.2–q24.3. Biochim. Biophys. Acta 1216: 360–368, 1993.

[68022] 6352. Nakano, K.; Takase, C.; Sakamoto, T.; Ohta, S.; Nakagawa, S.; Ariyama, T.; Inazawa, J.; Abe, T.; Matuda, S.: An unspliced cDNA for human dihydrolipoamide succinyltransferase: characterization and mapping of the gene to chromosome 14q24.2–q24.3. Biochem. Biophys. Res. Commun. 196: 527–533, 1993.

[68023] 6353. Patel, M. S.; Harris, R. A.: Mammalian alpha-keto acid dehydrogenase complexes: gene regulation and genetic defects. FASEB J. 9: 1164–1172, 1995.

- [68024] 6354.McKusick, V. A.: Heritable Disorders of Connective Tissue. St. Louis: C. V. Mosby (pub.) (4th ed.): 1972.
- [68025] 6355.Palmer, G.; Manen, D.; Bonjour, J.-P.; Caverzasio, J.: Characterization of the human Glvr-1 phosphate transporter/retrovirus receptor gene and promoter region. *Gene* 226: 25-33, 1999.
- [68026] 6356.Fan, N. C.; Peng, C.; Krisinger, J.; Leung, P. C. K.: The human gonadotropin-releasing hormone receptor gene: complete structure including multiple promoters, transcription initiation sites, and polyadenylation signals. *Molec. Cell. Endocr.* 107: R1-R8, 1995.
- [68027] 6357.Boghosian-Sell, L.; Comings, D. E.; Overhauser, J.: Tourette syndrome in a pedigree with a 7;18 translocation: identification of a YAC spanning the translocation breakpoint at 18q22.3. *Am. J. Hum. Genet.* 59:999-1005, 1996.
- [68028] 6358.Grosse, R.; Schoneberg, T.; Schultz, G.; Gudermann, T.: Inhibition of gonadotropin-releasing hormone receptor signaling by expression of a splice variant of the human receptor. *Molec. Endocr.* 11: 1305-1318, 1997.
- [68029] 6359.Kaiser, U. B.; Dushkin, H.; Altherr, M. R.; Beier, D. R.; Chin, W. W.: Chromosomal localization of the gonadotropin-releasing hormone receptor gene to human chromosome 4q13.1-q21.1 and mouse chromosome 5.

Genomics 20: 506–508, 1994.

- [68030] 6360.Kakar, S. S.; Musgrove, L. C.; Devor, D. C.; Sellers, J. C.; Neill, J. D.: Cloning, sequencing, and expression of human gonadotropin-releasing hormone (GnRH) receptor. *Biochem. Biophys. Res. Commun.* 189:289–295, 1992.
- [68031] 6361.Kakar, S. S.; Neill, J. D.: The human gonadotropin-releasing hormone receptor gene (GNRHR) maps to chromosome band 4q13. *Cytogenet. Cell Genet.* 70: 211–214, 1995.
- [68032] 6362.Kottler, M.–L.; Chauvin, S.; Lahlou, N.; Harris, C. E.; Johnston, C. J.; Lagarde, J.–P.; Bouchard, P.; Farid, N. R.; Counis, R.: A new compound heterozygous mutation of the gonadotropin-releasing hormone receptor (L314X, Q106R) in a woman with complete hypogonadotropic hypogonadism: chronic estrogen administration amplifies the gonadotropin defect. *J. Clin. Endocr. Metab.* 85: 3002–3008, 2000.
- [68033] 6363.Kottler, M.–L.; Counis, R.; Bouchard, P.: Mutations of the GnRH receptor gene: a new cause of autosomal-recessive hypogonadotropic hypogonadism. *Arch. Med. Res.* 30: 481–485, 1999.
- [68034] 6364.Kottler, M. L.; Lorenzo, F.; Bergametti, F.; Comeron, P.; Souchier, C.; Counis, R.: Subregional mapping of

the human gonadotropin-releasing hormone receptor (GnRH-R) gene to 4q between the markers D4S392 and D4S409. Hum. Genet. 96: 477–480, 1995.

- [68035] 6365. Layman, L. C.; Cohen, D. P.; Jin, M.; Xie, J.; Li, Z.; Reindollar, R. H.; Bolbolan, S.; Bick, D. P.; Sherins, R. R.; Duck, L. W.; Musgrove, L. C.; Sellers, J. C.; Neill, J. D.: Mutations in gonadotropin-releasing hormone receptor gene cause hypogonadotropic hypogonadism. (Letter) Nature-Genet. 18: 14–15, 1998.
- [68036] 6366. Leung, P. C. K.; Squire, J.; Peng, C.; Fan, N.; Hayden, M. R.; Olofsson, J. I.: Mapping of the gonadotropin-releasing hormone (GnRH) receptor gene to human chromosome 4q21.2 by fluorescence in situ hybridization. Mammalian Genome 6: 309–310, 1995.
- [68037] 6367. Mason, A. J.; Hayflick, J. S.; Zoeller, R. T.; Young, W. S., III; Phillips, H. S.; Nikolics, K.; Seeburg, P. H.: A deletion truncating the gonadotropin-releasing hormone gene is responsible for hypogonadism in the 'hpg' mouse. Science 234: 1366–1371, 1986.
- [68038] 6368. Morrison, N.; Sellar, R. E.; Boyd, E.; Eidne, K. A.; Connor, J. M.: Assignment of the gene encoding the human gonadotropin-releasing hormone receptor to 4q13.2–13.3 by fluorescence in situ hybridization. Hum. Genet. 93:

714–715, 1994.

- [68039] 6369. Pitteloud, N.; Boepple, P. A.; DeCruz, S.; Valkenburg, S. B.; Crowley, W. F., Jr.; Hayes, F. J.: The fertile eunuch variant of idiopathic hypogonadotropic hypogonadism: spontaneous reversal associated with a homozygous mutation in the gonadotropin-releasing hormone receptor. *J. Clin. Endocr. Metab.* 86: 2470–2475, 2001.
- [68040] 6370. Pralong, F. P.; Gomez, F.; Castillo, E.; Cotechia, S.; Abuin, L.; Aubert, M. L.; Portmann, L.; Gaillard, R. C.: Complete hypogonadotropic hypogonadism associated with a novel inactivating mutation of the gonadotropin-releasing hormone receptor. *J. Clin. Endocr. Metab.* 84: 3811–3816, 1999.
- [68041] 6371. Szende, B.; Srkalovic, G.; Timar, J.; Mulchahey, J. J.; Neill, J. D.; Lapis, K.; Csikos, A.; Szepeshazi, K.; Schally, A. V.: Localization of receptors for luteinizing hormone-releasing hormone in pancreatic and mammary cancer cells. *Proc. Nat. Acad. Sci.* 88: 4153–4156, 1991.
- [68042] 6372. Smith, M.; Herrell, S.; Lusher, M.; Lako, L.; Simpson, C.; Wiestner, A.; Skoda, R.; Ireland, M.; Strachan, T.: Genomic organisation of the human chordin gene and mutation screening of candidate Corneli de Lange syndrome genes. *Hum. Genet.* 105: 104–111, 1999.

- [68043] 6373.Spemann, H.; Mangold, H.: Ueber induktion von embryonalanlagen durch implantation artfremder organ- isatoren. Arch. Mikroskopische Anat. Entwick- lungsmechanik 100: 599–638, 1924.
- [68044] 6374.Zech, L.; Haglund, U.; Nilsson, K.; Klein, G.: Characteristic chromosomal abnormalities in biopsies and lymphoid-cell lines from patients with Burkitt and non-Burkitt lymphomas. Int. J. Cancer 17:47–56, 1976.
- [68045] 6375.Bahler, M.; Kehrer, I.; Gordon, L.; Stoffler, H.–E.; Olsen, A.S.: Physical mapping of human myosin–IXB (MYO9B), the human orthologue of the rat myosin myr 5, to chromosome 19p13.1. Genomics 43: 107–109, 1997.
- [68046] 6376.Miyajima, N.; Kadowaki, Y.; Fukushige, S.; Shimizu, S.; Semba, K.; Yamanashi, Y.; Matsubara, K.; Toyoshima, K.; Yamamoto, T.: Identification of two novel members of erbA superfamily by molecular cloning: the gene products of the two are highly related to each other. Nucleic Acids Res. 16: 11057–11073, 1988.
- [68047] 6377.Qiu, Y.; Krishnan, V.; Zeng, Z.; Gilbert, D. J.; Copeland, N. G.; Gibson, L.; Yang–Feng, T.; Jenkins, N. A.; Tsai, M.–J.; Tsai, S. Y.: Isolation, characterization, and chromosomal localization of mouse and human COUP–TF I and II genes. Genomics 29: 240–246, 1995.

- [68048] 6378. MacDonald, N. J.; Freije, J. M. P.; Stracke, M. L.; Manrow, R.E.; Steeg, P. S.: Site-directed mutagenesis of nm23-H1: mutation of proline 96 or serine 120 abrogates its motility inhibitory activity upon transfection into human breast carcinoma cells. *J. Biol. Chem.* 271:25107–25116, 1996.
- [68049] 6379. Okabe-Kado, J.; Kasukabe, T.; Honma, Y.; Hayashi, M.; Henzel, W.J.; Hozumi, M.: Identity of a differentiation inhibiting factor from mouse myeloid leukemia cells with NM23/nucleoside diphosphate kinase. *Biochem. Biophys. Res. Commun.* 182: 987–994, 1992.
- [68050] 6380. Postel, E. H.: Modulation of c-myc transcription by triple helix formation. *Ann. N.Y. Acad. Sci.* 660: 57–63, 1992.
- [68051] 6381. Postel, E. H.; Berberich, S. J.; Flint, S. J.; Ferrone, C. A.: Human c-myc transcription factor PuF identified as nm23-H2 nucleoside diphosphate kinase, a candidate suppressor of tumor metastasis. *Science* 261:478–480, 1993.
- [68052] 6382. Petek, E.; Windpassinger, C.; Vincent, J. B.; Cheung, J.; Boright, A. P.; Scherer, S. W.; Kroisel, P. M.; Wagner, K.: Disruption of a novel gene (IMMP2L) by a breakpoint in 7q31 associated with Tourette syndrome. *Am. J. Hum.*



Genet. 848–858, 2001.

- [68053] 6383.Kugler, S.; Plenz, G.; Muller, P. K.: Two additional 5–prime exons in the human vigilin gene distinguish it from the chicken gene and provide the structural basis for differential routes of gene expression. *Europ.J. Biochem.* 238: 410–417, 1996.
- [68054] 6384.LeBoeuf, R. C.; Xia, Y.–R.; Oram, J. F.; Lusis, A. J.: Mapping of the gene for high–density lipoprotein binding protein (Hdlbp) to proximal mouse chromosome 1. *Genomics* 23: 296–298, 1994.
- [68055] 6385.Plenz, G.; Kugler, S.; Schnittger, S.; Rieder, H.; Fonatsch, C.; Muller, P. K.: The human vigilin gene: identification, chromosomal localization and expression pattern. *Hum. Genet.* 93: 575–582, 1994.
- [68056] 6386.Xia, Y.–R.; Klisak, I.; Sparkes, R. S.; Oram, J.; Lusis, A. J.: Localization of the gene for high–density lipoprotein binding protein (HDLBP) to human chromosome 2q37. *Genomics* 16: 524–525, 1993.
- [68057] 6387.Kaelbling, M.; Eddy, R.; Shows, T. B.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Klinger, H. P.; O'Hara, B.: Localization of the human gene allowing infection by gibbon ape leukemia virus to human chromosome region 2q11–q14 and to the homologous region on mouse chro–

mosome 2. J. Virol. 65: 1743–1747, 1991.

- [68058] 6388.Kavanaugh, M. P.; Miller, D. G.; Zhang, W.; Law, W.; Kozak, S.L.; Kabat, D.; Miller, A. D.: Cell–surface receptors for gibbon apeleukemia virus and amphotropic murine retrovirus are inducible sodium–dependentphosphate symporters. Proc. Nat. Acad. Sci. 91: 7071–7075, 1994.
- [68059] 6389.O'Hara, B.; Johann, S. V.; Klinger, H. P.; Blair, D. G.; Robinson,H.; Dunn, K. J.; Sass, P.; Vitek, S. M.; Robins, T.: Characterizationof a human gene conferring sensitivity to infection by gibbon apeleukemia virus. Cell Growth Differ. 1: 119–127, 1990.
- [68060] 6390.Palmer, G.; Manen, D.; Bonjour, J.–P.; Caverzasio, J.: Species–specificmechanisms control the activity of the Pit1/PIT1 phosphate transporter gene promoter in mouse and human. Gene 279: 49–62, 2001.
- [68061] 6391.Hayashi, K.; Yano, H.; Hashida, T.; Takeuchi, R.; Takeda, O.; Asada,K.; Takahashi, E.; Kato, I.; Sobue, K.: Genomic structure of thehuman caldesmon gene. Proc. Nat. Acad. Sci. 89: 12122–12126, 1992.
- [68062] 6392.Humphrey, M. B.; Herrera–Sosa, H.; Gonzalez, G.; Lee, R.; Bryan,J.: Cloning of cDNAs encoding human caldesmons. Gene 112: 197–204,1992.
- [68063] 6393.Dickson, K. M.; Bergeron, J. J. M.; Shames, I.; Colby,

J.; Nguyen,D. T.; Chevet, E.; Thomas, D. Y.; Snipes, G. J.: Association of calnexinwith mutant peripheral myelin protein-22 ex vivo: a basis for 'gain-of-function'ER diseases. Proc. Nat. Acad. Sci. 99: 9852–9857, 2002.

[68064] 6394.Gray, P. W.; Byers, M. G.; Eddy, R. L.; Shows, T. B.: The assignmentof the calnexin gene to the q35 region of chromosome 5. (Abstract) HumanGenome Mapping Workshop 93 9 only, 1993.

[68065] 6395.Schrag, J. D.; Bergeron, J. J. M.; Li, Y.; Borisova, S.; Hahn,M.; Thomas, D. Y.; Cygler, M.: The structure of calnexin, an ER chaperoneinvolved in quality control of protein folding. Molec. Cell 8: 633–644,2001.

[68066] 6396.Tjoelker, L. W.; Seyfried, C. E.; Eddy, R. L., Jr.; Byers, M. G.;Shows, T. B.; Calderon, J.; Schreiber, R. B.; Gray, P. W.: Human,mouse, and rat calnexin cDNA cloning: identification of potentialcalcium binding motifs and gene localization to human chromosome 5. Biochemistry 33:3229–3236, 1994.

[68067] 6397.Richard, I.; Beckmann, J. S.: Molecular cloning of mouse canp3,the gene associated with limb-girdle muscular dystrophy 2A in human. MammalianGenome 7: 377–379, 1996.

[68068] 6398.Bataille, N.; Schmitt, N.; Aumercier–Maes, P.; Ollivier,

B.; Lucas-Heron, B.; Lestienne, P.: Molecular cloning of human calmitine, a mitochondrial calcium binding protein, reveals identity with calsequestrine. *Biochem. Biophys. Res. Commun.* 203: 1477–1482, 1994.

[68069] 6399. Fujii, J.; Willard, H. F.; MacLennan, D. H.: Characterization and localization to human chromosome 1 of human fast-twitch skeletal muscle calsequestrin gene. *Somat. Cell Molec. Genet.* 16: 185–189, 1990.

[68070] 6400. Chen, C.-S.; Bejeck, B. E.; Kersey, J. H.: A mapping study of 13 genes on human chromosome bands 4q11–q25. *Cytogenet. Cell Genet.* 69:260–265, 1995.

[68071] 6401. Fujiwara, Y.; Miwa, M.; Nogami, M.; Okumura, K.; Nobori, T.; Suzuki, T.; Ueda, M.: Genomic organization and chromosomal localization of the human casein gene family. *Hum. Genet.* 99: 368–373, 1997.

[68072] 6402. Rijnkels, M.; Meershoek, E.; de Boer, H. A.; Pieper, F. R.: Physical map and localization of the human casein gene locus. *Mammalian Genome* 8:285–286, 1997.

[68073] 6403. Voglino, G. F.; Ponzzone, A.: Polymorphism in human casein. *Nature N.B.* 238: 149 only, 1972.

[68074] 6404. Ferretti, L.; Leone, P.; Sgaramella, V.: Long range restriction analysis of the bovine casein genes. *Nucleic Acids Res.* 18: 6829–6833, 1990.

- [68075] 6405.Threadgill, D. W.; Womack, J. E.: Genomic analysis of the majorbovine milk protein genes. *Nucleic Acids Res.* 18: 6935–6942, 1990.
- [68076] 6406.Kurotaki, N.; Imaizumi, K.; Harada, N.; Masuno, M.; Kondoh, T.;Nagai, T.; Ohashi, H.; Naritomi, K.; Tsukahara, M.; Makita, Y.; Sugimoto,T.; Sonoda, T.; and 11 others: Haploinsufficiency of NSD1 causesSotos syndrome. *Nature Genet.* 30: 365–366, 2002.
- [68077] 6407.Whang–Peng, J.; Triche, T. J.; Knutsen, T.; Miser, J.; Kao–Shan,S.; Tsai, S.; Israel, M. A.: Cytogenetic characteri- zation of selectedsmall round cell tumors of childhood. *Cancer Genet. Cytogenet.* 21:185–208, 1986.
- [68078] 6408.Mount, L. A.; Reback, S.: Familial paroxysmal choreoathetosis:preliminary report on a hitherto unde- scribed clinical syndrome. *Arch.Neurol. Psychiat.* 44: 841–847, 1940.
- [68079] 6409.Lee, S.; Lin, M.; Mele, A.; Cao, Y.; Farmar, J.; Russo, D.; Redman,C.: Proteolytic processing of big endothelin–3 by the Kell bloodgroup protein. *Blood* 94: 1440–1450, 1999.
- [68080] 6410.Cyr, C.; Huebner, K.; Druck, T.; Kris, R.: Cloning and chromosomallocalization of a human endothelin ETA re- ceptor. *Biochem. Biophys.Res. Commun.* 181: 184–190,

1991.

- [68081] 6411.Hosoda, K.; Nakao, K.; Tamura, N.; Arai, H.; Ogawa, Y.; Suga, S.;Nakanishi, S.; Imura, H.: Organization, structure, chromosomal assignment,and expression of the gene encoding the human endothelin-A receptor. J.Biol. Chem. 267: 18797–18804, 1992.
- [68082] 6412.Tzourio, C.; El Amrani, M.; Poirier, O.; Nicaud, V.; Bousser, M.–G.;Alperovitch, A.: Association between migraine and endothelin typeA receptor (ETA –231 A/G) gene polymorphism. Neurology 56: 1273–1277,2001.
- [68083] 6413.Amiel, J.; Attie, T.; Jan, D.; Pelet, A.; Edery, P.; Bidaud, C.;Lacombe, D.; Tam, P.; Simeoni, J.; Flori, E.; Nihoul-Fekete, C.; Munnich,A.; Lyonnet, S.: Heterozygous endothelin receptor B (EDNRB) mutationsin isolated Hirschsprung disease. Hum. Molec. Genet. 5: 355–357,1996.
- [68084] 6414.Arai, H.; Nakao, K.; Takaya, K.; Hosoda, K.; Ogawa, Y.; Nakanishi,S.; Imura, H.: The human endothelin-B receptor gene: structural organizationand chromosomal assignment. J. Biol. Chem. 268: 3463–3470, 1993.
- [68085] 6415.Attie, T.; Till, M.; Pelet, A.; Amiel, J.; Edery, P.; Boutrand,L.; Munnich, A.; Lyonnet, S.: Mutation of the endothelin-receptorB gene in Waardenburg–Hirschsprung

disease. Hum. Molec. Genet. 4:2407–2409, 1995.

- [68086] 6416.Auricchio, A.; Griseri, P.; Carpentieri, M. L.; Betsos, N.; Staiano,A.; Tozzi, A.; Priolo, M.; Thompson, H.; Boccia-  
rdi, R.; Romeo, G.;Ballabio, A.; Ceccherini, I.: Double het-  
erozygosity for a RET substitutioninferring with splicing  
and an EDNRB missense mutation in Hirschsprungdisease.  
(Letter) Am. J. Hum. Genet. 64: 1216–1221, 1999.
- [68087] 6417.Carrasquillo, M. M.; McCallion, A. S.; Puffenberger,  
E. G.; Kashuk,C. S.; Nouri, N.; Chakravarti, A.: Genome-  
wide association studyand mouse model identify interac-  
tion between RET and EDNRB pathwaysin Hirschsprung  
disease. Nature Genet. 32: 237–244, 2002.
- [68088] 6418.Ceccherini, I.; Zhang, A. L.; Matera, I.; Yang, G.; De-  
voto, M.;Romeo, G.; Cass, D. T.: Interstitial deletion of the  
endothelin–Breceptor gene in the spotting lethal (sl) rat.  
Hum. Molec. Genet. 4:2089–2096, 1995.
- [68089] 6419.Chakravarti, A.: Endothelin receptor–mediated sig-  
naling in Hirschsprungdisease. Hum. Molec. Genet. 5:  
303–307, 1996.
- [68090] 6420.Elshourbagy, N. A.; Adamou, J. E.; Gagnon. A. W.;  
Wu, H.–L.; Pullen,M.; Nambi, P.: Molecular characterization  
of a novel human endothelinreceptor splice variant. J. Biol.  
Chem. 271: 25300–25307, 1996.

- [68091] 6421. Gariepy, C. E.; Cass, D. T.; Yanagisawa, M.: Null mutation of endothelin receptor type B gene in spotting lethal rats causes aganglionic megacolon and white coat color. *Proc. Nat. Acad. Sci.* 93: 867–872, 1996.
- [68092] 6422. Gariepy, C. E.; Ohuchi, T.; Williams, S. C.; Richardson, J. A.; Yanagisawa, M.: Salt-sensitive hypertension in endothelin-B receptor-deficient rats. *J. Clin. Invest.* 105: 925–933, 2000.
- [68093] 6423. Gariepy, C. E.; Williams, S. C.; Richardson, J. A.; Hammer, R. E.; Yanagisawa, M.: Transgenic expression of the endothelin-B receptor prevents congenital intestinal aganglionosis in a rat model of Hirschsprung disease. *J. Clin. Invest.* 102: 1092–1101, 1998.
- [68094] 6424. Gross, A.; Kunze, J.; Maier, R. F.; Stoltenberg-Didinger, G.; Grimmer, I.; Obladen, M.: Autosomal-recessive neural crest syndrome with albinism, black lock, cell migration disorder of the neurocytes of the gut, and deafness: ABCD syndrome. *Am. J. Med. Genet.* 56: 322–326, 1995.
- [68095] 6425. Hofstra, R. M. W.; Osinga, J.; Buys, C. H. C. M.: Mutations in Hirschsprung disease: when does a mutation contribute to the phenotype? *Europ. J. Hum. Genet.* 5: 180–185, 1997.



- [68096] 6426.Hosoda, K.; Hammer, R. E.; Richardson, J. A.; Baynash, A. G.;Cheung, J. C.; Giaid, A.; Yanagisawa, M.: Targeted and natural (piebald-lethal)mutations of endothelin-B receptor gene produce megacolon associatedwith spotted coat color in mice. *Cell* 79: 1267-1276, 1994.
- [68097] 6427.Charlton, P.; Guida, L.; Copeland, N.; Jenkins, N.; Munroe, D.;Greenberg, F.; Fiedorek, F. T.; Nicholls, R. D.: Genetic approachto function of the neuropeptide galanin. (Abstract) *Am. J. Hum. Genet.* 53(suppl.): A1137, 1993.
- [68098] 6428.Evans, H.; Baumgartner, M.; Shine, J.; Herzog, H.: Genomic organizationand localization of the gene encoding human preprogalanin. *Genomics* 18:473-477, 1993.
- [68099] 6429.Evans, H. F.; Shine, J.: Human galanin: molecular cloning reveals a unique structure. *Endocrinology* 129: 1682-1684, 1991.
- [68100] 6430.Harris, G. W.: Neural control of the pituitary gland. *Physiol.Rev.* 28: 139-179, 1948.
- [68101] 6431.Holmes, F. E.; Mahoney, S.; King, V. R.; Bacon, A.; Kerr, N. C.H.; Pachnis, V.; Curtis, R.; Priestley, J. V.; Wynn-ick, D.: Targetedisruption of the galanin gene reduces the number of sensory neuronsand their regenerative capacity. *Proc. Nat. Acad. Sci.* 97: 11563-11568,2000.
- [68102] 6432.Lopez, F. J.; Merchenthaler, I.; Ching, M.; Wisniewski,

M. G.;Negro-Vilar, A.: Galanin: a hypothalamic-hypophysiotropic hormonemodulating reproductive functions. Proc. Nat. Acad. Sci. 88: 4508–4512,1991.

[68103] 6433.Lundkvist, J.; Land, T.; Kahl, U.; Bedecs, K.; Bartfai, T.: cDNAsequence, ligand binding, and regulation of galanin/GMAP in mousebrain. Neurosci. Lett. 200: 121–124, 1995.

[68104] 6434.O'Meara, G.; Coumis, U.; Ma, S. Y.; Kehr, J.; Mahoney, S.; Bacon,A.; Allen, S. J.; Holmes, F.; Kahl, U.; Wang, F. H.; Kearns, I. R.;Ove-Ogren, S.; Dawbarn, D.; Mufson, E. J.; Davies, C.; Dawson, G.;Wynick, D.: Galanin regulates the postnatal survival of a subsetof basal forebrain cholinergic neurons. Proc. Nat. Acad. Sci. 97:11569–11574, 2000.

[68105] 6435.Rattan, S.: Role of galanin in the gut. Gastroenterology 100:1762–1768, 1991.

[68106] 6436.Schmidt, W. E.; Kratzin, H.; Eckart, K.; Dreves, D.; Mundkowski,G.; Clemens, A.; Katsoulis, S.; Schafer, H.; Gallwitz, B.; Creutzfeldt,W.: Isolation and primary structure of pituitary human galanin, a30-residue nonamidated neuropeptide. Proc. Nat. Acad. Sci. 88: 11435–11439,1991.

[68107] 6437.Steiner, R. A.; Hohmann, J. G.; Holmes, A.; Wrenn, C.

C.; Cadd, G.; Jureus, A.; Clifton, D. K.; Luo, M.; Gutshall, M.; Ma, S. Y.; Mufson, E. J.; Crawley, J. N.: Galanin transgenic mice display cognitive and neurochemical deficits characteristic of Alzheimer's disease. *Proc. Nat. Acad. Sci.* 98: 4184–4189, 2001.

[68108] 6438. Wynick, D.; Small, C. J.; Bacon, A.; Holmes, F. E.; Norman, M.; Ormandy, C. J.; Kilic, E.; Kerr, N. C. H.; Ghatei, M.; Talamantes, F.; Bloom, S. R.; Pachnis, V.: Galanin regulates prolactin release and lactotroph proliferation. *Proc. Nat. Acad. Sci.* 95: 12671–12676, 1998.

[68109] 6439. Puech, A.; Saint-Jore, B.; Funke, B.; Gilbert, D. J.; Sirotkin, H.; Copeland, N. G.; Jenkins, N. A.; Kucherlapati, R.; Morrow, B.; Skoultschi, A. I.: Comparative mapping of the human 22q11 chromosomal region and the orthologous region in mice reveals complex changes in gene organization. *Proc. Nat. Acad. Sci.* 94: 14608–14613, 1997.

[68110] 6440. van Hille, B.; Vanek, M.; Richener, H.; Green, J. R.; Bilbe, G.: Cloning and tissue distribution of subunits C, D, and E of the human vacuolar H(+)-ATPase. *Biochem. Biophys. Res. Commun.* 97: 15–21, 1993.

[68111] 6441. Lewis, J.; Dickson, D. W.; Lin, W.-L.; Chisholm, L.; Corral, A.; Jones, G.; Yen, S.-H.; Sahara, N.; Skipper, L.; Yager, D.; Eckman, C.; Hardy, J.; Hutton, M.; McGowan, E.:

Enhanced neurofibrillarydegeneration in transgenic mice expressing mutant tau and APP. Science 293:1487–1491, 2001.

[68112] 6442.Denker, S. P.; Huang, D. C.; Orlowski, J.; Furthmayr, H.; Barber,D. L.: Direct binding of the Na–H exchanger NHE1 to ERM proteinsregulates the cortical cytoskeleton and cell shape independently ofH(+) translocation. Molec. Cell 6: 1425–1436, 2000.

[68113] 6443.Dudley, C. R. K.; Giuffra, L. A.; Tippet, P.; Kidd, K. K.; Reeder,S. T.: The Na<sup>+</sup>/H<sup>+</sup> antiporter: a 'melt' polymorphism allows regionalmapping to the short arm of chromosome 1. Hum. Genet. 86: 79–83,1990.

[68114] 6444.Franchi, A.; Perucca–Lostanlen, D.; Pouyssegur, J.: Functionalexpression of a human Na<sup>+</sup>/H<sup>+</sup> antiporter gene transfected into antiporter–deficientmouse L cells. Proc. Nat. Acad. Sci. 83: 9388–9392, 1986.

[68115] 6445.Lifton, R. P.; Sardet, C.; Pouyssegur, J.; Lalouel, J.–M.: Cloningof the human genomic amiloride–sensitive Na<sup>+</sup>/H<sup>+</sup> antiporter gene, identificationof genetic polymorphisms, and localization on the genetic map of chromosome1p. Genomics 7: 131–135, 1990.

[68116] 6446.Mattei, M.–G.; Galloni, M.; Sardet, C.; Franchi, A.; Counillon,L.; Passage, E.; Pouyssegur, J.: Localization of

the antiporter gene(APNH) and chromosomal homology between human 1p, mouse 4 and Chinesehamster 2q.

(Abstract) Cytogenet. Cell Genet. 51: 1041, 1989.

[68117] 6447.Mattei, M.-G.; Sardet, C.; Franchi, A.; Pouyssegur, J.: Chromosomal mapping of the amiloride-sensitive  $\text{Na}^+/\text{H}^+$  antiporter gene. (Abstract) Cytogenet. Cell Genet. 46: 658-659, 1987.

[68118] 6448.Mattei, M.-G.; Sardet, C.; Franchi, A.; Pouyssegur, J.: The human amiloride-sensitive  $\text{Na}^+/\text{H}^+$  antiporter: localization to chromosome 1 by in situ hybridization. Cytogenet. Cell Genet. 48: 6-8, 1988.

[68119] 6449.Mendoza, S. A.: The  $\text{Na}^+/\text{H}^+$  antiport is a mediator of cell proliferation. Acta Paediat. Scand. 76: 545-547, 1987.

[68120] 6450.Morahan, G.; Rakar, S.: Localization of the mouse  $\text{Na}^+/\text{H}^+$  exchanger gene on distal chromosome 4. Genomics 15: 231-232, 1993.

[68121] 6451.Sardet, C.; Franchi, A.; Pouyssegur, J.: Molecular cloning, primary structure, and expression of the human growth factor-activatable  $\text{Na}^+/\text{H}^+$  antiporter. Cell 56: 271-280, 1989.

[68122] 6452.Davare, M. A.; Avdonin, V.; Hall, D. D.; Peden, E. M.; Burette, A.; Weinberg, R. J.; Horne, M. C.; Hoshi, T.; Hell, J.

W.: A beta-2adrenergic receptor signaling complex assembled with the Ca(2+) channelCa(V)1.2. Science 293: 98–101, 2001.

- [68123] 6453.Jacobs, P. A.; Brunton, M.; Frackiewicz, A.; Newton, M.; Cook,P. J. L.; Robson, E. B.: Studies on a family with three cytogeneticmarkers. Ann. Hum. Genet. 33: 325–336, 1970.
- [68124] 6454.Brown, A.; Bernier, G.; Mathieu, M.; Rossant, J.; Kothary, R.:The mouse dystonia musculorum gene is a neural isoform of bullouspemphigoid antigen 1. Nature Genet. 10: 301–306, 1995.
- [68125] 6455.Brown, A.; Dalpe, G.; Mathieu, M.; Kothary, R.: Cloning and characterizationof the neural isoforms of human dystonin. Genomics 29: 777–780,1995.
- [68126] 6456.Copeland, N. G.; Gilbert, D. J.; Li, K.; Sawamura, D.; Giudice,G. J.; Chu, M.–L.; Jenkins, N. A.; Uitto, J.: Chromosomal localizationof mouse bullous pemphigoid antigens, BPAG1 and BPAG2: identificationof a new region of homology between mouse and human chromosomes. Genomics 15:180–181, 1993.
- [68127] 6457.Diaz, L. A.; Ratrie, H., III; Saunders, W. S.; Futamura, S.; Squiquera,H. L.; Anhalt, G. J.; Giudice, G. J.: Isolation of a human epidermalcDNA corresponding to the 180–kD

autoantigen recognized by bullous pemphigoid and herpes gestationis sera: immunolocalization of this protein to the hemidesmosome. *J. Clin. Invest.* 86: 1088–1094, 1990.

[68128] 6458. Guo, L.; Degenstein, L.; Dowling, J.; Yu, Q.-C.; Wollmann, R.; Perman, B.; Fuchs, E.: Gene targeting of BPAG1: abnormalities in mechanical strength and cell migration in stratified epithelia and neurologic degeneration. *Cell* 81: 233–243, 1995.

[68129] 6459. Minoshima, S.; Amagai, M.; Kudoh, J.; Fukuyama, R.; Hashimoto, T.; Nishikawa, T.; Shimizu, N.: Localization of the human gene for 230-kDa bullous pemphigoid autoantigen to the pter-q15 region of chromosome 6. (Abstract) *Cytogenet. Cell Genet.* 58: 1914–1915, 1991.

[68130] 6460. Minoshima, S.; Amagai, M.; Kudoh, J.; Fukuyama, R.; Hashimoto, T.; Nishikawa, T.; Shimizu, N.: Localization of the human gene for 230-kDa bullous pemphigoid autoantigen (BPAG1) to chromosome 6 pter-q15. *Cytogenet. Cell Genet.* 57: 30–32, 1991.

[68131] 6461. Ryynanen, M.; Knowlton, R. G.; Kero, M.; Sawamura, D.; Li, K.; Giudice, G. J.; Diaz, L. A.; Uitto, J.: Bullous pemphigoid antigens (BPAGs): identification of RFLPs in human BPAG1 and BPAG2, and exclusion as candidate genes in a large kindred with dominant epidermolysis bullosa sim-

plex. Genomics 11: 1025–1029, 1991.

- [68132] 6462.Sawamura, D.; Nomura, K.; Sugita, Y.; Mattei, M.–G.; Chu, M.–L.; Knowlton, R.; Uitto, J.: Bullous pemphigoid antigen (BPAG1): cDNA cloning and mapping of the gene to the short arm of human chromosome 6. Genomics 8: 722–726, 1990.
- [68133] 6463.Stanley, J. R.; Tanaka, T.; Mueller, S.; Klaus–Kovtun, V.; Roop, D.: Isolation of complementary DNA for bullous pemphigoid antigen by use of patients' autoantibodies. J. Clin. Invest. 82: 1864–1870, 1988.
- [68134] 6464.Tamai, K.; Sawamura, D.; Do, H. C.; Tamai, Y.; Li, K.; Uitto, J.: The human 230–kD bullous pemphigoid antigen gene (BPAG1): exon–intron organization and identification of regulatory tissue specific elements in the promoter region. J. Clin. Invest. 92: 814–822, 1993.
- [68135] 6465.Yang, Y.; Bauer, C.; Strasser, G.; Wollman, R.; Julien, J.–P.; Fuchs, E.: Integrators of the cytoskeleton that stabilize microtubules. Cell 98:229–238, 1999.
- [68136] 6466.Tashian, R. E.: Genetics of the mammalian carbonic anhydrases. Adv.Genet. 30: 321–356, 1992.
- [68137] 6467.Montgomery, J. C.; Shows, T. B.; Venta, P. J.; Tashian, R. E.: Gene for novel human carbonic anhydrase (CA) isozyme on chromosome 16 is unlinked to the



CA1/CA2/CA3 gene cluster. (Abstract) Am. J. Hum. Genet. 41: A229 only, 1987.

- [68138] 6468. Montgomery, J. C.; Venta, P. J.; Eddy, R. L.; Fukushima, Y.-S.; Shows, T. B.; Tashian, R. E.: Characterization of the human gene for a newly discovered carbonic anhydrase, CA VII, and its localization to chromosome 16. Genomics 11: 835–848, 1991.
- [68139] 6469. Venta, P. J.; Montgomery, J. C.; Tashian, R. E.: Molecular genetics of carbonic anhydrase isozymes. Isozymes: Curr. Top. Biol. Med. Res. 14: 59–72, 1987.
- [68140] 6470. Blake, N. M.: Genetic variants of carbonic anhydrase in the Asian–Pacific area. Ann. Hum. Biol. 5: 557–568, 1978.
- [68141] 6471. Blake, N. M.; Kirk, R. L.: Widespread distribution of variant forms of carbonic anhydrase in Australian aborigines. Med. J. Aust. 1: 183–185, 1978.
- [68142] 6472. Butterworth, P.; Barlow, J.; Konialis, C.; Povey, S.; Edwards, Y. H.: The assignment of human erythrocyte carbonic anhydrase CA1 to chromosome 8. (Abstract) Cytogenet. Cell Genet. 40: 597 only, 1985.
- [68143] 6473. Carter, N. D.: Carbonic anhydrase isozymes in *Cavia porcellus*, *Cavia aperea* and their hybrids. Comp. Biochem. Physiol. B 43: 743–747, 1972.

- [68144] 6474.Carter, N. D.: Carbonic anhydrase II polymorphism in Africa. *Hum.Hered.* 22: 539–541, 1972.
- [68145] 6475.Carter, N. D.; Tashian, R. E.; Huntsman, R. G.; Sacker, L.: Characterization of two new variants of red cell carbonic anhydrase in the British population: Ca le Portsmouth and Ca le Hull. *Am. J. Hum. Genet.* 24:330–338, 1972.
- [68146] 6476.Davis, M. B.; West, L. F.; Barlow, J. H.; Butterworth, P. H. W.; Lloyd, J. C.; Edwards, Y. H.: Regional localization of carbonic anhydrase genes CA1 and CA3 on human chromosome 8. *Somat. Cell Molec. Genet.* 13:173–178, 1987.
- [68147] 6477.Davis, M. B.; West, L. F.; Butterworth, P.; Edwards, Y. H.: The assignment of human carbonic anhydrases CA1 and CA3 to chromosome 8q13–22. (Abstract) 7th Int. Cong. Hum. Genet., Berlin 616 only, 1986.
- [68148] 6478.DeSimone, J.; Linde, M.; Tashian, R. E.: Evidence for linkage of carbonic anhydrase isozyme genes in the pig-tailed macaque, *Macaca nemestrina*. *Nature N.B.* 242: 55–56, 1973.
- [68149] 6479.Dodgson, S. J.; Tashian, R. E.; Gross, G.; Carter, N. D.: *The Carbonic Anhydrases: Cellular Physiology and Molecular Genetics*. New York: Plenum, 1991.
- [68150] 6480.Edwards, Y. H.; Barlow, J. H.; Konialis, C. P.; Povey,

S.; Butterworth, P. H. W.: Assignment of the gene determining human carbonic anhydrase, CAI, to chromosome 8. *Ann. Hum. Genet.* 50: 123–129, 1986.

[68151] 6481. Eicher, E. M.; Stern, R. H.; Womack, J. E.; Davisson, M. T.; Roderick, T. H.; Reynolds, S. C.: Evolution of mammalian carbonic anhydrase loci by tandem duplication: close linkage of Car-1 and Car-2 to the centromere region of chromosome 3 of the mouse. *Biochem. Genet.* 14:651–660, 1976.

[68152] 6482. Goriki, K.; Tashian, R. E.; Stroup, S. K.; Yu, Y.-S. L.; Henriksson, D. M.: Chemical characterization of a new Japanese variant of carbonic anhydrase I, Ca 2 (Nagasaki 1) (76 arg-to-gln). *Biochem. Genet.* 17:449–460, 1979.

[68153] 6483. Hewett-Emmett, D.; Tashian, R. E.: Functional diversity, conservation and convergence in the evolution of the alpha-, beta-, and gamma-carbonic anhydrase gene families. *Molec. Phylogenet. Evol.* 5: 50–77, 1996.

[68154] 6484. Kere, J.; Ruutu, T.; Davies, K. A.; Roninson, I. B.; Watkins, P. C.; Winqvist, R.; de la Chapelle, A.: Chromosome 7 long arm deletion in myeloid disorders: a narrow breakpoint region in 7q22 defined by molecular mapping. *Blood* 73: 230–234, 1989.

[68155] 6485. Neufeld, E. J.: Personal Communication. Boston,

Mass. 2/21/1995.

- [68156] 6486. Neufeld, E. J.; Skalnik, D. G.; Lievens, P. M.-J.; Orkin, S. H.: Human CCAAT displacement protein is homologous to the Drosophila homeoprotein, cut. *Nature Genet.* 1: 50–55, 1992.
- [68157] 6487. Scherer, S. W.; Neufeld, E. J.; Lievens, P. M.-J.; Orkin, S. H.; Kim, J.; Tsui, L.-C.: Regional localization of the CCAAT displacement protein gene (CUTL1) to 7q22 by analysis of somatic cell hybrids. *Genomics* 15:695–696, 1993.
- [68158] 6488. Snyder, S. R.; Wang, J.; Waring, J. F.; Ginder, G. D.: Identification of CCAAT displacement protein (CDP/cut) as a locus-specific repressor of major histocompatibility complex gene expression in human tumor cells. *J. Biol. Chem.* 276: 5323–5330, 2001.
- [68159] 6489. Zeng, W. R.; Scherer, S. W.; Koutsilieris, M.; Huizenga, J. J.; Filteau, F.; Tsui, L.-C.; Nepveu, A.: Loss of heterozygosity and reduced expression of the CUTL1 gene in uterine leiomyomas. *Oncogene* 14:2355–2365, 1997.
- [68160] 6490. Blackburn, M. R.; Datta, S. K.; Kellems, R. E.: Adenosine deaminase-deficient mice generated using a two-stage genetic engineering strategy exhibit a combined immunodeficiency. *J. Biol. Chem.* 273: 5093–5100, 1998.

- [68161] 6491. Blaese, R. M.; Culver, K. W.; Miller, A. D.; Carter, C. S.; Fleisher, T.; Clerici, M.; Shearer, G.; Chang, L.; Chiang, Y.; Tolstoshev, P.; Greenblatt, J. J.; Rosenberg, S. A.; Klein, H.; Berger, M.; Mullen, C. A.; Ramsey, W. J.; Muul, L.; Morgan, R. A.; Anderson, W. F.: T lymphocyte-directed gene therapy for ADA-SCID: initial trial results after 4 years. *Science* 270: 475-480, 1995.
- [68162] 6492. Bollinger, M. E.; Arredondo-Vega, F. X.; Santisteban, I.; Schwarz, K.; Hershfield, M. S.; Lederman, H. M.: Hepatic dysfunction as a complication of adenosine deaminase deficiency. *New Eng. J. Med.* 334:1367-1371, 1996.
- [68163] 6493. Bonthron, D. T.; Markham, A. F.; Ginsburg, D.; Orkin, S. H.: Identification of a point mutation in the adenosine deaminase gene responsible for immunodeficiency. *J. Clin. Invest.* 76: 894-897, 1985.
- [68164] 6494. Bordignon, C.; Notarangelo, L. D.; Nobili, N.; Ferrari, G.; Casorati, G.; Panina, P.; Mazzolari, E.; Maggioni, D.; Rossi, C.; Servida, P.; Ugazio, A. G.; Mavilio, F.: Gene therapy in peripheral blood lymphocytes and bone marrow for ADA: immunodeficient patients. *Science* 270:470-475, 1995.
- [68165] 6495. Bortin, M. M.; Rimm, A. A. (eds.): Severe combined immunodeficiency disease: characterization of the disease

and results of transplantation. J.A.M.A. 238:591–600, 1977.

- [68166] 6496.Boss, G. R.; Thompson, L. F.; O'Connor, R. D.; Ziering, R. W.;Seegmiller, J. E.: Ecto-5-prime-nucleotidase deficiency: associationwith adenosine deaminase deficiency and nonassociation with deoxyadenosinetoxicity. Clin. Immun. Immunopath. 19: 1–7, 1981.
- [68167] 6497.Bottini, N.; De Luca, D.; Saccucci, P.; Fiumara, A.; Elia, M.;Porfirio, M. C.; Lucarelli, P.; Curatolo, P.: Autism: evidence ofassociation with adenosine deaminase genetic polymorphism. Neurogenetics 3:111–113, 2001.
- [68168] 6498.Chen, S.–H.; Ochs, H. D.; Scott, C. R.: Adenosine deaminase deficiency:disappearance of adenine deoxynucleotides from a patient's erythrocytesafter successful marrow transplantation. J. Clin. Invest. 62: 1386–1389,1978.
- [68169] 6499.Chen, S.–H.; Ochs, H. D.; Scott, C. R.; Giblett, E. R.: Adenosinedeaminase and nucleoside phosphorylase activity in patients with immunodeficiencysyndromes. Clin. Immun. Immunopath. 13: 156–160, 1979.
- [68170] 6500.Chen, S.–H.; Scott, C. R.; Giblett, E. R.: Adenosine deaminase:demonstration of a 'silent' gene associated with combined immunodeficiencydisease. Am. J. Hum.

Genet. 26: 103–107, 1974.

- [68171] 6501.Cohen, A.; Hirschhorn, R.; Horowitz, S. D.; Rubinstein, A.; Polmar, S. H.; Hong, R.; Martin, D. W., Jr.: Deoxyadenosine triphosphate as a potentially toxic metabolite in adenosine deaminase deficiency. Proc.Nat. Acad. Sci. 75: 472–476, 1978.
- [68172] 6502.Cook, P. J. L.; Hopkinson, D. A.; Robson, E. B.: The linkage relationships of adenosine deaminase. Ann. Hum. Genet. 34: 187–188, 1970.
- [68173] 6503.Creagan, R. P.; Tischfield, J. A.; Nichols, E. A.; Ruddle, F.H.: Autosomal assignment of the gene for the form of adenosine deaminase which is deficient in patients with combined immunodeficiency syndrome.(Letter) Lancet II: 1449 only, 1973.
- [68174] 6504.Culver, K. W.: Gene Therapy: A Primer for Physicians. Larchmont, New York: Mary Ann Liebert, Inc. , 1996.
- [68175] 6505.Daddona, P. E.; Kelley, W. N.: Human adenosine deaminase: stoichiometry of the adenosine deaminase-binding protein complex. Biochim. Biophys. Acta 580: 302–311, 1979.
- [68176] 6506.Daddona, P. E.; Mitchell, B. S.; Meuwissen, H. J.; Davidson, B.L.; Wilson, J. M.; Koller, C. A.: Adenosine deaminase deficiency with normal immune function: an

acidic enzyme mutation. J. Clin. Invest. 72: 483–492, 1983.

[68177] 6507. Detter, J. C.; Stamatoyannopoulos, G.; Giblett, E. R.; Motulsky, A. G.: Adenosine deaminase: racial distribution and report of a new phenotype. J. Med. Genet. 7: 356–357, 1970.

[68178] 6508. Dissing, J.; Knudsen, B.: Adenosine–deaminase deficiency and combined immunodeficiency syndrome. (Letter) Lancet II: 1316, 1972.

[68179] 6509. Dissing, J.; Knudsen, J. B.: A new red cell adenosine deaminase phenotype in man. Hum. Hered. 19: 375–377, 1969.

[68180] 6510. Dyminski, J. W.; Daoud, A.; Lampkin, B. C.; Limouze, S.; Donofrio, J.; Coleman, M. S.; Hutton, J. J.: Immunological and biochemical profiles in response to transfusion therapy in adenosine deaminase–deficient patient with severe combined immunodeficiency disease. Clin. Immun. Immunopath. 14: 307–326, 1979.

[68181] 6511. Egashira, M.; Ariga, T.; Kawamura, N.; Miyoshi, O.; Niikawa, N.; Sakiyama, Y.: Visible integration of the adenosine deaminase (ADA) gene into the recipient genome after gene therapy. Am. J. Med. Genet. 75: 314–317, 1998.

[68182] 6512. Ellis, N. A.; Lennon, D. J.; Proytcheva, M.; Alhadeff,



B.; Henderson, E. E.; German, J.: Somatic intragenic recombination within the mutated locus BLM can correct the high SCE phenotype of Bloom syndrome cells. *Am. J. Hum. Genet.* 57: 1019–1027, 1995.

[68183] 6513. Ferrari, G.; Rossini, S.; Giavazzi, R.; Maggioni, D.; Nobili, N.; Soldati, M.; Ungers, G.; Mavilio, F.; Gilboa, E.; Bordignon, C.: An in vivo model of somatic cell gene therapy for human severe combined immunodeficiency. *Science* 251: 1363–1366, 1991.

[68184] 6514. Giblett, E. R.; Anderson, J. E.; Cohen, F.; Pollara, B.; Meuwissen, H. J.: Adenosine-deaminase deficiency in two patients with severely impaired cellular immunity. *Lancet* I: 1067–1069, 1972.

[68185] 6515. Hart, S. L.; Lane, A. B.; Jenkins, T.: Partial adenosine deaminase deficiency: another family from southern Africa. *Hum. Genet.* 74: 307–312, 1986.

[68186] 6516. Herbschleb-Voogt, E.; Scholten, J.-W.; Meera Khan, P.: Basic defect in the expression of adenosine deaminase in ADA SCID disease. II. Deficiency of ADA-CRM detected in heterozygote human-Chinese hamster cell hybrids. *Hum. Genet.* 63: 121–125, 1983.

[68187] 6517. Hershfield, M. S.: PEG-ADA: an alternative to haploidentical bone marrow transplantation and an adjunct to

gene therapy for adenosinedeaminase deficiency. Hum. Mutat. 5: 107–112, 1995.

[68188] 6518.Hershfield, M. S.; Buckley, R. H.; Greenberg, M. L.; Melton, A.L.; Schiff, R.; Hatem, C.; Kurtzberg, J.; Markert, M. L.; Kobayashi,R. H.; Kobayashi, A. L.; Abuchowski, A.: Treatment of adenosine deaminasedeficiency with polyethylene glycol–modified adenosine deaminase. NewEng. J. Med. 316: 589–596, 1987.

[68189] 6519.Hershfield, M. S.; Kredich, N. M.: S–adenosylhomocysteine hydrolaseis an adenosine–binding protein: a target for adenosine toxicity. Science 202:757–760, 1978.

[68190] 6520.Presley, J. F.; Ward, T. H.; Pfeifer, A. C.; Siggia, E. D.; Phair,R. D.; Lippincott–Schwartz, J.: Dissection of COPI and Arf1 dynamicsin vivo and role in Golgi membrane transport. Nature 417: 187–193,2002.

[68191] 6521.McGuire, R. E.; Daiger, S. P.; Green, E. D.: Localization andcharacterization of the human ADP–ribosylation factor 5 (ARF5) gene. Genomics 41:481–484, 1997.

[68192] 6522.Tsuchiya, M.; Price, S. R.; Tsai, S.–C.; Moss, J.; Vaughan, M.: Molecular identification of ADP–ribosylation factor mRNAs and their expression in mammalian cells. J. Biol. Chem. 266: 2772–2777, 1991.

- [68193] 6523.Chang, C.-Y.; Wu, D.-A.; Lai, C.-C.; Miller, W. L.; Chung, B.-C.: Cloning and structure of the human adrenodoxin gene. DNA 7: 609–615,1988.
- [68194] 6524.Chang, C.-Y.; Wu, D.-A.; Mohandas, T. K.; Chung, B.-C.: Structure,sequence, chromosomal location, and evolution of the human ferredoxingene family. DNA Cell Biol. 9: 205–212, 1990.
- [68195] 6525.Lachman, H. M.; Papolos, D. F.; Saito, T.; Yu, Y. M.; Szumlanski,C. L.; Weinshilboum, R. M.: Human catechol-O-methyltransferase pharmacogenetics:description of a functional polymorphism and its potential applicationto neuropsychiatric disorders. Pharmacogenetics 6: 243–250, 1996.
- [68196] 6526.Tiihonen, J.; Hallikainen, T.; Lachman, H.; Saito, T.; Volavka,J.; Kauhanen, J.; Salonen, J. T.; Ryyanen, O.-P.; Koulu, M.; Karvonen,M. K.; Pohjalainen, T.; Syvalahti, E.; Hietala, J.: Association betweenthe functional variant of the catechol-O-methyltransferase (COMT)gene and type 1 alcoholism. Molec. Psychiat. 4: 286–289, 1999.
- [68197] 6527.Bohren, K. M.; Bullock, B.; Wermuth, B.; Gabbay, K. H.: The aldo-ketoreductase superfamily: cDNAs and deduced amino acid sequences of humanaldehyde and aldose reductases. J. Biol. Chem. 264: 9547–9551, 1989.

- [68198] 6528.Charlesworth, D.: Starch–gel electrophoresis of four enzymes from human red blood cells: glyceraldehyde–3–phosphate dehydrogenase, fructoaldolase, glyoxalase II and sorbitol dehydrogenase. *Ann. Hum. Genet.* 35: 477–484, 1972.
- [68199] 6529.Frade, R.; Balbo, M.; Barel, M.: RB18A, whose gene is localized on chromosome 17q12–q21.1, regulates in vivo p53 transactivating activity. *Cancer Res.* 60: 6585–6589, 2000.
- [68200] 6530.Wolffe, A. P.: Transcriptional control: sinful repression. *Nature* 387:16–17, 1997.
- [68201] 6531.Wu, L.; Aster, J. C.; Blacklow, S. C.; Lake, R.; Artavanis–Tsakonas, S.; Griffin, J. D.: MAML1, a human homologue of *Drosophila* mastermind, is a transcriptional co–activator for NOTCH receptors. *Nature Genet.* 26:484–489, 2000.
- [68202] 6532.Inohara, N.; Koseki, T.; del Peso, L.; Hu, Y.; Yee, C.; Chen, S.; Carrio, R.; Merino, J.; Liu, D.; Ni, J.; Nunez, G.: Nod1, an Apaf–1–like activator of caspase–9 and nuclear factor–kappa–B. *J. Biol. Chem.* 274:14560–14567, 1999.
- [68203] 6533.Charroux, B.; Pellizzoni, L.; Perkinson, R. A.; Shevchenko, A.; Mann, M.; Dreyfuss, G.: Gemin3: a novel DEAD box protein that interacts with SMN, the spinal mus–

cular atrophy gene product, and is a component of gems.

J. Cell Biol. 147: 1181–1193, 1999.

[68204] 6534. Grundhoff, A. T.; Kremmer, E.; Tureci, O.; Gleden, A.; Gindorf, C.; Atz, J.; Mueller-Lantzsch, N.; Schubach, W. H.; Grasser, F. A.: Characterization of DP103, a novel DEAD box protein that binds to the Epstein-Barr virus nuclear proteins EBNA2 and EBNA3C. J. Biol. Chem. 274: 19136–19144, 1999.

[68205] 6535. Ruteshouser, E. C.; Ashworth, L. K.; Huff, V.: Absence of PPP2R1A mutations in Wilms tumor. Oncogene 20: 2050–2054, 2001.

[68206] 6536. Everett, A. D.; Xue, C.; Stoops, T.: Developmental expression of protein phosphatase 2A in the kidney. J. Am. Soc. Nephrol. 10: 1737–1745, 1999.

[68207] 6537. Hemmings, B. A.; Adams-Pearson, C.; Maurer, F.; Muller, P.; Goris, J.; Merlevede, W.; Hofsteenge, J.; Stone, S. R.: Alpha- and beta-forms of the 65-kDa subunit of protein phosphatase 2A have a similar 39 amino acid repeating structure. Biochemistry 29: 3166–3173, 1990.

[68208] 6538. Hendrix, P.; Turowski, P.; Mayer-Jaekel, R. E.; Goris, J.; Hofsteenge, J.; Merlevede, W.; Hemmings, B. A.: Analysis of subunit isoforms in protein phosphatase 2A holoenzymes from rabbit and *Xenopus*. J. Biol. Chem. 268:

7330–7337, 1993.

- [68209] 6539.Hunt, S. L.; Hsuan, J. J.; Totty, N.; Jackson, R. J.: unr, a cellularcytoplasmic RNA-binding protein with five cold-shock domains, is requiredfor internal initiation of translation of human rhinovirus RNA. *GenesDev.* 13: 437–448, 1999.
- [68210] 6540.Matsuda, S.; Katsumata, R.; Okuda, T.; Yamamoto, T.; Miyazaki,K.; Senga, T.; Machida, K.; Thant, A. A.; Nakatsugawa, S.; Hamaguchi,M.: Molecular cloning and characterization of human MAWD, a novelprotein containing WD-40 repeats frequently overexpressed in breast-cancer. *Cancer Res.* 60: 13–17, 2000.
- [68211] 6541.Levy, C.; Nechushtan, H.; Razin, E.: A new role for the STAT3inhibitor, PIAS3: a repressor of microphthalmia transcription factor. *J.Biol. Chem.* 277: 1962–1966, 2002.
- [68212] 6542.Ueki, N.; Seki, N.; Yano, K.; Saito, T.; Masuho, Y.; Muramatsu,M.: Isolation and chromosomal assignment of a human gene encodingprotein inhibitor of activated STAT3 (PIAS3). *J. Hum. Genet.* 44:193–196, 1999.
- [68213] 6543.Pasteris, N. G.; Trask, B. J.; Sheldon, S.; Gorski, J. L.: Discordantphenotype of two overlapping deletions involving the PAX3 gene inchromosome 2q35. *Hum. Molec. Genet.* 2: 953–959, 1993.

- [68214] 6544. Beeson, D.; Brydson, M.; Betty, M.; Jeremiah, S.; Povey, S.; Vincent, A.; Newsom-Davis, J.: Primary structure of the human muscle acetylcholinereceptor cDNA cloning of the gamma and epsilon subunits. *Europ. J. Biochem.* 215: 229–238, 1993.
- [68215] 6545. Ohno, K.; Hutchinson, D. O.; Milone, M.; Brengman, J. M.; Bouzat, C.; Sine, S. M.; Engel, A. G.: Congenital myasthenic syndrome caused by prolonged acetylcholine receptor channel openings due to a mutation in the M2 domain of the epsilon subunit. *Proc. Nat. Acad. Sci.* 92: 758–762, 1995.
- [68216] 6546. Ohno, K.; Quiram, P. A.; Milone, M.; Wang, H.-L.; Harper, M. C.; Pruitt, J. N., II; Brengman, J. M.; Pao, L.; Fischbeck, K. H.; Crawford, T. O.; Sine, S. M.; Engel, A. G.: Congenital myasthenic syndromes due to heteroallelic nonsense/missense mutations in the acetylcholinereceptor epsilon subunit gene: identification and functional characterization of six new mutations. *Hum. Molec. Genet.* 6: 753–766, 1997.
- [68217] 6547. Ohno, K.; Wang, H.-L.; Milone, M.; Bren, N.; Brengman, J. M.; Nakano, S.; Quiram, P.; Pruitt, J. N.; Sine, S. M.; Engel, A. G.: Congenital myasthenic syndrome caused by decreased agonist binding affinity due to a mutation in the

acetylcholine receptor epsilon subunit. *Neuron* 17:157–170, 1996.

- [68218] 6548.Sieb, J. P.; Dorfler, P.; Tzartos, S.; Wewer, U. M.; Ruegg, M.A.; Meyer, D.; Baumann, I.; Lindemuth, R.; Jakschik, J.; Ries, F.: Congenital myasthenic syndromes in two kinships with end-plate acetylcholinereceptor and utrophin deficiency. *Neurology* 50: 54–61, 1998.
- [68219] 6549.Sieb, J. P.; Kraner, S.; Rauch, M.; Steinlein, O. K.: Immature end-plates and utrophin deficiency in congenital myasthenic syndrome caused by epsilon-AChR subunit truncating mutations. *Hum. Genet.* 107:160–164, 2000.
- [68220] 6550.Witzemann, V.; Schwarz, H.; Koenen, M.; Berberich, C.; Villarroel, A.; Wernig, A.; Brenner, H. R.; Sakmann, B.: Acetylcholine receptorepsilon-subunit deletion causes muscle weakness and atrophy in juvenile and adult mice. *Proc. Nat. Acad. Sci.* 93: 13286–13291, 1996.
- [68221] 6551.Bartels, C. F.; Zelinski, T.; Lockridge, O.: Mutation at codon 322 in the human acetylcholinesterase (ACHE) gene accounts for Y blood group polymorphism. *Am. J. Hum. Genet.* 52: 928–936, 1993.
- [68222] 6552.Coates, P. M.; Simpson, N. E.: Genetic variation in human erythrocyte acetylcholinesterase. *Science* 175: 1466–1467, 1972.



- [68223] 6553.Ehrlich, G.; Viegas-Pequignot, E.; Ginzberg, D.; Sindel, L.; Soreq,H.; Zakut, H.: Mapping the human acetylcholinesterase gene to chromosome7q22 by fluorescent in situ hybridization coupled with selective PCRamplification from a somatic hybrid cell panel and chromosome-sortedDNA libraries. *Genomics* 13: 1192–1197, 1992.
- [68224] 6554.Feng, G.; Krejci, E.; Molgo, J.; Cunningham, J. M.; Massoulie,J.; Sanes, J. R.: Genetic analysis of collagen Q: roles in acetylcholinesteraseand butyrylcholinesterase assembly and in synaptic structure and function. *J.Cell Biol.* 144: 1349–1360, 1999.
- [68225] 6555.Getman, D. K.; Eubanks, J. H.; Camp, S.; Evans, G. A.; Taylor,P.: The human gene encoding acetylcholinesterase is located on the long arm of chromosome 7. *Am. J. Hum. Genet.* 51: 170–177, 1992.
- [68226] 6556.Lapidot-Lifson, Y.; Prody, C. A.; Ginzberg, D.; Meytes, D.; Zakut,H.; Soreq, H.: Coamplification of human acetylcholinesterase andbutyrylcholinesterase genes in blood cells: correlation with variousleukemias and abnormal megakaryocytopoiesis. *Proc. Nat. Acad. Sci.* 86:4715–4719, 1989.
- [68227] 6557.Meshorer, E.; Erb, C.; Gazit, R.; Pavlovsky, L.; Kaufer, D.; Friedman,A.; Glick, D.; Ben-Arie, N.; Soreq, H.: Alter-

native splicing and neuritic mRNA translocation under long-term neuronal hypersensitivity. *Science* 295:508–512, 2002.

- [68228] 6558. Rachinsky, T. L.; Crenshaw, E. B., III; Taylor, P.: Assignment of the gene for acetylcholinesterase to distal mouse chromosome 5. *Genomics* 14:511–514, 1992.
- [68229] 6559. Rotundo, R. L.; Gomez, A. M.; Fernandez-Valle, C.; Randall, W. R.: Allelic variants of acetylcholinesterase: genetic evidence that all acetylcholinesterase forms in avian nerves and muscles are encoded by a single gene. *Proc. Nat. Acad. Sci.* 85: 7805–7809, 1988.
- [68230] 6560. Shapira, M.; Tur-Kaspa, I.; Bosgraaf, L.; Livni, N.; Grant, A. D.; Grisaru, D.; Korner, M.; Ebstein, R. P.; Soreq, H.: A transcription-activating polymorphism in the ACHE promoter associated with acute sensitivity to anti-acetylcholinesterases. *Hum. Molec. Genet.* 9: 1273–1281, 2000.
- [68231] 6561. Drayna, D. T.; McLean, J. W.; Wion, K. L.; Trent, J. M.; Drabkin, H. A.; Lawn, R. M.: Human apolipoprotein D gene: gene sequence, chromosomal localization, and homology to the alpha-2mu-globulin superfamily. *DNA* 6:199–204, 1987.
- [68232] 6562. Fielding, P. E.; Fielding, C. J.: A cholesteryl ester

transfercomplex in human plasma. Proc. Nat. Acad. Sci. 77: 3327–3330, 1980.

[68233] 6563.Graham, C. A.; McGrew, W. C.: Menstrual synchrony in female undergraduates living on a coeducational campus. Psychoneuroendocrinology 5: 245–252, 1980.

[68234] 6564.Kamboh, M. I.; Albers, J. J.; Majumder, P. P.; Ferrell, R. E.: Genetic studies of human apolipoproteins. IX. Apolipoprotein D polymorphism and its relation to serum lipoprotein lipid levels. Am. J. Hum. Genet. 45:147–154, 1989.

[68235] 6565.McClintock, M. K.: Menstrual synchrony and suppression. Nature 229:244–245, 1971.

[68236] 6566.McClintock, M. K.: Estrous synchrony and its mediation by airborne chemical communication (*Rattus norvegicus*). Horm. Behav. 10: 264–276, 1978.

[68237] 6567.Quadagno, D. M.; Shubeita, H. E.; Deck, J.; Francoeur, D.: Influence of male social contacts, exercise and all-female living conditions on the menstrual cycle. Psychoneuroendocrinology 6: 239–244, 1981.

[68238] 6568.Warden, C. H.; Diep, A.; Taylor, B. A.; Lusi, A. J.: Localization of the gene for apolipoprotein D on mouse chromosome 16. Genomics 12:851–852, 1992.

[68239] 6569.Zeng, C.; Spielman, A. I.; Vowels, B. R.; Leyden, J. J.;

Biemann,K.; Preti, G.: A human axillary odorant is carried by apolipoproteinD. *Proc. Nat. Acad. Sci.* 93: 6626–6630, 1996.

[68240] 6570.Hopkinson, D. A.; Coppock, J. S.; Muhlemann, M. F.; Edwards, Y.H.: The detection and differentiation of the products of the humancarbonic anhydrase loci, Ca I and Ca II, using fluorogenic substrates. *Ann.Hum. Genet.* 38: 155–162, 1974.

[68241] 6571.Cleutjens, C. B. J. M.; van Eekelen, C. C. E. M.; van Dekken, H.;Smit, E. M. E.; Hagemeijer, A.; Wagner, M. J.; Wells, D. E.; Trapman,J.: The human C/EBP–delta (CRP3/CELF) gene: structure and chromosomallocalization. *Genomics* 16: 520–523, 1993.

[68242] 6572.Williams, S. C.; Cantwell, C. A.; Johnson, P. F.: A family ofC/EBP–related proteins capable of forming covalently linked leucinezipper dimers in vitro. *Genes Dev.* 5: 1553–1567, 1991.

[68243] 6573.Al–Awqati, Q.; Preisig, P. A.: Size does matter: will knockoutof p21(WAF1/CIP1) save the kidney by limiting compensatory renal growth?(Commentary) *Proc. Nat. Acad. Sci.* 96: 10551–10553, 1999.

[68244] 6574.Chedid, M.; Michieli, P.; Lengel, C.; Huppi, K.; Givol, D.: Asingle nucleotide substitution at codon 31 (ser/arg)

defines a polymorphism in a highly conserved region of the p53-inducible gene WAF1/CIP1. *Oncogene* 9:3021–3024, 1994.

- [68245] 6575. Cheng, T.; Rodrigues, N.; Shen, H.; Yang, Y.; Dom-bkowski, D.; Sykes, M.; Scadden, D. T.: Hematopoietic stem cell quiescence maintained by p21(cip1/waf1). *Science* 287: 1804–1808, 2000.
- [68246] 6576. Harper, J. W.; Adami, G. R.; Wei, N.; Keyomarsi, K.; Elledge, S. J.: The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases. *Cell* 75: 805–816, 1993.
- [68247] 6577. Huppi, K.; Siwarski, D.; Dosik, J.; Michieli, P.; Chedid, M.; Reed, S.; Mock, B.; Givol, D.; Mushinski, J. F.: Molecular cloning, sequencing, chromosomal localization and expression of mouse p21 (Waf1). *Oncogene* 9:3017–3020, 1994.
- [68248] 6578. Megyesi, J.; Price, P. M.; Tamayo, E.; Sfirstein, R. L.: The lack of a functional p21(WAF1/CIP1) gene ameliorates progression to chronic renal failure. *Proc. Nat. Acad. Sci.* 96: 10830–10835, 1999.
- [68249] 6579. Mousses, S.; Ozcelik, H.; Lee, P. D.; Malkin, D.; Bull, S. B.; Andrulis, I. L.: Two variants of the CIP1/WAF1 gene occur together and are associated with human cancer.

Hum. Molec. Genet. 4: 1089–1092,1995.

- [68250] 6580.Iwata, N.; Tsubuki, S.; Takaki, Y.; Shirotani, K.; Lu, B.; Gerard,N. P.; Gerard, C.; Hama, E.; Lee, H.–J.; Saido, T. C.: Metabolicregulation of brain A–beta by neprilysin. Sci–ence 292: 1550–1552,2001.
- [68251] 6581.Lu, P.–J.; Wulf, G.; Zhou, X. Z.; Davies, P.; Lu, K. P.: Theprolyl isomerase Pin1 restores the function of Alzheimer–associatedphosphorylated tau protein. Nature 399: 784–788, 1999.
- [68252] 6582.Kageoka, T.; Hewett–Emmett, D.; Stroup, S. K.; Yu, Y.–S. L.; Tashian,R. E.: Amino acid substitution and chemical characterization of aJapanese variant of carbonic anhydrase I: CA I Hiroshima–1 (86 asp–to–gly). Biochem.Genet. 19: 535–549, 1981.
- [68253] 6583.Kendall, A. G.; Tashian, R. E.: Erythrocyte carbonic anhydraseI: inherited deficiency in humans. Science 197: 471–472, 1977.
- [68254] 6584.Lindskog, S.; Henderson, L. E.; Kannan, K. K.; Liljas, A.; Nyman,P. O.; Strandberg, B.: Carbonic anhydrase.In: Boyer, P. D.: TheEnzymes. New York: Academic Press (pub.) 5: 1971. Pp. 587–665.
- [68255] 6585.McGeer, P. L.; McGeer, E. G.: Polymorphisms in inflammatory genesand the risk of Alzheimer disease. Arch.

Neurol. 58: 1790–1792,2001.

- [68256] 6586.Modi, W. S.; Masuda, A.; Yamada, M.; Oppenheim, J. J.; Matsushima,K.; O'Brien, S. J.: Chromosomal localization of the human interleukin1 alpha (IL-1-alpha) gene. Genomics 2: 310–314, 1988.
- [68257] 6587.Mosley, B.; Urdal, D. L.; Prickett, K. S.; Larsen, A.; Cosman,D.; Conlon, P. J.; Gillis, S.; Dower, S. K.: The interleukin-1 receptorbinds the human interleukin-1-alpha precursor but not the interleukin-1-betaprecursor. J. Biol. Chem. 262: 2941–2944, 1987.
- [68258] 6588.Bailly, S.; di Giovine, F. S.; Blakemore, A. I. F.; Duff, G. W.: Genetic polymorphism of human interleukin-1-alpha. Europ. J. Immun. 23:1240–1245, 1993.
- [68259] 6589.Boultwood, J.; Breckon, G.; Birch, D.; Cox, R.: Chromosomal localizationof murine interleukin-1 alpha and beta genes. Genomics 5: 481–485,1989.
- [68260] 6590.Cox, A.; Camp, N. J.; Cannings, C.; di Giovine, F. S.; Dale, M.;Worthington, J.; John, S.; Ollier, W. E. R.; Silman, A. J.; Duff,G. W.: Combined sib-TDT and TDT provide evidence for linkage of theinterleukin-1 gene cluster to erosive rheumatoid arthritis. Hum.Molec. Genet. 8: 1707–1713, 1999.
- [68261] 6591.Cox, A.; Camp, N. J.; Nicklin, M. J. H.; di Giovine, F.

S.; Duff, G. W.: An analysis of linkage disequilibrium in the interleukin-1 gene cluster, using a novel grouping method for multiallelic markers. *Am. J. Hum. Genet.* 62: 1180–1188, 1998.

[68262] 6592. Diehl, S. R.; Wang, Y.; Brooks, C. N.; Burmeister, J. A.; Califano, J. V.; Wang, S.; Schenkein, H. A.: Linkage disequilibrium of interleukin-1 genetic polymorphisms with early-onset periodontitis. *J. Periodont.* 70:418–430, 1999.

[68263] 6593. Du, Y.; Dodel, R. C.; Eastwood, B. J.; Bales, K. R.; Gao, F.; Lohmuller, F.; Muller, U.; Kurz, A.; Zimmer, R.; Evans, R. M.; Hake, A.; Gasser, T.; Oertel, W. H.; Griffin, W. S. T.; Paul, S. M.; Farlow, M. R.: Association of an interleukin 1-alpha polymorphism with Alzheimer's disease. *Neurology* 55: 480–484, 2000.

[68264] 6594. Furutani, Y.; Notake, M.; Fukui, T.; Ohue, M.; Nomura, H.; Yamada, M.; Nakamura, S.: Complete nucleotide sequence of the gene for human interleukin 1 alpha. *Nucleic Acids Res.* 14: 3167–3179, 1986.

[68265] 6595. Gray, P. W.; Glaister, D.; Chen, E.; Goeddel, D. V.; Pennica, D.: Two interleukin 1 genes in the mouse: cloning and expression of the cDNA for murine interleukin 1-beta. *J. Immun.* 137: 3644–3648, 1986.

[68266] 6596. Grimaldi, L. M. E.; Casadei, V. M.; Ferri, C.; Veglia, F.;



Licastro,F.; Annoni, G.; Biunno, I.; De Bellis, G.; Sorbi, S.; Mariani, C.;Canal, N.; Griffin, W. S. T.: Association of early-onset Alzheimer'sdisease with an interleukin-1-alpha gene polymorphism. *Ann. Neurol.* 47:361-365, 2000.

[68267] 6597.Hogquist, K. A.; Nett, M. A.; Unanue, E. R.; Chaplin, D. D.:Interleukin 1 is processed and released during apoptosis. *Proc. Nat.Acad. Sci.* 88: 8485-8489, 1991.

[68268] 6598.Hurwitz, A.; Loukides, J.; Ricciarelli, E.; Botero, L.; Katz,E.; McAllister, J. M.; Garcia, J. E.; Rohan, R.; Adashi, E. Y.; Hernandez,E. R.: Human intraovarian interleukin-1 (IL-1) system: highly compartmentalizedand hormonally dependent regulation of the genes encoding IL-1, itsreceptor, and its receptor antagonist. *J. Clin. Invest.* 89: 1746-1754,1992.

[68269] 6599.Ki, C.-S.; Na, D. L.; Kim, D. K.; Kim, H. J.; Kim, J.-W.: Lackof association of the interleukin-1-alpha gene polymorphism with Alzheimer'sdisease in a Korean population. (Letter) *Ann. Neurol.* 49: 817-818,2001.

[68270] 6600.Kolsch, H.; Ptok, U.; Bagli, M.; Papassotiropoulos, A.; Schmitz,S.; Barkow, K.; Kockler, M.; Rao, M. L.; Maier, W.; Heun, R.: Genepolymorphisms of interleukin-1-alpha influence the course of Alzheimer'sdisease. (Letter) *Ann.*

Neurol. 49: 818–819, 2001.

- [68271] 6601.Kornman, K. S.; Crane, A.; Wang, H.–Y.; di Giovine, F. S.; Newman,M. G.; Pirk, F. W.; Wilson, T. G., Jr.; Higginbottom, F. L.; Duff,G. W.: The interleukin–1 genotype as a severity factor in adult periodontaldisease. J. Clin. Periodont. 24: 72–77, 1997.
- [68272] 6602.Lafage, M.; Maroc, N.; Dubreuil, P.; de Waal Malefijt, R.; Pebusque,M.–J.; Carcassonne, Y.; Mannoni, P.: The human interleukin–1–alphagene is located on the long arm of chromosome 2 at band q13. Blood 73:104–107, 1989.
- [68273] 6603.Lord, P. C. W.; Wilmoth, L. M. G.; Mizel, S. B.; McCall, C. E.: Expression of interleukin–1 alpha and beta genes by human bloodpolymorphonuclear leukocytes. J. Clin. Invest. 87: 1312–1321, 1991.
- [68274] 6604.Murphy, G. M., Jr.; Claassen, J. D.; DeVoss, J. J.; Pascoe, N.;Taylor, J.; Tinklenberg, J. R.; Yesavage, J. A.: Rate of cognitivedecline in AD is accelerated by the interleukin–1–alpha –889 \*1 allele. Neurology 56:1595–1597, 2001.
- [68275] 6605.Stamey, T. A.; Yang, N.; Hay, A. R.; McNeal, J. E.; Freiha, F.S.; Redwine, E.: Prostate–specific antigen as a serum marker foradenocarcinoma of the prostate. New Eng. J. Med. 317: 909–916, 1987.
- [68276] 6606.Mallinson, G.; Soo, K. S.; Schall, T. J.; Pisacka, M.;

Anstee,D. J.: Mutations in the erythrocyte chemokine receptor (Duffy) gene:the molecular basis of the Fy(a)/Fy(b) antigens and identificationof a deletion in the Duffy gene of an apparently healthy individualwith the Fy(a-b-) phenotype. Brit. J. Haemat. 90: 823-829, 1995.

[68277] 6607.Miller, L. H.; Mason, S. J.; Clyde, D. F.; McGinnis, M. H.: Theristance factor to Plasmodium vivax in blacks: the Duffy blood groupgenotype, FyFy. New Eng. J. Med. 295: 302-304, 1976.

[68278] 6608.Miller, L. H.; Mason, S. J.; Dvorak, J. A.: Erythrocyte receptors of Plasmodium knowlesi malaria: Duffy blood group determinants. Science 189:561-562, 1975.

[68279] 6609.Nance, W. E.; Conneally, M.; Kang, K. W.; Reed, T. E.; Schroder,J.; Rose, S.: Genetic linkage analysis of human hemoglobin variants. Am.J. Hum. Genet. 22: 453-459, 1970.

[68280] 6610.Nichols, M. E.; Rubinstein, P.; Barnwell, J.; Rodriguez de Cordoba,S.; Rosenfield, R. E.: A new human Duffy blood group specificitydefined by a murine monoclonal antibody: immunogenetics and associationwith susceptibility to Plasmodium vivax. J. Exp. Med. 166: 776-785,1987.

[68281] 6611.Olsson, M. L.; Smythe, J. S.; Hansson, C.; Poole, J.;

Mallinson,G.; Jones, J.; Avent, N. D.; Daniels, G.: The Fy(x) phenotype is associated with a missense mutation in the Fy(b) allele predicting Arg89Cys in the Duffy glycoprotein. Brit. J. Haemat. 103: 1184–1191, 1998.

[68282] 6612. Palmer, C. G.; Christian, J. C.; Merritt, A. D.: Partial trisomy 1 due to a 'shift' and probable location of the Duffy (Fy) locus. Am. J. Hum. Genet. 29: 371–377, 1977.

[68283] 6613. Parasol, N.; Reid, M.; Rios, M.; Castilho, L.; Harari, I.; Kosower, N. S.: A novel mutation in the coding sequence of the FY\*B allele of the Duffy chemokine receptor gene is associated with an altered erythrocyte phenotype. Blood 92: 2237–2243, 1998.

[68284] 6614. Pasvol, G.; Wilson, R. J. M.: The interaction of malaria parasites with red blood cells. Brit. Med. Bull. 38: 133–140, 1982.

[68285] 6615. Peiper, S. C.; Wang, Z.; Neote, K.; Martin, A. W.; Showell, H. J.; Conklyn, M. J.; Ogborne, K.; Hadley, T. J.; Lu, Z.; Hesselgesser, J.; Horuk, R.: The Duffy antigen/receptor for chemokines (DARC) is expressed in endothelial cells of Duffy negative individuals who lack the erythrocyte receptor. J. Exp. Med. 181: 1311–1317, 1995.

[68286] 6616. Ritter, H.: Zur formalen Genetik des Duffy-systems. Untersuchung von 247 Familien. Humangenetik 4: 59–61,

1967.

- [68287] 6617. Robson, E. B.; Cook, P. J. L.; Corney, G.; Hopkinson, D. A.; Noades, J.; Cleghorn, T. E.: Linkage data on Rh, PGM, PGD, peptidase C and Fy from family studies. *Ann. Hum. Genet.* 36: 393–399, 1973.
- [68288] 6618. Szabo, M. C.; Soo, K. S.; Zlotnik, A.; Schall, T. J.: Chemokine class differences in binding to the Duffy antigen-erythrocyte chemokine receptor. *J. Biol. Chem.* 270: 25348–25351, 1995.
- [68289] 6619. Tamasauskas, D.; Powell, V.; Saksela, K.; Yazdanbakhsh, K.: A homologous naturally occurring mutation in Duffy and CCR5 leading to reduced receptor expression. *Blood* 97: 3651–3654, 2001.
- [68290] 6620. Tang, T.; Owen, J. D.; Du, J.; Walker, C. L.; Richmond, A.: Molecular cloning and characterization of a mouse gene with homology to the Duffy-antigen receptor for chemokines. *DNA Seq.* 9: 129–143, 1999.
- [68291] 6621. Tournamille, C.; Kim Le Van, C.; Gane, P.; Le Pennec, P. Y.; Roubinet, F.; Babinet, J.; Cartron, J. P.; Colin, Y.: Arg89Cys substitution results in very low membrane expression of the Duffy antigen/receptor for chemokines in Fy(x) individuals. *Blood* 92: 2147–2156, 1998. Note: Erratum. *Blood* 95: 2753 only, 2000.

- [68292] 6622.Tournamille, C.; Le Van Kim, C.; Gane, P.; Cartron, J.-P.; Colin,Y.: Molecular basis and PCR-DNA typing of the Fya/fyb blood group polymorphism. Hum. Genet. 95: 407-410, 1995.
- [68293] 6623.Weitkamp, L. R.: Personal Communication. Rochester, N. Y. 1972.
- [68294] 6624.Zimmerman, P. A.; Woolley, I.; Masinde, G. L.; Miller, S. M.;McNamara, D. T.; Hazlett, F.; Mgone, C. S.; Alpers, M. P.; Genton,B.; Boatin, B. A.; Kazura, J. W.: Emergence of FY\*A(null) in a Plasmodiumvivax-endemic region of Papua New Guinea. Proc. Nat. Acad. Sci. 96:13973-13977, 1999.
- [68295] 6625.Anderson, S. E.; McKenzie, J. L.; McLoughlin, K.; Beard, M. E.J.; Hart, D. N. J.: The inheritance of abnormal sialoglycoproteins found in a Gerbich negative individual. Pathology 18: 407-412, 1986.
- [68296] 6626.Anstee, D. J.; Parsons, S. F.; Ridgwell, K.; Tanner, M. J. A.;Merry, A. H.; Thomson, E. E.; Judson, P. A.; Johnson, P.; Bates, S.;Fraser, I. D.: Two individuals with elliptocytic red cells apparentlylack three minor erythrocyte membrane sialoglycoproteins. Biochem.J. 218: 615-619, 1984.
- [68297] 6627.Barnes, R.; Lewis, T. L. T.: A rare antibody (anti-Ge) causinghemolytic disease of the newborn. Lancet II: 1285-1286, 1961.

- [68298] 6628.Bennett, V.: The membrane skeleton of human erythrocytes and its implications for more complex cells. *Annu. Rev. Biochem.* 54: 273–304, 1985.
- [68299] 6629.Bloomfield, L.; Rowe, G. P.; Green, C.: The Webb (Wb) antigen in South Wales donors. *Hum. Hered.* 36: 352–356, 1986.
- [68300] 6630.Booth, P. B.; McLoughlin, K.: The Gerbich blood group system, especially in Melanesians. *Vox Sang.* 22: 73–84, 1972.
- [68301] 6631.Cartron, J.-P.; Colin, Y.; Kudo, S.; Fukuda, M.: Molecular genetics of human erythrocyte sialoglycoproteins A, B, C, and D. In: Harris, J. R.: *Erythroid Cells. Blood Cell Biochemistry*. New York: Plenum Press (pub.) 1: 1990. Pp. 299–335.
- [68302] 6632.Chang, S.; Reid, M. E.; Conboy, J.; Kan, Y. W.; Mohandas, N.: Molecular characterization of erythrocyte glycophorin C variants. *Blood* 77:644–648, 1991.
- [68303] 6633.Ming, G.; Wong, S. T.; Henley, J.; Yuan, X.; Song, H.; Spitzer, N. C.; Poo, M.: Adaptation in the chemotactic guidance of nerve growth cones. *Nature* 417: 411–418, 2002.
- [68304] 6634.Ozcelik, T.; Rosenthal, A.; Francke, U.: Chromosomal mapping of brain-derived neurotrophic factor and neu-

rotrophin-3 genes in man and mouse. Genomics 10: 569–575, 1991.

[68305] 6635. Zuccato, C.; Ciammola, A.; Rigamonti, D.; Leavitt, B. R.; Goffredo, D.; Conti, L.; MacDonald, M. E.; Friedlander, R. M.; Silani, V.; Hayden, M. R.; Timmusk, T.; Sipione, S.; Cattaneo, E.: Loss of huntingtin-mediated BDNF gene transcription in Huntington's disease. Science 293: 493–498, 2001.

[68306] 6636. Ichimura-Ohshima, Y.; Morii, K.; Ichimura, T.; Araki, K.; Takahashi, Y.; Isobe, T.; Minoshima, S.; Fukuyama, R.; Shimizu, N.; Kuwano, R.: cDNA cloning and chromosome assignment of the gene for human brain 14–3–3 protein eta chain. J. Neurosci. Res. 31: 600–605, 1992.

[68307] 6637. Muratake, T.; Hayashi, S.; Ichikawa, T.; Kumanishi, T.; Ichimura, Y.; Kuwano, R.; Isobe, T.; Wang, Y.; Minoshima, S.; Shimizu, N.; Takahashi, Y.: Structural organization and chromosomal assignment of the human 14–3–3-eta chain gene (YWHAH). Genomics 36: 63–69, 1996.

[68308] 6638. Tommerup, N.; Leffers, H.: Assignment of the human genes encoding 14–3–3 eta (YWHAH) to 22q12, 14–3–3 zeta (YWHAZ) to 2p25.1–p25.2, and 14–3–3 beta (YWHA B) to 20q13.1 by in situ hybridization. Genomics



33:149–150, 1996.

- [68309] 6639.Watanabe, M.; Isobe, T.; Ichimura, T.; Kuwano, R.; Takahashi, Y.;Kondo, H.; Inoue, Y.: Molecular cloning of rat cDNAs for the zetaand theta subtypes of 14–3–3 protein and differential distributionsof their mRNAs in the brain. *Molec. Brain Res.* 25: 113–121, 1994.
- [68310] 6640.Yaffe, M. B.; Rittinger, K.; Volinia, S.; Caron, P. R.; Aitken,A.; Leffers, H.; Gamblin, S. J.; Smerdon, S. J.; Cantley, L. C.:The structural basis for 14–3–3:phosphopeptide binding specificity. *Cell* 91:961–971, 1997.
- [68311] 6641.Zupan, L. A.; Steffens, D. L.; Berry, C. A.; Landt, M.; Gross,R. W.: Cloning and expression of a human 14–3–3 protein mediatingphospholipolysis. *J. Biol. Chem.* 267: 8707–8710, 1992.
- [68312] 6642.Ben–Yosef, T.; Eden, A.; Benvenisty, N.: Characterization of murineBCAT genes: Bcat1, a c–Myc target, and its homolog, Bcat2. *MammalianGenome* 9: 595–597, 1998.
- [68313] 6643.Benvenisty, N.; Leder, A.; Kuo, A.; Leder, P.: An embryonicallyexpressed gene is a target for c–Myc regulation via the c–Myc–bindingsequence. *Genes Dev.* 6: 2513–2523, 1992.
- [68314] 6644.Eden, A.; Simchen, G.; Benvenisty, N.: Two yeast ho–

mologs of ECA39, a target for c-Myc regulation, code for cytosolic and mitochondrial branched-chain amino acid aminotransferases. *J. Biol. Chem.* 271:20242–20245, 1996.

[68315] 6645. Jones, C.; Moore, E. E.: Isolation of mutants lacking branched-chain amino acid transaminase. *Somat. Cell Genet.* 2: 235–243, 1976.

[68316] 6646. Jones, C.; Moore, E. E.: Assignment of the human gene complementing the auxotrophic marker TRANS-minus (BCT1) to chromosome 12. (Abstract) *Cytogenet. Cell Genet.* 25: 168 only, 1979.

[68317] 6647. Jones, C.; Moore, E. E.: Localization of a gene which complements branched-chain amino acid transaminase deficiency to the short arm of human chromosome 12. *Hum. Genet.* 66: 206–211, 1984.

[68318] 6648. Naylor, S. L.; Shows, T. B.: Branched-chain amino-transferase genes (BCT-1 and BCT-2) assigned to human chromosomes 12 and 19 using alpha-ketoacid selection media. (Abstract) *Cytogenet. Cell Genet.* 25: 191–192, 1979.

[68319] 6649. Naylor, S. L.; Shows, T. B.: Branched-chain amino-transferase deficiency in Chinese hamster cells complemented by two independent genes on human chromo-

somes 12 and 19. Somat. Cell Genet. 6: 641–652, 1980.

[68320] 6650.Schuldiner, O.; Eden, A.; Ben-Yosef, T.; Yanuka, O.; Simchen, G.;Benvenisty, N.: ECA39, a conserved gene regulated by c-Myc in mice,is involved in G1/S cell cycle regulation in yeast. Proc. Nat. Acad.Sci. 93: 7143–7148, 1996.

[68321] 6651.Tanaka, K.; Rosenberg, L. E.: Disorders of branched chain aminoacid and organic acid metabolism.In: Stanbury, J. B.; Wyngaarden,J. B.; Fredrickson, D. S.; Goldstein, J. L.; Brown, M. S.: The MetabolicBasis of Inherited Disease. New York: McGraw–Hill (pub.) (5thed.): 1983. Pp. 450–451.

[68322] 6652.Bassett, D. E., Jr.; Boguski, M. S.; Spencer, F.; Reeves, R.; Kim,S.; Weaver, T.; Hieter, P.: Genome cross-referencing and XREFdb:implications for the identification and analysis of genes mutatedin human disease. Nature Genet. 15: 339–344, 1997.

[68323] 6653.Goebel, H. H.; Anderson, J. R.; Hubner, C.; Oexle, K.; Warlo, I.: Congenital myopathy with excess of thin myofilaments. NeuromuscularDisord. 7: 160–168, 1997.

[68324] 6654.Cattanach, B. M.; Barr, J. A.; Evans, E. P.; Burtenshaw, M.; Beechey,C. V.; Leff, S. E.; Brannan, C. I.; Copeland, N. G.; Jenkins, N. A.;Jones, J.: A candidate mouse model for

Prader–Willi syndrome which shows an absence of Snrpn expression. *Nature Genet.* 2: 270–274, 1992.

- [68325] 6655. Colin, Y.; Rahuel, C.; London, J.; Romeo, P. H.; d'Auriol, L.; Galibert, F.; Cartron, J.–P.: Isolation of cDNA clones and complete amino acid sequence of human erythrocyte glycophorin C. *J. Biol. Chem.* 261: 229–233, 1986.
- [68326] 6656. Accolla, R. S.; Gross, N.; Carrel, S.; Corte, G.: Distinct forms of both alpha and beta subunits are present in the human Ia molecular pool. *Proc. Nat. Acad. Sci.* 78: 4549–4551, 1981.
- [68327] 6657. Kingsmore, S. F.; Snoddy, J.; Choubey, D.; Lengyel, P.; Seldin, M. F.: Physical mapping of a family of interferon-activated genes, serum amyloid P-component, and alpha-spectrin on mouse chromosome 1. *Immunogenetics* 30: 169–174, 1989.
- [68328] 6658. Thomas, K. R.; Folger, K. R.; Capecchi, M. R.: *Cell* 44: 419–428, 1986.
- [68329] 6659. Watanabe, K.; Kessler, C. A.; Bachurski, C. J.; Kanda, Y.; Richardson, B. D.; Stanek, J.; Handwerger, S.; Brar, A. K.: Identification of a decidua-specific enhancer on the human prolactin gene with two critical activator protein 1 (AP-1) binding sites. *Molec. Endocr.* 15: 638–653, 2001.
- [68330] 6660. Guillon, H.; de Massy, B.: An initiation site for mei–

otic crossing-over and gene conversion in the mouse. *Nature Genet.* 32: 296–299, 2002.

- [68331] 6661. Irsch, J.; Nitsch, S.; Hansmann, M.-L.; Rajewsky, K.; Tesch, H.; Diehl, V.; Jox, A.; Kuppers, R.; Radbruch, A.: Isolation of viable Hodgkin and Reed–Sternberg cells from Hodgkin disease tissues. *Proc. Nat. Acad. Sci.* 95: 10117–10122, 1998.
- [68332] 6662. Hanby, A. M.; Poulsom, R.; Singh, S.; Elia, G.; Jeffery, R. E.; Wright, N. A.: Spasmolytic polypeptide is a major antral peptide: distribution of the trefoil peptides human spasmolytic polypeptide and pS2 in the stomach. *Gastroenterology* 105: 1110–1116, 1993.
- [68333] 6663. Blankstein, J.; Faiman, C.; Reyes, F. I.; Schroeder, M. L.; Winter, J. S. D.: Adult-onset familial adrenal 21-hydroxylase deficiency. *Am. J. Med.* 68: 441–448, 1980.
- [68334] 6664. Hoeg, J. M.; Osborne, J. C., Jr.; Gregg, R. E.; Brewer, H. B., Jr.: Initial diagnosis of lipoprotein lipase deficiency in a 75-year-old man. *Am. J. Med.* 75: 889–892, 1983.
- [68335] 6665. Batischev, A. I.; Chernyak, N. B.; Torakev, Y. N.: Detection of a new abnormal variant of glucose-6-phosphate dehydrogenase in human red cells. *Bulleten Eksperimentalnoi Biologii i Meditsiny* 84: 728–731, 1977.

- [68336] 6666.Parolini, S.; Bottino, C.; Falco, M.; Augugliaro, R.; Giliani,S.; Franceschini, R.; Ochs, H. D.; Wolf, H.; Bonnefoy, J.-Y.; Biassoni,R.; Moretta, L.; Notarangelo, L. D.; Moretta, A.: X-linked lymphoproliferativedisease: 2B4 molecules displaying inhibitory rather than activatingfunction are responsible for the inability of natural killer cellsto kill Epstein-Barr virus-infected cells. *J. Exp. Med.* 192: 337-346,2000.
- [68337] 6667.Hawkins, J. R.; Taylor, A.; Goodfellow, P. N.; Migeon, C. J.;Smith, K. D.; Berkovitz, G. D.: Evidence for increased prevalenceof SRY mutations in XY females with complete rather than partial gonadaldysgenesis. *Am. J. Hum. Genet.* 51: 979-984, 1992.
- [68338] 6668.Burchell, B.; Nebert, D. W.; Nelson, D. R.; Bock, K. W.; Iyanagi,T.; Jansen, P. L. M.; Lancet, T.; Mulder, G. J.; Chowdhury, J. R.;Siest, G.; Tephly, T. R.; Mackenzie, P. I.: The UDP-glucuronosyltransferasegene superfamily: suggested nomenclature based on evolutionary divergence. *DNACell Biol.* 10: 487-494, 1991.
- [68339] 6669.Harrington, J. J.; Lieber, M. R.: The characterization of a mammalianDNA structure-specific endonuclease. *EMBO J.* 13: 1235-1246, 1994.
- [68340] 6670.Yi, T.; Cleveland, J. L.; Ihle, J. N.: Identification of

novel protein tyrosine phosphatases of hematopoietic cells by polymerase chain reaction amplification. Blood 78: 2222–2228, 1991.

[68341] 6671. Amano, M.; Mukai, H.; Ono, Y.; Chihara, K.; Matsui, T.; Hamajima, Y.; Okawa, K.; Iwamatsu, A.; Kaibuchi, K.: Identification of a putative target for rho as the serine–threonine kinase protein kinase N. Science 271:648–651, 1996.

[68342] 6672. Gorlatov, S. N.; Stadtman, T. C.: Human thioredoxin reductase from HeLa cells: selective alkylation of seleno–cysteine in the protein inhibits enzyme activity and reduction with NADPH influences affinity to heparin. Proc. Nat. Acad. Sci. 95: 8520–8525, 1998.

[68343] 6673. ten Dijke, P.; Franzen, P.; Yamashita, H.; Ichijo, H.; Heldin, C. H.; Miyazono, K.: Serine/threonine kinase receptors. Prog. Growth Factor Res. 5: 55–72, 1994.

[68344] 6674. MacGrogan, D.; Levy, A.; Bova, G. S.; Isaacs, W. B.; Bookstein, R.: Structure and methylation–associated silencing of a gene within a homozygously deleted region of human chromosome band 8p22. Genomics 35:55–65, 1996.

[68345] 6675. McPherron, A. C.; Lee, S.–J.: Suppression of body fat accumulation in myostatin–deficient mice. J. Clin. Invest.

109: 595–601, 2002.

- [68346] 6676.Klein, R. D.; Sherman, D.; Ho, W.–H.; Stone, D.; Bennett, G. L.;Moffat, B.; Vandlen, R.; Simmons, L.; Gu, Q.; Hongo, J.–A.; Devaux,B.; Poulsen, K.; Armanini, M.; Nozaki, C.; Asai, N.; Goddard, A.;Phillips, H.; Henderson, C. E.; Takahashi, M.; Rosenthal, A.: A GPI–linkedprotein that interacts with Ret to form a candidate neurturin receptor. *Nature* 387:717–721, 1997.
- [68347] 6677.Masuda, H.; Tanaka, K.; Takagi, M.; Ohgami, K.; Sakamaki, T.; Shibata,N.; Takahashi, K.: Molecular cloning and characterization of humannon–smooth muscle calponin. *J. Biochem.* 120: 415–424, 1996.
- [68348] 6678.Vlangos, C. N.; Das, P.; Patel, P. I.; Elsea, S. H.: Assignmentof developmentally regulated GTP–binding protein (DRG2) to human chromosomeband 17p11.2 with somatic cell hybrids and localization to the Smith–Magenissyndrome critical interval. *Cytogenet. Cell Genet.* 88: 283–285,2000.
- [68349] 6679.Peyrard, M.: Personal Communication. Stockholm, Sweden 1/7/1999.
- [68350] 6680.Ware, F. E.; Lehrman, M. A.: Expression cloning of a novel suppressorof the Lec15 and Lec35 glycosylation mutations of Chinese hamsterovary cells. *J. Biol. Chem.*



271: 13935–13938, 1996.

- [68351] 6681.Lowe, D. G.; Camerato, T. R.; Goeddel, D. V.: cDNA sequence of the human atrial natriuretic peptide clearance receptor. *Nucleic Acids Res.* 18: 3412 only, 1990.
- [68352] 6682.Porter, J. G.; Arfsten, A.; Fuller, F.; Miller, J. A.; Gregory, L. C.; Lewicki, J. A.: Isolation and functional expression of the human atrial natriuretic peptide clearance receptor cDNA. *Biochem. Biophys. Res. Commun.* 171: 796–803, 1990.
- [68353] 6683.Suda, M.; Ogawa, Y.; Tanaka, K.; Tamura, N.; Yasoda, A.; Takigawa, T.; Uehira, M.; Nishimoto, H.; Itoh, H.; Saito, Y.; Shiota, K.; Nakao, K.: Skeletal overgrowth in transgenic mice that overexpress brain natriuretic peptide. *Proc. Nat. Acad. Sci.* 95: 2337–2342, 1998.
- [68354] 6684.Boehm, J.; Orth, T.; Van Nguyen, P.; Soling, H. D.: Systemic lupus erythematosus is associated with increased auto-antibody titers against calreticulin and grp94, but calreticulin is not the Ro/SS-A antigen. *Europ. J. Clin. Invest.* 24: 248–257, 1994.
- [68355] 6685.Burns, K.; Duggan, B.; Atkinson, E. A.; Famulski, K. S.; Nemer, M.; Bleackley, R. C.; Michalak, M.: Modulation of gene expression by calreticulin binding to the glucocorticoid receptor. *Nature* 367:476–480, 1994.

- [68356] 6686.Telen, M. J.; Whitsett, C. F.: Erythrocyte acetylcholinesterase bears the Cartwright blood group antigens. (Abstract) Clin. Res. 40:170A only, 1992.
- [68357] 6687.Wjst, M.; Fischer, G.; Immervoll, T.; Jung, M.; Saar, K.; Rueschendorf, F.; Reis, A.; Ulbrecht, M.; Gomolka, M.; Weiss, E. H.; Jaeger, L.; Nickel, R.; and 14 others: A genome-wide search for linkage to asthma. Genomics 58: 1–8, 1999.
- [68358] 6688.Kalachikov, S.; Evgrafov, O.; Ross, B.; Winawer, M.; Barker–Cummings, C.; Boneschi, F. M.; Choi, C.; Morozov, P.; Das, K.; Teplitskaya, E.; Yu, A.; Cayanis, E.; Penchaszadeh, G.; Kottmann, A. H.; Pedley, T. A.; Hauser, W. A.; Ottman, R.; Gilliam, T. C.: Mutations in LIG1 cause autosomal–dominant partial epilepsy with auditory features. Nature Genet. 30: 335–341, 2002.
- [68359] 6689.Privalsky, M. L.; Ralston, R.; Bishop, J. M.: The membrane glycoprotein encoded by the retroviral oncogene v-erb-B is structurally related to tyrosine–specific protein kinases. Proc. Nat. Acad. Sci. 81:704–707, 1984.
- [68360] 6690.Gordon, M. S.; Kato, R. M.; Lansigan, F.; Thompson, A. A.; Wall, R.; Rawlings, D. J.: Aberrant B cell receptor signaling from B29(Ig–beta, CD79b) gene mutations of chronic lymphocytic leukemia B cells. Proc. Nat. Acad. Sci.

97: 5504–5509, 2000.

- [68361] 6691. Kuida, K.; Lippke, J. A.; Ku, G.; Harding, M. W.; Livingston, D. J.; Su, M. S.-S.; Flavell, R. A.: Altered cytokine export and apoptosis in mice deficient in interleukin-1-beta converting enzyme. *Science* 267:2000–2003, 1995.
- [68362] 6692. Morissette, J.; Rheaume, E.; Leblanc, J.-F.; Luu-The, V.; Labrie, F.; Simard, J.: Genetic linkage mapping of HSD3B1 and HSD3B2 encoding human types I and II 3-beta-hydroxysteroid dehydrogenase/delta-5-delta-4-isomerase close to D1S514 and the centromeric D1Z5 locus. *Cytogenet. Cell Genet.* 69: 59–62, 1995.
- [68363] 6693. Morrison, N.; Nickson, D. A.; McBride, M. W.; Mueller, U. W.; Boyd, E.; Sutcliffe, R. G.: Regional chromosomal assignment of human 3-beta-hydroxy-5-enesteroid dehydrogenase to 1p13.1 by non-isotopic in situ hybridisation. *Hum. Genet.* 87: 223–225, 1991.
- [68364] 6694. Race, R. R.; Sanger, R.: *Blood Groups in Man*. Oxford: Blackwell (pub.) (6th ed.): 1975. Pp. 92–138.
- [68365] 6695. Siebert, P. D.; Fukuda, M.: Molecular cloning of a human glycophorin B cDNA: nucleotide sequence and ge-

nomie relationship to glycophorinA. Proc. Nat. Acad. Sci. 84: 6735–6739, 1987.

- [68366] 6696. Kumar, B. V.; Aleman–Gomez, J. A.; Colwell, N.; Lopez–Candales, A.; Bosner, M. S.; Spilburg, C. A.; Lowe, M.; Lange, L. G.: Structure of the human pancreatic cholesterol esterase gene. Biochemistry 31:6077–6081, 1992.
- [68367] 6697. Lidberg, U.; Nilsson, J.; Stromberg, K.; Stenman, G.; Sahlin, P.; Enerback, S.; Bjursell, G.: Genomic organization, sequence analysis, and chromosomal localization of the human carboxyl ester lipase (CEL) gene and a CEL–like (CELL) gene. Genomics 13: 630–640, 1992.
- [68368] 6698. Galland, F.; Stefanova, M.; Pirisi, V.; Birnbaum, D.: Characterization of a murine glyceraldehyde–3–phosphate dehydrogenase pseudogene. Biochimie 72:759–762, 1990.
- [68369] 6699. Lidmer, A.–S.; Kannius, M.; Lundberg, L.; Bjursell, G.; Nilsson, J.: Molecular cloning and characterization of the mouse carboxylester lipase gene and evidence for expression in the lactating mammary gland. Genomics 29: 115–122, 1995.
- [68370] 6700. Nilsson, J.; Hellquist, M.; Bjursell, G.: The human carboxyl ester lipase–like (CELL) gene is ubiquitously expressed and contains a hypervariable region. Genomics 17:

416–422, 1993.

- [68371] 6701. Taylor, A. K.; Zambaux, J. L.; Klisak, I.; Mohandas, T.; Sparkes, R. S.; Schotz, M. C.; Lusi, A. J.: Carboxyl-ester lipase: a highly polymorphic locus on chromosome 9qter. *Genomics* 10: 425–431, 1991.
- [68372] 6702. Vanin, E. F.: Processed pseudogenes: characteristics and evolution. *Annu. Rev. Genet.* 19: 253–272, 1985.
- [68373] 6703. Gardell, S. J.; Craik, C. S.; Clauser, E.; Goldsmith, E. J.; Stewart, C. B.; Graf, M.; Rutter, W. J.: A novel rat carboxypeptidase, CPA2: characterization, molecular cloning, and evolutionary implications on substrate specificity in the carboxypeptidase gene family. *J. Biol. Chem.* 263: 17828–17836, 1988.
- [68374] 6704. Lowe, N.; Edwards, Y. H.; Edwards, M.; Butterworth, P. H. W.: Physical mapping of the human carbonic anhydrase gene cluster on chromosome 8. *Genomics* 10: 882–888, 1991.
- [68375] 6705. Marriq, C.; Gulian, J. M.; Laurent, G.: Cleavage by cyanogen bromide of carbonic anhydrase from human erythrocyte B. *Biochim. Biophys. Acta* 221: 662–664, 1970.
- [68376] 6706. Moore, M. J.; Deutsch, H. F.; Ellis, F. R.: Human carbonic anhydrase. IX. Inheritance of variant erythrocyte forms. *Am. J. Hum. Genet.* 25: 29–35, 1973.

- [68377] 6707.Omoto, K.: Carbonic anhydrase-I polymorphism in a Philippineaboriginal population. Am. J. Hum. Genet. 31: 747-750, 1979.
- [68378] 6708.Omoto, K.; Ueda, S.; Goriki, K.; Takahashi, N.; Misawa, S.; Pagaran,I. G.: Population genetic studies of the Philippine Negritos. III.Identification of the carbonic anhydrase-1 variant with CA(1) Guam. Am.J. Hum. Genet. 33: 105-111, 1981.
- [68379] 6709.Tashian, R. E.: The esterases and carbonic anhydrases of humanerythrocytes.In: Yunis, J. J.: Biochemical Methods in Red Cell Genetics. New York: Academic Press (pub.) 1969. Pp. 307-336.
- [68380] 6710.Tashian, R. E.; Carter, N. D.: Biochemical genetics of carbonicanhydrase. Adv. Hum. Genet. 7: 1-56, 1976.
- [68381] 6711.Tashian, R. E.; Goodman, M.; Headings, V. E.; Desimone, J.; Ward,R. H.: Genetic variation and evolution in the red cell carbonic anhydraseisozymes of Macaque monkeys. Biochem. Genet. 5: 183-200, 1971.
- [68382] 6712.Tashian, R. E.; Plato, C. C.; Shows, T. B.: Inherited variantof erythrocyte carbonic anhydrase in Micronesians from Guam and Saipan. Science 140:53-54, 1963.
- [68383] 6713.Wagner, L. E.; Venta, P. J.; Tashian, R. E.: A human carbonicanhydrase I deficiency appears to be caused by a

destabilizing aminoacid substitution (246arg-to-his).

Isozyme Bull. 24: 35 only, 1991.

- [68384] 6714. Avramopoulos, D.; Cox, T.; Forrest, G. L.; Chakravarti, A.; Antonarakis, S. E.: Linkage mapping of the carbonyl reductase (CBR) gene on human chromosome 21 using a DNA polymorphism in the 3-prime untranslated region. *Genomics* 13: 447–448, 1992.
- [68385] 6715. Forrest, G. L.; Akman, S.; Krutzik, S.; Paxton, R. J.; Sparkes, R. S.; Doroshov, J.; Felsted, R. L.; Glover, C. J.; Mohandas, T.; Bachur, N. R.: Induction of a human carbonyl reductase gene located on chromosome 21. *Biochim. Biophys. Acta* 1048: 149–155, 1990.
- [68386] 6716. Lemieux, N.; Malfoy, B.; Forrest, G. L.: Human carbonyl reductase (CBR) localized to band 21q22.1 by high-resolution fluorescence in situ hybridization displays gene dosage effects in trisomy 21 cells. *Genomics* 15: 169–172, 1993.
- [68387] 6717. Watanabe, K.; Sugawara, C.; Ono, A.; Fukuzumi, Y.; Itakura, S.; Yamazaki, M.; Tashiro, H.; Osoegawa, K.; Soeda, E.; Nomura, T.: Mapping of a novel human carbonyl reductase, CBR3, and ribosomal pseudogene to human chromosome 21q22.2. *Genomics* 52: 95–100, 1998.
- [68388] 6718. Wei, J.; Dlouhy, S. R.; Hara, A.; Ghetti, B.; Hodes, M.

E.: Cloning a cDNA for carbonyl reductase (Cbr) from mouse cerebellum: murine genes that express Cbr map to chromosomes 16 and 11. *Genomics* 34:147–148, 1996.

[68389] 6719. Wermuth, B.; Bohren, K. M.; Heinemann, G.; von Wartburg, J.-P.; Gabbay, K. H.: Human carbonyl reductase: nucleotide sequence analysis of a cDNA and amino acid sequence of the encoded protein. *J. Biol. Chem.* 263: 16185–16188, 1988.

[68390] 6720. Yang-Feng, T. L.; Floyd-Smith, G.; Nemer, M.; Drouin, J.; Francke, U.: The pronatriodilatin gene is located on the distal short arm of human chromosome 1 and on mouse chromosome 4. *Am. J. Hum. Genet.* 37:1117–1128, 1985.

[68391] 6721. Ferrari, P.; Weidmann, P.; Ferrier, C.; Dietler, R.; Hollmann, R.; Pisoni, R. J.; Wey, J.; Shaw, S.: Dysregulation of atrial natriuretic factor in hypertension-prone man. *J. Clin. Endocr. Metab.* 71: 944–951, 1990.

[68392] 6722. Kishimoto, I.; Rossi, K.; Garbers, D. L.: A genetic model provides evidence that the receptor for atrial natriuretic peptide (guanylyl cyclase-A) inhibits cardiac ventricular myocyte hypertrophy. *Proc. Nat. Acad. Sci.* 98: 2703–2706, 2001.

[68393] 6723. Lowe, D. G.; Klisak, I.; Sparkes, R. S.; Mohandas, T.;



Goeddel,D. V.: Chromosomal distribution of three members of the human natriureticpeptide receptor/guanylyl cyclase gene family. *Genomics* 8: 304–312,1990.

[68394] 6724.AbdAlla, S.; Lothar, H.; Quitterer, U.: AT(1)–receptor heterodimersshow enhanced G–protein activation and altered receptor sequestration. *Nature* 407:94–98, 2000.

[68395] 6725.Bergsma, D. J.; Ellis, C.; Kumar, C.; Nuthulaganti, P.; Kersten,H.; Elshourbagy, N.; Griffin, E.; Stadel, J. M.; Aiyar, N.: Cloningand characterization of a human angiotensin II type 1 receptor. *Biochem.Biophys. Res. Commun.* 183: 989–995, 1992.

[68396] 6726.Bonnardeaux, A.; Davies, E.; Jeunemaitre, X.; Fery, I.; Charru,A.; Clauser, E.; Tiret, L.; Cambien, F.; Corvol, P.; Soubrier, F.: Angiotensin II type 1 receptor gene polymorphisms in human essentialhypertension. *Hypertension* 24: 63–69, 1994.

[68397] 6727.Curnow, K. M.; Pascoe, L.; White, P. C.: Genetic analysis of thehuman type–1 angiotensin II receptor. *Molec. Endocr.* 6: 1113–1118,1992.

[68398] 6728.Elton, T. S.; Stephan, C. C.; Taylor, G. R.; Kimball, M. G.; Martin,M. M.; Durand, J. N.; Oparil, S.: Isolation of two distinct typel angiotensin II receptor genes. *Biochem. Biophys. Res. Commun.* 184:1067–1073, 1992.

- [68399] 6729.Furuta, H.; Guo, D.-F.; Inagami, T.: Molecular cloning and sequencing of the gene encoding human angiotensin II type 1 receptor. *Biochem. Biophys. Res. Commun.* 183: 8-13, 1992.
- [68400] 6730.Gemmill, R. M.; Drabkin, H. A.: Report of The Second International Workshop on Human Chromosome 3 Mapping. *Cytogenet. Cell Genet.* 57:162-166, 1991.
- [68401] 6731.Guo, D.-F.; Furuta, H.; Mizukoshi, M.; Inagami, T.: The genomic organization of human angiotensin II type 1 receptor. *Biochem. Biophys. Res. Commun.* 200: 313-319, 1994.
- [68402] 6732.Harada, K.; Komuro, I.; Hayashi, D.; Sugaya, T.; Murakami, K.; Yazaki, Y.: Angiotensin II type 1a receptor is involved in the occurrence of reperfusion arrhythmias. *Circulation* 97: 315-317, 1998.
- [68403] 6733.Haywood, G. A.; Gullestad, L.; Katsuya, T.; Hutchinson, H. G.; Pratt, R. E.; Horiuchi, M.; Fowler, M. B.: AT(1) and AT(2) angiotensin receptor gene expression in human heart failure. *Circulation* 95:1201-1206, 1997.
- [68404] 6734.Herzig, T. C.; Jobe, S. M.; Aoki, H.; Molkentin, J. D.; Cowley, A. W., Jr.; Izumo, S.; Markham, B. E.: Angiotensin II type-1a receptor gene expression in the heart: AP-1 and GATA-4 participate in the response to pressure overload.

Proc. Nat. Acad. Sci. 94: 7543–7548, 1997.

- [68405] 6735. Ito, M.; Oliverio, M. I.; Mannon, P. J.; Best, C. F.; Maeda, N.; Smithies, O.; Coffman, T. M.: Regulation of blood pressure by type 1A angiotensin II receptor gene. Proc. Nat. Acad. Sci. 92: 3521–3525, 1995.
- [68406] 6736. Iyer, S. N.; Lu, D.; Katovich, M. J.; Raizada, M. K.: Chronic control of high blood pressure in the spontaneously hypertensive rat by delivery of angiotensin type 1 receptor antisense. Proc. Nat. Acad. Sci. 93: 9960–9965, 1996.
- [68407] 6737. Konishi, H.; Kuroda, S.; Inada, Y.; Fujisawa, Y.: Novel subtype of human angiotensin II type 1 receptor: cDNA cloning and expression. Biochem. Biophys. Res. Commun. 199: 467–474, 1994.
- [68408] 6738. Martens, J. R.; Reaves, P. Y.; Lu, D.; Katovich, M. J.; Berecek, K. H.; Bishop, S. P.; Raizada, M. K.; Gelband, C. H.: Prevention of renovascular and cardiac pathophysiological changes in hypertension by angiotensin II type 1 receptor antisense gene therapy. Proc. Nat. Acad. Sci. 95: 2664–2669, 1998.
- [68409] 6739. Martin, M. M.; Willardson, B. M.; Burton, G. F.; White, C. R.; McLaughlin, J. N.; Bray, S. M.; Ogilvie, J. W., Jr.; Elton, T. S.: Human angiotensin II type 1 receptor isoforms en-

coded by messengerRNA splice variants are functionally distinct. *Molec. Endocr.* 15:281–293, 2001.

- [68410] 6740.Mauzy, C. A.; Hwang, O.; Egloff, A. M.; Wu, L.–H.; Chung, F.–Z.: Cloning, expression, and characterization of a gene encoding thehuman angiotensin II type 1A receptor. *Biochem. Biophys. Res. Commun.* 186:277–284, 1992.
- [68411] 6741.Murphy, T. J.; Alexander, R. W.; Griendling, K. K.; Runge, M.S.; Bernstein, K. E.: Isolation of a cDNA encoding the vascular type–1angiotensin II receptor. *Nature* 351: 233–236, 1991.
- [68412] 6742.Paradis, P.; Dali–Youcef, N.; Paradis, F. W.; Thibault, G.; Nemer,M.: Overexpression of angiotensin II type I receptor in cardiomyocytesinduces cardiac hypertrophy and remodeling. *Proc. Nat. Acad. Sci.* 97:931–936, 2000.
- [68413] 6743.Sasaki, K.; Murohara, T.; Ikeda, H.; Sugaya, T.; Shimada, T.;Shintani, S.; Imaizumi, T.: Evidence for the importance of angiotensinII type 1 receptor in ischemia–induced angiogenesis. *J. Clin. Invest.* 109:603–611, 2002.
- [68414] 6744.Oliverio, M. I.; Kim, H–S.; Ito, M.; Le, T.; Audoly, L.; Best,C. F.; Hiller, S.; Kluckman, K.; Maeda, N.; Smithies, O.; Coffman,T. M.: Reduced growth, abnormal kidney structure, and type 2 (AT2)angiotensin receptor–mediated

blood pressure regulation in mice lacking both AT1A and AT1B receptors for angiotensin II. *Proc. Nat. Acad.Sci.* 95: 15496–15501, 1998.

[68415] 6745.Sasaki, K.; Yamano, Y.; Bardhan, S.; Iwai, N.; Murray, J. J.; Hasegawa, M.; Matsuda, Y.; Inagami, T.: Cloning and expression of a complementary DNA encoding a bovine adrenal angiotensin II type-1 receptor. *Nature* 351: 230–233, 1991.

[68416] 6746.Scott, A. F.: Personal Communication. Baltimore, Md. 3/20/2001.

[68417] 6747.Szpirer, C.; Riviere, M.; Szpirer, J.; Levan, G.; Guo, D. F.; Iwai, N.; Inagami, T.: Chromosomal assignment of human and rat hypertension candidate genes: type 1 angiotensin II receptor genes and the SA gene. *J.Hypertension* 11: 919–925, 1993.

[68418] 6748.Takayanagi, R.; Ohnaka, K.; Sakai, Y.; Nakao, R.; Yanase, T.; Haji, M.; Inagami, T.; Furuta, H.; Gou, D.-F.; Nakamura, M.; Nawata, H.: Molecular cloning, sequence analysis and expression of a cDNA encoding human type-1 angiotensin II receptor. *Biochem. Biophys. Res. Commun.* 183: 910–916, 1992.

[68419] 6749.Tsuchida, S.; Matsusaka, T.; Chen, X.; Okubo, S.; Nishimura, F.; Nishimura, H.; Fogo, A.; Utsunomiya, H.; In-

agami, T.; Ichikawa, I.: Murine double nullizygotes of the angiotensin type 1A and 1B receptorgenes duplicate severe abnormal phenotypes of angiotensinogen nullizygotes. *J.Clin. Invest.* 101: 755–760, 1998.

[68420] 6750.Erba, H. P.; Eddy, R.; Shows, T.; Kedes, L.; Gunning, P.: Structure, chromosome location, and expression of the human gamma-actin gene: differential evolution, location, and expression of the cytoskeletal beta- and gamma-actin genes. *Molec. Cell. Biol.* 8: 1775–1789, 1988.

[68421] 6751.Ueyama, H.; Inazawa, J.; Nishino, H.; Ohkubo, I.; Miwa, T.: FISH localization of human cytoplasmic actin genes ACTB to 7p22 and ACTG1 to 17q25 and characterization of related pseudogenes. *Cytogenet. Cell Genet.* 74: 221–224, 1996.

[68422] 6752.Beggs, A. H.; Byers, T. J.; Knoll, J. H. M.; Boyce, F. M.; Bruns, G. A. P.; Kunkel, L. M.: Cloning and characterization of two human skeletal muscle alpha-actinin genes located on chromosomes 1 and 11. *J. Biol. Chem.* 267: 9281–9288, 1992.

[68423] 6753.Beggs, A. H.; Phillips, H. A.; Kozman, H.; Mulley, J. C.; Wilton, S. D.; Kunkel, L. M.; Laing, N. G.: A (CA)<sub>n</sub> repeat polymorphism for the human skeletal muscle alpha-actinin gene ACTN2 and its localization on the linkage map of

chromosome 1. Genomics 13: 1314–1315, 1992.

- [68424] 6754.Mills, M. A.; Yang, N.; Weinberger, R. P.; Vander Woude, D. L.; Beggs, A. H.; Easteal, S.; North, K. N.: Differential expression of the actin-binding proteins, alpha-actinin-2 and -3, in different species: implications for the evolution of functional redundancy. Hum.Molec. Genet. 10: 1335–1346, 2001.
- [68425] 6755.Attisano, L.; Carcamo, J.; Ventura, F.; Weis, F. M. B.; Massague, J.; Wrana, J. L.: Identification of human activin and TGF-beta type I receptors that form heteromeric kinase complexes with type II receptors. Cell 75:671–680, 1993.
- [68426] 6756.Mathews, L. S.; Vale, W. W.: Expression cloning of an activin receptor, a predicted transmembrane serine kinase. Cell 65: 973–982, 1991.
- [68427] 6757.Matsuzaki, K.; Xu, J.; Wang, F.; McKeehan, W. L.; Krummen, L.; Kan, M.: A widely expressed transmembrane serine/threonine kinase that does not bind activin, inhibin, transforming growth factor beta, or bone morphogenic factor. J. Biol. Chem. 268: 12719–12723, 1993.
- [68428] 6758.Roijer, E.; Miyazono, K.; Astrom, A.-K.; Geurts van Kessel, A.; ten Dijke, P.; Stenman, G.: Chromosomal localization of three human genes encoding members of the

TGF-beta superfamily of type I serine/threonine kinase receptors. *Mammalian Genome* 9: 266–268, 1998.

[68429] 6759. ten Dijke, P.; Ichijo, H.; Franzen, P.; Schulz, P.; Saras, J.; Toyoshima, H.; Heldin, C.-H.; Miyazono, K.: Activin receptor-like kinases: a novel subclass of cell-surface receptors with predicted serine/threonine kinase activity. *Oncogene* 8: 2879–2887, 1993.

[68430] 6760. Chen, M.; Pan, Z.-Q.; Hurwitz, J.: Studies of the cloned 37-kDa subunit of activator 1 (replication factor C) of HeLa cells. *Proc. Nat. Acad. Sci.* 89: 5211–5215, 1992.

[68431] 6761. Ferrari, B.; Pavia, A. A.: Blood group antigens: synthesis of Ss antigenic peptides related to human glycophorin B. *Int. J. Peptide Protein Res.* 28: 456–461, 1986.

[68432] 6762. Huang, C.-H.; Blumenfeld, O. O.: Characterization of a genomic hybrid specifying the human erythrocyte antigen Dantu: Dantu gene is duplicated and linked to a delta glycophorin gene deletion. *Proc. Nat. Acad. Sci.* 85: 9640–9644, 1988.

[68433] 6763. Huang, C.-H.; Johe, K.; Moulds, J. J.; Siebert, P. D.; Fukuda, M.; Blumenfeld, O. O.: Delta-glycophorin (glycophorin B) gene deletion in two individuals homozygous for the S--s--U-- blood group phenotype. *Blood* 70: 1830–1835, 1987.



- [68434] 6764.Kudo, S.; Fukuda, M.: Structural organization of glycophorin A and B genes: glycophorin B gene evolved by homologous recombination at Alu repeat sequences. *Proc. Nat. Acad. Sci.* 86: 4619–4623, 1989.
- [68435] 6765.Marchesi, V. T.; Tillack, T. M.; Jackson, R. L.; Segrest, J. P.; Scott, R. E.: Chemical characterization and surface orientation of the major glycoprotein of the human erythrocyte membrane. *Proc. Nat. Acad. Sci.* 69: 1445–1449, 1972.
- [68436] 6766.Buettner, R.; Schaffler, A.; Arndt, H.; Rogler, G.; Nusser, J.; Zietz, B.; Enger, I.; Hugl, S.; Cuk, A.; Scholmerich, J.; Palitzsch, K.-D.: The trp64arg polymorphism of the beta-3-adrenergic receptor gene is not associated with obesity or type 2 diabetes mellitus in a large population-based Caucasian cohort. *J. Clin. Endocr. Metab.* 83:2892–2897, 1998.
- [68437] 6767.Clement, K.; Vaisse, C.; Manning, B. S. J.; Basdevant, A.; Guy-Grand, B.; Ruiz, J.; Silver, K. D.; Shuldiner, A. R.; Froguel, P.; Strosberg, A. D.: Genetic variation in the beta-3-adrenergic receptor and an increased capacity to gain weight in patients with morbid obesity. *New Eng. J. Med.* 333: 352–354, 1995.
- [68438] 6768.Elbein, S. C.; Hoffman, M.; Barrett, K.; Wegner, K.;

Miles, C.; Bachman, K.; Berkowitz, D.; Shuldiner, A. R.; Leppert, M. F.; Hasstedt, S.: Role of the beta-3-adrenergic receptor locus in obesity and noninsulin-dependent diabetes among members of Caucasian families with a diabetic sibling pair. *J. Clin. Endocr. Metab.* 81: 4422-4427, 1996.

[68439] 6769. Emorine, L. J.; Marullo, S.; Briand-Sutren, M.-M.; Patey, G.; Tate, K.; Delavie-Klutchko, C.; Strosberg, A. D.: Molecular characterization of the human beta-3-adrenergic receptor. *Science* 245: 1118-1121, 1989.

[68440] 6770. Festa, A.; Krugluger, W.; Shnawa, N.; Hopmeier, P.; Haffner, S.M.; Schernthaner, G.: Trp64Arg polymorphism of the beta-3-adrenergic receptor gene in pregnancy: association with mild gestational diabetes mellitus. *J. Clin. Endocr. Metab.* 84: 1695-1699, 1999.

[68441] 6771. Gagnon, J.; Mauriege, P.; Roy, S.; Sjostrom, D.; Chagnon, Y. C.; Dionne, F. T.; Oppert, J.-M.; Perusse, L.; Sjostrom, L.; Bouchard, C.: The trp64arg mutation of the beta-3 adrenergic receptor gene has no effect on obesity phenotypes in the Quebec Family Study and Swedish Obese Subjects cohorts. *J. Clin. Invest.* 98: 2086-2093, 1996.

[68442] 6772. Garcia-Rubi, E.; Starling, R. D.; Tchernof, A.; Matthews, D. E.; Walston, J. D.; Shuldiner, A. R.; Silver, K.;

Poehlman, E. T.; Calles–Escandon, J.: Trp64Arg variant of the beta–3–adrenoceptor and insulin resistance in obese postmenopausal women. *J. Clin. Endocr. Metab.* 83: 4002–4005, 1998.

[68443] 6773. Hoffstedt, J.; Poirier, O.; Thorne, A.; Lonnqvist, F.; Herrmann, S. M.; Cambien, F.; Arner, P.: Polymorphism of the human beta–3–adrenoceptor gene forms a well-conserved haplotype that is associated with moderate obesity and altered receptor function. *Diabetes* 48: 203–205, 1999.

[68444] 6774. Kim–Motoyama, H.; Yasuda, K.; Yamaguchi, T.; Yamada, N.; Katakura, T.; Shuldiner, A. R.; Akanuma, Y.; Ohashi, Y.; Yazaki, Y.; Kadowaki, T.: A mutation of the beta–3–adrenergic receptor is associated with visceral obesity but decreased serum triglyceride. *Diabetologia* 40: 469–472, 1997.

[68445] 6775. Mitchell, B. D.; Blangero, J.; Comuzzie, A. G.; Almasy, L. A.; Shuldiner, A. R.; Silver, K.; Stern, M. P.; McCluer, J. W.; Hixson, J. E.: A paired sibling analysis of the beta–3 adrenergic receptor and obesity in Mexican Americans. *J. Clin. Invest.* 101: 584–587, 1998.

[68446] 6776. Nagase, T.; Aoki, A.; Yamamoto, M.; Yasuda, H.; Kado, S.; Nishikawa, M.; Kugai, N.; Akatsu, T.; Nagata, N.:

Lack of association between the trp64arg mutation in the beta-3-adrenergic receptor gene and obesity in Japanese men: a longitudinal analysis. *J. Clin. Endocr. Metab.* 82:1284–1287, 1997.

- [68447] 6777. Shihara, N.; Yasuda, K.; Moritani, T.; Ue, H.; Adachi, T.; Tanaka, H.; Tsuda, K.; Seino, Y.: The association between Trp64Arg polymorphism of the beta-3-adrenergic receptor and autonomic nervous system activity. *J. Clin. Endocr. Metab.* 84: 1623–1627, 1999.
- [68448] 6778. Van Spronsen, A.; Nahmias, C.; Krief, S.; Briand-Sutren, M.-M.; Strosberg, A. D.; Emorine, L. J.: The promoter and intron/exon structure of the human and mouse beta-3-adrenergic-receptor genes. *Europ. J. Biochem.* 213: 1117–1124, 1993.
- [68449] 6779. Urhammer, S. A.; Hansen, T.; Borch-Johnsen, K.; Pedersen, O.: Studies of the synergistic effect of the trp/arg64 polymorphism of the beta-3-adrenergic receptor gene and the -3826 A-G variant of the uncoupling protein-1 gene on features of obesity and insulin resistance in a population-based sample of 379 young Danish subjects. *J. Clin. Endocr. Metab.* 85: 3151–3154, 2000.
- [68450] 6780. Walston, J.; Silver, K.; Bogardus, C.; Knowler, W. C.; Celi, F. S.; Austin, S.; Manning, B.; Strosberg, A. D.; Stern,

M. P.; Raben, N.; Sorkin, J. D.; Roth, J.; Shuldiner, A. R.:  
Time of onset of non-insulin-dependent diabetes mellitus  
and genetic variation in the beta-  
3-adrenergic-receptor gene. *New Eng. J. Med.* 333:  
343-347, 1995.

[68451] 6781. Walston, J.; Silver, K.; Hilfiker, H.; Andersen, R. E.;  
Seibert, M.; Beamer, B.; Roth, J.; Poehlman, E.; Shuldiner, A.  
R.: Insulin response to glucose is lower in individuals ho-  
mozygous for the arg64 variant of the beta-3-adrenergic  
receptor. *J. Clin. Endocr. Metab.* 85:4019-4022, 2000.

[68452] 6782. Hershfield, M. S.; Mitchell, B. S.: Immunodeficiency  
diseases caused by adenosine deaminase deficiency and  
purine nucleoside phosphorylase deficiency. In: Scriver, C.  
R.; Beaudet, A. L.; Sly, W. S.; Valle, D. (eds.): *The Metabolic  
and Molecular Bases of Inherited Disease*. Vol. 2. New  
York: McGraw-Hill (7th ed.): 1995. Pp. 1725-1768.

[68453] 6783. Hirschhorn, R.: Identification of two new missense  
mutations (R156C and S291L) in two ADA(-) SCID patients  
unusual for response to therapy with partial exchange  
transfusions. *Hum. Mutat.* 1: 166-168, 1992.

[68454] 6784. Hirschhorn, R.: Personal Communication. New York,  
N.Y. 1976.

[68455] 6785. Hirschhorn, R.; Borkowsky, W.; Jiang, C.-K.; Yang, D.

R.; Jenkins,T.: Two newly identified mutations (Thr233Ile and Leu152Met) in partiallyadenosine deaminase–deficient (ADA–) individuals that result in differingbiochemical and metabolic phenotypes. Hum. Genet. 100: 22–29, 1997.

[68456] 6786.Hirschhorn, R.; Chakravarti, V.; Puck, J.; Douglas, S. D.: Homozygosityfor a newly identified missense mutation in a patient with very severecombined immunodeficiency due to adenosine deaminase deficiency (ADA–SCID). Am.J. Hum. Genet. 49: 878–885, 1991.

[68457] 6787.Hirschhorn, R.; Ellenbogen, A.: Genetic heterogeneity in adenosinedeaminase (ADA) deficiency: five different mutations in five new patientswith partial ADA deficiency. Am. J. Hum. Genet. 38: 13–25, 1986.

[68458] 6788.Hirschhorn, R.; Ellenbogen, A.; Tzall, S.: Five missense mutationsat the adenosine deaminase locus (ADA) detected by altered restrictionfragments and their frequency in ADA–patients with severe combinedimmunodeficiency (ADA–SCID). Am. J. Med. Genet. 42: 201–207, 1992.

[68459] 6789.Hirschhorn, R.; Levytska, V.; Parkman, R.: A mutant form of adenosinedeaminase in severe combined immunodeficiency. (Abstract) J. Clin.Invest. 53: 33A, 1974.

[68460] 6790.Hirschhorn, R.; Martiniuk, F.; Roegner–Maniscalco,

V.; Ellenbogen, A.; Perignon, J.-L.; Jenkins, T.: Genetic heterogeneity in partial adenosine deaminase deficiency. *J. Clin. Invest.* 71: 1887–1892, 1983.

- [68461] 6791. Hirschhorn, R.; Papageorgiou, P. S.; Kesarwala, H. H.; Taft, L. T.: Amelioration of neurologic abnormalities after 'enzyme replacement' in adenosine deaminase deficiency. *New Eng. J. Med.* 303: 377–380, 1980.
- [68462] 6792. Hirschhorn, R.; Roegner, V.; Jenkins, T.; Seaman, C.; Piomelli, S.; Borkowsky, W.: Erythrocyte adenosine deaminase deficiency without immunodeficiency: evidence for an unstable mutant enzyme. *J. Clin. Invest.* 64: 1130–1139, 1979.
- [68463] 6793. Hirschhorn, R.; Tzall, S.; Ellenbogen, A.: Hot spot mutations in adenosine deaminase deficiency. *Proc. Nat. Acad. Sci.* 87: 6171–6175, 1990.
- [68464] 6794. Hirschhorn, R.; Tzall, S.; Ellenbogen, A.; Orkin, S. H.: Identification of a point mutation resulting in a heat-labile adenosine deaminase (ADA) in two unrelated children with partial ADA deficiency. *J. Clin. Invest.* 83: 497–501, 1989.
- [68465] 6795. Hirschhorn, R.; Vawter, G. F.; Kirkpatrick, J. A., Jr.; Rosen, F. S.: Adenosine deaminase deficiency: frequency and comparative pathology in autosomally recessive severe combined immunodeficiency. *Clin. Immun. Immunopath.*

14: 107–120, 1979.

- [68466] 6796.Hirschhorn, R.; Yang, D. R.; Israni, A.: An asp8-to-asn substitution results in the adenosine deaminase (ADA) genetic polymorphism (ADA2 allozyme): occurrence on different chromosomal backgrounds and apparent intragenic crossover. *Ann. Hum. Genet.* 58: 1–9, 1994.
- [68467] 6797.Hirschhorn, R.; Yang, D. R.; Israni, A.; Huie, M. L.; Ownby, D.R.: Somatic mosaicism for a newly identified splice-site mutation in a patient with adenosine deaminase-deficient immunodeficiency and spontaneous clinical recovery. *Am. J. Hum. Genet.* 55: 59–68, 1994.
- [68468] 6798.Hirschhorn, R.; Yang, D. R.; Puck, J. M.; Huie, M. L.; Jiang, C.-K.; Kurlandsky, L. E.: Spontaneous in vivo reversion to normal of an inherited mutation in a patient with adenosine deaminase deficiency. *Nature Genet.* 13: 290–295, 1996.
- [68469] 6799.Hong, R.; Galti, R.; Rathbun, J. C.; Good, R. A.: Thymic hypoplasia and thyroid dysfunction. *New Eng. J. Med.* 282: 470–474, 1970.
- [68470] 6800.Honig, J.; Martiniuk, F.; D'Eustachio, P.; Zamfirescu, C.; Desnick, R.; Hirschhorn, K.; Hirschhorn, L. R.; Hirschhorn, R.: Confirmation of the regional localization of



the genes for human acid alpha-glucosidase(GAA) and adenosine deaminase (ADA) by somatic cell hybridization. Ann.Hum. Genet. 48: 49–56, 1984.

[68471] 6801.Hopkinson, D. A.; Cook, P. J. L.; Harris, H.: Further data on the adenosine deaminase (ADA) polymorphism and a report of a new phenotype. Ann.Hum. Genet. 32: 361–368, 1969.

[68472] 6802.Hutton, J. J.; Wiginton, D. A.; Coleman, M. S.; Fuller, S. A.; Limouze, S.; Lampkin, B. C.: Biochemical and functional abnormalities in lymphocytes from an adenosine deaminase-deficient patient during enzyme replacement therapy. J. Clin. Invest. 68: 413–421, 1981.

[68473] 6803.Jenkins, T.; Rabson, A. R.; Nurse, G. T.; Lane, A. B.; Hopkinson, D. A.: Deficiency of adenosine deaminase not associated with severe combined immunodeficiency. J. Pediat. 89: 732–736, 1976.

[68474] 6804.Jhanwar, S. C.; Berkvens, T. M.; Breukel, C.; van Ormondt, H.; van der Eb, A. J.; Meera Khan, P.: Localization of human adenosine deaminase (ADA) gene sequences to the q12–q13.11 region of chromosome 20 by in situ hybridization. Cytogenet. Cell Genet. 50: 168–171, 1989.

[68475] 6805.Jiang, C.-K.; Hong, R.; Horowitz, S. D.; Kong, X.-P.; Hirschhorn, R.: An adenosine deaminase (ADA) allele con-

tains two newly identified deleterious mutations (Y97C and L106V) that interact to abolish enzyme activity. Hum. Molec. Genet. 6: 2271–2278, 1997.

[68476] 6806. Kaitila, I.; Rimoin, D. L.; Cederbaum, S. D.; Stiehm, E. R.; Lechman, R. S.: Chondroosseous histopathology in adenosine deaminase deficient combined immunodeficiency disease. Birth Defects Orig. Art. Ser. XII(6):115–121, 1976.

[68477] 6807. Kellems, R. E.; Yeung, C.-Y.; Ingolia, D. E.: Adenosine deaminase deficiency and severe combined immunodeficiencies. Trends Genet. 1:278–283, 1985.

[68478] 6808. Kenny, A. B.; Hitzig, W. H.: Bone marrow transplantation for severe combined immunodeficiency disease: reported from 1968–1977. Europ. J. Pediatr. 131: 155–176, 1979.

[68479] 6809. Koch, G.; Shows, T. B.: Somatic cell genetics of adenosine deaminase expression and severe combined immunodeficiency disease in humans. Proc. Nat. Acad. Sci. 77: 4211–4215, 1980.

[68480] 6810. Kredich, N. M.; Martin, D. W., Jr.: Role of 5-adenosylhomocysteine in adenosine-mediated toxicity in cultured mouse T-lymphoma cells. Cell 12:931–938, 1977.

- [68481] 6811. Levy, Y.; Hershfield, M. S.; Fernandez-Mejia, C.; Polmar, S. H.; Scudiery, D.; Berger, M.; Sorensen, R. U.: Adenosine deaminase deficiency with late onset of recurrent infections: response to treatment with polyethylene glycol-modified adenosine deaminase. *J. Pediat.* 113:312-317, 1988.
- [68482] 6812. Chashchin, V. L.; Lapko, V. N.; Adamovich, T. B.; Kirillova, N. M.; Lapko, A. G.; Akhrem, A. A.: The primary structure of hepatoferredoxin from bovine liver mitochondria. *Bioorg. Khim.* 12: 1286-1289, 1986.
- [68483] 6813. Jefcoate, C. R.; McNamara, B. C.; DiBartolomeis, M. J.: Control of steroid synthesis in adrenal fasciculata cells. *Endocr. Res.* 12:314-350, 1986.
- [68484] 6814. Maruya, N.; Hiwatashi, A.; Ichikawa, Y.; Yamano, T.: Purification and characterization of renal ferredoxin from bovine renal mitochondria. *J. Biochem.* 93: 1239-1247, 1983.
- [68485] 6815. Mittal, S.; Zhu, Y. Z.; Vickery, L. E.: Molecular cloning and sequence analysis of human placental ferredoxin. *Arch. Biochem. Biophys.* 264:383-391, 1988.
- [68486] 6816. Morel, Y.; Picado-Leonard, J.; Mohandas, T. K.; Miller, W. L.: Two highly homologous genes for adrenodoxin lie on human chromosomes 11 and 20. (Abstract)

Am. J. Hum. Genet. 41: A178 only, 1987.

- [68487] 6817. Morel, Y.; Picado-Leonard, J.; Wu, D.-A.; Chang, C.-Y.; Mohandas, T. K.; Chung, B.-C.; Miller, W. L.: Assignment of the functional gene for human adrenodoxin to chromosome 11q13-qter and of adrenodoxin pseudo-genes to chromosome 20cen-q13.1. Am. J. Hum. Genet. 43: 52-59, 1988.
- [68488] 6818. Okamura, T.; John, M. E.; Zuber, M. X.; Simpson, E. R.; Waterman, M. R.: Molecular cloning and amino acid sequence of the precursor form of bovine adrenodoxin: evidence for a previously unidentified COOH-terminal peptide. Proc. Nat. Acad. Sci. 82: 5705-5709, 1985.
- [68489] 6819. Picado-Leonard, J.; Voutilainen, R.; Kao, L.-C.; Chung, B.-C.; Strauss, J. F., III; Miller, W. L.: Human adrenodoxin: cloning of three cDNAs and cycloheximide enhancement in JEG-3 cells. J. Biol. Chem. 263: 3240-3244, 1988.
- [68490] 6820. Sparkes, R. S.; Klisak, I.; Miller, W. L.: Regional mapping of genes encoding human steroidogenic enzymes: P450<sub>scc</sub> to 15q23-q24; adrenodoxin to 11q22; adrenodoxin reductase to 17q24-q25; and P450<sub>c17</sub> to 10q24-q25. DNA Cell Biol. 10: 359-365, 1991.
- [68491] 6821. Lin, D.; Shi, Y.; Miller, W. L.: Cloning and sequence

of the human adrenodoxin reductase gene. Proc. Nat. Acad. Sci. 87: 8516–8520, 1990.

[68492] 6822. Solish, S. B.; Picado-Leonard, J.; Morel, Y.; Kuhn, R. W.; Mohandas, T. K.; Hanukoglu, I.; Miller, W. L.: Human adrenodoxin reductase: two mRNAs encoded by a single gene on chromosome 17cen–q25 are expressed in steroidogenic tissues. Proc. Nat. Acad. Sci. 85: 7104–7108, 1988.

[68493] 6823. Caron, K. M.; Smithies, O.: Extreme hydrops fetalis and cardiovascular abnormalities in mice lacking a functional adrenomedullin gene. Proc. Nat. Acad. Sci. 98: 615–619, 2001.

[68494] 6824. Ishimitsu, T.; Kojima, M.; Kangawa, K.; Hino, J.; Matsuo, H.; Kitamura, K.; Eto, T.; Matsuo, H.: Genomic structure of human adrenomedullin gene. Biochem. Biophys. Res. Commun. 203: 631–639, 1994.

[68495] 6825. Kitamura, K.; Sakata, J.; Kangawa, K.; Kojima, M.; Matsuo, H.; Eto, T.: Cloning and characterization of cDNA encoding a precursor for human adrenomedullin. Biochem. Biophys. Res. Commun. 194: 720–725, 1993.

[68496] 6826. Makino, Y.; Shibata, K.; Makino, I.; Kangawa, K.; Kawarabayashi, T.: Alteration of the adrenomedullin receptor components gene expression associated with the blood

pressure in pregnancy-induced hypertension. *J.Clin. Endocr. Metab.* 86: 5079–5082, 2001.

- [68497] 6827. McLatchie, L. M.; Fraser, N. J.; Main, M. J.; Wise, A.; Brown, J.; Thompson, N.; Solari, R.; Lee, M. G.; Foord, S. M.: RAMPs regulate the transport and ligand specificity of the calcitonin-receptor-like receptor. *Nature* 393: 333–339, 1998.
- [68498] 6828. Okazaki, T.; Ogawa, Y.; Tamura, N.; Mori, K.; Isse, N.; Aoki, T.; Rochelle, J. M.; Taketo, M. M.; Seldin, M. F.; Nakao, K.: Genomic organization, expression, and chromosomal mapping of the mouse adrenomedullin gene. *Genomics* 37: 395–399, 1996.
- [68499] 6829. Richards, A. M.; Nicholls, M. G.; Lewis, L.; Lainchbury, J. G.: Adrenomedullin. *Clin. Sci.* 91: 3–16, 1996.
- [68500] 6830. Udonio, T.; Takahashi, K.; Nakayama, M.; Yoshinoya, A.; Totsune, K.; Murakami, O.; Durlu, Y. K.; Tamai, M.; Shibahara, S.: Induction of adrenomedullin by hypoxia in cultured retinal pigment epithelial cells. *Invest. Ophthalm. Vis. Sci.* 42: 1080–1086, 2001.
- [68501] 6831. van Heyningen, V.; Jones, C.: Report of the committee on the genetic constitution of chromosome 11. In: Cuticchia, A. J.; Pearson, P. L.; Klinger, H. P. (eds.): Chromosome coordinating meeting, 1992. *Genome Priority Re-*

ports, Vol 1. Basel: S. Karger (pub.) 1993. Pp. 365–401.

- [68502] 6832.Ferguson-Smith, A. C.; Cattanach, B. M.; Barton, S. C.; Beechey, C. V.; Surani, M. A.: Embryological and molecular investigationsof parental imprinting on mouse chromosome 7. *Nature* 351: 667–670, 1991.
- [68503] 6833.Ohno, S.: Ancient linkage groups and frozen accidents. *Nature* 244:259–262, 1973.
- [68504] 6834.Tolan, D. R.; Niclas, J.; Bruce, B. D.; Lebo, R. V.: Evolutionaryimplications of the human aldolase–A, –B, –C, and –pseudogene chromosomelocations. *Am. J. Hum. Genet.* 41: 907–924, 1987.
- [68505] 6835.Buono, P.; Paolella, G.; Mancini, F. P.; Izzo, P.; Salvatore, F.: The complete nucleotide sequence of the gene coding for the humanaldolase C. *Nucleic Acids Res.* 16: 4733 only, 1988.
- [68506] 6836.Rocchi, M.; Vitale, E.; Covone, A.; Romeo, G.; Santamaria, R.;Buono, P.; Paolella, G.; Salvatore, F.: Assignment of human aldolaseC gene to chromosome 17, region centromere–q21.1. *Hum. Genet.* 82: 279–282, 1989.
- [68507] 6837.Rottmann, W. H.; Deselms, K. R.; Niclas, J.; Camerato, T.; Holman, P. S.; Green, C. J.; Tolan, D. R.: The complete amino acid sequenceof the human aldolase C isozyme derived from genomic clones. *Biochimie*

69:137–145, 1987.

- [68508] 6838. Bateman, J. B.; Kojis, T.; Heinzmann, C.; Klisak, I.; Diep, A.; Carper, D.; Nishimura, C.; Mohandas, T.; Sparkes, R. S.: Mapping of aldose reductase gene sequences to human chromosomes 1, 3, 7, 9, 11, and 13. *Genomics* 17: 560–565, 1993.
- [68509] 6839. Brown, L.; Hedge, P. J.; Markham, A. F.; Graham, A.: A human aldehyde dehydrogenase (aldose reductase) pseudogene: nucleotide sequence analysis and assignment to chromosome 3. *Genomics* 13: 465–468, 1992.
- [68510] 6840. Chung, S.; LaMendola, J.: Cloning and sequence determination of human placental aldose reductase gene. *J. Biol. Chem.* 264: 14775–14777, 1989.
- [68511] 6841. Graham, A.; Brown, L.; Hedge, P. J.; Gammack, A. J.; Markham, A. F.: Structure of the human aldose reductase gene. *J. Biol. Chem.* 266: 6872–6877, 1991.
- [68512] 6842. Graham, A.; Heath, P.; Morten, J. E. N.; Markham, A. F.: The human aldose reductase gene maps to chromosome region 7q35. *Hum. Genet.* 86: 509–514, 1991.
- [68513] 6843. Nishimura, C.; Matsuura, Y.; Kokai, Y.; Akera, T.; Carper, D.; Morjana, N.; Lyons, C.; Flynn, T. G.: Cloning and expression of human aldose reductase. *J. Biol. Chem.* 265: 9788–9792, 1990.



- [68514] 6844. Shah, V. O.; Dorin, R. I.; Sun, Y.; Braun, M.; Zager, P. G.: Aldosereductase gene expression is increased in diabetic nephropathy. *J.Clin. Endocr. Metab.* 82: 2294–2298, 1997.
- [68515] 6845. Roberds, S. L.; Anderson, J.; Basi, G.; Bienkowski, M. J.; Branstetter, D. G.; Chen, K. S.; Freedman, S. B.; Frigon, N. L.; Games, D.; Hu, K.; Johnson–Wood, K.; Kappenman, K. E.; and 20 others: BACE knockout mice are healthy despite lacking the primary beta–secretase activity in brain: implications for Alzheimer's disease therapeutics. *Hum. Molec. Genet.* 10: 1317–1324, 2001.
- [68516] 6846. Botto, M.; Hawkins, P. N.; Bickerstaff, M. C.; Herbert, J.; Bygrave, A. E.; McBride, A.; Hutchinson, W. L.; Tennent, G. A.; Walport, M. J.; Pepys, M. B.: Amyloid deposition is delayed in mice with targeted deletion of the serum amyloid P component gene. *Nature Med.* 3: 855–859, 1997.
- [68517] 6847. Floyd–Smith, G. A.; Whitehead, A. S.; Colten, H. R.; Francke, U.: Human serum amyloid P component (SAP) is located on the proximal long arm of chromosome 1. (Abstract) *Cytogenet. Cell Genet.* 40:631 only, 1985.
- [68518] 6848. Ionasescu, V.; Burns, T.; Searby, C.; Ionasescu, R.: Linkage between the loci for Duffy (FY) and serum amyloid P component (APCS) on human chromosome 1. *Cytogenet.*

Cell Genet. 45: 240–241, 1987.

[68519] 6849.Mantzouranis, E. C.; Dowton, S. B.; Whitehead, A. S.; Edge, M.D.; Bruns, G. A. P.; Colten, H. R.: Human serum amyloid P component:cDNA isolation, complete sequence of pre-serum amyloid P component,and localization of the gene to chromosome 1. J. Biol. Chem. 260:7752–7756, 1985.

[68520] 6850.Mortensen, R. F.; Le, P. T.; Taylor, B. A.: Mouse serum amyloidP–component (SAP) levels controlled by a locus on chromosome 1. Immunogenetics 22:367–375, 1985.

[68521] 6851.Pepys, M. B.; Herbert, J.; Hutchinson, W. L.; Tennent, G. A.; Lachmann,H. J.; Gallimore, J. R.; Lovat, L. B.; Bartfai, T.; Alanine, A.; Hertel,C.; Hoffmann, T.; Jakob–Roetne, R.; and 11 others: Targeted pharmacologicaldepletion of serum amyloid P component for treatment of human amyloidosis. Nature 417:254–259, 2002.

[68522] 6852.Prelli, F.; Pras, M.; Frangione, B.: The primary structure ofhuman tissue amyloid P component from a patient with primary idiopathicamyloidosis. J. Biol. Chem. 260: 12895–12898, 1985.

[68523] 6853.Whitehead, A. S.; Rits, M.; Michaelson, J.: Molecular geneticsof mouse serum amyloid P component (SAP):

cloning and gene mapping. *Immunogenetics* 28:388–390, 1988.

[68524] 6854.Woo, P.; O'Brien, J.; Robson, M.; Ansell, B. M.: A genetic marker for systemic amyloidosis in juvenile arthritis. *Lancet* I: 767–769, 1987.

[68525] 6855.Lenkkeri, U.; Kestila, M.; Lamerdin, J.; McCready, P.; Adamson, A.; Olsen, A.; Tryggvason, K.: Structure of the human amyloid-precursor-like protein gene APLP1 at 19q13.1. *Hum. Genet.* 102: 192–196, 1998.

[68526] 6856.Wasco, W.; Brook, J. D.; Tanzi, R. E.: The amyloid precursor-like protein (APLP) gene maps to the long arm of human chromosome 19. *Genomics* 15:237–239, 1993.

[68527] 6857.Wasco, W.; Bupp, K.; Magendantz, M.; Gusella, J. F.; Tanzi, R.E.; Solomon, F.: Identification of a mouse brain cDNA that encodes a protein related to the Alzheimer-associated amyloid beta-protein precursor. *Proc. Nat. Acad. Sci.* 89: 10758–10762, 1992.

[68528] 6858.Leach, R.; Ko, M.; Krawetz, S. A.: Assignment of amyloid-precursor-like protein 2 gene (APLP2) to 11q24 by fluorescent in situ hybridization. *Cytogenet. Cell Genet.* 87: 215–216, 1999.

[68529] 6859.von der Kammer, H.; Loffler, C.; Hanes, J.; Klaudiny, J.; Scheit, K. H.; Hansmann, I.: The gene for the amyloid

precursor-like protein APLP2 is assigned to human chromosome 11q23–25. *Genomics* 10: 308–311, 1994.

[68530] 6860. von Koch, C. S.; Lahiri, D. K.; Mammen, A. L.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Sisodia, S. S.: The mouse APLP2 gene: chromosomal localization and promoter characterization. *J. Biol. Chem.* 270: 25475–25480, 1995.

[68531] 6861. Yan, Y. C.; Bai, Y.; Wang, L.; Miao, S.; Koide, S. S.: Characterization of cDNA encoding a human sperm membrane protein related to A4 amyloid protein. *Proc. Nat. Acad. Sci.* 87: 2405–2408, 1990.

[68532] 6862. Yang, Y.; Martin, L.; Cuzin, F.; Mattei, M.-G.; Rasoulzadegan, M.: Genomic structure and chromosomal localization of the mouse CDEI-binding protein CDEBP (APLP2) gene and promoter sequences. *Genomics* 35: 24–29, 1996.

[68533] 6863. Wallukat, G.; Homuth, V.; Fischer, T.; Lindschau, C.; Horstkamp, B.; Jupner, A.; Baur, E.; Nissen, E.; Vetter, K.; Neichel, D.; Dudenhausen, J. W.; Haller, H.; Luft, F. C.: Patients with preeclampsia develop agonistic autoantibodies against the angiotensin AT-1 receptor. *J. Clin. Invest.* 103: 945–952, 1999.

[68534] 6864. Wang, W. Y. S.; Zee, R. Y. L.; Morris, B. J.: Association

of angiotensin II type 1 receptor gene polymorphism with essential hypertension. Clin.Genet. 51: 31–34, 1997.

[68535] 6865. Wilson, G. N.; Dasouki, M.; Barr, M., Jr.: Further delineation of the dup(3q) syndrome. Am. J. Med. Genet. 22: 117–123, 1985.

[68536] 6866. Amant, C.; Bauters, C.; Bodart, J.-C.; Lablanche, J.-M.; Grollier, G.; Danchin, N.; Hamon, M.; Richard, F.; Helbecque, N.; McFadden, E. P.; Amouyel, P.; Bertrand, M. E.: D allele of the angiotensin I-converting enzyme is a major risk factor for restenosis after coronary stenting. Circulation 96: 56–60, 1997.

[68537] 6867. Arbustini, E.; Grasso, M.; Fasani, R.; Klersy, C.; Diegoli, M.; Porcu, E.; Banchieri, N.; Fortina, P.; Danesino, C.; Specchia, G.: Angiotensin converting enzyme gene deletion allele is independently and strongly associated with coronary atherosclerosis and myocardial infarction. Brit. Heart J. 74: 584–591, 1995.

[68538] 6868. Berge, K. E.; Berg, K.: No effect of insertion/deletion polymorphism at the ACE locus on normal blood pressure level or variability. Clin.Genet. 45: 169–174, 1994.

[68539] 6869. Bohn, M.; Berge, K. E.; Bakken, A.; Erikssen, J.; Berg, K.: Insertion/deletion (I/D) polymorphism at the locus for angiotensin I-converting enzyme and myocardial infarction.

tion. Clin. Genet. 44: 292–297, 1993.

- [68540] 6870. Bohn, M.; Berge, K. E.; Bakken, A.; Erikssen, J.; Berg, K.: Insertion/deletion(I/D) polymorphism at the locus for angiotensin I-converting enzyme and parental history of myocardial infarction. Clin. Genet. 44:298–301, 1993.
- [68541] 6871. Brown, N. J.; Ray, W. A.; Snowden, M.; Griffin, M. R.: Black Americans have an increased rate of angiotensin converting enzyme inhibitor-associated angioedema. Clin. Pharmacol. Therapeutics 60: 8–13, 1996.
- [68542] 6872. Cambien, F.; Alhenc-Gelas, F.; Herbeth, B.; Andre, J. L.; Rakotovo, R.; Gonzales, M. F.; Allegrini, J.; Bloch, C.: Familial resemblance of plasma angiotensin-converting enzyme level: the Nancy study. Am. J. Hum. Genet. 43: 774–780, 1988.
- [68543] 6873. Cambien, F.; Poirier, O.; Lecerf, L.; Evans, A.; Cambou, J.-P.; Arveiler, D.; Luc, G.; Bard, J.-M.; Bara, L.; Ricard, S.; Tiret, L.; Amouyel, P.; Alhenc-Gelas, F.; Soubrier, F.: Deletion polymorphism in the gene for angiotensin-converting enzyme is a potent risk factor for myocardial infarction. Nature 359: 641–644, 1992.
- [68544] 6874. Doria, A.; Warram, J. H.; Krolewski, A. S.: Genetic predisposition to diabetic nephropathy: evidence for a role of the angiotensin I-converting enzyme gene. Diabetes 43:

690–695, 1994.

- [68545] 6875.Ehlers, M. R. W.; Fox, E. A.; Strydom, D. J.; Riordan, J. F.:Molecular cloning of human testicular angiotensin–converting enzyme:the testis isozyme is identical to the C-terminal half of endothelialangiotensin–converting enzyme. *Proc. Nat. Acad. Sci.* 86: 7741–7745,1989.
- [68546] 6876.Esther, C. R., Jr.; Marino, E. M.; Howard, T. E.; Machaud, A.;Corvol, P.; Capecchi, M. R.; Bernstein, K. E.: The critical roleof tissue angiotensin–converting enzyme as revealed by gene targetingin mice. *J. Clin. Invest.* 99: 2375–2385, 1997.
- [68547] 6877.Evans, A. E.; Poirier, O.; Kee, F.; Lecerf, L.; McCrum, E.; Falconer,T.; Crane, J.; O'Rourke, D. F.; Cambien, F.: Polymorphisms of theangiotensin–converting–enzyme gene in subjects who die from coronaryheart disease. *Quart. J. Med.* 87: 211–214, 1994.
- [68548] 6878.Exner, D. V.; Dries, D. L.; Domanski, M. J.; Cohn, J. N.: Lesserresponse to angiotensin–converting–enzyme inhibitor therapy in blackas compared with white patients with left ventricular dysfunction. *NewEng. J. Med.* 344: 1351–1357, 2001.
- [68549] 6879.Gardemann, A.; Fink, M.; Stricker, J.; Nguyen, Q. D.; Humme, J.;Katz, N.; Tillmanns, H.; Hehrlein, F. W.; Rau, M.;

Haberbosch, W.: ACE I/D gene polymorphism: presence of the ACE D allele increases the risk of coronary artery disease in younger individuals. *Atherosclerosis* 139:153–159, 1998.

[68550] 6880. Gayagay, G.; Yu, B.; Hambly, B.; Boston, T.; Hahn, A.; Celermajer, D. S.; Trent, R. J.: Elite endurance athletes and the ACE I allele: the role of genes in athletic performance. *Hum. Genet.* 103: 48–50, 1998.

[68551] 6881. Hagaman, J. R.; Moyer, J. S.; Bachman, E. S.; Sibony, M.; Magyar, P. L.; Welch, J. E.; Smithies, O.; Kregge, J. H.; O'Brien, D. A.: Angiotensin-converting enzyme and male fertility. *Proc. Nat. Acad. Sci.* 95: 2552–2557, 1998.

[68552] 6882. Hamdi, H. K.; Reznik, J.; Castellon, R.; Atilano, S. R.; Ong, J. M.; Udar, N.; Tavis, J. H.; Aoki, A. M.; Nesburn, A. B.; Boyer, D. S.; Small, K. W.; Brown, D. J.; Kenney, M. C.: Alu DNA polymorphism in ACE gene is protective for age-related macular degeneration. *Biochem. Biophys. Res. Commun.* 295: 668–672, 2002.

[68553] 6883. Howard, T. E.; Shai, S. Y.; Langford, K. G.; Martin, B. M.; Bernstein, K. E.: Transcription of testicular angiotensin-converting enzyme (ACE) is initiated within the 12th intron of the somatic ACE gene. *Molec. Cell. Biol.* 10: 4294–4302, 1990.



- [68554] 6884.Huang, W.; Gallois, Y.; Bouby, N.; Bruneval, P.; Heudes, D.; Belair,M.–F.; Krege, J. H.; Menteton, P.; Marre, M.; Smithies, O.; Alhenc–Gelas,F.: Genetically increased angiotensin I–converting enzyme level andrenal complications in the diabetic mouse. *Proc. Nat. Acad. Sci.* 98:13330–13334, 2001.
- [68555] 6885.Jeffery, S.; Malik, A. K. S.; Crosby, A.; Bland, M.; Eastwood,J. B.; Amoah–Danquah, J.; Acheampong, J. W.; Plange–Rhule, J.: Adominant relationship between the ACE D allele and serum ACE levelsin a Ghanaian population. (Letter) *J. Med. Genet.* 36: 869–870, 1999.
- [68556] 6886.Jeunemaitre, X.; Lifton, R. P.; Hunt, S. C.; Williams, R. R.;Lalouel, J.–M.: Absence of linkage between the an–giotensin convertingenzyme locus and human essential hypertension. *Nature Genet.* 1:72–75, 1992.
- [68557] 6887.Keavney, B.; McKenzie, C.; Parish, S.; Palmer, A.; Clark, S.;Youngman, L.; Delepine, M.; Lathrop, M.; Peto, R.; Collins, R.: Large–scale test of hypothesized associations between the angiotensin–convert–ing–enzymeinsertion/deletion polymorphism and myocar–dial infarction in about5000 cases and 6000 controls. *Lancet* 355: 434–442, 2000.
- [68558] 6888.Kehoe, P. G.; Russ, C.; McIlroy, S.; Williams, H.; Hol–

mans, P.;Holmes, C.; Liolitsa, D.; Vahidassr, D.; Powell, J.; McGleenon, B.;Liddell, M.; Plomin, R.; and 9 others: Variation in DCP1, encodingACE, is associated with susceptibility to Alzheimer disease. (Letter) NatureGenet. 21: 71–72, 1999.

- [68559] 6889.Keramatipour, M.; McConnell, R. S.; Kirkpatrick, P.; Tebbs, S.;Furlong, R. A.; Rubinsztein, D. C.: The ACE I allele is associatedwith increased risk for ruptured intracranial aneurysms. J. Med.Genet. 37: 498–500, 2000.
- [68560] 6890.Oliver, P. M.; Fox, J. E.; Kim, R.; Rockman, H. A.; Kim, H.–S.;Reddick, R. L.; Pandey, K. N.; Milgram, S. L.; Smithies, O.; Maeda,N.: Hypertension, cardiac hypertrophy, and sudden death in mice lackingnatriuretic peptide receptor A. Proc. Nat. Acad. Sci. 94: 14730–14735,1997.
- [68561] 6891.Oliver, P. M.; John, S. W. M.; Purdy, K. E.; Kim, R.; Maeda, N.;Goy, M. F.; Smithies, O.: Natriuretic peptide receptor 1 expressioninfluences blood pressures of mice in a dose–dependent manner. Proc.Nat. Acad. Sci. 95: 2547–2551, 1998.
- [68562] 6892.He, X.; Chow, D.; Martick, M. M.; Garcia, K. C.: Allosteric activationof a spring–loaded natriuretic peptide receptor dimer by hormone. Science 293:1657–1662, 2001.

- [68563] 6893. Jaubert, J.; Jaubert, F.; Martin, N.; Washburn, L. L.; Lee, B.K.; Eicher, E. M.; Guenet, J.-L.: Three new allelic mouse mutations that cause skeletal overgrowth involve the natriuretic peptide receptor C gene (Npr3). *Proc. Nat. Acad. Sci.* 96: 10278–10283, 1999.
- [68564] 6894. Lopez, M. J.; Wong, S. K.-F.; Kishimoto, I.; Dubois, S.; Mach, V.; Friesen, J.; Garbers, D. L.; Beuve, A.: Salt-resistant hypertension in mice lacking the guanylyl cyclase-A receptor for atrial natriuretic peptide. *Nature* 378: 65–68, 1995.
- [68565] 6895. Tweardy, D. J.; Cannizzaro, L. A.; Palumbo, A. P.; Shane, S.; Huebner, K.; Vantuinen, P.; Ledbetter, D. H.; Finan, J. B.; Nowell, P. C.; Rovera, G.: Molecular cloning and characterization of a cDNA for human granulocyte colony-stimulating factor (G-CSF) from a glioblastoma multiforme cell line and localization of the G-CSF gene to chromosome band 17q21. *Oncogene Res.* 1: 209–220, 1987.
- [68566] 6896. Dong, F.; Hoefsloot, L. H.; Schelen, A. M.; Broeders, L. C. A.M.; Meijer, Y.; Veerman, A. J. P.; Touw, I. P.; Lowenberg, B.: Identification of a nonsense mutation in the granulocyte-colony-stimulating factor receptor in severe congenital neutropenia. *Proc. Nat. Acad. Sci.* 91: 4480–4484, 1994.

- [68567] 6897.Dong, F.; Qiu, Y.; Yi, T.; Touw, I. P.; Larner, A. C.:  
The carboxylterminus of the granulocyte colony–  
stimulating factor receptor, truncated in patients with se–  
vere congenital neutropenia/acute myeloid leukemia, is re–  
quired for SH2–containing phosphatase–1 suppression of  
Stat activation. *J.Immun.* 167: 6447–6452, 2001.
- [68568] 6898.Fukunaga, R.; Seto, Y.; Mizushima, S.; Nagata, S.:  
Three different mRNAs encoding human granulocyte  
colony–stimulating factor receptor. *Proc.Nat. Acad. Sci.*  
87: 8702–8706, 1990.
- [68569] 6899.Inazawa, J.; Fukunaga, R.; Seto, Y.; Nakagawa, H.;  
Misawa, S.; Abe, T.; Nagata, S.: Assignment of the human  
granulocyte colony–stimulating factor receptor gene  
(CSF3R) to chromosome 1 at region p35–p34.3. *Genomics*  
10:1075–1078, 1991.
- [68570] 6900.Kostmann, R.: Infantile Genetic Agranulocytosis  
(agranulocytosis infantilis hereditaria): a new recessive  
lethal disease in man. Uppsala: Almqvist and Wiksells  
Boktryckeri (pub.) 1956.
- [68571] 6901.McLemore, M. L.; Poursine–Laurent, J.; Link, D. C.:  
Increased granulocyte colony–stimulating factor respon–  
siveness but normal resting granulopoiesis in mice carry–  
ing a targeted granulocyte colony–stimulating factor re–

ceptor mutation derived from a patient with severe congenital neutropenia. *J. Clin. Invest.* 102: 483–492, 1998.

[68572] 6902. Seto, Y.; Fukunaga, R.; Nagata, S.: Chromosomal gene organization of the human granulocyte colony-stimulating factor receptor. *J. Immun.* 148:259–266, 1992.

[68573] 6903. Tidow, N.; Pilz, C.; Teichmann, B.; Muller-Brechlin, A.; Germeshausen, M.; Kasper, B.; Rauprich, P.; Sykora, K.-W.; Welte, K.: Clinical relevance of point mutations in the cytoplasmic domain of the granulocyte colony-stimulating factor receptor gene in patients with severe congenital neutropenia. *Blood* 89: 2369–2375, 1997.

[68574] 6904. Tweardy, D. J.; Anderson, K.; Cannizzaro, L. A.; Steinman, R. A.; Croce, C. M.; Huebner, K.: Molecular cloning of cDNAs for the human granulocyte colony-stimulating factor receptor from HL-60 and mapping of the gene to chromosome region 1p32–34. *Blood* 79: 1148–1154, 1992.

[68575] 6905. Wong, W.-Y.; Williams, D.; Slovak, M. L.; Charak, B.; Mazumder, A.; Snyder, D.; Powars, D. R.; Byrnes, R. K.: Terminal acute myelogenous leukemia in a patient with congenital agranulocytosis. *Am. J. Hemat.* 43:133–138, 1993.

- [68576] 6906.D'Andrea, R.; Rayner, J.; Moretti, P.; Lopez, A.; Goodall, G. J.;Gonda, T. J.; Vadas, M.: A mutation of the common receptor subunitfor interleukin-3 (IL-3), granulocyte-macrophage colony-stimulatingfactor, and IL-5 that leads to ligand independence and tumorigenicity. Blood 83:2802-2808, 1994.
- [68577] 6907.Dirksen, U.; Hattenhorst, U.; Schneider, P.; Schroten, H.; Gobel,U.; Bocking, A.; Muller, K.-M.; Murray, R.; Burdach, S.: Defectiveexpression of granulocyte-macrophage colony-stimulating factor/interleukin-3/interleukin-5receptor common beta chain in children with acute myeloid leukemiaassociated with respiratory failure. Blood 92: 1097-1103, 1998.
- [68578] 6908.Dirksen, U.; Nishinakamura, R.; Groneck, P.; Hattenhorst, U.; Noguee,L.; Murray, R.; Burdach, S.: Human pulmonary alveolar proteinosisassociated with a defect in GM-CSF/IL-3/IL-5 receptor common betachain expression. J. Clin. Invest. 100: 2211-2217, 1997.
- [68579] 6909.Wada, K.; Yokotani, N.; Hunter, C.; Doi, K.; Wentholt, R. J.; Shimasaki,S.: Differential expression of two distinct forms of mRNA encodingmembers of a dipeptidyl aminopeptidase family. Proc. Nat. Acad. Sci. 89:197-201, 1992.

- [68580] 6910.Wada, K.; Zimmerman, K. L.; Adamson, M. C.; Yokotani, N.; Wenthold, R. J.; Kozak, C. A.: Genetic mapping of the mouse gene encoding dipeptidylaminopeptidase-like proteins. *Mammalian Genome* 4: 234–237, 1993.
- [68581] 6911.Yokotani, N.; Doi, K.; Wenthold, R. J.; Wada, K.: Non-conservation of a catalytic residue in a dipeptidyl aminopeptidase IV-related protein encoded by a gene on human chromosome 7. *Hum. Molec. Genet.* 2: 1037–1039, 1993.
- [68582] 6912.Chang, T.-M.; Neville, D. M., Jr.: Demonstration of diphtheria toxin receptors on surface membranes from both toxin-sensitive and toxin-resistant species. *J. Biol. Chem.* 253: 6866–6871, 1978.
- [68583] 6913.Creagan, R. P.; Chen, S.-H.; Ruddle, F. H.: Genetic analysis of the cell surface: association of human chromosome 5 with sensitivity to diphtheria toxin in mouse-human somatic cell hybrids. *Proc. Nat. Acad. Sci.* 72: 2237–2241, 1975.
- [68584] 6914.Fen, Z.; Dhadly, M. S.; Yoshizumi, M.; Hilkert, R. J.; Quertermous, T.; Eddy, R. L.; Shows, T. B.; Lee, M.-E.: Structural organization and chromosomal assignment of the gene encoding the human heparin-binding epidermal

growth factor-like growth factor/diphtheria toxin receptor. *Biochemistry* 32:7932–7938, 1993.

[68585] 6915. George, D. L.; Francke, U.: Regional mapping of human genes for hexosaminidase B and diphtheria toxin sensitivity on chromosome 5 using mouse X human hybrid cells. *Somat. Cell Genet.* 3: 629–638, 1977.

[68586] 6916. Gupta, R. S.; Siminovitch, L.: Isolation and characterization of mutants of human diploid fibroblasts resistant to diphtheria toxin. *Proc. Nat. Acad. Sci.* 75: 3337–3340, 1978.

[68587] 6917. Hayes, H.; Kaneda, Y.; Uchida, T.; Okada, Y.: Regional assignment of the gene for diphtheria toxin sensitivity using subchromosomal fragments in microcell hybrids. *Chromosoma* 96: 26–32, 1987.

[68588] 6918. Higashiyama, S.; Abraham, J. A.; Miller, J.; Fiddes, J. C.; Klagsbrun, M.: A heparin-binding growth factor secreted by macrophage-like cells that is related to EGF. *Science* 251: 936–939, 1991.

[68589] 6919. Wang, Q.; Curren, M. E.; Splawski, I.; Burn, T. C.; Millholland, J. M.; VanRaay, T. J.; Shen, J.; Timothy, K. W.; Vincent, G. M.; deJager, T.; Schwartz, P. J.; Towbin, J. A.; Moss, A. J.; Atkinson, D. L.; Landes, G. M.; Connors, T. D.; Keating, M. T.: Positional cloning of a novel potassium



channel gene: KVLQT1 mutations cause cardiac arrhythmias. *Nature Genet.* 12: 17–23, 1996.

[68590] 6920. Gorman, D. M.; Itoh, N.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Miyajima, A.: Chromosomal localization and organization of the murine genes encoding the beta subunits (AIC2A and AIC2B) of the interleukin 3, granulocyte/macrophage colony-stimulating factor, and interleukin 5 receptors. *J. Biol. Chem.* 267: 15842–15848, 1992.

[68591] 6921. Jenkins, B. J.; D'Andrea, R.; Gonda, T. J.: Activating point mutations in the common beta subunit of the human GM-CSF, IL-3 and IL-5 receptors suggest the involvement of beta subunit dimerization and cell type-specific molecules in signalling. *EMBO J.* 14: 4276–4287, 1995.

[68592] 6922. Kitamura, T.; Sato, N.; Arai, K.; Miyajima, A.: Expression cloning of the human IL-3 receptor cDNA reveals a shared beta subunit for the human IL-3 and GM-CSF receptors. *Cell* 66: 1165–1174, 1991.

[68593] 6923. Kondo, M.; Scherer, D. C.; Miyamoto, T.; King, A. G.; Akashi, K.; Sugamura, K.; Weissman, I. L.: Cell-fate conversion of lymphoid-committed progenitors by instructive actions of cytokines. *Nature* 407: 383–386, 2000.

- [68594] 6924.Robb, L.; Drinkwater, C. C.; Metcalf, D.; Li, R.; Kontgen, F.; Nicola, N. A.; Begley, C. G.: Hematopoietic and lung abnormalities in mice with a null mutation of the common beta subunit of the receptors for granulocyte-macrophage colony-stimulating factor and interleukins 3 and 5. *Proc. Nat. Acad. Sci.* 92: 9565–9569, 1995.
- [68595] 6925.Shen, Y.; Baker, E.; Callen, D. F.; Sutherland, G. R.; Willson, T. A.; Rakar, S.; Gough, N. M.: Localization of the human GM-CSF receptor beta chain gene (CSF2RB) to chromosome 22q12.2–q13.1. *Cytogenet. Cell Genet.* 61: 175–177, 1992.
- [68596] 6926.Tavernier, J.; Devos, R.; Cornelis, S.; Tuypens, T.; Van der Heyden, J.; Fiers, W.; Plaetinck, G.: A human high affinity interleukin-5 receptor (IL5R) is composed of an IL5-specific alpha chain and a beta chain shared with the receptor for GM-CSF. *Cell* 66: 1175–1184, 1991.
- [68597] 6927.Prohl, C.; Pelzer, W.; Diekert, K.; Kmita, H.; Bedkovics, T.; Kispal, G.; Lill, R.: The yeast mitochondrial carrier Leu5p and its human homologue Graves' disease protein are required for accumulation of coenzyme A in the matrix. *Molec. Cell Biol.* 21: 1089–1097, 2001.
- [68598] 6928.Rossi, E.; Zarrilli, R.; Zuffardi, O.: Regional assignment of the gene coding for a human Graves' disease au-

to antigen to 10q21.3–q22.1. Hum. Genet. 90: 653–654, 1993.

[68599] 6929. Zarrilli, R.; Oates, E. L.; McBride, O. W.; Lerman, M. I.; Chan, J. Y.; Santisteban, P.; Ursini, M. V.; Notkins, A. L.; Kohn, L. D.: Sequence and chromosomal assignment of a novel cDNA identified by immunoscreening of a thyroid expression library: similarity to a family of mitochondrial solute carrier proteins. Molec. Endocr. 3: 1498–1508, 1989.

[68600] 6930. Douglas, A. J.; Fox, M. F.; Abbott, C. M.; Hinks, L. J.; Sharpe, G.; Povey, S.; Thompson, R. J.: Structure and chromosomal localization of the human 2'-prime,3'-prime-cyclic nucleotide 3'-prime-phosphodiesterase gene. Ann. Hum. Genet. 56: 243–254, 1992.

[68601] 6931. Douglas, A. J.; Fox, M. F.; Hinks, L. J.; Povey, S.; Thompson, R. J.: Localization of the myelin specific enzyme 2'-prime,3'-prime-cyclic nucleotide 3'-prime-phosphohydrolase to 17q21. (Abstract) Cytogenet. Cell Genet. 58: 2004 only, 1991.

[68602] 6932. Kurihara, T.; Monoh, K.; Sakimura, K.; Takahashi, Y.: Alternative splicing of mouse brain

2-prime,3-prime-cyclic nucleotide

3-prime-phosphodiesterasemRNA. Biochem. Biophys.

Res. Commun. 170: 1074–1081, 1990.

[68603] 6933.Monoh, K.; Kurihara, T.; Takahashi, Y.; Ichikawa, T.;

Kumanishi,T.; Hayashi, S.; Minoshima, S.; Shimizu, N.:

Structure, expressionand chromosomal localization of the  
gene encoding human

2-prime,3-prime-cyclic-nucleotide3-prime-phosphodiesterase. Gene 129: 297–301, 1993.

[68604] 6934.Bernier, L.; Colman, D. R.; D'Eustachio, P.: Chromo-

somal locationsof genes encoding 2-prime,3-prime cyclic  
nucleotide 3-prime-phosphodiesteraseand glial fibrillary  
acidic protein in the mouse. J. Neurosci. Res. 20:497–504,  
1988.

[68605] 6935.Bifulco, M.; Laezza, C.; Stingo, S.; Wolff, J.:

2-prime,3-prime-cyclicnucleotide

3-prime-phosphodiesterase: a membrane-bound, micro-  
tubule-associatedprotein and membrane anchor for tubu-  
lin. Proc. Nat. Acad. Sci. 99:1807–1812, 2002.

[68606] 6936.Sprinkle, T. J.; Kouri, R. E.; Fain, P. D.; Stoming, T.

A.; Whitney,J. B., III: Chromosomal mapping of the human  
CNP gene using a meioticcrossover DNA panel, PCR, and  
allele-specific probes. Genomics 16:542–545, 1993.

- [68607] 6937. Sprinkle, T. J.; Lanclos, K. D.; Lapp, D. F.: Assignment of the human 2'-prime,3'-prime-cyclic nucleotide 3'-prime-phosphohydrolase gene to chromosome 17. *Genomics* 13: 877-880, 1992.
- [68608] 6938. Vogel, U. S.; Thompson, R. J.: Molecular structure, localization, and possible functions of the myelin-associated enzyme 2'-prime,3'-prime-cyclic nucleotide 3'-prime-phosphodiesterase. *J. Neurochem.* 50: 1667-1677, 1988.
- [68609] 6939. Bibb, J. A.; Chen, J.; Taylor, J. R.; Svenningsson, P.; Nishi, A.; Snyder, G. L.; Yan, Z.; Sagawa, Z. K.; Ouimet, C. C.; Nairn, A. C.; Nestler, E. J.; Greengard, P.: Effects of chronic exposure to cocaine are regulated by the neuronal protein Cdk5. *Nature* 410:376-380, 2001.
- [68610] 6940. Bibb, J. A.; Snyder, G. L.; Nishi, A.; Yan, Z.; Meijer, L.; Fienberg, A. A.; Tsai, L.-H.; Kwon, Y. T.; Girault, J.-A.; Czernik, A. J.; Huganir, R. L.; Hemmings, H. C., Jr.; Nairn, A. C.; Greengard, P.: Phosphorylation of DARPP-32 by Cdk5 modulates dopamine signalling in neurons. *Nature* 402:669-671, 1999.
- [68611] 6941. Ohshima, T.; Nagle, J. W.; Pant, H. C.; Joshi, J. B.; Kozak, C. A.; Brady, R. O.; Kulkarni, A. B.: Molecular cloning and chromosomal mapping of the mouse cyclin-

dependent kinase 5 gene. *Genomics* 28:585–588, 1995.

- [68612] 6942. Ohshima, T.; Ward, J. M.; Huh, C.-G.; Longenecker, G.; Veeranna; Pant, H. C.; Brady, R. O.; Martin, L. J.; Kulka-rni, A. B.: Targeted disruption of the cyclin-dependent kinase 5 gene results in abnormal corticogenesis, neuronal pathology and perinatal death. *Proc. Nat. Acad. Sci.* 93: 11173–11178, 1996.
- [68613] 6943. Patrick, G. N.; Zukerberg, L.; Nikolic, M.; de la Monte, S.; Dikkes, P.; Tsai, L.-H.: Conversion of p35 to p25 deregulates Cdk5 activity and promotes neurodegeneration. *Nature* 402: 615–622, 1999.
- [68614] 6944. Demetrick, D. J.; Matsumoto, S.; Hannon, G. J.; Okamoto, K.; Xiong, Y.; Zhang, H.; Beach, D. H.: Chromosomal mapping of the genes for the human cell cycle proteins cyclin C (CCNC), cyclin E (CCNE), p21(CDKN1) and KAP (CDKN3). *Cytogenet. Cell Genet.* 69: 190–192, 1995.
- [68615] 6945. Inaba, T.; Matsushime, H.; Valentine, M.; Roussel, M. F.; Sherr, C. J.; Look, A. T.: Genomic organization, chromosomal localization, and independent expression of human cyclin D genes. *Genomics* 13:565–574, 1992.
- [68616] 6946. Kim, H. A.; Pomeroy, S. L.; Whoriskey, W.; Pawlitzky, I.; Benowitz, L. I.; Sicinski, P.; Stiles, C. D.; Roberts, T. M.: A developmentally regulated switch directs regenerative

growth of Schwann cells through cyclin D1. *Neuron* 26: 405–416, 2000.

- [68617] 6947. Xiong, Y.; Menninger, J.; Beach, D.; Ward, D. C.: Molecular cloning and chromosomal mapping of CCND genes encoding human D-type cyclins. *Genomics* 13:575–584, 1992.
- [68618] 6948. Motokura, T.; Keyomarsi, K.; Kronenberg, H. M.; Arnold, A.: Cloning and characterization of human cyclin D3, a cDNA closely related in sequence to the PRAD1/cyclin D1 proto-oncogene. *J. Biol. Chem.* 267:20412–20415, 1992.
- [68619] 6949. Motokura, T.; Yi, H. F.; Kronenberg, H. M.; McBride, O. W.; Arnold, A.: Assignment of the human cyclin D3 gene (CCND3) to chromosome 6p–q13. *Cytogenet. Cell Genet.* 61: 5–7, 1992.
- [68620] 6950. Wang, Z.; Sicinski, P.; Weinberg, R. A.; Zhang, Y.; Ravid, K.: Characterization of the mouse cyclin D3 gene: exon/intron organization and promoter activity. *Genomics* 35: 156–163, 1996.
- [68621] 6951. Schwartz, J.-C. D.; Zhang, X.; Fedorov, A. A.; Nathenson, S. G.; Almo, S. C.: Structural basis for co-stimulation by the human CTLA-4/B7-2 complex. *Nature* 410: 604–608, 2001.

- [68622] 6952. Stamper, C. C.; Zhang, Y.; Tobin, J. F.; Erbe, D. V.; Ikemizu, S.; Davis, S. J.; Stahl, M. L.; Seehra, J.; Somers, W. S.; Mosyak, L.: Crystal structure of the B7-1/CTLA-4 complex that inhibits human immune responses. *Nature* 410: 608-611, 2001. Note: Erratum: *Nature* 411: 617 only, 2001.
- [68623] 6953. Allenspach, E. J.; Cullinan, P.; Tong, J.; Tang, Q.; Tesciuba, A. G.; Cannon, J. L.; Takahashi, S. M.; Morgan, R.; Burkhardt, J. K.; Sperling, A. I.: ERM-dependent movement of CD43 defines a novel protein complex distal to the immunological synapse. *Immunity* 15: 739-750, 2001.
- [68624] 6954. Bonilha, V. L.; Rodriguez-Boulan, E.: Polarity and developmental regulation of two PDZ proteins in the retinal pigment epithelium. *Invest. Ophthalmol. Vis. Sci.* 42: 3274-3282, 2001.
- [68625] 6955. Gould, K. L.; Bretscher, A.; Esch, F. S.; Hunter, T.: cDNA cloning and sequencing of the protein-tyrosine kinase substrate, ezrin, reveals homology to band 4.1. *EMBO J.* 8: 4133-4142, 1989.
- [68626] 6956. Majander-Nordenswan, P.; Sainio, M.; Turunen, O.; Jaaskelainen, J.; Carpen, O.; Kere, J.; Vaheri, A.: Genomic structure of the human ezrin gene. *Hum. Genet.* 103: 662-665, 1998.



- [68627] 6957. Pakkanen, R.; Vaheri, A.: Cytovillin and other microvillar proteins of human choriocarcinoma cells. *J. Cell. Biochem.* 41: 1–12, 1989.
- [68628] 6958. Rao, P. H.; Murty, V. V. V. S.; Gaidano, G.; Hauptschein, R.; Dalla-Favera, R.; Chaganti, R. S. K.: Sub-regional mapping of 8 single copy loci to chromosome 6 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 66: 272–273, 1994.
- [68629] 6959. Roumier, A.; Olivo-Marin, J. C.; Arpin, M.; Michel, F.; Martin, M.; Mangeat, P.; Acuto, O.; Dautry-Varsat, A.; Alcover, A.: The membrane-microfilament linker ezrin is involved in the formation of the immunological synapse and in T cell activation. *Immunity* 15: 715–728, 2001.
- [68630] 6960. Turunen, O.; Winqvist, R.; Pakkanen, R.; Grzeschik, K.-H.; Wahlstrom, T.; Vaheri, A.: Cytovillin, a microvillar Mr 75,000 protein: cDNA sequence, prokaryotic expression, and chromosomal localization. *J. Biol. Chem.* 264: 16727–16732, 1989.
- [68631] 6961. Winqvist, R.; Turunen, O.; Pakkanen, R.; Grzeschik, K.-H.; Wahlstrom, T. and Vaheri, A.: Localization of the cytovillin gene to region q22–q27 of human chromosome 6. (Abstract) *Cytogenet. Cell Genet.* 51: 1108–1109, 1989.
- [68632] 6962. Brunet, J.-F.; Dosseto, M.; Denizot, F.; Mattei, M.-G.;

Clark, W. R.; Haqqi, T. M.; Ferrier, P.; Nabholz, M.; Schmitt-Verhulst, A.-M.; Luciani, M.-F.; Golstein, P.: The inducible cytotoxic T-lymphocyte-associated gene transcript CTLA-1 sequence and gene localization to mouse chromosome 14. *Nature* 322: 268-271, 1986.

[68633] 6963. Crosby, J. L.; Bleackley, R. C.; Nadeau, J. H.: A complex of serine protease genes expressed preferentially in cytotoxic T-lymphocytes closely linked to the T-cell receptor alpha- and delta-chain genes on mouse chromosome 14. *Genomics* 6: 252-259, 1990.

[68634] 6964. Dahl, C. A.; Bach, F. H.; Chan, W.; Huebner, K.; Russo, G.; Croce, C. M.; Herfurth, T.; Cairns, J. S.: Isolation of a cDNA clone encoding a novel form of granzyme B from human NK cells and mapping to chromosome 14. *Hum. Genet.* 84: 465-470, 1990.

[68635] 6965. Sheffield, V. C.; Stone, E. M.; Alward, W. L. M.; Drack, A. V.; Johnson, A. T.; Streb, L. M.; Nichols, B. E.: Genetic linkage of familial open angle glaucoma to chromosome 1q21-q31. *Nature Genet.* 4:47-50, 1993.

[68636] 6966. Stone, E. M.; Fingert, J. H.; Alward, W. L. M.; Nguyen, T. D.; Polansky, J. R.; Sunden, S. L. F.; Nishimura, D.; Clark, A. F.; Nystuen, A.; Nichols, B. E.; Mackey, D. A.; Ritch, R.; Kalenak, J. W.; Craven, E. R.; Sheffield, V. C.: Identification

of a gene that causes primaryopen angle glaucoma. Science 275: 668–670, 1997.

[68637] 6967.Wiggs, J. L.; Vollrath, D.: Molecular and clinical evaluationof a patient hemizygous for TIGR/MYOC. Arch. Ophthal. 119: 1674–1678,2001.

[68638] 6968.Yoon, S.–J. K.; Kim, H.–S.; Moon, J.–I.; Lim, J. M.; Joo, C.–K.: Mutations of the TIGR/MYOC gene in primary open–angle glaucoma inKorea. (Letter) Am. J. Hum. Genet. 64: 1775–1778, 1999.

[68639] 6969.Muragaki, Y.; Mattei, M.–G.; Yamaguchi, N.; Olsen, B. R.; Ninomiya,Y.: The complete primary structure of the human alpha–1(VIII) chainand assignment of its gene (COL8A1) to chromosome 3. Europ. J. Biochem. 197:615–622, 1991.

[68640] 6970.Muragaki, Y.; Shiota, C.; Inoue, M.; Ooshima, A.; Olsen, B. R.;Ninomiya, Y.: Alpha–1(VIII)–collagen gene transcripts encode a short–chaincollagen polypeptide and are expressed by various epithelial, endothelialand mes–enchymal cells in newborn mouse tissues. Europ. J. Biochem. 207:895–902, 1992.

[68641] 6971.Odink, K.; Cerletti, N.; Bruggen, J.; Clerc, R. G.; Tarc–say, L.;Zwadlo, G.; Gerhards, G.; Schlegel, R.; Sorg, C.: Two calcium–bindingproteins in infiltrate macrophages of

rheumatoid arthritis. *Nature* 330:80–82, 1987.

- [68642] 6972.Schafer, B. W.; Wicki, R.; Engelkamp, D.; Mattei, M.–G.; Heizmann,C. W.: Isolation of a YAC clone covering a cluster of nine S100 genes on human chromosome 1q21: rationale for a new nomenclature of the S100 calcium-binding protein family. *Genomics* 25: 638–643, 1995.
- [68643] 6973.van Heyningen, V.; Emslie, E.; Dorin, J. R.: Related calcium binding proteins map to the same sub-region of chromosome 1q and to an extended region of synteny on mouse chromosome 3. (Abstract) *Cytogenet.Cell Genet.* 51: 1095, 1989.
- [68644] 6974.van Heyningen, V.; Hayward, C.; Fletcher, J.; McAuley, C.: Tissue localization and chromosomal assignment of a serum protein which tracks the cystic fibrosis gene. *Nature* 315: 513–515, 1985.
- [68645] 6975.Welsh, M. J.; Liedtke, C. M.: Chloride and potassium channels in cystic fibrosis airway epithelia. *Nature* 322: 467–470, 1986.
- [68646] 6976.Wilkinson, M. M.; Busuttil, A.; Hayward, C.; Brock, D. J. H.; Dorin, J. R.; van Heyningen, V.: Expression pattern of two related cystic fibrosis-associated calcium-binding proteins in normal and abnormal tissues. *J. Cell Sci.* 91: 221–230, 1988.

- [68647] 6977.Wilson, G. B.; Fudenberg, H. H.; Jahn, T. L.: Studies on cysticfibrosis using isoelectric focusing. I. An assay for detection ofcystic fibrosis homozygotes and heterozygote carriers from serum.Pediat. Res. 9: 635–640, 1975.
- [68648] 6978.van Heyningen, V.; Emslie, E.; Dorin, J. R.: Related calcium bindingproteins map to the same sub–region of chromosome 1q and to an extendedregion of syntenry on mouse chromosome 3. (Abstract) Cytogenet. CellGenet. 51: 1095 only, 1989.
- [68649] 6979.Buonavista, N.; Balzano, C.; Pontarotti, P.; Le Paslier, D.; Golstein,P.: Molecular linkage of the human CTLA4 and CD28 Ig–superfamilygenes in yeast artificial chromo–somes. Genomics 13: 856–861, 1992.
- [68650] 6980.Howard, T. A.; Rochelle, J. M.; Seldin, M. F.: Cd28 and Ctla–4,two related members of the Ig supergene fam–ily, are tightly linkedon proximal mouse chromosome 1. Immunogenetics 33: 74–76, 1991.
- [68651] 6981.Albertson, D. G.; Ylstra, B.; Segraves, R.; Collins, C.; Dairkee,S. H.; Kowbel, D.; Kuo, W.– L.; Gray, J. W.; Pinkel, D.: Quantitativemapping of amplicon structure by array CGH identifies CYP24 as a candidateoncogene. Nature Genet. 25: 144–146, 2000.
- [68652] 6982.Chen, K.–S.; Prahl, J. M.; DeLuca, H. F.: Isolation and

expression of human 1,25-dihydroxyvitamin D<sub>3</sub>  
24-hydroxylase cDNA. *Proc. Nat. Acad. Sci.* 90:  
4543–4547, 1993.

[68653] 6983. Hahn, C. N.; Baker, E.; Laslo, P.; May, B. K.; Omdahl, J. L.; Sutherland, G. R.: Localization of the human vitamin D 24-hydroxylase gene (CYP24) to chromosome 20q13.2–q13.3. *Cytogenet. Cell Genet.* 62: 192–193, 1993.

[68654] 6984. Labuda, M.; Lemieux, N.; Tihy, F.; Prinster, C.; Glorieux, F. H.: Human 25-hydroxyvitamin D 24-hydroxylase cytochrome P450 subunit maps to a different chromosomal location than that of pseudovitamin D-deficient rickets. *J. Bone Miner. Res.* 8: 1397–1406, 1993.

[68655] 6985. Malas, S.; Peters, J.; Abbott, C.: The genes for endothelin 3, vitamin D 24-hydroxylase, and melanocortin 3 receptor map to distal mouse chromosome 2, in the region of conserved synteny with human chromosome 20. *Mammalian Genome* 5: 577–579, 1994.

[68656] 6986. Ohyama, Y.; Noshiro, M.; Okuda, K.: Cloning and expression of cDNA encoding 25-hydroxyvitamin D(3) 24-hydroxylase. *FEBS Lett.* 278:195–198, 1991.

[68657] 6987. Mendel, D. B.; Khavari, P. A.; Conley, P. B.; Graves, M. K.; Hansen, L. P.; Admon, A.; Crabtree, G. R.: Character–

ization of a cofactor that regulates dimerization of a mammalian homeodomain protein. *Science* 254:1762–1767, 1991.

[68658] 6988. Hogenesch, J. B.; Chan, W. K.; Jackiw, V. H.; Brown, R. C.; Gu, Y.-Z.; Pray-Grant, M.; Perdew, G. H.; Bradfield, C. A.: Characterization of a subset of the basic-helix-loop-helix-PAS superfamily that interacts with components of the dioxin signaling pathway. *J. Biol. Chem.* 272:8581–8593, 1997.

[68659] 6989. Wang, G. L.; Jiang, B.-H.; Rue, E. A.; Semenza, G. L.: Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O(2) tension. *Proc. Nat. Acad. Sci.* 92: 5510–5514, 1995.

[68660] 6990. Le Beau, M. M.; Lemons, R. S.; Carrino, J. J.; Pette-nati, M. J.; Souza, L. M.; Diaz, M. O.; Rowley, J. D.: Chromosomal localization of the human G-CSF gene to 17q11 proximal to the breakpoint of the t(15;17) in acute promyelocytic leukemia. *Leukemia* 1: 795–799, 1987.

[68661] 6991. Haaparanta, T.; Uitto, J.; Ruoslahti, E.; Engvall, E.: Molecular cloning of the cDNA encoding human laminin A chain. *Matrix* 11: 151–160, 1991.

[68662] 6992. Gallagher, P. G.; Forget, B. G.: Structure, organization, and expression of the band 7.2b gene, a candidate

gene for hereditary hydrocytosis. *J. Biol. Chem.* 270: 26358–26363, 1995.

[68663] 6993. Gallagher, P. G.; Romana, M.; Lieman, J. H.; Ward, D. C.: cDNA structure, tissue-specific expression, and chromosomal localization of the murine band 7.2b gene. *Blood* 86: 359–365, 1995.

[68664] 6994. Gallagher, P. G.; Upender, M.; Ward, D. C.; Forget, B. G.: The gene for human erythrocyte membrane protein band 7.2 (EPB72) maps to 9q33–q34 centromeric to the Philadelphia chromosome translocation breakpoint region. *Genomics* 18: 167–169, 1993.

[68665] 6995. Hiebl-Dirschmied, C. M.; Entler, B.; Glotzmann, C.; Maurer-Fogy, I.; Stratowa, C.; Prohaska, R.: Cloning and nucleotide sequence of cDNA encoding human erythrocyte band 7 integral membrane protein. *Biochim. Biophys. Acta* 1090: 123–124, 1991.

[68666] 6996. Pilz, A.; Prohaska, R.; Peters, J.; Abbott, C.: Genetic linkage analysis of the Ak1, Col5a1, Epb7.2, Fpgs, Grp78, Pbx3, and Notch1 genes in the region of mouse chromosome 2 homologous to human chromosome 9q. *Genomics* 21: 104–109, 1994.

[68667] 6997. Unfried, I.; Entler, B.; Prohaska, R.: The organization of the gene (EPB72) encoding the human erythrocyte band



7 integral membraneprotein (protein 7.2b). Genomics 30: 521–528, 1995.

- [68668] 6998. Westberg, J. A.; Entler, B.; Prohaska, R.; Schroder, J. P.: The gene coding for erythrocyte protein band 7.2b (EPB72) is located in band q34.1 of human chromosome 9. Cytogenet. Cell Genet. 63: 241–243, 1993.
- [68669] 6999. Zhu, Y.; Paszty, C.; Turetsky, T.; Tsai, S.; Kuypers, F. A.; Lee, G.; Cooper, P.; Gallagher, P. G.; Stevens, M. E.; Rubin, E.; Mohandas, N.; Mentzer, W. C.: Stomatocytosis is absent in 'stomatin'-deficient murine red blood cells. Blood 93: 2404–2410, 1999.
- [68670] 7000. de la Chapelle, A.; Sistonen, P.; Lehvaslaiho, H.; Ikkala, E.; Juvonen, E.: Familial erythrocytosis genetically linked to erythropoietin receptor gene. Lancet 341: 82–84, 1993.
- [68671] 7001. Juvonen, E.; Ikkala, E.; Fyhrquist, F.; Ruutu, T.: Autosomal dominant erythrocytosis caused by increased sensitivity to erythropoietin. Blood 78: 3066–3069, 1991.
- [68672] 7002. Craig, J. E.; Baird, P. N.; Healey, D. L.; McNaught, A. I.; McCartney, P. J.; Rait, J. L.; Dickinson, J. L.; Roe, L.; Fingert, J. H.; Stone, E. M.; Mackey, D. A.: Evidence for genetic heterogeneity within eight glaucoma families, with the GLC1A Gln368STOP mutation being an important pheno-

typic modifier. *Ophthalmology* 108: 1607–1620, 2001.

[68673] 7003.Crombie, A. L.; Cullen, J. F.: Hereditary glaucoma: occurrence in five generations of an Edinburgh family. *Brit. J. Ophthal.* 48:143–147, 1964.

[68674] 7004.Metcalf, D.: The granulocyte–macrophage colony–stimulating factors. *Science* 229:16–22, 1985.

[68675] 7005.Nagata, S.; Tsuchiya, M.; Asano, S.; Kaziro, Y.; Yamazaki, T.; Yamamoto, O.; Hirata, Y.; Kubota, N.; Oheda, M.; Nomura, H.; Ono, M.: Molecular cloning and expression of cDNA for the human granulocyte colony–stimulating factor. *Nature* 319: 415–418, 1986.

[68676] 7006.Nagata, S.; Tsuchiya, M.; Asano, S.; Yamamoto, O.; Hirata, Y.; Kubota, N.; Oheda, M.; Nomura, H.; Yamazaki, T.: The chromosomal gene structure and two mRNAs for human granulocyte colony–stimulating factor. *EMBO J.* 5: 575–581, 1986.

[68677] 7007.Petit, I.; Szyper–Kravitz, M.; Nagler, A.; Lahav, M.; Peled, A.; Habler, L.; Ponomaryov, T.; Taichman, R. S.; Arenzana–Seisdedos, F.; Fujii, N.; Sandbank, J.; Zipori, D.; Lapidot, T.: G-CSF induces stem cell mobilization by decreasing bone marrow SDF–1 and up–regulating CXCR4. *Nature Immun.* 3: 687–694, 2002.

[68678] 7008.Simmers, R. N.; Smith, J.; Shannon, M. F.; Wong, G.;

Lopez, A.F.; Baker, E.; Sutherland, G. R.; Vadas, M. A.: Localization of the human G-CSF gene to the region of a breakpoint in the translocation typical of acute promyelocytic leukemia. *Hum. Genet.* 78: 134–136, 1988.

[68679] 7009. Souza, L. M.; Boone, T. C.; Gabrilove, J.; Lai, P. H.; Zsebo, K.M.; Murdock, D. C.; Chazin, V. R.; Bruszewski, J.; Lee, H.; Chen, K. K.; Barendt, J.; Platzer, E.; Moore, M. A. S.; Mertelsmann, R.; Welte, K.: Recombinant human granulocyte colony-stimulating factor: effects on normal and leukemic myeloid cells. *Science* 232: 61–65, 1986.

[68680] 7010. Sudo, S.; Yamada, H.; Kikuchi, K.; Sumie, A.; Yamashita, Y.; Tumura, N.; Kawaguchi, I.; Fujimoto, S.; Kato, A.; Yamaguchi, J.: A case of ovarian carcinoma with production of granulocyte colony-stimulating factor. *Brit. J. Haemat.* 92: 137–139, 1996.

[68681] 7011. Dong, F.; Brynes, R. K.; Tidow, N.; Welte, K.; Lowenberg, B.; Touw, I. P.: Mutations in the gene for the granulocyte colony-stimulating-factor receptor in patients with acute myeloid leukemia preceded by severe congenital neutropenia. *New Eng. J. Med.* 333: 487–493, 1995.

[68682] 7012. Scott, M. P.: Vertebrate homeobox gene nomenclature. *Cell* 71: 551–553, 1992.

[68683] 7013. Carr, D. W.; Hausken, Z. E.; Fraser, I. D.; Stofko–

Hahn, R. E.; Scott, J. D: Association of the type II cAMP-dependent protein kinase with a human thyroid RII-anchoring protein: cloning and characterization of the RII-binding domain. *J. Biol. Chem.* 267: 13376–13382, 1992.

[68684] 7014. Carr, D. W.; Stofko-Hahn, R. E.; Fraser, I. D.; Bishop, S. M.; Acott, T. S.; Brennan, R. G.; Scott, J. D.: Interaction of the regulatory subunit (RII) of cAMP-dependent protein kinase with RII-anchoring proteins occurs through an amphipathic helix binding motif. *J. Biol. Chem.* 266: 14188–14192, 1991.

[68685] 7015. Rubino, D.; Driggers, P.; Arbit, D.; Kemp, L.; Miller, B.; Coso, O.; Pagliai, K.; Gray, K.; Gutkind, S.; Segars, J.: Characterization of Brx, a novel Dbl family member that modulates estrogen receptor action. *Oncogene* 16: 2513–2526, 1998.

[68686] 7016. Sterpetti, P.; Hack, A. A.; Bashar, M. P.; Park, B.; Cheng, S.-D.; Knoll, J. H. M.; Urano, T.; Feig, L. A.; Toksoz, D.: Activation of the Lbc Rho exchange factor proto-oncogene by truncation of an extended C terminus that regulates transformation and targeting. *Molec. Cell. Biol.* 19: 1334–1345, 1999.

[68687] 7017. Toksoz, D.; Williams, D. A.: Novel human oncogene lbc detected by transfection with distinct homology re-

gions to signal transduction products. *Oncogene* 9: 621–628, 1994.

- [68688] 7018. Hirai, H.; Tanaka, K.; Yoshie, O.; Ogawa, K.; Kenmotsu, K.; Takamori, Y.; Ichimasa, M.; Sugamura, K.; Nakamura, M.; Takano, S.; Nagata, K.: Prostaglandin D2 selectively induces chemotaxis in T helper type 2 cells, eosinophils, and basophils via seven-transmembrane receptor CRTH2. *J. Exp. Med.* 193: 255–261, 2001.
- [68689] 7019. Margolis, R. L.; O'Hearn, E.; Rosenblatt, A.; Willour, V.; Holmes, S. E.; Franz, M. L.; Callahan, C.; Hwang, H. S.; Troncoso, J. C.; Ross, C. A.: A disorder similar to Huntington's disease is associated with a novel CAG repeat expansion. *Ann. Neurol.* 50: 373–380, 2001.
- [68690] 7020. Lacy, S. E.; Bonnemann, C. G.; Buzney, E. A.; Kunkel, L. M.: Identification of FLRT1, FLRT2, and FLRT3: a novel family of transmembrane leucine-rich repeat proteins. *Genomics* 62: 417–426, 1999.
- [68691] 7021. Andre, P.; Biassoni, R.; Colonna, M.; Cosman, D.; Lanier, L. L.; Long, E. O.; Lopez-Botet, M.; Moretta, A.; Moretta, L.; Parham, P.; Trowsdale, J.; Vivier, E.; Wagtmann, N.; Wilson, M. J.: New nomenclature for MHC receptors. *Nature Immun.* 2: 661 only, 2001.
- [68692] 7022. Fenster, S. D.; Chung, W. J.; Zhai, R.; Cases–

Langhoff, C.; Voss, B.; Garner, A. M.; Kaempfer, U.; Kindler, S.; Gundelfinger, E. D.; Garner, C. C.: Piccolo, a presynaptic zinc finger protein structurally related to Bassoon. *Neuron* 25: 203–214, 2000.

[68693] 7023. Cowan, C. A.; Henkemeyer, M.: The SH2/SH3 adaptor Grb4 transduces B-ephrin reverse signals. *Nature* 413: 174–179, 2001.

[68694] 7024. Hayashi, T.; Huang, J.; Deeb, S. S.: RINX(VSX1), a novel homeobox gene expressed in the inner nuclear layer of the adult retina. *Genomics* 67: 128–139, 2000.

[68695] 7025. Ohtoshi, A.; Justice, M. J.; Behringer, R. R.: Isolation and characterization of Vsx1, a novel mouse CVC paired-like homeobox gene expressed during embryogenesis and in the retina. *Biochem. Biophys. Res. Commun.* 286: 133–140, 2001.

[68696] 7026. Semina, E. V.; Mintz-Hittner, H. A.; Murray, J. C.: Isolation and characterization of a novel human paired-like homeodomain-containing transcription factor gene, VSX1, expressed in ocular tissues. *Genomics* 63: 289–293, 2000.

[68697] 7027. Jones, J. M.; Morrell, J. C.; Gould, S. J.: Identification and characterization of HAOX1, HAOX2, and HAOX3, three human peroxisomal  $\alpha$ -hydroxy acid oxidases. *J. Biol. Chem.* 275: 12590–12597, 2000.

- [68698] 7028.Kohler, S. A.; Menotti, E.; Kuhn, L. C.: Molecular cloning of mouse glycolate oxidase: high evolutionary conservation and presence of an iron-responsive element-like sequence in the mRNA. *J. Biol.Chem.* 274: 2401–2407, 1999.
- [68699] 7029.Williams, E.; Cregeen, D.; Rumsby, G.: Identification and expression of a cDNA for human glycolate oxidase. *Biochim. Biophys. Acta* 1493:246–248, 2000.
- [68700] 7030.Horikoshi, N.; Cong, J.; Kley, N.; Shenk, T.: Isolation of differentially expressed cDNAs from p53-dependent apoptotic cells: activation of the human homologue of the *Drosophila* peroxidase gene. *Biochem.Biophys. Res. Commun.* 261: 864–869, 1999.
- [68701] 7031.Weiler, S. R.; Taylor, S. M.; Deans, R. J.; Kan-Mitchell, J.; Mitchell, M. S.; Trent, J. M.: Assignment of a human melanoma associated gene MG50 (D2S448) to chromosome 2p25.3 by fluorescence in situ hybridization. *Genomics* 22:243–244, 1994.
- [68702] 7032.Deng, H.; Unutmaz, D.; KewalRamani, V. N.; Littman, D. R.: Expression cloning of new receptors used by simian and human immunodeficiency viruses. *Nature* 388: 296–300, 1997.
- [68703] 7033.Liao, F.; Alkhatib, G.; Peden, K. W. C.; Sharma, G.;

Berger, E.A.; Farber, J. M.: STRL33, a novel chemokine receptor-like protein, functions as a fusion cofactor for both macrophage-tropic and T cellline-tropic HIV-1. *J. Exp. Med.* 185: 2015–2023, 1997.

[68704] 7034. Matloubian, M.; David, A.; Engel, S.; Ryan, J. E.; Cyster, J. G.: A transmembrane CXC chemokine is a ligand for HIV-coreceptor Bonzo. *Nature Immun.* 1: 298–304, 2000.

[68705] 7035. Betz, R.; Gray, S. G.; Ekstrom, C.; Larsson, C.; Ekstrom, T. J.: Human histone deacetylase 2, HDAC2 (human RPD3), is localized to 6q21 by radiation hybrid mapping. *Genomics* 52: 245–246, 1998.

[68706] 7036. Inouye, C. J.; Seto, E.: Relief of YY1-induced transcriptional repression by protein-protein interaction with the nuclear phosphoprotein B23. *J. Biol. Chem.* 269: 6506–6510, 1994.

[68707] 7037. Randhawa, G. S.; Bell, D. W.; Testa, J. R.; Feinberg, A. P.: Identification and mapping of human histone acetylation modifier gene homologues. *Genomics* 51: 262–269, 1998.

[68708] 7038. Yang, W.-M.; Inouye, C.; Zeng, Y.; Bearss, D.; Seto, E.: Transcriptional repression by YY1 is mediated by interaction with a mammalian homolog of the yeast global regulator RPD3. *Proc. Nat. Acad. Sci.* 93: 12845–12850, 1996.



- [68709] 7039.Yarden, R. I.; Brody, L. C.: BRCA1 interacts with components of the histone deacetylase complex. *Proc. Nat. Acad. Sci.* 96: 4983–4988, 1999.
- [68710] 7040.Fedele, M.; Benvenuto, G.; Pero, R.; Majello, B.; Battista, S.; Lembo, F.; Vollono, E.; Day, P. M.; Santoro, M.; Lania, L.; Bruni, C. B.; Fusco, A.; Chiariotti, L.: A novel member of the BTB/POZ family, PATZ, associates with the RNF4 RING finger protein and acts as a transcriptional repressor. *J. Biol. Chem.* 275: 7894–7901, 2000.
- [68711] 7041.Kobayashi, A.; Yamagiwa, H.; Hoshino, H.; Muto, A.; Sato, K.; Morita, M.; Hayashi, N.; Yamamoto, M.; Igarashi, K.: A combinatorial code for gene expression generated by transcription factor Bach2 and MAZR (MAZ-related factor) through the BTB/POZ domain. *Molec. Cell. Biol.* 20:1733–1746, 2000.
- [68712] 7042.McAleer, M. A.; Breen, M. A.; White, N. L.; Matthews, N.: pABC11 (also known as MOAT-C and MRP5), a member of the ABC family of proteins, has anion transporter activity but does not confer multidrug resistance when overexpressed in human embryonic kidney 293 cells. *J. Biol. Chem.* 274: 23541–23548, 1999.
- [68713] 7043.Oguri, T.; Isobe, T.; Suzuki, T.; Nishio, K.; Fujiwara, Y.; Katoh, O.; Yamakido, M.: Increased expression of the

MRP5 gene is associated with exposure to platinum drugs in lung cancer. *Int. J. Cancer* 86:95–100, 2000.

- [68714] 7044. Suzuki, T.; Nishio, K.; Sasaki, H.; Kurokawa, H.; Saito–Ohara, F.; Ikeuchi, T.; Tanabe, S.; Terada, M.; Saijo, N.: cDNA cloning of a short type of multidrug resistance protein homologue, SMRP, from a human lung cancer cell line. *Biochem. Biophys. Res. Commun.* 238:790–794, 1997.
- [68715] 7045. Suzuki, T.; Sasaki, H.; Kuh, H.–J.; Agui, M.; Tatsumi, Y.; Tanabe, S.; Terada, M.; Saijo, N.; Nishio, K.: Detailed structural analysis on both human MRP5 and mouse mrp5 transcripts. *Gene* 242: 167–173, 2000.
- [68716] 7046. Wijnholds, J.; Mol, C. A. A. M.; van Deemter, L.; de Haas, M.; Scheffer, G. L.; Baas, F.; Beijnen, J. H.; Scheper, R. J.; Hatse, S.; De Clercq, E.; Balzarini, J.; Borst, P.: Multidrug–resistance protein 5 is a multispecific organic anion transporter able to transport nucleotide analogs. *Proc. Nat. Acad. Sci.* 97: 7476–7481, 2000.
- [68717] 7047. Bebenek, K.; Tissier, A.; Frank, E. G.; McDonald, J. P.; Prasad, R.; Wilson, S. H.; Woodgate, R.; Kunkel, T. A.: 5–prime–deoxyribosephosphate lyase activity of human DNA polymerase  $\iota$  in vitro. *Science* 291:2156–2159, 2001.

- [68718] 7048.Johnson, R. E.; Washington, M. T.; Haracska, L.; Prakash, S.; Prakash,L.: Eukaryotic polymerases iota and zeta act sequentially to bypassDNA lesions. *Nature* 406: 1015–1019, 2000.
- [68719] 7049.McDonald, J. P.; Rasic–Otrin, V.; Epstein, J. A.; Broughton, B.C.; Wang, X.; Lehmann, A. R.; Wolgemuth, D. J.; Woodgate, R.: Novelhuman and mouse homologs of *Saccharomyces cerevisiae* DNA polymeraseeta. *Genomics* 60: 20–30, 1999.
- [68720] 7050.Dermaut, B.; Theuns, J.; Sleegers, K.; Hasegawa, H.; Van den Broeck,M.; Vennekens, K.; Corsmit, E.; St. George–Hyslop, P.; Cruts, M.;van Duijn, C. M.; Van Broeckhoven, C.: The gene encoding nicastrin,a major gamma–secretase component, modifies risk for familial early–onsetAlzheimer disease in a Dutch population–based sample. *Am. J. Hum.Genet.* 70: 1568–1574, 2002.
- [68721] 7051.Feldman, R. G.; Chandler, K. A.; Levy, L. L.; Glaser, G. H.: FamilialAlzheimer's disease. *Neurology* 13: 811–824, 1963.
- [68722] 7052.Foncin, J.–F.; Salmon, D.; Supino–Viterbo, V.; Feld–man, R. G.;Macchi, G.; Mariotti, P.; Scoppetta, C.; Caruso, G.; Bruni, A. C.: Demence presenile d'Alzheimer transmise dans une famille etendue. *Rev.Neurol. (Paris)* 141:

194–202, 1985.

- [68723] 7053.Hiltunen, M.; Mannermaa, A.; Thompson, D.; Easton, D.; Pirskanen,M.; Helisalmi, S.; Koivisto, A. M.; Lehtovirta, M.; Ryyananen, M.;Soininen, H.: Genome-wide linkage disequilibrium mapping of late-onsetAlzheimer's disease in Finland. *Neurology* 57: 1663–1668, 2001.
- [68724] 7054.Kehoe, P.; Wavrant–De Vrieze, F.; Crook, R.; Wu, W. S.; Holmans,P.; Fenton, I.; Spurlock, G.; Norton, N.; Williams, H.; Williams,N.; Lovestone, S.; Perez–Tur, J.; Hutton, J.; and 10 others: Afull genome scan for late onset Alzheimer disease. *Hum. Molec. Genet.* 8:237–245, 1999.
- [68725] 7055.Kopan, R.; Goate, A.: Aph–2/nicastrin: an essential componentof gamma–secretase and regulator of Notch signaling and presenilinlocalization. *Neuron* 33: 321–324, 2002.
- [68726] 7056.Yu, G.; Nishimura, M.; Arawaka, S.; Levitan, D.; Zhang, L.; Tandon,A.; Song, Y.–Q.; Rogaeva, E.; Chen, F.; Kawarai, T.; Supala, A.; Levesque,L.; and 18 others: Nicas–trin modulates presenilin–mediated notch/glp–1signal transduction and beta–APP processing. *Nature* 407: 48–54,2000.
- [68727] 7057.Zubenko, G. S.; Hughes, H. B.; Stiffler, J. S.; Hurtt, M. R.; Kaplan,B. B.: A genome survey for novel Alzheimer dis–

ease risk loci: results at 10-cM resolution. *Genomics* 50: 121–128, 1998.

[68728] 7058. Tateishi, S.; Sakuraba, Y.; Masuyama, S.; Inoue, H.; Yamaizumi, M.: Dysfunction of human Rad18 results in defective postreplication repair and hypersensitivity to multiple mutagens. *Proc. Nat. Acad. Sci.* 97: 7927–7932, 2000.

[68729] 7059. Arakawa, H.; Hauschild, J.; Buerstedde, J.-M.: Requirement of the activation-induced deaminase (AID) gene for immunoglobulin gene conversion. *Science* 295: 1301–1306, 2002.

[68730] 7060. Fagarasan, S.; Kinoshita, K.; Muramatsu, M.; Ikuta, K.; Honjo, T.: In situ class switching and differentiation to IgA producing cells in the gut lamina propria. *Nature* 413: 639–643, 2001.

[68731] 7061. Muramatsu, M.; Kinoshita, K.; Fagarasan, S.; Yamada, S.; Shinkai, Y.; Honjo, T.: Class switch recombination and hypermutation require activation-induced cytidine deaminase (AID), a potential RNA editing enzyme. *Cell* 102: 553–563, 2000.

[68732] 7062. Muramatsu, M.; Sankaranand, V. S.; Anant, S.; Sugai, M.; Kinoshita, K.; Davidson, N. O.; Honjo, T.: Specific expression of activation-induced cytidine deaminase (AID), a

novel member of the RNA-editing deaminase family in germinal center B cells. *J. Biol. Chem.* 274: 18470–18476, 1999.

- [68733] 7063. Rowley, D. R.; Tindall, D. J.: Responses of NBT-II bladder carcinoma cells to conditioned medium from normal fetal urogenital sinus. *Cancer Res.* 47: 2955–2960, 1987.
- [68734] 7064. Kasaian, M. T.; Whitters, M. J.; Carter, L. L.; Lowe, L. D.; Jussif, J. M.; Deng, B.; Johnson, K. A.; Witek, J. S.; Senices, M.; Konz, R. F.; Wurster, A. L.; Donaldson, D. D.; Collins, M.; Young, D. A.; Grusby, M. J.: IL-21 limits NK cell responses and promotes antigen-specific T cell activation: a mediator of the transition from innate to adaptive immunity. *Immunity* 16: 559–569, 2002.
- [68735] 7065. Pashmforoush, M.; Pomies, P.; Peterson, K. L.; Kubalak, S.; Ross, J., Jr.; Hefti, A.; Aebi, U.; Beckerle, M. C.; Chien, K. R.: Adult mice deficient in actinin-associated LIM-domain protein reveal a developmental pathway for right ventricular cardiomyopathy. *Nature Med.* 7: 591–597, 2001.
- [68736] 7066. Zheng, P.; Eastman, J.; Vande Pol, S.; Pimplikar, S. W.: PAT1, a microtubule-interacting protein, recognizes the basolateral sorting signal of amyloid precursor protein.

Proc. Nat. Acad. Sci. 95: 14745–14750,1998.

- [68737] 7067.Aoyama, T.; Yamano, S.; Waxman, D. J.; Lapenson, D. P.; Meyer,U. A.; Fischer, V.; Tyndale, R.; Inaba, T.; Kalow, W.; Gelboin, H.V.; Gonzalez, F. J.: Cytochrome P-450 hPCN3, a novel cytochrome P-450IIIA product that is differentially expressed in adult human liver:cDNA and deduced amino acid sequence and distinct specificities ofcDNA-expressed hPCN1 and hPCN3 for the metabolism of steroid hormonesand cyclosporine. J. Biol. Chem. 264: 10388–10395, 1989.
- [68738] 7068.Jounaidi, Y.; Guzelian, P. S.; Maurel, P.; Vilarem, M.-J.: Sequenceof the 5-prime-flanking region of CYP3A5: comparative analysis withCYP3A4 and CYP3A7. Biochem. Biophys. Res. Commun. 205: 1741–1747,1994.
- [68739] 7069.Kuehl, P.; Zhang, J.; Lin, Y.; Lamba, J.; Assem, M.; Schuetz, J.;Watkins, P. B.; Daly, A.; Wrighton, S. A.; Hall, S. D.; Maurel, P.;Relling, M.; Brimer, C.; Yasuda, K.; Venkataramanan, R.; Strom, S.;Thummel, K.; Boguski, M. S.; Schuetz, E.: Sequence diversity in CYP3A promoters and characterization of the genetic basis of polymorphic-CYP3A5 expression. Nature Genet. 27: 383–391, 2001.
- [68740] 7070.Paulussen, A.; Lavrijsen, K.; Bohets, H.; Hendrickx, J.; Verhasselt,P.; Luyten, W.; Konings, F.; Armstrong, M.:

Two linked mutations in transcriptional regulatory elements of the CYP3A5 gene constitute the major genetic determinant of polymorphic activity in humans. *Pharmacogenetics* 10:415–424, 2000.

- [68741] 7071. Yamakoshi, Y.; Kishimoto, T.; Sugimura, K.; Kawashima, H.: Human prostate CYP3A5: identification of a unique 5-prime-untranslated sequence and characterization of purified recombinant protein. *Biochem. Biophys. Res. Commun.* 260: 676–681, 1999.
- [68742] 7072. De Valck, D.; Jin, D. Y.; Heyninck, K.; Van de Craen, M.; Contreras, R.; Fiers, W.; Jeang, K. T.; Beyaert, R.: The zinc finger protein A20 interacts with a novel anti-apoptotic protein which is cleaved by specific caspases. *Oncogene* 18: 4182–4190, 1999.
- [68743] 7073. Gachon, F.; Peleraux, A.; Thebault, S.; Dick, J.; Lemasson, I.; Devaux, C.; Mesnard, J.-M.: CREB-2, a cellular CRE-dependent transcription repressor, functions in association with tax as an activator of the human T-cell leukemia virus type 1 promoter. *J. Virol.* 72: 8332–8337, 1998.
- [68744] 7074. Ling, L.; Goeddel, D. V.: T6BP, a TRAF6-interacting protein involved in IL-1 signaling. *Proc. Nat. Acad. Sci.* 97: 9567–9572, 2000.



- [68745] 7075.Ozaki, K.; Kikly, K.; Michalovich, D.; Young, P. R.; Leonard, W.J.: Cloning of a type I cytokine receptor most related to the IL-2receptor beta chain. *Proc. Nat. Acad. Sci.* 97: 11439–11444, 2000.
- [68746] 7076.Parrish–Novak, J.; Dillon, S. R.; Nelson, A.; Hammond, A.; Sprecher,C.; Gross, J. A.; Johnston, J.; Madden, K.; Xu, W.; West, J.; Schrader,S.; Burkhead, S.; and 26 others: Interleukin 21 and its receptorare involved in NK cell expansion and regulation of lymphocyte function. *Nature* 408:57–63, 2000.
- [68747] 7077.Boyartchuk, V. L.; Ashby, M. N.; Rine, J.: Modulation of Ras anda–factor function by carboxyl–terminal proteolysis. *Science* 275:1796–1800, 1997.
- [68748] 7078.Freije, J. M. P.; Blay, P.; Pendas, A. M.; Cadinanos, J.; Crespo,P.; Lopez–Otin, C.: Identification and chromosomal location of twohuman genes encoding enzymes potentially involved in proteolytic maturationof farnesylated proteins. *Genomics* 58: 270–280, 1999.
- [68749] 7079.Otto, J. C.; Kim, E.; Young, S. G.; Casey, P. J.: Cloning andcharacterization of a mammalian prenyl protein–specific protease. *J.Biol. Chem.* 274: 8379–8382, 1999.
- [68750] 7080.Reither, A.; Hehlmann, R.; Goldman, J. M.; Cross, N. C.: The 8p11myeloproliferative syndrome. *Med. Klin.* 94:

207–210, 1999.

- [68751] 7081. Bartholin, L.; Maguer-Satta, V.; Hayette, S.; Martel, S.; Gadoux, M.; Bertrand, S.; Corbo, L.; Lamadon, C.; Morera, A.-M.; Magaud, J.-P.; Rimokh, R.: FLRG, an activin-binding protein, is a new target of TGF-beta transcription activation through Smad proteins. *Oncogene* 20:5409–5419, 2001.
- [68752] 7082. Rimokh, R.; Berger, F.; Delsol, G.; Charrin, C.; Bertheas, M. F.; Ffrench, M.; Garoscio, M.; Felman, P.; Coiffier, C.; Bryon, P. A.: Rearrangement and overexpression of the BCL-1/PRAD-1 gene in intermediatelymphocytic lymphomas and in t(11q13)-bearing leukemias. *Blood* 81:3063–3067, 1993.
- [68753] 7083. Bahou, W. F.; Campbell, A. D.; Wicha, M. S.: cDNA cloning and molecular characterization of MSE55, a novel human serum constituent protein that displays bone marrow stromal/endothelial cell-specific expression. *J. Biol. Chem.* 267: 13986–13992, 1992.
- [68754] 7084. Burbelo, P. D.; Drechsel, D.; Hall, A.: A conserved binding motif defines numerous candidate target proteins for both Cdc42 and Rac GTPases. *J. Biol. Chem.* 270: 29071–29074, 1995.
- [68755] 7085. Bak, M.; Hansen, C.; Henriksen, K. F.; Tommerup, N.:

The human hedgehog-interacting protein gene: structure and chromosome mapping to 4q31.21–q31.3. *Cytogenet. Cell Genet.* 92: 300–303, 2001.

- [68756] 7086. Chuang, P.-T.; McMahon, A. P.: Vertebrate hedgehog signalling modulated by induction of a hedgehog-binding protein. *Nature* 397:617–621, 1999.
- [68757] 7087. Bacher, N.; Zisman, Y.; Berent, E.; Livneh, E.: Isolation and characterization of PKC-L, a new member of the protein kinase C-related gene family specifically expressed in lung, skin, and heart. *Molec. Cell. Biol.* 11: 126–133, 1991.
- [68758] 7088. Quan, T.; Fisher, G. J.: Cloning and characterization of the human protein kinase C- $\eta$  promoter. *J. Biol. Chem.* 274: 28566–28574, 1999.
- [68759] 7089. Gong, T.-W. L.; Burmeister, M.; Lomax, M. I.: The novel gene D4Mit1 maps to mouse chromosome 4 and human chromosome 1p36. *Mammalian Genome* 7:790–791, 1996.
- [68760] 7090. Nangaku, M.; Sato-Yoshitake, R.; Okada, Y.; Noda, Y.; Takemura, R.; Yamazaki, H.; Hirokawa, N.: KIF1B, a novel microtubule plus end-directed monomeric motor protein for transport of mitochondria. *Cell* 79:1209–1220, 1994.

- [68761] 7091.Saito, M.; Hayashi, Y.; Suzuki, T.; Tanaka, H.; Hozumi, I.; Tsuji,S.: Linkage mapping of the gene for Charcot–Marie–Tooth disease type2 to chromosome 1p (CMT2A) and the clinical features of CMT2A. *Neurology* 49:1630–1635, 1997.
- [68762] 7092.Yang, H. W.; Chen, Y. Z.; Takita, J.; Soeda, E.; Piao, H. Y.; Hayashi,Y.: Genomic structure and mutational analysis of the human KIF1Bgene which is homozygously deleted in neuroblastoma at chromosome1p36.2. *Oncogene* 20: 5075–5083, 2001.
- [68763] 7093.Suzuki, Y.; Demoliere, C.; Kitamura, D.; Takeshita, H.; Deuschle,U.; Watanabe, T.: HAX–1, a novel intracellular protein, localizedon mitochondria, directly associates with HS1, a substrate of Srcfamily tyrosine kinases. *J. Immun.* 158: 2736–2744, 1997.
- [68764] 7094.Dell'Angelica, E. C.; Puertollano, R.; Mullins, C.; Aguilar, R.C.; Vargas, J. D.; Hartnell, L. M.; Bonifacino, J. S.: GGAs: a familyof ADP ribosylation factor–binding proteins related to adaptors andassociated with the Golgi complex. *J. Cell Biol.* 149: 81–93, 2000.
- [68765] 7095.Hirst, J.; Lui, W. W. Y.; Bright, N. A.; Totty, N.; Seaman, M.N. J.; Robinson, M. S.: A family of proteins with gamma–adaptin andVHS domains that facilitate trafficking

between the trans-Golgi network and the vacuole/lysosome. *J. Cell Biol.* 149: 67–69, 2000.

[68766] 7096. Ma, Z.; Morris, S. W.; Valentine, V.; Li, M.; Herbrick, J.-A.; Cui, X.; Bouman, D.; Li, Y.; Mehta, P. K.; Nizetic, D.; Kaneko, Y.; Chan, G. C. F.; Chan, L. C.; Squire, J.; Scherer, S. W.; Hitzler, J. K.: Fusion of two novel genes, RBM15 and MKL1, in the t(1;22)(p13;q13) of acute megakaryoblastic leukemia. *Nature Genet.* 28: 220–221, 2001.

[68767] 7097. Mercher, T.; Coniat, M. B.-L.; Monni, R.; Mauchauffe, M.; Khac, F. N.; Gressin, L.; Mugneret, F.; Leblanc, T.; Dastugue, N.; Berger, R.; Bernard, O. A.: Involvement of a human gene related to the *Drosophila* spen gene in the recurrent t(1;22) translocation of acute megakaryocytic leukemia. *Proc. Nat. Acad. Sci.* 98: 5776–5779, 2001.

[68768] 7098. Wang, D.-Z.; Chang, P. S.; Wang, Z.; Sutherland, L.; Richardson, J. A.; Small, E.; Krieg, P. A.; Olson, E. N.: Activation of cardiac gene expression by myocardin, a transcriptional cofactor for serum response factor. *Cell* 105: 851–862, 2001.

[68769] 7099. Blumberg, H.; Conklin, D.; Xu, W.; Grossmann, A.; Brender, T.; Carollo, S.; Eagan, M.; Foster, D.; Haldeman, B. A.; Hammond, A.; Haugen, H.; Jelinek, L.; and 14 others: Interleukin 20: discovery, receptor identification, and role

in epidermal function. Cell 104:9–19, 2001.

- [68770] 7100.Huang, E. Y.; Madireddi, M. T.; Gopalkrishnan, R. V.; Leszczyniecka,M.; Su, Z.; Lebedeva, I. V.; Kang, D.; Jiang, H.; Lin, J. J.; Alexandre,D.; Chen, Y.; Vozhilla, N.: and 9 others: Genomic structure, chromosomallocalization and expression profile of a novel melanoma differentiationas-sociated (mda-7) gene with cancer specific growth sup-pressing andapoptosis inducing properties. Oncogene 20: 7051–7063, 2001.
- [68771] 7101.Jiang, H.; Lin, J. J.; Su, Z.–Z.; Goldstein, N. I.; Fisher, P.B.: Subtraction hybridization identifies a novel melanoma differentiationassociated gene, mda-7, modu-lated during human melanoma differentiation,growth and progression. Oncogene 11: 2477–2486, 1995.
- [68772] 7102.Jiang, H.; Su, Z.–Z.; Lin, J. J.; Goldstein, N. I.; Young, C. S.H.; Fisher, P. B.: The melanoma differentiation associ-ated gene mda-7suppresses cancer cell growth. Proc. Nat. Acad. Sci. 93: 9160–9165,1996.
- [68773] 7103.Su, Z.; Madireddi, M. T.; Lin, J. J.; Young, C. S. H.; Ki-tada,S.; Reed, J. C.; Goldstein, N. I.; Fisher, P. B.: The can-cer growthsuppressor gene mda-7 selectively induces apoptosis in human breastcancer cells and inhibits tumor growth in nude mice. Proc. Nat. Acad.Sci. 95:

14400–14405, 1998.

- [68774] 7104.Hsieh, S.–L.; Campbell, R. D.: Evidence that gene G7a in the humanmajor histocompatibility complex encodes valyl–tRNA synthetase. *Biochem.J.* 278: 809–816, 1991. Note: Erratum: *Biochem. J.* 281: 879 only,1992.
- [68775] 7105.Hsieh, S.–L.; Kendall, E.; Milner, C.; Cross, S.; Cheng, J.; Khanna,A.; Olaversen, M.; Campbell, R. D.: Cloning of the human valyl–tRNA synthetase gene and its location in the major histocompatibility complex on chromosome 6. (Abstract) *Cytogenet. Cell Genet.* 58: 1912 only,1991.
- [68776] 7106.Mansour, S. J.; Skaug, J.; Zhao, X.–H.; Giordano, J.; Scherer,S. W.; Melancon, P.: p200 ARF–GEP1: a Golgi–localized guanine nucleotideexchange protein whose Sec7 domain is targeted by the drug brefeldinA. *Proc. Nat. Acad. Sci.* 96: 7968–7973, 1999.
- [68777] 7107.Morinaga, N.; Moss, J.; Vaughan, M.: Cloning and expression of a cDNA encoding a bovine brain brefeldin A–sensitive guanine nucleotide–exchangeprotein for ADP–ribosylation factor. *Proc. Nat. Acad. Sci.* 94: 12926–12931,1997.
- [68778] 7108.Morinaga, N.; Tsai, S.–C.; Moss, J.; Vaughan, M.: Isolation of a brefeldin A–inhibited guanine nucleotide–exchange protein for ADPribosylation factor (ARF) 1 and

ARF3 that contains a Sec7-like domain. *Proc.Nat. Acad. Sci.* 93: 12856–12860, 1996.

- [68779] 7109.Togawa, A.; Morinaga, N.; Ogasawara, M.; Moss, J.; Vaughan, M.: Purification and cloning of a brefeldin A-inhibited guanine nucleotide-exchange protein for ADP-ribosylation factors. *J. Biol. Chem.* 274: 12308–12315, 1999.
- [68780] 7110.Mitsui, S.; Tsuruoka, N.; Yamashiro, K.; Nakazato, H.; Yamaguchi, N.: A novel form of human neuropsin, a brain-related serine protease, is generated by alternative splicing and is expressed preferentially in human adult brain. *Europ. J. Biochem.* 260: 627–634, 1999.
- [68781] 7111.Underwood, L. J; Tanimoto, H.; Wang, Y.; Shigemasa, K.; Parmley, T. H.; O'Brien, T. J.: Cloning of tumor-associated differentially expressed gene-14, a novel serine protease overexpressed by ovarian carcinoma. *Cancer Res.* 59: 4435–4439, 1999.
- [68782] 7112.Yoshida, S.; Taniguchi, M.; Hirata, A.; Shiosaka, S.: Sequence analysis and expression of human neuropsin cDNA and gene. *Gene* 213:9–16, 1998.
- [68783] 7113.Markert, M. L.; Hershfield, M. S.; Schiff, R. I.; Buckley, R.H.: Adenosine deaminase and purine nucleoside phosphorylase deficiencies: evaluation of therapeutic in-



terventions in eight patients. J. Clin.Immun. 7: 389–399, 1987.

[68784] 7114.Ewen, M. E.; Xing, Y.; Lawrence, J. B.; Livingston, D. M.: Molecularcloning, chromosomal mapping, and expression of the cDNA for p107,a retinoblastoma gene product-related protein. Cell 66: 1155–1164,1991.

[68785] 7115.Huppi, K.; Siwarski, D.; Mock, B. A.; Dosik, J.; Hamel, P. A.:Molecular cloning, chromosomal mapping, and expression of the mousep107 gene. Mammalian Genome 7: 353–355, 1996.

[68786] 7116.Kim, K. K.; Soonpaa, M. H.; Wang, H.; Field, L. J.: Developmentalexpression of p107 mRNA and evidence for alternative splicing of thep107 (RBL1) gene product. Genomics 28: 520–529, 1995.

[68787] 7117.LeCouter, J. E.; Kablar, B.; Hardy, W. R.; Ying, C.; Megeney, L.A.; May, L. L.; Rudnicki, M. A.: Strain-dependent myeloid hyperplasia,growth deficiency, and accelerated cell cycle in mice lacking theRb-related p107 gene. Molec. Cell. Biol. 18: 7455–7465, 1998.

[68788] 7118.Morohashi, K.; Sogawa, K.; Omura, T.; Fujii-Kuriyama, Y.: Genestructure of human cytochrome P-450(SCC), cholesterol desmolase. J.Biochem. 101: 879–887, 1987.

- [68789] 7119.Youngblood, G. L.; Nesbitt, M. N.; Payne, A. H.: The structuralgenes encoding P450SCC and P450AROM are closely linked on mouse chromosome9. *Endocrinology* 125: 2784–2786, 1989.
- [68790] 7120.Barrard, B. A.; Lottspeich, F.; Braun, A.; Barde, Y. A.; Mallet,J.: cDNA cloning and complete sequence of porcine choline acetyltransferase:in vitro translation of the corresponding RNA yields an active protein. *Proc.Nat. Acad. Sci.* 84: 9280–9284, 1987.
- [68791] 7121.Chireux, M. A.; Le Van Thai, A.; Weber, M. J.: Human choline acetyltransferasegene: localization of alternative first exons. *J. Neurosci. Res.* 40:427–438, 1995.
- [68792] 7122.Cohen–Haguenauer, O.; Brice, A.; Berrard, S.; Van Cong, N.; Mallet,J.; Frezal, J.: Localization of the choline acetyltransferase (CHAT)gene to human chromosome 10. *Genomics* 6: 374–378, 1990.
- [68793] 7123.Misawa, H.; Ishii, K.; Deguchi, T.: Gene expression of mouse cholineacetyltransferase: alternative splicing and identification of a highlyactive promoter region. *J. Biol. Chem.* 267: 20392–20399, 1992.
- [68794] 7124.Ohno, K.; Tsujino, A.; Brengman, J. M.; Harper, C. M.; Bajzer,Z.; Udd, B.; Beyring, R.; Robb, S.; Kirkham, F. J.; Engel, A. G.:Choline acetyltransferase mutations cause

myasthenic syndrome associated with episodic apnea in humans. Proc. Nat. Acad. Sci. 98: 2017–2022, 2001.

[68795] 7125. Strauss, W. L.; Kemper, R. R.; Jayakar, P.; Kong, C. F.; Hersh, L. B.; Hilt, D. C.; Rabin, M.: Human choline acetyltransferase gene maps to region 10q11–q22.2 by in situ hybridization. Genomics 9:396–398, 1991.

[68796] 7126. Toussaint, J. L.; Geoffroy, V.; Schmitt, M.; Werner, A.; Garnier, J. M.; Simoni, P.; Kempf, J.: Human choline acetyltransferase (CHAT): partial gene sequence and potential control regions. Genomics 12:412–416, 1992.

[68797] 7127. Viegas-Pequignot, E.; Berrard, S.; Brice, A.; Apiou, F.; Mallet, J.: Localization of a 900-bp-long fragment of the human choline acetyltransferase gene to 10q11.2 by non-radioactive in situ hybridization. Genomics 9:210–212, 1991.

[68798] 7128. Bonner, T. I.; Modi, W. S.; Seunemann, H. N.; O'Brien, S. J.: Chromosomal mapping of five human genes encoding muscarinic acetylcholine receptors. (Abstract) Cytogenet. Cell Genet. 58: 1850–1851, 1991.

[68799] 7129. Anand, R.; Lindstrom, J.: Chromosomal localization of seven neuronal nicotinic acetylcholine receptor subunit genes in humans. Genomics 13:962–967, 1992.

[68800] 7130. Bessis, A.; Simon-Chazottes, D.; Devillers-Thiery,

A.; Guenet, J.-L.; Changeux, J.-P.: Chromosomal localization of the mouse genes coding for alpha-2, alpha-3, alpha-4 and beta-2 subunits of neuronal nicotinic acetylcholine receptor. *FEBS Lett.* 264: 48-52, 1990.

[68801] 7131. Boulter, J.; O'Shea-Greenfield, A.; Duvoisin, R. M.; Connolly, J. G.; Wada, E.; Jensen, A.; Gardner, P. D.; Ballivet, M.; Deneris, E. S.; McKinnon, D.; Heinemann, S.; Patrick, J.: Alpha3, alpha5 and beta4: three members of the rat neuronal nicotinic acetylcholine receptor-related gene family form a gene cluster. *J. Biol. Chem.* 265: 4472-4482, 1990.

[68802] 7132. Duga, S.; Solda, G.; Asselta, R.; Bonati, M. T.; Dalpra, L.; Malcovati, M.; Tenchini, M. L.: Characterization of the genomic structure of the human neuronal nicotinic acetylcholine receptor CHRNA5/A3/B4 gene cluster and identification of novel intragenic polymorphisms. *J. Hum. Genet.* 46: 640-648, 2001.

[68803] 7133. Elliott, K. J.; Ellis, S. B.; Berckhan, K. J.; Urrutia, A.; Chavez-Noriega, L. E.; Johnson, E. C.; Velicelebi, G.; Harpold, M. M.: Comparative structure of human neuronal alpha-2-alpha-7 and beta-2-beta-4 nicotinic acetylcholine receptor subunits and functional expression of the alpha-2, alpha-3, alpha-4, alpha-7, beta-2, and beta-4 subunits. *J. Molec. Neurosci.* 7: 217-228, 1996.

- [68804] 7134. Armstrong, E.; Partanen, J.; Cannizzaro, L.; Huebner, K.; Alitalo, K.: Localization of the fibroblast growth factor receptor-4 gene to chromosome region 5q33-qter. *Genes Chromosomes Cancer* 4: 94–98, 1992.
- [68805] 7135. Bange, J.; Prechtel, D.; Cheburkin, Y.; Specht, K.; Harbeck, N.; Schmitt, M.; Knyazeva, T.; Muller, S.; Gartner, S.; Sures, I.; Wang, H.; Imyanitov, E.; Haring, H.-U.; Knayzev, P.; Iacobelli, S.; Hofler, H.; Ullrich, A.: Cancer progression and tumor cell motility are associated with the FGFR4 Arg388 allele. *Cancer Res.* 62: 840–847, 2002.
- [68806] 7136. Holtrich, U.; Brauninger, A.; Strebhardt, K.; Rubsamens-Waigmann, H.: Two additional protein-tyrosine kinases expressed in human lung: fourth member of the fibroblast growth factor receptor family and an intracellular protein-tyrosine kinase. *Proc. Nat. Acad. Sci.* 88: 10411–10415, 1991.
- [68807] 7137. Kostrzewa, M.; Muller, U.: Genomic structure and complete sequence of the human FGFR4 gene. *Mammalian Genome* 9: 131–135, 1998.
- [68808] 7138. Partanen, J.; Makela, T. P.; Eerola, E.; Korhonen, J.; Hirvonen, H.; Claesson-Welsh, L.; Alitalo, K.: FGFR-4, a novel acidic fibroblast growth factor receptor with a distinct expression pattern. *EMBO J.* 10: 1347–1354, 1991.

- [68809] 7139.Scott, A. F.: Personal Communication. Baltimore, Md. 10/12/1999.
- [68810] 7140.Vainikka, S.; Partanen, J.; Bellosta, P.; Coulier, F.; Basilico,C.; Jaye, M.; Alitalo, K.: Fibroblast growth factor receptor-4 shows novel features in genomic structure, ligand binding and signal transduction. EMBOJ. 11: 4273-4280, 1992.
- [68811] 7141.Warrington, J. A.; Bailey, S. K.; Armstrong, E.; Aprelikova, O.;Alitalo, K.; Dolganov, G. M.; Wilcox, A. S.; Sikela, J. M.; Wolfe,S. F.; Lovett, M.; Wasmuth, J. J.: A radiation hybrid map of 18 growthfactor, growth factor receptor, hormone receptor, or neurotransmitterreceptor genes on the distal region of the long arm of chromosome5. Genomics 13: 803-808, 1992.
- [68812] 7142.Becker, P. E.; Von Knorre, G. V.: Myositis ossificans progressiva. Ergeb.Inn. Med. Kinderheilk. 27: 1-31, 1968.
- [68813] 7143.Beratis, N. G.; Kaffe, S.; Aron, A. M.; Hirschhorn, K.: Alkalinephosphatase activity in cultured skin fibroblasts from fibrodysplasiaossificans progressiva. J. Med. Genet. 13: 307-309, 1976.
- [68814] 7144.Cohen, R. B.; Hahn, G. V.; Tabas, J. A.; Peeper, J.; Levitz, C.L.; Sando, A.; Sando, N.; Zasloff, M.; Kaplan, F. S.: The naturalhistory of heterotopic ossification in patients

who have fibrodysplasia ossificans progressiva: a study of forty-four patients. J. Bone Joint Surg. 75A: 215–219, 1993.

- [68815] 7145. Connor, J. M.: Soft Tissue Ossification. New York: Springer-Verlag(pub.) 1983.
- [68816] 7146. Connor, J. M.; Evans, D. A. P.: Genetic aspects of fibrodysplasia ossificans progressiva. J. Med. Genet. 19: 35–39, 1982.
- [68817] 7147. Connor, J. M.; Evans, D. A. P.: Fibrodysplasia ossificans progressiva: the clinical features and natural history of 34 patients. J. Bone Joint Surg. 64: 76–83, 1982.
- [68818] 7148. Connor, J. M.; Skirton, H.; Lunt, P. W.: A three generation family with fibrodysplasia ossificans progressiva. J. Med. Genet. 30: 687–689, 1993.
- [68819] 7149. Daltroff, G.; Lutz, P.; Bellocq, P.; Christmann, D.; Flamant, F.; Lenoir, G.; Clavert, J. M.; Levy, J. M.: Fibromatose et fibrodysplasie ossifiante progressive: une erreur diagnostique evitable. Arch. Fr. Pediat. 49: 441–444, 1992.
- [68820] 7150. Feldman, G.; Li, M.; Martin, S.; Urbanek, M.; Urtizberea, J. A.; Fardeau, M.; LeMerrer, M.; Connor, J. M.; Triffitt, J.; Smith, R.; Muenke, M.; Kaplan, F. S.; Shore, E. M.: Fibrodysplasia ossificans progressiva, a heritable disorder

of severe heterotopic ossification, maps to human chromosome 4q27–31. *Am. J. Hum. Genet.* 66: 128–135, 2000.

[68821] 7151. Janoff, H. B.; Muenke, M.; Johnson, L. O.; Rosenberg, A.; Shore, E. M.; Okereke, E.; Zasloff, M.; Kaplan, F. S.: Fibrodysplasia ossificans progressiva in two half-sisters: evidence for maternal mosaicism. *Am. J. Med. Genet.* 61: 320–324, 1996.

[68822] 7152. Digicaylioglu, M.; Lipton, S. A.: Erythropoietin-mediated neuroprotection involves cross-talk between Jak2 and NF-kappa-B signalling cascades. *Nature* 412: 641–647, 2001.

[68823] 7153. Clancy, D. J.; Gems, D.; Harshman, L. G.; Oldham, S.; Stocker, H.; Hafen, E.; Leivers, S. J.; Partridge, L.: Extension of life-span by loss of CHICO, a *Drosophila* insulin receptor substrate protein. *Science* 292: 104–106, 2001.

[68824] 7154. Chai, K. X.; Chen, L.-M.; Chao, J.; Chao, L.: Kallistatin: a novel human serine proteinase inhibitor: molecular cloning, tissue distribution, and expression in *Escherichia coli*. *J. Biol. Chem.* 268: 24498–24505, 1993.

[68825] 7155. Chai, K. X.; Ward, D. C.; Chao, J.; Chao, L.: Molecular cloning, sequence analysis, and chromosomal localization of the human protease inhibitor 4 (kallistatin) gene (PI4). *Genomics* 23: 370–378, 1994.



- [68826] 7156.Zhou, G. X.; Chao, L.; Chao, J.: Kallistatin: a novel human tissue kallikrein inhibitor; purification, characterization, and reactive center sequence. *J. Biol. Chem.* 267: 25873–25880, 1992.
- [68827] 7157.Christmanson, L.; Rorsman, F.; Stenman, G.; Westermark, P.; Betsholtz, C.: The human islet amyloid polypeptide (IAPP) gene: organization, chromosomal localization and functional identification of a promoter region. *FEBS Lett.* 267: 160–166, 1990.
- [68828] 7158.Cockburn, D. C.; Holt, S. M.; Roberts, A. N.; Cooper, G. J. S.; Reid, K. B. M.; Boyd, Y.: Localization of the amylin locus to chromosome 12. (Abstract) *Cytogenet. Cell Genet.* 51: 977, 1989.
- [68829] 7159.Cooper, G. J.; Willis, A. C.; Clark, A.; Turner, R. C.; Sim, R. B.; Reid, K. B.: Purification and characterization of a peptide from amyloid-rich pancreases of type 2 diabetic patients. *Proc. Nat. Acad. Sci.* 84: 8628–8632, 1987.
- [68830] 7160.Cooper, G. J. S.; Leighton, B.; Dimitriadis, G. D.; Parry-Billings, M.; Kowalchuk, J. M.; Howland, K.; Rothbard, J. B.; Willis, A. C.; Reid, K. B. M.: Amylin found in amyloid deposits in human type 2 diabetes mellitus may be a hormone that regulates glycogen metabolism in skeletal muscle. *Proc. Nat. Acad. Sci.* 85: 7763–7766, 1988.

- [68831] 7161.Fan, Y.-S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.M.; Sanke, T.; Steiner, D. F.; Shows, T. B.; Bell, G. I.: Localization of the human islet amyloid polypeptide gene (IAP) to chromosome 12p12.3.(Abstract) Cytogenet. Cell Genet. 51: 997, 1989.
- [68832] 7162.Harvey, A. M.: Medical students on the march: Brown, MacCallum and Opie. Johns Hopkins Med. J. 134: 330–345, 1974.
- [68833] 7163.Hoovers, J. M. N.; Redeker, E.; Speleman, F.; Hoppener, J. W. M.; Bhola, S.; Blik, J.; van Roy, N.; Leschot, N. J.; Westerveld, A.; Mannens, M.: High-resolution chromosomal localization of the human calcitonin/CGRP/IAPP gene family members. Genomics 15: 525–529, 1993.
- [68834] 7164.Hoppener, J. W. M.; Oosterwijk, C.; van Hulst, K. L.; Verbeek, J. S.; Capel, P. J. A.; de Koning, E. J. P.; Clark, A.; Jansz, H.S.; Lips, C. J. M.: Molecular physiology of the islet amyloid polypeptide(IAPP)/amylin gene in man, rat, and transgenic mice. J. Cell. Biochem. 55S:39–53, 1994.
- [68835] 7165.Jaikaran, E. T. A. S.; Clark, A.: Islet amyloid and type 2 diabetes: from molecular misfolding to islet pathophysiology. Biochim. Biophys. Acta 1537: 179–203, 2001.
- [68836] 7166.Janson, J.; Soeller, W. C.; Roche, P. C.; Nelson, R. T.; Torchia, A. J.; Kreutter, D. K.; Butler, P. C.: Spontaneous di-

abetes mellitus in transgenic mice expressing human islet amyloid polypeptide. *Proc. Nat. Acad. Sci.* 93: 7283–7288, 1996.

- [68837] 7167. Johnson, K. H.; O'Brien, T. D.; Betsholtz, C.; Westermark, P.: Islet amyloid, islet-amyloid polypeptide, and diabetes mellitus. *New Eng. J. Med.* 321: 513–518, 1989.
- [68838] 7168. Lorenzo, A.; Razzaboni, B.; Weir, G. C.; Yankner, B. A.: Pancreatic islet cell toxicity of amylin associated with type-2 diabetes mellitus. *Nature* 368: 756–760, 1994.
- [68839] 7169. Mosselman, S.; Hoppener, J. W. M.; Lips, C. J. M.; Jansz, H. S.: The complete islet amyloid polypeptide precursor is encoded by two exons. *FEBS Lett.* 247: 154–158, 1989.
- [68840] 7170. Mosselman, S.; Hoppener, J. W. M.; Zandberg, J.; van Mansfeld, A. D. M.; Geurts van Kessel, A. H. M.; Lips, C. J. M.; Jansz, H. S.: Islet amyloid polypeptide: identification and chromosomal localization of the human gene. *FEBS Lett.* 239: 227–232, 1988.
- [68841] 7171. Nishi, M.; Sanke, T.; Seino, S.; Eddy, R. L.; Fan, Y.-S.; Byers, M. G.; Shows, T. B.; Bell, G. I.; Steiner, D. F.: Human islet amyloid polypeptide gene: complete nucleotide sequence, chromosomal localization, and evolutionary history. *Molec. Endocr.* 3: 1775–1781, 1989.

- [68842] 7172.Opie, E. L.: On the relation of the chronic interstitial pancreatitisto the islands of Langerhans and to diabetes mellitus. J. Exp. Med. 5:397–428, 1900.
- [68843] 7173.Bergsagel, P. L.; Chesi, M.; Nardini, E.; Brents, L. A.; Kirby,S. L.; Kuehl, W. M.: Promiscuous translocations into immunoglobulinheavy chain switch regions in multiple myeloma. Proc. Nat. Acad.Sci. 93: 13931–13936, 1996.
- [68844] 7174.Chen, L.; Li, C.; Qiao, W.; Xu, X.; Deng, C.: A ser365–cys mutationof fibroblast growth factor receptor 3 in mouse downregulates Ihh/PTHrPsignals and causes severe achondroplasia. Hum. Molec. Genet. 10:457–465, 2001.
- [68845] 7175.Clausen, J. O.; Hansen, T.; Bjorbaek, C.; Echwald, S. M.; Urhammer,S. A.; Rasmussen, S.; Andersen, C. B.; Hansen, L.; Almind, K.; Winther,K.; Haraldsdottir, J.; Borch–Johnsen, K.; Pedersen, O.: Insulin resistance:interactions between obesity and a common variant of insulin receptorsubstrate–1. Lancet 346: 397–402, 1995.
- [68846] 7176.Esposito, D. L.; Mammarella, S.; Ranieri, A.; Loggia, F. D.; Capani,F.; Consoli, A.; Mariani–Costantini, R.; Caramia, F. G.; Cama, A.;Battista, P.: Deletion of gly723 in the insulin receptor substrate–1of a patient with nonin–

sulin-dependent diabetes mellitus. Hum. Mutat.

7:364–366, 1996.

[68847] 7177.Hribal, M. L.; Federici, M.; Porzio, O.; Lauro, D.; Borboni, P.; Accili, D.; Lauro, R.; Sesti, G.: The gly-to-arg(972) amino acid polymorphism in insulin receptor substrate-1 affects glucose metabolism in skeletal muscle cells. J. Clin. Endocr. Metab. 85: 2004–2013, 2000.

[68848] 7178.Kido, Y.; Burks, D. J.; Withers, D.; Bruning, J. C.; Kahn, C.R.; White, M. F.; Accili, D.: Tissue-specific insulin resistance in mice with mutations in the insulin receptor, IRS-1, and IRS-2. J. Clin. Invest. 105: 199–205, 2000.

[68849] 7179.Kulkarni, R. N.; Winnay, J. N.; Daniels, M.; Bruning, J. C.; Flier, S. N.; Hanahan, D.; Kahn, C. R.: Altered function of insulin receptor substrate-1-deficient mouse islets and cultured beta-cell lines. J. Clin. Invest. 104: R69–R75, 1999.

[68850] 7180.Laakso, M.; Malkki, M.; Kekalainen, P.; Kuusisto, J.; Deeb, S.S.: Insulin receptor substrate-1 variants in non-insulin-dependent diabetes. J. Clin. Invest. 94: 1141–1146, 1994.

[68851] 7181.Myers, M. G., Jr.; Sun, X. J.; White, M. F.: The IRS-1 signaling system. Trends Biochem. Sci. 19: 289–293, 1994.

- [68852] 7182. Nishiyama, M.; Inazawa, J.; Ariyama, T.; Nakamura, Y.; Matsufuji, S.; Furusaka, A.; Tanaka, T.; Hayashi, S.; Wands, J. R.: The human insulin receptor substrate-1 gene (IRS1) is localized on 2q36. *Genomics* 20:139–141, 1994.
- [68853] 7183. Nishiyama, M.; Wands, J. R.: Cloning and increased expression of an insulin receptor substrate-1-like gene in human hepatocellular carcinoma. *Biochem. Biophys. Res. Commun.* 183: 280–285, 1992.
- [68854] 7184. Ogata, N.; Chikazu, D.; Kubota, N.; Terauchi, Y.; Tobe, K.; Azuma, Y.; Ohta, T.; Kadowaki, T.; Nakamura, K.; Kawaguchi, H.: Insulin receptor substrate-1 in osteoblast is indispensable for maintaining bone turnover. *J. Clin. Invest.* 105: 935–943, 2000.
- [68855] 7185. Ogihara, T.; Isobe, T.; Ichimura, T.; Taoka, M.; Funaki, M.; Sakoda, H.; Onishi, Y.; Inukai, K.; Anai, M.; Fukushima, Y.; Kikuchi, M.; Yazaki, Y.; Oka, Y.; Asano, T.: 14-3-3 protein binds to insulin receptor substrate-1, one of the binding sites of which is in the phosphotyrosine binding domain. *J. Biol. Chem.* 272: 25267–25274, 1997.
- [68856] 7186. Porzio, O.; Federici, M.; Hribal, M. L.; Lauro, D.; Accili, D.; Lauro, R.; Borboni, P.; Sesti, G.: The gly972-to-arginine amino acid polymorphism in IRS-1 impairs insulin secre-

tion in pancreatic beta cells. *J. Clin. Invest.* 104: 357–364, 1999.

- [68857] 7187. Stoffel, M.; Espinosa, R., III; Keller, S. R.; Lienhard, G. E.; Le Beau, M. M.; Bell, G. I.: Human insulin receptor substrate–1 gene (IRS1): chromosomal localization to 2q35–q36.1 and identification of a simple tandem repeat DNA polymorphism. *Diabetologia* 36: 335–337, 1993.
- [68858] 7188. Sun, X. J.; Rothenberg, P.; Kahn, C. R.; Backer, J. M.; Araki, E.; Wilden, P. A.; Cahill, D. A.; Goldstein, B. J.; White, M. F.: Structure of the insulin receptor substrate IRS–1 defines a unique signal transduction protein. *Nature* 352: 73–77, 1991.
- [68859] 7189. Diaz, M. O.; Bohlander, S.: Nomenclature of the human interferon genes. *J. Interferon Res.* 13: 443–444, 1993.
- [68860] 7190. Henco, K.; Brosius, J.; Fujisawa, A.; Fujisawa, J.-I.; Haynes, J. R.; Hochstadt, J.; Kovacic, T.; Pasek, M.; Schamblock, A.; Schmid, J.; Todokoro, K.; Walchli, M.; Nagata, S.; Weissmann, C.: Structural relationship of human interferon alpha genes and pseudogenes. *J. Molec. Biol.* 185: 227–260, 1985.
- [68861] 7191. Olopade, O. I.; Bohlander, S. K.; Pomykala, H.; Maltepe, E.; VanMelle, E.; Le Beau, M. M.; Diaz, M. O.: Map–

ping of the shortest region of overlap of deletions of the short arm of chromosome 9 associated with human neoplasia. *Genomics* 14: 437–443, 1992.

- [68862] 7192. Hogervorst, F.; Kuikman, I.; Geurts van Kessel, A.; Sonnenberg, A.: Molecular cloning of the human alpha-6 integrin subunit: alternative splicing of alpha-6 mRNA and chromosomal localization of the alpha-6 and beta-4 genes. *Europ. J. Biochem.* 199: 425–433, 1991.
- [68863] 7193. Chavanas, S.; Gache, Y.; Vailly, J.; Kanitakis, J.; Pulkkinen, L.; Uitto, J.; Ortonne, J.-P.; Meneguzzi, G.: Splicing modulation of integrin beta-4 pre-mRNA carrying a branch point mutation underlies epidermolysis bullosa with pyloric atresia undergoing spontaneous amelioration with ageing. *Hum. Molec. Genet.* 8: 2097–2105, 1999.
- [68864] 7194. Inoue, M.; Tamai, K.; Shimizu, H.; Owaribe, K.; Nakama, T.; Hashimoto, T.; McGrath, J. A.: A homozygous missense mutation in the cytoplasmic tail of beta-4 integrin, G931D, that disrupts hemidesmosome assembly and underlies non-Herlitz junctional epidermolysis bullosa without pyloric atresia? *J. Invest. Derm.* 114: 1061–1064, 2000.
- [68865] 7195. Mellerio, J. E.; Pulkkinen, L.; McMillan, J. R.; Lake, B. D.; Horn, H. M.; Tidman, M. J.; Harper, J. I.; McGrath, J. A.;



Uitto, J.; Eady, R. A. J.: Pyloric atresia–junctional epidermolysis bullosa syndrome: mutations in the integrin beta–4 gene (ITGB4) in two unrelated patients with mild disease. *Brit. J. Derm.* 139: 862–871, 1998.

[68866] 7196. Carayannopoulos, M. O.; Chi, M. M.–Y.; Cui, Y.; Pingsterhaus, J. M.; McKnight, R. A.; Mueckler, M.; Devaskar, S. U.; Moley, K. H.: GLUT8 is a glucose transporter responsible for insulin–stimulated glucose uptake in the blastocyst. *Proc. Nat. Acad. Sci.* 97: 7313–7318, 2000.

[68867] 7197. Doege, H.; Schurmann, A.; Bahrenberg, G.; Brauers, A.; Joost, H. G.: GLUT8, a novel member of the sugar transport facilitator family with glucose transport activity. *J. Biol. Chem.* 275: 16275–16280, 2000.

[68868] 7198. Ibberson, M.; Uldry, M.; Thorens, B.: GLUTX1, a novel mammalian glucose transporter expressed in the central nervous system and insulin–sensitive tissues. *J. Biol. Chem.* 275: 4607–4612, 2000.

[68869] 7199. Kool, M.; de Haas, M.; Scheffer, G. L.; Scheper, R. J.; van Eijk, M. J.; Juijn, J. A.; Baas, F.; Borst, P.: Analysis of expression of cMOAT (MRP2), MRP3, MRP4, and MRP5, homologues of the multidrug resistance–associated protein gene (MRP1), in human cancer cell lines. *Cancer Res.* 57: 3537–3547, 1997.

- [68870] 7200.Lee, K.; Belinsky, M. G.; Bell, D. W.; Testa, J. R.; Kruh, G. D.: Isolation of MOAT-B, a widely expressed multidrug resistance-associated protein/canalicular multispecific organic anion transporter-related transporter. *Cancer Res.* 58: 2741–2747, 1998.
- [68871] 7201.Schuetz, J. D.; Connelly, M. C.; Sun, D.; Paibir, S. G.; Flynn, P. M.; Srinivas, R. V.; Kumar, A.; Fridland, A.: MRP4: a previously unidentified factor in resistance to nucleoside-based antiviral drugs. *Nature Med.* 5: 1048–1051, 1999.
- [68872] 7202.Mahlknecht, U.; Schnittger, S.; Ottmann, O. G.; Schoch, C.; Mosebach, M.; Hiddemann, W.; Hoelzer, D.: Chromosomal organization and localization of the human histone deacetylase 5 gene (HDAC5). *Biochim. Biophys. Acta* 1493: 342–348, 2000.
- [68873] 7203.McKinsey, T. A.; Zhang, C.-L.; Lu, J.; Olson, E. N.: Signal-dependent nuclear export of a histone deacetylase regulates muscle differentiation. *Nature* 408:106–111, 2000.
- [68874] 7204.Bruce, H. A.; Margolis, R. L.: FOXP2: novel exons, splice variants, and CAG repeat length stability. *Hum. Genet.* 111: 136–144, 2002.
- [68875] 7205.Enard, W.; Przeworski, M.; Fisher, S. E.; Lai, C. S. L.;

Wiebe,V.; Kitano, T.; Monaco, A. P.; Paabo, S.: Molecular evolution ofFOXP2, a gene involved in speech and language. Nature 418: 869–872,2002.

- [68876] 7206.Chesney, J.; Mitchell, R.; Benigni, F.; Bacher, M.; Spiegel, L.;Al–Abed, Y.; Han, J. H.; Metz, C.; Bucala, R.: An inducible geneproduct for 6–phosphofructo–2–kinase with an AU–rich instability element:role in tumor cell glycolysis and the Warburg effect. Proc. Nat.Acad. Sci. 96: 3047–3052, 1999.
- [68877] 7207.Manzano, A.; Rosa, J. L.; Ventura, F.; Perez, J. X.; Nadal, M.;Estivill, X.; Ambrosio, S.; Gil, J.; Bartrons, R.: Molecular cloning,expression, and chromosomal localization of a ubiquitously expressedhuman 6–phosphofructo–2–kinase/fructose–2,6–bisphosphatase gene (PFKFB3). Cytogenet.Cell Genet. 83: 214–217, 1998.
- [68878] 7208.Nicholl, J.; Hamilton, J. A.; Sutherland, G. R.; Sutherland, R.L.; Watts, C. K. W.: The third human isoform of 6–phosphofructo–2–kinase/fructose–2,6–bisphosphatase( PFKFB3) map position 10p14–p15. Chromosome Res. 5: 150 only, 1997.
- [68879] 7209.Sakai, A.; Kato, M.; Fukasawa, M.; Ishiguro, M.; Furuya, E.; Sakakibara,R.: Cloning of cDNA encoding for a novel isozyme of fructose

6-phosphate,2-kinase/fructose2,6-bisphosphatase from human placenta. J. Biochem. 119: 506-511,1996.

[68880] 7210.Newbury, D. F.; Bonora, E.; Lamb, J. A.; Fisher, S. E.; Lai, C.S. L.; Baird, G.; Jannoun, L.; Slonims, V.; Stott, C. M.; Merricks,M. J.; Bolton, P. F.; Bailey, A. J.; Monaco, A. P.; InternationalMolecular Genetic Study of Autism Consortium: FOXP2 is not a majorsusceptibility gene for autism or specific language impairment. Am.J. Hum. Genet. 70: 1318-1327, 2002.

[68881] 7211.Sakakibara, R.; Kato, M.; Okamura, N.; Nakagawa, T.; Komada, Y.;Tominaga, N.; Shimojo, M.; Fukasawa, M.: Characterization of a humanplacental fructose-6-phosphate,2-kinase/fructose-2,6-bisphosphatase. J.Biochem. 122: 122-128, 1997.

[68882] 7212.Manzano, A.; Perez, J. X.; Nadal, M.; Estivill, X.; Lange, A.;Bartrons, R.: Cloning, expression and chromosomal localization ofa human testis 6-phosphofructo-2-kinase/fructose-2,6-bisphosphatase gene. Gene 229: 83-89, 1999.

[68883] 7213.Larsen, M.; Ressler, S. J.; Gerdes, M. J.; Lu, B.; Byron, M.; Lawrence,J. B.; Rowley, D. R.: The WFDC1 gene encoding ps20 localizes to 16q24,a region of LOH in multiple cancers. Mammalian Genome 11: 767-773,2000.

- [68884] 7214.Larsen, M.; Ressler, S. J.; Lu, B.; Gerdes, M. J.; McBride, L.;Dang, T. D.; Rowley, D. R.: Molecular cloning and expression of ps20growth inhibitor: a novel WAP-type 'four-disulfide core' domain proteinexpressed in smooth muscle. *J. Biol. Chem.* 273: 4574–4584, 1998.
- [68885] 7215.Rowley, D. R.: Characterization of a fetal urogenital sinus mesenchymalcell line U4F: secretion of a negative growth factor regulatory activity. *InVitro Cell. Dev. Biol.* 28A: 29–38, 1992.
- [68886] 7216.Rowley, D. R.; Dang, T. D.; Larsen, M.; Gerdes, M. J.; McBride,L.; Lu, B.: Purification of a novel protein (ps20) from urogenitalsinus mesenchymal cells with growth inhibitory properties in vitro. *J.Biol. Chem.* 270: 22058–22065, 1995.
- [68887] 7217.Chen, G.; Cizeau, J.; Vande Velde, C.; Park, J. H.; Bozek, G.;Bolton, J.; Shi, L.; Dubik, D.; Greenberg, A.: Nix and nip3 forma subfamily of pro-apoptotic mitochondrial proteins. *J. Biol. Chem.* 274:7–10, 1999.
- [68888] 7218.Matsushima, M.; Fujiwara, T.; Takahashi, E.; Minaguchi, T.; Eguchi,Y.; Tsujimoto, Y.; Suzumori, K.; Nakamura, Y.: Isolation, mapping,and functional analysis of a novel human cDNA (BNIP3L) encoding aprotein homologous to human NIP3. *Genes Chromosomes Cancer* 21:

230–235,1998.

- [68889] 7219.Yasuda, M.; Han, J.; Dionne, C. A.; Boyd, J. M.; Chinadurai, G.: BNIP3- $\alpha$ : a human homolog of mitochondrial proapoptotic proteinBNIP3. *Cancer Res.* 59: 533–537, 1999.
- [68890] 7220.Hildebrand, J. D.; Taylor, J. M.; Parsons, T. J.: An SH3 domain-containingGTPase-activating protein for Rho and Cdc42 associates with focaladhesion kinase. *Molec. Cell. Biol.* 16: 3169–3178, 1996.
- [68891] 7221.Smith, D. E.; Renshaw, B. R.; Ketchem, R. R.; Kubin, M.; Garka,K. E.; Sims, J. E.: Four new members expand the interleukin-1 superfamily. *J.Biol. Chem.* 275: 1169–1175, 2000.
- [68892] 7222.Kumar, S.; McDonnell, P. C.; Lehr, R.; Tierney, L.; Tzimas, M.N.; Griswold, D. E.; Capper, E. A.; Tal-Singer, R.; Wells, G. I.;Doyle, M. L.; Young, P. R.: Identification and initial characterizationof four novel members of the interleukin-1 family. *J. Biol. Chem.* 275:10308–10314, 2000.
- [68893] 7223.Borkhardt, A.; Bojesen, S.; Haas, O. A.; Fuchs, U.; Bartelheimer,D.; Loncarevic, I. F.; Bohle, R. M.; Harbott, J.; Repp, R.; Jaeger,U.; Viehmann, S.; Henn, T.; Korth, P.; Scharr, D.; Lampert, F.: Thehuman GRAF gene is fused to MLL in a unique t(5;11)(q31;q23) and bothalleles are dis-

rupted in three cases of myelodysplastic syndrome/ acutemyeloid leukemia with a deletion 5q. Proc. Nat. Acad. Sci. 97: 9168–9173,2000.

- [68894] 7224.Krystal, G. W.; Armstrong, B. C.; Battey, J. F.: N-myc mRNA formsan RNA–RNA duplex with endogenous anti-sense transcripts. Molec. Cell.Biol. 10: 4180–4191, 1990.
- [68895] 7225.Torrents, D.; Estevez, R.; Pineda, M.; Fernandez, E.; Lloberas,J.; Shi, Y.–B.; Zorzano, A.; Palacin, M.: Identification and characterizationof a membrane protein (y(+)-L amino acid transporter–1) that associateswith 4F2hc to encode the amino acid transport activity y(+)-L: a candidategene for lysinuric protein intolerance. J. Biol. Chem. 273: 32437–32445,1998.
- [68896] 7226.Cohen–Salmon, M.; Frenz, D.; Liu, W.; Verpy, E.; Voegeling, S.;Petit, C.: Fdp, a new fibrocyte–derived protein related to MIA/CD–RAP,has an in vitro effect on the early differentiation of the inner ear mesenchyme. J. Biol. Chem. 275: 40036–40041, 2000.
- [68897] 7227.Rendtorff, N. D.; Frodin, M.; Attie–Bitach, T.; Veke-mans, M.; Tommerup,N.: Identification and characterization of an inner ear–expressedhuman melanoma inhibitory activity (MIA)–like gene (MIAL) with a frequent–polymorphism that abolishes translation. Genomics 71:

40–52, 2001.

- [68898] 7228. Robertson, N. G.; Heller, S.; Lin, J. S.; Resendes, B. L.; Weremowicz, S.; Denis, C. S.; Bell, A. M.; Hudspeth, A. J.; Morton, C. C.: A novel conserved cochlear gene, OTOR: identification, expression analysis, and chromosomal mapping. *Genomics* 66: 242–248, 2000.
- [68899] 7229. Haft, C. R.; de la Luz Sierra, M.; Bafford, R.; Lesniak, M. A.; Barr, V. A.; Taylor, S. I.: Human orthologs of yeast vacuolar protein sorting proteins Vps26, 29, and 35: assembly into multimeric complexes. *Molec. Biol. Cell* 11: 4105–4116, 2000.
- [68900] 7230. Busfield, S. J.; Comrack, C. A.; Yu, G.; Chickering, T. W.; Smutko, J. S.; Zhou, H.; Leiby, K. R.; Holmgren, L. M.; Gearing, D. P.; Pan, Y.: Identification and gene organization of three novel members of the IL-1 family on human chromosome 2. *Genomics* 66: 213–216, 2000.
- [68901] 7231. Mulero, J. J.; Pace, A. M.; Nelken, S. T.; Loeb, D. B.; Correa, T. R.; Drmanac, R.; Ford, J. E.: IL1HY1: a novel interleukin-1 receptor antagonist gene. *Biochem. Biophys. Res. Commun.* 263: 702–706, 1999.
- [68902] 7232. Pratt, W. S.; Crawley, S.; Hicks, J.; Ho, J.; Nash, M.; Kim, Y. S.; Gum, J. R.; Swallow, D. M.: Multiple transcripts of MUC3: evidence for two genes, MUC3A and MUC3B.



Biochem. Biophys. Res. Commun. 275:916–923, 2000.

- [68903] 7233.Ishida, N.; Ito, M.; Yoshioka, S.; Sun-Wada, G.-H.; Kawakita, M.: Functional expression of human Golgi CMP-sialic acid transporter in the Golgi complex of a transporter-deficient Chinese hamster ovary cell mutant. J. Biochem. 124: 171–178, 1998.
- [68904] 7234.Kasof, G. M.; Gomes, B. C.: Livin, a novel inhibitor of apoptosis protein family member. J. Biol. Chem. 276: 3238–3246, 2001.
- [68905] 7235.Lin, J.-H.; Deng, G.; Huang, Q.; Morser, J.: KIAP, a novel member of the inhibitor of apoptosis protein family. Biochem. Biophys. Res. Commun. 279: 820–831, 2000.
- [68906] 7236.Vucic, D.; Stennicke, H. R.; Pisabarro, M. T.; Salvesen, G. S.; Dixit, V. M.: ML-IAP, a novel inhibitor of apoptosis that is preferentially expressed in human melanomas. Curr. Biol. 10: 1359–1366, 2000.
- [68907] 7237.Kobayashi, S.; Akiyama, T.; Nata, K.; Abe, M.; Tajima, M.; Shervani, N. J.; Unno, M.; Matsuno, S.; Sasaki, H.; Takasawa, S.; Okamoto, H.: Identification of a receptor for Reg (regenerating gene) protein, a pancreatic beta-cell regeneration factor. J. Biol. Chem. 275:10723–10726, 2000.
- [68908] 7238.Van Hul, W.; Wuyts, W.; Hendrickx, J.; Speleman, F.;

Wauters, J.; De Boulle, K.; Van Roy, N.; Bossuyt, P.; Willems, P. J.: Identification of a third EXT-like gene (EXTL3) belonging to the EXT gene family. *Genomics* 47:230–237, 1998.

[68909] 7239. Dowler, S.; Currie, R. A.; Downes, C. P.; Alessi, D. R.: DAPP1: a dual adaptor for phosphotyrosine and 3-phosphoinositides. *Biochem. J.* 342: 7–12, 1999.

[68910] 7240. Marshall, A. J.; Niino, H.; Lerner, C. G.; Yun, T. J.; Thomas, S.; Disteche, C. M.; Clark, E. A.: A novel B lymphocyte-associated adaptor protein, Bam32, regulates antigen receptor signaling downstream of phosphatidylinositol 3-kinase. *J. Exp. Med.* 191: 1319–1331, 2000.

[68911] 7241. Di Fiore, P. P.; Pelicci, P. G.; Sorkin, A.: EH: a novel protein-protein interaction domain potentially involved in intracellular sorting. *Trends Biochem. Sci.* 22: 411–413, 1997.

[68912] 7242. Haider, N. B.; Searby, C.; Galperin, E.; Mintz, L.; Horowitz, M.; Stone, E. M.; Sheffield, V. C.: Evaluation and molecular characterization of EHD1, a candidate gene for Bardet-Biedl syndrome 1 (BBS1). *Gene* 240:227–232, 1999.

[68913] 7243. Mintz, L.; Galperin, E.; Pasmanik-Chor, M.; Tulzinsky, S.; Bromberg, Y.; Kozak, C. A.; Joyner, A.; Fein, A.;

Horowitz, M.: EHD1--an EH-domain-containing protein with a specific expression pattern. *Genomics* 59: 66-76, 1999.

[68914] 7244. Bouju, S.; Pietu, G.; Le Cunff, M.; Cros, N.; Malzac, P.; Pellissier, J.-F.; Pons, F.; Leger, J.-J.; Auffray, C.; Dechesne, C. A.: Exclusion of muscle specific actinin-associated LIM protein (ALP) gene from 4q35 facioscapulohumeral muscular dystrophy (FSHD) candidate genes. *Neuromusc. Disord.* 9: 3-10, 1999.

[68915] 7245. Black, B. E.; Levesque, L.; Holaska, J. M.; Wood, T. C.; Paschal, B. M.: Identification of an NTF2-related factor that binds Ran-GTP and regulates nuclear protein export. *Molec. Cell. Biol.* 19: 8616-8624, 1999.

[68916] 7246. Katahira, J.; Straber, K.; Podtelejnikov, A.; Mann, M.; Jung, J. U.; Hurt, E.: The Mex67p-mediated nuclear mRNA export pathway is conserved from yeast to human. *EMBO J.* 18: 2593-2609, 1999.

[68917] 7247. Ossareh-Nazari, B.; Maison, C.; Black, B. E.; Levesque, L.; Paschal, B. M.; Dargemont, C.: RanGTP-binding protein NXT1 facilitates nuclear export of different classes of RNA in vitro. *Molec. Cell. Biol.* 20: 4562-4571, 2000.

[68918] 7248. Scott, A. F.: Personal Communication. Baltimore, Md.

4/3/2001.

- [68919] 7249.Pati, D.; Meistrich, M. L.; Plon, S. E.: Human Cdc34 and Rad6Bubiquitin-conjugating enzymes target repressors of cyclic AMP-inducedtranscription for proteolysis. *Molec. Cell. Biol.* 19: 5001-5013,1999.
- [68920] 7250.Courjal, F.; Louason, G.; Speiser, P.; Katsaros, D.; Zeillinger,R.; Theillet, C.: Cyclin gene amplification and overexpression inbreast and ovarian cancers: evidence for the selection of cyclin D1in breast and cyclin E in ovarian tumors. *Int. J. Cancer* 69: 247-253,1996.
- [68921] 7251.Strohmaier, H.; Spruck, C. H.; Kaiser, P.; Won, K.-A.; Sangfelt,O.; Reed, S. I.: Human F-box protein hCdc4 targets cyclin E for proteolysisand is mutated in a breast cancer cell line. *Nature* 413: 316-322,2001.
- [68922] 7252.Matsuda, S.; Iriyama, C.; Yokozaki, S.; Ichigotani, Y.; Shirafuji,N.; Yamaki, K.; Hayakawa, T.; Hamaguchi, M.: Cloning and sequencingof a novel human gene that encodes a putative target protein of Nesh-SH3. *J.Hum. Genet.* 46: 483-486, 2001.
- [68923] 7253.Jager, D.; Stockert, E.; Jager, E.; Gure, A. O.; Scanlan, M. J.;Knuth, A.; Old, L. J.; Chen, Y.-T.: Serological cloning of a melanocyterab guanosine 5-prime-triphosphate-binding protein and a chromo-

somecondensation protein from a melanoma complementary DNA library. *CancerRes.* 60: 3584–3591, 2000.

[68924] 7254.Loftus, S. K.; Larson, D. M.; Baxter, L. L.; Antonellis, A.; Chen,Y.; Wu, X.; Jiang, Y.; Bittner, M.; Hammer, J. A., III; Pavan, W.J.: Mutation of melanosome protein RAB38 in chocolate mice. *Proc.Nat. Acad. Sci.* 99: 4471–4476, 2002.

[68925] 7255.Matsuyoshi, N.; Tanaka, T.; Toda, K.; Imamura, S.: Identificationof novel cadherins expressed in human melanoma cells. *J. Invest.Derm.* 108: 908–913, 1997.

[68926] 7256.Chen, K.–S.; DeLuca, H. F.: Isolation and characterization ofa novel cDNA from HL–60 cells treated with 1,25–dihydroxyvitamin D–3. *Biochim.Biophys. Acta* 1219: 26–32, 1994.

[68927] 7257.Ludwig, D. L.; Kotanides, H.; Le, T.; Chavkin, D.; Bohlen, P.;Witte, L.: Cloning, genetic characterization, and chromosomal mappingof the mouse VDUP1 gene. *Gene* 269: 103–112, 2001.

[68928] 7258.Yun, C. H. C.; Oh, S.; Zizak, M.; Steplock, D.; Tsao, S.; Tse,C.–M.; Weinman, E. J.; Donowitz, M.: cAMP–mediated inhibition ofthe epithelial brush border Na(+)/H(+) exchanger, NHE3, requires anassociated regulatory protein. *Proc. Nat. Acad. Sci.* 94: 3010–3015,1997.

- [68929] 7259.Li, Y.; Chin, L.-S.; Weigel, C.; Li, L.: Spring, a novel RINGfinger protein that regulates synaptic vesicle exocytosis. *J. Biol.Chem.* 276: 40824–40833, 2001.
- [68930] 7260.Reymond, A.; Meroni, G.; Fantozzi, A.; Merla, G.; Cairo, S.; Luzi,L.; Riganelli, D.; Zanaria, E.; Messali, S.; Cainarca, S.; Guffanti,A.; Minucci, S.; Pelicci, P. G.; Bal-labio, A.: The tripartite motiffamily identifies cell compartments. *EMBO J.* 20: 2140–2151, 2001.
- [68931] 7261.Scardigli, R.; Schuurmans, C.; Gradwohl, G.; Guillemot, F.: Crossregulationbetween neurogenin2 and pathways specifying neuronal identity in thespinal cord. *Neuron* 31: 203–217, 2001.
- [68932] 7262.Boles, K. S.; Mathew, P. A.: Molecular cloning of CS1, a novelhuman natural killer cell receptor belonging to the CD2 subset ofthe immunoglobulin superfamily. *Immunogenetics* 52: 302–307, 2001.
- [68933] 7263.Bouchon, A.; Cella, M.; Grierson, H. L.; Cohen, J. I.; Colonna,M.: Cutting edge: activation of NK cell-mediated cytotoxicity bya SAP-independent receptor of the CD2 family. *J. Immun.* 167: 5517–5521,2001.
- [68934] 7264.Habas, R.; Kato, Y.; He, X.: Wnt/Frizzled activation of Rho regulatesvertebrate gastrulation and requires a novel Formin homology proteinDaam1. *Cell* 107: 843–854,

2001.

- [68935] 7265. Peters, C. S.; Liang, X.; Li, S.; Kannan, S.; Peng, Y.; Taub, R.; Diamond, R. H.: ATF-7, a novel bZIP protein, interacts with the PRL-1 protein-tyrosine phosphatase. *J. Biol. Chem.* 276: 13718–13726, 2001.
- [68936] 7266. White, J. H.; McIlhinney, R. A. J.; Wise, A.; Ciruela, F.; Chan, W.-Y.; Emson, P. C.; Billinton, A.; Marshall, F. H.: The GABA-B receptor interacts directly with the related transcription factors CREB2 and ATFx. *Proc. Nat. Acad. Sci.* 97: 13967–13972, 2000.
- [68937] 7267. Futai, E.; Kubo, T.; Sorimachi, H.; Suzuki, K.; Maeda, T.: Molecular cloning of PalBH, a mammalian homologue of the *Aspergillus* atypical calpain PalB. *Biochim. Biophys. Acta* 1517: 316–319, 2001.
- [68938] 7268. Huang, Y.; Wang, K. K. W.: The calpain family and human disease. *Trends Molec. Med.* 7: 355–362, 2001.
- [68939] 7269. Lee, H.-J.; Sorimachi, H.; Jeong, S.-Y.; Ishiura, S.; Suzuki, K.: Molecular cloning and characterization of a novel tissue-specific calpain predominantly expressed in digestive tract. *Biol. Chem.* 379: 175–183, 1998.
- [68940] 7270. Yoshikawa, Y.; Mukai, H.; Hino, F.; Asada, K.; Kato, I.: Isolation of two novel genes, down-regulated in gastric cancer. *Jpn. J. Cancer Res.* 91: 459–463, 2000.

- [68941] 7271.Burgess, D. L.; Davis, C. F.; Gefrides, L. A.; Noebels, J. L.:Identification of three novel Ca(2+) channel gamma subunit genes revealsmolecular diversification by tandem and chromosome duplication. *GenomeRes.* 9: 1204–1213, 1999.
- [68942] 7272.Burgess, D. L.; Gefrides, L. A.; Foreman, P. J.; Noebels, J. L.: A cluster of three novel Ca(2+) channel gamma subunit genes on chromosome19q13.4: evolution and expression profile of the gamma subunit genefamily. *Genomics* 71: 339–350, 2001.
- [68943] 7273.Chu, P.–J.; Robertson, H. M.; Best, P. M.: Calcium channel gammasubunits provide insights into the evolution of this gene family. *Gene* 280:37–48, 2001.
- [68944] 7274.Chen, X.; Wen, S.; Fukuda, M. N.; Gava, N. R.; Hsu, D.; Akama,T. O.; Yang–Feng, T.; Shen, C. K. J.: Human ITCH is a coregulatorof the hematopoietic transcription factor NF–E2. *Genomics* 73: 238–241,2001.
- [68945] 7275.D'Andrea, A. D.; Serhan, C. N.: Relieving the Itch. *Nature Genet.* 18:97–99, 1998.
- [68946] 7276.Perry, W. L.; Hustad, C. M.; Swing, D. A.; O'Sullivan, T. N.; Jenkins,N. A.; Copeland, N. G.: The itchy locus encodes a novel ubiquitinprotein ligase that is disrupted in a18H mice. *Nature Genet.* 18:143–146, 1998.



- [68947] 7277. Qiu, L.; Joazeiro, C.; Fang, N.; Wang, H.-Y.; Elly, C.; Altman, Y.; Fang, D.; Hunter, T.; Liu, Y.-C.: Recognition and ubiquitination of Notch by Itch, a Hect-type E3 ubiquitin ligase. *J. Biol. Chem.* 275:35734–35737, 2000.
- [68948] 7278. Winberg, G.; Matskova, L.; Chen, F.; Plant, P.; Rotin, D.; Gish, G.; Ingham, R.; Ernberg, I.; Pawson, T.: Latent membrane protein 2A of Epstein–Barr virus binds WW domain E3 protein–ubiquitin ligases that ubiquitinate B-cell tyrosine kinases. *Molec. Cell. Biol.* 20:8526–8535, 2000.
- [68949] 7279. Bradley, K. A.; Mogridge, J.; Mourez, M.; Collier, R. J.; Young, J. A. T.: Identification of the cellular receptor for anthrax toxin. *Nature* 414:160–161, 2001.
- [68950] 7280. Dragon, F.; Pogacic, V.; Filipowicz, W.: In vitro assembly of human H/ACA small nucleolar RNPs reveals unique features of U17 and telomerase RNAs. *Molec. Cell. Biol.* 20: 3037–3048, 2000.
- [68951] 7281. Tollervey, D.; Kiss, T.: Function and synthesis of small nucleolar RNAs. *Curr. Opin. Cell Biol.* 9: 337–342, 1997.
- [68952] 7282. Pogacic, V.; Dragon, F.; Filipowicz, W.: Human H/ACA small nucleolar RNPs and telomerase share evolutionarily conserved proteins NHP2 and NOP10. *Molec. Cell. Biol.* 20: 9028–9040, 2000.

- [68953] 7283.Ichtchenko, K.; Nguyen, T.; Sudhof, T. C.: Structures, alternativesplicing, and neurexin binding of multiple neuroligins. J. Biol.Chem. 271: 2676–2682, 1996.
- [68954] 7284.Hammarsund, M.; Wilson, W.; Corcoran, M.; Merup, M.; Einhorn, S.;Grander, D.; Sangfelt, O.: Identification and characterization oftwo novel human mitochondrial elongation factor genes, hEFG2 and hEFG1,phylogenetically conserved through evolution. Hum. Genet. 109: 542–550,2001.
- [68955] 7285.Ishibashi, K.; Suzuki, M.; Sasaki, S.; Imai, M.: Identificationof a new multigene four–transmembrane family (MS4A) related to CD20,HTm4 and beta subunit of the high–affinity IgE receptor. Gene 264:87–93, 2001.
- [68956] 7286.Liang, Y.; Tedder, T. F.: Identification of a CD20–, Fc–epsilon–RI–beta–relatedgene family: sixteen new MS4A family members expressed in human andmouse. Genomics 72: 119–127, 2001.
- [68957] 7287.Ishii, H.; Vecchione, A.; Murakumo, Y.; Baldassarre, G.; Numata,S.; Trapasso, F.; Alder, H.; Baffa, R.; Croce, C. M.: FEZ1/LZTS1gene at 8p22 suppresses cancer cell growth and regulates mitosis. Proc.Nat. Acad. Sci. 98: 10374–10379, 2001.
- [68958] 7288.Mahon, M. J.; Donowitz, M.; Yun, C. C.; Segre, G. V.:

Na<sup>+</sup>/H<sup>+</sup> exchanger regulatory factor 2 directs parathyroid hormone 1 receptor signalling. *Nature* 417:858–861, 2002.

- [68959] 7289. Poulat, F.; de Santa Barbara, P.; Desclozeaux, M.; Soullier, S.; Moniot, B.; Bonneaud, N.; Boizet, B.; Berta, P. The human testis determining factor SRY binds a nuclear factor containing PDZ protein interaction domains. *J. Biol. Chem.* 272: 7167–7172, 1997.
- [68960] 7290. Yun, C.-H. C.; Lamprecht, G.; Forster, D. V.; Sidor, A.: NHE3 kinase A regulatory protein E3KARP binds the epithelial brush border Na<sup>(+)</sup>/H<sup>(+)</sup> exchanger NHE3 and the cytoskeletal protein ezrin. *J. Biol. Chem.* 273: 25856–25863, 1998.
- [68961] 7291. Hall, R. A.; Ostedgaard, L. S.; Premont, R. T.; Blitzer, J. T.; Rahman, N.; Welsh, M. J.; Lefkowitz, R. J.: A C-terminal motif found in the beta-2-adrenergic receptor, P2Y1 receptor and cystic fibrosis transmembrane conductance regulator determines binding to the Na<sup>(+)</sup>/H<sup>(+)</sup> exchanger regulatory factor family of PDZ proteins. *Proc. Nat. Acad. Sci.* 95: 8496–8501, 1998.
- [68962] 7292. Druck, T.; Podolski, J.; Byrski, T.; Wyrwicz, L.; Zajaczk, S.; Kata, G.; Borowka, A.; Lubinski, J.; Huebner, K.: The DIRC1 gene at chromosome 2q33 spans a familial

RCC-associated t(2;3)(q33;q21) chromosome translocation. J. Hum. Genet. 46: 583–589, 2001.

[68963] 7293. Podolski, J.; Zajaczek, S.; Byrski, T.; Druck, T.; Zimonjic, D.B.; Popescu, N. C.; Lubinski, J.; Huebner, K.: Characterization of a familial RCC-associated t(2;3)(q33;q21) chromosome translocation. J. Hum. Genet. in-press, 2001.

[68964] 7294. Bruick, R. K.; McKnight, S. L.: A conserved family of prolyl-4-hydroxylases that modify HIF. Science 294: 1337–1340, 2001.

[68965] 7295. Epstein, A. C. R.; Gleadle, J. M.; McNeill, L. A.; Hewitson, K.S.; O'Rourke, J.; Mole, D. R.; Mukherji, M.; Metzen, E.; Wilson, M.I.; Dhanda, A.; Tian, Y.-M.; Masson, N.; Hamilton, D. L.; Jaakkola, P.; Barstead, R.; Hodgkin, J.; Maxwell, P. H.; Pugh, C. W.; Schofield, C. J.; Ratcliffe, P. J.: C. elegans EGL-9 and mammalian homologs define a family of dioxygenases that regulate HIF by prolyl hydroxylation. Cell 107:43–54, 2001.

[68966] 7296. Dupuy, D.; Aubert, I.; Duperat, V. G.; Petit, J.; Taine, L.; Stef, M.; Bloch, B.; Arveiler, B.: Mapping, characterization, and expression analysis of the SM-20 human homologue, C1orf12, and identification of a novel related gene, SCAND2. Genomics 69: 348–354, 2000.

- [68967] 7297.Ritter, J. K.; Crawford, J. M.; Owens, I. S.: Cloning of two human liver bilirubin UDP-glucuronosyltransferase cDNAs with expression in COS-1 cells. *J. Biol. Chem.* 266: 1043-1047, 1991.
- [68968] 7298.Rheaume, E.; Leblanc, J. F.; Lachance, Y.; Labrie, F.; Simard, J.: Detection of frequent BglII polymorphism by polymerase chain reaction and TaqI restriction fragment length polymorphism for 3-beta-hydroxysteroid dehydrogenase/delta-5-delta-4 isomerase at the human HSDB3 locus (1p11-p13). *Hum. Genet.* 87: 753-754, 1991.
- [68969] 7299.Reynolds, D. S.; Gurley, D. S.; Austen, K. F.: Cloning and characterization of the novel gene for mast cell carboxypeptidase A. *J. Clin. Invest.* 89:273-282, 1992.
- [68970] 7300.Yamaguchi, N.; Benayahu, P. D.; van der Rest, M.; Nishimura, Y.: The cloning and sequencing of alpha-1(VIII) collagen cDNAs demonstrate that type VIII collagen is a short chain collagen and contains triple-helical and carboxyl-terminal non-triple-helical domains similar to those of type X collagen. *J. Biol. Chem.* 264: 16022-16029, 1989.
- [68971] 7301.Bleul, C. C.; Farzan, M.; Choe, H.; Parolin, C.; Clark-Lewis, I.; Sodroski, J.; Springer, T. A.: The lymphocyte

chemoattractant SDF-1 is a ligand for LESTR/fusin and blocks HIV-1 entry. *Nature* 382:829-833, 1996.

- [68972] 7302. Caruz, A.; Samsom, M.; Alonso, J. M.; Alcami, J.; Baleux, F.; Virelizier, J. L.; Parmentier, M.; Arenzana-Seisdedos, F.: Genomic organization and promoter characterization of human CXCR4 gene. *FEBS Lett.* 426:271-278, 1998.
- [68973] 7303. Clapham, P. R.; Blanc, D.; Weiss, R. A.: Specific cell surface requirements for the infection of CD4-positive cells by human immunodeficiency virus types 1 and 2 and by Simian immunodeficiency virus. *Virology* 181:703-715, 1991.
- [68974] 7304. Dragic, T.; Alizon, M.: Different requirements for membrane fusion mediated by the envelopes of human immunodeficiency virus types 1 and 2. *J. Virol.* 67: 2355-2359, 1993.
- [68975] 7305. Federspiel, B.; Melhado, I. G.; Duncan, A. M. V.; Delaney, A.; Schappert, K.; Clark-Lewis, I.; Jirik, F. R.: Molecular cloning of the cDNA and chromosomal localization of the gene for a putative seven-transmembrane segment (7-TMS) receptor isolated from human spleen. *Genomics* 16:707-712, 1993.
- [68976] 7306. Feng, Y.; Broder, C. C.; Kennedy, P. E.; Berger, E. A.:

HIV-1 entry cofactor: functional cDNA cloning of a seven-transmembrane, G protein-coupled receptor. *Science* 272: 872–876, 1996.

[68977] 7307. Hendrix, C. W.; Flexner, C.; MacFarland, R. T.; Giandomenico, C.; Fuchs, E. J.; Redpath, E.; Bridger, G.; Henson, G. W.: Pharmacokinetics and safety of AMD-3100, a novel antagonist of the CXCR-4 chemokine receptor, in human volunteers. *Antimicrob. Agents Chemother.* 44:1667–1673, 2000.

[68978] 7308. Herzog, H.; Hort, Y. J.; Shine, J.; Selbie, L. A.: Molecular cloning, characterization, and localization of the human homolog to the reported bovine NPY Y3 receptor: lack of NPY binding and activation. *DNA Cell Biol.* 12: 465–471, 1993.

[68979] 7309. Jazin, E. E.; Yoo, H.; Blomqvist, A. G.; Yee, F.; Weng, G.; Walker, M. W.; Salon, J.; Larhammar, D.; Wahlestedt, C.: A proposed bovine neuropeptide Y (NPY) receptor cDNA clone, or its human homologue, confers neither NPY binding sites nor NPY responsiveness on transfected cells. *Regul. Pept.* 47: 247–258, 1993.

[68980] 7310. Liotta, L. A.: An attractive force in metastasis. *Nature* 410:24–25, 2001.

[68981] 7311. Loetscher, M.; Geiser, T.; O'Reilly, T.; Zwahlen, R.;

Baggiolini,M.; Moser, B.: Cloning of a human seven-transmembrane domain receptor,LESTR, that is highly expressed in leukocytes. J. Biol. Chem. 269:232–237, 1994.

[68982] 7312.Lu, M.; Grove, E. A.; Miller, R. J.: Abnormal development ofthe hippocampal dentate gyrus in mice lacking the CXCR4 chemokinereceptor. Proc. Nat. Acad. Sci. 99: 7090–7095, 2002.

[68983] 7313.Ma, Q.; Jones, D.; Borghesani, P. R.; Segal, R. A.; Nagasawa,T.; Kishimoto, T.; Bronson, R. T.; Springer, T. A.: Impaired B-lymphopoiesis,myelopoiesis, and derailed cerebellar neuron migration in CXCR4- andSDF-1-deficient mice. Proc. Nat. Acad. Sci. 95: 9448–9453, 1998.

[68984] 7314.Muller, A.; Homey, B.; Soto, H.; Ge, N.; Catron, D.; Buchanan,M. E.; McClanahan, T.; Murphy, E.; Yuan, W.; Wagner, S. N.; Barrera,J. L.; Mohar, A.; Verastegui, E.; Zlotnik, A.: Involvement of chemokinereceptors in breast cancer metastasis. Nature 410: 50–56, 2001.

[68985] 7315.Nagasawa, T.; Hirota, S.; Tachibana, K.; Takakura, N.; Nishikawa,S.; Kitamura, Y.; Yoshida, N.; Kikutani, H.; Kishimoto, T.: Defectsof B-cell lymphopoiesis and bone-marrow myelopoiesis in mice lackingthe CXC chemokine PBSF/SDF-1. Nature 382: 635–638, 1996.



- [68986] 7316.Oberlin, E.; Amara, A.; Bachelerie, F.; Bessia, C.; Virelizier,J.-L.; Arenzana-Seisdedos, F.; Schwartz, O.; Heard, J.-M.; Clark-Lewis,I.; Legler, D. F.; Loetscher, M.; Baggiolini, M.; Moser, B.: TheCXC chemokine SDF-1 is the ligand for LESTR/fusin and prevents infectionby T-cell-line-adapted HIV-1. *Nature* 382: 833-835, 1996.
- [68987] 7317.Peled, A.; Petit, I.; Kollet, O.; Magid, M.; Ponomaryov, T.; Byk,T.; Nagler, A.; Ben-Hur, H.; Many, A.; Shultz, L.; Lider, O.; Alon,R.; Zipori, D.; Lapidot, T.: Dependence of human stem cell engraftmentand repopulation of NOD/SCID mice on CXCR4. *Science* 283: 845-848,1999.
- [68988] 7318.Rimland, J.; Xin, W.; Sweetnam, P.; Saijoh, K.; Nestler, E. J.;Duman, R. S.: Sequence and expression of a neuropeptide Y receptorcDNA. *Molec. Pharm.* 40: 869-875, 1991.
- [68989] 7319.Tachibana, K.; Hirota, S.; Iizasa, H.; Yoshida, H.; Kawabata,K.; Kataoka, Y.; Kitamura, Y.; Matsushima, K.; Yoshida, N.; Nishikawa,S.; Kishimoto, T.; Nagasawa, T.: The chemokine receptor CXCR4 isessential for vascular-ization of the gastrointestinal tract. *Nature* 393:591-594, 1998.
- [68990] 7320.Wegner, S. A.; Ehrenberg, P. K.; Chang, G.; Dayhoff, D. E.; Sleeker,A. L.; Michael, N. L.: Genomic organization

and functional characterization of the chemokine receptor CXCR4, a major entry co-receptor for human immunodeficiency virus type 1. *J. Biol. Chem.* 273: 4754–4760, 1998.

[68991] 7321. Weiner, D. B.; Huebner, K.; Williams, W. V.; Greene, M. I.: Human genes other than CD4 facilitate HIV-1 infection of murine cells. *Pathobiology* 59:361–371, 1991.

[68992] 7322. Szosteki, C.; Guldner, H. H.; Netter, H. J.; Will, H.: Isolation and characterization of cDNA encoding a human nuclear antigen predominantly recognized by autoantibodies from patients with primary biliary cirrhosis. *J. Immun.* 145: 4338–4347, 1990.

[68993] 7323. Freeman, G. J.; Disteché, C. M.; Gribben, J. G.; Adler, D. A.; Freedman, A. S.; Dougery, J.; Nadler, L. M.: The gene for B7, a costimulatory signal for T-cell activation, maps to chromosomal region 3q13.3–3q21. *Blood* 79:489–494, 1992.

[68994] 7324. Reeves, R. H.; Patch, D.; Sharpe, A. H.; Borriello, F.; Freeman, G. J.; Edelhoff, S.; Disteché, C.: The costimulatory genes Cd80 and Cd86 are linked on mouse chromosome 16 and human chromosome 3. *Mammalian Genome* 8: 581–582, 1997.

[68995] 7325. Selvakumar, A.; Mohanraj, B. K.; Eddy, R. L.; Shows, T. B.; White, P. C.; Dupont, B.: Genomic organization and

chromosomal location of the human gene encoding the B-lymphocyte activation antigen B7. *Immunogenetics* 36:175–181, 1992.

- [68996] 7326. Ha, H.; Kubagawa, H.; Burrows, P. D.: Molecular cloning and expression pattern of a human gene homologous to the murine mb-1 gene. *J. Immun.* 148:1526–1531, 1992.
- [68997] 7327. Hashimoto, S.; Mohrenweiser, H. W.; Gregersen, P. K.; Chiorazzi, N.: Chromosomal localization, genomic structure, and allelic polymorphism of the human CD79a (Ig-alpha/mb-1) gene. *Immunogenetics* 40: 287–295, 1994.
- [68998] 7328. Reth, M.: Antigen receptors on B lymphocytes. *Annu. Rev. Immun.* 10:97–121, 1992.
- [68999] 7329. Liang, Y.; Buckley, T. R.; Tu, L.; Langdon, S. D.; Tedder, T. F.: Structural organization of the human MS4A gene cluster on chromosome 11q12. *Immunogenetics* 53: 357–368, 2001.
- [69000] 7330. Tedder, T. F.; Disteche, C. M.; Louie, E.; Adler, D. A.; Croce, C. M.; Schlossman, S. F.; Saito, H.: The gene that encodes the human CD20 (B1) differentiation antigen is located on chromosome 11 near the t(11;14)(q13;q32) translocation site. *J. Immun.* 142: 2555–2559, 1989.

- [69001] 7331.Tedder, T. F.; Klejman, G.; Schlossman, S. F.; Saito, H.: Structure of the gene encoding the human B lymphocyte differentiation antigen CD20 (B1). *J. Immun.* 142: 2560–2568, 1989.
- [69002] 7332.Tedder, T. F.; Streuli, M.; Schlossman, S. F.; Saito, H.: Isolation and structure of a cDNA encoding the B1 (CD20) cell-surface antigen of human B lymphocytes. *Proc. Nat. Acad. Sci.* 85: 208–212, 1988.
- [69003] 7333.Kaplan, F. S.; Tabas, J. A.; Zasloff, M. A.: Fibrodysplasia ossificans progressiva: a clue from the fly? *Calcif. Tissue Int.* 47: 117–125, 1990.
- [69004] 7334.Hopken, U. E.; Lu, D.; Gerard, N. P.; Gerard, C.: The C5a chemoattractant receptor mediates mucosal defense to infection. *Nature* 383: 86–89, 1996.
- [69005] 7335.Chen, H.; Jawahar, S.; Qian, Y.; Duong, Q.; Chan, G.; Parker, A.; Meyer, J. M.; Moore, K. J.; Chayen, S.; Gross, D. J.; Glaser, B.; Permutt, M. A.; Fricker, L. D.: Missense polymorphism in the human carboxypeptidase E gene alters enzymatic activity. *Hum. Mutat.* 18:120–131, 2001.
- [69006] 7336.Hall, C.; Manser, E.; Spurr, N. K.; Lim, L.: Assignment of the human carboxypeptidase E (CPE) gene to chromosome 4. *Genomics* 15:461–463, 1993.
- [69007] 7337.Manser, E.; Fernandez, D.; Loo, L.; Goh, P. Y.; Mon-

fries, C.; Hall, C.; Lim, L.: Human carboxypeptidase E: isolation and characterization of the cDNA, sequence conservation, expression and processing in vitro. *Biochem. J.* 267: 517–525, 1990.

[69008] 7338. Kas, K.; Schoenmakers, E. F. P. M.; Van de Ven, W. J. M.: Physical map location of the human carboxypeptidase M gene (CPM) distal to D12S375 and proximal to D12S8 at chromosome 12q15. *Genomics* 30:403–405, 1995.

[69009] 7339. Rehli, M.; Krause, S. W.; Kreutz, M.; Andreesen, R.: Carboxypeptidase M is identical to the MAX.1 antigen and its expression is associated with monocyte to macrophage differentiation. *J. Biol. Chem.* 270:15644–15649, 1995.

[69010] 7340. Tan, F.; Chan, S. J.; Steiner, D. F.; Schilling, J. W.; Skidgel, R. A.: Molecular cloning and sequencing of the cDNA for human membrane-bound carboxypeptidase M: comparison with carboxypeptidases A, B, H, and N. *J. Biol. Chem.* 264: 13165–13170, 1989.

[69011] 7341. Gold, P.; Freedman, S. O.: Demonstration of tumor-specific antigens in human colonic carcinomata by immunological tolerance and absorption techniques. *J. Exp. Med.* 121: 439–462, 1965.

[69012] 7342. Kamarck, M. E.; Elting, J. J.; Hart, J. T.; Goebel, S. J.; Rae, P. M. M.; Nothdurft, M. A.; Nedwin, J. J.; Barnett, T. R.:

Carcinoembryonic antigen family: expression in a mouse L-cell transfectant and characterization of a partial cDNA in bacteriophage lambda-gt11. *Proc. Nat. Acad. Sci.* 84: 5350–5354, 1987.

[69013] 7343. Nishi, M.; Inazawa, J.; Inoue, K.; Nakagawa, H.; Taniwaki, M.; Misawa, S.; Oikawa, S.; Nakazato, H.; Abe, T.: Regional chromosomal assignment of carcinoembryonic antigen gene (CEA) to chromosome 19 at band q13.2. *Cancer Genet. Cytogenet.* 54: 77–81, 1991.

[69014] 7344. Oikawa, S.; Nakazato, H.; Kosaki, G.: Primary structure of human carcinoembryonic antigen (CEA) deduced from cDNA sequence. *Biochem. Biophys. Res. Commun.* 142: 511–528, 1987.

[69015] 7345. Thompson, J.; Zimmermann, W.: The carcinoembryonic antigen gene family: structure, expression and evolution. *Tumor Biol.* 9: 63–83, 1988.

[69016] 7346. Thompson, J. A.; Pande, H.; Paxton, R. J.; Shively, L.; Padma, A.; Simmer, R. L.; Todd, C. W.; Riggs, A. D.; Shively, J. E.: Molecular cloning of a gene belonging to the carcinoembryonic antigen gene family and discussion of a domain model. *Proc. Nat. Acad. Sci.* 84: 2965–2969, 1987.

[69017] 7347. Willcocks, T. C.; Craig, I. W.: Characterization of the genomic organization of human carcinoembryonic antigen

(CEA): comparison with other family members and sequence analysis of 5-prime controlling region. Genomics 8: 492–500, 1990.

[69018] 7348. Willcocks, T. C.; Craig, S. P.; Coates, D.; Craig, I. W.: Coding sequences for carcinoembryonic antigen (CEA) assigned to human chromosome 19q13. (Abstract) Cytogenet. Cell Genet. 46: 716 only, 1987.

[69019] 7349. Zimmer, R.; Thomas, P.: Mutations in the carcinoembryonic antigen gene in colorectal cancer patients: implications on liver metastasis. Cancer Res. 61: 2822–2826, 2001.

[69020] 7350. Zimmermann, W.; Ortlieb, B.; Friedrich, R.; von Kleist, S.: Isolation and characterization of cDNA clones encoding the human carcinoembryonic antigen reveal a highly conserved repeating structure. Proc. Nat. Acad. Sci. 84: 2960–2964, 1987.

[69021] 7351. Zimmermann, W.; Weber, B.; Ortlieb, B.; Rudert, F.; Schempp, W.; Fiebig, H.-H.; Shively, J. E.; von Kleist, S.; Thompson, J. A.: Chromosomal localization of the carcinoembryonic antigen gene family and differential expression in various tumors. Cancer Res. 48: 2550–2554, 1988.

[69022] 7352. Walker, E. S.: Familial paroxysmal dystonic choreoathetosis: a neurologic disorder simulating psychiatric ill-

ness. Johns Hopkins Med. J. 148: 108–113, 1981.

- [69023] 7353. Schuback, D.; Kramer, P.; Ozelius, L.; Holmgren, G.; Forsgren, L.; Kyllerman, M.; Wahlstrom, J.; Craft, C. M.; Nygaard, T.; Brin, M.; de Leon, D.; Bressman, S.; Moskowitz, C. B.; Burke, R. E.; Sanner, G.; Drugge, U.; Gusella, J. F.; Fahn, S.; Breakefield, X. O.: Dopamine beta-hydroxylase gene excluded in four subtypes of hereditary dystonia. Hum. Genet. 87: 311–316, 1991.
- [69024] 7354. Arahata, K.; Hayashi, Y. K.; Mizuno, Y.; Yoshida, M.; Ozawa, E.: Dystrophin-associated glycoprotein and dystrophin co-localisation at sarcolemma in Fukuyama congenital muscular dystrophy. (Letter) Lancet 342: 623–624, 1993.
- [69025] 7355. Campanelli, J. T.; Roberds, S. L.; Campbell, K. P.; Scheller, R. H.: A role for dystrophin-associated glycoproteins and utrophin in agrin-induced AChR clustering. Cell 77: 663–674, 1994.
- [69026] 7356. Cao, W.; Henry, M. D.; Borrow, P.; Yamada, H.; Elder, J. H.; Ravkov, E. V.; Nichol, S. T.; Compans, R. W.; Campbell, K. P.; Oldstone, M. B. A.: Identification of alpha-dystroglycan as a receptor for lymphocytic choriomeningitis virus and Lassa fever virus. Science 282: 2079–2081, 1998.



- [69027] 7357.Ottman, R.; Risch, N.; Hauser, W. A.; Pedley, T. A.; Lee, J. H.;Barker-Cummings, C.; Lustenberger, A.; Nagle, K. J.; Lee, K. S.; Scheuer,M. L.; Neystat, M.; Susser, M.; Wilhelmssen, K. C.: Localization of a gene for partial epilepsy to chromosome 10q. *Nature Genet.* 10:56–60, 1995.
- [69028] 7358.Stogmann, E.; Zimprich, A.; Baumgartner, C.; Aull-Watschinger,S.; Holtt, V.; Zimprich, F.: A functional polymorphism in the prodynorphin gene promoter is associated with temporal lobe epilepsy. *Ann. Neurol.* 51:260–263, 2002.
- [69029] 7359.Summar, M. L.; Phillips, J. A., III; Battey, J.; Castiglione, C.M.; Kidd, K. K.; Maness, K. J.; Weiffenbach, B.; Gravius, T. C.:Linkage relationships of human arginine vasopressin–neurophysin–II and oxytocin–neurophysin–I to prodynorphin and other loci on chromosome 20. *Molec. Endocr.* 4: 947–950, 1990.
- [69030] 7360.Zimprich, A.; Kraus, J.; Woltje, M.; Mayer, P.; Rauch, E.; Holtt,V.: An allelic variation in the human prodynorphin gene promoter alters stimulus-induced expression. *J. Neurochem.* 74: 472–477, 2000.
- [69031] 7361.Craig, S. P.; Day, I. N. M.; Thompson, R. J.; Craig, I. W.: Localization of human neurone-specific enolase to chromosome 12p13. (Abstract) *Cytogenet.Cell Genet.* 51:

980 only, 1989.

- [69032] 7362.Craig, S. P.; Day, I. N. M.; Thompson, R. J.; Craig, I. W.: Localisation of neurone-specific enolase (ENO2) to 12p13. Cytogenet. Cell Genet. 54:71–73, 1990.
- [69033] 7363.Feo, S.; Oliva, D.; Barbieri, G.; Xu, W.; Fried, M.; Giallongo, A.: The gene for the muscle-specific enolase is on the short arm of human chromosome 17. Genomics 6: 192–194, 1990.
- [69034] 7364.Grzeschik, K.-H.: Assignment of human genes: beta-glucuronidase to chromosome 7, adenylate kinase-1 to 9, a second enzyme with enolase activity to 12, and mitochondrial IDH to 15. Cytogenet. Cell Genet. 16:142–148, 1976. Note: Alternate: Birth Defects Orig. Art. Ser. 12(7):142–148, 1976.
- [69035] 7365.Herbschleb-Voogt, E.; Monteba-van Heuvel, M.; Wijnen, L. M. M.; Westerveld, A.; Pearson, P. L.; Meera Khan, P.: Chromosomal assignment and regional localization of CS, ENO-2, GAPDH, LDH-B, PEP-B and TPI in man-rodent cell hybrids. Cytogenet. Cell Genet. 22: 482–486, 1978.
- [69036] 7366.Hinks, L. J.; Day, I. N. M.: Further studies of enolase loci. (Abstract) Cytogenet. Cell Genet. 58: 1854 only, 1991.
- [69037] 7367.Law, M. L.; Kao, F.: Regional mapping of the gene coding for enolase-2 on human chromosome 12. J. Cell

Sci. 53: 245–254, 1982.

- [69038] 7368. Mattei, J. F.; Baeteman, M. A.; Mattei, M. G.; Ardissonne, J. P.; Giraud, F.: Regional assignments of CS and ENO2 on chromosome 12. (Abstract) Cytogenet. Cell Genet. 32: 297 only, 1982.
- [69039] 7369. Oliva, D.; Cali, L.; Feo, S.; Giallongo, A.: Complete structure of the human gene encoding neuron-specific enolase. Genomics 10:157–165, 1991.
- [69040] 7370. Jenkins, N. A.; Justice, M. J.; Gilbert, D. J.; Chu, M.-L.; Copeland, N. G.: Nidogen/entactin (Nid) maps to the proximal end of mouse chromosome 13 linked to beige (bg) and identifies a new region of homology between mouse and human chromosomes. Genomics 9: 401–403, 1991.
- [69041] 7371. Olavesen, M. G.; Bentley, E.; Mason, R. V. F.; Stephens, R. J.; Ragoussis, J.: Fine mapping of 39 ESTs on human chromosome 6p23–p25. Genomics 46:303–306, 1997.
- [69042] 7372. Boyd, C. D.; Toth-Fejel, S.; Gadi, I. K.; Litt, M.; Condon, M. R.; Kolbe, M.; Hagen, I. K.; Kurkinen, M.; Mackenzie, J. W.; Magenis, E.: The genes coding for human pro alpha-1(IV) collagen and pro alpha-2(IV) collagen are both located at the end of the long arm of chromosome 13. Am.

J. Hum. Genet. 42: 309–314, 1988.

[69043] 7373. Brazel, D.; Pollner, R.; Oberbaumer, I.; Kuhn, K.: Human basementmembrane collagen (type IV): the amino acid sequence of the alpha-2(IV)chain and its comparison with the alpha-1(IV) chain reveals deletions in the alpha-1(IV) chain. Europ. J. Biochem. 172: 35–42, 1988.

[69044] 7374. Griffin, C. A.; Emanuel, B. S.; Hansen, J. R.; Cavenee, W. K.; Myers, J. C.: Human collagen genes encoding basement membrane alpha-1(IV) and alpha-2(IV) chains map to the distal long arm of chromosome 13. Proc. Nat. Acad. Sci. 84: 512–516, 1987.

[69045] 7375. Haniel, A.; Welge-Lussen, U.; Kuhn, K.; Poschl, E.: Identification and characterization of a novel transcriptional silencer in the human collagen type IV gene COL4A2. J. Biol. Chem. 270: 11209–11215, 1995.

[69046] 7376. Killen, P. D.; Francomano, C. A.; Yamada, Y.; Modi, W. S.; O'Brien, S. J.: Partial structure of the human alpha-2(IV) collagen chain and chromosomal localization of the gene (COL4A2). Hum. Genet. 77: 318–324, 1987.

[69047] 7377. Solomon, E.; Hall, V.; Kurkinen, M.: The human alpha-2(IV) collagen gene, COL4A2, is syntenic with the alpha-1(IV) gene, COL4A1, on chromosome 13. Ann. Hum. Genet. 51: 125–127, 1987.

- [69048] 7378.Davis, R. C.; Xia, Y.; Mohandas, T.; Schotz, M. C.; Lysis, A. J.: Assignment of the human pancreatic colipase gene to chromosome 6p21.1to pter. *Genomics* 10: 262–265, 1991.
- [69049] 7379.Sims, H. F.; Lowe, M. E.: The human colipase gene: isolation,chromosomal location, and tissue-specific expression. *Biochemistry* 31:7120–7125, 1992.
- [69050] 7380.Alper, C. A.; Marcus, D.; Raum, D.; Petersen, B. H.; Spira, T.J.: Genetic polymorphism in C8 beta-chains: evidence for two unlinkedgenetic loci for the eighth component of human complement (C8). *J.Clin. Invest.* 72: 1526–1531, 1983.
- [69051] 7381.Densen, P.; Brown, E. J.; O'Neill, G. J.; Tedesco, F.; Clark, R.A.; Frank, M. M.; Webb, D.; Myers, J.: Inherited deficiency of C8in a patient with recurrent meningococcal infections: further evidencefor a dysfunctional C8 molecule and nonlinkage to the HLA system. *J.Clin. Immun.* 3: 90–99, 1983.
- [69052] 7382.Giraldo, G.; Degos, L.; Beth, E.; Sasportes, M.; Marcelli, A.;Gharbi, R.; Day, N. K.: C8 deficiency in a family with xerodermapigmentosum: lack of linkage to HLA region. *Clin. Immun. Immunopath.* 8:377–384, 1977.
- [69053] 7383.Jasin, H. E.: Absence of the eighth component of

complement in association with systemic lupus erythematosus-like disease. J. Clin. Invest. 60: 709–715, 1977.

[69054] 7384. Kolb, W. P.; Muller-Eberhard, H. J.: The membrane attack mechanism of complement: the three polypeptide chain structure of the eighth component (C8). J. Exp. Med. 143: 1131–1139, 1976.

[69055] 7385. Komatsu, M.; Imaoka, K.; Satoh, M.; Mikami, H.: Hereditary C8- $\alpha$ - $\gamma$  deficiency associated with dwarfism in the rabbit. J. Hered. 81: 413–417, 1990.

[69056] 7386. Komatsu, M.; Yamamoto, K.; Kawashima, T.; Migita, S.: Genetic deficiency of the  $\alpha$ - $\gamma$ -subunit of the eighth complement component in the rabbit. J. Immun. 134: 2607–2609, 1985.

[69057] 7387. Marcus, D.; Spira, T. J.; Petersen, B. H.; Raum, D.; Alper, C. A.: There are two unlinked genetic loci for human C8. (Abstract) Molec. Immun. 19: 1385 only, 1982.

[69058] 7388. Matthews, N.; Stark, J. M.; Harper, P. S.; Doran, J.; Jones, D. M.: Recurrent meningococcal infections associated with a functional deficiency of the C8 component of human complement. Clin. Exp. Immun. 39: 53–59, 1980.

[69059] 7389. Merritt, A. D.; Petersen, B. H.; Biegel, A. A.; Meyers, D. A.; Brooks, G. F.; Hodes, M. E.: Chromosome 6: linkage of the eighth component of complement (C8) to the histo-

compatibility region (HLA). Cytogenet.Cell Genet. 16: 331–334, 1976.

[69060] 7390.Michelotti, G. A.; Snider, J. V.; Sodetz, J. M.: Genomic organization of human complement protein C8–alpha and further examination of its linkage to C8–beta. Hum. Genet. 95: 513–518, 1995.

[69061] 7391.Nakamura, S.; Ohue, O.; Abe, K.: Genetic polymorphism of human complement component C81 in the Japanese population. Hum. Genet. 72:344–347, 1986.

[69062] 7392.Pericak–Vance, M. A.; Elston, R. C.; Spira, T. J.; Band, J.: Segregation and linkage analysis of immunochemical C8 levels in a family with C8 beta–chain deficiency. (Abstract) Am. J. Hum. Genet. 34:109A only, 1982.

[69063] 7393.Petersen, B. H.; Graham, J. A.; Brooks, G. F.: Human deficiency of the eighth component of complement: the requirement of C8 for serum *Neisseria gonorrhoeae* bactericidal activity. J. Clin. Invest. 57:283–290, 1976.

[69064] 7394.Pickering, R. J.; Rynes, R. I.; LoCascio, N.; Monahan, J. B.; Sodetz, J. M.: Identification of the alpha–gamma subunit of the eighth component of complement (C8) in a patient with systemic lupus erythematosus and absent C8 activity: patient and family studies. Clin. Immunol.Immunopath. 23: 323–334, 1982.

- [69065] 7395.Rittner, C.; Hargesheimer, W.; Mollenhauer, E.: Population and formal genetics of the human C81(alpha-gamma) polymorphism. Hum.Genet. 67: 166-169, 1984.
- [69066] 7396.Rittner, C.; Hargesheimer, W.; Stradmann, B.; Bertrams, J.; Baur, M. P.; Petersen, B. H.: Human C81 (alpha-gamma) polymorphism: detection in the alpha-gamma subunit on SDS-PAGE, formal genetics and linkage relationship. Am. J. Hum. Genet. 38: 482-491, 1986.
- [69067] 7397.Rogde, S.; Gedde-Dahl, T., Jr.; Teisberg, P.; Jonassen, R.; Hoyheim, B.; Olaisen, B.: Linkage and association studies with C8A and C8BRFLPs on chromosome 1. Ann. Hum. Genet. 56: 233-242, 1992.
- [69068] 7398.Rogde, S.; Mevag, B.; Olaisen, B.; Gedde-Dahl, T., Jr.; Teisberg, P.: Structural genes for complement factor C8 on chromosome 1. (Abstract) Cytogenet. Cell Genet. 37: 571 only, 1984.
- [69069] 7399.Rogde, S.; Mevag, B.; Teisberg, P.; Gedde-Dahl, T., Jr.; Tedesco, F.; Olaisen, B.: Genetic polymorphism of complement component C8. Hum.Genet. 70: 211-216, 1985.
- [69070] 7400.Rogde, S.; Olaisen, B.; Gedde-Dahl, T., Jr.; Teisberg, P.: Two complement component C8 loci are localized between PGM1 and Rh on chromosome 1. (Abstract) Cyto-



genet. Cell Genet. 40: 734–735, 1985.

- [69071] 7401.Rogde, S.; Olaisen, B.; Gedde–Dahl, T., Jr.; Teisberg, P.: The C8A and C8B loci are closely linked on chromosome 1. Ann. Hum. Genet. 50:139–144, 1986.
- [69072] 7402.Tedesco, F.; Densen, P.; Villa, M. A.; Petersen, B. H.; Sirchia, G.: Two types of dysfunctional eighth component of complement (C8) molecules in C8 deficiency in man: reconstitution of normal C8 from the mixture of two abnormal C8 molecules. J. Clin. Invest. 71: 183–191, 1983.
- [69073] 7403.Quax, W.; Meera Khan, P.; Quax–Jeuken, Y.; Bloemendal, H.: The human desmin and vimentin genes are located on different chromosomes. Gene 38:189–196, 1985.
- [69074] 7404.Reynolds, F. H., Jr.; Todaro, G. J.; Fryling, C.; Stephenson, J. R.: Human transforming growth factors induce tyrosine phosphorylation of EGF receptors. Nature 292: 259–262, 1981.
- [69075] 7405.Shimizu, N.; Behzadian, M. A.; Shimizu, Y.: Genetics of cell surface receptors for bioactive polypeptides: binding of epidermal growth factor is associated with the presence of human chromosome 7 in human–mouse cell hybrids. Proc. Nat. Acad. Sci. 77: 3600–3604, 1980.
- [69076] 7406.Sibilia, M.; Fleischmann, A.; Behrens, A.; Stingl, L.;

Carroll,J.; Watt, F. M.; Schlessinger, J.; Wagner, E. F.: The EGF receptor provides an essential survival signal for SOS-dependent skin tumor development. *Cell* 102: 211–220, 2000.

[69077] 7407.Silver, J.; Whitney, J. B., III; Kozak, C.; Hollis, G.; Kirsch,I.: Erbb is linked to the alpha-globin locus on mouse chromosome11. *Molec. Cell. Biol.* 5: 1784–1786, 1985.

[69078] 7408.Spurr, N. K.; Goodfellow, P. N.; Solomon, E.; Parkar, M.; Vennstrom,B.; Bodmer, W. F.: Mapping of cellular oncogenes; erb B on chromosome7. (Abstract) *Cytogenet. Cell Genet.* 37: 590 only, 1984.

[69079] 7409.Spurr, N. K.; Solomon, E.; Jansson, M.; Sheer, D.; Goodfellow,P. N.; Bodmer, W. F.; Vennstrom, B.: Chromosomal localisation of the human homologues to the oncogenes erbA and B. *EMBO J.* 3: 159–163,1984.

[69080] 7410.Thaug, C.; West, K.; Clark, B. J.; McKie, L.; Morgan, J. E.;Arnold, K.; Nolan, P. M.; Peters, J.; Hunter, A. J.; Brown, S. D.M.; Jackson, I. J.; Cross, S. H.: Novel ENU-induced eye mutations in the mouse: models for human eye disease. *Hum. Molec. Genet.* 11:755–767, 2002.

[69081] 7411.Ullrich, A.; Coussens, L.; Hayflick, J. S.; Dull, T. J.; Gray,A.; Tam, A. W.; Lee, J.; Yarden, Y.; Libermann, T. A.;

Schlessinger, J.; Downward, J.; Mayes, E. L. V.; Whittle, N.; Waterfield, M. D.; Seeburg, P. H.: Human epidermal growth factor receptor cDNA sequence and aberrant expression of the amplified gene in A431 epidermoid carcinoma cells. *Nature* 309: 418–425, 1984.

[69082] 7412. Verveer, P. J.; Wouters, F. S.; Reynolds, A. R.; Bastiaens, P. I. H.: Quantitative imaging of lateral ErbB1 receptor signal propagation in the plasma membrane. *Science* 290: 1567–1570, 2000.

[69083] 7413. Wakeling, E. L.; Abu-Amero, S. N.; Stanier, P.; Preece, M. A.; Moore, G. E.: Human EGFR, a candidate gene for the Silver-Russell syndrome, is biallelically expressed in a wide range of fetal tissues. *Europ. J. Hum. Genet.* 6: 158–164, 1998.

[69084] 7414. Yang, E.-B.; Wang, D.-F.; Mack, P.; Cheng, L.-Y.: Genistein, a tyrosine kinase inhibitor, reduces EGF-induced EGF receptor internalization and degradation in human hepatoma HepG2 cells. *Biochem. Biophys. Res. Commun.* 224: 309–317, 1996.

[69085] 7415. Baumann, C. A.; Ribon, V.; Kanzaki, M.; Thurmond, D. C.; Mora, S.; Shigematsu, S.; Bickel, P. E.; Pessin, J. E.; Saltiel, A. R.: CAP defines a second signalling pathway required for insulin-stimulated glucose transport. *Nature*

407: 202–207, 2000.

- [69086] 7416. Bickel, P. E.; Scherer, P. E.; Schnitzer, J. E.; Oh, P.; Lisanti, M. P.; Lodish, H. F.: Flotillin and epidermal surface antigen define a new family of caveolae-associated integral membrane proteins. *J. Biol. Chem.* 272: 13793–13802, 1997.
- [69087] 7417. Cho, Y.-J.; Chema, D.; Moskow, J. J.; Cho, M.; Schroeder, W. T.; Overbeek, P.; Buchberg, A. M.; Duvic, M.: Epidermal surface antigen (MS17S1) is highly conserved between mouse and human. *Genomics* 27: 251–258, 1995.
- [69088] 7418. Kayes, L. M.; Schroeder, W. T.; Marchuk, D. A.; Collins, F. S.; Riccardi, V. M.; Duvic, M.; Stephens, K.: The gene for a novel epidermal antigen maps near the neurofibromatosis 1 gene. *Genomics* 14: 369–376, 1992.
- [69089] 7419. Schroeder, W. T.; Siciliano, M. J.; Stewart-Galetka, S. L.; Duvic, M.: The human gene for an epidermal surface antigen (M17S1) is located at 17q11–12. *Genomics* 11: 481–482, 1991.
- [69090] 7420. Schroeder, W. T.; Stephens, K.; Stewart-Galetka, S.; Riccardi, V.; Duvic, M.: The gene for an epidermal surface antigen is in close proximity to the locus for von Recklinghausen neurofibromatosis. (Abstract) *Clin. Res.* 39: 323A only, 1991.

- [69091] 7421.Cohn, R. D.; Henry, M. D.; Michele, D. E.; Barresi, R.; Saito,F.; Moore, S. A.; Flanagan, J. D.; Skwarchuk, M. W.; Robbins, M. E.;Mendell, J. R.; Williamson, R. A.; Campbell, K. P.: Disruption ofDag1 in differentiated skeletal muscle reveals a role for dystroglycanin muscle regeneration. *Cell* 110: 639–648, 2002.
- [69092] 7422.Matsuoka, R.; Yoshida, M. C.; Furutani, Y.; Imamura, S.; Kanda,N.; Yanagisawa, M.; Masaki, T.; Takao, A.: Human smooth muscle myosinheavy chain gene mapped to chromosomal region 16q12. *Am. J. Med.Genet.* 46: 61–67, 1993.
- [69093] 7423.Apperley, J. F.; Gardembas, M.; Melo, J. V.; Russell-Jones, R.;Bain, B. J.; Baxter, E. J.; Chase, A.; Chessells, J. M.; Colombat,M.; Dearden, C. E.; Dimitrijevic, S.; Mahon, F.-X.; Marin, D.; Nikolova,Z.; Olavarria, E.; Silberman, S.; Schultheis, B.; Cross, N. C. P.;Goldman, J. M.: Response to imatinib mesylate inpatients with chronicmyeloproliferative diseases with rearrangements of the platelet-derivedgrowth factor receptor beta. *New Eng. J. Med.* 347: 481–487, 2002.
- [69094] 7424.Matsuoka, R.; Yoshida, M. C.; Kanda, N.; Furutani, Y.; Bruns, G.;Yanagisawa, M.; Masaki, T.; Takao, A.: Human smooth muscle myosinheavy-chain gene mapped to

chromosomal region 16q12.1–q12.2. (Abstract) Cytogenet. Cell Genet. 58: 2000–2001, 1991.

- [69095] 7425. Spear, P. G.: A welcome mat for leprosy and Lassa fever. Science 282:1999–2000, 1998.
- [69096] 7426. Tinsley, J. M.; Blake, D. J.; Zuellig, R. A.; Davies, K. E.: Increasing complexity of the dystrophin-associated protein complex. Proc. Nat. Acad. Sci. 91: 8307–8313, 1994.
- [69097] 7427. Williamson, R. A.; Henry, M. D.; Daniels, K. J.; Hrstka, R. F.; Lee, J. C.; Sunada, Y.; Ibraghimov-Beskrovnaya, O.; Campbell, K. P.: Dystroglycan is essential for early embryonic development: disruption of Reichert's membrane in Dag1-null mice. Hum. Molec. Genet. 6:831–841, 1997.
- [69098] 7428. Yamada, H.; Denzer, A. J.; Hori, H.; Tanaka, T.; Anderson, L. V. B.; Fujita, S.; Fukuta-Ohi, H.; Shimizu, T.; Ruegg, M. A.; Matsumura, K.: Dystroglycan is a dual receptor for agrin and laminin-2 in Schwann cell membrane. J. Biol. Chem. 271: 23418–23423, 1996.
- [69099] 7429. Cote, P. D.; Moukhles, H.; Lindenbaum, M.; Carbonetto, S.: Chimeric mice deficient in dystroglycans develop muscular dystrophy and have disrupted myoneural synapses. Nature Genet. 23: 338–342, 1999.
- [69100] 7430. Gee, S. H.; Montanaro, F.; Lindenbaum, M. H.; Car-

bonetto, S.:Dystroglycan- $\alpha$ , a dystrophin-associated glycoprotein, is a functional agrin receptor. *Cell* 77: 675–686, 1994.

[69101] 7431.Gorecki, D. C.; Derry, J. M. J.; Barnard, E. A.: Dystroglycan:brain localisation and chromosome mapping in the mouse. *Hum. Molec.Genet.* 3: 1589–1597, 1994.

[69102] 7432.Hayashi, Y. K.; Ogawa, M.; Tagawa, K.; Noguchi, S.; Ishihara, T.;Nonaka, I.; Arahata, K.: Selective deficiency of  $\alpha$ -dystroglycanin Fukuyama-type congenital muscular dystrophy. *Neurology* 57: 115–121,2001.

[69103] 7433.Henry, M. D.; Campbell, K. P.: A role for dystroglycan in basementmembrane assembly. *Cell* 95: 859–970, 1998.

[69104] 7434.Ibraghimov-Beskrovnaya, O.; Ervasti, J. M.; Leveille, C. J.; Slaughter,C. A.; Sernett, S. W.; Campbell, K. P.: Primary structure of dystrophin-associatedglycoproteins linking dystrophin to the extracellular matrix. *Nature* 355:696–702, 1992.

[69105] 7435.Ibraghimov-Beskrovnaya, O.; Milatovich, A.; Ozcelik, T.; Yang,B.; Francke, U.; Campbell, K. P.: Dystroglycan: tissue distribution,human muscle cDNA, genomic structure and chromosome mapping. (Abstract) *Am.J. Hum. Genet.* 51 (suppl.): A130 only, 1992.

- [69106] 7436.Ibraghimov-Beskrovnaya, O.; Milatovich, A.; Ozcelik, T.; Yang,B.; Koepnick, K.; Francke, U.; Campbell, K. P.: Human dystroglycan:skeletal muscle cDNA, genomic structure, origin of tissue specific isoforms and chromosomal localization. Hum. Molec. Genet. 2: 1651–1657,1993.
- [69107] 7437.Ma, J.; Nastuk, M. A.; McKechnie, B. A.; Fallon, J. R.: The agrin receptor: localization in the postsynaptic membrane, interaction with agrin, and relationship to the acetylcholine receptor. J. Biol. Chem. 268:25108–25117, 1993.
- [69108] 7438.Matsumura, K.; Nonaka, I.; Campbell, K. P.: Abnormal expression of dystrophin-associated proteins in Fukuyama-type congenital muscular dystrophy. Lancet 341: 521–522, 1993.
- [69109] 7439.Matsumura, K.; Tome, F. M. S.; Collin, H.; Azibi, K.; Chaouch,M.; Kaplan, J.-C.; Fardeau, M.; Campbell, K. P.: Deficiency of the 50K dystrophin-associated glycoprotein in severe childhood autosomal recessive muscular dystrophy. Nature 359: 320–322, 1992.
- [69110] 7440.Matsumura, K.; Tome, F. M. S.; Ionasescu, V.; Ervasti, J. M.; Anderson, R. D.; Romero, N. B.; Simon, D.; Re-can, D.; Kaplan, J.-C.; Fardeau, M.; Campbell, K. P.: Deficiency of dystrophin-associated proteins in Duchenne



muscular dystrophy patients lacking COOH-terminal domains of dystrophin. J. Clin. Invest. 92: 866–871, 1993.

- [69111] 7441. Michele, D. E.; Barresi, R.; Kanagawa, M.; Saito, F.; Cohn, R.D.; Satz, J. S.; Dollar, J.; Nishino, I.; Kelley, R. I.; Somer, H.; Straub, V.; Mathews, K. D.; Moore, S. A.; Campbell, K. P.: Post-translational disruption of dystroglycan–ligand interactions in congenital muscular dystrophies. Nature 418: 417–422, 2002.
- [69112] 7442. Moore, S. A.; Saito, F.; Chen, J.; Michele, D. E.; Henry, M. D.; Messing, A.; Cohn, R. D.; Ross–Barta, S. E.; Westra, S.; Williamson, R. A.; Hoshi, T.; Campbell, K. P.: Deletion of brain dystroglycan recapitulates aspects of congenital muscular dystrophy. Nature 418: 422–425, 2002.
- [69113] 7443. Rambukkana, A.; Yamada, H.; Zanazzi, G.; Mathus, T.; Salzer, J.L.; Yurchenco, P. D.; Campbell, K. P.; Fischetti, V. A.: Role of  $\alpha$ -dystroglycan as a Schwann cell receptor for *Mycobacterium leprae*. Science 282: 2076–2078, 1998.
- [69114] 7444. Sealock, R.; Froehner, S. C.: Dystrophin–associated proteins and synapse formation: is  $\alpha$ -dystroglycan the agrin receptor? Cell 77: 617–619, 1994.

- [69115] 7445. Yamada, H.; Saito, F.; Fukuta-Ohi, H.; Zhong, D.; Hase, A.; Arai, K.; Okuyama, A.; Maekawa, R.; Shimizu, T.; Matsumura, K.: Processing of beta-dystroglycan by matrix metalloproteinase disrupts the link between the extracellular matrix and cell membrane via the dystroglycan complex. *Hum. Molec. Genet.* 10: 1563–1569, 2001.
- [69116] 7446. Blake, D. J.; Love, D. R.; Tinsley, J.; Morris, G. E.; Turley, H.; Gatter, K.; Dickson, G.; Edwards, Y. H.; Davies, K. E.: Characterization of a 4.8kb transcript from the Duchenne muscular dystrophy locus expressed in schwannoma cells. *Hum. Molec. Genet.* 1: 103–109, 1992.
- [69117] 7447. Golub, T. R.; Barker, G. F.; Lovett, M.; Gilliland, D. G.: Fusion of PDGF receptor beta to a novel ets-like gene, tel, in chronic myelomonocytic leukemia with t(5;12) chromosomal translocation. *Cell* 77: 307–316, 1994.
- [69118] 7448. Brissenden, J. E.; Ullrich, A.; Francke, U.: Chromosomal mapping of loci for insulin-like growth factors I and II and for epidermal growth factor in man. (Abstract) *Am. J. Hum. Genet.* 36: 133S only, 1984.
- [69119] 7449. Brissenden, J. E.; Ullrich, A.; Francke, U.: Human chromosomal mapping of genes for insulin-like growth factors I and II and epidermal growth factor. *Nature* 310: 781–784, 1984.

- [69120] 7450.Carpenter, G.; Cohen, S.: Epidermal growth factor. Ann. Rev.Biochem. 48: 193–216, 1979.
- [69121] 7451.Cohen, S.: Isolation of a mouse submaxillary gland protein acceleratingincisor eruption and eyelid opening in the new-born animal. J. Biol.Chem. 237: 1555–1562, 1962.
- [69122] 7452.Gray, A.; Dull, T. J.; Ullrich, A.: Nucleotide sequence of epidermalgrowth factor cDNA predicts a 128,000–molecular weight protein precursor. Nature 303:722–725, 1983.
- [69123] 7453.Morton, C. C.; Byers, M. G.; Nakai, H.; Bell, G. I.; Shows, T.B.: Human genes for insulin–like growth factors I and II and epidermalgrowth factor are located on 12q22–q24.1, 11p15, and 4q25–q27, respectively. Cyto–genet.Cell Genet. 41: 245–249, 1986.
- [69124] 7454.Sassone–Corsi, P.; Mizzen, C. A.; Cheung, P.; Crosjo, C.; Monaco,L.; Jacquot, S.; Hanauer, A.; Allis, C. D.: Re–quirement of Rsk–2for epidermal growth factor–activated phosphorylation of histone H3. Science 285:886–891, 1999.
- [69125] 7455.Scott, A. F.: Personal Communication. Baltimore, Md. 10/11/1999.
- [69126] 7456.Shahbazi, M.; Pravica, V.; Nasreen, N.; Fakhoury, H.;

Fryer, A.A.; Strange, R. C.; Hutchinson, P. E.; Osborne, J. E.; Lear, J. T.; Smith, A. G.; Hutchinson, I. V.: Association between functional polymorphism in EGF gene and malignant melanoma. *Lancet* 359: 397–401, 2002.

[69127] 7457. Smith, J.; Cook, E.; Fotheringham, I.; Pheby, S.; Derbyshire, R.; Eaton, M. A. W.; Doel, M.; Lilley, D. M. J.; Pardon, J. F.; Patel, T.; Lewis, H.; Bell, L. D.: Chemical synthesis and cloning of a gene for human beta-urogastrone. *Nucleic Acids Res.* 10: 4467–4482, 1982.

[69128] 7458. Sudhof, T. C.; Russell, D. W.; Goldstein, J. L.; Brown, M. S.; Sanchez-Pescador, R.; Bell, G. I.: Cassette of eight exons shared by genes for LDL receptor and EGF precursor. *Science* 228: 893–895, 1985.

[69129] 7459. Tsutsumi, O.; Kurachi, H.; Oka, T.: A physiological role of epidermal growth factor in male reproductive function. *Science* 233: 975–977, 1986.

[69130] 7460. Urdea, M. S.; Merryweather, J. P.; Mullenbach, G. T.; Coit, D.; Heberlein, U.; Valenzuela, P.; Barr, P. J.: Chemical synthesis of a gene for human epidermal growth factor urogastrone and its expression in yeast. *Proc. Nat. Acad. Sci.* 80: 7461–7465, 1983.

[69131] 7461. Zabel, B. U.; Eddy, R. L.; Lalley, P. A.; Scott, J.; Bell, G. I.; Shows, T. B.: Chromosomal locations of the human

and mouse genes for precursors of epidermal growth factor and the beta subunit of nerve growth factor. *Proc. Nat. Acad. Sci.* 82: 469–473, 1985.

[69132] 7462. Aden, D. P.; Knowles, B. B.: Cell surface antigens coded for by the human chromosome 7. *Immunogenetics* 3: 209–211, 1976.

[69133] 7463. Carlin, C. R.; Aden, D. P.; Knowles, B. B.: S6 is the human receptor for epidermal growth factor (EGF). (Abstract) *Cytogenet. Cell Genet.* 32:256 only, 1982.

[69134] 7464. Carlin, C. R.; Knowles, B. B.: Identity of human epidermal growth factor (EGF) receptor with glycoprotein SA-7: evidence for differential phosphorylation of the two components of the EGF receptor from A431 cells. *Proc. Nat. Acad. Sci.* 79: 5026–5030, 1982.

[69135] 7465. Carpenter, G.: Properties of the receptor for epidermal growth factor. *Cell* 37: 357–358, 1984.

[69136] 7466. Chen, B.; Bronson, R. T.; Klamann, L. D.; Hampton, T. G.; Wang, J.; Green, P. J.; Magnuson, T.; Douglas, P. S.; Morgan, J. P.; Neel, B. G.: Mice mutant for *Egfr* and *Shp2* have defective cardiac semilunar valvulogenesis. *Nature Genet.* 24: 296–299, 2000.

[69137] 7467. Ishii, H.; Baffa, R.; Numata, S.-I.; Murakumo, Y.; Rattan, S.; Inoue, H.; Mori, M.; Fidanza, V.; Alder, H.; Croce, C.

M.: The FEZ1 gene at chromosome 8p22 encodes a leucine-zipper protein, and its expression is altered in multiple human tumors. *Proc. Nat. Acad. Sci.* 96: 3928–3933, 1999.

- [69138] 7468. Kuroki, T.; Trapasso, F.; Shiraishi, T.; Alder, H.; Mimori, K.; Mori, M.; Croce, C. M.: Genetic alterations of the tumor suppressor gene WWOX in esophageal squamous cell carcinoma. *Cancer Res.* 62:2258–2260, 2002.
- [69139] 7469. Li, G.; Hu, N.; Goldstein, A. M.; Tang, Z.-Z.; Roth, M. J.; Wang, Q.-H.; Dawsey, S. M.; Han, X.-Y.; Ding, T.; Huang, J.; Giffen, C.; Taylor, P. R.; Emmert-Buck, M. R.: Allelic loss on chromosome bands 13q11–q13 in esophageal squamous cell carcinoma. *Genes Chromosomes Cancer* 31: 390–397, 2001.
- [69140] 7470. Lo, H. S.; Hu, N.; Gere, S.; Lu, N.; Su, H.; Goldstein, A. M.; Taylor, P. R.; Lee, M. P.: Identification of somatic mutations of the RNF6 gene in human esophageal squamous cell carcinoma. *Cancer Res.* 62: 4191–4193, 2002.
- [69141] 7471. Su, P.-H.; Hou, J.-W.; Hwu, W.-L.; Wu, M.-H.; Wang, J.-K.; Wang, T.-R.: Congenital contractural arachnodactyly (Beals syndrome). *Acta Paediat.* 41: 59–62, 2000.
- [69142] 7472. Viljoen, D.: Congenital contractural arachnodactyly (Beals syndrome). *J. Med. Genet.* 31: 640–643, 1994.

- [69143] 7473.Viljoen, D.; Ramesar, R.; Behari, D.: Beals syndrome: clinical and molecular investigations in a kindred of Indian descent. *Clin.Genet.* 39: 181–188, 1991.
- [69144] 7474.Wang, M.; Clericuzio, C. L.; Godfrey, M.: Familial occurrence of typical and severe lethal congenital contractural arachnodactyly caused by missplicing of exon 34 of fibrillin-2. *Am. J. Hum. Genet.* 59:1027–1034, 1996.
- [69145] 7475.Wang, M.; Price, C. E.; Han, J.; Cisler, J.; Imaizumi, K.; VanThienen, M. N.; DePaepe, A.; Godfrey, M.: Recurrent mis-splicing of fibrillin exon 32 in two patients with neonatal Marfan syndrome. *Hum.Molec. Genet.* 4: 607–613, 1995.
- [69146] 7476.Wang, M.; Tsipouras, P.; Godfrey, M.: Fibrillin-2 (FBN2) mutation in congenital contractural arachnodactyly. (Abstract) *Am. J. Hum.Genet.* 57: A231, 1995.
- [69147] 7477.Zhang, H.; Apfelroth, S. D.; Hu, W.; Davis, E. C.; Sanguineti, C.; Bonadio, J.; Mecham, R. P.; Ramirez, F.: Structure and expression of fibrillin-2, a novel microfibrillar component preferentially located in elastic matrices. *J. Cell Biol.* 124: 855–863, 1994.
- [69148] 7478.Zhang, H.; Hu, W.; Ramirez, F.: Developmental expression of fibrillin genes suggests heterogeneity of extracellular microfibrils. *J. Cell Biol.* 129: 1165–1176, 1995.

- [69149] 7479. Beck, C.; Moulard, B.; Steinlein, O.; Guipponi, M.; Vallee, L.; Montpied, P.; Baldy-Moulinier, M.; Malafosse, A.: A nonsense mutation in the alpha-4 subunit of the nicotinic acetylcholine receptor (CHRNA4) cosegregates with 20q-linked benign neonatal familial convulsions (EBN1). *Neurobiol. Dis.* 1: 95–99, 1994.
- [69150] 7480. Singh, N. A.; Charlier, C.; Stauffer, D.; DuPont, B. R.; Leach, R. J.; Melis, R.; Ronen, G. M.; Bjerre, I.; Quattlebaum, T.; Murphy, J. V.; McHarg, M. L.; Gagnon, D.; Rosales, T. O.; Peiffer, A.; Anderson, V. E.; Leppert, M.: A novel potassium channel gene, KCNQ2, is mutated in an inherited epilepsy of newborns. *Nature Genet.* 18: 25–29, 1998.
- [69151] 7481. Charlier, C.; Singh, N. A.; Ryan, S. G.; Lewis, T. B.; Reus, B. E.; Leach, R. J.; Leppert, M.: A pore mutation in a novel KQT-like potassium channel gene in an idiopathic epilepsy family. *Nature Genet.* 18: 53–55, 1998.
- [69152] 7482. Ryan, S. G.; Wiznitzer, M.; Hollman, C. H.; Torres, M. C.; Szekeresova, M.; Schneider, S.: Benign familial neonatal convulsions: evidence for clinical and genetic heterogeneity. *Ann. Neurol.* 29: 469–473, 1991.
- [69153] 7483. Earnshaw, W. C.: When is a centromere not a kinetochore? *J. Cell Sci.* 99: 1–4, 1991.
- [69154] 7484. Fowler, K. J.; Newson, A. J.; MacDonald, A. C.; Kalit-



sis, P.; Lyu, M. S.; Kozak, C. A.; Choo, K. H. A.: Chromosomal localization of mouse Cenpa gene. *Cytogenet. Cell Genet.* 79: 298–301, 1997.

[69155] 7485. Howman, E. V.; Fowler, K. J.; Newson, A. J.; Redward, S.; MacDonald, A. C.; Kalitsis, P.; Choo, K. H. A.: Early disruption of centromeric chromatin organization in centromere protein A (Cenpa) null mice. *Proc. Nat. Acad. Sci.* 97: 1148–1153, 2000.

[69156] 7486. Palmer, D. K.; O'Day, K.; Trong, H. L.; Charbonneau, H.; Margolis, R. L.: Purification of the centromere-specific protein CENP-A and demonstration that it is a distinctive histone. *Proc. Nat. Acad. Sci.* 88: 3734–3738, 1991.

[69157] 7487. Sullivan, K. F.; Hechenberger, M.; Masri, K.: Human CENP-A contains a histone H3 related histone fold domain that is required for targeting to the centromere. *J. Cell Biol.* 127: 581–592, 1994.

[69158] 7488. Earnshaw, W. C.; Sullivan, K. F.; Machlin, P. S.; Cooke, C. A.; Kaiser, D. A.; Pollard, T. D.; Rothfield, N. F.; Cleveland, D. W.: Molecular cloning of cDNA for CENP-B, the major human centromere autoantigen. *J. Cell Biol.* 104: 817–829, 1987.

[69159] 7489. Seki, N.; Saito, T.; Kitagawa, K.; Masumoto, H.; Okazaki, T.; Hori, T.-A.: Mapping of the human cen-

tromere protein B gene (CENPB) to chromosome 20p13 by fluorescence in situ hybridization. *Genomics* 24:187–188, 1994.

[69160] 7490. Sugimoto, K.; Yata, H.; Himeno, M.: Mapping of the human CENP-B gene to chromosome 20 and the CENP-C gene to chromosome 12 by a rapid cycle DNA amplification procedure. *Genomics* 17: 240–242, 1993.

[69161] 7491. Earnshaw, W.; Bordwell, B.; Marino, C.; Rothfield, N.: Three human chromosomal autoantigens are recognized by sera from patients with anti-centromere antibodies. *J. Clin. Invest.* 77: 426–430, 1986.

[69162] 7492. Earnshaw, W. C.; Ratrie, H., III; Stetten, G.: Visualization of centromere proteins CENP-B and CENP-C on a stable dicentric chromosome in cytological spreads. *Chromosoma* 98: 1–12, 1989.

[69163] 7493. Earnshaw, W. C.; Rothfield, N.: Identification of a family of human centromere proteins using autoimmune sera from patients with scleroderma. *Chromosoma* 91: 313–321, 1985.

[69164] 7494. Jones, C.; Nguyen, L.; Burkin, D.; McGrew, J.; Tomkiel, J.; Earnshaw, W.: Localization of centromere autoantigen C (CENPC) to human chromosome 12. (Abstract) Human Genome Mapping Workshop 93 25 only, 1993.

- [69165] 7495.McKay, S.; Thomson, E.; Cooke, H.: Sequence homologies and linkagegroup conservation of the human and mouse Cenpc genes. *Genomics* 22:36–40, 1994.
- [69166] 7496.Page, S. L.; Earnshaw, W. C.; Choo, K. H. A.; Shaffer, L. G.:Further evidence that CENP–C is a necessary component of active centromeres:studies of a dic(X;15) with simultaneous immunofluorescence and FISH. *Hum.Molec. Genet.* 4: 289–294, 1995.
- [69167] 7497.Saitoh, H.; Tomkiel, J.; Cooke, C. A.; Ratrie, H., III; Maurer,M.; Rothfield, N. F.; Earnshaw, W. C.: CENP–C, an autoantigen inscleroderma, is a component of the human inner kinetochore plate. *Cell* 70:115–125, 1992.
- [69168] 7498.Gaynor, R. B.; Muchardt, C.; Xia, Y.; Klisak, I.; Mohandas, T.;Sparkes, R. S.; Lysis, A. J.: Localization of the gene for the DNA–bindingprotein AP–2 to human chromosome 6p22.3–pter. *Genomics* 10: 1100–1102,1991.
- [69169] 7499.Schorle, H.; Meier, P.; Buchert, M.; Jaenisch, R.; Mitchell, P.J.: Transcription factor AP–2 essential for cranial closure and craniofacialdevelopment. *Nature* 381: 235–238, 1996.
- [69170] 7500.Warren, G.; Gordon, M.; Siracusa, L. D.; Buchberg, A. M.; Williams,T.: Physical and genetic localization of the gene encoding the AP–2transcription factor to mouse

chromosome 13. *Genomics* 31: 234–237, 1996.

[69171] 7501. Williams, T.; Admon, A.; Luscher, B.; Tjian, R.: Cloning and expression of AP-2, a cell-type-specific transcription factor that activates inducible enhancer elements. *Genes Dev.* 2: 1557–1569, 1988.

[69172] 7502. Williamson, J. A.; Bosher, J. M.; Skinner, A.; Sheer, D.; Williams, T.; Hurst, H. C.: Chromosomal mapping of the human and mouse homologues of two new members of the AP-2 family of transcription factors. *Genomics* 35: 262–264, 1996.

[69173] 7503. Zhang, J.; Hagopian–Donaldson, S.; Serbedzija, G.; Elsemore, J.; Plehn–Dujowich, D.; McMahon, A. P.; Flavell, R. A.; Williams, T.: Neural tube, skeletal and body wall defects in mice lacking transcription factor AP-2. *Nature* 381: 238–241, 1996.

[69174] 7504. Agarwal, V. R.; Ashanullah, C. I.; Simpson, E. R.; Bulun, S. E.: Alternatively spliced transcripts of the aromatase cytochrome P450 (CYP19) gene in adipose tissue of women. *J. Clin. Endocr. Metab.* 82: 70–74, 1997.

[69175] 7505. Nalabolu, S. R.; Shukla, H.; Nallur, G.; Parimoo, S.; Weissman, S. M.: Genes in a 220-kb region spanning the TNF cluster in human MHC. *Genomics* 31: 215–222, 1996.

[69176] 7506. Gao, B.; Guo, J.; She, C.; Shu, A.; Yang, M.; Tan, Z.;

Yang, X.;Guo, S.; Feng, G.; He, L.: Mutations in IHH, encoding Indian hedgehog,cause brachydactyly type A-1. Nature Genet. 28: 386-388, 2001.

[69177] 7507.DeChiara, T. M.; Kimble, R. B.; Poueymirou, W. T.; Rojas, J.; Masiakowski,P.; Valenzuela, D. M.; Yancopoulos, G. D.: Ror2, encoding a receptor-liketyrosine kinase, is required for cartilage and growth plate development. NatureGenet. 24: 271-274, 2000.

[69178] 7508.Oldridge, M.; Fortuna, A. M.; Maringa, M.; Propping, P.; Mansour,S.; Pollitt, C.; DeChiara, T. M.; Kimble, R. B.; Valenzuela, D. M.;Yancopoulos, G. D.; Wilkie, A. O. M.: Dominant mutations in ROR2,encoding an orphan receptor tyrosine kinase, cause brachydactyly typeB. Nature Genet. 24: 375-378, 2000.

[69179] 7509.De Windt, L. J.; Lim, H. W.; Bueno, O. F.; Liang, Q.; Delling,U.; Braz, J. C.; Glascock, B. J.; Kimball, T. F.; del Monte, F.; Hajjar,R. J.; Molkentin, J. D.: Targeted inhibition of calcineurin attenuatescardiac hypertrophy in vivo. Proc. Nat. Acad. Sci. 98: 3322-3327,2001.

[69180] 7510.Fuentes, J. J.; Genesca, L.; Kingsbury, T. J.; Cunningham, K. W.;Perez-Riba, M.; Estivill, X.; de la Luna, S.: DSCR1, overexpressedin Down syndrome, is an inhibitor of calcineurin-mediated signalingpathways. Hum. Molec.

Genet. 9: 1681–1690, 2000.

- [69181] 7511.Giri, P.; Higuchi, S.; Kincaid, R. L.: Chromosomal mapping of the human genes for the calmodulin-dependent protein phosphatase (calcineurin) catalytic subunit. Biochem. Biophys. Res. Commun. 181: 252–258, 1991.
- [69182] 7512.Guerini, D.; Klee, C. B.: Cloning of human calcineurin A: evidence for two isozymes and identification of a polyproline structural domain. Proc. Nat. Acad. Sci. 86: 9183–9187, 1989.
- [69183] 7513.Leinwand, L. A.: Calcineurin inhibition and cardiac hypertrophy: a matter of balance. Proc. Nat. Acad. Sci. 98: 2947–2949, 2001.
- [69184] 7514.Malleret, G.; Haditsch, U.; Genoux, D.; Jones, M. W.; Bliss, T.V. P.; Vanhoose, A. M.; Weitlauf, C.; Kandel, E. R.; Winder, D. G.; Mansuy, I. M.: Inducible and reversible enhancement of learning, memory, and long-term potentiation by genetic inhibition of calcineurin. Cell 104:675–686, 2001.
- [69185] 7515.Mansuy, I. M.; Mayford, M.; Jacob, B.; Kandel, E. R.; Bach, M.E.: Restricted and regulated overexpression reveals calcineurin as a key component in the transition from short-term to long-term memory. Cell 92:39–49, 1998.
- [69186] 7516.Rothermel, B. A.; McKinsey, T. A.; Vega, R. B.; Nicol,

R. L.; Mammen,P.; Yang, J.; Antos, C. L.; Shelton, J. M.; Bassel–Duby, R.; Olson,E. N.; Williams, R. S.: Myocyte–enriched calcineurin–interactingprotein, MCIP1, inhibits cardiac hypertrophy in vivo. Proc. Nat.Acad. Sci. 98: 3328–3333, 2001.

[69187] 7517.Seitz, D. P.; Pasha, M. K.; Singh, B.; Chu, A.; Sharma, R. K.: Localization and characterization of calcineurin in bovine eye. Invest.Ophthal. Vis. Sci. 43: 15–21, 2002.

[69188] 7518.Wang, M. G.; Yi, H.; Guerini, D.; Klee, C. B.; McBride, O. W.: Calcineurin A alpha (PPP3CA), calcineurin A beta (PPP3CB) and calcineurinB (PPP3R1) are located on human chromosomes 4, 10q21–q22 and 2p16–p15respectively. Cytogenet. Cell Genet. 72: 236–241, 1996.

[69189] 7519.Winder, D. G.; Mansuy, I. M.; Osman, M.; Moallem, T. M.; Kandel,E. R.: Genetic and pharmacological evidence for a novel, intermediatephase of long–term potentiation suppressed by calcineurin. Cell 92:25–37, 1998.

[69190] 7520.Bueno, O. F.; Wilkins, B. J.; Tymitz, K. M.; Glascock, B. J.; Kimball,T. F.; Lorenz, J. N.; Molkentin, J. D.: Impaired cardiac hypertrophicresponse in calcineurin A–beta–deficient mice. Proc. Nat. Acad. Sci. 99:4586–4591, 2002.

[69191] 7521.Guerini, D.; Krinks, M. H.; Sikela, J. M.; Hahn, W. E.;

Klee, C.B.: Isolation and sequence of a cDNA clone for human calcineurinB, the  $\text{Ca}(2+)$ -binding subunit of the  $\text{Ca}(2+)$ /calmodulin-stimulated protein phosphatase. DNA 8: 675-682, 1989.

- [69192] 7522. Wang, M. G.; Yi, H.; Guerini, D.; Klee, C. B.; McBride, O. W.: Calcineurin A alpha (PPP3CA), calcineurin A beta (PPP3CB) and calcineurinB (PPP3R1) are located on human chromosomes 4, 10q21-q22 and 2p16-p15 respectively. Cytogenet. Cell Genet. 72: 236-241, 1996.
- [69193] 7523. Ferrari, S.; Calabretta, B.; deRiel, J. K.; Battini, R.; Ghezzi, F.; Lauret, E.; Griffin, C.; Emanuel, B. S.; Gurrieri, F.; Baserga, R.: Structural and functional analysis of a growth-regulated gene, the human calcyclin. J. Biol. Chem. 262: 8325-8332, 1987.
- [69194] 7524. Alevizaki, M.; Stevenson, J. C.; Girgis, S. I.; MacIntyre, I.; Legon, S.: Altered calcitonin gene in a young patient with osteoporosis. Brit. Med. J. 298: 1215-1216, 1989.
- [69195] 7525. Amara, S. G.; Jonas, V.; Rosenfeld, M. G.; Ong, E. S.; Evans, R. M.: Alternative RNA processing in calcitonin gene expression generates mRNAs encoding different polypeptide products. Nature 298: 240-244, 1982.
- [69196] 7526. Aiyar, N.; Rand, K.; Elshourbagy, N. A.; Zeng, Z.;



Adamou, J. E.;Bergsma, D. J.; Li, Y.: A cDNA encoding the calcitonin gene-relatedpeptide type 1 receptor. J. Biol. Chem. 271: 11325–11329, 1996.

[69197] 7527.Fluhmann, B.; Muff, R.; Hunziker, W.; Fischer, J. A.; Born, W.: A human orphan calcitonin receptor-like structure. Biochem. Biophys.Res. Commun. 206: 341–347, 1995.

[69198] 7528.Foord, S. M.; Craig, R. K.: Isolation and characterisation of a human calcitonin-gene-related-peptide receptor. Europ. J. Biochem. 170:373–379, 1987.

[69199] 7529.Foord, S. M.; Marshall, F. H.: RAMPs: accessory proteins for seven transmembrane domain receptors. Trends Pharm. Sci. 20: 184–187,1999.

[69200] 7530.Kamitani, S.; Asakawa, M.; Shimekake, Y.; Kuwasako, K.; Nakahara,K.; Sakata, T.: The RAMP2/CRLR complex is a functional adrenomedullin receptor in human endothelial and vascular smooth muscle cells. FEBS Lett. 448: 111–114, 1999.

[69201] 7531.Nakazawa, I.; Nakajima, T.; Harada, H.; Ishigami, T.; Umemura,S.; Emi, M.: Human calcitonin receptor-like receptor for adrenomedullin:genomic structure, eight single-nucleotide polymorphisms, and haplotype analysis. J. Hum. Genet. 46: 132–136, 2001.

- [69202] 7532.Dolmetsch, R. E.; Pajvani, U.; Fife, K.; Spotts, J. M.; Greenberg,M. E.: Signaling to the nucleus by an L-type calcium channel–calmodulincomplex through the MAP kinase pathway. *Science* 294: 333–339, 2001.
- [69203] 7533.McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba,Canada 2/14/1992.
- [69204] 7534.Powers, P. A.; Gregg, R. G.; Hogan, K.: Linkage mapping of thehuman gene for the alpha–1 subunit of the cardiac DHP–sensitive Ca(2+)channel (CACNL1A1) to chromosome 12p13.2–pter using a dinucleotiderepeat. *Genomics* 14: 206–207, 1992.
- [69205] 7535.Powers, P. A.; Gregg, R. G.; Lalley, P. A.; Liao, M.; Hogan, K.: Assignment of the human gene for the alpha–1 subunit of the cardiacDHP–sensitive Ca(2+) channel (CCHL1A1) to chromosome 12p12–pter. *Genomics* 10:835–839, 1991.
- [69206] 7536.Yang, R.; Morosetti, R.; Koeffler, H. P.: Characterization ofa second human cyclin A that is highly expressed in testis and in several leukemic cell lines. *Cancer Res.* 57: 913–920, 1997.
- [69207] 7537.Sartor, H.; Ehlert, F.; Grzeschik, K.–H.; Muller, R.; Adolph, S.: Assignment of two human cell cycle genes, CDC25C and CCNB1, to 5q31and 5q12, respectively. *Ge–*

nomics 13: 911–912, 1992.

- [69208] 7538.Geng, Y.; Whoriskey, W.; Park, M. Y.; Bronson, R. T.; Medema, R.H.; Li, T.; Weinberg, R. A.; Sicinski, P.: Rescue of cyclin D1 deficiency by knockin cyclin E. *Cell* 97: 767–777, 1999.
- [69209] 7539.Hinchcliffe, E. H.; Li, C.; Thompson, E. A.; Maller, J. L.; Sluder, G.: Requirement of Cdk2–cyclin E activity for repeated centrosome reproduction in *Xenopus* egg extracts. *Science* 283: 851–854, 1999.
- [69210] 7540.Lew, D. J.; Dulic, V.; Reed, S. I.: Isolation of three novel human cyclins by rescue of G1 cyclin (cln) function in yeast. *Cell* 66:1197–1206, 1991.
- [69211] 7541.Li, H.; Lahti, J. M.; Valentine, M.; Saito, M.; Reed, S. I.; Look, A. T.; Kidd, V. J.: Molecular cloning and chromosomal localization of the human cyclin C (CCNC) and cyclin E (CCNE) genes: deletion of the CCNC gene in human tumors. *Genomics* 32: 253–259, 1996.
- [69212] 7542.Moberg, K. H.; Bell, D. W.; Wahrer, D. C. R.; Haber, D. A.; Hariharan, I. K.: Archipelago regulates cyclin E levels in *Drosophila* and is mutated in human cancer cell lines. *Nature* 413: 311–316, 2001.
- [69213] 7543.Sheaff, R. J.; Groudine, M.; Gordon, M.; Roberts, J. M.; Clurman, B. E.: Cyclin E–CDK2 is a regulator of

p27(Kip1). *Genes Dev.* 11:1464–1478, 1997.

[69214] 7544.Akoulitchev, S.; Chuikov, S.; Reinberg, D.: TFIIH is negatively regulated by cdk8-containing mediator complexes. *Nature* 407: 102–106, 2000.

[69215] 7545.Hanson, R. D.; Hohn, P. A.; Popescu, N. C.; Ley, T. J.: A cluster of hematopoietic serine protease genes is found on the same chromosomal band as the human alpha/delta T-cell receptor locus. *Proc. Nat. Acad.Sci.* 87: 960–963, 1990.

[69216] 7546.Harper, K.; Mattei, M.–G.; Simon, D.; Suzan, M.; Guenet, J.–L.; Haddad, P.; Sasportes, M.; Golstein, P.: Proximity of the CTLA–1 serine esterase and Tcr(alpha) loci in mouse and man. *Immunogenetics* 28:439–444, 1988.

[69217] 7547.Klein, J. L.; Shows, T. B.; Dupont, B.; Trapani, J. A.: Genomic organization and chromosomal assignment for a serine protease gene(CSP–B), expressed by human cytotoxic lymphocytes. *Genomics* 5: 110–117, 1989.

[69218] 7548.Lin, C. C.; Meier, M.; Sorensen, O.; Sasi, R.; Tainaka, T.; Bleackley, R. C.: Chromosome localization of two human serine protease genes to region 14q11.2–q12 by in situ hybridization. *Cytogenet. Cell Genet.* 53:169–171, 1990.

[69219] 7549.Motyka, B.; Korbitt, G.; Pinkoski, M. J.; Heibin, J. A.;

Caputo,A.; Hobman, M.; Barry, M.; Shostak, I.; Sawchuk, T.; Holmes, C. F.B.; Gauldie, J.; Bleackley, R. C.: Mannose 6-phosphate/insulin-like growth factor II receptor is a death receptor for granzyme B during cytotoxic T cell-induced apoptosis. *Cell* 103: 491–500, 2000.

[69220] 7550.Miles, J. S.; Spurr, N. K.; Gough, A. C.; Jowett, T.; McLaren,A. W.; Brook, J. D.; Wolf, C. R.: A novel human cytochrome P450 gene(P450IIB): chromosomal localization and evidence for alternative splicing. *NucleicAcids Res.* 16: 5783–5795, 1988.

[69221] 7551.Santisteban, I.; Povey, S.; Shephard, E. A.; Phillips, I. R.:The major phenobarbital-inducible cytochrome P-450 gene subfamily(P450IIB) mapped to the long arm of human chromosome 19. *Ann. Hum.Genet.* 52: 129–135, 1988.

[69222] 7552.Trask, B.; Fertitta, A.; Christensen, M.; Youngblom, J.; Bergmann,A.; Copeland, A.; de Jong, P.; Mohrenweiser, H.; Olsen, A.; Carrano,A.; Tynan, K.: Fluorescence in situ hybridization mapping of human chromosome 19: cytogenetic band location of 540 cosmids and 70 genes or DNA markers. *Genomics* 15: 133–145, 1993.

[69223] 7553.Barletta, C.; Batticane, N.; Ragusa, R. M.; Leube, R.; Peschle,C.; Romano, V.: Subchromosomal localization of

two human cyto kerat ingenes (KRT4 and KRT15) by in situ hybridization. Cytogenet. Cell Genet. 54: 148–150, 1990.

- [69224] 7554. Barletta, C.; Batticane, N.; Ragusa, R. M.; Leube, R. E.; Franke, W. W.; Peschle, C.; Romano, V.: Sub-chromosomal localization of human cyto keratin 4, 15, and 19 genes. (Abstract) Cytogenet. Cell Genet. 51:958 only, 1989.
- [69225] 7555. Romano, V.; Bosco, P.; Costa, G.; Leube, R.; Franke, W. W.; Rocchi, M.; Romeo, G.: Chromosomal assignment of cyto keratin genes. (Abstract) Cytogenet. Cell Genet. 46: 683 only, 1987.
- [69226] 7556. Romano, V.; Bosco, P.; Rocchi, M.; Costa, G.; Leube, R. E.; Franke, W. W.; Romeo, G.: Chromosomal assignments of human type I and type II cyto keratin genes to different chromosomes. Cytogenet. Cell Genet. 48:148–151, 1988.
- [69227] 7557. Schrijver, H. M.; Crusius, J. B. A.; Uitdehaag, B. M. J.; Garcia Gonzalez, M. A.; Kostense, P. J.; Polman, C. H.; Pena, A. S.: Association of interleukin–1–beta and interleukin–1 receptor antagonist genes with disease severity in MS. Neurology 52: 595–599, 1999.
- [69228] 7558. Terasaki, P. I.; Park, M. S.; Opelz, G.; Ting, A.: Multiple sclerosis and high incidence of a B-lymphocyte antigen. Science 193: 1245–1247, 1976.

- [69229] 7559.Feldman, R. D.; Hegele, R. A.: G-protein polymorphisms and maternal/neonatalmetabolism: still a weight for the answer. (Commentary) *Lancet* 355:1201–1202, 2000.
- [69230] 7560.Shitoh, K.; Konishi, F.; Iijima, T.; Ohdaira, T.; Sakai, K.; Kanazawa,K.; Miyaki, M.: A novel case of a sporadic desmoid tumour with mutationof the beta catenin gene. *J. Clin. Path.* 52: 695–696, 1999.
- [69231] 7561.Cartron, P.–F.; Oliver, L.; Martin, S.; Moreau, C.; LeCabellec,M.–T.; Jezequel, P.; Meflah, K.; Vallette, F. M.: The expressionof a new variant of the pro-apoptotic molecule Bax, Bax–psi, is correlatedwith an increased survival of glioblastoma multiforme patients. *Hum.Molec. Genet.* 11: 675–687, 2002.
- [69232] 7562.Holland, E. C.; Celestino, J.; Dai, C.; Schaefer, L.; Sawaya,R. E.; Fuller, G. N.: Combined activation of Ras and Akt in neuralprogenitors induces glioblastoma formation in mice. *Nature Genet.* 25:55–57, 2000.
- [69233] 7563.Mollenhauer, J.; Wiemann, S.; Scheurlen, W.; Korn, B.; Hayashi,Y.; Wilgenbus, K. K.; van Deimling, A.; Poustka, A.: DMBT1, a newmember of the SRCR superfamily, on chromosome 10q25.3–26.1 is deletedin malignant brain tumours. *Nature Genet.* 17: 32–39, 1997.

- [69234] 7564.Efstratiadis, A.; Posakony, J. W.; Maniatis, T.; Lawn, R. M.; O'Connell, C.; Spritz, R. A.; DeRiel, J. K.; Forget, B. G.; Weissman, S. M.; Slightom, J. L.; Blechl, A. E.; Smithies, O.; Baralle, F. E.; Shoulders, C. C.; Proudfoot, N. J.: The structure and evolution of the human beta-globin gene family. *Cell* 21: 653–668, 1980.
- [69235] 7565.Gutersohn, A.; Naber, C.; Muller, N.; Erbel, R.; Siffert, W.: G protein beta-3 subunit 825 TT genotype and post-pregnancy weight retention. *Lancet* 355: 1240–1241, 2000.
- [69236] 7566.Hegele, R. A.; Anderson, C.; Young, T. K.; Connelly, P. W.: G-protein beta-3 subunit gene splice variant and body fat distribution in Nunavut Inuit. *Genome Res.* 9: 972–977, 1999.
- [69237] 7567.Hocher, B.; Slowinski, T.; Stolze, T.; Pleschka, A.; Neumayer, H.-H.; Halle, H.: Association of maternal G protein beta-3 subunit 825T allele with low birthweight. *Lancet* 355: 1241–1242, 2000.
- [69238] 7568.Jin, Y.; Dietz, H. C.; Montgomery, R. A.; Bell, W. R.; McIntosh, I.; Coller, B.; Bray, P. F.: Glanzmann thrombasthenia: cooperation between sequence variants in C1s during splice site selection. *J. Clin. Invest.* 98: 1745–1754, 1996.



- [69239] 7569. Levine, M. A.; Modi, W. S.; O'Brien, S. J.: Chromosomal localization of the genes encoding two forms of the G-protein beta polypeptide, beta-1 and beta-3, in man. *Genomics* 8: 380-386, 1990.
- [69240] 7570. Levine, M. A.; Smallwood, P. M.; Moen, P. T., Jr.; Helman, L. J.; Ahn, T. G.: Molecular cloning of beta-3 subunit, a third form of the G protein beta-subunit polypeptide. *Proc. Nat. Acad. Sci.* 87:2329-2333, 1990.
- [69241] 7571. Liu, W.; Qian, C.; Francke, U.: Silent mutation induces exon skipping of fibrillin-1 gene in Marfan syndrome. (Letter) *Nature Genet.* 16:328-329, 1997.
- [69242] 7572. Modi, W. S.; Levine, M. A.; Seuanez, H.; O'Brien, S. J.: Chromosomal localization of the gene encoding a third form of the beta subunit of GTP-binding regulatory proteins. (Abstract) *Cytogenet. Cell Genet.* 51:1046 only, 1989.
- [69243] 7573. Pietruck, F.; Moritz, A.; Montemurro, M.; Sell, A.; Busch, S.; Roskopf, D.; Virchow, S.; Esche, H.; Brockmeyer, N.; Jakobs, K. H.; Siffert, W.: Selectively enhanced cellular signaling by G(i) proteins in essential hypertension: G-alpha(i2), G-alpha(i3), G-beta(2) are not mutated. *Circ. Res.* 79: 974-983, 1996.
- [69244] 7574. Siffert, W.; Forster, P.; Jockel, K.-H.; Mvere, D. A.;

Brinkmann,B.; Naber, C.; Crookes, R.; Du P. Heyns, A.; Eppelen, J. T.; Fridley,J.; Freedman, B. I.; Muller, N.; and 15 others: Worldwide ethnicdistribution of the G protein beta-3 subunit 825T allele and its associationwith obesity in Caucasian, Chinese, and black African individuals. J.Am. Soc. Nephrol. 10: 1921-1030, 1999.

[69245] 7575.Siffert, W.; Rosskopf, D.; Moritz, A.; Wieland, T.; Kaldenberg-Stasch,S.; Kettler, N.; Hartung, K.; Beckmann, S.; Jakobs, K. H.: EnhancedG protein activation in immortalized lymphoblasts from patients withessential hypertension. J. Clin. Invest. 96: 759-766, 1995.

[69246] 7576.Siffert, W.; Rosskopf, D.; Siffert, G.; Busch, S.; Moritz, A.;Erbel, R.; Sharma, A. M.; Ritz, E.; Wichmann, H.-E.; Jakobs, K. H.;Horsthemke, B.: Association of a human G-protein beta-3 subunit variantwith hypertension. Nature Genet. 18: 45-48, 1998.

[69247] 7577.Stallings-Mann, M. L.; Ludwiczak, R. L.; Klinger, K. W.; Rottman,F.: Alternative splicing of exon 3 of the human growth hormone receptoris the result of an unusual genetic polymorphism. Proc. Nat. Acad.Sci. 93: 12394-12399, 1996.

[69248] 7578.Chang, C.; Kokontis, J.; Liao, S. S.; Chang, Y.: Isolation andcharacterization of human TR3 receptor: a mem-

ber of steroid receptorsuperfamily. J. Steroid Biochem. 34: 391–395, 1989.

[69249] 7579.DeGroot, L. J.: Personal Communication. Chicago, Ill. 12/19/1991.

[69250] 7580.Forman, B. M.; Umesono, K.; Chen, J.; Evans, R. M.: Unique responsepathways are established by allosteric interactions among nuclearhormone receptors. Cell 81: 541–550, 1995.

[69251] 7581.Labelle, Y.; Bussieres, J.; Courjal, F.; Goldring, M. B.: TheEWS/TEC fusion protein encoded by the t(9;22) chromosomal translocationin human chondrosarcomas is a highly potent transcriptional activator. Oncogene 18:3303–3308, 1999.

[69252] 7582.Li, H.; Kolluri, S. K.; Gu, J.; Dawson, M. I.; Cao, X.; Hobbs,P. D; Lin, B.; Chen, G.; Lu, J.; Lin, F.; Xie, Z.; Fontana, J. A.;Reed, J. C.; Zhang, X.: Cytochrome c release and apoptosis inducedby mitochondrial targeting of orphan receptor TR3 nuclear. Science 289:1159–1164, 2000.

[69253] 7583.Nakai, A.; Kartha, S.; Sakurai, A.; Toback, F. G.; DeGroot, L.J.: A human early response gene homologous to murine nur77 and ratNGFI-B, and related to the nuclear receptor superfamily. Molec. Endocr. 4: 1438–1443, 1990.

- [69254] 7584. Perlmann, T.; Jansson, L.: A novel pathway for vitamin A signaling mediated by RXR heterodimerization with NGFI-B and NURR1. *Genes Dev.* 9:769–782, 1995.
- [69255] 7585. Ryseck, R.-P.; Macdonald-Bravo, H.; Mattei, M. G.; Siegfried, R.L.; Bravo, R.: Structure, mapping and expression of a growth factor inducible gene encoding a putative nuclear hormonal binding receptor. *EMBO J.* 8: 3327–3335, 1989.
- [69256] 7586. Wu, Q.; Dawson, M. I.; Zheng, Y.; Hobbs, P. D.; Agadir, A.; Jong, L.; Li, Y.; Liu, R.; Lin, B.; Zhang, X. K.: Inhibition of trans-retinoic acid-resistant human breast cancer cell growth by retinoid X receptor-selective retinoids. *Molec. Cell. Biol.* 17: 6598–6608, 1997.
- [69257] 7587. Youn, H.-D.; Sun, L.; Prywes, R.; Liu, J. O.: Apoptosis of T cells mediated by  $Ca^{2+}$ -induced release of the transcription factor MEF2. *Science* 286: 790–793, 1999.
- [69258] 7588. Friedman, E.; Gejman, P. V.; Martin, G. A.; McCormick, F.: Nonsense mutations in the C-terminal SH2 region of the GTPase activating protein (GAP) gene in human tumours. *Nature Genet.* 5: 242–247, 1993.
- [69259] 7589. Hsieh, C. L.; Francke, U.: The gene for GTPase activating protein (GAP) is on human chromosome 5q and mouse chromosome 13. (Abstract) *Cytogenet. Cell Genet.*

51: 1016 only, 1989.

- [69260] 7590.Hsieh, C. L.; Vogel, U. S.; Dixon, R. A.; Francke, U.: Chromosomelocalization and cDNA sequence of murine and human genes for ras p21GTPase activating protein (GAP). *Somat. Cell Molec. Genet.* 15: 579–590,1989.
- [69261] 7591.Lemons, R. S.; Espinosa, R., III; Rebentisch, M.; McCormick, F.;Ladner, M.; Le Beau, M. M.: Chromosomal localization of the geneencoding GTPase-activating protein (RASA) to human chromosome 5, bandsq13–q15. *Genomics* 6: 383–385, 1990.
- [69262] 7592.Mitsudomi, T.; Friedman, E.; Gejman, P. V.; McCormick, F.; Gazdar,A. F.: Genetic analysis of the catalytic domain of the GAP gene inhuman lung cancer cell lines. *Hum. Genet.* 93: 27–31, 1994.
- [69263] 7593.Trahey, M.; Wong, G.; Halenbeck, R.; Rubinfeld, B.; Martin, G.A.; Ladner, M.; Long, C. M.; Crosier, W. J.; Watt, K.; Koths, K.;McCormick, F.: Molecular cloning of two types of GAP complementaryDNA from human placenta. *Science* 242: 1697–1700, 1988.
- [69264] 7594.Blatt, C.; Eversole–Cire, P.; Cohn, V. H.; Zollman, S.; Fournier,R. E. K.; Mohandas, L. T.; Nesbitt, M.; Lugo, T.; Jones, D. T.; Reed,R. R.; Weiner, L. P.; Sparkes, R. S.; Simon, M. I.: Chromosomal localizationof genes encoding

guanine nucleotide-binding protein subunits in mouse and human. *Proc. Nat. Acad. Sci.* 85: 7642–7646, 1988.

- [69265] 7595. Allen, B.; Ostrer, H.; Stein, J.; Stein, G.: Histone gene clusters map to chromosomes 1 and 6. (Abstract) *Cytogenet. Cell Genet.* 51:950 only, 1989.
- [69266] 7596. Carozzi, N.; Marashi, F.; Plumb, M.; Zimmerman, S.; Zimmerman, A.; Coles, L. S.; Wells, J. R. E.; Stein, G.; Stein, J.: Clustering of human H1 and core histone genes. *Science* 224: 1115–1117, 1984.
- [69267] 7597. Chandler, M. E.; Kedes, L. H.; Cohn, R. H.; Yunis, J. J.: Genes coding for histone proteins in man are located on the distal end of chromosome 7. *Science* 205: 908–910, 1979.
- [69268] 7598. Clark, S. J.; Krieg, P. A.; Wells, J. R. E.: Isolation of a clone containing human histone genes. *Nucleic Acids Res.* 9: 1583–1597, 1981.
- [69269] 7599. Delange, R. J.; Smith, E. L.: Histones: structure and function. *Ann. Rev. Biochem.* 40: 279–314, 1971.
- [69270] 7600. Felsenfeld, G.: Chromatin as an essential part of the transcriptional mechanism. *Nature* 355: 219–224, 1992.
- [69271] 7601. Green, L.; Van Antwerpen, R.; Stein, J.; Stein, G.; Tripputi, P.; Emanuel, B.; Selden, J.; Croce, C.: A major human histone gene cluster on the long arm of chromosome

1. Science 226: 838–840, 1984.

[69272] 7602.Heintz, N.; Zernik, M.; Roeder, R. G.: The structure of the human histone genes: clustered but not tandemly repeated. Cell 24: 661–668, 1981.

[69273] 7603.Hentschel, C. C.; Birnstiel, M. L.: The organization and expression of histone gene families. Cell 25: 301–313, 1981.

[69274] 7604.Kedes, L.; Maxson, R.: Histone gene organization: paradigm lost. Nature 294:11–12, 1981.

[69275] 7605.Lichtler, A. C.; Sierra, F.; Clark, S.; Wells, J. R. E.; Stein, J. L.; Stein, G. S.: Multiple H4 histone mRNAs of HeLa cells are encoded in different genes. Nature 298: 195–198, 1982.

[69276] 7606.McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba, Canada 1989.

[69277] 7607.Sierra, F.; Lichtler, A.; Marashi, F.; Rickles, R.; Van Dyke, T.; Clark, S.; Wells, J.; Stein, G.; Stein, J.: Organization of human histone genes. Proc. Nat. Acad. Sci. 79: 1795–1799, 1982.

[69278] 7608.Steffensen, D. M.: Human histone genes mapped to chromosome 7.(Abstract) Cytogenet. Cell Genet. 25: 211 only, 1979.

[69279] 7609.Szabo, P.; Yu, L. C.; Borun, T.; Varicchio, F.; Sinis-

calco, M.;Prensky, W.: Localization of the histone genes in man. Cytogenet.Cell Genet. 22: 359–363, 1978.

[69280] 7610.Tripputi, P.; Emanuel, B. S.; Croce, C. M.; Green, L. G.; Stein,G. S.; Stein, J. L.: Human histone genes map to multiple chromosomes. Proc.Nat. Acad. Sci. 83: 3185–3188, 1986.

[69281] 7611.Yunis, J. J.; Chandler, M. E.: Localization of histone genes to bands 7q32–36 in man and the homologous chromosome segments in chimpanzee, gorilla, and orangutan. (Abstract) Cytogenet. Cell Genet. 25:220 only, 1979.

[69282] 7612.Doyle, J.; Hoffman, S.; Ucla, C.; Reith, W.; Mach, B.; Stubbs,L.: Locations of human and mouse genes encoding the RFX1 and RFX2transcription factor proteins. Genomics 35: 227–230, 1996.

[69283] 7613.Pugliatti, L.; Derre, J.; Berger, R.; Ucla, C.; Reith, W.; Mach,B.: The genes for MHC class II regulatory factors RFX1 and RFX2 arelocated on the short arm of chromosome 19. Genomics 13: 1307–1310,1992.

[69284] 7614.Reith, W.; Ucla, C.; Barras, E.; Gaud, A.; Durand, B.; Herrero–Sanchez,C.; Kobr, M.; Mach, B.: RFX1, a transactivator of hepatitis B virusenhancer I, belongs to a novel family of homodimeric and heterodimericDNA–binding



proteins. *Molec. Cell. Biol.* 14: 1230–1244, 1994.

[69285] 7615.Claesson, L.; Larhammar, D.; Rask, L.; Peterson, P. A.: cDNA clone for the human invariant gamma chain of class II histocompatibility antigens and its implications for the protein structure. *Proc. Nat. Acad. Sci.* 80: 7395–7399, 1983.

[69286] 7616.Claesson–Welsh, L.; Barker, P. E.; Larhammar, D.; Rask, L.; Ruddle, F. H.; Peterson, P. A.: The gene encoding the human class II antigen-associated gamma chain is located on chromosome 5. *Immunogenetics* 20: 89–93, 1984.

[69287] 7617.Driessen, C.; Bryant, R. A.; Lennon–Dumenil, A. M.; Villadangos, J. A.; Bryant, P. W.; Shi, G. P.; Chapman, H. A.; Ploegh, H. L.: Cathepsin S controls the trafficking and maturation of MHC class II molecules in dendritic cells. *J. Cell Biol.* 147: 775–790, 1999.

[69288] 7618.Genuardi, M.; Saunders, G. F.: Localization of the HLA class II-associated invariant chain gene to human chromosome band 5q32. *Immunogenetics* 28:53–56, 1988.

[69289] 7619.Bergsma, D. J.; Eder, C.; Gross, M.; Kersten, H.; Sylvester, D.; Appelbaum, E.; Cusimano, D.; Livi, G. P.; McLaughlin, M. M.; Kasyan, K.; Porter, T. G.; Silverman, C.;

Dunnington, D.; Hand, A.; Prichett, W. P.; Bossard, M. J.; Brandt, M.; Levy, M. A.: The cyclophilin multigene family of peptidyl-prolyl isomerases: characterization of three separate human isoforms. *J. Biol. Chem.* 266: 23204–23214, 1991.

[69290] 7620. Bowles, K. R.; Zintz, C.; Abraham, S. E.; Brandon, L.; Bowles, N. E.; Towbin, J. A.: Genomic characterization of the human peptidyl-prolyl-cis-trans-isomerase, mitochondrial precursor gene: assessment of its role in familial dilated cardiomyopathy. *Hum. Genet.* 105: 582–586, 1999.

[69291] 7621. Bachmaier, K.; Krawczyk, C.; Kozieradzki, I.; Kong, Y.-Y.; Sasaki, T.; Oliveira-dos-Santos, A.; Mariathasan, S.; Bouchard, D.; Wakeham, A.; Itie, A.; Le, J.; Ohashi, P. S.; Sarosi, I.; Nishina, H.; Lipkowitz, S.; Penninger, J. M.: Negative regulation of lymphocyte activation and autoimmunity by the molecular adaptor Cbl-b. *Nature* 403: 211–216, 2000.

[69292] 7622. Chiang, Y. J.; Kole, H. K.; Brown, K.; Naramura, M.; Fukuhara, S.; Hu, R.-J.; Jang, I. K.; Gutkind, J. S.; Shevach, E.; Gu, H.: Cbl-b regulates the CD28 dependence of T-cell activation. *Nature* 403: 216–220, 2000.

[69293] 7623. Keane, M. M.; Rivero-Lezcano, O. M.; Mitchell, J. A.;

Robbins, K. C.; Lipkowitz, S.: Cloning and characterization of cbl-b: a SH3binding protein with homology to the c-cbl proto-oncogene. *Oncogene* 10:2367–2377, 1995.

[69294] 7624. Komeda, K.; et al; et al: Establishment of two substrains, diabetes-prone and nondiabetic, from Long-Evans Tokushima Lean (LETL) rats. *Endocr.J.* 45: 737–744, 1998.

[69295] 7625. Yokoi, N.; Komeda, K.; Wang, H.-Y.; Yano, H.; Kitada, K.; Saitoh, Y.; Seino, Y.; Yasuda, K.; Serikawa, T.; Seino, S.: Cblb is a major susceptibility gene for rat type 1 diabetes mellitus. *Nature Genet.* 31:391–394, 2002.

[69296] 7626. Bathori, G.; Parolini, I.; Tombola, F.; Szabo, I.; Messina, A.; Oliva, M.; De Pinto, V.; Lisanti, M.; Sargiacomo, M.; Zoratti, M.: Porin is present in the plasma membrane where it is concentrated in caveolae and caveolae-related domains. *J. Biol. Chem.* 274: 29607–29612, 1999.

[69297] 7627. Huizing, M.; Ruitenbeek, W.; Thinnen, F. P.; DePinto, V.; Wendel, U.; Trijbels, F. J. M.; Smit, L. M. E.; Ter Laak, H. J.; Van Den Heuvel, L. P.: Deficiency of the voltage-dependent anion channel: a novel cause of mitochondriopathy. *Pediatr. Res.* 39: 760–765, 1996.

[69298] 7628. Chang, P. W. G.; Tsui, S. K. W.; Liew, C.; Lee, C.; Waye, M. M. Y.; Fung, K.: Isolation, characterization, and

chromosomal mapping of a novel cDNA clone encoding human selenium binding protein. *J. Cell. Biochem.* 64: 217–224, 1997.

- [69299] 7629. Lanfear, J.; Fleming, J.; Walker, M.; Harrison, P.: Different patterns of regulation of the genes encoding the closely related 56kDa selenium- and acetaminophen-binding proteins in normal tissues and during carcinogenesis. *Carcinogenesis* 14: 335–340, 1993.
- [69300] 7630. Creutz, C. E.; Tomsig, J. L.; Snyder, S. L.; Gautier, M.-C.; Skouri, F.; Beisson, J.; Cohen, J.: The copines, a novel class of C2 domain-containing, calcium-dependent, phospholipid-binding proteins conserved from *Paramecium* to humans. *J. Biol. Chem.* 273: 1393–1402, 1998.
- [69301] 7631. Caudell, E. G.; Caudell, J. J.; Tang, C.-H.; Yu, T.-K.; Frederick, M. J.; Grimm, E. A.: Characterization of human copine III as a phosphoprotein with associated kinase activity. *Biochemistry* 39: 13034–13043, 2000.
- [69302] 7632. Ito, M.; Yuan, C.-X.; Okano, H. J.; Darnell, R. B.; Roeder, R. G.: Involvement of the TRAP220 component of the TRAP/SMCC coactivator complex in embryonic development and thyroid hormone action. *Molec. Cell* 5: 683–693, 2000.
- [69303] 7633. Yuan, C.-X.; Ito, M.; Fondell, J. D.; Fu, Z.-Y.; Roeder,

R. G.: The TRAP220 component of a thyroid hormone receptor-associated protein (TRAP) coactivator complex interacts directly with nuclear receptors in a ligand-dependent fashion. *Proc. Nat. Acad. Sci.* 95:

7939–7944, 1998. Note: Erratum: *Proc. Nat. Acad. Sci.* 95: 14584 only, 1998.

[69304] 7634. Zhu, Y.; Qi, C.; Jain, S.; Le Beau, M. M.; Espinosa, R., III; Atkins, G. B.; Lazar, M. A.; Yeldandi, A. V.; Rao, M. S.; Reddy, J. K.: Amplification and overexpression of peroxisome proliferator-activated receptor binding protein (PBP/PPARBP) gene in breast cancer. *Proc. Nat. Acad. Sci.* 96:10848–10853, 1999.

[69305] 7635. Zhu, Y.; Qi, C.; Jain, S.; Rao, M. S.; Reddy, J. K.: Isolation and characterization of PBP, a protein that interacts with peroxisome proliferator-activated receptor. *J. Biol. Chem.* 272: 25500–25506, 1997.

[69306] 7636. Abrahamson, M.; Olafsson, I.; Palsdottir, A.; Ulvsback, M.; Lundwall, A.; Jensson, O.; Grubb, A.: Structure and expression of the human cystatin C gene. *Biochem. J.* 268: 287–294, 1990.

[69307] 7637. Balbin, M.; Abrahamson, M.: SstII polymorphic sites in the promoter region of the human cystatin C gene. *Hum. Genet.* 87: 751–752, 1991.

- [69308] 7638.Barrett, A. J.; Davies, M. E.; Grubb, A.: The place of human gamma-trace(cystatin C) amongst the cysteine proteinase inhibitors. *Biochem.Biophys. Res. Commun.* 120: 631-636, 1984.
- [69309] 7639.Gopal Rao, V. V.; Schnittger, S.; Abrahamson, M.; Hansmann, I.: Cystatin-C (CST3), the candidate gene for the hereditary cystatin-Camyloid angiopathy (HCCAA) maps to or close to human chromosome 20p11.22.(Abstract) *Cytogenet. Cell Genet.* 58: 2029, 1991.
- [69310] 7640.Grubb, A.; Lofberg, H.: Human gamma-trace, a basic microprotein:amino acid sequence and presence in the adenohypophysis. *Proc. Nat.Acad. Sci.* 79: 3024-3027, 1982.

- [69311] 7641.Huh, C.; Nagle, J. W.; Kozak, C. A.; Abrahamson, M.; Karlsson,S.: Structural organization, expression and chromosomal mapping of the mouse cystatin-C-encoding gene (Cst3). *Gene* 152: 221–226, 1995.
- [69312] 7642.Lofberg, H.; Grubb, A.; Davidsson, L.; Kjellander, B.; Stromblad,L.-G.; Tibblin, S.; Olsson, S.-O.: Occurrence of gamma-trace in the calcitonin-producing C-cells of simian thyroid gland. *Acta Endocr.* 104:69–76, 1983.
- [69313] 7643.Lofberg, H.; Grubb, A. O.: Quantitation of gamma-trace in human biological fluids: indications for production in the central nervous system. *Scand. J. Clin. Lab. Invest.* 39: 619–626, 1979.
- [69314] 7644.Galarneau, L.; Drouin, R.; Belanger, L.: Assignment of the fetoprotein transcription factor gene (FTF) to human chromosome band 1q32.11 by in situ hybridization. *Cytogenet. Cell Genet.* 82: 269–270, 1998.
- [69315] 7645.Li, M.; Xie, Y.-H.; Kong,Y.-Y.; Wu, X.; Zhu, L.; Wang, Y.: Cloning and characterization of a novel human hepatocyte transcription factor, hB1F, which binds and activates enhancer II of hepatitis B virus. *J.Biol. Chem.* 273: 29022–29031, 1998.
- [69316] 7646.Kadereit, S.; Gewert, D. R.; Galabru, J.; Hovanessian, A. G.; Meurs,E. F.: Molecular cloning of two new inter-

feron-induced, highly related nuclear phosphoproteins. J. Biol. Chem. 268: 24432–24441, 1993.

[69317] 7647. Welsh, G. I.; Kadereit, S.; Coccia, E. M.; Hovanessian, A. G.; Meurs, E. F.: Colocalization within the nucleolus of two highly related IFN-induced human nuclear phosphoproteins with nucleolin. Exp. Cell Res. 250: 62–74, 1999.

[69318] 7648. Pietu, G.; Alibert, O.; Guichard, V.; Lamy, B.; Bois, F.; Leroy, E.; Mariage-Samson, R.; Houlgatte, R.; Soularue, P.; Auffray, C.: Novel gene transcripts preferentially expressed in human muscles revealed by quantitative hybridization of a high density cDNA array. Genome Res. 6: 492–503, 1996.

[69319] 7649. Xia, H.; Winokur, S. T.; Kuo, W.-L.; Altherr, M. R.; Bredt, D. S.: Actinin-associated LIM protein: identification of a domain interaction between PDZ and spectrin-like repeat motifs. J. Cell Biol. 139: 507–515, 1997.

[69320] 7650. Carim, L.; Sumoy, L.; Andreu, N.; Estivill, X.; Escarceller, M.: Identification and expression analysis of C15orf3, a novel gene on chromosome 15q21.1–q21.2. Cytogenet. Cell Genet. 88: 330–332, 2000.

[69321] 7651. MacLellan, W. R.; Xiao, G.; Abdellatif, M.; Schneider, M. D.: A novel Rb- and p300-binding protein inhibits transactivation by MyoD. Molec. Cell. Biol. 20: 8903–8915,



2000.

- [69322] 7652.Miyake, S.; Sellers, W. R.; Safran, M.; Li, X.; Zhao, W.; Grossman,S. R.; Gan, J.; DeCaprio, J. A.; Adams, P. D.; Kaelin, W. G., Jr.: Cells degrade a novel inhibitor of differentiation with E1A-likeproperties upon exiting the cell cycle. *Molec. Cell. Biol.* 20: 8889–8902,2000.
- [69323] 7653.Schweizer, A.; Valdenaire, O.; Koster, A.; Lang, Y.; Schmitt, G.;Lenz, B.; Bluethmann, H.; Rohrer, J.: Neonatal lethality in micedeficient in XCE, a novel member of the endothelin–converting enzymeand neutral endopeptidase family. *J. Biol. Chem.* 274: 20450–20456,1999.
- [69324] 7654.Valdenaire, O.; Richards, J. G.; Faull, R. L. M.; Schweizer, A.: XCE, a new member of the endothelin–converting enzyme and neutralendopeptidase family, is preferentially expressed in the CNS. *Molec.Brain Res.* 64: 211–221, 1999.
- [69325] 7655.Valdenaire, O.; Rohrbacher, E.; Langeveld, A.; Schweizer, A.; Meijers,C.: Organization and chromosomal localization of the human ECEL1(XCE) gene encoding a zinc metallopeptidase involved in the nervouscontrol of respiration. *Biochem. J.* 346: 611–616, 2000.
- [69326] 7656.Hengst, U.; Albrecht, H.; Hess, D.; Monard, D.: The phosphatidylethanolamine–bindingprotein is the proto–

type of a novel family of serine protease inhibitors. *J. Biol. Chem.* 276: 535–540, 2001.

- [69327] 7657. Hori, N.; Chae, K.; Murakawa, K.; Matoba, R.; Fukushima, A.; Okubo, K.; Matsubara, K.: A human cDNA sequence homologue of bovine phosphatidylethanolamine-binding protein. *Gene* 140: 293–294, 1994.
- [69328] 7658. Moore, C.; Perry, A. C. F.; Love, S.; Hall, L.: Sequence analysis and immunolocalisation of phosphatidylethanolamine binding protein (PBP) in human brain tissue. *Molec. Brain Res.* 37: 74–78, 1996.
- [69329] 7659. Schoentgen, F.; Jolles, P.: From structure to function: possible biological roles of a new widespread protein family binding hydrophobic ligands and displaying a nucleotide binding site. *FEBS Lett.* 369: 22–26, 1995.
- [69330] 7660. Seddiqi, N.; Bollengier, F.; Alliel, P. M.; Perin, J.-P.; Bonnet, F.; Bucquoy, S.; Jolles, P.; Schoentgen, F.: Amino acid sequence of the Homo sapiens brain 21–23-kDa protein (neuropolypeptide h3), comparison with its counterparts from Rattus norvegicus and Bos taurus species, and expression of its mRNA in different tissues. *J. Molec. Evol.* 39: 655–660, 1994.
- [69331] 7661. Tohdoh, N.; Tojo, S.; Agui, H.; Ojika, K.: Sequence

homology of rat and human HCNP precursor proteins, bovine phosphatidylethanolamine-binding protein and rat 23-kDa protein associated with the opioid-binding protein. *Molec. Brain Res.* 30: 381–384, 1995.

- [69332] 7662. Yeung, K.; Seitz, T.; Li, S.; Janosch, P.; McFerran, B.; Kaiser, C.; Fee, F.; Katsanakis, K. D.; Rose, D. W.; Mischak, H.; Sedivy, J. M.; Kolch, W.: Suppression of Raf-1 kinase activity and MAP kinase signalling by RKIP. *Nature* 401: 173–177, 1999.
- [69333] 7663. Heinemann, T.; Bulwin, G. C.; Randall, J.; Schnieders, B.; Sandhoff, K.; Volk, H. D.; Milford, E.; Gullans, S. R.; Utku, N.: Genomic organization of the gene coding for TIRC7, a novel membrane protein essential for T cell activation. *Genomics* 57: 398–406, 1999.
- [69334] 7664. Li, Y.-P.; Chen, W.; Liang, Y.; Li, E.; Stashenko, P.: ATP6i-deficient mice exhibit severe osteopetrosis due to loss of osteoclast-mediated extracellular acidification. *Nature Genet.* 23: 447–451, 1999.
- [69335] 7665. Li, Y.-P.; Chen, W.; Stashenko, P.: Molecular cloning and characterization of a putative novel human osteoclast-specific 116-kDa vacuolar proton pump subunit. *Biochem. Biophys. Res. Commun.* 218: 813–821, 1996.
- [69336] 7666. Scimeca, J.-C.; Franchi, A.; Trojani, C.; Parrinello, H.;

Grosgeorge, J.; Robert, C.; Jaillon, O.; Poirier, C.; Gaudray, P.; Carle, G. F.: The gene encoding the mouse homologue of the human osteoclast-specific 116-kDa V-ATPase subunit bears a deletion in osteosclerotic (oc/oc) mutants. *Bone* 26: 207–213, 2000.

- [69337] 7667. Sobacchi, C.; Frattini, A.; Orchard, P.; Porras, O.; Tezcan, I.; Andolina, M.; Babul-Hirji, R.; Baric, I.; Canham, N.; Chitayat, D.; Dupuis-Girod, S.; Ellis, I.; and 21 others: The mutational spectrum of human malignant autosomal recessive osteopetrosis. *Hum. Molec. Genet.* 10: 1767–1773, 2001.
- [69338] 7668. Utku, N.; Heinemann, T.; Tullius, S. G.; Bulwin, G. C.; Beinke, S.; Blumberg, R. S.; Beato, F.; Randall, J.; Kojima, R.; Busconi, L.; Robertson, E. S.; Schulein, R.; Volk, H. D.; Milford, E. L.; Gullans, S. R.: Prevention of acute allograft rejection by antibody targeting of TIRC7, a novel T cell membrane protein. *Immunity* 9: 509–518, 1998.
- [69339] 7669. Blum, R.; Feick, P.; Puype, M.; Vandekerckhove, J.; Klengel, R.; Nastainczyk, W.; Schulz, I.: Tmp21 and p24A, two type I proteins enriched in pancreatic microsomal membranes, are members of a protein family involved in vesicular trafficking. *J. Biol. Chem.* 271: 17183–17189, 1996.

- [69340] 7670.Horer, J.; Blum, R.; Feick, P.; Nastainczyk, W.; Schulz, I.: Acomparative study of rat and human Tmp21 (p23) reveals the pseudogene-likefeatures of human Tmp21-II. DNA Seq. 10: 121-126, 1999.
- [69341] 7671.Sherrington, R.; Rogaev, E. I.; Liang, Y.; Rogaeva, E. A.; Levesque,G.; Ikeda, M.; Chi, H.; Lin, C.; Li, G.; Holman, K.; Tsuda, T.; Mar,L.; Foncin, J.-F.; Bruni, A. C.; Montesi, M. P.; Sorbi, S.; Rainero,I.; Pinessi, L.; Nee, L.; Chumakov, I.; Pollen, D.; Brookes, A.; Sanseau,P.; Polinsky, R. J.; Wasco, W.; Da Silva, H. A. R.; Haines, J. L.;Pericak-Vance, M. A.; Tanzi, R. E.; Roses, A. D.; Fraser, P. E.; Rommens,J. M.; St George-Hyslop, P. H.: Cloning of a gene bearing missensemutations in early-onset familial Alzheimer's disease. Nature 375:754-760, 1995.
- [69342] 7672.Bohne, J.; Cole, S. E.; Sune, C.; Lindman, B. R.; Ko, V. D.; Vogt,T. F.; Garcia-Blanco, M. A.: Expression analysis and mapping of themouse and human transcriptional regulator CA150. Mammalian Genome 11:930-933, 2000.
- [69343] 7673.Sune, C.; Garcia-Blanco, M. A.: Transcriptional co-factor CA150regulates RNA polymerase II elongation in a TATA-box-dependent manner. Molec.Cell. Biol. 19: 4719-4728, 1999.
- [69344] 7674.Sune, C.; Garcia-Blanco, M. A.: Transcriptional trans

activation by human immunodeficiency virus type 1 Tat requires specific coactivators that are not basal factors. *J. Virol.* 69: 3098–3107, 1995.

[69345] 7675. Sune, C.; Hayashi, T.; Liu, Y.; Lane, W. S.; Young, R. A.; Garcia-Blanco, M. A.: CA150, a nuclear protein associated with the RNA polymerase II holoenzyme, is involved in Tat-activated human immunodeficiency virus type 1 transcription. *Molec. Cell. Biol.* 17: 6029–6039, 1997.

[69346] 7676. Postma, A. V.; Bezzina, C. R.; de Vries, J. F.; Wilde, A. A. M.; Moorman, A. F. M.; Mannens, M. M. A. M.: Genomic organisation and chromosomal localisation of two members of the KCND ion channel family, KCND2 and KCND3. *Hum. Genet.* 106: 614–619, 2000.

[69347] 7677. Zhu, X.-R.; Wulf, A.; Schwarz, M.; Isbrandt, D.; Pongs, O.: Characterization of human Kv4.2 mediating a rapidly-inactivating transient voltage-sensitive K<sup>+</sup> current. *Receptors Channels* 6: 387–400, 1999.

[69348] 7678. Dilks, D.; Ling, H.-P.; Cockett, M.; Sokol, P.; Numan, R.: Cloning and expression of the human Kv4.3 potassium channel. *J. Neurophysiol.* 81: 1974–1977, 1999.

[69349] 7679. Dixon, J. E.; Shi, W.; Wang, H.-S.; McDonald, C.; Yu, H.; Wymore, R. S.; Cohen, I. S.; McKinnon, D.: Role of the Kv4.3 K<sup>+</sup> channel in ventricular muscle: a molecular corre-

late for the transient outward current. *Circ. Res.* 79: 659–668, 1996.

- [69350] 7680. Kong, W.; Po, S.; Yamagishi, T.; Ashen, M. D.; Stetten, G.; Tomaselli, G. F.: Isolation and characterization of the human gene encoding *Isl*(to): further diversity by alternative mRNA splicing. *Am. J. Physiol.* 275:H1963–H1970, 1998.
- [69351] 7681. Tseng, G.-N.: Molecular structure of cardiac *Isl*(to) channels: *Kv4.2*, *Kv4.3*, and other possibilities? *Cardiovasc. Res.* 41: 16–18, 1999.
- [69352] 7682. Zhu, X. R.; Wulf, A.; Schwarz, M.; Isbrandt, D.; Pongs, O.: Characterization of human *Kv4.2* mediating a rapidly-inactivating transient voltage-sensitive  $K^+$  current. *Receptors Channels* 6: 387–400, 1999.
- [69353] 7683. van den Eynde, B. J.; Gaugler, B.; Probst-Kepper, M.; Michaux, L.; Devuyst, O.; Lorge, F.; Weynants, P.; Boon, T.: A new antigen recognized by cytolytic T lymphocytes on a human kidney tumor results from reverse strand transcription. *J. Exp. Med.* 190: 1793–1799, 1999.
- [69354] 7684. Lai, C.-H.; Chou, C.-Y.; Ch'ang, L.-Y.; Liu, C.-S.; Lin, W.: Identification of novel human genes evolutionarily conserved in *Caenorhabditis elegans* by comparative proteomics. *Genome Res.* 10: 703–713, 2000.

- [69355] 7685.Shamsadin, R.; Adham, I. M.; von Beust, G.; Engel, W.: Molecularcloning, expression and chromosome location of the human pelota genePELO. Cytogenet. Cell Genet. 90: 75–78, 2000.
- [69356] 7686.Kile, B. T.; Viney, E. M.; Willson, T. A.; Brodnicki, T. C.; Cancilla,M. R.; Herlihy, A. S.; Croker, B. A.; Baca, M.; Nicola, N. A.; Hilton,D. J.; Alexander, W. S.: Cloning and characterization of the genesencoding the ankyrin repeat and SOCS box-containing proteins Asb-1,Asb-2, Asb-3, and Asb-4. Gene 258: 31–41, 2000.
- [69357] 7687.Gaugler, B.; Brouwenstijn, N.; Vantomme, V.; Szikora, J.-P.; Vander Spek, C. W.; Patard, J.-J.; Boon, T.; Schrier, P.; Van den Eynde,B. J.: A new gene coding for an antigen recognized by autologouscytolytic T lymphocytes on a human renal carcinoma. Immunogenetics 44:323–330, 1996.
- [69358] 7688.Miyata, Y.; Akashi, M.; Nishida, E.: Molecular cloning and characterizationof a novel member of the MAP kinase superfamily. Genes Cells 4:299–309, 1999.
- [69359] 7689.Liu, Y.; Corcoran, M.; Rasool, O.; Ivanova, G.; Ibbotson, R.; Grander,D.; Iyengar, A.; Baranova, A.; Kashuba, V.; Merup, M.; Wu, X.; Gardiner,A.; and 12 others: Cloning of two candidate tumor suppressor geneswithin a 10 kb re-



gion on chromosome 13q14, frequently deleted in chronic lymphocytic leukemia. *Oncogene* 15: 2463–2473, 1997.

[69360] 7690. Wolf, S.; Mertens, D.; Schaffner, C.; Korz, C.; Dohner, H.; Stilgenbauer, S.; Lichter, P.: B-cell neoplasia associated gene with multiple splicing (BCMS): the candidate B-CLL gene on 13q14 comprises more than 560kb covering all critical regions. *Hum. Molec. Genet.* 10: 1275–1285, 2001.

[69361] 7691. Agarwal, A. K.; Garg, A.: A novel heterozygous mutation in peroxisome proliferator-activated receptor-gamma gene in a patient with familial partial lipodystrophy. *J. Clin. Endocr. Metab.* 87: 408–411, 2002.

[69362] 7692. Muto, T.; Muramatsu, M.; Taniwaki, M.; Kinoshita, K.; Honjo, T.: Isolation, tissue distribution, and chromosomal localization of the human activation-induced cytidine deaminase (AID) gene. *Genomics* 68:85–88, 2000.

[69363] 7693. Geck, P.; Maffini, M. V.; Szelei, J.; Sonnenschein, C.; Soto, A.M.: Androgen-induced proliferative quiescence in prostate cancer cells: the role of AS3 as its mediator. *Proc. Nat. Acad. Sci.* 97:10185–10190, 2000.

[69364] 7694. Kas, K.; Finger, E.; Grall, F.; Gu, X.; Akbarali, Y.; Boltax, J.; Weiss, A.; Oettgen, P.; Kapeller, R.; Libermann, T. A.: ESE-3, a novel member of an epithelium-specific Ets transcription factor subfamily, demonstrates different tar-

get gene specificity from ESE-1. J. Biol.Chem. 275: 2986–2998, 2000.

[69365] 7695.Kleinbaum, L. A.; Duggan, C.; Ferreira, E.; Coffey, G. P.; Buttice,G.; Burton, F. H.: Human chromosomal localization, tissue/tumor expression,and regulatory function of the ets family gene EHF. Biochem. Biophys.Res. Commun. 264: 119–126, 1999.

[69366] 7696.Davidson, J. D.; Riley, B.; Burrigh, E.N.; Duvick, L.A.; Zoghbi,H.Y.; Orr, H. T.: Identification and characterization of an ataxin-1-interactingprotein: A1Up, a ubiquitin-like nuclear protein. Hum. Molec. Genet. 9:2305–2312, 2000.

[69367] 7697.Zhao, L.; Gregoire, F.; Sul, H. S.: Transient induction of ENC-1,a kelch-related actin-binding protein, is required for adipocyte differentiation. J.Biol. Chem. 275: 16845–16850, 2000.

[69368] 7698.Hansen, L. L.; Jorgensen, R.; Justesen, J.: Assignment of thehuman mitochondrial translational release factor 1 (MTRF1) to chromosome13q14.1–q14.3 and of the human mitochondrial ribosome recycling factor(MRRF) to chromosome 9q32–q34.1 with radiation hybrid mapping. Cytogenet.Cell Genet. 88: 91–92, 2000.

[69369] 7699.Zhang, Y.; Spremulli, L. L.: Identification and cloning of humanmitochondrial translational release factor 1 and

the ribosome recycling factor. *Biochim. Biophys. Acta* 1443: 245–250, 1998.

[69370] 7700. Stelnicki, E. J.; Arbeit, J.; Cass, D. L.; Saner, C.; Harrison, M.; Largman, C.: Modulation of the human homeobox genes PRX-2 and HOXB13 in scarless fetal wounds. *J. Invest. Derm.* 111: 57–63, 1998.

[69371] 7701. Zeltser, L.; Desplan, C.; Heintz, N.: Hoxb-13: a new Hox gene in a distant region of the HOXB cluster maintains colinearity. *Development* 122: 2475–2484, 1996.

[69372] 7702. Rajagopalan, S.; Fu, J.; Long, E. O.: Cutting edge: induction of IFN- $\gamma$  production but not cytotoxicity by the killer cell Ig-like receptor KIR2DL4 (CD158d) in resting NK cells. *J. Immun.* 167: 1877–1881, 2001.

[69373] 7703. Selvakumar, A.; Steffens, U.; Dupont, B.: NK cell receptor gene of the KIR family with two IG domains but highest homology to KIR receptors with three IG domains. *Tissue Antigens* 48: 285–294, 1996.

[69374] 7704. Pende, D.; Biassoni, R.; Cantoni, C.; Verdiani, S.; Falco, M.; di Donato, C.; Accame, L.; Bottino, C.; Moretta, A.; Moretta, L.: The natural killer cell receptor specific for HLA-A allotypes: a novel member of the p58/p70 family of inhibitory receptors that is characterized by three immunoglobulin-like domains and is expressed as a

140-kDdisulphide-linked dimer. J. Exp. Med. 184: 505–518, 1996.

[69375] 7705.Beadling, C.; Johnson, K. W.; Smith, K. A.: Isolation of interleukin2-induced immediate-early genes. Proc. Nat. Acad. Sci. 90: 2719–2723,1993.

[69376] 7706.Gong, R.; Yu, L.; Zhang, H.; Tu, Q.; Zhao, Y.; Yang, J.; Xu, Y.;Zhao, S.: Assignment of human GADD45G to chromosome 9q22.1–q22.3by radiation hybrid mapping. Cytogenet. Cell Genet. 88: 95–96, 2000.

[69377] 7707.Nakayama, K.; Hara, T.; Hibi, M.; Hirano, T.; Miyajima, A.: Anovel oncostatin M-inducible gene OIG37 forms a gene family with MyD118and GADD45 and negatively regulates cell growth. J. Biol. Chem. 274:24766–24772, 1999.

[69378] 7708.Suzuki, M.; Watanabe, T. K.; Fujiwara, T.; Nakamura, Y.; Takahashi,E.; Tanigami, A.: Molecular cloning, expression, and mapping of anovel human cDNA, GRP17, highly homologous to human gadd45 and murineMyD118. J. Hum. Genet. 44: 300–303, 1999.

[69379] 7709.Zhang, X.; Sun, H.; Danila, D. C.; Johnson, S. R.; Zhou, Y.; Swearingen,B.; Klibanski, A.: Loss of expression of GADD45-gamma, a growth inhibitorygene, in human pituitary adenomas: implications for tumorigenesis. J.Clin.

Endocr. Metab. 87: 1262–1267, 2002.

- [69380] 7710. Jones, S. D.; van der Flier, A.; Sonnenberg, A.: Genomic organization of the human alpha-3 integrin subunit gene. *Biochem. Biophys. Res. Commun.* 248: 896–898, 1998.
- [69381] 7711. Takada, Y.; Murphy, E.; Pil, P.; Chen, C.; Ginsberg, M. H.; Hemler, M. E.: Molecular cloning and expression of the cDNA for alpha-3 subunit of human alpha-3/beta-1 (VLA-3), an integrin receptor for fibronectin, laminin, and collagen. *J. Cell Biol.* 115: 257–266, 1991.
- [69382] 7712. Tsuji, T.; Hakomori, S.; Osawa, T.: Identification of human galactoprotein b3, an oncogenic transformation-induced membrane glycoprotein, as VLA-3 alpha subunit: the primary structure of human integrin alpha-3. *J. Biochem.* 109: 659–665, 1991.
- [69383] 7713. Hudson, J. W.; Chen, L.; Fode, C.; Binkert, C.; Dennis, J. W.: Sak kinase gene structure and transcriptional regulation. *Gene* 241: 65–73, 2000.
- [69384] 7714. Karn, T.; Holtrich, U.; Wolf, G.; Hock, B.; Strebhardt, K.; Rubsamen-Waigmann, H.: Human SAK related to the PLK/polo family of cell cycle kinases shows high mRNA expression in testis. *Oncol. Reports* 4: 505–510, 1997.
- [69385] 7715. McMahon, H. T.; Missler, M.; Li, C.; Sudhof, T. C.:

Complexins:cytosolic proteins that regulate SNAP receptor function. Cell 83:111–119, 1995.

[69386] 7716.Reim, K.; Mansour, M.; Varoqueaux, F.; McMahon, H. T.; Sudhof, T. C.; Brose, N.; Rosenmund, C.: Complexins regulate a late step in  $\text{Ca}^{2+}$ –dependent neurotransmitter release. Cell 104: 71–81, 2001.

[69387] 7717.Tokumaru, H.; Umayahara, K.; Pellegrini, L. L.; Ishizuka, T.; Saisu, H.; Betz, H.; Augustine, G. J.; Abe, T.: SNARE complex oligomerization by synaphin/complexin is essential for synaptic vesicle exocytosis. Cell 104:421–432, 2001.

[69388] 7718.Donzeau, M.; Kaldi, K.; Adam, A.; Paschen, S.; Wanner, G.; Guiard, B.; Bauer, M. F.; Neupert, W.; Brunner, M.: Tim23 links the inner and outer mitochondrial membranes. Cell 101: 401–412, 2000.

[69389] 7719.Machesky, L. M.; Insall, R. H.: Scar1 and the related Wiskott–Aldrich syndrome protein, WASP, regulate the actin cytoskeleton through the Arp2/3 complex. Curr. Biol. 8: 1347–1356, 1998.

[69390] 7720.Miki, H.; Suetsugu, S.; Takenawa, T.: WAVE, a novel WASP–family protein involved in actin reorganization induced by Rac. EMBO J. 17:6932–6941, 1998.

[69391] 7721.Scott, A. F.: Personal Communication. Baltimore, Md.

10/23/2001.

- [69392] 7722.McGaughran, J. M.; Ward, H. B.; Evans, D. G.: WAGR syndrome and multiple exostoses in a patient with del(11)(p11.2p14.2). *J. Med.Genet.* 32: 823–824, 1995.
- [69393] 7723.Wuyts, W.; Van Hul, W.; Wauters, J.; Nemtsova, M.; Reyniers, E.; Van Hul, E. V.; De Boulle, K.; de Vries, B. B.; Hendrickx, J.; Herrygers, I.; Bossuyt, P.; Balemans, W.; Fransen, E.; Vits, L.; Coucke, P.; Nowak, N. J.; Shows, T. B.; Mallet, L.; van den Ouweland, A. M.; McGaughran, J.; Hall-ley, D. J.; Willems, P. J.: Positional cloning of a gene involved in hereditary multiple exostoses. *Hum. Molec. Genet.* 5: 1547–1557, 1996.
- [69394] 7724.Clark, M. E.; Kelner, G. S.; Turbeville, L. A.; Boyer, A.; Arden, K. C.; Maki, R. A.: ADAMTS9, a novel member of the ADAM-TS/metallospontingene family. *Genomics* 67: 343–350, 2000.
- [69395] 7725.Marie-Cardine, A.; Kirchgessner, H.; Bruyns, E.; Shevchenko, A.; Mann, M.; Autschbach, F.; Ratnofsky, S.; Meuer, S.; Schraven, B.: SHP2-interacting transmembrane adaptor protein (SIT), a novel disulfide-linked dimer regulating human T cell activation. *J. Exp. Med.* 189: 1181–1194, 1999.
- [69396] 7726.Creasy, C. L.; Chernoff, J.: Cloning and characteriza-

tion of a human protein kinase with homology to Ste20. J. Biol. Chem. 270:21695–21700, 1995.

[69397] 7727. Taylor, L. K.; Wang, H.-C. R.; Erikson, R. L.: Newly identified stress-responsive protein kinases, Krs-1 and Krs-2. Proc. Nat. Acad. Sci. 93: 10099–10104, 1996.

[69398] 7728. Lennard, A.; Gorman, P.; Carrier, M.; Griffiths, S.; Scotney, H.; Sheer, D.; Solari, R.: Cloning and chromosome mapping of the human interleukin-1 receptor antagonist gene. Cytokine 4: 83–89, 1992.

[69399] 7729. Mansfield, J. C.; Holden, H.; Tarlow, J. K.; Di Giovine, F. S.; McDowell, T. L.; Wilson, A. G.; Holdsworth, C. D.; Duff, G. W.: Novel genetic association between ulcerative colitis and the anti-inflammatory cytokine interleukin-1 receptor antagonist. Gastroenterology 106:637–642, 1994.

[69400] 7730. Patterson, D.; Jones, C.; Hart, I.; Bleskan, J.; Berger, R.; Geyer, D.; Eisenberg, S. P.; Smith, M. F., Jr.; Arend, W. P.: The human interleukin-1 receptor antagonist (IL1RN) gene is located in the chromosome 2q14 region. Genomics 15: 173–176, 1993.

[69401] 7731. Steinkasserer, A.; Spurr, N. K.; Cox, S.; Jeggo, P.; Sim, R. B.: The human IL-1 receptor antagonist gene (IL1RN) maps to chromosome 2q14–q21, in the region of



the IL-1-alpha and IL-1-beta loci. *Genomics* 13:654-657, 1992.

[69402] 7732. Tarlow, J. K.; Blakemore, A. I. F.; Lennard, A.; Solari, R.; Hughes, H. N.; Steinkasserer, A.; Duff, G. W.: Polymorphism in human IL-1 receptor antagonist gene intron 2 is caused by variable numbers of an 86-bp tandem repeat. *Hum. Genet.* 91: 403-404, 1993.

[69403] 7733. Tarlow, J. K.; Clay, F. E.; Cork, M. J.; Blakemore, A. I. F.; McDonagh, A. J. G.; Messenger, A. G.; Duff, G. W.: Severity of alopecia areata is associated with a polymorphism in the interleukin-1 receptor antagonist gene. *J. Invest. Dermatol.* 103: 387-390, 1994.

[69404] 7734. Schild, D.; Brake, A. J.; Kiefer, M. C.; Young, D.; Barr, P. J.: Cloning of three human multifunctional de novo purine biosynthetic genes by functional complementation of yeast mutations. *Proc. Nat. Acad. Sci.* 87: 2916-2920, 1990.

[69405] 7735. Garrow, T. A.; Brenner, A. A.; Whitehead, V. M.; Chen, X.-N.; Duncan, R. G.; Korenberg, J. R.; Shane, B.: Cloning of human cDNAs encoding mitochondrial and cytosolic serine hydroxymethyltransferases and chromosomal localization. *J. Biol. Chem.* 268: 11910-11916, 1993.

[69406] 7736. Kao, F.-T.; Chasin, L. A.; Puck, T. T.: Genetics of so-

matic mammalian cells. X. Complementation analysis of glycine-requiring mutants. *Proc. Nat. Acad. Sci.* 64: 1284–1291, 1969.

[69407] 7737. Stover, P. J.; Chen, L. H.; Suh, J. R.; Stover, D. M.; Keyomarsi, K.; Shane, B.: Molecular cloning, characterization, and regulation of the human mitochondrial serine hydroxymethyltransferase gene. *J. Biol. Chem.* 272: 1842–1848, 1997.

[69408] 7738. Abbal, M.; Moennarid, C.; Cambon-Thomsen, A.; Tkaczuk, J.; Ohayon, E.; Mauff, G.: A new BF variant (BF S11) with information for orientation of MHC class III genes. *Immunogenetics* 26: 320–322, 1987.

[69409] 7739. Agarwal, D. P.; Goedde, H. W.; Benkmann, H.-G.; Flatz, G.; Rahimi, A. G.; Kaifie, A.; Delbruck, H.: Genetic polymorphism of C3 and serum levels of immunoglobulins, C3, C4 components of complement and C3-proactivator in four different populations of Afghanistan. *Hum. Genet.* 33: 67–72, 1976.

[69410] 7740. Albert, E. D.; Rittner, C.; Grosse-Wilde, H.; Netzel, B.; Scholz, S.: Recombination frequency and linkage disequilibrium between HL-A and Bf. In: Kissmeyer-Nielsen, F.: *Histocompatibility Testing 1975*. Copenhagen: Munksgaard (pub.) 1975. Pp. 941–944.

- [69411] 7741.Allen, F. H., Jr.: Linkage of HL-A and GBG. Vox Sang. 27: 382-384,1974.
- [69412] 7742.Alper, C. A.: Inherited structural polymorphism in human C2: evidence for genetic linkage between C2 and Bf. J. Exp. Med. 144: 1111-1115,1976.
- [69413] 7743.Alper, C. A.; Boenisch, T.; Watson, L.: Genetic polymorphism in human glycine-rich-beta glycoprotein. J. Exp. Med. 135: 68-80, 1972.
- [69414] 7744.Alper, C. A.; Goodkofsky, I.; Lepow, I. H.: The relationship of glycine-rich beta-glycoprotein to factor B in the properdin system and to cobra-binding protein of human serum. J. Exp. Med. 137: 424-437,1973.
- [69415] 7745.Arnason, A.; Larsen, B.; Marshall, W. H.; Edwards, J. H.; Mackintosh, P.; Olaisen, B.; Teisberg, P.: Very close linkage between HLA-B and Bf inferred from allelic association. Nature 268: 527-528, 1977.
- [69416] 7746.Bender, K.; Mayerova, A.; Frank, R.; Hiller, C.; Wienker, T.: Haplotype analysis of the linkage group HLA-A: HLA-B: Bf and its bearing on the interpretation of the linkage disequilibrium. Hum. Genet. 36:191-196, 1977.
- [69417] 7747.Benkman, H. G.; Goedde, H. W.; Agarwal, D. P.; Flatz, G.; Rahimi, A.; Kaifia, R. S.; Delbruck, H.: Properdin factor B polymorphism in Afghanistan. Hum. Hered. 30:

39–43, 1980.

- [69418] 7748.Bertrams, J.; Mauff, G.: Another family with a silent allele of properdin factor B polymorphism (BF\*QO). *Hum. Genet.* 70: 321–323, 1985.
- [69419] 7749.Campbell, R. D.: The molecular genetics and polymorphism of C2 and factor B. *Brit. Med. Bull.* 43: 37–49, 1987.
- [69420] 7750.Board, P. G.: Biochemical genetics of glutathione-S-transferase in man. *Am. J. Hum. Genet.* 33: 36–43, 1981.
- [69421] 7751.Rochelle, J. M.; Watson, M. L.; Oakey, R. J.; Seldin, M. F.: A linkage map of mouse chromosome 19: definition of comparative mapping relationships with human chromosomes 10 and 11 including the MEN1 locus. *Genomics* 14: 26–31, 1992.
- [69422] 7752.Seldin, M. F.; Saunders, A. M.; Rochelle, J. M.; Howard, T. A.: A proximal mouse chromosome 9 linkage map that further defines linkage groups homologous with segments of human chromosomes 11, 15, and 19. *Genomics* 9: 678–685, 1991.
- [69423] 7753.Baulac, S.; Huberfeld, G.; Gourfinkel-An, I.; Mitropoulou, G.; Beranger, A.; Prud'homme, J.-F.; Baulac, M.; Brice, A.; Bruzzone, R.; LeGuern, E.: First genetic evi-

dence of GABA(A) receptor dysfunction in epilepsy: a mutation in the gamma-2-subunit gene. *Nature Genet.* 28:46–48, 2001.

- [69424] 7754. Buckwalter, M. S.; Lossie, A. C.; Scarlett, L. M.; Camper, S. A.: Localization of the human chromosome 5q genes *Gabra-1*, *Gabrg-2*, *l1-4*, *l1-5*, and *lrf-1* on mouse chromosome 11. *Mammalian Genome* 3:604–607, 1992.
- [69425] 7755. Harkin, L. A.; Bowser, D. N.; Dibbens, L. M.; Singh, R.; Phillips, F.; Wallace, R. H.; Richards, M. C.; Williams, D. A.; Mulley, J. C.; Berkovic, S. F.; Scheffer, I. E.; Petrou, S.: Truncation of the GABA-A-receptor gamma-2 subunit in a family with generalized epilepsy with febrile seizures plus. *Am. J. Hum. Genet.* 70: 530–536, 2002.
- [69426] 7756. Italian League Against Epilepsy Genetic Collaborative Group: Concordance of clinical forms of epilepsy in families with several affected members. *Epilepsia* 34: 819–826, 1993.
- [69427] 7757. Jensen, K. B.; Dredge, B. K.; Stefani, G.; Zhong, R.; Buckanovich, R. J.; Okano, H. J.; Yang, Y. Y. L.; Darnell, R. B.: Nova-1 regulates neuron-specific alternative splicing and is essential for neuronal viability. *Neuron* 25: 359–371, 2000.
- [69428] 7758. Kananura, C.; Haug, K.; Sander, T.; Runge, U.; Gu,

W.; Hallmann, K.; Rebstock, J.; Heils, A.; Steinlein, O. K.: A splice-site mutation in GABRG2 associated with childhood absence epilepsy and febrile convulsions. *Arch. Neurol.* 59: 1137–1141, 2002.

[69429] 7759. Olsen, R. W.; DeLorey, T. M.; Gordey, M.; Kang, M.-H.: GABA receptor function and epilepsy. *Adv. Neurol.* 79: 499–510, 1999.

[69430] 7760. Wallace, R. H.; Marini, C.; Petrou, S.; Harkin, L. A.; Bowser, D. N.; Panchal, R. G.; Williams, D. A.; Sutherland, G. R.; Mulley, J. C.; Scheffer, I. E.; Berkovic, S. F.: Mutant GABA(A) receptor gamma-2-subunit in childhood absence epilepsy and febrile seizures. *Nature Genet.* 28: 49–52, 2001.

[69431] 7761. Ozcelik, T.; Suedhof, T. C.; Francke, U.: The genes for inositol 1,4,5-triphosphate receptors 1 (ITPR1) and 3 (ITPR3) are localized on human chromosomes 3p and 6pter-p21, respectively. (Abstract) *Cytogenet. Cell Genet.* 58: 1880, 1991.

[69432] 7762. Aschauer, H.; Sanguanserm, T.; Braunitzer, G.: Embryonale Haemoglobine des Menschen: Die Primärstruktur der zeta-Ketten (Human embryonic haemoglobins: the primary structure of the zeta chains). *Hoppe Seyler's Z. Physiol. Chem.* 362: 1159–1162,

1981.

- [69433] 7763.Black, J. A.: Human zeta hemoglobin chain. (Letter) Nature 261:348 only, 1976.
- [69434] 7764.Capp, G. L.; Rigas, D. A.; Jones, R. T.: Evidence for a new hemoglobinchain (zeta chain). Nature 228: 278–280, 1970.
- [69435] 7765.Capp, G. L.; Rigas, D. A.; Jones, R. T.: Hemoglobin Portland 1:a new human hemoglobin unique in structure. Science 157: 65–66,1967.
- [69436] 7766.Chung, S.–W.; Wong, S. C.; Clarke, B. J.; Patterson, M.; Walker,W. H. C.; Chui, D. H. K.: Human embryonic zeta–globin chains in adultpatients with alpha–thalassemias. Proc. Nat. Acad. Sci. 81: 6188–6191,1984.
- [69437] 7767.Clegg, J. B.; Gagnon, J.: Structure of the zeta chain of humanembryonic hemoglobin. Proc. Nat. Acad. Sci. 78: 6076–6080, 1981.
- [69438] 7768.Felice, A. E.; Cleek, M. P.; Marino, E. M.; McKie, K. M.; McKie,V. C.; Chang, B. K.; Huisman, T. H. J.: Different zeta globin genedeletions among black Americans. Hum. Genet. 73: 221–224, 1986.
- [69439] 7769.Goodbourn, S. E. Y.; Higgs, D. R.; Clegg, J. B.; Weatherall, D.J.: Molecular basis of length polymorphism in the human zeta–globingene complex. Proc. Nat. Acad.

Sci. 80: 5022–5026, 1983.

- [69440] 7770.Hecht, F.; Jones, R. T.; Koler, R. D.: Newborn infants with HbPortland 1, an indicator of alpha-chain deficiency. Ann. Hum. Genet. 31:215–218, 1968.
- [69441] 7771.Higgs, D. R.; Pressley, L.; Aldridge, B.; Clegg, J. B.; Weatherall,D. J.; Cao, A.; Hadjiminias, M. G.; Kattamis, C.; Metaxatou–Mavromati,A.; Rachmilewitz, E. A.; Sophocleous, T.: Genetic and molecular diversityin nondeletion Hb H disease. Proc. Nat. Acad. Sci. 78: 5833–5837,1981.
- [69442] 7772.Woodrooffe, M. N.; Tunnacliffe, A.; Pym, B.; Goodfellow, P. N.;Walsh, F. S.: Human muscle cell surface antigen 16–3A5 is encodedby a gene on chromosome 11. Somat. Cell Molec. Genet. 10: 535–540,1984.
- [69443] 7773.Hasegawa, M.; Fujimoto, M.; Poe, J. C.; Steeber, D. A.; Lowell,C. A.; Tedder, T. F.: A CD19–dependent signaling pathway regulatesautoimmunity in Lyn–deficient mice. J. Immun. 167: 2469–2478, 2001.
- [69444] 7774.Hasegawa, T.; Hasegawa, Y.; Aso, T.; Koto, S.; Nagai, T.; Tsuchiya,Y.; Kim, K. C.; Ohashi, H.; Wakui, K.; Fukushima, Y.: HDR syndrome(hypoparathyroidism, sensorineural deafness, renal dysplasia) associatedwith del(10)(p13). Am. J. Med. Genet. 73: 416–418, 1997.
- [69445] 7775.Yamada, N.; Makino, Y.; Clark, R. A.; Pearson, D. W.;



Mattei, M.-G.; Guenet, J.-L.; Ohama, E.; Fujino, I.; Miyawaki, A.; Furuichi, T.; Mikoshiba, K.: Human inositol 1,4,5-triphosphate type-1 receptor, InsP3R1: structure, function, regulation of expression and chromosomal localization. *Biochem. J.* 302: 781-790, 1994.

[69446] 7776. Maranto, A. R.: Primary structure, ligand binding, and localization of the human type 3 inositol 1,4,5-trisphosphate receptor expressed in intestinal epithelium. *J. Biol. Chem.* 269: 1222-1230, 1994.

[69447] 7777. Ozcelik, T.; Suedhof, T. C.; Francke, U.: The genes for inositol 1,4,5-triphosphate receptors 1 (ITPR1) and 3 (ITPR3) are localized on human chromosomes 3p and 6pter-p21, respectively. (Abstract) *Cytogenet. Cell Genet.* 58: 1880 only, 1991.

[69448] 7778. Yamamoto-Hino, M.; Sugiyama, T.; Hikichi, K.; Mattei, M. G.; Hasegawa, K.; Sekine, S.; Sakurada, K.; Miyawaki, A.; Furuichi, T.; Hasegawa, M.; Mikoshiba, K.: Cloning and characterization of human type 2 and type 3 inositol 1,4,5-triphosphate receptors. *Receptors Channels* 2:9-22, 1994.

[69449] 7779. Acquati, F.; Malgaretti, N.; Hauptschein, R.; Rao, P.; Gaidano, G.; Taramelli, R.: A 2-Mb YAC contig linking the plasminogen-apoprotein(a) gene family to the insulin-like

growth factor 2 receptor (IGF2R) gene on the telomeric region of chromosome 6 (6q26–q27). *Genomics* 22:664–666, 1994.

[69450] 7780. Barlow, D. P.; Stoger, R.; Herrmann, B. G.; Saito, K.; Schweifer, N.: The mouse insulin-like growth factor type-2 receptor is imprinted and closely linked to the Tme locus. *Nature* 349: 84–87, 1991.

[69451] 7781. DeChiara, T. M.; Robertson, E. J.; Efstratiadis, A.: Parental imprinting of the mouse insulin-like growth factor II gene. *Cell* 64:849–859, 1991.

[69452] 7782. De Souza, A. T.; Hankins, G. R.; Washington, M. K.; Fine, R. L.; Orton, T. C.; Jirtle, R. L.: Frequent loss of heterozygosity on 6q at the mannose 6-phosphate/insulin-like growth factor II receptor locus in human hepatocellular tumors. *Oncogene* 10: 1725–1729, 1995.

[69453] 7783. De Souza, A. T.; Hankins, G. R.; Washington, M. K.; Orton, T. C.; Jirtle, R. L.: M6P/IGF2R gene is mutated in human hepatocellular carcinomas with loss of heterozygosity. *Nature Genet.* 11: 447–449, 1995.

[69454] 7784. Feinberg, A. P.: Genomic imprinting and gene activation in cancer. *Nature Genet.* 4: 110–113, 1993.

[69455] 7785. Haig, D.; Graham, C.: Genomic imprinting and the

strange case of the insulin-like growth factor II receptor.  
Cell 64: 1045–1046, 1991.

- [69456] 7786. Haig, D.; Westoby, M.: Parent-specific gene expression and the triploid endosperm. *Am. Nat.* 134: 147–155, 1989.
- [69457] 7787. Kalscheuer, V. M.; Mariman, E. C.; Schepens, M. T.; Rehder, H.; Ropers, H.-H.: The insulin-like-growth factor type-2 receptor gene is imprinted in the mouse but not in humans. *Nature Genet.* 5: 74–78, 1993.
- [69458] 7788. Kiess, W.; Blickenstaff, G. D.; Sklar, M. M.; Thomas, C. L.; Nissley, S. P.; Sahagian, G. G.: Biochemical evidence that the type II insulin-like growth factor receptor is identical to the cation-independent mannose 6-phosphate receptor. *J. Biol. Chem.* 263: 9339–9344, 1988.
- [69459] 7789. Killian, J. K.; Oka, Y.; Jang, H.-S.; Fu, X.; Waterland, R. A.; Sohda, T.; Sakaguchi, S.; Jirtle, R. L.: Mannose 6-phosphate/insulin-like growth factor 2 receptor (M6P/IGF2R) variants in American and Japanese populations. *Hum. Mutat.* 18: 25–31, 2001.
- [69460] 7790. Kornfeld, S.; Mellman, I.: The biogenesis of lysosomes. *Annu. Rev. Cell Biol.* 5: 483–525, 1989.
- [69461] 7791. Lau, M. M. H.; Stewart, C. E. H.; Liu, Z.; Bhatt, H.; Rotwein, P.; Stewart, C. L.: Loss of the imprinted

IGF2/cation-independentmannose 6-phosphate receptor results in fetal overgrowth and perinatallethality. Genes Dev. 8: 2953–2963, 1994.

- [69462] 7792.Laureys, G.; Barton, D. E.; Ullrich, A.; Francke, U.: Chromosomalmapping of the gene for the type II insulin-like growth factor receptor/cation-independentmannose 6-phosphate receptor in man and mouse. Genomics 3: 224–229,1988.
- [69463] 7793.MacDonald, R. G.; Pfeffer, S. R.; Coussens, L.; Tepper, M. A.;Brocklebank, C. M.; Mole, J. E.; Anderson, J. K.: A single receptorbinds both insulin-like growth factor II and mannose–6-phosphate. Science 239:1134–1137, 1988.
- [69464] 7794.Morgan, D. O.; Edman, J. D.; Standring, D. N.; Fried, V. A.; Smith,M. C.; Roth, R. A.; Rutter, W. J.: Insulin-like growth factor IIreceptor as a multifunctional binding protein. Nature 329: 301–307,1987.
- [69465] 7795.Morris, S. W.; Kirstein, M. N.; Valentine, M. B.; Dittmer, K. G.;Shapiro, D. N.; Saltman, D. L.; Look, A. T.: Fusion of a kinase gene,ALK, to a nucleolar protein gene, NPM, in non-Hodgkin's lymphoma. Science 263:1281–1284, 1994.
- [69466] 7796.Tait, J. F.; Frankenberry, D. A.; Miao, C. H.; Killary, A.

M.;Adler, D. A.; Disteché, C. M.: Chromosomal localization of the humanannexin III (ANX3) gene. Genomics 10: 441–448, 1991.

[69467] 7797.Tait, J. F.; Smith, C.; Xu, L.; Cookson, B. T.: Structure andpolymorphisms of the human annexin III (ANX3) gene. Genomics 18:79–86, 1993.

[69468] 7798.Grundmann, U.; Amann, E.; Abel, K.–J.; Kupper, H. A.: Isolationand expression of cDNA coding for a new member of the phospholipaseA2 inhibitor family. Behring Inst. Mitt. 82: 59–67, 1988.

[69469] 7799.Hauptmann, R.; Maurer–Fogy, I.; Krystek, E.; Bodo, G.; Andree,H.; Reutelingsperger, C. P. M.: Vascular anticoagulant beta, a novelhuman Ca(2+)/phospholipid binding protein that inhibits coagulationand phospholipase A–2 activity: its molecular cloning, expressionand comparison with VAC–alpha. Europ. J. Biochem. 185: 63–71, 1989.

[69470] 7800.Tait, J. F.; Smith, C.; Frankenberry, D. A.; Miao, C. H.; Adler,D. A.; Disteché, C. M.: Chromosomal mapping of the human annexinIV (ANX4) gene. Genomics 12: 313–318, 1992.

[69471] 7801.Bilezikian, J. P.; Morishima, A.; Bell, J.; Grumbach, M. M.: Increasedbone mass as a result of estrogen therapy in a man with aromatasedeficiency. New Eng. J. Med. 339:

599–603, 1998.

- [69472] 7802.Oldridge, M.; Temple, I. K. Santos, H. G.; Gibbons, R. J.; Mustafa,Z.; Chapman, K. E.; Loughlin, J.; Wilkie, A. O. M.: Brachydactylytype B: linkage to chromosome 9q22 and evidence for genetic heterogeneity. *Am.J. Hum. Genet.* 64: 578–585, 1999.
- [69473] 7803.Breimer, L. H.; MacIntyre, I.; Zaidi, M.: Peptides from the calcitonin genes: molecular genetics, structure and function. *Biochem. J.* 255:377–390, 1988.
- [69474] 7804.Schultz, D.; Mikala, G.; Yatani, A.; Engle, D. B.; Iles, D. E.;Segers, B.; Sinke, R. J.; Weghuis, D. O.; Klockner, U.; Wakamori,M.; Wang, J.–J.; Melvin, D.; Varadi, G.; Schwartz, A.: Cloning, chromosomal localization, and functional expression of the alpha–1 subunit of the L–type voltage–dependent calcium channel from normal human heart. *Proc.Nat. Acad. Sci.* 90: 6228–6232, 1993.
- [69475] 7805.Soldatov, N. M.: Genomic structure of human L–type Ca(2+) channel. *Genomics* 22:77–87, 1994.
- [69476] 7806.Sun, W.; McPherson, J. D.; Hoang, D. Q.; Wasmuth, J. J.; Evans,G. A.; Montal, M.: Mapping of a human brain voltage–gated calcium channel to human chromosome 12p13–pter. *Genomics* 14: 1092–1094,1992.
- [69477] 7807.Tsien, R. W.; Ellinor, P. T.; Horne, W. A.: Molecular

diversity of voltage-dependent  $\text{Ca}^{2+}$  channels. Trends Pharm. Sci. 12: 349–354, 1991.

[69478] 7808. Chin, H.; Kozak, C. A.; Kim, H.-L.; Mock, B.; McBride, O. W.: A brain L-type calcium channel  $\alpha$ -1 subunit gene (CCHL1A2) maps to mouse chromosome 14 and human chromosome 3. Genomics 11: 914–919, 1991.

[69479] 7809. Jinnah, H. A.; Yitta, S.; Drew, T.; Kim, B. S.; Visser, J. E.; Rothstein, J. D.: Calcium channel activation and self-biting in mice. Proc. Nat. Acad. Sci. 96: 15228–15232, 1999.

[69480] 7810. Mori, Y.; Friedrich, T.; Kim, M.-S.; Mikami, A.; Nakai, J.; Ruth, P.; Bosse, E.; Hofmann, F.; Flockerzi, V.; Furuichi, T.; Mikoshiba, K.; Imoto, K.; Tanabe, T.; Numa, S.: Primary structure and functional expression from complementary DNA of a brain calcium channel. Nature 350: 398–402, 1991.

[69481] 7811. Pennartz, C. M. A.; de Jeu, M. T. G.; Bos, N. P. A.; Schaap, J.; Geurtsen, A. M. S.: Diurnal modulation of pacemaker potentials and calcium current in the mammalian circadian clock. Nature 416: 286–290, 2002.

[69482] 7812. Platzer, J.; Engel, J.; Schrott-Fischer, A.; Stephan, K.; Bova, S.; Chen, H.; Zheng, H.; Striessnig, J.: Congenital deafness and sinoatrial node dysfunction in mice lacking

class D L-type  $\text{Ca}^{2+}$  channels. Cell 102: 89–97, 2000.

[69483] 7813. Seino, S.; Chen, L.; Seino, M.; Blondel, O.; Takeda, J.; Johnson, J. H.; Bell, G. I.: Cloning of the  $\alpha$ -1 subunit of a voltage-dependent calcium channel expressed in pancreatic beta-cells. Proc. Nat. Acad. Sci. 89: 584–588, 1992.

[69484] 7814. Seino, S.; Yamada, Y.; Espinosa, R., III; Le Beau, M. M.; Bell, G. I.: Assignment of the gene encoding the  $\alpha$ -1 subunit of the neuroendocrine/brain-type calcium channel (CACNL1A2) to human chromosome 3, band p14.3. Genomics 13: 1375–1377, 1992.

[69485] 7815. Powers, P. A.; Liu, S.; Hogan, K.; Gregg, R. G.: Skeletal muscle and brain isoforms of a beta-subunit of human voltage-dependent calcium channels are encoded by a single gene. J. Biol. Chem. 267: 22967–22972, 1992.

[69486] 7816. Gregg, R. G.; Messing, A.; Strube, C.; Beurg, M.; Moss, R.; Behan, M.; Sukhareva, M.; Haynes, S.; Powell, J. A.; Coronado, R.; Powers, P. A.: Absence of the beta subunit (cchb1) of the skeletal muscle dihydropyridine receptor alters expression of the  $\alpha$ -1 subunit and eliminates excitation-contraction coupling. Proc. Nat. Acad. Sci. 93: 13961–13966, 1996.

[69487] 7817. Iles, D. E.; Segers, B.; Sengers, R. C. A.; Monsieurs, K.; Heytens, L.; Halsall, P. J.; Hopkins, P. M.; Ellis, F. R.;



Hall–Curran, J.L.; Stewart, A. D.; Wieringa, B.: Genetic mapping of the beta–1–and gamma–subunits of the human skeletal muscle L–type voltage–dependent calcium channel on chromosome 17q and exclusion as candidate genes for malignant hyperthermia susceptibility. *Hum. Molec. Genet.* 2:863–868, 1993.

[69488] 7818. Pragnell, M.; Sakamoto, J.; Jay, S. D.; Campbell, K. P.: Cloning and tissue–specific expression of the brain calcium channel beta–subunit. *FEBS Lett.* 291: 253–258, 1991.

[69489] 7819. Bulun, S. E.: Aromatase deficiency in women and men: would you have predicted the phenotypes? *J. Clin. Endocr. Metab.* 81: 867–871, 1996.

[69490] 7820. Carani, C.; Qin, K.; Simoni, M.; Faustini–Fustini, M.; Serpente, S.; Boyd, J.; Korach, K. S.; Simpson, E. R.: Effect of testosterone and estradiol in a man with aromatase deficiency. *New Eng. J. Med.* 337:91–95, 1997.

[69491] 7821. Chen, S.; Besman, M. J.; Sparkes, R. S.; Zollman, S.; Klisak, I.; Mohandas, T.; Hall, P. F.; Shively, J. E.: Human aromatase: cDNA cloning, Southern blot analysis, and assignment of the gene to chromosome 15. *DNA* 7: 27–38, 1988.

[69492] 7822. Chen, S.; Shively, J. E.; Nakajin, S.; Shinoda, M.; Hall,

P. F.: Amino terminal sequence analysis of human placenta aromatase. *Biochem. Biophys. Res. Commun.* 135: 713–719, 1986.

[69493] 7823. Conte, F. A.; Grumbach, M. M.; Ito, Y.; Fisher, C. R.; Simpson, E. R.: A syndrome of female pseudohermaphroditism, hypergonadotrophic hypogonadism, and multicystic ovaries associated with missense mutations in the gene encoding aromatase (P450arom). *J. Clin. Endocr. Metab.* 78:1287–1292, 1994.

[69494] 7824. Corbin, C. J.; Graham-Lorence, S.; McPhaul, M.; Mason, J. I.; Mendelson, C. R.; Simpson, E. R.: Isolation of a full-length cDNA insert encoding human aromatase system cytochrome P-450 and its expression in nonsteroidogenic cells. *Proc. Nat. Acad. Sci.* 85: 8948–8952, 1988.

[69495] 7825. Deladoey, J.; Fluck, C.; Bex, M.; Yoshimura, N.; Harada, N.; Mullis, P. E.: Aromatase deficiency caused by a novel P450(arom) gene mutation: impact of absent estrogen production on serum gonadotropin concentration in a boy. *J. Clin. Endocr. Metab.* 84: 4050–4054, 1999.

[69496] 7826. Ellis, J. A.; Stebbing, M.; Harrap, S. B.: Significant population variation in adult male height associated with the Y chromosome and the aromatase gene. *J. Clin. Endocr. Metab.* 86: 4147–4150, 2001.

- [69497] 7827. Evans, C. T.; Ledesma, D. B.; Schulz, T. Z.; Simpson, E. R.; Mendelson, C. R.: Isolation and characterization of a complementary DNA specific for human aromatase-system cytochrome P-450 mRNA. *Proc. Nat. Acad. Sci.* 83: 6387-6391, 1986.
- [69498] 7828. Fisher, C. R.; Graves, K. H.; Parlow, A. F.; Simpson, E. R.: Characterization of mice deficient in aromatase (ArKO) because of targeted disruption of the *cyp19* gene. *Proc. Nat. Acad. Sci.* 95: 6965-6970, 1998.
- [69499] 7829. George, F. W.; Matsumine, H.; McPhaul, M. J.; Somes, R. G., Jr.; Wilson, J. D.: Inheritance of the henney feathering trait in the Golden Campine chicken: evidence for allelism with the gene that causes henney feathering in the Sebright Bantam. *J. Hered.* 81: 107-110, 1990.
- [69500] 7830. George, F. W.; Wilson, J. D.: Pathogenesis of the henney feathering trait in the Sebright Bantam chicken. *J. Clin. Invest.* 66: 57-65, 1980.
- [69501] 7831. Harada, N.: Cloning of a complete cDNA encoding human aromatase: immunochemical identification and sequence analysis. *Biochem. Biophys. Res. Commun.* 156: 725-732, 1988.
- [69502] 7832. Harada, N.; Ogawa, H.; Shozu, M.; Yamada, K.: Genetic studies to characterize the origin of the mutation in

placental aromatase deficiency. Am. J. Hum. Genet. 51: 666–672, 1992.

- [69503] 7833. Harada, N.; Ogawa, H.; Shozu, M.; Yamada, K.; Suhara, K.; Nishida, E.; Takagi, Y.: Biochemical and molecular genetic analyses on placental aromatase (P-450-AROM) deficiency. J. Biol. Chem. 267: 4781–4785, 1992.
- [69504] 7834. Hemsell, D. L.; Edman, C. D.; Marks, J. F.; Siiteri, P. K.; MacDonald, P. C.: Massive extraglandular aromatization of plasma androstenedione resulting in feminization of a prepubertal boy. J. Clin. Invest. 60: 455–464, 1977.
- [69505] 7835. Ito, Y.; Fisher, C. R.; Conte, F. A.; Grumbach, M. M.; Simpson, E. R.: Molecular basis of aromatase deficiency in an adult female with sexual infantilism and polycystic ovaries. Proc. Nat. Acad. Sci. 90: 11673–11677, 1993.
- [69506] 7836. Leiberman, E.; Zachmann, M.: Familial adrenal feminization probably due to increased steroid aromatization. Hormone Res. 37: 96–102, 1992.
- [69507] 7837. Leshin, M.; Baron, J.; George, F. W.; Wilson, J. D.: Increased estrogen formation and aromatase activity in fibroblasts cultured from the skin of chickens with the Henny feathering trait. J. Biol. Chem. 256: 4341–4344, 1981.

- [69508] 7838.Leshin, M.; George, F. W.; Wilson, J. D.: Increased estrogensynthesis in the Sebright bantam is due to a mutation that causesincreased aromatase activity. Trans. Assoc. Am. Phys. 94: 97–105,1981.
- [69509] 7839.Mango, D.; Montemurro, A.; Scirpa, P.; Bompiani, A.; Menini, E.: Four cases of pregnancy with low estrogen production due to placentalezymatic deficiency. Europ. J. Obstet. Gynec. Reprod. Biol. 8:65–71, 1978.
- [69510] 7840.McTernan, P. G.; Anderson. L. A.; Anwar, A. J.; Eggo, M. C.; Crocker,J.; Barnett, A. H.; Stewart, P. M.; Kumar, S.: Glucocorticoid regulationof P450 aromatase activity in human adipose tissue: gender and sitedifferences. J. Clin. Endocr. Metab. 87: 1327–1336, 2002.
- [69511] 7841.Morishima, A.; Grumbach, M. M.; Simpson, E. R.; Fisher, C.; Qin,K.: Aromatase deficiency in male and female siblings caused by a novel mutation and the physiological role of estrogens. J. Clin.Endocr. Metab. 80: 3689–3698, 1995.
- [69512] 7842.Mullis, P. E.; Yoshimura, N.; Kuhlmann, B.; Lippuner, K.; Jaeger,P.; Harada, H.: Aromatase deficiency in a female who is compoundheterozygote for two new point mutations in the P450(arom) gene: impactof estrogens on hypergonadotropic hypogonadism, multicystic ovaries,and

bone densitometry in childhood. *J. Clin. Endocr. Metab.* 82:1739–1745, 1997.

- [69513] 7843. Phornphutkul, C.; Okubo, T.; Wu, K.; Harel, Z.; Tracy, T. F., Jr.; Pinar, H.; Chen, S.; Gruppuso, P. A.; Goodwin, G.: Aromatase P450 expression in a feminizing adrenal adenoma presenting as isosexual precocious puberty. *J. Clin. Endocr. Metab.* 86: 649–652, 2001.
- [69514] 7844. Robertson, K. M.; O'Donnell, L.; Jones, M. E. E.; Meachem, S. J.; Boon, W. C.; Fisher, C. R.; Graves, K. H.; McLachlan, R. I.; Simpson, E. R.: Impairment of spermatogenesis in mice lacking a functional aromatase (cyp 19) gene. *Proc. Nat. Acad. Sci.* 96: 7986–7991, 1999.
- [69515] 7845. Sebastian, S.; Bulun, S. E.: A highly complex organization of the regulatory region of the human CYP19 (aromatase) gene revealed by the Human Genome Project. *J. Clin. Endocr. Metab.* 86: 4600–4602, 2001.
- [69516] 7846. Mohr, J.; Eiberg, H.: Colton blood groups: indication of linkage with the Kidd (Jk) system as support for assignment to chromosome 7. *Clin. Genet.* 11: 372–374, 1977.
- [69517] 7847. Everman, D. B.; Bartels, C. F.; Yang, Y.; Yanamandra, N.; Goodman, F. R.; Mendoza-Londono, J. R.; Savarirayan, R.; White, S. M.; Graham, J. M., Jr.; Gale, R. P.; Svarch, E.; Newman, W. G.; Kleckers, A. R.; Francomano, C. A.; Govin-

daiah, V.; Singh, L.; Morrison, S.; Thomas, J. T.; Warman, M. L.: The mutational spectrum of brachydactyly type C. Am. J. Med. Genet. 112: 291–296, 2002.

[69518] 7848. Galjaard, R. J. H.; van der Ham, L. I.; Posch, N. A. S.; Dijkstra, P. F.; Oostra, B. A.; Hovius, S. E. R.; Timmenga, E. J. F.; Sonneveld, G. J.; Hoogeboom, A. J. M.; Heutink, P.: Differences in complexity of isolated brachydactyly type C cannot be attributed to locus heterogeneity alone. Am. J. Med. Genet. 98: 256–262, 2001.

[69519] 7849. Haws, D. V.: Inherited brachydactyly and hypoplasia of the bones of the extremities. Ann. Hum. Genet. 26: 201–212, 1963.

[69520] 7850. Polinkovsky, A.; Robin, N. H.; Thomas, J. T.; Irons, M.; Lynn, A.; Goodman, F. R.; Reardon, W.; Kant, S. G.; Brunner, H. G.; vander Burgt, I.; Chitayat, D.; McGaughran, J.; Donnai, D.; Luyten, F. P.; Warman, M. L.: Mutations in CDMP1 cause autosomal dominant brachydactyly type C. (Letter) Nature Genet. 17: 18–19, 1997.

[69521] 7851. Polymeropoulos, M. H.; Ide, S. E.; Magyari, T.; Francomano, C. A.: Brachydactyly type C gene maps to human chromosome 12q24. Genomics 38: 45–50, 1996.

[69522] 7852. Robin, N. H.; Gunay-Aygun, M.; Polinkovsky, A.; Warman, M. L.; Morrison, S.: Clinical and locus hetero-

geneity in brachydactyly type C. Am. J. Med. Genet. 68: 369–377, 1997.

[69523] 7853. Storm, E. E.; Huynh, T. V.; Copeland, N. G.; Jenkins, N. A.; Kingsley, D. M.; Lee, S. J.: Limb alterations in brachypodism mice due to mutations in a new member of the TGF- $\beta$  superfamily. Nature 368: 639–643, 1994.

[69524] 7854. Dayhoff, M. O.: Atlas of Protein Sequence and Structure. Hormones, active peptides and toxins. Washington: National Biomedical Research Foundation (pub.) 5: 1972. Pp. D205 only.

[69525] 7855. Edbrooke, M. R.; Parker, D.; McVey, J. H.; Riley, J. H.; Sorenson, G. D.; Pettengill, O. S.; Craig, R. K.: Expression of the human calcitonin/CGRP gene in lung and thyroid carcinoma. EMBO J. 4: 715–724, 1985.

[69526] 7856. Girgis, S. I.; Macdonald, D. W. R.; Stevenson, J. C.; Bevis, P. J. R.; Lynch, C.; Wimalawansa, S. J.; Self, C. H.; Morris, H. R.; MacIntyre, I.: Calcitonin gene-related peptide: potent vasodilator and major product of calcitonin gene. Lancet II: 14–16, 1985.

[69527] 7857. Goltzman, D.; Mitchell, J.: Interaction of calcitonin and calcitonin gene-related peptide at receptor sites in target tissues. Science 227: 1343–1345, 1985.

[69528] 7858. Henry, I.; Grandjouan, S.; Barichard, F.; Huerre-



Jeanpierre, C.;Junien, C.: Mitotic deletions of 11p15.5 in two different tumorsindicate that the CALCA locus is distal to the PTH locus. Cytogenet.Cell Genet. 50: 155–157, 1989.

[69529] 7859.Hillyard, C. J.; Myers, C.; Abeyasekera, G.; Stevenson, J. C.;Craig, R. K.; MacIntyre, I.: Katalcalcin: a new plasma calcium–loweringhormone. Lancet I: 846–848, 1983.

[69530] 7860.Hoovers, J. M. N.; Redeker, E.; Speleman, F.; Hoppener, J. W.M.; Bhola, S.; Blik, J.; van Roy, N.; Leschot, N. J.; Westerveld,A.; Mannens, M.: High–resolution chromosomal localization of thehuman calcitonin/CGRP/IAPP gene family members. Genomics 15: 525–529,1993.

[69531] 7861.Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.; Adema, G.J.; Geurts van Kessel, A. H. M.; Lips, C. J. M.; Jansz, H. S.: Athird human CALC (pseudo)gene on chromosome 11. FEBS Lett. 233:57–63, 1988.

[69532] 7862.Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.; Bakker, E.;Pearson, P. L.; Geurts van Kessel, A. H. M.; Jansz, H. S.; Lips, C.J. M.: Localization of the polymorphic human calcitonin gene on chromosome11. Hum. Genet. 66: 309–312, 1984.

[69533] 7863.Jacobs, J. W.; Goodman, R. H.; Chin, W. W.; Dee, P.

C.; Habener, J. F.; Bell, N. H.; Potts, J. T., Jr.: Calcitonin messenger RNA encodes multiple polypeptides in a single precursor. *Science* 213: 457–459, 1981.

[69534] 7864. Jonas, V.; Lin, C. R.; Kawashima, E.; Semon, D.; Swanson, L. W.; Mermoud, J.-J.; Evans, R. M.; Rosenfeld, M. G.: Alternative RNA processing events in human calcitonin/calcitonin gene-related peptide gene expression. *Proc. Nat. Acad. Sci.* 82: 1994–1998, 1985.

[69535] 7865. Kittur, S. D.; Hoppener, J. W. M.; Antonarakis, S. E.; Daniels, J. D. J.; Meyers, D. A.; Maestri, N. E.; Jansen, M.; Korneluk, R. G.; Nelkin, B. D.; Kazazian, H. H., Jr.: Linkage map of the short arm of human chromosome 11: location of the genes for catalase, calcitonin, and insulin-like growth factor II. *Proc. Nat. Acad. Sci.* 82: 5064–5067, 1985.

[69536] 7866. MacIntyre, I.; Hillyard, C. J.; Murphy, P. K.; Reynolds, J. J.; Gaines-Das, R. E.; Craig, R. K.: A second plasma calcium-lowering peptide from the human calcitonin precursor. *Nature* 300: 460–462, 1982.

[69537] 7867. Mathe, A. A.; Agren, H.; Lindstrom, L.; Theodorsson, E.: Increased concentration of calcitonin gene-related peptide in cerebrospinal fluid of depressed patients: a possible trait marker of major depressive disorder. *Neurosci. Lett.*

182: 138–142, 1994.

[69538] 7868.Neher, R.; Riniker, B.; Rittel, W.; Zuber, H.: Thyrocalcitonin.II. Struktur von alpha-Thyrocalcitonin. *Helv. Chim. Acta* 51: 917–924,1968.

[69539] 7869.New, H. V.; Mudge, A. W.: Calcitonin gene-related peptide regulatesmuscle acetylcholine receptor synthesis. *Nature* 323: 809–811, 1986.

[69540] 7870.Przepiorka, D.; Baylin, S. B.; McBride, D. W.; Testa, J. R.; deBustros, A.; Nelkin, B. D.: The human calcitonin gene is locatedon the short arm of chromosome 11. *Biochem. Biophys. Res. Commun.* 120:493–499, 1984.

[69541] 7871.Rosenfeld, M. G.; Lin, C. R.; Amara, S. G.; Stolarsky, L.; Roos,B. A.; Ong, E. S.; Evans, R. M.: Calcitonin mRNA polymorphism: peptideswitching associated with alternative RNA splicing events. *Proc.Nat. Acad. Sci.* 79: 1717–1721, 1982.

[69542] 7872.Vora, S.; Seaman, C.; Durham, S.; Piomelli, S.: Isozymes of humanphosphofructokinase: identification and subunit structural characterizationof a new system. *Proc. Nat. Acad. Sci.* 77: 62–66, 1980.

[69543] 7873.Pizon, V.; Chardin, P.; Lerosey, I.; Olofsson, B.; Tavian, A.: Human cDNAs RAP1 and RAP2 homologous to the *Drosophila* gene *Dras3*encode proteins closely related to

ras in the 'effector' region. *Oncogene* 3:201–204, 1988.

[69544] 7874.Mimmack, M. L.; Ryan, M.; Baba, H.; Navarro-Ruiz, J.; Iritani, S.; Faull, R. L. M.; McKenna, P. J.; Jones, P. B.; Arai, H.; Starkey, M.; Emson, P. C.; Bahn, S.: Gene expression analysis in schizophrenia: reproducible up-regulation of several members of the apolipoprotein L family located in a high-susceptibility locus for schizophrenia on chromosome 22. *Proc. Nat. Acad. Sci.* 99: 4680–4685, 2002.

[69545] 7875.Schmauss, C.; McAllister, G.; Ohosone, Y.; Hardin, J. A.; Lerner, M. R.: A comparison of snRNP-associated Sm-autoantigens: human N, rat N and human B/B-prime. *Nucleic Acids Res.* 17: 1733–1743, 1989.

[69546] 7876.Kovacs, G.; Kung, H.: Nonhomologous chromatid exchange in hereditary and sporadic renal cell carcinomas. *Proc. Nat. Acad. Sci.* 88: 194–198, 1991.

[69547] 7877.Montini, E.; Rugarli, E. I.; Van de Vosse, E.; Andolfi, G.; Mariani, M.; Puca, A. A.; Consalez, G. G.; den Dunnen, J. T.; Ballabio, A.; Franco, B.: A novel human serine–threonine phosphatase related to the *Drosophila* retinal degeneration C (rdgC) gene is selectively expressed in sensory neurons of neural crest origin. *Hum. Molec. Genet.* 6:1137–1145, 1997.

[69548] 7878.Johnson, M. D.; Tho, S. P. T.; Behzadian, A.; Mc–

Donough, P. G.: Molecular scanning of Yq11 (interval 6) in men with Sertoli–cell–only syndrome. *Am. J. Obstet. Gynec.* 161: 1732–1737, 1989.

[69549] 7879. Bolger, G.; Michaeli, T.; Martins, T.; St. John, T.; Steiner, B.; Rodgers, L.; Riggs, M.; Wigler, M.; Ferguson, K.: A family of human phosphodiesterases homologous to the dunce learning and memory gene product of *Drosophila melanogaster* are potential targets for antidepressant–drugs. *Molec. Cell. Biol.* 13: 6558–6571, 1993.

[69550] 7880. Argyrokastritis, A.; Kamakari, S.; Kapsetaki, M.; Kritis, A.; Talianidis, I.; Moschonas, N. K.: Human hepatocyte nuclear factor–4 (hHNF–4) gene maps to 20q12–q13.1 between PLCG1 and D20S17. *Hum. Genet.* 99:233–236, 1997.

[69551] 7881. Covert, D. D.; Le, T. T.; Morris, G. E.; Man, N. T.; Kralewski, M.; Sendtner, M.; Burghes, A. H. M.: Does the survival motor neuron protein (SMN) interact with Bcl–2? (Letter) *J. Med. Genet.* 37: 536–539, 2000.

[69552] 7882. Nizet, V.; Ohtake, T.; Lauth, X.; Trowbridge, J.; Rudisill, J.; Dorschner, R. A.; Pestonjamas, V.; Piraino, J.; Huttner, K.; Gallo, R. L.: Innate antimicrobial peptide protects the skin from invasive bacterial infection. *Nature* 414: 454–457, 2001.

- [69553] 7883.Weaver, D. R.; Rivkees, S. A.; Carlson, L. L.; Reppert, S. M.:Localization of melatonin receptors in mammalian brain.In: Klein,D. C.; Moore, R. Y.; Reppert, S. M.:  
Suprachiasmatic Nucleus: TheMind's Clock. New York: Oxford Press (pub.) 1991.
- [69554] 7884.Bohm, H.; Gross, B.; Gaestel, M.; Bommer, U.-A.; Ryffel, G.; Bielka,H.: The 5-prime-untranslated region of p23 mRNA from the Ehrlichascites tumor is involved in translation control of the growth relatedprotein p23.  
Biomed. Biochim. Acta 50: 1193-1203, 1991.
- [69555] 7885.Wei, Y.-F.; Robins, P.; Carter, K.; Caldecott, K.; Pappin, D. J.C.; Yu, G.-L.; Wang, R.-P.; Shell, B. K.; Nash, R. A.; Schar, P.;Barnes, D. E.; Haseltine, W. A.; Lindahl, T.:  
Molecular cloning andexpression of human cDNAs encoding a novel DNA ligase IV and DNA ligaseIII, an enzyme active in DNA repair and recombination. Molec. Cell.Biol. 15: 3206-3216, 1995.
- [69556] 7886.Pickard, R. T.; Striffler, B. A.; Kramer, R. M.; Sharp, J. D.:Molecular cloning of two new human paralogs of 85-kDa cytosolic phospholipaseA2. J. Biol. Chem. 274: 8823-8831, 1999.
- [69557] 7887.Hegele, R. A.; Cao, H.; Harris, S. B.; Zinman, B.; Hanley, A.J. G.; Anderson, C. M.: Peroxisome proliferator-acti-

vated receptor–gamma2P12A and type 2 diabetes in Canadian Oji–Cree. *J. Clin. Endocr. Metab.* 85:2014–2019, 2000.

[69558] 7888.Chua, A. O.; Chizzonite, R.; Desai, B. B.; Truitt, T. P.; Nunes,P.; Minetti, L. J.; Warriar, R. R.; Presky, D. H.; Levine, J. F.;Gately, M. K.; Gubler, U.: Expression cloning of a human IL–12 receptorcomponent: a new member of the cytokine receptor superfamily withstrong homology to gp130. *J. Immun.* 153: 128–136, 1994.

[69559] 7889.Wedemeyer, N.; Peoples, R.; Himmelbauer, H.; Lehrach, H.; Francke,U.; Wanker, E. E.: Localization of the human HIP1 gene close to theelastin (ELN) locus on 7q11.23. *Genomics* 46: 313–315, 1997.

[69560] 7890.Zetterstrom, R. H.; Solomin, L.; Jansson, L.; Hoffer, B. J.; Olson,L.; Perlmann, T.: Dopamine neuron agenesis in *Nurr1*–deficient mice. *Science* 276:248–250, 1997.

[69561] 7891.Gong, D.–W.; Monemdjou, S.; Gavrilova, O.; Leon, L. R.; Marcus–Samuels,B.; Chou, C. J.; Everett, C.; Kozak, L. P.; Li, C.; Deng, C.; Harper,M.–E.; Reitman, M. L.: Lack of obesity and normal response to fastingand thyroid hormone in mice lacking uncoupling protein–3. *J. Biol.Chem.* 275: 16251–16257, 2000.

[69562] 7892.Palmiter, R. D.; Cole, T. B.; Findley, S. D.: ZnT–2, a

mammalian protein that confers resistance to zinc by facilitating vesicular sequestration. *EMBO J.* 15: 1784–1791, 1996.

[69563] 7893. Bryan, J.; Kane, R. E.: Actin gelation in sea urchin egg extracts. *Methods Cell Biol.* 25: 175–199, 1982.

[69564] 7894. Dosanjh, M. K.; Collins, D. W.; Fan, W.; Lennon, G. G.; Albala, J. S.; Shen, Z.; Schild, D.: Isolation and characterization of RAD51C, a new human member of the RAD51 family of related genes. *Nucleic Acids Res.* 26: 1179–1184, 1998.

[69565] 7895. Carpino, N.; Wisniewski, D.; Strife, A.; Marshak, D.; Kobayashi, R.; Stillman, B.; Clarkson, B.: p62(dok): a constitutively tyrosine-phosphorylated, GAP-associated protein in chronic myelogenous leukemia progenitor cells. *Cell* 88: 197–204, 1997.

[69566] 7896. Bittner, R. E.; Anderson, L. V. B.; Burkhardt, E.; Bashir, R.; Vafiadaki, E.; Ivanova, S.; Raffelsberger, T.; Maerk, I.; Hoyer, H.; Jung, M.; Karbasiyan, M.; Storch, M.; Lassmann, H.; Moss, J. A.; Davison, K.; Harrison, R.; Bushby, K. M. D.; Reis, A.: Dysferlin deletion in SJL mice (SJL-Dysf) defines a natural model for limb girdle muscular dystrophy 2B. (Letter) *Nature Genet.* 23: 141–142, 1999.



- [69567] 7897.Koi, M.; Johnson, L. A.; Kalikin, L. M.; Little, P. F. R.; Nakamura,Y.; Feinberg, A. P.: Tumor cell growth arrest caused by subchromosomaltransferable DNA fragments from chromosome 11. *Science* 260: 361–364,1993.
- [69568] 7898.Makoff, A.; Pilling, C.; Harrington, K.; Emson, P.: Human metabotropicglutamate receptor type 7: molecular cloning and mRNA distributionin the CNS. *Molec. Brain Res.* 40: 165–170, 1996.
- [69569] 7899.Esterbauer, H.; Oberkofler, H.; Krempler, F.; Patsch, W.: Humanperoxisome proliferator activated receptor gamma coactivator 1 (PPARGC1)gene: cDNA sequence, genomic organization, chromosomal localization,and tissue expression. *Genomics* 62: 98–102, 1999.
- [69570] 7900.Lietzke, S. E.; Bose, S.; Cronin, T.; Klarlund, J.; Chawla, A.;Czech, M. P.; Lambright, D. G.: Structural basis of 3–phosphoinositiderecognition by pleckstrin homology domains. *Molec. Cell* 6: 385–394,2000.
- [69571] 7901.Xu, F.; Xia, W.; Luo, R. Z.; Peng, H.; Zhao, S.; Dai, J.; Long,Y.; Zou, L.; Le, W.; Liu, J.; Parlow, A. F.; Hung, M.–C.; Bast, R.C., Jr.; Yu, Y.: The human ARHI tumor suppressor gene inhibits lactationand growth in transgenic mice. *Cancer Res.* 60: 4913–4920, 2000.
- [69572] 7902.Tsuchida, K.; Arai, K. Y.; Kuramoto, Y.; Yamakawa,

N.; Hasegawa,Y.; Sugino, H.: Identification and characterization of a novel follistatin-likeprotein as a binding protein for the TGF-beta family. J. Biol. Chem. 275:40788-40796, 2000.

[69573] 7903.Takatsu, H.; Yoshino, K.; Nakayama, K.: Adaptor gamma-ear homologydomain conserved in gamma-adaptin and GGA proteins that interact withgamma-synergin. Biochem. Biophys. Res. Commun. 271: 719-725, 2000.

[69574] 7904.Burbelo, P. D.; Snow, D. M.; Bahou, W.; Spiegel, S.: MSE55, aCdc42 effector protein, induces long cellular extensions in fibroblasts. Proc.Nat. Acad. Sci. 96: 9083-9088, 1999.

[69575] 7905.Tsavalier, L.; Shapero, M. H.; Morkowski, S.; Laus, R.: Trp-p8,a novel prostate-specific gene, is up-regulated in prostate cancerand other malignancies and shares high homology with transient receptorpotential calcium channel proteins. Cancer Res. 61: 3760-3769, 2001.

[69576] 7906.Rafi, M. A.; Luzi, P.; Chen, Y. Q.; Wenger, D. A.: A large deletiontogether with a point mutation in the GALC gene is a common mutantallele in patients with infantile Krabbe disease. Hum. Molec. Genet. 4:1285-1289, 1995.

[69577] 7907.Ohta, T.; Michel, J. J.; Schottelius, A. J.; Xiong, Y.:

ROC1, a homolog of APC11, represents a family of cullin partners with an associated ubiquitin ligase activity. *Molec. Cell* 3: 535–541, 1999.

- [69578] 7908. Andreev, J.; Simon, J.-P.; Sabatini, D. D.; Kam, J.; Plowman, G.; Randazzo, P. A.; Schlessinger, J.: Identification of a new Pyk2 target protein with Arf-GAP activity. *Molec. Cell. Biol.* 19: 2338–2350, 1999.
- [69579] 7909. Burden, S.; Yarden, Y.: Neuregulins and their receptors: a versatile signaling module in organogenesis and oncogenesis. *Neuron* 18: 847–855, 1997.
- [69580] 7910. Busfield, S. J.; Michnick, D. A.; Chickering, T. W.; Revett, T. L.; Ma, J.; Woolf, E. A.; Comrack, C. A.; Dussault, B. J.; Woolf, J.; Goodearl, A. D. J.; Gearing, D. P.: Characterization of a neuregulin-related gene, Don-1, that is highly expressed in restricted regions of the cerebellum and hippocampus. *Molec. Cell. Biol.* 17: 4007–4014, 1997.
- [69581] 7911. Carraway, K. L., III; Weber, J. L.; Unger, M. J.; Ledesma, J.; Yu, N.; Gassmann, M.; Lai, C.: Neuregulin-2, a new ligand of ErbB3/ErbB4-receptor tyrosine kinases. *Nature* 387: 512–516, 1997.
- [69582] 7912. Chang, H.; Riese, D. J., II; Gilbert, W.; Stern, D. F.; McMahon, U. J.: Ligands for ErbB-family receptors encoded by a neuregulin-like gene. *Nature* 387: 509–512, 1997.

- [69583] 7913. Ring, H. Z.; Chang, H.; Guilbot, A.; Brice, A.; LeGuern, E.; Francke, U.: The human neuregulin-2 (NRG2) gene: cloning, mapping and evaluation as a candidate for the autosomal recessive form of Charcot-Marie-Tooth disease linked to 5q. *Hum. Genet.* 104: 326-332, 1999.
- [69584] 7914. Yamada, K.; Ichino, N.; Nishii, K.; Sawada, H.; Higashiyama, S.; Ishiguro, H.; Nagatsu, T.: Characterization of the human NTAK gene structure and distribution of the isoforms for rat NTAK mRNA. *Gene* 255:15-24, 2000.
- [69585] 7915. Hu, R.-J.; Lee, M. P.; Connors, T. D.; Johnson, L. A.; Burn, T. C.; Su, K.; Landes, G. M.; Feinberg, A. P.: A 2.5-Mb transcript map of a tumor-suppressing subchromosomal transferable fragment from 11p15.5, and isolation and sequence analysis of three novel genes. *Genomics* 46:9-17, 1997.
- [69586] 7916. Stephan, D.; Bon, C.; Holzwarth, J. A.; Galvan, M.; Pruss, R. M.: Human metabotropic glutamate receptor 1: mRNA distribution, chromosomal localization and functional expression of two splice variants. *Neuropharmacology* 35:1649-1660, 1996.
- [69587] 7917. Chen, M. S.; Huber, A. B.; van der Haar, M. E.; Frank, M.; Schnell, L.; Spillmann, A. A.; Christ, F.; Schwab, M. E.:

Nogo-A is a myelin-associated neurite outgrowth inhibitor and an antigen for monoclonal antibody IN-1. *Nature* 403: 434–439, 2000.

- [69588] 7918. Fournier, A. E.; GrandPré, T.; Strittmatter, S. M.: Identification of a receptor mediating Nogo-66 inhibition of axonal regeneration. *Nature* 409: 341–346, 2001.
- [69589] 7919. GrandPré, T.; Li, S.; Strittmatter, S. M.: Nogo-66 receptor antagonist peptide promotes axonal regeneration. *Nature* 417: 547–551, 2002.
- [69590] 7920. GrandPré, T.; Nakamura, F.; Vartanian, T.; Strittmatter, S. M.: Identification of the Nogo inhibitor of axon regeneration as a reticulon protein. *Nature* 403: 439–444, 2000.
- [69591] 7921. Prinjha, R.; Moore, S. E.; Vinson, M.; Blake, S.; Morrow, R.; Christie, G.; Michalovich, D.; Simmons, D. L.; Walsh, F. S.: Inhibitor of neurite outgrowth in humans. (Letter) *Nature* 403: 383–384, 2000.
- [69592] 7922. Spillmann, A. A.; Bandtlow, C. E.; Lottspeich, F.; Keller, F.; Schwab, M. E.: Identification and characterization of a bovine neurite growth inhibitor (bNI-220). *J. Biol. Chem.* 273: 19283–19293, 1998.
- [69593] 7923. Yang, J.; Yu, L.; Bi, A. D.; Zhao, S.-Y.: Assignment of the human reticulon 4 gene (RTN4) to chromosome

2p14–2p13 by radiation hybridmapping. Cytogenet. Cell Genet. 88: 101–102, 2000.

- [69594] 7924.Korner, C. G.; Wahle, E.: Poly(A) tail shortening by a mammalianpoly(A)–specific 3–prime–exoribonuclease. J. Biol. Chem. 272: 10448–10456,1997.
- [69595] 7925.Korner, C. G.; Wormington, M.; Muckenthaler, M.; Schneider, S.;Dehlin, E.; Wahle, E.: The deadenylating nuclease (DAN) is involvedin poly(A) tail removal during the meiotic maturation of Xenopus oocytes. EMBOJ. 17: 5427–5437, 1998.
- [69596] 7926.Lofberg, H.; Stromblad, L.–G.; Grubb, A. O.; Olsson, S.–O.: Demonstrationof gamma–trace in normal and neoplastic endocrine A–cells of the pancreaticislets: an immunohistochemical study in monkey, rat and man. Biomed.Res. 2: 527–535, 1981.
- [69597] 7927.Merz, G. S.; Benedikz, E.; Schwenk, V.; Johansen, T. E.; Vogel,L. K.; Rushbrook, J. I.; Wisniewski, H. M.: Human cystatin C formsan inactive dimer during intracellular trafficking in transfectedCHO cells. J. Cell. Physiol. 173: 423–432, 1997.
- [69598] 7928.Schnittger, S.; Gopal Rao, V. V. N.; Abrahamson, M.; Hansmann,I.: Cystatin C (CST3), the candidate gene for hereditary cystatinC amyloid angiopathy (HCCAA), and

other members of the cystatin gene family are clustered on chromosome 20p11.2. *Genomics* 16: 50–55, 1993.

[69599] 7929. Shi, G.-P.; Sukhova, G. K.; Grubb, A.; Ducharme, A.; Rhode, L.H.; Lee, R. T.; Ridker, P. M.; Libby, P.; Chapman, H. A.: CystatinC deficiency in human atherosclerosis and aortic aneurysms. *J. Clin. Invest.* 104: 1191–1197, 1999.

[69600] 7930. Ai, Y.; Jenkins, N. A.; Copeland, N. G.; Gilbert, D. J.; Bergsma, D. J.; Stambolian, D.: Mouse galactokinase: isolation, characterization, and location on chromosome 11. *Genome Res.* 5: 53–59, 1995.

[69601] 7931. Ai, Y.; Zheng, Z.; O'Brien-Jenkins, A.; Bernard, D. J.; Wynshaw-Boris, T.; Ning, C.; Reynolds, R.; Segal, S.; Huang, K.; Stambolian, D.: A mouse model of galactose-induced cataracts. *Hum. Molec. Genet.* 9: 1821–1827, 2000.

[69602] 7932. Bergsma, D. J.; Ai, Y.; Skach, W. R.; Nesburn, K.; Anoaia, E.; VanHorn, S.; Stambolian, D.: Fine structure of the human galactokinase *GALK1* gene. *Genome Res.* 6: 980–985, 1996.

[69603] 7933. Croce, C. M.; Huebner, K.; Koprowski, H.: Chromosome assignment of the T-antigen gene of simian virus 40 in African green monkey cells transformed by adeno 7-SV 40 hybrid. *Proc. Nat. Acad. Sci.* 71: 4116–4119, 1974.

[69604] 7934. de Jonge, A. J. R.; de Smit, S.; Kroos, M. A.; Reuser,

A. J. J.: Cotransfer of syntenic human genes into mouse cells using isolated metaphase chromosomes or cellular DNA. Hum. Genet. 69: 32–38, 1985.

- [69605] 7935. Elsevier, S. M.; Kucherlapati, R. S.; Nichols, E. A.; Willecke, K.; Creagan, R. P.; Giles, R. E.; McDougall, J. K.; Ruddle, F. H.: Assignment and regional localization of a gene coding for galactokinase to human chromosome 17q21–22. Birth Defects Orig. Art. Ser. 11(3):117–119, 1975. Note: Alternate: Cytogenet. Cell Genet. 14: 287–289, 1975.
- [69606] 7936. Klobutcher, L. A.; Ruddle, F. H.: Phenotype stabilization and integration of transferred material in chromosome-mediated gene transfer. Nature 280: 657–660, 1979.
- [69607] 7937. Kolosha, V.; Anoaia, E.; de Cespedes, C.; Gitzelmann, R.; Shih, L.; Casco, T.; Saborio, M.; Trejos, R.; Buist, N.; Tedesco, T.; Skach, W.; Mitelmann, O.; Ledee, D.; Huang, K.; Stambolian, D.: Novel mutations in 13 probands with galactokinase deficiency. Hum. Mutat. 15: 447–453, 2000.
- [69608] 7938. Okajima, K.; Yazaki, M.; Wada, Y.: Thymidine kinase activity in individuals with galactokinase deficiency. (Letter) Am. J. Hum. Genet. 41: 503–504, 1987.
- [69609] 7939. Okano, Y.; Asada, M.; Fujimoto, A.; Ohtake, A.; Mu-



rayama, K.;Hsiao, K.-J.; Choeh, K.; Yang, Y.; Cao, Q.; Reichardt, J. K. V.; Niihira,S.; Imamura, T.; Yamano, T.: A genetic factor for age-related cataract:identification and characterization of a novel galactokinase variant,'Osaka,' in Asians. Am. J. Hum. Genet. 68: 1036-1042, 2001.

[69610] 7940.Ruddle, F. H.: Personal Communication. New Haven, Conn. 5/4/1982.

[69611] 7941.Schoen, R. C.; Cox, S. H.; Wagner, R. P.: Thymidine-kinase activityof cultured cells from individuals with inherited galactokinase deficiency. Am.J. Hum. Genet. 36: 815-822, 1984.

[69612] 7942.Scott, A. F.: Personal Communication. Baltimore, Md. 2/20/2001.

[69613] 7943.Schmidt, A.; Wolde, M.; Thiele, C.; Fest, W.; Kratzin, H.; Podtelejnikov,A. V.; Witke, W.; Huttner, W. B.; Soling, H.-D.: Endophilin I mediatesynaptic vesicle formation by transfer of arachidonate to lysophosphatidicacid. Nature 401: 133-141, 1999.

[69614] 7944.Aiba, A.; Chen, C.; Herrup, K.; Rosenmund, C.; Stevens, C. F.;Tonegawa, S.: Reduced hippocampal long-term potentiation and context-specificdeficit in associative learning in mGluR1 mutant mice. Cell 79:365-375, 1994.

- [69615] 7945.Aiba, A.; Kano, M.; Chen, C.; Stanton, M. E.; Fox, G. D.; Herrup, K.; Zwingman, T. A.; Tonegawa, S.: Deficient cerebellar long-term depression and impaired motor learning in mGluR1 mutant mice. *Cell* 79:377–388, 1994.
- [69616] 7946.Conquet, F.; Bashir, Z. I.; Davies, C. H.; Daniel, H.; Ferraguti, F.; Bordi, F.; Franz-Bacon, K.; Reggiani, A.; Matarese, V.; Conde, F.; Collingridge, G. L.; Crepel, F.: Motor deficit and impairment of synaptic plasticity in mice lacking mGluR1. *Nature* 372: 237–243, 1994.
- [69617] 7947.Ganesh, S.; Amano, K.; Yamakawa, K.: Assignment of the gene GRM1 coding for metabotropic glutamate receptor 1 to human chromosome band 6q24 by in situ hybridization. *Cytogenet. Cell Genet.* 88: 314–315, 2000.
- [69618] 7948.Ichise, T.; Kano, M.; Hashimoto, K.; Yanagihara, D.; Nakao, K.; Shigemoto, R.; Katsuki, M.; Aiba, A.: mGluR1 in cerebellar Purkinje cells essential for long-term depression, synapse elimination, and motor coordination. *Science* 288: 1832–1835, 2000.
- [69619] 7949.Kunishima, N.; Shimada, Y.; Tsuji, Y.; Sato, T.; Yamamoto, M.; Kumasaka, T.; Nakanishi, S.; Jingami, H.; Morikawa, K.: Structural basis of glutamate recognition by a dimeric metabotropic glutamate receptor. *Nature* 407: 971–977, 2000.

- [69620] 7950.Okamoto, T.; Sekiyama, N.; Otsu, M.; Shimada, Y.; Sato, A.; Nakanishi,S.; Jingami, H.: Expression and purification of the extracellularligand binding region of metabotropic glutamate receptor subtype 1. *J.Biol. Chem.* 273: 13089–13096, 1998.
- [69621] 7951.Smitt, P. S.; Kinoshita, A.; De Leeuw, B.; Moll, W.; Coesmans,M.; Jaarsma, D.; Henzen–Logmans, S.; Vecht, C.; De Zeeuw, C.; Sekiyama,N.; Nakanishi, S.; Shigemoto, R.: Paraneoplastic cerebellar ataxiadue to autoantibodies against a glutamate receptor. *New Eng. J. Med.* 342:21–27, 2000.
- [69622] 7952.Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hiro-sawa, M.;Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Predictionof the coding sequences of unidentified human genes. XII. The complete sequences of 100 new cDNA clones from brain which code for large proteinsin vitro. *DNA Res.* 5: 355–364, 1998.
- [69623] 7953.Chadwick, B. P.; Kidd, T.; Sgouros, J.; Ish–Horowicz, D.; Frischauf,A.–M.: Cloning, mapping and expression of UBL3, a novel ubiquitin–likegene. *Gene* 233: 189–195, 1999.
- [69624] 7954.Hodges, M.; Tissot, C.; Freemont, P. S. Protein regulation:tag wrestling with relatives of ubiquitin. *Curr. Biol.*

8: R749–R752,1998.

- [69625] 7955.Bora, R. S.; Kanamori, A.; Hirabayashi, Y.: Assignment of a putative acetyl–CoA transporter gene (Acatn) to mouse chromosome band 3E1–E3 by in situ hybridization. *Cytogenet. Cell Genet.* 83: 78–79, 1998.
- [69626] 7956.Andree, B.; Hillemann, T.; Kessler–Ieckson, G.; Schmitt–John, T.; Jockusch, H.; Arnold, H.–H.; Brand, T.: Isolation and characterization of the novel Popeye gene family expressed in skeletal muscle and heart. *Dev.Biol.* 223: 371–382, 2000.
- [69627] 7957.Reese, D. E.; Bader, D. M.: Cloning and expression of hbves, a novel and highly conserved mRNA expressed in the developing and adult heart and skeletal muscle in the human. *Mammalian Genome* 10: 913–915,1999.
- [69628] 7958.Reese, D. E.; Zavaljevski, M.; Streiff, N. L.; Bader, D.: Bves: a novel gene expressed during coronary blood vessel development. *Dev.Biol.* 209: 159–171, 1999.
- [69629] 7959.Kowal, R. C.; Jolsin, J. M.; Olson, E. N.; Schultz, R. A.: Assignment of fibulin–5 (FBLN5) to human chromosome 14q31 by in situ hybridization and radiation hybrid mapping. *Cytogenet. Cell Genet.* 87: 2–3, 1999.
- [69630] 7960.Nakamura, T.; Lozano, P. R.; Ikeda, Y.; Iwanaga, Y.; Hinek, A.; Minamisawa, S.; Cheng, C.–F.; Kobuke, K.; Dal–

ton, N.; Takada, Y.; Tashiro, K.; Ross, J., Jr.; Honjo, T.; Chien, K. R.: Fibulin-5/DANCE is essential for elastogenesis in vivo. *Nature* 415: 171–175, 2002.

[69631] 7961. Nakamura, T.; Ruiz-Lozano, P.; Lindner, V.; Yabe, D.; Taniwaki, M.; Furukawa, Y.; Kobuke, K.; Tashiro, K.; Lu, Z.; Andon, N. L.; Schaub, R.; Matsumori, A.; Sasayama, S.; Chien, K. R.; Honjo, T.: DANCE, a novel secreted RGD protein expressed in developing, atherosclerotic, and balloon-injured arteries. *J. Biol. Chem.* 274: 22476–22483, 1999.

[69632] 7962. Yanagisawa, H.; Davis, E. C.; Starcher, B. C.; Ouchi, T.; Yanagisawa, M.; Richardson, J. A.; Olson, E. N.: Fibulin-5 is an elastin-binding protein essential for elastic fibre development in vivo. *Nature* 415: 168–171, 2002.

[69633] 7963. Dong, F.; Feldmesser, M.; Casadevall, A.; Rubin, C. S.: Molecular characterization of a cDNA that encodes six isoforms of a novel murine A kinase anchor protein. *J. Biol. Chem.* 273: 6533–6541, 1998.

[69634] 7964. Dent, A. L.; Yewdell, J.; Puvion-Dutilleul, F.; Koken, M. H.; de The, H.; Staudt, L. M.: LYSP100 associated nuclear domains (LANDs): description of a new class of subnuclear structures and their relationship to PML nuclear bodies. *Blood* 88: 1423–1426, 1996.

[69635] 7965. Seeler, J. S.; Marchio, A.; Sitterlin, D.; Transy, C.; De-

jean,A.: Interaction of SP100 with HP1 proteins: a link between the promyelocytic leukemia-associated nuclear bodies and the chromatin compartment. *Proc.Nat. Acad. Sci.* 95: 7316–7321, 1998.

[69636] 7966.Scanlan, M. J.; Chen, Y.–T.; Williamson, B.; Gure, A. O.; Stockert,E.; Gordan, J. D.; Tureci, O.; Sahin, U.; Pfrendschuh, M.; Old, L.J.: Characterization of human colon cancer antigens recognized by autologous antibodies. *Int. J. Cancer* 76: 652–658, 1998.

[69637] 7967.Halford, S.; Dulai, K. S.; Daw, S. C.; Fitzgibbon, J.; Hunt, D.M.: Isolation and chromosomal localization of two human CDP–diacylglycerol synthase (CDS) genes. *Genomics* 54: 140–144, 1998.

[69638] 7968.Heacock, A. M.; Uhler, M. D.; Agranoff, B. W.: Cloning of CDP–diacylglycerol synthase from a human neuronal cell line. *J. Neurochem.* 67: 2200–2203, 1996.

[69639] 7969.Weeks, R.; Dowhan, W.; Shen, H.; Balantac, N.; Meengs, B.; Nudelman,E.; Leung, D. W.: Isolation and expression of an isoform of human CDP–diacylglycerol synthase cDNA. *DNA Cell Biol.* 16: 281–289, 1997.

[69640] 7970.Volta, M.; Bulfone, A.; Gattuso, C.; Rossi, E.; Mariani, M.; Consalez,G. G.; Zuffardi, O.; Ballabio, A.; Banfi, S.; Franco, B.: Identification and characterization of CDS2, a

mammalian homolog of the *Drosophila* CDP-diacylglycerol synthase gene. *Genomics* 55: 68–77, 1999.

[69641] 7971. Borsani, G.; DeGrandi, A.; Ballabio, A.; Bulfone, A.; Bernard, L.; Banfi, S.; Gattuso, C.; Mariani, M.; Dixon, M.; Donnai, D.; Metcalfe, K.; Winter, R.; Robertson, M.; Axton, R.; Brown, A.; van Heyningen, V.; Hanson, I.: EYA4, a novel vertebrate gene related to *Drosophila* eyes absent. *Hum. Molec. Genet.* 8: 11–23, 1999.

[69642] 7972. Lepperdinger, G.; Strobl, B.; Kreil, G.: HYAL2, a human gene expressed in many cells, encodes a lysosomal hyaluronidase with a novel type of specificity. *J. Biol. Chem.* 273: 22466–22470, 1998.

[69643] 7973. Kanamori, A.; Nakayama, J.; Fukuda, M. N.; Stallcup, W. B.; Sasaki, K.; Fukuda, M.; Hirabayashi, Y.: Expression cloning and characterization of a cDNA encoding a novel membrane protein required for the formation of O-acetylated ganglioside: a putative acetyl-CoA transporter. *Proc. Nat. Acad. Sci.* 94: 2897–2902, 1997.

[69644] 7974. Holmes, M.; Turner, J.; Fox, A.; Chisholm, O.; Crossley, M.; Chong, B.: hFOG-2, a novel zinc finger protein, binds the co-repressor mCtBP2 and modulates GATA-mediated activation. *J. Biol. Chem.* 274: 23491–23498, 1999.

- [69645] 7975.Svensson, E. C.; Huggins, G. S.; Lin, H.; Clendenin, C.; Jiang,F.; Tufts, R.; Dardik, F. B.; Leiden, J. M.: A syndrome of tricuspidatresia in mice with a targeted mutation of the gene encoding Fog-2. NatureGenet. 25: 353-356, 2000.
- [69646] 7976.Svensson, E. C.; Tufts, R. L.; Polk, C. E.; Leiden, J. M.: Molecularcloning of FOG-2: a modulator of transcription factor GATA-4 in cardiomyocytes. Proc.Nat. Acad. Sci. 96: 956-961, 1999.
- [69647] 7977.Tevosian, S. G.; Deconinck, A. E.; Tanaka, M.; Schinke, M.; Litovsky,S. H.; Izumo, S.; Fujiwara, Y.; Orkin, S. H.: FOG-2, a cofactor forGATA transcription factors, is essential for heart morphogenesis anddevelopment of coronary vessels from epicardium. Cell 101: 729-739,2000.
- [69648] 7978.Behrends, S.; Kazmierczak, B.; Steenpass, A.; Knauf, B.; Bullerdiek,J.; Scholz, H.; Eiberg, H.: Assignment of GUCY1B2, the gene codingfor the beta-2 subunit of human guanylyl cyclase to chromosomal band13q14.3 between markers D13S168 and D13S155. Genomics 55: 126-127,1999.
- [69649] 7979.Malterer, A.; Gupta, G.; Danziger, R. S.: Assignment of GUCY1B2,the human homologue of a candidate gene



for hypertension, to chromosomebands 13q14.2–q14.3 by in situ hybridization. Cytogenet. Cell Genet. 85:256–257, 1999.

- [69650] 7980.Yuen, P. S. T.; Potter, L. R.; Garbers, D. L.: A new form of guanylylcyclase is preferentially expressed in rat kidney. Biochemistry 29:10872–10878, 1990.
- [69651] 7981.Paavola, P.; Horelli-Kuitunen, N.; Palotie, A.; Peltonen, L.:Characterization of a novel gene, PNUTL2, on human chromosome 17q22–q23and its exclusion as the Meckel syndrome gene. Genomics 55: 122–125,1999.
- [69652] 7982.Tedesco, F.; Roncelli, L.; Petersen, B. H.; Agnello, V.; Sodetz,J. M.: Two distinct abnormalities in patients with C8–alpha–gammadeficiency: low level of C8–beta chain and presence of dysfunctionalC8–alpha–gamma subunit. J. Clin. Invest. 86: 884–888, 1990.
- [69653] 7983.Savov, A.; Angelicheva, D.; Balassopoulou, A.; Jordanova, A.;Noussia-Arvanitakis, S.; Kalaydjieva, L.: Double mutant alleles:are they rare? Hum. Molec. Genet. 4: 1169–1171, 1995.
- [69654] 7984.Theriault, A.; Boyd, E.; Whaley, K.; Sodetz, J. M.; Connor, J.M.: Regional chromosomal assignment of genes encoding the alpha andbeta subunits of human complement protein c8 to 1p32. (Abstract) Cytogenet.Cell Genet.

58: 1864 only, 1991.

- [69655] 7985.Theriault, A.; Boyd, E.; Whaley, K.; Sodetz, J. M.; Connor, J.M.: Regional chromosomal assignment of genes encoding the alpha and beta subunits of human complement protein C8 to 1p32. Hum. Genet. 88:703–704, 1992.
- [69656] 7986.Zhang, L.; Rittner, C.; Sodetz, J. M.; Schneider, P. M.; Kaufmann,T.: The eighth component of human complement: molecular basis of C8A (C81) polymorphism. Hum. Genet. 96: 281–284, 1995.
- [69657] 7987.Bahary, N.; Zorich, G.; Pachter, J. E.; Leibel, R. L.; Friedman,J. M.: Molecular genetic linkage maps of mouse chromosomes 4 and 6. Genomics 11: 33–47, 1991.
- [69658] 7988.Arnaout, M. A.; Gupta, S. K.; Pierce, M. W.; Tenen, D. G.: Amino acid sequence of the alpha subunit of human leukocyte adhesion receptor Mo1 (complement receptor type 3). J. Cell Biol. 106: 2153–2158, 1988.
- [69659] 7989.Arnaout, M. A.; Remold–O'Donnell, E.; Pierce, M. W.; Harris, P.; Tenen, D. G.: Molecular cloning of the alpha-subunit of human and guinea pig leukocyte adhesion glycoprotein Mo1: chromosomal localization and homology to the alpha-subunits of integrins. Proc. Nat. Acad.Sci. 85: 2776–2780, 1988.
- [69660] 7990.Callen, D. F.; Chen, L. Z.; Nancarrow, J.; Whitmore, S.

A.; Apostolou,S.; Thompson, A. D.; Lane, S. A.; Stallings, R. L.; Hildebrand, C.E.; Harris, P. G.; Sutherland, G. R.: Current state of the physical map of human chromosome 16. (Abstract) Cytogenet. Cell Genet. 58:1998 only, 1991.

[69661] 7991. Corbi, A. L.; Kishimoto, T. K.; Miller, L. J.; Springer, T. A.: The human leukocyte adhesion glycoprotein Mac-1 (complement receptor type 3, CD11b) alpha subunit: cloning, primary structure, and relation to the integrins, von Willebrand factor and factor B. J. Biol. Chem. 263:12403–12411, 1988.

[69662] 7992. Corbi, A. L.; Larson, R. S.; Kishimoto, T. K.; Springer, T. A.; Morton, C. C.: Chromosomal location of the genes encoding the leukocyte adhesion receptors LFA-1, Mac-1 and p150,95: identification of a gene cluster involved in cell adhesion. J. Exp. Med. 167: 1597–1607, 1988.

[69663] 7993. Pierce, M. W.; Remold-O'Donnell, E.; Todd, R. F., III; Arnaout, M. A.: N-terminal sequence of human leukocyte glycoprotein Mo1: conservation across species and homology to platelet IIb/IIIa. Biochim. Biophys. Acta 874: 368–371, 1986.

[69664] 7994. Simon, D. I.; Chen, Z.; Seifert, P.; Edelman, E. R.; Balantyne, C. M.; Rogers, C.: Decreased neointimal formation in Mac-1  $-/-$  mice reveals a role for inflammation in vas-

cular repair after angioplasty. *J.Clin. Invest.* 105: 293–300, 2000.

- [69665] 7995.Springer, T. A.; Teplow, D. B.; Dreyer, W. J.: Sequence homology of the LFA-1 and Mac-1 leukocyte adhesion glycoproteins and unexpected relation to leukocyte interferon. *Nature* 314: 540–542, 1985.
- [69666] 7996.Cartier, M.; Breitman, M. L.; Tsui, L.-C.: A frameshift mutation in the gamma-E-crystallin gene of the Elo mouse. *Nature Genet.* 2:42–45, 1992.
- [69667] 7997.Petersen, M. B.; Slaugenhaupt, S. A.; Lewis, J. G.; Warren, A.C.; Chakravarti, A.; Antonarakis, S. E.: A genetic linkage map of 27 markers on human chromosome 21. *Genomics* 9: 407–419, 1991.
- [69668] 7998.Amagai, M.; Klaus-Kovtun, V.; Stanley, J. R.: Autoantibodies against a novel epithelial cadherin in pemphigus vulgaris, a disease of cell adhesion. *Cell* 67: 869–877, 1991.
- [69669] 7999.Koch, P. J.; Mahoney, M. G.; Ishikawa, H.; Pulkkinen, L.; Uitto, J.; Schultz, L.; Murphy, G. F.; Whitaker-Menezes, D.; Stanley, J.R.: Targeted disruption of the pemphigus vulgaris antigen (desmoglein 3) gene in mice causes loss of keratinocyte cell adhesion with a phenotype similar to pemphigus vulgaris. *J. Cell Biol.* 137: 1091–1102, 1997.

- [69670] 8000.Wang, Y.; Amagai, M.; Minoshima, S.; Sakai, K.; Green, K. J.; Nishikawa,T.; Shimizu, N.: The human genes for desmogleins (DSG1 and DSG3)are located in a small region on chromosome 18q12. *Genomics* 20:492–495, 1994.
- [69671] 8001.Arnemann, J.; Spurr, N. K.; Magee, A. I.; Buxton, R. S.: The humangene (DSG2) coding for HDGC, a second member of the desmoglein subfamilyof the desmosomal cadherins, is, like DSG1 coding for desmoglein DGI,assigned to chromosome 18. *Genomics* 13: 484–486, 1992.
- [69672] 8002.Koch, P. J.; Goldschmidt, M. D.; Walsh, M. J.; Zimbelmann, R.;Franke, W. W.: Complete amino acid sequence of the epidermal desmogleinprecursor polypeptide and identification of a second type of desmogleingene. *Europ. J. Cell Biol.* 55: 200–208, 1991.
- [69673] 8003.Inskip, A.; Elexperu–Camiruaga, J.; Buxton, N.; Dias, P. S.; MacIntosh,J.; Campbell, D.; Jones, P. W.; Yengi, L.; Talbot, J. A.; Strange,R. C.; Fryer, A. A.: Identification of polymorphism at the glutathioneS–transferase, GSTM3 locus: evidence for linkage with GSTM1\*A. *Biochem.J.* 312: 713–716, 1995.
- [69674] 8004.Van Cong, N.; Laisney, V.; Gross, M. S.; Frezal, J.:

Glutathione-S-transferases--tissuesdistribution, number of loci, polymorphism, chromosome localization.(Abstract) Cytogenet. Cell Genet. 37: 554, 1984.

- [69675] 8005.Benham, F. J.; Povey, S.: Members of the human glyceraldehyde-3-phosphatedehydrogenase-related gene family map to dispersed chromosomal locations. Genomics 5:209-214, 1989.
- [69676] 8006.Bruns, G. A. P.; Gerald, P. S.: Human glyceraldehyde-3-phosphatedehydrogenase in man-rodent somatic cell hybrids. Science 192: 54-56,1976.
- [69677] 8007.Edwards, Y. H.; Clark, P.; Harris, H.: Isozymes of glyceraldehyde-3-phosphatedehydrogenase in man and other mammals. Ann. Hum. Genet. 40: 67-77,1976.
- [69678] 8008.Goodfellow, P. N.; Davies, K. E.; Ropers, H.-H.: Report of thecommittee on the genetic constitution of the X and Y chromosomes. Cytogenet.Cell Genet. 40: 296-352, 1985.
- [69679] 8009.Law, M. L.; Kao, F.-T.: Induced segregation of human syntenicgenes by 5-bromodeoxyuridine plus near-visible light. Somat. CellGenet. 4: 465-476, 1978.
- [69680] 8010.Lebherz, H. G.; Rutter, W. J.: Glyceraldehyde-3-phosphate dehydrogenasevariants in phyletically diverse organisms. Science 157: 1198-1199,1967.

- [69681] 8011.Piechaczyk, M.; Blanchard, J. M.; Riaad-el Sabouty, S.; Dani,C.; Marty, L.; Jeanteur, P.: Unusual abundance of glyceraldehyde3-phosphate dehydrogenase pseudogenes in vertebrate genomes. *Nature* 312:469-471, 1984.
- [69682] 8012.Rethore, M.-O.; Junien, C.; Malpuech, G.; Baccichetti, C.; Tenconi,R.; Kaplan, J.-C.; de Romeuf, J.; Lejeune, J.: Localisation du gene de la glyceraldehyde-3-phosphate dehydrogenase (G3PD) sur le segment distal du bras court de chromosome 12. *Ann. Genet.* 19: 140-142,1976.
- [69683] 8013.Rivas, F.; Vaca, G.; Zuniga, G.; Gonzalez, R. M.; Ruiz, C.; Rivera,H.; Moller, M.; Cantu, J. M.: 46,XX,-12,+der(12),rcp(3;12)(p25.1;p13.31)patkaryotype in a girl: probably subregional assignment of glyceraldehyde-3-phosphatedehydrogenase locus to 12p13.1-p13.31 by exclusion. *Ann. Genet.* 28:189-192, 1985.
- [69684] 8014.Serville, F.; Junien, C.; Kaplan, J. C.; Gachet, M.; Cadoux, J.;Broustet, A.: Gene dosage effect for human triosephosphate isomerase and glyceraldehyde-3-phosphate dehydrogenase in partial trisomy 12p13 and trisomy 18p. *Hum. Genet.* 45: 63-69, 1978.
- [69685] 8015.Tso, J. Y.; Sun, X.-H.; Kao, T.; Reece, K. S.; Wu, R.: Isolation and characterization of rat and human glycer-

aldehyde-3-phosphate dehydrogenase cDNAs: genomic complexity and molecular evolution of the gene. *Nucleic Acids Res.* 13: 2485-2502, 1985.

- [69686] 8016. Hashimoto, S.; Chiorazzi, N.; Gregersen, P. K.: The complete sequence of the human CD79b (Ig-beta/B29) gene: identification of a conserved exon/intron organization, immunoglobulin-like regulatory regions, and allelic polymorphism. *Immunogenetics* 40: 145-149, 1994.
- [69687] 8017. Hashimoto, S.; Gregersen, P. K.; Chiorazzi, N.: The human Ig-beta cDNA sequence, a homologue of murine B29, is identical in B cell and plasma cell lines producing all the human Ig isotypes. *J. Immun.* 150:491-498, 1993.
- [69688] 8018. Wood, W. J., Jr.; Thompson, A. A.; Korenberg, J.; Chen, X.; May, W.; Wall, R.; Denny, C. T.: Isolation and chromosomal mapping of the human immunoglobulin-associated B29 gene (IGB). *Genomics* 16:187-192, 1993.
- [69689] 8019. Okabe, I.; Nussbaum, R. L.: Identification and chromosomal mapping of the mouse inositol polyphosphate 1-phosphatase gene. *Genomics* 30:358-360, 1995.
- [69690] 8020. Woodcock, E. A.; Wang, B. H.; Arthur, J. F.; Lennard, A.; Matkovich, S. J.; Du, X.-J.; Brown, J. H.; Hannan, R. D.: Inositol polyphosphate 1-phosphatase is a novel antihypertrophic factor. *J. Biol. Chem.* 277:22734-22742, 2002.



- [69691] 8021.York, J. D.; Veile, R. A.; Donis-Keller, H.; Majerus, P. W.: Cloning, heterologous expression, and chromosomal localization of human inositol polyphosphate 1-phosphatase. *Proc. Nat. Acad. Sci.* 90: 5833–5837, 1993.
- [69692] 8022.Janne, P. A.; Dutra, A. S.; Dracopoli, N. C.; Charnas, L. R.; Puck, J. M.; Nussbaum, R. L.: Localization of the 75-kDa inositol polyphosphate-5-phosphatase (INPP5B) to human chromosome band 1p34. *Cytogenet. Cell Genet.* 66:164–166, 1994.
- [69693] 8023.Janne, P. A.; Rochelle, J. M.; Martin-DeLeon, P. A.; Stambolian, D.; Seldin, M. F.; Nussbaum, R. L.: Mapping of the 75-kDa inositol polyphosphate-5-phosphatase (Inpp5b) to distal mouse chromosome 4 and its exclusion as a candidate gene for dysgenetic lens. *Genomics* 28:280–285, 1995.
- [69694] 8024.Ross, T. S.; Jefferson, A. B.; Mitchell, C. A.; Majerus, P. W.: Cloning and expression of human 75-kDa inositol polyphosphate-5-phosphatase. *J. Biol. Chem.* 266: 20283–20289, 1991.
- [69695] 8025.Matsumoto, M.; Nakagawa, T.; Inoue, T.; Nagata, E.; Tanaka, K.; Takano, H.; Minowa, O.; Kuno, J.; Sakakibara, S.; Yamada, M.; Yoneshima, H.; Miyawaki, A.; Fukuichi, T.;

Furuichi, T.; Okano, H.; Mikoshiba, K.; Noda, T.: Ataxia and epileptic seizures in mice lacking type 1 inositol

1,4,5-trisphosphate receptor. *Nature* 379: 168–171, 1996.

[69696] 8026. Nucifora, F. C., Jr.; Li, S.-H.; Danoff, S.; Ullrich, A.; Ross, C. A.: Molecular cloning of a cDNA for the human inositol 1,4,5-trisphosphate receptor type 1, and the identification of a third alternatively spliced variant. *Molec. Brain Res.* 32: 291–296, 1995.

[69697] 8027. Bradham, D. M.; Igarashi, A.; Potter, R. L.; Groten-dorst, G. R.: Connective tissue growth factor: a cysteine-rich mitogen secreted by human vascular endothelial cells is related to the SRC-induced immediate early gene product CEF-10. *J. Cell Biol.* 114: 1285–1294, 1991.

[69698] 8028. Aguilar-Salinas, C. A.; Reyes-Rodriguez, E.; Ordonez-Sanchez, M. L.; Torres, M. A.; Ramirez-Jimenez, S.; Dominguez-Lopez, A.; Martinez-Francois, J. R.; Velasco-Perez, M. L.; Alpizar, M.; Garcia-Garcia, E.; Gomez-Perez, F.; Rull, J.; Tusie-Luna, M. T.: Early-onset type 2 diabetes: metabolic and genetic characterization in the Mexican population. *J. Clin. Endocr. Metab.* 86: 220–226, 2001.

[69699] 8029. Der Kinderen, D. J.; Koten, J. W.; Tan, K. E. W. P.; Beemer, F. A.; Van Romunde, L. K. J.; Den Otter, W.:

Parental age in sporadic hereditary retinoblastoma. *Am. J. Ophthalmol.* 110: 605–609, 1990.

[69700] 8030. Orphanides, G.; Wu, W.-H.; Lane, W. S.; Hampsey, M.; Reinberg, D.: The chromatin-specific transcription elongation factor FACT comprises human SPT16 and SSRP1 proteins. *Nature* 400: 284–288, 1999.

[69701] 8031. Li, H.; Chen, J.; Huang, A.; Stinson, J.; Heldens, S.; Foster, J.; Dowd, P.; Gurney, A. L.; Wood, W. I.: Cloning and characterization of IL-17B and IL-17C, two new members of the IL-17 cytokine family. *Proc. Nat. Acad. Sci.* 97: 773–778, 2000.

[69702] 8032. Shi, Y.; Ullrich, S. J.; Zhang, J.; Connolly, K.; Grzegorzewski, K. J.; Barber, M. C.; Wang, W.; Wathen, K.; Hodge, V.; Fisher, C. L.; Olsen, H.; Ruben, S. M.; Knyazev, I.; Cho, Y. H.; Kao, V.; Wilkinson, K. A.; Carrell, J. A.; Ebner, R.: A novel cytokine receptor–ligand pair: identification, molecular characterization, and in vivo immunomodulatory activity. *J. Biol. Chem.* 275: 19167–19176, 2000.

[69703] 8033. Friedman, L. S.; Ostermeyer, E. A.; Lynch, E. D.; Welcsh, P.; Szabo, C. I.; Meza, J. E.; Anderson, L. A.; Dowd, P.; Lee, M. K.; Rowell, S. E.; Ellison, J.; Boyd, J.; King, M.-C.: 22 genes from chromosome 17q21: cloning, sequencing, and characterization of mutations in breast cancer families

and tumors. *Genomics* 25: 256–163, 1995.

[69704] 8034.Linial, M.; Miller, K.; Scheller, R. H.: VAT-1: an abundant membraneprotein from Torpedo cholinergic synaptic vesicles. *Neuron* 2: 1265–1273,1989.

[69705] 8035.Smith, T. M.; Lee, M. K.; Szabo, C. I.; Jerome, N.; McEuen, M.;Taylor, M.; Hood, L.; King, M.–C.: Complete genomic sequence andanalysis of 117 kb of human DNA containing the gene BRCA1. *GenomeRes.* 6: 1029–1049, 1996.

[69706] 8036.Johnstone, R. W.; Tommerup, N.; Hansen, C.; Vissing, H.; Shi, Y.: Structural organization, tissue expression, and chromosomal localizationof Ciao 1, a functional modulator of the Wilms' tumor suppressor,WT1. *Immunogenetics* 49: 900–905, 1999.

[69707] 8037.Johnstone, R. W.; Wang, J.; Tommerup, N.; Vissing, H.; Roberts,T.; Shi, Y.: Ciao 1 is a novel WD40 protein that interacts with thetumor suppressor protein WT1. *J. Biol. Chem.* 273: 10880–10887, 1998.

[69708] 8038.Huebner, K.; Cannizzaro, L. A.; Nakamura, T.; Hillova, J.; Mariage–Samson,R.; Hecht, F.; Hill, M.; Croce, C. M.: A rearranged transforminggene, tre, is made up of human sequences derived from chromosome regions5q, 17q and 18q. *Oncogene* 3: 449–455, 1988.

- [69709] 8039.Nakamura, T.; Hillova, J.; Mariage-Samson, R.; Hill, M.: Molecularcloning of a novel oncogene generated by DNA recombination duringtransfection. *Oncogene Res.* 2: 357–370, 1988.
- [69710] 8040.Nakamura, T.; Hillova, J.; Mariage-Samson, R.; Onno, M.; Huebner,K.; Cannizzaro, L. A.; Boghosian-Sell, L.; Croce, C. M.; Hill, M.: A novel transcriptional unit of the tre oncogene widely expressedin human cancer cells. *Oncogene* 7: 733–741, 1992.
- [69711] 8041.Chen, H.; Rossier, C.; Morris, M. A.; Scott, H. S.; Gos, A.; Bairoch,A.; Antonarakis, S. E.: A testis-specific gene, TPTE, encodes a putativetransmembrane tyrosine phosphatase and maps to the pericentromericregion of human chromosomes 21 and 13, and to chromosomes 15, 22,and Y. *Hum. Genet.* 105: 399–409, 1999.
- [69712] 8042.Guipponi, M.; Tapparel, C.; Jousson, O.; Scamuffa, N.; Mas, C.;Rossier, C.; Hutter, P.; Meda, P.; Lyle, R.; Raymond, A.; Antonarakis,S. E.: The murine orthologue of the Golgi-localized TPTE proteinprovides clues to the evolutionary history of the human TPTE genefamily. *Hum. Genet.* 109: 569–575, 2001.
- [69713] 8043.Guipponi, M.; Yaspo, M.–L.; Riesselman, L.; Chen, H.; De Sario,A.; Roizes, G.; Antonarakis, S. E.: Genomic

structure of a copy of the human TPTE gene which encompasses 87 kb on the short arm of chromosome 21. *Hum. Genet.* 107: 127–131, 2000.

- [69714] 8044. Gruber, A. D.; Pauli, B. U.: Molecular cloning and biochemical characterization of a truncated, secreted member of the human family of  $\text{Ca}^{2+}$ -activated  $\text{Cl}^-$  channels. *Biochim. Biophys. Acta* 1444: 418–423, 1999.
- [69715] 8045. Bause, E.; Bieberich, E.; Rolfs, A.; Volker, C.; Schmidt, B.: Molecular cloning and primary structure of Man(9)-mannosidase from human kidney. *Eur. J. Biochem.* 217: 535–540, 1993.
- [69716] 8046. Tremblay, L. O.; Campbell Dyke, N.; Herscovics, A.: Molecular cloning, chromosomal mapping and tissue-specific expression of a novel human  $\alpha$ -1,2-mannosidase gene involved in N-glycan maturation. *Glycobiology* 8: 585–595, 1998.
- [69717] 8047. Ueno, M.; Kimura, N.; Nakashima, K.; Saito-Ohara, F.; Inazawa, J.; Taga, T.: Genomic organization, sequence and chromosomal localization of the mouse *Tbr2* gene and a comparative study with *Tbr1*. *Gene* 254: 29–35, 2000.
- [69718] 8048. Baker, E.; Crawford, J.; Sutherland, G. R.; Freeman, C.; Parish, C. R.; Hulett, M. D.: Human HPA endoglycosidase heparanase. *Chromosome Res.* 7: 319 only, 1999.

- [69719] 8049.Hulett, M. D.; Freeman, C.; Hamdorf, B. J.; Baker, R. T.; Harris, M. J.; Parish, C. R.: Cloning of mammalian heparanase, an important enzyme in tumor invasion and metastasis. *Nature Med.* 5: 803–809, 1999.
- [69720] 8050.Kussie, P. H.; Hulmes, J. D.; Ludwig, D. L.; Patel, S.; Navarro, E. C.; Seddon, A. P.; Giorgio, N. A.; Bohlen, P.: Cloning and functional expression of a human heparanase gene. *Biochem. Biophys. Res. Commun.* 261:183–187, 1999.
- [69721] 8051.Toyoshima, M.; Nakajima, M.: Human heparanase: purification, characterization, cloning, and expression. *J. Biol. Chem.* 274: 24153–24160, 1999.
- [69722] 8052.Vlodavsky, I.; Friedmann, Y.; Elkin, M.; Aingorn, H.; Atzmon, R.; Ishai-Michaeli, R.; Bitan, M.; Pappo, O.; Peretz, T.; Michal, I.; Spector, L.; Pecker, I.: Mammalian heparanase: gene cloning, expression and function in tumor progression and metastasis. *Nature Med.* 5:793–802, 1999.
- [69723] 8053.Ikeda, A.; Ikeda, S.; Gridley, T.; Nishina, P. M.; Nagert, J.K.: Neural tube defects and neuroepithelial cell death in *Tulp3* knockout mice. *Hum. Molec. Genet.* 10: 1325–1334, 2001.
- [69724] 8054.Nishina, P. M.; North, M. A.; Ikeda, A.; Yan, Y.; Nag-

gert, J. K.: Molecular characterization of a novel tubby gene family member, TULP3, in mouse and humans. *Genomics* 54: 215–220, 1998.

[69725] 8055. Wu, X.; Kekuda, R.; Huang, W.; Fei, Y.-J.; Leibach, F. H.; Chen, J.; Conway, S. J.; Ganapathy, V.: Identity of the organic cation transporter OCT3 as the extraneuronal monoamine transporter (uptake-2) and evidence for the expression of the transporter in the brain. *J. Biol. Chem.* 273: 32776–32786, 1998.

[69726] 8056. Seki, N.; Ohira, M.; Nagase, T.; Ishikawa, K.; Miyajima, N.; Nakajima, D.; Nomura, N.; Ohara, O.: Characterization of cDNA clones in size-fractionated cDNA libraries from human brain. *DNA Res.* 4: 345–349, 1997.

[69727] 8057. Biassoni, R.; Cantoni, C.; Falco, M.; Verdiani, S.; Bottino, C.; Vitale, M.; Conte, R.; Poggi, A.; Moretta, A.; Moretta, L.: The human leukocyte antigen (HLA)-C-specific 'activatory' or 'inhibitory' natural killer cell receptors display highly homologous extracellular domains but differ in their transmembrane and intracytoplasmic portions. *J. Exp. Med.* 183: 645–650, 1996.

[69728] 8058. Bottino, C.; Sivori, S.; Vitale, M.; Cantoni, C.; Falco, M.; Pende, D.; Morelli, L.; Augugliaro, R.; Semenzato, G.; Biassoni, R.; Moretta, L.; Moretta, A.: A novel surface



molecule homologous to the p58/p50 family of receptors is selectively expressed on a subset of human natural killer cells and induces both triggering of cell functions and proliferation. *Europ. J. Immun.* 26: 1816–1824, 1996.

[69729] 8059. Sumoy, L.; Pluvinet, R.; Andreu, N.; Estivill, X.; Escarceller, M.: PACSIN 3 is a novel SH3 domain cytoplasmic adapter protein of the pacsin-syndapin-FAP52 gene family. *Gene* 262: 199–205, 2001.

[69730] 8060. Hubener, C.; Mincheva, A.; Lichter, P.; Schraven, B.; Bruyns, E.: Genomic organization and chromosomal localization of the human gene encoding the T-cell receptor-interacting molecule (TRIM). *Immunogenetics* 51: 154–158, 2000.

[69731] 8061. Hubener, C.; Mincheva, A.; Lichter, P.; Schraven, B.; Bruyns, E.: Complete sequence, genomic organization, and chromosomal localization of the human gene encoding the SHP2-interacting transmembrane adaptor protein (SIT). *Immunogenetics* 53: 337–341, 2001.

[69732] 8062. Bruyns, E.; Marie-Cardine, A.; Kirchgessner, H.; Sagolla, K.; Shevchenko, A.; Mann, M.; Autschbach, F.; Bensussan, A.; Meuer, S.; Schraven, B.: T cell receptor (TCR) interacting molecule (TRIM), a novel disulfide-linked dimer associated with the TCR-CD3-zeta complex, recruits in-

tracellular signaling proteins to the plasma membrane. J. Exp. Med. 188: 561–575, 1998.

- [69733] 8063. Zhao, R.; Qi, Y.; Chen, J.; Zhao, Z. J.: FYVE–DSP2, a FYVE domain–containing dual specificity protein phosphatase that dephosphorylates phosphatidylinositol(3)-phosphate. Exp. Cell Res. 265: 329–338, 2001.
- [69734] 8064. Cui, X.; De Vivo, I.; Slany, R.; Miyamoto, A.; Firestein, R.; Cleary, M. L.: Association of SET domain and myotubularin–related proteins modulates growth control. Nature Genet. 18: 331–337, 1998.
- [69735] 8065. Kahyo, T.; Nishida, T.; Yasuda, H.: Involvement of PIAS1 in the sumoylation of tumor suppressor p53. Molec. Cell 8: 713–718, 2001.
- [69736] 8066. Liu, B.; Liao, J.; Rao, X.; Kushner, S. A.; Chung, C. D.; Chang, D. D.; Shuai, K.: Inhibition of Stat1–mediated gene activation by PIAS1. Proc. Nat. Acad. Sci. 95: 10626–10631, 1998.
- [69737] 8067. Valdez, B. C.; Henning, D.; Perlaky, L.; Busch, R. K.; Busch, H.: Cloning and characterization of Gu/RH–II binding protein. Biochem. Biophys. Res. Commun. 234: 335–340, 1997.
- [69738] 8068. Mansour, S. J.; Herbrick, J.–A.; Scherer, S. W.; Melancon, P.: Human GBF1 is a ubiquitously expressed gene of

the Sec7 domain family mapping to 10q24. Genomics 54: 323–327, 1998.

[69739] 8069. Heisenberg, C.-P.; Tada, M.; Rauch, G.-J.; Saude, L.; Concha, M.L.; Geisler, R.; Stemple, D. L.; Smith, J. C.; Wilson, S. W.: Silberblick/Wnt11 mediates convergent extension movements during zebrafish gastrulation. Nature 405:76–81, 2000.

[69740] 8070. Lako, M.; Strachan, T.; Bullen, P.; Wilson, D. I.; Robson, S. C.; Lindsay, S.: Isolation, characterisation and embryonic expression of WNT11, a gene which maps to 11q13.5 and has possible roles in the development of skeleton, kidney and lung. Gene 219: 101–110, 1998.

[69741] 8071. Pandur, P.; Lasche, M.; Eisenberg, L. M.; Kuhl, M.: Wnt-11 activation of a non-canonical Wnt signalling pathway is required for cardiogenesis. Nature 418:636–641, 2002.

[69742] 8072. Frye, R. A.: Characterization of five human cDNAs with homology to the yeast SIR2 gene: Sir2-like proteins (sirtuins) metabolize NAD and may have protein ADP-ribosyltransferase activity. Biochem. Biophys. Res. Commun. 260: 273–279, 1999.

[69743] 8073. Imai, S.; Armstrong, C. M.; Kaeblerlein, M.; Guarente, L.: Transcriptional silencing and longevity protein Sir2 is an

NAD-dependent histone deacetylase. *Nature* 403:795–800, 2000.

- [69744] 8074. Kimura, A.; Umehara, T.; Horikoshi, M.: Chromosomal gradient of histone acetylation established by Sas2p and Sir2p functions as a shield against gene silencing. *Nature Genet.* 15 Oct.: , 2002. Note: Advance Electronic Publication.
- [69745] 8075. Aula, N.; Salomaki, P.; Timonen, R.; Verheijen, F.; Mancini, G.; Mansson, J.-E.; Aula, P.; Peltonen, L.: The spectrum of SLC17A5 gene mutations resulting in free sialic acid-storage diseases indicates some genotype-phenotype correlation. *Am. J. Hum. Genet.* 67: 832–840, 2000.
- [69746] 8076. Biancheri, R.; Verbeek, E.; Rossi, A.; Gaggero, R.; Roccatagliata, L.; Gatti, R.; van Diggelen, O.; Verheijen, F. W.; Mancini, G. M. S.: An Italian severe Salla disease variant associated with a SLC17A5 mutation earlier described in infantile sialic acid storage disease. *Clin. Genet.* 61: 443–447, 2002.
- [69747] 8077. Belinsky, M. G.; Bain, L. J.; Balsara, B. B.; Testa, J. R.; Kruh, G. D.: Characterization of MOAT-C and MOAT-D, new members of the MRP/cMOAT subfamily of transporter proteins. *J. Nat. Cancer Inst.* 90:1735–1741, 1998.

- [69748] 8078. Fromm, M. F.; Leake, B.; Roden, D. M.; Wilkinson, G. R.; Kim, R. B.: Human MRP3 transporter: identification of the 5-prime flanking region, genomic organization and alternative splice variants. *Biochim. Biophys. Acta* 1415: 369–374, 1999.
- [69749] 8079. Kiuchi, Y.; Suzuki, H.; Hirohashi, T.; Tyson, C. A.; Sugiyama, Y.: cDNA cloning and inducible expression of human multidrug resistance associated protein 3 (MRP3). *FEBS Lett.* 433: 149–152, 1998.
- [69750] 8080. König, J.; Rost, D.; Cui, Y.; Keppler, D.: Characterization of the human multidrug resistance protein isoform MRP3 localized to the basolateral hepatocyte membrane. *Hepatology* 29: 1156–1163, 1999.
- [69751] 8081. Kool, M.; van der Linden, M.; de Haas, M.; Scheffer, G. L.; de Vree, J. M. L.; Smith, A. J.; Jansen, G.; Peters, G. J.; Ponne, N.; Scheper, R. J.; Oude Elferink, R. P. J.; Baas, F.; Borst, P.: MRP3, an organic anion transporter able to transport anti-cancer drugs. *Proc. Nat. Acad. Sci.* 96: 6914–6919, 1999.
- [69752] 8082. Ortiz, D. F.; Li, S.; Iyer, R.; Zhang, X.; Novikoff, P.; Arias, I. M.: MRP3, a new ATP-binding cassette protein localized to the canalicular domain of the hepatocyte. *Am. J. Physiol.* 276: G1493–G1500, 1999.

- [69753] 8083.Hjalt, T. A.; Murray, J. C.: The human BARX2 gene: genomic structure, chromosomal localization, and single nucleotide polymorphisms. *Genomics* 62:456–459, 1999.
- [69754] 8084.Jones, F. S.; Kioussi, C.; Copertino, D. W.; Kallunki, P.; Holst, B. D.; Edelman, G. M.: Barx2, a new homeobox gene of the Bar class, is expressed in neural and craniofacial structures during development. *Proc. Nat. Acad. Sci.* 94: 2632–2637, 1997.
- [69755] 8085.Arking, D. E.; Krebsova, A.; Macek, M., Sr.; Macek, M., Jr.; Arking, A.; Mian, I. S.; Fried, L.; Hamosh, A.; Dey, S.; McIntosh, I.; Dietz, H. C.: Association of human aging with a functional variant of klotho. *Proc. Nat. Acad. Sci.* 99: 856–861, 2002.
- [69756] 8086.Koh, N.; Fujimori, T.; Nishiguchi, S.; Tamori, A.; Shiomi, S.; Nakatani, T.; Sugimura, K.; Kishimoto, T.; Kinoshita, S.; Kuroki, T.; Nabeshima, Y.: Severely reduced production of Klotho in human chronic renal failure kidney. *Biochem. Biophys. Res. Commun.* 280:1015–1020, 2001.
- [69757] 8087.Kuro-o, M.; Matsumura, Y.; Aizawa, H.; Kawaguchi, H.; Suga, T.; Utsugi, T.; Ohyama, Y.; Kurabayashi, M.; Kaname, T.; Kume, E.; Iwasaki, H.; Iida, A.; Shiraki-Iida, T.; Nishikawa, S.; Nagai, R.; Nabeshima, Y.: Mutation of the mouse klotho gene leads to a syndrome resembling age-

ing. Nature 390: 45–51, 1997.

- [69758] 8088. Matsumura, Y.; Aizawa, H.; Shiraki-Iida, T.; Nagai, R.; Kuro-o, M.; Nabeshima, Y.: Identification of the human klotho gene and its two transcripts encoding membrane and secreted klotho protein. Biochem. Biophys. Res. Commun. 242: 626–630, 1998.
- [69759] 8089. Mori, K.; Yahata, K.; Mukoyama, M.; Suganami, T.; Makino, H.; Nagae, T.; Masuzaki, H.; Ogawa, Y.; Sugawara, A.; Nabeshima, Y.; Nakao, K.: Disruption of klotho gene causes an abnormal energy homeostasis in mice. Biochem. Biophys. Res. Commun. 278: 665–670, 2000.
- [69760] 8090. Saito, Y.; Nakamura, T.; Ohyama, Y.; Suzuki, T.; Iida, A.; Shiraki-Iida, T.; Kuro-o, M.; Nabeshima, Y.; Kurabayashi, M.; Nagai, R.: In vivo klotho gene delivery protects against endothelial dysfunction in multiple risk factor syndrome. Biochem. Biophys. Res. Commun. 276: 767–772, 2000.
- [69761] 8091. Little, N. A.; Hastie, N. D.; Davies, R. C.: Identification of WTAP, a novel Wilms' tumour 1–associating protein. Hum. Molec. Genet. 9: 2231–2239, 2000.
- [69762] 8092. Vazquez, F.; Hastings, G.; Ortega, M.–A.; Lane, T. F.; Oikemus, S.; Lombardo, M.; Iruela-Arispe, M. L.: METH–1, a human ortholog of ADAMTS–1, and METH–2 are mem–

bers of a new family of proteins with angio-inhibitory activity. *J. Biol. Chem.* 274: 23349–23357, 1999.

- [69763] 8093. Georgiadis, K. E.; Hirohata, S.; Seldin, M. F.; Apte, S. S.: ADAM-TS8, a novel metalloprotease of the ADAM-TS family located on mouse chromosome 9 and human chromosome 11. *Genomics* 62: 312–315, 1999.
- [69764] 8094. Sugawara, M.; Nakanishi, T.; Fei, Y.-J.; Huang, W.; Ganapathy, M. E.; Leibach, F. H.; Ganapathy, V.: Cloning of an amino acid transporter with functional characteristics and tissue expression pattern identical to that of system A. *J. Biol. Chem.* 275: 16473–16477, 2000.
- [69765] 8095. Fedi, P.; Bafico, A.; Soria, A. N.; Burgess, W. H.; Miki, T.; Bottaro, D. P.; Kraus, M. H.; Aaronson, S. A.: Isolation and biochemical characterization of the human Dkk-1 homologue, a novel inhibitor of mammalian Wnt signaling. *J. Biol. Chem.* 274: 19465–19472, 1999.
- [69766] 8096. Krupnik, V. E.; Sharp, J. D.; Jiang, C.; Robison, K.; Chickering, T. W.; Amaravadi, L.; Brown, D. E.; Guyot, D.; Mays, G.; Leiby, K.; Chang, B.; Duong, T.; Goodearl, A. D. J.; Gearing, D. P.; Sokol, S. Y.; McCarthy, S. A.: Functional and structural diversity of the human Dickkopf gene family. *Gene* 238: 301–313, 1999.
- [69767] 8097. Roessler, E.; Du, Y.; Glinka, A.; Dutra, A.; Niehrs, C.;



Muenke,M.: The genomic structure, chromosome location, and analysis of thehuman DKK1 head inducer gene as a candidate for holoprosencephaly. Cytogenet.Cell Genet. 89: 220–224, 2000.

[69768] 8098.Gatignol, A.; Buckler–White, A.; Berkhout, B.; Jeang, K. T.: Characterizationof a human TAR RNA–binding protein that activates the HIV–1 LTR. Science 251:1597–1600, 1991.

[69769] 8099.Gatignol, A.; Duarte, M.; Daviet, L.; Chang, Y.–N.; Jeang, K.–T.: Sequential steps in Tat trans–activation of HIV–1 mediated throughcellular DNA, RNA, and protein binding factors. Gene Expr. 5: 217–228,1996.

[69770] 8100.Kozak, C. A.; Gatignol, A.; Graham, K.; Jeang, K. T.; McBride,O. W.: Genetic mapping in human and mouse of the locus encoding TRBP,a protein that binds the TAR region of the human immunodeficiencyvirus (HIV–1). Genomics 25: 66–72, 1995.

[69771] 8101.Bauer, M. F.; Gempel, K.; Reichert, A. S.; Rappold, G. A.; Lichtner,P.; Gerbitz, K. D.; Neupert, W.; Brunner, M.; Hofmann, S.: Geneticand structural characterization of the human mitochondrial inner membranetranslocase. J. Molec. Biol. 289: 69–82, 1999.

[69772] 8102.Bomer, U.; Rassow, J.; Zufall, N.; Pfanner, N.; Meijer,

M.; Maarse, A. C.: The preprotein translocase of the inner mitochondrial membrane: evolutionary conservation of targeting and assembly of Tim17. *J. Molec. Biol.* 262: 389–395, 1996.

[69773] 8103. Phornphutkul, C.; Anikster, Y.; Huizing, M.; Braun, P.; Brodie, C.; Chou, J. Y.; Gahl, W. A.: The promoter of a lysosomal membrane transporter gene, CTNS, binds Sp-1, shares sequences with the promoter of an adjacent gene, CARKL, and causes cystinosis if mutated in a critical region. *Am. J. Hum. Genet.* 69: 712–721, 2001.

[69774] 8104. Lebre, A.-S.; Jamot, L.; Takahashi, J.; Spasskey, N.; Leprince, C.; Ravise, N.; Zander, Fujigasaki, H.; Kussel-Andermann, P.; Duyckaerts, C.; Camonis, J. H.; Brice, A.: Ataxin-7 interacts with a Cbl-associated protein that it recruits into neuronal intranuclear inclusions. *Hum. Molec. Genet.* 10: 1201–1213, 2001.

[69775] 8105. Lin, W.-H.; Chiu, K. C.; Chang, H.-M.; Lee, K.-C.; Tai, T.-Y.; Chuang, L.-M.: Molecular scanning of the human sorbin and SH3-domain-containing-1 (SORBS1) gene: positive association of the T228A polymorphism with obesity and type 2 diabetes. *Hum. Molec. Genet.* 10: 1753–1760, 2001.

[69776] 8106. Scott, A. F.: Personal Communication. Baltimore, Md.

9/13/2000.

- [69777] 8107. Nishi, M.; Mizushima, A.; Nakagawara, K.; Takeshima, H.: Characterization of human junctophilin subtype genes. *Biochem. Biophys. Res. Commun.* 273:920–927, 2000.
- [69778] 8108. Takeshima, H.; Komazaki, S.; Nishi, M.; Iino, M.; Kangawa, K.: Junctophilins: a novel family of junctional membrane complex proteins. *Molec. Cell* 6: 11–22, 2000.
- [69779] 8109. Holmes, S. E.; O'Hearn, E.; Rosenblatt, A.; Callahan, C.; Hwang, H. S.; Ingersoll–Ashworth, R. G.; Fleisher, A.; Stevanin, G.; Brice, A.; Potter, N. T.; Ross, C. A.; Margolis, R. L.: A repeat expansion in the gene encoding junctophilin–3 is associated with Huntington disease–like 2. *Nature Genet.* 29: 377–378, 2001. Note: Erratum: *Nature Genet.* 30: 123 only, 2002.
- [69780] 8110. Chavez, R. A.; Gray, A. T.; Zhao, B. B.; Kindler, C. H.; Mazurek, M. J.; Mehta, Y.; Forsayeth, J. R.; Yost, C. S.: TWIK–2, a new weak inward rectifying member of the tandem pore domain potassium channel family. *J. Biol. Chem.* 274: 7887–7892, 1999.
- [69781] 8111. Gray, A. T.; Kindler, C. H.; Sampson, E. R.; Yost, C. S.: Assignment of KCNK6 encoding the human weak inward rectifier potassium channel TWIK–2 to chromosome

band 19q13.1 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 84: 190–191, 1999.

[69782] 8112. Pountney, D. J.; Gulkarov, I.; Vega-Saenz de Miera, E.; Holmes, D.; Saganich, M.; Rudy, B.; Artman, M.; Coetzee, W. A.: Identification and cloning of TWIK-originated similarity sequence (TOSS): a novel human 2-pore K(+) channel principal subunit. *FEBS Lett.* 450: 191–196, 1999.

[69783] 8113. Salinas, M.; Reyes, R.; Lesage, F.; Fosset, M.; Heurteaux, C.; Romey, G.; Lazdunski, M.: Cloning of a new mouse two-P domain channel subunit and a human homologue with a unique pore structure. *J. Biol. Chem.* 274: 11751–11760, 1999.

[69784] 8114. Barbetti, F.; Rocchi, M.; Bossolasco, M.; Cordera, R.; Sbraccia, P.; Finelli, P.; Consalez, G. G.: The human skeletal muscle glycogenin gene: cDNA, tissue expression, and chromosomal localization. *Biochem. Biophys. Res. Commun.* 220: 72–77, 1996.

[69785] 8115. Machesky, L. M.; Reeves, E.; Wientjes, F.; Mattheyse, F. J.; Grogan, A.; Totty, N. F.; Burlingame, A. L.; Hsuan, J. J.; Segal, A. W.: Mammalian actin-related protein 2/3 complex localizes to regions of lamellipodial protrusion and is composed of evolutionarily conserved proteins. *Biochem. J.* 328: 105–112, 1997.

- [69786] 8116.Luo, J.; Nikolaev, A. Y.; Imai, S.; Chen, D.; Su, F.; Shiloh, A.;Guarente, L.; Gu, W.: Negative control of p53 by Sir2- $\alpha$  promotes cell survival under stress. *Cell* 107: 137–148, 2001.
- [69787] 8117.Bulfone, A.; Smiga, S. M.; Shimamura, K.; Peterson, A.; Puellas,L.; Rubenstein, J. L. R.: T-brain-1: a homolog of Brachyury whose expression defines molecularly distinct domains within the cerebral cortex. *Neuron* 15: 63–78, 1995.
- [69788] 8118.Makoff, A.; Lelchuk, R.; Ozer, M.; Harrington, K.; Emson, P.:Molecular characterization and localization of human metabotropic glutamate receptor type 4. *Molec. Brain Res.* 37: 239–248, 1996.
- [69789] 8119.Pekhletski, R.; Gerlai, R.; Overstreet, L. S.; Huang, X. P.; Agopyan,N.; Slater, N. T.; Abramow-Newerly, W.; Roder, J. C.; Hampson, D.R.: Impaired cerebellar synaptic plasticity and motor performance in mice lacking the mGluR4 subtype of metabotropic glutamate receptor. *J.Neurosci.* 16: 6364–6373, 1996.
- [69790] 8120.Barbon, A.; Ferraboli, S.; Barlati, S.: Assignment of the human metabotropic glutamate receptor gene GRM7 to chromosome 3p26.1–p25.2 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 88: 288 only, 2000.

- [69791] 8121.Shore, D.; Squire, M.; Nasmyth, K. A.: Characterization of two genes required for the position-effect control of yeast mating-type genes. *EMBO J.* 3: 2817–2823, 1984.
- [69792] 8122.Suka, N.; Luo, K.; Grunstein, M.: Sir2p and Sas2p oppositely regulate acetylation of yeast histone H4 lysine16 and spreading of heterochromatin. *Nature Genet.* 15 Oct.: , 2002. Note: Advance Electronic Publication.
- [69793] 8123.Tanny, J. C.; Dowd, G. J.; Huang, J.; Hilz, H.; Moazed, D.: An enzymatic activity in the yeast Sir2 protein that is essential for gene silencing. *Cell* 99: 735–745, 1999.
- [69794] 8124.Vaziri, H.; Dessain, S. K.; Eaton, E. N.; Imai, S.-I.; Frye, R.A.; Pandita, T. K.; Guarente, L.; Weinberg, R. A.: hSIR2–SIRT1 functions as an NAD-dependent p53 deacetylase. *Cell* 107: 149–159, 2001.
- [69795] 8125.Verhaagh, S.; Schweifer, N.; Barlow, D. P.; Zwart, R.: Cloning of the mouse and human solute carrier 22a3 (Slc22a3/SLC22A3) identifies a conserved cluster of three organic cation transporters on mouse chromosome 17 and human 6q26–q27. *Genomics* 55: 209–218, 1999.
- [69796] 8126.Raich, N.; Mattei, M. G.; Romeo, P. H.; Beaupain, D.: PHTF, a novel atypical homeobox gene on chromosome 1p13, is evolutionarily conserved. *Genomics* 59: 108–109, 1999.

- [69797] 8127.Liu, J.; Shworak, N. W.; Sinay, P.; Schwartz, J. J.; Zhang, L.;Fritze, L. M.; Rosenberg, R. D.: Expression of heparan sulfate D-glucosaminyl3-O-sulfotransferase isoforms reveals novel substrate specificities. J.Biol. Chem. 274: 5185–5192, 1999.
- [69798] 8128.Shworak, N. W.; Liu, J.; Petros, L. M.; Zhang, L.; Kobayashi, M.;Copeland, N. G.; Jenkins, N. A.; Rosenberg, R. D.: Multiple isoformsof heparan sulfate D-glucosaminyl 3-O-sulfotransferase: isolation,characterization, and expression of human cDNAs and identificationof distinct genomic loci. J. Biol. Chem. 274: 5170–5184, 1999.
- [69799] 8129.McIlhatton, M. A.; Burrows, J. F.; Donaghy, P. G.; Chanduloy, S.;Johnston, P. G.; Russell, S. E. H.: Genomic organization, complexsplicing pattern and expression of a human septin gene on chromosome17q25.3. Oncogene 20: 5930–5939, 2001.
- [69800] 8130.Osaka, M.; Rowley, J. D.; Zeleznik-Le, N. J.: MSF (MLL septin-likefusion), a fusion partner gene of MLL, in a therapy-related acutemyeloid leukemia with a t(11;17)(q23;q25). Proc. Nat. Acad. Sci. 96:6428–6433, 1999.
- [69801] 8131.Russell, S. E. H.; McIlhatton, M. A.; Burrows, J. F.; Donaghy,P. G.; Chanduloy, S.; Petty, E. M.; Kalikin, L. M.;

Church, S. W.; McIlroy, S.; Harkin, D. P.; Keilty, G. W.; Cranston, A. N.; Weissenbach, J.; Hickey, I.; Johnston, P. G.: Isolation and mapping of a human septin gene to a region on chromosome 17q, commonly deleted in sporadic epithelial ovarian tumors. *Cancer Res.* 60: 4729–4734, 2000.

[69802] 8132. Karpinski, B. A.; Morle, G. D.; Huggenvik, J.; Uhler, M. D.; Leiden, J. M.: Molecular cloning of human CREB-2: an ATF/CREB transcription factor that can negatively regulate transcription from the cAMP response element. *Proc. Nat. Acad. Sci.* 89: 4820–4824, 1992.

[69803] 8133. Kasukabe, T.; Kobayashi, H.; Kaneko, Y.; Okabe-Kado, J.; Honma, Y.: Identity of human normal counterpart (MmTRA1b) of mouse leukemogenesis-associated gene (MmTRA1a) product as a plasma membrane phospholipid scramblase and chromosome mapping of the human Mm-TRA1b/phospholipid scramblase gene. *Biochem. Biophys. Res. Commun.* 249: 449–455, 1998.

[69804] 8134. Zhou, Q.; Sims, P. J.; Wiedmer, T.: Identity of a conserved motif in phospholipid scramblase that is required for  $\text{Ca}^{2+}$ -accelerated transbilayer movement of membrane phospholipids. *Biochemistry* 37: 2356–2360, 1998.

[69805] 8135. Zhou, Q.; Zhao, J.; Stout, J. G.; Luhm, R. A.; Wiedmer,



T.; Sims, P. J.: Molecular cloning of human plasma membrane phospholipid scramblase: a protein mediating trans-bilayer movement of plasma membrane phospholipids. *J. Biol. Chem.* 272: 18240–18244, 1997.

[69806] 8136. Robinson, R. C.; Turbedsky, K.; Kaiser, D. A.; Marchand, J.-B.; Higgs, H. N.; Choe, S.; Pollard, T. D.: Crystal structure of Arp2/3 complex. *Science* 294: 1679–1684, 2001.

[69807] 8137. Volkmann, N.; Amann, K. J.; Stoilova-McPhie, S.; Egile, C.; Winter, D. C.; Hazelwood, L.; Heuser, J. E.; Li, R.; Pollard, T. D.; Hanein, D.: Structure of Arp2/3 complex in its activated state and in actin filament branch junctions. *Science* 293: 2456–2459, 2001.

[69808] 8138. Welch, M. D.; DePace, A. H.; Verma, S.; Iwamatsu, A.; Mitchison, T. J.: The human Arp2/3 complex is composed of evolutionarily conserved subunits and is localized to cellular regions of dynamic actin filament assembly. *J. Cell Biol.* 138: 375–384, 1997.

[69809] 8139. Michel, J. J.; Xiong, Y.: Human CUL-1, but not other cullin family members, selectively interacts with SKP1 to form a complex with SKP2 and cyclin A. *Cell Growth Differ.* 9: 435–449, 1998.

[69810] 8140. Du, M.; Sansores-Garcia, L.; Zu, Z.; Wu, K. K.:

Cloning and expression analysis of a novel salicylate suppressible gene, Hs-CUL-3, a member of cullin/Cdc53 family. *J. Biol. Chem.* 273: 24289–24292, 1998.

[69811] 8141. Bao, S.; Shen, X.; Shen, K.; Liu, Y.; Wang, X.-F.: The mammalian Rad24 homologous to yeast *Saccharomyces cerevisiae* Rad24 and *Schizosaccharomyces pombe* Rad17 is involved in DNA damage checkpoint. *Cell Growth Diff.* 9:961–967, 1998.

[69812] 8142. Bao, S.; Chang, M.-S.; Auclair, D.; Sun, Y.; Wang, Y.; Wong, W.-K.; Zhang, J.; Liu, Y.; Qian, X.; Sutherland, R.; Magi-Galluzi, C.; Weisberg, E.; Cheng, E. Y. S.; Hao, L.; Sasaki, H.; Campbell, M. S.; Kraeft, S.-K.; Loda, M.; Lo, K.-M.; Chen, L. B.: Hrad17, a human homologue of the *Schizosaccharomyces pombe* checkpoint gene rad17, is overexpressed in colon carcinoma. *Cancer Res.* 59: 2023–2028, 1999.

[69813] 8143. Bluysen, H. A. R.; Naus, N. C.; van Os, R. I.; Jaspers, I.; Hoeijmakers, J. H. J.; de Klein, A.: Human and mouse homologues of the *Schizosaccharomyces pombe* rad17+ cell cycle checkpoint control gene. *Genomics* 55: 219–228, 1999.

[69814] 8144. Loijens, J. C.; Anderson, R. A.: Type I phosphatidylinositol-4-phosphate 5-kinases are distinct members of

this novel lipid kinase family. J.Biol. Chem. 271:  
32937–32943, 1996.

- [69815] 8145.Xie, Y.; Zhu, L.; Zhao, G.: Assignment of type I phosphatidylinositol-4-phosphate5-kinase (PIP5K1A) to human chromosome bands 1q22–q24 by in situ hybridization. Cytogenet.Cell Genet. 88: 197–199, 2000.
- [69816] 8146.Seelig, H. P.; Moosbrugger, I.; Ehrfeld, H.; Fink, T.; Renz, M.;Genth, E.: The major dermatomyositis-specific Mi-2 autoantigen isa presumed helicase involved in transcriptional activation. ArthritisRheum. 38: 1389–1399, 1995.
- [69817] 8147.Wade, P. A.; Geronne, A.; Jones, P. L.; Ballestar, E.; Aubry, F.;Wolffe, A. P.: Mi-2 complex couples DNA methylation to chromatinremodelling and histone deacetylation. Nature Genet. 23: 62–66,1999.
- [69818] 8148.Zhang, Y.; LeRoy, G.; Seelig, H.–P.; Lane, W. S.; Reinberg, D.: The dermatomyositis-specific autoantigen Mi2 is a component of acomplex containing histone deacetylase and nucleosome remodeling activities. Cell 95:279–289, 1998.
- [69819] 8149.Aihara, T.; Miyoshi, Y.; Koyama, K.; Suzuki, M.; Takahashi, E.;Monden, M.; Nakamura, Y.: Cloning and mapping of SMARCA5 encodinghSNF2H, a novel human

homologue of Drosophila ISWI. Cytogenet. CellGenet. 81: 191–193, 1998.

- [69820] 8150.Bochar, D. A.; Savard, J.; Wang, W.; Lafleur, D. W.; Moore, P.;Cote, J.; Shiekhattar, R.: A family of chromatin remodeling factorsrelated to Williams syndrome transcription factor. Proc. Nat. Acad.Sci. 97: 1038–1043, 2000.
- [69821] 8151.Bozhenok, L.; Wade, P. A.; Varga–Weisz, P.: WSTF–ISWI chromatinremodeling complex targets heterochromatic replication foci. EMBOJ. 21: 2231–2241, 2002.
- [69822] 8152.LeRoy, G.; Orphanides, G.; Lane, W. S.; Reinberg, D.: Requirementof RSF and FACT for transcription of chromatin templates in vitro. Science 282:1900–1904, 1998.
- [69823] 8153.Poot, R. A.; Dellaire, G.; Hulsmann, B. B.; Grimaldi, M. A.; Corona,D. F. V.; Becker, P. B.; Bickmore, W. A.; Varga–Weisz, P. D.: HuCHRAC,a human ISWI chromatin remodelling complex contains hACF1 and twonovel histone-fold proteins. EMBO J. 19: 3377–3387, 2000.
- [69824] 8154.Alberati–Giani, D.; Cesura, A. M.; Broger, C.; Warren, W. D.; Rover,S.; Malherbe, P.: Cloning and functional expression of human kynurenine3–monooxygenase. FEBS Lett. 410: 407–412, 1997.
- [69825] 8155.Bounous, G.; Kongshavn, P. A. L.; Taveroff, A.; Gold, P.: Evolutionarytraits in human milk proteins. Med. Hy–

potheses 27: 133–140, 1988.

- [69826] 8156.Davies, M. S.; West, L. F.; Davis, M. B.; Povey, S.; Craig, R.K.: The gene for human alpha-lactalbumin is assigned to chromosome12q13. *Ann. Hum. Genet.* 51: 183–188, 1987.
- [69827] 8157.Dayhoff, M. O.: *Atlas of Protein Sequence and Structure. Lactalbuminand Lysozymes.* Washington: National Biomedical Research Foundation(pub.) 5: 1972. Pp. D133–D140.
- [69828] 8158.Hakansson, A.; Zhivotovsky, B.; Orrenius, S.; Sabharwal, H.; Svanborg,C.: Apoptosis induced by a human milk protein. *Proc. Nat. Acad.Sci.* 92: 8064–8068, 1995.
- [69829] 8159.Hall, L.; Davies, M. S.; Craig, R. K.: The construction, identificationand characterisation of plasmids containing human alpha-lactalbumincDNA sequences. *Nucleic Acids Res.* 9: 65–84, 1981.
- [69830] 8160.Hall, L.; Emery, D. C.; Davies, M. S.; Parker, D.; Craig, R. K.: Organization and sequence of the human alpha-lactalbumin gene. *Biochem.J.* 242: 735–742, 1987.
- [69831] 8161.Stacey, A.; Schnieke, A.; Kerr, M.; Scott, A.; McKee, C.; Cottingham,I.; Binas, B.; Wilde, C.; Colman, A.: Lactation is disrupted by alpha-lactalbumindeficiency and can be restored by human alpha-lactalbumin gene replace-

mentin mice. *Proc. Nat. Acad. Sci.* 92: 2835–2839, 1995.

[69832] 8162. Boyer, S. H.; Fainer, D. C.; Watson–Williams, E. J.: Lactate dehydrogenase variant from human blood: evidence for molecular subunits. *Science* 141:642–643, 1963.

[69833] 8163. Ben–Neriah, Y.; Bauskin, A. R.: Leukocytes express a novel gene encoding a putative transmembrane protein–kinase devoid of an extracellular domain. (Letter) *Nature* 333: 672–676, 1988.

[69834] 8164. Krolewski, J. J.; Lee, R.; Eddy, R.; Shows, T. B.; Dalla–Favera, R.: Identification and chromosomal mapping of new human tyrosine kinase genes. *Oncogene* 5: 277–282, 1990.

[69835] 8165. Liao, X.; Zhou, R.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.: Receptor tyrosine kinase gene Tyro3 maps to mouse chromosome 2, closely linked to Ltk. *Mammalian Genome* 7: 395–396, 1996.

[69836] 8166. Toyoshima, H.; Kozutsumi, H.; Maru, Y.; Hagiwara, K.; Furuya, A.; Mioh, H.; Hanai, N.; Takaku, F.; Yazaki, Y.; Hirai, H.: Differently spliced cDNAs of human leukocyte tyrosine kinase receptor tyrosine kinase predict receptor proteins with and without a tyrosine kinase domain and a soluble receptor protein. *Proc. Nat. Acad. Sci.*

90:5404–5408, 1993.

- [69837] 8167.Knobeloch, K.–P.; Wright, M. D.; Ochsenbein, A. F.; Liesenfeld,O.; Lohler, J.; Zinkernagel, R. M.; Horak, I.; Orinska, Z.: Targetedinactivation of the tetraspanin CD37 impairs T–cell–dependent B–cellresponse under suboptimal costimulatory conditions. *Molec. Cell.Biol.* 20: 5363–5369, 2000.
- [69838] 8168.Virtaneva, K. I.; Angelisova, P.; Baumruker, T.; Horejsi, V.; Nevanlinna,H.; Schroder, J.: The genes for CD37, CD53, and R2, all members ofa novel gene family, are located on different chromosomes. *Immunogenetics* 37:461–465, 1993.
- [69839] 8169.Delmas, B.; Gelfi, J.; L'Haridon, R.; Vogel, L. K.; Sjostrom, H.;Noren, O.; Laude, H.: Aminopeptidase N is a major receptor for theenteropathogenic coronavirus TGEV. *Nature* 357: 417–420, 1992.
- [69840] 8170.Kruse, T. A.; Bolund, L.; Grzeschik, K.–H.; Ropers, H. H.; Olsen,J.; Sjostrom, H.; Noren, O.: Assignment of the human aminopeptidaseN (peptidase E) gene to chromosome 15q13–qter. *FEBS Lett.* 239: 305–308,1988.
- [69841] 8171.Look, A. T.; Ashmun, R. A.; Shapiro, L. H.; Peiper, S. C.: Humanmyeloid plasma membrane glycoprotein CD13 (gp150) is identical toaminopeptidase N. *J. Clin. Invest.*

83: 1299–1307, 1989.

[69842] 8172.Look, A. T.; Peiper, S. C.; Rebentisch, M. B.; Ashmun, R. A.; Roussel, M. F.; Lemons, R. S.; Le Beau, M. M.; Rubin, C. M.; Sherr, C. J.:Molecular cloning, expression, and chromosomal localization of the gene encoding a human myeloid membrane antigen (gp150). *J. Clin. Invest.* 78: 914–921, 1986.

[69843] 8173.Watt, V. M.; Willard, H. F.: The human aminopeptidase N gene:isolation, chromosome localization, and DNA polymorphism analysis. *Hum.Genet.* 85: 651–654, 1990.

[69844] 8174.Yeager, C. L.; Ashmun, R. A.; Williams, R. K.; Cardelli, C.B.; Shapiro, L. H.; Look, A. T.; Holmes, K. V.: Human aminopeptidase N is a receptor for human coronavirus 229E. *Nature* 357: 420–422, 1992.

[69845] 8175.Yoshimura, K.; Toibana, A.; Nakahama, K.: Human lysozyme: sequencing of a cDNA, and expression and secretion by *Saccharomyces cerevisiae*. *Biochem.Biophys. Res. Commun.* 150: 794–801, 1988.

[69846] 8176.Salmikangas, P.; Mykkanen, O.-M.; Gronholm, M.; Heiska, L.; Kere, J.; Carpen, O.: Myotilin, a novel sarcomeric protein with two Ig-like domains, is encoded by a candidate gene for limb-girdle muscular dystrophy. *Hum.Molec. Genet.* 8: 1329–1336, 1999.



- [69847] 8177. Hauser, M. A.; Horrigan, S. K.; Salmikangas, P.; Torian, U. M.; Viles, K. D.; Dancel, R.; Tim, R. W.; Taivainen, A.; Bartoloni, L.; Gilchrist, J. M.; Stajich, J. M.; Gaskell, P. C.; Gilbert, J. R.; Vance, J. M.; Pericak-Vance, M. A.; Carpen, O.; Westbrook, C. A.; Speer, M. C.: Myotilin is mutated in limb girdle muscular dystrophy 1A. *Hum. Molec. Genet.* 9: 2141–2147, 2000.
- [69848] 8178. Riegman, P. H. J.; Vlietstra, R. J.; van der Korput, J. A. G. M.; Romijn, J. C.; Trapman, J.: Characterization of the prostate-specific antigen gene: a novel human kallikrein-like gene. *Biochem. Biophys. Res. Commun.* 159: 95–102, 1989.
- [69849] 8179. Santoro, M.; Carlomagno, F.; Hay, I. D.; Herrmann, M. A.; Grieco, M.; Melillo, R.; Pierotti, M. A.; Bongarzone, I.; Della Porta, G.; Berger, N.; Peix, J. L.; Paulin, C.; Fabien, N.; Vecchio, G.; Jenkins, R. B.; Fusco, A.: Ret oncogene activation in human thyroid neoplasms is restricted to the papillary cancer subtype. *J. Clin. Invest.* 89: 1517–1522, 1992.
- [69850] 8180. Santoro, M.; Carlomagno, F.; Romano, A.; Bottaro, D. P.; Dathan, N. A.; Grieco, M.; Fusco, A.; Vecchio, G.; Matoskova, B.; Kraus, M. H.; Di Fiore, P. P.: Activation of RET as a dominant transforming gene by germline mutations of MEN2A and MEN2B. *Science* 267: 381–383, 1995.

- [69851] 8181.Schuchardt, A.; D'Agati, V.; Larsson-Blomberg, L.; Costantini,F.; Pachnis, V.: Defects in the kidney and enteric nervous system of mice lacking the tyrosine kinase receptor Ret. *Nature* 367: 380–383,1994.
- [69852] 8182.Shirahama, S.; Ogura, K.; Takami, H.; Ito, K.; Tohsen, T.; Miyauchi,A.; Nakamura, Y.: Mutational analysis of the RET proto-oncogene in 71 Japanese patients with medullary thyroid carcinoma. *J. Hum. Genet.* 43:101–106, 1998.
- [69853] 8183.Seri, M.; Yin, L.; Barone, A.; Bolino, A.; Celli, I.; Bocciardi,R.; Pasini, B.; Ceccherini, I.; Lerone, M.; Kristoffersson, U.; Larsson,L. T.; Casasa, J. M.; Cass, D. T.; Abramowicz, M. J.; Vanderwinden,J.-M.; Kravcenkiene, I.; Baric, I.; Silengo, M.; Martucciello, G.;Romeo, G.: Frequency of RET mutations in long- and short-segment Hirschsprung disease. *Hum. Mutat.* 9: 243–249, 1997.
- [69854] 8184.Takahashi, M.; Buma, Y.; Hiai, H.: Isolation of ret proto-oncogene cDNA with an amino-terminal signal sequence. *Oncogene* 4: 805–806,1989.
- [69855] 8185.Takahashi, M.; Buma, Y.; Iwamoto, T.; Inaguma, Y.; Ikeda, H.;Hiai, H.: Cloning and expression of the ret proto-oncogene encoding a tyrosine kinase with two po-

tential transmembrane domains. *Oncogene* 3:571–578, 1988.

[69856] 8186. Takahashi, M.; Ritz, J.; Cooper, G. M.: Activation of a novel human transforming gene, *ret*, by DNA rearrangement. *Cell* 42: 581–588, 1985.

[69857] 8187. Tessitore, A.; Sinisi, A. A.; Pasquali, D.; Cardone, M.; Vitale, D.; Bellastella, A.; Colantuoni, V.: A novel case of multiple endocrineneoplasia type 2A associated with two *de novo* mutations of the *RET* protooncogene. *J. Clin. Endocr. Metab.* 84: 3522–3527, 1999.

[69858] 8188. van Heyningen, V.: One gene—four syndromes. *Nature* 367: 319–320, 1994.

[69859] 8189. Xue, F.; Yu, H.; Maurer, L. H.; Memoli, V. A.; Nutile-McMenemey, N.; Schuster, M. K.; Bowden, D. W.; Mao, J.; Noll, W. W.: Germline *RET* mutations in MEN 2A and FMTC and their detection by simple DNA diagnostic tests. *Hum. Molec. Genet.* 3: 635–638, 1994.

[69860] 8190. Yin, L.; Ceccherini, I.; Pasini, B.; Matera, I.; Bicocchi, M. P.; Barone, V.; Bocciardi, R.; Kaariainen, H.; Weber, D.; Devoto, M.; Romeo, G.: Close linkage with the *RET* protooncogene and boundaries of deletion mutations in autosomal dominant Hirschsprung disease. *Hum. Molec. Genet.* 2: 1803–1808, 1993.

- [69861] 8191.Feller, S. M.; Ren, R. B.; Hanafusa, H.; Baltimore, D.: SH2 and SH3 domains as molecular adhesives: the interactions of crk and abl. *Trends Biochem. Sci.* 19: 453–458, 1994.
- [69862] 8192.Fioretos, T.; Heisterkamp, N.; Groffen, J.; Benjes, S.; Morris, C.: CRK proto-oncogene maps to human chromosome band 17p13. *Oncogene* 8:2853–2855, 1993.
- [69863] 8193.Matsuda, M.; Tanaka, S.; Nagata, S.; Kojima, A.; Kurata, T.; Shibuya, M.: Two species of human CRK cDNA encode proteins with distinct biological activities. *Molec. Cell. Biol.* 12: 3482–3489, 1992.
- [69864] 8194.Reichman, C. T.; Mayer, B. J.; Keshav, S.; Hanafusa, H.: The product of the cellular crk gene consists primarily of SH2 and SH3 regions. *Cell Growth Differ.* 3: 451–460, 1992.
- [69865] 8195.Schuuring, E.; Verhoeven, E.; Mooi, W. J.; Michalides, R. J. A.M.: Identification and cloning of two overexpressed genes, U21B31/PRAD1 and EMS1, within the amplified chromosome 11q13 region in human carcinomas. *Oncogene* 7:355–361, 1992.
- [69866] 8196.van Damme, H.; Brok, H.; Schuuring-Scholtes, E.; Schuuring, E.: The redistribution of cortactin into cell-matrix contact sites in human carcinoma cells with 11q13

amplification is associated with both overexpression and post-translational modification. *J. Biol.Chem.* 272: 7374–7380, 1997.

- [69867] 8197. Benz-Lemoine, E.; Brizard, A.; Huret, J.-L.; Babin, P.; Guilhot, F.; Couet, D.; Tanzer, J.: Malignant histiocytosis: a specific t(2;5)(p23;q35) translocation? Review of the literature. *Blood* 72: 1045–1047, 1988.
- [69868] 8198. Boulton, J.; Rack, K.; Kelly, S.; Madden, J.; Sakaguchi, A. Y.; Wang, L.-M.; Oscier, D. G.; Buckle, V. J.; Wainscoat, J. S.: Loss of both CSF1R (FMS) alleles in patients with myelodysplasia and chromosome 5 deletion. *Proc. Nat. Acad. Sci.* 88: 6176–6180, 1991.
- [69869] 8199. De Qi Xu; Guilhot, S.; Galibert, F.: Restriction fragment length polymorphism of the human c-fms gene. *Proc. Nat. Acad. Sci.* 82: 2862–2865, 1985.
- [69870] 8200. Eccles, M. R.: Genes encoding the platelet-derived growth factor (PDGF) receptor and colony-stimulating factor 1 (CSF-1) receptor are physically associated in mice as in humans. *Gene* 108: 285–288, 1991.
- [69871] 8201. Gisselbrecht, S.; Fichelson, S.; Sola, B.; Bordereaux, D.; Hampe, A.; Andre, C.; Galibert, F.; Tambourin, P. E.: Frequent c-fms activation by proviral insertion in mouse myeloblastic leukaemias. *Nature* 329: 259–261, 1987.

- [69872] 8202.Groffen, J.; Heisterkamp, N.; Spurr, N. K.; Dana, S. L.; Wasmuth, J. J.; Stephenson, J. R.: Regional assignment of the human c-fms oncogene to band q34 of chromosome 5. (Abstract) Cytogenet. Cell Genet. 37: 484 only, 1984.
- [69873] 8203.Hampe, A.; Gobet, M.; Sherr, C. J.; Galibert, F.: Nucleotide sequence of the feline retroviral oncogene v-fms shows unexpected homology with oncogenes encoding tyrosine-specific protein kinases. Proc. Nat. Acad. Sci. 81: 85-89, 1984.
- [69874] 8204.Hampe, A.; Shamon, B.-M.; Gobet, M.; Sherr, C. J.; Galibert, F.: Nucleotide sequence and structural organization of the human FMS proto-oncogene. Oncogene Res. 4: 9-17, 1989.
- [69875] 8205.How, G.-F.; Venkatesh, B.; Brenner, S.: Conserved linkage between the puffer fish (*Fugu rubripes*) and human genes for platelet-derived growth factor receptor and macrophage colony-stimulating factor receptor. Genome Res. 6: 1185-1191, 1996.
- [69876] 8206.Hibbs, M. L.; Tarlinton, D. M.; Armes, J.; Grail, D.; Hodgson, G.; Maglitta, R.; Stacker, S. A.; Dunn, A. R.: Multiple defects in the immune system of Lyn-deficient mice, culminating in autoimmune disease. Cell 83: 301-311, 1995.

- [69877] 8207. Webster, M. A.; Cardiff, R. D.; Muller, W. J.: Induction of mammary epithelial hyperplasias and mammary tumors in transgenic mice expressing a murine mammary tumor virus/activated c-src fusion gene. *Proc. Nat. Acad. Sci.* 92: 7849–7853, 1995.
- [69878] 8208. Yamanashi, Y.; Fukushige, S.-I.; Semba, K.; Sukegawa, J.; Miyajima, N.; Matsubara, K.-I.; Yamamoto, T.; Toyoshima, K.: The yes-related cellular gene lyn encodes a possible tyrosine kinase similar to p56(lck). *Molec. Cell. Biol.* 7: 237–243, 1987.
- [69879] 8209. Berger, A.; Rosenthal, D.; Spiegel, S.: Sphingolipophosphocholine, a signaling molecule which accumulates in Niemann-Pick disease type A, stimulates DNA-binding activity of the transcription activator protein AP-1. *Proc. Nat. Acad. Sci.* 92: 5885–5889, 1995.
- [69880] 8210. Bohmann, D.; Bos, T. J.; Admon, A.; Nishimura, T.; Vogt, P. K.; Tjian, R.: Human proto-oncogene c-jun encodes a DNA binding protein with structural and functional properties of transcription factor AP-1. *Science* 238: 1386–1392, 1987.
- [69881] 8211. Bos, T. J.; Bohmann, D.; Tsuchie, H.; Tjian, R.; Vogt, P. K.: v-jun encodes a nuclear protein with enhancer binding properties of AP-1. *Cell* 52: 705–712, 1988.

- [69882] 8212. Haluska, F. G.; Huebner, K.; Isobe, M.; Nishimura, T.; Croce, C.M.; Vogt, P. K.: Localization of the human JUN protooncogene to chromosomal region 1p31–32. *Proc. Nat. Acad. Sci.* 85: 2215–2218, 1988.
- [69883] 8213. Hattori, K.; Angel, P.; Le Beau, M. M.; Karin, M.: Structure and chromosomal localization of the functional intronless human JUN protooncogene. *Proc. Nat. Acad. Sci.* 85: 9148–9152, 1988.
- [69884] 8214. Lamph, W. W.; Wamsley, P.; Sassone-Corsi, P.; Verma, I. M.: Induction of proto-oncogene JUN/AP-1 by serum and TPA. *Nature* 334: 629–631, 1988.
- [69885] 8215. Marx, J. L.: 'Jun' is bustin' out all over. (Research News). *Science* 242: 1377–1378, 1988.
- [69886] 8216. Mattei, M. G.; Simon-Chazottes, D.; Hirai, S.-I.; Ryssek, R.-P.; Galcheva-Gargova, Z.; Guenet, J. L.; Mattei, J. F.; Bravo, R.; Yaniv, M.: Chromosomal localization of the three members of the jun proto-oncogene family in mouse and man. *Oncogene* 5: 151–156, 1990.
- [69887] 8217. Shaulian, E.; Karin, M.: AP-1 as a regulator of cell life and death. *Nature Cell Biol.* 4: E131–E136, 2002.
- [69888] 8218. Shaulian, E.; Schreiber, M.; Piu, F.; Beeche, M.; Wagner, E. F.; Karin, M.: The mammalian UV response: c-Jun induction is required for exit from p53-imposed growth



arrest. *Cell* 103: 897–907, 2000.

[69889] 8219. Mattei, M. G.; Simon–Chazottes, D.; Hirai, S.; Ryseck, R. P.; Galcheva–Gargova, Z.; Guenet, J. L.; Mattei, J. F.; Bravo, R.; Yaniv, M.: Chromosomal localization of the three members of the jun proto–oncogene family in mouse and man. *Oncogene* 5: 151–156, 1990.

[69890] 8220. Passegue, E.; Jochum, W.; Behrens, A.; Ricci, R.; Wagner, E. F.: JunB can substitute for Jun in mouse development and cell proliferation. *Nature Genet.* 30: 158–166, 2002.

[69891] 8221. Passegue, E.; Jochum, W.; Schorpp–Kistner, M.; Mohle–Steinlein, U.; Wagner, E. F.: Chronic myeloid leukemia with increased granulocyte progenitors in mice lacking JunB expression in the myeloid lineage. *Cell* 104: 21–32, 2001.

[69892] 8222. Phinney, D. G.; Tseng, S. W.; Ryder, K.: Complex genetic organization of junB: multiple blocks of flanking evolutionarily conserved sequence at the murine and human junB loci. *Genomics* 28: 228–234, 1995.

[69893] 8223. Sullivan, M.; Olsen, A. S.; Houslay, M. D.: Genomic organisation of the human cyclic AMP–specific phosphodiesterase PDE4C gene and its chromosomal localisation to 19p13.1, between RAB3A and JUND. *Cell. Signal.* 11:

735–742, 1999.

- [69894] 8224. Weitzman, J. B.; Fiette, L.; Matsuo, K.; Yaniv, M.: JunD protects cells from p53-dependent senescence and apoptosis. *Molec. Cell* 6:1109–1119, 2000.
- [69895] 8225. DeKoter, R. P.; Lee, H.-J.; Singh, H.: PU.1 regulates expression of the interleukin-7 receptor in lymphoid progenitors. *Immunity* 16:297–309, 2002.
- [69896] 8226. DeKoter, R. P.; Singh, H.: Regulation of B lymphocyte and macrophage development by graded expression of PU.1. *Science* 288: 1439–1441, 2000.
- [69897] 8227. Li, S.-L.; Valente, A. J.; Zhao, S.-J.; Clark, R. A.: PU.1 is essential for p47(phox) promoter activity in myeloid cells. *J. Biol. Chem.* 272: 17802–17809, 1997.
- [69898] 8228. Moreau-Gachelin, F.; Tavittian, A.; Tambourin, P.: Spi-1 is a putative oncogene in virally induced murine erythroleukaemias. *Nature* 331:277–280, 1988.
- [69899] 8229. Ray, D.; Cline, S.; Tavittian, A.; Moreau-Gachelin, F.: The human homologue of the putative proto-oncogene Spi-1: characterization and expression in tumors. *Oncogene* 5: 663–667, 1990.
- [69900] 8230. Tondravi, M. M.; McKercher, S. R.; Anderson, K.; Erdmann, J. M.; Quiroz, M.; Maki, R.; Teitelbaum, S. L.: Osteopetrosis in mice lacking haematopoietic transcription

factor PU.1 Nature 386: 81–84, 1997.

[69901] 8231.Xie, C.–W.; Sayah, D.; Chen, Q.–S.; Wei, W.–Z.; Smith, D.; Liu,X.: Deficient long–term memory and long–lasting long–term potentiationin mice with a targeted deletion of neurotrophin–4 gene. Proc. Nat.Acad. Sci. 97: 8116–8121, 2000.

[69902] 8232.Steiglitz, B. M.; Greenspan, D. S.: Assignment of the mouse Pcolce2gene, which encodes procollagen C–proteinase enhancer protein 2, to chromosome 9 and localization of PCOLCE2 to human chromosome 3q23. Cytogenet.Cell Genet. 95: 244–245, 2001.

[69903] 8233.Xu, H.; Acott, T. S.; Wirtz, M. K.: Identification and expressionof a novel type I procollagen C–proteinase enhancer protein gene fromthe glaucoma candidate region on 3q21–q24. Genomics 66: 264–273,2000.

[69904] 8234.Ke, N.; Godzik, A.; Reed, J. C.: Bcl–B, a novel Bcl–2 family memberthat differentially binds and regulates Bax and Bak. J. Biol. Chem. 276:12481–12484, 2001.

[69905] 8235.Lee, R.; Chen, J.; Matthews, C. P.; McDougall, J. K.; Neiman, P.E.: Characterization of NR13–related human cell death regulator,Boo/Diva, in normal and cancer tissues. Biochim. Biophys. Acta 1520:187–194, 2001.

[69906] 8236.Naumann, U.; Weit, S.; Wischhusen, J.; Weller, M.:

Diva/Boo is a negative regulator of cell death in human glioma cells. FEBS Lett. 505:23–26, 2001.

[69907] 8237.Zhang, H.; Holzgreve, W.; De Geyter, C.: Bcl2-L-10, a novel anti-apoptotic member of the Bcl-2 family, blocks apoptosis in the mitochondria death pathway but not in the death receptor pathway. Hum. Molec. Genet.

10:2329–2339, 2001.

[69908] 8238.Brand, S. H.; Castle, J. D.: SCAMP-37, a new marker within the general cell surface recycling system. EMBO J.

12: 3753–3761, 1993.

[69909] 8239.Singleton, D. R.; Wu, T. T.; Castle, J. D.: Three mammalian SCAMPs (secretory carrier membrane proteins) are highly related products of distinct genes having similar subcellular distributions. J. Cell Sci. 110: 2099–2107,

1997.

[69910] 8240.Yamashita, A.; Ohnishi, T.; Kashima, I.; Taya, Y.;

Ohno, S.: Human SMG-1, a novel phosphatidylinositol

3-kinase-related protein kinase, associates with components of the mRNA surveillance complex and is involved in

the regulation of nonsense-mediated mRNA decay.

Genes Dev. 15: 2215–2228, 2001.

[69911] 8241.Spataro, V.; Toda, T.; Craig, R.; Seeger, M.; Dubiel,

W.; Harris, A. L.; Norbury, C.: Resistance to diverse drugs

and ultraviolet light conferred by overexpression of a novel human 26 S proteasome subunit. *J. Biol. Chem.* 272: 30470–30475, 1997.

- [69912] 8242. Yao, T.; Cohen, R. E.: A cryptic protease couples deubiquitination and degradation by the proteasome. *Nature* 1 Sep, 2002. Note: Advance Electronic Publication.
- [69913] 8243. Kamberov, E.; Makarova, O.; Roh, M.; Liu, A.; Karnak, D.; Straight, S.; Margolis, B.: Molecular cloning and characterization of Pals, proteins associated with mLin-7. *J. Biol. Chem.* 275: 11425–11431, 2000.
- [69914] 8244. Tseng, T.-C.; Marfatia, S. M.; Bryant, P. J.; Pack, S.; Zhuang, A.; O'Brien, J. E.; Lin, L.; Hanada, T.; Chishti, A. H.: VAM-1: a new member of the MAGUK family binds to human Veli-1 through a conserved domain. *Biochim. Biophys. Acta* 1518: 249–259, 2001.
- [69915] 8245. Wei, X.; Malicki, J. M.: *nagie oko*, encoding a MAGUK-family protein, is essential for cellular patterning of the retina. *Nature Genet.* 31: 150–157, 2002. Note: Erratum: *Nature Genet.* 31: 439 only, 2002.
- [69916] 8246. Esser, V.; Russell, D. W.: Transport-deficient mutations in the low density lipoprotein receptor: alterations in the cysteine-rich and cysteine-poor regions of the protein block intracellular transport. *J. Biol. Chem.* 263:

13276–13281, 1988.

- [69917] 8247.Fumeron, F.; Grandchamp, B.; Fricker, J.; Krempf, M.; Wolf, L.–M.;Khayat, M.–C.; Boiffard, O.; Apfelbaum, M.: Presence of the FrenchCanadian deletion in a French patient with familial hypercholesterolemia.(Letter) New Eng. J. Med. 326: 69 only, 1992.
- [69918] 8248.Gilbert, W.: Genes–in–pieces revisited. Science 228: 823–824,1985.
- [69919] 8249.Goldfarb, L. G.; Petersen, R. B.; Tabaton, M.; Brown, P.; LeBlanc,A. C.; Montagna, P.; Cortelli, P.; Julien, J.; Vital, C.; Pendelbury,W. W.; Haltia, M.; Wills, P. R.; and 9 others: Fatal familial insomniaand familial Creutzfeldt–Jakob disease: disease phenotype determinedby a DNA polymorphism. Science 258: 806–808, 1992.
- [69920] 8250.Graadt van Roggen, F.; van der Westhuyzen, D. R.; Marais, A. D.;Gevers, W.; Coetzee, G. A.: Low density lipoprotein receptor foundermutations in Afrikaner familial hypercholesterolaemic patients: acomparison of two geographical areas. Hum. Genet. 88: 204–208, 1991.
- [69921] 8251.Gudnason, V.; King–Underwood, L.; Seed, M.; Sun, X.–M.; Soutar,A. K.; Humphries, S. E.: Identification of recurrent and novel mutationsin exon 4 of the LDL receptor gene in patients with familial hypercholesterolemia in the

United Kingdom. Arteriosclerosis Thromb. 13: 56–63, 1993.

- [69922] 8252. Gudnason, V.; Sigurdsson, G.; Nissen, H.; Humphries, S. E.: Commonfounder mutation in the LDL receptor gene causing familial hypercholesterolemia in the Icelandic population. Hum. Mutat. 10: 36–44, 1997.
- [69923] 8253. Henderson, H. E.; Berger, G. M. B.; Marais, A. D.: A new LDLreceptor gene deletion mutation in the South African population. Hum. Genet. 80: 371–374, 1988.
- [69924] 8254. Hobbs, H. H.: Personal Communication. Dallas, Tex. 12/1/1990.
- [69925] 8255. Hobbs, H. H.; Brown, M. S.; Russell, D. W.; Davignon, J.; Goldstein, J. L.: Deletion in the gene for the low-density-lipoprotein receptor in a majority of French Canadians with familial hypercholesterolemia. New Eng. J. Med. 317: 734–737, 1987.
- [69926] 8256. Hobbs, H. H.; Lehrman, M. A.; Yamamoto, T.; Russell, D. W.: Polymorphism and evolution of Alu sequences in the human low density lipoprotein receptor gene. Proc. Nat. Acad. Sci. 82: 7651–7655, 1985.
- [69927] 8257. Hobbs, H. H.; Leitersdorf, E.; Goldstein, J. L.; Brown, M. S.; Russell, D. W.: Multiple CRM- mutations in familial hypercholesterolemia: evidence for 13 alleles, including

four deletions. *J. Clin. Invest.* 81:909–917, 1988.

[69928] 8258.Horsthemke, B.; Dunning, A.; Humphries, S.: Identification of deletions in the human low density lipoprotein receptor gene. *J. Med. Genet.* 24: 144–147, 1987.

[69929] 8259.Horsthemke, B.; Kessling, A. M.; Seed, M.; Wynn, V.; Williamson, R.; Humphries, S. E.: Identification of a deletion in the low density lipoprotein (LDL) receptor gene in a patient with familial hypercholesterolaemia. *Hum. Genet.* 71: 75–78, 1985.

[69930] 8260.Jensen, H. K.; Jensen, T. G.; Faergeman, O.; Jensen, L. G.; Andresen, B. S.; Corydon, M. J.; Andreassen, P. H.; Hansen, P. S.; Heath, F.; Bolund, L.; Gregersen, N.: Two mutations in the same low-density lipoprotein receptor allele act in synergy to reduce receptor function in heterozygous familial hypercholesterolemia. *Hum. Mutat.* 9: 437–444, 1997.

[69931] 8261.Jensen, J. M.; Kruse, T. A.; Brorholt-Petersen, J. U.; Christiansen, T. M.; Jensen, H. K.; Kolvraa, S.; Faergeman, O.: Linking genotype to aorto-coronary atherosclerosis: a model using familial hypercholesterolemia and aorto-coronary calcification. *Ann. Hum. Genet.* 63: 511–520, 1999.

[69932] 8262.Kajinami, K.; Fujita, H.; Koizumi, J.; Mabuchi, H.;



Takeda, R.;Ohta, M.: Genetically determined mild type of familial hypercholesterolemia including normocholesterolemic patients: FH-Tonami-2. Circulation 80(suppl. 2): 278 only, 1989.

[69933] 8263.Kajinami, K.; Mabuchi, H.; Itoh, H.; Michishita, I.; Takeda, M.;Wakasugi, T.; Koizumi, J.; Takeda, R.: New variant of low densitylipoprotein receptor gene FH-Tonami. Arteriosclerosis 8: 187-192,1988.

[69934] 8264.Podos, S. D.; Reddy, P.; Ashkenas, J.; Krieger, M.: LDLC encodes a brefeldin A-sensitive, peripheral Golgi protein required for normal Golgi function. J. Cell Biol. 127: 679-691, 1994.

[69935] 8265.Suvorova, E. S.; Kurten, R. C.; Lupashin, V. V.: Identification of a human orthologue of Sec34p as a component of the cis-Golgi vesicle tethering machinery. J. Biol. Chem. 276: 22810-22818, 2001.

[69936] 8266.Whyte, J. R. C.; Munro, S.: The Sec34/35 Golgi transport complex is related to the exocyst, defining a family of complexes involved in multiple steps of membrane traffic. Dev. Cell 1: 527-537, 2001.

[69937] 8267.Haataja, L.; Groffen, J.; Heisterkamp, N.: Identification of a novel Rac3-interacting protein C1D. Int. J. Molec. Med. 1: 665-670,1998.

- [69938] 8268.Nehls, P.; Keck, T.; Greferath, R.; Spiess, E.; Glaser, T.; Rothbarth,K.; Stammer, H.; Werner, D.: cDNA cloning, recombinant expressionand characterization of polypeptides with exceptional DNA affinity. *NucleicAcids Res.* 26: 1160–1166, 1998.
- [69939] 8269.Rothbarth, K.; Hunziker, A.; Stammer, H.; Werner, D.: Promoterof the gene encoding the 16 kDa DNA-binding and apoptosis-inducingC1D protein. *Biochim. Biophys. Acta* 1518: 271–275, 2001.
- [69940] 8270.Zamir, I.; Dawson, J.; Lavinsky, R. M.; Glass, C. K.; Rosenfeld,M. G.; Lazar, M. A.: Cloning and characteriza-tion of a corepressorand potential component of the nuclear hormone receptor repressioncomplex. *Proc. Nat. Acad. Sci.* 94: 14400–14405, 1997.
- [69941] 8271.Bera, T. K.; Lee, S.; Salvatore, G.; Lee, B.; Pastan, I.: MRP8,a new member of ABC transporter superfamily, identified by EST databasemining and gene prediction program, is highly expressed in breastcancer. *Molec. Med.* 7: 509–516, 2001.
- [69942] 8272.Tammur, J.; Prades, C.; Arnould, I.; Rzhetsky, A.; Hutchinson,A.; Adachi, M.; Schuetz, J. D.; Swoboda, K. J.; Ptacek, L. J.; Rosier,M.; Dean, M.; Allikmets, R.: Two new genes from the human ATP-bindingcassette transporter

superfamily, ABCC11 and ABCC12, tandemly duplicated on chromosome 16q12. *Gene* 273: 89–96, 2001.

- [69943] 8273. Yabuuchi, H.; Shimizu, H.; Takayanagi, S.; Ishikawa, T.: Multiple splicing variants of two new human ATP-binding cassette transporters, ABCC11 and ABCC12. *Biochem. Biophys. Res. Commun.* 288: 933–939, 2001.
- [69944] 8274. Hoefler, G.; Forstner, M.; McGuinness, M. C.; Hulla, W.; Hiden, M.; Krisper, P.; Kenner, L.; Ried, T.; Lengauer, C.; Zechner, R.; Moser, H. W.; Chen, G. L.: cDNA cloning of the human peroxisomal enoyl-CoA hydratase: 3-hydroxyacyl-CoA dehydrogenase bifunctional enzyme and localization to chromosome 3q26.3–3q28: a free left Alu arm is inserted in the 3-prime noncoding region. *Genomics* 19: 60–67, 1994.
- [69945] 8275. Bera, T. K.; Iavarone, C.; Kumar, V.; Lee, S.; Lee, B.; Pastan, I.: MRP9, an unusual truncated member of the ABC transporter superfamily, is highly expressed in breast cancer. *Proc. Nat. Acad. Sci.* 99: 6997–7002, 2002.
- [69946] 8276. Nagle, D. L.; McGrail, S. H.; Vitale, J.; Woolf, E. A.; Dussault, B. J., Jr.; DiRocco, L.; Holmgren, L.; Montagno, J.; Bork, P.; Huszar, D.; Fairchild-Huntress, V.; Ge, P.; Keilty, J.; Ebell, C.; Baldini, L.; Gilchrist, J.; Burr, P.; Carlson, G. A.; Moore, K. J.: The mahogany protein is a receptor in–

volved in suppression of obesity. *Nature* 398:148–151, 1999.

- [69947] 8277. Zhang, Y.; Ng, H.-H.; Erdjument-Bromage, H.; Tempst, P.; Bird, A.; Reinberg, D.: Analysis of the NuRD subunits reveals a histone deacetylase core complex and a connection with DNA methylation. *Genes Dev.* 13: 1924–1935, 1999.
- [69948] 8278. Strobl, B.; Wechselberger, C.; Beier, D. R.; Lepperdinger, G.: Structural organization and chromosomal localization of Hyal2, a gene encoding a lysosomal hyaluronidase. *Genomics* 53: 214–219, 1998.
- [69949] 8279. Pineda, M.; Fernandez, E.; Torrents, D.; Estevez, R.; Lopez, C.; Camps, M.; Lloberas, J.; Zorzano, A.; Palacin, M.: Identification of a membrane protein, LAT-2, that co-expresses with 4F2 heavy chain, an L-type amino acid transport activity with broad specificity for small and large zwitterionic amino acids. *J. Biol. Chem.* 274: 19738–19744, 1999.
- [69950] 8280. Alexander, W. S.; Rakar, S.; Robb, L.; Farley, A.; Willson, T. A.; Zhang, J.-G.; Hartley, L.; Kikuchi, Y.; Kojima, T.; Nomura, H.; Hasegawa, M.; Maeda, M.; Fabri, L.; Jachno, K.; Nash, A.; Metcalf, D.; Nicola, N. A.; Hilton, D. J.: Suckling defect in mice lacking the soluble haemopoietin receptor

NR6. Curr. Biol. 9: 605–608, 1999.

- [69951] 8281.Elson, G. C. A.; Graber, P.; Losberger, C.; Herren, S.; Gretener,D.; Menoud, L. N.; Wells, T. N. C.; Kosco–Vilbois, M. H.; Gauchat,J.–F.: Cytokine–like factor–1, a novel soluble protein, shares homologywith members of the cytokine type I receptor family. J. Immun. 161:1371–1379, 1998.
- [69952] 8282.Hinderlich, S.; Stasche, R.; Zeitler, R.; Reutter, W.: A bifunctionalenzyme catalyzes the first two steps in N–acetylneuraminic acid biosynthesisof rat liver: purification and characterization of UDP–N–acetylglucosamine2–epimerase/N–acetylmannosamine kinase. J. Biol. Chem. 272: 24313–24318,1997.
- [69953] 8283.Huizing, M.; Anikster, Y.: Personal Communication. Bethesda,Md. 1/10/2000.
- [69954] 8284.Kayashima, T.; Matsuo, H.; Satoh, A.; Ohta, T.; Yoshiura, K.; Matsumoto,N.; Nakane, Y.; Niikawa, N.; Kishino, T.: Nonaka myopathy is causedby mutations in the UDP–N–acetylglu–cosamine–2–epimerase/N–acetylmannosaminekinase gene (GNE). J. Hum. Genet. 47: 77–79, 2002.
- [69955] 8285.Keppler, O. T.; Hinderlich, S.; Langner, J.; Schwartz–Al–

biez, R.; Reutter, W.; Pawlita, M.: UDP-GlcNAc

2-epimerase: a regulator of cell surface sialylation. *Science* 284: 1372–1376, 1999.

[69956] 8286. Nonaka, I.; Sunohara, N.; Ishiura, S.; Satoyoshi, E.: Familial distal myopathy with rimmed vacuole and lamellar (myeloid) body formation. *J. Neurol. Sci.* 51: 141–155, 1981.

[69957] 8287. Schwarzkopf, M.; Knobloch, K.-P.; Rohde, E.; Hinderlich, S.; Wiechens, N.; Lucka, L.; Horak, I.; Reutter, W.; Horstkorte, R.: Sialylation is essential for early development in mice. *Proc. Nat. Acad. Sci.* 99: 5267–5270, 2002.

[69958] 8288. Stasche, R.; Hinderlich, S.; Weise, C.; Effertz, K.; Lucka, L.; Moormann, P.; Reutter, W.: A bifunctional enzyme catalyzes the first two steps in N-acetylneuraminic acid biosynthesis of rat liver: molecular cloning and functional expression of UDP-N-acetyl-glucosamine 2-epimerase/N-acetylmannosamine kinase. *J. Biol. Chem.* 272: 24319–24324, 1997.

[69959] 8289. Carter, M. G.; Johns, M. A.; Zeng, X.; Zhou, L.; Zink, M. C.; Mankowski, J. L.; Donovan, D. M.; Baylin, S. B.: Mice deficient in the candidate tumor suppressor gene *Hic1* exhibit developmental defects of structures affected in the Miller–Dieker syndrome. *Hum. Molec. Genet.* 9:

413–419,2000.

- [69960] 8290.Chong, S. S.; Tanigami, A.; Roschke, A. V.; Ledbetter, D. H.:14–3–3epsilon has no homology to LIS1 and lies telomeric to it on chromosome 17p13.3 outside the Miller–Dieker syndrome chromosome region. *GenomeRes.* 6: 735–741, 1996.
- [69961] 8291.Grimm, C.; Sporle, R.; Schmid, T. E.; Adler, I.–D.; Adamski, J.;Schughart, K.; Graw, J.: Isolation and embryonic expression of thenovel mouse gene Hic1, the homologue of HIC1, a candidate gene forthe Miller–Dieker syndrome. *Hum. Molec. Genet.* 8: 697–710, 1999.
- [69962] 8292.Makos Wales, M.; Biel, M. A.; El Deiry, W.; Nelkin, B. D.; Issa,J.–P.; Cavenee, W. K.; Kuerbitz, S. J.; Baylin, S. B.: p53 activatesexpression of HIC–1, a new candidate tumour suppressor gene on 17p13.3. *NatureMed.* 1: 570–577, 1995.
- [69963] 8293.Smith, D. J.; Salmi, M.; Bono, P.; Hellman, J.; Leu, T.; Jalkanen,S.: Cloning of vascular adhesion protein 1 reveals a novel multifunctionaladhesion molecule. *J. Exp. Med.* 188: 17–27, 1998.
- [69964] 8294.Zhang, X.; McIntire, W. S.: Cloning and sequencing of a copper–containing,topa quinone–containing monoamine oxidase from human placenta. *Gene*

179:279–286, 1996.

- [69965] 8295.Lomako, J.; Mazuruk, K.; Lomako, W. M.; Alonso, M. D.; Whelan, W. J.; Rodriguez, I. R.: The human intron-containing gene for glycogenin maps to chromosome 3, band q24. *Genomics* 33: 519–522, 1996.
- [69966] 8296.Tanaka, T.; Tsujimura, T.; Takeda, K.; Sugihara, A.; Maekawa, A.; Terada, N.; Yoshida, N.; Akira, S.: Targeted disruption of ATF4 discloses its essential role in the formation of eye lens fibres. *Genes Cells* 3:801–810, 1998.
- [69967] 8297.Maas, S.; Gerber, A. P.; Rich, A.: Identification and characterization of a human tRNA-specific adenosine deaminase related to the ADAR family of pre-mRNA editing enzymes. *Proc. Nat. Acad. Sci.* 96: 8895–8900, 1999.
- [69968] 8298.Viskupic, E.; Cao, Y.; Zhang, W.; Cheng, C.; DePaoli-Roach, A.A.; Roach, P. J.: Rabbit skeletal muscle glycogenin. Molecular cloning and production of fully functional protein in *Escherichia coli*. *J. Biol. Chem.* 267: 25759–25763, 1992.
- [69969] 8299.Hosokawa, Y.; Matsumoto, A.; Oka, J.; Itakura, H.; Yamaguchi, K.: Isolation and characterization of a cDNA for rat liver cysteine dioxygenase. *Biochem. Biophys. Res. Commun.* 168: 473–478, 1990.
- [69970] 8300.Jeremiah, S.; McCann, K. P.; Williams, A. C.; Rams-



den, D. B.; Pilz, A. J.; Fox, M. F.; Povey, S.: Chromosomal localisation of genes coding for human and mouse liver cytosolic cysteine dioxygenase. *Ann. Hum. Genet.* 60: 29–33, 1996.

[69971] 8301. McCann, K. P.; Akbari, M. T.; Williams, A. C.; Ramsden, D. B.: Human cysteine dioxygenase type I: primary structure derived from base sequencing of cDNA. *Biochim. Biophys. Acta* 1209: 107–110, 1994.

[69972] 8302. Ramsden, D. B.; Kapadi, A.; Fitch, N. J. S.; Farmer, M. J.; Bennett, P.; Williams, A. C.: Human cysteine dioxygenase type I (CDO-I; EC1.13.11.20): 5-prime flanking region and intron-exon structure of the gene. *Molec. Path.* 50: 269–271, 1997.

[69973] 8303. Fox, M. F.; Lahbib, F.; Pratt, W.; Attwood, J.; Gum, J.; Kim, Y.; Swallow, D. M.: Regional localization of the intestinal mucin gene MUC3 to chromosome 7q22. *Ann. Hum. Genet.* 56: 281–287, 1992.

[69974] 8304. Kyo, K.; Muto, T.; Nagawa, H.; Lathrop, G. M.; Nakamura, Y.: Association of distinct variants of the intestinal mucin gene MUC3A with ulcerative colitis and Crohn's disease. *J. Hum. Genet.* 46: 5–20, 2001.

[69975] 8305. Hill, A. V. S.; Gentile, B.; Bonnardot, J. M.; Roux, J.; Weatherall, D. J.; Clegg, J. B.: Polynesian origins and affini-

ties: globin genevariants in eastern Polynesia. *Am. J. Hum. Genet.* 40: 453–463, 1987.

[69976] 8306.Zou, Y.-R.; Kottmann, A. H.; Kuroda, M.; Taniuchi, I.; Littman, D. R.: Function of the chemokine receptor CXCR4 in haematopoiesis and in cerebellar development. *Nature* 393: 595–599, 1998.

[69977] 8307.Laurent, P.; Clerc, P.; Mattei, M.-G.; Forgez, P.; Dumont, X.; Ferrara, P.; Caput, D.; Rostene, W.: Chromosomal localization of mouse and human neurotensin receptor genes. *Mammalian Genome* 5:303–306, 1994.

[69978] 8308.Le, F.; Groshan, K.; Zeng, X. P.; Richelson, E.: Characterization of the genomic structure, promoter region, and a tetranucleotide repeat polymorphism of the human neurotensin receptor gene. *J. Biol. Chem.* 272:1315–1322, 1997.

[69979] 8309.Vincent, J.-P.: Neurotensin receptors: binding properties, transduction pathways, and structure. *Cell. Molec. Neurobiol.* 15: 501–512, 1995.

[69980] 8310.Vita, N.; Laurent, P.; Lefort, S.; Chalon, P.; Dumont, X.; Kaghad, M.; Gully, D.; Le Fur, G.; Ferrara, P.; Caput, D.: Cloning and expression of a complementary DNA encoding a high affinity human neurotensin receptor. *FEBS Lett.* 317: 139–142, 1993.

- [69981] 8311.Jones, K. R.; Reichardt, L. F.: Molecular cloning of a human gene that is a member of the nerve growth factor family. *Proc. Nat. Acad.Sci.* 87: 8060–8064, 1990.
- [69982] 8312.Berkemeier, L. R.; Ozcelik, T.; Francke, U.; Rosenthal, A.: Human chromosome 19 contains the neurotrophin–5 gene locus and three related genes that may encode novel acidic neurotrophins. *Somat. Cell Molec.Genet.* 18: 233–245, 1992.
- [69983] 8313.Ibanez, C. F.: Neurotrophin–4: the odd one out in the neurotrophin family. *Neurochem. Res.* 21: 787–793, 1996.
- [69984] 8314.Ip, N. Y.; Ibanez, C. F.; Nye, S. H.; McClain, J.; Jones, P. F.; Gies, D. R.; Belluscio, L.; Le Beau, M. M.; Espinosa, R., III; Squinto, S. P.; Persson, H.; Yancopoulos, G. D.: Mammalian neurotrophin–4: structure, chromosomal localization, tissue distribution, and receptor specificity. *Proc. Nat. Acad. Sci.* 89: 3060–3064, 1992.
- [69985] 8315.Hill, A. V. S.; Nicholls, R. D.; Thein, S. L.; Higgs, D. R.: Recombination within the human embryonic zeta–globin locus: a common zeta–zeta chromosome produced by gene conversion of the psi–zeta gene. *Cell* 42:809–819, 1985.
- [69986] 8316.Housman, D.: Personal Communication. Boston,

Mass. 1979.

- [69987] 8317.Kamuzora, H.; Lehmann, H.: Human embryonic haemoglobins including a comparison by homology of the human zeta and alpha chains. *Nature* 256:511–513, 1975.
- [69988] 8318.Melderis, H.; Steinheider, G.; Ostertag, W.: Evidence for a unique kind of alpha-type globin chain in early mammalian embryos. *Nature* 250:774–776, 1974.
- [69989] 8319.Pressley, L.; Higgs, D. R.; Clegg, J. B.; Weatherall, D. J.: Gene deletions in alpha-thalassemia prove that the 5-prime zeta locus is functional. *Proc. Nat. Acad. Sci.* 77: 3586–3589, 1980.
- [69990] 8320.Proudfoot, N. J.; Shander, M. H. M.; Manley, J. L.; Gefer, M.L.; Maniatis, T.: Structure and in vitro transcription of human globin genes. *Science* 209: 1329–1336, 1980.
- [69991] 8321.Randhawa, Z. I.; Jones, R. T.; Lie-Injo, L. E.: Separation of the tryptic peptides and cyanogen bromide fragments of the human embryonic zeta chains of hemoglobin Portland I and II by reverse phase high performance liquid chromatography. *Hemoglobin* 8: 463–482, 1984.
- [69992] 8322.Whitney, J. B., III; Russell, E. S.: Linkage of genes for adult alpha-globin and embryonic alpha-like globin chains. *Proc. Nat. Acad. Sci.* 77: 1087–1090, 1980.

- [69993] 8323. Bluteau, O.; Jeannot, E.; Bioulac-Sage, P.; Marques, J. M.; Blanc, J.-F.; Bui, H.; Beaudoin, J.-C.; Franco, D.; Balabaud, C.; Laurent-Puig, P.; Zucman-Rossi, J.: Bi-allelic inactivation of TCF1 in hepatic adenomas. *Nature Genet.* 32: 312–315, 2002.
- [69994] 8324. Foster, J. H.; Donohue, T. A.; Berman, M. M.: Familial liver-cell adenomas and diabetes mellitus. *New Eng. J. Med.* 299: 239–241, 1978.
- [69995] 8325. Quintrell, N.; Lebo, R.; Varmus, H.; Bishop, J. M.; Pettenati, M. J.; Le Beau, M. M.; Diaz, M. O.; Rowley, J. D.: Identification of a human gene (HCK) that encodes a protein-tyrosine kinase and is expressed in hemopoietic cells. *Molec. Cell. Biol.* 7: 2267–2275, 1987.
- [69996] 8326. Ziegler, S. F.; Marth, J. D.; Lewis, D. B.; Perlmutter, R. M.: Novel protein-tyrosine kinase gene (hck) preferentially expressed in cells of hematopoietic origin. *Molec. Cell. Biol.* 7: 2276–2285, 1987.
- [69997] 8327. Honda, H.; Inaba, T.; Suzuki, T.; Oda, H.; Ebihara, Y.; Tsuiji, K.; Nakahata, T.; Ishikawa, T.; Yazaki, Y.; Hirai, H.: Expression of E2A-HLF chimeric protein induced T-cell apoptosis, B-cell maturation arrest, and development of acute lymphoblastic leukemia. *Blood* 93: 2780–2790, 1999.

- [69998] 8328.Hunger, S. P.: Chromosomal translocations involving the E2A gene in acute lymphoblastic leukemia: clinical features and molecular pathogenesis. *Blood* 87:1211–1224, 1996.
- [69999] 8329.Inaba, T.; Roberts, W. M.; Shapiro, L. H.; Jolly, K. W.; Raimondi, S. C.; Smith, S. D.; Look, A. T.: Fusion of the leucine zipper gene HLF to the E2A gene in human acute B-lineage leukemia. *Science* 257:531–534, 1992.
- [70000] 8330.Smith, K. S.; Rhee, J. W.; Naumovski, L.; Cleary, M. L.: Disrupted differentiation and oncogenic transformation of lymphoid progenitors in E2A–HLF transgenic mice. *Molec. Cell. Biol.* 19: 4443–4451, 1999.
- [70001] 8331.Acampora, D.; D'Esposito, M.; Faiella, A.; Pannese, M.; Migliaccio, E.; Morelli, F.; Stornaiuolo, A.; Nigro, V.; Simeone, A.; Boncinelli, E.: The human HOX gene family. *Nucleic Acids Res.* 17: 10385–10402, 1989.
- [70002] 8332.Balling, R.; Mutter, G.; Gruss, P.; Kessel, M.: Cranio-facial abnormalities induced by ectopic expression of the homeobox gene Hox–1.1 in transgenic mice. *Cell* 58: 337–347, 1989.
- [70003] 8333.Bucan, M.; Yang–Feng, T.; Colberg–Poley, A. M.; Wolgemuth, D. J.; Guenet, J.–L.; Francke, U.; Lehrach, H.: Genetic and cytogenetic localisation of the homeo box

containing genes on mouse chromosome 6 and human chromosome 7. EMBO J. 5: 2899–2905, 1986.

[70004] 8334.Ferguson-Smith, A. C.; Fienberg, A.; Ruddle, F. H.: Isolation, chromosomal localization, and nucleotide sequence of the human HOX1.4 homeobox. Genomics 5: 250–258, 1989.

[70005] 8335.Gaunt, S. J.; Singh, P. B.: Homeogene expression patterns and chromosomal imprinting. Trends Genet. 6: 208–212, 1990.

[70006] 8336.Gehring, W. J.: The homeo box: a key to the understanding of development? Cell 40:3–5, 1985.

[70007] 8337.Joyner, A. L.; Lebo, R. V.; Kan, Y. W.; Tjian, R.; Cox, D. R.; Martin, G. R.: Comparative chromosome mapping of a conserved homeobox region in mouse and human. Nature 314: 173–175, 1985.

[70008] 8338.McGinnis, W.; Garber, R. L.; Wirz, J.; Kuroiwa, A.; Gehring, W.J.: A homologous protein-coding sequence in Drosophila homeotic genes and its conservation in other metazoans. Cell 37: 403–408, 1984.

[70009] 8339.McGinnis, W.; Levine, M. S.; Hafen, E.; Kuroiwa, A.; Gehring, W.J.: A conserved DNA sequence in homeotic genes of the Drosophila Antennapedia and bithorax complexes. Nature 308: 428–433, 1984.

- [70010] 8340.Ohno, S.: Evolution by Gene Duplication. Heidelberg: Springer(pub.) 1970.
- [70011] 8341.Rabin, M.; Ferguson-Smith, A.; Hart, C. P.; Ruddle, F. H.: Cognatehomeo-box loci mapped on homologous human and mouse chromosomes. Proc.Nat. Acad. Sci. 83: 9104-9108, 1986.
- [70012] 8342.Rabin, M.; Hart, C. P.; Ferguson-Smith, A.; McGinnis, W.; Levine,M.; Ruddle, F. H.: Two homoeo box loci mapped in evolutionarily relatedmouse and human chromosomes. Nature 314: 175-178, 1985.
- [70013] 8343.Rieger, R.; Michaelis, A.; Green, M. M.: Glossary of Geneticsand Cytogenetics. New York: Springer-Verlag (pub.) 1976. Pp.281 only.
- [70014] 8344.Schughart, K.; Kappen, C.; Ruddle, F. H.: Duplication of largegenomic regions during the evolution of vertebrate homeobox genes. Proc.Nat. Acad. Sci. 86: 7067-7071, 1989.
- [70015] 8345.Scott, M. P.: Vertebrate homeobox gene nomenclature. (Letter) Cell 71:551-553, 1992.
- [70016] 8346.Greer, J. M.; Puetz, J.; Thomas, K. R.; Capecchi, M. R.: Maintenanceof functional equivalence during paralogous Hox gene evolution. Nature 403:661-665, 2000.
- [70017] 8347.Apiou, F.; Flagiello, D.; Cillo, C.; Malfoy, B.; Poupon,



M.-F.;Dutrillaux, B.: Fine mapping of human HOX gene clusters. *Cytogenet.Cell Genet.* 73: 114–115, 1996.

- [70018] 8348.Chan, H. Y. E.; Warrick, J. M.; Gray-Board, G. L.; Paulson, H.L.; Bonini, N. M.: Mechanisms of chaperone suppression of polyglutaminedisease: selectivity, synergy and modulation of protein solubilityin *Drosophila*. *Hum. Molec. Genet.* 9: 2811–2820, 2000.
- [70019] 8349.Dunah, A. W.; Jeong, H.; Griffin, A.; Kim, Y.-M.; Standaert, D.G.; Hersch, S. M.; Mouradian, M. M.; Young, A. B.; Tanese, N.; Krainc,D.: Sp1 and TAFII130 transcriptional activity disrupted in earlyHuntington's disease. *Science* 296: 2238–2243, 2002.
- [70020] 8350.Man in't Veld, A. J.; Boomsma, F.; Moleman, P.; Schalekamp, M.A. D. H.: Congenital dopamine-beta-hydroxylase deficiency: a novelorthostatic syndrome. *Lancet I*: 183–188, 1987.
- [70021] 8351.Robertson, D.; Goldberg, M. R.; Onrot, J.; Hollister, A. S.; Wiley,R.; Thompson, J. G., Jr.; Robertson, R. M.: Isolated failure of autonomicnoradrenergic neurotransmission: evidence for impaired beta-hydroxylationof dopamine. *New Eng. J. Med.* 314: 1494–1497, 1986.
- [70022] 8352.Ogawa, O.; McNoe, L. A.; Eccles, M. R.; Morison, I. M.; Reeve,A. E.: Human insulin-like growth factor type I

and type II receptors are not imprinted. *Hum. Molec. Genet.* 2: 2163–2165, 1993.

[70023] 8353. Oshima, A.; Nolan, C. M.; Kyle, J. W.; Grubb, J. H.; Sly, W. S.: The human cation-independent mannose 6-phosphate receptor: cloning and sequence of the full-length cDNA and expression of functional receptor in COS cells. *J. Biol. Chem.* 263: 2553–2562, 1988.

[70024] 8354. Puertollano, R.; Aguilar, R. C.; Gorshkova, I.; Crouch, R. J.; Bonifacino, J. S.: Sorting of mannose 6-phosphate receptors mediated by the GGAs. *Science* 292: 1712–1716, 2001.

[70025] 8355. Rao, P. H.; Murty, V. V. V. S.; Gaidano, G.; Hauptschein, R.; Dalla-Favera, R.; Chaganti, R. S. K.: Sub-regional mapping of 8 single-copy loci to chromosome 6 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 66: 272–273, 1994.

[70026] 8356. Roth, R. A.: Structure of the receptor for insulin-like growth factor II: the puzzle amplified. *Science* 239: 1269–1271, 1988.

[70027] 8357. Sleutels, F.; Zwart, R.; Barlow, D. P.: The non-coding Air RNAs are required for silencing autosomal imprinted genes. *Nature* 415: 810–813, 2002.

[70028] 8358. Souza, R. F.; Appel, R.; Yin, J.; Wang, S.; Smolinski,

K. N.; Abraham, J. M.; Zou, T.-T.; Shi, Y.-Q.; Lei, J.; Cottrell, J.; Cymes, K.; Biden, K.; Simms, L.; Leggett, B.; Lynch, P. M.; Frazier, M.; Powell, S. M.; Harpaz, N.; Sugimura, H.; Young, J.; Meltzer, S. J.: Microsatellite instability in the insulin-like growth factor II receptor gene in gastrointestinal tumours. (Letter) *Nature Genet.* 14:255–257, 1996.

[70029] 8359. Szebeny, G.; Rotwein, P.: The mouse insulin-like growth factor II/cation-independent mannose 6-phosphate (IGF-II/MPR) receptor gene: molecular cloning and genomic organization. *Genomics* 19: 120–129, 1994.

[70030] 8360. Tong, P. Y.; Tollefsen, S. E.; Kornfeld, S.: The cation-independent mannose 6-phosphate receptor binds insulin-like growth factor II. *J. Biol. Chem.* 263: 2585–2588, 1988.

[70031] 8361. Waheed, A.; Braulke, T.; Junghans, U.; von Figura, K.: Mannose 6-phosphate/insulin like growth factor II receptor: the two types of ligands bind simultaneously to one receptor at different sites. *Biochem. Biophys. Res. Commun.* 152: 1248–1254, 1988.

[70032] 8362. Xu, Y.; Goodyer, C. G.; Deal, C.; Polychronakos, C.: Functional polymorphism in the parental imprinting of the human IGF2R gene. *Biochem. Biophys. Res. Commun.* 197:

747–754, 1993.

- [70033] 8363. Young, L. E.; Fernandes, K.; McEvoy, T. G.; Butterwith, S. C.; Gutierrez, C. G.; Carolan, C.; Broadbent, P. J.; Robinson, J. J.; Wilmut, I.; Sinclair, K. D.: Epigenetic change in IGF2R is associated with fetal overgrowth after sheep embryo culture. *Nature Genet.* 27:153–154, 2001.
- [70034] 8364. Zhu, Y.; Doray, B.; Poussu, A.; Lehto, V.-P.; Kornfeld, S.: Binding of GGA2 to the lysosomal enzyme sorting motif of the mannose 6-phosphate receptor. *Science* 292: 1716–1718, 2001.
- [70035] 8365. Brown, C. W.; Houston-Hawkins, D. E.; Woodruff, T. K.; Matzuk, M. M.: Insertion of *Inhbb* into the *Inhba* locus rescues the *Inhba*-null phenotype and reveals new activin functions. *Nature Genet.* 25: 453–457, 2000.
- [70036] 8366. Ferguson, C. A.; Tucker, A. S.; Christensen, L.; Lau, A. L.; Matzuk, M. M.; Sharpe, P. T.: Activin is an essential early mesenchymal signal in tooth development that is required for patterning of the murine dentition. *Genes Dev.* 12: 2636–2649, 1998.
- [70037] 8367. Burger, H. G.; Igarashi, M.; Baird, D.; Mason, T.; Bardin, W.; McLachlan, R.; Chappel, S.; Miyamoto, K.; de Jong, F.; Moudgal, A.; Demoulin, A.; Nieschlag, E.; de Kretser, D.; Robertson, D.; Findlay, J.; Sasamoto, S.; Forage,

R.; Schwartz, N.; Fukuda, M.; Steinberger, A.; Hasegawa, Y.; Tanabe, K.; Ling, N.; Ying, S.-Y.: Inhibin: definition and nomenclature, including related substances. (Letter) J. Clin. Endocr. Metab. 66: 885–886, 1988.

[70038] 8368. Lumpkin, M. D.; Moltz, J. H.; Yu, W. H.; Samson, W. K.; McCann, S. M.: Purification of FSH-releasing factor: its dissimilarity from LHRH of mammalian, avian, and piscine origin. Brain Res. Bull. 18: 175–178, 1987.

[70039] 8369. Matzuk, M. M.; Kumar, T. R.; Vassalli, A.; Bickenbach, J. R.; Roop, D. R.; Jaenisch, R.; Bradley, A.: Functional analysis of activins during mammalian development. Nature 374: 354–356, 1995.

[70040] 8370. Mellor, S. L.; Cranfield, M.; Ries, R.; Pedersen, J.; Cancilla, B.; de Kretser, D.; Groome, N. P.; Mason, A. J.; Risbridger, G. P.: Localization of activin beta(A)-, beta(B)-, and beta(C)-subunits in human prostate and evidence for formation of new activin heterodimers of beta(C)-subunit. J. Clin. Endocr. Metab. 85: 4851–4858, 2000.

[70041] 8371. Murata, M.; Eto, Y.; Shibai, H.; Sakai, M.; Muramatsu, M.: Erythroid differentiation factor is encoded by the same mRNA as that of the inhibin beta-A chain. Proc. Nat. Acad. Sci. 85: 2434–2438, 1988.

[70042] 8372. You, L.; Kruse, F. E.: Differential effect of activin A

and BMP-7 on myofibroblast differentiation and the role of the Smad signaling pathway. *Invest. Ophthalm. Vis. Sci.* 43: 72–81, 2002.

[70043] 8373. Lacy, D. A.; Wang, Z.-E.; Symula, D. J.; McArthur, C. J.; Rubin, E. M.; Frazer, K. A.; Locksley, R. M.: Faithful expression of the human 5q31 cytokine cluster in transgenic mice. *J. Immun.* 164: 4569–4574, 2000.

[70044] 8374. Taniyama, Y.; Kuroki, R.; Omura, F.; Seko, C.; Kikuchi, M.: Evidence for intramolecular disulfide bond shuffling in the folding of mutant human lysozyme. *J. Biol. Chem.* 266: 6456–6461, 1991.

[70045] 8375. Loots, G. G.; Locksley, R. M.; Blankespoor, C. M.; Wang, Z. E.; Miller, W.; Rubin, E. M.; Frazer, K. A.: Identification of a coordinate regulator of interleukins 4, 13, and 5 by cross-species sequence comparisons. *Science* 288: 136–140, 2000.

[70046] 8376. McKenzie, A. N. J.; Culpepper, J. A.; de Waal Malefyt, R.; Briere, F.; Punnonen, J.; Aversa, G.; Sato, A.; Dang, W.; Cocks, B. G.; Menon, S.; de Vries, J. E.; Banchereau, J.; Zurawski, G.: Interleukin 13, a T-cell-derived cytokine that regulates human monocyte and B-cell function. *Proc. Nat. Acad. Sci.* 90: 3735–3739, 1993.

[70047] 8377. Minty, A.; Chalon, P.; Derocq, J.-M.; Dumont, X.;

Guillemot, J.-C.;Kaghad, M.; Labit, C.; Leplatois, P.; Li-  
aun, P.; Miloux, B.; Minty,C.; Casellas, P.; Loison, G.;  
Lupker, J.; Shire, D.; Ferrara, P.;Caput, D.: Interleukin-13  
is a new human lymphokine regulating inflammatoryand  
immune responses. Nature 362: 248-250, 1993.

[70048] 8378.Morgan, J. G.; Dolganov, G. M.; Robbins, S. E.; Hin-  
ton, L. M.;Lovett, M.: The selective isolation of novel cD-  
NAs encoded by theregions surrounding the human inter-  
leukin 4 and 5 genes. NucleicAcids Res. 20: 5173-5179,  
1992.

[70049] 8379.Paul, W. E.: Personal Communication. Bethesda, Md.  
3/6/1993.

[70050] 8380.Punnonen, J.; Aversa, G.; Cocks, B. G.; McKenzie, A.  
N. J.; Menon,S.; Zurawski, G.; de Waal Malefyt, R.; de Vries,  
J. E.: Interleukin13 induces interleukin 4-independent  
IgG4 and IgE synthesis and CD23expression by human B  
cells. Proc. Nat. Acad. Sci. 90: 3730-3734,1993.

[70051] 8381.Smirnov, D. V.; Smirnova, M. G.; Korobko, V. G.;  
Frolova, E. I.: Tandem arrangement of human genes for  
interleukin-4 and interleukin-13:resemblance in their or-  
ganization. Gene 155: 277-281, 1995.

[70052] 8382.Wills-Karp, M.; Luyimbazi, J.; Xu, X.; Schofield, B.;  
Neben, T.Y.; Karp, C. L.; Donaldson, D. D.: Interleukin-13:

central mediator of allergic asthma. *Science* 282: 2258–2261, 1998.

- [70053] 8383. Zhu, Z.; Homer, R. J.; Wang, Z.; Chen, Q.; Geba, G. P.; Wang, J.; Zhang, Y.; Elias, J. A.: Pulmonary expression of interleukin-13 causes inflammation, mucus hypersecretion, subepithelial fibrosis, physiologic abnormalities, and eotaxin production. *J. Clin. Invest.* 103:779–788, 1999.
- [70054] 8384. Zurawski, G.; de Vries, J. E.: Interleukin 13 elicits a subset of the activities of its close relative interleukin 4. *Stem Cells* 12:169–174, 1994.
- [70055] 8385. Li, C.; Lai, C.; Sigman, D. S.; Gaynor, R. B.: Cloning of a cellular factor, interleukin binding factor, that binds to NFAT-like motifs in the human immunodeficiency virus long terminal repeat. *Proc. Nat. Acad. Sci.* 88: 7739–7743, 1991.
- [70056] 8386. Li, C.; Lysis, A. J.; Sparkes, R.; Nirula, A.; Gaynor, R.: Characterization and chromosomal mapping of the gene encoding the cellular DNA binding protein ILF. *Genomics* 13: 665–671, 1992.
- [70057] 8387. Boone, C.; Chen, T.-R.; Ruddle, F. H.: Assignment of three human genes to chromosomes (LDH-A to 11, TK to 17 and IDH to 20) and evidence for translocation between human and mouse chromosomes in somatic cell hybrids.



Proc. Nat. Acad. Sci. 68: 510–514, 1972.

- [70058] 8388.Chen, S.–H.; Fossum, B. L. G.; Giblett, E. R.: Genetic variation of the soluble form of NADP–dependent isocitric dehydrogenase in man. Am.J. Hum. Genet. 24: 325–329, 1972.
- [70059] 8389.Creagan, R. P.; Carritt, B.; Chen, S.–H.; Kucherlapati, R. S.;McMorris, F. A.; Ricciuti, F.; Tan, Y. H.; Tischfield, J. A.; Ruddle,F. H.: Chromosome assignments of genes in man using mouse–human somatic cell hybrids: cytoplasmic isocitrate dehydrogenase (IDH 1) and malate dehydrogenase (MDH 1) to chromosome 2. Am. J. Hum. Genet. 26: 604–613,1974.
- [70060] 8390.Glass, I. A.; Swindlehurst, C. A.; Aitken, D. A.; McCrea, W.; Boyd,E.: Interstitial deletion of the long arm of chromosome 2 with normal levels of isocitrate dehydrogenase. J. Med. Genet. 26: 127–130,1989.
- [70061] 8391.Henderson, N. S.: Isozymes of isocitrate dehydrogenase: subunit structure and intracellular location. J. Exp. Zool. 158: 263–273,1965.
- [70062] 8392.Henderson, N. S.: Intracellular location and genetic control of isozymes of NADP–dependent isocitrate dehydrogenase and malate dehydrogenase. Ann.N.Y. Acad. Sci. 151: 429–440, 1968.

- [70063] 8393.Narahara, K.; Kimura, S.; Kikkawa, K.; Takahashi, Y.; Wakita, Y.;Kasai, R.; Nagai, S.; Nishibayashi, Y.; Kimoto, H.: Probable assignmentof soluble isocitrate dehydrogenase (IDH-1) to 2q33.3. Hum. Genet. 71:37-40, 1985.
- [70064] 8394.Ruddle, F. H.: Linkage analysis in man by somatic cell genetics. Nature 242:165-169, 1973.
- [70065] 8395.Shows, T. B.: . (Abstract) 4th Int. Cong. Hum. Genet., Paris 165,1971.
- [70066] 8396.Shows, T. B.: Genetics of human-mouse somatic cell hybrids: linkageof human genes for isocitrate dehydrogenase and malate dehydrogenase. Biochem.Genet. 7: 193-204, 1972.
- [70067] 8397.Turner, B. M.; Fisher, R. A.; Garthwaite, E.; Whale, R. J.; Harris,H.: An account of two new ICD-S variants not detectable in red bloodcells. Ann. Hum. Genet. 37: 469-476, 1974.
- [70068] 8398.Van Cong, N.: Personal Communication. Paris, France 1976.
- [70069] 8399.Weil, D.; Van Cong, N.; Finaz, C.; Rebourcet, R.; Cochet, C.;de Grouchy, J.; Frezal, J.: Localisation regionale des genes humainsIDH-S, MDH-S, PGK, alpha-GAL, G6PD par l'hybridation cellulaire interspecifique. Hum.Genet. 36: 205-211, 1977.

- [70070] 8400. Auron, P. E.; Webb, A. C.; Rosenwasser, L. J.; Mucci, S. F.; Rich, A.; Wolff, S. M.; Dinarello, C. A.: Nucleotide sequence of human monocyte interleukin 1 precursor cDNA. *Proc. Nat. Acad. Sci.* 81:7907–7911, 1984.
- [70071] 8401. Winter, H.; Labreze, C.; Chapalain, V.; Surleve-Bazeille, J. E.; Mercier, M.; Rogers, M. A.; Taieb, A.; Schweizer, J.: A variable monilethrix phenotype associated with a novel mutation, glu402lys, in the helix termination motif of the type II hair keratin hHb1. *J. Invest. Derm.* 111: 169–172, 1998.
- [70072] 8402. Francke, U.; Foellmer, B. E.; Haynes, B. F.: Chromosome mapping of human cell surface molecules: monoclonal anti-human lymphocyte antibodies 4F2, A3D8, and A1G3 define antigens controlled by different regions of chromosome 11. *Somat. Cell Genet.* 9: 333–344, 1983.
- [70073] 8403. Gottesdiener, K. M.; Karpinski, B. A.; Lindsten, T.; Strominger, J. L.; Jones, N. H.; Thompson, C. B.; Leiden, J. M.: Isolation and structural characterization of the human 4F2 heavy-chain gene, an inducible gene involved in T-lymphocyte activation. *Molec. Cell Biol.* 8: 3809–3819, 1988.
- [70074] 8404. Haynes, B. F.; Hemler, M. E.; Mann, D. L.; Eisenbarth, G. S.; Shelhamer, J.; Mostowski, H. S.; Thomas, C. A.; Stro-

mingler, J. L.; Fauci, A.S.: Characterization of a monoclonal antibody (4F2) that binds to human monocytes and to a subset of activated lymphocytes. J. Immun. 126:1409–1414, 1981.

[70075] 8405. Hemler, M. E.; Strominger, J. L.: Characterization of antigen recognized by the monoclonal antibody (4F2): different molecular forms on human T and B lymphoblastoid cell lines. J. Immun. 129: 623–628, 1982.

[70076] 8406. Lindsten, T.; June, C. H.; Thompson, C. B.; Leiden, J. M.: Regulation of 4F2 heavy-chain gene expression during normal human T-cell activation can be mediated by multiple distinct molecular mechanisms. Molec. Cell. Biol. 8: 3820–3826, 1988.

[70077] 8407. Lumadue, J. A.; Glick, A. B.; Ruddle, F. H.: Cloning, sequence analysis, and expression of the large subunit of the human lymphocyte activation antigen 4F2. Proc. Nat. Acad. Sci. 84: 9204–9208, 1987.

[70078] 8408. Mastroberardino, L.; Spindler, B.; Pfeiffer, R.; Skelly, P. J.; Loffing, J.; Shoemaker, C. B.; Verrey, F.: Amino-acid transport by heterodimers of 4F2hc/CD98 and members of a permease family. Nature 395:288–291, 1998.

[70079] 8409. Copeland, N. G.; Hutchison, K. W.; Jenkins, N. A.: Excision of the DBA ecotropic provirus in dilute coat-color

revertants of mice occurs by homologous recombination involving the viral LTRs. *Cell* 33:379–387, 1983.

[70080] 8410.El-Husseini, A. E.; Vincent, S. R.: Cloning and characterization of a novel RING finger protein that interacts with class V myosins. *J.Biol. Chem.* 274: 19771–19777, 1999.

[70081] 8411.Engle, L. J.; Kennett, R. H.: Cloning, analysis, and chromosomal localization of myosin (MYH12): the human homologue to the mouse dilute gene. *Genomics* 19: 407–416, 1994.

[70082] 8412.Espresifco, E. M.; Coling, D. E.; Tsakraklides, V.; Krogh, K.; Wolenski, J. S.; Kalinec, G.; Kachar, B.: Localization of myosin-V in the centrosome. *Proc. Nat. Acad. Sci.* 95: 8636–8641, 1998.

[70083] 8413.Jenkins, N. A.; Copeland, N. G.; Taylor, B. A.; Lee, B. K.: Dilute(d) coat colour mutation of DNA/2J mice is associated with the site of integration of an ecotropic MuLV genome. *Nature* 293: 370–374, 1981.

[70084] 8414.Jones, J. M.; Huang, J.-D.; Mermall, V.; Hamilton, B. A.; Mooseker, M. S.; Escayg, A.; Copeland, N. G.; Jenkins, N. A.; Meisler, M. H.: The mouse neurological mutant flailar expresses a novel hybrid gene derived by exon shuffling between *Gnb5* and *Myo5a*. *Hum. Molec. Genet.*

9:821–828, 2000.

- [70085] 8415.Karcher, R. L.; Roland, J. T.; Zappacosta, F.; Huddleston, M. J.;Annan, R. S.; Carr, S. A.; Gelfand, V. I.: Cell cycle regulationof myosin-V by calcium/calmodulin-dependent protein kinase II. *Science* 293:1317–1320, 2001.
- [70086] 8416.Dooley, T. P.; Huang, Z.: Genomic organization and DNA sequencesof two human phenol sulfotransferase genes (STP1 and STP2) on theshort arm of chromosome 16. *Biochem. Biophys. Res. Commun.* 228:134–140, 1996.
- [70087] 8417.Her, C.; Raftogianis, R.; Weinshilboum, R. M.: Human phenol sulfotransferaseSTP2 gene: molecular cloning, structural characterization, and chromosomallocalization. *Genomics* 33: 409–420, 1996.
- [70088] 8418.Jones, A. L.; Hagen, M.; Coughtrie, M. W. H.; Roberts, R. C.; Glatt,H.: Human platelet phenolsulfotransferases: cDNA cloning, stableexpression in V79 cells and identification of a novel allelic variantof the phenol-sulfating form. *Biochem. Biophys. Res. Commun.* 208:855–862, 1995.
- [70089] 8419.Barbosa, M. D. F. S.; Nguyen, Q. A.; Tchernev, V. T.; Ashley, J.A.; Detter, J. C.; Blaydes, S. M.; Brandt, S. J.; Chotai, D.; Hodgman,C.; Solari, R. C. E.; Lovett, M.;

Kingsmore, S. F.: Identification of the homologous beige and Chediak–Higashi syndrome genes. *Nature* 382:262–265, 1996.

[70090] 8420. Fukai, K.; Oh, J.; Karim, M. A.; Moore, K. J.; Kandil, H. H.; Ito, H.; Burger, J.; Spritz, R. A.: Homozygosity mapping of the gene for Chediak–Higashi syndrome to chromosome 1q42–q44 in a segment of conserved synteny that includes the mouse beige locus (bg). *Am. J. Hum. Genet.* 59:620–624, 1996.

[70091] 8421. Jackson, I. J.: Homologous pigmentation mutations in human, mouse and other model organisms. *Hum. Molec. Genet.* 6: 1613–1624, 1997.

[70092] 8422. Karim, M. A.; Nagle, D. L.; Kandil, H. H.; Burger, J.; Moore, K. J.; Spritz, R. A.: Mutations in the Chediak–Higashi syndrome gene (CHS1) indicate requirement for the complete 3801 amino acid CHS protein. *Hum. Molec. Genet.* 6: 1087–1089, 1997.

[70093] 8423. Nagle, D. L.; Karim, M. A.; Woolf, E. A.; Holmgren, L.; Bork, P.; Misumi, D. J.; McGrail, S. H.; Dussault, B. J., Jr.; Perou, C. M.; Boissy, R. E.; Duyk, G. M.; Spritz, R. A.; Moore, K. J.: Identification and mutation analysis of the complete gene for Chediak–Higashi syndrome. *Nature–Genet.* 14: 307–311, 1996.

- [70094] 8424. Perou, C. M.; Moore, K. J.; Nagle, D. L.; Misumi, D. J.; Woolf, E. A.; McGrail, S. H.; Holmgren, L.; Brody, T. H.; Dussault, B. J., Jr.; Monroe, C. A.; Duyk, G. M.; Pryor, R. J.; Li, L.; Justice, M. J.; Kaplan, J.: Identification of the murine beige gene by YAC complementation and positional cloning. *Nature Genet.* 13: 303–308, 1996.
- [70095] 8425. Baysal, B. E.; Potkin, S. G.; Farr, J. E.; Higgins, M. J.; Korcz, J.; Gollin, S. M.; James, M. R.; Evans, G. A.; Richard, C. W., III: Bipolar affective disorder partially cosegregates with a balanced t(9;11)(p24;q23.1) chromosomal translocation in a small pedigree. *Am. J. Med. Genet.* 81: 81–91, 1998.
- [70096] 8426. Baysal, B. E.; Willett-Brozick, J. E.; Badner, J. A.; Corona, W.; Ferrell, R. E.; Nimgaonkar, V. L.; Detera-Wadleigh, S. D.: A mannosyltransferase gene at 11q23 is disrupted by a translocation breakpoint that cosegregates with bipolar affective disorder in a small family. *Neurogenetics* 4:43–53, 2002.
- [70097] 8427. Borg, J.-P.; Marchetto, S.; Le Bivic, A.; Ollendorff, V.; Jaulin-Bastard, F.; Saito, H.; Fournier, E.; Adelaide, J.; Margolis, B.; Birnbaum, D.: ERBIN: a basolateral PDZ protein that interacts with the mammalian ERBB2/HER2 receptor. *Nature Cell Biol.* 2: 407–414, 2000.



- [70098] 8428.Dustin, M. L.; Olszowy, M. W.; Holdorf, A. D.; Li, J.; Bromley,S.; Desai, N.; Widder, P.; Rosenberger, F.; van der Merwe, P. A.;Allen, P. M.; Shaw, A. S.: A novel adaptor protein orchestrates receptorpatterning and cytoskeletal polarity in T-cell contacts. *Cell* 94:667–677, 1998.
- [70099] 8429.Kirsch, K. H.; Georgescu, M.–M.; Ishimaru, S.; Hanafusa, H.: CMS:an adapter molecule involved in cytoskeletal rearrangements. *Proc.Nat. Acad. Sci.* 96: 6211–6216, 1999.
- [70100] 8430.Shih, N.–Y.; Li, J.; Karpitskii, V.; Nguyen, A.; Dustin, M. L.;Kanagawa, O.; Miner, J. H.; Shaw, A. S.: Congenital nephrotic syndromein mice lacking CD2–associated protein. *Science* 286: 312–315, 1999.
- [70101] 8431.Loughney, K.; Hill, T. R.; Florio, V. A.; Uher, L.; Rosman, G.J.; Wolda, S. L.; Jones, B. A.; Howard, M. L.; McAllister–Lucas, L.M.; Sonnenburg, W. K.; Francis, S. H.; Corbin, J. D.; Beavo, J. A.;Ferguson, K.: Isolation and characterization of cDNAs encoding PDE5A,a human cGMP–binding, cGMP–specific 3–prime,5–prime–cyclic nucleotidephosphodiesterase. *Gene* 216: 139–147, 1998.
- [70102] 8432.Kim, J. M.; Sato, N.; Yamada, M.; Arai, K.; Masai, H.: Growthregulation of the expression of mouse cDNA and gene encoding a serine/threoninekinase related to Sac–

charomyces cerevisiae CDC7 essential for G(1)/Stransition: structure, chromosomal localization, and expression of mouse gene for S. cerevisiae CDC7-related kinase. J. Biol. Chem. 273:23248–23257, 1998.

[70103] 8433.Wei, M.–H.; Latif, F.; Bader, S.; Kashuba, V.; Chen, J.–Y.; Duh,F.–M.; Sekido, Y.; Lee, C.–C.; Geil, L.; Kuzmin, I.; Zabarovsky, E.;Klein, G.; Zbar, B.; Minna, J. D.; Lerman, M. I.: Construction of a 600–kilobase cosmid clone contig and generation of a transcriptional map surrounding the lung cancer tumor suppressor gene (TSG) locus on human chromosome 3p21.3: progress toward the isolation of a lung–cancer TSG. Cancer Res. 56: 1487–1492, 1996.

[70104] 8434.Dolphin, C. T.; Beckett, D. J.; Janmohamed, A.; Cullingford, T.E.; Smith, R. L.; Shephard, E. A.; Phillips, I. R.: The flavin–containing monooxygenase 2 gene (FMO2) of humans, but not of other primates, encodes a truncated, nonfunctional protein. J. Biol. Chem. 273:30599–30607, 1998.

[70105] 8435.Williams, D. E.; Kelly, J.; Dutchuk, M.: Flavin–containing monooxygenase in pulmonary microsomes of primates. In: Ingelman–Sundberg, M.; Gustafsson, J.–A.; Orrenius, S. (eds.): Drug Metabolizing Enzymes: Genetics, Regulation and Toxicology: Proceedings of the 8th In–

ternational Symposium on Microsomes and Drug Oxidations. Stockholm: Karolinska Institutet 1990. P. 173.

[70106] 8436. Gelb, B. D.; Zhang, J.; Cotter, P. D.; Gershin, I. F.; Desnick, R. J.: Physical mapping of the human connexin 40 (GJA5), flavin-containing monooxygenase 5, and natriuretic peptide receptor A genes on 1q21. *Genomics* 39:409–411, 1997.

[70107] 8437. Overby, L. H.; Buckpitt, A. R.; Lawton, M. P.; Atta-Asafo-Adjei, E.; Schulze, J.; Philpot, R. M.: Characterization of flavin-containing monooxygenase 5 (FMO5) cloned from human and guinea pig: evidence that the unique catalytic properties of FMO5 are not confined to the rabbit ortholog. *Arch. Biochem. Biophys.* 317: 275–284, 1995.

[70108] 8438. Poindexter, K.; Nelson, N.; DuBose, R. F.; Black, R. A.; Cerretti, D. P.: The identification of seven metalloproteinase-disintegrin (ADAM) genes from genomic libraries. *Gene* 237: 61–70, 1999.

[70109] 8439. Li-Hawkins, J.; Lund, E. G.; Turley, S. D.; Russell, D. W.: Disruption of the oxysterol 7- $\alpha$ -hydroxylase gene in mice. *J. Biol. Chem.* 275:16536–16542, 2000.

[70110] 8440. Setchell, K. D. R.; Schwarz, M.; O'Connell, N. C.; Lund, E. G.; Davis, D. L.; Lathe, R.; Thompson, H. R.; Tyson, R. W.; Sokol, R. J.; Russell, D. W.: Identification of a new in-

born error in bileacid synthesis: mutation of the oxysterol 7- $\alpha$ -hydroxylase gene causes severe neonatal liver disease. *J. Clin. Invest.* 102: 1690–1703, 1998.

- [70111] 8441. Granadino, B.; Gallardo, M. E.; Lopez-Rios, J.; Sanz, R.; Ramos, C.; Ayuso, C.; Bovolenta, P.; Rodriguez de Cordoba, S.: Genomic cloning, structure, expression pattern, and chromosomal location of the human *SIX3* gene. *Genomics* 55: 100–105, 1999.
- [70112] 8442. Loosli, F.; Winkler, S.; Wittbrodt, J.: *Six3* overexpression initiates the formation of ectopic retina. *Genes Dev.* 13: 649–654, 1999.
- [70113] 8443. Oliver, G.; Mailhos, A.; Wehr, R.; Copeland, N. G.; Jenkins, N. A.; Gruss, P.: *Six3*, a murine homologue of the *sine oculis* gene, demarcates the most anterior border of the developing neural plate and is expressed during eye development. *Development* 121: 4045–4055, 1995.
- [70114] 8444. Pasquier, L.; Dubourg, C.; Blayau, M.; Lazaro, L.; Le Marec, B.; David, V.; Odent, S.: A new mutation in the *six*-domain of *SIX3* gene causes holoprosencephaly. *Europ. J. Hum. Genet.* 8: 797–800, 2000.
- [70115] 8445. Roessler, E.; Muenke, M.: Holoprosencephaly: a paradigm for the complex genetics of brain development. *J. Inherit. Metab. Dis.* 21: 481–497, 1998.

- [70116] 8446.Gunther, T.; Chen, Z.-F.; Kim, J.; Priemel, M.; Rueger, J. M.;Amling, M.; Moseley, J. M.; Martin, T. J.; Anderson, D. J.; Karsenty,G.: Genetic ablation of parathyroid glands reveals another sourceof parathyroid hormone. Nature 406: 199–203, 2000.
- [70117] 8447.Kanemura, Y.; Hiraga, S.; Arita, N.; Ohnishi, T.; Izumoto, S.;Mori, K.; Matsumura, H.; Yamasaki, M.; Fushiki, S.; Yoshimine, T.: Isolation and expression analysis of a novel human homologue ofthe Drosophila glial cells missing (gcm) gene. FEBS Lett. 442: 151–156,1999.
- [70118] 8448.Schell, U.; Wienberg, J.; Kohler, A.; Bray-Ward, P.; Ward, D. E.;Wilson, W. G.; Allen, W. P.; Lebel, R. R.; Sawyer, J. R.; Campbell,P. L.; Aughton, D. J.; Punnett, H. H.; Lammer, E. J.; Kao, F.-T.;Ward, D. C.; Muenke, M.: Molecular characterization of breakpointsin patients with holoprosencephaly and definition of the HPE2 criticalregion 2p21. Hum. Molec. Genet. 5: 223–229, 1996.
- [70119] 8449.Bouillet, P.; Metcalf, D.; Huang, D. C. S.; Tarlinton, D. M.; Kay,T. W. H.; Kontgen, F.; Adams, J. M.; Strasser, A.: Proapoptotic Bcl-2relative Bim required for certain apoptotic responses, leukocyte homeostasis,and to preclude autoimmunity. Science 286: 1735–1738, 1999.
- [70120] 8450.Bouillet, P.; Purton, J. F.; Godfrey, D. I.; Zhang, L.-C.;

Coultas,L.; Puthalakath, H.; Pellegrini, M.; Cory, S.; Adams, J. M.; Strasser,A.: BH3-only Bcl-2 family member Bim is required for apoptosis of autoreactive thymocytes. *Nature* 415: 922-926, 2002.

[70121] 8451.Bouillet, P.; Zhang, L. C.; Huang, D. C. S.; Webb, G. C.; Bottema,C. D. K.; Shore, P.; Eyre, H. J.; Sutherland, G. R.; Adams, J. M.: Gene structure, alternative splicing, and chromosomal localization of pro-apoptotic Bcl-2 relative Bim. *Mammalian Genome* 12: 163-168,2001.

[70122] 8452.Murray, S.; Halford, S.; Ebenezer, N. D.; Gregory-Evans, C. Y.;Bhattacharya, S. S.: Assignment of BCL2L11 to human chromosome band2p13 with somatic cell and radiation hybrids. *Cytogenet. Cell Genet.* 92:353, 2001.

[70123] 8453.O'Connor, L.; Strasser, A.; O'Reilly, L. A.; Hausmann, G.; Adams,J. M.; Cory, S.; Huang, D. C.: Bim: a novel member of the Bcl-2 family that promotes apoptosis. *EMBO J.* 17: 384-395, 1998.

[70124] 8454.Puthalakath, H.; Huang, D. C. S.; O'Reilly, L. A.; King, S. M.;Strasser, A.: The proapoptotic activity of the Bcl-2 family memberBim is regulated by interaction with the dynein motor complex. *Molec.Cell* 3: 287-296, 1999.

[70125] 8455.Custer, M.; Spindler, B.; Verrey, F.; Murer, H.; Biber, J.: Identification of a new gene product (diphor-1) regu-

lated by dietary phosphate. Am.J. Physiol. 273:  
F801–F806, 1997.

[70126] 8456.Kocher, O.; Comella, N.; Gilchrist, A.; Pal, R.; Tognazzi, K.;Brown, L. F.; Knoll, J. H. M.: PDZK1, a novel PDZ domain-containingprotein up-regulated in carcinomas and mapped to chromosome 1q21,interacts with cMOAT (MRP2), the multidrug resistance-associated protein. Lab.Invest. 79: 1161–1170, 1999.

[70127] 8457.Kocher, O.; Comella, N.; Tognazzi, K.; Brown, L. F.: Identificationand partial characterization of PDZK1: a novel protein containingPDZ interaction domains. Lab. Invest. 78: 117–125, 1998.

[70128] 8458.Wang, S.; Yue, H.; Derin, R. B.; Guggino, W. B.; Li, M.: Accessoryprotein facilitated CFTR–CFTR interaction, a molecular mechanism topotentiate the chloride channel activity. Cell 103: 169–179, 2000.

[70129] 8459.White, K. E.; Biber, J.; Murer, H.; Econs, M. J.: A PDZ domain-containingprotein with homology to Diphor-1 maps to human chromosome 1q21. Ann.Hum. Genet. 62: 287–290, 1998.

[70130] 8460.Pastorek, J.; Pastorekova, S.; Callebaut, I.; Mornon, J. P.; Zelnik,V.; Opavsky, R.; Zat'ovicova, M.; Liao, S.; Portetelle, D.; Stanbridge,E. J.; Zavada, J.; Burny, A.;

Kettmann, R.: Cloning and characterization of MN, a human tumor-associated protein with a domain homologous to carbonic anhydrase and a putative helix-loop-helix DNA binding segment. *Oncogene* 9:2877–2888, 1994.

[70131] 8461. Matsumoto-Taniura, N.; Pirollet, F.; Monroe, R.; Gerace, L.; Westendorf, J. M.: Identification of novel M phase phosphoproteins by expression cloning. *Molec. Biol. Cell* 7: 1455–1469, 1996.

[70132] 8462. Doi, T.; Minami, T.; Itoh, M.; Aburatani, H.; Kawabe, Y.; Kodama, T.; Kondo, N.; Satoh, Y.; Asayama, T.; Imanishi, T.: An alternative form of nucleolysin binds to a T-cluster DNA in the silencer element of platelet factor 4 gene. *Biochem. Biophys. Res. Commun.* 235: 625–630, 1997.

[70133] 8463. Kawakami, A.; Tian, Q.; Duan, X.; Streuli, M.; Schlossman, S. F.; Anderson, P.: Identification and functional characterization of a TIA-1-related nucleolysin. *Proc. Nat. Acad. Sci.* 89: 8681–8685, 1992.

[70134] 8464. Roebroek, A. J. M.; Contreras, B.; Pauli, I. G. L.; Van de Ven, W. J. M.: cDNA cloning, genomic organization, and expression of the human RTN2 gene, a member of a gene family encoding reticulons. *Genomics* 51:98–106, 1998.

[70135] 8465. Kiriakidou, M.; Driscoll, D. A.; Lopez-Guisa, J. M.;



Strauss, J.F., III: Cloning and expression of primate Daxx cDNAs and mapping of the human gene to chromosome 6p21.3 in the MHC region. *DNA Cell Biol.* 16: 1289–1298, 1997.

[70136] 8466. Yang, X.; Khosravi-Far, R.; Chang, H. Y.; Baltimore, D.: Daxx, a novel Fas-binding protein that activates JNK and apoptosis. *Cell* 89:1067–1076, 1997.

[70137] 8467. Parlati, F.; McNew, J. A.; Fukuda, R.; Miller, R.; Sollner, T.H.; Rothman, J. E.: Topological restriction of SNARE-dependent membrane fusion. *Nature* 407: 194–198, 2000.

[70138] 8468. Beckstead, R.; Ortiz, J. A.; Sanchez, C.; Prokopenko, S. N.; Chambon, P.; Losson, R.; Bellen, H. J.: Bonus, a *Drosophila* homolog of TIF1 proteins, interacts with nuclear receptors and can inhibit beta-FTZ-F1-dependent transcription. *Molec. Cell* 7: 753–765, 2001.

[70139] 8469. Le Douarin, B.; Zechel, C.; Garnier, J.-M.; Lutz, Y.; Tora, L.; Pierrat, B.; Heery, D.; Gronemeyer, H.; Chambon, P.; Losson, R.: The N-terminal part of TIF1, a putative mediator of the ligand-dependent activation function (AF-2) of nuclear receptors, is fused to B-raf in the oncogenic protein T18. *EMBO J.* 14: 2020–2033, 1995.

[70140] 8470. Thenot, S.; Henriquet, C.; Rochefort, H.; Cavailles,

V.: Differential interaction of nuclear receptors with the putative human transcriptional coactivator hTIF1. *J. Biol. Chem.* 272: 12062–12068, 1997.

- [70141] 8471. Feral, C.; Mattei, M. G.; Pawlak, A.; Guellaen, G.: Chromosomal localization of three human poly(A)–binding protein genes and four related pseudogenes. *Hum. Genet.* 105: 347–353, 1999.
- [70142] 8472. Houg, A. K.; Maggini, L.; Clement, C. Y.; Reed, G. L.: Identification and structure of activated–platelet protein–1, a protein with RNA–binding domain motifs that is expressed by activated platelets. *Europ. J. Biochem.* 243: 209–218, 1997.
- [70143] 8473. Yang, H.; Duckett, C. S.; Lindsten, T.: iPABP, an inducible poly(A)–binding protein detected in activated human T cells. *Molec. Cell. Biol.* 15: 6770–6776, 1995.
- [70144] 8474. Liu, T.; DeCostanzo, A. J.; Liu, X.; Wang, H.; Hallagan, S.; Moon, R. T.; Malbon, C. C.: G protein signaling from activated rat frizzled–1 to the beta–catenin–Lef–Tcf pathway. *Science* 292: 1718–1722, 2001.
- [70145] 8475. Sagara, N.; Toda, G.; Hirai, M.; Terada, M.; Katoh, M.: Molecular cloning, differential expression, and chromosomal localization of human frizzled–1, frizzled–2, and frizzled–7. *Biochem. Biophys. Res. Commun.* 252:

117–122, 1998.

- [70146] 8476. Tanaka, S.; Akiyoshi, T.; Mori, M.; Wands, J. R.; Sugimachi, K.: A novel frizzled gene identified in human esophageal carcinoma mediates APC/beta-catenin signals. *Proc. Nat. Acad. Sci.* 95: 10164–10169, 1998.
- [70147] 8477. Winklbauer, R.; Medina, A.; Swain, R. K.; Steinbeisser, H.: Frizzled-7 signalling controls tissue separation during *Xenopus* gastrulation. *Nature* 413: 856–860, 2001.
- [70148] 8478. Messina, A.; Oliva, M.; Rosato, C.; Huizing, M.; Ruitenbeek, W.; van den Heuvel, L. P.; Forte, M.; Rocchi, M.; De Pinto, V.: Mapping of the human voltage-dependent action channel isoforms 1 and 2 reconsidered. *Biochem. Biophys. Res. Commun.* 255: 707–710, 1999.
- [70149] 8479. Parnet, P.; Garka, K. E.; Bonnert, T. P.; Dower, S. K.; Sims, J. E.: IL-1Rrp is a novel receptor-like molecule similar to the type I interleukin-1 receptor and its homologues T1/ST2 and IL-1R AcP. *J. Biol. Chem.* 271: 3967–3970, 1996.
- [70150] 8480. Torigoe, K.; Ushio, S.; Okura, T.; Kobayashi, S.; Tanai, M.; Kunikata, T.; Murakami, T.; Sanou, O.; Kojima, H.; Fujii, M.; Ohta, T.; Ikeda, M.; Ikegami, H.; Kurimoto, M.: Purification and characterization of the human interleukin-18 receptor. *J. Biol. Chem.* 272: 25737–25742, 1997.

- [70151] 8481.Akiba, H.; Atsuta, M.; Yagita, H.; Okumura, K.: Identification of rat OX40 ligand by molecular cloning. Biochem. Biophys. Res. Commun. 251:131–136, 1998.
- [70152] 8482.Baum, P. R.; Gayle, R. B., III; Ramsdell, F.; Srinivasan, S.; Sorensen, R. A.; Watson, M. L.; Seldin, M. F.; Baker, E.; Sutherland, G. R.; Clifford, K. N.; Alderson, M. R.; Goodwin, R. G.; Fanslow, W. C.: Molecular characterization of murine and human OX40/OX40 ligand systems: identification of a human OX40 ligand as the HTLV-1-regulated protein gp34. EMBO J. 13: 3992–4001, 1994.
- [70153] 8483.Godfrey, W. R.; Fagnoni, F. F.; Harara, M. A.; Buck, D.; Engleman, E. G.: Identification of a human OX-40 ligand, a costimulator of CD4+ T cells with homology to tumor necrosis factor. J. Exp. Med. 180:757–762, 1994.
- [70154] 8484.Malmstrom, V.; Shipton, D.; Singh, B.; Al-Shamkhani, A.; Puklavec, M. J.; Barclay, A. N.; Powrie, F.: CD134L expression on dendritic cells in the mesenteric lymph nodes drives colitis in T cell-restored SCID mice. J. Immun. 166: 6972–6981, 2001.
- [70155] 8485.Miura, S.; Ohtani, K.; Numata, N.; Niki, M.; Ohbo, K.; Ina, Y.; Gojobori, T.; Tanaka, Y.; Tozawa, H.; Nakamura, M.; Sugamura, K.: Molecular cloning and characterization of a novel glycoprotein, gp34, that is specifically induced by

the human T-cell leukemia virus type I transactivator p40-tax. *Molec. Cell. Biol.* 11: 1313–1325, 1991.

[70156] 8486.Cretney, E.; Takeda, K.; Yagita, H.; Glaccum, M.; Peschon, J. J.; Smyth, M. J.: Increased susceptibility to tumor initiation and metastasis in TNF-related apoptosis-inducing ligand-deficient mice. *J. Immun.* 168:1356–1361, 2002.

[70157] 8487.Degli-Esposti, M. A.; Dougall, W. C.; Smolak, P. J.; Waugh, J.Y.; Smith, C. A.; Goodwin, R. G.: The novel receptor TRAIL-R4 induces NF- $\kappa$ B and protects against TRAIL-mediated apoptosis, yet retains an incomplete death domain. *Immunity* 7: 813–820, 1997.

[70158] 8488.Nagashima, M.; Shiseki, M.; Miura, K.; Hagiwara, K.; Linke, S.P.; Pedoux, R.; Wang, X. W.; Yokota, J.; Riabowol, K.; Harris, C.C.: DNA damage-inducible gene p33<sup>ING2</sup> negatively regulates cell proliferation through acetylation of p53. *Proc. Nat. Acad. Sci.* 98: 9671–9676, 2001.

[70159] 8489.Shimada, Y.; Saito, A.; Suzuki, M.; Takahashi, E.; Horie, M.: Cloning of a novel gene (ING1L) homologous to ING1, a candidate tumor suppressor. *Cytogenet. Cell Genet.* 83: 232–235, 1998.

[70160] 8490.Bardoni, B.; Giglio, S.; Schenck, A.; Rocchi, M.; Mandel, J. L.: Assignment of NUFIP1 (nuclear FMRP interacting

protein 1) gene to chromosome 13q14 and assignment of a pseudogene to chromosome 6q12. *Cytogenet. Cell Genet.* 89: 11–13, 2000.

- [70161] 8491. Bardoni, B.; Schenck, A.; Mandel, J. L.: A novel RNA-binding nuclear protein that interacts with the fragile X mental retardation (FMR1) protein. *Hum. Molec. Genet.* 8: 2557–2566, 1999.
- [70162] 8492. Blagitko, N.; Schulz, U.; Schinzel, A. A.; Ropers, H.-H.; Kalscheuer, V. M.: Gamma-2-COP, a novel imprinted gene on chromosome 7q32, defines a new imprinting cluster in the human genome. *Hum. Molec. Genet.* 8: 2387–2396, 1999.
- [70163] 8493. Maw, M. A.; Corbeil, D.; Koch, J.; Hellwig, A.; Wilson-Wheeler, J. C.; Bridges, R. J.; Kumaramanickavel, G.; John, S.; Nancarrow, D.; Roper, K.; Weigmann, A.; Huttner, W. B.; Denton, M. J.: A frameshift mutation in prominin (mouse)-like 1 causes human retinal degeneration. *Hum. Molec. Genet.* 9: 27–34, 2000.
- [70164] 8494. Miraglia, S.; Godfrey, W.; Yin, A. H.; Atkins, K.; Warnke, R.; Holden, J. T.; Bray, R. A.; Waller, E. K.; Buck, D. W.: A novel five-transmembrane hematopoietic stem cell antigen: isolation, characterization, and molecular cloning. *Blood* 90: 5013–5021, 1997.

- [70165] 8495. Yin, A. H.; Miraglia, S.; Zanjani, E. D.; Almeida-Porada, G.; Ogawa, M.; Leary, A. G.; Olweus, J.; Kearney, J.; Buck, D. W.: AC133, a novel marker for human hematopoietic stem and progenitor cells. *Blood* 90:5002–5012, 1997.
- [70166] 8496. Taylor, S. S.; Ha, E.; McKeon, F.: The human homologue of Bub3 is required for kinetochore localization of Bub1 and a Mad3/Bub1-related protein kinase. *J. Cell Biol.* 142: 1–11, 1998.
- [70167] 8497. Bonne, S.; van Hengel, J.; van Roy, F.: Assignment of the plakophilin-2 gene (PKP2) and a plakophilin-2 pseudogene (PKP2P1) to human chromosome bands 12p11 and 12p13, respectively, by in situ hybridization. *Cytogenet. Cell Genet.* 88: 286–287, 2000.
- [70168] 8498. Mertens, C.; Kuhn, C.; Franke, W. W.: Plakophilins 2a and 2b: constitutive proteins of dual location in the karyoplasm and the desmosomal plaque. *J. Cell Biol.* 135: 1009–1025, 1996.
- [70169] 8499. Schmidt, A.; Langbein, L.; Pratzel, S.; Rode, M.; Rackwitz, H.-R.; Franke, W. W.: Plakophilin 3—a novel cell-type-specific desmosomal plaque protein. *Differentiation* 64: 291–306, 1999.
- [70170] 8500. Bergstein, I.; Eisenberg, L. M.; Bhalerao, J.; Jenkins,

N. A.; Copeland, N. G.; Osborne, M. P.; Bowcock, A. M.; Brown, A. M. C.: Isolation of two novel WNT genes, WNT14 and WNT15, one of which (WNT15) is closely linked to WNT3 on human chromosome 17q21. *Genomics* 46:450–458, 1997.

[70171] 8501. Hartmann, C.; Tabin, C. J.: Wnt-14 plays a pivotal role in inducing synovial joint formation in the developing appendicular skeleton. *Cell* 104:341–351, 2001.

[70172] 8502. Saitoh, T.; Hirai, M.; Katoh, M.: Molecular cloning and characterization of WNT3A and WNT14 clustered in human chromosome 1q42 region. *Biochem. Biophys. Res. Commun.* 284: 1168–1175, 2001.

[70173] 8503. Habuchi, T.; Luscombe, M.; Elder, P. A.; Knowles, M. A.: Structure and methylation-based silencing of a gene (DBCCR1) within a candidate bladder cancer tumor suppressor region at 9q32–q33. *Genomics* 48:277–288, 1998.

[70174] 8504. Habuchi, T.; Yoshida, O.; Knowles, M. A.: A novel candidate tumour suppressor locus at 9q32–33 in bladder cancer: localization of the candidate region within a single 840 kb YAC. *Hum. Molec. Genet.* 6:913–919, 1997.

[70175] 8505. Nishiyama, H.; Takahashi, T.; Kakehi, Y.; Habuchi, T.; Knowles, M. A.: Homozygous deletion at the 9q32–33



candidate tumor suppressor locus in primary human bladder cancer. *Genes Chromosomes Cancer* 26:171–175, 1999.

- [70176] 8506. Murphy, M.; Pykett, M. J.; Harnish, P.; Zang, K. D.; George, D.L.: Identification and characterization of genes differentially expressed in meningiomas. *Cell Growth Differ.* 4: 715–722, 1993.
- [70177] 8507. Oh, Y.; Nagalla, S. R.; Yamanaka, Y.; Kim, H.-S.; Wilson, E.; Rosenfeld, R. G.: Synthesis and characterization of insulin-like growth factor-binding protein (IGFBP)-7. *J. Biol. Chem.* 271: 30322–30325, 1996.
- [70178] 8508. Swisshelm, K.; Ryan, K.; Tsuchiya, K.; Sager, R.: Enhanced expression of an insulin growth factor-like binding protein (mac25) in senescent human mammary epithelial cells and induced expression with retinoic acid. *Proc. Nat. Acad. Sci.* 92: 4472–4476, 1995.
- [70179] 8509. Yamauchi, T.; Umeda, F.; Masakado, M.; Isaji, M.; Mizushima, S.; Nawata, H.: Purification and molecular cloning of prostacyclin-stimulating factor from serum-free conditioned medium of human diploid fibroblast cells. *Biochem. J.* 303: 591–598, 1994.
- [70180] 8510. Ayala-Madrigal, M. L.; Doerr, S.; Ramirez-Duenas, M. L.; Hansmann, I.: Assignment of KPNA4 and KPNB1 en-

coding karyopherin alpha 4 and beta 1 to human chromosome bands 11q22 and 17q21 respectively, by in situ hybridization. *Cytogenet. Cell Genet.* 89: 258–259, 2000.

[70181] 8511. Bayliss, R.; Littlewood, T.; Stewart, M.: Structural basis for the interaction between FxFG nucleoporin repeats and importin-beta in nuclear trafficking. *Cell* 102: 99–108, 2000.

[70182] 8512. Chi, N. C.; Adam, E. J. H.; Adam, S. A.: Sequence and characterization of cytoplasmic nuclear protein import factor p97. *J. Cell Biol.* 130:265–274, 1995.

[70183] 8513. Gorlich, D.; Kostka, S.; Kraft, R.; Dingwall, C.; Laskey, R. A.; Hartmann, E.; Prehn, S.: Two different subunits of importin cooperate to recognize nuclear localization signals and bind them to the nuclear envelope. *Curr. Biol.* 5: 383–392, 1995.

[70184] 8514. Kutay, U.; Izaurralde, E.; Bischoff, F. R.; Mattaj, I. W.; Gorlich, D.: Dominant-negative mutants of importin-beta block multiple pathways of import and export through the nuclear pore complex. *EMBO J.* 16:1153–1163, 1997.

[70185] 8515. Matsuda, Y.; Hamatani, K.; Itoh, M.; Takahashi, E.; Araki, R.; Abe, M.: Localization of the importin-beta gene to mouse chromosome 11D and rat chromosome 10q32.1. *Genomics* 36: 213–215, 1996.

- [70186] 8516.Thornton, C.; Snowden, M. A.; Carling, D.: Identification of anovel AMP-activated protein kinase beta subunit isoform that is highlyexpressed in skeletal muscle. J. Biol. Chem. 273: 12443-12450, 1998.
- [70187] 8517.Woods, A.; Cheung, P. C. F.; Smith, F. C.; Davison, M. D.; Scott,J.; Beri, R. K.; Carling, D.: Characterization of AMP-activated proteinkinase beta and gamma subunits: assembly of the heterotrimeric complexin vitro. J. Biol. Chem. 271: 10282-10290, 1996.
- [70188] 8518.Gao, G.; Fernandez, C. S.; Stapleton, D.; Auster, A. S.; Widmer,J.; Dyck, J. R. B.; Kemp, B. E.; Witters, L. A.: Non-catalytic beta-and gamma-subunit isoforms of the 5-prime-AMP-activated protein kinase. J.Biol. Chem. 271: 8675-8681, 1996.
- [70189] 8519.Cheung, P. C. F.; Salt, I. P.; Davies, S. P.; Hardie, D. G.; Carling,D.: Characterization of AMP-activated protein kinase gamma-subunitisoforms and their role in AMP binding. Biochem. J. 346: 659-669,2000.
- [70190] 8520.Gollob, M. H.; Green, M. S.; Tang, A. S.-L.; Gollob, T.; Karibe,A.; Al Sayegh, A. H.; Ahmad, F.; Lozado, R.; Shah, G.; Fananapazir,L.; Bachinski, L. L.; Roberts, R.: Identification of a gene responsiblefor familial Wolff-Parkinson-White syndrome. New Eng. J. Med.

344:1823–1831, 2001. Note: Erratum: New Eng. J. Med.  
345: 552 only, 2001.

- [70191] 8521.Lang, T.; Yu, L.; Qiang, T.; Jiang, J.; Chen, Z.; Xin, Y.; Liu,G.; Zhao, S.: Molecular cloning, genomic organization, and mapping of PRKAG2, a heart abundant gamma-2 subunit of 5-prime-AMP-activated protein kinase, to human chromosome 7q36. *Genomics* 70: 258–263,2000.
- [70192] 8522.Sinha, S. C.; Nair, M.; Gambhir, D. S.; Mohan, J. C.; Kaul, U.A.; Arora, R.: Genetically transmitted ventricular pre-excitation in a family with hypertrophic cardiomyopathy. *Indian Heart J.* 52:76–78, 2000.
- [70193] 8523.Lee, S.-J.: Expression of growth/differentiation factor 1 in the nervous system: conservation of a bicistronic structure. *Proc. Nat.Acad. Sci.* 88: 4250–4254, 1991.
- [70194] 8524.Yamagoe, S.; Akasaka, T.; Uchida, T.; Hachiya, T.; Okabe, T.; Yamakawa,Y.; Arai, T.; Mizuno, S.; Suzuki, K.: Expression of a neutrophil chemotactic protein LECT2 in human hepatocytes revealed by immunochemical studies using polyclonal and monoclonal antibodies to a recombinant LECT2. *Biochem. Biophys. Res. Commun.* 237: 116–120, 1997.
- [70195] 8525.Yamagoe, S.; Kameoka, Y.; Hashimoto, K.; Mizuno, S.; Suzuki, K.: Molecular cloning, structural characteriza-

tion, and chromosomal mapping of the human LECT2 gene. *Genomics* 48: 324–329, 1998.

[70196] 8526. Yamagoe, S.; Mizuno, S.; Suzuki, K.: Molecular cloning of human and bovine LECT2 having a neutrophil chemotactic activity and its specific expression in the liver. *Biochim. Biophys. Acta* 1396: 105–113, 1998.

[70197] 8527. Yamagoe, S.; Watanabe, T.; Mizuno, S.; Suzuki, K.: The mouse LECT2 gene: cloning of cDNA and genomic DNA, structural characterization and chromosomal localization. *Gene* 216: 171–178, 1998.

[70198] 8528. Yamagoe, S.; Yamakawa, Y.; Matsuo, Y.; Minowada, J.; Mizuno, S.; Suzuki, K.: Purification and primary amino acid sequence of a novel neutrophil chemotactic factor LECT2. *Immun. Lett.* 52: 9–13, 1996.

[70199] 8529. Kullak-Ublick, G.-A.; Beuers, U.; Meier, P. J.; Domdey, H.; Paumgartner, G.: Assignment of the human organic anion transporting polypeptide (OATP) gene to chromosome 12p12 by fluorescence in situ hybridization. *J. Hepatol.* 25: 985–987, 1996.

[70200] 8530. Kullak-Ublick, G. A.; Hagenbuch, B.; Stieger, B.; Schteingart, C. D.; Hofmann, A. F.; Wolkoff, A. W.; Meier, P. J.: Molecular and functional characterization of an organic anion transporting polypeptide cloned from human liver.

:Gastroenterology 109: 1274–1282, 1995.

- [70201] 8531.Ohyama, C.; Smith, P. L.; Angata, K.; Fukuda, M. N.; Lowe, J. B.;Fukuda, M.: Molecular cloning and expression of GDP-D-mannose-4,6-dehydratase,a key enzyme for fucose metabolism defective in Lec13 cells. J. Biol.Chem. 273: 14582–14587, 1998.
- [70202] 8532.El-Husseini, A. E.-D.; Schnell, E.; Chetkovich, D. M.; Nicoll,R. A.; Bredt, D. S.: PSD-95 involvement in maturation of excitatorysynapses. Science 290: 1364–1368, 2000.
- [70203] 8533.El-Husseini, A. E.-D.; Schnell, E.; Dakoji, S.; Sweeney, N.; Zhou,Q.; Prange, O.; Gauthier-Campbell, C.; Aguilera-Moreno, A.; Nicoll,R. A.; Bredt, D. S.: Synaptic strength regulated by palmitate cyclinon PSD-95. Cell 108: 849–863, 2002.
- [70204] 8534.Kim, E.; Cho, K.-O.; Rothschild, A.; Sheng, M.: Heteromultimerizationand NMDA receptor-clustering activity of Chapsyn-110, a member ofthe PSD-95 family of proteins. Neuron 17: 103–113, 1996.
- [70205] 8535.Kim, E.; Niethammer, M.; Rothschild, A.; Jan, Y. N.; Sheng, M.: Clustering of Shaker-type K<sup>+</sup> channels by interaction with a familyof membrane-associated guanylate kinases. Nature 378: 85–88, 1995.

- [70206] 8536.Kistner, U.; Wenzel, B. M.; Veh, R. W.; Cases–Langhoff, C.; Garner,A. M.; Appeltauer, U.; Voss, B.; Gundelfinger, E. D.; Garner, C. C.: SAP90, a rat presynaptic protein related to the product of the Drosophilatumor suppressor gene, dLg–A. J. Biol. Chem. 268: 4580–4583, 1993.
- [70207] 8537.Migaud, M.; Charlesworth, P.; Dempster, M.; Webster, L. C.; Watabe,A. M.; Makhinson, M.; He, Y.; Ramsay, M. F.; Morris, R. G. M.; Morrison,J. H.; O'Dell, T. J.; Grant, S. G. N.: Enhanced long–term potentiationand impaired learning in mice with mutant postsynaptic density–95protein. Nature 396: 433–439, 1998.
- [70208] 8538.Sattler, R.; Xiong, Z.; Lu, W.–Y.; Hafner, M.; MacDonald, J. F.;Tymianski, M.: Specific coupling of NMDA receptor activation to nitricoxide neurotoxicity by PSD–95 protein. Science 284: 1845–1848, 1999.
- [70209] 8539.Stathakis, D. G.; Hoover, K. B.; You, Z.; Bryant, P. J.: Humanpostsynaptic density–95 (PSD95): location of the gene (DLG4) and possiblefunction in nonneural as well as in neural tissues. Genomics 44:71–82, 1997.
- [70210] 8540.Strippoli, P.; Petrini, M.; Lenzi, L.; Carinci, P.; Zannotti, M.: The murine DSCR1–like (Down syndrome candidate region 1) gene family:conserved synteny with the hu–

man orthologous genes. *Gene* 257: 223–232, 2000.

[70211] 8541. Yang, J.; Rothermel, B.; Vega, R. B.; Frey, N.; McKinsey, T. A.; Olson, E. N.; Bassel-Duby, R.; Williams, R. S.: Independent signals control expression of the calcineurin inhibitory proteins MCIP1 and MCIP2 in striated muscles. *Circ. Res.* 87: 61e–68e, 2000.

[70212] 8542. Zheng, B.; Larkin, D. W.; Albrecht, U.; Sun, Z. S.; Sage, M.; Eichele, G.; Lee, C. C.; Bradley, A.: The mPer2 gene encodes a functional component of the mammalian circadian clock. *Nature* 400: 169–173, 1999.

[70213] 8543. Nakamura, H.; Sudo, T.; Tsuiki, H.; Miyake, H.; Morisaki, T.; Sasaki, J.; Masuko, N.; Kochi, M.; Ushio, Y.; Saya, H.: Identification of a novel human homolog of the *Drosophila* dlg, P-dlg, specifically expressed in the gland tissues and interacting with p55. *FEBS Lett.* 433: 63–67, 1998.

[70214] 8544. Abrieu, A.; Magnaghi-Jaulin, L.; Kahana, J. A.; Peter, M.; Castro, A.; Vigneron, S.; Lorca, T.; Cleveland, D. W.; Labbe, J.-C.: Mps1 is a kinetochore-associated kinase essential for the vertebrate mitotic checkpoint. *Cell* 106: 83–93, 2001.

[70215] 8545. Fisk, H. A.; Winey, M.: The mouse Mps1p-like kinase regulates centrosome duplication. *Cell* 106: 95–104, 2001.



- [70216] 8546.Lauze, E.; Stoelcker, B.; Luca, F. C.; Weiss, E.; Schutz, A. R.; Winey, M.: Yeast spindle pole body duplication gene MPS1 encodes an essential dual specificity protein kinase. EMBO J. 14: 1655–1663,1995.
- [70217] 8547.Lindberg, R. A.; Fischer, W. H.; Hunter, T.: Characterization of a human protein threonine kinase isolated by screening an expression library with antibodies to phosphotyrosine. Oncogene 8: 351–359,1993.
- [70218] 8548.Funderburgh, J. L.; Perchellet, A. L.; Swiergiel, J.; Conrad, G.W.; Justice, M. J.: Keratocan (Kera), a corneal keratan sulfate proteoglycan, maps to the distal end of mouse chromosome 10. Genomics 52: 110–111,1998.
- [70219] 8549.Liu, C.-Y.; Shiraishi, A.; Kao, C. W.-C.; Converse, R. L.; Funderburgh, J. L.; Corpuz, L. M.; Conrad, G. W.; Kao, W. W.-Y.: The cloning of mouse keratocan cDNA and genomic DNA and the characterization of its expression during eye development. J. Biol. Chem. 273: 22584–22588,1998.
- [70220] 8550.Tasheva, E. S.; Funderburgh, J. L.; Funderburgh, M. L.; Corpuz, L. M.; Conrad, G. W.: Structure and sequence of the gene encoding human keratocan. DNA Seq. 10: 67–74, 1999.
- [70221] 8551.Tasheva, E. S.; Pettenati, M.; Von Kap-Her, C.; Con-

rad, G. W.: Assignment of keratocan gene (KERA) to human chromosome band 12q22 by in situ hybridization. *Cytogenet. Cell Genet.* 88: 244–245, 2000.

[70222] 8552. Bruick, R. K.: Expression of the gene encoding the proapoptotic Nip3 protein is induced by hypoxia. *Proc. Nat. Acad. Sci.* 97: 9082–9087, 2000.

[70223] 8553. Vaughan, K. T.; Mikami, A.; Paschal, B. M.; Holzbaur, E. L. F.; Hughes, S. M.; Echeverri, C. J.; Moore, K. J.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.; Vallee, R. B.: Multiple mouse chromosomal loci for dynein-based motility. *Genomics* 36: 29–38, 1996.

[70224] 8554. Soyombo, A. A.; Hofmann, S. L.: Molecular cloning and expression of palmitoyl-protein thioesterase 2 (PPT2), a homolog of lysosomal palmitoyl-protein thioesterase with a distinct substrate specificity. *J. Biol. Chem.* 272: 27456–27463, 1997.

[70225] 8555. Richter-Cook, N. J.; Dever, T. E.; Hensold, J. O.; Merrick, W. C.: Purification and characterization of a new eukaryotic protein translation factor: eukaryotic initiation factor 4H. *J. Biol. Chem.* 273: 7579–7587, 1998.

[70226] 8556. Hoogenraad, C. C.; Eussen, B. H. J.; Langeveld, A.; van Haperen, R.; Winterberg, S.; Wouters, C. H.; Grosveld, F.; De Zeeuw, C. I.; Galjart, N.: The murine CYLN2 gene:

genomic organization, chromosomelocalization, and comparison to the human gene that is located within the 7q11.23 Williams syndrome critical region. *Genomics* 53: 348–358, 1998.

[70227] 8557. Araujo, H.; Danziger, N.; Cordier, J.; Glowinski, J.; Chneiweiss, H.: Characterization of PEA-15, a major substrate for protein kinase C in astrocytes. *J. Biol. Chem.* 268: 5911–5920, 1993.

[70228] 8558. Bera, T. K.; Guzman, R. C.; Miyamoto, S.; Panda, D. K.; Sasaki, M.; Hanyu, K.; Enami, J.; Nandi, S.: Identification of a mammary transforming gene (MAT1) associated with mouse mammary carcinogenesis. *Proc. Nat. Acad. Sci.* 91: 9789–9793, 1994.

[70229] 8559. Condorelli, G.; Vigliotta, G.; Iavarone, C.; Caruso, M.; Tocchetti, C. G.; Andreozzi, F.; Cafieri, A.; Tecce, M. F.; Formisano, P.; Beguinot, L.; Beguinot, F.: PED/PEA-15 gene controls glucose transport and is overexpressed in type 2 diabetes mellitus. *EMBO J.* 17: 3858–3866, 1998.

[70230] 8560. Danziger, N.; Yokoyama, M.; Jay, T.; Cordier, J.; Glowinski, J.; Chneiweiss, H.: Cellular expression, developmental regulation, and phylogenetic conservation of PEA-15, the astrocytic major phosphoprotein and protein kinase C substrate. *J. Neurochem.* 64: 1016–1025, 1995.

- [70231] 8561. Estelles, A.; Yokoyama, M.; Nothias, F.; Vincent, J.-D.; Glowinski, J.; Vernier, P.; Chneiweiss, H.: The major astrocytic phosphoprotein PEA-15 is encoded by two mRNAs conserved on their full length in mouse and human. *J. Biol. Chem.* 271: 14800–14806, 1996.
- [70232] 8562. Hwang, S.; Kuo, W.-L.; Cochran, J. F.; Guzman, R. C.; Tsukamoto, T.; Bandyopadhyay, G.; Myambo, K.; Collins, C. C.: Assignment of HMAT1, the human homolog of the murine mammary transforming gene (MAT1) associated with tumorigenesis, to 1q21.1, a region frequently gained in human breast cancers. *Genomics* 42: 540–542, 1997.
- [70233] 8563. Wolford, J. K.; Bogardus, C.; Ossowski, V.; Prochazka, M.: Molecular characterization of the human PEA15 gene on 1q21–q22 and association with type 2 diabetes mellitus in Pima Indians. *Gene* 241: 143–148, 2000.
- [70234] 8564. Aurrand-Lions, M.; Galland, F.; Bazin, H.; Zakharyev, V. M.; Imhof, B. A.; Naquet, P.: Vanin-1, a novel GPI-linked perivascular molecule involved in thymus homing. *Immunity* 5: 391–405, 1996.
- [70235] 8565. Galland, F.; Malergue, F.; Bazin, H.; Mattei, M. G.; Aurrand-Lions, M.; Theillet, C.; Naquet, P.: Two human genes related to murine vanin-1 are located on the long

arm of human chromosome 6. *Genomics* 53:203–213, 1998.

- [70236] 8566. Martin, F.; Malergue, F.; Pitari, G.; Philippe, J. M.; Philips, S.; Chabret, C.; Granjeaud, S.; Mattei, M. G.; Mungall, A. J.; Naquet, P.; Galland, F.: Vanin genes are clustered (human 6q22–24 and mouse 10A2B1) and encode isoforms of pantetheinase ectoenzymes. *Immunogenetics* 53:296–306, 2001.
- [70237] 8567. Pitari, G.; Malergue, F.; Martin, F.; Philippe, J. M.; Massucci, M. T.; Chabret, C.; Maras, B.; Dupre, S.; Naquet, P.; Galland, F.: Pantetheinase activity of membrane-bound vanin-1: lack of free cysteamine in tissues of vanin-1 deficient mice. *FEBS Lett.* 483: 149–154, 2000.
- [70238] 8568. Hendrich, B.; Bird, A.: Identification and characterization of a family of mammalian methyl-CpG binding proteins. *Molec. Cell. Biol.* 18:6538–6547, 1998.
- [70239] 8569. Rasooly, R. S.: Personal Communication. Baltimore, Md. 2/23/1999.
- [70240] 8570. Zhang, Y.; Ng, H.-H.; Erdjument-Bromage, H.; Tempst, P.; Bird, A.; Reinberg, D.: Analysis of the NuRD subunits reveals a histone deacetylase core complex and a connection with DNA methylation. *Genes Dev.* 13:1924–1935, 1999.

- [70241] 8571. Bellacosa, A.; Cicchillitti, L.; Schepis, F.; Riccio, A.; Yeung, A. T.; Matsumoto, Y.; Golemis, E. A.; Genuardi, M.; Neri, G.: MED1, a novel human methyl-CpG-binding endonuclease, interacts with DNA mismatch repair protein MLH1. *Proc. Nat. Acad. Sci.* 96: 3969–3974, 1999.
- [70242] 8572. Boland, C. R.; Thibodeau, S. N.; Hamilton, S. R.; Sidransky, D.; Eshleman, J. R.; Burt, R. W.; Meltzer, S. J.; Rodriguez-Bigas, M. A.; Fodde, R.; Ranzani, G. N.; Srivastava, S.: A National Cancer Institute workshop on microsatellite instability for cancer detection and familial predisposition: development of international criteria for the determination of microsatellite instability in colorectal cancer. *Cancer Res.* 58: 5248–5257, 1998.
- [70243] 8573. Hendrich, B.; Hardeland, U.; Ng, H.-H.; Jiricny, J.; Bird, A.: The thymine glycosylase MBD4 can bind to the product of deamination at methylated CpG sites. *Nature* 401: 301–304, 1999.
- [70244] 8574. Millar, C. B.; Guy, J.; Sansom, O. J.; Selfridge, J.; MacDougall, E.; Hendrich, B.; Keightley, P. D.; Bishop, S. M.; Clarke, A. R.; Bird, A.: Enhanced CpG mutability and tumorigenesis in MBD4-deficient mice. *Science* 297: 403–405, 2002.
- [70245] 8575. Riccio, A.; Aaltonen, L. A.; Godwin, A. K.; Loukola,

A.; Percesepe,A.; Salovaara, R.; Masciullo, V.; Genuardi M.; Paravatou–Petsotas,M.; Bassi, D. E.; Ruggeri, B. A.; Klein–Szanto, A. J. P.; Testa, J.R.; Neri, G.; Bellacosa, A.: The DNA repair gene MBD4 (MED1) is mutated in human carcinomas with microsatellite instability. (Letter) *Nature Genet.* 23: 266–268, 1999.

[70246] 8576.Duncan, L. M.; Deeds, J.; Hunter, J.; Shao, J.; Holmgren, L. M.; Woolf, E. A.; Tepper, R. I.; Shyjan, A. W.: Down–regulation of the novel gene melastatin correlates with potential for melanoma metastasis. *Cancer Res.* 58: 1515–1520, 1998.

[70247] 8577.Fang, D.; Setaluri, V.: Expression and up–regulation of alternatively spliced transcripts of melastatin, a melanoma metastasis–related gene, in human melanoma cells. *Biochem. Biophys. Res. Commun.* 279: 53–61, 2000.

[70248] 8578.Hunter, J. J.; Shao, J.; Smutko, J. S.; Dussault, B. J.; Nagle,D. L.; Woolf, E. A.; Holmgren, L. M.; Moore, K. J.; Shyjan, A. W.: Chromosomal localization and genomic characterization of the mouse melastatin gene (*Mlsn1*). *Genomics* 54: 116–123, 1998.

[70249] 8579.Xu, X. Z.; Moebius, F.; Gill, D. L.; Montell, C.: Regulation of melastatin, a TRP–related protein, through interaction with a cytoplasmic isoform. *Proc. Nat. Acad. Sci.* 98:

10692–10697, 2001.

- [70250] 8580. Kalitsis, P.; Earle, E.; Fowler, K. J.; Choo, K. H. A.: Bub3 genedisruption in mice reveals essential mitotic spindle checkpoint functionduring early embryogenesis. *Genes Dev.* 14: 2277–2282, 2000.
- [70251] 8581. Ardley, H. C.; Moynihan, T. P.; Thompson, J.; Leek, J. P.; Markham, A. F.; Robinson, P. A.: Rapid isolation of genomic clones for individualmembers of human multigene families: identification and localisationof UBE2L4, a novel member of a ubiquitin conjugating enzyme dispersedgene family. *Cytogenet. Cell Genet.* 79: 188–192, 1997.
- [70252] 8582. Moynihan, T. P.; Cole, C. G.; Dunham, I.; O'Neil, L.; Markham, A. F.; Robinson, P. A.: Fine–mapping, genomic organization, and transcriptanalysis of the human ubiquitin–conjugating enzyme gene UBE2L3. *Genomics* 51:124–127, 1998.
- [70253] 8583. Hu, M. C.–T.; Qiu, W. R.; Wang, Y.–P.; Hill, D.; Ring, B. D.; Scully, S.; Bolon, B.; DeRose, M.; Luethy, R.; Simonet, W. S.; Arakawa, T.; Danilenko, D. M.: FGF–18, a novel member of the fibroblast growthfactor family, stimulates hepatic and intestinal proliferation. *Molec. Cell. Biol.* 18: 6063–6074, 1998.
- [70254] 8584. Baens, M.; Chaffanet, M.; Cassiman, J.–J.; van den



Berghe, H.;Marynen, P.: Construction and evaluation of a hncDNA library of human12 transcribed sequences derived from a somatic cell hybrid. *Genomics* 16:214–218, 1993.

[70255] 8585.Fearnley, I. M.; Finel, M.; Skehel, J. M.; Walker, J. E.: NADH:ubiquinoneoxidoreductase from bovine heart mitochondria. *Biochem. J.* 278:821–829, 1991.

[70256] 8586.Baens, M.; Chaffanet, M.; Cassiman, J.-J.; van den Berghe, H.;Marynen, P.: Construction and evaluation of a hncDNA library of human12p transcribed sequences derived from a somatic cell hybrid. *Genomics* 16:214–218, 1993.

[70257] 8587.Kawasaki, H.; Springett, G. M.; Toki, S.; Canales, J. J.; Harlan,P.; Blumenstiel, J. P.; Chen, E. J.; Bany, I. A.; Mochizuki, N.; Ashbacher,A.; Matsuda, M.; Housman, D. E.; Graybiel, A. M.: A Rap guanine nucleotideexchange factor enriched highly in the basal ganglia. *Proc. Nat.Acad. Sci.* 95: 13278–13283, 1998.

[70258] 8588.Erle, D. J.; Sheppard, D.; Breuss, J.; Ruegg, C.; Pytela, R.:Novel integrin alpha and beta subunit cDNAs identified in airway epithelialcells and lung leukocytes using the polymerase chain reaction. *Am.J. Resp. Cell Molec. Biol.* 5: 170–177, 1991.

- [70259] 8589.Hibi, K.; Yamakawa, K.; Ueda, R.; Horio, Y.; Murata, Y.; Tamari,M.; Uchida, K.; Takahashi, T.; Nakamura, Y.; Takahashi, T.: Aberrantupregulation of a novel integrin alpha subunit gene at 3p21.3 in smallcell lung cancer. *Oncogene* 9: 611–619, 1994.
- [70260] 8590.Palmer, E. L.; Ruegg, C.; Ferrando, R.; Pytela, R.; Sheppard, D.: Sequence and tissue distribution of the integrin alpha-9 subunit,a novel partner of beta-1 that is widely distributed in epitheliaand muscle. *J. Cell Biol.* 123: 1289–1297, 1993.
- [70261] 8591.Yamakawa, K.; Takahashi, T.; Horio, Y.; Murata, Y.; Takahashi,E.; Hibi, K.; Yokoyama, S.; Ueda, R.; Takahashi, T.; Nakamura, Y.: Frequent homozygous deletions in lung cancer cell lines detectedby a DNA marker located at 3p21.3–p22. *Oncogene* 8: 327–330, 1993.
- [70262] 8592.La Starza, R.; Wlodarska, I.; Aventin, A.; Falzetti, D.; Crescenzi,B.; Martelli, M. F.; Van den Berghe, H.; Mecucci, C.: Molecular delineationof 13q deletion boundaries in 20 patients with myeloid malignancies. *Blood* 91:231–237, 1998.
- [70263] 8593.Favre, B.; Fontao, L.; Koster, J.; Shafaatian, R.; Jaunin, F.;Saurat, J.–H.; Sonnenberg, A.; Borradori, L.: The hemidesmosomalprotein bullous pemphigoid antigen 1

and the integrin beta-4 subunit bind to ERBIN: molecular cloning of multiple alternative splice variants of ERBIN and analysis of their tissue expression. *J. Biol. Chem.* 276:32427–32436, 2001.

- [70264] 8594. Agnello, V.; Abel, G.; Elfahal, M.; Knight, G. B.; Zhang, Q.-X.: Hepatitis C virus and other flaviviridae viruses enter cells via low density lipoprotein receptor. *Proc. Nat. Acad. Sci.* 96: 12766–12771, 1999.
- [70265] 8595. Allen, J. M.; Thompson, G. R.; Myant, N. B.; Steiner, R.; Oakley, C. M.: Cardiovascular complications of homozygous familial hypercholesterolaemia. *Brit. Heart J.* 44: 361–368, 1980.
- [70266] 8596. Benlian, P.; Amselem, S.; Loux, N.; Pastier, D.; Giraud, G.; deGennes, J. L.; Turpin, G.; Monnier, L.; Rieu, D.; Douste-Blazy, P.; Dastugue, B.; Goossens, M.; Junien, C.: A LDL receptor gene homozygous mutation: PCR amplification, direct genomic sequencing, associated haplotype, rapid screening for frequency. *Ann. Genet.* 33: 65–69, 1990.
- [70267] 8597. Bertolini, S.; Lelli, N.; Coviello, D. A.; Ghisellini, M.; Masturzo, P.; Tiozzo, R.; Elicio, N.; Gaddi, A.; Calandra, S.: A large deletion in the LDL receptor gene—the cause of familial hypercholesterolemia in three Italian families: a

study that dates back to the 17th century(FH-Pavia). Am. J. Hum. Genet. 51: 123–134, 1992.

[70268] 8598.Betard, C.; Kessling, A. M.; Roy, M.; Chamberland, A.; Lussier-Cacan,S.; Davignon, J.: Molecular genetic evidence for a founder effectin familial hypercholesterolemia among French Canadians. Hum. Genet. 88:529–536, 1992.

[70269] 8599.Boehnke, M.; Arnheim, N.; Li, H.; Collins, F. S.: Fine-structuregenetic mapping of human chromosomes using the polymerase chain reactionon single sperm: experimental design considerations. Am. J. Hum.Genet. 45: 21–32, 1989.

[70270] 8600.Brown, M. S.; Goldstein, J. L.: Receptor-mediated endocytosis:insights from the lipoprotein receptor system. Proc. Nat. Acad. Sci. 76:3330–3337, 1979.

[70271] 8601.Defesche, J. C.; van Diermen, D. E.; Lansberg, P. J.; Lamping,R. J.; Reymer, P. W. A.; Hayden, M. R.; Kastelein, J. J. P.: SouthAfrican founder mutations in the low-density lipoprotein receptorgene causing familial hypercholesterolemia in the Dutch population. Hum.Genet. 92: 567–570, 1993.

[70272] 8602.Defesche, J. C.; van de Ree, M. A.; Kastelein, J. J. P.; van Diermen,D. E.; Janssens, N. W. E.; van Doormaal, J. J.;

Hayden, M. R.: Detection of the pro664-to-leu mutation in the low-density lipoprotein receptor and its relation to lipoprotein(a) levels in patients with familial hypercholesterolemia of Dutch ancestry from The Netherlands and Canada. Clin.Genet. 42: 273–280, 1992.

[70273] 8603.Chen, H. I.; Einbond, A.; Kwak, S.-J.; Linn, H.; Koepf, E.; Peterson, S.; Kelly, J. W.; Sudol, M.: Characterization of the WW domain of human Yes-associated protein and its polyproline containing ligands. J.Biol. Chem. 272: 17070–17077, 1997.

[70274] 8604.Chen, H. I.; Sudol, M.: The WW domain of Yes-associated protein binds a proline-rich ligand that differs from the consensus established for Src homology 3-binding modules. Proc. Nat. Acad. Sci. 92: 7819–7823, 1995.

[70275] 8605.Hoggard, N.; Brintell, B.; Howell, A.; Weissenbach, J.; Varley, J.: Allelic imbalance on chromosome 1 in human breast cancer. II. Microsatellite repeat analysis. Genes Chromosomes Cancer 12: 24–31, 1995.

[70276] 8606.White, G. R. M.; Varley, J. M.; Heighway, J.: Isolation and characterization of a human homologue of the latrophilin gene from a region of 1p31.1 implicated in breast cancer. Oncogene 17: 3513–3519, 1998.

- [70277] 8607.Tommasi, S.; Dammann, R.; Jin, S.-G.; Zhang, X.; Avruch, J.; Pfeifer, G. P.: RASSF3 and NORE1: identification and cloning of two human homologues of the putative tumor suppressor gene RASSF1. *Oncogene* 21:2713–2720, 2002.
- [70278] 8608.Yao, R.; Wang, Y.; You, M.: Chromosome mapping and sequence variation of the murine Ras effector gene Nore1. *Cytogenet. Cell Genet.* 95:126–128, 2001.
- [70279] 8609.Heyer, B. S.; Warsowe, J.; Solter, D.; Knowles, B. B.; Ackerman, S. L.: New member of the Snf1/AMPK kinase family, Melk, is expressed in the mouse egg and preimplantation embryo. *Molec. Reprod. Dev.* 47:148–156, 1997.
- [70280] 8610.Merrill, R. A.; Plum, L. A.; Kaiser, M. E.; Clagett-Dame, M.: A mammalian homolog of unc-53 is regulated by all-trans retinoic acid in neuroblastoma cells and embryos. *Proc. Nat. Acad. Sci.* 99: 3422–3427, 2002.
- [70281] 8611.van Hille, B.; Richener, H.; Evans, D. B.; Green, J. R.; Bilbe, G.: Identification of two subunit A isoforms of the vacuolar H(+)-ATPase in human osteoclastoma. *J. Biol. Chem.* 268: 7075–7080, 1993.
- [70282] 8612.van Hille, B.; Richener, H.; Green, J. R.; Bilbe, G.: The ubiquitous VA68 isoform of subunit A of the vacuolar

H(+)-ATPase is highly expressed in human osteoclasts.

Biochem. Biophys. Res. Commun. 214: 1108–1113, 1995.

[70283] 8613. Boyhan, A.; Casimir, C. M.; French, J. K.; Teahan, C. G.; Segal, A. W.: Molecular cloning and characterization of grancalcin, a novel EF-hand calcium-binding protein abundant in neutrophils and monocytes. J. Biol. Chem. 267: 2928–2933, 1992.

[70284] 8614. Teahan, C. G.; Totty, N. F.; Segal, A. W.: Isolation and characterization of grancalcin, a novel 28 kDa EF-hand calcium-binding protein from human neutrophils. Biochem. J. 286: 549–554, 1992.

[70285] 8615. Denning, G.; Jamieson, L.; Maquat, L. E.; Thompson, E. A.; Fields, A. P.: Cloning of a novel phosphatidylinositol kinase-related kinase: characterization of the human SMG-1 RNA surveillance protein. J. Biol. Chem. 276: 22709–22714, 2001.

[70286] 8616. Diaz-Meco, M. T.; Municio, M. M.; Sanchez, P.; Lozano, J.; Moscat, J.: Lambda-interacting protein, a novel protein that specifically interacts with the zinc finger domain of the atypical protein kinase C isotype lambda/iota and stimulates its kinase activity in vitro and in vivo. Molec. Cell. Biol. 16: 105–114, 1996.

[70287] 8617. Kondo, M.; Ji, L.; Kamibayashi, C.; Tomizawa, Y.;

Randle, D.; Sekido, Y.; Yokota, J.; Kashuba, V.; Zabarovsky, E.; Kuzmin, I.; Lerman, M.; Roth, J.; Minna, J. D.: Overexpression of candidate tumor suppressor gene FUS1 isolated from the 3p21.3 homozygous deletion region leads to G1 arrest and growth inhibition of lung cancer cells. *Oncogene* 20:6258–6262, 2001.

[70288] 8618. Deshpande, K. L.; Seubert, P. H.; Tillman, D. M.; Farkas, W. R.; Katze, J. R.: Cloning and characterization of cDNA encoding the rabbit RNA–guanine transglycosylase 60–kilodalton subunit. *Arch. Biochem. Biophys.* 326: 1–7, 1996.

[70289] 8619. Wilson, S. M.; Bhattacharyya, B.; Rachel, R. A.; Coppola, V.; Tessarollo, L.; Householder, D. B.; Fletcher, C. F.; Miller, R. J.; Copeland, N. G.; Jenkins, N. A.: Synaptic defects in ataxia mice result from a mutation in *Usp14*, encoding a ubiquitin–specific protease. *Nature Genet.* 7 Oct, 2002. Note: Advance Electronic Publication.

[70290] 8620. Cunningham, S. A.; Arrate, M. P.; Rodriguez, J. M.; Bjercke, R. J.; Vanderslice, P.; Morris, A. P.; Brock, T. A.: A novel protein with homology to the junctional adhesion molecule: characterization of leukocyte interactions. *J. Biol. Chem.* 275: 34750–34756, 2000.

[70291] 8621. Liang, T. W.; Chiu, H. H.; Gurney, A.; Sidle, A.; Tu–



mas, D. B.;Schow, P.; Foster, J.; Klassen, T.; Dennis, K.; DeMarco, R. A.; Pham,T.; Frantz, G.; Fong, S.: Vascular endothelial-junctional adhesionmolecule (VE-JAM)/JAM 2 interacts with T, NK, and dendritic cellsthrough JAM 3. J. Immun. 168: 1618–1626, 2002.

- [70292] 8622.Palmeri, D.; van Zante, A.; Huang, C.-C.; Hemmerich, S.; Rosen,S. D.: Vascular endothelial junction-associated molecule, a novelmember of the immunoglobulin superfamily, is localized to intercellularboundaries of endothelial cells. J. Biol. Chem. 275: 19139–19145,2000.
- [70293] 8623.Arrate, M. P.; Rodriguez, J. M.; Tran, T. T.; Brock, T. A.; Cunningham,S. A.: Cloning of human junctional adhesion molecule 3 (JAM3) andits identification as the JAM2 counter-receptor. J. Biol. Chem. 276:45826–45832, 2001.
- [70294] 8624.Heng, Y. M.; Fox, M.; Sablitzky, F.: Assignment of the murinedef-3 gene (Rbm6) to chromosome 9F1–F2 and its pseudogenes Rbm6-ps1and Rbm6-ps2 to chromosome 1 by in situ hybridisation. Cytogenet.Cell. Genet. 89: 238–239, 2000.
- [70295] 8625.Garrett, R. M.; Bellissimo, D. B.; Rajagopalan, K. V.: Molecularcloning of human liver sulfite oxidase. Biochim. Biophys. Acta 1262:147–149, 1995.
- [70296] 8626.Johnson, J. L.; Coyne, K. E.; Garrett, R. M.; Zabet,

M.-T.; Dorche,C.; Kisker, C.; Rajagopalan, K. V.: Isolated sulfite oxidase deficiency:identification of 12 novel SUOX mutations in 10 patients.(Abstract) Hum.Mutat. 20: 74 only, 2002.

- [70297] 8627.Chen, Y. Q.; Rafi, M. A.; de Gala, G.; Wenger, D. A.: Cloningand expression of cDNA encoding human galacto-cerebrosidase, the enzymedeficient in globoid cell leukodystrophy. Hum. Molec. Genet. 2:1841-1845, 1993.
- [70298] 8628.Chen, Y. Q.; Wenger, D. A.: Galactocerebrosidase from human urine:purification and partial characteriza-tion. Biochim. Biophys. Acta 1170:53-61, 1993.
- [70299] 8629.Fiumara, A.; Pavone, L.; Siciliano, L.; Tine, A.; Parano, E.; Innico,G.: Late-onset globoid cell leukodystro-phy: report on 7 new patients. Child'sNerv. Syst. 6: 194-197, 1990.
- [70300] 8630.Furuya, H.; Kukita, Y.; Nagano, S.; Sakai, Y.; Ya-mashita, Y.; Fukuyama,H.; Inatomi, Y.; Saito, Y.; Koike, R.; Tsuji, S.; Fukumaki, Y.; Hayashi,K.; Kobayashi, T.: Adult onset globoid cell leukodystrophy (Krabbedisease): analy-sis of galactosylceramidase cDNA from four Japanesepa-tients. Hum. Genet. 100: 450-456, 1997.
- [70301] 8631.Kodama, S.; Igisu, H.; Siegel, D. A.; Suzuki, K.: Gly-cosylceramidesynthesis in the developing spinal cord and

kidney of the twitchermouse, an enzymatically authentic model of human Krabbe disease. *J.Neurochem.* 39: 1314–1318, 1982.

[70302] 8632.Luzi, P.; Rafi, M. A.; Victoria, T.; Baskin, G. B.; Wenger, D.A.: Characterization of the rhesus monkey galactocerebrosidase (GALC)cDNA and gene and identification of the mutation causing globoid cellleukodystrophy (Krabbe disease) in this primate. *Genomics* 42: 319–324,1997.

[70303] 8633.Luzi, P.; Rafi, M. A.; Wenger, D. A.: Structure and organizationof the human galactocerebrosidase (GALC) gene. *Genomics* 26: 407–409,1995.

[70304] 8634.Wang, K.; Zhou, B.; Kuo, Y.–M.; Zemansky, J.; Gitschier, J.: Anovel member of a zinc transporter family is defective in acrodermatitisenteropathica. *Am. J. Hum. Genet.* 71: 66–73, 2002.

[70305] 8635.Freeman, B. C.; Yamamoto, K. R.: Disassembly of transcriptionalregulatory complexes by molecular chaperones. *Science* 296: 2232–2235,2002.

[70306] 8636.Johnson, J. L.; Beito, T. G.; Krco, C. J.; Toft, D. O.: Characterizationof a novel 23–kilodalton protein of unactive progesterone receptorcomplexes. *Molec. Cell. Biol.* 14: 1956–1963, 1994.

- [70307] 8637.Smith, D. F.; Faber, L. E.; Toft, D. O.: Purification of unactivated progesterone receptor and identification of novel receptor-associated proteins. *J. Biol. Chem.* 265: 3996–4003, 1990.
- [70308] 8638.Patterson, C. E.; Gao, J.; Rooney, A. P.; Davis, E. C.: Genomic organization of mouse and human 65 kDa FK506-binding protein genes and evolution of the FKBP multigene family. *Genomics* 79: 881–889, 2002.
- [70309] 8639.Patterson, C. E.; Schaub, T.; Coleman, E. J.; Davis, E. C.: Developmental regulation of FKBP65: an ER localized extracellular matrix-binding protein. *Molec. Biol. Cell* 11: 3925–3935, 2000.
- [70310] 8640.Cai, H.; Wang, Y.; McCarthy, D.; Wen, H.; Borchelt, D. R.; Price, D. L.; Wong, P. C.: BACE1 is the major beta-secretase for generation of A-beta peptides by neurons. *Nature Neurosci.* 4: 233–234, 2001.
- [70311] 8641.Meyaard, L.; Adema, G. J.; Chang, C.; Woollatt, E.; Sutherland, G. R.; Lanier, L. L.; Phillips, J. H.: LAIR-1, a novel inhibitory receptor expressed on human mononuclear leukocytes. *Immunity* 7:283–290, 1997.
- [70312] 8642.Sathish, J. G.; Johnson, K. G.; Fuller, K. J.; LeRoy, F. G.; Meyaard, L.; Sims, M. J.; Matthews, R. J.: Constitutive association of SHP-1 with leukocyte-associated Ig-like re-

ceptor-1 in human T cells. J.Immun. 166: 1763-1770, 2001.

[70313] 8643.Xu, M.; Zhao, R.; Zhao, Z. J.: Identification and characterization of leukocyte-associated Ig-like receptor-1 as a major anchor protein of tyrosine phosphatase SHP-1 in hematopoietic cells. J. Biol. Chem. 275:17440-17446, 2000.

[70314] 8644.Hofmann, R. M.; Pickart, C. M.: Noncanonical MMS2-encoded ubiquitin-conjugating enzyme functions in assembly of novel polyubiquitin chains for DNA repair. Cell 96: 645-653, 1999.

[70315] 8645.Rothofsky, M. L.; Lin, S. L.: CROC-1 encodes a protein which mediates transcriptional activation of the human FOS promoter. Gene 195:141-149, 1997.

[70316] 8646.Sancho, E.; Vila, M. R.; Sanchez-Pulido, L.; Lozano, J. J.; Paciucci, R.; Nadal, M.; Fox, M.; Harvey, C.; Bercovich, B.; Loukili, N.; Ciechanover, A.; Lin, S. L.; Sanz, F.; Estivill, X.; Valencia, A.; Thomson, T.M.: Role of UEV-1, an inactive variant of the E2 ubiquitin-conjugating enzymes, in in vitro differentiation and cell cycle behavior of HT-29-M6 intestinal mucosecretory cells. Molec. Cell. Biol. 18: 576-589, 1998.

[70317] 8647.Charles, C. H.; Yoon, J. K.; Simske, J. S.; Lau, L. F.:

Genomic structure, cDNA sequence, and expression of gly96, a growth factor-inducible immediate-early gene encoding a short-lived glycosylated protein. *Oncogene* 8:797–801, 1993.

[70318] 8648. Kondratyev, A. D.; Chung, K.-N.; Jung, M. O.: Identification and characterization of a radiation-inducible glycosylated human early-response gene. *Cancer Res.* 56: 1498–1502, 1996.

[70319] 8649. Pietzsch, A.; Buchler, C.; Aslanidis, C.; Schmitz, G.: Identification and characterization of a novel monocyte/macrophage differentiation-dependent gene that is responsive to lipopolysaccharide, ceramide, and lysophosphatidylcholine. *Biochem. Biophys. Res. Commun.* 235: 4–9, 1997.

[70320] 8650. Pietzsch, A.; Buchler, C.; Schmitz, G.: Genomic organization, promoter cloning, and chromosomal localization of the Dif-2 gene. *Biochem. Biophys. Res. Commun.* 245: 651–657, 1998.

[70321] 8651. Schafer, H.; Trauzold, A.; Siegel, E. G.; Folsch, U. R.; Schmidt, W. E.: PRG1: a novel early-response gene transcriptionally induced by pituitary adenylate cyclase activating polypeptide in a pancreatic carcinoma cell line. *Cancer Res.* 56: 2641, 1996.

- [70322] 8652.Wu, M. X.; Ao, Z.; Prasad, K. V. S.; Wu, R.; Schlossman, S. F.: IEX-1L, an apoptosis inhibitor involved in NF-kappa-B-mediated cell survival. *Science* 281: 998-1001, 1998.
- [70323] 8653.De Strooper, B.; Konig, G.: A firm base for drug development. *Nature* 402:471-472, 1999.
- [70324] 8654.Fan, W.; Bennett, B. D.; Babu-Khan, S.; Luo, Y.; Louis, J.-C.; McCaleb, M.; Citron, M.; Vassar, R.; Richards, W. G.: Response to Saunders et al. (1999). *Science* 286: 1255a, 1999. Note: Electronic Publication.
- [70325] 8655.Haniu, M.; Denis, P.; Young, Y.; Mendiaz, E. A.; Fuller, J.; Hui, J. O.; Bennett, B. D.; Kahn, S.; Ross, S.; Burgess, T.; Katta, V.; Rogers, G.; Vassar, R.; Citron, M.: Characterization of Alzheimer's beta-secretase protein BACE: a pepsin family member with unusual properties. *J. Biol. Chem.* 275: 21099-21106, 2000.
- [70326] 8656.Hong, L.; Koelsch, G.; Lin, X.; Wu, S.; Terzyan, S.; Ghosh, A.K.; Zhang, X. C.; Tang, J.: Structure of the protease domain of memapsin2 (beta-secretase) complexed with inhibitor. *Science* 290: 150-153, 2000.
- [70327] 8657.Hussain, I.; Powell, D.; Howlett, D. R.; Tew, D. G.; Meek, T. D.; Chapman, C.; Gloger, I. S.; Murphy, K. E.; Southan, C. D.; Ryan, D.M.; Smith, T. S.; Simmons, D. L.;

Walsh, F. S.; Dingwall, C.; Christie, G.: Identification of a novel aspartic protease (asp2) as beta-secretase  
Molec. Cell Neurosci. 14: 419–427, 1999.

[70328] 8658. Parker, A. E.; Van de Weyer, I.; Laus, M. C.; Verhaselt, P.; Luyten, W. H. M. L.: Identification of a human homologue of the *Schizosaccharomyces pombe* rad17+ checkpoint gene. J. Biol. Chem. 273: 18340–18346, 1998. Note: Erratum: J. Biol. Chem. 274: 24438–24439, 1999.

[70329] 8659. von Deimling, F.; Scharf, J. M.; Liehr, T.; Rothe, M.; Kelter, A.-R.; Albers, P.; Dietrich, W. F.; Kunkel, L. M.; Wernert, N.; Wirth, B.: Human and mouse RAD17 genes: identification, localization, genomic structure and histological expression pattern in normal testis and seminoma. Hum. Genet. 105: 17–27, 1999.



- [70330] 8660.Serra-Pages, C.; Medley, Q. G.; Tang, M.; Hart, A.; Streuli, M.: Liprins, a family of LAR transmembrane protein-tyrosine phosphatase-interacting proteins. J. Biol. Chem. 273: 15611-15620, 1998.
- [70331] 8661.Thomas, M. K.; Yao, K.-M.; Tenser, M. S.; Wong, G. G.; Habener, J. F.: Bridge-1, a novel PDZ-domain coactivator of E2A-mediated regulation of insulin gene transcription. Molec. Cell. Biol. 19: 8492-8504, 1999.
- [70332] 8662.Watanabe, T. K.; Saito, A.; Suzuki, M.; Fujiwara, T.; Takahashi, E.; Slaughter, C. A.; DeMartino, G. N.; Hendil, K. B.; Chung, C. H.; Tanahashi, N.; Tanaka, K.: cDNA cloning and characterization of a human proteasomal modulator subunit, p27 (PSMD9). Genomics 50: 241-250, 1998.
- [70333] 8663.Kennedy, J.; Rossi, D. L.; Zurawski, S. M.; Vega, F., Jr.; Kastelein, R. A.; Wagner, J. L.; Hannum, C. H.; Zlotnick, A.: Mouse IL-17: a cytokine preferentially expressed by alpha beta TCR + CD4-CD8-T cells. J. Interferon Cytokine Res. 16: 611-617, 1996.
- [70334] 8664.Kotake, S.; Udagawa, N.; Takahashi, N.; Matsuzaki, K.; Itoh, K.; Ishiyama, S.; Saito, S.; Inoue, K.; Kamatani, N.; Gillespie, M. T.; Martin, T. J.; Suda, T.: IL-17 in synovial fluids from patients with rheumatoid arthritis is a potent stimulator of osteoclastogenesis. J. Clin. Invest. 103:

1345–1352, 1999.

[70335] 8665.Rouvier, E.; Luciani, M.–F.; Mattei, M.–G.; Denizot, F.; Golstein, P.: CTLA–8, cloned from an activated T cell, bearing AU–rich messengerRNA instability sequences, and homologous to a Herpesvirus saimirigene. J. Immun. 150: 5445–5456, 1993.

[70336] 8666.Yao, Z.; Painter, S. L.; Fanslow, W. C.; Ulrich, D.; Macduff, B.M.; Spriggs, M. K.; Armitage, R. J.: Human IL–17: a novel cytokinederived from T cells. J. Immun. 155: 5483–5486, 1995.

[70337] 8667.Higuti, T.; Tsurumi, C.; Kawamura, Y.; Tsujita, H.; Osaka, F.;Yoshihara, Y.; Tani, I.; Tanaka, K.; Ichihara, A.: Molecular cloningof cDNA for the import precursor of human coupling factor 6 of H(+)-ATPsynthase in mitochondria. Biochem. Biophys. Res. Commun. 178: 793–799,1991.

[70338] 8668.Javed, A. A.; Ogata, K.; Sanadi, D. R.: Human mitochondrial ATPsynthase: cloning cDNA for the nuclear–encoded precursor of couplingfactor 6. Gene 97: 307–310, 1991.

[70339] 8669.Webster, K. A.; Oliver, N. A.; Wallace, D. C.: Assignment of anoligomycin–resistance locus to human chromosome 10. Somat. Cell Genet. 8:223–244, 1982.

- [70340] 8670.Parker, A. E.; Van de Weyer, I.; Laus, M. C.; Oostveen, I.; Yon,J.; Verhasselt, P.; Luyten, W. H. M. L.: A human homologue of Schizosaccharomycespombe rad1+ checkpoint gene encodes an exonuclease. J. Biol. Chem. 273:18332–18339, 1998.
- [70341] 8671.Barbon, A.; Ferraboli, S.; Barlati, S.: Assignment of the humanmetabotropic glutamate receptor gene GRM4 to chromosome 6 band p21.3by radiation hybrid mapping. Cytogenet. Cell Genet. 88: 210 only,2000.
- [70342] 8672.Morita, R.; Miyazaki, E.; Fong, C. G.; Chen, X.–N.; Korenberg,J. R.; Delgado–Escueta, A. V.; Yamakawa, K.: JH8, a gene highly homologousto the mouse jerky gene, maps to the region for childhood absenceepilepsy on 8q24. Biochem. Biophys. Res. Commun. 248: 307–314, 1998.
- [70343] 8673.Toth, M.; Grimsby, J.; Buzsaki, G.; Donovan, G. P.: Epilepticseizures caused by inactivation of a novel gene, jerky, related tocentromere binding protein–B in transgenic mice. Nature Genet. 11:71–75, 1995. Note: Erratum: Nature Genet. 12: 110 only, 1996.
- [70344] 8674.Zeng, Z.; Kyaw, H.; Gakenheimer, K. R.; Augustus, M.; Fan, P.;Zhang, X.; Su, K.; Carter, K. C.; Li, Y.: Cloning, mapping, and tissuedistribution of a human homologue of

the mouse jerky gene product. *Biochem. Biophys. Res. Commun.* 236: 389–395, 1997.

- [70345] 8675. Kimura, M.; Matsuda, Y.; Eki, T.; Yoshioka, T.; Okumura, K.; Hanaoka, F.; Okano, Y.: Assignment of STK6 to human chromosome 20q13.2–q13.3 and a pseudogene STK6P to 1q41–q42. *Cytogenet. Cell Genet.* 79: 201–203, 1997.
- [70346] 8676. Bryan, J.; Edwards, R.; Matsudaira, P.; Otto, J.; Wulfschlegel, J.: Fascin, an echinoid actin-bundling protein, is a homolog of the *Drosophila* singed gene product. *Proc. Nat. Acad. Sci.* 90: 9115–9119, 1993.
- [70347] 8677. Dressel, U.; Thormeyer, D.; Altincicek, B.; Paululat, A.; Eggert, M.; Schneider, S.; Tenbaum, S. P.; Renkawitz, R.; Banihmad, A.: Alien, a highly conserved protein with characteristics of a corepressor for members of the nuclear hormone receptor superfamily. *Molec. Cell Biol.* 19: 3383–3394, 1999.
- [70348] 8678. Schaefer, L.; Beermann, M. L.; Miller, J. B.: Coding sequence, genomic organization, chromosomal localization, and expression pattern of the signalosome component Cops2: the mouse homologue of *Drosophila* alien. *Genomics* 56: 310–316, 1999.
- [70349] 8679. Scott, A. F.: Personal Communication. Baltimore, Md.

11/8/2000.

- [70350] 8680. Seeger, M.; Kraft, R.; Ferrell, K.; Dawadschargal, B.-O.; Dumdey, R.; Schade, R.; Gordon, C.; Naumann, M.; Dubiel, W.: A novel protein complex involved in signal transduction possessing similarities to 26S proteasome subunits. *FASEB J.* 12: 469–478, 1998.
- [70351] 8681. Barbosa, M. D.; Johnson, S. A.; Achey, K.; Gutierrez, M. J.; Wakeland, E. K.; Zerial, M.; Kingsmore, S. F.: The Rab protein family: genetic mapping of six Rab genes in the mouse. *Genomics* 30: 439–444, 1995.
- [70352] 8682. Vitelli, R.; Chiariello, M.; Lattero, D.; Bruni, C. B.; Bucci, C.: Molecular cloning and expression analysis of the human Rab7 GTP-ase complementary deoxyribonucleic acid. *Biochem. Biophys. Res. Commun.* 229: 887–890, 1996.
- [70353] 8683. Wang, L.; Mizzen, C.; Ying, C.; Candau, R.; Barlev, N.; Brownell, J.; Allis, C. D.; Berger, S. L.: Histone acetyltransferase activity is conserved between yeast and human GCN5 and is required for complementation of growth and transcriptional activation. *Molec. Cell. Biol.* 17: 519–527, 1997.
- [70354] 8684. Xu, W.; Edmondson, D. G.; Evrard, Y. A.; Wakamiya, M.; Behringer, R. R.; Roth, S. Y.: Loss of Gcn5l2 leads to in-

creased apoptosis and mesodermal defects during mouse development. *Nature Genet.* 26: 229–232, 2000.

- [70355] 8685. Xu, W.; Edmondson, D. G.; Roth, S. Y.: Mammalian GCN5 and P/CAF acetyltransferases have homologous amino-terminal domains important for recognition of nucleosomal substrates. *Molec. Cell. Biol.* 18:5659–5669, 1998.
- [70356] 8686. Yang, X.-J.; Ogryzko, V. V.; Nishikawa, J.; Howard, B. H.; Nakatani, Y.: A p300/CBP-associated factor that competes with the adenoviral oncoprotein E1A. *Nature* 382: 319–324, 1996.
- [70357] 8687. Miao, D.; He, B.; Karaplis, A. C.; Goltzman, D.: Parathyroid hormone is essential for normal fetal bone formation. *J. Clin. Invest.* 109:1173–1182, 2002.
- [70358] 8688. Betticher, D. C.; Thatcher, N.; Altermatt, H. J.; Hoban, P.; Ryder, W. D. J.; Heighway, J.: Alternate splicing produces a novel cyclin D1 transcript. *Oncogene* 11: 1005–1011, 1995.
- [70359] 8689. Chesi, M.; Bergsagel, P. L.; Brents, L. A.; Smith, C. M.; Gerhard, D. S.; Kuehl, W. M.: Dysregulation of cyclin D1 by translocation into an IgH gamma switch region in two multiple myeloma cell lines. *Blood* 88:674–681, 1996.
- [70360] 8690. Gailani, M. R.; Petty, E. M.; Horsthemke, B.; Arnold,

A.; Marx, S. J.; Bale, A. E.: Physical mapping of chromosome 11q12–13 by pulsed-field gel electrophoresis (PFGE). (Abstract) *Cytogenet. Cell Genet.* 58:1959, 1991.

- [70361] 8691. Geng, Y.; Yu, Q.; Sicinska, E.; Das, M.; Bronson, R. T.; Sicinski, P.: Deletion of the p27(Kip1) gene restores normal development in cyclin D1-deficient mice. *Proc. Nat. Acad. Sci.* 98: 194–199, 2001.
- [70362] 8692. Hayette, S.; Gadoux, M.; Martel, S.; Bertrand, S.; Tigaud, I.; Magaud, J.-P.; Rimokh, R.: FLRG (follistatin-related gene), a new target of chromosomal rearrangement in malignant blood disorders. *Oncogene* 16:2949–2954, 1998.
- [70363] 8693. Tajima, T.; Kitagawa, H.; Yokoya, S.; Tachibana, K.; Adachi, M.; Nakae, J.; Suwa, S.; Katoh, S.; Fujieda, K.: A novel missense mutation of mineralocorticoid receptor gene in one Japanese family with a renal form of pseudo-hypoaldosteronism type 1. *J. Clin. Endocr. Metab.* 85:4690–4694, 2000.
- [70364] 8694. Viemann, M.; Peter, M.; Lopez-Siguero, J. P.; Simic-Schleicher, G.; Sippell, W. G.: Evidence for genetic heterogeneity of pseudohypoaldosteronism type 1: identification of a novel mutation in the human mineralocorticoid receptor in one sporadic case and no mutations in two autosomal

mal dominant kindreds. *J. Clin. Endocr. Metab.* 86: 2056–2059, 2001.

- [70365] 8695. Lee, Y. L.; Helman, L.; Hoffman, T.; Laborda, J.: dlk, pG2 and Pref-1 mRNAs encode similar proteins belonging to the EGF-like superfamily. Identification of polymorphic variants of this RNA. *Biochim. Biophys. Acta* 1261: 223–232, 1995.
- [70366] 8696. Smas, C. M.; Sul, H. S.: Pref-1, a protein containing EGF-like repeats, inhibits adipocyte differentiation. *Cell* 73: 725–734, 1993.
- [70367] 8697. Takada, S.; Paulsen, M.; Tevendale, M.; Tsai, C.-E.; Kelsey, G.; Cattanach, B. M.; Ferguson-Smith, A. C.: Epigenetic analysis of the Dlk1-Gtl2 imprinted domain on mouse chromosome 12: implications for imprinting control from comparison with Igf2-H19. *Hum. Molec. Genet.* 11:77–86, 2002.
- [70368] 8698. Wylie, A. A.; Murphy, S. K.; Orton, T. C.; Jirtle, R. L.: Novel imprinted DLK1/GTL2 domain on human chromosome 14 contains motifs that mimic those implicated in IGF2/H19 regulation. *Genome Res.* 10:1711–1718, 2000.
- [70369] 8699. Okumura, K.; Nogami, M.; Taguchi, H.; Hisamatsu, H.; Tanaka, K.: The genes for the alpha-type HC3 (PMSA2) and beta-type HC5 (PMSB1) subunits of human protea-



somes map to chromosomes 6q27 and 7p12–p13 by fluorescence in situ hybridization. *Genomics* 27: 377–379, 1995.

- [70370] 8700. Tamura, T.; Osaka, F.; Kawamura, Y.; Higuchi, T.; Ishida, N.; Nothwang, H. G.; Tsurumi, C.; Tanaka, K.; Ichihara, A. : *J. Molec. Biol.* 244:1117–1124, 1994.
- [70371] 8701. Coux, O.; Tanaka, K.; Goldberg, A. L.: Structure and function of the 20S and 26S proteasomes. *Ann. Rev. Biochem.* 65: 801–847, 1996.
- [70372] 8702. DeMartino, G. N.; Orth, K.; McCullough, M. L.; Lee, L. W.; Munn, T. Z.; Moomaw, C. R.; Dawson, P. A.; Slaughter, C. A.: The primary structures of four subunits of the human, high molecular weight proteinase, macropain (proteasome), are distinct but homologous. *Biochim. Biophys. Acta* 1079: 29–38, 1991.
- [70373] 8703. Cruz, M.; Elenich, L. A.; Smolarek, T. A.; Menon, A. G.; Monaco, J. J.: DNA sequence, chromosomal localization, and tissue expression of the mouse proteasome subunit Lmp10 (Psm10) gene. *Genomics* 45:618–622, 1997.
- [70374] 8704. Goldberg, A. L.; Rock, K. L.: Proteolysis, proteasomes and antigen presentation. *Nature* 357: 375–379, 1992.
- [70375] 8705. Larsen, F.; Gundersen, G.; Lopez, R.; Prydz, H.: CpG

islands as gene markers in the human genome. *Genomics* 13: 1095–1107, 1992.

[70376] 8706. Larsen, F.; Solheim, J.; Kristensen, T.; Kolsto, A.-B.; Prydz, H.: A tight cluster of five unrelated human genes on chromosome 16q22.1. *Hum. Molec. Genet.* 2: 1589–1595, 1993.

[70377] 8707. DeVry, C. G.; Clarke, S.: Assignment of the protein L-isoaspartate(D-aspartate) O-methyltransferase gene (PCMT1) to human chromosome bands 6q24–q25 with radiation hybrid mapping. *Cytogenet. Cell Genet.* 84:130–131, 1999.

[70378] 8708. Farrar, C.; Clarke, S.: Altered levels of S-adenosylmethionine and S-adenosylhomocysteine in the brains of L-isoaspartyl (D-aspartyl)O-methyltransferase-deficient mice. *J. Biol. Chem.* 277: 27856–27863, 2002.

[70379] 8709. Ingrosso, D.; Kagan, R. M.; Clarke, S.: Distinct C-terminal sequences of isozymes I and II of the human erythrocyte L-isoaspartyl/D-aspartyl protein methyltransferase. *Biochem. Biophys. Res. Commun.* 175: 351–358, 1991.

[70380] 8710. Kim, E.; Lowenson, J. D.; Clarke, S.; Young, S. G.: Phenotypic analysis of seizure-prone mice lacking L-

isoaspartate (D-aspartate)O-methyltransferase. J. Biol. Chem. 274: 20671–20678, 1999.

[70381] 8711.Lowenson, J. D.; Kim, E.; Young, S. G.; Clarke, S.: Limited accumulationof damaged proteins in L-isoaspartyl (D-aspartyl) O-methyltransferase-deficientmice. J. Biol. Chem. 276: 20695–20702, 2001.

[70382] 8712.MacLaren, D. C.; Kagan, R. M.; Clarke, S.: Alternative splicingof the human isoaspartyl protein carboxyl methyltransferase leadsto the generation of a C-terminal –RDEL sequence in isozyme II. Biochem.Biophys. Res. Commun. 185: 277–283, 1992.

[70383] 8713.MacLaren, D. C.; O'Connor, C. M.; Xia, Y.–R.; Mehra–bian, M.; Klisak,I.; Sparkes, R. S.; Clarke, S.; Lusi, A. J.: The L-isoaspartyl/D-aspartylprotein methyltransferase gene (PCMT1) maps to human chromosome 6q22.3–6q24and the syntenic region of mouse chromosome 10. Genomics 14: 852–856,1992.

[70384] 8714.Ota, I. M.; Gilbert, J. M.; Clarke, S.: Two major isozymes ofthe protein D-aspartyl/L-isoaspartyl methyltransferase from humanerythrocytes. Biochem. Biophys. Res. Commun. 151: 1136–1143, 1988.

[70385] 8715.Hunter, A. G. W.; Clifford, B.; Cox, D. M.: The characteristicphysiognomy and tissue specific karyotype dis–

tribution in the Pallister–Killiansyndrome. Clin. Genet. 28: 47–53, 1985.

[70386] 8716.Astrom, A.–K.; Voz, M. L.; Kas, K.; Roijer, E.; Wedell, B.; Mandahl,N.; Van de Ven, W.; Mark, J.; Stenman, G.: Conserved mechanism ofPLAG1 activation in salivary gland tumors with and without chromosome8q12 abnormalities: identification of SII as a new fusion partner gene. Cancer Res. 59: 918–923, 1999.

[70387] 8717.Bullerdiek, J.; Bartnitzke, S.; Weinberg, M.; Chilla, R.; Haubrich,J.; Schloot, W.: Rearrangements of chromosome region 12q13–q15 inpleomorphic adenomas of the human salivary gland (PSA). Cytogenet.Cell Genet. 45: 187–190, 1987.

[70388] 8718.Bullerdiek, J.; Raabe, G.; Bartnitzke, S.; Boschen, C.; Schloot,W.: Structural rearrangements of chromosome #8 involving 8q12--a primary event in pleomorphic adenomas of the parotid gland. Genetica 72:85–92, 1987.

[70389] 8719.Mark, J.; Dahlenfors, R.: Cytogenetical observations in 100 humanbenign pleomorphic adenomas: specificity of the chromosomal aberrationsand their relationship to sites of localized oncogenes. AnticancerRes. 6: 299–308, 1986.

[70390] 8720.Stenman, G.; Sandros, J.; Mark, J.; Nordkvist, A.:

High p21(RAS)expression levels correlate with chromosome 8 rearrangements in benign human mixed salivary gland tumors. *Genes Chromosomes Cancer* 1: 59–66, 1989.

[70391] 8721.Hinds, P. W.; Dowdy, S. F.; Eaton, E. N.; Arnold, A.; Weinberg, R. A.: Function of a human cyclin gene as an oncogene. *Proc. Nat. Acad. Sci.* 91: 709–713, 1994.

[70392] 8722.Komatsu, H.; Iida, S.; Yamamoto, K.; Mikuni, C.; Nitta, M.; Takahashi, T.; Ueda, R.; Seto, M.: A variant chromosome translocation at 11q13 identifying PRAD1/cyclin D1 as the BCL-1 gene. *Blood* 84: 1226–1231, 1994.

[70393] 8723.Kong, S.; Amos, C. I.; Luthra, R.; Lynch, P. M.; Levin, B.; Frazier, M. L.: Effects of cyclin D1 polymorphism on age of onset of hereditary nonpolyposis colorectal cancer. *Cancer Res.* 60: 249–252, 2000.

[70394] 8724.Kong, S.; Wei, Q.; Amos, C. I.; Lynch, P. M.; Zong, J.; Frazier, M. L.: Cyclin D1 polymorphism and increased risk of colorectal cancer at young age. *J. Nat. Cancer Inst.* 93: 1106–1108, 2001.

[70395] 8725.Ma, C.; Papermaster, D.; Cepko, C. L.: A unique pattern of photoreceptor degeneration in cyclin D1 mutant mice. *Proc. Nat. Acad. Sci.* 95: 9938–9943, 1998.

[70396] 8726.Motokura, T.; Bloom, T.; Kim, H. G.; Juppner, H.; Ru-

derman, J.V.; Kronenberg, H. M.; Arnold, A.: A novel cyclin encoded by a bcl1-linked candidate oncogene. *Nature* 350: 512–515, 1991.

[70397] 8727. Muller, H.; Lukas, J.; Schneider, A.; Warthoe, P.; Bartek, J.; Eilers, M.; Strauss, M.: Cyclin D1 expression is regulated by the retinoblastoma protein. *Proc. Nat. Acad. Sci.* 91: 2945–2949, 1994.

[70398] 8728. Richard, C. W., III; Withers, D. A.; Meeker, T. C.; Maurer, S.; Evans, G. A.; Myers, R. M.; Cox, D. R.: A radiation hybrid map of the proximal long arm of human chromosome 11 containing the multiple endocrine neoplasia type I (MEN-1) and bcl-1 disease loci. *Am. J. Hum. Genet.* 49: 1189–1196, 1991.

[70399] 8729. Rimokh, R.; Berger, F.; Bastard, C.; Klein, B.; French, M.; Archimbaud, E.; Rouault, J. P.; Santa Lucia, B.; Duret, L.; Vuillaume, M.; et al.: Rearrangement of CCND1 (BCL1/PRAD1) 3-prime untranslated region in mantle-cell lymphomas and t(11q13)-associated leukemias. *Blood* 83:3689–3696, 1994.

[70400] 8730. Rimokh, R.; Berger, F.; Delsol, G.; Charrin, C.; Bertheas, M.F.; French, M.; Garoscio, M.; Felman, P.; Coiffier, C.; Bryon, P.A.: Rearrangement and overexpression of the BCL-1/PRAD-1 gene in intermediate lymphocytic lym-

phomas and in t(11q13)-bearing leukemias. Blood 81:3063–3067, 1993.

- [70401] 8731.Rosenberg, C. L.; Kim, H. G.; Shows, T. B.; Kronenberg, H. M.;Arnold, A.: Rearrangement and overexpression of D11S287E, a candidate oncogene on chromosome 11q13 in benign parathyroid tumors. Oncogene 6:449–453, 1991.
- [70402] 8732.Sicinski, P.; Donaher, J. L.; Parker, S. B.; Li, T.; Fazeli, A.;Gardner, H.; Haslam, S. Z.; Bronson, R. T.; Elledge, S. J.; Weinberg,R. A.: Cyclin D1 provides a link between development and oncogenesis in the retina and breast. Cell 82: 621–630, 1995.
- [70403] 8733.Szepetowski, P.; Perucca–Lostanlen, D.; Gaudray, P.: Mapping genes according to their amplification status in tumor cells: contribution to the map of 11q13. Genomics 16: 745–750, 1993.
- [70404] 8734.Clark, T. G.; Conway, S. J.; Scott, I. C.; Labosky, P. A.; Winnier,G.; Bundy, J.; Hogan, B. L. M.; Greenspan, D. S.: The mammalian Tollid–like1 gene, Tll1, is necessary for normal septation and positioning of the heart. Development 126: 126:–2631–2642, 1999.
- [70405] 8735.Scott, I. C.; Clark, T. G.; Takahara, K.; Hoffman, G. G.; Eddy,R. L.; Haley, L. L.; Shows, T. B.; Greenspan, D. S.:

Assignment of TLL1 and TLL2, which encode human BMP-1/tolloid-related metalloproteases, to chromosomes 4q32-q33 and 10q23-q24 and assignment of murine Tll2 to chromosome 19. *Cytogenet. Cell Genet.* 86: 64-65, 1999.

[70406] 8736. Takahara, K.; Brevard, R.; Hoffman, G. G.; Suzuki, N.; Greenspan, D. S.: Characterization of a novel gene product (mammalian tolloid-like) with high sequence similarity to mammalian tolloid/bone morphogenetic protein-1. *Genomics* 34: 157-165, 1996.

[70407] 8737. Bauer, W. O.; Nanda, I.; Beck, G.; Schmid, M.; Jakob, F.: Human puromycin-sensitive aminopeptidase: cloning of 3-prime UTR, evidence for a polymorphism at aa 140 and refined chromosomal localization to 17q21. *Cytogenet. Cell Genet.* 92: 221-224, 2001.

[70408] 8738. Huber, G.; Thompson, A.; Gruninger, F.; Mechler, H.; Hochstrasser, R.; Hauri, H.-P.; Malherbe, P.: cDNA cloning and molecular characterization of human brain metalloprotease MP100: a beta-secretase candidate? *J. Neurochem.* 72: 1215-1223, 1999.

[70409] 8739. Osada, T.; Sakaki, Y.; Takeuchi, T.: Puromycin-sensitive aminopeptidase gene (Psa) maps to mouse chromosome 11. *Genomics* 56: 361-362, 1999.



- [70410] 8740.Schonlein, C.; Loffler, J.; Huber, G.: Purification and characterization of a novel metalloprotease from human brain with the ability to cleave substrates derived from the N-terminus of beta-amyloid protein. *Biochem.Biophys. Res. Commun.* 201: 45-53, 1994.
- [70411] 8741.Thompson, M. W.; Tobler, A.; Fontana, A.; Hersh, L. B.: Cloning and analysis of the gene for the human puromycin-sensitive aminopeptidase. *Biochem.Biophys. Res. Commun.* 258: 234-240, 1999.
- [70412] 8742.Tobler, A. R.; Constam, D. B.; Schmitt-Graff, A.; Malipiero, U.; Schlapbach, R.; Fontana, A.: Cloning of the human puromycin-sensitive aminopeptidase and evidence for expression in neurons. *J. Neurochem.* 68:889-897, 1997.
- [70413] 8743.Zhang, Y.; Cai, X.; Schlegelberger, B.; Zheng, S.: Assignment of human putative tumor suppressor genes ST13 (alias SNC6) and ST14 (alias SNC19) to human chromosome bands 22q13 and 11q24-q25 by in situ hybridization. *Cytogenet. Cell Genet.* 83: 56-57, 1998.
- [70414] 8744.Friedrich, R.; Fuentes-Prior, P.; Ong, E.; Coombs, G.; Hunter, M.; Oehler, R.; Pierson, D.; Gonzalez, R.; Huber, R.; Bode, W.; Madison, E. L.: Catalytic domain structures of MT-SP1/matriptase, a matrix-degrading transmembrane

serine proteinase. J. Biol. Chem. 277: 2160–2168,2002.

- [70415] 8745.Lin, C.–Y.; Wang, J.–K.; Torri, J.; Dou, L.; Sang, Q. A.; Dickson,R. B.: Characterization of a novel, membrane–bound, 80–kDa matrix–degradingprotease from human breast cancer cells: monoclonal antibody produc–tion,isolation, and localization. J. Biol. Chem. 272: 9147–9152, 1997.
- [70416] 8746.Gacy, A. M.; Goellner, G. M.; Spiro, C.; Chen, X.; Gupta, G.;Bradbury, E. M.; Dyer, R. B.; Mikesell, M. J.; Yao, J. Z.; Johnson,A. J.; Richter, A.; Melancon, S. B.; McMurray, C. T.: GAA instabilityin Friedreich's ataxia shares a com–mon, DNA–directed and intraallelicmechanism with other trinucleotide diseases. Molec. Cell 1: 583–593,1998.
- [70417] 8747.Gordon, D. M.; Kogan, M.; Knight, S. A. B.; Dancis, A.; Pain,D.: Distinct roles for two N–terminal cleaved do–mains in mitochondrialimport of the yeast frataxin ho–molog, Yfh1p. Hum. Molec. Genet. 10:259–269, 2001.
- [70418] 8748.Koenig, M.; Mandel, J.–L.: Deciphering the cause of Friedreichataxia. Curr. Opin. Neurobiol. 7: 689–694, 1997.
- [70419] 8749.Montermini, L.; Rodius, F.; Pianese, L.; Molto, M. D.; Cossee,M.; Campuzano, V.; Cavalcanti, F.; Monticelli, A.; Palau, F.; Gyapay,G.; Wenhert, M.; Zara, F.; Patel, P. I.; Co–

cozza, S.; Koenig, M.; Pandolfo, M.: The Friedreich ataxia critical region spans a 150-kb interval on chromosome 9q13. *Am. J. Hum. Genet.* 57: 1061–1067, 1995.

[70420] 8750. Pandolfo, M.: Personal Communication. Montreal, Canada 4/28/1997.

[70421] 8751. Ristow, M.; Pfister, M. F.; Yee, A. J.; Schubert, M.; Michael, L.; Zhang, C.-Y.; Ueki, K.; Michael, M. D., II; Lowell, B. B.; Kahn, C. R.: Frataxin activates mitochondrial energy conversion and oxidative phosphorylation. *Proc. Nat. Acad. Sci.* 97: 12239–12243, 2000.

[70422] 8752. Sakamoto, N.; Chastain, P. D.; Parniewski, P.; Ohshima, K.; Pandolfo, M.; Griffith, J. D.; Wells, R. D.: Sticky DNA: self-association properties of long GAA–TTC repeats in R–R–Y triplex structures from Friedreich's ataxia. *Molec. Cell* 3: 465–475, 1999.

[70423] 8753. Santos, M. M.; Ohshima, K.; Pandolfo, M.: Frataxin deficiency enhances apoptosis in cells differentiating into neuroectoderm. *Hum. Molec. Genet.* 10: 1935–1944, 2001.

[70424] 8754. Shoichet, S. A.; Baumer, A. T.; Stamenkovic, D.; Sauer, H.; Pfeiffer, A. F. H.; Kahn, C. R.; Muller-Wieland, D.; Richter, C.; Ristow, M.: Frataxin promotes antioxidant defense in a thiol-dependent manner resulting in diminished

malignant transformation in vitro. Hum. Molec.Genet. 11: 815–821, 2002.

[70425] 8755.Wilson, R. B.; Roof, D. M.: Respiratory deficiency due to loss of mitochondrial DNA in yeast lacking the frataxin homologue. NatureGenet. 16: 352–357, 1997.

[70426] 8756.Zuhlke, C.; Laccone, F.; Cossee, M.; Kohlschutter, A.; Koenig, M.; Schwinger, E.: Mutation of the start codon in the FRDA1 gene: linkage analysis of three pedigrees with the ATG to ATT transversion points to a unique common ancestor. Hum. Genet. 103: 102–105, 1998.

[70427] 8757.Fernandez–Gonzalez, A.; La Spada, A. R.; Treadaway, J.; Higdon, J. C.; Harris, B. S.; Sidman, R. L.; Morgan, J. I.; Zuo, J.: Purkinje cell degeneration (pcd) phenotypes caused by mutations in the axotomy–induced gene, Nna1. Science 295: 1904–1906, 2002.

[70428] 8758.Harris, A.; Morgan, J. I.; Pecot, M.; Soumare, A.; Osborne, A.; Soares, H. D.: Regenerating motor neurons express Nna1, a novel ATP/GTP–binding protein related to zinc carboxypeptidases. Molec. Cell. Neurosci. 16: 578–596, 2000.

[70429] 8759.Yuan, C.–X.; Ito, M.; Fondell, J. D.; Fu, Z.–Y.; Roeder, R. G.: The TRAP220 component of a thyroid hormone receptor–associated protein (TRAP) coactivator complex in–

teracts directly with nuclear receptors in a ligand-dependent fashion. *Proc. Nat. Acad. Sci.* 95: 7939–7944, 1998.

- [70430] 8760. Zhang, J.; Fondell, J. D.: Identification of mouse TRAP100: a transcriptional coregulatory factor for thyroid hormone and vitamin D receptors. *Molec. Endocr.* 13: 1130–1140, 1999.
- [70431] 8761. Csoka, T. B.; Frost, G. I.; Heng, H. H. Q.; Scherer, S. W.; Mohapatra, G.; Stern, R.: The hyaluronidase gene HYAL1 maps to chromosome 3p21.2–p21.3 in human and 9F1–F2 in mouse, a conserved candidate tumor suppressor locus. *Genomics* 48: 63–70, 1998.
- [70432] 8762. Frost, G. I.; Csoka, T. B.; Wong, T.; Stern, R.: Purification, cloning, and expression of human plasma hyaluronidase. *Biochem. Biophys. Res. Commun.* 236: 10–15, 1997.
- [70433] 8763. Laurent, T. C.; Fraser, J. R. E.: Hyaluronan. *FASEB J.* 6: 2397–2404, 1992.
- [70434] 8764. Zegerman, P.; Bannister, A. J.; Kouzarides, T.: The putative tumour suppressor Fus-2 is an N-acetyltransferase. *Oncogene* 19: 161–163, 2000.
- [70435] 8765. Brodbeck, J.; Davies, A.; Courtney, J.-M.; Meir, A.; Balaguero, N.; Canti, C.; Moss, F. J.; Page, K. M.; Pratt, W.

S.; Hunt, S. P.; Barclay, J.; Rees, M.; Dolphin, A. C.: The ducky mutation in Cacna2d2 results in altered Purkinje cell morphology and is associated with the expression of a truncated alpha-2/delta-2 protein with abnormal function. J. Biol. Chem. 277: 7684–7693, 2002.

[70436] 8766. Gao, B.; Sekido, Y.; Maximov, A.; Saad, M.; Forgacs, E.; Latif, F.; Wei, M. H.; Lerman, M.; Lee, J.-H.; Perez-Reyes, E.; Bezprozvanny, I.; Minna, J. D.: Functional properties of a new voltage-dependent calcium channel alpha-2/delta auxiliary subunit gene (CACNA2D2). J. Biol. Chem. 275: 12237–12242, 2000.

[70437] 8767. Angrand, P.-O.; Apiou, F.; Stewart, A. F.; Dutrillaux, B.; Losson, R.; Chambon, P.: NSD3, a new SET domain-containing gene, maps to 8p12 and is amplified in human breast cancer cell lines. Genomics 74:79–88, 2001.

[70438] 8768. Stec, I.; van Ommen, G.-J. B.; den Dunnen, J. T.: WHSC1L1, on human chromosome 8p11.2, closely resembles WHSC1 and maps to a duplicated region shared with 4p16.3. Genomics 76: 5–8, 2001.

[70439] 8769. Drabkin, H. A.; West, J. D.; Hotfilder, M.; Heng, Y. M.; Erickson, P.; Calvo, R.; Dalmau, J.; Gemmill, R. M.; Sablitzky, F.: DEF-3(g16/NY-LU-12), an RNA binding protein from the 3p21.3 homozygous deletion region in SCLC.

Oncogene 18: 2589–2597, 1999.

- [70440] 8770. Gure, A. O.; Altorki, N. K.; Stockert, E.; Scanlan, M. J.; Old, L. J.; Chen, Y.-T.: Human lung cancer antigens recognized by autologous antibodies: definition of a novel cDNA derived from the tumor suppressor gene locus on chromosome 3p21.3. *Cancer Res.* 58: 1034–1041, 1998.
- [70441] 8771. Timmer, T.; Terpstra, P.; van den Berg, A.; Veldhuis, P. M. J. F.; Elst, A. T.; Voutsinas, G.; Hulsbeek, M. M. F.; Draaijers, T. G.; Looman, M. W. G.; Kok, K.; Naylor, S. L.; Buys, C. H. C. M.: A comparison of genomic structures and expression patterns of two closely related flanking genes in a critical lung cancer region at 3p21.3. *Europ. J. Hum. Genet.* 7: 478–486, 1999.
- [70442] 8772. Singh, S.; Chao, L. Y.; Mishra, R.; Davies, J.; Saunders, G. F.: Missense mutation at the C-terminus of PAX6 negatively modulates homeodomain function. *Hum. Molec. Genet.* 10: 911–918, 2001.
- [70443] 8773. Singh, S.; Mishra, R.; Arango, N. A.; Deng, J. M.; Behringer, R. R.; Saunders, G. F.: Iris hypoplasia in mice that lack the alternatively spliced Pax6(5a) isoform. *Proc. Nat. Acad. Sci.* 99: 6812–6815, 2002.
- [70444] 8774. Singh, S.; Tang, H. K.; Lee, J.-Y.; Saunders, G. F.: Truncation mutations in the transactivation region of PAX6

result in dominant-negative mutants. *J. Biol. Chem.* 273: 21531–21541, 1998.

- [70445] 8775. Ton, C. C. T.; Miwa, H.; Saunders, G. F.: Small eye (Sey): cloning and characterization of the murine homolog of the human aniridia gene. *Genomics* 13:251–256, 1992.
- [70446] 8776. Walther, C.; Gruss, P.: Pax-6, a murine paired box gene, is expressed in the developing CNS. *Development* 113: 1435–1449, 1991.
- [70447] 8777. Wawersik, S.; Maas, R. L.: Vertebrate eye development as modeled in *Drosophila*. *Hum. Molec. Genet.* 9: 917–925, 2000.
- [70448] 8778. Xu, W.; Kozak, C. A.; Desnick, R. J.: Uroporphyrinogen-III synthase: molecular cloning, nucleotide sequence, expression of a mouse full-length cDNA, and its localization on mouse chromosome 7. *Genomics* 26: 556–562, 1995.
- [70449] 8779. Xu, W.; Warner, C. A.; Desnick, R. J.: Congenital erythropoietic porphyria: identification and expression of 10 mutations in the uroporphyrinogen III synthase gene. *J. Clin. Invest.* 95: 905–912, 1995.
- [70450] 8780. Winfield, S. L.; Tayebi, N.; Martin, B. M.; Ginns, E. I.; Sidransky, E.: Identification of three additional genes contiguous to the glucocerebrosidase locus on chromosome



1q21: implications for Gaucher disease. *GenomeRes.* 7: 1020–1026, 1997.

- [70451] 8781.Kawasawa, Y.; Kume, K.; Izumi, T.; Shimizu, T.: Mammalian PSP24s(alpha and beta isoforms) are not responsive to lysophosphatidic acid in mammalian expression systems. *Biochem. Biophys. Res. Commun.* 276:957–964, 2000.
- [70452] 8782.Kawasawa, Y.; Kume, K.; Nakade, S.; Haga, H.; Izumi, T.; Shimizu, T.: Brain-specific expression of novel G-protein-coupled receptors, with homologies to *Xenopus* PSP24 and human GPR45. *Biochem. Biophys. Res. Commun.* 276: 952–956, 2000.
- [70453] 8783.Lee, D. K.; George, S. R.; Cheng, R.; Nguyen, T.; Liu, Y.; Brown, M.; Lynch, K. R.; O'Dowd, B. F.: Identification of four novel human G protein-coupled receptors expressed in the brain. *Molec. BrainRes.* 86: 13–22, 2001.
- [70454] 8784.Cikos, S.; Gregor, P.; Koppel, J.: Cloning of a novel biogenic amine receptor-like G protein-coupled receptor expressed in human brain. *Biochim. Biophys. Acta* 1521: 66–72, 2001.
- [70455] 8785.Bascom, R. A.; Srinivasan, S.; Nussbaum, R. L.: Identification and characterization of golgin-84, a novel Golgi integral membrane protein with a cytoplasmic coiled-coil

domain. J. Biol. Chem. 274:2953–2962, 1999. Note: Erratum: J. Biol. Chem 274: 12950 only, 1999.

[70456] 8786. Ishizaka, Y.; Ochiai, M.; Tahira, T.; Sugimura, T.; Nagao, M.: Activation of the ret-II oncogene without a sequence encoding a transmembrane domain and transforming activity of two ret-II oncogene products differing in carboxy-termini due to alternative splicing. Oncogene 4: 789–794, 1989.

[70457] 8787. Klugbauer, S.; Demidchik, E. P.; Lengfelder, E.; Rabes, H. M.: Detection of a novel type of RET rearrangement (PTC5) in thyroid carcinomas after Chernobyl and analysis of the involved RET-fused gene RFG5. Cancer Res. 58: 198–203, 1998.

[70458] 8788. Jiang, J. C.; Kirchman, P. A.; Zagulski, M.; Hunt, J.; Jazwinski, S. M.: Homologs of the yeast longevity gene LAG1 in Caenorhabditis elegans and human. Genome Res. 8: 1259–1272, 1998.

[70459] 8789. Koivisto, U.-M.; Hubbard, A. L.; Mellman, I.: A novel cellular phenotype for familial hypercholesterolemia due to a defect in polarized targeting of LDL receptor. Cell 105: 575–585, 2001.

[70460] 8790. Koivisto, U.-M.; Turtola, H.; Aalto-Setälä, K.; Top, B.; Frants, R. R.; Kovanen, P. T.; Syvänen, A.-C.; Kontula, K.:

The familial hypercholesterolemia (FH)–North Karelia mutation of the low density lipoprotein receptor gene deletes seven nucleotides of exon 6 and is a common cause of FH in Finland. *J. Clin. Invest.* 90: 219–228, 1992.

[70461] 8791. Koivisto, U.–M.; Viikari, J. S.; Kontula, K.: Molecular characterization of minor gene rearrangements in Finnish patients with heterozygous familial hypercholesterolemia: identification of two common missense mutations (Gly823–to–Asp and Leu380–to–His) and eight rare mutations of the LDL receptor gene. *Am. J. Hum. Genet.* 57: 789–797, 1995.

[70462] 8792. Kotze, M. J.; Langenhoven, E.; Warnich, L.; du Plessis, L.; Retief, A. E.: The molecular basis and diagnosis of familial hypercholesterolaemia in South African Afrikaners. *Ann. Hum. Genet.* 55: 115–121, 1991.

[70463] 8793. Kotze, M. J.; Theart, L.; Peeters, A.; Langenhoven, E.: A *de novo* duplication in the low density lipoprotein receptor gene. *Hum. Mutat.* 6: 181–183, 1995.

[70464] 8794. Kotze, M. J.; Warnich, L.; Langenhoven, E.; du Plessis, L.; Retief, A. E.: An exon 4 mutation identified in the majority of South African familial hypercholesterolaemics. *J. Med. Genet.* 27: 298–302, 1990.

[70465] 8795. Landsberger, D.; Meiner, V.; Reshef, A.; Levy, Y.; van

der Westhuyzen, D. R.; Coetzee, G. A.; Leitersdorf, E.: A nonsense mutation in the LDL receptor gene leads to familial hypercholesterolemia in the Druzesect. *Am. J. Hum. Genet.* 50: 427–433, 1992.

[70466] 8796. Langlois, S.: Personal Communication. Vancouver, British Columbia, Canada 1989.

[70467] 8797. Langlois, S.; Kastelein, J. J. P.; Hayden, M. R.: Characterization of six partial deletions in the low-density-lipoprotein (LDL) receptor gene causing familial hypercholesterolemia (FH). *Am. J. Hum. Genet.* 43:60–68, 1988.

[70468] 8798. Lee, W. K.; Haddad, L.; Macleod, M. J.; Dorrance, A. M.; Wilson, D. J.; Gaffney, D.; Dominiczak, M. H.; Packard, C. J.; Day, I. N.; Humphries, S. E.; Dominiczak, A. F.: Identification of a common low density lipoprotein receptor mutation (C163Y) in the West of Scotland. *J. Med. Genet.* 35: 573–578, 1998.

[70469] 8799. Lehrman, M. A.; Goldstein, J. L.; Russell, D. W.; Brown, M. S.: Duplication of seven exons in LDL receptor gene caused by Alu–Alu recombination in a subject with familial hypercholesterolemia. *Cell* 48:827–835, 1987.

[70470] 8800. Lehrman, M. A.; Russell, D. W.; Goldstein, J. L.; Brown, M. S.: Alu–Alu recombination deletes splice accep–

tor sites and produces secreted low density lipoprotein receptor in a subject with familial hypercholesterolemia. J. Biol. Chem. 262: 3354–3361, 1987.

- [70471] 8801. Lehrman, M. A.; Russell, D. W.; Goldstein, J. L.; Brown, M. S.: Exon–Alu recombination deletes 5 kilobases from the low density lipoprotein receptor gene, producing a null phenotype in familial hypercholesterolemia. Proc. Nat. Acad. Sci. 83: 3679–3683, 1986.
- [70472] 8802. Lehrman, M. A.; Schneider, W. J.; Sudhof, T. C.; Brown, M. S.; Goldstein, J. L.; Russell, D. W.: Mutation in LDL receptor: Alu–Alu recombination deletes exons encoding transmembrane and cytoplasmic domains. Science 227: 140–146, 1985.
- [70473] 8803. Leitersdorf, E.; Hobbs, H. H.: Personal Communication. Dallas, Tex. 12/1990.
- [70474] 8804. Leitersdorf, E.; Hobbs, H. H.; Fourie, A. M.; Jacobs, M.; vander Westhuyzen, D. R.; Coetzee, G. A.: Deletion in the first cysteine–rich repeat of low–density lipoprotein receptor impairs its transport but not lipoprotein binding in fibroblasts from a subject with familial hypercholesterolemia. Proc. Nat. Acad. Sci. 85: 7912–7916, 1988.
- [70475] 8805. Leitersdorf, E.; Tobin, E. J.; Davignon, J.; Hobbs, H. H.: Common low–density lipoprotein receptor mutations in

the French Canadian population. J. Clin. Invest. 85: 1014–1023, 1990.

- [70476] 8806. Leitersdorf, E.; van der Westhuyzen, D. R.; Coetzee, G. A.; Hobbs, H. H.: Two common low density lipoprotein receptor gene mutations cause familial hypercholesterolemia in Afrikaners. J. Clin. Invest. 84:954–961, 1989.
- [70477] 8807. Lelli, N.; Ghisellini, M.; Calandra, S.; Gaddi, A.; Ciarrocchi, A.; Coviello, D. A.; Bertolini, S.: Duplication of exons 13, 14 and 15 of the LDL-receptor gene in a patient with heterozygous familial hypercholesterolemia. Hum. Genet. 86: 359–362, 1991.
- [70478] 8808. Leren, T. P.; Solberg, K.; Rodningen, O. K.; Tonstad, S.; Ose, L.: Two founder mutations in the LDL receptor gene in Norwegian familial hypercholesterolemia subjects. Atherosclerosis 111: 175–182, 1994.
- [70479] 8809. Li, H.; Gyllenstein, U. B.; Cui, X.; Saiki, R. K.; Erlich, H. A.; Arnheim, N.: Amplification and analysis of DNA sequences in single human sperm and diploid cells. Nature 335: 414–417, 1988.
- [70480] 8810. Downes, G. B.; Gilbert, D. J.; Copeland, N. G.; Gautam, N.; Jenkins, N. A.: Chromosomal mapping of five mouse G protein gamma subunits. Genomics 57:173–176, 1999.

- [70481] 8811.Modarressi, M. H.; Taylor, K. E.; Wolfe, J.: Cloning, characterization, and mapping of the gene encoding the human G protein gamma-2 subunit. *Biochem. Biophys. Res. Commun.* 272: 610–615, 2000.
- [70482] 8812.Yu, Y.; Zhang, C.; Zhou, G.; Wu, S.; Qu, X.; Wei, H.; Xing, G.; Dong, C.; Zhai, Y.; Wan, J.; Ouyang, S.; Li, L., Zhang, S.; Zhou, K.; Zhang, Y.; Wu, C.; He, F.: Gene expression profiling in human fetal liver and identification of tissue- and developmental-stage-specific genes through compiled expression profiles and efficient cloning of full-length cDNAs. *Genome Res.* 11: 1392–1403, 2001.
- [70483] 8813.Ericsson, J.; Greene, J. M.; Carter, K. C.; Shell, B. K.; Duan, D. R.; Florence, C.; Edwards, P. A.: Human geranylgeranyl diphosphate synthase: isolation of the cDNA, chromosomal mapping and tissue expression. *J. Lipid Res.* 39: 1731–1739, 1998.
- [70484] 8814.Kainou, T.; Kawamura, K.; Tanaka, K.; Matsuda, H.; Kawamukai, M.: Identification of the GGPS1 genes encoding geranylgeranyl diphosphate synthases from mouse and human. *Biochim. Biophys. Acta* 1437: 333–340, 1999.
- [70485] 8815.Kuzuguchi, T.; Morita, Y.; Sagami, I.; Sagami, H.; Ogura, K.: Human geranylgeranyl diphosphate synthase: cDNA cloning and expression. *J. Biol. Chem.* 274:

5888–5894, 1999.

[70486] 8816.Vicent, D.; Maratos-Flier, E.; Kahn, C. R.: The branch point enzyme of the mevalonate pathway for protein prenylation is overexpressed in the ob/ob mouse and induced by adipogenesis. *Molec. Cell. Biol.* 20:2158–2166, 2000.

[70487] 8817.Chen, C.-K.; Zhang, K.; Church-Kopish, J.; Huang, W.; Zhang, H.; Chen, Y.-J.; Frederick, J. M.; Baehr, W.: Characterization of human GRK7 as a potential cone opsin kinase. *Molec. Vision* 7: 305–313, 2001.

[70488] 8818.Weiss, E. R.; Ducceschi, M. H.; Horner, T. J.; Li, A.; Craft, C.M.; Osawa, S.: Species-specific differences in expression of G-protein-coupled receptor kinase (GRK) 7 and GRK1 in mammalian cone photoreceptor cells: implications for cone cell phototransduction. *J. Neurosci.* 21: 9175–9184, 2001.

[70489] 8819.Borregaard, N.; Cowland, J. B.: Granules of the human neutrophilic polymorphonuclear leukocyte. *Blood* 89: 3503–3521, 1997.

[70490] 8820.Chang, K. S.; Schroeder, W.; Siciliano, M. J.; Thompson, L. H.; McCredie, K.; Beran, M.; Freireich, E. J.; Liang, J. C.; Trujillo, J. M.; Stass, S. A.: The localization of the human myeloperoxidase gene is in close proximity to the



translocation breakpoint in acute promyelocytic leukemia. *Leukemia* 1: 458–462, 1987.

- [70491] 8821. DeLeo, F. R.; Goedken, M.; McCormick, S. J.; Nauseef, W. M.: A novel form of hereditary myeloperoxidase deficiency linked to endoplasmic reticulum/proteasome degradation. *J. Clin. Invest.* 101: 2900–2909, 1998.
- [70492] 8822. Eiserich, J. P.; Baldus, S.; Brennan, M.-L.; Ma, W.; Zhang, C.; Tousson, A.; Castro, L.; Lusa, A. J.; Nauseef, W. M.; White, C. R.; Freeman, B. A.: Myeloperoxidase, a leukocyte-derived vascular NO oxidase. *Science* 296: 2391–2394, 2002.
- [70493] 8823. Inazawa, J.; Inoue, K.; Nishigaki, H.; Tsuda, S.; Taniguchi, M.; Misawa, S.; Abe, T.: Assignment of the human myeloperoxidase gene (MPO) to bands q21.3–q23 of chromosome 17. *Cytogenet. Cell Genet.* 50: 135–136, 1989.
- [70494] 8824. Johnson, K.; Gemperlein, I.; Hudson, S.; Shane, S.; Rovera, G.: Complete nucleotide sequence of the human myeloperoxidase gene. *Nucleic Acids Res.* 17: 7985–7986, 1989.
- [70495] 8825. Kizaki, M.; Miller, C. W.; Selsted, M. E.; Koeffler, H. P.: Myeloperoxidase (MPO) gene mutation in hereditary MPO deficiency. *Blood* 83: 1935–1940, 1994.
- [70496] 8826. Klebanoff, S. J.: Myeloperoxidase. *Proc. Assoc. Am.*

Physicians 111:383–389, 1999.

- [70497] 8827.Kudoh, J.; Minoshima, S.; Hashinaka, K.; Nishio, C.; Yamada, M.; Shimizu, Y.; Shimizu, N.: Assignment of the myeloperoxidase gene MPO to human chromosome 17 using somatic cell hybrids and flow-sorted chromosomes. Jpn. J. Hum. Genet. 33: 315–324, 1988.
- [70498] 8828.Kudoh, J.; Minoshima, S.; Hashinaka, K.; Nishio, C.; Yamada, M.; Shimizu, Y.; Shimizu, N.: Assignment of the myeloperoxidase (MPO) gene to human chromosome 17. (Abstract) Cytogenet. Cell Genet. 46:641–642, 1987.
- [70499] 8829.Law, D. J.; Prasad, M. A.; King, S. E.; Spranger, K. D.; Lee, Y. H.; Fox, R. E.; Collins, E. E.; Gebuhr, T. C.; Miller, D. E.; Petty, E. M.: Localization of the human estrogen-responsive finger protein (EFP) gene (ZNF147) within a YAC contig containing the myeloperoxidase (MPO) gene. Genomics 28: 361–363, 1995.
- [70500] 8830.Williams, M.; Lyu, M.-S.; Yang, Y.-L.; Lin, E. P.; Dunbrack, R.; Birren, B.; Cunningham, J.; Hunter, K.: ler5, a novel member of the slow-kinetics immediate-early genes. Genomics 55: 327–334, 1999.
- [70501] 8831.Arce, I.; Roda-Navarro, P.; Montoya, M. C.; Hernanz-Falcon, P.; Puig-Kroger, A.; Fernandez-Ruiz, E.: Molecular and genomic characterization of human DLEC, a novel

member of the C-type lectin receptor gene family preferentially expressed on monocyte-derived dendritic cells. *Europ.J. Immun.* 31: 2733–2740, 2001.

- [70502] 8832.Dzionic, A.; Sohma, Y.; Nagafune, J.; Cella, M.; Colonna, M.; Facchetti, F.; Gunther, G.; Johnston, I.; Lanza-vecchia, A.; Nagasaka, T.; Okada, T.; Vermi, W.; Winkels, G.; Yamamoto, T.; Zysk, M.; Yamaguchi, Y.; Schmitz, J.: BDCA-2, a novel plasmacytoid dendritic cell-specific type II C-type lectin, mediates antigen capture and is a potent inhibitor of interferon alpha/beta induction. *J. Exp. Med.* 194: 1823–1834, 2001.
- [70503] 8833.Antonarakis, S. E.: Personal Communication. Baltimore, Md. 3/25/2002.
- [70504] 8834.McKemy, D. D.; Neuhausser, W. M.; Julius, D.: Identification of a cold receptor reveals a general role for TRP channels in thermosensation. *Nature* 416:52–58, 2002.
- [70505] 8835.Peier, A. M.; Moqrich, A.; Hergarden, A. C.; Reeve, A. J.; Andersson, D. A.; Story, G. M.; Earley, T. J.; Dragoni, I.; McIntyre, P.; Bevan, S.; Patapoutian, A.: A TRP channel that senses cold stimuli and menthol. *Cell* 108:705–715, 2002.
- [70506] 8836.Ashery-Padan, R.; Marquardt, T.; Zhou, X.; Gruss, P.: Pax6 activity in the lens primordium is required for lens formation and for correct placement of a single retina in

the eye. *Genes Dev.* 14: 2701–2711,2000.

[70507] 8837.Azuma, N.; Yamaguchi, Y.; Handa, H.; Hayakawa, M.; Kanai, A.; Yamada,M.: Missense mutation in the alternative splice region of the PAX6gene in eye anomalies. *Am. J. Hum. Genet.* 65: 656–663, 1999.

[70508] 8838.Roginski, R. S.; Mohan Raj, B. K.; Finkernagel, S. W.; Sciorra,L. J.: Assignment of an ionotropic glutamate receptor–like gene (GRINL1A)to human chromosome 15q22.1 by in situ hybridization. *Cytogenet.Cell Genet.* 93: 143–144, 2001.

[70509] 8839.Wydner, K. S.; Mohan Raj, B. K.; Sciorra, L. J.; Roginski, R. S.: The mouse orthologue of the human ionotropic glutamate receptor–likegene (GRINL1A) maps to mouse chromosome 9. *Cytogenet. Cell Genet.* 95:240–241, 2001.

[70510] 8840.Akiyama, H.; Hiraki, Y.; Noda, M.; Shigeno, C.; Ito, H.; Nakamura,T.: Molecular cloning and biological activity of a novel Ha–Ras suppressorgene predominantly expressed in skeletal muscle, heart, brain, andbone marrow by differential display using clonal mouse EC cells, ATDC5. *J.Biol. Chem.* 274: 32192–32197, 1999.

[70511] 8841.Ito, H.; Akiyama, H.; Shigeno, C.; Nakamura, T.: Isolation, characterization,and chromosome mapping of a human A–C1 Ha–Ras suppressor gene (HRASLS). *Cyto–*

genet.Cell Genet. 93: 36–39, 2001.

- [70512] 8842.Nakamura, T.; Yamazaki, Y.; Saiki, Y.; Moriyama, M.; Largaespada,D. A.; Jenkins, N. A.; Copeland, N. G.: Evi9 encodes a novel zincfinger protein that physically interacts with BCL6, a known humanB–cell proto–oncogene product. Molec. Cell Biol. 20: 3178–3186,2000.
- [70513] 8843.Saiki, Y.; Yamazaki, Y.; Yoshida, M.; Katoh, O.; Nakamura, T.:Human EVI9, a homologue of the mouse myeloid leukemia gene, is expressedin the hematopoietic progenitors and down–regulated during myeloiddifferentiation of HL60 cells. Genomics 70: 387–391, 2000.
- [70514] 8844.Satterwhite, E.; Sonoki, T.; Willis, T. G.; Harder, L.; Nowak,R.; Arriola, E. L.; Liu, H.; Price, H. P.; Gesk, S.; Steinemann, D.;Schlegelberger, B.; Oscier, D. G.; Siebert, R.; Tucker, P. W.; Dyer,M. J. S.: The BCL11 gene family: involvement of BCL11A in lymphoidmalignancies. Blood 98: 3413–3420, 2001.
- [70515] 8845.Frohlich, O.; Po, C.; Young, L. G.: Organization of the humangene encoding the epididymis–specific EP2 protein variants and itsrelationship to defensin genes. Biol. Reprod. 64: 1072–1079, 2001.
- [70516] 8846.Hamil, K. G.; Sivashanmugam, P.; Richardson, R. T.; Grossman, G.;Ruben, S. M.; Mohler, J. L.; Petrusz, P.;

O'rand, M. G.; French, F.S.; Hall, S. H.: HE2-beta and HE2-gamma, new members of an epididymis-specific-family of androgen-regulated proteins in the human. *Endocrinology* 141:1245-1253, 2000.

[70517] 8847.Jia, H. P.; Schutte, B. C.; Schudy, A.; Linzmeier, R.; Guthmiller, J. M.; Johnson, G. K.; Tack, B. F.; Mitros, J. P.; Rosenthal, A.; Ganz, T.; McCray, P. B., Jr.: Discovery of new human beta-defensins using a genomics-based approach. *Gene* 263: 211-218, 2001.

[70518] 8848.Li, P.; Chan, H. C.; He, B.; So, S. C.; Chung, Y. W.; Shang, Q.; Zhang, Y.-D.; Zhang, Y.-L.: An antimicrobial peptide gene found in the male reproductive system of rats. *Science* 291: 1783-1785, 2001.

[70519] 8849.Osterhoff, C.; Kirchhoff, C.; Krull, N.; Ivell, R.: Molecular cloning and characterization of a novel human sperm antigen (HE2) specifically expressed in the proximal epididymis. *Biol. Reprod.* 50:516-525, 1994.

[70520] 8850.Carim-Todd, L.; Escarceller, M.; Estivill, X.; Sumoy, L.: Cloning of the novel gene TM6SF1 reveals conservation of clusters of paralogous genes between human chromosomes 15q24-q26 and 19p13.3-p12. *Cytogenet. Cell Genet.* 90: 255-260, 2000.

[70521] 8851.Nagase, T.; Kikuno, R.; Ohara, O.: Prediction of the

coding sequences of unidentified human genes. XXI. The complete sequences of 60 new cDNA clones from brain which code for large proteins. DNA Res. 8:179–187, 2001.

- [70522] 8852. Gress, T. M.; Wallrapp, C.; Frohme, M.; Muller-Pillasch, F.; Lacher, U.; Friess, H.; Buchler, M.; Adler, G.; Hohenheisel, J. D.: Identification of genes with specific expression in pancreatic cancer by cDNA representational difference analysis. Genes Chromosomes Cancer 19: 97–103, 1997.
- [70523] 8853. Wallrapp, C.; Hahnel, S.; Muller-Pillasch, F.; Burghardt, B.; Iwamura, T.; Ruthenburger, M.; Lerch, M. M.; Adler, G.; Gress, T. M.: A novel transmembrane serine protease (TMPRSS3) overexpressed in pancreatic cancer. Cancer Res. 60: 2602–2606, 2000.
- [70524] 8854. Kiss, H.; Kedra, D.; Kiss, C.; Kost-Alimova, M.; Yang, Y.; Klein, G.; Imreh, S.; Dumanski, J. P.: The LZTFL1 gene is a part of a transcriptional map covering 250 kb within the common eliminated region 1 (C3CER1) in 3p21.3. Genomics 73: 10–19, 2001.
- [70525] 8855. Quentmeier, H.; Drexler, H. G.; Fleckenstein, D.; Zaborski, M.; Armstrong, A.; Sims, J. E.; Lyman, S. D.: Cloning of human thymic stromal lymphopoietin (TSLP) and signaling mechanisms leading to proliferation.

Leukemia 15:1286–1292, 2001.

[70526] 8856.Soumelis, V.; Reche, P. A.; Kanzler, H.; Yuan, W.; Edward, G.;Homey, B.; Gilliet, M.; Ho, S.; Antonenko, S.; Lauerma, A.; Smith,K.; Gorman, D.; Zurawski, S.; Abrams, J.; Menon, S.; McClanahan, T.;de Waal–Malefyt, R.; Bazan, F.; Kastelein, R. A.; Liu, Y.–J.: Humanepithelial cells trigger dendritic cell–mediated allergic inflammationby producing TSLP. Nature Immun. 3: 673–680, 2002.

[70527] 8857.Guo, D.; Hasham, S.; Kuang, S.–Q.; Vaughan, C. J.; Boerwinkle,E.; Chen, H.; Abuelo, D.; Dietz, H. C.; Basson, C. T.; Shete, S. S.;Milewicz, D. M.: Familial thoracic aortic aneurysms and dissections:genetic heterogeneity with a major locus mapping to 5q13–14. Circulation 103:2461–2468, 2001.

[70528] 8858.Vaughan, C. J.; Casey, M.; He, J.; Veuglers, M.; Henderson, K.;Guo, D.; Campagna, R.; Roman, M. J.; Milewicz, D. M.; Devereux, R.B.; Basson, C. T.: Identification of a chromosome 11q23.2–q24 locusfor familial aortic aneurysm disease, a genetically heterogeneousdisorder. Circulation 103: 2469–2475, 2001.

[70529] 8859.Tapon, N.; Harvey, K. F.; Bell, D. W.; Wahrer, D. C. R.; Schiripo,T. A.; Haber, D. A.; Hariharan, I. K.: salvador promotes both cellcycle exit and apoptosis in Drosophila and



is mutated in human cancer cell lines. *Cell* 110: 467–478, 2002.

[70530] 8860. Valverde, P.: Cloning, expression, and mapping of hWW45, a novel human WW domain-containing gene. *Biochem. Biophys. Res. Commun.* 276:990–998, 2000.

[70531] 8861. Wang, X.; McLachlan, J.; Zamore, P. D.; Hall, T. M. T.: Modular recognition of RNA by a human Pumilio-homology domain. *Cell* 110:501–512, 2002.

[70532] 8862. Ballinger, C. A.; Connell, P.; Wu, Y.; Hu, Z.; Thompson, L. J.; Yin, L.-Y.; Patterson, C.: Identification of CHIP, a novel tetratricopeptide repeat-containing protein that interacts with heat shock proteins and negatively regulates chaperone functions. *Molec. Cell. Biol.* 19:4535–4545, 1999.

[70533] 8863. Jiang, J.; Ballinger, C. A.; Wu, Y.; Dai, Q.; Cyr, D. M.; Hohfeld, J.; Patterson, C.: CHIP is a U-box-dependent E3 ubiquitin ligase: identification of Hsc70 as a target for ubiquitylation. *J. Biol. Chem.* 276: 42938–42944, 2001.

[70534] 8864. Gaide, O.; Martinon, F.; Micheau, O.; Bonnet, D.; Thome, M.; Tschopp, J.: Carma1, a CARD-containing binding partner of Bcl10, induces Bcl10 phosphorylation and NF- $\kappa$ B activation. *FEBS Lett.* 496: 121–127, 2001.

[70535] 8865. McAllister-Lucas, L. M.; Inohara, N.; Lucas, P. C.; Ru-

land, J.; Benito, A.; Li, Q.; Chen, S.; Chen, F. F.; Yamaoka, S.; Verma, I.M.; Mak, T. W.; Nunez, G.: Bimp1, a MAGUK family member linking protein kinase C activation to Bcl10-mediated NF-kappa-B induction. *J. Biol. Chem.* 276: 30589–30597, 2001.

[70536] 8866. Wang, L.; Guo, Y.; Huang, W.-J.; Ke, X.; Poyet, J.-L.; Manji, G.A.; Merriam, S.; Glucksmann, M. A.; DiStefano, P. S.; Alnemri, E.S.; Bertin, J.: CARD10 is a novel caspase recruitment domain/membrane-associated guanylate kinase family member that interacts with BCL10 and activates NF-kappa-B. *J. Biol. Chem.* 276: 21405–21409, 2001.

[70537] 8867. Wagner, D. S.; Gan, L.; Klein, W. H.: Identification of a differentially expressed RNA helicase by gene trapping. *Biochem. Biophys. Res. Commun.* 262: 677–684, 1999.

[70538] 8868. Takeuchi, T.; Shuman, M. A.; Craik, C. S.: Reverse biochemistry: use of macromolecular protease inhibitors to dissect complex biological processes and identify a membrane-type serine protease in epithelial cancer and normal tissue. *Proc. Nat. Acad. Sci.* 96: 11054–11061, 1999.

[70539] 8869. Loux, N.; Benlian, P.; Pastier, D.; Boileau, C.; Cambou, J. P.; Monnier, L.; Percheron, C.; Junien, C.: Recurrent mutation at aa792 in the LDL receptor gene in a French patient. *Hum. Genet.* 87: 373–375, 1991.

- [70540] 8870. Le Beau, M. M.; Lemons, R. S.; Rosner, G. L.; Carrino, J. C.; Reid, M. S.; Chisholm, R. L.; Diaz, M. O.; Weil, S. C.: Chromosomal localization of the gene encoding myeloperoxidase. (Abstract) *Cytogenet. Cell Genet.* 46: 645, 1987.
- [70541] 8871. De Deken, X.; Wang, D.; Many, M.-C.; Costagliola, S.; Libert, F.; Vassart, G.; Dumont, J. E.; Miot, F.: Cloning of two human thyroid cDNAs encoding new members of the NADPH oxidase family. *J. Biol. Chem.* 275: 23227–23233, 2000.
- [70542] 8872. Lacroix, L.; Nocera, M.; Mian, C.; Caillou, B.; Virion, A.; Dupuy, C.; Filetti, S.; Bidart, J. M.; Schlumberger, M.: Expression of nicotinamide adenine dinucleotide phosphate oxidase flavoprotein DUOX genes and proteins in human papillary and follicular thyroid carcinomas. *Thyroid* 11:1017–1023, 2001.
- [70543] 8873. Dupuy, C.; Ohayon, R.; Valent, A.; Noel-Hudson, M.-S.; Deme, D.; Virion, A.: Purification of a novel flavoprotein involved in the thyroid NADPH oxidase: cloning of the porcine and human cDNAs. *J. Biol. Chem.* 274: 37265–37269, 1999.
- [70544] 8874. Moreno, J. C.; Bikker, H.; Kempers, M. J. E.; van Trotsenburg, A. S. P.; Baas, F.; de Vijlder, J. J. M.; Vulsma, T.; Ris-Stalpers, C.: Inactivating mutations in the gene for thy-

roid oxidase 2 (THOX2) and congenital hypothyroidism.

New Eng. J. Med. 347: 95–102, 2002.

- [70545] 8875. Okajima, T.; Fukumoto, S.; Miyazaki, H.; Ishida, H.; Kiso, M.; Furukawa, K.; Urano, T.; Furukawa, K.: Molecular cloning of a novel  $\alpha$ -2,3-sialyltransferase (ST3Gal VI) that sialylates type II lactosamine structures on glycoproteins and glycolipids. J. Biol. Chem. 274:11479–11486, 1999.
- [70546] 8876. Richardson, J.; Cvekl, A.; Wistow, G.: Pax-6 is essential for lens-specific expression of zeta-crystallin. Proc. Nat. Acad. Sci. 92:4676–4680, 1995.
- [70547] 8877. Sander, M.; Neubuser, A.; Kalamaras, J.; Ee, H. C.; Martin, G.R.; German, M. S.: Genetic analysis reveals that PAX6 is required for normal transcription of pancreatic hormone genes and islet development. Genes Dev. 11: 1662–1673, 1997.
- [70548] 8878. Taniguchi, A.; Kaneta, R.; Morishita, K.; Matsumoto, K.: Gene structure and transcriptional regulation of human Gal beta-1,4(3)GlcNac  $\alpha$ -2,3-sialyltransferase VI (hST3Gal VI) gene in prostate cancer cell line. Biochem. Biophys. Res. Commun. 287: 1148–1156, 2001.
- [70549] 8879. Schwientek, T.; Nomoto, M.; Levery, S. B.; Merks, G.; van Kessel, A. G.; Bennett, E. P.; Hollingsworth, M. A.;

Clausen, H.: Control of O-glycan branch formation: molecular cloning of human cDNA encoding a novel beta-1,6-N-acetylglucosaminyltransferase forming core 2 and-core 4. *J. Biol. Chem.* 274: 4504–4512, 1999.

[70550] 8880. Chano, T.; et al; et al: Isolation, characterization and mapping of the mouse and human RB1CC1 genes. *Gene* (in-press), 2002.

[70551] 8881. Chano, T.; Kontani, K.; Teramoto, K.; Okabe, H.; Ikegawa, S.: Truncating mutations of RB1CC1 in human breast cancers. *Nature Genet.* 31:285–288, 2002.

[70552] 8882. Kaneko, M.; Kudo, T.; Iwasaki, H.; Ikehara, Y.; Nishihara, S.; Nakagawa, S.; Sasaki, K.; Shiina, T.; Inoko, H.; Saitou, N.; Narimatsu, H.: Alpha-1,3-fucosyltransferase (sic) IX (Fuc-TIX) is very highly conserved between human and mouse; molecular cloning, characterization and tissue distribution of human Fuc-TIX. *FEBS Lett.* 452: 237–242, 1999.

[70553] 8883. Kaneko, M.; Kudo, T.; Iwasaki, H.; Shiina, T.; Inoko, H.; Kozaki, T.; Saitou, N.; Narimatsu, H.: Assignment of the human alpha-1,3-fucosyltransferase IX gene (FUT9) to chromosome band 6q16 by in situ hybridization. *Cytogenet. Cell Genet.* 86: 329–330, 1999.

[70554] 8884. Lane, J. D.; Lucocq, J.; Pryde, J.; Barr, F. A.; Woodman,

P. G.;Allan, V. J.; Lowe, M.: Caspase-mediated cleavage of the stackingprotein GRASP65 is required for Golgi fragmentation during apoptosis. *J.Cell Biol.* 156: 495–509, 2002.

[70555] 8885.Sutterlin, C.; Hsu, P.; Mallabiabarrena, A.; Malhotra, V.: Fragmentationand dispersal of the pericentriolar Golgi complex is required forentry into mitosis in mammalian cells. *Cell* 109: 359–369, 2002.

[70556] 8886.Davis, A.; Cowell, J. K.: Mutations in the PAX6 gene in patientswith hereditary aniridia. *Hum. Molec. Genet.* 2: 2093–2097, 1993.

[70557] 8887.Gronskov, K.; Rosenberg, T.; Sand, A.; Brondum-Nielsen, K.: Mutationalanalysis of PAX6: 16 novel mutations including 5 missense mutationswith a mild aniridia phenotype. *Europ. J. Hum. Genet.* 7: 274–286,1999.

[70558] 8888.Halder, G.; Callaerts, P.; Gehring, W. J.: Induction of ectopiceyes by targeted expression of the eyeless gene in *Drosophila*. *Science* 267:1788–1792, 1995.

[70559] 8889.Hanson, I.; Brown, A.; van Heyningen, V.: A new PAX6 mutationin familial aniridia. *J. Med. Genet.* 32: 488–489, 1995.

[70560] 8890.Hanson, I.; Van Heyningen, V.: Pax6: more than meets the eye. *TrendsGenet.* 11: 268–272, 1995.

- [70561] 8891.Hanson, I. M.; Fletcher, J. M.; Jordon, T.; Brown, A.; Taylor,D.; Adams, R. J.; Punnett, H. H.; van Heyningen, V.: Mutations atthe PAX6 locus are found in heterogeneous anterior segment malformationsincluding Peters' anomaly. *Nature Genet.* 6: 168–173, 1994.
- [70562] 8892.Heins, N.; Malatesta, P.; Cecconi, F.; Nakafuku, M.; Tucker, K.L.; Hack, M. A.; Chapouton, P.; Barde, Y.–A.; Gotz, M.: Glial cellsgenerate neurons: the role of the transcription factor Pax6. *NatureNeurosci.* 5: 308–315, 2002.
- [70563] 8893.Holmstrom, G. E.; Reardon, W. P.; Baraitser, M.; Elston, J. S.;Taylor, D. S.: Heterogeneity in dominant anterior segment malformations.. *Brit.J. Ophthal.* 75: 591–597, 1991.
- [70564] 8894.Kioussi, C.; O'Connell, S.; St-Onge, L.; Treier, M.; Gleiberman,A. S.; Gruss, P.; Rosenfeld, M. G.: Pax6 is essential for establishingventral–dorsal cell boundaries in pituitary gland development. *Proc.Nat. Acad. Sci.* 96: 14378–14382, 1999.
- [70565] 8895.Kleinjan, D. A.; Seawright, A.; Schedl, A.; Quinlan, R. A.; Danes,S.; van Heyningen, V.: Aniridia–associated translocations, DNasehypersensitivity, sequence comparison and transgenic analysis redefinethe functional domain of PAX6. *Hum. Molec. Genet.* 10: 2049–2059,2001.

- [70566] 8896.Lauderdale, J. D.; Wilensky, J. S.; Oliver, E. R.; Walton, D.S.; Glaser, T.: 3-prime deletions cause aniridia by preventing PAX6gene expression. *Proc. Nat. Acad. Sci.* 97: 13755–13759, 2000.
- [70567] 8897.Marquardt, T.; Ashery-Padan, R.; Andrejewski, N.; Scardigli, R.;Guillemot, F.; Gruss, P.: Pax6 is required for the multipotent stateof retinal progenitor cells. *Cell* 105: 43–55, 2001.
- [70568] 8898.Croze, E.; Russell-Harde, D.; Wagner, T. C.; Pu, H.; Pfeffer, L.M.; Perez, H. D.: The human type I interferon receptor: identificationof the interferon beta-specific receptor-associated phosphoprotein. *J.Biol. Chem.* 271: 33165–33168, 1996.
- [70569] 8899.Croze, E.; Usacheva, A.; Asarnow, D.; Minshall, R.D.; Perez, H.D.;Colamonici, O.: Receptor for activated C-kinase (RACK-1), a WD motif-containingprotein, specifically associates withthe human type I IFN receptor. *J.Immun.* 165: 5127–5132, 2000.
- [70570] 8900.Domanski, P.; Witte, M.; Kellum, M.; Rubinstein, M.; Hackett, R.;Pitha, P.; Colamonici, O. R.: Cloning and expression of a long formof the beta subunit of the interferon alpha/beta receptor that isrequired for signaling. *J. Biol. Chem.* 270: 21606–21611, 1995.



- [70571] 8901.Lutfalla, G.; Holland, S. J.; Cinato, E.; Monneron, D.; Reboul,J.; Rogers, N. C.; Smith, J. M.; Stark, G. R.; Gardiner, K.; Mogensen,K. E.; Kerr, I. M.; Uze, G.: Mutant U5A cells are complemented byan interferon-alpha/beta receptor subunit generated by alternativeprocessing of a new member of a cytokine receptor gene cluster. EMBOJ. 14: 5100-5108, 1995.
- [70572] 8902.Platanias, L. C.; Uddin, S.; Domanski, P.; Colamonici, O. R.:Differences in interferon alpha and beta signaling: interferon betaselectively induces the interaction of the alpha and beta(L) subunitsof the type I interferon receptor. J. Biol. Chem. 271: 23630-23633,1996.
- [70573] 8903.Raz, R.; Cheung, K.; Ling, L.; Levy, D. E.: Three distinct locion human chromosome 21 contribute to interferon-alpha/beta responsiveness. Somat.Cell Molec. Genet. 21: 139-145, 1995.
- [70574] 8904.Klocke, R.; Augustin, A.; Ronsiek, M.; Stief, A.; van der Putten,H.; Jockusch, H.: Dynamin genes Dnm1 and Dnm2 are located on proximalmouse chromosomes 2 and 9, respectively. Genomics 41: 290-292, 1997.
- [70575] 8905.Marks, B.; Stowell, M. H. B.; Vallis, Y.; Mills, I. G.; Gibson,A.; Hopkins, C. R.; McMahon, H. T.: GTPase activity of dynamin andresulting conformation change are essen-

tial for endocytosis. *Nature* 410:231–235, 2001.

- [70576] 8906. Newman–Smith, E. D.; Shurland, D.–L.; van der Blik, A. M.: Assignment of the dynamin–1 gene (DNM1) to human chromosome 9q34 by fluorescence in situ hybridization and somatic cell hybrid analysis. *Genomics* 41:286–289, 1997.
- [70577] 8907. Sontag, J.–M.; Fykse, E. M.; Ushkaryov, Y.; Liu, J.–P.; Robinson, P. J.; Sudhof, T. C.: Differential expression and regulation of multiple dynamins. *J. Biol. Chem.* 269: 4547–4554, 1994.
- [70578] 8908. Sweitzer, S. M.; Hinshaw, J. E.: Dynamin undergoes a GTP–dependent conformational change causing vesiculation. *Cell* 93: 1021–1029, 1998.
- [70579] 8909. van der Blik, A. M.; Redelmeier, T. E.; Damke, H.; Tisdale, E. J.; Meyerowitz, E. M.; Schmid, S. L.: Mutations in human dynamin block an intermediate stage in coated vesicle formation. *J. Cell Biol.* 122: 553–563, 1993.
- [70580] 8910. Diatloff–Zito, C.; Gordon, A. J. E.; Duchaud, E.; Merlin, G.: Isolation of an ubiquitously expressed cDNA encoding human dynamin II, a member of the large GTP–binding protein family. *Gene* 163:301–306, 1995.
- [70581] 8911. Finch, J. L.; Webb, G. C.; Evdokiou, A.; Cowled, P. A.: Chromosomal localization of the human urothelial

tetraspan gene, UPK1B, to 3q13.3–q21 and detection of a Taq1 polymorphism. *Genomics* 40: 501–503, 1997.

[70582] 8912. Yu, J.; Lin, J.-H.; Wu, X.-R.; Sun, T.-T.: Uroplakins Ia and Ib, two major differentiation products of bladder epithelium, belong to a family of four transmembrane domain (4TM) proteins. *J. Cell Biol.* 125:171–182, 1994.

[70583] 8913. Svaren, J.; Sevetson, B. R.; Apel, E. D.; Zimonjic, D. B.; Popescu, N. C.; Milbrandt, J.: NAB2, a corepressor of NGFI-A (Egr-1) and Krox20, is induced by proliferative and differentiative stimuli. *Molec. Cell Biol.* 16: 3545–3553, 1996.

[70584] 8914. Colley, W. C.; Altshuler, Y. M.; Sue-Ling, C. K.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Branch, K. D.; Tsirka, S. E.; Bollag, R. J.; Bollag, W. B.; Frohman, M. A.: Cloning and expression analysis of murine phospholipase D1. *Biochem. J.* 326: 745–753, 1997.

[70585] 8915. Bentz, H.; Nathan, R. M.; Rosen, D. M.; Armstrong, R. M.; Thompson, A. Y.; Segarini, P. R.; Mathews, M. C.; Dasch, J. R.; Piez, K. A.; Seyedin, S. M.: Purification and characterization of a unique osteoinductive factor from bovine bone. *J. Biol. Chem.* 264: 20805–20810, 1989.

[70586] 8916. Madisen, L.; Neubauer, M.; Plowman, G.; Rosen, D.; Segarini, P.; Dasch, J.; Thompson, A.; Ziman, J.; Bentz, H.;

Purchio, A. F.: Molecularcloning of a novel bone-forming compound: osteoinductive factor. DNACell Biol. 9: 303–309, 1990.

- [70587] 8917.Fischer, G.; Perez–Rodriguez, M.; Arguello, J. R.; Cox, S. T.;McWhinnie, A.; Travers, P. J.; Madrigal, J. A.: Three novel MICBalleles. Tissue Antigens 55: 166–170, 2000.
- [70588] 8918.Groh, V.; Bahram, S.; Bauer, S.; Herman, A.; Beauchamp, M.; Spies,T.: Cell stress–regulated human major histocompatibility complexclass I gene expressed in gastrointestinal epithelium. Proc. Nat.Acad. Sci. 93: 12445–12450, 1996.
- [70589] 8919.Nakai, A.; Tanabe, M.; Kawazoe, Y.; Inazawa, J.; Morimoto, R. I.;Nagata, K.: HSF4, a new member of the human heat shock factor familywhich lacks properties of a transcriptional activator. Molec. Cell.Biol. 17: 469–481, 1997.
- [70590] 8920.Uchida, K.; Yoshimura, A.; Inazawa, J.; Yanagisawa, K.; Osada,H.; Masuda, A.; Saito, T.; Takahashi, T.; Miyajima, A.; Takahashi,T.: Molecular cloning of CISH, chromosome assignment to 3p21.3, andanalysis of expression in fetal and adult tissues. Cytogenet. CellGenet. 78: 209–212, 1997.

- [70591] 8921.Yoshimura, A.; Ohkubo, T.; Kiguchi, T.; Jenkins, N. A.; Gilbert,D. J.; Copeland, N. G.; Hara, T.; Miyajima, A.: A novel cytokine-inducible gene CIS, encodes an SH2-containing protein that binds to tyrosine-phosphorylated interleukin 3 and erythropoietin receptors. EMBO J. 14: 2816–2826,1995.
- [70592] 8922.Chen, H.; Antonarakis, S. E.: The SH3D1A gene maps to human chromosome 21q22.1–q22.2. Cytogenet. Cell Genet. 78: 213–215, 1997.
- [70593] 8923.Pucharcos, C.; Estivill, X.; de la Luna, S.: Intersectin 2, a new multimodular protein involved in clathrin-mediated endocytosis. FEBS Lett. 478: 43–51, 2000.
- [70594] 8924.Szepietowski, P.; Simon, M.-P.; Grosgeorge, J.; Huebner, K.; Bastard,C.; Evans, G. A.; Tsujimoto, Y.; Birnbaum, D.; Theillet, C.; Gaudray,P.: Localization of 11q13 loci with respect to regional chromosomal breakpoints. Genomics 12: 738–744, 1992.
- [70595] 8925.Wang, T. C.; Cardiff, R. D.; Zukerberg, L.; Lees, E.; Arnold,A.; Schmidt, E. V.: Mammary hyperplasia and carcinoma in MMTV-cyclinD1 transgenic mice. Nature 369: 669–671, 1994.
- [70596] 8926.Xiong, Y.; Connelly, T.; Futcher, B.; Beach, D.: Human D-type cyclin. Cell 65: 691–699, 1991.

- [70597] 8927.Yu, Q.; Geng, Y.; Sicinski, P.: Specific protection against breastcancers by cyclin D1 ablation. *Nature* 411: 1017–1021, 2001.
- [70598] 8928.Zatyka, M.; da Silva, N. F.; Clifford, S. C.; Morris, M. R.; Wiesener, M. S.; Eckardt, K.–U.; Houlston, R. S.; Richards, F. M.; Latif, F.; Maher, E. R.: Identification of cyclin D1 and other novel targets for the von Hippel–Lindau tumor suppressor gene by expression array analysis and investigation of cyclin D1 genotype as a modifier in von Hippel–Lindau disease. *Cancer Res.* 62: 3803–3811, 2002.
- [70599] 8929.Skoldberg, F.; Grimelius, L.; Woodward, E. R.; Rorsman, F.; VanSchothorst, E. W.; Winqvist, O.; Karlsson, F. A.; Akerstrom, G.; Kampe, O.; Husebye, E. S.: A family with hereditary extra–adrenal paragangliomas without evidence for mutations in the von Hippel–Lindau disease or ret genes. *Clin. Endocr.* 48: 11–16, 1998.
- [70600] 8930.Gibbons, B.; Scott, D.; Hungerford, J. L.; Cheung, K. L.; Harrison, C.; Attard–Montalto, S.; Evans, M.; Birch, J. M.; Kingston, J. E.: Retinoblastoma in association with the chromosome breakage syndromes Fanconi's anaemia and Bloom's syndrome: clinical and cytogenetic findings. *Clin. Genet.* 47: 311–317, 1995.
- [70601] 8931.Girardet, A.; McPeck, M. S.; Leeflang, E. P.; Munier,

F.; Arnheim, N.; Claustres, M.; Pellestor, F.: Meiotic segregation analysis of RB1 alleles in retinoblastoma pedigrees by use of single-sperm typing. *Am. J. Hum. Genet.* 66: 167–175, 2000.

[70602] 8932. Godbout, R.; Dryja, T. P.; Squire, J.; Gallie, B. L.; Phillips, R. A.: Somatic inactivation of genes on chromosome 13 is a common event in retinoblastoma. *Nature* 304: 451–453, 1983.

[70603] 8933. Goodrich, D. W.; Wang, N. P.; Qian, Y.-W.; Lee, E. Y.-H. P.; Lee, W.-H.: The retinoblastoma gene product regulates progression through the G1 phase of the cell cycle. *Cell* 67: 293–302, 1991.

[70604] 8934. Grace, E.; Drennan, J.; Colver, D.; Gordon, R. R.: The 13q deletion syndrome. *J. Med. Genet.* 8: 351–357, 1971.

[70605] 8935. Green, A. R.; Wyke, J. A.: Anti-oncogenes: a subset of regulatory genes involved in carcinogenesis? *Lancet* II: 475–477, 1985.

[70606] 8936. Greger, V.; Kerst, S.; Messmer, E.; Hopping, W.; Passarge, E.; Horsthemke, B.: Application of linkage analysis to genetic counselling in families with hereditary retinoblastoma. *J. Med. Genet.* 25: 217–221, 1988.

[70607] 8937. Greger, V.; Passarge, E.; Horsthemke, B.: Somatic mosaicism in a patient with bilateral retinoblastoma. *Am. J.*

Hum. Genet. 46:1187–1193, 1990.

- [70608] 8938.Hagstrom, S. A.; Dryja, T. P.: Mitotic recombination map of 13cen–13q14derived from an investigation of loss of heterozygosity in retinoblastomas. Proc.Nat. Acad. Sci. 96: 2952–2957, 1999.
- [70609] 8939.Hall, J. G.: Personal Communication. Vancouver, British Columbia,Canada 5/29/1993.
- [70610] 8940.Hanahan, D.; Weinberg, R. A.: The hallmarks of cancer. Cell 100:57–70, 2000.
- [70611] 8941.Harbour, J. W.: Molecular basis of low–penetrance retinoblastoma. Arch.Ophthal. 119: 1699–1704, 2001.
- [70612] 8942.Harbour, J. W.; Lai, S.–L.; Whang–Peng, J.; Gazdar, A. F.; Minna,J. D.; Kaye, F. J.: Abnormalities in structure and expression ofthe human retinoblastoma gene in SCLC. Science 241: 353–357, 1988.
- [70613] 8943.Hensel, C.; Hsieh, C.–L.; Lee, W.–H.; Pam–Lee, E.; Gazdar, A.;Sakaguchi, A. Y.; Naylor, S. L.: Allele loss and lack of expressionof the RB–1 locus in small cell lung cancer. (Abstract) Am. J. Hum.Genet. 43: A25, 1988.
- [70614] 8944.Henson, J. W.; Schnitker, B. L.; Correa, K. M.; von Diemling,A.; Fassbender, F.; Xu, H.–J.; Benedict, W. F.; Yandell, D. W.; Louis,D. N.: The retinoblastoma gene is involved in malignant progressionof astrocytomas. Ann.



Neurol. 36: 714–721, 1994.

- [70615] 8945.Higgins, M. J.; Hansen, M. F.; Cavenee, W. K.; Lalande, M.: Molecular detection of chromosomal translocations that disrupt the putative retinoblastoma susceptibility locus. *Molec. Cell. Biol.* 9: 1–5, 1989.
- [70616] 8946.Hoegerman, S. F.: Chromosome 13 long arm interstitial deletion may result from maternal inverted insertion. *Science* 205: 1035–1036, 1979.
- [70617] 8947.Hogg, A.; Bia, B.; Onadim, Z.; Cowell, J. K.: Molecular mechanisms of oncogenic mutations in tumours from patients with bilateral and unilateral retinoblastoma. *Proc. Nat. Acad. Sci.* 90: 7351–7355, 1993.
- [70618] 8948.Honavar, S. G.; Shields, C. L.; Shields, J. A.; Demirci, H.; Naduvilath, T. J.: Intraocular surgery after treatment of retinoblastoma. *Arch. Ophthalmol.* 119: 1613–1621, 2001.
- [70619] 8949.Honavar, S. G.; Singh, A. D.; Shields, C. L.; Meadows, A. T.; Demirci, H.; Cater, J.; Shields, J. A.: Postenucleation adjuvant therapy in high-risk retinoblastoma. *Arch. Ophthalmol.* 120: 923–931, 2002.
- [70620] 8950.Hong, F. D.; Huang, H.-J. S.; To, H.; Young, L.-J. S.; Oro, A.; Bookstein, R.; Lee, E. Y.-H. P.; Lee, W.-H.: Structure of the human retinoblastoma gene. *Proc. Nat. Acad. Sci.* 86: 5502–5506, 1989. 100. Horowitz, J. M.; Park,

S.-H.; Bogenmann, E.; Cheng, J.-C.; Yandell, D. W.; Kaye, F. J.; Minna, J. D.; Dryja, T. P.; Weinberg, R. A.: Frequent inactivation of the retinoblastoma anti-oncogene is restricted to a subset of human tumor cells. *Proc. Nat. Acad. Sci.* 87: 2775–2779, 1990. 101. Horowitz, J. M.; Park, S. H.; Yandell, D. W.; Weinberg, R. A.: Involvement of the retinoblastoma gene in the genesis of various human tumors.: In: Kavenee, W.; Hastie, N.; Stanbridge, E.: *Recessive Oncogenes and Tumor Suppression: Current Communications in Molecular Biology*. Cold Spring Harbor, New York: Cold Spring Harbor Laboratory Press (pub.) 1989. Pp. 101–108. 102. Horsthemke, B.; Greger, V.; Barnert, H. J.; Hopping, W.; Passarge, E.: Detection of submicroscopic deletions and a DNA polymorphism at the retinoblastoma locus. *Hum. Genet.* 76: 257–261, 1987. 103. Horsthemke, B.; Greger, V.; Becher, R.; Passarge, E.: Mechanism of i(6p) formation in retinoblastoma tumor cells. *Cancer Genet. Cytogenet.* 37: 95–102, 1989. 104. Hsieh, J.-K.; Chan, F. S. G.; O'Connor, D. J.; Mitnacht, S.; Zhong, S.; Lu, X.: RB regulates the stability and the apoptotic function of p53 via MDM2. *Molec. Cell* 3: 181–193, 1999. 105. Huang, H.-J. S.; Yee, J.-K.; Shew, J.-Y.; Chen, P.-L.; Bookstein, R.; Friedmann, T.; Lee, E. Y.-H. P.; Lee, W.-H.: Suppression of the–

neoplastic phenotype by replacement of the RB gene in human cancer cells. *Science* 242: 1563–1566, 1988.106. Janson, M.; Nordenskjold, M.: A constitutional mutation within the retinoblastoma gene detected by PFGE. *Clin. Genet.* 45: 5–10, 1994.107. Jensen, R. D.; Miller, R. W.: Retinoblastoma: epidemiologic characteristics. *New Eng. J. Med.* 285: 307–311, 1971.108. Kimchi, A.; Wang, X.-F.; Weinberg, R. A.; Cheifetz, S.; Massague, J.: Absence of TGF- $\beta$  receptors and growth inhibitory responses in retinoblastoma cells. *Science* 240: 196–199, 1988.109. Kitchin, F. D.; Ellsworth, R. M.: Pleiotropic effects of the gene for retinoblastoma. *J. Med. Genet.* 11: 244–246, 1974.110. Kivela, T.: Trilateral retinoblastoma: a meta-analysis of hereditary retinoblastoma associated with primary ectopic intracranial retinoblastoma. *J. Clin. Oncol.* 17: 1829–1837, 1999.111. Kivela, T.; Asko-Seljavaara, S.; Pihkala, U.; Hovi, L.; Heikkonen, J.: Sebaceous carcinoma of the eyelid associated with retinoblastoma. *Ophthalmology* 108: 1124–1128, 2001.112. Klutz, M.; Brockmann, D.; Lohmann, D. R.: A parent-of-origin effect in two families with retinoblastoma is associated with a distinct splice mutation in the RB1 gene. *Am. J. Hum. Genet.* 71: 174–179, 2002.113. Knight, L. A.; Gardner, H. A.; Gallie, B.

L.: Familial retinoblastoma: segregation of chromosome 13 in four families. *Am. J. Hum. Genet.* 32:194–201, 1980.

114. Knudson, A. G.: Hereditary cancer, oncogenes and anti-oncogenes. *Cancer Res.* 45: 1437–1443, 1985.

115. Knudson, A. G., Jr.: Mutation and cancer: statistical study of retinoblastoma. *Proc. Nat. Acad. Sci.* 68: 820–823, 1971.

116. Knudson, A. G., Jr.: Genetics of human cancer. *Annu. Rev. Genet.* 20:231–251, 1986.

117. Knudson, A. G., Jr.; Hethcote, H. W.; Brown, B. W.: Mutation and childhood cancer: a probabilistic model for the incidence of retinoblastoma. *Proc. Nat. Acad. Sci.* 72: 5116–5120, 1975.

118. Knudson, A. G., Jr.; Meadows, A. T.; Nichols, W. W.; Hill, R.: Chromosomal deletion and retinoblastoma. *New Eng. J. Med.* 295:1120–1123, 1976.

119. Laquis, S. J.; Rodriguez-Galindo, C.; Wilson, M. W.; Fleming, J. C.; Haik, B. G.: Retinoblastoma in a patient with an X;13 translocation and facial abnormalities consistent with 13q-syndrome. *Am. J. Ophthalm.* 133:285–287, 2002.

120. Lee, W.-H.; Bookstein, R.; Hong, F.; Young, L.-J.; Shew, J.-Y.; Lee, E. Y.-H. P.: Human retinoblastoma susceptibility gene: cloning, identification, and sequence. *Science* 235: 1394–1399, 1987.

121. Lee, W.-H.; Shew, J.-Y.; Hong, F. D.; Sery, T. W.; Donoso, L. A.; Young, L.-J.;

Bookstein, R.; Lee, E. Y.-H. P.: The retinoblastoma susceptibility gene encodes a nuclear phosphoprotein associated with DNA binding activity. *Nature* 329: 642–645, 1987.122. Lele, K. P.; Penrose, L. S.; Stallard, H. B.: Chromosome deletion in a case of retinoblastoma. *Ann. Hum. Genet.* 27: 171–174, 1963.123. Lemieux, N.; Messier, P. E.; Jacob, J. L.; Milot, J.; Richer, C. L.: Precise cytogenetic localization of the Rb locus at subband 13q14.11 by ultrastructural detection after immunochemical chromosome banding. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A27, 1989.124. Liu, Z.; Song, Y.; Bia, B.; Cowell, J. K.: Germline mutations in the RB1 gene in patients with hereditary retinoblastoma. *Genes Chromosomes Cancer* 14: 277–284, 1995.125. Lohmann, D. R.; Brandt, B.; Hopping, W.; Passarge, E.; Horsthemke, B.: Spectrum of small length germline mutations in the RB1 gene. *Hum. Molec. Genet.* 3: 2187–2193, 1994.126. Lohmann, D. R.; Brandt, B.; Hopping, W.; Passarge, E.; Horsthemke, B.: The spectrum of RB1 germ-line mutations in hereditary retinoblastoma. *Am. J. Hum. Genet.* 58: 940–949, 1996.127. Lohmann, D. R.; Gerick, M.; Brandt, B.; Oelschlager, U.; Lorenz, B.; Passarge, E.; Horsthemke, B.: Constitutional RB1-gene mutations in patients with isolated unilateral retinoblastoma.

Am. J. Hum. Genet. 61: 282–294, 1997.128. Lomazzi, M.; Moroni, M. C.; Jensen, M. R.; Frittoli, E.; Helin, K.: Suppression of the p53- or pRB-mediated G1 checkpoint is required for E2F-induced S-phase entry. Nature Genet 31: 190–194, 2002.129. Lueder, G. T.; Judisch, G. F.; Wen, B.-C.: Heritable retinoblastoma and pinealoma. Arch. Ophthalmol. 109: 1707–1709, 1991.130. Luo, R. X.; Postigo, A. A.; Dean, D. C.: Rb interacts with histone deacetylase to repress transcription. Cell 92: 463–473, 1998.131. Maat-Kievit, J. A.; Oepkes, D.; Hartwig, N. G.; Vermeij-Keers, C.; van Kamp, I. L.; van de Kamp, J. J. P.: A large retinoblastoma detected in a fetus at 21 weeks of gestation. Prenatal Diag. 13: 377–384, 1993.132. MacKay, C. J.; Abramson, D. H.; Ellsworth, R. M.: Metastatic patterns of retinoblastoma. Arch. Ophthalmol. 102: 391–396, 1984.133. Macklin, M. T.: A study of retinoblastoma in Ohio. Am. J. Hum. Genet. 12: 1–43, 1960.134. Macklin, M. T.: Inheritance of retinoblastoma in Ohio. Arch. Ophthalmol. 62: 842–851, 1959.135. Manchester, P. T., Jr.: Retinoblastoma among offspring of adult survivors. Arch. Ophthalmol. 65: 546–549, 1961.136. Mancini, D.; Singh, S.; Ainsworth, P.; Rodenhiser, D.: Constitutively methylated CpG dinucleotides as mutation hot spots in the retinoblastoma gene (RB1). Am. J. Hum. Genet.

61: 80–87, 1997.137. Mancini, M. A.; Shan, B.; Nickerson, J. A.; Penman, S.; Lee, W.-H.: The retinoblastoma gene product is a cell cycle-dependent, nuclear matrix-associated protein. *Proc. Nat. Acad. Sci.* 91: 418–422, 1994.138. Marino, S.; Vooijs, M.; van der Gulden, H.; Jonker, J.; Berns, A.: Induction of medulloblastomas in p53-null mutant mice by somatic inactivation of Rb in the external granular layer cells of the cerebellum. *Genes Dev.* 14: 994–1004, 2000.139. Matsunaga, E.: Recurrence risks to relatives of patients with retinoblastoma. *Jpn. J. Ophthalm.* 22: 313–319, 1978.140. Matsunaga, E.: Almost synchronous appearance of bilateral retinoblastomas. (Letter) *Am. J. Med. Genet.* 11: 485–487, 1982.141. Matsunaga, E.: Retinoblastoma: mutational mosaicism or host resistance? *Am. J. Med. Genet.* 8: 375–387, 1981.142. Matsunaga, E.: Hereditary retinoblastoma: host resistance and second primary tumors. *J. Nat. Cancer Inst.* 65: 47–51, 1980.143. Matsunaga, E.: Hereditary retinoblastoma: delayed mutation or host resistance? *Am. J. Hum. Genet.* 30: 406–425, 1978.144. Matsunaga, E.; Minoda, K.; Sasaki, M. S.: Parental age and seasonal variation in the births of children with sporadic retinoblastoma: a mutation-epidemiologic study. *Hum. Genet.* 84: 155–158, 1990.145.

Michalova, K.; Kloucek, F.; Musilova, J.: Deletion of 13q  
 in two patients with retinoblastoma, one probably due to  
 13q- mosaicism in the mother. Hum. Genet. 61: 264-266,  
 1982.146. Moll, A. C.; Imhof, S. M.; Schouten-Van  
 Meeteren, A. Y. N.; Kuik, D. J.; Hofman, P.; Boers, M.: Sec-  
 ond primary tumors in hereditary retinoblastoma: a regis-  
 ter-based study, 1945-1997. Is there an age effect on ra-  
 diation-related risk? Ophthalmology 108:  
 1109-1114, 2001.147. Motegi, T.: Lymphocyte chromo-  
 some survey in 42 patients with retinoblastoma: effort to  
 detect 13q14 deletion mosaicism. Hum. Genet.  
 58:168-173, 1981.148. Motegi, T.: High rate of detection  
 of 13q14 deletion mosaicism among retinoblastoma pa-  
 tients (using more extensive methods). Hum. Genet. 61:  
 95-97, 1982.149. Motegi, T.; Kaga, M.; Yanagawa, Y.;  
 Kadowaki, H.; Watanabe, K.; Inoue, A.; Komatsu, M.; Min-  
 oda, K.: A recognizable pattern of the midface of  
 retinoblastoma patients with interstitial deletion of 13q.  
 Hum. Genet. 64: 160-162, 1983.150. Motegi, T.; Komatsu,  
 M.; Minoda, K.: Is the interstitial deletion of 13q in  
 retinoblastoma patients not transmissible? (Letter)  
 Hum. Genet. 64: 205, 1983.151. Motegi, T.; Komatsu, M.;  
 Nakazato, Y.; Ohuchi, M.; Minoda, K.: Retinoblastoma in a



boy with a de novo mutation of a 13/18 translocation: the assumption that the retinoblastoma locus is at 13q141, particularly at the distal portion of it. Hum. Genet. 60: 193–195, 1982.152. Munier, F.; Spence, M. A.; Pescia, G.; Balmer, A.; Gailloud, C.; Thonney, F.; van Melle, G.; Rutz, H. P.: Paternal selection favoring mutant alleles of the retinoblastoma susceptibility gene. Hum. Genet. 89:508–512, 1992.153. Munier, F. L.; Wang, M. X.; Spence, M. A.; Thonney, F.; Balmer, A.; Pescia, G.; Donoso, L. A.; Murphree, A. L.: Pseudo low penetrance in retinoblastoma: fortuitous familial aggregation of sporadic cases caused by independently derived mutations in two large pedigrees. Arch. Ophthalmol. 111: 1507–1511, 1993.154. Murphree, A. L.; Benedict, W. F.: Retinoblastoma: clues to human oncogenesis. Science 223: 1028–1033, 1984.155. Naumova, A.; Hansen, M.; Strong, L.; Jones, P. A.; Hadjistilianou, D.; Mastrangelo, D.; Griegel, S.; Rajewsky, M. F.; Shields, J.; Donoso, L.; Wang, M.; Sapienza, C.: Concordance between parental origin of chromosome 13q loss and chromosome 6p duplication in sporadic retinoblastoma. Am. J. Hum. Genet. 54: 274–281, 1994.156. Naumova, A.; Sapienza, C.: The genetics of retinoblastoma, revisited. Am. J. Hum. Genet. 54:

264–273, 1994.157. Nevins, J. R.: The Rb/E2F pathway and cancer. *Hum. Molec. Genet.* 10:699–703, 2001.158. Nichols, W. W.; Miller, R. C.; Sobel, M.; Hoffman, E.; Sparkes, R. S.; Mohandas, T.; Veomett, I.; Davis, J. R.: Further observations on a 13qXp translocation associated with retinoblastoma. *Am. J. Ophthalmol.* 89:621–627, 1980.159. Nielsen, S. J.; Schneider, R.; Bauer, U.–M.; Bannister, A. J.; Morrison, A.; O'Carroll, D.; Firestein, R.; Cleary, M.; Jenuwein, T.; Herrera, R. E.; Kouzarides, T.: Rb targets histone H3 methylation and HP1 to promoters. *Nature* 412: 561–565, 2001.160. Nirankari, M. S.; Gulati, G. C.; Chaddah, M. R.: Retinoblastoma: genetics and report of a family. *Am. J. Ophthalmol.* 53: 523–532, 1962.161. Noorani, H. Z.; Khan, H. N.; Gallie, B. L.; Detsky, A. S.: Cost comparison of molecular versus conventional screening of relatives at risk for retinoblastoma. *Am. J. Hum. Genet.* 59: 301–307, 1996.162. Nussbaum, R.; Puck, J.: Recurrence risks for retinoblastoma: a model for autosomal dominant disorders with complex inheritance. *J. Pediatr. Ophthalmol.* 13: 89–98, 1976.163. Onadim, Z.; Hogg, A.; Baird, P. N.; Cowell, J. K.: Oncogenic point mutations in exon 20 of the RB1 gene in families showing incomplete penetrance and mild expression of the retinoblastoma phenotype. *Proc. Nat. Acad. Sci.*

89: 6177–6181, 1992.164. Onadim, Z.; Woolford, A. J.; Kingston, J. E.; Hungerford, J.L.: The RB1 gene mutation in a child with ectopic intracranial retinoblastoma. *Brit.J. Cancer* 76: 1405–1409, 1997.165. Ono, T.; Yoshida, M. C.: Chromosomal assignment of retinoblastoma1 gene (RB1) to mouse 14D3 and rat 15q12 by fluorescence in situ hybridization. *Jpn.J. Genet.* 68: 617–621, 1993.166. Orye, E.; Benoit, Y.; Coppieters, R.; Jeannin, P.; Ver-cruysse,C.; Delaey, J.; Delbeke, M.–J.: A case of retinoblas-toma, associatedwith histiocytosis–X and mosaicism of a deleted D–group chromosome(13q14–q31). *Clin. Genet.* 22: 37–39, 1982.167. Orye, E.; Delbeke, M. J.; Vanden-abeele, B.: Retinoblastoma andlong arm deletion of chro-mosome 13. Attempts to define the deletedsegment. *Clin. Genet.* 5: 457–464, 1974.168. Orye, E.; Delbeke, M. J.; Vandenabeele, B.: Retinoblastoma andD–chromosome deletions. (Letter) *Lancet* II: 1376, 1971.169. Otterson, G. A.; Modi, S.; Nguyen, K.; Coxon, A. B.; Kaye, F.J.: Temper-ature–sensitive RB mutations linked to incomplete pene-tranceof familial retinoblastoma in 12 families. *Am. J. Hum. Genet.* 65:1040–1046, 1999.170. Otterson, G. W.; Chen, W.; Coxon, A. B.; Khleif, S. N.; Kaye,F. J.: Incomplete penetrance of familial retinoblastoma linked togerm–line

mutations that result in partial loss of RB function.

Proc.Nat. Acad. Sci. 94: 12036–12040, 1997.171. Pendergrass, T. W.; Davis, S.: Incidence of retinoblastoma in the United States. Arch. Ophthal. 98: 1204–1210, 1980.172. Pennaneach, V.; Salles–Passador, I.; Munshi, A.; Brickner, H.; Regazzoni, K.; Dick, F.; Dyson, N.; Chen, T.–T.; Wang, J. Y. J.; Fotedar, R.; Fotedar, A.: The large subunit of replication factor C promotes cell survival after DNA damage in an LxCxE motif– and Rb–dependent manner. Molec. Cell 7: 715–727, 2001.173. Riccardi, V. M.; Hittner, H. M.; Francke, U.; Pippin, S.; Holmquist, G. P.; Kretzer, F. L.; Ferrell, R.: Partial triplication and deletion of 13q: study of a family presenting with bilateral retinoblastomas. Clin.Genet. 15: 332–345, 1979.174. Rivera, H.; Turleau, C.; de Grouchy, J.; Junien, C.; Despoisse, S.; Zucker, J.–M.: Retinoblastoma–del(13q14): report of two patients, one with a trisomic sib due to maternal insertion; gene–dosage effect for esterase D. Hum. Genet. 59: 211–214, 1981.175. Sakai, T.; Ohtani, N.; McGee, T. L.; Robbins, P. D.; Dryja, T.P.: Oncogenic germ–line mutations in Sp1 and ATF sites in the human retinoblastoma gene. Nature 353: 83–86, 1991.176. Sakai, T.; Ohtani, N.; McGee, T. L.; Robbins, P. D.; Dryja, T.P.: Oncogenic germ–line mutations in

Sp1 and ATF sites in the human retinoblastoma gene. *Nature* 353: 83–86, 1991.177. Sakai, T.; Toguchida, J.; Ohtani, N.; Yandell, D. W.; Rapaport, J. M.; Dryja, T. P.: Allele-specific hypermethylation of the retinoblastoma tumor-suppressor gene. *Am. J. Hum. Genet.* 48: 880–888, 1991.178. Schappert-Kimmijser, J.; Hemmes, G. D.; Nijland, R.: The heredity of retinoblastoma. *Ophthalmologica* 151: 197–213, 1966.179. Scheffer, H.; te Meerman, G. J.; Kruize, Y. C. M.; van den Berg, A. H. M.; Penninga, D. P.; Tan, K. E. W. P.; der Kinderen, D. J.; Buys, C. H. C. M.: Linkage analysis of families with hereditary retinoblastoma: nonpenetrance of mutation, revealed by combined use of markers within and flanking the RB1 gene. *Am. J. Hum. Genet.* 45: 252–260, 1989.180. Schimke, R. N.; Lowman, J.; Cowan, G.: Retinoblastoma and osteogenic sarcoma in sibs. *Cancer* 34: 2077–2079, 1974.181. Schubert, E. L.; Strong, L. C.; Hansen, M. F.: A splicing mutation in RB1 in low penetrance retinoblastoma. *Hum. Genet.* 100: 557–563, 1997.182. Shields, C. L.; Honavar, S.; Shields, J. A.; Demirci, H.; Meadows, A. T.: Vitrectomy in eyes with unsuspected retinoblastoma. *Ophthalmology* 107: 2250–2255, 2000.183. Shio, Y.; Yamamoto, T.; Yamaguchi, N.: Negative regulation of Rb expression by the

p53 gene product. *Proc. Nat. Acad. Sci.* 89:5206–5210, 1992.184. Shroeder, W. T.; Chao, L.-Y.; Dao, D. D.; Strong, L. C.; Pathak, S.; Riccardi, V.; Lewis, W. H.; Saunders, G. F.: Nonrandom loss of maternal chromosome 11 alleles in Wilms tumors. *Am. J. Hum. Genet.* 40:413–420, 1987.185. Sippel, K. C.; Fraioli, R. E.; Smith, G. D.; Schalkoff, M. E.; Sutherland, J.; Gallie, B. L.; Dryja, T. P.: Frequency of somatic and germ-line mosaicism in retinoblastoma: implications for genetic counseling. *Am. J. Hum. Genet.* 62: 610–619, 1998.186. Smith, S. M.; Sorsby, A.: Retinoblastoma: some genetic aspects. *Ann. Hum. Genet.* 23: 50–58, 1958.187. Sparkes, R. S.: The genetics of retinoblastoma. *Biochim. Biophys. Acta* 780: 95–118, 1985.188. Sparkes, R. S.; Muller, H.; Klisak, I.; Abram, J. A.: Retinoblastoma with 13q; chromosomal deletion associated with maternal paracentric inversion of 13q. *Science* 203: 1027–1029, 1979.189. Sparkes, R. S.; Murphree, A. L.; Lingua, R. W.; Sparkes, M. C.; Field, L. L.; Funderburk, S. J.; Benedict, W. F.: Gene for hereditary retinoblastoma assigned to human chromosome 13 by linkage to esterase D. *Science* 219: 971–973, 1983.190. Sparkes, R. S.; Sparkes, M. C.; Wilson, M. G.; Towner, J. W.; Benedict, W.; Murphree, A. L.; Yunis, J. J.: Regional as-

signment of genes for human esterase D and retinoblastoma to chromosome band 13q14. *Science* 208: 1042–1044, 1980.191. Sparkes, R. S.; Sparkes, M. C.; Wilson, M. G.; Towner, J. W.; Benedict, W.; Murphree, A. L.; Yunis, J. J.: Regional assignment of genes for human esterase D and retinoblastoma to chromosome band 13q14. (Abstract) *Cytogenet. Cell Genet.* 25: 209, 1979.192. Squire, J.; Gallie, B. L.; Phillips, R. A.: A detailed analysis of chromosomal changes in heritable and non-heritable retinoblastoma. *Hum. Genet.* 70: 291–301, 1985.193. Squire, J.; Phillips, R. A.; Boyce, S.; Godbout, R.; Rogers, B.; Gallie, B. L.: Isochromosome 6p, a unique chromosomal abnormality in retinoblastoma: verification by standard staining techniques, new densitometric methods, and somatic cell hybridization. *Hum. Genet.* 66:46–53, 1984.194. Stallard, H. B.: The conservative treatment of retinoblastoma. *Trans. Ophthalm. Soc.* 82: 473, 1962.195. Stone, J. C.; Crosby, J. L.; Kozak, C. A.; Schievella, A. R.; Bernards, R.; Nadeau, J. H.: The murine retinoblastoma homolog maps to chromosome 14 near Es-10. *Genomics* 5: 70–75, 1989.196. Strong, L. C.; Riccardi, V. M.; Ferrell, R. E.; Sparkes, R. S.: Familial retinoblastoma and chromosome 13 deletion transmitted via an insertional translocation.

tion. Science 213: 1501–1503, 1981.197. Taylor, A. I.: Dq-, Dr and retinoblastoma. Humangenetik 10:209–217, 1970.198. Thomas, D. M.; Carty, S. A.; Piscopo, D. M.; Lee, J.-S.; Wang, W.-F.; Forrester, W. C.; Hinds, P. W.: The retinoblastoma protein acts as a transcriptional coactivator required for osteogenic differentiation. Molec. Cell 8: 303–316, 2001.199. Toguchida, J.; Ishizaki, K.; Sasaki, M. S.; Nakamura, Y.; Ikenaga, M.; Kato, M.; Sugimoto, M.; Koutoura, Y.; Yamamuro, T.: Preferential mutation of paternally derived RB gene as the initial event in sporadic osteosarcoma. Nature 338: 156–158, 1989.200. Toguchida, J.; McGee, T. L.; Paterson, J. C.; Eagle, J. R.; Tucker, S.; Yandell, D. W.; Dryja, T. P.: Complete genomic sequence of the human retinoblastoma susceptibility gene. Genomics 17: 535–543, 1993.201. Turleau, C.; de Grouchy, J.; Chavin-Colin, F.; Despoisses, S.; Leblanc, A.: Two cases of del(13q)-retinoblastoma and two cases of partial trisomy due to a familial insertion. Ann. Genet. 26: 158–160, 1983.202. Turleau, C.; de Grouchy, J.; Chavin-Colin, F.; Junien, C.; Seger, J.; Schlienger, P.; Leblanc, A.; Haye, C.: Cytogenetic forms of retinoblastoma: their incidence in a survey of 66 patients. Cancer Genet. Cytogenet. 16:321–334, 1985.203. Verma, R. S.; Ramesh, K. H.; Sa-



monte, R. V.; Conte, R. A.: Mapping the homolog of the human Rb1 gene to chromosome 14 of higher primates. *Mammalian Genome* 7: 591–592, 1996.204. Vogel, F.: Genetics of retinoblastoma. *Modern Trends in Ophthalmology*. (pub.) 1968.205. Vogel, F.: Genetics of retinoblastoma. In: *Genetic Counseling*. Heidelberg University, Science Library. Trans. by Sabine Kurth. New York: Springer Verlag (pub.) 1969.206. Vogel, F.: The genetics of retinoblastoma. *Hum. Genet.* 52:1–54, 1979.207. Warburg, M.: Retinoblastoma. In: Goldberg, M. F.: *Genetic and Metabolic Eye Disease*. Boston: Little, Brown and Co. (pub.) 1974. Pp. 447–461.208. Weichselbaum, R. R.; Beckett, M.; Diamond, A.: Some retinoblastomas, osteosarcomas, and soft tissue sarcomas may share a common etiology. *Proc. Nat. Acad. Sci.* 85: 2106–2109, 1988.209. Weichselbaum, R. R.; Nove, J.; Little, J. B.: Fibroblasts from a deletion type retinoblastoma patient are abnormally x-ray sensitive. *Nature* 266:726–727, 1977.210. Weinberg, R. A.: The retinoblastoma protein and cell cycle control. *Cell* 81:323–330, 1995.211. Whyte, P.; Buchkovich, K. J.; Horowitz, J. M.; Friend, S. H.; Raybuck, M.; Weinberg, R. A.; Harlow, E.: Association between an oncogene and an anti-oncogene: the adenovirus E1A proteins bind to the

retinoblastoma gene product. *Nature* 334: 124–129, 1988.212. Wiggs, J.; Nordenskjold, M.; Yandell, D.; Rapaport, J.; Grondin, V.; Janson, M.; Werelius, B.; Petersen, R.; Craft, A.; Riedel, K.; Liberfarb, R.; Walton, D.; Wilson, W.; Dryja, T. P.: Prediction of the risk of hereditary retinoblastoma, using DNA polymorphisms within the retinoblastoma gene. *New Eng. J. Med.* 318: 151–157, 1988.213. Wilson, M. G.; Ebbin, A. J.; Towner, J. W.; Spencer, W. H.: Chromosomal anomalies in patients with retinoblastoma. *Clin. Genet.* 12:1–8, 1977.214. Wilson, M. G.; Melnyk, J.; Towner, J. W. J.: Retinoblastoma and deletion D(14) syndrome. *J. Med. Genet.* 6: 322–327, 1969.215. Wilson, M. G.; Towner, J. W.; Fujimoto, A.: Retinoblastoma and D-chromosome deletions. *Am. J. Hum. Genet.* 25: 57–61, 1973.216. Windle, J. J.; Albert, D. M.; O'Brien, J. M.; Marcus, D. M.; Distecche, C. M.; Bernards, R.; Mellon, P. L.: Retinoblastoma in transgenic mice. *Nature* 343: 665–669, 1990.217. Yandell, D. W.; Campbell, T. A.; Dayton, S. H.; Petersen, R.; Walton, D.; Little, J. B.; McConkie-Rosell, A.; Buckley, E.; Dryja, T.: Oncogenic point mutations in the human retinoblastoma gene: their application to genetic counseling. *New Eng. J. Med.* 321: 1689–1695, 1989.218. Yokota, J.; Akiyama, T.; Fung, Y.-K. T.; Benedict, W. F.;

Namba,Y.; Hanaoka, M.; Wada, M.; Terasaki, T.; Shimamoto, Y.; Sugimura,T.; Terada, M.: Altered expression of the retinoblastoma (RB) gene in small-cell carcinoma of the lung. *Oncogene* 3: 471–475, 1988.219. Zeschnigk, M.; Lohmann, D.; Horsthemke, B.: A PCR test for the detection of hypermethylated alleles at the retinoblastoma locus. *J.Med. Genet.* 36: 793–794, 1999.220. Zhang, H. S.; Postigo, A. A.; Dean, D. C.: Active transcriptional repression by the Rb–E2F complex mediates G1 arrest triggered by p16(INK4a), TGF–beta, and contact inhibition. *Cell* 97: 53–61, 1999.221. Zhu, X.; Dunn, J. M.; Phillips, R. A.; Goddard, A. D.; Paton, K. E.; Becker, A.; Gallie, B. L.: Preferential germline mutation of the paternal allele in retinoblastoma. *Nature* 340: 312–313, 1989.

[70621] 8951. Baens, M.; Aerssens, J.; Van Zand, K.; Van den Berghe, H.; Marynen, P.: Isolation and regional assignment of human chromosome 12p cDNAs. *Genomics* 29:44–52, 1995.

[70622] 8952. Mao, S.; Neale, G. A. M.; Goorha, R. M.: T-cell oncogene rhombotin-2 interacts with retinoblastoma-binding protein 2. *Oncogene* 14: 1531–1539, 1997.

[70623] 8953. Baldi, A.; Boccia, V.; Claudio, P. P.; De Luca, A.; Giordano, A.: Genomic structure of the human retinoblas-

toma-related Rb2/p130 gene. Proc.Nat. Acad. Sci. 93: 4629–4632, 1996.

- [70624] 8954. Mayol, X.; Grana, X.; Baldi, A.; Sang, N.; Hu, Q.; Giordano, A.: Cloning of a new member of the retinoblastoma gene family (pRb2) which binds to the E1A transforming domain. *Oncogene* 8: 2561–2566, 1993.
- [70625] 8955. Yeung, R. S.; Bell, D. W.; Testa, J. R.; Mayol, X.; Baldi, A.; Grana, X.; Klinga-Levan, K.; Knudson, A. G.; Giordano, A.: The retinoblastoma-related gene, RB2, maps to human chromosome 16q12 and rat chromosome 19. *Oncogene* 8: 3465–3468, 1993.
- [70626] 8956. Benbrook, D.; Lernhardt, E.; Pfahl, M.: A new retinoic acid receptor identified from a hepatocellular carcinoma. (Letter) *Nature* 333: 669–672, 1988.
- [70627] 8957. Brand, N.; Petkovich, M.; Krust, A.; Chambon, P.; de The, H.; Marchio, A.; Tiollais, P.; Dejean, A.: Identification of a second human retinoic acid receptor. (Letter) *Nature* 332: 850–853, 1988.
- [70628] 8958. Dejean, A.; Bougueleret, L.; Grzeschik, K.-H.; Tiollais, P.: Hepatitis B virus DNA integration in a sequence homologous to v-erb-A and steroid receptor genes in a hepatocellular carcinoma. *Nature* 322: 70–72, 1986.
- [70629] 8959. de The, H.; del Mar Vivanco-Ruiz, M.; Tiollais, P.;

Stunnenberg,H.; Dejean, A.: Identification of a retinoic acid responsive element in the retinoic acid receptor beta gene. *Nature* 343: 177–180, 1990.

- [70630] 8960.Aruffo, A.; Seed, B.: Molecular cloning of a CD28 cDNA by a high-efficiency COS cell expression system. *Proc. Nat. Acad. Sci.* 84: 8573–8577, 1987.
- [70631] 8961.Lafage-Pochitaloff, M.; Costello, R.; Couez, D.; Simonetti, J.; Mannoni, P.; Mawas, C.; Olive, D.: Human CD28 and CTLA-4 Ig superfamily genes are located on chromosome 2 at bands q33–q34. *Immunogenetics* 31:198–201, 1990.
- [70632] 8962.Lee, K. P.; Taylor, C.; Petryniak, B.; Turka, L. A.; June, C. H.; Thompson, C. B.: The genomic organization of the CD28 gene: implications for the regulation of CD28 mRNA expression and heterogeneity. *J.Immun.* 145: 344–352, 1990.
- [70633] 8963.Lessleruer, W.; Gmunder, H.; Bohlen, P.: Purification and N-terminal amino acid sequence of the human T90/44 (CD28) antigen. *Immunogenetics* 27:388–391, 1988.
- [70634] 8964.Okkenhaug, K.; Wu, L.; Garza, K. M.; La Rose, J.; Khoo, W.; Odermatt, B.; Mak, T. W.; Ohashi, P. S.; Rottapel, R.: A point mutation in CD28 distinguishes proliferative signals from survival signals. *Nature Immun.* 2: 325–332,

2001.

- [70635] 8965. Dear, T. N.; Colledge, W. H.; Carlton, M. B. L.; Lavenir, I.; Larson, T.; Smith, A. J. H.; Warren, A. J.; Evans, M. J.; Sofroniew, M. V.; Rabbitts, T. H.: The Hox11 gene is essential for cell survival during spleen development. *Development* 121: 2909–2915, 1995.
- [70636] 8966. Dear, T. N.; Sanchez-Garcia, I.; Rabbitts, T. H.: The HOX11 gene encodes a DNA-binding nuclear transcription factor belonging to a distinct family of homeobox genes. *Proc. Nat. Acad. Sci.* 90: 4431–4435, 1993.
- [70637] 8967. Dube, I. D.; Kamel-Reid, S.; Yuan, C. C.; Lu, M.; Wu, X.; Corpus, G.; Raimondi, S. C.; Crist, W. M.; Carroll, A. J.; Minowada, J.; Baker, J. B.: A novel human homeobox gene lies at the chromosome 10 breakpoint in lymphoid neoplasias with chromosomal translocation t(10;14). *Blood* 78:2996–3003, 1991.
- [70638] 8968. Hatano, M.; Roberts, C. W. M.; Minden, M.; Crist, W. M.; Korsmeyer, S. J.: Deregulation of a homeobox gene, HOX11, by the t(10;14) in T cell leukemia. *Science* 253: 79–82, 1991.
- [70639] 8969. Kagan, J.; Finan, J.; Letofsky, J.; Besa, E. C.; Nowell, P. C.; Croce, C. M.: Alpha-chain locus of the T-cell antigen receptor is involved in the t(10;14) chromosome translocation.

tion of T-cell acutelymphocytic leukemia. Proc. Nat. Acad. Sci. 84: 4543–4546, 1987.

[70640] 8970.Kennedy, M. A.; Gonzalez–Sarmiento, R.; Kees, U. R.; Lampert, F.;Dear, N.; Boehm, T.; Rabbitts, T. H.: HOX11, a homeobox–containingT–cell oncogene on human chromosome 10q24. Proc. Nat. Acad. Sci. 88:8900–8904, 1991.

[70641] 8971.Lu, M.; Gong, Z. Y.; Shen, W. F.; Ho, A. D.: The TCL–3 proto–oncogenealtered by chromosomal translocation in T–cell leukemia codes fora homeobox protein. EMBO J. 10: 2905–2910, 1991.

[70642] 8972.Roberts, C. W. M.; Shutter, J. R.; Korsmeyer, S. J.: Hox11 controlsthe genesis of the spleen. Nature 368: 747–750, 1994.

[70643] 8973.Zutter, M.; Hockett, R. D.; Roberts, C. W. M.; McGuire, E. A.;Bloomstone, J.; Morton, C. C.; Deaven, L. L.; Crist, W. M.; Carroll,A. J.; Korsmeyer, S. J.: The t(10;14)(q24;q11) of T–cell acute lymphoblasticleukemia juxtaposes the delta T–cell receptor with TCL3, a conservedand activated locus at 10q24. Proc. Nat. Acad. Sci. 87: 3161–3165,1990.

[70644] 8974.Alarcon, B.; Regueiro, J. R.; Arnaiz–Villena, A.; Terhorst, C.: Familial defect in the surface expression of the

T-cell receptor-CD3complex. New Eng. J. Med. 319:  
1203-1208, 1988.

[70645] 8975.Caplan, S.; Zeligler, S.; Wang, L.; Baniyash, M.: Cell-  
surface-expressedT-cell antigen-receptor epsilon chain is  
associated with the cytoskeleton. Proc.Nat. Acad. Sci. 92:  
4768-4772, 1995.

[70646] 8976.Clevers, H.; Alarcon, B.; Wileman, T.; Terhorst, C.:  
The T cellreceptor/CD3 complex: a dynamic protein en-  
semble. Annu. Rev. Immun. 6:629-662, 1988.

[70647] 8977.Grakoui, A.; Bromley, S. K.; Sumen, C.; Davis, M. M.;  
Shaw, A.S.; Allen, P. M.; Dustin, M. L.: The immunological  
synapse: a molecularmachine controlling T cell activation.  
Science 285: 221-227, 1999.

[70648] 8978.Krummel, M. F.; Sjaastad, M. D.; Wulfing, C.; Davis,  
M. M.: Differentialclustering of CD4 and CD3-zeta during  
T cell recognition. Science 289:1349-1352, 2000.

[70649] 8979.Weissman, A. M.; Baniyash, M.; Hou, D.; Samelson, L.  
E.; Burgess,W. H.; Klausner, R. D.: Molecular cloning of the  
zeta chain of theT cell antigen receptor. Science 239:  
1018-1021, 1988.

[70650] 8980.Weissman, A. M.; Hou, D.; Orloff, D. G.; Modi, W. S.;  
Seuanez,H.; O'Brien, S. J.; Klausner, R. D.: Molecular  
cloning and chromosomallocalization of the human T-cell



receptor zeta chain: distinction from the molecular CD3 complex. *Proc. Nat. Acad. Sci.* 85: 9709–9713, 1988.

- [70651] 8981. Weissman, A. M.; Samelson, L. E.; Klausner, R. D.: A new subunit of the human T-cell antigen receptor complex. *Nature* 324: 480–482, 1986.
- [70652] 8982. Zeng, Z.-H.; Castano, A. R.; Segelke, B. W.; Stura, E. A.; Peterson, P. A.; Wilson, I. A.: Crystal structure of mouse CD1: an MHC-like fold with a large hydrophobic binding groove. *Science* 277: 339–345, 1997.
- [70653] 8983. Audhya, T.; Schlesinger, D. H.; Goldstein, G.: Isolation and complete amino acid sequence of human thymopoietin and splenin. *Proc. Nat. Acad. Sci.* 84: 3545–3549, 1987.
- [70654] 8984. Berger, R.; Theodor, L.; Shoham, J.; Gokkel, E.; Brok-Simoni, F.; Avraham, K. B.; Copeland, N. G.; Jenkins, N. A.; Rechavi, G.; Simon, A. J.: The characterization and localization of the mouse thymopoietin/lamina-associated polypeptide 2 gene and its alternatively spliced products. *Genome Res.* 6: 361–370, 1996.
- [70655] 8985. Harris, C. A.; Andryuk, P. J.; Cline, S. W.; Mathew, S.; Siekierka, J. J.; Goldstein, G.: Structure and mapping of the human thymopoietin (TMPO) gene and relationship of human TMPO beta to rat lamin-associated polypeptide 2. *Ge-*

nomics 28: 198–205, 1995.

[70656] 8986. Eschenfeldt, W. H.; Berger, S. L.: The human prothymosin alpha gene is polymorphic and induced upon growth stimulation: evidence using a cloned cDNA. *Proc. Nat. Acad. Sci.* 83: 9403–9407, 1986.

[70657] 8987. Goodall, G. J.; Dominguez, F.; Horecker, B. L.: Molecular cloning of cDNA for human prothymosin alpha. *Proc. Nat. Acad. Sci.* 83: 8926–8928, 1986.

[70658] 8988. Haritos, A. A.; Goodall, G. J.; Horecker, B. L.: Prothymosin alpha: isolation and properties of the major immunoreactive form of thymosin alpha-1 in rat thymus. *Proc. Nat. Acad. Sci.* 81: 1008–1011, 1984.

[70659] 8989. Manrow, R. E.; Leone, A.; Krug, M. S.; Eschenfeldt, W. H.; Berger, S. L.: The human prothymosin alpha gene family contains several processed pseudogenes lacking deleterious lesions. *Genomics* 13: 319–331, 1992.

[70660] 8990. Szabo, P.; Panneerselvam, C.; Clinton, M.; Frangou-Lazaridis, M.; Weksler, D.; Whittington, E.; Macera, M. J.; Grzeschik, K.-H.; Selvakumar, A.; Horecker, B. L.: Prothymosin alpha-gene in humans: organization of its promoter region and localization to chromosome 2. *Hum. Genet.* 90: 629–634, 1993.

[70661] 8991. Demczuk, S.; Aledo, R.; Zucman, J.; Delattre, O.;

Desmaze, C.; Dauphinot, L.; Jalbert, P.; Rouleau, G. A.; Thomas, G.; Aurias, A.: Cloning of a balanced translocation breakpoint in the DiGeorge syndrome critical region and isolation of a novel potential adhesion receptor gene in its vicinity. *Hum. Molec. Genet.* 4: 551–558, 1995.

[70662] 8992. Jourdan-Le Saux, C.; Tomsche, A.; Ujfalusi, A.; Jia, L.; Csiszar, K.: Central nervous system, uterus, heart, and leukocyte expression of the LOXL3 gene, encoding a novel lysyl oxidase-like protein. *Genomics* 74:211–218, 2001.

[70663] 8993. Maki, J. M.; Kivirikko, K. I.: Cloning and characterization of a fourth human lysyl oxidase isoenzyme. *Biochem. J.* 355: 381–387, 2001.

[70664] 8994. Jaju, R. J.; Haas, O. A.; Neat, M.; Harbott, J.; Saha, V.; Boulwood, J.; Brown, J. M.; Pirc-Danoewinata, H.; Krings, B. W.; Muller, U.; Morris, S. W.; Wainscoat, J. S.; Kearney, L.: A new recurrent translocation, t(5;11)(q35p15.5), associated with del(5q) in childhood acute myeloid leukemia. *Blood* 94: 773–780, 1999.

[70665] 8995. Kurotaki, N.; Harada, N.; Yoshiura, K.; Sugano, S.; Niikawa, N.; Matsumoto, N.: Molecular characterization of NSD1, a human homologue of the mouse Nsd1 gene. *Gene* 279: 197–204, 2001.

[70666] 8996. Wang, X.; Yeh, S.; Wu, G.; Hsu, C.-L.; Wang, L.; Chi-

ang, T.; Yang, Y.; Guo, Y.; Chang, C.: Identification and characterization of a novel androgen receptor coregulator ARA267- $\alpha$  in prostate cancer cells. *J. Biol. Chem.* 276: 40417–40423, 2001.

[70667] 8997. Riewald, M.; Petrovan, R. J.; Donner, A.; Mueller, B. M.; Ruf, W.: Activation of endothelial cell protease activated receptor 1 by the protein C pathway. *Science* 296: 1880–1882, 2002.

[70668] 8998. Jankowski, S. A.; De Jong, P.; Meltzer, P. S.: Genomic structure of SAS, a member of the transmembrane 4 superfamily amplified in human sarcomas. *Genomics* 25: 501–506, 1995.

[70669] 8999. Meltzer, P. S.; Jankowski, S. A.; Dal Cin, P.; Sandberg, A. A.; Paz, I. B.; Coccia, M. A.; Smith, S. H.: Identification and cloning of a novel amplified DNA sequence in human malignant fibrous histiocyte derived from a region of chromosome 12 frequently rearranged in soft tissue tumors. (Abstract) *Cytogenet. Cell Genet.* 58: 1979 only, 1991.

[70670] 9000. Peschle, C.; Mavilio, F.; Sposi, N. M.; Giampaolo, A.; Care, A.; Bottero, L.; Bruno, M.; Mastroberardino, G.; Gastaldi, R.; Testa, M. G.; Alimena, G.; Amadori, S.; Mandelli, F.: Translocation and rearrangement of c-myc into

immunoglobulin alpha heavy chain locus in primary cells from acute lymphocytic leukemia. *Proc. Nat. Acad. Sci.* 81: 5514–5518, 1984.

[70671] 9001. Shao, X.; Tarnasky, H. A.; Schalles, U.; Oko, R.; van der Hoorn, F. A.: Interactional cloning of the 84-kDa major outer dense fiber protein Odf84: leucine zippers mediate associations of Odf84 and Odf27. *J. Biol. Chem.* 272: 6105–6113, 1997.

[70672] 9002. Shao, X.; van der Hoorn, F. A.: Self-interaction of the major 27-kilodalton outer dense fiber protein is in part mediated by a leucine zipper domain in the rat. *Biol. Reprod.* 55: 1343–1350, 1996.

[70673] 9003. Ammer, H.; Henschen, A.; Lee, C.-H.: Isolation and amino-acid sequence analysis of human sperm protamines P1 and P2: occurrence of two forms of protamine P2. *Biol. Chem. Hoppe-Seyler* 367: 515–522, 1986.

[70674] 9004. Cho, C.; Willis, W. D.; Goulding, E. H.; Jung-Ha, H.; Choi, Y.-C.; Hecht, N. B.; Eddy, E. M.: Haploinsufficiency of protamine-1 or -2 causes infertility in mice. *Nature Genet.* 28: 82–86, 2001.

[70675] 9005. Choudhary, S. K.; Wykes, S. M.; Kramer, J. A.; Mohamed, A. N.; Koppitch, F.; Nelson, J. E.; Krawetz, S. A.: A haploid expressed gene cluster exists as a single chro-

matin domain in human sperm. J. Biol. Chem. 270:8755–8762, 1995.

- [70676] 9006.Domenjoud, L.; Nussbaum, G.; Adham, I. M.; Greeske, G.; Engel,W.: Genomic sequences of human protamines whose genes, PRM1 and PRM2,are clustered. Genomics 8: 127–133, 1990.
- [70677] 9007.Engel, W.: Personal Communication. Goettingen, Germany 5/17/1990.
- [70678] 9008.Hecht, N. B.; Kleene, K. C.; Yelick, P. C.; Johnson, P. A.; Pravtcheva,D. D.; Ruddle, F. H.: Mapping of haploid expressed genes: genes forboth mouse protamines are located on chromosome 16. Somat. Cell Molec.Genet. 12: 203–208, 1986.
- [70679] 9009.Krawetz, S. A.; Herfort, M. H.; Hamerton, J. L.; Pon, R. T.; Dixon,G. H.: Chromosomal localization and structure of the human P1 protaminegene. Genomics 5: 639–645, 1989.
- [70680] 9010.Reeves, R. H.; Gallahan, D.; O'Hara, B. F.; Callahan, R.; Gearhart,J. D.: Genetic mapping of Prm–1, Igl–1, Smst, Mtv–6, Sod–1, and Ets–2and localization of the Down syndrome region on mouse chromosome 16. Cytogenet.Cell Genet. 44: 76–81, 1987.
- [70681] 9011.Reeves, R. H.; Gearhart, J. D.; Hecht, N. B.; Yelick, P.;

Johnson,P.; O'Brien, S. J.: The gene encoding protamine 1 is located on chromosome16 in humans and near the proximal end of mouse chromosome 16, whereit is tightly linked to protamine 2. (Abstract) Cytogenet. Cell Genet. 46:680 only, 1987.

[70682] 9012.Reeves, R. H.; Gearhart, J. D.; Hecht, N. B.; Yelick, P.; Johnson,P.; O'Brien, S. J.: The gene encoding protamine 1 is located on humanchromosome 16, and near the proximal end of mouse chromosome 16 whereit is tightly linked to the gene encoding protamine 2. J. Hered. 80:442–446, 1989.

[70683] 9013.Viguie, F.; Domenjoud, L.; Rousseau–Merck, M.–F.; Dadoune, J.–P.;Chevaillier, P.: Chromosomal localization of the human protaminegenes, PRM1 and PRM2, to 16p13.3 by in situ hybridization. Hum. Genet. 85:171–174, 1990.

[70684] 9014.de The, H.; Marchio, A.; Tiollais, P.; Dejean, A.: A novel steroidthyroid hormone receptor–related gene inappropriately expressed inhuman hepatocellular carcinoma. Nature 330: 667–670, 1987.

[70685] 9015.Clevers, H. C.; Dunlap, S.; Wileman, T. E.; Terhorst, C.: HumanCD3–epsilon gene contains three minixons and is transcribed froma non–TATA promoter. Proc. Nat. Acad. Sci. 85: 8156–8160, 1988.

- [70686] 9016.DeJarnette, J. B.; Sommers, C. L.; Huang, K.; Woodside, K. J.;Emmons, R.; Katz, K.; Shores, E. W.; Love, P. E.: Specific requirementfor CD3–epsilon in T cell development. *Proc. Nat. Acad. Sci.* 95:14909–14914, 1998.
- [70687] 9017.Gold, D. P.; Puck, J. M.; Pettey, C. L.; Cho, M.; Coligan, J.;Woody, J. N.; Terhorst, C.: Isolation of cDNA clones encoding the20K non-glycosylated polypeptide chain of the human T–cell receptor/T3complex. *Nature* 321: 431–434, 1986.
- [70688] 9018.Le Deist, F.; Thoenes, G.; Corado, J.; Lisowska–GrosPierre, B.;Fischer, A.: Immunodeficiency with low expression of the T cell receptor/CD3complex: effect on T lymphocyte activation. *Europ. J. Immun.* 21:1641–1647, 1991.
- [70689] 9019.Soudais, C.; de Villartay, J.–P.; Le Deist, F.; Fischer, A.; Lisowska–GrosPierre,B.: Independent mutations of the human CD3–epsilon gene resultingin a T cell receptor/CD3 complex immunodeficiency. *Nature Genet.* 3:77–81, 1993.
- [70690] 9020.Thoenes, G.; Soudais, C.; Le Deist, F.; Griscelli, C.; Fischer,A.; Lisowska–GrosPierre, B.: Structural analysis of low TCR–CD3 complexexpression in T cells of an immunodeficient patient. *J. Biol. Chem.* 267:487–493, 1992.



- [70691] 9021.Wong, S.; Moore, S.; Orisio, S.; Millward, A.; De-  
maine, A. G.:Susceptibility to type I diabetes in women is  
associated with theCD3 epsilon locus on chromosome 11.  
Clin. Exp. Immun. 83: 69–73,1991.
- [70692] 9022.DeMartino, G. N.; Proske, R. J.; Moomaw, C. R.;  
Strong, A. A.;Song, X.; Hisamatsu, H.; Tanaka, K.; Slaugh-  
ter, C. A.: Identification,purification, and characterization  
of a PA700–dependent activatorof the proteasome. J. Biol.  
Chem. 271: 3112–3118, 1996.
- [70693] 9023.Ohana, B.; Moore, P. A.; Ruben, S. M.; Southgate, C.  
D.; Green,M. R.; Rosen, C. A.: The type 1 human immun-  
odeficiency virus Tatbinding protein is a transcriptional  
activator belonging to an additionalfamily of evolutionarily  
conserved genes. Proc. Nat. Acad. Sci. 90:138–142, 1993.
- [70694] 9024.Patel, A.; Rochelle, J. M.; Jones, J. M.; Sumegi, J.; Uhl,  
G. R.;Seldin, M. F.; Meisler, M. H.; Gregor, P.: Mapping of  
the taurinetransporter gene to mouse chromosome 6 and  
to the short arm of humanchromosome 3. Genomics 25:  
314–317, 1995.
- [70695] 9025.Ramamoorthy, S.; Leibach, F. H.; Mahesh, V. B.; Han,  
H.; Yang–Feng,T.; Blakely, R. D.; Ganapathy, V.: Functional  
characterization andchromosomal localization of a cloned  
taurine transporter from humanplacenta. Biochem. J. 300:

893–900, 1994.

- [70696] 9026.Kreczel, W.; Ghyselinck, N.; Samad, T. A.; Dupe, V.; Kastner, P.; Borrelli, E.; Chambon, P.: Impaired locomotion and dopamine signaling in retinoid receptor mutant mice. *Science* 279: 863–867, 1998.
- [70697] 9027.Lotan, R.; Xu, X.-C.; Lippman, S. M.; Ro, J. Y.; Lee, J. S.; Lee, J. J.; Hong, W. K.: Suppression of retinoic acid receptor-beta in premalignant oral lesions and its up-regulation by isotretinoin. *New Eng. J. Med.* 332: 1405–1410, 1995.
- [70698] 9028.Mattei, M.-G.; de The, H.; Mattei, J.-F.; Marchio, A.; Tiollais, P.; Dejean, A.: Assignment of the human hap retinoic acid receptor RAR-beta gene to the p24 band of chromosome 3. *Hum. Genet.* 80: 189–190, 1988.
- [70699] 9029.Nadeau, J. H.; Compton, J. G.; Giguere, V.; Rossant, J.; Varmuza, S.: Close linkage of retinoic acid receptor genes with homeobox- and keratin-encoding genes on paralogous segments of mouse chromosomes 11 and 15. *Mammalian Genome* 3: 202–208, 1992.
- [70700] 9030.Samad, A.; Kreczel, W.; Chambon, P.; Borrelli, E.: Regulation of dopaminergic pathways by retinoids: activation of the D2 receptor promoter by members of the retinoic acid receptor-retinoid X receptor family. *Proc. Nat.*

Acad. Sci. 94: 14349–14354, 1997.

- [70701] 9031. Flagiello, D.; Apiou, F.; Gibaud, A.; Poupon, M. F.; Dutrillaux, B.; Malfoy, B.: Assignment of the genes for cellular retinoic acid binding protein 1 (CRABP1) and 2 (CRABP2) to human chromosome band 15q24 and 1q21.3, respectively, by in situ hybridization. *Cytogenet. Cell Genet.* 76: 17–18, 1997.
- [70702] 9032. Geurts van Kessel, A.; de Leeuw, H.; Dekker, E. J.; Rijks, L.; Spurr, N.; Ledbetter, D.; Kootwijk, E.; Vaessen, M. J.: Localization of the cellular retinoic acid binding protein (CRABP) gene relative to the acute promyelocytic leukemia-associated breakpoint on human chromosome 15. *Hum. Genet.* 87: 201–204, 1991.
- [70703] 9033. MacGregor, T. M.; Copeland, N. G.; Jenkins, N. A.; Giguere, V.: The murine gene for cellular retinoic acid-binding protein type II: genomic organization, chromosomal localization, and post-transcriptional regulation by retinoic acid. *J. Biol. Chem.* 267: 7777–7783, 1992.
- [70704] 9034. Ong, D. E.: Cellular retinoid-binding proteins. *Arch. Derm.* 123: 1693–1695, 1987.
- [70705] 9035. Vaessen, M. J.; Rijks, L.; Dekker, E. J.; Kootwijk, E.; Bootsma, D.; Westerveld, A.; Geurts van Kessel, A. H. M.: Localization of the human cellular retinoic acid-binding

protein (CRABP) on chromosome 15. (Abstract) Cytogenet. Cell Genet. 51: 1094 only, 1989.

[70706] 9036. Elder, J. T.; Astrom, A.; Pettersson, U.; Voorhees, J. J.; Trent, J. M.: Assignment of the human CRABP-II gene to chromosome 1q21 by nonisotopic in situ hybridization. Hum. Genet. 89: 487–490, 1992.

[70707] 9037. Flagiello, D.; Apiou, F.; Gibaud, A.; Poupon, M. F.; Dutrillaux, B.; Malfoy, B.: Assignment of the genes for cellular retinoic acid binding protein 1 (CRABP1) and 2 (CRABP2) to human chromosome bands 15q24 and 1q21.3, respectively, by in situ hybridization. Cytogenet. Cell Genet. 76: 17–18, 1997.

[70708] 9038. Giguere, V.; Lyn, S.; Yip, P.; Siu, C.-H.; Amin, S.: Molecular cloning of cDNA encoding a second cellular retinoic acid-binding protein. Proc. Nat. Acad. Sci. 87: 6233–6237, 1990.

[70709] 9039. Dieterich, W.; Ehnis, T.; Bauer, M.; Donner, P.; Volta, U.; Riecken, E. O.; Schuppan, D.: Identification of tissue transglutaminase as the autoantigen of celiac disease. Nature Med. 3: 797–801, 1997.

[70710] 9040. Gentile, V.; Davies, P. J. A.; Baldini, A.: The human tissue transglutaminase gene maps on chromosome 20q12 by in situ fluorescence hybridization. Genomics

20:295–297, 1994.

- [70711] 9041. Gentile, V.; Saydak, M.; Chiocca, E. A.; Akande, O.; Birckbichler, P. J.; Lee, K. N.; Stein, J. P.; Davies, P. J. A.: Isolation and characterization of cDNA clones to mouse macrophage and human endothelial cell tissue transglutaminases. *J. Biol. Chem.* 266: 478–483, 1991.
- [70712] 9042. Lu, S.; Saydak, M.; Gentile, V.; Stein, J. P.; Davies, P. J. A.: Isolation and characterization of the human tissue transglutaminase gene promoter. *J. Biol. Chem.* 270: 9748–9756, 1995.
- [70713] 9043. Wang, M.; Kim, I.-G.; Steinert, P. M.; McBride, O. W.: Assignment of the human transglutaminase 2 (TGM2) and transglutaminase 3 (TGM3) genes to chromosome 20q11.2. *Genomics* 23: 721–722, 1994.
- [70714] 9044. Dodd, J.; Morton, S. B.; Karagogeos, D.; Yamamoto, M.; Jessell, T. M.: Spatial regulation of axonal glycoprotein expression on subset of embryonic spinal neurons. *Neuron* 1: 105–116, 1988.
- [70715] 9045. Freigang, J.; Proba, K.; Leder, L.; Diederichs, K.; Sonderegger, P.; Welte, W.: The crystal structure of the ligand binding module of axonin-1/TAG-1 suggests a zipper mechanism for neural cell adhesion. *Cell* 101: 425–433, 2000.

- [70716] 9046.Kenwrick, S.; Leversha, M.; Rooke, L.; Hasler, T.; Sonderegger,P.: Localization of the human TAX-1 gene to 1q32.1: a region implicated in microcephaly and Van der Woude syndrome. *Hum. Molec. Genet.* 2:1461–1462, 1993.
- [70717] 9047.Kozlov, S. V.; Giger, R. J.; Hasler, T.; Korvatska, E.; Schorderet,D. F.; Sonderegger, P.: The human TAX1 gene encoding the axon-associated cell adhesion molecule TAG-1/axonin-1: genomic structure and basic promoter. *Genomics* 30: 141–148, 1995.
- [70718] 9048.Rickman, D. S.; Tyagi, R.; Zhu, X.-X.; Bobek, M. P.; Song, S.; Blaivas, M.; Misek, D. E.; Israel, M. A.; Kurnit, D. M.; Ross, D.A.; Kish, P. E.; Hanash, S. M.: The gene for the axonal cell adhesion molecule TAX-1 is amplified and aberrantly expressed in malignant gliomas. *Cancer Res.* 61: 2162–2168, 2001.
- [70719] 9049.Tsiotra, P. C.; Karagogeos, D.; Theodorakis, K.; Michaelisis, T.M.; Modi, W. S.; Furley, A. J.; Jessell, T. M.; Papamatheakis, J.: Isolation of the cDNA and chromosomal localization of the gene (TAX1) encoding the human axonal glycoprotein TAG-1. *Genomics* 18: 562–567, 1993.
- [70720] 9050.Artavanis-Tsakonas, S.; Matsuono, K.; Fortini, M.: Notch signaling. *Science* 268:225–232, 1995.

- [70721] 9051.Brou, C.; Logeat, F.; Gupta, N.; Bessia, C.; LeBail, O.; Doedens, J. R.; Cumano, A.; Roux, P.; Black, R. A.; Israel, A.: A novel proteolytic cleavage involved in Notch signaling: the role of the disintegrin-metalloprotease TACE. *Molec. Cell* 5: 207–216, 2000.
- [70722] 9052.Bruckner, K.; Perez, L.; Clausen, H.; Cohen, S.: Glycosyltransferase activity of Fringe modulates Notch–Delta interactions. *Nature* 406:411–415, 2000.
- [70723] 9053.Ackerman, M. J.; Clapham, D. E.: Ion channels—basic science and clinical disease. *New Eng. J. Med.* 336: 1575–1586, 1997.
- [70724] 9054.Ackerman, M. J.; Schroeder, J. J.; Berry, R.; Schaid, D. J.; Porter, C. B. J.; Michels, V. V.; Thibodeau, S. N.: A novel mutation in KVLQT1 is the molecular basis of inherited long QT syndrome in a near-drowning patient's family. *Pediatr. Res.* 44: 148–153, 1998.
- [70725] 9055.Ackerman, M. J.; Tester, D. J.; Porter, C. J.; Edwards, W. D.: Molecular diagnosis of the inherited long-QT syndrome in a woman who died after near-drowning. *New Eng. J. Med.* 341: 1121–1125, 1999.
- [70726] 9056.Barlow, D. P.: Box: KVLQT1 complexities in Beckwith–Wiedeman (sic) syndrome. *Nature Genet.* 15: 114 only, 1997.

- [70727] 9057.Barlow, J. B.; Bosman, C. K.; Cochrane, J. W. C.: Congenital cardiacarrhythmia. Lancet II: 531 only, 1964.
- [70728] 9058.Baudouy, P.; Andreassian, B.; Attuel, P.; Greze, M.; Soulie, J.;Fruchaud, J.: Syndrome de Romano–Ward et stellectomie gauche: revuegenerale a propos d'un nouveau cas. Arch. Mal. Coeur 70: 645–652,1977.
- [70729] 9059.Benhorin, J.; Kalman, Y. M.; Medina, A.; Towbin, J.; Rave–Harel,N.; Dyer, T. D.; Blangero, J.; MacCluer, J. W.; Kerem, B.: Evidenceof genetic heterogeneity in the long QT syndrome. Science 260: 1960–1962,1993.
- [70730] 9060.Bhandari, A. K.; Scheinman, M.: The long QT syndrome. Mod. ConceptsCardiovasc. Dis. 54: 45–50, 1985.
- [70731] 9061.Bonduelle, M.: Personal Communication. Brussels, Belgium 5/30/1993.
- [70732] 9062.Casimiro, M. C.; Knollmann, B. C.; Ebert, S. N.; Vary, J. C.,Jr.; Greene, A. E.; Franz, M. R.; Grinberg, A.; Huang, S. P.; Pfeifer,K.: Targeted disruption of the Kcnq1 gene produces a mouse modelof Jervell and Lange–Nielsen syndrome. Proc. Nat. Acad. Sci. 98:2526–2531, 2001.
- [70733] 9063.Chen, Q.; Zhang, D.; Gingell, R. L.; Moss, A. J.; Napolitano,C.; Priori, S. G.; Schwartz, P. J.; Kehoe, E.; Robinson, J. L.; Schulze–Bahr,E.; Wang, Q.; Towbin, J. A.: Homozygous deletion in KVLQT1 associatedwith Jervell



and Lange–Nielsen syndrome. *Circulation* 99:  
1344–1347,1999.

- [70734] 9064.Cleary, M. A.; van Raamsdonk, C. D.; Levorse, J.; Zheng, B.; Bradley,A.; Tilghman, S. M.: Disruption of an imprinted gene cluster by atargeted chromosomal translocation in mice. *Nature Genet.* 29: 78–82,2001.
- [70735] 9065.Curran, M.; Atkinson, D.; Timothy, K.; Vincent, G. M.; Moss, A.J.; Leppert, M.; Keating, M.: Locus heterogeneity of autosomal dominantlong QT syndrome. *J. Clin. Invest.* 92: 799–803, 1993.
- [70736] 9066.DeSilvey, D. L.; Moss, A. J.: Primidone in the treatment of thelong QT syndrome: QT shortening and ventricular arrhythmia suppression. *Ann.Intern. Med.* 93: 53–54, 1980.
- [70737] 9067.DiSegni, E.; David, D.; Katzenstein, M.; Klein, H. O.; Kaplinsky,E.; Levy, M. J.: Permanent overdrive pacing for the suppression ofrecurrent ventricular tachycardia in a newborn with long QT syndrome. *J.Electrocardiol.* 13: 189–192, 1980.
- [70738] 9068.Donger, C.; Denjoy, I.; Berthet, M.; Neyroud, N.; Cruaud, C.;Bennaceur, M.; Chivoret, G.; Schwartz, K.; Coumel, P.; Guicheney,P.: KVLQT1 C-terminal missense mutation causes a forme fruste long–QTsyndrome. *Circu-*

lation 96: 2778–2781, 1997.

[70739] 9069.Engel, J. R.; et al.; et al.: Epigenotype–phenotype correlations in Beckwith–Wiedemann syndrome. *J. Med. Genet.* 37: 921–926, 2000.

[70740] 9070.Fitzpatrick, G. V.; Soloway, P. D.; Higgins, M. J.: Regional loss of imprinting and growth deficiency in mice with a targeted deletion of KvDMR1. *Nature Genet.* 9 Sept, 2002. Note: Advance Electronic Publication.

[70741] 9071.Flugelman, M. Y.; Pollack, S.; Hammerman, H.; Riss, E.; Barzilai, D.: Congenital prolongation of Q–T interval: a family study of three generations. *Cardiology* 69: 170–174, 1982.

[70742] 9072.Furberg, C.; Hornell, H.: Familial QT prolongation and risk of sudden death. *Acta Paediat. Scand.* 64: 777–782, 1975.

[70743] 9073.Gale, G. E.; Bosman, C. K.; Tucker, R. B. K.; Barlow, J. B.: Hereditary prolongation of Q–T interval: study of two families. *Brit. Heart J.* 32: 505–509, 1970.

[70744] 9074.Gamstorp, I.; Nilsen, R.; Westling, H.: Congenital cardiac arrhythmia. (Letter) *Lancet* II: 965 only, 1964.

[70745] 9075.Garza, L. A.; Vick, R. L.; Nora, J. J.; McNamara, D. G.: Heritable Q–T prolongation without deafness. *Circulation* 41: 39–48, 1970.

- [70746] 9076.Giuffre, R. M.; Hejtmancik, J. F.; McCabe, E. R. B.; Towbin, J.A.: Long QT (Romano–Ward) syndrome: molecular genetic evidence against tight HLA linkage. (Abstract) Am. J. Hum. Genet. 47 (suppl.): A180only, 1990.
- [70747] 9077.Gohl, K.; Feistel, H.; Weikl, A.; Bachmann, K.; Wolf, F.: Congenital myocardial sympathetic dysinnervation (CMSD)—a structural defect of idiopathic long QT syndrome. PACE 14: 1544–1553, 1991.
- [70748] 9078.Hashiba, K.: Hereditary QT prolongation syndrome in Japan: genetic analysis and pathological findings of the conduction system. Jpn.Circ. J. 42: 1133–1150, 1978.
- [70749] 9079.Horn, C. A.; Beekman, R. H.; Dick, M., II; Lacina, S. J.: The congenital long QT syndrome: an unusual cause of childhood seizures. Am.J. Dis. Child. 140: 659–661, 1986.
- [70750] 9080.Itoh, S.; Munemura, S.; Satoh, H.: A study of the inheritance pattern of Romano–Ward syndrome: prolonged Q–T interval, syncope, and sudden death. Clin. Pediat. 21: 20–24, 1982.
- [70751] 9081.Bulle, F.; Mattei, M. G.; Siegrist, S.; Pawlak, A.; Passage, E.; Chobert, M. N.; Laperche, Y.; Guellaen, G.: Assignment of the human gamma–glutamyl transferase gene to the long arm of chromosome 22. Hum.Genet. 76: 283–286, 1987.

[70752] 9082. Miller, W. L.: Phenotypic heterogeneity associated with the splicing mutation in congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *J. Clin. Endocr. Metab.* 82: 1304 only, 1997.100. Miller, W. L.: Congenital adrenal hyperplasia. (Letter) *New Eng. J. Med.* 317: 1413–1414, 1987.101. Morel, Y.; Andre, J.; Uring-Lambert, B.; Hauptmann, G.; Betuel, H.; Tossi, M.; Forest, M. G.; David, M.; Bertrand, J.; Miller, W. L.: Rearrangements and point mutations of P450c21 genes are distinguished in five restriction endonuclease haplotypes identified by a new probing strategy in 57 families with congenital adrenal hyperplasia. *J. Clin. Invest.* 83: 527–536, 1989.102. Mornet, E.; Boue, J.; Raux-Demay, M.; Couillin, P.; Oury, J. F.; Dumez, Y.; Dausset, J.; Cohen, D.; Boue, A.: First trimester prenatal diagnosis of 21-hydroxylase deficiency by linkage analysis to HLA-DNA probes and by 17-hydroxyprogesterone determination. *Hum. Genet.* 73: 358–364, 1986.103. Mornet, E.; Crete, P.; Kuttann, F.; Raux-Demay, M.-C.; Boue, J.; White, P. C.; Boue, A.: Distribution of deletions and seven point mutations on CYP21B genes in three clinical forms of steroid 21-hydroxylase deficiency. *Am. J. Hum. Genet.* 48: 79–88, 1991.104. Mulaikal, R. M.; Migeon, C. J.; Rock, J. A.: Fertil-

ity rates in female patients with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *New Eng. J. Med.* 316: 178–182, 1987.105. Murtaza, L.; Sibert, J. R.; Hughes, I.; Balfour, I. C.: Congenital adrenal hyperplasia—a clinical and genetic survey: are we detecting male salt-losers? *Arch. Dis. Child.* 55: 622–625, 1980.106. Murtaza, L. M.; Hughes, I. A.; Sibert, J. R.; Balfour, I. C.: HLA and congenital adrenal hyperplasia. (Letter) *Lancet* II: 524 only, 1978.107. New, M. I.; Dupont, B.; Pang, S.; Pollack, M.; Levine, L. S.: An update of congenital adrenal hyperplasia. *Recent Prog. Horm. Res.* 37: 105–181, 1981.108. New, M. I.; Dupont, B.; Pollack, M. S.; Levine, L. S.: The biochemical basis for genotyping 21-hydroxylase deficiency. *Hum. Genet.* 58:123–127, 1981.109. New, M. I.; Levine, L. S.: Recent advances in 21-hydroxylase deficiency. *Ann. Rev. Med.* 35: 649–663, 1984.110. New, M. I.; Lorenzen, F.; Lerner, A. J.; Kohn, B.; Oberfield, S. E.; Pollack, M. S.; Dupont, B.; Stoner, E.; Levy, D. J.; Pang, S.; Levine, L. S.: Genotyping steroid 21-hydroxylase deficiency: hormonal reference data. *J. Clin. Endocr. Metab.* 57: 320–326, 1983.111. New, M. I.; Wilson, R. C.: Steroid disorders in children: congenital adrenal hyperplasia and apparent mineralocorticoid excess. *Proc. Nat. Acad. Sci.* 96:

12790–12797, 1999.112. Nikoshkov, A.; Lajic, S.; Holst, M.; Wedell, A.; Luthman, H.: Synergistic effect of partially inactivating mutations in steroid 21-hydroxylase deficiency. *J. Clin. Endocr. Metab.* 82: 194–199, 1997.113. Nimkarn, S.; Cerame, B. I.; Wei, J.-Q.; Domic, M.; Zunec, R.; Brkljacic, L.; Skrabic, V.; New, M. I.; Wilson, R. C.: Congenital adrenal hyperplasia (21-hydroxylase deficiency) without demonstrable genetic mutations. *J. Clin. Endocr. Metab.* 84: 378–381, 1999.114. Nordenstrom, A.; Marcus, C.; Axelson, M.; Wedell, A.; Ritzen, E. M.: Failure of cortisone acetate treatment in congenital adrenal hyperplasia because of defective 11-beta-hydroxysteroid dehydrogenase reductase activity. *J. Clin. Endocr. Metab.* 84: 1210–1213, 1999.115. Nordenstrom, A.; Thilen, A.; Hagenfeldt, L.; Larsson, A.; Wedell, A.: Genotyping is a valuable diagnostic complement to neonatal screening for congenital adrenal hyperplasia due to steroid 21-hydroxylase deficiency. *J. Clin. Endocr. Metab.* 84: 1505–1509, 1999.116. Olney, R. C.; Mougey, E. B.; Wang, J.; Shulman, D. I.; Sylvester, J. E.: Using real-time, quantitative PCR for rapid genotyping of the steroid 21-hydroxylase gene in a north Florida population. *J. Clin. Endocr. Metab.* 87: 735–741, 2002.117. Orta-Flores, Z.;

Cantu, J. M.; Dominguez, O. V.: Reciprocal interactions of progesterone and 17-alpha-OH-progesterone as exogenous substrates for rat adrenal 21-hydroxylase. *J. Steroid Biochem.* 7:761-767, 1976.118.

Owerbach, D.; Crawford, Y. M.; Draznin, M. B.: Direct analysis of CYP21B genes in 21-hydroxylase deficiency using polymerase chain reaction amplification. *Molec. Endocr.* 4: 125-131, 1990.119.

Owerbach, D.; Sherman, L.; Ballard, A.-L.; Azziz, R.: Pro453-to-ser mutation in CYP21 is associated with non-classic steroid 21-hydroxylase deficiency. *Molec. Endocr.* 6: 1211-1215, 1992.120.

Pang, S.; Murphey, W.; Levine, L. S.; Spence, D. A.; Leon, A.; LaFranchi, S.; Surve, A. S.; New, M. I.: A pilot newborn screening program for congenital adrenal hyperplasia in Alaska. *J. Clin. Endocr. Metab.* 55: 413-420, 1982.121.

Pang, S.; Pollack, M. S.; Loo, M.; Green, O.; Nussbaum, R.; Clayton, G.; Dupont, B.; New, M. I.: Pitfalls of prenatal diagnosis of 21-hydroxylase deficiency congenital adrenal hyperplasia. *J. Clin. Endocr. Metab.* 61:89-97, 1985.122.

Partanen, J.; Campbell, R. D.: Substitution of ile172-to-asn in the steroid 21-hydroxylase B (P450c21B) gene in a Finnish patient with the simple virilizing form of congenital adrenal hyperplasia. *Hum. Genet.* 87: 716-720, 1991.123.

Paulino, L. C.; Araujo, M.; Guerra, G., Jr.; Marini, S. H.; DeMello, M. P.: Mutation distribution and CYP21/C4 locus variability in Brazilian families with the classical form of the 21-hydroxylase deficiency. *Acta Paediat.* 88: 275–83, 1999.124. Petersen, F.; Knudsen, F. U.; Nielsen, M. D.; Mikkelsen, M.: Congenital adrenal hyperplasia associated with a balanced 13–18 translocation. *Europ. J. Pediat.* 133: 283–285, 1980.125. Petersen, K. E.; Svejgaard, A.; Nielsen, M. D.; Dissing, J.: Heterozygotes and cryptic patients in families of patients with congenital adrenal hyperplasia (21-hydroxylase deficiency): HLA and glyoxalase typing and hormonal studies. *Hormone Res.* 16: 151–159, 1982.126. Pollack, M. S.; Keenan, B.; Christiansen, F. T.; Cobain, T. J.; Dawkins, R. L.; Clayton, G.: The immunological detection of a 21-OH deficiency mutation HLA supertype. *Am. J. Hum. Genet.* 38: 688–698, 1986.127. Pollack, M. S.; Levine, L. S.; O'Neill, G. J.; Pang, S.; Lorenzen, F.; Kohn, B.; Rondanini, G. F.; Chi-umello, G.; New, M. I.; Dupont, B.: HLA linkage and B14, DR1, BfS haplotype association with the genes for late onset and cryptic 21-hydroxylase deficiency. *Am. J. Hum. Genet.* 33: 540–550, 1981.128. Pollack, M. S.; Levine, L. S.; Pang, S.; Owens, R. P.; Nitowsky, H. M.; Mau-



rer, D.; New, M. I.; Duchon, M.; Merkatz, I. R.; Sachs, G.; Dupont, B.: Prenatal diagnosis of congenital adrenal hyperplasia (21-hydroxylase deficiency) by HLA typing. Lancet I: 1107–1108, 1979.129. Pollack, M. S.; New, M. I.; O'Neill, G. J.; Levine, L. S.; Callaway, C.; Pang, S.; Cacciari, E.; Mantero, F.; Cassio, A.; Scaroni, C.; Chiumello, G.; Ron-  
danini, G. F.; Gargantini, L.; Giovannelli, G.; Viridis, R.; Bar-  
tolotta, E.; Migliori, C.; Pintor, C.; Tato, L.; Barboni, F.; Dupont, B.: HLA genotypes and HLA-linked genetic mark-  
ers in Italian patients with classical 21-hydroxylase defi-  
ciency. Hum. Genet. 58:331–337, 1981.130. Prader, A.: Die Haeufigkeit des kongenitalen adrenogenitalen Syn-  
droms. Helv. Paediat. Acta 13: 426–431, 1958.131.  
Prader, A.; Anders, G. J. P. A.; Habich, H.: Zur Genetik  
des kongenitalen adrenogenitalen Syndroms (virilisierende  
Nebennierenhyperplasia). Helv. Paediat. Acta 17: 271–284,  
1962.132. Pucholt, V.; Fitzsimmons, J. S.; Gelsthorpe, K.;  
Reynolds, M. A.; Milner, R. D. G.: Location of the gene for  
21-hydroxylase deficiency. J. Med. Genet. 17: 447–452,  
1980.133. Qazi, Q. H.; Thompson, M. W.: Incidence of  
salt-losing form of congenital virilizing adrenal hyperpla-  
sia. Arch. Dis. Child. 47:302–303, 1972.134. Ravichan-  
dran, R.; Lafferty, F.; McGinniss, M. J.; Taylor, H. C.: Con-

genital adrenal hyperplasia presenting as massive adrenal incidentalomas in the sixth decade of life: report of two patients with 21-hydroxylase deficiency. *J. Clin. Endocrin. Metab.* 81: 1776–1779, 1996. 135. Reindollar, R. H.; Lewis, J. B.; White, P. C.; Fernhoff, P. M.; McDonough, P. G.; Whitney, J. B., III: Prenatal diagnosis of 21-hydroxylase deficiency by the complementary deoxyribonucleic acid probe for cytochrome P-450(C-21OH). *Am. J. Obstet. Gynec.* 158: 545–547, 1988. 136. Rodrigues, N. R.; Dunham, I.; Yu, C. Y.; Carroll, M. C.; Porter, R. R.; Campbell, R. D.: Molecular characterization of the HLA-linked steroid 21-hydroxylase B gene from an individual with congenital adrenal hyperplasia. *EMBO J.* 6: 1653–1661, 1987. 137. Rosenmann, A.; Schumert, Z.; Theodor, R.; Cohen, T.; Brautbar, C.: Amniotic 17-alpha hydroxyprogesterone and HLA typing for the prenatal diagnosis of 21-alpha hydroxylase deficiency–congenital adrenal hyperplasia. *Am. J. Med. Genet.* 6: 295–300, 1980. 138. Rumsby, G.; Carroll, M. C.; Porter, R. R.; Grant, D. B.; Hjelm, M.: Deletion of the steroid 21-hydroxylase and complement C4 genes in congenital adrenal hyperplasia. *J. Med. Genet.* 23: 204–209, 1986. 139. Schneider, P. M.; Carroll, M. C.; Alper, C. A.;

Rittner, C.; Whitehead, A. S.; Yunis, E. J.; Colten, H. R.: Polymorphism of the human complement C4 and steroid 21-hydroxylase genes: restriction fragment length polymorphisms revealing structural deletions, homoduplications, and size variants. *J. Clin. Invest.* 78: 650–657, 1986.140. Sherman, S. L.; Aston, C. E.; Morton, N. E.; Speiser, P. W.; New, M. I.: A segregation and linkage study of classical and nonclassical 21-hydroxylase deficiency. *Am. J. Hum. Genet.* 42: 830–838, 1988.141. Sinnott, P.; Collier, S.; Costigan, C.; Dyer, P. A.; Harris, R.; Strachan, T.: Genesis by meiotic unequal crossover of a de novo deletion that contributes to a steroid 21-hydroxylase deficiency. *Proc. Nat. Acad. Sci.* 87: 2107–2111, 1990.142. Sinnott, P. J.; Dyer, P. A.; Price, D. A.; Harris, R.; Strachan, T.: 21-hydroxylase deficiency families with HLA identical affected and unaffected sibs. *J. Med. Genet.* 26: 10–17, 1989.143. Slominski, A.; Ermak, G.; Mihm, M.: ACTH receptor, CYP11A1, CYP17 and CYP21A2 genes are expressed in skin. *J. Clin. Endocr. Metab.* 81: 2746–2749, 1996.144. Sobel, D. O.; Gutai, J. P.; Jones, J. C.; Wagener, D. D.; Smith, W.: Detection of heterozygote of 21 hydroxylase deficiency. (Letter) *Lancet* I: 47 only, 1980.145. Speiser, P. W.; Dupont, B.; Rubinstein, P.; Piazza, A.;

Kastelan,A.; New, M. I.: High frequency of nonclassical steroid 21-hydroxylase deficiency. *Am. J. Hum. Genet.* 37: 650-667, 1985.146. Speiser, P. W.; Dupont, J.; Zhu, D.; Serrat, J.; Buegeleisen,M.; Tusie-Luna, M.-T.; Lesser, M.; New, M. I.; White, P. C.: Disease expression and molecular genotype in congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *J. Clin. Invest.* 90: 584-595,1992.147. Speiser, P. W.; New, M. I.: Genetics of steroid 21-hydroxylase deficiency. *Trends Genet.* 1: 275-278, 1985.148. Speiser, P. W.; New, M. I.; Tannin, G. M.; Pickering, D.; Yang,S. Y.; White, P. C.: Genotype of Yupik Eskimos with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *Hum. Genet.* 88: 647-648,1992.149. Speiser, P. W.; New, M. I.; White, P. C.: Molecular genetic analysis of nonclassic steroid 21-hydroxylase deficiency associated with HLA-B14,DR1. *New Eng. J. Med.* 319: 19-23, 1988.150. Speiser, P. W.; New, M. I.; White, P. C.: Clinical and genetic characterization of nonclassic 21-hydroxylase deficiency. *Endocr.Res.* 15: 257-276, 1989.151. Spiro, R. P.; Christian, S. L.; Ledbetter, D. H.; New, M. I.; Wilson, R. C.; Roizen, N.; Rosenfield, R. L.: Intrauterine growth retardation associated with maternal uniparental disomy for chromosome 6 unmasked

by congenital adrenal hyperplasia. *Pediat. Res.* 46: 510–513, 1999.152. Stikkelbroeck, N. M. M. L.; Otten, B. J.; Pasic, A.; Jager, G.J.; Sweep, C. G. J. (Fred); Noordam, K.; Hermus, A. R. M. M.: Highprevalence of testicular adrenal rest tumors, impaired spermatogenesis, and Leydig cell failure in adolescent and adult males with congenital adrenal hyperplasia. *J. Clin. Endocr. Metab.* 86: 5721–5728, 2001.153. Tajima, T.; Fujieda, K.; Fujii–Kuriyama, Y.: De novo mutation causes steroid 21–hydroxylase deficiency in one family of HLA–identical affected and unaffected individuals. *J. Clin. Endocr. Metab.* 77:86–89, 1993.154. Tajima, T.; Fujieda, K.; Nakae, J.; Toyoura, T.; Shimozawa, K.; Kusuda, S.; Goji, K.; Nagashima, T.; Cutler, G. B., Jr.: Molecular basis of nonclassical steroid 21–hydroxylase deficiency detected by neonatal mass screening in Japan. *J. Clin. Endocr. Metab.* 82: 2350–2356, 1997.155. Tajima, T.; Fujieda, K.; Nakayama, K.; Fujii–Kuriyama, Y.: Molecular analysis of patient and carrier genes with congenital steroid 21–hydroxylase deficiency by using polymerase chain reaction and single strand conformation polymorphism. *J. Clin. Invest.* 92: 2182–2190, 1993.156. Thilen, A.; Larsson, A.: Congenital adrenal hyperplasia in Sweden 1969–1986: prevalence, symptoms and age at diag-

nosis. *Acta Paediat.Scand.* 79: 168–175, 1990.157.

Travitz, J.; Metzger, D. L.: Antenatal treatment for classic 21-hydroxylase forms of congenital adrenal hyperplasia and the issues. *Genet.Med.* 1: 224–230, 1999.158.

Tusie-Luna, M.-T.; Speiser, P. W.; Dumin, M.; New, M. I.; White, P. C.: A mutation (pro30-to-leu) in CYP21 represents a potential nonclassic steroid 21-hydroxylase deficiency allele. *Molec. Endocr.* 5:685–692, 1991.159.

Tusie-Luna, M.-T.; White, P. C.: Gene conversions and unequal crossovers between CYP21 (steroid 21-hydroxylase gene) and CYP21P involve different mechanisms. *Proc. Nat. Acad. Sci.* 92: 10796–10800, 1995.160.

Tusie-Luna, M. T.; Traktman, P.; White, P. C.: Determination of functional effects of mutations in the steroid 21-hydroxylase gene (CYP21) using recombinant vaccinia virus. *J. Biol. Chem.* 265: 20916–20922, 1990.161.

Urabe, K.; Kimura, A.; Harada, F.; Iwanaga, T.; Sasazuki, T.: Gene conversion in steroid 21-hydroxylase genes. *Am. J. Hum. Genet.* 46:1178–1186, 1990.162.

Urban, M. D.; Lee, P. A.; Migeon, C. J.: Adult height and fertility in men with congenital virilizing adrenal hyperplasia. *New Eng. J. Med.* 299: 1392–1396, 1978.163.

Warsof, S. L.; Larsen, J. W.; Kent, S. G.; Rosenbaum, K. N.; August, G. P.; Migeon, C. J.; Schul-

man, J. D.: Prenatal diagnosis of congenital adrenal hyperplasia. *Obstet. Gynec.* 55: 751–754, 1980.164. Webb, T.; Mackintosh, P.; Wells, L. J.: Cytogenetic evidence for the localisation of the gene for congenital adrenal hyperplasia. *Clin. Genet.* 17: 349–354, 1980.165. Wedell, A.: Molecular genetics of congenital adrenal hyperplasia (21-hydroxylase deficiency): implications for diagnosis, prognosis and treatment. *Acta Paediat.* 87: 159–164, 1998.166. Wedell, A.: An update on the molecular genetics of congenital adrenal hyperplasia: diagnostic and therapeutic aspects. *J. Pediatr. Endocr. Metab.* 11: 581–589, 1998.167. Wedell, A.; Luthman, H.: Steroid 21-hydroxylase deficiency: two additional mutations in salt-wasting disease and rapid screening of disease-causing mutations. *Hum. Molec. Genet.* 2: 499–504, 1993.168. Wedell, A.; Ritzen, E. M.; Haglund-Stengler, B.; Luthman, H.: Steroid 21-hydroxylase deficiency: three additional mutated alleles and establishment of phenotype-genotype relationships of common mutations. *Proc. Nat. Acad. Sci.* 89: 7232–7236, 1992.169. Wedell, A.; Thilen, A.; Ritzen, E. M.; Stengler, B.; Luthman, H.: Mutational spectrum of the steroid 21-hydroxylase gene in Sweden: implications for genetic diagnosis and association

with disease malformation. *J.Clin. Endocr. Metab.* 78: 1145–1152, 1994.170. Wenzel, U.; Schneider, M.; Zachmann, M.; Knorr–Murset, G.; Weber, A.; Prader, A.: Intelligence of patients with congenital adrenal hyperplasia due to 21–hydroxylase deficiency, their parents and unaffected siblings. *Helv. Paediat. Acta* 33: 11–16, 1978.171. Werder, E. A.; Siebenmann, R. E.; Knorr–Murset, G.; Zimmermann, A.; Sizonenko, P. C.; Theintz, P.; Girard, J.; Zachmann, M.; Prader, A.: The incidence of congenital adrenal hyperplasia in Switzerland – a survey of patients born in 1960 to 1974. *Helv. Paediat. Acta* 35: 5–11, 1980.172. Werkmeister, J. W.; New, M. I.; Dupont, B.; White, P. C.: Frequent deletion and duplication of the steroid 21–hydroxylase genes. *Am.J. Hum. Genet.* 39: 461–469, 1986.173. White, P. C.; Chaplin, D. D.; Weis, J. H.; Dupont, B.; New, M. I.; Seidman, J. G.: Two steroid 21–hydroxylase genes are located in the murine S region. *Nature* 312: 465–467, 1984.174. White, P. C.; Grossberger, D.; Onufer, B. J.; Chaplin, D. D.; New, M. I.; Dupont, B.; Strominger, J. L.: Two genes encoding steroid 21–hydroxylase are located near the genes encoding the fourth component of complement in man. *Proc. Nat. Acad. Sci.* 82: 1089–1093, 1985.175. White, P. C.;



New, M.; Dupont, B.: Congenital adrenal hyperplasia. (Letter) *New Eng. J. Med.* 317: 1414–1415, 1987.176.

White, P. C.; New, M. I.; Dupont, B.: Congenital adrenal hyperplasia. *NewEng. J. Med.* 316: 1519–1524, 1987.177.

White, P. C.; New, M. I.; Dupont, B.: HLA-linked congenital adrenal hyperplasia results from a defective gene encoding a cytochromeP-450 specific for steroid 21-hydroxylation. *Proc. Nat. Acad. Sci.* 81:7505–7509, 1984.178.

White, P. C.; New, M. I.; Dupont, B.: Structure of human steroid21-hydroxylase genes. *Proc. Nat. Acad. Sci.* 83: 5111–5115, 1986.179.

White, P. C.; Tusie-Luna, M.-T.; New, M. I.; Speiser, P. W.: Mutations in steroid 21-hydroxylase (CYP21). *Hum. Mutat.* 3: 373–378, 1994.180.

White, P. C.; Vitek, A.; Dupont, B.; New, M. I.: Characterization of frequent deletions causing steroid 21-hydroxylase deficiency. *Proc. Nat. Acad. Sci.* 85: 4436–4440, 1988.181.

Winqvist, O.; Karlsson, F. A.; Kampe, O.: 21-Hydroxylase, a major autoantigen in idiopathic Addison's disease. *Lancet* 339: 1559–1562, 1992.182.

Witchel, S. F.; Bhamidipati, D. K.; Hoffman, E. P.; Cohen, J. B.: Phenotypic heterogeneity associated with the splicing mutation in congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *J. Clin. En-*

docr. Metab. 81: 4081–4088, 1996.183. Witchel, S. F.; Lee, P. A.; Suda–Hartman, M.; Trucco, M.; Hoffman, E. P.: Evidence for a heterozygote advantage in congenital adrenalhyperplasia due to 21–hydroxylase deficiency. J. Clin. Endocr. Metab. 82:2097–2101, 1997.184. Wu, D.–A.; Chung, B.: Mutations of P450c21 (steroid 21–hydroxylase)at cys–428, val–281, and ser–268 result in complete, partial, or no loss of enzymatic activity, respectively. J. Clin. Invest. 88: 519–523, 1991.185. Wudy, S. A.; Dorr, H. G.; Solleder, C.; Djalali, M.; Homoki, J.: Profiling steroid hormones in amniotic fluid of midpregnancy by routine stable isotope dilution/gas chromatography–mass spectrometry: Reference values and concentrations in fetuses at risk for 21–hydroxylase deficiency. J. Clin. Endocr. Metab. 84: 2724–2728, 1999.186. Wudy, S. A.; Homoki, J.; Teller, W. M.: Successful prenatal treatment of congenital adrenal hyperplasia due to 21–hydroxylase deficiency. Europ. J. Pediat. 153: 556–559, 1994.187. Yang, S. Y.; Levine, L. S.; Zachmann, M.; New, M. I.; Prader, A.; Oberfield, S. E.; O'Neill, G. J.; Pollack, M. S.; Dupont, B.: Mapping of the 21–hydroxylase deficiency gene within the HLA linkage group. Transplant. Proc. 10: 753–755, 1978.188. Zachmann, M.; Prader, A.: Unusual heterozygotes of con–

genitaladrenal hyperplasia due to 21-hydroxylase deficiency confirmed byHLA tissue typing. Acta Endocr. 92: 542–546, 1979.

[70753] 9083.Uchida, S.; Kwon, H. M.; Yamauchi, A.; Preston, A. S.; Marumo,F.; Handler, J. S.: Molecular cloning of the cDNA for an MDCK cellNa(+)- and Cl(-)-dependent taurine transporter that is regulated byhypertonicity. Proc. Nat. Acad. Sci. 89: 8230–8234, 1992.

[70754] 9084.Sakaguchi, A. Y.; Zabel, B. U.; Grzeschik, K. H.; Law, M. L.;Ellis, R. W.; Skolnick, E. M.; Naylor, S. L.: Regional localizationof two human cellular Kirsten ras genes on chromosomes 6 and 12. Molec.Cell. Biol. 4: 989–993, 1984.

[70755] 9085.Neyroud, N.; Tesson, F.; Denjoy, I.; Leibovici, M.; Donger, C.;Barhanin, J.; Faure, S.; Gary, F.; Coumel, P.; Petit, C.; Schwartz,K.; Guicheney, P.: A novel mutation in the potassium channel geneKVLQT1 causes the Jervell and Lange–Nielsen cardioauditory syndrome. NatureGenet. 15: 186–189, 1997.

[70756] 9086.Pervaiz, S.; Brew, K.: Homology of beta-lactoglobulin, serum retinol-bindingprotein, and protein HC. Science 228: 335–337, 1985.

[70757] 9087.Barber, G. N.; Edelhoff, S.; Katze, M. G.; Disteche, C. M.: Chromosomalassignment of the interferon-inducible

double-stranded RNA-dependent protein kinase (PRKR) to human chromosome 2p21-p22 and mouse chromosome 17 E2. Genomics 16: 765-767, 1993.

[70758] 9088. Kuhen, K. L.; Shen, X.; Carlisle, E. R.; Richardson, A. L.; Weier, H.-U. G.; Tanaka, H.; Samuel, C. E.: Structural organization of the human gene (PKR) encoding an interferon-inducible RNA-dependent protein kinase (PKR) and differences from its mouse homolog. Genomics 36: 197-201, 1996.

[70759] 9089. Kuhen, K. L.; Shen, X.; Samuel, C. E.: Mechanism of interferon action: sequence of the human interferon-inducible RNA-dependent protein kinase (PKR) deduced from genomic clones. Gene 178: 191-193, 1996.

[70760] 9090. Hansen, R. S.; Wijmenga, C.; Luo, P.; Stanek, A. M.; Canfield, T. K.; Weemaes, C. M. R.; Gartler, S. M.: The DNMT3B DNA methyltransferase gene is mutated in the ICF immunodeficiency syndrome. Proc. Nat. Acad. Sci. 96: 14412-14417, 1999.

[70761] 9091. Hassan, K. M. A.; Norwood, T.; Gimelli, G.; Gartler, S. M.; Hansen, R. S.: Satellite 2 methylation patterns in normal and ICF syndrome cells and association of hypomethylation with advanced replication. Hum. Genet. 109: 452-462, 2001.

- [70762] 9092. Shirohzu, H.; Kubota, T.; Kumazawa, A.; Sado, T.; Chijiwa, T.; Inagaki, K.; Suetake, I.; Tajima, S.; Wakui, K.; Miki, Y.; Hayashi, M.; Fukushima, Y.; Sasaki, H.: Three novel DNMT3B mutations in Japanese patients with ICF syndrome. *Am. J. Med. Genet.* 112: 31–37, 2002.
- [70763] 9093. Bonifaci, N.; Moroianu, J.; Radu, A.; Blobel, G.: Karyopherin beta-2 mediates nuclear import of a mRNA binding protein. *Proc. Nat. Acad. Sci.* 94: 5055–5060, 1997.
- [70764] 9094. Pollard, V. W.; Michael, W. M.; Nakielnny, S.; Siomi, M. C.; Wang, F.; Dreyfuss, G.: A novel receptor-mediated nuclear protein import pathway. *Cell* 86: 985–994, 1996.
- [70765] 9095. Salinas, M.; Duprat, F.; Heurteaux, C.; Hugnot, J.-P.; Lazdunski, M.: New modulatory alpha subunits for mammalian Shab K(+) channels. *J. Biol. Chem.* 272: 24371–24379, 1997.
- [70766] 9096. Kas, K.; Roijer, E.; Voz, M.; Meyen, E.; Stenman, G.; Van de Ven, W. J. M.: A 2-Mb YAC contig and physical map covering the chromosome 8q12 breakpoint cluster region in pleomorphic adenomas of the salivary glands. *Genomics* 43: 349–358, 1997.
- [70767] 9097. Kas, K.; Voz, M. L.; Roijer, E.; Astrom, A.-K.; Meyen, E.; Stenman, G.; Van de Ven, W. J. M.: Promoter swapping

between the genes for a novel zinc finger protein and beta-catenin in pleiomorphic adenomas with t(3;8)(p21;q12) translocations. *Nature Genet.* 15: 170–174, 1997.

[70768] 9098. Szabo, G.; Dallmann, G.; Muller, G.; Patthy, L.; Soller, M.; Varga, L.: A deletion in the myostatin gene causes the compact (Cmpt) hypermuscular mutation in mice. *Mammalian Genome* 9: 671–672, 1998.

[70769] 9099. Zimmers, T. A.; Davies, M. V.; Koniaris, L. G.; Haynes, P.; Esquela, A. F.; Tomkinson, K. N.; McPherron, A. C.; Wolfman, N. M.; Lee, S.-J.: Induction of cachexia in mice by systemically administered myostatin. *Science* 296:1486–1488, 2002.

[70770] 9100. Fransen, M.; Terlecky, S. R.; Subramani, S.: Identification of a human PTS1 receptor docking protein directly required for peroxisomal protein import. *Proc. Nat. Acad. Sci.* 95: 8087–8092, 1998.

[70771] 9101. Albertini, M.; Rehling, P.; Erdmann, R.; Girzalsky, W.; Kiel, J. A. K.; Veenhuis, M.; Kunau, W.-H.: Pex14p, a peroxisomal membrane protein binding both receptors of the two PTS-dependent import pathways. *Cell* 89:83–92, 1997.

[70772] 9102. Cohen, P.; Rylatt, D. B.; Nimmo, G. A.: The hormonal

control of glycogen metabolism: the amino acid sequence at the phosphorylation site of protein phosphatase inhibitor-1. FEBS Lett. 76: 182–186, 1977.

[70773] 9103. Helps, N. R.; Street, A. J.; Elledge, S. J.; Cohen, P. T.: Cloning of the complete coding region for human protein phosphatase inhibitor 2 using the two hybrid system and expression of inhibitor 2 in E. coli. FEBS Lett. 340: 93–98, 1994.

[70774] 9104. Huang, F. L.; Glinsmann, W. H.: Separation and characterization of two phosphorylase phosphatase inhibitors from rabbit skeletal muscle. Europ. J. Biochem. 70: 419–426, 1976.

[70775] 9105. Majer, M.; Mott, D. M.; Mochizuki, H.; Rowles, J. C.; Pedersen, O.; Knowler, W. C.; Bogardus, C.; Prochazka, M.: Association of the glycogen synthase locus on 19q13 with NIDDM in Pima Indians. Diabetologia 39: 314–321, 1996.

[70776] 9106. Permana, P. A.; Mott, D. M.: Genetic analysis of human type 1 protein phosphatase inhibitor 2 in insulin-resistant Pima Indians. Genomics 41: 110–114, 1997.

[70777] 9107. Prochazka, M.; Mochizuki, H.; Baier, L. J.; Cohen, P. T. W.; Bogardus, C.: Molecular and linkage analysis of type-1 protein phosphatase catalytic beta-subunit gene: lack of evidence for its major role in insulin resistance in

Pima Indians. *Diabetologia* 38: 461–466, 1995.

- [70778] 9108.Sakagami, H.; Kondo, H.: Molecular cloning of the cDNA for ratphosphatase inhibitor-2 and its wide gene expression in the centralnervous system. *J. Chem. Neuroanat.* 8: 259–266, 1995.
- [70779] 9109.Sanseau, P.; Jackson, A.; Alderton, R. P.; Beck, S.; Senger, G.;Sheer, D.; Kelly, A.; Trowsdale, J.: Cloning and characterizationof human phosphatase inhibitor-2 (IPP-2) sequences. *Mammalian Genome* 5:490–496, 1994.
- [70780] 9110.Charest, D. L.; Mordret, G.; Harder, K. W.; Jirik, F.; Pelech,S. L.: Molecular cloning, expression, and characterization of thehuman mitogen-activated protein kinase p44erk1. *Molec. Cell. Biol.* 13:4679–4690, 1993.
- [70781] 9111.Durand, B.; Sperisen, P.; Emery, P.; Barras, E.; Zuferey, M.;Mach, B.; Reith, W.: RFXAP, a novel subunit of the RFX DNA bindingcomplex is mutated in MHC class II deficiency. *EMBO J.* 16: 1045–1055,1997.
- [70782] 9112.Mach, B.: Personal Communication. Geneva, Switzerland 10/28/1998.
- [70783] 9113.Nekrep, N.; Jabrane-Ferrat, N.; Peterlin, B. M.: Mutations inthe bare lymphocyte syndrome define critical steps in the assemblyof the regulatory factor X complex. *Molec. Cell Biol.* 20: 4455–4461,2000.



- [70784] 9114.Schwartz, R. S.: The case of the bare lymphocyte syndrome: trackingdown faulty transcription factors. (Editorial) New Eng. J. Med. 337:781–783, 1997.
- [70785] 9115.Steimle, V.; Durand, B.; Barras, E.; Zufferey, M.; Hadam, M. R.;Mach, B.; Reith, W.: A novel DNA binding–regulatory factor is mutatedin primary MHC class II deficiency (bare lymphocyte syndrome). GenesDev. 9: 1021–1032, 1995.
- [70786] 9116.Touraine, J. L.; Betuel, H.: Immunodeficiency diseases and expressionof HLA antigens. Hum. Immun. 2: 147–153, 1981.
- [70787] 9117.Villard, J.; Lisowska–GrosPierre, B.; van den Elsen, P.; Fischer,A.; Reith, W.; Mach, B.: Mutation of RFXAP, a regulator of MHC classII genes, in primary MHC class II deficiency. New Eng. J. Med. 337:748–753, 1997.
- [70788] 9118.Peijnenburg, A.; Van Eggermond, M. C. J. A.; Van den Berg, R.;Sanal, O.; Vossen, J. M. J. J.; Van den Elsen, P. J.: Molecular analysisof an MHC class II deficiency patient reveals a novel mutation inthe RFX5 gene. Immunogenetics 49: 338–345, 1999.
- [70789] 9119.Kim, I.; Kim, J.–H.; Ryu, Y. S.; Jung, S. H.; Nah, J. J.; Koh,G. Y.: Characterization and expression of a novel alternatively splicedhuman angiopoietin–2. J. Biol. Chem.

275: 18550–18556, 2000.

- [70790] 9120. Maisonpierre, P. C.; Suri, C.; Jones, P. F.; Bartunkova, S.; Wiegand, S. J.; Radziejewski, C.; Compton, D.; McClain, J.; Aldrich, T. H.; Papadopoulos, N.; Daly, T. J.; Davis, S.; Sato, T. N.; Yancopoulos, G. D.: Angiopoietin-2, a natural antagonist for Tie2 that disrupts in vivo angiogenesis. *Science* 277: 55–60, 1997.
- [70791] 9121. Tanaka, S.; Mori, M.; Sakamoto, Y.; Makuuchi, M.; Sugimachi, K.; Wands, J. R.: Biologic significance of angiopoietin-2 expression in human hepatocellular carcinoma. *J. Clin. Invest.* 103: 341–345, 1999.
- [70792] 9122. Chow, V. T. K.; Quek, H. H.: HEP-COP, a novel human gene whose product is highly homologous to the alpha-subunit of the yeast coatamer protein complex. *Gene* 169: 223–227, 1996.
- [70793] 9123. Quek, H. H.; Chow, V. T. K.: Molecular and cellular studies of the human homolog of the 160-kD alpha-subunit of the coatamer protein complex. *DNA Cell Biol.* 16: 275–280, 1997.
- [70794] 9124. Quek, H. H.; Chow, V. T. K.: Genomic organization and mapping of the human HEP-COP gene (COPA) to 1q. *Cytogenet. Cell Genet.* 76: 139–143, 1997.
- [70795] 9125. Leffers, H.; Nielsen, M. S.; Andersen, A. H.; Honore,

B.; Madsen,P.; Vandekerckhove, J.; Celis, J. E.: Identification of two humanrho GDP dissociation inhibitor proteins whose overexpression leadsto disruption of the actin cytoskeleton. *Exp. Cell Res.* 209: 165–174,1993.

[70796] 9126.Wagner, T.; Tommerup, N.; Wirth, J.; Leffers, H.; Zimmer, J.; Back,E.; Weissenbach, J.; Scherer, G.: A somatic cell hybrid panel fordistal 17q: GDIA1 maps to 17q25.3. *Cytogenet. Cell Genet.* 76: 172–175,1997.

[70797] 9127.Sanicola, M.; Hession, C.; Worley, D.; Carmillo, P.; Ehrenfels,C.; Walus, L.; Robinson, S.; Jaworski, G.; Wei, H.; Tizard, R.; Whitty,A.; Pepinsky, R. B.; Cate, R. L.: Glial cell line–derived neurotrophicfactor–dependent RET activation can be mediated by two different cell–surfaceaccessory proteins. *Proc. Nat. Acad. Sci.* 94: 6238–6243, 1997.

[70798] 9128.Suvanto, P.; Wartiovaara, K.; Lindahl, M.; Arumae, U.; Moshnyakov,M.; Horelli–Kuitunen, N.; Airaksinen, M. S.; Palotie, A.; Sariola,H.; Saarma, M.: Cloning, mRNA distribution and chromosomal localisationof the gene for glial cell lined–derived neurotrophic factor receptorbeta, a homologue to GDNFR–alpha. *Hum. Molec. Genet.* 6: 1267–1273,1997.

[70799] 9129.Vanhorne, J. B.; Gimm, O.; Myers, S. M.; Kaushik, A.; von Deimling,A.; Eng, C.; Mulligan, L. M.: Cloning and

characterization of the human GFRA2 locus and investigation of the gene in Hirschsprung disease. Hum.Genet. 108: 409–415, 2001.

- [70800] 9130.Katsanis, N.; Yaspo, M.-L.; Fisher, E. M. C.: Identification and mapping of a novel human gene, HRMT1L1, homologous to the rat protein arginine N-methyltransferase 1 (PRMT1) gene. Mammalian Genome 8:526–529, 1997.
- [70801] 9131.Lin, W. J.; Gary, J. D.; Yang, M. C.; Clarke, S.; Herschman, H.R.: The mammalian intermediate-early TIS21 protein and the leukemia-associated BTG1 protein interact with a protein-arginine N-methyltransferase. J.Biol. Chem. 271: 15034–15044, 1996.
- [70802] 9132.Scott, H. S.; Antonarakis, S. E.; Lalioti, M. D.; Rossier, C.; Silver, P. A.; Henry, M. F.: Identification and characterization of two putative human arginine methyltransferases (HRMT1L1 and HRMT1L2). Genomics 48:330–340, 1998.
- [70803] 9133.Herberg, J. A.; Sgouros, J.; Jones, T.; Copeman, J.; Humphray, S. J.; Sheer, D.; Cresswell, P.; Beck, S.; Trowsdale, J.: Genomic analysis of the Tapasin gene, located close to the TAP loci in the MHC. Europ. J. Immun. 28: 459–467, 1998.
- [70804] 9134.Mayer, W. E.; Klein, J.: Is tapasin a modified Mhc class I molecule? Immunogenetics 53:719–723, 2001.

- [70805] 9135. Michalova, V.; Murray, B. W.; Sultmann, H.; Klein, J.: A contigmap of the Mhc class I genomic region in the zebrafish reveals ancient synteny. *J. Immun.* 164: 5296–5305, 2000.
- [70806] 9136. Ortmann, B.; Copeman, J.; Lehner, P. J.; Sadasivan, B.; Herberg, J. A.; Grandea, A. G.; Riddell, S. R.; Tampe, R.; Spies, T.; Trowsdale, J.; Cresswell, P.: A critical role for tapasin in the assembly and function of multimeric MHC class I–TAP complexes. *Science* 277: 1306–1309, 1997.
- [70807] 9137. Sadasivan, B.; Lehner, P. J.; Ortmann, B.; Spies, T.; Cresswell, P.: Roles for calreticulin and a novel glycoprotein, tapasin, in the interaction of MHC class I molecules with TAP. *Immunity* 5: 103–114, 1996.
- [70808] 9138. Teng, M. S.; Stephens, R.; Du Pasquier, L.; Freeman, T.; Lindquist, J. A.; Trowsdale, J.: A human TAPBP (TAPASIN)-related gene, TAPBP-R. *Europ. J. Immun.* 32: 1059–1068, 2002.
- [70809] 9139. Yabe, T.; Kawamura, S.; Sato, M.; Kashiwase, K.; Tanaka, H.; Ishikawa, Y.; Asao, Y.; Oyama, J.; Tsuruta, K.; Tokunaga, K.; Tadokoro, K.; Juji, T.: A subject with a novel type I bare lymphocyte syndrome has tapasin deficiency due to deletion of 4 exons by Alu-mediated recombination. *Blood* 100: 1496–1498, 2002.

- [70810] 9140.Blank, V.; Kim, M. J.; Andrews, N. C.: Human MAFG is a functional partner for p45 NF-E2 in activating globin gene expression. *Blood* 89:3925–3935, 1997.
- [70811] 9141.Blank, V.; Knoll, J. H. M.; Andrews, N. C.: Molecular characterization and localization of the human MafG gene. *Genomics* 44: 147–149, 1997.
- [70812] 9142.Kimura, K.; Ito, M.; Amano, M.; Chihara, K.; Fukata, Y.; Nakafuku, M.; Yamamori, B.; Feng, J.; Nakano, T.; Okawa, K.; Iwamatsu, A.; Kaibuchi, K.: Regulation of myosin phosphatase by Rho and Rho-associated kinase (Rho-kinase). *Science* 273: 245–248, 1996.
- [70813] 9143.Takahashi, N.; Ito, M.; Tanaka, J.; Nakano, T.; Kaibuchi, K.; Odai, H.; Takemura, K.: Localization of the gene coding for myosin phosphatase, target subunit 1 (MYPT1) to human chromosome 12q15–q21. *Genomics* 44:150–152, 1997.
- [70814] 9144.Ishikawa, S.; Takahashi, T.; Ogawa, M.; Nakamura, Y.: Genomic structure of the human PLCD1 (phospholipase C delta 1) locus on 3p22–p21.3. *Cytogenet. Cell Genet.* 78: 58–60, 1997.
- [70815] 9145.Brown, A. M.; Willi, S. M.; Argyropoulos, G.; Garvey, W. T.: A novel missense mutation, R70W, in the human uncoupling protein 3 gene in a family with type 2 diabetes.

(Abstract) Hum. Mutat. 13: 508only, 1999. Note: Full article online.

- [70816] 9146.Sato, N.; Arai, K.; Masai, H.: Human and Xenopus cDNAs encoding budding yeast Cdc7-related kinases: in vitro phosphorylation of MCM subunits by a putative human homologue of Cdc7. EMBO J. 16: 4340–4351, 1997.
- [70817] 9147.Liu, J.; Dalmau, J.; Szabo, A.; Rosenfeld, M.; Huber, J.; Furneaux, H.: Paraneoplastic encephalomyelitis antigens bind to the AU-rich elements of mRNA. Neurology 45: 544–550, 1995.
- [70818] 9148.Sakai, K.; Gofuku, M.; Kitagawa, Y.; Ogasawara, T.; Hirose, G.; Yamazaki, M.; Koh, C.-S.; Yanagisawa, N.; Steinman, L.: A hippocampal protein associated with paraneoplastic neurologic syndrome and small cell lung carcinoma. Biochem. Biophys. Res. Commun. 199: 1200–1208, 1994.
- [70819] 9149.Van Tine, B. A.; Knops, J. F.; Butler, A.; Deloukas, P.; Shaw, G. M.; King, P. H.: Localization of HuC (ELAVL3) to chromosome 19p13.2 by fluorescence in situ hybridization utilizing a novel tyramide labeling technique. Genomics 53: 296–299, 1998.
- [70820] 9150.Whitmore, T. E.; Maurer, M. F.; Sexson, S.; Raymond, F.; Conklin, D.; Deisher, T. A.: Assignment of fibroblast

growth factor 18 (FGF18) to human chromosome 5q34 by use of radiation hybrid mapping and fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 90: 231–233, 2000.

[70821] 9151. Sherrington, R.; Rogaev, E. I.; Liang, Y.; Rogaeva, E. A.; Levesque, G.; Ikeda, M.; Chi, H.; Lin, C.; Li, G.; Holman, K.; Tsuda, T.; Mar.; and 21 others: Cloning of a gene bearing missense mutations in early-onset familial Alzheimer's disease. *Nature* 375: 754–760, 1995.

[70822] 9152. Verdi, J. M.; Bashirullah, A.; Goldhawk, D. E.; Kubu, C. J.; Jamali, M.; Meakin, S. O.; Lipshitz, H. D.: Distinct human NUMB isoforms regulate differentiation vs. proliferation in the neuronal lineage. *Proc. Nat. Acad. Sci.* 96: 10472–10476, 1999.

[70823] 9153. Zhong, W.; Feder, J. N.; Jiang, M.-M.; Jan, L. Y.; Jan, Y. N.: Asymmetric localization of a mammalian Numb homolog during mouse cortical neurogenesis. *Neuron* 17: 43–53, 1996.

[70824] 9154. Thresher, R. J.; Vitaterna, M. H.; Miyamoto, Y.; Kazantsev, A.; Hsu, D. S.; Petit, C.; Selby, C. P.; Dawut, L.; Smithies, O.; Takahashi, J. S.; Sancar, A.: Role of mouse cryptochrome blue-light photoreceptor in circadian photoresponses. *Science* 282: 1490–1494, 1998.

[70825] 9155. Strehl, S.; LaSalle, J. M.; Lalande, M.: High-resolution



analysis of DNA replication domain organization across an R/G-band boundary. *Molec. Cell. Biol.* 17: 6157–6166, 1997.

- [70826] 9156. Stathakis, D. G.; Lee, D.; Bryant, P. J.: Fine-scale physical map of the 11q21 region surrounding the human DLG2 locus, the gene encoding Chapsyn-110. *Genomics* 54: 186–188, 1998.
- [70827] 9157. Chow, V. T. K.; Lee, S. S.: DENN, a novel human gene differentially expressed in normal and neoplastic cells. *DNA Seq.* 6: 263–273, 1996.
- [70828] 9158. Chow, V. T. K.; Lim, K. M.; Lim, D.: The human DENN gene: genomic organization, alternative splicing, and localization to chromosome 11p11.21–p11.22. *Genome* 41: 543–552, 1998.
- [70829] 9159. Peyrard, M.; Seroussi, E.; Sandberg-Nordqvist, A.-C.; Xie, Y.-G.; Han, F.-Y.; Fransson, I.; Collins, J.; Dunham, I.; Kost-Alimova, M.; Imreh, S.; Dumanski, J. P.: The human LARGE gene from 22q12.3–q13.1 is a new, distinct member of the glycosyltransferase gene family. *Proc. Nat. Acad. Sci.* 96: 598–603, 1999.
- [70830] 9160. Gerlai, R.; Roder, J. C.; Hampson, D. R.: Altered spatial learning and memory in mice lacking the mGluR4 subtype of metabotropic glutamate receptor. *Behav. Neurosci.*

112: 525–532, 1998.

- [70831] 9161. Chatterjee, T. K.; Eapen, A.; Kanis, A. B.; Fisher, R. A.: Genomic organization, 5-prime-flanking region, and chromosomal localization of the human RGS3 gene. *Genomics* 45: 429–433, 1997.
- [70832] 9162. Druey, K. M.; Blumer, K. J.; Kang, V. H.; Kehrl, J. H.: Inhibition of G-protein-mediated MAP kinase activation by a new mammalian gene family. *Nature* 379: 742–746, 1996.
- [70833] 9163. Oettgen, P.; Alani, R. M.; Barcinski, M. A.; Brown, L.; Akbarali, Y.; Boltax, J.; Kunsch, C.; Munger, K.; Liebermann, T. A.: Isolation and characterization of a novel epithelium-specific transcription factor, ESE-1, a member of the Ets family. *Molec. Cell. Biol.* 17: 4419–4433, 1997.
- [70834] 9164. Oettgen, P.; Barcinski, M.; Boltax, J.; Stolt, P.; Akbarali, Y.; Libermann, T. A.: Genomic organization of the human ELF3 (ESE-1/ESX) gene, a member of the Ets transcription factor family, and identification of a functional promoter. *Genomics* 55: 358–362, 1999.
- [70835] 9165. Oettgen, P.; Carter, K. C.; Augustus, M.; Barcinski, M.; Boltax, J.; Kunsch, C.; Libermann, T. A.: The novel epithelial-specific Ets transcription factor gene ESX maps to human chromosome 1q32.1. *Genomics* 45: 456–457,

1997.

- [70836] 9166. Lunn, C. A.; Fan, X.; Dalie, B.; Miller, K.; Zavodny, P. J.; Narula, S. K.; Lundell, D.: Purification of ADAM 10 from bovine spleen as a TNF $\alpha$  convertase. *FEBS Lett.* 400: 333–335, 1997.
- [70837] 9167. Tymms, M. J.; Ng, A. Y. N.; Thomas, R. S.; Schutte, B. C.; Zhou, J.; Eyre, H. J.; Sutherland, G. R.; Seth, A.; Rosenberg, M.; Papas, T.; Debouck, C.; Kola, I.: A novel epithelial-expressed ETS gene, ELF3: human and murine cDNA sequences, murine genomic organization, human mapping to 1q32.2 and expression in tissues and cancer. *Oncogene* 15:2449–2462, 1997.
- [70838] 9168. Wolfsberg, T. G.; Primakoff, P.; Myles, D. G.; White, J. M.: ADAM, a novel family of membrane proteins containing a disintegrin and metalloprotease domain: multipotential functions in cell–cell and cell–matrix interactions. *J. Cell Biol.* 131: 275–278, 1995.
- [70839] 9169. Yamazaki, K.; Mizui, Y.; Tanaka, I.: Radiation hybrid mapping of human ADAM10 gene to chromosome 15. *Genomics* 45: 457–459, 1997.
- [70840] 9170. Choi, D.-S.; Handa, M.; Young, H.; Gordon, A. S.; Diamond, I.; Messing, R. O.: Genomic organization and expression of the mouse equilibrative, nitrobenzylthioino-

sine-sensitive nucleoside transporter1 (ENT1) gene.

Biochem. Biophys. Res. Commun. 277: 200–208, 2000.

[70841] 9171.Coe, I. R.; Griffiths, M.; Young, J. D.; Baldwin, S. A.; Cass, C. E.: Assignment of the human equilibrative nucleoside transporter(hENT1) to 6p21.1–p21.2. Genomics 45: 459–460, 1997.

[70842] 9172.Griffiths, M.; Beaumont, N.; Yao, S. Y. M.; Sundaram, M.; Boumah, C. E.; Davies, A.; Kwong, F. Y. P.; Coe, I.; Cass, C. E.; Young, J.D.; Baldwin, S. A.: Cloning of a human nucleoside transporter implicated in the cellular uptake of adenosine and chemotherapeutic drugs. Nature Med. 3: 89–94, 1997.

[70843] 9173.Hu, S.–I.; Carozza, M.; Klein, M.; Nantermet, P.; Luk, D.; Crowl, R. M.: Human HtrA, an evolutionarily conserved serine protease identified as a differentially expressed gene product in osteoarthritic cartilage. J.Biol. Chem. 273: 34406–34412, 1998.

[70844] 9174.Zumbrunn, J.; Trueb, B.: Localization of the gene for a serineprotease with IGF-binding domain (PRSS11) to human chromosome 10q25.3–q26.2. Genomics 45:461–462, 1997.

[70845] 9175.Zumbrunn, J.; Trueb, B.: Primary structure of a putative serineprotease specific for IGF-binding proteins. FEBS

Lett. 398: 187–192,1996.

- [70846] 9176.D'Esposito, M.; Strazzullo, M.; Cuccurese, M.; Spal-  
luto, C.; Rocchi,M.; D'Urso, M.; Ciccodicola, A.: Identifica-  
tion and assignment ofthe human transient receptor po-  
tential channel 6 gene TRPC6 to chromosome11q21–q22.  
Cytogenet. Cell Genet. 83: 46–47, 1998.
- [70847] 9177.Hofmann, T.; Obukhov, A. G.; Schaefer, M.; Harte-  
neck, C.; Gudermann,T.; Schultz, G.: Direct activation of  
human TRPC6 and TRPC3 channelsby diacylglycerol. Na-  
ture 397: 259–263, 1999.
- [70848] 9178.Garcia, C. K.; Brown, M. S.; Pathak, R. K.; Goldstein,  
J. L.:cDNA cloning of MCT2, a second monocarboxylate  
transporter expressedin different cells than MCT1. J. Biol.  
Chem. 270: 1843–1849, 1995.
- [70849] 9179.Lin, R.–Y.; Vera, J. C.; Chaganti, R. S. K.; Golde, D.  
W.: Humanmonocarboxylate transporter 2 (MCT2) is a  
high affinity pyruvate transporter. J.Biol. Chem. 273:  
28959–28965, 1998.
- [70850] 9180.Ong, O. C.; Hu, K.; Rong, H.; Lee, R. H.; Fung, B.  
K.–K.: Genestructure and chromosome localization of the  
G–gamma–c subunit ofhuman cone G–protein (GNGT2).  
Genomics 44: 101–109, 1997.
- [70851] 9181.Ong, O. C.; Yamane, H. K.; Phan, K. B.; Fong, H. K.;

Bok, D.; Lee, R. H.; Fung, B. K.-K.: Molecular cloning and characterization of the G protein gamma subunit of cone photoreceptors. *J. Biol. Chem.* 270:8495–8500, 1995.

[70852] 9182. Tasheva, E. S.; Pettenati, M.; Von Kap-Her, C.; Conrad, G. W.: Assignment of mimecan gene (OGN) to human chromosome band 9q22 by in situ hybridization. *Cytogenet. Cell Genet.* 88: 326–327, 2000.

[70853] 9183. Park, S. H.; Ryu, S. H.; Suh, P. G.; Kim, H.: Assignment of human PLD2 to chromosome band 17p13.1 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 82: 225 only, 1998.

[70854] 9184. Pucharcos, C.; Fuentes, J.-J.; Casas, C.; de la Luna, S.; Alcantara, S.; Arbones, M. L.; Soriano, E.; Estivill, X.; Prichard, M.: Alu-splice cloning of human intersectin (ITSN), a putative multivalent binding protein expressed in proliferating and differentiating neurons and overexpressed in Down syndrome. *Europ. J. Hum. Genet.* 7: 704–712, 1999.

[70855] 9185. Fumagalli, P.; Accarino, M.; Egeo, A.; Scartezzini, P.; Rappazzo, G.; Pizzuti, A.; Avvantaggiato, V.; Simeone, A.; Arrigo, G.; Zuffardi, O.; Ottolenghi, S.; Taramelli, R.: Human NRD convertase: a highly conserved metalloendopeptidase expressed at specific sites during development and

in adult tissues. *Genomics* 47: 238–245, 1998.

- [70856] 9186.Kanezaki, R.; Toki, T.; Yokoyama, M.; Yomogida, K.; Sugiyama, K.; Yamamoto, M.; Igarashi, K.; Ito, E.: Transcription factor BACH1 is recruited to the nucleus by its novel alternative spliced isoform. *J. Biol. Chem.* 276: 7278–7284, 2001.
- [70857] 9187.Katz, P.; Whalen, G.; Kehrl, J. H.: Differential expression of a novel protein kinase in human B lymphocytes: preferential localization in the germinal center. *J. Biol. Chem.* 269: 16802–16809, 1994.
- [70858] 9188.Pombo, C. M.; Kehrl, J. H.; Sanchez, I.; Katz, P.; Avruch, J.; Zon, L. I.; Woodgett, J. R.; Force, T.; Kyriakis, J. M.: Activation of the SAPK pathway by the human STE20 homologue germinal centre kinase. *Nature* 377:750–754, 1995.
- [70859] 9189.Ren, M.; Zeng, J.; De Lemos–Chiarandini, C.; Rosenfeld, M.; Adesnik, M.; Sabatini, D. D.: In its active form, the GTP-binding protein rab8 interacts with a stress-activated protein kinase. *Proc. Nat. Acad. Sci.* 93: 5151–5155, 1996.
- [70860] 9190.Bonthron, D. T.; Hayward, B. E.; Moran, V.; Strain, L.: Characterization of TH1 and CTSZ, two non-imprinted genes downstream of GNAS1 in chromosome 20q13. *Hum. Genet.* 107: 165–175, 2000.

- [70861] 9191.Santamaria, I.; Velasco, G.; Pendas, A. M.; Fueyo, A.; Lopez-Otin,C.: Cathepsin Z, a novel human cysteine protease with a short propeptidedomain and a unique chromosomal location. J. Biol. Chem. 273: 16816–16823,1998.
- [70862] 9192.Jacquemin, P.; Chen, Z.; Martial, J. A.; Davidson, I.: Genomicstructure and chromosomal mapping of the mouse transcription factorTEF-5 (Tead3) gene. Mammalian Genome 10: 632–634, 1999.
- [70863] 9193.Sparks, A. B.; Hoffman, N. G.; McConnell, S. J.; Fowlkes, D. M.;Kay, B. K.: Cloning of ligand targets: systematic isolation of SH3domain-containing proteins. Nature Biotech. 14: 741–744, 1996.
- [70864] 9194.Blau, N.; Niederwieser, A.; Shmerling, D. H.: Peptiduria presumablycaused by aminopeptidase-P deficiency: a new inborn error of metabolism. J.Inherit. Metab. Dis. 11 (suppl): 240–242, 1988.
- [70865] 9195.Cottrell, G. S.; Hooper, N. M.; Turner, A. J.: Cloning, expression,and characterization of human cytosolic aminopeptidase P: a singlemanganese(II)–dependent enzyme. Biochemistry 39: 15121–15128, 2000.
- [70866] 9196.Sprinkle, T. J.; Caldwell, C.; Ryan, J. W.: Cloning, chromosomalsublocalization of the human soluble



aminopeptidase P gene (XPNPEP1) to 10q25.3 and conservation of the putative proton shuttle and metalligand binding sites with XPNPEP2. Arch. Biochem. Biophys. 378:51–56, 2000.

[70867] 9197. Vanhoof, G.; Goossens, F.; Juliano, M. A.; Juliano, L.; Hendriks, D.; Schatteman, K.; Lin, A. H.; Scharpe, S.: Isolation and sequence analysis of a human cDNA clone (XPNPEPL) homologous to X-prolyl aminopeptidase (aminopeptidase P). Cytogenet. Cell Genet. 78: 275–280, 1997.

[70868] 9198. Vergas Romero, C.; Neudorfer, I.; Mann, K.; Schafer, W.: Purification and amino acid sequence of aminopeptidase P from pig kidney. Europ. J. Biochem. 229: 262–269, 1995.

[70869] 9199. Hegele, R. A.; Connelly, P. W.; Scherer, S. W.; Hanley, A. J. G.; Harris, S. B.; Tsui, L.-C.; Zinman, B.: Paraoxonase-2 gene (PON2) G148 variant associated with elevated fasting plasma glucose in noninsulin-dependent diabetes mellitus. J. Clin. Endocr. Metab. 82: 3373–3377, 1997.

[70870] 9200. Heinecke, J. W.; Lusis, A. J.: Paraoxonase-gene polymorphisms associated with coronary heart disease: support for the oxidative damage hypothesis? (Letter) Am. J. Hum. Genet. 62: 20–24, 1998.

- [70871] 9201. Primo-Parmo, S. L.; Hsu, C.; Law, D. J.; La Du, B. N.: Location and arrangement of three paraoxonase genes: PON1, PON2, and PON3, on human chromosome 7. (Abstract) Am. J. Hum. Genet. 59 (suppl.):A406, 1996.
- [70872] 9202. Primo-Parmo, S. L.; Sorenson, R. C.; Teiber, L.; La Du, B. N.: The human serum paraoxonase/arylesterase gene (PON1) is one member of multigene family. Genomics 33: 498–507, 1996.
- [70873] 9203. Prochazka, M.; Thompson, D. B.; Scherer, S. W.; Tsui, L.-C.; Knowler, W. C.; Bennett, P. H.; Bogardus, C.: Linkage and association of insulin resistance and NIDDM with markers at 7q21.3–q22.1 in the Pima Indians. (Abstract) Diabetes 44 (suppl.): 42A, 1995.
- [70874] 9204. Sanghera, D. K.; Aston, C. E.; Saha, N.; Kamboh, M. I.: DNA polymorphisms in two paraoxonase genes (PON1 and PON2) are associated with the risk of coronary heart disease. Am. J. Hum. Genet. 62: 36–44, 1998.
- [70875] 9205. Chang, H. Y.; Nishitoh, H.; Yang, X.; Ichijo, H.; Baltimore, D.: Activation of apoptosis signal-regulating kinase 1 (ASK1) by the adapter protein Daxx. Science 281: 1860–1863, 1998.
- [70876] 9206. Geleziunas, R.; Xu, W.; Takeda, K.; Ichijo, H.; Greene, W. C.: HIV-1 Nef inhibits ASK1-dependent death signalling

providing a potential mechanism for protecting the infected host cell. *Nature* 410: 834–838, 2001.

[70877] 9207. Ichijo, H.; Nishida, E.; Irie, K.; ten Dijke, P.; Saitoh, M.; Moriguchi, T.; Takagi, M.; Matsumoto, K.; Miyazono, K.; Gotoh, Y.: Induction of apoptosis by ASK1, a mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science* 275: 90–94, 1997.

[70878] 9208. Nishitoh, H.; Saitoh, M.; Mochida, Y.; Takeda, K.; Nakano, H.; Rothe, M.; Miyazono, K.; Ichijo, H.: ASK1 is essential for JNK/SAPK activation by TRAF2. *Molec. Cell* 2: 389–395, 1998.

[70879] 9209. Fuchs, M.; Muller, T.; Lerch, M. M.; Ullrich, A.: Association of human protein-tyrosine phosphatase kappa with members of the armadillo family. *J. Biol. Chem.* 271: 16712–16719, 1996.

[70880] 9210. Yang, Y.; Gil, M. C.; Choi, E. Y.; Park, S. H.; Pyun, K. H.; Ha, H.: Molecular cloning and chromosomal localization of a human gene homologous to the murine R-PTP-kappa, a receptor-type protein tyrosine phosphatase. *Gene* 186: 77–82, 1997.

[70881] 9211. Zhang, Y.; Siebert, R.; Matthiesen, P.; Yang, Y.; Ha, H.; Schlegelberger, B.: Cytogenetical assignment and physical mapping of the human R-PTP-kappa gene (PTPRK) to

the putative tumor suppressor gene region 6q22.2–q22.3  
Genomics 51:309–311, 1998.

- [70882] 9212.Hospital, V.; Prat, A.; Joulie, C.; Cherif, D.; Day, R.; Cohen,P.: Human and rat testis express two mRNA species encoding variantsof NRD convertase, a metalloendopeptidase of the insulinase family. Biochem.J. 327: 773–779, 1997.
- [70883] 9213.Pierotti, A. R.; Prat, A.; Chesneau, V.; Gaudoux, F.; Leseney,A.–M.; Foulon, T.; Cohen, P.: N–Arginine dibasic convertase, a metalloendopeptidaseas a prototype of a class of processing enzymes. Proc. Nat. Acad.Sci. 91: 6078–6082, 1994.
- [70884] 9214.Tomasetto, C.; Regnier, C.; Moog–Lutz, C.; Mattei, M. G.; Chenard,M. P.; Lidereau, R.; Basset, P.; Rio, M. C.: Identification of fournovel human genes amplified and overexpressed in breast carcinomaand localized to the q11–q21.3 region of chromosome 17. Genomics 28:367–376, 1995.
- [70885] 9215.Xu, Z.–H.; Otterness, D. M.; Freimuth, R. R.; Carlini, E. J.;Wood, T. C.; Mitchell, S.; Moon, E.; Kim, U.–J.; Xu, J.–P.; Siciliano,M. J.; Weinshilboum, R. M.: Human 3–prime–phosphoadenosine 5–prime–phosphosulfatesynthetase 1 (PAPSS1) and

PAPSS2: gene cloning, characterization and chromosomal localization. *Biochem. Biophys. Res. Commun.* 268: 437–444, 2000.

- [70886] 9216. Chalmers, I. J.; Hofler, H.; Atkinson, M. J.: Mapping of a cadherin gene cluster to a region of chromosome 5 subject to frequent allelic loss in carcinoma. *Genomics* 57: 160–163, 1999.
- [70887] 9217. Suzuki, S.; Sano, K.; Tanihara, H.: Diversity of the cadherin family: evidence for eight new cadherins in nervous tissue. *Cell Regul.* 2: 261–270, 1991.
- [70888] 9218. Shimoyama, Y.; Gotoh, M.; Terasaki, T.; Kitajima, M.; Hirohashi, S.: Isolation and sequence analysis of human cadherin-6 complementary DNA for the full coding sequence and its expression in human carcinoma cells. *Cancer Res.* 55: 2206–2211, 1995.
- [70889] 9219. Anderson, L. V. B.; Davison, K.; Moss, J. A.; Young, C.; Cullen, M. J.; Walsh, J.; Johnson, M. A.; Bashir, R.; Britton, S.; Keers, S.; Argov, Z.; Mahjneh, I.; Fougere, F.; Beckmann, J. S.; Bushby, K. M. D.: Dysferlin is a plasma membrane protein and is expressed early in human development. *Hum. Molec. Genet.* 8: 855–861, 1999.
- [70890] 9220. Gong, B.; Almasan, A.: Genomic organization and transcriptional regulation of human Apo2/TRAIL gene.

Biochem. Biophys. Res. Commun. 278:747–752, 2000.

[70891] 9221.Ohira, M.; Seki, N.; Nagase, T.; Ishikawa, K.; Nomura, N.; Ohara,O.: Characterization of a human homolog (BACH1) of the mouse Bach1gene encoding a BTB–basic leucine zipper transcription factor andits mapping to chromosome 21q22.1. Genomics 47: 300–306, 1998.

[70892] 9222.Oyake, T.; Itoh, K.; Motohashi, H.; Hayashi, N.; Hoshino, H.; Nishizawa,M.; Yamamoto, M.; Igarashi, K.: Bach proteins belong to a novel familyof BTB–basic leucine zipper transcription factors that interact withMafK and regulate transcription through the NF–E2 site. Molec. Cell.Biol. 16: 6083–6095, 1996.

[70893] 9223.Johnson, K. R.; Smith, L.; Johnson, D. K.; Rhodes, J.; Rinchik,E. M.; Thayer, M.; Lewis, E. J.: Mapping of the ARIX homeodomaingene to mouse chromosome 7 and human chromosome 11q13. Genomics 33:527–531, 1996.

[70894] 9224.Merscher, S.; Bekri, S.; de Leeuw, B.; Pedeutour, F.; Grosgeorge,J.; Show, S. T. D.; Mullenbach, R.; et al.: A 5,5 megabase high resolutionintegrated map of distal 11q13. Genomics 39: 340–347, 1997.

[70895] 9225.Morin, X.; Cremer, H.; Hirsch, M.–R.; Kapur, R. P.; Goridis, C.;Brunet, J.–F.: Defects in sensory and autonomic ganglia and absenceof locus coeruleus in mice deficient

for the homeobox gene Phox2a. *Neuron* 18:411–423, 1997.

[70896] 9226. Pattyn, A.; Morin, X.; Goridis, C.; Brunet, J. F.: Expression and interaction of the two closely related homeobox genes PHOX2a and PHOX2b during neurogenesis. *Development* 124: 4065–4075, 1997.

[70897] 9227. Zellmer, E.; Zhang, Z.; Greco, D.; Rhodes, J.; Cassel, S.; Lewis, E. J.: A homeodomain protein selectively expressed in noradrenergic tissue regulates transcription of neurotransmitter biosynthetic genes. *J. Neurosci.* 15: 8109–8120, 1995.

[70898] 9228. Ghanshani, S.; Coleman, M.; Gustavsson, P.; Wu, A. C.-L.; Gargus, J. J.; Gutman, G. A.; Dahl, N.; Mohrenweiser, H.; Chandy, K. G.: Human calcium-activated potassium channel gene KCNN4 maps to chromosome 19q13.2 in the region deleted in Diamond-Blackfan anemia. *Genomics* 51:160–161, 1998.

[70899] 9229. Ishii, T. M.; Silvia, C.; Hirschberg, B.; Bond, C. T.; Adelman, J. P.; Maylie, J.: A human intermediate conductance calcium-activated potassium channel. *Proc. Nat. Acad. Sci.* 94: 11651–11656, 1997.

[70900] 9230. Joiner, W. J.; Wang, L.-Y.; Tang, M. D.; Kaczmarek, L. K.: hSK4, a member of a novel subfamily of calcium-acti-

vated potassium channels. Proc.Nat. Acad. Sci. 94: 11013–11018, 1997.

[70901] 9231.Logsdon, N. J.; Kang, J.; Togo, J. A.; Christian, E. P.; Aiyar,J.: A novel gene, hKCa4, encodes the calcium-activated potassiumchannel in human T lymphocytes. J. Biol. Chem. 272: 32723–32726,1997.

[70902] 9232.Matsuoka, H.; Iwata, N.; Ito, M.; Shimoyama, M.; Nagata, A.; Chihara,K.; Takai, S.; Matsui, T.: Expression of a kinase-defective Eph-likereceptor in the normal human brain. Biochem. Biophys. Res. Commun. 235:487–492, 1997.

[70903] 9233.Tang, X. X.; Zhao, H.; Robinson, M. E.; Cohen, B.; Cnaan, A.; London,W.; Cohn, S. L.; Cheung, N.-K. V.; Brodeur, G. M.; Evans, A. E.; Ikegaki,N.: Implications of EPHB6, EFNB2, and EFNB3 expressions in humanneuroblastoma. Proc. Nat. Acad. Sci. 97: 10936–10941, 2000.

[70904] 9234.Meyers, R.; Cantley, L. C.: Cloning and characterization of awortmannin-sensitive human phosphatidylinositol 4-kinase. J. Biol.Chem. 272: 4384–4390, 1997.

[70905] 9235.Nakanishi, S.; Catt, K. J.; Balla, T.: A wortmannin-sensitivephosphatidylinositol 4-kinase that regulates hormone-sensitive poolsof inositolphospholipids. Proc. Nat. Acad. Sci. 92: 5317–5321, 1995.



- [70906] 9236.Saito, T.; Seki, N.; Ishii, H.; Ohira, M.; Hayashi, A.; Kozuma,S.; Hori, T.: Complementary DNA cloning and chromosomal mapping of a novel phosphatidylinositol kinase gene. DNA Res. 4: 301–305, 1997.
- [70907] 9237.Suzuki, K.; Hirano, H.; Okutomi, K.; Suzuki, M.; Kuga, Y.; Fujiwara,T.; Kanemoto, N.; Isono, K.; Horie, M.: Identification and characterization of a novel human phosphatidylinositol 4–kinase. DNA Res. 4: 273–280,1997.
- [70908] 9238.Miura, Y.; Miyake, K.; Yamashita, Y.; Shimazu, R.; Copeland, N.G.; Gilbert, D. J.; Jenkins, N. A.; Inazawa, J.; Abe, T.; Kimoto,M.: Molecular cloning of a human RP105 homologue and chromosomal localization of the mouse and human RP105 genes (Ly64 and LY64). Genomics 38:299–304, 1996.
- [70909] 9239.Miura, Y.; Shimazu, R.; Miyake, K.; Akashi, S.; Ogata, H.; Yamashita,Y.; Narisawa, Y.; Kimoto, M.: RP105 is associated with MD–1 and transmits an activation signal in human B cells. Blood 92: 2815–2822, 1998.
- [70910] 9240.Castrop, J.; van Norren, K.; Clevers, H.: A gene family of HMG–box transcription factors with homology to TCF–1. Nucleic Acids Res. 20:611 only, 1992.
- [70911] 9241.Duval, A.; Busson–Leconiat, M.; Berger, R.; Hamelin, R.: Assignment of the TCF–4 (TCF7L2) to human chromo–

some band 10q25.3. Cytogenet.Cell Genet. 88: 264–265, 2000.

[70912] 9242.Duval, A.; Gayet, J.; Zhou, X.–P.; Iacopetta, B.; Thomas, G.; Hamelin,R.: Frequent frameshift mutations of the TCF–4 gene in colorectal cancers with microsatellite instability. Cancer Res. 59: 4213–4215,1999.

[70913] 9243.Duval, A.; Rolland, S.; Tubacher, E.; Bui, H.; Thomas, G.; Hamelin,R.: The human T–cell transcription factor–4 gene: structure, extensive characterization of alternative splicings, and mutational analysis in colorectal cancer cell lines. Cancer Res. 60: 3872–3879, 2000.

[70914] 9244.Korinek, V.; Barker, N.; Moerer, P.; van Donselaar, E.; Huls, G.; Peters, P. J.; Clevers, H.: Depletion of epithelial stem–cell compartments in the small intestine of mice lacking Tcf–4. Nature Genet. 19:379–383, 1998.

[70915] 9245.Ando, A.; Kikuti, Y. Y.; Abe, K.; Shigenari, A.; Kawata, H.; Ikemura,T.; Kimura, M.; Inoko, H.: cDNA cloning, northern hybridization, and mapping analysis of a putative GDS–related protein gene at the centromeric ends of the human and mouse MHC regions. Immunogenetics 49:354–356, 1999.

[70916] 9246.Isomura, M.; Okui, K.; Fujiwara, T.; Shin, S.; Nakamura, Y.: Isolation and mapping of RAB2L, a human cDNA

that encodes a protein homologous to RalGDS. *Cytogenet. Cell Genet.* 74: 263–265, 1996.

- [70917] 9247. Pirozzi, G.; McConnell, S. J.; Uveges, A. J.; Carter, J. M.; Sparks, A. B.; Kay, B. K.; Fowlkes, D. M.: Identification of novel human WW domain-containing proteins by cloning of ligand targets. *J. Biol. Chem.* 272: 14611–14616, 1997.
- [70918] 9248. Chambers, J. A.; Gardner, E.; Hauptmann, R.; Ponder, B. A.; Mulligan, L. M.: TaqI polymorphisms at the annexin VIII locus (ANX8). *Hum. Molec. Genet.* 1: 550 only, 1992.
- [70919] 9249. Chang, K.-S.; Wang, G.; Freireich, E. J.; Daly, M.; Naylor, S. L.; Trujillo, J. M.; Stass, S. A.: Specific expression of the annexin VIII gene in acute promyelocytic leukemia. *Blood* 79: 1802–1810, 1992.
- [70920] 9250. Fernandez, M. P.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Morgan, R. O.: The genetic origin of mouse annexin VIII. *Mammalian Genome* 9: 8–14, 1998.
- [70921] 9251. Sarkar, A.; Yang, P.; Fan, Y.-H.; Mu, Z. M.; Hauptmann, R.; Adolf, G. R.; Stass, S. A.; Chang, K.-S.: Regulation of the expression of annexin VIII in acute promyelocytic leukemia. *Blood* 84: 279–286, 1994.
- [70922] 9252. Waterham, H. R.; Koster, J.; Romeijn, G. J.; Hennekam, R. C. M.; Vreken, P.; Andersson, H. C.; FitzPatrick,

D. R.; Kelley, R. I.; Wanders, R. J. A.: Mutations in the 3-beta-hydroxysterol delta-24-reductase gene cause desmosterolosis, an autosomal recessive disorder of cholesterol biosynthesis. *Am. J. Hum. Genet.* 69: 685-694, 2001.

[70923] 9253. Lechner, C.; Zahalka, M. A.; Giot, J.-F.; Moller, N. P. H.; Ullrich, A.: ERK6, a mitogen-activated protein kinase involved in C2C12 myoblast differentiation. *Proc. Nat. Acad. Sci.* 93: 4355-4359, 1996.

[70924] 9254. Li, Z.; Jiang, Y.; Ulevitch, R. J.; Han, J.: The primary structure of p38-gamma: a new member of p38 group of MAP kinases. *Biochem. Biophys. Res. Commun.* 228: 334-340, 1996.

[70925] 9255. Hibi, M.; Lin, A.; Smeal, T.; Minden, A.; Karin, M.: Identification of an oncoprotein- and UV-responsive protein kinase that binds and potentiates the c-Jun activation domain. *Genes Dev.* 7: 2135-2148, 1993.

[70926] 9256. Kallunki, T.; Su, B.; Tsigelny, I.; Sluss, H. K.; Derijard, B.; Moore, G.; Davis, R.; Karin, M.: JNK2 contains a specificity-determining region responsible for efficient c-Jun binding and phosphorylation. *Genes Dev.* 8: 2996-3007, 1994.

[70927] 9257. Sluss, H. K.; Barrett, T.; Derijard, B.; Davis, R. J.:

Signaltransduction by tumor necrosis factor mediated by JNK protein kinases. *Molec.Cell Biol.* 14: 8376–8384, 1994.

[70928] 9258.Mohit, A. A.; Martin, J. H.; Miller, C. A.: p49(3F12) kinase:a novel MAP kinase expressed in a subset of neurons in the human nervoussystem. *Neuron* 14: 67–78, 1995.

[70929] 9259.Yang, D. D.; Kuan, C. Y.; Whitmarsh, A. J.; Rincon, M.; Zheng,T. S.; Davis, R. J.; Rakic, P.; Flavell, R. A.: Absence of excitotoxicity–inducedapoptosis in the hippocampus of mice lacking the Jnk3 gene. *Nature* 389:865–870, 1997.

[70930] 9260.Yoshida, S.; Fukino, K.; Harada, H.; Nagai, H.; Imoto, I.; Inazawa,J.; Takahashi, H.; Teramoto, A.; Emi, M.: The c–Jun NH2–terminalkinase 3 (JNK3) gene: genomic structure, chromosomal assignment, andloss of expression in brain tumors. *J. Hum. Genet.* 46: 182–187,2001.

[70931] 9261.Carpenter, N. J.; Filipovich, A.; Blaese, R. M.; Carey, T. L.;Berkel, A. I.: Variable immunodeficiency with abnormal condensationof the heterochromatin of chromosomes 1, 9, and 16. *J. Pediat.* 112:757–760, 1988.

[70932] 9262.Bunger, M. K.; Wilsbacher, L. D.; Moran, S. M.; Clendenin, C.;Radcliffe, L. A.; Hogenesch, J. B.; Simon, M. C.;

Takahashi, J. S.;Bradfield, C. A.: Mop3 is an essential component of the master circadianpacemaker in mammals. Cell 103: 1009–1017, 2000.

[70933] 9263.Ikeda, M.; Nomura, M.: cDNA cloning and tissue-specific expressionof a novel basic helix–loop–helix/PAS protein (BMAL1) and identificationof alternatively spliced variants with alternative translation initiationsite usage. Biochem. Biophys. Res. Commun. 233: 258–264, 1997.

[70934] 9264.Shaw, P. J.; Tononi, G.; Greenspan, R. J.; Robinson, D. F.: Stressresponse genes protect against lethal effects of sleep deprivationin Drosophila. Nature 417: 287–291, 2002.

[70935] 9265.Wolting, C. D.; McGlade, C. J.: Cloning and chromosomal localizationof a new member of the bHLH/PAS transcription factor family. MammalianGenome 9: 463–468, 1998.

[70936] 9266.Faucheu, C.; Diu, A.; Chan, A. W. E.; Blanchet, A.–M.; Miossec,C.; Herve, F.; Collard–Dutilleul, V.; Gu, Y.; Aldape, R. A.; Lippke,J. A.; Rocher, C.; Su, M. S.–S.; Livingston, D. J.; Hercend, T.; Lalanne,J.–L.: A novel human protease similar to the interleukin–1–beta convertingenzyme induces apoptosis in transfected cells. EMBO J. 14: 1914–1922,1995.

- [70937] 9267.Kamens, J.; Paskind, M.; Hugunin, M.; Talanian, R. V.; Allen, H.; Banach, D.; Bump, N.; Hackett, M.; Johnston, C. G.; Li, P.; Mankovich, J. A.; Terranova, M.; Ghayur, T.: Identification and characterization of ICH-2, a novel member of the interleukin-1-beta-converting enzyme family of cysteine proteases. *J. Biol. Chem.* 270: 15250-15256, 1995.
- [70938] 9268.Munday, N. A.; Vaillancourt, J. P.; Ali, A.; Casano, F. J.; Miller, D. K.; Molineaux, S. M.; Yamin, T.-T.; Yu, V. L.; Nicholson, D. W.: Molecular cloning and pro-apoptotic activity of ICE(rel)II and ICE(rel)III, members of the ICE/CED-3 family of cysteine proteases. *J. Biol. Chem.* 270:15870-15876, 1995.
- [70939] 9269.Wang, S.; Miura, M.; Jung, Y.; Zhu, H.; Li, E.; Yuan, J.: Murine caspase-11, an ICE-interacting protease, is essential for the activation of ICE. *Cell* 92: 501-509, 1998.
- [70940] 9270.Liburd, N.; Ghosh, M.; Riazuddin, S.; Naz, S.; Khan, S.; Ahmed, Z.; Riazuddin, S.; Liang, Y.; Menon, P. S. N.; Smith, T.; Smith, A.C. M.; Chen, K.-S.; Lupski, J. R.; Wilcox, E. R.; Potocki, L.; Friedman, T. B.: Novel mutations of MYO15A associated with profound deafness in consanguineous families and moderately severe hearing loss in a patient with Smith-Magenis syndrome. *Hum. Genet.* 109:

535–541, 2001.

- [70941] 9271. Carlomagno, F.; Chang–Claude, J.; Dunning, A. M.; Ponder, B. A.J.: Determination of the frequency of the common 675del5 Nijmegenbreakage syndrome mutation in the German population: no association with risk of breast cancer. *Genes Chromosomes Cancer* 25: 393–395, 1999.
- [70942] 9272. Kleier, S.; Herrmann, M.; Wittwer, B.; Varon, R.; Reis, A.; Horst, J.: Clinical presentation and mutation identification in the NBS1 gene in a boy with Nijmegen breakage syndrome. *Clin. Genet.* 57: 384–387, 2000.
- [70943] 9273. Lim, D.–S.; Kim, S.–T.; Xu, B.; Maser, R. S.; Lin, J.; Petrini, J. H. J.; Kastan, M. B.: ATM phosphorylates p95/nbs1 in an S–phase checkpoint pathway. *Nature* 404: 613–617, 2000.
- [70944] 9274. Lombard, D. B.; Guarente, L.: Nijmegen breakage syndrome disease protein and MRE11 at PML nuclear bodies and meiotic telomeres. *Cancer Res.* 60: 2331–2334, 2000.
- [70945] 9275. Maser, R. S.; Zinkel, R.; Petrini, J. H. J.: An alternative mode of translation permits production of a variant NBS1 protein from the common Nijmegen breakage syndrome allele. *Nature Genet.* 27: 417–421, 2001.



- [70946] 9276.Matsuura, S.; Tauchi, H.; Nakamura, A.; Kondo, N.; Sakamoto, S.; Endo, S.; Smeets, D.; Solder, B.; Belohradsky, B. H.; Der Kaloustian, V. M.; Oshimura, M.; Isomura, M.; Nakamura, Y.; Komatsu, K.: Positional cloning of the gene for Nijmegen breakage syndrome. *Nature Genet.* 19:179–181, 1998.
- [70947] 9277.Skelly, R. H.; Korbonits, M.; Grossman, A.; Besser, G. M.; Monson, J. P.; Geddes, J. F.; Burrin, J. M.: Expression of the pituitary transcription factor Ptx-1, but not that of the trans-activating factor Prop-1, is reduced in human corticotroph adenomas and is associated with decreased alpha-subunit secretion. *J. Clin. Endocr. Metab.* 85:2537–2542, 2000.
- [70948] 9278.Morgan, R. O.; Bell, D. W.; Testa, J. R.; Fernandez, M.-P.: Human annexin 31 genetic mapping and origin. *Gene* 227: 33–38, 1999.
- [70949] 9279.Morgan, R. O.; Fernandez, M.-P.: Expression profile and structural divergence of novel human annexin 31. *FEBS Lett.* 434: 300–304, 1998.
- [70950] 9280.Nadler, M. J. S.; Hermosura, M. C.; Inabe, K.; Perraud, A.-L.; Zhu, Q.; Stokes, A. J.; Kurosaki, T.; Kinet, J.-P.; Penner, R.; Scharenberg, A. M.; Fleig, A.: LTRPC7 is a Mg-ATP-regulated divalent cation channel required for cell vi-

ability. *Nature* 411: 590–595, 2001. Note: Erratum *Nature* 412:660 only, 2001.

[70951] 9281. Runnels, L. W.; Yue, L.; Clapham, D. E.: TRP-PLIK, a bifunctional protein with kinase and ion channel activities. *Science* 291: 1043–1047, 2001.

[70952] 9282. Schlingmann, K. P.; Weber, S.; Peters, M.; Nejsum, L. N.; Vitzthum, H.; Klingel, K.; Kratz, M.; Haddad, E.; Ristoff, E.; Dinour, D.; Syrrou, M.; Nielsen, S.; Sassen, M.; Waldegger, S.; Seyberth, H. W.; Konrad, M.: Hypomagnesemia with secondary hypocalcemia is caused by mutations in TRPM6, a new member of the TRPM gene family. *Nature Genet.* 31:166–170, 2002.

[70953] 9283. Walder, R. Y.; Shalev, H.; Brennan, T. M. H.; Carmi, R.; Elbedour, K.; Scott, D. A.; Hanauer, A.; Mark, A. L.; Patil, S.; Stone, E. M.; Sheffield, V. C.: Familial hypomagnesemia maps to chromosome 9q, not to the X chromosome: genetic linkage mapping and analysis of a balanced translocation breakpoint. *Hum. Molec. Genet.* 6: 1491–1497, 1997.

[70954] 9284. Brohmann, H.; Pinnecke, S.; Hoyer-Fender, S.: Identification and characterization of new cDNAs encoding outer dense fiber proteins of rat sperm. *J. Biol. Chem.* 272: 10327–10332, 1997.

- [70955] 9285.Scott, A. F.: Personal Communication. Baltimore, Md. 9/10/1997.
- [70956] 9286.Shao, X.; Murthy, S.; Demetrick, D. J.; van der Hoorn, F. A.:Human outer dense fiber gene, ODF2, localizes to chromosome 9q34. Cytogenet.Cell Genet. 83: 221–223, 1998.
- [70957] 9287.Trachtulec, Z.; Hamvas, R. M. J.; Forejt, J.; Lehrach, H. R.; Vincek,V.; Klein, J.: Linkage of TATA-binding protein and proteasome subunitC5 genes in mice and humans reveals syntenic conserved between mammals and invertebrates. Genomics 44: 1–7, 1997.
- [70958] 9288.Calabrese, G.; Sallese, M.; Stornaiuolo, A.; Stuppia, L.; Palka,G.; De Blasi, A.: Chromosome mapping of the human arrestin (SAG),beta-arrestin 2 (ARRB2), and beta-adrenergic receptor kinase 2 (ADRBK2)genes. Genomics 23: 286–288, 1994.
- [70959] 9289.Porteous, S.; Torban, E.; Cho, N.-P.; Cunliffe, H.; Chua, L.;McNoe, L.; Ward, T.; Souza, C.; Gus, P.; Giugliani, R.; Sato, T.;Yun, K.; Favor, J.; Sicotte, M.; Goodyer, P.; Eccles, M.: Primaryrenal hypoplasia in humans and mice with PAX2 mutations: evidenceof increased apoptosis in fetal kidneys of Pax2(1Neu) +/- mutant mice. Hum.Molec. Genet. 9: 1–11, 2000.

- [70960] 9290.Johnson, L.; Mercer, K.; Greenbaum, D.; Bronson, R. T.; Crowley,D.; Tuveson, D. A.; Jacks, T.: Somatic activation of the K-ras oncogene causes early onset lung cancer in mice. *Nature* 410: 1111–1116, 2001.
- [70961] 9291.Kozma, S. C.; Bogaard, M. E.; Buser, K.; Saurer, S. M.; Bos, J.L.; Groner, B.; Hynes, N. E.: The human c-Kirsten ras gene is activated by a novel mutation in codon 13 in the breast carcinoma cell line MDA-MB231. *Nucleic Acids Res.* 15: 5963–5971, 1987.
- [70962] 9292.Laghi, L.; Orbetegli, O.; Bianchi, P.; Zerbi, A.; Di Carlo, V.; Boland, C. R.; Malesci, A.: Common occurrence of multiple K-RAS mutations in pancreatic cancers with associated precursor lesions and in biliary cancers. *Oncogene* 21: 4301–4306, 2002.
- [70963] 9293.Lee, K.-H.; Lee, J.-S.; Suh, C.; Kim, S.-W.; Kim, S.-B.; Lee, J.-H.; Lee, M.-S.; Park, M.-Y.; Sun, H.-S.; Kim, S.-H.: Clinicopathologic significance of the K-ras gene codon 12 point mutation in stomach cancer: an analysis of 140 cases. *Cancer* 75: 2794–2801, 1995.
- [70964] 9294.Liu, E.; Hjelle, B.; Morgan, R.; Hecht, F.; Bishop, J. M.: Mutations of the Kirsten-ras proto-oncogene in human preleukaemia. *Nature* 330:186–188, 1987.
- [70965] 9295.McBride, O. W.; Swand, D. C.; Tronick, S. R.; Gol, R.;

Klimanis,D.; Moore, D. E.; Aaronson, S. A.: Regional chromosomal localization of N-ras, K-ras-1, K-ras-2 and myb oncogenes in human cells. *Nucleic Acids Res.* 11: 8221-8236, 1983.

[70966] 9296. McCoy, M. S.; Toole, J. J.; Cunningham, J. M.; Chang, E. H.; Lowy, D. R.; Weinberg, R. A.: Characterization of a human colon/lung carcinoma oncogene. *Nature* 302: 79-81, 1983.

[70967] 9297. McGrath, J. P.; Capon, D. J.; Smith, D. H.; Chen, E. Y.; Seeburg, P. H.; Goeddel, D. V.; Levinson, A. D.: Structure and organization of the human Ki-ras proto-oncogene and a related processed pseudogene. *Nature* 304:501-506, 1983.

[70968] 9298. Motojima, K.; Urano, T.; Nagata, Y.; Shiku, H.; Tsurifune, T.; Kanematsu, T.: Detection of point mutations in the Kirsten-ras oncogene provides evidence for the multicentricity of pancreatic carcinoma. *Ann. Surg.* 217: 138-143, 1993.

[70969] 9299. Muller, R.; Slamon, D. J.; Adamson, E. D.; Tremblay, J. M.; Muller, D.; Cline, M. J.; Verma, I. M.: Transcription of c-onc genes c-ras(Ki) and c-fms during mouse development. *Molec. Cell. Biol.* 3: 1062-1069, 1983.

[70970] 9300. Nakano, H.; Yamamoto, F.; Neville, C.; Evans, D.;

Mizuno, T.;Perucho, M.: Isolation of transforming sequences of two human lungcarcinomas: structural and functional analysis of the activated c-K-rasoncogenes. Proc. Nat. Acad. Sci. 81: 71–75, 1984.

[70971] 9301.O'Brien, S. J.; Nash, W. G.; Goodwin, J. L.; Lowry, D. R.; Chang,E. H.: Dispersion of the ras family of transforming genes to fourdifferent chromosomes in man. Nature 302: 839–842, 1983.

[70972] 9302.O'Connell, P.; Leppert, M.; Hoff, M.; Kumlin, E.; Thomas, W.;Cai, G.; Law, M.; White, R.: A linkage map for human chromosome 12.(Abstract) Am. J. Hum. Genet. 37: A169 only, 1985.

[70973] 9303.Ohashi, H.; Ishikiriya, S.; Fukushima, Y.: New diagnostic methodfor Pallister–Killian syndrome: detection of i(12p) in interphasenuclei of buccal mucosa by fluorescence in situ hybridization. Am.J. Med. Genet. 45: 123–128, 1993.

[70974] 9304.Otori, K.; Oda, Y.; Sugiyama, K.; Hasebe, T.; Mukai, K.; Fujii,T.; Tajiri, H.; Yoshida, S.; Fukushima, S.; Esumi, H.: High frequencyof K–ras mutations in human colorectal hyperplastic polyps. Gut 40:660–663, 1997.

[70975] 9305.Peltomaki, P.; Knuutila, S.; Ritvanen, A.; Kaitila, I.; de laChapelle, A.: Pallister–Killian syndrome: cytogenetic and

molecular studies. Clin. Genet. 31: 399–405, 1987.

[70976] 9306. Pfeifer, G. P.: A new verdict for an old convict. Nature Genet. 29:3–4, 2001.

[70977] 9307. Porta, M.; Malats, N.; Jarrod, M.; Grimalt, J. O.; Rifa, J.; Carrato, A.; Guarner, L.; Salas, A.; Santiago-Silva, M.; Corominas, J. M.; Andreu, M.; Real, F. X.: Serum concentrations of organochlorine compounds and K-ras mutations in exocrine pancreatic cancer. Lancet 354: 2125–2129, 1999.

[70978] 9308. Pulciani, S.; Santos, E.; Lauver, A. V.; Long, L. K.; Aaronson, S. A.; Barbacid, M.: Oncogene in solid human tumors. Nature 300:539–542, 1982.

[70979] 9309. Rodenhuis, S.; van de Wetering, M. L.; Mooi, W. J.; Evers, S. G.; van Zandwijk, N.; Bos, J. L.: Mutational activation of the K-RAS oncogene: a possible pathogenetic factor in adenocarcinoma of the lung. New Eng. J. Med. 317: 929–935, 1987.

[70980] 9310. Bertazzoni, U.; Scovassi, A. I.; Brun, G. M.: Chick-embryo DNA polymerase gamma: identity of gamma-polymerases purified from nuclei and mitochondria. Europ. J. Biochem. 81: 237–248, 1977.

[70981] 9311. Chang, L.-S.; Zhao, L.; Zhu, L.; Chen, M.-L.; Lee, M. Y. W. T.: Structure of the gene for the catalytic subunit of

human DNA polymerase delta (POLD1). Genomics 28: 411–419, 1995.

[70982] 9312. Chung, D. W.; Zhang, J.; Tan, C.-K.; Davie, E. W.; So, A. G.; Downey, K. M.: Primary structure of the catalytic subunit of human DNA polymerase delta and chromosomal location of the gene. Proc. Nat. Acad. Sci.

88:11197–11201, 1991.

[70983] 9313. Dresler, S. L.; Gowans, B. J.; Robinson-Hill, R. M.; Hunting, D. J.: Involvement of DNA polymerase delta in DNA repair synthesis in human fibroblasts at late times after ultraviolet irradiation. Biochemistry 27:6379–6383, 1988.

[70984] 9314. Goldsby, R. E.; Singh, M.; Preston, B. D.: Mouse DNA polymerase delta gene (Pold1) maps to chromosome 7. Mammalian Genome 9: 92–93, 1998.

[70985] 9315. Kamath-Loeb, A. S.; Johansson, E.; Burgers, P. M. J.; Loeb, L. A.: Functional interaction between the Werner syndrome protein and DNA polymerase delta. Proc. Nat. Acad. Sci. 97: 4603–4608, 2000.

[70986] 9316. Kemper, R. R.; Ahn, E. R.; Zhang, P.; Lee, M. Y. W. T.; Rabin, M.: Human DNA polymerase delta gene maps to region 19q13.3–q13.4 by in situ hybridization. Genomics 14: 205–206, 1992.



- [70987] 9317.Lee, M. Y. W. T.: Isolation of multiple forms of DNA polymerase  $\delta$ : evidence of proteolytic modification during isolation. *Biochemistry* 27:5188–5193, 1988.
- [70988] 9318.Lee, M. Y. W. T.; Toomey, N. L.: Human placental DNA polymerase  $\delta$ : identification of a 170-kilodalton polypeptide by activity staining and immunoblotting. *Biochemistry* 26: 1076–1085, 1987.
- [70989] 9319.Lestienne, P.: Evidence for a direct role of the DNA polymerase  $\gamma$  in the replication of the human mitochondrial DNA in vitro. *Biochem. Biophys. Res. Commun.* 146: 1146–1153, 1987.
- [70990] 9320.Nishida, C.; Reinhard, P.; Linn, S.: DNA repair synthesis in human fibroblasts requires DNA polymerase  $\delta$ . *J. Biol. Chem.* 263:501–510, 1988.
- [70991] 9321.Syvaoja, J.; Linn, S.: Characterization of a large form of DNA polymerase  $\delta$  from HeLa cells that is insensitive to proliferating cell nuclear antigen. *J. Biol. Chem.* 264: 2489–2497, 1989.
- [70992] 9322.Syvaoja, J.; Suomensari, S.; Nishida, C.; Goldsmith, J. S.; Chui, G. S. J.; Jain, S.; Linn, S.: DNA polymerases  $\alpha$ ,  $\delta$ , and  $\epsilon$ : three distinct enzymes from HeLa cells. *Proc. Nat. Acad. Sci.* 87:6664–6668, 1990.
- [70993] 9323.Yang, C.-L.; Chang, L.-S.; Zhang, P.; Hao, H.; Zhu,

L.; Toomey, N. L.; Lee, M. Y. W. T.: Molecular cloning of the cDNA for the catalytic subunit of human DNA polymerase delta. *Nucleic Acids Res.* 20: 735–745, 1992.

[70994] 9324. Zhao, L.; Chang, L.-S.: The human POLD1 gene: identification of an upstream activator sequence, activation by Sp1 and Sp3, and cell cycle regulation. *J. Biol. Chem.* 272: 4869–4882, 1997.

[70995] 9325. Lind, B.; van Solinge, W. W.; Schwartz, M.; Thorsen, S.: Splice site mutation in the human protein C gene associated with venous thrombosis: demonstration of exon skipping by ectopic transcript analysis. *Blood* 82: 2423–2432, 1993.

[70996] 9326. Domenjoud, L.; Fronia, C.; Uhde, F.; Engel, W.: Sequence of human protamine 2 cDNA. *Nucleic Acids Res.* 16: 7733 only, 1988.

[70997] 9327. Funke, B.; Epstein, J. A.; Kochilas, L. K.; Lu, M. M.; Pandita, R. K.; Liao, J.; Bauerndistel, R.; Schuler, T.; Schorle, H.; Brown, M. C.; Adams, J.; Morrow, B. E.: Mice overexpressing genes from the 22q11 region deleted in velocardio-facial syndrome/DiGeorge syndrome have middle and inner ear defects. *Hum. Molec. Genet.* 10: 2549–2556, 2001.

[70998] 9328. Halford, S.; Wilson, D. I.; Daw, S. C. M.; Roberts, C.;

Wadey,R.; Kamath, S.; Wickremasinghe, A.; Burn, J.; Goodship, J.; Mattei,M.-G.; Moorman, A. F. M.; Scambler, P. J.: Isolation of a gene expressed during early embryogenesis from the region of 22q11 commonly deleted in DiGeorge syndrome.. Hum. Molec. Genet. 2: 1577–1582, 1993.

[70999] 9329.Jerome, L. A.; Papaioannou, V. E.: DiGeorge syndrome phenotype in mice mutant for the T-box gene, Tbx1. Nature Genet. 27: 286–291, 2001.

[71000] 9330.Santos, E.; Martin-Zanca, D.; Reddy, E. P.; Pierotti, M. A.; DellaPorta, G.; Barbacid, M.: Malignant activation of a K-ras oncogene in lung carcinoma but not in normal tissue of the same patient. Science 223:661–664, 1984.

[71001] 9331.Schinzel, A.: Tetrasomy 12p (Pallister-Killian syndrome). J. Med. Genet. 28: 122–125, 1991.

[71002] 9332.Shimizu, K.; Birnbaum, D.; Ruley, M. A.; Fasano, O.; Suard, Y.; Edlund, L.; Taparowsky, E.; Goldfarb, M.; Wigler, M.: Structure of the Ki-ras gene of the human lung carcinoma cell line Calu-1. Nature 304:497–500, 1983.

[71003] 9333.Sidransky, D.; Tokino, T.; Hamilton, S. R.; Kinzler, K. W.; Levin, B.; Frost, P.; Vogelstein, B.: Identification of RAS oncogene mutations in the stool of patients with curable colorectal tumors. Science 256:102–105, 1992.

[71004] 9334.Smit, V. T. H. B. M.; Boot, A. J. M.; Smits, A. M. M.;

Fleuren, G. J.; Cornelisse, C. J.; Bos, J. L.: KRAS codon 12 mutations occur very frequently in pancreatic adenocarcinomas. *Nucleic Acids Res.* 16:7773–7782, 1988.

[71005] 9335. Soukup, S.; Neidich, K.: Prenatal diagnosis of Pallister-Killian syndrome. *Am. J. Med. Genet.* 35: 526–528, 1990.

[71006] 9336. Speleman, F.; Leroy, J. G.; Van Roy, N.; De Paepe, A.; Suijkerbuijk, R.; Brunner, H.; Looijenga, L.; Verschraegen-Spae, M.-R.; Orye, E.: Pallister-Killian syndrome: characterization of the isochromosome 12p by fluorescent in situ hybridization. *Am. J. Med. Genet.* 41:381–387, 1991.

[71007] 9337. Takeda, S.; Ichii, S.; Nakamura, Y.: Detection of K-ras mutation in sputum by mutant-allele-specific amplification (MASA). *Hum. Mutat.* 2:112–117, 1993.

[71008] 9338. Weinberg, R. A.: Fewer and fewer oncogenes. *Cell* 30: 3–4, 1982.

[71009] 9339. Wenger, S. L.; Boone, L. Y.; Steele, M. W.: Mosaicism in Pallister (12p) syndrome. *Am. J. Med. Genet.* 35: 523–525, 1990.

[71010] 9340. Wingo, P. A.; Ries, L. A. G.; Giovino, G. A.; Miller, D. S.; Rosenberg, H. M.; Shopland, D. R.; Thun, M. J.; Edwards, B. K.: Annual report to the nation on the status of cancer, 1973–1996, with a special section on lung cancer and to–

bacco smoking. J. Nat. Cancer Inst. 91: 675–690,1999.

- [71011] 9341.Yanez, L.; Groffen, J.; Valenzuela, D. M.: c-K-ras mutationsin human carcinomas occur preferentially in codon 12. Oncogene 1:315–318, 1987.
- [71012] 9342.Zakowski, M. F.; Wright, Y.; Ricci, A., Jr.: Pericardial agenesisand focal aplasia cutis in tetrasomy 12p (Pallister–Killian syndrome). Am.J. Med. Genet. 42: 323–325, 1992.
- [71013] 9343.Zhang, J.; Marynen, P.; Devriendt, K.; Fryns, J.–P.; Van den Berghe,H.; Cassiman, J.–J.: Molecular analysis of the isochromosome 12pin the Pallister–Killian syndrome: construction of a mouse–human hybridcell line containing an i(12p) as the sole human chromosome. Hum.Genet. 83: 359–363, 1989.
- [71014] 9344.Zhang, Z.; Wang, Y.; Vikis, H. G.; Johnson, L.; Liu, G.; Li, J.;Anderson, M. W.; Sills, R. C.; Hong, H. L.; Devereux, T. R.; Jacks,T.; Guan, K.–L.; You, M.: Wildtype Kras2 can inhibit lung carcinogenesisin mice. Nature Genet. 29: 25–33, 2001.
- [71015] 9345.Adams, J. M.; Gerondakis, S.; Webb, E.; Corcoran, L. M.; Cory,S.: Cellular myc oncogene is altered by chromo–some translocationto an immunoglobulin locus in murine plasmacytomas and is rearrangedsimilarly in human

Burkitt lymphomas. Proc. Nat. Acad. Sci. 80:1982–1986, 1983.

[71016] 9346. Alitalo, K.; Schwab, M.; Lin, C. C.; Varmus, H. E.; Bishop, J.M.: Homogeneously staining chromosomal regions contain amplified copies of an abundantly expressed cellular oncogene (c-myc) in malignant neuroendocrine cells from a human colon carcinoma. Proc. Nat. Acad. Sci. 80: 1707–1711, 1983.

[71017] 9347. Atchley, W. R.; Fitch, W. M.: Myc and Max: molecular evolution of a family of proto-oncogene products and their dimerization partner. Proc. Nat. Acad. Sci. 92: 10217–10221, 1995.

[71018] 9348. Battey, J.; Moulding, C.; Taub, R.; Murphy, W.; Stewart, T.; Potter, H.; Lenoir, G.; Leder, P.: The human c-myc oncogene: structural consequences of translocation into the IgH locus in Burkitt lymphoma. Cell 34:779–787, 1983.

[71019] 9349. Beimling, P.; Benter, T.; Sander, T.; Moelling, K.: Isolation and characterization of the human cellular myc gene product. Biochemistry 24:6349–6355, 1985.

[71020] 9350. Bernard, O.; Cory, S.; Gerondakis, S.; Webb, E.; Adams, J. M.: Sequence of the murine and human cellular myc oncogenes and two modes of myc transcription re-

sulting from chromosome translocation in Blymphoid tumours. EMBO J. 2: 2375–2383, 1983.

[71021] 9351. Bernheim, A.; Berger, R.; Lenoir, G.: Cytogenetic studies on African Burkitt's lymphoma cell lines: t(8;14), t(2;8) and t(8;22) translocations. Cancer Genet. Cytogenet. 3: 307–315, 1981.

[71022] 9352. Bhatia, K.; Huppi, K.; Spangler, G.; Siwarski, D.; Iyer, R.; Magrath, I.: Point mutations in the c-Myc transactivation domain are common in Burkitt's lymphoma and mouse plasmacytomas. Nature Genet. 5: 56–61, 1993.

[71023] 9353. Colby, W. W.; Chen, E. Y.; Smith, D. H.; Levinson, A. D.: Identification and nucleotide sequence of a human locus homologous to the v-myc oncogene of avian myelocytomatosis virus MC29. Nature 301: 722–725, 1983.

[71024] 9354. Cole, M. D.: The myc oncogene: its role in transformation and differentiation. Annu. Rev. Genet. 20: 361–384, 1986.

[71025] 9355. Collins, S.; Groudine, M.: Amplification of endogenous myc-related DNA sequences in a human myeloid leukaemia cell line. Nature 298: 679–681, 1982.

[71026] 9356. Croce, C. M.; Thierfelder, W.; Erikson, J.; Nishikura, K.; Finan, J.; Lenoir, G. M.; Nowell, P. C.: Transcriptional activation of an unrearranged and untranslocated c-myc

oncogene by translocation of a C-lambda locus in Burkitt lymphoma cells. Proc. Nat. Acad. Sci. 80:6922–6926, 1983.

[71027] 9357. Dalla-Favera, R.; Bregni, M.; Erikson, J.; Patterson, D.; Gallo, R. C.; Croce, C. M.: Human c-myc onc gene is located on the region of chromosome 8 that is translocated in Burkitt lymphoma cells. Proc. Nat. Acad. Sci. 79: 7824–7827, 1982.

[71028] 9358. Dalla-Favera, R.; Gelmann, E. P.; Martinotti, S.; Franchini, G.; Papas, T. S.; Gallo, R. C.; Wong-Staal, F.: Cloning and characterization of different human sequences related to the onc gene (v-myc) of avian myelocytomatosis virus (MC29). Proc. Nat. Acad. Sci. 79: 6497–6501, 1982.

[71029] 9359. Duesberg, P. H.: Cancer genes: rare recombinants instead of activated oncogenes (a review). Proc. Nat. Acad. Sci. 84: 2117–2124, 1987.

[71030] 9360. Dyson, E.; Sucov, H. M.; Kubalak, S. W.; Schmid-Schonbein, G. W.; DeLano, F. A.; Evans, R. M.; Ross, J., Jr.; Chien, K. R.: Atrial-like phenotype is associated with embryonic ventricular failure in retinoid X receptor alpha -/- mice. Proc. Nat. Acad. Sci. 92: 7386–7390, 1995.

[71031] 9361. Gampe, R. T., Jr.; Montana, V. G.; Lambert, M. H.; Miller, A. B.; Bledsoe, R. K.; Milburn, M. V.; Kliewer, S. A.;



Willson, T. M.; Xu, H. E.: Asymmetry in the PPAR-gamma/RXR-alpha crystal structure reveals the molecular basis of heterodimerization among nuclear receptors. *Molec. Cell* 5: 545-555, 2000.

[71032] 9362. Gruber, P. J.; Kubalak, S. W.; Pexieder, T.; Sucov, H. M.; Evans, R. M.; Chien, K. R.: RXR-alpha deficiency confers genetic susceptibility for aortic sac, conotruncal, atrioventricular cushion, and ventricular muscle defects in mice. *J. Clin. Invest.* 98: 1332-1343, 1996.

[71033] 9363. Heyman, R. A.; Mangelsdorf, D. J.; Dyck, J. A.; Stein, R. B.; Eichele, G.; Evans, R. M.; Thaller, C.: 9-cis retinoic acid is a high affinity ligand for the retinoid X receptor. *Cell* 68: 397-406, 1992.

[71034] 9364. Hoopes, C. W.; Taketo, M.; Ozato, K.; Liu, Q.; Howard, T. A.; Linney, E.; Seldin, M. F.: Mapping the mouse *Rxr* loci encoding nuclear retinoid X receptors *Rxr*-alpha, *Rxr*-beta, and *Rxr*-gamma. *Genomics* 14: 611-617, 1992.

[71035] 9365. Jones, K. A.; Fitzgibbon, J.; Woodward, K. J.; Goudie, D.; Ferguson-Smith, M. A.; Povey, S.; Wolfe, J.; Solomon, E.: Localization of the retinoid X receptor alpha gene (*RXRA*) to chromosome 9q34. *Ann. Hum. Genet.* 57: 195-201, 1993.

[71036] 9366. Li, M.; Indra, A. K.; Warot, X.; Brocard, J.; Messaddeq,

N.; Kato,S.; Metzger, D.; Chambon, P.: Skin abnormalities generated by temporallycontrolled RXR-alpha mutations in mouse epidermis. *Nature* 407: 633-636,2000.

[71037] 9367.Lu, T. T.; Makishima, M.; Repa, J. J.; Schoonjans, K.; Kerr, T.A.; Auwerx, J.; Mangelsdorf, D. J.: Molecular basis for feedbackregulation of bile acid synthesis by nuclear receptors. *Molec. Cell* 6:507-515, 2000.

[71038] 9368.Mangelsdorf, D. J.; Ong, E. S.; Dyck, J. A.; Evans, R. M.: Nuclearreceptor that identifies a novel retinoic acid response pathway. *Nature* 345:224-229, 1990.

[71039] 9369.Mangelsdorf, D. J.; Umesono, K.; Kliewer, S. A.; Borgmeyer, U.;Ong, E. S.; Evans, R. M.: A direct repeat in the cellular retinol-bindingprotein type II gene confers differential regulation by RXR and RAR. *Cell* 66:555-561, 1991.

[71040] 9370.Repa, J. J.; Turley, S. D.; Lobaccaro, J.-M. A.; Medina, J.; Li,L.; Lustig, K.; Shan, B.; Heyman, R. A.; Dletschy, J. M.; Mangelsdorf,D. J.: Regulation of absorption and ABC1-mediated efflux of cholesterolby RXR heterodimers. *Science* 289: 1524-1529, 2000.

[71041] 9371.Tontonoz, P.; Hu, E.; Spiegelman, B. M.: Stimulation of adipogenesisin fibroblasts by PPAR-gamma-2, a lipid-activated transcription factor. *Cell* 79:1147-1156, 1994.

- [71042] 9372.Willy, P. J.; Umesono, K.; Ong, E. S.; Evans, R. M.; Heyman, R.A.; Mangelsdorf, D. J.: LXR, a nuclear receptor that defines a distinct retinoid response pathway. *Genes Dev.* 9: 1033–1045, 1995.
- [71043] 9373.Wan, Y.–J. Y.; An, D.; Cai, Y.; Repa, J. J.; Chen, T. H.–P.; Flores, M.; Postic, C.; Magnuson, M. A.; Chen, J.; Chien, K. R.; French, S.; Mangelsdorf, D. J.; Sucov, H. M.: Hepatocyte–specific mutation establishes retinoid X receptor alpha as a heterodimeric integrator of multiple physiological processes in the liver. *Molec. Cell. Biol.* 20: 4436–4444, 2000.
- [71044] 9374.Zhou, M. D.; Sucov, H. M.; Evans, R. M.; Chien, K. R.: Retinoid–dependent pathways suppress myocardial cell hypertrophy. *Proc. Nat. Acad. Sci.* 92:7391–7395, 1995.
- [71045] 9375.Fitzgibbon, J.; Gillett, G. T.; Woodward, K. J.; Boyle, J. M.; Wolfe, J.; Povey, S.: Mapping of RXRB to human chromosome 6p21.3. *Ann.Hum. Genet.* 57: 203–209, 1993.
- [71046] 9376.Fleischhauer, K.; McBride, O. W.; DiSanto, J. P.; Ozato, K.; Yang, S. Y.: Cloning and chromosome mapping of human retinoid X receptor beta: selective amino acid sequence conservation of a nuclear hormone receptor in mammals. *Hum. Genet.* 90: 505–510, 1993.
- [71047] 9377.Nagata, T.; Weiss, E. H.; Abe, K.; Kitagawa, K.; Ando,

A.; Yara-Kikuti, Y.; Seldin, M. F.; Ozato, K.; Inoko, H.; Taketo, M.: Physical mapping of the retinoid X receptor B gene in mouse and human. *Immunogenetics* 41:83–90, 1995.

[71048] 9378. Yu, V. C.; Delsert, C.; Andersen, B.; Holloway, J. M.; Devary, O. V.; Naar, A. M.; Kim, S. Y.; Boutin, J.-M.; Glass, C. K.; Rosenfeld, M. G.: RXR-beta: a coregulator that enhances binding of retinoic acid, thyroid hormone, and vitamin D receptors to their cognate response elements. *Cell* 67: 1251–1266, 1991.

[71049] 9379. Graycar, J. L.; Miller, D. A.; Arrick, B. A.; Lyons, R. M.; Moses, H. L.; Derynck, R.: Human transforming growth factor-beta-3: recombinant expression, purification, and biological activities in comparison with transforming growth factors-beta-1 and beta-2. *Molec. Endocr.* 3:1977–1986, 1989.

[71050] 9380. Kaartinen, V.; Voncken, J. W.; Shuler, C.; Warburton, D.; Bu, D.; Heisterkamp, N.; Groffen, J.: Abnormal lung development and cleft palate in mice lacking TGF-beta-3 indicates defects of epithelial-mesenchymal interaction. *Nature Genet.* 11: 415–421, 1995.

[71051] 9381. Lee, B.-S.; Nowak, R. A.: Human leiomyoma smooth muscle cells show increased expression of transforming

growth factor-beta-3 (TGF-beta-3) and altered responses to the antiproliferative effects of TGF-beta. J.Clin. Endocr. Metab. 86: 913-920, 2001.

- [71052] 9382. Moren, A.; Ichijo, H.; Miyazono, K.: Molecular cloning and characterization of the human and porcine transforming growth factor-beta type III receptors. Biochem. Biophys. Res. Commun. 189: 356-362, 1992.
- [71053] 9383. Proetzel, G.; Pawlowski, S. A.; Wiles, M. V.; Yin, M.; Boivin, G. P.; Howles, P. N.; Ding, J.; Ferguson, M. W. J.; Doetschman, T.: Transforming growth factor-beta-3 is required for secondary palate fusion. Nature Genet. 11: 409-414, 1995.
- [71054] 9384. ten Dijke, P.; Geurts van Kessel, A. H. M.; Foulkes, J. G.; LeBeau, M. M.: Transforming growth factor type beta-3 maps to human chromosome 14, region q23-q24. Oncogene 3: 721-724, 1988.
- [71055] 9385. ten Dijke, P.; Hansen, P.; Iwata, K. K.; Pieler, C.; Foulkes, J. G.: Identification of another member of the transforming growth factor type beta gene family. Proc. Nat. Acad. Sci. 85: 4715-4719, 1988.
- [71056] 9386. Engel, W.; Keime, S.; Kremling, H.; Hameister, H.; Schluter, G.: The genes for protamine 1 and 2 (PRM1 and PRM2) and transition protein 2 (TNP2) are closely linked in

the mammalian genome. *Cytogenet. CellGenet.* 61: 158–159, 1992.

[71057] 9387.Nelson, J. E.; Krawetz, S. A.: Linkage of human spermatid-specific basic nuclear protein genes: definition and evolution of the P1-to-P2-to-TP2 locus. *J. Biol. Chem.* 268: 2932–2936, 1993.

[71058] 9388.Schluter, G.; Kremling, H.; Engel, W.: The gene for human transition protein 2: nucleotide sequence, assignment to the protamine gene cluster, and evidence for its low expression. *Genomics* 14: 377–383, 1992.

[71059] 9389.Clapham, J. C.; Arch, J. R. S.; Chapman, H.; Haynes, A.; Lister, C.; Moore, G. B. T.; Piercy, V.; Carter, S. A.; Lehner, I.; Smith, S. A.; Beeley, L. J.; Godden, R. J.; and 15 others: Mice overexpressing human uncoupling protein-3 in skeletal muscle are hyperphagic and lean. *Nature* 406: 415–418, 2000.

[71060] 9390.Dalgaard, L. T.; Sorensen, T. I. A.; Drivsholm, T.; Borch-Johnsen, K.; Andersen, T.; Hansen, T.; Pedersen, O.: A prevalent polymorphism in the promoter of the UCP3 gene and its relationship to body mass index and long term body weight change in the Danish population. *J. Clin. Endocr. Metab.* 86: 1398–1402, 2001.

[71061] 9391.Kim, P. K. M.; Dutra, A. S.; Chandrasekharappa, S.

C.; Puck, J.M.: Genomic structure and mapping of human FADD, an intracellular mediator of lymphocyte apoptosis. *J. Immun.* 157: 5461–5466, 1996.

[71062] 9392. Lavedan, C.; Leroy, E.; Torres, R.; Dehejia, A.; Dutra, A.; Buchholtz, S.; Nussbaum, R. L.; Polymeropoulos, M. H.: Genomic organization and expression of the human beta-synuclein gene (SNCB) *Genomics* 54:173–175, 1998.

[71063] 9393. Lorenz, B.; Migliaccio, C.; Lichtner, P.; Meyer, C.; Strom, T.M.; D'Urso, M.; Becker, J.; Ciccodicola, A.; Meitinger, T.: Cloning and gene structure of the rod cGMP phosphodiesterase delta subunit gene (PDED) in man and mouse. *Europ. J. Hum. Genet.* 6: 283–290, 1998.

[71064] 9394. Hymowitz, S. G.; Christinger, H. W.; Fuh, G.; Ultsch, M.; O'Connell, M.; Kelley, R. F.; Ashkenazi, A.; de Vos, A. M.: Triggering cell death: the crystal structure of Apo2L/TRAIL in a complex with death receptor 5. *Molec. Cell* 4: 563–571, 1999.

[71065] 9395. Nitsch, R.; Bechmann, I.; Deisz, R. A.; Haas, D.; Lehmann, T. N.; Wendling, U.; Zipp, F.: Human brain-cell death induced by tumour-necrosis-factor-related apoptosis-inducing ligand (TRAIL). *Lancet* 356: 827–828, 2000.

[71066] 9396. Pitti, R. M.; Marsters, S. A.; Ruppert, S.; Donahue, C.

J.; Moore,A.; Ashkenazi, A.: Induction of apoptosis by Apo-2 ligand, a newmember of the tumor necrosis factor cytokine family. J. Biol. Chem. 271:12687-12690, 1996.

[71067] 9397.Wiley, S. R.; Schooley, K.; Smolak, P. J.; Din, W. S.; Huang, C.-P.;Nicholl, J. K.; Sutherland, G. R.; Smith, T. D.; Rauch, C.; Smith,C. A.; Goodwin, R. G.: Identification and characterization of a newmember of the TNF family that induces apoptosis. Immunity 3: 673-682,1995.

[71068] 9398.Lai, C. S. L.; Fisher, S. E.; Hurst, J. A.; Levy, E. R.; Hodgson,S.; Fox, M.; Jeremiah, S.; Povey, S.; Jamison, D. C.; Green, E. D.;Vargha-Khadem, F.; Monaco, A. P.: The SPCH1 region on human 7q31:genomic characterization of the critical interval and localizationof translocations associated with speech and language disorder. Am.J. Hum. Genet. 67: 357-368, 2000.

[71069] 9399.Lai, C. S. L.; Fisher, S. E.; Hurst, J. A.; Vargha-Khadem, F.;Monaco, A. P.: A forkhead-domain gene is mutated in a severe speechand language disorder. Nature 413: 519-523, 2001.

[71070] 9400.Lyu, M. S.; Park, D. J.; Rhee, S. G.; Kozak, C. A.: Genetic mappingof the human and mouse phospholipase C genes. Mammalian Genome 7:501-504, 1996.

[71071] 9401.Shimohama, S.; Kamiya, S.; Fujii, M.; Ogawa, T.;



Kanamori, M.;Kawamata, J.; Imura, T.; Taniguchi, T.; Yagi-sawa, H.: Mutation inthe pleckstrin homology domain of the human phospholipase C-delta-1gene is associated with loss of function. *Biochem. Biophys. Res.Comm.* 245: 722-728, 1998.

[71072] 9402.Iwabuchi, K.; Bartel, P. L.; Li, B.; Marraccino, R.; Fields, S.: Two cellular proteins that bind to wild-type but not mutant p53. *Proc.Nat. Acad. Sci.* 91: 6098-6102, 1994.

[71073] 9403.Iwabuchi, K.; Li, B.; Massa, H. F.; Trask, B. J.; Date, T.; Fields,S.: Stimulation of p53-mediated transcriptional activation by thep53-binding proteins, 53BP1 and 53BP2. *J. Biol. Chem.* 273: 26061-26068,1998.

[71074] 9404.Naumovski, L.; Cleary, M. L.: The p53-binding protein 53BP2 alsointeracts with Bcl2 and impedes cell cycle progression at G2/M. *Molec.Cell Biol.* 16: 3884-3892, 1996.

[71075] 9405.Samuels-Lev, Y.; O'Connor, D. J.; Bergamaschi, D.; Trigiante, G.;Hsieh, J.-K.; Zhong, S.; Campargue, I.; Naumovski, L.; Crook, T.;Lu, X.: ASPP proteins specifically stimulate the apoptotic functionof p53. *Molec. Cell* 8: 781-794, 2001.

[71076] 9406.Yang, J.-P.; Ono, T.; Sonta, S.; Kawabe, T.; Okamoto,

T.: Assignment of p53 binding protein (TP53BP2) to human chromosome band 1q42.1 by in situ hybridization. *Cytogenet. Cell Genet.* 78: 61–62, 1997.

[71077] 9407. Crawford, M. J.; Lanctot, C.; Tremblay, J. J.; Jenkins, N.; Gilbert, D.; Copeland, N.; Beatty, B.; Drouin, J.: Human and murine PTX1/Ptx1 gene maps to the region for Treacher Collins syndrome. *Mammalian Genome* 8: 841–845, 1997.

[71078] 9408. Lamonerie, T.; Tremblay, J. J.; Lanctot, C.; Thierrien, M.; Gauthier, Y.; Drouin, J.: Ptx1, a bicoid-related homeobox transcription factor involved in transcription of the pro-opiomelanocortin gene. *Genes Dev.* 10: 1284–1295, 1996.

[71079] 9409. Shang, J.; Li, X.; Ring, H. Z.; Clayton, D. A.; Francke, U.: Backfoot, a novel homeobox gene, maps to human chromosome 5 (BFT) and mouse chromosome 13 (Bft). *Genomics* 40: 108–113, 1997.

[71080] 9410. Szeto, D. P.; Rodriguez-Esteban, C.; Ryan, A. K.; O'Connell, S. M.; Liu, F.; Kioussi, C.; Gleiberman, A. S.; Izpisua-Belmonte, J. C.; Rosenfeld, M. G.: Role of the Bicoid-related homeodomain factor Pitx1 in specifying hindlimb morphogenesis and pituitary development. *Genes Dev.* 13: 484–494, 1999.

- [71081] 9411.Cohen, M. E.; Yin, M.; Paznekas, W. A.; Schertzer, M.; Wood, S.;Jabs, E. W.: Human SLUG gene organization, expression, and chromosomemap location on 8q. *Genomics* 51: 468–471, 1998.
- [71082] 9412.Inukai, T.; Inoue, A.; Kurosawa, H.; Goi, K.; Shinjyo, T.; Ozawa,K.; Mao, M.; Inaba, T.; Look, A. T.: SLUG, a ces-1-related zinc fingertranscription factor gene with anti-apoptotic activity, is a downstreamtarget of the E2A-HLF oncoprotein. *Molec. Cell* 4: 343–352, 1999.
- [71083] 9413.Nieto, A. M.; Sargent, M. G.; Wilkinson, D. G.; Cooke, J.: Control of cell behavior during vertebrate development by Slug, a zinc finger gene. *Science* 264: 835–839, 1994.
- [71084] 9414.Glover, M. T.; Atherton, D. J.: Transient zinc deficiency in two full-term breast-fed siblings associated with low maternal breastmilk zinc concentration. *Pediat. Derm.* 5: 10–13, 1988.
- [71085] 9415.Kuramoto, Y.; Igarashi, Y.; Tagami, H.: Acquired zinc deficiency in breast-fed infants. *Semin. Derm.* 10: 309–312, 1991.
- [71086] 9416.Kury, S.; Devilder, M.-C.; Avet-Loiseau, H.; Dreno, B.; Moisan, J.-P.: Expression pattern, genomic structure and evaluation of the human SLC30A4 gene as a candidate for acrodermatitis enteropathica. *Hum.Genet.* 109:

178–185, 2001.

- [71087] 9417.Nakano, A.; Nakano, H.; Hanada, K.; Nomura, K.; Uitto, J.: ZNT4 gene is not responsible for acrodermatitis enteropathica in Japanese families. (Letter) Hum. Genet. 110: 201–202, 2002.
- [71088] 9418.Agulnick, A. D.; Taira, M.; Breen, J. J.; Tanaka, T.; Dawid, I.B.; Westphal, H.: Interactions of the LIM-domain-binding factor Ldb1 with LIM homeodomain proteins. Nature 384: 270–272, 1996.
- [71089] 9419.Bach, I.; Carriere, C.; Ostendorff, H. P.; Andersen, B.; Rosenfield, M. G.: A family of LIM domain-associated cofactors confer transcriptional synergism between LIM and Otx homeodomain proteins. Genes Dev. 11:1370–1380, 1997.
- [71090] 9420.Jurata, L. W.; Gill, G. N.: Functional analysis of the nuclear LIM domain interactor NL1. Molec. Cell. Biol. 17: 5688–5698, 1997.
- [71091] 9421.Jurata, L. W.; Kenny, D. A.; Gill, G. N.: Nuclear LIM interactor, a rhombotin and LIM homeodomain interacting protein, is expressed early in neuronal development. Proc. Nat. Acad. Sci. 93: 11693–11698, 1996.
- [71092] 9422.Semina, E. V.; Altherr, M. R.; Murray, J. C.: Cloning and chromosomal localization of two novel human genes

encoding LIM-domain binding factors CLIM1 and CLIM2/LDB1/NL1. Mammalian Genome 9: 921–924, 1998.

- [71093] 9423. Yamashita, T.; Agulnick, A. D.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Westphal, H.: Genomic structure and chromosomal localization of the mouse LIM domain-binding protein 1 gene, Ldb1. Genomics 48:87–92, 1998.
- [71094] 9424. Ahmad, M.; Srinivasula, S. M.; Wang, L.; Talanian, R. V.; Litwack, G.; Fernandes-Alnemri, T.; Alnemri, E. S.: CRADD, a novel human apoptotic adaptor molecule for caspase-2, and FasL/tumor necrosis factor receptor-interacting protein RIP. Cancer Res. 57: 615–619, 1997.
- [71095] 9425. Duan, H.; Dixit, V. M.: RAIDD is a new 'death' adaptor molecule. Nature 385:86–89, 1997.
- [71096] 9426. Horvat, S.; Medrano, J. F.: A 500-kb YAC and BAC contig encompassing the high-growth deletion in mouse chromosome 10 and identification of the murine Raidd/Cradd gene in the candidate region. Genomics 54:159–164, 1998.
- [71097] 9427. Foley, B. T.; Moehring, J. M.; Moehring, T. J.: Mutations in the elongation factor 2 gene which confer resistance to diphtheria toxin and Pseudomonas exotoxin A: genetic and biochemical analyses. J. Biol. Chem. 270:

23218–23225, 1995.

- [71098] 9428.Mattheakis, L. C.; Sor, F.; Collier, R. J.: Diphthamide synthesis in *Saccharomyces cerevisiae*: structure of the DPH2 gene. *Gene* 132:149–154, 1993.
- [71099] 9429.Schultz, D. C.; Balasara, B. R.; Testa, J. R.; Godwin, A. K.:Cloning and localization of a human diphthamide biosynthesis-like protein-2 gene, DPH2L2. *Genomics* 52: 186–191, 1998.
- [71100] 9430.Abe, R.; Sakashita, E.; Yamamoto, K.; Sakamoto, H.: Two different RNA binding activities for the AU-rich element and the poly(A) sequence of the mouse neuronal protein mHuC. *Nucleic Acids Res.* 24: 4895–4901, 1996.
- [71101] 9431.Rhim, H.; Savagner, P.; Thibaudieu, G.; Thiery, J. P.; Pavan, W.J.: Localization of a neural crest transcription factor, Slug, to mouse chromosome 16 and human chromosome 8. *Mammalian Genome* 8:872–873, 1997.
- [71102] 9432.Savagner, P.; Yamada, K. M.; Thiery, J. P.: The zinc-finger protein Slug causes desmosome dissociation, an initial and necessary step for growth factor-induced epithelial-mesenchymal transition. *J. Cell Biol.* 137: 1403–1419, 1997.
- [71103] 9433.Winter, H.; Rogers, M. A.; Gebhardt, M.; Wollina, U.; Boxall, L.; Chitayat, D.; Babul-Hirji, R.; Stevens, H. P.; Zlo-

togorski, A.; Schweizer, J.: A new mutation in the type II hair cortex keratin hHb1 involved in the inherited hair disorder monilethrix. *Hum. Genet.* 101: 165–169, 1997.

[71104] 9434. Buckanovich, R. J.; Posner, J. B.; Darnell, R. B.: Nova, the paraneoplastic Ri antigen, is homologous to an RNA-binding protein and is specifically expressed in the developing motor system. *Neuron* 11: 657–672, 1993.

[71105] 9435. Buckanovich, R. J.; Yang, Y. Y.; Darnell, R. B.: The onconeural antigen Nova-1 is a neuron-specific RNA-binding protein, the activity of which is inhibited by paraneoplastic antibodies. *J. Neurosci.* 16: 1114–1122, 1996.

[71106] 9436. Prestigiacomo, C. J.; Balmaceda, C.; Dalmau, J.: Anti-Ri-associated paraneoplastic opsoclonus-ataxia syndrome in a man with transitional cell carcinoma: a case report. *Cancer* 91: 1423–1428, 2001.

[71107] 9437. Anguita, J.; Chalfant, M. L.; Civan, M. M.; Coca-Prados, M.: Molecular cloning of the human volume-sensitive chloride conductance regulatory protein, pl(Cln), from ocular ciliary epithelium. *Biochem. Biophys. Res. Commun.* 208: 89–95, 1995.

[71108] 9438. Buyse, G.; De Greef, C.; Raeymaekers, L.; Droogmans, G.; Nilius, B.; Eggermont, J.: The ubiquitously expressed pl(Cln) protein forms homomeric complexes in

vitro. Biochem. Biophys. Res. Commun. 218:822–827, 1996.

[71109] 9439. Nagl, U. O.; Erdel, M.; Schmarda, A.; Seri, M.; Pinggera, G. M.; Gschwentner, M.; Duba, C.; Galletta, L. J. V.; Deetjen, P.; Utermann, G.; Paulmichl, M.: Chromosomal localization of the genes (CLNS1A and CLNS1B) coding for the swelling-dependent chloride channel I(Cln). Genomics 38:438–441, 1996.

[71110] 9440. Schwartz, R. S.; Rybicki, A. C.; Nagel, R. L.: Molecular cloning and expression of a chloride channel-associated protein pI(Cln) in human young red blood cells: association with actin. Biochem. J. 327:609–616, 1997.

[71111] 9441. Muzio, M.; Ni, J.; Feng, P.; Dixit, V. M.: IRAK (Pelle) family member IRAK-2 and MyD88 as proximal mediators of IL-1 signaling. Science 278:1612–1615, 1997.

[71112] 9442. Takeuchi, O.; Hoshino, K.; Akira, S.: Cutting edge: TLR2-deficient and MyD88-deficient mice are highly susceptible to Staphylococcus aureus infection. J. Immun. 165: 5392–5396, 2000.

[71113] 9443. Eggertsen, G.; Olin, M.; Andersson, U.; Ishida, H.; Kubota, S.; Hellman, U.; Okuda, K.-I.; Bjorkhem, I.: Molecular cloning and expression of rabbit sterol 12- $\alpha$ -hydroxylase. J. Biol. Chem. 271:



32269–32275,1996.

- [71114] 9444.Gafvels, M.; Olin, M.; Chowdhary, B. P.; Raudsepp, T.; Andersson,U.; Persson, B.; Jansson, M.; Bjorkhem, I.; Eggertsen, G.: Structureand chromosomal assignment of the sterol 12- $\alpha$ -hydroxylase gene(CYP8B1) in human and mouse: eukaryotic cytochrome P-450 gene devoidof introns. *Genomics* 56: 184–196, 1999.
- [71115] 9445.Zhang, M.; Chiang, J. Y. L.: Transcriptional regulation of thehuman sterol 12- $\alpha$ -hydroxylase gene (CYP8B1): roles of hepatocytenuclear factor 4- $\alpha$  in mediating bile acid repression. *J. Biol.Chem.* 276: 41690–41699, 2001.
- [71116] 9446.Brinkman, A.; van der Flier, S.; Kok, E. M.; Dorssers, L. C. J.: BCAR1, a human homologue of the adapter protein p130Cas, and antiestrogenresistance in breast cancer cells. *J. Nat. Cancer Inst.* 92: 112–120,2000.
- [71117] 9447.Brugge, J. S.: Casting light on focal adhesions. *Nature Genet.* 19:309–311, 1998.
- [71118] 9448.Dorssers, L. C. J.; van Agthoven, T.; Dekker, A.; van Agthoven,T. L. A.; Kok, E. M.: Induction of antiestrogen resistance in humanbreast cancer cells by random insertional mutagenesis using defectiveretroviruses: identification of bcar-1, a common integration site. *Molec.Endocr.*

7: 870–878, 1993.

- [71119] 9449. Honda, H.; Oda, H.; Nakamoto, T.; Honda, Z.; Sakai, R.; Suzuki, T.; Saito, T.; Nakamura, K.; Nakao, K.; Ishikawa, T.; Katsuki, M.; Yazaki, Y.; Hirai, H.: Cardiovascular anomaly, impaired actin bundling and resistance to Src-induced transformation in mice lacking p130-Cas. *Nature Genet.* 19: 361–365, 1998.
- [71120] 9450. Kanner, S. B.; Reynolds, A. B.; Wang, H. C.; Vines, R. R.; Parsons, J. T.: The SH2 and SH3 domains of pp60src direct stable association with tyrosine phosphorylated proteins p130 and p110. *EMBO J.* 10: 1689–1698, 1991.
- [71121] 9451. Kuo, C. T.; Morrissey, E. E.; Anandappa, R.; Sigrist, K.; Lu, M. M.; Parmacek, M. S.; Soudais, C.; Leiden, J. M.: GATA4 transcription factor is required for ventral morphogenesis and heart tube formation. *Genes Dev.* 11: 1048–1060, 1997.
- [71122] 9452. Reynolds, A. B.; Kanner, S. B.; Wang, H. C.; Parsons, J. T.: Stable association of activated pp60src with two tyrosine-phosphorylated cellular proteins. *Molec. Cell. Biol.* 9: 3951–3958, 1989.
- [71123] 9453. Sakai, R.; Iwamatsu, A.; Hirano, N.; Ogawa, S.; Tanaka, T.; Mano, H.; Yazaki, Y.; Hirai, H.: A novel signaling molecule, p130, forms stable complexes in vivo with

v-Crk and v-Src in a tyrosine phosphorylation-dependent manner. EMBO J. 13: 3748–3756, 1994.

- [71124] 9454. van der Flier, S.; Brinkman, A.; Look, M. P.; Kok, E. M.; Meijer-van Gelder, M. E.; Klijn, J. G. M.; Dorssers, L. C. J.; Foekens, J. A.: Bcr1/p130Cas protein and primary breast cancer: prognosis and response to tamoxifen treatment. J. Nat. Cancer Inst. 92: 120–127, 2000.
- [71125] 9455. Yoshida, K.; Taga, T.; Saito, M.; Suematsu, S.; Kumanogoh, A.; Tanaka, T.; Fujiwara, H.; Hirata, M.; Yamagami, T.; Nakahata, T.; Hirabayashi, T.; Yoneda, Y.; Tanaka, K.; Wang, W. Z.; Mori, C.; Shiota, K.; Yoshida, N.; Kishimoto, T.: Targeted disruption of gp130, a common signal transducer for the interleukin 6 family of cytokines, leads to myocardial and hematological disorders. Proc. Nat. Acad. Sci. 93: 407–411, 1996.
- [71126] 9456. Hirose, T.; Smith, R. J.; Jetten, A. M.: ROR-gamma: the third member of ROR/RZR orphan receptor subfamily that is highly expressed in skeletal muscle. Biochem. Biophys. Res. Commun. 205: 1976–1983, 1994.
- [71127] 9457. Kurebayashi, S.; Ueda, E.; Sakaue, M.; Patel, D. D.; Medvedev, A.; Zhang, F.; Jetten, A. M.: Retinoid-related orphan receptor gamma (ROR-gamma) is essential for lymphoid organogenesis and controls apoptosis during thy-

mopoiesis. Proc. Nat. Acad. Sci. 97: 10132–10137, 2000.

[71128] 9458. Medvedev, A.; Chistokhina, A.; Hirose, T.; Jetten, A. M.: Genomic structure and chromosomal mapping of the nuclear orphan receptor ROR–gamma(RORC) gene. Genomics 46: 93–102, 1997.

[71129] 9459. Soengas, M. S.; Capodieci, P.; Polsky, D.; Mora, J.; Esteller, M.; Opitz–Araya, X.; McCombie, R.; Herman, J. G.; Gerald, W. L.; Lazebnik, Y. A.; Cordon–Cardo, C.; Lowe, S. W.: Inactivation of the apoptosis effector Apaf–1 in malignant melanoma. Nature 409: 207–211, 2001.

[71130] 9460. Srinivasula, S. M.; Ahmad, M.; Fernandes–Alnemri, T.; Alnemri, E. S.: Autoactivation of procaspase–9 by Apaf–1-mediated oligomerization. Molec. Cell 1: 949–957, 1998.

[71131] 9461. Yoshida, H.; Kong, Y.–Y.; Yoshida, R.; Elia, A. J.; Hakem, A.; Hakem, R.; Penninger, J. M.; Mak, T. W.: Apaf1 is required for mitochondrial pathways of apoptosis and brain development. Cell 94: 739–750, 1998.

[71132] 9462. Zou, H.; Henzel, W. J.; Liu, X.; Lutschg, A.; Wang, X.: APAF–1, a human protein homologous to *C. elegans* CED–4, participates in cytochrome c-dependent activation of caspase–3. Cell 90: 405–413, 1997.

[71133] 9463. Biervert, C.; Schroeder, B. C.; Kubisch, C.; Berkovic,

S. F.; Propping, P.; Jentsch, T. J.; Steinlein, O. K.: A potassium channel mutation in neonatal human epilepsy. *Science* 279: 406–409, 1998.

[71134] 9464. Biervert, C.; Steinlein, O. K.: Structural and mutational analysis of KCNQ2, the major gene locus for benign familial neonatal convulsions. *Hum. Genet.* 104: 234–240, 1999.

[71135] 9465. Dedek, K.; Kunath, B.; Kananura, C.; Reuner, U.; Jentsch, T. J.; Steinlein, O. K.: Myokymia and neonatal epilepsy caused by a mutation in the voltage sensor of the KCNQ2 K(+) channel. *Proc. Nat. Acad. Sci.* 98: 12272–12277, 2001.

[71136] 9466. del Giudice, E. M.; Coppola, G.; Scuccimarra, G.; Cirillo, G.; Bellini, G.; Pascotto, A.: Benign familial neonatal convulsions (BFNC) resulting from mutation of the KCNQ2 voltage sensor. *Europ. J. Hum. Genet.* 8: 994–997, 2000.

[71137] 9467. Niederreither, K.; Abu-Abed, S.; Schuhbaur, B.; Petkovich, M.; Chambon, P.; Dolle, P.: Genetic evidence that oxidative derivatives of retinoic acid are not involved in retinoid signaling during mouse development. *Nature Genet.* 31: 84–88, 2002.

[71138] 9468. White, J. A.; Beckett, B.; Scherer, S. W.; Herbrick, J.-A.; Petkovich, M.: P450RAI (CYP26A) maps to human

chromosome 10q23–q24 and mouse chromosome 19C2–3. *Genomics* 48: 270–272, 1998.

- [71139] 9469. White, J. A.; Beckett–Jones, B.; Guo, Y.–D.; Dilworth, F. J.; Bonasoro, J.; Jones, G.; Petkovich, M.: cDNA cloning of human retinoic acid–metabolizing enzyme (hP450RAI) identifies a novel family of cytochromes P450 (CYP26). *J. Biol. Chem.* 272: 18538–18541, 1997.
- [71140] 9470. Chardin, P.; Paris, S.; Antonny, B.; Robineau, S.; Beraud–Dufour, S.; Jackson, C. L.; Chabre, M.: A human exchange factor for ARF contains Sec7– and pleckstrin–homology domains. *Nature* 384: 481–484, 1996.
- [71141] 9471. Perletti, L.; Talarico, D.; Trecca, D.; Ronchetti, D.; Fracchiolla, N. S.; Maiolo, A. T.; Neri, A.: Identification of a novel gene, PSD, adjacent to NFKB2/lyt–10, which contains Sec7 and pleckstrin–homology domains. *Genomics* 46: 251–259, 1997.
- [71142] 9472. Biunno, I.; Appierto, V.; Cattaneo, M.; Leone, B. E.; Balzano, G.; Socci, C.; Saccone, S.; Letizia, A.; Valle, G. D.; Sgaramella, V.: Isolation of a pancreas–specific gene located on human chromosome 14q31: expression analysis in human pancreatic ductal carcinomas. *Genomics* 46: 284–286, 1997.
- [71143] 9473. Donoviel, D. B.; Bernstein, A.: SEL–1L maps to hu–

man chromosome14, near the insulin-dependent diabetes mellitus locus 11. Genomics 56:232–233, 1999.

[71144] 9474. Grant, B.; Greenwald, I.: The *Caenorhabditis. elegans* sel-1 gene, a negative regulator of lin-12 and glp-1, encodes a predicted extracellular protein. Genetics 143: 237–247, 1996.

[71145] 9475. Grant, B.; Greenwald, I.: Structure, function and expression of SEL-1, a negative regulator of LIN-12 and GLP-1 in *C. elegans*. Development 124:637–644, 1997.

[71146] 9476. Frohling, S.; Nakabayashi, K.; Scherer, S. W.; Dohner, H.; Dohner, K.: Mutation analysis of the origin recognition complex subunit 5 (ORC5L) gene in adult patients with myeloid leukemias exhibiting deletion of chromosome band 7q22. Hum. Genet. 108: 304–309, 2001.

[71147] 9477. Ishiai, M.; Dean, F. B.; Okumura, K.; Abe, M.; Moon, K.-Y.; Amin, A. A.; Kagotani, K.; Taguchi, H.; Murakami, Y.; Hanaoka, F.; O'Donnell, M.; Hurwitz, J.; Eki, T.: Isolation of human and fission yeast homologues of the budding yeast origin recognition complex subunit ORC5: human homologue (ORC5L) maps to 7q22. Genomics 46: 294–298, 1997.

[71148] 9478. Quintana, D. G.; Thome, K. C.; Hou, Z.; Ligon, A. H.; Morton, C. C.; Dutta, A.: ORC5L, a new member of the hu-

man origin recognition complex, is deleted in uterine leiomyomas and malignant myeloid diseases. *J. Biol. Chem.* 273: 27137–27145, 1998.

[71149] 9479. Ben Porath, I.; Kozak, C. A.; Benvenisty, N.: Chromosomal mapping of Tmp (Emp1), Xmp (Emp2), and Ymp (Emp3), genes encoding membrane proteins related to Pmp22. *Genomics* 49: 443–447, 1998.

[71150] 9480. Chen, Y.; Medvedev, A.; Ruzanov, P.; Marvin, K. W.; Jetten, A. M.: cDNA cloning, genomic structure, and chromosome mapping of the human epithelial membrane protein CL-20 gene (EMP1), a member of the PMP22 family. *Genomics* 41: 40–48, 1997.

[71151] 9481. Liehr, T.; Kühlenbaumer, G.; Wulf, P.; Taylor, V.; Suter, U.; VanBroeckhoven, C.; Lupski, J. R.; Claussen, U.; Rautenstrauss, B.: Regional localization of the human epithelial membrane protein genes 1, 2, and 3 (EMP1, EMP2, EMP3) to 12p12.3, 16p13.2, and 19q13.3. *Genomics* 58: 106–108, 1999.

[71152] 9482. Marvin, K. W.; Fujimoto, W.; Jetten, A. M.: Identification and characterization of a novel squamous cell-associated gene related to PMP22. *J. Biol. Chem.* 270: 28910–28916, 1995.

[71153] 9483. Ruegg, C. L.; Wu, H.; Fagnoni, F. F.; Engleman, E. G.;



Laus, R.: B4B, a novel growth-arrest gene, is expressed by a subset of progenitor/pre-B lymphocytes negative for cytoplasmic mu-chain. *J. Immun.* 157: 72-80, 1996.

[71154] 9484. Taylor, V.; Suter, U.: Epithelial membrane protein-2 and epithelial membrane protein-3: two novel members of the peripheral myelin protein 22 gene family. *Gene* 175: 115-120, 1996.

[71155] 9485. Masiakowski, P.; Carroll, R. D.: A novel family of cell surface receptors with tyrosine kinase-like domain. *J. Biol. Chem.* 267:26181-26190, 1992.

[71156] 9486. Adelmant, G.; Begue, A.; Stehelin, D.; Laudet, V.: A functional Rev-erb-alpha responsive element located in the human Rev-erb-alpha promoter mediates a repressing activity. *Proc. Nat. Acad. Sci.* 93:3553-3558, 1996.

[71157] 9487. Chawla, A.; Lazar, M. A.: Induction of Rev-ErbA alpha, an orphan receptor encoded on the opposite strand of the alpha-thyroid hormone receptor gene, during adipocyte differentiation. *J. Biol. Chem.* 268:16265-16269, 1993.

[71158] 9488. Lazar, M. A.; Hodin, R. A.; Darling, D. S.; Chin, W. W.: A novel member of the thyroid/steroid hormone receptor family is encoded by the opposite strand of the rat c-erbA-alpha transcriptional unit. *Molec. Cell. Biol.* 9:

1128–1136, 1989.

- [71159] 9489.Miyajima, N.; Horiuchi, R.; Shibuya, S.; Matsubara, K.; Toyoshima,K.; Yamamoto, T.: Two erbA homologs encoding proteins with differentT(3) binding capacities are transcribed from opposite DNA strands of the same genetic locus. *Cell* 57: 31–39, 1989.
- [71160] 9490.Preitner, N.; Damiola, F.; Lopez–Molina, L.; Zakany, J.; Duboule,D.; Albrecht, U.; Schibler, U.: The orphan nuclear receptor REV–ERB–alphacontrols circadian transcription within the positive limb of the mammalian circadian oscillator. *Cell* 110: 251–260, 2002.
- [71161] 9491.Zhao, Q.; Khorasanizadeh, S.; Miyoshi, Y.; Lazar, M. A.; Rastinejad,F.: Structural elements of an orphan nuclear receptor–DNA complex. *Molec.Cell* 1: 849–861, 1998.
- [71162] 9492.Gregorini, A.; Sahin, F. I.; Lillington, D. M.; Meerabux, J.; Saha,V.; McCullagh, P.; Bocci, M.; Menevse, S.; Papa, S.; Young, B. D.: Gene BR140, which is related to AF10 and AF17, maps to chromosomeband 3p25. *Genes Chromosomes Cancer* 17: 269–272, 1996.
- [71163] 9493.Thompson, K. A.; Wang, B.; Argraves, W. S.; Giannotti, F. G.; Schranck,D. P.; Ruoslahti, E.: BR140, a novel zinc–finger protein with homologyto the TAF250 subunit of TFIID. *Biochem. Biophys. Res. Commun.*

1143–1152,1994.

- [71164] 9494.Saito, T.; Seki, N.; Yamauchi, M.; Tsuji, S.; Hayashi, A.; Kozuma,S.; Hori, T.: Structure, chromosomal location, and expression profileof EXTR1 and EXTR2, new members of the multiple exostoses gene family. *Biochem.Biophys. Res. Commun.* 243: 61–66, 1998.
- [71165] 9495.Wuyts, W.; Van Hul, W.: Characterization and genomic localizationof the mouse *Extl2* gene. *Cytogenet. Cell Genet.* 89: 185–188, 2000.
- [71166] 9496.Wuyts, W.; Van Hul, W.; Hendrickx, J.; Speleman, F.; Wauters, J.;De Boule, K.; Van Roy, N.; Van Agtmael, T.; Bossuyt, P.; Willems,P. J.: Identification and characterization of a novel member of theEXT gene family, EXTL2. *Europ. J. Hum. Genet.* 5: 382–389, 1997.
- [71167] 9497.Dick, T.; Ray, K.; Salz, H. K.; Chia, W.: Cytoplasmic dynein (*ddlc1*)mutations cause morphogenetic defects and apoptotic cell death in*Drosophila melanogaster*. *Molec. Cell. Biol.* 16: 1966–1977, 1996.
- [71168] 9498.Matsushita, M.; Endo, Y.; Taira, S.; Sato, Y.; Fujita, T.; Ichikawa,N.; Nakata, M.; Misuochi, T.: A novel human lectin with collagen–and fibrinogen–like domains which functions as an opsonin. *J. Biol.Chem.* 271: 2448–2454, 1996.

- [71169] 9499. Haag, F.; Koch-Nolte, F.; Kuhl, M.; Lorenzen, S.; Thiele, H.-G.: Premature stop codons inactivate the RT6 genes of the human and chimpanzee species. *J. Molec. Biol.* 243: 537–546, 1994.
- [71170] 9500. Koch-Nolte, F.; Haag, F.; Kuhl, M.; van Heyningen, V.; Hoovers, J.; Grzeschik, K.-H.; Singh, S.; Thiele, H.-G.: Assignment of the human RT6 gene to 11q13 by PCR screening of somatic cell hybrids and in situ hybridization. *Genomics* 18: 404–406, 1993.
- [71171] 9501. Koch-Nolte, F.; Kuhl, M.; Haag, F.; Cetkovic-Cvrlje, M.; Leiter, E. H.; Thiele, H.-G.: Assignment of the human and mouse genes for muscle ecto mono (ADP-ribose) transferase to a conserved linkage group on human chromosome 11p15 and mouse chromosome 7. *Genomics* 36: 215–216, 1996.
- [71172] 9502. Okazaki, I. J.; Zolkiewska, A.; Nightingale, M. S.; Moss, J.: Immunological and structural conservation of mammalian skeletal muscle glycosylphosphatidylinositol-linked ADP-ribosyltransferases. *Biochemistry* 33: 12828–12836, 1994.
- [71173] 9503. Prochazka, M.; Gaskins, H. R.; Leiter, E. H.; Koch-Nolte, F.; Haag, F.; Thiele, H.-G.: Chromosomal localization, DNA polymorphism and expression of Rt-6, the

mouse homologue of rat T-lymphocyte differentiation-marker RT6. Immunogenetics 33: 152–156, 1991.

[71174] 9504.Banfi, S.; Borsani, G.; Rossi, E.; Bernard, L.; Guffanti, A.; Rubboli,F.; Marchitiello, A.; Giglio, S.; Coluccia, E.; Zollo, M.; Zuffardi,O.; Ballabio, A.: Identification and mapping of human cDNAs homologousto Drosophila mutant genes through EST database searching. NatureGenet. 13: 167–174, 1996.

[71175] 9505.Duncan, M. K.; Kos, L.; Jenkins, N. A.; Gilbert, D. J.; Copeland,N. G.; Tomarev, S. I.: Eyes absent: a gene family found in severalmetazoan phyla. Mammalian Genome 8: 479–485, 1997. Note: Erratum:Mammalian Genome 8: 877 only, 1997.

[71176] 9506.Xu, P.–X.; Cheng, J.; Epstein, J. A.; Maas, R. L.: Mouse Eya genesare expressed during limb tendon development and encode a transcriptionalactivation function. Proc. Nat. Acad. Sci. 94: 11974–11979, 1997.

[71177] 9507.Zimmerman, J. E.; Bui, Q. T.; Steingrimsson, E.; Nagle, D. L.;Fu, W.; Genin, A.; Spinner, N. B.; Copeland, N. G.; Jenkins, N. A.;Bucan, M.; Bonini, N. M.: Cloning and characterization of two vertebratehomologs of the Drosophila eyes absent gene. Genome Res. 7: 128–141,1997.

[71178] 9508.Morrissey, E. E.; Tang, Z.; Sigrist, K.; Lu, M. M.; Jiang,

F.; Ip, H. S.; Parmacek, M. S.: GATA6 regulates HNF4 and is required for differentiation of visceral endoderm in the mouse embryo. *Genes Dev.* 12:3579–3590, 1998.

[71179] 9509. Suzuki, E.; Evans, T.; Lowry, J.; Truong, L.; Bell, D. W.; Testa, J. R.; Walsh, K.: The human GATA-6 gene: structure, chromosomal location, and regulation of expression by tissue-specific and mitogen-responsive signals. *Genomics* 38: 283–290, 1996.

[71180] 9510. Meurs, E.; Chong, K.; Galabru, J.; Shaun, N.; Thomas, S. B.; Kerr, I. M.; Williams, B. R. G.; Hovanessian, A. G.: Molecular cloning and characterization of the human double-stranded RNA-activated protein kinase induced by interferon. *Cell* 62: 379–390, 1990.

[71181] 9511. Dunnick, W.; Shell, B. E.; Dery, C.: DNA sequences near the site of reciprocal recombination between a c-myc oncogene and an immunoglobulin switch region. *Proc. Nat. Acad. Sci.* 80: 7269–7273, 1983.

[71182] 9512. Bashir, R.; Strachan, T.; Keers, S.; Stephenson, A.; Mahjneh, I.; Marconi, G.; Nashef, L.; Bushby, K. M. D.: A gene for autosomal recessive limb-girdle muscular dystrophy maps to chromosome 2p. *Hum. Molec. Genet.* 3: 455–457, 1994.

[71183] 9513. Levy, G. G.; Nichols, W. C.; Lian, E. C.; Foroud, T.;

McClintick, J. N.; McGee, B. M.; Yang, A. Y.; Slemieniak, D. R.; Stark, K. R.; Gruppo, R.; Sarode, R.; Shurin, S. B.; Chandrasekaran, V.; Stabler, S. P.; Sabio, H.; Bouhassira, E. E.; Upshaw, J. D., Jr.; Ginsburg, D.; Tsai, H.-M.: Mutations in a member of the ADAMTS gene family cause thrombotic thrombocytopenic purpura. *Nature* 413: 488–494, 2001.

[71184] 9514. Amrani, A.; Verdaguer, J.; Serra, P.; Tafuro, S.; Tan, R.; Santamaria, P.: Progression of autoimmune diabetes driven by avidity maturation of a T-cell population. *Nature* 406: 739–742, 2000.

[71185] 9515. Bowcock, A. M.; Kavathas, P.; Margolskee, R. F.; Herzenberg, L.; Cavalli-Sforza, L. L.: An RFLP associated with pcDLeu2–14, a human T-cell differentiation antigen CD8 (Leu2) cDNA mapped to 2p12. *Nucleic Acids Res.* 14: 7817 only, 1986.

[71186] 9516. Bruns, G.; Kavathas, P.; Shiloh, Y.; Sakai, K.; Schwaber, J.; Latt, S. A.; Herzenberg, L. A.: The human T cell antigen Leu–2 (T8) is encoded on chromosome 2. *Hum. Genet.* 70: 311–314, 1985.

[71187] 9517. Giblin, P.; Ledbetter, J. A.; Kavathas, P.: A secreted form of the human lymphocyte cell surface molecule CD8 arises from alternative splicing. *Proc. Nat. Acad. Sci.* 86: 998–1002, 1989.

- [71188] 9518.Kavathas, P.; Sukhatme, V. P.; Herzenberg, L. A.; Parnes, J. R.: Isolation of the gene encoding the human T-lymphocyte differentiation antigen Leu-2 (T8) by gene transfer and cDNA subtraction. *Proc. Nat.Acad. Sci.* 81: 7688–7692, 1984.
- [71189] 9519.Ledbetter, J. A.; Evans, R. L.; Lipinski, M.; Cunningham-Rundles,C.; Good, R. A.; Herzenberg, L. A.: Evolutionary conservation of surface molecules that distinguish T lymphocyte helper-inducer and cytotoxic-suppressor subpopulations in mouse and man. *J. Exp. Med.* 153:310–323, 1981.
- [71190] 9520.Ledbetter, J. A.; Seaman, W. E.; Tsu, T. T.; Herzenberg, L. A.: Lyt-2 and Lyt-3 antigens are on two different polypeptide subunits linked by disulfide bonds: relationship of subunits to T cell cytolytic activity. *J. Exp. Med.* 153: 1503–1516, 1981.
- [71191] 9521.Leishman, A. J.; Naidenko, O. V.; Attinger, A.; König, F.; Lena,C. J.; Xiong, Y.; Chang, H.-C.; Reinherz, E.; Kronenberg, M.; Cheroutre,H.: T cell responses modulated through interaction between CD8-alpha-alpha and the nonclassical MHC class I molecule, TL. *Science* 294: 1936–1939, 2001.
- [71192] 9522.Littman, D. R.: The structure of the CD4 and CD8



genes. *Annu.Rev. Immun.* 5: 561–584, 1987.

[71193] 9523.Littman, D. R.; Thomas, Y.; Maddon, P. J.; Chess, L.; Axel, R.: The isolation and sequence of the gene encoding T8: a molecule defining functional classes of T lymphocytes. *Cell* 40: 237–246, 1985.

[71194] 9524.Mecucci, C.; Van Den Berghe, H.: OKT8–positive T–cell lymphoma associated with a chromosome rearrangement t(2;17) possibly involving the T8 locus. (Letter) *New Eng. J. Med.* 313: 185–186, 1985.

[71195] 9525.Nakauchi, H.; Nolan, G. P.; Hsu, C.; Huang, H. S.; Kavathas, P.; Herzenberg, L. A.: Molecular cloning of Lyt–2, a membrane glycoprotein marking a subset of mouse T lymphocytes: molecular homology to its human counterpart, Leu–2/T8, and to immunoglobulin variable regions. *Proc.Nat. Acad. Sci.* 82: 5126–5130, 1985.

[71196] 9526.Lindsay, E. A.; Vitelli, F.; Su, H.; Morishima, M.; Huynh, T.; Pramparo, T.; Jurecic, V.; Ogunrinu, G.; Sutherland, H. F.; Scambler, P. J.; Bradley, A.; Baldini, A.: Tbx1 haploinsufficiency in the DiGeorges syndrome region causes aortic arch defects in mice. *Nature* 410:97–101, 2001.

[71197] 9527.Merscher, S.; Funke, B.; Epstein, J. A.; Heyer, J.; Puech, A.; Lu, M. M.; Xavier, R. J.; Demay, M. B.; Russell, R. G.; Factor, S.; Tokooya, K.; St. Jore, B.; and 12 others: TBX1

is responsible for cardiovascular defects in velo-cardio-facial/DiGeorge syndrome. *Cell* 104:619–629, 2001.

- [71198] 9528. Erikson, J.; Finger, L.; Sun, L.; ar-Rushdi, A.; Nishikura, K.; Minowada, J.; Finan, J.; Emanuel, B. S.; Nowell, P. C.; Croce, C.M.: Deregulation of c-myc by translocation of the alpha-locus of the T-cell receptor in T-cell leukemias. *Science* 232: 884–886, 1986.
- [71199] 9529. Erikson, J.; Nishikura, K.; ar-Rushdi, A.; Finan, J.; Emanuel, B.; Lenoir, G.; Nowell, P. C.; Croce, C. M.: Translocation of an immunoglobulin kappa locus to a region 3-prime of an unrearranged c-myc oncogene enhances c-myc transcription. *Proc. Nat. Acad. Sci.* 80:7581–7585, 1983.
- [71200] 9530. Feng, X.-H.; Liang, Y.-Y.; Liang, M.; Zhai, W.; Lin, X.: Direct interaction of c-Myc with Smad2 and Smad3 to inhibit TGF-beta-mediated induction of the CDK inhibitor p15(Ink4B). *Molec. Cell* 9: 133–143, 2002.
- [71201] 9531. Grandori, C.; Mac, J.; Siebelt, F.; Ayer, D. E.; Eisenman, R.N.: Myc-Max heterodimers activate a DEAD box gene and interact with multiple E box-related sites in vivo. *EMBO J.* 15: 4344–4357, 1996.
- [71202] 9532. Hamlyn, P. H.; Rabbitts, T. H.: Translocation joins c-myc and immunoglobulin gamma-1 genes in a Burkitt

lymphoma revealing a third exon in the c-myc oncogene.  
Nature 304: 135–139, 1983.

- [71203] 9533. Hayday, A. C.; Gillies, S. D.; Saito, H.; Wood, C.; Wiman, K.; Hayward, W. S.; Tonegawa, S.: Activation of a translocated human c-myc gene by an enhancer in the immunoglobulin heavy-chain locus. Nature 307:334–340, 1984.
- [71204] 9534. He, T.-C.; Sparks, A. B.; Rago, C.; Hermeking, H.; Zawel, L.; da Costa, L. T.; Morin, P. J.; Vogelstein, B.; Kinzler, K. W.: Identification of c-MYC as a target of the APC pathway. Science 281: 1509–1512, 1998.
- [71205] 9535. Heim, S.; Mitelman, F.: Nineteen of 26 cellular oncogenes precisely localized in the human genome map to one of the 83 bands involved in primary cancer-specific rearrangements. Hum. Genet. 75: 70–72, 1987.
- [71206] 9536. Henderson, A.; Ripley, S.; Heller, M.; Kieff, E.: Chromosome site for Epstein-Barr virus DNA in a Burkitt tumor cell line and in lymphocytes growth-transformed in vitro. Proc. Nat. Acad. Sci. 80:1987–1991, 1983.
- [71207] 9537. Jain, M.; Arvanitis, C.; Chu, K.; Dewey, W.; Leonhardt, E.; Trinh, M.; Sundberg, C. D.; Bishop, J. M.; Felsher, D. W.: Sustained loss of a neoplastic phenotype by brief inactivation of MYC. Science 297:102–104, 2002.

- [71208] 9538.Leder, P.: Personal Communication. Boston, Mass. 10/1/1982.
- [71209] 9539.Lee, T. C.; Li, L.; Philipson, L.; Ziff, E. B.: Myc re-  
pressestranscription of the growth arrest gene gas1. Proc.  
Nat. Acad. Sci. 94:12886–12891, 1997.
- [71210] 9540.Leven, D.: Disentangling the MYC web.  
(Commentary) Proc. Nat.Acad. Sci. 99: 5757–5759, 2002.
- [71211] 9541.Lin, C. S.; Goldthwait, D. A.; Samols, D.: Identifica-  
tion ofAlu transposition in human lung carcinoma cells.  
Cell 54: 153–159,1988.
- [71212] 9542.Mager, D. L.; Goodchild, N. L.: Homologous recom-  
bination between the LTRs of a human retrovirus-like ele-  
ment causes a 5-kb deletion in two siblings. Am. J. Hum.  
Genet. 45: 848–854, 1989.
- [71213] 9543.Magrath, I.; Erikson, J.; Whang-Peng, J.; Sieverts, H.;  
Armstrong, G.; Benjamin, D.; Triche, T.; Alabaster, O.;  
Croce, C. M.: Synthesis of kappa light chains by cell lines  
containing an 8;22 chromosomal translocation derived  
from a male homosexual with Burkitt's lymphoma. Science  
222:1094–1098, 1983.
- [71214] 9544.Maguire, R. T.; Robins, T. S.; Thorgeirsson, S. S.;  
Heilman, C.A.: Expression of cellular myc and mos genes  
in undifferentiated B cell lymphomas of Burkitt and non-

Burkitt types. Proc. Nat. Acad.Sci. 80: 1947–1950, 1983.

- [71215] 9545.Manolov, G.; Manolova, Y.: Marker band in one chromosome 14 fromBurkitt lymphomas. Nature 237: 33–34, 1972.
- [71216] 9546.Marcu, K. B.; Harris, L. J.; Stanton, L. W.; Erikson, J.; Watt,R.; Croce, C. M.: Transcriptionally active c–myc oncogene is containedwithin NIARD, a DNA sequence associated with chromosome translocationsin B–cell neoplasia. Proc. Nat. Acad. Sci. 80: 519–523, 1983.
- [71217] 9547.Menssen, A.; Hermeking, H.: Characterization of the c–MYC–regulatedtranscriptome by SAGE: identification and analysis of c–MYC targetgenes. Proc. Nat. Acad. Sci. 99: 6274–6279, 2002.
- [71218] 9548.Mitelman, F.: Catalog of Chromosome Aberrations in Cancer. New York: Alan R. Liss (pub.) (2nd ed.): 1985.
- [71219] 9549.Morse, B.; Rotherg, P. G.; South, V. J.; Spandorfer, J. M.; Astrin,S. M.: Insertional mutagenesis of the myc locus by a LINE–1 sequencein a human breast carcinoma. Nature 333: 87–90, 1988.
- [71220] 9550.Murphy, W.; Sarid, J.; Taub, R.; Vasicek, T.; Battey, J.; Lenoir,G.; Leder, P.: A translocated human c–myc oncogene is altered ina conserved coding sequence. Proc. Nat. Acad. Sci. 83: 2939–2943,1986.

- [71221] 9551.Nishikura, K.; ar-Rushdi, A.; Erikson, J.; Watt, R.; Rovera, G.;Croce, C. M.: Differential expression of the normal and of the translocatedhuman c-myc oncogenes in B cells. *Proc. Nat. Acad. Sci.* 80: 4822–4826,1983.
- [71222] 9552.Pasqualucci, L.; Neumeister, P.; Goossens, T.; Nan-jangud, G.;Chaganti, R. S. K.; Kuppers, R.; Dalla-Favera, R.: Hypermutationof multiple proto-oncogenes in B-cell diffuse large-cell lymphomas. *Nature* 412:341–346, 2001.
- [71223] 9553.Pelengaris, S.; Khan, M.; Evan, G. I.: Suppression of Myc-inducedapoptosis in beta cells exposes multiple oncogenic properties of Mycand triggers carcinogenic progression. *Cell* 109: 321–334, 2002.
- [71224] 9554.Persson, H.; Hennighausen, L.; Taub, R.; DeGrado, W.; Leder, P.: Antibodies to human c-myc oncogene prod-uct: evidence of an evolutionarilyconserved protein in-duced during cell proliferation. *Science* 225:687–693, 1984.
- [71225] 9555.Persson, H.; Leder, P.: Nuclear localization and DNA bindingproperties of a protein expressed by human c-myc oncogene. *Science* 225:718–721, 1984.
- [71226] 9556.Peukert, K.; Staller, P.; Schneider, A.; Carmichael, G.; Hanel,F.; Eilers, M.: An alternative pathway for gene regu-lation by Myc. *EMBOJ.* 16: 5672–5686, 1997.

- [71227] 9557.Saito, H.; Hayday, A. C.; Wiman, K.; Hayward, W. S.; Tonegawa,S.: Activation of the c-myc gene by translocation: a model for translational control. Proc. Nat. Acad. Sci. 80: 7476–7480, 1983.
- [71228] 9558.Brookman, K. W.; Tebbs, R. S.; Allen, S. A.; Tucker, J. D.; Swiger,R. R.; Lamerdin, J. E.; Carrano, A. V.; Thompson, L. H.: Isolation and characterization of mouse Xrcc1, a DNA repair gene affecting ligation. Genomics 22:180–188, 1994.
- [71229] 9559.Lamerdin, J. E.; Montgomery, M. A.; Stilwagen, S. A.; Scheidecker,L. K.; Tebbs, R. S.; Brookman, K. W.; Thompson, L. H.; Carrano, A.V.: Genomic sequence comparison of the human and mouse XRCC1 DNA repair gene regions. Genomics 25: 547–554, 1995.
- [71230] 9560.Thompson, L. H.; Bachinski, L. L.; Stallings, R. L.; Dolf, G.;Weber, C. A.; Westerveld, A.; Siciliano, M. J.: Complementation of repair gene mutations on the hemizygous chromosome 9 in CHO: a third repair gene on human chromosome 19. Genomics 5: 670–679, 1989.
- [71231] 9561.Whitehouse, C. J.; Taylor, R. M.; Thistlethwaite, A.; Zhang, H.;Karimi-Busheri, F.; Lasko, D. D.; Weinfeld, M.; Caldecott, K. W.:XRCC1 stimulates human polynucleotide kinase activity at damaged DNA termini and accelerates

DNA single-strand break repair. *Cell* 104:107–117, 2001.

- [71232] 9562. Gao, Y.; Sun, Y.; Frank, K. M.; Dikkes, P.; Fujiwara, Y.; Seidl, K. J.; Sekiguchi, J. M.; Rathbun, G. A.; Swat, W.; Wang, J.; Bronson, R. T.; Malynn, B. A.; Bryans, M.; Zhu, C.; Chaudhuri, J.; Davidson, L.; Ferrini, R.; Stamato, T.; Orkin, S. H.; Greenberg, M. E.; Alt, F. W.: A critical role for DNA end-joining proteins in both lymphogenesis and neurogenesis. *Cell* 95: 891–902, 1998.
- [71233] 9563. Giaccia, A. J.; Denko, N.; MacLaren, R.; Mirman, D.; Waldren, C.; Hart, I.; Stamato, T. D.: Human chromosome 5 complements the DNA double-strand break-repair deficiency and gamma-ray sensitivity of the XR-1 hamster variant. *Am. J. Hum. Genet.* 47: 459–469, 1990.
- [71234] 9564. Li, Z.; Otevrel, T.; Gao, Y.; Cheng, H.-L.; Seed, B.; Stamato, T. D.; Taccioli, G. E.; Alt, F. W.: The XRCC4 gene encodes a novel protein involved in DNA double-strand break repair and V(D)J recombination. *Cell* 83:1079–1089, 1995.
- [71235] 9565. Otevrel, T.; Stamato, T. D.: Regional localization of the XRCC4 human radiation repair gene. *Genomics* 27: 211–214, 1995.
- [71236] 9566. Blunt, T.; Taccioli, G. E.; Priestley, A.; Hafezparast, M.; McMillan, T.; Liu, J.; Cole, C. C.; White, J.; Alt, F. W.;



Jackson, S. P.; Schurr, E.; Lehmann, A. R.; Jeggo, P. A.: A YAC contig encompassing the XRCC5(Ku80) DNA repair gene and complementation defective cells by YAC protoplast fusion. *Genomics* 30: 320–328, 1995.

[71237] 9567. Fuentes, J.-J.; Pritchard, M. A.; Planas, A. M.; Bosch, A.; Ferrer, I.; Estivill, X.: A new human gene from the Down syndrome critical region encodes a proline-rich protein highly expressed in fetal brain and heart. *Hum. Molec. Genet.* 4: 1935–1944, 1995.

[71238] 9568. Magnussen, K.: Beitrag zur Genetik und Histologie eines isolierten Augen albinismus beim Kaninchen. *Z. Morph. Anthropol.* 44: 127–135, 1952.

[71239] 9569. Sukhatme, V. P.; Sizer, K. C.; Vollmer, A. C.; Hunkapiller, T.; Parnes, J. R.: The T cell differentiation antigen leu-2/T8 is homologous to immunoglobulin and T cell receptor variable regions. *Cell* 40: 591–597, 1985.

[71240] 9570. Sakaguchi, A. Y.; Lalley, P. A.; Naylor, S. L.: Human and mouse cellular myc protooncogenes reside on chromosomes involved in numerical and structural aberrations in cancer. *Somat. Cell Genet.* 9: 391–405, 1983.

[71241] 9571. Chen, D. J.; Marrone, B. L.; Nguyen, T.; Stackhouse, M.; Zhao, Y.; Siciliano, M. J.: Regional assignment of a human DNA repair gene (XRCC5) to 2q35 by x-ray hybrid

mapping. *Genomics* 21: 423–427, 1994.

[71242] 9572.Kirchgessner, C. U.; Patil, C. K.; Evans, J. W.; Cuomo, C. A.; Fried, L. M.; Carter, T.; Oettinger, M. A.; Brown, J. M.: DNA-dependent kinase (p350) as a candidate gene for the murine SCID defect. *Science* 267:1178–1183, 1995.

[71243] 9573.Oakley, C. E.; Oakley, B. R.: Identification of gamma-tubulin, a new member of the tubulin superfamily encoded by mipA gene of *Aspergillus nidulans*. *Nature* 338: 662–664, 1989.

[71244] 9574.Rommens, J. M.; Durocher, F.; McArthur, J.; Tonin, P.; LeBlanc, J.-F.; Allen, T.; Samson, C.; Ferri, L.; Narod, S.; Morgan, K.; Simard, J.: Generation of a transcription map at the HSD17B locus centromeric to BRCA1 at 17q21. *Genomics* 28: 530–542, 1995.

[71245] 9575.Simerly, C.; Wu, G.-J.; Zoran, S.; Ord, T.; Rawlins, R.; Jones, J.; Navara, C.; Gerrity, M.; Rinehart, J.; Binor, Z.; Asch, R.; Schatten, G.: The paternal inheritance of the centrosome, the cell's microtubule-organizing center, in humans, and the implications for infertility. *Nature Med.* 1:47–52, 1995.

[71246] 9576.Stearns, T.; Evans, L.; Kirschner, M.: Gamma-tubulin is a highly conserved component of the centrosome. *Cell* 65: 825–836, 1991.

- [71247] 9577.Wise, D. O.; Krahe, R.; Oakley, B. R.: The gamma-tubulin gene family in humans. *Genomics* 67: 164–170, 2000.
- [71248] 9578.Zheng, Y.; Jung, M. K.; Oakley, B. R.: Gamma-tubulin is present in *Drosophila melanogaster* and *Homo sapiens* and is associated with the centrosome. *Cell* 65: 817–823, 1991.
- [71249] 9579.Bettaieb, A.; Fromont, P.; Rodet, M.; Godeau, B.; Duedari, N.; Bierling, P.: Br(b), a platelet alloantigen involved in neonatal alloimmune thrombocytopenia. *Vox Sang.* 60: 230–234, 1991.
- [71250] 9580.Carlsson, L. E.; Santoso, S.; Spitzer, C.; Kessler, C.; Greinacher, A.: The alpha-2 gene coding sequence T807/A873 of the platelet collagen receptor integrin alpha-2/beta-1 might be a genetic risk factor for the development of stroke in younger patients. *Blood* 93: 3583–3586, 1999.
- [71251] 9581.Chen, D. J.; Park, M. S.; Campbell, E.; Oshimura, M.; Liu, P.; Zhao, Y.; White, B. F.; Siciliano, M. J.: Assignment of a human DNA double-strand break repair gene (XRCC5) to chromosome 2. *Genomics* 13:1088–1094, 1992.
- [71252] 9582.Difilippantonio, M. J.; Zhu, J.; Chen, H. T.; Meffre, E.; Nussenzweig, M. C.; Max, E. E.; Ried, T.; Nussenzweig, A.:

DNA repair protein Ku80 suppresses chromosomal aberrations and malignant transformation. *Nature* 404:510–514, 2000.

- [71253] 9583. Jeggo, P. A.; Hafezparast, M.; Thompson, A. F.; Broughton, B. C.; Kaur, G. P.; Zdzienicka, M. Z.; Athwal, R. S.: Localization of a DNA repair gene (XRCC5) involved in double-strand-break rejoining to human chromosome 2. *Proc. Nat. Acad. Sci.* 89: 6423–6427, 1992.
- [71254] 9584. Taccioli, G. E.; Gottlieb, T. M.; Blunt, T.; Priestley, A.; Demengeot, J.; Mizuta, R.; Lehmann, A. R.; Alt, F. W.; Jackson, S. P.; Jeggo, P. A.: Ku80: product of the XRCC5 gene and its role in DNA repair and V(D)J recombination. *Science* 265: 1442–1445, 1994.
- [71255] 9585. Ohkuchi, A.; Shiraishi, H.; Minakami, H.; Eguchi, Y.; Izumi, A.; Sato, I.: Fetus with long QT syndrome manifested by tachyarrhythmia: a case report. *Prenat. Diag.* 19: 990–992, 1999.
- [71256] 9586. Pacia, S. V.; Devinsky, O.; Luciano, D. J.; Vazquez, B.: The prolonged QT syndrome presenting as epilepsy: a report of two cases and literature review. *Neurology* 44: 1408–1410, 1994.
- [71257] 9587. Piippo, K.; Swan, H.; Pasternack, M.; Chapman, H.; Paavonen, K.; Viitasalo, M.; Toivonen, L.; Kontula, K.: A

founder mutation of the potassium channel KCNQ1 in long QT syndrome: implications for estimation of disease prevalence and molecular diagnostics. *J. Am. Coll. Cardiol.* 37:562–568, 2001.

- [71258] 9588. Pony, J. C.; Matheyses, M.; Daubert, J. C.; Fourdilis, M.; Gouffault, J.: Le syndrome QT long–syncope familial: deux observations de syndrome Romano et Ward. *Arch. Mal. Coeur* 70: 1105–1114, 1977.
- [71259] 9589. Priori, S. G.; Schwartz, P. J.; Napolitano, C.; Bianchi, L.; Dennis, A.; De Fusco, M.; Brown, A. M.; Casari, G.: A recessive variant of the Romano–Ward Long–QT syndrome? *Circulation* 97: 2420–2425, 1998.
- [71260] 9590. Romano, C.: Congenital cardiac arrhythmia. (Letter) *Lancet* i:658–659, 1965.
- [71261] 9591. Romano, C.; Gemme, G.; Pongiglione, R.: Aritmie cardiache rare dell'età pediatrica. II. Accessi sincopali per fibrillazione ventricolare parossistica. (Presentazione del primo caso della letteratura pediatrica italiana.) *Clin. Pediat.* 45: 656–683, 1963.
- [71262] 9592. Roy, N.; Kahlem, P.; Dausse, E.; Bennaceur, M.; Faure, S.; Weissenbach, J.; Komajda, M.; Denjoy, I.; Coumel, P.; Schwartz, K.; Guicheney, P.: Exclusion of HRAS from long QT locus. (Letter) *Nature Genet.* 8:113–114, 1994.

- [71263] 9593.Roy, P. R.; Emanuel, R.; Ismail, S. A.; Tayib, M. H.: Hereditaryprolongation of the Q-T interval: genetic observations and managementin three families with 12 affected members. *Am. J. Cardiol.* 37:237-243, 1976.
- [71264] 9594.Russell, M. W.; Dick, M., II; Collins, F. S.; Brody, L. C.: KVLQT1mutations in three families with familial or sporadic long QT syndrome. *Hum.Molec. Genet.* 5: 1319-1324, 1996.
- [71265] 9595.Schmitt, N.; Schwarz, M.; Peretz, A.; Abitbol, I.; Attali, B.;Pongs, O.: A recessive C-terminal Jervell and Lange-Nielsen mutationof the KCNQ1 channel impairs subunit assembly. *EMBO J.* 19: 332-340,2000.
- [71266] 9596.Schwartz, P. J.: Cardiac sympathetic innervation and the suddeninfant death syndrome: a possible pathogenetic link. *Am. J. Med.* 60:167-172, 1976.
- [71267] 9597.Schwartz, P. J.; Priori, S. G.; Bloise, R.; Napolitano, C.; Ronchetti,E.; Piccinini, A.; Goj, C.; Breithardt, G.; Schulze-Bahr, E.; Wedekind,H.; Nastoli, J.: Molecular diagnosis in a child with sudden infantdeath syndrome. (Letter) *Lancet* 358: 1342-1343, 2001.
- [71268] 9598.Schwartz, P. J.; Stramba-Badiale, M.; Segantini, A.; Austoni,P.; Bosi, G.; Giorgetti, R.; Grancini, F.; Marni, E. D.; Perticone,F.; Rosti, D.; Salice, P.: Prolongation of the QT

interval and the sudden infant death syndrome. *New Eng. J. Med.* 338: 1709–1714, 1998.

[71269] 9599. Shalaby, F. Y.; Levesque, P. C.; Yang, W.-P.; Little, W. A.; Conder, M. L.; Jenkins-West, T.; Blannar, M. A.: Dominant-negative KvLQT1 mutations underlie the LQT1 form of long QT syndrome. *Circulation* 96:1733–1736, 1997.

[71270] 9600. Shimizu, W.; Kurita, T.; Matsuo, K.; Suyama, K.; Aihara, N.; Kamakura, S.; Towbin, J. A.; Shimomura, K.: Improvement of repolarization abnormalities by a K<sup>+</sup> channel opener in the LQT1 form of congenital long-QT syndrome. *Circulation* 97:1581–1588, 1998.

[71271] 9601. Singer, P. A.; Crampton, R. S.; Bass, N. H.: Familial Q-T prolongation syndrome: convulsive seizures and paroxysmal ventricular fibrillation. *Arch. Neurol.* 31: 64–66, 1974.

[71272] 9602. Smolinich, N. J.; Day, C. D.; Fitzpatrick, G. V.; Caldwell, G. M.; Lossie, A. C.; Cooper, P. R.; Smallwood, A. C.; Joyce, J. A.; Schofield, P. N.; Reik, W.; Nicholls, R. D.; Weksberg, R.; Driscoll, D. J.; Maher, E. R.; Shows, T. B.; Higgins, M. J.: A maternally methylated CpG island in KvLQT1 is associated with an antisense paternal transcript and loss of imprinting in Beckwith-Wiedemann syndrome. *Proc. Nat. Acad. Sci.* 96: 8064–8069, 1999.

- [71273] 9603.Splawski, I.; Timothy, K. W.; Vincent, G. M.; Atkinson, D. L.; Keating, M. T.: Molecular basis of the long-QT syndrome associated with deafness. *New Eng. J. Med.* 336: 1562–1567, 1997.
- [71274] 9604.Campbell, H. D.; Webb, G. C.; Young, I. G.: A human homologue of the *Drosophila melanogaster* sluggish-A (proline oxidase) gene maps to 22q11.2, and is a candidate gene for type-I hyperproliferative anaemia. *Hum. Genet.* 101: 69–74, 1997.
- [71275] 9605.Slominski, A.; Ermak, G.; Mihm, M.: ACTH receptor, CYP11A1, CYP17 and CYP21A2 genes are expressed in skin. *J. Clin. Endocr. Metab.* 81:2746–2749, 1996.
- [71276] 9606.Touraine, J. L.; Betuel, H.; Souillet, G.; Jeune, M.: Combined immunodeficiency disease associated with absence of cell-surface HLA-A and -B antigens. *J. Pediatr.* 93: 47–51, 1978.
- [71277] 9607.Wolf, H. M.; Hauber, I.; Gulle, H.; Thon, V.; Eggenbauer, H.; Fischer, M. B.; Fiala, S.; Eibl, M. M.: Brief report: Twin boys with major histocompatibility complex class II deficiency but inducible immune responses. *New Eng. J. Med.* 332: 86–90, 1995.
- [71278] 9608.Dupuis, S.; Dargemont, C.; Fieschi, C.; Thomassin, N.; Rosenzweig, S.; Harris, J.; Holland, S. M.; Schreiber, R.



D.; Casanova, J.-L.: Impairment of mycobacterial but not viral immunity by a germline human STAT1 mutation. *Science* 293: 300–303, 2001.

[71279] 9609. Findlay, K. A. B.; Kaptein, E.; Visser, T. J.; Burchell, B.: Characterization of the uridine diphosphate–glucuronosyltransferase–catalyzing thyroid hormone glucuronidation in man. *J. Clin. Endocr. Metab.* 85: 2879–2883, 2000.

[71280] 9610. Wooster, R.; Sutherland, L.; Ebner, T.; Clarke, D.; Da Cruz e Silva, O.; Burchell, B.: Cloning and stable expression of a new member of the human liver phenol/bilirubin:UDP–glucuronosyltransferase cDNA family. *Biochem. J.* 278: 465–469, 1991.

[71281] 9611. Tanaka, S.; Yamashita, S.; Hosaka, K.: Cloning and expression of human cDNA encoding phosphatidylinositol transfer protein beta. *Biochim. Biophys. Acta* 1259: 199–202, 1995.

[71282] 9612. Choi, Y.-H.; Kim, K.-B.; Kim, H.-H.; Hong, G.-S.; Kwon, Y.-K.; Chung, C.-W.; Park, Y.-M.; Shen, Z.-J.; Kim, B. J.; Lee, S.-Y.; Jung, Y.-K.: FLASH coordinates NF- $\kappa$ B activity via TRAF2. *J. Biol. Chem.* 276: 25073–25077, 2001.

[71283] 9613. Imai, Y.; Kimura, T.; Murakami, A.; Yajima, N.; Saka-

maki, K.; Yonehara,S.: The CED-4-homologous protein FLASH is involved in Fas-mediatedactivation of caspase-8 during apoptosis. *Nature* 398: 777-785, 1999.

[71284] 9614.Cossee, M.; Durr, A.; Schmitt, M.; Dahl, N.; Trouillas, P.; Allinson,P.; Kostrzewa, M.; Nivelon-Chevallier, A.; Gustavson, K.-H.; Kohlschutter,A.; Muller, U.; Mandel, J.-L.; and 11 others: Friedreich's ataxia:point mutations and clinical presentation of compound heterozygotes. *Ann.Neurol.* 45: 200-206, 1999.

[71285] 9615.De Castro, M.; Garcia-Planells, J.; Monros, E.; Canizares, J.;Vazquez-Manrique, R.; Vilchez, J. J.; Urtasun, M.; Lucus, M.; Navarro,G.; Izquierdo, G.; Molto, M. D.; Palau, F.: Genotype and phenotypeanalysis of Friedreich's ataxia compound heterozygous patients. *Hum.Genet.* 106: 86-92, 2000.

[71286] 9616.Delatycki, M. B.; Paris, D.; Gardner, R. J. M.; Forshaw, K.; Nicholson,G. A.; Nassif, N.; Williamson, R.; Forrest, S. M.: Sperm DNA analysisin a Friedreich ataxia premutation carrier suggests both meiotic andmitotic expansion in the FRDA gene. *J. Med. Genet.* 35: 713-716,1998.

[71287] 9617.Duclos, F.; Boschert, U.; Sirugo, G.; Mandel, J.-L.; Hen, R.;Koenig, M.: Gene in the region of the Friedreich ataxia locus encodesa putative transmembrane protein

expressed in the nervous system. *Proc.Nat. Acad. Sci.* 90: 109–113, 1993.

- [71288] 9618.Doi, A.; Shiosaka, T.; Takaoka, Y.; Yanagisawa, K.; Fujita, S.: Molecular cloning of the cDNA encoding A+U-rich element RNA bindingfactor. *Biochim. Biophys. Acta* 1396: 51–56, 1998.
- [71289] 9619.Kamei, D.; Tsuchiya, N.; Yamazaki, M.; Meguro, H.; Yamada, M.:Two forms of expression and genomic structure of the human heterogeneousnuclear ribonucleoprotein D-like JKTBP gene (HNRPDL). *Gene* 228:13–22, 1999.
- [71290] 9620.Tsuchiya, N.; Kamei, D.; Takano, A.; Matsui, T.; Yamada, M.: Cloningand characterization of a cDNA encoding a novel heterogeneous nuclearribonucleoprotein-like protein and its expression in myeloid leukemiacells. *J. Biochem.* 123: 499–507, 1998.
- [71291] 9621.Ma, Y. H.; Betard, C.; Roy, M.; Davignon, J.; Kessling, A. M.: Identification of a second 'French Canadian' LDL receptor gene deletionand development of a rapid method to detect both deletions. *Clin.Genet.* 36: 219–228, 1989.
- [71292] 9622.Mandelstam, M.; Chakir, K.; Shevtsov, S.; Golubkov, V.; Skobeleva,N.; Lipovetsky, B.; Konstantinov, V.; Denisenko, A.; Gaitskhoki, V.;Schwartz, E.: Prevalence of Lithuanian mutation among St. PetersburgJews with famil–

ial hypercholesterolemia. Hum. Mutat. 12: 255–258,1998.

[71293] 9623.Meiner, V.; Landsberger, D.; Berkman, N.; Reshef, A.; Segal, P.; Seftel, H. C.; van der Westhuyzen, D. R.; Jeenah, M. S.; Coetzee, G. A.; Leitersdorf, E.: A common Lithuanian mutation causing familial hypercholesterolemia in Ashkenazi Jews. Am. J. Hum. Genet. 49: 443–449,1991.

[71294] 9624.Miyake, Y.; Tajima, S.; Funahashi, T.; Yamamura, T.; Yamamoto, A.: A point mutation of low-density-lipoprotein receptor causing rapid degradation of the receptor. Europ. J. Biochem. 210: 1–7,1992.

[71295] 9625.Moorjani, S.; Roy, M.; Torres, A.; Betard, C.; Gagne, C.; Lambert, M.; Brun, D.; Davignon, J.; Lupien, P.: Mutations of low-density-lipoprotein-receptor gene, variation in plasma cholesterol, and expression of coronary heart disease in homozygous familial hypercholesterolaemia. Lancet 341:1303–1306, 1993.

[71296] 9626.Oppenheim, A.; Friedlander, Y.; Dann, E. J.; Berkman, N.; Schwartz, S. P.; Leitersdorf, E.: Hypercholesterolemia in five Israeli Christian-Arab kindreds is caused by the 'Lebanese' allele at the low density lipoprotein receptor gene locus and by an additional independent major factor. Hum. Genet. 88: 75–84, 1991.

[71297] 9627.Pisciotta, L.; Cantafora, A.; De Stefano, F.; Langheim,

S.; Calandra, S.; Bertolini, S.: A 'de novo' mutation of the LDL-receptor gene as the cause of familial hypercholesterolemia. *Biochim. Biophys. Acta* 1587: 7–11, 2002.

[71298] 9628. Rodningen, O. K.; Rosby, O.; Tonstad, S.; Ose, L.; Berg, K.; Leren, T. P.: A 9.6 kilobase deletion in the low density lipoprotein receptor gene in Norwegian familial hypercholesterolemia subjects. *Clin. Genet.* 42:288–295, 1992.

[71299] 9629. Rubinsztein, D. C.; Coetzee, G. A.; Marais, A. D.; Leitersdorf, E.; Seftel, H. C.; van der Westhuyzen, D. R.: Identification and properties of the proline-644-to-leucine mutant LDL receptor in South Africans of Indian origin. *J. Lipid Res.* 33: 1647–1655, 1992.

[71300] 9630. Rudiger, N. S.; Heinsvig, E. M.; Hansen, F. A.; Faergeman, O.; Bolund, L.; Gregersen, N.: DNA deletions in the low density lipoprotein (LDL) receptor gene in Danish families with familial hypercholesterolemia. *Clin. Genet.* 39: 451–462, 1991.

[71301] 9631. Ruffner, D. E.; Sprung, C. N.; Minghetti, P. P.; Gibbs, P. E. M.; Dugaiczyk, A.: Invasion of the human albumin-alpha-fetoprotein gene family by Alu, Kpn, and two novel repetitive DNA elements. *Molec. Biol. Evol.* 4: 1–9, 1987.

[71302] 9632. Russell, D. W.; Lehrman, M. A.; Sudhof, T. C.; Ya-

mamoto, T.; Davis, C. G.; Hobbs, H. H.; Brown, M. S.; Goldstein, J. L.: The LDL receptor in familial hypercholesterolemia: use of human mutations to dissect a membrane protein. Cold Spring Harbor Symp. Quant. Biol. 51: 811–819, 1986.

[71303] 9633. Russell, D. W.; Schneider, W. J.; Yamamoto, T.; Luskey, K. L.; Brown, M. S.; Goldstein, J. L.: Domain map of the LDL receptor: sequence homology with the epidermal growth factor precursor. Cell 37: 577–585, 1984.

[71304] 9634. Schuster, H.; Manke, C.; Fischer, J.; Keller, C.; Wolfram, G.; Zollner, N.: Identification of the valine 408 to methionine mutation in the LDL receptor in a Greek patient with homozygous familial hypercholesterolemia. Clin. Genet. 48: 90–92, 1995.

[71305] 9635. Schuster, H.; Ostwald, P.; Keller, P.; Wolfram, G.; Keller, C.: Identification of the serine–156 to leucine mutation in the low-density lipoprotein receptor in a German family with familial hypercholesterolemia. Clin. Investig. 71: 172–175, 1993.

[71306] 9636. Slagel, V.; Flemington, E.; Traina-Dorge, V.; Bradshaw, H.; Deininger, P.: Clustering and subfamily relationships of the Alu family in the human genome. Molec. Biol. Evol. 4: 19–29, 1987. 100. Soutar, A. K.; Knight, B. L.; Pa-

tel, D. D.: Identification of a point mutation in growth repeat C of the low density lipoprotein-receptor gene in a patient with homozygous familial hypercholesterolemia that affects ligand binding and intracellular movement of receptors. *Proc. Nat. Acad. Sci.* 86: 4166–4170, 1989.101.

Soutar, A. K.; McCarthy, S. N.; Seed, M.; Knight, B. L.: Relationship between apolipoprotein(a) phenotype, lipoprotein(a) concentration in plasma, and low density lipoprotein receptor function in a large kindred with familial hypercholesterolemia due to the pro664-to-leu mutation in the LDL receptor gene. *J. Clin. Invest.* 88: 483–492, 1991.102.

Steyn, K.; Weight, M. J.; Dando, B. R.; Christopher, K. J.; Rossouw, J. E.: The use of low density lipoprotein receptor activity of lymphocytes to determine the prevalence of familial hypercholesterolaemia in a rural South African community. *J. Med. Genet.* 26: 32–36, 1989.103.

Sudhof, T. C.; Goldstein, J. L.; Brown, M. S.; Russell, D. W.: The LDL receptor gene: a mosaic of exons shared with different proteins. *Science* 228: 815–822, 1985.104.

Sudhof, T. C.; Russell, D. W.; Goldstein, J. L.; Brown, M. S.; Sanchez-Pescador, R.; Bell, G. I.: Cassette of eight exons shared by genes for LDL receptor and EGF precursor. *Science* 228: 893–895, 1985.105.

Sun, X.-M.;

Patel, D. D.; Bhatnager, D.; Knight, B. L.; Soutar, A. K.: Characterization of a splice-site mutation in the gene for the LDL receptor associated with an unpredictably severe clinical phenotype in English patients with heterozygous FH. *Arterioscler. Thromb. Vasc. Biol.* 15: 219–227, 1995.106. Takahashi, M.; Ikeda, U.; Takahashi, S.; Hattori, H.; Iwasaki, T.; Ishihara, M.; Egashira, T.; Honma, S.; Asano, Y.; Shimada, K.: A novel mutation in exon 2 of the low-density lipoprotein-receptor gene in a patient with homozygous familial hypercholesterolemia. (Letter) *Clin. Genet.* 59: 290–292, 2001.107. Taylor, R.; Bryant, J.; Gudnason, V.; Sigurdsson, G.; Humphries, S.: A study of familial hypercholesterolaemia in Iceland using RFLPs. *J. Med. Genet.* 26: 494–498, 1989.108. Thiart, R.; Scholtz, C. L.; Vergotine, J.; Hoogendijk, C. F.; de Villiers, J. N. P.; Nissen, H.; Brusgaard, K.; Gaffney, D.; Hoffs, M. S.; Vermaak, W. J. H.; Kotze, M. J.: Predominance of a 6 bp deletion in exon 2 of the LDL receptor gene in Africans with familial hypercholesterolaemia. *J. Med. Genet.* 37: 514–519, 2000.109. Tolleshaug, H.; Goldstein, J. L.; Schneider, W. J.; Brown, M. S.: Posttranslational processing of the LDL receptor and its genetic disruption in familial hypercholesterolemia. *Cell* 30: 715–724, 1982.110. Top, B.; Koele-



man, B. P. C.; Gevers Leuven, J. A.; Havekes, L.M.; Frants, R. R.: Rearrangements in the LDL receptor gene in Dutch-familial hypercholesterolemic patients and the presence of a common 4 kb deletion. *Atherosclerosis* 83: 127–136, 1990.111. Top, B.; Uitterlinden, A. G.; van der Zee, A.; Kastelein, J.J. P.; Gevers Leuven, J. A.; Havekes, L. M.; Frants, R. R.: Absence of mutations in the promoter region of the low density lipoprotein receptor gene in a large number of familial hypercholesterolaemia patients as revealed by denaturing gradient gel electrophoresis. *Hum.Genet.* 89: 561–565, 1992.112. Torrington, M.; Botha, J. L.: Familial hypercholesterolaemia and church affiliation. (Letter) *Lancet* II: 1120 only, 1981.113. Ullu, E.; Tschudi, C.: Alu sequences are processed 7SL RNA genes. *Nature* 312:171–172, 1984.114. Varret, M.; Rabes, J.-P.; Collod-Beroud, G.; Junien, C.; Boileau, C.; Beroud, C.: Software and database for the analysis of mutations in the human LDL receptor gene. *Nucleic Acids Res.* 25: 172–180, 1997.115. Vergopoulos, A.; Bajari, T.; Jouma, M.; Knoblauch, H.; Aydin, A.; Bähring, S.; Mueller-Myhsok, B.; Dresel, A.; Joubran, R.; Luft, F. C.; Schuster, H.: A xanthomatosis-susceptibility gene may exist in a Syrian family with familial hypercholesterolemia. *Europ. J.Hum. Genet.*

5: 315–323, 1997.116. Vergotine, J.; Thiart, R.; Langenhoven, E.; Hillermann, R.; DeJong, G.; Kotze, M. J.: Prenatal diagnosis of familial hypercholesterolemia: importance of DNA analysis in the high-risk South African population. *Genet.Counsel.* 12: 121–127, 2001.117. Vuorio, A. F.; Turtola, H.; Piilahti, K.–M.; Repo, P.; Kanninen, T.; Kontula, K.: Familial hypercholesterolemia in the Finnish NorthKarelia: a molecular, clinical, and genealogical study. *Arterioscler.Thromb. Vasc. Biol.* 17: 3127–3138, 1997.118. Wilson, D. J.; Gahan, M.; Haddad, L.; Heath, K.; Whittall, R.A.; Williams, R. R.; Humphries, S. E.; Day, I. N. M.: A World WideWeb site for low-density lipoprotein receptor gene mutations in familial hypercholesterolemia: sequence-based, tabular, and direct submission data handling. *Am. J. Cardiol.* 81: 1509–1511, 1998.119. Yamakawa, K.; Okafuji, T.; Iwamura, Y.; Yuzawa, K.; Satoh, J.; Hattori, N.; Yamanouchi, Y.; Yanagi, H.; Kawai, K.; Tsuchiya, S.; Russell, D. W.; Hamaguchi, H.: TaqI polymorphism in the LDL receptor gene and a TaqI 1.5-kb band associated with familial hypercholesterolemia. *Hum.Genet.* 80: 1–5, 1988.120. Yamakawa, K.; Takada, K.; Yanagi, H.; Tsuchiya, S.; Kawai, K.; Nakagawa, S.; Kajiyama, G.; Hamaguchi, H.: Three novel partial deletions of the low-density

lipoprotein (LDL) receptor gene in familial hypercholesterolemia. Hum.Genet. 82: 317–321, 1989.121. Yamamoto, T.; Davis, C. G.; Brown, M. S.; Schneider, W. J.; Casey, M. L.; Goldstein, J. L.; Russell, D. W.: The human LDL receptor: a cysteine-rich protein with multiple Alu sequences in its mRNA. Cell 39:27–38, 1984.122. Zuliani, G.; Hobbs, H. H.: Personal Communication. Dallas, Tex. 1990.

[71307] 9637. Ishibashi, K.; Suzuki, M.; Sasaki, S.; Imai, M.: Identification of a new multigene four transmembrane family (MS4A) related to CD20, HTm4 and beta subunit of the high affinity IgE receptor. Gene 264:87–93, 2001.

[71308] 9638. Liang, Y.; Tedder, T. F.: Identification of a CD20-, Fc-epsilon-RI-beta-, and HTm4-related gene family: sixteen new MS4A family members expressed in human and mouse. Genomics 72: 119–127, 2001.

[71309] 9639. Nagase, T.; Kikuno, R.; Hattori, A.; Kondo, Y.; Okumura, K.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XIX. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. DNA Res. 7: 347–355, 2000.

[71310] 9640. Gingras, M. C.; Lapillonne, H.; Margolin, J. F.: CFFM4: a new member of the CD20/Fc-epsilon-RI-beta

family. *Immunogenetics* 53:468–476, 2001.

- [71311] 9641.O'Carroll, D.; Scherthan, H.; Peters, A. H. F. M.; Opravil, S.; Haynes, A. R.; Laible, G.; Rea, S.; Schmid, M.; Lebersorger, A.; Jerratsch, M.; Sattler, L.; Mattei, M. G.; Denny, P.; Brown, S. D. M.; Schweizer, D.; Jenuwein, T.: Isolation and characterization of Suv39h2, a second histone H3 methyltransferase gene that displays testis-specific expression. *Molec. Cell. Biol.* 20: 9423–9433, 2000.
- [71312] 9642.Davis, R. S.; Wang, Y.–H.; Kubagawa, H.; Cooper, M. D.: Identification of a family of Fc receptor homologs with preferential B cell expression. *Proc. Nat. Acad. Sci.* 98: 9772–9777, 2001.
- [71313] 9643.Xu, M.; Zhao, R.; Zhao, Z. J.: Molecular cloning and characterization of SPAP1, an inhibitory receptor. *Biochem. Biophys. Res. Commun.* 280:768–775, 2001.
- [71314] 9644.Drewes, G.; Ebner, A.; Preuss, U.; Mandelkow, E. M.; Mandelkow, E.: MARK, a novel family of protein kinases that phosphorylate microtubule-associated proteins and trigger microtubule disruption. *Cell* 89: 297–308, 1997.
- [71315] 9645.Nagase, T.; Kikuno, R.; Ishikawa, K.; Hirose, M.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XVII. The complete sequences of 100

new cDNA clones from brain which code for large proteins in vitro. DNA Res. 7: 143–150, 2000.

- [71316] 9646. Yoshikawa, T.; Sanders, A. R.; Esterling, L. E.; Detera-Wadleigh, S. D.: Multiple transcriptional variants and RNA editing in C18orf1, a novel gene with LDLRA and transmembrane domains on 18p11.2. Genomics 47:246–257, 1998.
- [71317] 9647. Yoshikawa, T.; Sanders, A. R.; Esterling, L. E.; Overharser, J.; Garnes, J. A.; Lennon, G.; Grewal, R.; Detera-Wadleigh, S. D.: Isolation of chromosome 18-specific brain transcripts as positional candidates for bipolar disorder. Am. J. Med. Genet. (Neuropsych. Genet.) 74:140–149, 1997.
- [71318] 9648. Anneren, C.; Reedquist, K. A.; Bos, J. L.; Welsh, M.: GTK, a Src-related tyrosine kinase, induces nerve growth factor-independent neurite outgrowth in PC12 cells through activation of the Rap1 pathway: relationship to Shb tyrosine phosphorylation and elevated levels of focal adhesion kinase. J. Biol. Chem. 275: 29153–29161, 2000.
- [71319] 9649. Cance, W. G.; Craven, R. J.; Bergman, M.; Xu, L.; Altitalo, K.; Liu, E. T.: Rak, a novel nuclear tyrosine kinase expressed in epithelial cells. Cell Growth Differ. 5: 1347–1355, 1994.

- [71320] 9650.Cance, W. G.; Craven, R. J.; Weiner, T. M.; Liu, E. T.: Novel protein kinases expressed in human breast cancer. *Int. J. Cancer* 54:571–577, 1993.
- [71321] 9651.Lee, J.; Wang, Z.; Luoh, S.–M.; Wood, W. I.; Scadden, D. T.: Cloning of FRK, a novel human intracellular SRC–like tyrosine kinase–encoding gene. *Gene* 138: 247–251, 1994.
- [71322] 9652.Scott, A. F.: Personal Communication. Baltimore, Md. 12/19/2001.
- [71323] 9653.Cao, L.; Zhang, L.; Ruiz–Lazano, P.; Yang, Q.; Chien, K. R.; Graham, R. M.; Zhou, M.: A novel putative protein–tyrosine phosphatase contains a BRO1–like domain and suppresses Ha–ras–mediated transformation. *J. Biol. Chem.* 273: 21077–21083, 1998.
- [71324] 9654.Toyooka, S.; Ouchida, M.; Jitsumori, Y.; Tsukuda, K.; Sakai, A.; Nakamura, A.; Shimizu, N.; Shimizu, K.: HD–PTP: a novel protein tyrosine phosphatase gene on human chromosome 3p21.3. *Biochem. Biophys. Res. Commun.* 278: 671–678, 2000.
- [71325] 9655.Betz, A.; Thakur, P.; Junge, H. J.; Ashery, U.; Rhee, J.–S.; Scheuss, V.; Rosenmund, C.; Rettig, J.; Brose, N.: Functional interaction of the active zone proteins Munc13–1 and RIM1 in synaptic vesicle priming. *Neuron* 30: 183–196, 2001.

- [71326] 9656.Castillo, P. E.; Schoch, S.; Schmitz, F.; Sudhof, T. C.; Malenka, R. C.: RIM1- $\alpha$  is required for presynaptic long-term potentiation. *Nature* 415:327–330, 2002.
- [71327] 9657.Coppola, T.; Magnin-Luthi, S.; Perret-Menoud, V.; Gattesco, S.; Schiavo, G.; Regazzi, R.: Direct interaction of the Rab3 effector RIM with  $\text{Ca}^{2+}$  channels, SNAP-25, and synaptotagmin. *J. Biol. Chem.* 276:32756–32762, 2001.
- [71328] 9658.Schoch, S.; Castillo, P. E.; Jo, T.; Mukherjee, K.; Geppert, M.; Wang, Y.; Schmitz, F.; Malenka, R. C.; Sudhof, T. C.: RIM1- $\alpha$  forms a protein scaffold for regulating neurotransmitter release at the active zone. *Nature* 415: 321–326, 2002.
- [71329] 9659.Wang, Y.; Sugita, S.; Sudhof, T. C.: The RIM/NIM family of neuronal C-2 domain proteins: interactions with Rab3 and a new class of Src homology 3 domain proteins. *J. Biol. Chem.* 275: 20033–20044, 2000.
- [71330] 9660.Cha, S. H.; Sekine, T.; Kusuhashira, H.; Yu, E.; Kim, J. Y.; Kim, D. K.; Sugiyama, Y.; Kanai, Y.; Endou, H.: Molecular cloning and characterization of multispecific organic anion transporter 4 expressed in the placenta. *J. Biol. Chem.* 275: 4507–4512, 2000.
- [71331] 9661.Vreugde, S.; Erven, A.; Kros, C. J.; Marcotti, W.; Fuchs, H.; Kurima, K.; Wilcox, E. R.; Friedman, T. B.; Grif-

fith, A. J.; Balling, R.; de Angelis, M. H.; Avraham, K. B.; Steel, K. P.: Beethoven, a mouse model for dominant, progressive hearing loss DFNA36. *Nature Genet.* 30: 257–258, 2002.

[71332] 9662. Grunder, S.; Geissler, H.-S.; Bassler, E.-L.; Ruppersberg, J. P.: A new member of acid-sensing ion channels from pituitary gland. *Neuroreport* 11:1607–1611, 2000.

[71333] 9663. Tatarelli, C.; Linnenbach, A.; Mimori, K.; Croce, C. M.: Characterization of the human TESTIN gene localized in the FRA7G region at 7q31.2. *Genomics* 68:1–12, 2000.

[71334] 9664. Tobias, E. S.; Hurlstone, A. F. L.; MacKenzie, E.; McFarlane, R.; Black, D. M.: The TES gene at 7q31.1 is methylated in tumours and encodes a novel growth-suppressing LIM domain protein. *Oncogene* 20:2844–2853, 2001.

[71335] 9665. Mizuno, Y.; Thompson, T. G.; Guyon, J. R.; Lidov, H. G. W.; Brosius, M.; Imamura, M.; Ozawa, E.; Watkins, S. C.; Kunkel, L. M.: Desmuslin, an intermediate filament protein that interacts with alpha-dystrobrevin and desmin. *Proc. Nat. Acad. Sci.* 98: 6156–6161, 2001.

[71336] 9666. Ferdinandusse, S.; Mulders, J.; IJlst, L.; Denis, S.; Dacremont, G.; Waterham, H. R.; Wanders, R. J. A.: Molecular cloning and expression of human carnitine octanoyl-transferase: evidence for its role in the peroxisomal beta-



oxidation of branched-chain fatty acids. *Biochem. Biophys. Res. Commun.* 263: 213–218, 1999.

- [71337] 9667. Nestel, F. P.; Colwill, K.; Harper, S.; Pawson, T.; Anderson, S.K.: RS cyclophilins: identification of an NK-TR(1)-related cyclophilin. *Gene* 180:151–155, 1996.
- [71338] 9668. Gaynor, E. C.; Mondesert, G.; Grimme, S. J.; Reed, S. I.; Orlean, P.; Emr, S. D.: MCD4 encodes a conserved endoplasmic reticulum membrane protein essential for glycosylphosphatidylinositol anchor synthesis in yeast. *Molec. Biol. Cell* 10: 627–648, 1999.
- [71339] 9669. Hong, Y.; Maeda, Y.; Watanabe, R.; Ohishi, K.; Mishkind, M.; Riezman, H.; Kinoshita, T.: Pig-n, a mammalian homologue of yeast Mcd4p, is involved in transferring phosphoethanolamine to the first mannose of the glycosylphosphatidylinositol. *J. Biol. Chem.* 274: 35099–35106, 1999.
- [71340] 9670. Gao, J.; Yu, L.; Zhang, P.; Jiang, J.; Chen, J.; Peng, J.; Wei, Y.; Zhao, S.: Cloning and characterization of human and mouse mitochondrial elongation factor G, GFM and Gfm, and mapping of GFM to human chromosome 3q25.1–q26.2. *Genomics* 74: 109–114, 2001.
- [71341] 9671. Clark, K. L.; Zeng, Z.; Langford, A. L.; Bowen, S. M.; Todd, S. C.: PGRL is a major CD81-associated protein on

lymphocytes and distinguishes a new family of cell surface proteins. *J. Immun.* 167: 5115–5121, 2001.

[71342] 9672. Stipp, C. S.; Kolesnikova, T. V.; Hemler, M. E.: EWI-2 is a major CD9 and CD81 partner and member of a novel Ig protein subfamily. *J. Biol. Chem.* 276: 40545–40554, 2001.

[71343] 9673. Miura, K.; Jacques, K. M.; Stauffer, S.; Kubosaki, A.; Zhu, K.; Hirsch, D. S.; Resau, J.; Zheng, Y.; Randazzo, P. A.: ARAP1: a point of convergence for Arf and Rho signaling. *Molec. Cell* 9: 109–119, 2002.

[71344] 9674. Dumoutier, L.; Lejeune, D.; Colau, D.; Renauld, J. C.: Cloning and characterization of IL-22 binding protein, a natural antagonist of IL-10-related T cell-derived inducible factor/IL 22. *J. Immun.* 166: 7090–7095, 2001.

[71345] 9675. Kotenko, S. V.; Izotova, L. S.; Mirochnitchenko, O. V.; Esterova, E.; Dickensheets, H.; Donnelly, R. P.; Pestka, S.: Identification, cloning, and characterization of a novel soluble receptor that binds IL-22 and neutralizes its activity. *J. Immun.* 166: 7096–7103, 2001.

[71346] 9676. Xu, W.; Presnell, S. R.; Parrish-Novak, J.; Kindsvogel, W.; Jaspers, S.; Chen, Z.; Dillon, S. R.; Gao, Z.; Gilbert, T.; Madden, K.; Schlutsmeyer, S.; Yao, L.; and 11 others: A soluble class II cytokine receptor, IL-22RA2, is a naturally

occurring IL-22 antagonist. Proc. Nat. Acad.Sci. 98: 9511–9516, 2001.

- [71347] 9677.Hicar, M. D.; Liu, Y.; Allen, C. E.; Wu, L.-C.: Structure of the human zinc finger protein HIVEP3: molecular cloning, expression, exon–intron structure, and comparison with paralogous genes HIVEP1 and HIVEP2. Genomics 71:89–100, 2001.
- [71348] 9678.Bates, E. E. M.; Fridman, W. H.; Mueller, C. G. F.: The ADAMDEC1(decysin) gene structure: evolution by duplication in a metalloprotease gene cluster on chromosome 8p12. Immunogenetics 54: 96–105, 2002.

- [71349] 9679. Bridges, L. C.; Tani, P. H.; Hanson, K. R.; Roberts, C. M.; Judkins, M. B.; Bowditch, R. D.: The lymphocyte metalloprotease MDC-L (ADAM28) is a ligand for the integrin  $\alpha$ -4/ $\beta$ -1. *J. Biol. Chem.* 277:3784–3792, 2002.
- [71350] 9680. Jury, J. A.; Perry, A. C. F.; Hall, L.: Identification, sequence analysis and expression of transcripts encoding a putative metalloproteinase, eMDC II, in human and macaque epididymis. *Molec. Hum. Reprod.* 5:1127–1134, 1999.
- [71351] 9681. Roberts, C. M.; Tani, P. H.; Bridges, L. C.; Laszik, Z.; Bowditch, R. D.: MDC-L, a novel metalloprotease disintegrin cysteine-rich protein family member expressed by human lymphocytes. *J. Biol. Chem.* 274:29251–29259, 1999.
- [71352] 9682. Koontz, J. I.; Soreng, A. L.; Nucci, M.; Kuo, F. C.; Pauwels, P.; van den Berghe, H.; Cin, P. D.; Fletcher, J. A.; Sklar, J.: Frequent fusion of the JAZF1 and JJAZ1 genes in endometrial stromal tumors. *Proc. Nat. Acad. Sci.* 98: 6348–6353, 2001.
- [71353] 9683. Fitzgerald, K. A.; Palsson-McDermott, E. M.; Bowie, A. G.; Jefferies, C. A.; Mansell, A. S.; Brady, G.; Brint, E.; Dunne, A.; Gray, P.; Harte, M. T.; McMurray, D.; Smith, D. E.; Sims, J. E.; Bird, T. A.; O'Neill, L. A. J.: Mal

(MyD88–adapter–like) is required for Toll–likereceptor–4 signal transduction. *Nature* 413: 78–83, 2001.

[71354] 9684.Horng, T.; Barton, G. M.; Medzhitov, R.: TIRAP: an adapter molecule in the Toll signaling pathway. *Nature Immun.* 2: 835–841, 2001.

[71355] 9685.Mirzayans, F.; Pearce, W. G.; MacDonald, I. M.; Walter, M. A.: Mutation of the PAX6 gene in patients with autosomal dominant keratitis. *Am.J. Hum. Genet.* 57: 539–548, 1995.

[71356] 9686.Dose, A. C.; Burnside, B.: Cloning and chromosomal localization of a human class III myosin. *Genomics* 67: 333–342, 2000.

[71357] 9687.Walsh, T.; Walsh, V.; Vreugde, S.; Hertzano, R.; Shahin, H.; Haika, S.; Lee, M. K.; Kanaan, M.; King, M.–C.; Avraham, K. B.: From flies' eyes to our ears: mutations in a human class III myosin cause progressive nonsyndromic hearing loss DFNB30. *Proc. Nat. Acad. Sci.* 99: 7518–7523, 2002.

[71358] 9688.Baron, M.: Genetics of schizophrenia and the new millennium: progress and pitfalls. *Am. J. Hum. Genet.* 68: 299–312, 2001.

[71359] 9689.Chakravarti, A.: A compelling genetic hypothesis for a complex disease: PRODH2/DGCR6 variation leads to

schizophrenia susceptibility. *Proc.Nat. Acad. Sci.* 99: 4755–4756, 2002.

- [71360] 9690.Gogos, J. A.; Santha, M.; Takacs, Z.; Beck, K. D.; Luine, V.; Lucas, L. R.; Nadler, J. V.; Karayiorgou, M.: The gene encoding prolinedehydrogenase modulates sensorimotor gating in mice. *Nature Genet.* 21:434–439, 1999.
- [71361] 9691.Liu, H.; Heath, S. C.; Sobin, C.; Roos, J. L.; Galke, B. L.; Blundell, M. L.; Lenane, M.; Robertson, B.; Wijsman, E. M.; Rapoport, J. L.; Gogos, J. A.; Karayiorgou, M.: Genetic variation at the 22q11 *PRODH2/DGCR6* locus presents an unusual pattern and increases susceptibility to schizophrenia. *Proc. Nat. Acad. Sci.* 99: 3717–3722, 2002.
- [71362] 9692.Scott, A.: Personal Communication. Baltimore, Md. 4/2002.
- [71363] 9693.Plager, D. A.; Loegering, D. A.; Weiler, D. A.; Checkel, J. L.; Wagner, J. M.; Clarke, N. J.; Naylor, S.; Page, S. M.; Thomas, L.L.; Akerblom, I.; Cocks, B.; Stuart, S.; Gleich, G. J.: A novel and highly divergent homolog of human eosinophil granule major basic protein. *J.Biol. Chem.* 274: 14464–14473, 1999.
- [71364] 9694.Leo, C. P.; Hsu, S. Y.; McGee, E. A.; Salanova, M.; Hsueh, A. J.W.: DEFT, a novel death effector domain–

containing molecule predominantly expressed in testicular germ cells. *Endocrinology* 139: 4839–4848, 1998.

- [71365] 9695. Stegh, A. H.; Schickling, O.; Ehret, A.; Scaffidi, C.; Peterhansel, C.; Hofmann, T. G.; Grummt, I.; Krammer, P. H.; Peter, M. E.: DEDD, a novel death effector domain-containing protein, targeted to the nucleolus. *EMBO J.* 17: 5974–5986, 1998.
- [71366] 9696. Charest, A.; Lane, K.; McMahon, K.; Housman, D. E.: Association of a novel PDZ domain-containing peripheral Golgi protein with the Q-SNARE (Q-soluble N-ethylmaleimide-sensitive fusion protein (NSF) attachment protein receptor) protein syntaxin 6. *J. Biol. Chem.* 276:29456–29465, 2001.
- [71367] 9697. Cheng, J.; Moyer, B. D.; Milewski, M.; Loffing, J.; Ikeda, M.; Mickle, J. E.; Cutting, G. R.; Li, M.; Stanton, B. A.; Guggino, W. B.: A Golgi-associated PDZ domain protein modulates cystic fibrosis transmembrane regulator plasma membrane expression. *J. Biol. Chem.* 277:3520–3529, 2002.
- [71368] 9698. Neudauer, C. L.; Joberty, G.; Macara, I. G.: PIST: a novel PDZ/coiled-coil domain binding partner for the Rho-family GTPase TC10. *Biochem. Biophys. Res. Commun.* 280: 541–547, 2001.

- [71369] 9699.Yao, R.; Maeda, T.; Takada, S.; Noda, T.: Identification of aPDZ domain containing Golgi protein, GOPC, as an interaction partner of frizzled. *Biochem. Biophys. Res. Commun.* 286: 771–778, 2001.
- [71370] 9700.Kelter, A.-R.; Herchenbach, J.; Wirth, B.: The transcription factor–like nuclear regulator (TFNR) contains a novel 55–amino–acid motif repeated nine times and maps closely to SMN1. *Genomics* 70: 315–326, 2000.
- [71371] 9701.Schramm, L.; Pendergrast, P. S.; Sun, Y.; Hernandez, N.: Different human TFIIIB activities direct RNA polymerase III transcription from TATA–containing and TATA–less promoters. *Genes Dev.* 14: 2650–2663, 2000.
- [71372] 9702.Cabart, P.; Murphy, S.: BRFU, a TFIIIB–like factor, is directly recruited to the TATA–box of polymerase III small nuclear RNA gene promoters through its interaction with TATA–binding protein. *J. Biol.Chem.* 276: 43056–43064, 2001.
- [71373] 9703.Schramm, L.; Pendergrast, P. S.; Sun, Y.; Hernandez, N.: Different human TFIIIB activities direct RNA polymerase III transcription from TATA–containing and TATA–less promoters. *Genes Dev.* 14: 2650–2663, 2000.
- [71374] 9704.Kobayashi, K.; Kuroda, S.; Fukata, M.; Nakamura, T.; Nagase, T.; Nomura, N.; Matsuura, Y.; Yoshida–Kubomura,



N.; Iwamatsu, A.; Kaibuchi, K.: p140Sra-1 (specifically Rac1-associated protein) is a novel specific target for Rac1 small GTPase. *J. Biol. Chem.* 273: 291–295, 1998.

[71375] 9705. Schenck, A.; Bardoni, B.; Moro, A.; Bagni, C.; Mandel, J. L.: A highly conserved protein family interacting with the fragile X mental retardation protein (FMRP) and displaying selective interactions with FMRP-related proteins FXR1P and FXR2P. *Proc. Nat. Acad. Sci.* 98: 8844–8849, 2001.

[71376] 9706. Lopez-Rios, J.; Gallardo, M. E.; Rodriguez de Cordoba, S.; Bovolenta, P.: Six9 (Optx2), a new member of the Six gene family of transcription factors, is expressed at early stages of vertebrate ocular and pituitary development. *Mech. Dev.* 83: 155–159, 1999.

[71377] 9707. Oukka, M.; Kim, S. T.; Lugo, G.; Sun, J.; Wu, L.-C.; Glimcher, L. H.: A mammalian homolog of *Drosophila* schnurri, KRC, regulates TNF receptor-driven responses and interacts with TRAF2. *Molec. Cell* 9: 121–131, 2002.

[71378] 9708. Walder, R. Y.; Landau, D.; Meyer, P.; Shalev, H.; Tsolia, M.; Borochowitz, Z.; Boettger, M. B.; Beck, G. E.; Englehardt, R. K.; Carmi, R.; Sheffield, V. C.: Mutation of TRPM6 causes familial hypomagnesemia with secondary hypocalcemia. *Nature Genet.* 31: 171–174, 2002.

[71379] 9709. Liang, J. C.; Chang, K. S.; Schroeder, W.; Siciliano,

M.; Trujillo, J.; Stass, S.: The human myeloperoxidase gene locates on chromosome 17q22–24 and is translocated in acute promyelocytic leukemia. (Abstract) *Am. J. Hum. Genet.* 41: A226, 1987.

[71380] 9710. Liang, J. C.; Chang, K. S.; Schroeder, W. T.; Freireich, E. J.; Stass, S. A.; Trujillo, J. M.: The myeloperoxidase gene is translocated from chromosome 17 to 15 in a patient with acute promyelocytic leukemia. *Cancer Genet. Cytogenet.* 30: 103–107, 1988.

[71381] 9711. Miki, T.; Weil, S. C.; Rosner, G. L.; Reid, M. S.; Kidd, K. K.: An MPO cDNA clone identifies an RFLP with *Pst*I. *Nucleic Acids Res.* 16:1649, 1988.

[71382] 9712. Morishita, K.; Kubota, N.; Asano, S.; Kaziro, Y.; Nagata, S.: Molecular cloning and characterization of cDNA for human myeloperoxidase. *J. Biol. Chem.* 262: 3844–3851, 1987.

[71383] 9713. Murao, S.-I.; Stevens, F. J.; Ito, A.; Huberman, E.: Myeloperoxidase: a myeloid cell nuclear antigen with DNA-binding properties. *Proc. Nat. Acad. Sci.* 85: 1232–1236, 1988.

[71384] 9714. Nauseef, W.; Cogley, M.; McCormick, S.: Effect of the R569W missense mutation on the biosynthesis of myeloperoxidase. *J. Biol. Chem.* 271:9546–9549, 1996.

- [71385] 9715.Nauseef, W. M.; Olsson, I.; Arnljots, K.: Biosynthesis and processing of myeloperoxidase--a marker for myeloid cell differentiation. *Europ.J. Haemat.* 40: 97-110, 1988.
- [71386] 9716.Reynolds, W. F.; Hiltunen, M.; Pirskanen, M.; Mannermaa, A.; Helisalmi, S.; Lehtovirta, M.; Alafuzoff, I.; Soininen, H.: MPO and APOE epsilon-4 polymorphisms interact to increase risk for AD in Finnish males. *Neurology* 55:1284-1290, 2000.
- [71387] 9717.Robinson, T. J.; Morris, D. J.; Ledbetter, D. H.: Chromosomal assignment and regional localization of myeloperoxidase in the mouse. *Cytogenet.Cell Genet.* 53: 83-86, 1990.
- [71388] 9718.Romano, M.; Dri, P.; Dadalt, L.; Patriarca, P.; Baralle, F. E.: Biochemical and molecular characterization of hereditary myeloproliferative deficiency. *Blood* 90: 4126-4134, 1997.
- [71389] 9719.van Tuinen, P.; Johnson, K. R.; Ledbetter, S. A.; Nussbaum, R.L.; Rovera, G.; Ledbetter, D. H.: Localization of myeloperoxidase to the long arm of human chromosome 17: relationship to the 15;17 translocation of acute promyelocytic leukemia. *Oncogene* 1: 319-322, 1987.
- [71390] 9720.Weil, S. C.; Rosner, G. L.; Reid, M. S.; Chisholm, R. L.; Farber, N. M.; Spitznagel, J. K.; Swanson, M. S.: cDNA

cloning of human myeloperoxidase:decrease in myeloperoxidase mRNA upon induction of HL-60 cells. *Proc.Nat. Acad. Sci.* 84: 2057-2061, 1987.

[71391] 9721.Weil, S. C.; Rosner, G. L.; Reid, M. S.; Chisholm, R. L.; Lemons,R. S.; Swanson, M. S.; Carrino, J. J.; Diaz, M. O.; Le Beau, M. M.: Translocation and rearrangement of myeloperoxidase gene in acute promyelocytic leukemia. *Science* 240: 790-792, 1988.

[71392] 9722.Yamada, M.; Hur, S.-J.; Hashinaka, K.; Tsuneoka, K.; Saeki, T.;Nishio, C.; Sakiyama, F.; Tsunasawa, S.: Isolation and characterization of a cDNA coding for human myeloperoxidase. *Arch. Biochem. Biophys.* 255:147-155, 1987.

[71393] 9723.Zaki, S. R.; Austin, G. E.; Chan, W. C.; Conaty, A. L.; Trusler,S.; Trappier, S.; Lindsey, R. B.; Swan, D. C.: Chromosomal localization of the human myeloperoxidase gene by in situ hybridization using oligonucleotide probes. *Genes Chromosomes Cancer* 2: 266-270, 1990.

[71394] 9724.Saiardi, A.; Erdjument-Bromage, H.; Snowman, A. M.; Tempst, P.;Snyder, S. H.: Synthesis of diphosphoinositol pentakisphosphate by a newly identified family of higher inositol polyphosphate kinases. *Curr.Biol.* 9: 1323-1326, 1999.

- [71395] 9725.Saiardi, A.; Nagata, E.; Luo, H. R.; Snowman, A. M.; Snyder, S.H.: Identification and characterization of a novel inositol hexakisphosphatekinase. *J. Biol. Chem.* 276: 39179–39185, 2001.
- [71396] 9726.Morrison, B. H.; Bauer, J. A.; Hu, J.; Grane, R. W.; Ozdemir, A.M.; Chawla–Sarkar, M.; Gong, B.; Almasan, A.; Kalvakolanu, D. V.;Lindner, D. J.: Inositol hexakisphosphate kinase 2 sensitizes ovariancarcinoma cells to multiple cancer therapeutics. *Oncogene* 21: 1882–1889,2002.
- [71397] 9727.Gasdaska, P. Y.; Berggren, M. M.; Berry, M. J.; Powis, G.: Cloning,sequencing, and functional expression of a novel human thioredoxinreductase. *FEBS Lett.* 442: 105–111, 1999.
- [71398] 9728.Kawai, H.; Ota, T.; Suzuki, F.; Tatsuka, M.: Molecular cloningof mouse thioredoxin reductases. *Gene* 242: 321–330, 2000.
- [71399] 9729.Miranda–Vizuite, A.; Damdimopoulos, A. E.; Spyrou, G.: cDNA cloning,expression and chromosomal localization of the mouse mitochondrialthioredoxin reductase gene. *Biochim. Biophys. Acta* 1447: 113–118,1999.
- [71400] 9730.Miranda–Vizuite, A.; Damdimopoulos, A. E.; Pedrajas, J. R.; Gustafsson,J.–A.; Spyrou, G.: Human mitochondrial thioredoxin reductase: cDNAcloning, expression, and

genomic organization. *Europ. J. Biochem.* 261:405–412, 1999.

[71401] 9731. Nishida, K.; Yoshida, Y.; Itoh, M.; Fukada, T.; Ohtani, T.; Shirogane, T.; Atsumi, T.; Takahashi–Tezuka, M.; Ishihara, K.; Hibi, M.; Hirano, T.: Gab–family adapter proteins act downstream of cytokine and growth factor receptors and T– and B–cell antigen receptors. *Blood* 93:1809–1816, 1999.

[71402] 9732. Zhao, C.; Yu, D.–H.; Shen, R.; Feng, G.–S.: Gab2, a new pleckstrin homology domain–containing adapter protein, acts to uncouple signaling from ERK kinase to Elk–1. *J. Biol. Chem.* 274: 19649–19654, 1999.

[71403] 9733. Brakenhoff, R. H.; Gerretsen, M.; Knippels, E. M. C.; van Dijk, M.; van Essen, H.; Weghuis, D. O.; Sinke, R. J.; Snow, G. B.; van Dongen, G. A. M. S.: The human E48 antigen, highly homologous to the murine Ly–6 antigen ThB, is a GPI–anchored molecule apparently involved in keratinocyte cell–cell adhesion. *J. Cell Biol.* 129:1677–1689, 1995.

[71404] 9734. Brakenhoff, R. H.; van Dijk, M.; Rood–Knippels, E. M. C.; Snow, G. B.: A gain of novel tissue specificity in the human Ly–6 gene E48. *J. Immun.* 159: 4879–4886, 1997.

[71405] 9735. Loewen, S. K.; Ng, A. M. L.; Yao, S. Y. M.; Cass, C. E.;

Baldwin, S. A.; Young, J. D.: Identification of amino acid residues responsible for the pyrimidine and purine nucleoside specificities of human concentrative Na<sup>+</sup> nucleoside cotransporters hCNT1 and hCNT2. *J. Biol. Chem.* 274:24475–24484, 1999.

[71406] 9736. Ritzel, M. W. L.; Yao, S. Y. M.; Huang, M.-Y.; Elliott, J. F.; Cass, C. E.; Young, J. D.: Molecular cloning and functional expression of cDNAs encoding a human Na<sup>(+)</sup>-nucleoside cotransporter (hCNT1). *Am. J. Physiol.* 272: C707–C714, 1997.

[71407] 9737. Ritzel, M. W. L.; Yao, S. Y. M.; Ng, A. M. L.; Mackey, J. R.; Cass, C. E.; Young, J. D.: Molecular cloning, functional expression and chromosomal localization of a cDNA encoding a human Na<sup>+</sup>/nucleoside cotransporter (hCNT2) selective for purine nucleosides and uridine. *Molec. Membrane Biol.* 15: 203–211, 1998.

[71408] 9738. Lescure, A.; Gautheret, D.; Carbon, P.; Krol, A.: Novel selenoproteins identified in silico and in vivo by using a conserved RNA structural motif. *J. Biol. Chem.* 274: 38147–38154, 1999.

[71409] 9739. Berghs, S.; Aggujaro, D.; Dirkx, R., Jr.; Maksimova, E.; Stabach, P.; Hermel, J.-M.; Zhang, J.-P.; Philbrick, W.; Slepnev, V.; Ort, T.; Slimena, M.: Beta-IV spectrin, a new

spectrin localized at axoninitial segments and nodes of Ranvier in the central and peripheralnervous system. J. Cell Biol. 151: 985–1001, 2000.

[71410] 9740.Bock, G. R.; Frank, M. P.; Steel, K. P.; Deol, M. S.: The quiveringmutant mouse: hereditary deafness of central origin. Acta Otolaryng. 96:371–377, 1983.

[71411] 9741.Deol, M. S.; Frank, M. P.; Steel, K. P.; Bock, G. R.: Geneticdeafness of central origin. Brain Res. 258: 177–179, 1983.

[71412] 9742.Parkinson, N. J.; Olsson, C. L.; Hallows, J. L.; McKee-Johnson,J.; Keogh, B. P.; Noben-Trauth, K.; Kujawa, S. G.; Tempel, B. L.:Mutant beta-spectrin 4 causes auditory and motor neuropathies in quiveringmice. Nature Genet. 29: 61–65, 2001.

[71413] 9743.Tse, W. T.; Tang, J.; Jin, O.; Korsgren, C.; John, K. M.; Kung,A. L.; Gwynn, B.; Peters, L. L.; Lux, S. E.: A new spectrin, beta-IV,has a major truncated isoform that associates with promyelocytic leukemia protein nuclear bodies and the nuclear matrix. J. Biol. Chem. 276:23974–23985, 2001.

[71414] 9744.Yoon, C. H.; Les, E. P.: Quivering, a new first chromosome mutationin mice. J. Hered. 48: 176–180, 1957.

[71415] 9745.Petek, E.; Windpassinger, C.; Egger, H.; Kroisel, P.



M.; Wagner,K.: Localization of the human anterior gradient-2 gene (AGR2) to chromosome band 7p21.3 by radiation hybrid mapping and fluorescence in situ hybridisation. *Cytogenet. Cell Genet.* 89: 141–142, 2000.

[71416] 9746. Thompson, D. A.; Weigel, R. J.: hAG-2, the human homologue of the *Xenopus laevis* cement gland gene XAG-2, is coexpressed with estrogen receptor in breast cancer cell lines. *Biochem. Biophys. Res. Commun.* 251:111–116, 1998.

[71417] 9747. Saitoh, T.; Katoh, M.: Molecular cloning and characterization of human WNT5B on chromosome 12p13.3 region. *Int. J. Oncol.* 19:347–351, 2001.

[71418] 9748. Miyazaki, K.; Matsuda, S.; Ichigotani, Y.; Takenouchi, Y.; Hayashi, K.; Fukuda, Y.; Nimura, Y.; Hamaguchi, M.: Isolation and characterization of a novel human gene (NESH) which encodes a putative signaling molecule similar to e3B1 protein. *Biochim. Biophys. Acta* 1493: 237–241, 2000.

[71419] 9749. Nosaka, K.; Onozuka, M.; Kakazu, N.; Hibi, S.; Nishimura, H.; Nishino, H.; Abe, T.: Isolation and characterization of a human thiamine pyrophosphokinase cDNA. *Biochem. Biophys. Acta* 1517: 293–297, 2001.

[71420] 9750. Nosaka, K.; Onozuka, M.; Nishino, H.; Nishimura, H.;

Kawasaki,Y.; Ueyama, H.: Molecular cloning and expression of a mouse thiaminpyrophosphokinase cDNA. J. Biol. Chem. 274: 34129–34133, 1999.

[71421] 9751.Zhao, R.; Gao, F.; Goldman, I. D.: Molecular cloning of humanthiamin pyrophosphokinase. Biochim. Biophys. Acta 1517: 320–322,2001.

[71422] 9752.Reboul, J.; Gardiner, K.; Monneron, D.; Uze, G.; Lutfalla, G.:Comparative genomic analysis of the interferon/interleukin–10 receptorgene cluster. Genome Res. 9: 242–250, 1999.

[71423] 9753.Chen, Q.; Ghilardi, N.; Wang, H.; Baker, T.; Xie, M.–H.; Gurney,A.; Grewal, I. S.; de Sauvage, F. J.: Development of Th1–type immuneresponses requires the type 1 cytokine receptor TCCR. Nature 407:916–920, 2000.

[71424] 9754.Sprecher, C. A.; Grant, F. J.; Baumgartner, J. W.; Presnell, S.R.; Schrader, S. K.; Yamagiwa, T.; Whitmore, T. E.; O'Hara, P. J.;Foster, D. F.: Cloning and characterization of a novel class I cytokinereceptor. Biochem. Biophys. Res. Comm. 246: 82–90, 1998.

[71425] 9755.Yoshida, H.; Hamano, S.; Senaldi, G.; Covey, T.; Faggioni, R.;Mu, S.; Xia, M.; Wakeham, A. C.; Nishina, H.; Potter, J.; Saris, C.J. M.; Mak, T. W.: WSX–1 is required for the initiation of Th1 responsesand resistance to L. major in–

fection. *Immunity* 15: 569–578, 2001.

[71426] 9756. Autieri, M. V.; Carbone, C. J.: 14-3-3-Gamma interacts with and is phosphorylated by multiple protein kinase C isoforms in PDGF-stimulated human vascular smooth muscle cells. *DNA Cell Biol.* 18: 555–564, 1999.

[71427] 9757. Yeh, W.-C.; de la Pompa, J. L.; McCurrach, M. E.; Shu, H.-B.; Elia, A. J.; Shahinian, A.; Ng, M.; Wakeham, A.; Khoo, W.; Mitchell, K.; El-Deiry, W. S.; Lowe, S. W.; Goeddel, D. V.; Mak, T. W.: FADD: essential for embryo development and signaling from some, but not all, inducers of apoptosis. *Science* 279: 1954–1958, 1998.

[71428] 9758. Zhang, J.; Cado, D.; Chen, A.; Kabra, N. H.; Winoto, A.: Fas-mediated apoptosis and activation-induced T-cell proliferation are defective in mice lacking FADD/Mort1. *Nature* 392: 296–300, 1998.

[71429] 9759. Torres, R.; Polymeropoulos, M. H.: Genomic organization and localization of the human CRMP-1 gene. *DNA Res.* 5: 393–395, 1998.

[71430] 9760. Koyama, K.; Sudo, K.; Nakamura, Y.: Isolation of 115 human chromosome 8-specific expressed-sequence tags by exon amplification. *Genomics* 26: 245–253, 1995.

[71431] 9761. Katahira, J.; Sugiyama, H.; Inoue, N.; Horiguchi, Y.; Matsuda, M.; Sugimoto, N.: *Clostridium perfringens* en-

terotoxin utilizes two structurally related membrane proteins as functional receptors *in vivo*. *J. Biol. Chem.* 272: 26652–26658, 1997.

- [71432] 9762. Furuse, M.; Fujita, K.; Hiragi, T.; Fujimoto, K.; Tsukita, S.: Claudin-1 and -2: novel integral membrane proteins localizing at tight junctions with no sequence similarity to occludin. *J. Cell Biol.* 141: 1539–1550, 1998.
- [71433] 9763. Peacock, R. E.; Keen, T. J.; Inglehearn, C. F.: Analysis of a human gene homologous to rat ventral prostate.1 protein. *Genomics* 46:443–449, 1997.
- [71434] 9764. Black, J. L., III; Lennon, V. A.: Identification and cloning of putative human neuronal voltage-gated calcium channel  $\gamma$ -2 and  $\gamma$ -3 subunits: neurologic implications. *Mayo Clin. Proc.* 74:357–361, 1999.
- [71435] 9765. Chen, L.; Chetkovich, D. M.; Petralla, R. S.; Sweeney, N. T.; Kawasaki, Y.; Wenthold, R. J.; Brecht, D. S.; Nicoli, R. A.: Stargazin regulates synaptic targeting of AMPA receptors by two distinct mechanisms. *Nature* 408:936–943, 2000.
- [71436] 9766. Letts, V. A.; Felix, R.; Biddlecome, G. H.; Arikath, J.; Mahaffey, C. L.; Valenzuela, A.; Bartlett, F. S., II; Mori, Y.; Campbell, K. P.; Frankel, W. N.: The mouse stargazer gene encodes a neuronal  $\text{Ca}^{2+}$ -channel  $\gamma$  subunit. *Na-*

ture Genet. 19: 340–347, 1998.

- [71437] 9767. Biffo, S.; Sanvito, F.; Costa, S.; Preve, L.; Pignatelli, R.; Spinardi, L.; Marchisio, P. C.: Isolation of a novel beta-4 integrin-binding protein (p27BBP) highly expressed in epithelial cells. *J. Biol. Chem.* 272:30314–30321, 1997.
- [71438] 9768. Sanvito, F.; Arrigo, G.; Zuffardi, O.; Agnelli, M.; Marchisio, P. C.; Biffo, S.: Localization of p27 beta-4 binding protein gene (ITGB4BP) to human chromosome region 20q11.2. *Genomics* 52: 111–112, 1998.
- [71439] 9769. Si, K.; Chaudhuri, J.; Chevesich, J.; Maitra, U.: Molecular cloning and functional expression of a human cDNA encoding translation initiation factor 6. *Proc. Nat. Acad. Sci.* 94: 14285–14290, 1997.
- [71440] 9770. Liu, Z.; Shen, J.; Carbrey, J. M.; Mukhopadhyay, R.; Agre, P.; Rosen, B. P.: Arsenite transport by mammalian aquaglyceroporins AQP7 and AQP9. *Proc. Nat. Acad. Sci.* 99: 6053–6058, 2002.
- [71441] 9771. Nuber, U.; Schwarz, S.; Kaiser, P.; Schneider, R.; Scheffner, M.: Cloning of human ubiquitin-conjugating enzymes UbCH6 and UbCH7 (E2-F1) and characterization of their interaction with E6-AP and RSP5. *J. Biol. Chem.* 271: 2795–2800, 1996.
- [71442] 9772. Fuentes, J. J.; Pritchard, M. A.; Estivill, X.: Genomic

organization, alternative splicing, and expression patterns of the DSCR1 (Down syndrome candidate region 1) gene. Genomics 44: 358–361, 1997.

- [71443] 9773. Kingsbury, T. J.; Cunningham, K. W.: A conserved family of calcineurin regulators. Genes Dev. 14: 1595–1604, 2000.
- [71444] 9774. Rothermel, B.; Vega, R. B.; Yang, J.; Wu, H.; Bassel-Duby, R.; Williams, R. S.: A protein encoded within the Down syndrome critical region is enriched in striated muscles and inhibits calcineurin signaling. J. Biol. Chem. 275: 8719–8725, 2000.
- [71445] 9775. Pata, I.; Tensing, K.; Metspalu, A.: A human cDNA encoding the homologue of NADH:ubiquinone oxidoreductase subunit B13. Biochim. Biophys. Acta 1350: 115–118, 1997.
- [71446] 9776. Russell, M. W.; du Manoir, S.; Collins, F. S.; Brody, L. C.: Cloning of the human NADH:ubiquinone oxidoreductase subunit B13: localization to chromosome 7q32 and identification of a pseudogene on 11p15. Mammalian Genome 8: 60–61, 1997.
- [71447] 9777. Perez Jurado, L. A.; Wang, Y.-K.; Peoples, R.; Coloma, A.; Cruces, J.; Francke, U.: A duplicated gene in the breakpoint regions of the 7q11.23 Williams–Beuren

syndrome deletion encodes the initiator binding protein TFII-I and BAP-135, a phosphorylation target of BTK. Hum.Molec. Genet. 7: 325-334, 1998.

- [71448] 9778. Yang, W.; Desiderio, S.: BAP-135, a target for Bruton's tyrosine kinase in response to B cell receptor engagement. Proc. Nat. Acad.Sci. 94: 604-609, 1997.
- [71449] 9779. Lee, J. W.; Choi, H.-S.; Gyuris, J.; Brent, R.; Moore, D. D.: Two classes of proteins dependent on either the presence or absence of thyroid hormone for interaction with the thyroid hormone receptor. Molec.Endocr. 9: 243-254, 1995.
- [71450] 9780. Burnatowska-Hledin, M. A.; Spielman, W. S.; Smith, W. L.; Shi, P.; Meyer, J. M.; Dewitt, D. L.: Expression cloning of an AVP-activated, calcium-mobilizing receptor from rabbit kidney medulla. Am. J. Physiol. 268:F1198-F1210, 1995.
- [71451] 9781. Byrd, P. J.; Stankovic, T.; McConville, C. M.; Smith, A. D.; Cooper, P. R.; Taylor, A. M. R.: Identification and analysis of expression of human VACM-1, a cullin gene family member located on chromosome 11q22-23. Genome Res. 7: 71-75, 1997.
- [71452] 9782. Venturini, L.; You, J.; Stadler, M.; Galien, R.; Lallemand, V.; Koken, M. H. M.; Mattei, M. G.; Ganser, A.;

Chambon, P.; Losson, R.; de The, H.: TIF1-gamma, a novel member of the transcriptional intermediary factor 1 family. *Oncogene* 18: 1209–1217, 1999.

[71453] 9783. Lesage, F.; Guillemare, E.; Fink, M.; Duprat, F.; Lazdunski, M.; Romey, G.; Barhanin, J.: TWIK-1, a ubiquitous human weakly inward rectifying K<sup>+</sup> channel with a novel structure. *EMBO J.* 15: 1004–1011, 1996.

[71454] 9784. Lesage, F.; Mattei, M.-G.; Fink, M.; Barhanin, J.; Lazdunski, M.: Assignment of the human weak inward rectifier K<sup>+</sup> channel TWIK-1 gene to chromosome 1q42–q43. *Genomics* 34: 153–155, 1996.

[71455] 9785. Holzmann, K.; Poltl, A.; Sauermann, G.: A novel spliced transcript of human CLAPS2 encoding a protein alternative to clathrin adaptor protein AP17. *Gene* 220: 39–44, 1998.

[71456] 9786. Winterpacht, A.; Ende, S.; Enklaar, T.; Fuhry, M.; Zabel, B.: Human CLAPS2 encoding AP17, a small chain of the clathrin-associated protein complex: cDNA cloning and chromosomal assignment to 19q13.2–q13.3. *Cytogenet. Cell Genet.* 75: 132–135, 1996.

[71457] 9787. Rodriguez, A. M.; Rodin, D.; Nomura, H.; Morton, C. C.; Weremowicz, S.; Schneider, M. C.: Identification, localization, and expression of two novel human genes similar



to deoxyribonuclease I. *Genomics* 42:507–513, 1997.

[71458] 9788.Zeng, Z.; Parmelee, D.; Hyaw, H.; Coleman, T. A.; Su, K.; Zhang, J.; Gentz, R.; Ruben, S.; Rosen, C.; Li, Y.: Cloning and characterization of a novel human DNase. *Biochem. Biophys. Res. Commun.* 231: 499–504, 1997.

[71459] 9789.Alderson, M. R.; Smith, C. A.; Tough, T. W.; Davis-Smith, T.; Armitage, R. J.; Falk, B.; Roux, E.; Baker, E.; Sutherland, G. R.; Din, W. S.; Goodwin, R. G.: Molecular and biological characterization of human 4–1BB and its ligand. *Europ. J. Immun.* 24: 2219–2227, 1994.

[71460] 9790.Kwon, B. S.; Weissman, S. M.: cDNA sequences of two inducible T-cell genes. *Proc. Nat. Acad. Sci.* 86: 1963–1967, 1989.

[71461] 9791.Loo, D. T.; Chalupny, N. J.; Bajorath, J.; Shuford, W. W.; Mittler, R. S.; Aruffo, A.: Analysis of 4–1BBL and laminin binding to murine 4–1BB, a member of the tumor necrosis factor receptor superfamily, and comparison with human 4–1BB. *J. Biol. Chem.* 272: 6448–6456, 1997.

[71462] 9792.Schwarz, H.; Arden, K.; Lotz, M.: CD137, a member of the tumor necrosis factor receptor family, is located on chromosome 1p36, in a cluster of related genes, and colocalizes with several malignancies. *Biochem. Biophys. Res. Commun.* 235: 699–703, 1997.

- [71463] 9793.Schwarz, H.; Blanco, F. J.; von Kempis, J.; Valbracht, J.; Lotz,M.: ILA, a member of the human nerve growth factor/tumor necrosisfactor receptor family, regulates T-lymphocyte proliferation and survival. *Blood* 87:2839–2845, 1996.
- [71464] 9794.Schwarz, H.; Tuckwell, J.; Lotz, M.: A receptor induced by lymphocyteactivation (ILA): a new member of the human nerve-growth-factor/tumor-necrosis-factorreceptor family. *Gene* 134: 295–298, 1993.
- [71465] 9795.Deere, M.; Johnson, J.; Garza, S.; Harrison, W. R.; Yoon, S.-J.;Elder, F. F. B.; Kucherlapati, R.; Hook, M.; Hecht, J. T.: Characterizationof human DSPG3, a small dermatan sulfate proteoglycan. *Genomics* 38:399–404, 1996.
- [71466] 9796.Shinomura, T.; Kimata, K.; Oike, Y.; Noro, A.; Hirose, N.; Tanabe,K.; Suzuki, S.: The occurrence of three different proteoglycan speciesin chick embryo cartilage: Isolation and characterization of a secondproteoglycan (PG-Lb) and its precursor form. *J. Biol. Chem.* 258:9314–9322, 1983.
- [71467] 9797.Feuchter-Murthy, A. E.; Freeman, J. D.; Mager, D. L.: Splicingof a human endogenous retrovirus to a novel

phospholipase A2 related gene. *Nucleic Acids Res.* 21: 135–143, 1993.

[71468] 9798. Kowalski, P. E.; Freeman, J. D.; Mager, D. L.: Inter-genic splicing between a HERV-H endogenous retrovirus and two adjacent human genes. *Genomics* 57:371–379, 1999.

[71469] 9799. Baylin, S. B.; Herman, J. G.; Graff, J. R.; Vertino, P. M.; Issa, J. P.: Alterations in DNA methylation: a fundamental aspect of neoplasia. *Adv. Cancer Res.* 72: 141–196, 1998.

[71470] 9800. Lei, H.; Oh, S. P.; Okano, M.; Juttermann, R.; Goss, K. A.; Jaenisch, R.; Li, E.: De novo DNA cytosine methyltransferase activities in mouse embryonic stem cells. *Development* 122: 3195–3205, 1996.

[71471] 9801. Okano, M.; Xie, S.; Li, E.: Cloning and characterization of a family of novel mammalian DNA (cytosine-5) methyltransferases. (Letter) *Nature Genet.* 19: 219–220, 1998.

[71472] 9802. Robertson, K. D.; Uzvolgyi, E.; Liang, G.; Talmadge, C.; Sumegi, J.; Gonzales, F. A.; Jones, P. A.: The human DNA methyltransferases (DNMTs) 1, 3a and 3b: coordinate mRNA expression in normal tissues and overexpression in tumors. *Nucleic Acids Res.* 27: 2291–2298, 1999.

[71473] 9803. Xie, S.; Wang, Z.; Okano, M.; Nogami, M.; Li, Y.; He,

W.-W.; Okumura,K.; Li, E.: Cloning, expression and chromosome locations of the humanDNMT3 gene family. Gene 236: 87–95, 1999.

[71474] 9804.Jiang, R.; Lan, Y.; Chapman, H. D.; Shawber, C.; Norton, C. R.;Serreze, D. V.; Weinmaster, G.; Gridley, T.: Defects in limb, craniofacial,and thymic development in Jagged2 mutant mice. Genes Dev. 12: 1046–1057,1998.

[71475] 9805.Lan, Y.; Jiang, R.; Shawber, C.; Weinmaster, G.; Gridley, T.:The Jagged2 gene maps to chromosome 12 and is a candidate for theIgl and sm mutations. Mammalian Genome 8: 875–876, 1997.

[71476] 9806.Lanford, P. J.; Lan, Y.; Jiang, R.; Lindsell, C.; Weinmaster, G.;Gridley, T.; Kelley, M. W.: Notch signalling pathway mediates haircell development in mammalian cochlea. Nature Genet. 21: 289–292,1999.

[71477] 9807.Luo, B.; Aster, J. C.; Hasserjian, R. P.; Kuo, F.; Sklar, J.:Isolation and functional analysis of a cDNA for human Jagged2, a geneencoding a ligand for the Notch1 receptor. Molec. Cell. Biol. 17:6057–6067, 1997.

[71478] 9808.Misaki, Y.; Pruijn, G. J. M.; van der Kemp, A. W. C. M.; van Venrooij,W. J.: The 56K autoantigen is identical to human annexin XI. J.Biol. Chem. 269: 4240–4246, 1994.

[71479] 9809.Morgan, R. O.; Bell, D. W.; Testa, J. R.; Fernandez, M.

P.: Genomiclocations of ANX11 and ANX13 and the evolutionary genetics of humanannexins. *Genomics* 48: 100–110, 1998.

[71480] 9810.Ono, T.; Kawabe, T.; Sonta, S.; Okamoto, T.: Assignment of MARK3alias KP78 to human chromosome band 14q32.3 by in situ hybridization. *Cytogenet.Cell Genet.* 79: 101–102, 1997.

[71481] 9811.Parsa, I.: Loss of Mr 78,000 marker in chemically induced transplantablecarcinomas and primary carcinoma of human pancreas. *Cancer Res.* 48:2265–2272, 1988.

[71482] 9812.Peng, C.–Y.; Graves, P. R.; Ogg, S.; Thoma, R. S.; Byrnes, M. J.,III; Wu, Z.; Stephenson, M. T.; Piwnica–Worms, H.: C–TAK1 proteinkinase phosphorylates human Cdc25C on serine 216 and promotes 14–3–3protein binding. *Cell Growth Differ.* 9: 197–208, 1998.

[71483] 9813.Brubaker, K.; Cowley, S. M.; Huang, K.; Loo, L.; Yochum, G. S.;Ayer, D. E.; Eisenman, R. N.; Radhakrishnan, I.: Solution structureof the interacting domains of the Mad–Sin3 complex: implications forrecruitment of a chromatin–modifying complex. *Cell* 103: 655–665,2000.

[71484] 9814.Jin, D.–Y.; Kozak, C. A.; Pangilinan, F.; Spencer, F.; Green, E.D.; Jeang, K.–T.: Mitotic checkpoint locus MAD1L1 maps to human chromosome7p22 and mouse

chromosome 5. *Genomics* 55: 363–364, 1999.

[71485] 9815.Jin, D.-Y.; Spencer, F.; Jeang, K.-T.: Human T cell leukemia virus type 1 oncoprotein Tax targets the human mitotic checkpoint protein MAD1. *Cell* 93: 81–91, 1998.

[71486] 9816.Tsukasaki, K.; Miller, C. W.; Greenspun, E.; Es-haghian, S.; Kawabata, H.; Fujimoto, T.; Tomonaga, M.; Sawyers, C.; Said, J. W.; Koeffler, H. P.: Mutations in the mitotic check point gene, MAD1L1, in human cancers. *Oncogene* 20: 3301–3305, 2001.

[71487] 9817.Kimura, M.; Kotani, S.; Hattori, T.; Sumi, N.; Yoshioka, T.; Todokoro, T.; Okano, Y.: Cell cycle-dependent expression and spindle pole localization of a novel human protein kinase Aik, related to Aurora of *Drosophila* and yeast Ipl1. *J. Biol. Chem.* 272: 13766–13771, 1997.

[71488] 9818.Goustin, A. S.: Personal Communication. Detroit, Mich. 8/17/1998.

[71489] 9819.Harrington, L.; McPhail, T.; Mar, V.; Zhou, W.; Oulton, R.; Bass, M. B.; Arruda, I.; Robinson, M. O.: A mammalian telomerase-associated protein. *Science* 275: 973–976, 1997.

[71490] 9820.Saito, T.; Matsuda, Y.; Suzuki, T.; Hayashi, A.; Yuan, X.; Saito, M.; Nakayama, J.; Hori, T.; Ishikawa, F.: Comparative gene mapping of the human and mouse TEP1 genes,

which encode one protein component of telomerases. *Genomics* 46: 46–50, 1997.

[71491] 9821. Coggins, K. G.; Latour, A.; Nguyen, M. S.; Audoly, L.; Coffman, T. M.; Koller, B. H.: Metabolism of PGE<sub>2</sub> by prostaglandin dehydrogenase is essential for remodeling the ductus arteriosus. (Letter) *Nature Med.* 8: 91–92, 2002.

[71492] 9822. Ensor, C. M.; Yang, J. Y.; Okita, R. T.; Tai, H. H.: Cloning and sequence analysis of the cDNA for human placental NAD(+)-dependent 15-hydroxyprostaglandin dehydrogenase. *J. Biol. Chem.* 265: 14888–14891, 1990.

[71493] 9823. Krook, M.; Marekov, L.; Jornvall, H.: Purification and structural characterization of placental NAD(+)-linked 15-hydroxyprostaglandin dehydrogenase: the primary structure reveals the enzyme to belong to the short-chain alcohol dehydrogenase family. *Biochemistry* 29: 738–743, 1990.

[71494] 9824. Chadwick, B. P.; Frischauf, A.-M.: The CD39-like gene family: identification of three new human members (CD39L2, CD39L3, and CD39L4), their murine homologues, and a member of the gene family from *Drosophila melanogaster*. *Genomics* 50: 357–367, 1998.

[71495] 9825. Enjyoji, K.; Sevigny, J.; Lin, Y.; Frenette, P. S.; Christie, P. D.; van Esch, J. S., II; Imai, M.; Edelberg, J. M.;

Rayburn, H.; Lech, M.; Beeler, D. L.; Csizmadia, E.; Wagner, D. D.; Robson, S. C.; Rosenberg, R. D.: Targeted disruption of cd39/ATP diphosphohydrolase results in disordered hemostasis and thromboregulation. *Nature Med.* 5: 1010–1017, 1999.

[71496] 9826. Granstein, R. D.: The skinny on CD39 in immunity and inflammation. *Nature Med.* 8: 336–338, 2002.

[71497] 9827. Gray, I. C.; Fallowfield, J.; Ford, S.; Nobile, C.; Volpi, E. V.; Spurr, N. K.: An integrated physical and genetic map spanning chromosome band 10q24. *Genomics* 43: 85–88, 1997.

[71498] 9828. Kaczmarek, E.; Koziak, K.; Seigniny, J.; Siegel, J. B.; Anrather, J.; Beaudoin, A. R.; Bach, F. H.; Robson, S. C.: Identification and characterization of CD39/vascular ATP diphosphohydrolase. *J. Biol. Chem.* 271: 33116–33122, 1996.

[71499] 9829. Maliszewski, C. R.; Delespesse, G. L.; Schoenborn, M. A.; Armitage, R. J.; Fanslow, W. C.; Nakajima, T.; Baker, E.; Sutherland, G. R.; Poindexter, K.; Birks, C.; Alpert, A.; Friend, D.; Gimpel, S. D.; Gayle, R. B., III: The CD39 lymphoid cell activation antigen. Molecular cloning and structural characterization. *J. Immun.* 153: 3574–3583, 1994.

[71500] 9830. Mizumoto, N.; Kumamoto, T.; Robson, S. C.; Sevi-



gny, J.; Matsue, H.; Enjoji, K.; Takashima, A.: CD39 is the dominant Langerhans cell-associated ecto-NTPDase: modulatory roles in inflammation and immune responsiveness. *Nature Med.* 8: 358–365, 2002.

[71501] 9831. Wolff, K.; Winkelmann, R. K.: Ultrastructural localization of nucleoside triphosphatase in Langerhans cells. *J. Invest. Derm.* 48:50–54, 1967.

[71502] 9832. Kieffer, L. J.; Seng, T. W.; Li, W.; Osterman, D. G.; Handschumacher, R. E.; Bayney, R. M.: Cyclophilin-40, a protein with homology to the P59 component of the steroid receptor complex: cloning of the cDNA and further characterization. *J. Biol. Chem.* 268: 12303–12310, 1993.

[71503] 9833. Kieffer, L. J.; Thalhammer, T.; Handschumacher, R. E.: Isolation and characterization of a 40-kDa cyclophilin-related protein. *J. Biol. Chem.* 267: 5503–5507, 1992.

[71504] 9834. Ratajczak, T.; Woollatt, E.; Kumar, P.; Ward, B. K.; Minchin, R. F.; Baker, E.: Cyclophilin 40 (PPID) gene map position 4q31.3. *Chromosome Res.* 5: 151 only, 1997.

[71505] 9835. Yokoi, H.; Shimizu, Y.; Anazawa, H.; Lefebvre, C. A.; Korneluk, R. G.; Ikeda, J.-E.: The structure and complete nucleotide sequence of the human cyclophilin 40 (PPID) gene. *Genomics* 35: 448–455, 1996.

[71506] 9836. Pizutti, A.; Novelli, G.; Ratti, A.; Amati, F.; Mari, A.;

Calabrese,G.; Nicolis, S.; Silani, V.; Marino, B.; Scarlato, G.; Ottolenghi,S.; Dallapiccola, B.: UFD1L, a developmentally expressed ubiquitination gene, is deleted in CATCH 22 syndrome. Hum. Molec. Genet. 6: 259–265,1997.

[71507] 9837.Yamagishi, H.; Garg, V.; Matsuoka, R.; Thomas, T.; Srivastava,D.: A molecular pathway revealing a genetic basis for human cardiac and craniofacial defects. Science 283: 1158–1161, 1999.

[71508] 9838.Gong, W.; Emanuel, B. S.; Collins, J.; Kim, D. H.; Wang, Z.; Chen,F.; Zhang, G.; Roe, B.; Budarf, M. L.: A transcription map of the DiGeorge and velo–cardio–facial syndrome minimal critical region on 22q11. Hum. Molec. Genet. 5: 789–800, 1996.

[71509] 9839.Gong, W.; Emanuel, B. S.; Galili, N.; Kim, D. H.; Roe, B.; Driscoll,D. A.; Budarf, M. L.: Structural and mutational analysis of a conserved gene (DGSI) from the minimal DiGeorge syndrome critical region. Hum.Molec. Genet. 6: 267–276, 1997.

[71510] 9840.Lindsay, E. A.; Botta, A.; Jurecic, V.; Carattini–Rivera, S.; Cheah,Y.–C.; Rosenblatt, H. M.; Bradley, A.; Baldini, A.: Congenital heart disease in mice deficient for the DiGeorge syndrome region. Nature 401:379–383, 1999.

[71511] 9841.Lindsay, E. A.; Harvey, E. L.; Scambler, P. J.; Baldini,

A.: ES2, a gene deleted in DiGeorge syndrome, encodes a nuclear protein and is expressed during early mouse development, where it shares an expression domain with a Goosecoid-like gene. *Hum. Molec. Genet.* 7: 629–635, 1998.

[71512] 9842. Fukunaga, R.; Hunter, T.: MNK1, a new MAP kinase-activated protein kinase, isolated by a novel expression screening method for identifying protein kinase substrates. *EMBO J.* 16: 1921–1933, 1997.

[71513] 9843. Pages, G.; Guerin, S.; Grall, D.; Bonino, F.; Smith, A.; Anjuere, F.; Auberger, P.; Pouyssegur, J.: Defective thymocyte maturation in p44 MAP kinase (Erk 1) knockout mice. *Science* 286: 1374–1378, 1999.

[71514] 9844. Stefanovsky, V. Y.; Pelletier, G.; Hannan, R.; Gagnon-Kugler, T.; Rothblum, L. I.; Moss, T.: An immediate response of ribosomal transcription to growth factor stimulation in mammals is mediated by ERK phosphorylation of UBF. *Molec. Cell* 8: 1063–1073, 2001.

[71515] 9845. Mengus, G.; May, M.; Carre, L.; Chambon, P.; Davidson, I.: Human TAF(II)135 potentiates transcriptional activation by the AF-2s of the retinoic acid, vitamin D3, and thyroid hormone receptors in mammalian cells. *Genes Dev.* 11: 1381–1395, 1997.

- [71516] 9846. Shimohata, T.; Nakajima, T.; Yamada, M.; Uchida, C.; Onodera, O.; Naruse, S.; Kimura, T.; Koide, R.; Nozaki, K.; Sano, Y.; Ishiguro, H.; Sakoe, K.; and 16 others: Expanded polyglutamine stretches interact with TAF(II)130, interfering with CREB-dependent transcription. *Nature Genet.* 26: 29–36, 2000.
- [71517] 9847. Ray, M. E.; Su, Y. A.; Meltzer, P. S.; Trent, J. M.: Isolation and characterization of genes associated with chromosome-6 mediated tumor suppression in human malignant melanoma. *Oncogene* 12: 2527–2533, 1996.
- [71518] 9848. Ray, M. E.; Wistow, G.; Su, Y. A.; Meltzer, P. S.; Trent, J. M.: AIM1, a novel non-lens member of the beta-gamma-crystallin superfamily, is associated with the control of tumorigenicity in human malignant melanoma. *Proc. Nat. Acad. Sci.* 94: 3229–3234, 1997.
- [71519] 9849. Teichmann, U.; Ray, M. E.; Ellison, J.; Graham, C.; Wistow, G.; Meltzer, P. S.; Trent, J. M.; Pavan, W. J.: Cloning and tissue expression of the mouse ortholog of AIM1, a beta-gamma-crystallin superfamily member. *Mammalian Genome* 9: 715–720, 1998.
- [71520] 9850. Trent, J. M.; Stanbridge, E. J.; McBride, H. L.; Meese, E. U.; Casey, G.; Araujo, D. E.; Witkowski, C. M.; Nagle, R. B.: Tumorigenicity in human melanoma cell lines controlled

by introduction of human chromosome 6. *Science* 247: 568–571, 1990.

[71521] 9851. Nakamura, Y.; Miura, K.; Fujino, Y.; Iwao, H.; Ogita, S.; Yamanaka, S.: Evolution, structure, and expression of GNPI/oscillin orthologous genes. *Genomics* 68: 179–186, 2000.

[71522] 9852. Rogers, M. J.; Ohgi, T.; Plumbridge, J.; Soll, D.: Nucleotide sequences of the *E. coli* *nagE* and *nagB* genes: the structural genes for the N-acetylglucosamine transport protein of the bacterial phosphoenolpyruvate:sugar phosphotransferase system and for glucosamine 6-phosphate deaminase. *Gene* 62:197–207, 1988.

[71523] 9853. Shevchenko, V.; Hogben, M.; Ekong, R.; Parrington, J.; Lai, F.A.: The human glucosamine-6-phosphate deaminase gene: cDNA cloning and expression, genomic organization and chromosomal localization. *Gene* 216:31–38, 1998.

[71524] 9854. Weidanz, J. A.; Campbell, P.; DeLucas, L. J.; Jin, J.; Moore, D.; Roden, L.; Yu, H.; Heilmann, E.; Vezza, A. C.: Glucosamine 6-phosphate deaminase in normal human erythrocytes. *Brit. J. Haemat.* 91: 72–79, 1995.

[71525] 9855. Bladergroen, B. A.; Strik, M. C. M.; Bovenschen, N.; van Berkum, O.; Scheffer, G. L.; Meijer, C. J. L. M.; Hack, C.

E.; Kummer, J.A.: The granzyme B inhibitor, protease inhibitor 9, is mainly expressed by dendritic cells and at immune-privileged sites. *J. Immun.* 166:3218–3225, 2001.

[71526] 9856. Eyre, H. J.; Sun, J.; Sutherland, G. R.; Bird, P.: Chromosomal mapping of the gene (PI9) encoding the intracellular serpin proteinase inhibitor 9 to 6p25 by fluorescence in situ hybridization. *Genomics* 37:406–408, 1996.

[71527] 9857. Krieg, S. A.; Krieg, A. J.; Shapiro, D. J.: A unique downstream estrogen responsive unit mediates estrogen induction of proteinase inhibitor-9, a cellular inhibitor of IL-1-beta-converting enzyme (caspase1). *Molec. Endocr.* 15: 1971–1982, 2001.

[71528] 9858. Sun, J.; Bird, C. H.; Sutton, V.; McDonald, L.; Coughlin, P. B.; De Jong, T. A.; Trapani, J. A.; Bird, P. I.: A cytosolic granzyme B inhibitor related to the viral apoptotic regulator cytokine response modifier A is present in cytotoxic lymphocytes. *J. Biol. Chem.* 271:27802–27809, 1996.

[71529] 9859. Steimle, V.; Durand, B.; Barras, E.; Zuffrey, M.; Hadam, M. R.; Mach, B.; Reith, W.: A novel DNA binding regulatory factor is mutated in primary MHC class II deficiency (bare lymphocyte syndrome). *Genes Dev.* 9: 1021–1032, 1995.

[71530] 9860. Teumer, J.; Tseng, H.; Green, H.: The human ba-

sonuclin gene. *Gene* 188:1–7, 1997.

- [71531] 9861. Tseng, H.; Green, H.: Basonuclin: a keratinocyte protein with multiple paired zinc fingers. *Proc. Nat. Acad. Sci.* 89: 10311–10315, 1992.
- [71532] 9862. Tseng, H.; Green, H.: Association of basonuclin with ability of keratinocytes to multiply and with absence of terminal differentiation. *J. Cell Biol.* 126: 495–506, 1994.
- [71533] 9863. Gibson, L.; Holmgreen, S. P.; Huang, D. C. S.; Bernard, O.; Copeland, N. G.; Jenkins, N. A.; Sutherland, G. R.; Baker, E.; Adams, J. M.; Cory, S.: bcl-w, a novel member of the bcl-2 family, promotes cell survival. *Oncogene* 13: 665–675, 1996.
- [71534] 9864. Ross, A. J.; Waymire, K. G.; Moss, J. E.; Parlow, A. F.; Skinner, M. K.; Russell, L. D.; MacGregor, G. R.: Testicular degeneration in Bclw-deficient mice. *Nature Genet.* 18: 251–256, 1998.
- [71535] 9865. Freedman, M. S.; Lucas, R. J.; Soni, B.; von Schantz, M.; Munoz, M.; David-Gray, Z.; Foster, R.: Regulation of mammalian circadian behavior by non-rod, non-cone, ocular photoreceptors. *Science* 284:502–504, 1999.
- [71536] 9866. Griffin, E. A., Jr.; Staknis, D.; Weitz, C. J.: Light-independent role of CRY1 and CRY2 in the mammalian circadian clock. *Science* 286:768–771, 1999.

- [71537] 9867.Hsu, D. S.; Zhao, X.; Zhao, S.; Kazantsev, A.; Wang, R.-P.; Todo,T.; Wei, Y.-F.; Sancar, A.: Putative human blue-light photoreceptorshCRY1 and hCRY2 are flavoproteins. *Biochemistry* 35: 13871–13877,1996.
- [71538] 9868.Kobayashi, K.; Kanno, S.; Smit, B.; van der Horst, G. T. J.; Takao,M.; Yasui, A.: Characterization of photolyase/blue-light receptorhomologs in mouse and human cells. *Nucleic Acids Res.* 26: 5086–5092,1998.
- [71539] 9869.Kume, K.; Zylka, M. J.; Sriram, S.; Shearman, L. P.; Weaver, D.R.; Jin, X.; Maywood, E. S.; Hastings, M. H.; Reppert, S. M.: mCRY1and mCRY2 are essential components of the negative limb of the circadianclock feedback loop. *Cell* 98: 193–205, 1999.
- [71540] 9870.Lucas, R. J.; Freedman, M. S.; Munoz, M.; Garcia-Fernandez, J.-M.;Foster, R. G.: Regulation of the mammalian pineal by non-rod, non-cone,ocular photoreceptors. *Science* 284: 505–507, 1999.
- [71541] 9871.Okamura, H.; Miyake, S.; Sumi, Y.; Yamaguchi, S.; Yasui, A.; Muijtjens,M.; Hoeijmakers, J. H. J.; van der Horst, G. T. J.: Photic inductionof mPer1 and mPer2 in Cry-deficient mice lacking a biological clock. *Science* 286:2531–2534, 1999.
- [71542] 9872.Reick, M.; Garcia, J. A.; Dudley, C.; McKnight, S. L.:



NPAS2:an analog of clock operative in the mammalian forebrain. Science 293:506–509, 2001.

- [71543] 9873.Todo, T.; Ryo, H.; Yamamoto, K.; Toh, H.; Inui, T.; Ayaki, H.;Nomura, T.; Ikenaga, M.: Similarity among the Drosophila (6–4)photolyase,a human photolyase homolog, and the DNA photolyase--blue-light photoreceptorfamily. Science 272: 109–112, 1996.
- [71544] 9874.van der Horst, G. T. J.; Muijtjens, M.; Kobayashi, K.; Takano,R.; Kanno, S.; Takao, M.; de Wit, J.; Verkerk, A.; Eker, A. P. M.;van Leenen, D.; Buijs, R.; Bootsma, D.; Hoeijmakers, J. H. J.; Yasui,A.: Mammalian Cry1 and Cry2 are essential for maintenance of circadianrhythms. Nature 398: 627–630, 1999.
- [71545] 9875.van der Spek, P. J.; Kobayashi, K.; Bootsma, D.; Takao, M.; Eker,A. P. M.; Yasui, A.: Cloning, tissue expression, and mapping of a human photolyase homolog with similarity to plant blue-light receptors. Genomics 37:177–182, 1996.
- [71546] 9876.Yagita, K.; Tamanini, F.; van der Horst, G. T. J.; Okamura, H.: Molecular mechanisms of the biological clock in cultured fibroblasts. Science 292:278–281, 2001.
- [71547] 9877.Villard, J.; Reith, W.; Barras, E.; Gos, A.; Morris, M. A.; Antonarakis,S. E.; Van den Elsen, P. J.; Mach, B.: Analy-

sis of mutations and chromosomal localisation of the gene encoding RFX5, a novel transcription factor affected in major histocompatibility complex class II deficiency.

Hum.Mutat. 10: 430–435, 1997.

[71548] 9878. Sipila, L.; Szatanik, M.; Vainionpää, H.; Ruotsalainen, H.; Myllylä, R.; Guenet, J.–L.: The genes encoding mouse lysyl hydroxylase isoforms map to chromosomes 4, 5, and 9. Mammalian Genome 11: 1132–1134, 2000.

[71549] 9879. Szpirer, C.; Szpirer, J.; Riviere, M.; Vanvooren, P.; Valtavaara, M.; Myllylä, R.: Localization of the gene encoding a novel isoform of lysyl hydroxylase. Mammalian Genome 8: 707–708, 1997.

[71550] 9880. Valtavaara, M.; Papponen, H.; Pirttilä, A.–M.; Hiltunen, K.; Helander, H.; Myllylä, R.: Cloning and characterization of a novel human lysyl hydroxylase isoform highly expressed in pancreas and muscle. J. Biol.Chem. 272: 6831–6834, 1997.

[71551] 9881. Hellborg, F.; Qian, W.; Mendez–Vidal, C.; Asker, C.; Kost–Alimova, M.; Wilhelm, M.; Imreh, S.; Wiman, K. G.: Human wig–1, a p53 target gene that encodes a growth inhibitory zinc finger protein. Oncogene 20: 5466–5474, 2001.

[71552] 9882. Varmeh–Ziaie, S.; Okan, I.; Wang, Y.; Magnusson, K.

P.; Warthoe,P.; Strauss, M.; Wiman, K. G.: Wig-1, a new p53-induced gene encoding a zinc finger protein. *Oncogene* 15: 2699-2704, 1997.

- [71553] 9883.Modregger, J.; Ritter, B.; Witter, B.; Paulsson, M.; Plomann, M.: All three PACSIN isoforms bind to endocytic proteins and inhibit endocytosis. *J. Cell Sci.* 113: 4511-4521, 2000.
- [71554] 9884.Fardaei, M.; Rogers, M. T.; Thorpe, H. M.; Larkin, K.; Hamshire, M. G.; Harper, P. S.; Brook, J. D.: Three proteins, MBNL, MBLL and MBXL, co-localize in vivo with nuclear foci of expanded-repeat transcripts in DM1 and DM2 cells. *Hum. Molec. Genet.* 11: 805-814, 2002.
- [71555] 9885.Miller, J. W.; Urbinati, C. R.; Teng-umnuay, P.; Stenberg, M. G.; Byrne, B. J.; Thornton, C. A.; Swanson, M. S.: Recruitment of human muscleblind proteins to (CUG)<sub>n</sub> expansions associated with myotonic dystrophy. *EMBO J.* 19: 4439-4448, 2000.
- [71556] 9886.Kutty, R. K.; Kutty, G.; Samuel, W.; Duncan, T.; Bridges, C. C.; El-Sherbeeney, A.; Nagineni, C. N.; Smith, S. B.; Wiggert, B.: Molecular characterization and developmental expression of NORPEG, a novel gene induced by retinoic acid. *J. Biol. Chem.* 276: 2831-2840, 2001.
- [71557] 9887.Aapola, U.; Shibuya, K.; Scott, H. S.; Ollila, J.; Vihi-

nen, M.;Heino, M.; Shintani, A.; Kawasaki, K.; Minoshima, S.; Krohn, K.; Antonarakis,S. E.; Shimizu, N.; Kudoh, J.; Peterson, P.: Isolation and initial characterization of a novel zinc finger gene, DNMT3L, on 21q22.3,related to the cysteine-5-methyltransferase 3 gene family. *Genomics* 65:293-298, 2000.

[71558] 9888.Bourc'his, D.; Xu, G.-L.; Lin, C.-S.; Bollman, B.; Bestor, T. H.: Dnmt3L and the establishment of maternal genomic imprints. *Science* 294:2536-2539, 2001.

[71559] 9889.Monney, L.; Sabatos, C.; Gaglia, J. L.; Ryu, A.; Waldner, H.; Chernova,T.; Manning, S.; Greenfield, E. A.; Coyle, A. J.; Sobel, R. A.; Freeman,G. J.; Kuchroo, V. K.: Th1-specific cell surface protein regulates macrophage activation and severity of an autoimmune disease. *Nature* 415:536-541, 2002.

[71560] 9890.Jourdan-Le Saux, C.; Le Saux, O.; Donlon, T.; Boyd, C. D.; Csiszar,K.: The human lysyl oxidase-related gene (LOXL2) maps between markersD8S280 and D8S278 on chromosome 8p21.2-p21.3. *Genomics* 51: 305-307,1998.

[71561] 9891.Jourdan-Le Saux, C.; Tronecker, H.; Bogic, L.; Bryant-Greenwood,G. D.; Boyd, C. D.; Csiszar, K.: The LOXL2 gene encodes a new lysyloxidase-like protein and

is expressed at high levels in reproductivetissues. J. Biol. Chem. 274: 12939–12944, 1999.

- [71562] 9892.Ikeda, K.; Sato, M.; Tsutsumi, O.; Tsuchiya, F.; Tsuneizumi, M.;Emi, M.; Imoto, I.; Inazawa, J.; Muramatsu, M.; Inoue, S.: Promoteranalysis and chromosomal mapping of human EBAG9 gene. Biochem. Biophys.Res. Commun. 273: 654–660, 2000.
- [71563] 9893.Nakashima, M.; Sonoda, K.; Watanabe, T.: Inhibition of cell growthand induction of apoptotic cell death by the human tumor-associatedantigen RCAS1. Nature Med. 5: 938–942, 1999.
- [71564] 9894.Kim, H.–S.; Choi, J.–Y.; Jung, A.–R.; Jang, K.–L.; Lee, W.–H.;Choi, W.–C.; Crow, T. J.; Hyun, B.–H.: Assignment of the human RhoHP1(ARHD) to chromosome 11q14.3 by radiation hybrid mapping. Cytogenet.Cell Genet. 89: 53 only, 2000.
- [71565] 9895.Shimizu, F.; Watanabe, T. K.; Okuno, S.; Omori, Y.; Fujiwara, T.;Takahashi, E.; Nakamura, Y.: Isolation of a novel human cDNA (rhoHP1)homologous to rho genes. Biochim. Biophys. Acta 1351: 13–16, 1997.
- [71566] 9896.Nagasaki, K.; Maass, N.; Manabe, T.; Hanzawa, H.; Tsukada, T.;Kikuchi, K.; Yamaguchi, K.: Identification of a novel gene, DAM1,amplified at chromosome 1p13.3–21

region in human breast cancer celllines. *Cancer Lett.* 140: 219–226, 1999.

[71567] 9897. Neubauer, G.; King, A.; Rappsilber, J.; Calvio, C.; Watson, M.; Ajuh, P.; Sleeman, J.; Lamond, A.; Mann, M.: Mass spectrometry and EST–database searching allows characterization of the multi–protein spliceosome complex. *Nature Genet.* 20: 46–50, 1998.

[71568] 9898. Rader, K.; Boyer, A. D.; Farquhar, M. G.; Arden, K. C.: Assignment of ankyrin repeat, family A (RFXANK–like) 2 (ANKRA2) to human chromosome 5q12–q13 by radiation hybrid mapping and somatic cell hybrid PCR. *Cytogenet. Cell Genet.* 89: 164–165, 2000.

[71569] 9899. Rader, K.; Orlando, R. A.; Lou, X.; Farquhar, M. G.: Characterization of ANKRA, a novel ankyrin repeat protein that interacts with the cytoplasmic domain of megalin. *J. Am. Soc. Nephrol.* 11: 2167–2178, 2000.

[71570] 9900. Nakagawa, H.; Koyama, K.; Murata, Y.; Morito, M.; Akiyama, T.; Nakamura, Y.: EB3, a novel member of the EB1 family preferentially expressed in the central nervous system, binds to a CNS–specific APC homologue. *Oncogene* 19: 210–216, 2000.

[71571] 9901. Dunbar, D. R.; Shibasaki, Y.; Dobbie, L.; Andersson, B.; Brookes, A. J.: In situ hybridisation mapping of genomic

clones for five human respiratory chain complex I genes. Cytogenet. Cell Genet. 78: 21–24, 1997.

[71572] 9902. Emahazion, T.; Brookes, A. J.: Mapping of the ND-UFA2, NDUFA6, NDUFA7, NDUF8, and NDUF8 electron transport chain genes by intron-based radiation hybrid mapping. Cytogenet. Cell Genet. 82: 114 only, 1998.

[71573] 9903. Ton, C.; Hwang, D. M.; Dempsey, A. A.; Liew, C.-C.: Identification and primary structure of five human NADH-ubiquinone oxidoreductase subunits. Biochem. Biophys. Res. Commun. 241: 589–594, 1997.

[71574] 9904. Walker, J. E.; Arizmendi, J. M.; Dupuis, A.; Fearnley, I. M.; Finel, M.; Medd, S. M.; Pilkington, S. J.; Runswick, M. J.; Skehel, J. M.: Sequences of 20 subunits of NADH:ubiquinone oxidoreductase from bovine heart mitochondria. J. Molec. Biol. 226: 1051–1072, 1992.

[71575] 9905. de Sury, R.; Martinez, P.; Procaccio, V.; Lunardi, J.; Issartel, J.-P.: Genomic structure of the human NDUF8 gene coding for the iron-sulfur TYKY subunit of the mitochondrial NADH:ubiquinone oxidoreductase. Gene 215: 1–10, 1998.

[71576] 9906. Loeffen, J.; Smeitink, J.; Triepels, R.; Smeets, R.; Schuelke, M.; Sengers, R.; Trijbels, F.; Hamel, B.; Mullaart, R.; van den Heuvel, L.: The first nuclear-encoded complex I

mutation in a patient with Leigh syndrome. *Am. J. Hum. Genet.* 63: 1598–1608, 1998.

[71577] 9907. Procaccio, V.; Depetris, D.; Soularue, P.; Mattei, M.-G.; Lunardi, J.; Issartel, J.-P.: cDNA sequence and chromosomal localization of the NDUF58 human gene coding for the 23 kDa subunit of the mitochondrial complex I.

*Biochim. Biophys. Acta* 1351: 37–41, 1997.

[71578] 9908. Cheng, H. F.; Jiang, M. J.; Chen, C. L.; Liu, S. M.; Wong, L. P.; Lomasney, J. W.; King, K.: Cloning and identification of amino acid residues of human phospholipase C delta 1 essential for catalysis. *J. Biol. Chem.* 270:

5495–5505, 1995.

[71579] 9909. Inhorn, R. C.; Aster, J. C.; Roach, S. A.; Slapak, C. A.; Soiffer, R.; Tantravani, R.; Stone, R. M.: A syndrome of lymphoblastic lymphoma, eosinophilia, and myeloid hyperplasia/malignancy associated with

t(8;13)(p11;q11): description of a distinctive clinicopathologic entity. *Blood* 85:1881–1887, 1995.

[71580] 9910. Michaux, L.; Mecucci, C.; Pereira Velloso, E. R.; Dierlamm, J.; Criel, A.; Louwagie, A.; van Orshoven, A.; Van den Berghe, H.: About the t(8;13)(p11;q12) clinico-pathologic entity. (Letter) *Blood* 87:1658–1659, 1996.

[71581] 9911. Reiter, A.; Sohal, J.; Kulkarni, S.; Chase, A.; Macdon-



ald, D. H.C.; Aguiar, R. C. T.; Goncalves, C.; Hernandez, J. M.; Jennings, B.A.; Goldman, J. M.; Cross, N. C. P.: Consistent fusion of ZNF198 to the fibroblast growth factor receptor-1 in the t(8;13)(p11;q12) myeloproliferative syndrome. *Blood* 92: 1735-1742, 1998.

[71582] 9912. Smedley, D.; Hamoudi, R.; Clark, J.; Warren, W.; Abdul-Rauf, M.; Somers, G.; Venter, D.; Fagan, K.; Cooper, C.; Shipley, J.: The t(8;13)(p11;q11-12) rearrangement associated with an atypical myeloproliferative disorder fuses the fibroblast growth factor receptor 1 gene to a novel gene R-AMP. *Hum. Molec. Genet.* 7: 637-642, 1998.

[71583] 9913. Still, I. H.; Cowell, J. K.: The t(8;13) atypical myeloproliferative disorder: further analysis of the ZNF198 gene and lack of evidence for multiple genes disrupted on chromosome 13. (Letter) *Blood* 92: 1456-1458, 1998.

[71584] 9914. Rousseau, D.; Gingras, A.-C.; Pause, A.; Sonenberg, N.: The eIF4E-binding proteins 1 and 2 are negative regulators of cell growth. *Oncogene* 13: 2415-2420, 1996.

[71585] 9915. Tsukiyama-Kohara, K.; Vidal, S. M.; Gingras, A.-C.; Glover, T.W.; Hanash, S. M.; Heng, H.; Sonenberg, N.: Tissue distribution, genomic structure, and chromosome mapping of mouse and human eukaryotic initiation factor 4E-binding proteins 1 and 2. *Genomics* 38:

353–363,1996.

- [71586] 9916.Meluh, P. B.; Koshland, D.: Suppressors of MIF2, a putative centromereprotein gene in *Saccharomyces cerevisiae*. (Abstract) *Molec. Biol.Cell* 6 (supp.): 360a only, 1995.
- [71587] 9917.Meluh, P. B.; Koshland, D.: Evidence that the MIF2 gene of *Saccharomycescerevisiae* encodes a centromere protein with homology to the mammaliancentromere protein CENP–C. *Molec. Biol. Cell* 6: 793–807, 1995.
- [71588] 9918.Cooper, E. C.; Aldape, K. D.; Abosch, A.; Barbaro, N. M.; Berger,M. S.; Peacock, W. S.; Jan, Y. N.; Jan, L. Y.: Colocalization andcoassembly of two human brain M-type potassium channel subunits thatare mutated in epilepsy. *Proc. Nat. Acad. Sci.* 97: 4914–4919, 2000.
- [71589] 9919.Wang, H.–S.; Pan, Z.; Shi, W.; Brown, B. S.; Wymore, R. S.; Cohen,I. S.; Dixon, J. E.; McKinnon, D.: KCNQ2 and KCNQ3 potassium channelsubunits: molecular correlates of the M–channel. *Science* 282: 1890–1893,1998.
- [71590] 9920.Yang, W.–P.; Levesque, P. C.; Little, W. A.; Conder, M. L.; Ramakrishnan,P.; Neubauer, M. G.; Blana, M. A.: Functional expression of twoKvLQT1–related potassium channels responsible for an inherited idiopathic epilepsy. *J. Biol. Chem.* 273: 19419–19423, 1998.

- [71591] 9921.Cecconi, F.; Alvarez-Bolado, G.; Meyer, B. I.; Roth, K. A.; Gruss,P.: Apaf1 (CED-4 homolog) regulates programmed cell death in mammaliandevlopment. Cell 94: 727-737, 1998.
- [71592] 9922.Hahn, C.; Hirsch, B.; Jahnke, D.; Durkop, H.; Stein, H.: Threenew types of Apaf-1 in mammalian cells. Biochem. Biophys. Res. Commun. 261:746-749, 1999.
- [71593] 9923.Honarpour, N.; Du, C.; Richardson, J. A.; Hammer, R. E.; Wang,X.; Herz, J.: Adult Apaf-1-deficient mice exhibit male infertility. Dev.Biol. 218: 248-258, 2000.
- [71594] 9924.Honarpour, N.; Gilbert, S. L.; Lahn, B. T.; Wang, X.; Herz, J.: Apaf-1 deficiency and neural tube closure defects are found in fogmice. Proc. Nat. Acad. Sci. 98: 9683-9687, 2001.
- [71595] 9925.Kim, H.; Jung, Y. K.; Kwon, Y. K.; Park, S. H.: Assign-ment ofapoptotic protease activating factor-1 gene (APAF1) to human chromosomeband 12q23 by fluo-escence in situ hybridization. Cytogenet. CellGenet. 87: 252-253, 1999.
- [71596] 9926.Kawai, J.; Suzuki, H.; Hara, A.; Hirose, K.; Watanabe, S.: Humanand mouse chromosomal mapping of Stac, a neuron-specific protein withan SH3 domain. Genomics 47: 140-142, 1998.

- [71597] 9927.Suzuki, H.; Kawai, J.; Taga, C.; Yaoi, T.; Hara, A.; Hirose, K.; Hayashizaki, Y.; Watanabe, S.: Stac, a novel neuron-specific protein with cysteine-rich and SH3 domains. *Biochem. Biophys. Res. Commun.* 229:902–909, 1996.
- [71598] 9928.Bradshaw, T.; Graves, T.; Biewald, T.: Personal Communication. St.Louis, Mo. 1/29/1998.
- [71599] 9929.Daga, A.; Micol, V.; Hess, D.; Aebersold, R.; Attardi, G.: Molecular characterization of the transcription termination factor from human mitochondria. *J. Biol. Chem.* 268: 8123–8130, 1993.
- [71600] 9930.Fernandez-Silva, P.; Martinez-Azorin, F.; Micol, V.; Attardi, G.: The human mitochondrial transcription termination factor (mTERF) is a multizipper protein but binds to DNA as a monomer, with evidence pointing to intramolecular leucine zipper interactions. *EMBO J.* 16:1066–1079, 1997.
- [71601] 9931.Hugnot, J. P.; Pedeutour, F.; Le Calvez, C.; Grosgeorge, J.; Passage, E.; Fontes, M.; Lazdunski, M.: The human inward rectifying K(+) channel Kir 2.2 (KCNJ12) gene: gene structure, assignment to chromosome 17p11.1, and identification of a simple tandem repeat polymorphism. *Genomics* 39:113–116, 1997.
- [71602] 9932.Namba, N.; Mori, R.; Tanaka, H.; Kondo, I.; Narahara,

K.; Seino,Y.: The inwardly rectifying potassium channel subunit Kir2.2v (KCNJN1)maps to 17p11.2–p11.1. Cyto-genet. Cell Genet. 79: 85–87, 1997.

[71603] 9933.Wible, B. A.; De Biasi, M.; Majumder, K.; Taglialatela, M.; Brown,A. M.: Cloning and functional expression of an inwardly rectifyingK(+) channel from human atrium. Circ. Res. 76: 343–350, 1995.

[71604] 9934.Imataka, H.; Olsen, H. S.; Sonenberg. N.: A new translationalregulator with homology to eukaryotic trans-lation initiation factor4G. EMBO J. 16: 817–825, 1997.

[71605] 9935.Levy–Strumpf, N.; Deiss, L. P.; Berissi, H.; Kimchi, A.: DAP–5,a novel homolog of eukaryotic translation initiation factor 4G isolatedas a putative modulator of gamma inter-feron–induced programmed cellddeath. Molec. Cell. Biol. 17: 1615–1625, 1997.

[71606] 9936.Shaughnessy, J. D., Jr.; Jenkins, N. A.; Copeland, N. G.: cDNAcloning, expression analysis, and chromosomal localization of a genewith high homology to wheat eIF–(iso)4F and mammalian eIF–4G. Genomics 39:192–197, 1997.

[71607] 9937.Yamanaka, S.; Poksay, K. S.; Arnold, K. S.; Innerarity, T. L.:A novel translational repressor mRNA is edited ex-tensively in liverscontaining tumors caused by the trans–

gene expression of the apoB mRNA-editing enzyme. *Genes Dev.* 11: 321–333, 1997.

- [71608] 9938. Fan, W.; Christensen, M.; Eichler, E.; Zhang, X.; Lennon, G.: Cloning, sequencing, gene organization, and localization of the human ribosomal protein RPL23A gene. *Genomics* 46: 234–239, 1997.
- [71609] 9939. Nomi, M.; Oishi, I.; Kani, S.; Suzuki, H.; Matsuda, T.; Yoda, A.; Kitamura, M.; Itoh, K.; Takeuchi, S.; Takeda, K.; Akira, S.; Ikeya, M.; Takada, S.; Minami, Y.: Loss of mRor1 enhances the heart and skeletal abnormalities in mRor2-deficient mice: redundant and pleiotropic functions of mRor1 and mRor2 receptor tyrosine kinases. *Molec. Cell. Biol.* 21: 8329–8335, 2001.
- [71610] 9940. Westendorf, J. M.; Rao, P. N.; Gerace, L.: Cloning of cDNAs for M-phase phosphoproteins recognized by the MPM2 monoclonal antibody and determination of the phosphorylated epitope. *Proc. Nat. Acad. Sci.* 91: 714–718, 1994.
- [71611] 9941. Wang, X. S.; Diener, K.; Jannuzzi, D.; Trollinger, D.; Tan, T.-H.; Lichenstein, H.; Zukowski, M.; Yao, Z.: Molecular cloning and characterization of a novel protein kinase with a catalytic domain homologous to mitogen-activated protein kinase kinase kinase. *J. Biol. Chem.* 271:

31607–31611, 1996.

[71612] 9942.Chuaqui, R. F.; Englert, C. R.; Strup, S. E.; Vocke, C. D.; Zhuang,Z.; Duray, P. H.; Bostwick, D. G.; Linehan, W. M.; Liotta, L. A.;Emmert–Buck, M. R.: Identification of a novel transcript up–regulatedin a clinically aggressive prostate carcinoma. *Urology* 50: 302–307,1997.

[71613] 9943.Cole, K. A.; Chuaqui, R. F.; Katz, K.; Pack, S.; Zhuang, Z.; Cole,C. E.; Lyne, J. C.; Linehan, W. M.; Liotta, L. A.; Emmert–Buck, M.R.: cDNA sequencing and analysis of POV1 (PB39): a novel gene up–regulatedin prostate cancer. *Genomics* 51: 282–287, 1998.

[71614] 9944.Bono, P.; Salmi, M.; Smith, D. J.; Leppanen, I.; Horelli–Kuitunen,N.; Palotie, A.; Jalkanen, S.: Isolation, structural characterization,and chromosomal mapping of the mouse vascular adhesion protein–1 geneand promoter. *J. Immun.* 161: 2953–2960, 1998.

[71615] 9945.Morris, N. J.; Ducret, A.; Aebersold, R.; Ross, S. A.; Keller,S. R.; Lienhard, G. E.: Membrane amine oxidase cloning and identificationas a major protein in the adipocyte plasma membrane. *J. Biol. Chem.* 272:9388–9392, 1997.

[71616] 9946.Bromme, D.; Rossi, A. B.; Smeekens, S. P.; Anderson, D. C.; Payan,D. G.: Human bleomycin hydrolase: molecular

cloning, sequencing, functional expression, and enzymatic characterization. *Biochemistry* 35:6706–6714, 1996.

- [71617] 9947. Cloos, J.; Nieuwenhuis, E. J. C.; Boomsma, D. I.; Kuik, D. J.; van der Sterre, M. L. T.; Arwert, F.; Snow, G. B.; Braakhuis, B. J. M.: Inherited susceptibility to bleomycin-induced chromatid breaks in cultured peripheral blood lymphocytes. *J. Nat. Cancer Inst.* 91:1125–1130, 1999.
- [71618] 9948. Farrer, L. A.; Abraham, C. R.; Haines, J. L.; Rogaeva, E. A.; Song, Y.; McGraw, W. T.; Brindle, N.; Premkumar, S.; Scott, W. K.; Yamaoka, L. H.; Saunders, A. M.; Roses, A. D.; Auerbach, S. A.; Sorbi, S.; Duara, R.; Pericak-Vance, M. A.; St. George-Hyslop, P. H.: Association between bleomycin hydrolase and Alzheimer's disease in Caucasians. *Ann. Neurol.* 44: 808–811, 1998.
- [71619] 9949. Ferrando, A. A.; Pendas, A. M.; Llano, E.; Velasco, G.; Lidereau, R.; Lopez-Otin, C.: Gene characterization, promoter analysis, and chromosomal localization of human bleomycin hydrolase. *J. Biol. Chem.* 272:33298–33304, 1997.
- [71620] 9950. Ferrando, A. A.; Velasco, G.; Campo, E.; Lopez-Otin, C.: Cloning and expression analysis of human bleomycin hydrolase, a cysteine proteinase involved in chemotherapy resistance. *Cancer Res.* 56: 1746–1750, 1996.



- [71621] 9951.Haston, C. K.; Amos, C. I.; King, T. M.; Travis, E. L.: Inheritance of susceptibility to bleomycin-induced pulmonary fibrosis in the mouse. *Cancer Res.* 56: 2596–2601, 1996.
- [71622] 9952.Hsu, T. C.; Johnston, D. A.; Cherry, L. M.; Ramkissoon, D.; Schantz, S. P.; Jessup, J. M.; Winn, R. J.; Shirley, L.; Furlong, C.: Sensitivity to genotoxic effects of bleomycin in humans: possible relationship to environmental carcinogenesis. *Int. J. Cancer* 43: 403–409, 1989.
- [71623] 9953.Lazo, J. S.; Humphreys, C. J.: Lack of metabolism as the biochemical basis of bleomycin-induced pulmonary toxicity. *Proc. Nat. Acad. Sci.* 80:3064–3068, 1983.
- [71624] 9954.Montoya, S. E.; Aston, C. E.; DeKosky, S. T.; Kamboh, M. I.; Lazo, J. S.; Ferrell, R. E.: Bleomycin hydrolase is associated with risk of sporadic Alzheimer's disease. (Letter) *Nature Genet.* 18: 211–212, 1998. Note: Erratum: *Nature Genet.* 19: 404 only, 1998.
- [71625] 9955.Montoya, S. E.; Ferrell, R. E.; Lazo, J. S.: Genomic structure and genetic mapping of the human neutral cysteine protease bleomycin hydrolase. *Cancer Res.* 57: 4191–4195, 1997.
- [71626] 9956.Zheng, W.; Johnston, S. A.; Joshua-Tor, L.: The unusual active site of Gal6/bleomycin hydrolase can act as a

carboxypeptidase, aminopeptidase, and peptide ligase.

Cell 93: 103–109, 1998.

- [71627] 9957. Xu, X.-N.; Screaton, G. R.; Gotch, F. M.; Dong, T.; Tan, R.; Almond, N.; Walker, B.; Stebbings, R.; Kent, K.; Nagata, S.; Stott, J. E.; McMichael, A. J.: Evasion of cytotoxic T lymphocyte (CTL) responses by Nef-dependent induction of Fas ligand (CD95L) expression on simian immunodeficiency virus-infected cells. *J. Exp. Med.* 186: 7–16, 1997.
- [71628] 9958. Huang, L. J.; Durick, K.; Weiner, J. A.; Chun, J.; Taylor, S. S.: Identification of a novel protein kinase A anchoring protein that binds both type I and type II regulatory subunits. *J. Biol. Chem.* 272: 8057–8064, 1997.
- [71629] 9959. Lin, R.-Y.; Moss, S. B.; Rubin, C. S.: Characterization of S-AKAP84, a novel developmentally regulated A kinase anchor protein of male germ cells. *J. Biol. Chem.* 270: 27804–27811, 1995.
- [71630] 9960. Trendelenburg, G.; Hummel, M.; Riecken, E.-O.; Hanski, C.: Molecular characterization of AKAP149, a novel A kinase anchor protein with a KH domain. *Biochem. Biophys. Res. Commun.* 225: 313–319, 1996.
- [71631] 9961. Moshous, D.; Callebaut, I.; de Chasseval, R.; Corneo, B.; Cavazzana-Calvo, M.; Le Deist, F.; Tezcan, I.; Sanal, O.; Bertrand, Y.; Philippe, N.; Fischer, A.; de Villartay, J.-P.:

Artemis, a novel DNA double-strandbreak repair/V(D)J recombination protein, is mutated in human severe combined immune deficiency. *Cell* 105: 177–186, 2001.

[71632] 9962.Chang, K.; Hanaoka, K.; Kumada, M.; Takuwa, Y.: Molecular cloning and functional analysis of a novel P2 nucleotide receptor. *J. Biol.Chem.* 270: 26152–26158, 1995.

[71633] 9963.Communi, D.; Parmentier, M.; Boeynaems, J.–M.: Cloning, functional expression and tissue distribution of the human P2Y6 receptor. *Biochem.Biophys. Res. Commun.* 222: 303–308, 1996.

[71634] 9964.Maier, R.; Glatz, A.; Mosbacher, J.; Bilbe, G.: Cloning of P2Y6 cDNAs and identification of a pseudogene: comparison of P2Y receptor subtype expression in bone and brain tissues. *Biochem. Biophys. Res.Comm.* 237: 297–302, 1997.

[71635] 9965.Pidlaoan, L. V.; Jin, J.; Sandhu, A. K.; Athwal, R. S.; Kunapuli, S. P.: Colocalization of P2Y2 and P2Y6 receptor genes at human chromosome 11q13.3–14.1. *Somat. Cell Molec. Genet.* 23: 291–296, 1997.

[71636] 9966.Wang, B.; Kishihara, K.; Zhang, D.; Hara, H.; Nomoto, K.: Molecular cloning and characterization of a novel human receptor protein tyrosine phosphatase gene, hPTP-J: down-regulation of gene expression by PMA and calcium

ionophore in Jurkat T lymphoma cells. *Biochem. Biophys. Res. Commun.* 231: 77–81, 1997.

[71637] 9967. Wang, H.; Lian, Z.; Lerch, M. M.; Chen, Z.; Xie, W.; Ullrich, A.: Characterization of PCP-2, a novel receptor protein tyrosine phosphatase of the MAM domain family. *Oncogene* 12: 2555–2562, 1996.

[71638] 9968. Chinnaiyan, A. M.; O'Rourke, K.; Tewari, M.; Dixit, V. M.: FADD, a novel death domain-containing protein, interacts with the death domain of Fas and initiates apoptosis. *Cell* 81: 505–512, 1995.

[71639] 9969. Kabra, N. H.; Kang, C.; Hsing, L. C.; Zhang, J.; Winoto, A.: T cell-specific FADD-deficient mice: FADD is required for early T cell development. *Proc. Nat. Acad. Sci.* 98: 6307–6312, 2001.

[71640] 9970. Yanagisawa, K.; Sakakibara, Y.; Suiko, M.; Takami, Y.; Nakayama, T.; Nakajima, H.; Takayanagi, K.; Natori, Y.; Liu, M.-C.: cDNA cloning, expression, and characterization of the human bifunctional ATP sulfurylase/adenosine 5'-prime-phosphosulfate kinase enzyme. *Biosci. Biotech. Biochem.* 62: 1037–1040, 1998.

[71641] 9971. Sahin, U.; Tureci, O.; Schmitt, H.; Cochlovius, B.; Johannes, T.; Schmits, R.; Stenner, F.; Luo, G.; Schobert, I.; Pfreundschuh, M.: Human neoplasms elicit multiple spe-

cific immune responses in the autologous host. Proc. Nat. Acad. Sci. 92: 11810–11813, 1995.

[71642] 1992. Tureci, O.; Sahin, U.; Vollmar, E.; Siemer, S.; Gottert, E.; Seitz, G.; Parkkila, A. K.; Shah, G. N.; Grubb, J. H.; Pfrendrich, M.; Sly, W. S.: Human carbonic anhydrase XII: cDNA cloning, expression, and chromosomal localization of a carbonic anhydrase gene that is overexpressed in some renal cell cancers. Proc. Nat. Acad. Sci. 95:7608–7613, 1998.

[71643] 1993. Herrscher, R. F.; Kaplan, M. H.; Lelsz, D. L.; Das, C.; Scheuermann, R.; Tucker, P. W.: The immunoglobulin heavy-chain matrix-associating regions are bound by Bright: a B cell-specific trans-activator that describes a new DNA-binding protein family. Genes Dev. 9: 3067–3082, 1995.

[71644] 1994. Kortschak, R. D.; Reimann, H.; Zimmer, M.; Eyre, H. J.; Saint, R.; Jenne, D. E.: The human dead ringer/bright homolog, DRIL1: cDNA cloning, gene structure, and mapping to D19S886, a marker on 19p13.3 that is strictly linked to the Peutz-Jeghers syndrome. Genomics 51:288–292, 1998.

[71645] 1995. Kamei, M.; Webb, G. C.; Young, I. G.; Campbell, H. D.: SOLH, a human homologue of the Drosophila

melanogaster small optic lobes gene is a member of the calpain and zinc-finger gene families and maps to human chromosome 16p13.3 near CATM (cataract with microphthalmia). *Genomics* 51:197–206, 1998.

[71646] 1996. Eriksson, I.; Sandback, D.; Ek, B.; Lindahl, U.; Kjellen, L.: cDNA cloning and sequencing of mouse mastocytoma glucosaminyl N-deacetylase/N-sulfotransferase, an enzyme involved in the biosynthesis of heparin. *J. Biol. Chem.* 269:10438–10443, 1994.

[71647] 1997. Forsberg, E.; Pejler, G.; Ringvall, M.; Lunderius, C.; Tomasini-Johansson, B.; Kusche-Gullberg, M.; Eriksson, I.; Ledin, J.; Hellman, L.; Kjellen, L.: Abnormal mast cells in mice deficient in a heparin-synthesizing enzyme. *Nature* 400: 773–776, 1999.

[71648] 1998. Humphries, D. E.; Lanciotti, J.; Karlinsky, J. B.: cDNA cloning, genomic organization and chromosomal localization of human heparan glucosaminyl N-deacetylase/N-sulphotransferase-2. *Biochem. J.* 332:303–307, 1998.

[71649] 1999. Humphries, D. E.; Wong, G. W.; Friend, D. S.; Gurish, M. F.; Qiu, W.-T.; Huang, C.; Sharpe, A. H.; Stevens, R. L.: Heparin is essential for the storage of specific granule proteases in mast cells. *Nature* 400:769–772, 1999.

- [71650] 9980.Orellana, A.; Hirschberg, C. B.; Wei, Z.; Swiedler, S. J.; Ishihara,M.: Molecular cloning and expression of a glycosaminoglycan N-acetylglucosaminylN-deacetylase/N-sulfotransferase from a heparin-producing cell line. *J.Biol. Chem.* 269: 2270-2276, 1994.
- [71651] 9981.Zehnder, J. L.; Galli, S. J.: Mast-cell heparin demystified. *Nature* 400:714-715, 1999.
- [71652] 9982.Lavedan, C.; Buchholtz, S.; Auburger, G.; Albin, R. L.; Athanassiadou,A.; Blancato, J.; Burguera, J. A.; Ferrell, R. E.; Kostic, V.; Leroy,E.; Leube, B.; Mota-Vieira, L.; and 9 others: Absence of mutationin the beta- and gamma-synuclein genes in familial autosomal dominantParkinson's disease. *DNA Research* 5: 401-402, 1998.
- [71653] 9983.Leek, J. P.; Hamlin, P. J.; Wilton, J.; Lench, N. J.: Assignmentof the Rab13 gene (RAB13) to human chromosome band 12q13 by in situhybridization. *Cytogenet. Cell Genet.* 79: 210-211, 1997.
- [71654] 9984.Novick, P.; Zerial, M.: The diversity of Rab proteins in vesicletransport. *Curr. Opin. Cell Biol.* 9: 496-504, 1997.
- [71655] 9985.Salminen, A.; Novick, P. J.: A ras-like protein is required fora post-Golgi event in yeast secretion. *Cell* 49:

527–538, 1987.

- [71656] 9986.Zahraoui, A.; Joberty, G.; Arpin, M.; Fontaine, J. J.; Hellio,R.; Tavitian, A.; Louvard, D.: A small rab GTPase is distributedin cytoplasmic vesicles in nonpolarized cells but colocalizes withthe tight junction marker ZO–1 in polarized epithelial cells. *J.Cell. Biol.* 124: 101–115, 1994.
- [71657] 9987.Liu, X.–L.; Wazer, D. E.; Watanabe, K.; Band, V.: Identificationof a novel serine protease–like gene, the expression of which is down–regulatedduring breast cancer progression. *Cancer Res.* 56: 3371–3379, 1996.
- [71658] 9988.Polikoff, D.; Kuo, W.–L.; Cochran, J. F.; Wernick, M.; Kowbel,D.; Myambo, K.; Collins, C. C.: Assignment of protease, serine–like1 (PRSSL1) to human chromosome 19q13 by in situ hybridization andradiation hybrid mapping. *Cytogenet. Cell Genet.* 79: 147–148, 1997.
- [71659] 9989.Aurich–Costa, J.; Cadel, S.; Gouzy, C.; Foulon, T.; Cherif, D.;Cohen, P.: Assignment of the aminopeptidase B gene (RNPEP) to humanchromosome 1 band q32 by in situ hybridization. *Cytogenet. Cell Genet.* 79:143–144, 1997.
- [71660] 9990.Cadel, S.; Foulon, T.; Viron, A.; Balogh, A.; Midol–Monnet, S.;Noel, N.; Cohen, P.: Aminopeptidase B from the rat testis is a bifunctionalenzyme structurally related to leukotriene–A4 hydrolase. *Proc. Nat.Acad. Sci.* 94:



2963–2968, 1997.

- [71661] 9991.Cadel, S.; Pierotti, A. R.; Foulton, T.; Creminon, C.; Barre, N.;Segretain, D.; Cohen, P.: Aminopeptidase–B in the rat testes: isolation,functional properties and cellular localization in the seminiferoustubules. *Molec. Cell. Endocr.* 110: 149–160, 1995.
- [71662] 9992.Hopsu, V. K.; Kantonen, U. M.; Glenner, G. G.: A peptidase fromrat tissues selectively hydrolyzing N–terminal arginine and lysineresidues. *Life Sci.* 3: 1449–1453, 1964.
- [71663] 9993.Ershova, G.; Derre, J.; Chetelin, S.; Nancy, V.; Berger, R.; Kaplan,J.; Munnich, A.; de Gunzburg, J.: cDNA sequence, genomic organizationand mapping of PDE6D, the human gene encoding the delta subunit ofthe cGMP phosphodiesterase of retinal rod cells to chromosome 2q36. *Cytogenet.Cell Genet.* 79: 139–141, 1997.
- [71664] 9994.Florio, S. K.; Prusti, R. K.; Beavo, J. A.: Solubilization ofmembrane–bound rod phosphodiesterase by the rod phosphodiesteraserecombinant delta subunit. *J. Biol. Chem.* 271: 24036–24047, 1996.
- [71665] 9995.Li, N.; Florio, S. K.; Pettenati, M. J.; Rao, P. N.; Beavo, J.A.; Baehr, W.: Characterization of human and mouse rod cGMP phosphodiesterasedelta subunit (PDE6D) and chro–

mosomal localization of the human gene. Genomics 49:76–82, 1998.

- [71666] 9996.Jacquemin, P.; Martial, J. A.; Davidson, I.: Human TEF-5 is preferentially expressed in placenta and binds to multiple functional elements of the human chorionic somatomammotropin-B gene enhancer. J. Biol. Chem. 272:12928–12937, 1997.
- [71667] 9997.Jiang, S.-W.; Wu, K.; Eberhardt, N. L.: Human placental TEF-5 transactivates the human chorionic somatomammotropin gene enhancer. Molec.Endocr. 13: 879–889, 1999.
- [71668] 9998.Enattah, N. S.; Sahi, T.; Savilahti, E.; Terwilliger, J. D.; Peltonen, L.; Jarvela, I.: Identification of a variant associated with adult-type hypolactasia. Nature Genet. 30: 233–237, 2002.
- [71669] 9999.Jarvela, I.; Enattah, N. S.; Kokkonen, J.; Varilo, T.; Savilahti, E.; Peltonen, L.: Assignment of the locus for congenital lactase deficiency to 2q21, in the vicinity of but separate from the lactase-phlorizin hydrolase gene. Am. J. Hum. Genet. 63: 1078–1085, 1998.
- [71670] 10000.Kotaka, M.; Kostin, S.; Ngai, S.; Chan, K.; Lau, Y.; Lee, S. M.Y.; Li, H.; Ng, E. K. O.; Schaper, J.; Tsui, S. K. W.; Fung, K.; Lee, C.; Waye, M. M. Y.: Interaction of hCLIM1, an

Enigma family protein,with alpha-actinin 2. J. Cell.

Biochem. 78: 558–565, 2000.

[71671] 10001.Kotaka, M.; Ngai, S.-M.; Garcia-Barcelo, M.; Tsui, S. K. W.; Fung,K.-P.; Lee, C.-Y.; Waye, M. M. Y.: Characteri-  
zation of the human36-kDa carboxyl terminal LIM domain  
protein (hCLIM1). J. Cell. Biochem. 72:279–285, 1999.

[71672] 10002.Bach, I.: The LIM domain: regulation by association.  
Mech. Dev. 91:5–17, 2000.

[71673] 10003.Guy, P. M.; Kenny, D. A.; Gill, G. N.: The PDZ do-  
main of the LIMprotein enigma binds to beta-  
tropomyosin. Molec. Biol. Cell 10:1973–1984, 1999.

[71674] 10004.Kuroda, S.; Tokunaga, C.; Kiyohara, Y.; Higuchi, O.;  
Konishi, H.;Mizuno, K.; Gill, G. N.; Kikkawa, U.: Protein-  
protein interactionof zinc finger LIM domains with protein  
kinase C. J. Biol. Chem. 271:31029–31032, 1996.

[71675] 10005.Wu, R.-Y.; Gill, G. N.: LIM domain recognition of a  
tyrosine-containingtight turn. J. Biol. Chem. 269:  
25085–25090, 1994.

[71676] 10006.Rudnick, A.; Ling, T. Y.; Odagiri, H.; Rutter, W. J.;  
German, M.S.: Pancreatic beta cells express a diverse set  
of homeobox genes. Proc.Nat. Acad. Sci. 91:  
12203–12207, 1994.

[71677] 10007.ten Berge, D.; Brouwer, A.; El Bahi, S.; Guenet, J.-L.;

Robert,B.; Meijlink, F.: Mouse *Alx3*: an *aristaless*-like homeobox gene expressed during embryogenesis in ectomesenchyme and lateral plate mesoderm. *Dev.Biol.* 199: 11–25, 1998.

[71678] 10008.Khoja, H.; Wang, G.; Ng, C.-T. L.; Tucker, J.; Brown, T.; Shyamala,V.: Cloning of CCRL1, an orphan seven transmembrane receptor related to chemokine receptors, expressed abundantly in the heart. *Gene* 246:229–238, 2000.

[71679] 10009.Schweickart, V. L.; Epp, A.; Raport, C. J.; Gray, P. W.: CCR11 is a functional receptor for the monocyte chemoattractant protein family of chemokines. *J. Biol. Chem.* 275: 9550–9556, 2000. Note:Erratum: *J. Biol. Chem.* 276: 856 only, 2001.

[71680] 10010.Launay, P.; Fleig, A.; Perraud, A.-L.; Scharenberg, A. M.; Penner,R.; Kinet, J.-P.: TRPM4 is a  $\text{Ca}^{2+}$ -activated nonselective cation channel mediating cell membrane depolarization. *Cell* 109: 397–407,2002.

[71681] 10011.Xu, X.-Z. S.; Moebius, F.; Gill, D. L.; Montell, C.: Regulation of melastatin, a TRP-related protein, through interaction with a cytoplasmic isoform. *Proc. Nat. Acad. Sci.* 98: 10692–10697, 2001.

[71682] 10012.Astrin, K. H.; Warner, C. A.; Yoo, H.-W.; Goodfel-

low, P. J.; Tsai, S.-F.; Desnick, R. J.: Regional assignment of the human uroporphyrinogen III synthase (UROS) gene to chromosome 10q25.2-q26.3. *Hum. Genet.* 87:18-22, 1991.

[71683] 10013. Boulechfar, S.; Da Silva, V.; Deybach, J.-C.; Nordmann, Y.; Grandchamp, B.; de Verneuil, H.: Heterogeneity of mutations in the uroporphyrinogen III synthase gene in congenital erythropoietic porphyria. *Hum. Genet.* 88:320-324, 1992.

[71684] 10014. Frank, J.; Wang, X.; Lam, H.-M.; Aita, V. M.; Jugert, F. K.; Goerz, G.; Merk, H. F.; Poh-Fitzpatrick, M. B.; Christiano, A. M.: C73R is a hotspot mutation in the uroporphyrinogen III synthase gene in congenital erythropoietic porphyria. *Ann. Hum. Genet.* 62: 225-230, 1998.

[71685] 10015. Shady, A. A.; Colby, B. R.; Cunha, L. F.; Astrin, K. H.; Bishop, D. F.; Desnick, R. J.: Congenital erythropoietic porphyria: identification and expression of eight novel mutations in the uroporphyrinogen III synthase gene. *Brit. J. Haemat.* 117: 980-987, 2002.

[71686] 10016. Tanigawa, K.; Takamura, N.; Yamashita, S.: Congenital erythropoietic porphyria. *Nippon Rinsho* 53: 1422-1426, 1995.

[71687] 10017. Tsai, S.-F.; Bishop, D. F.; Desnick, R. J.: Human

uroporphyrinogenIII synthase: molecular cloning, nucleotide sequence, and expression of a full-length cDNA. Proc. Nat. Acad. Sci. 85: 7049–7053, 1988.

- [71688] 10018. Warner, C. A.; Yoo, H.-W.; Roberts, A. G.; Desnick, R. J.: Congenital erythropoietic porphyria: identification and expression of exonic mutations in the uroporphyrinogen III synthase gene. J. Clin. Invest. 89:693–700, 1992.
- [71689] 10019. Warner, C. A.; Yoo, H. W.; Tsai, S.-F.; Roberts, A. G.; Desnick, R. J.: Congenital erythropoietic porphyria: characterization of the genomic structure and identification of mutations in the uroporphyrinogenIII synthase gene. (Abstract) Am. J. Hum. Genet. 47 (suppl.): A83 only, 1990.
- [71690] 10020. Lapsys, N. M.; Layfield, R.; Baker, E.; Callen, D. F.; Sutherland, G. R.; Abedinia, M.; Nixon, P. F.; Mattick, J. S.: Chromosomal location of the human transketolase gene. Cytogenet. Cell Genet. 61: 274–275, 1992.
- [71691] 10021. Gladyshev, V. N.; Liu, A.; Novoselov, S. V.; Krysan, K.; Sun, Q.-A.; Kryukov, V. M.; Kryukov, G. V.; Lou, M. F.: Identification and characterization of a new mammalian glutaredoxin (thioltransferase), Grx2. J. Biol. Chem. 276: 30374–30380, 2001.
- [71692] 10022. Lundberg, M.; Johansson, C.; Chandra, J.; Enoks-

son, M.; Jacobsson, G.; Ljung, J.; Johansson, M.; Holmgren, A.: Cloning and expression of a novel human glutaredoxin (Grx2) with mitochondrial and nuclear isoforms. *J. Biol. Chem.* 276: 26269–26275, 2001.

- [71693] 10023. Ungar, D.; Oka, T.; Brittle, E. E.; Vasile, E.; Lupashin, V. V.; Chatterton, J. E.; Heuser, J. E.; Krieger, M.; Waters, M. G.: Characterization of a mammalian Golgi-localized protein complex, COG, that is required for normal Golgi morphology and function. *J. Cell Biol.* 157: 405–415, 2002.
- [71694] 10024. Walter, D. M.; Paul, K. S.; Waters, M. G.: Purification and characterization of a novel 13 S hetero oligomeric protein complex that stimulates *in vitro* Golgi transport. *J. Biol. Chem.* 273: 29565–29576, 1998.
- [71695] 10025. Baloh, R. H.; Tansey, M. G.; Golden, J. P.; Creedon, D. J.; Heuckeroth, R. O.; Keck, C. L.; Zimonjic, D. B.; Popescu, N. C.; Johnson, E. M., Jr.; Milbrandt, J.: TrnR2, a novel receptor that mediates neurturin and GDNF signaling through Ret. *Neuron* 18: 793–802, 1997.
- [71696] 10026. Meuwissen, R. L. J.; Meerts, I.; Hoovers, J. M. N.; Leschot, N. J.; Heyting, C.: Human synaptonemal complex protein 1 (SCP1): isolation and characterization of the cDNA and chromosomal localization of the gene. *Genomics*

39: 377–384, 1997.

- [71697] 10027. Meuwissen, R. L. J.; Offenberg, H. H.; Dietrich, A. J. J.; Riesewijk, A.; van Iersel, M.; Heyting, C.: A coiled-coil related protein specific for synapsed regions of meiotic prophase chromosomes. *EMBO J.* 11:5091–5100, 1992.
- [71698] 10028. Levedakou, E. N.; He, M.; Baptist, E. W.; Craven, R. J.; Cance, W. G.; Welcsh, P. L.; Simmons, A.; Naylor, S. L.; Leach, R. L.; Lewis, T. B.; Bowcock, A.; Liu, E. T.: Two novel human serine/threonine kinases with homologies to the cell cycle regulating *Xenopus* MO15, and NIMA kinases: cloning and characterization of their expression pattern. *Oncogene* 9: 1977–1988, 1994.
- [71699] 10029. Kondoh, N.; Nishina, Y.; Tsuchida, J.; Koga, M.; Tanaka, H.; Uchida, K.; Inazawa, J.; Taketo, M.; Nozaki, M.; Nojima, H.; Matsumiya, K.; Namiki, M.; Okuyama, A.; Nishimune, Y.: Assignment of synaptonemal complex protein 1 (SCP1) to human chromosome 1p13 by fluorescence in situ hybridization and its expression in the testis. *Cytogenet. Cell Genet.* 78: 103–104, 1997.
- [71700] 10030. Blondel, O.; Vandecasteele, G.; Gastineau, M.; Leclerc, S.; Dahmoune, Y.; Langlois, M.; Fischmeister, R.: Molecular and functional characterization of a 5-HT(4) receptor cloned from human atrium. *FEBS Lett.*



412:465–474, 1997.

- [71701] 10031.Claeysen, S.; Faye, P.; Sebben, M.; Lemaire, S.; Bockaert, J.; Dumuis, A.; Taviaux, S.: Assignment of 5–hydroxytryptamine receptor(HTR4) to human chromosome 5 bands q31–to–q33 by in situ hybridization. *Cytogenet.Cell Genet.* 78: 133–134, 1997.
- [71702] 10032.Dumuis, A.; Bouhelal, R.; Sebben, M.; Cory, R.; Bockaert, J.:A nonclassical 5–hydroxytryptamine receptor positively coupled withadenylate cyclase in the central nervous system. *Molec. Pharm.* 34:880–887, 1988.
- [71703] 10033.Eglen, R. M.; Wong, E. H. F.; Dumuis, A.; Bockaert, J.: Central5–HT4 receptors. *Trends Pharm. Sci.* 16: 391–398, 1995.
- [71704] 10034.Isomura, T.; Tamiya–Koizumi, K.; Suzuki, M.; Yoshida, S.; Taniguchi,M.; Matsuyama, M.; Ishigaki, T.; Sakuma, S.; Takahashi, M.: RFP isa DNA binding protein associated with the nuclear matrix. *NucleicAcids Res.* 20: 5305–5310, 1992.
- [71705] 10035.Szpirer, C.; Szpirer, J.; Riviere, M.; Tazi, R.; Pontarotti, P.: Mapping of the Olf89 and Rfp genes to the rat genome: comparisonwith the mouse and human and new insights into the evolution of therodent genome. *Cytogenet. Cell Genet.* 78: 137–139, 1997.

- [71706] 10036.Lim, D.-S.; Kirsch, D. G.; Canman, C. E.; Ahn, J.-H.; Ziv, Y.;Newman, L. S.; Darnell, R. B.; Shiloh, Y.; Kastan, M. B.: ATM bindsto beta-adaptin in cytoplasmic vesicles. Proc. Nat. Acad. Sci. 95:10146–10151, 1998.
- [71707] 10037.Newman, L. S.; McKeever, M. O.; Okano, H. J.; Darnell, R. B.:Beta-NAP, a cerebellar degeneration antigen, is a neuron-specificvesicle coat protein. Cell 82: 773–783, 1995.
- [71708] 10038.Simpson, F.; Bright, N. A.; West, M. A.; Newman, L. S.; Darnell,R. B.; Robinson, M. S.: A novel adaptor-related protein complex. J.Cell. Biol. 133: 749–760, 1996.
- [71709] 10039.Luo, J.; Sladek, R.; Bader, J. A.; Matthyssen, A.; Rossant, J.;Giguere, V.: Placental abnormalities in mouse embryos lacking orphannuclear receptor ERR-beta. Nature 388: 778–782, 1997.
- [71710] 10040.Adachi, O.; Kawai, T.; Takeda, K.; Matsumoto, M.; Tsutsui, H.;Sakagami, M.; Nakanishi, K.; Akira, S.: Tar-geted disruption of theMyD88 gene results in loss of IL-1- and IL-18-mediated function. Immunity 143–150,1998.
- [71711] 10041.Bonnert, T. P.; Garka, K. E.; Parnet, P.; Sonoda, G.; Testa, J.R.; Sims, J. E.: The cloning and characterization of human MyD88:a member of an IL-1 receptor related fam-

ily. FEBS Lett. 402: 81–84,1997.

- [71712] 10042.Hardiman, G.; Jenkins, N. A.; Copeland, N. G.; Gilbert, D. J.;Garcia, D. K.; Naylor, S. L.; Kastelein, R. A.; Bazan, J. F.: Geneticstructure and chromosomal mapping of MyD88. Genomics 45: 332–339,1997.
- [71713] 10043.Hayashi, F.; Smith, K. D.; Ozinsky, A.,; Hawn, T. R.; Yi, E. C.;Goodlett, D. R.; Eng, J. K.; Akira, S.; Underhill, D. M.; Aderem,A.: The innate immune response to bacterial flagellin is mediatedby Toll–like receptor 5. Nature 410: 1099–1103, 2001.
- [71714] 10044.Kawai, T.; Adachi, O.; Ogawa, T.; Takeda, K.; Akira, S.: Unresponsivenessof MyD88–deficient mice to endo–toxin. Immunity 11: 115–122, 1999.
- [71715] 10045.Lord, K. A.; Hoffman–Liebermann, B.; Liebermann, D. A.: Complexityof the immediate early response of myeloid cells to terminal differentiationand growth arrest includes ICAM–1, Jun–B and histone variants. Oncogene 5:387–396, 1990.
- [71716] 10046.Medzhitov, R.; Preston–Hurlburt, P.; Kopp, E.; Staden, A.; Chen,C.; Ghosh, S.; Janeway, C. A., Jr.: MyD88 is an adaptor protein inthe hToll/Il–1 receptor family signaling pathways. Molec. Cell 2:253–258, 1998.
- [71717] 10047.Ge, K.; Guermah, M.; Yuan, C.–X.; Ito, M.; Wallberg,

A. E.; Spiegelman, B. M.; Roeder, R. G.: Transcription coactivator TRAP220 is required for PPAR- $\gamma$ -2-stimulated adipogenesis. *Nature* 417: 563–567, 2002.

[71718] 10048. Greene, M. E.; Blumberg, B.; McBride, O. W.; Yi, H. F.; Kronquist, K.; Kwan, K.; Hsieh, L.; Greene, G.; Nimer, S. D.: Isolation of the human peroxisome proliferator activated receptor gamma cDNA: expression in hematopoietic cells and chromosomal mapping. *Gene Expr.* 4: 281–299, 1995.

[71719] 10049. Hara, M.; Alcoser, S. Y.; Qadir, A.; Beiswenger, K. K.; Cox, N. J.; Ehrmann, D. A.: Insulin resistance is attenuated in women with polycystic ovary syndrome with the Pro12Ala polymorphism in the PPAR- $\gamma$  gene. *J. Clin. Endocr. Metab.* 87: 772–775, 2002.

[71720] 10050. Harris, S. G.; Phipps, R. P.: Prostaglandin D<sub>2</sub>, its metabolite 15-d-PGJ<sub>2</sub>, and peroxisome proliferator activated receptor- $\gamma$  agonists induce apoptosis in transformed, but not normal, human T lineage cells. *Immunology* 105: 23–34, 2002.

[71721] 10051. Hasstedt, S. J.; Ren, Q.-F.; Teng, K.; Elbein, S. C.: Effect of the peroxisome proliferator-activated receptor- $\gamma$ -2 Pro12Ala variant on obesity, glucose homeosta-

sis, and blood pressure in members of familial type 2 diabetic kindreds. *J. Clin. Endocr. Metab.* 86:536–541, 2001.

[71722] 10052. Dabovic, B.; Chen, Y.; Colarossi, C.; Obata, H.; Zambuto, L.; Perle, M. A.; Rifkin, D. B.: Bone abnormalities in latent TGF- $\beta$  binding protein (Ltbp)-3-null mice indicate a role for Ltbp-3 in modulating TGF- $\beta$  bioavailability. *J. Cell Biol.* 156: 227–232, 2002.

[71723] 10053. Li, X.; Yin, W.; Perez-Jurado, L.; Bonadio, J.; Francke, U.: Mapping of human and murine genes for latent TGF- $\beta$  binding protein-2 (LTBP2). *Mammalian Genome* 6: 42–45, 1995.

[71724] 10054. Sawicki, M.; Arnold, E.; Ebrahimi, S.; Duell, T.; Jin, S.; Wood, T.; Chakrabarti, R.; Peters, J.; Wan, Y.; Samara, G.; Weier, H.-U. G.; Udar, N.; Passaro, E., Jr.; Srivatsan, E. S.: A transcript map encompassing the multiple endocrine neoplasia type-1 (MEN1) locus on chromosome 11q13. *Genomics* 42: 405–412, 1997.

[71725] 10055. Yin, W.; Smiley, E.; Germiller, J.; Mechan, R. P.; Florer, J. B.; Wenstrup, R. J.; Bonadio, J.: Isolation of a novel latent transforming growth factor- $\beta$  binding protein gene (LTBP-3). *J. Biol. Chem.* 270:10147–10160, 1995.

[71726] 10056. Moren, A.; Olofsson, A.; Stenman, G.; Sahlin, P.; Kanzaki, T.; Claesson-Welsh, L.; ten Dijke, P.; Miyazono,

K.; Heldin, C.-H.: Identification and characterization of LTBP-2, a novel latent transforming growth factor-beta-binding protein. *J. Biol. Chem.* 269: 32469-32478, 1994.

- [71727] 10057. Scherer, S. W.; Duvoisin, R. M.; Kuhn, R.; Heng, H. H. Q.; Belloni, E.; Tsui, L.-C.: Localization of two metabotropic glutamate receptor genes, GRM3 and GRM8, to human chromosome 7q. *Genomics* 31: 230-233, 1996.
- [71728] 10058. Scherer, S. W.; Soder, S.; Duvoisin, R. M.; Huizenga, J. J.; Tsui, L.-C.: The human metabotropic glutamate receptor 8 (GRM8) gene: a disproportionately large gene located at 7q31.3-q32.1. *Genomics* 44: 232-236, 1997.
- [71729] 10059. Wu, S.; Wright, R. A.; Rockey, P. K.; Burgett, S. G.; Arnold, J. S.; Rosteck, P. R., Jr.; Johnson, B. G.; Schoepp, D. D.; Belagaje, R. M.: Group III human metabotropic glutamate receptors 4, 7 and 8: molecular cloning, functional expression, and comparison of pharmacological properties in RGT cells. *Molec. Brain Res.* 53: 88-97, 1998.
- [71730] 10060. Meckelein, B.; Rohan de Silva, H. A.; Roses, A. D.; Rao, P. N.; Pettenati, M. J.; Xu, P.-T.; Hodge, R.; Gluckman, M. J.; Abraham, C. R.: Human endopeptidase (THOP1) is localized on chromosome 19 within the linkage region for the late-onset Alzheimer disease AD2 locus. *Genomics*

31: 246–249, 1996.

- [71731] 10061.Papastoitsis, G.; Siman, R.; Scott, R.; Abraham, C. R.: Identification of a metalloprotease from Alzheimer's disease brain able to degrade the beta-amyloid precursor protein and generate amyloidogenic fragments. *Biochemistry* 33:192–199, 1994.
- [71732] 10062.Torres, M. P.; Prange, C.; Lennon, G.: Human endopeptidase 24.15 (THOP1) is localized on chromosome 19p13.3 and is excluded from the linkage region for late-onset Alzheimer disease. *Genomics* 53: 239–240, 1998.
- [71733] 10063.Bram, R. J.; Crabtree, G. R.: Calcium signalling in T cells stimulated by a cyclophilin B-binding protein. *Nature* 371: 355–358, 1994.
- [71734] 10064.Bram, R. J.; Valentine, V.; Shapiro, D. N.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.: The gene for calcium-modulating cyclophilin ligand (CAMLG) is located on human chromosome 5q23 and a syntenic region of mouse chromosome 13. *Genomics* 31: 257–260, 1996.
- [71735] 10065.Bussemakers, M. J. G.; van Bokhoven, A.; Voller, M.; Smit, F. P.; Schalken, J. A.: The genes for the calcium-dependent cell adhesion molecules P- and E-cadherin are tandemly arranged in the human genome. *Biochem. Biophys. Res. Commun.* 203: 1291–1294, 1994.

- [71736] 10066.Carmeliet, P.; Lampugnani, M.-G.; Moons, L.; Breviario, F.; Compernelle,V.; Bono, F.; Balconi, G.; Spagnuolo, R.; Oosthuyse, B.; Dewerchin,M.; Zanetti, A.; Angelillo, A.; and 11 others: Targeted deficiencyof cytosolic truncation of the VE-cadherin gene in mice impairs VEGF-mediatedendothelial survival and angiogenesis. *Cell* 98: 147–157, 1999.
- [71737] 10067.Huber, P.; Dalmon, J.; Engiles, J.; Breviario, F.; Gory, S.; Siracusa,L. D.; Buchberg, A. M.; Dejana, E.: Genomic structure and chromosomal mapping of the mouse VE-cadherin gene (*Cdh5*). *Genomics* 32: 21–28,1996.
- [71738] 10068.Salomon, D.; Ayalon, O.; Patel-King, R.; Hynes, R. O.; Geiger,B.: Extrajunctional distribution of N-cadherin in cultured humanendothelial cells. *J. Cell Sci.* 102: 7–17, 1992.
- [71739] 10069.Carmeliet, P.; Moons, L.; Luttun, A.; Vincenti, V.; Compernelle,V.; De Mol, M.; Wu, Y.; Bono, F.; Devy, L.; Beck, H.; Scholz, D.;Acker, T.; and 17 others: Synergism between vascular endothelialgrowth factor and placental growth factor contributes to angiogenesisand plasma extravasation in pathological conditions. *Nature Med.* 7:575–583, 2001.
- [71740] 10070.Luttun, A.; Tjwa, M.; Moons, L.; Wu, Y.; Angelillo-



Scherrer, A.; Liao, F.; Nagy, J. A.; Hooper, A.; Priller, J.; De Klerck, B.; Compernelle, V.; Daci, E.; and 10 others: Revascularization of ischemic tissues by PlGF treatment, and inhibition of tumor angiogenesis, arthritis and atherosclerosis by anti-Flt1. *Nature Med.* 8: 831–840, 2002.

[71741] 10071. Maglione, D.; Guerriero, V.; Viglietto, G.; Delli-Bovi, P.; Persico, M. G.: Isolation of a human placenta cDNA coding for a protein related to the vascular permeability factor. *Proc. Nat. Acad. Sci.* 88: 9267–9271, 1991.

[71742] 10072. Mattei, M.-G.; Borg, J.-P.; Rosnet, O.; Marme, D.; Birnbaum, D.: Assignment of vascular endothelial growth factor (VEGF) and placenta growth factor (PLGF) genes to human chromosome 6p12–p21 and 14q24–q31 regions, respectively. *Genomics* 32: 168–169, 1996.

[71743] 10073. Le Coniat, M.; Choi, D.-S.; Maroteaux, L.; Launay, J.-M.; Berger, R.: The 5-HT<sub>2B</sub> receptor gene maps to 2q36.3–2q37.1. *Genomics* 32: 172–173, 1996.

[71744] 10074. Nebigil, C. G.; Choi, D.-S.; Dierich, A.; Hickel, P.; Le Meur, M.; Messaddeq, N.; Launay, J.-M.; Maroteaux, L.: Serotonin 2B receptor is required for heart development. *Proc. Nat. Acad. Sci.* 97: 9508–9513, 2000.

[71745] 10075. Goshima, Y.; Nakamura, F.; Strittmatter, P.; Strittmatter, S. M.: Collapsin-induced growth cone col-

lapse mediated by an intracellular protein related to UNC-33. *Nature* 376: 509–514, 1995.

- [71746] 10076. Charrin, S.; Le Naour, F.; Oualid, M.; Billard, M.; Faure, G.; Hanash, S. M.; Boucheix, C.; Rubinstein, E.: The major CD9 and CD81 molecular partner: identification and characterization of the complexes. *J. Biol. Chem.* 276: 14329–14337, 2001.
- [71747] 10077. Nagase, T.; Kikuno, R.; Ishikawa, K.; Hirose, M.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XVI. The complete sequences of 150 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.* 7: 65–73, 2000.
- [71748] 10078. Orlicky, D. J.; Berry, R.; Sikela, J. M.: Human chromosome 1 localization of the gene for a prostaglandin F<sub>2</sub>-alpha receptor negative regulatory protein. *Hum. Genet.* 97: 655–658, 1996.
- [71749] 10079. Stipp, C. S.; Orlicky, D.; Hemler, M. E.: FPRP, a major, highly stoichiometric, highly specific CD81- and CD9-associated protein. *J. Biol. Chem.* 276: 4853–4862, 2001.
- [71750] 10080. Oliver, G.; Wehr, R.; Jenkins, N. A.; Copeland, N. G.; Cheyette, B. N. R.; Hartenstein, V.; Zipursky, S. L.; Gruss, P.: Homeobox genes and connective tissue patterning. De-

velopment 121: 693–705, 1995.

- [71751] 10081.Casellas, R.; Jankovic, M.; Meyer, G.; Gazumyan, A.; Luo, Y.; Roeder, R. G.; Nussenzweig, M. C.: OcaB is required for normal transcription and V(D)J recombination of a subset of immunoglobulin kappa genes. *Cell* 110:575–585, 2002.
- [71752] 10082.Gstaiger, M.; Knoepfel, L.; Georgiev, O.; Schaffner, W.; Hovens, C. M.: A B-cell coactivator of octamer-binding transcription factors. *Nature* 373:360–362, 1995.
- [71753] 10083.Junker, S.; Brondum-Nielsen, K.; Newell, J. W.; Matthias, P.; Tommerup, N.: Assignment of the human gene for Oct-binding factor-1 (OBF1), a B-cell-specific coactivator of octamer-binding transcription factors 1 and 2, to 11q23.1 by somatic cell hybridization and in situ hybridization. *Genomics* 33:143–145, 1996.
- [71754] 10084.Urness, L. D.; Sorensen, L. K.; Li, D. Y.: Arteriovenous malformations in mice lacking activin receptor-like kinase-1. *Nature Genet.* 26:328–331, 2000.
- [71755] 10085.Jantti, J.; Lahdenranta, J.; Olkkonen, V. M.; Soderlund, H.; Keranen, S.: SEM1, a homologue of the split hand/split foot malformation candidate gene Dss1, regulates exocytosis and pseudohyphal differentiation in yeast. *Proc. Nat. Acad. Sci.* 96: 909–914, 1999.

- [71756] 10086.Han, L. Wong, D.; Dhaka, A.; Afar, D.; White, M.; Xie, W.; Herschman,H.; Witte, O.; Colicelli, J.: Protein binding and signaling properties of RIN1 suggest a unique effector function. *Proc. Nat. Acad. Sci.* 94:4954–4959, 1997.
- [71757] 10087.Leffers, H.; Madsen, P.; Rasmussen, H. H.; Honore, B.; Andersen,A. H.; Walbum, E.; Vandekerckhove, J.; Celis, J. E.: Molecular cloning and expression of the transformation sensitive epithelial marker stratifin:a member of a protein family that has been involved in the protein kinase C signalling pathway. *J. Molec. Biol.* 231: 982–998, 1993.
- [71758] 10088.Jaspers, M.; Marynen, P.; Aly, M. S.; Cuppens, H.; Hilliker, C.;Cassiman, J.–J.: Localization of the gene encoding the alpha–2 subunit of the human VLA–2 receptor to chromosome 5q23–31. *Somat. Cell Molec.Genet.* 17: 505–511, 1991.
- [71759] 10089.Hammond, J. W.; Potter, M.; Wilcken, B.; Truscott, R.: Siblings with gamma–glutamyltransferase deficiency. *J. Inherit. Metab. Dis.* 18:82–83, 1995.
- [71760] 10090.Heisterkamp, N.; Groffen, J.: Duplication of the bcr and gamma–glutamyltranspeptidase genes. *Nucleic Acids Res.* 16: 8045–8056, 1988.
- [71761] 10091.Laperche, Y.; Bulle, F.; Aissani, T.; Chobert, M. N.; Aggerbeck,M.; Hanoune, J.; Guellaen, G.: Molecular

cloning and nucleotide sequence of rat kidney gamma-glutamyl transpeptidase cDNA. *Proc. Nat. Acad. Sci.* 83: 937–941, 1986.

[71762] 10092. O'Daley, S.: An abnormal sulphhydryl compound in urine. (Abstract) *Irish J. Med. Sci.* 7: 578–579, 1968.

[71763] 10093. Pawlak, A.; Lahuna, O.; Bulle, F.; Suzuki, A.; Ferry, N.; Siegrist, S.; Chikhi, N.; Chobert, M. N.; Guellaen, G.; Laperche, Y.: Gamma-glutamyl transpeptidase: a single copy gene in the rat and a multigene family in the human genome. *J. Biol. Chem.* 263: 9913–9916, 1988.

[71764] 10094. Rouleau, G. A.; Bazanowski, A.; Cohen, E. H.; Guellaen, G.; Gusella, J. F.: Gamma-glutamyl transferase locus (GGT) displays a PvuII polymorphism. *Nucleic Acids Res.* 16: 11848 only, 1988.

[71765] 10095. Sakamuro, D.; Yamazoe, M.; Matsuda, Y.; Kangawa, K.; Taniguchi, N.; Matsuo, H.; Yoshikawa, H.; Ogasawara, N.: The primary structure of human gamma-glutamyl transpeptidase. *Gene* 73: 1–9, 1988.

[71766] 10096. Schulman, J. D.; Goodman, S. I.; Mace, J. W.; Patrick, A. D.; Tietze, F.; Butler, E. J.: Glutathionuria: inborn error of metabolism due to tissue deficiency of gamma-glutamyl transpeptidase. *Biochem. Biophys. Res. Commun.* 65: 68–74, 1975.

- [71767] 10097. Annabi, B.; Hiraiwa, H.; Mansfield, B. C.; Lei, K.-J.; Ubagai, T.; Polymeropoulos, M. H.; Moses, S. W.; Parvari, R.; HersHKovitz, E.; Mandel, H.; Fryman, M.; Chou, J. Y.: The gene for glycogen-storage disease type 1b maps to chromosome 11q23. *Am. J. Hum. Genet.* 62:400–405, 1998.
- [71768] 10098. Jaeken, J.; Goemans, N.; Fryns, J.-P.; Francois, I.; de Zegher, F.: Association of hyperprolinaemia type I and heparin cofactor II deficiency with CATCH 22 syndrome: evidence for a contiguous gene syndrome locating the proline oxidase gene. *J. Inherit. Metab. Dis.* 19:275–277, 1996.
- [71769] 10099. Harvey, C. B.; Hollox, E. J.; Poulter, M.; Wang, Y.; Rossi, M.; Auricchio, S.; Iqbal, T. H.; Cooper, B. T.; Barton, R.; Sarner, M.; Korpela, R.; Swallow, D. M.: Lactase haplotype frequencies in Caucasians: association with the lactase persistence/non-persistence polymorphism. *Ann. Hum. Genet.* 62: 215–223, 1998.
- [71770] 10100. Harvey, C. B.; Pratt, W. S.; Islam, I.; Whitehouse, D. B.; Swallow, D. M.: DNA polymorphisms in the lactase gene: linkage disequilibrium across the 70-kb region. *Eur. J. Hum. Genet.* 3: 27–41, 1995.
- [71771] 10101. Hollox, E. J.; Poulter, M.; Zvarik, M.; Ferak, V.; Krause, A.; Jenkins, T.; Saha, N.; Kozlov, A. I.; Swallow, D.

M.: Lactase haplotypediversity in the Old World. *Am. J. Hum. Genet.* 68: 160–172, 2001.

- [71772] 10102.Murano, S.; Thweatt, R.; Reis, R. J. S.; Jones, R. A.; Moerman,E. J.; Goldstein, S.: Diverse gene sequences are overexpressed inWerner syndrome fibroblasts undergoing premature replicative senescence. *Molec.Cell. Biol.* 11: 3905–3914, 1991.
- [71773] 10103.Li, S.; Strelow, A.; Fontana, E. J.; Wesche, H.: IRAK–4: a novelmember of the IRAK family with the properties of an IRAK–kinase. *Proc.Nat. Acad. Sci.* 99: 5567–5572, 2002.
- [71774] 10104.Scanlan, M. J.; Gordon, J. D.; Williamson, B.; Stock–ert, E.; Bander,N. H.; Jongeneel, V.; Gure, A. O.; Jager, D.; Jager, E.; Knuth, A.;Chen, Y.–T.; Old, L. J.: Antigens recog–nized by autologous antibodyin patients with renal–cell carcinoma. *Int. J. Cancer* 83: 456–464,1999.
- [71775] 10105.Scott, A. F.: Personal Communication. Baltimore, Md. 4/25/2002.
- [71776] 10106.Suzuki, N.; Suzuki, S.; Duncan, G. S.; Millar, D. G.; Wada, T.;Mirtsos, C.; Takada, H.; Wakeham, A.; Itie, A.; Li, S.; Penninger,J. M.; Wesche, H.; Ohashi, P. S.; Mak, T. W.; Yeh, W.–C.: Severeimpairment of interleukin–1 and Toll–like receptor signalling in micelacking IRAK–4. *Nature*

416: 750–754, 2002.

[71777] 10107. Pennarun, G.; Chapelin, C.; Escudier, E.; Bridoux, A.-M.; Dastot, F.; Cacheux, V.; Goossens, M.; Amselem, S.; Duriez, B.: The human dynein intermediate chain 2 gene (DNAI2): cloning, mapping, expression pattern, and evaluation as a candidate for primary ciliary dyskinesia.

Hum. Genet. 107: 642–649, 2000.

[71778] 10108. Amerik, A. Y.; Petukhova, G. V.; Grigorenko, V. G.; Lykov, I. P.; Yarovoi, S. V.; Lipkin, V. M.; Gorbalenya, A. E.: Cloning and sequence analysis of cDNA for a human homolog of eubacterial ATP-dependent Lon proteases. FEBS Lett. 340: 25–28, 1994.

[71779] 10109. Wang, N.; Gottesman, S.; Willingham, M. C.; Gottesman, M. M.; Maurizi, M. R.: A human mitochondrial ATP-dependent protease that is highly homologous to bacterial Lon protease. Proc. Nat. Acad. Sci. 90: 11247–11251, 1993.

[71780] 10110. Popovici, C.; Mattei, M.-G.; Rattner, J. B.; Birnbaum, D.; Pebusque, M.-J.: Assignment of the centrosomal protein 110 gene (Cep110) to mouse chromosome bands 2B–C1 by in situ hybridization. Cytogenet. Cell Genet. 89: 216–217, 2000.

[71781] 10111. Castagnola, P.; Gennari, M.; Morello, R.; Tonachini,



L.; Marin,O.; Gaggero, A.; Cancedda, R.: Cartilage associated protein (CASP)is a novel developmentally regulated chick embryo protein. *J. CellSci.* 110: 1351–1359, 1997.

[71782] 10112.Morello, R.; Tonachini, L.; Monticone, M.; Viggiano, L.; Rocchi,M.; Cancedda, R.; Castagnola, P.: cDNA cloning, characterizationand chromosome mapping of Crtap encoding the mouse cartilage associatedprotein. *Matrix Biol.* 18: 319–324, 1999.

[71783] 10113.Tonachini, L.; Morello, R.; Monticone, M.; Skaug, J.; Scherer,S. W.; Cancedda, R.; Castagnola, P.: cDNA cloning, characterizationand chromosome mapping of the gene encoding human cartilage associatedprotein (CRTAP). *Cytogenet. Cell Genet.* 87: 191–194, 1999.

[71784] 10114.Shibanuma, M.; Mashimo, J.; Mita, A.; Kuroki, T.; Nose, K.: Cloningfrom a mouse osteoblastic cell line of a set of transforming–growth–factor–beta–1–regulated genes, one of which seems to encode a follistatin–relatedpolypeptide. *Europ. J. Biochem.* 217: 13–19, 1993.

[71785] 10115.Tanaka, M.; Ozaki, S.; Osakada, F.; Mori, K.; Okubo, M.; Nakao,K.: Cloning of follistatin–related protein as a novel autoantigenin systemic rheumatic diseases. *Int. Immun.* 10: 1305–1314, 1998.

[71786] 10116.Zwijssen, A.; Blockx, H.; van Arnhem, W.; Willems, J.;

Fransen,L.; Devos, K.; Raymackers, J.; van de Voorde, A.; Slegers, H.: Characterization of a rat C6 glioma–secreted follistatin–related protein (FRP): cloning and sequence of the human homologue. *Europ. J. Biochem.* 225: 937–946,1994.

[71787] 10117.Fang, M.; Jaffrey, S. R.; Sawa, A.; Ye, K.; Luo, X.; Snyder, S.H.: Dexas1: a G protein specifically coupled to neuronal nitric oxide synthase via CAPON. *Neuron* 28: 183–193, 2000.

[71788] 10118.Kemppainen, R. J.; Behrend, E. N.: Dexamethasone rapidly induces a novel Ras superfamily member–related gene in AtT–20 cells. *J. Biol.Chem.* 273: 3129–3131, 1998.

[71789] 10119.Tu, Y.; Wu, C.: Cloning, expression and characterization of a novel human Ras–related protein that is regulated by glucocorticoid hormone. *Biochim. Biophys. Acta* 1489: 452–456, 1999.

[71790] 10120.Jaffrey, S. R.; Snowman, A. M.; Eliasson, M. J. L.; Cohen, N. A.; Snyder, S. H.: CAPON: a protein associated with neuronal nitric oxide synthase that regulates its interactions with PSD95. *Neuron* 115–124,1998.

[71791] 10121.Thuresson, A.–C.; Astrom, J.; Astrom, A.; Gronvik, K.–O.; Virtanen,A.: Multiple forms of poly(A) polymerases

in human cells. *Proc.Nat. Acad. Sci.* 91: 979–983, 1994.

- [71792] 10122.Yamauchi, T.; Sugimoto, J.; Hatakeyama, T.; Asakawa, S.; Shimizu,N.; Isobe, M.: Assignment of the human poly(A) polymerase (PAP) gene to chromosome 14q32.1–q32.2 and isolation of a polymorphic CA repeat-sequence. *J. Hum. Genet.* 44: 253–255, 1999.
- [71793] 10123.Boles, K. S.; Nakajima, H.; Colonna, M.; Chuang, S. S.; Stepp,S. E.; Bennett, M.; Kumar, V.; Mathew, P. A.: Molecular characterization of a novel human natural killer cell receptor homologous to mouse 2B4. *Tissue Antigens* 54: 27–34, 1999.
- [71794] 10124.Fort, M. M.; Cheung, J.; Yen, D.; Li, J.; Zurawski, S. M.; Lo,S.; Menon, S.; Clifford, T.; Hunte, B.; Lesley, R.; Muchamuel, T.;Hurst, S. D.; Zurawski, G.; Leach, M. W.; Gorman, D. M.; Rennick,D. M.: IL-25 induces IL-4, IL-5, and IL-13 and Th2-associated pathologies in vivo. *Immunity* 15: 985–995, 2001.
- [71795] 10125.Scott, A. F.: Personal Communication. Baltimore, Md. 2/16/2001.
- [71796] 10126.Mori, T.; Fukuda, Y.; Kuroda, H.; Matsumura, T.; Ota, S.; Sugimoto,T.; Nakamura, Y.; Inazawa, J.: Cloning and characterization of a novel Rab-family gene, Rab36, within the region at 22q11.2 that is homozygously deleted

in malignant rhabdoid tumors. Biochem. Biophys. Res. Commun. 254: 594–600, 1999.

[71797] 10127. Zhou, J.-Y.; Fogelgren, B.; Wang, Z.; Roe, B. A.; Biegel, J. A.: Isolation of genes from the rhabdoid tumor deletion region in chromosome band 22q11.2. Gene 241: 133–141, 2000.

[71798] 10128. Sood, R.; Makalowska, I.; Carpten, J. D.; Robbins, C. M.; Stephan, D. A.; Connors, T. D.; Morgenbesser, S. D.; Su, K.; Pinkett, H. W.; Graham, C. L.; Quesenberry, M. I.; Baxevanis, A. D.; Klinger, K. W.; Trent, J. M.; Bonner, T. I.: The human RGL (RalGDS-like) gene: cloning, expression analysis and genomic organization. Biochim. Biophys. Acta 1491: 285–288, 2000.

[71799] 10129. Acquati, F.; Accarino, M.; Nucci, C.; Fumagalli, P.; Jovine, L.; Ottolenghi, S.; Taramelli, R.: The gene encoding DRAP (BACE2), a glycosylated transmembrane protein of the aspartic protease family, maps to the Down syndrome critical region. FEBS Lett. 468: 59–64, 2000.

[71800] 10130. Bennett, B. D.; Babu-Khan, S.; Loeloff, R.; Louis, J.-C.; Curran, E.; Citron, M.; Vassar, R.: Expression analysis of BACE2 in brain and peripheral tissues. J. Biol. Chem. 275: 20647–20651, 2000.

[71801] 10131. Saunders, A. J.; Kim, T.-W.; Tanzi, R. E.: BACE maps

to chromosome 11 and a BACE homolog, BACE2, reside in the obligate Down syndrome region of chromosome 21. Science 286: 1255A only, 1999.

- [71802] 10132. Solans, A.; Estivill, X.; de la Luna, S.: A new aspartyl protease on 21q22.3, BACE2, is highly similar to Alzheimer's amyloid precursor protein beta-secretase. Cytogenet. Cell Genet. 89: 177–184, 2000.
- [71803] 10133. Xin, H.; Stephens, J. C.; Duan, X.; Harrowe, G.; Kim, E.; Grieshammer, U.; Kingsley, C.; Giese, K.: Identification of a novel aspartic-like protease differentially expressed in human breast cancer cell lines. Biochim. Biophys. Acta 1501: 125–137, 2000.
- [71804] 10134. Kuang, W. W.; Thompson, D. A.; Hoch, R. V.; Weigel, R. J.: Differential screening and suppression subtractive hybridization identified genes differentially expressed in an estrogen receptor-positive breast carcinoma cell line. Nucleic Acids Res. 26: 1116–1123, 1998.
- [71805] 10135. Adamec, J.; Rusnak, F.; Owen, W. G.; Naylor, S.; Benson, L. M.; Gacy, A. M.; Isaya, G.: Iron-dependent self-assembly of recombinant yeast frataxin: implications for Friedreich ataxia. Am. J. Hum. Genet. 67: 549–562, 2000.
- [71806] 10136. Campuzano, V.; Montermini, L.; Lutz, Y.; Cova, L.; Hindelang, C.; Jiralerspong, S.; Trottier, Y.; Kish, S. J.;

Faucheux, B.; Trouillas, P.; Authier, F. J.; Durr, A.; Mandel, J.-L.; Vescovi, A.; Pandolfo, M.; Koenig, M.: Frataxin is reduced in Friedreich ataxia patients and is associated with mitochondrial membranes. *Hum. Molec. Genet.* 6:1771–1780, 1997.

[71807] 10137. Cavadini, P.; O'Neill, H. A.; Benada, O.; Isaya, G.: Assembly and iron-binding properties of human frataxin, the protein deficient in Friedreich ataxia. *Hum. Molec. Genet.* 11: 217–227, 2002.

[71808] 10138. Chamberlain, S.; Pook, M.; Carvajal, J.; Doudney, K.; Hillermann, R.: Frataxin fracas. (Letter) *Nature Genet.* 15: 337–338, 1997.

[71809] 10139. Cossee, M.; Campuzano, V.; Koutnikova, H.; Fischbeck, K.; Mandel, J.-L.; Koenig, M.; Bidichandani, S. I.; Patel, P. I.; Molte, M. D.; Canizares, J.; De Frutos, R.; Piane, L.; Cavalcanti, F.; Monticelli, A.; Coccozza, S.; Monttermini, L.; Pandolfo, M.: Frataxin fracas. (Letter) *Nature Genet.* 15: 337–338, 1997.

[71810] 10140. Bartles, J. R.; Wierda, A.; Zheng, L.: Identification and characterization of espin, an actin-binding protein localized to the F-actin-rich junctional plaques of Sertoli cell ectoplasmic specializations. *J. Cell Sci.* 109:1229–1239, 1996.

- [71811] 10141. Bartles, J. R.; Zheng, L.; Li, A.; Wierda, A.; Chen, B.: Smallespin: a third actin-bundling protein and potential forked protein ortholog in brush border microvilli. *J. Cell Biol.* 143: 107–119, 1998.
- [71812] 10142. Chen, B.; Li, A.; Wang, D.; Wang, M.; Zheng, L.; Bartles, J. R.: Espin contains an additional actin-binding site in its N terminus and is a major actin-bundling protein of the Sertoli cell-spermatid ectoplasmic specialization junctional plaque. *Molec. Biol. Cell* 10:4327–4339, 1999.
- [71813] 10143. Zheng, L.; Sekerkova, G.; Vranich, K.; Tilney, L. G.; Mugnaini, E.; Bartles, J. R.: The deaf jerker mouse has a mutation in the gene encoding the espin actin-bundling proteins of hair cell stereocilia and lacks espins. *Cell* 102: 377–385, 2000.
- [71814] 10144. Illarioshkin, S. N.; Ivanova-Smolenskaya, I. A.; Tanaka, H.; Poleshchuk, V. V.; Markova, E. D.; Tsuji, S.: Refined genetic location of the chromosome 2p-linked progressive muscular dystrophy gene. *Genomics* 42:345–348, 1997.
- [71815] 10145. Illarioshkin, S. N.; Ivanova-Smolenskaya, I. A.; Tanaka, H.; Vereshchagin, N. V.; Markova, E. D.; Poleshchuk, V. V.; Lozhnikova, S. M.; Sukhorukov, V. S.; Limborska, S. A.; Slominsky, P. A.; Bulayeva, K. B.; Tsuji, S.:

Clinical and molecular analysis of a large family with threedistinct phenotypes of progressive muscular dystrophy. *Brain* 119:1895–1909, 1996.

[71816] 10146.McNally, E. M.; Ly, C. T.; Rosenmann, H.; Rosenbaum, S. M.; Jiang,W.; Anderson, L. V. B.; Soffer, D.; Argov, Z.: Splicing mutationin dysferlin produces limb–girdle muscular dystrophy with inflammation. *Am.J. Med. Genet.* 91: 305–312, 2000.

[71817] 10147.Weiler, T.; Greenberg, C. R.; Nylén, E.; Halliday, W.; Morgan,K.; Eggertson, D.; Wrogemann, K.: Limb–girdle muscular dystrophyand Miyoshi myopathy in an aboriginal Canadian kindred map to LGMD2Band segregate with the same haplotype. *Am. J. Hum. Genet.* 59: 872–878,1996.

[71818] 10148.Crosbie, R. H.; Lim, L. E.; Moore, S. A.; Hirano, M.; Hays, A.P.; Maybaum, S. W.; Collin, H.; Dovico, S. A.; Stolle, C. A.; Fardeau,M.; Tome, F. M. S.; Campbell, K. P.: Molecular and genetic characterizationof sarcospan: insights into sarcoglycan–sarcospan interactions. *Hum.Molec. Genet.* 9: 2019–2027, 2000.

[71819] 10149.Augustin, I.; Betz, A.; Herrmann, C.; Jo, T.; Brose, N.: Differentialexpression of two novel Munc13 proteins in rat brain. *Biochem. J.* 337:363–371, 1999.

[71820] 10150.Augustin, I.; Rosenmund, C.; Sudhof, T. C.; Brose,



N.: Munc13-1 is essential for fusion competence of glutamatergic synaptic vesicles. *Nature* 400:457–461, 1999.

- [71821] 10151. Rhee, J.-S.; Betz, A.; Pyott, S.; Reim, K.; Varoqueaux, F.; Augustin, I.; Hesse, D.; Sudhof, T. C.; Takahashi, M.; Rosenmund, C.; Brose, N.: Beta phorbol ester- and diacylglycerol-induced augmentation of transmitter release is mediated by Munc13s and not by PKCs. *Cell* 108:121–133, 2002.
- [71822] 10152. Rosenmund, C.; Sigler, A.; Augustin, I.; Reim, K.; Brose, N.; Rhee, J.-S.: Differential control of vesicle priming and short-term plasticity by Munc13 isoforms. *Neuron* 33: 411–424, 2002.
- [71823] 10153. Song, Y.; Ailenberg, M.; Silverman, M.: Cloning of a novel gene in the human kidney homologous to rat munc13s: its potential role in diabetic nephropathy. *Kidney Int.* 53: 1689–1695, 1998.
- [71824] 10154. Ji, Y.; Walkowicz, M. J.; Buiting, K.; Johnson, D. K.; Tarvin, R. E.; Rinchik, E. M.; Horsthemke, B.; Stubbs, L.; Nicholls, R. D.: The ancestral gene for transcribed, low-copy repeats in the Prader-Willi/Angelman region encodes a large protein implicated in protein trafficking, which is deficient in mice with neuromuscular and spermiogenic abnormalities. *Hum. Molec. Genet.* 8: 533–542, 1999.

- [71825] 10155.Manda, R.; Kohno, T.; Matsuno, Y.; Takenoshita, S.; Kuwano, H.;Yokota, J.: Identification of genes (SPON2 and C20orf2) differentiallyexpressed between cancerous and noncancerous lung cells by mRNA differentialdisplay. Genomics 61: 5–14, 1999.
- [71826] 10156.Zhang, Y.; Heidebrecht, H.–J.; Rott, A.; Schlegelberger, B.; Parwaresch,R.: Assignment of human proliferation associated p100 gene (C20orf1)to human chromosome band 20q11.2 by in situ hybridization. Cytogenet.Cell Genet. 84: 182–183, 1999.
- [71827] 10157.Parker, N. J.; Begley, C. G.; Smith, P. J.; Fox, R. M.: Molecularcloning of a novel human gene (D11S4896E) at chromosomal region 11p15.5. Genomics 37:253–256, 1996.
- [71828] 10158.Sabbioni, S.; Veronese, A.; Trubia, M.; Taramelli, R.; Barbanti–Brodano,G.; Croce, C. M.; Negrini, M.: Exon structure and promoter identificationof STIM1 (alias GOK), a human gene causing growth arrest of the humantumor cell lines G401 and RD. Cytogenet. Cell. Genet. 86: 214–218,1999.
- [71829] 10159.Boudin, H.; Doan, A.; Xia, J.; Shigemoto, R.; Huganir, R. L.; Worley,P.; Craig, A. M.: Presynaptic clustering of mGluR7a requires thePICK1 PDZ domain binding

site. *Neuron* 28: 485–497, 2000.

- [71830] 10160.Dev, K. K.; Nishimune, A.; Henley, J. M.; Nakanishi, S.: The proteinkinase C- $\alpha$  binding protein PICK1 interacts with short but not long form alternative splice variants of AMPA receptor subunits. *Neuropharmacology* 38:635–644, 1999.
- [71831] 10161.Staudinger, J.; Zhou, J.; Burgess, R.; Elledge, S. J.; Olson, E.N.: PICK1: a perinuclear binding protein and substrate for proteinkinase C isolated by the yeast two-hybrid system. *J. Cell Biol.* 128:263–271, 1995.
- [71832] 10162.Takeya, R.; Takeshige, K.; Sumimoto, H.: Interaction of the PDZ domain of human PICK1 with class I ADP-ribosylation factors. *Biochem.Biophys. Res. Commun.* 267: 149–155, 2000.
- [71833] 10163.Xia, J.; Zhang, X.; Staudinger, J.; Huganir, R. L.: Clustering of AMPA receptors by the synaptic PDZ domain-containing protein PICK1. *Neuron* 22:179–187, 1999.
- [71834] 10164.Blencowe, B. J.; Bauren, G.; Eldridge, A. G.; Issner, R.; Nickerson, J. A.; Rosonina, E.; Sharp, P. A.: The SRm160/300 splicing coactivator subunits. *RNA* 6: 111–120, 2000.
- [71835] 10165.Sawada, Y.; Miura, Y.; Umeki, K.; Tamaoki, T.; Fujinaga, K.; Ohtaki, S.: Cloning and characterization of a

novel RNA-binding protein SRL300 with RS domains.

Biochim. Biophys. Acta 1492: 191–195, 2000.

- [71836] 10166. Hirose, K.; Morita, M.; Ema, M.; Mimura, J.; Hamada, H.; Fujii, H.; Saijo, Y.; Gotoh, O.; Sogawa, K.; Fujii-Kuriyama, Y.: cDNA cloning and tissue-specific expression of a novel basic helix-loop-helix/PAS factor (Arnt2) with close sequence similarity to the aryl hydrocarbon receptor nuclear translocator (Arnt). *Molec. Cell. Biol.* 16: 1706–1713, 1996.
- [71837] 10167. Keith, B.; Adelman, D. M.; Simon, M. C.: Targeted mutation of the murine aryl hydrocarbon receptor nuclear translocator 2 (Arnt2) gene reveals partial redundancy with Arnt. *Proc. Nat. Acad. Sci.* 98:6692–6697, 2001.
- [71838] 10168. Michaud, J. L.; DeRossi, C.; May, N. R.; Holdener, B. C.; Fan, C.-M.: ARNT2 acts as the dimerization partner of SIM1 for the development of the hypothalamus. *Mech. Dev.* 90: 253–261, 2000.
- [71839] 10169. Wende, H.; Volz, A.; Ziegler, A.: Extensive gene duplications and a large inversion characterize the human leukocyte receptor cluster. *Immunogenetics* 51:703–713, 2000.
- [71840] 10170. Fukunaga-Johnson, N.; Lee, S. W.; Liebert, M.; Grossman, H. B.: Molecular analysis of a gene, BB1, over-

expressed in bladder and breast carcinoma. *Anticancer Res.* 16: 1085–1090, 1996.

- [71841] 10171. Saito, H.; Papaconstantinou, J.; Sato, H.; Goldstein, S.: Regulation of a novel gene encoding a lysyl oxidase-related protein in cellular adhesion and senescence. *J. Biol. Chem.* 272: 8157–8160, 1997.
- [71842] 10172. Hsu, S. Y.; Liang, S.-G.; Hsueh, A. J. W.: Characterization of two LGR genes homologous to gonadotropin and thyrotropin receptors with extracellular leucine-rich repeats and a G protein-coupled, seven-transmembrane region. *Molec. Endocr.* 12: 1830–1845, 1998.
- [71843] 10173. Loh, E. D.; Broussard, S. R.; Kolakowski, L. F.: Molecular characterization of a novel glycoprotein hormone G-protein-coupled receptor. *Biochem. Biophys. Res. Commun.* 282: 757–764, 2001.
- [71844] 10174. Loh, E. D.; Broussard, S. R.; Liu, Q.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Kolakowski, L. F., Jr. Chromosomal localization of GPR48, a novel glycoprotein hormone receptor like GPCR, in human and mouse with radiation hybrid and interspecific backcross mapping. *Cytogenet. Cell Genet.* 89: 2–5, 2000.
- [71845] 10175. Giffon, T.; Lepourcelet, M.; Pichon, L.; Jezequel, P.;

Bouric,P.; Carn, G.; Pontarotti, P.; Le Gall, J.-Y.; David, V.: Cloning of a human homologue of the mouse Tctex-5 gene within the MHC class I region. *Immunogenetics* 44: 331-339, 1996.

[71846] 10176. Lepourcelet, M.; Andrieux, N.; Giffon, T.; Pichon, L.; Hampe, A.; Galibert, F.; Mosser, J.: Systematic sequencing of the human HLA-A/HLA-F region: establishment of a cosmid contig and identification of a new gene cluster within 37 kb of sequence. *Genomics* 37: 316-326, 1996.

[71847] 10177. Zhang, J.; Zhang, L.; Zhao, S.; Lee, E. Y. C.: Identification and characterization of the human HCG V gene product as a novel inhibitor of protein phosphatase-1. *Biochemistry* 37: 16728-16734, 1998.

[71848] 10178. Hayakawa, A.; Matsuda, Y.; Daibata, M.; Nakamura, H.; Sano, K.: Genomic organization, tissue expression, and cellular localization of AF3p21, a fusion partner of MLL in therapy-related leukemia. *Genes Chromosomes Cancer* 30: 364-374, 2001.

[71849] 10179. Sano, K.; Hayakawa, A.; Piao, J.-H.; Kosaka, Y.; Nakamura, H.: Novel SH3 protein encoded by the AF3p21 gene is fused to the mixed lineage leukemia protein in a therapy-related leukemia with t(3;11)(p21;q23). *Blood* 95:1066-1068, 2000.

- [71850] 10180. Alagramam, K. N.; Murcia, C. L.; Kwon, H. Y.; Pawlowski, K. S.; Wright, C. G.; Woychik, R. P.: The mouse Ames waltzer hearing-loss mutant is caused by mutation of Pcdh15, a novel protocadherin gene. *Nature Genet.* 27: 99–102, 2001.
- [71851] 10181. Alagramam, K. N.; Yuan, H.; Kuehn, M. H.; Murcia, C. L.; Wayne, S.; Srisailpathy, C. R. S.; Lowry, R. B.; Knaus, R.; Van Laer, L.; Bernier, F. P.; Schwartz, S.; Lee, C.; Morton, C. C.; Mullins, R. F.; Ramesh, A.; Van Camp, G.; Hagemen, G. S.; Woychik, R. P.; Smith, R. J. H.: Mutations in the novel protocadherin PCDH15 cause Usher syndrome type 1F. *Hum. Molec. Genet.* 10: 1709–1718, 2001.
- [71852] 10182. Cao, H.; Hegele, R. A.: Identification of single-nucleotide polymorphisms in the human LPIN1 gene. *J. Hum. Genet.* 47: 370–372, 2002.
- [71853] 10183. Peterfy, M.; Phan, J.; Xu, P.; Reue, K.: Lipodystrophy in the fld mouse results from mutation of a new gene encoding a nuclear protein, lipin. *Nature Genet.* 27: 121–124, 2001.
- [71854] 10184. Reue, K.; Xu, P.; Wang, X.-P.; Slavin, B. G.: Adipose tissue deficiency, glucose intolerance, and increased atherosclerosis result from mutation in the mouse fatty liver dystrophy (fld) gene. *J. Lipid Res.* 41: 1067–1076,

2000.

- [71855] 10185.Shakhov, A. N.; Rubtsov, A. V.; Lyakhov, I. G.; Tumanov, A. V.;Nedospasov, S. A.: SPLASH (PLA(2)IID), a novel member of phospholipaseA2 family, is associated with lymphotoxin-deficiency. *Genes Immun.* 1:191–199, 2000.
- [71856] 10186.Bodine, S. C.; Latres, E.; Baumhueter, S.; Lai, V. K.–M.; Nunez,L.; Clarke, B. A.; Poueymirou, W. T.; Panaro, F. J.; Na, E.; Dharmarajan,K.; Pan, Z.–Q.; Valenzuela, D. M.; DeChiara, T. M.; Stitt, T. N.;Yancopoulos, G. D.; Glass, D. J.: Identification of ubiquitin ligasesrequired for skeletal muscle atrophy. *Science* 294: 1704–1708, 2001.
- [71857] 10187.Domanski, T. L.; Finta, C.; Halpert, J. R.; Zaphiropoulos, P. G.: cDNA cloning and initial characterization of CYP3A43, a novel humancytochrome P450. *Molec. Pharm.* 59: 386–392, 2001.
- [71858] 10188.Gellner, K.; Eiselt, R.; Hustert, E.; Arnold, H.; Koch, I.; Haberl,M.; Deglmann, C. J.; Burk, O.; Buntefuss, D.; Escher, S.; Bishop,C.; Koebe, H.–G.; Brinkmann, U.; Klenk, H.–P.; Kleine, K.; Meyer,U. A.; Wojnowski, L.: Genomic organization of the human CYP3A locus:identification of a new, inducible CYP3A gene. *Pharmacogenetics* 11:111–121, 2001.



- [71859] 10189. Westlind, A.; Malmebo, S.; Johansson, I.; Otter, C.; Andersson, T. B.; Ingelman-Sundberg, M.; Oscarson, M.: Cloning and tissue distribution of a novel human cytochrome P450 of the CYP3A subfamily, CYP3A43. *Biochem. Biophys. Res. Commun.* 281: 1349–1355, 2001.
- [71860] 10190. Holt, J. R.; Gillespie, S. K. H.; Provance, D. W., Jr.; Shah, K.; Shokat, K. M.; Corey, D. P.; Mercer, J. A.; Gillespie, P. G.: A chemical-genetic strategy implicates myosin-1c in adaptation by hair cells. *Cell* 108:371–381, 2002.
- [71861] 10191. Nakayama, T.; Yaoi, T.; Kuwajima, G.: Localization and subcellular distribution of N-copine in mouse brain. *J. Neurochem.* 72: 373–379, 1999.
- [71862] 10192. Nakayama, T.; Yaoi, Y.; Yasui, M.; Kuwajima, G.: N-copine: a novel two C2-domain-containing protein with neuronal activity-regulated expression. *FEBS Lett.* 428: 80–84, 1998.
- [71863] 10193. Runnels, L. W.; Yue, L.; Clapham, D. E.: The TRPM7 channel is inactivated by PIP(2) hydrolysis. *Nature Cell Biol.* 4: 329–336, 2002.
- [71864] 10194. Scott, A. F.: Personal Communication. Baltimore, Md. 3/8/2001.
- [71865] 10195. Chen, D.; Guo, J.; Miki, T.; Tachibana, M.; Gahl, W. A.: Molecular cloning of two novel rab genes from human

melanocytes. *Gene* 174:129–134, 1996.

- [71866] 10196. Centner, T.; Yano, J.; Kimura, E.; McElhinny, A. S.; Pelin, K.; Witt, C. C.; Bang, M.-L.; Trombitas, K.; Granzier, H.; Gregorio, C.C.; Sorimachi, H.; Labeit, S.: Identification of muscle specific ring finger proteins as potential regulators of the titin kinase domain. *J. Molec. Biol.* 306: 717–726, 2001.
- [71867] 10197. Scott, A. F.: Personal Communication. Baltimore, Md. 7/20/2001.
- [71868] 10198. Kirikoshi, H.; Koike, J.; Sagara, N.; Saitoh, T.; Tokuhara, M.; Tanaka, K.; Sekihara, H.; Hirai, M.; Katoh, M.: Molecular cloning and genomic structure of human Frizzled-3 at chromosome 8p21. *Biochem. Biophys. Res. Commun.* 271: 8–14, 2000.
- [71869] 10199. Sala, C. F.; Formenti, E.; Terstappen, G. C.; Caricasole, A.: Identification, gene structure, and expression of human frizzled-3 (FZD3). *Biochem. Biophys. Res. Commun.* 273: 27–34, 2000.
- [71870] 10200. Eggenschwiler, J. T.; Espinoza, E.; Anderson, K. V.: Rab23 is an essential negative regulator of the mouse Sonic hedgehog signalling pathway. *Nature* 412: 194–198, 2001.
- [71871] 10201. Zhang, Q.-H.; Ye, M.; Wu, X.-Y.; Ren, S.-X.; Zhao,

M.; Zhao, C.-J.; Fu, G.; Shen, Y.; Fan, H.-Y.; Lu, G.; Zhong, M.; Xu, X.-R.; and 9 others: Cloning and functional analysis of cDNAs with open reading frames for 300 previously undefined genes expressed in CD34+ hematopoietic stem/progenitor cells. *Genome Res.* 10: 1546–1560, 2000.

[71872] 10202. Saitoh, T.; Hirai, M.; Katoh, M.: Molecular cloning and characterization of human Frizzled-8 gene on chromosome 10p11.2. *Int. Oncol.* 18:991–996, 2001.

[71873] 10203. Wittenberger, T.; Schaller, H. C.; Hellebrand, S.: An expressed sequence tag (EST) data mining strategy succeeding in the discovery of new G-protein coupled receptors. *J. Molec. Biol.* 307: 799–813, 2001.

[71874] 10204. Communi, D.; Gonzalez, N. S.; Detheux, M.; Brezillon, S.; Lannoy, V.; Parmentier, M.; Boeynaems, J.-M.: Identification of a novel human ADP receptor coupled to G(i). *J. Biol. Chem.* 276: 41479–41485, 2001.

[71875] 10205. Chen, H.; Ross, C. A.; Wang, N.; Huo, Y.; MacKinnon, D. F.; Potash, J. B.; Simpson, S. G.; McMahon, F. J.; DePaulo, J. R., Jr.; McInnis, M. G.: NEDD4L on human chromosome 18q21 has multiple forms of transcripts and is a homologue of the mouse Nedd4-2 gene. *Europ. J. Hum. Genet.* 9:922–930, 2001.

[71876] 10206. Erdeniz, N.; Rothstein, R.: Rsp5, a ubiquitin-pro-

tein ligase, is involved in degradation of the single-stranded-DNA binding protein Rfa1 in *Saccharomyces cerevisiae*. *Molec. Cell. Biol.* 20: 224–232, 2000.

[71877] 10207. Lu, Q. R.; Park, J. K.; Noll, E.; Chan, J. A.; Alberta, J.; Yuk, D.; Alzamora, M. G.; Louis, D. N.; Stiles, C. D.; Rowitch, D. H.; Black, P. M.: Oligodendrocyte lineage genes (OLIG) as molecular markers for human glial brain tumors. *Proc. Nat. Acad. Sci.* 98: 10851–10856, 2001.

[71878] 10208. Riggins, G. J.; Thiagalingam, S.; Rozenblum, E.; Weinstein, C. L.; Kern, S. E.; Hamilton, S. R.; Willson, J. K. V.; Markowitz, S. D.; Kinzler, K. W.; Vogelstein, B.: Mad-related genes in the human. *Nature Genet.* 13: 347–349, 1996.

[71879] 10209. Harteneck, C.; Wedel, B.; Koesling, D.; Malkewitz, J.; Bohme, E.; Schultz, G.: Molecular cloning and expression of a new alpha-subunit of soluble guanylyl cyclase. Interchangeability of the alpha-subunit of the enzyme. *FEBS Lett.* 292: 217–222, 1991.

[71880] 10210. Yu, F.; Warburton, D.; Wellington, S.; Danziger, R. S.: Assignment of GUCIA2, the gene coding for the alpha-2 subunit of soluble guanylyl cyclase, to position 11q21–q22 on human chromosome 11. *Genomics* 33: 334–336, 1996.

[71881] 10211. Abe, K.; Yamamura, K.; Suzuki, M.: Molecular and

embryological characterization of a new transgene-induced null allele of mouse Brachyury locus. *Mammalian Genome* 11: 238–240, 2000.

[71882] 10212. European Consortium on MEN1: Mapping of the gene encoding the  $\beta$ 56-subunit of protein phosphatase 2A (PPP2R5B) to a 0.5-Mb region of chromosome 11q13 and its exclusion as a candidate gene for multiple endocrine neoplasia type 1 (MEN1). *Hum. Genet.* 100: 481–485, 1997.

[71883] 10213. Arsenijevic, D.; Onuma, H.; Pecqueur, C.; Raimbault, S.; Manning, B. S.; Miroux, B.; Couplan, E.; Alves-Guerra, M.-C.; Goubern, M.; Surwit, R.; Bouillard, F.; Richard, D.; Collins, S.; Ricquier, D.: Disruption of the uncoupling protein-2 gene in mice reveals a role in immunity and reactive oxygen species production. *Nature Genet.* 26:435–439, 2000.

[71884] 10214. Bouchard, C.; Perusse, L.; Chagnon, Y. C.; Warden, C.; Ricquier, D.: Linkage between markers in the vicinity of the uncoupling protein 2 gene and resting metabolic rate in humans. *Hum. Molec. Genet.* 6:1887–1889, 1997.

[71885] 10215. Brauner, P.; Nibbelink, M.; Flachs, P.; Vitkova, I.; Kopecky, P.; Mertelikova, I.; Janderova, L.; Penicaud, L.; Casteilla, L.; Plavka, R.; Kopecky, J.: Fast decline of

hematopoiesis and uncoupling protein2 content in human liver after birth: location of the protein in Kupffercells. *Pediat. Res.* 49: 440–447, 2001.

[71886] 10216.Esterbauer, H.; Schneitler, C.; Oberkofler, H.; Ebenbichler, C.; Paulweber, B.; Sandhofer, F.; Ladurner, G.; Hell, E.; Strosberg, A.D.; Patsch, J. R.; Krempler, F.; Patsch, W.: A common polymorphism in the promoter of UCP2 is associated with decreased risk of obesity in middle-aged humans. *Nature Genet.* 28: 178–183, 2001.

[71887] 10217.Fleury, C.; Neverova, M.; Collins, S.; Raimbault, S.; Champigny, O.; Levi-Meyrueis, C.; Bouillaud, F.; Seldin, M. F.; Surwit, R. S.; Ricquier, D.; Warden, C. H.: Uncoupling protein-2: a novel gene linked to obesity and hyperinsulinemia. *Nature Genet.* 15: 269–272, 1997.

[71888] 10218.Flier, J. S.; Lowell, B. B.: Obesity research springs a proton leak. *Nature Genet.* 15: 223–224, 1997.

[71889] 10219.Millet, L.; Vidal, H.; Andreelli, F.; Larrouy, D.; Riou, J.-P.; Ricquier, D.; Laville, M.; Langin, D.: Increased uncoupling protein-2 and -3 mRNA expression during fasting in obese and lean humans. *J.Clin. Invest.* 100: 2665–2670, 1997.

[71890] 10220.Edwards, Y. H.; Putt, W.; Lekoape, K. M.; Stott, D.; Fox, M.; Hopkinson, D. A.; Sowden, J.: The human homolog

T of the mouse T (Brachyury) gene: gene structure, cDNA sequence, and assignment to chromosome 6q27. *Genome Res.* 6: 226–233, 1996.

- [71891] 10221. Morrison, K.; Papapetrou, C.; Attwood, J.; Hol, F.; Lynch, S. A.; Sampath, A.; Hamel, B.; Burn, J.; Sowden, J.; Stott, D.; Mariman, E.; Edwards, Y. H.: Genetic mapping of the human homologue (T) of mouse T (Brachyury) and a search for allele association between human T and spina bifida. *Hum. Molec. Genet.* 5: 669–674, 1996.
- [71892] 10222. Papapetrou, C.; Drummond, F.; Reardon, W.; Winter, R.; Spitz, L.; Edwards, Y. H.: A genetic study of the human T gene and its exclusion as a major candidate gene for sacral agenesis with anorectal atresia. *J. Med. Genet.* 36: 208–213, 1999.
- [71893] 10223. Shields, D. C.; Ramsbottom, D.; Donoghue, C.; Pinjon, E.; Kirke, P. N.; Molloy, A. M.; Edwards, Y. H.; Mills, J. L.; Mynett-Johnson, L.; Weir, D. G.; Scott, J. M.; Whitehead, A. S.: Association between historically high frequencies of neural tube defects and the human T homologue of mouse T (Brachyury). *Am. J. Med. Genet.* 92: 206–211, 2000.
- [71894] 10224. Speer, M. C.; Melvin, E. C.; Viles, K. D.; Bauer, K. A.; Rampersaud, E.; Drake, C.; George, T. M.; Enterline, D. S.; Mackey, J. F.; Worley, G.; Gilbert, J. R.; Nye, J. S.; NTD Col-

laborative Group: T locus shows no evidence for linkage disequilibrium or mutation in American Caucasian neural tube defect families. *Am. J. Med. Genet.* 110: 215–218, 2002.

- [71895] 10225. Trembath, D.; Sherbondy, A. L.; Vandyke, D. C.; Shaw, G. M.; Todoroff, K.; Lammer, E. J.; Finnell, R. H.; Marker, S.; Lerner, G.; Murray, J. C.: Analysis of select folate pathway genes, PAX3, and human Tin in a midwestern neural tube defect population. *Teratology* 59: 331–341, 1999.
- [71896] 10226. Bellomo, D.; Headrick, J. P.; Silins, G. U.; Paterson, C. A.; Thomas, P. S.; Gartside, M.; Mould, A.; Cahill, M. M.; Tonks, I. D.; Grimmond, S. M.; Townson, S.; Wells, C.; Little, M.; Cummings, M. C.; Hayward, N. K.; Kay, G. F.: Mice lacking the vascular endothelial growth factor-B gene (Vegfb) have smaller hearts, dysfunctional coronary vasculature, and impaired recovery from cardiac ischemia. *Circ. Res.* 86: e29–e35, 2000.
- [71897] 10227. Gerace, L.; Cirenei, N.; Cappelletti, M.; Petraroli, R.; Sebastiani, F.; Marziliano, N.: Assignment of the mouse Vegfb gene to mouse chromosome 19B by in situ hybridization. *Cytogenet. Cell Genet.* 95: 242–243, 2001.
- [71898] 10228. Grimmond, S.; Lagercrantz, J.; Drinkwater, C.;



Silins, G.; Townson, S.; Pollock, P.; Gotley, D.; Carson, E.; Rakar, S.; Nordenskjold, M.; Ward, L.; Hayward, N.; Weber, G.: Cloning and characterization of a novel human gene related to vascular endothelial growth factor. *Genome Res.* 6: 124–131, 1996.

[71899] 10229. Olofsson, B.; Pajusola, K.; Kaipainen, A.; von Euler, G.; Joukov, V.; Saksela, O.; Orpana, A.; Pettersson, R. F.; Alitalo, K.; Eriksson, U.: Vascular endothelial growth factor B, a novel growth factor for endothelial cells. *Proc. Nat. Acad. Sci.* 93: 2576–2581, 1996.

[71900] 10230. Olofsson, B.; Pajusola, K.; von Euler, G.; Chilov, D.; Alitalo, K.; Eriksson, U.: Genomic organization of the mouse and human genes for vascular endothelial growth factor B (VEGF-B) and characterization of a second splice isoform. *J. Biol. Chem.* 271: 19310–19317, 1996.

[71901] 10231. Gunduz, M.; Ouchida, M.; Fukushima, K.; Hanafusa, H.; Etani, T.; Nishioka, S.; Nishizaki, K.; Shimizu, K.: Genomic structure of the human ING1 gene and tumor-specific mutations detected in head and neck squamous cell carcinomas. *Cancer Res.* 60: 3143–3146, 2000.

[71902] 10232. Hasegawa, H.; Kiyokawa, E.; Tanaka, S.; Nagashima, K.; Gotoh, N.; Shibuya, M.; Kurata, T.; Matsuda, M.: DOCK180, a major CRK-binding protein, alters cell

morphology upon translocation to the membrane.

Molec.Cell Biol. 16: 1770–176, 1996.

[71903] 10233.Savill, J.: Phagocytic docking without shocking. Nature 442–443,1998.

[71904] 10234.Takai, S.; Hasegawa, H.; Kiyokawa, E.; Yamada, K.; Kurata, T.;Matsuda, M.: Chromosomal mapping of the gene encoding DOCK180, amajor Crk–binding protein, to 10q26.13–q26.3 by fluorescence in situhybridization. Genomics 35: 403–404, 1996.

[71905] 10235.Wu, Y.–C.; Horvitz, H. R.: C. elegans phagocytosis and cell–migrationprotein CED–5 is similar to human DOCK180. Nature 392: 501–504,1998.

[71906] 10236.Alexander, J. M.; Bikkal, H. A.; Zervas, N. T.; Laws, E. R., Jr.;Klibanski, A.: Tumor–specific expression and alternate splicing ofmessenger ribonucleic acid encoding activin/transforming growth factor–betareceptors in human pituitary adenomas. J. Clin. Endocr. Metab. 81:783–790, 1996.

[71907] 10237.Su, G. H.; Bansal, R.; Murphy, K. M.; Montgomery, E.; Yeo, C. J.;Hruban, R. H.; Kern, S. E.: ACVR1B (ALK4, activin receptor type 1B)gene mutations in pancreatic carcinoma. Proc. Nat. Acad. Sci. 98:3254–3257, 2001.

[71908] 10238.Xu, J.; Matsuzaki, K.; McKeehan, K.; Wang, F.; Kan,

M.; McKeehan, W. L.: Genomic structure and cloned cDNAs predict that four variants in the kinase domain of serine/threonine kinase receptors arise by alternative splicing and poly(A) addition. *Proc. Nat. Acad. Sci.* 91:7957–7961, 1994.

[71909] 10239. Zhou, Y.; Sun, H.; Danila, D. C.; Johnson, S. R.; Sigai, D. P.; Zhang, X.; Klibanski, A.: Truncated activin type I receptor Alk4 isoforms are dominant negative receptors inhibiting activin signaling. *Molec. Endocr.* 14: 2066–2075, 2000.

[71910] 10240. Mann, S. S.; Pettenati, M. J.; von Kapherr, C.; Hart, T. C.: Reassignment of peptidyl prolyl isomerase-like 1 gene (PPIL1) to human chromosome region 6p21.1 by radiation hybrid mapping and fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 83: 228–229, 1998.

[71911] 10241. Ozaki, K.; Fujiwara, T.; Kawai, A.; Shimizu, F.; Takami, S.; Okuno, S.; Takeda, S.; Shimada, Y.; Nagata, M.; Watanabe, T.; Takaichi, A.; Takahashi, E.; Nakamura, Y.; Shin, S.: Cloning, expression and chromosomal mapping of a novel cyclophilin-related gene (PPIL1) from human fetal brain. *Cytogenet. Cell Genet.* 72: 242–245, 1996.

[71912] 10242. Graef, I. A.; Chen, F.; Chen, L.; Kuo, A.; Crabtree, G. R.: Signal transduced by  $\text{Ca}^{2+}$ /calcineurin and

NFATc3/c4 pattern the developing vasculature. *Cell* 105: 863–875, 2001.

- [71913] 10243. Wang, M. G.; Yi, H.; Guerini, D.; Klee, C. B.; McBride, O. W.: Calcineurin A alpha (PPP3CA), calcineurin A beta (PPP3CB) and calcineurin B (PPP3R1) are located on human chromosomes 4, 10q21–q22 and 2p16–p15, respectively. *Cytogenet. Cell Genet.* 72: 236–241, 1996.
- [71914] 10244. Zeng, H.; Chattarji, S.; Barbarosie, M.; Rondi-Reig, L.; Philpot, B. D.; Miyakawa, T.; Bear, M. F.; Tonegawa, S.: Forebrain-specific calcineurin knockout selectively impairs bidirectional synaptic plasticity and working/episodic-like memory. *Cell* 107: 617–629, 2001.
- [71915] 10245. Kolodrubetz, D.; Burgum, A.: Sequence and genetic analysis of NHP2: a moderately abundant high mobility group-like nuclear protein with an essential function in *Saccharomyces cerevisiae*. *Yeast* 7: 79–90, 1991.
- [71916] 10246. Saito, H.; Fujiwara, T.; Shin, S.; Okui, K.; Nakamura, Y.: Cloning and mapping of a human novel cDNA (NHP2L1) that encodes a protein highly homologous to yeast nuclear protein NHP2. *Cytogenet. Cell Genet.* 72: 191–193, 1996.
- [71917] 10247. Mooseker, M. S.; Cheney, R. E.: Unconventional myosins. *Annu. Rev. Cell Dev. Biol.* 11: 633–675, 1995.

- [71918] 10248.Bement, W. M.; Wirth, J. A.; Mooseker, M. S.: Cloning and mRNA expression of human unconventional myosin-IC: a homologue of amoeboid myosins-I with a single IQ motif and an SH3 domain. *J. Molec. Biol.* 243:356–363, 1994.
- [71919] 10249.Bement, W. M.; Hasson, T.; Wirth, J. A.; Cheney, R. E.; Mooseker, M. S.: Identification and overlapping expression of multiple unconventional myosin genes in vertebrate cell types. *Proc. Nat. Acad. Sci.* 91:6549–6553, 1994. Erratum: *Proc. Nat. Acad. Sci.* 91: 11767, 1994.
- [71920] 10250.Crozet, F.; Amraoui, A. E.; Blanchard, S.; Lenoir, M.; Ripoll, C.; Vago, P.; Hamel, C.; Fizames, C.; Levi-Acobas, F.; Depetris, D.; Mattei, M.-G.; Weil, D.; Pujol, R.; Petit, C.: Cloning of the genes encoding two murine and human cochlear unconventional type I myosins. *Genomics* 40:332–341, 1997.
- [71921] 10251.Goppelt, A.; Stelzer, G.; Lottspeich, F.; Meisterernst, M.: A mechanism for repression of class II gene transcription through specific binding of NC2 to TBP-promoter complexes via heterodimeric histone fold domains. *EMBO J.* 15: 3105–3116, 1996.
- [71922] 10252.Inostroza, J. A.; Mermeistein, F. H.; Ha, I.; Lane, W. S.; Reinberg, D.: Dr1, a TATA-binding protein-associated

phosphoprotein and inhibitor of class II gene transcription. Cell 70: 477–489, 1992.

[71923] 10253. Kamada, K.; Shu, F.; Chen, H.; Malik, S.; Stelzer, G.; Roeder, R. G.; Meisterernst, M.; Burley, S. K.: Crystal structure of negative cofactor 2 recognizing the TBP–DNA transcription complex. Cell 106:71–81, 2001.

[71924] 10254. Mermelstein, F.; Yeung, K.; Cao, J.; Inostroza, J. A.; Erdjument-Bromage, H.; Egelson, K.; Landsman, D.; Levitt, P.; Tempst, P.; Reinberg, D.: Requirement of a corepressor for Dr1-mediated repression of transcription. Genes Dev. 10: 1033–1048, 1996.

[71925] 10255. Purello, M.; Di Pietro, C.; Rapisarda, A.; Viola, A.; Corsaro, C.; Motta, S.; Grzeschik, K.–H.; Sichel, G.: Genomic localization of the human gene encoding Dr1, a negative modulator of transcription of class II and class III genes. Cytogenet. Cell Genet. 75: 186–189, 1996.

[71926] 10256. Rozet, J.–M.; Gerber, S.; Perrault, I.; Camuzat, A.; Calvas, P.; Viegas-Pequignot, E.; Molina-Gomes, D.; Le Paslier, D.; Chumakov, I.; Munnich, A.; Kaplan, J.: Structure and physical mapping of DR1, a TATA-binding protein-associated phosphoprotein gene, to chromosome 1p22.1 and its exclusion in Stargardt disease (STGD). Genomics 36:554–556, 1996.

- [71927] 10257.Willy, P. J.; Kobayashi, R.; Kadonaga, J. T.: A basal transcriptionfactor that activates or represses transcription. *Science* 290: 982–984,2000.
- [71928] 10258.Yeung, K.; Kim, S.; Reinberg, D.: Functional dissection of a humanDr1–DRAP1 repressor complex. *Molec. Cell. Biol.* 17: 36–45, 1997.
- [71929] 10259.Dong, L. Q.; Du, H.; Porter, S. G.; Kolakowski, L. F., Jr.; Lee,A. V.; Mandarino, J.; Fan, J.; Yee, D.; Liu, F.: Cloning, chromosomelocalization, expression, and characterization of an Src homology2 and pleckstrin homology domain–containing insulin receptor bindingprotein hGrb10–gamma. *J. Biol. Chem.* 272: 29104–29112, 1997.
- [71930] 10260.Margolis, B.; Silvennoinen, O.; Comoglio, F.; Roonprapunt, C.;Skolnik, E.; Ullrich, A.; Schlessinger, J.: High-efficiency expression/cloningof epidermal growth factor–receptor–binding proteins with Src homology2 domains. *Proc. Nat. Acad. Sci.* 89: 8894–8898, 1992.
- [71931] 10261.Skolnik, E. Y.; Margolis, B.; Mohammadi, M.; Lowenstein, E.; Fischer,R.; Drepps, A.; Ullrich, A.; Schlessinger, J.: Cloning of PI3 kinase–associated p85 utilizing a novel method for expression/cloning oftarget proteins for receptor tyrosine kinases. *Cell* 65: 83–90, 1991.
- [71932] 10262.Tanaka, S.; Mori, M.; Akiyoshi, T.; Tanaka, Y.; Ma–

fune, K.; Wands, J. R.; Sugimachi, K.: A novel variant of human Grb7 is associated with invasive esophageal carcinoma. *J. Clin. Invest.* 102: 821–827, 1998.

[71933] 10263. Blagitko, N.; Mergenthaler, S.; Schulz, U.; Wollmann, H. A.; Craigen, W.; Eggermann, T.; Ropers, H.-H.; Kalscheuer, V. M.: Human GRB10 is imprinted and expressed from the paternal and maternal allele in a highly tissue- and isoform-specific fashion. *Hum. Molec. Genet.* 9: 1587–1595, 2000.

[71934] 10264. Frantz, J. D.; Giorgetti-Peraldi, S.; Ottinger, E. A.; Shoelson, S. E.: Human GRB-IR-beta/GRB10: splice variants of an insulin and growth factor receptor-binding protein with PH and SH2 domains. *J. Biol. Chem.* 272: 2659–2667, 1997.

[71935] 10265. Hannula, K.; Lipsanen-Nyman, M.; Kontiokari, T.; Kere, J.: A narrow segment of maternal uniparental disomy of chromosome 7q31-qter in Silver-Russell syndrome delimits a candidate gene region. *Am. J. Hum. Genet.* 68: 247–253, 2001.

[71936] 10266. Jerome, C. A.; Scherer, S. W.; Tsui, L.-C.; Gietz, R. D.; Triggs-Raine, B.: Assignment of growth factor receptor-bound protein 10 (GRB10) to human chromosome 7p11.2–p12. *Genomics* 40: 215–216, 1997.



- [71937] 10267.Liu, F.; Roth, R. A.: Grb-IR: a SH2-domain-containing protein that binds to the insulin receptor and inhibits its function. *Proc.Nat. Acad. Sci.* 92: 10287-10291, 1995.
- [71938] 10268.Ooi, J.; Yajnik, V.; Immanuel, D.; Gordon, M.; Moskow, J. J.; Buchberg, A. M.; Margolis, B.: The cloning of Grb10 reveals a new family of SH2 domain proteins. *Onco-gene* 10: 1621-1630, 1995.
- [71939] 10269.Daly, R. J.; Sanderson, G. M.; Janes, P. W.; Sutherland, R. L.: Cloning and characterization of GRB14, a novel member of the GRB7 gene family. *J. Biol. Chem.* 271: 12502-12510, 1996.
- [71940] 10270.Hakala, B. E.; White, C.; Recklies, A. D.: Human cartilage gp-39, a major secretory product of articular chondrocytes and synovial cells, is a mammalian member of a chitinase protein family. *J. Biol. Chem.* 268:25803-25810, 1993.
- [71941] 10271.Rehli, M.; Krause, S. W.; Andreesen, R.: Molecular characterization of the gene for human cartilage gp-39 (CHI3L1), a member of the chitinase protein family and marker for late stages of macrophage differentiation. *Genomics* 43:221-225, 1997.
- [71942] 10272.Dry, K.; Kenwrick, S.; Rosenthal, A.; Platzer, M.: The

complete sequence of the human locus for NgCAM-related cell adhesion molecule reveals a novel alternative exon in chick and man and conserved genomic organization for the L1 subfamily. *Gene* 273: 115–122, 2001.

[71943] 10273. Grumet, M.; Mauro, V.; Burgoon, M. P.; Edelman, G. M.; Cunningham, B. A.: Structure of a new nervous system glycoprotein, Nr-CAM, and its relationship to subgroups of neural cell adhesion molecules. *J. Cell. Biol.* 113: 1399–1412, 1991.

[71944] 10274. Kayyem, J. F.; Roman, J. M.; de la Rosa, E. J.; Schwarz, U.; Dreyer, W. J.: Bravo/Nr-CAM is closely related to the cell adhesion molecules L1 and Ng-CAM and has a similar heterodimer structure. *J. Cell. Biol.* 118: 1259–1270, 1992.

[71945] 10275. Lane, R. P.; Chen, X.-N.; Yamakawa, K.; Vielmetter, J.; Korenberg, J. R.; Dreyer, W. J.: Characterization of a highly conserved human homolog to the chicken neural cell surface protein Bravo/Nr-CAM that maps to chromosome band 7q31. *Genomics* 35: 456–465, 1996.

[71946] 10276. Wang, B.; Williams, H.; Du, J.-S.; Terrett, J.; Kenwick, S.: Alternative splicing of human NrCAM in neural and nonneural tissues. *Molec. Cell. Neurosci.* 10: 287–295, 1998.

- [71947] 10277. Damen, J. E.; Liu, L.; Rosten, P.; Humphries, R. K.; Jefferson, A. B.; Majerus, P. W.; Krystal, G.: The 145-kDa protein induced to associate with Shc by multiple cytokines is an inositol tetrakisphosphate and phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase. *Proc. Nat. Acad. Sci.* 93: 1689–1693, 1996.
- [71948] 10278. Drayer, A. L.; Pesesse, X.; De Smedt, F.; Woscholski, R.; Parker, P.; Erneux, C.: Cloning and expression of a human placenta inositol 1,3,4,5-tetrakisphosphate and phosphatidylinositol 3,4,5-trisphosphate 5-phosphatase. *Biochem. Biophys. Res. Commun.* 225: 243–249, 1996.
- [71949] 10279. Helgason, C. D.; Damen, J. E.; Rosten, P.; Grewal, R.; Sorensen, P.; Chappel, S. M.; Borowski, A.; Jirik, F.; Krystal, G.; Humphries, R. K.: Targeted disruption of SHIP leads to hemopoietic perturbations, lung pathology, and a shortened life span. *Genes Dev.* 12: 1610–1620, 1998.
- [71950] 10280. Huber, M.; Helgason, C. D.; Damen, J. E.; Liu, L.; Humphries, R. K.; Krystal, G.: The src homology 2-containing inositol phosphatase (SHIP) is the gatekeeper of mast cell degranulation. *Proc. Nat. Acad. Sci.* 95: 11330–11335, 1998.
- [71951] 10281. Kavanaugh, W. M.; Pot, D. A.; Chin, S. M.; Deuter-Reinhard, M.; Jefferson, A. B.; Norris, F. A.; Masiarz, F. R.;

Cousens, L. S.; Majerus, P. W.; Williams, L. T.: Multiple forms of an inositol polyphosphate 5-phosphatase form signaling complexes with Shc and Grb2. *Curr. Biol.* 6: 438–445, 1996.

- [71952] 10282. Lioubin, M. N.; Algate, P. A.; Tsai, S.; Carlberg, K.; Aebersold, R.; Rohrschneider, L. R.: p150(Ship), a signal transduction molecule with inositol polyphosphate 5-phosphatase activity. *Genes Dev.* 10:1084–1095, 1996.
- [71953] 10283. Liu, Q.; Amgen EST Program; Dumont, D. J.: Molecular cloning and chromosomal localization in human and mouse of the SH2-containing inositol phosphatase, INPP5D (SHIP). *Genomics* 39: 109–112, 1997.
- [71954] 10284. Liu, Q.; Shalaby, F.; Jones, J.; Bouchard, D.; Dumont, D. J.: The SH2-containing inositol polyphosphate 5-phosphatase, Ship, is expressed during hematopoiesis and spermatogenesis. *Blood* 91: 2753–2759, 1998.
- [71955] 10285. Takeshita, S.; Namba, N.; Zhao, J. J.; Jiang, Y.; Genant, H. K.; Silva, M. J.; Brodt, M. D.; Helgason, C. D.; Kalesnikoff, J.; Rauh, M. J.; Humphries, R. K.; Krystal, G.; Teitelbaum, S. L.; Ross, F. P.: SHIP-deficient mice are severely osteoporotic due to increased numbers of hyper-resorptive osteoclasts. *Nature Med.* 8: 943–949, 2002.
- [71956] 10286. Wang, J.-W.; Howson, J. M.; Ghansah, T.; Desponts,

C.; Ninos, J. M.; May, S. L.; Nguyen, K. H. T.; Toyama-Sorimachi, N.; Kerr, W.G.: Influence of SHIP on the NK repertoire and allogeneic bone marrow transplantation. *Science* 295: 2094–2097, 2002.

[71957] 10287. Ware, M. D.; Rosten, P.; Damen, J. E.; Liu, L.; Humphries, R.K.; Krystal, G.: Cloning and characterization of human SHIP, the 145-kD inositol 5-phosphatase that associates with SHC after cytokine stimulation. *Blood* 88: 2833–2840, 1996.

[71958] 10288. Cates, C. A.; Michael, R. L.; Stayrook, K. R.; Harvey, K. A.; Burke, Y. D.; Randall, S. K.; Crowell, P. L.; Crowell, D. N.: Prenylation of oncogenic human PTP(CAAX) protein tyrosine phosphatases. *Cancer Lett.* 110: 49–55, 1996.

[71959] 10289. Montagna, M.; Serova, O.; Sylla, B. S.; Feunteun, J.; Lenoir, G.M.: A 100-kb physical and transcriptional map around the EDH17B2 gene: identification of three novel genes and a pseudogene of a human homologue of the rat PRL-1 tyrosine phosphatase. *Hum. Genet.* 96: 532–538, 1995.

[71960] 10290. Zeng, Q.; Hong, W.; Tan, Y. H.: Mouse PRL-2 and PRL-3, two potentially prenylated protein tyrosine phosphatases homologous to PRL-1. *Biochem. Biophys. Res. Commun.* 244: 421–427, 1998.

- [71961] 10291.Muneer, S.; Ramalingam, V.; Wyatt, R.; Schultz, R. A.; Minna, J.D.; Kamibayashi, C.: Genomic organization and mapping of the gene encoding the PP2A B56-gamma regulatory subunit. *Genomics* 79: 344–348, 2002.
- [71962] 10292.Deveraux, Q.; Jensen, C.; Rechsteiner, M.: Molecular cloning and expression of a 26 S protease subunit enriched in dileucine repeats. *J.Biol. Chem.* 270: 23726–23729, 1995.
- [71963] 10293.Deveraux, Q.; Ustrell, V.; Pickart, C.; Rechsteiner, M.: A 26S protease subunit that binds ubiquitin conjugates. *J. Biol. Chem.* 269:7059–7061, 1994.
- [71964] 10294.Hu, R.-J.; Lee, M. P.; Johnson, L. A.; Feinberg, A. P.: A novel human homologue of yeast nucleosome assembly protein, 65 kb centromeric to the p57(KIP2) gene, is biallelically expressed in fetal and adult tissues. *Hum. Molec. Genet.* 5: 1743–1748, 1996.
- [71965] 10295.Abderrahim, H.; Jaramillo-Babb, V. L.; Zhou, Z.; Vollrath, D.: Characterization of the murine TIGR/myocilin gene. *Mammalian Genome* 9:673–675, 1998.
- [71966] 10296.Adam, M. F.; Belmouden, A.; Binisti, P.; Brezin, A. P.; Valtot, F.; Bechetoille, A.; Dascotte, J.-C.; Copin, B.; Gomez, L.; Chaventre, A.; Bach, J.-F.; Garchon, H.-J.: Recurrent mutations in a single exon encoding the evolution-

arily conserved olfactomedin-homology domain of TIGR in familial open-angle glaucoma. *Hum. Molec. Genet.* 6: 2091–2097, 1997.

- [71967] 10297. Baird, P. N.; Dickinson, J.; Craig, J. E.; Mackey, D. A.: The Taa1 restriction enzyme provides a simple means to identify the Q368STOP mutation of the myocilin gene in primary open angle glaucoma. *Am.J. Ophthalm.* 131: 510–511, 2001.
- [71968] 10298. Rodriguez, P.; Munroe, D.; Prawitt, D.; Chu, L. L.; Bric, E.; Kim, J.; Reid, L. H.; Davies, C.; Nakagama, H.; Loebbert, R.; Winterpacht, A.; Petruzzi, M.-J.; Higgins, M. J.; Nowak, N.; Evans, G.; Shows, T.; Weissman, B. E.; Zabel, B.; Housman, D. E.; Pelletier, J.: Functional characterization of human nucleosome assembly protein-2 (NAP1L4) suggests a role as a histone chaperone. *Genomics* 44: 253–265, 1997.
- [71969] 10299. Stegmaier, K.; Pendse, S.; Barker, G. F.; Bray-Ward, P.; Ward, D. C.; Montgomery, K. T.; Krauter, K. S.; Reynolds, C.; Sklar, J.; Donnelly, M.; Bohlander, S. K.; Rowley, J. D.; Sallan, S. E.; Gilliland, D. G.; Golub, T. R.: Frequent loss of heterozygosity at the TEL gene locus in acute lymphoblastic leukemia of childhood. *Blood* 86: 38–44, 1995.

- [71970] 10300. Arakawa, H.; Nagase, H.; Hayashi, N.; Fujiwara, T.; Ogawa, M.; Shin, S.; Nakamura, Y.: Molecular cloning and expression of a novel human gene that is highly homologous to human FK506-binding protein 12 kDa (hFKBP-12) and characterization of two alternatively spliced transcripts. *Biochem. Biophys. Res. Commun.* 200: 836–843, 1994.
- [71971] 10301. Xin, H.-B.; Senbonmatsu, T.; Cheng, D.-S.; Wang, Y.-X. Copello, J. A.; Ji, G.-J.; Collier, M. L.; Deng, K.-Y.; Jeyakumar, L. H.; Magnuson, M. A.; Inagami, T.; Kotlikoff, M. I.; Fleischer, S.: Oestrogen protects FKBP12.6 null mice from cardiac hypertrophy. *Nature* 416: 334–337, 2002.
- [71972] 10302. Stein, R.; Mori, N.; Matthews, K.; Lo, L.-C.; Anderson, D. J.: The NGF-inducible SCG10 mRNA encodes a novel membrane-bound protein present in growth cones and abundant in developing neurons. *Neuron* 1: 463–476, 1988.
- [71973] 10303. Dong, J.-T.; Isaacs, W. B.; Barrett, J. C.; Isaacs, J. T.: Genomic organization of the human KAI1 metastasis-suppressor gene. *Genomics* 41: 25–32, 1997.
- [71974] 10304. Dong, J.-T.; Lamb, P. W.; Rinker-Schaeffer, C. W.; Vukanovic, J.; Ichikawa, T.; Isaacs, J. T.; Barrett, J. C.: KAI1, a metastasis suppressor gene for prostate cancer on hu-



man chromosome 11p11.2. Science 268:884–886, 1995.

- [71975] 10305. Guo, X.-Z.; Friess, H.; Di Mola, F. F.; Heinicke, J.-M.; Abou-Shady, M.; Graber, H. U.; Baer, H. U.; Zimmermann, A.; Korc, M.; Buchler, M. W.: KAI1, a new metastasis suppressor gene, is reduced in metastatic hepatocellular carcinoma. Hepatology 28: 1481–1488, 1998.
- [71976] 10306. Mashimo, T.; Watabe, M.; Hirota, S.; Hosobe, S.; Miura, K.; Tegtmeyer, P. J.; Rinker-Shaeffer, C. W.; Watabe, K.: The expression of the KAI1 gene, a tumor metastasis suppressor, is directly activated by p53. Proc. Nat. Acad. Sci. 95: 11307–11311, 1998.
- [71977] 10307. Miyazaki, T.; Kato, H.; Shitara, Y.; Yoshikawa, M.; Tajima, K.; Masuda, N.; Shouji, H.; Tsukada, K.; Nakajima, T.; Kuwano, H.: Mutation and expression of the metastasis suppressor gene KAI1 in esophageal squamous cell carcinoma. Cancer 89: 955–962, 2000.
- [71978] 10308. Gupta, P.; Soyombo, A. A.; Atashband, A.; Wisniewski, K. E.; Shelton, J. M.; Richardson, J. A.; Hammer, R. E.; Hofmann, S. L.: Disruption of PPT1 or PPT2 causes neuronal ceroid lipofuscinosis in knockout mice. Proc. Nat. Acad. Sci. 98: 13566–13571, 2001.
- [71979] 10309. Ardell, M. D.; Aragon, I.; Oliveira, L.; Porche, G. E.; Burke, E.; Pittler, S. J.: The beta subunit of human rod pho-

photoreceptor cGMP-gated cation channel is generated from a complex transcription unit. FEBS Lett. 389: 213–218, 1996.

[71980] 10310. Ardell, M. D.; Makhija, A. K.; Oliveira, L.; Miniou, P.; Viegas-Pequignot, E.; Pittler, S. J.: cDNA, gene structure, and chromosomal localization of human GAR1 (CNCG3L), a homolog of the third subunit of bovine photoreceptor-cGMP-gated channel. Genomics 28: 32–38, 1995.

[71981] 10311. Bareil, C.; Hamel, C. P.; Delague, V.; Arnaud, B.; Demaille, J.; Claustres, M.: Segregation of a mutation in CNGB1 encoding the beta-subunit of the rod cGMP-gated channel in a family with autosomal recessive retinitis pigmentosa. Hum. Genet. 108: 328–334, 2001.

[71982] 10312. Bird, A. C.: Retinal photoreceptor dystrophies. LI. Edward Jackson Memorial Lecture. Am. J. Ophthalmol. 119: 543–562, 1995.

[71983] 10313. Chen, T.-Y.; Illing, M.; Molday, L. L.; Hsu, Y.-T.; Yau, K.-W.; Molday, R. S.: Subunit 2 (or beta) of retinal rod cGMP-gated cation channel is a component of the 240-kDa channel-associated protein and mediates  $\text{Ca}^{2+}$ -calmodulin modulation. Proc. Nat. Acad. Sci. 91: 11757–11761, 1994.

[71984] 10314. Chen, T. Y.; Peng, Y. W.; Dhallan, R. S.; Ahamed, B.;

Reed, R.R.; Yau, K. W.: A new subunit of the cyclic nucleotide-gated cation channel in retinal rods. *Nature* 362: 764–767, 1993.

[71985] 10315. Korschen, H. G.; Beyermann, M.; Muller, F.; Heck, M.; Vantler, M.; Koch, K.-W.; Kellner, R.; Wolfrum, U.; Bode, C.; Hofmann, K. P.; Kaupp, U. B.: Interaction of glutamic-acid-rich proteins with the cGMP signalling pathway in rod photoreceptors. *Nature* 400: 761–766, 1999.

[71986] 10316. Neufeld, E. F.: Personal Communication. Los Angeles, Calif. 7/24/1987.

[71987] 10317. Blair, H. J.; Ho, M.; Monaco, A. P.; Fisher, S.; Craig, I. W.; Boyd, Y.: High-resolution comparative mapping of the proximal region of the mouse X chromosome. *Genomics* 28: 305–310, 1995.

[71988] 10318. Cox, J. P. D.; Yamamoto, K.; Christie, P. T.; Wooding, C.; Feest, T.; Flinter, F. A.; Goodyer, P. R.; Leumann, E.; Neuhaus, T.; Reid, C.; Williams, P. F.; Wrong, O.; Thakker, R. V.: Renal chloride channel, CLCN5, mutations in Dent's disease. *J. Bone Min. Res.* 14: 1536–1542, 1999.

[71989] 10319. Craig, I.: Personal Communication. Oxford, England 1/31/1995.

[71990] 10320. Devuyst, O.; Christie, P. T.; Courtoy, P. J.; Beauwens, R.; Thakker, R. V.: Intra-renal and subcellular

distribution of the human chloride channel, CLC-5, reveals a pathophysiological basis for Dent's disease. *Hum.Molec. Genet.* 8: 247–257, 1999.

[71991] 10321.Dutzler, R.; Campbell, E. B.; Cadene, M.; Chait, B.

T.; MacKinnon,R.: X-ray structure of a CIC chloride channel at 3.0 angstrom reveals the molecular basis of anion selectivity. *Nature* 415: 287–294, 2002.

[71992] 10322.Fisher, S. E.; Black, G. C. M.; Lloyd, S. E.; Hatchwell,

E.; Wrong,O.; Thakker, R. V.; Craig, I. W.: Isolation and partial characterization of a chloride channel gene which is expressed in kidney and is a candidate for Dent's disease (an X-linked hereditary nephrolithiasis). *Hum.Molec. Genet.* 3: 2053–2059, 1994.

[71993] 10323.Fisher, S. E.; Van Bakel, I.; Lloyd, S. E.; Pearce, S. H.

S.; Thakker,R. V.; Craig, I. W.: Cloning and characterization of CLCN5, the human kidney chloride channel gene implicated in Dent disease (an X-linked hereditary nephrolithiasis). *Genomics* 29: 598–606, 1995.

[71994] 10324.Gunther, W.; Luchow, A.; Cluzeaud, F.; Vandewalle,

A.; Jentsch,T. J.: CLC-5, the chloride channel mutated in Dent's disease, colocalizes with the proton pump in endocytotically active kidney cells. *Proc.Nat. Acad. Sci.* 95: 8075–8080, 1998.

- [71995] 10325. Igarashi, T.; Hayakawa, H.; Shiraga, H.; Kawato, H.; Yan, K.; Kawaguchi, H.; Yamanaka, T.; Tsuchida, S.; Akagi, K.: Hypercalciuria and nephrocalcinosis in patients with idiopathic low molecular weight proteinuria in Japan: is the disease identical to Dent's disease in the United Kingdom? *Nephron*. 69:242–247, 1995.
- [71996] 10326. Lloyd, S. E.; Gunther, W.; Pearce, S. H. S.; Thomson, A.; Bianchi, M. L.; Bosio, M.; Craig, I. W.; Fisher, S. E.; Scheinman, S. J.; Wrong, O.; Jentsch, T. J.; Thakker, R. V.: Characterisation of renal chloride channel, CLCN5, mutations in hypercalciuric nephrolithiasis (kidney stones) disorders. *Hum. Molec. Genet.* 6: 1233–1239, 1997.
- [71997] 10327. Lloyd, S. E.; Pearce, S. H. S.; Fisher, S. E.; Steinmeyer, K.; Schwappach, B.; Scheinman, S. J.; Harding, B.; Bolino, A.; Devoto, M.; Goodyer, P.; Rigden, S. P. A.; Wrong, O.; Jentsch, T. J.; Craig, I. W.; Thakker, R. V.: A common molecular basis for three inherited kidney stone diseases. *Nature* 370: 445–449, 1996.
- [71998] 10328. Lloyd, S. E.; Pearce, S. H. S.; Gunther, W.; Kawaguchi, H.; Igarashi, T.; Jentsch, T. J.; Thakker, R. V.: Idiopathic low molecular weight proteinuria associated with hypercalciuric nephrocalcinosis in Japanese children is due to mutations of the renal chloride channel (CLCN5).

J.Clin. Invest. 99: 967–974, 1997.

- [71999] 10329.Oudet, C.; Martin–Coignard, D.; Pannetier, S.; Praud, E.; Champion,G.; Hanauer, A.: A second family with XLRH displays the mutationS244L in the CLCN5 gene. Hum. Genet. 99: 781–784, 1997.
- [72000] 10330.Piwon, N.; Gunther, W.; Schwake, M.; Bosl, M. R.; Jentsch, T.J.: CIC–5 Cl(–)–channel disruption impairs endocytosis in a mousemodel for Dent's disease. Nature 408: 369–373, 2000.
- [72001] 10331.Schurman, S. J.; Norden, A. G. W.; Scheinman, S. J.: X–linkedrecessive nephrolithiasis: presentation and diagnosis in children. J.Pediat. 132: 859–862, 1998.
- [72002] 10332.Cabeza–Arvelaiz, Y.; Shih, L.–C. N.; Hardman, N.; Asselbergs, F.;Bilbe, G.; Schmitz, A.; White, B.; Siciliano, M. J.; Lachman, L. B.: Cloning and genetic organization of the human kinesin light–chain(KLC) gene. DNA Cell Biol. 12: 881–892, 1993.
- [72003] 10333.Pearce, S. H. S.; Cheetham, T.; Imrie, H.; Vaidya, B.; Barnes,N. D.; Bilous, R. W.; Carr, D.; Meeran, K.; Shaw, N. J.; Smith, C.S.; Toft, A. D.; Williams, G.; Kendall–Taylor, P.: A common and recurrent13–bp deletion in the autoimmune regulator gene in British kindredswith autoimmune polyendocrinopathy type 1. Am. J. Hum. Genet.

63:1675–1684, 1998.

- [72004] 10334.Mitnick, M.; Reichlin, S.: Enzymatic synthesis of thyrotropin–releasing hormone (TRH) by hypothalamic 'TRH synthetase.'. *Endocrinology* 91:1145–1153, 1972.
- [72005] 10335.Battini, R.; Ferrari, S.; Kaczmarek, L.; Calabretta, B.; Chen, S.–T.; Baserga, R.: Molecular cloning of a cDNA for a human ADP/ATP carrier which is growth–regulated. *J. Biol. Chem.* 262: 4355–4359, 1987.
- [72006] 10336.Chen, S.–T.; Chang, C.–D.; Huebner, K.; Ku, D.–H.; McFarland, M.; DeRiel, J. K.; Baserga, R.; Wurzel, J.: A human ADP/ATP translocase gene has seven pseudogenes and localizes to chromosome X. *Somat.Cell Molec. Genet.* 16: 143–149, 1990.
- [72007] 10337.Ku, D.–H.; Kagan, J.; Chen, S.–T.; Chang, C.–D.; Baserga, R.; Wurzel, J.: The human fibroblast adenine nucleotide translocator gene: molecular cloning and sequence. *J. Biol. Chem.* 265: 16060–16063, 1990.
- [72008] 10338.Schiebel, K.; Mertz, A.; Winkelmann, M.; Nagaraja, R.; Rappold, G.: Localization of the adenine nucleotide translocase gene ANT2 to chromosome Xq24–q25 with tight linkage to DXS425. *Genomics* 24:605–606, 1994.
- [72009] 10339.Lu, Q. R.; Sun, T.; Zhu, Z.; Ma, N.; Garcia, M.; Stiles, C. D.; Rowitch, D. H.: Common developmental requirement

for Olig function indicates a motor neuron/oligodendrocyte connection. Cell 109: 75–86, 2002.

- [72010] 10340. Lu, Q. R.; Yuk, D.; Alberta, J. A.; Zhu, Z.; Pawlitzky, I.; Chan, J.; McMahon, A. P.; Stiles, C. D.; Rowitch, D. H.: Sonic hedgehog-regulated oligodendrocyte lineage genes encoding bHLH proteins in the mammalian central nervous system. Neuron 25: 317–329, 2000.
- [72011] 10341. Raff, M. C.; Miller, R. H.; Noble, M.: A glial progenitor cell that develops in vitro into an astrocyte or an oligodendrocyte depending on culture medium. Nature 303: 390–396, 1983.
- [72012] 10342. Zhou, Q.; Anderson, D. J.: The bHLH transcription factors OLIG2 and OLIG1 couple neuronal and glial subtype specification. Cell 109: 61–73, 2002.
- [72013] 10343. Zhou, Q.; Wang, S.; Anderson, D. J.: Identification of a novel family of oligodendrocyte lineage-specific basic helix–loop–helix transcription factors. Neuron 25: 331–343, 2000.
- [72014] 10344. Guimaraes, M. J.; Peterson, D.; Vicari, A.; Cocks, B. G.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Ferrick, D. A.; Kastelein, R. A.; Bazan, J. F.; Zlotnik, A.: Identification of a novel selD homolog from eukaryotes, bacteria, and archaea: is there an autoregulatory mechanism in selenocys–



teine metabolism? Proc. Nat. Acad. Sci. 93:15086–15091, 1996.

[72015] 10345.Koesters, R.; Adams, V.; Betts, D.; Moos, R.; Schmid, M.; Siermann,A.; Hassam, S.; Weitz, S.; Lichter, P.; Heitz, P. U.; von Knebel Doeberitz,M.; Briner, J.: Human eukaryotic initiation factor EIF2C1 gene: cDNAsequence, genomic organization, localization to chromosomal bands1q34–p35, and expression. Genomics 61: 210–218, 1999.

[72016] 10346.Martinez, J.; Patkaniowska, A.; Urlaub, H.; Luhrmann, R.; Tuschli,T.: Single–stranded antisense siR–NAs guide target RNA cleavage inRNAi. Cell 110: 563–574, 2002.

[72017] 10347.Suetsugu, S.; Miki, H.; Takenawa, T.: Identification of two humanWAVE/SCAR homologues as general actin regulatory molecules which associatewith the Arp2/3 complex. Biochem. Biophys. Res. Commun. 260: 296–302,1999.

[72018] 10348.Myers, L. C.; Gustafsson, c. M.; Bushnell, D. A.; Lui, M.; Erdjument–Bromage,H.; Tempst, P.; Kornberg, R. D.: The Med proteins of yeast and theirfunction through the RNA polymerase II carboxy–terminal domain. GenesDev. 12: 45–54, 1998.

- [72019] 10349.Hanaoka, E.; Ozaki, T.; Ohira, M.; Nakamura, Y.; Suzuki, M.; Takahashi,E.; Moriya, H.; Nakagawara, A.; Sakiyama, S.: Molecular cloning and expression analysis of the human DA41 gene and its mapping to chromosome9q21.2–q21.3. *J. Hum. Genet.* 45: 188–191, 2000.
- [72020] 10350.Ozaki, T.; Hishiki, T.; Toyama, Y.; Yuasa, S.; Nakagawara, A.;Sakiyama, S.: Identification of a new cellular protein that can interactspecifically with DAN. *DNA Cell Biol.* 16: 985–991, 1997.
- [72021] 10351.Venkateswarlu, K.; Gunn–Moore, F.; Oatey, P. B.; Tavaré, J. M.;Cullen, P. J.: Nerve growth factor– and epidermal growth factor–stimulatedtranslocation of the ADP–ribosylation factor–exchange factor GRP1to the plasma membrane of PC12 cells requires activation of phosphatidylinositol3–kinase and the GRP1 pleckstrin homology domain. *Biochem. J.* 335:139–146, 1998.
- [72022] 10352.Dammann, R.; Li, C.; Yoon, J.–H.; Chin, P. L.; Bates, S.; Pfeifer,G. P.: Epigenetic inactivation of a RAS association domain familyprotein from the lung tumour suppressor locus 3p21.3. *Nature Genet.* 25:315–319, 2000.
- [72023] 10353.Dreijerink, K.; Braga, E.; Kuzmin, I.; Geil, L.; Duh, F.–M.; Angeloni,D.; Zbar, B.; Lerman, M. I.; Stanbridge, E. J.; Minna, J. D.; Protopopov,A.; Li, J.; Kashuba, V.; Klein, G.;

Zabarovsky, E. R.: The candidate tumor suppressor gene, RASSF1A, from human chromosome 3p21.3 is involved in kidney tumorigenesis. *Proc. Nat. Acad. Sci.* 98: 7504–7509, 2001.

[72024] 10354. Harada, K.; Toyooka, S.; Maitra, A.; Maruyama, R.; Toyooka, K.O.; Timmons, C. F.; Tomlinson, G. E.; Masetrangelo, D.; Hay, R. J.; Minna, J. D.; Gazdar, A. F.: Aberrant promoter methylation and silencing of the RASSF1A gene in pediatric tumors and cell lines. *Oncogene* 21:4345–4349, 2002.

[72025] 10355. Sekido, Y.; Ahmadian, M.; Wistuba, I. I.; Latif, F.; Bader, S.; Wei, M.-H.; Duh, F.-M.; Gazdar, A. F.; Lerman, M. I.; Minna, J. D.: Cloning of a breast cancer homozygous deletion junction narrows the region of search for a 3p21.3 tumor suppressor gene. *Oncogene* 16:3151–3157, 1998.

[72026] 10356. Dann, C. E.; Hsieh, J.-C.; Rattner, A.; Sharma, D.; Nathans, J.; Leahy, D. J.: Insights into Wnt binding and signalling from the structures of two Frizzled cysteine-rich domains. *Nature* 412: 86–90, 2001.

[72027] 10357. Hoang, B.; Moos, M., Jr.; Vukicevic, S.; Luyten, F. P.: Primary structure and tissue distribution of FRZB, a novel protein related to *Drosophila* frizzled, suggest a role in skeletal morphogenesis. *J. Biol. Chem.* 271: 26131–26137,

1996.

- [72028] 10358.Leyns, L.; Bouwmeester, T.; Kim, S.-H.; Piccolo, S.; De Robertis, E. M.: Frzb-1 is a secreted antagonist of Wnt signaling expressed in the Spemann organizer. *Cell* 88: 747-756, 1997.
- [72029] 10359.Schumann, H.; Holtz, J.; Zerkowski, H.-R.; Hatzfeld, M.: Expression of secreted frizzled related proteins 3 and 4 in human ventricular myocardium correlates with apoptosis related gene expression. *Cardiovasc.Res.* 45: 720-728, 2000.
- [72030] 10360.Bouchon, A.; Dietrich, J.; Colonna, M.: Cutting edge: inflammatory responses can be triggered by TREM-1, a novel receptor expressed on neutrophils and monocytes. *J. Immun.* 164: 4991-4995, 2000.
- [72031] 10361.Bouchon, A.; Facchetti, F.; Weigand, M. A.; Colonna, M.: TREM-1 amplifies inflammation and is a crucial mediator of septic shock. *Nature* 410:1103-1107, 2001.
- [72032] 10362.Yu, Y.; Xu, F.; Peng, H.; Fang, X.; Zhao, S.; Li, Y.; Cuevas, B.; Kuo, W.-L.; Gray, J. W.; Siciliano, M.; Mills, G. B.; Bast, R. C., Jr.: NOEY2 (ARHI), an imprinted putative tumor suppressor gene in ovarian and breast carcinomas. *Proc. Nat. Acad. Sci.* 96: 214-219, 1999.

- [72033] 10363.Walowsky, C.; Fitzhugh, D. J.; Castano, I. B.; Ju, J. Y.; Levin, N. A.; Christman, M. F.: The topoisomerase-related function gene TRF4 affects cellular sensitivity to the antitumor agent camptothecin. *J. Biol. Chem.* 274: 7302–7308, 1999.
- [72034] 10364.Wang, Z.; Castano, I. B.; De Las Penas, A.; Adams, C.; Christman, M. F.: Pol kappa: a DNA polymerase required for sister chromatid cohesion. *Science* 289: 774–779, 2000.
- [72035] 10365.Clark, J.; Lu, Y.-J.; Sidhar, S. K.; Parker, C.; Gill, S.; Smedley, D.; Hamoudi, R.; Linehan, W. M.; Shipley, J.; Cooper, C. S.: Fusion of splicing factor genes PSF and NonO (p54-nrb) to the TFE3 gene in papillary renal cell carcinoma. *Oncogene* 15: 2233–2239, 1997.
- [72036] 10366.Patton, J. G.; Porro, E. B.; Galceran, J.; Tempst, P.; Nadal-Ginard, B.: Cloning and characterization of PSF, a novel pre-mRNA splicing factor. *Genes Dev.* 7: 393–406, 1993.
- [72037] 10367.Piquemal, D.; Joulia, D.; Balaguer, P.; Basset, A.; Marti, J.; Combes, T.: Differential expression of the RTP/Drg1/Ndr1 gene product in proliferating and growth arrested cells. *Biochim. Biophys. Acta* 1450:364–373, 1999.
- [72038] 10368.LeClerc, S.; Palaniswami, R.; Xie, B.; Govdan, M. V.:

Molecularcloning and characterization of a factor that binds the human glucocorticoidreceptor gene and represses its expression. J. Biol. Chem. 266:17333–17340, 1991.

[72039] 10369.Tikoo, A.; Czekay, S.; Viars, C.; White, S.; Heath, J. K.; Arden,K.; Maruta, H.: p190–A, a human tumor suppressor gene, maps to thechromosomal region 19q13.3 that is reportedly deleted in some gliomas. Gene 257:23–31, 2000.

[72040] 10370.Boccaccio, I.; Glatt–Deeley, H.; Watrin, F.; Roeckel, N.; Lalande,M.; Muscatelli, F.: The human MAGEL2 gene and its mouse homologueare paternally expressed and mapped to the Prader–Willi region. Hum.Molec. Genet. 8: 2497–2505, 1999.

[72041] 10371.Lee, S.; Kozlov, S.; Hernandez, L.; Chamberlain, S. J.; Brannan,C. I.; Stewart, C. L.; Wevrick, R.: Expression and imprinting ofMAGEL2 suggest a role in Prader–Willi syndrome and the homologousmurine imprinting phenotype. Hum. Molec. Genet. 9: 1813–1819, 2000.

[72042] 10372.Wang, A. H.; Bertos, N. R.; Vezmar, M.; Pelletier, N.; Crosato,M.; Heng, H. H.; Th'ng, J.; Han, J.; Yang, X.–J.: HDAC4, a humanhistone deacetylase related to yeast HDA1, is a transcriptional corepressor. Molec.Cell. Biol.

19: 7816–7827, 1999.

- [72043] 10373. Nishimura, Y.; Hayashi, M.; Inada, H.; Tanaka, T.: Molecular cloning and characterization of mammalian homologues of vesicle-associated membrane protein-associated (VAMP-associated) proteins. *Biochem. Biophys. Res. Commun.* 254: 21–26, 1999.
- [72044] 10374. Weir, M. L.; Klip, A.; Trimble, W. S.: Identification of a human homologue of the vesicle-associated membrane protein (VAMP)-associated protein of 33 kDa (VAP-33): a broadly expressed protein that binds to VAMP. *Biochem. J.* 333: 247–251, 1998.
- [72045] 10375. Perez Jurado, L. A.; Wang, Y.-K.; Francke, U.; Cruces, J.: TBL2, a novel transducin family member in the WBS deletion: characterization of the complete sequence, genomic structure, transcriptional variants and the mouse ortholog. *Cytogenet. Cell Genet.* 86: 277–284, 1999.
- [72046] 10376. Akhmanova, A.; Hoogenraad, C. C.; Drabek, K.; Stepanova, T.; Dortland, B.; Verkerk, T.; Vermeulen, W.; Burgering, B. M.; De Zeeuw, C. I.; Grosveld, F.; Galjart, N.: CLASPs are CLIP-115 and -170 associating proteins involved in the regional regulation of microtubule dynamics in motile fibroblasts. *Cell* 104: 923–935, 2001.
- [72047] 10377. Cattanach, B. M.; Barr, J. A.; Beechey, C. V.; Martin,

J.; Noebels, J.; Jones, J.: A candidate model for Angelman syndrome in the mouse. *Mammalian Genome* 8: 472–478, 1997.

[72048] 10378. Dhar, M.; Webb, L. S.; Smith, L.; Hauser, L.; Johnson, D.; West, D. B.: A novel ATPase on mouse chromosome 7 is a candidate gene for increased body fat. *Physiol. Genomics* 4: 93–100, 2000.

[72049] 10379. Halleck, M. S.; Lawler, J. F., Jr.; Blackshaw, S.; Gao, L.; Nagarajan, P.; Hacker, C.; Pyle, S.; Newman, J. T.; Nakanishi, Y.; Ando, H.; Weinstock, D.; Williamson, P.; Schlegel, R. A.: Differential expression of putative transbilayer amphipath transporters. *Physiol. Genomics* 1: 139–150, 1999.

[72050] 10380. Fitzgibbon, J.; Neat, M. J.; Foot, N.; Hill, A. S.; Lister, T. A.; Gupta, R. K.: Assignment of brain acid-soluble protein 1 (BASP1) to human chromosome 5p15.1–p14, differential expression in human cancer cell lines as a result of alterations in gene dosage. *Cytogenet. Cell Genet.* 89: 147–149, 2000.

[72051] 10381. Mosevitsky, M. I.; Capony, J. P.; Skladchikova, G. Y.; Novitskaya, V. A.; Plekhanov, A. Y.; Zakharov, V. V.: The BASP1 family of myristoylated proteins abundant in axonal termini: primary structure analysis and physico-chemical



properties. *Biochimie* 79: 373–384, 1997.

[72052] 10382.Park, S.; Kim, Y.-I.; Kim, B.; Seong, C.; Oh, Y.; Baek, K.; Yoon,J.: Characterization of bovine and human cDNAs encoding NAP-22 (22kDa neuronal tissue-enriched acidic protein) homologs. *Molec. Cells* 8:471–477, 1998.

[72053] 10383.Shichijo, S.; Nakao, M.; Imai, Y.; Takasu, H.; Kawamoto, M.; Niiya,F.; Yang, D.; Toh, Y.; Yamana, H.; Itoh, K.: A gene encoding antigenicpeptides of human squamous cell carcinoma recognized by cytotoxicT lymphocytes. *J. Exp. Med.* 187: 277–288, 1998.

[72054] 10384.Valenta, R.; Natter, S.; Seiberler, S.; Wichlas, S.; Maurer, D.;Hess, M.; Pavelka, M.; Grote, M.; Ferreira, F.; Szepfalusi, Z.; Valent,P.; Stingl, G.: Molecular characterization of an autoallergen, Homs 1, identified by serum IgE from atopic dermatitis patients. *J.Invest. Derm.* 111: 1178–1183, 1998.

[72055] 10385.Loftus, B. J.; Kim, U.-J.; Sneddon, V. P.; Kalush, F.; Brandon,R.; Fuhrmann, J.; Mason, T.; Crosby, M. L.; Barnstead, M.; Cronin,L.; Mays, A. D.; Cao, Y.; Xu, R. X.; Kang, H.-L.; Mitchell, S.; Eichler,E. E.; Harris, P. C.; Venter, J. C.; Adams, M. D.: Genome duplicationsand other features in 12 Mb of DNA sequence from human chromosome16p and 16q. *Genomics* 60: 295–308, 1999.

- [72056] 10386.Brauner–Osborne, H.; Krogsgaard–Larsen, P.: Sequence and expression pattern of a novel human orphan G–protein–coupled receptor, GPRC5B, a family C receptor with a short amino–terminal domain. *Genomics* 65:121–128, 2000.
- [72057] 10387.Robbins, M. J.; Michalovich, D.; Hill, J.; Calver, A. R.; Medhurst, A. D.; Gloger, I.; Sims, M.; Middlemiss, D. N.; Pangalos, M. N.: Molecular cloning and characterization of two novel retinoic acid–inducible orphan G–protein–coupled receptors (GPRC5B and GPRC5C). *Genomics* 67:8–18, 2000.
- [72058] 10388.Autieri, M. V.; Feuerstein, G. Z.; Yue, T.–L.; Ohlstein, E. H.; Douglas, S. A.: Use of differential display to identify differentially expressed mRNAs induced by rat carotid artery balloon angioplasty. *Lab. Invest.* 72: 656–661, 1995.
- [72059] 10389.Autieri, M. V.; Haines, D. S.; Romanic, A. M.; Ohlstein, E. H.: Expression of 14–3–3–gamma in injured arteries and growth factor– and cytokine–stimulated human vascular smooth muscle cells. *Cell Growth Diff.* 7: 1453–1460, 1996.
- [72060] 10390.Horie, M.; Suzuki, M.; Takahashi, E.; Tanigami, A.: Cloning, expression, and chromosomal mapping of the

human 14-3-3gamma gene (YWHAG) to 7q11.23. Genomics 60:241-243, 1999.

[72061] 10391.Morrison, D.: 14-3-3: modulators of signaling proteins? Science 266:56-57, 1994.

[72062] 10392.Brazas, R.; Ganem, D.: A cellular homolog of hepatitis delta antigen:implications for viral replication and evolution. Science 274: 90-94,1996.

[72063] 10393.Long, M.; de Souza, S. J.; Gilbert, W.: Delta-interacting proteinA and the origin of hepatitis delta antigen. Science 276: 824-825,1997.

[72064] 10394.Robertson, H. D.: How did replicating and coding RNAs first gettogether? Science 274: 66-67, 1996.

[72065] 10395.Asada, H.; Kawamura, Y.; Maruyama, K.; Kume, H.; Ding, R.-G.; Kanbara,N.; Kuzume, H.; Sanbo, M.; Yagi, T.; Obata, K.: Cleft palate anddecreased brain gamma-aminobutyric acid in mice lacking the 67-kDaisoform of glutamic acid decarboxylase. Proc. Nat. Acad. Sci. 94:6496-6499, 1997.

[72066] 10396.Brilliant, M. H.; Szabo, G.; Katarova, Z.; Kozak, C. A.; Glaser,T. M.; Greenspan, R. J.; Housman, D. E.: Sequences homologous toglutamic acid decarboxylase cDNA are present on mouse chromosomes2 and 10. Genomics 6: 115-122, 1990.

- [72067] 10397.Bu, D.-F.; Tobin, A. J.: The exon-intron organization of the genes(GAD1 and GAD2) encoding two human glutamate decarboxylases (GAD-67and GAD-65) suggests that they derive from a common ancestral GAD. *Genomics* 21:222-228, 1994.
- [72068] 10398.Condie, B. G.; Bain, G.; Gottlieb, D. I.; Capecchi, M. R.: Cleftpalate in mice with a targeted mutation in the gamma-aminobutyricacid-producing enzyme glutamic acid decarboxylase 67. *Proc. Nat.Acad. Sci.* 94: 11451-11455, 1997.
- [72069] 10399.Erlander, M. G.; Tillakaratne, N. J. K.; Feldblum, S.; Patel, N.;Tobin, A. J.: Two genes encode distinct glutamate decarboxylases. *Neuron* 7:91-100, 1991.
- [72070] 10400.Kelly, C. D.; Edwards, Y.; Johnstone, A. P.; Harfst, E.; Nogradi,A.; Nussey, S. S.; Povey, S.; Carter, N. D.: Nucleotide sequenceand chromosomal assignment of a cDNA encoding the large isoform ofhuman glutamate decarboxylase. *Ann. Hum. Genet.* 56: 255-265, 1992.
- [72071] 10401.Krishnamoorthy, K. S.: Pyridoxine-dependency seizure: reportof a rare presentation. *Ann. Neurol.* 13: 103-104, 1983.
- [72072] 10402.Sparkes, R. S.; Kaufman, D. L.; Heinzmann, C.; Tobin, A. J.; Mohandas,T.: Brain glutamate decarboxylase

(GAD) gene assigned to human chromosome2 by somatic cell hybrid analysis. (Abstract) Cytogenet. Cell Genet. 46:696 only, 1987.

- [72073] 10403.Kedra, D.; Seroussi, E.; Fransson, I.; Trifunovic, J.; Clark, M.; Lagercrantz, J.; Blennow, E.; Mehlin, H.; Duman-ski, J.: The germinalcenter kinase gene and a novel CDC25-like gene are located in the vicinity of the PYGM gene on 11q13. Hum. Genet. 100: 611-619, 1997.
- [72074] 10404.Hirsch, D. S.; Pirone, D. M.; Burbelo, P. D.: A new family of Cdc42 effector proteins, CEPs, function in fibroblast and epithelial cell shape changes. J. Biol. Chem. 276: 875-883, 2001.
- [72075] 10405.Joberty, G.; Perlungher, R. R.; Macara, I. G.: The Borgs, a new family of Cdc42 and TC10 GTPase-interacting proteins. Molec. Cell. Biol. 19: 6585-6597, 1999.
- [72076] 10406.Yang, Z.-Q.; Imoto, I.; Fukuda, Y.; Pimkhaokham, A.; Shimada, Y.; Imamura, M.; Sugano, S.; Nakamura, Y.; Inazawa, J.: Identification of a novel gene, GASC1, within an amplicon at 9p23-24 frequently detected in esophageal cancer cell lines. Cancer Res. 60: 4735-4739, 2000.
- [72077] 10407.LeCouter, J.; Kowalski, J.; Foster, J.; Hass, P.; Zhang, Z.; Dillard-Telm, L.; Frantz, G.; Rangell, L.; DeGuzman, L.; Keller, G.-A.; Peale, F.; Gurney, P.; Hillan, K. J.;

Ferrara, N.: Identification of an angiogenic mitogen selective for endocrine gland endothelium. *Nature* 412: 877–884, 2001.

- [72078] 10408. Li, M.; Bullock, C. M.; Knauer, D. J.; Ehlert, F. J.; Zhou, Q. Y.: Identification of two prokineticin cDNAs: recombinant proteins potently contract gastrointestinal smooth muscle. *Molec. Pharm.* 59:692–698, 2001.
- [72079] 10409. Sun, Q.-A.; Wu, Y.; Zappacosta, F.; Jeang, K.-T.; Lee, B. J.; Hatfield, D. L.; Gladyshev, V. N.: Redox regulation of cell signaling by selenocysteine in mammalian thioredoxin reductases. *J. Biol. Chem.* 274: 24522–24530, 1999.
- [72080] 10410. Nagase, T.; Nakayama, M.; Nakajima, D.; Kikuno, R.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XX. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.* 8: 85–95, 2001.
- [72081] 10411. Telford, E. A. R.; Moynihan, L. M.; Markham, A. F.; Lench, N. J.: Isolation and characterisation of a cDNA encoding the precursor for a novel member of the acyl-CoA dehydrogenase gene family. *Biochim. Biophys. Acta* 1446: 371–376, 1999.
- [72082] 10412. Jaquemar, D.; Schenker, T.; Trueb, B.: An ankyrin-

like protein with transmembrane domains is specifically lost after oncogenic transformation of human fibroblasts. J. Biol. Chem. 274: 7325–7333, 1999.

- [72083] 10413. Schenker, T.; Trueb, B.: Down-regulated proteins of mesenchymal tumor cells. Exp. Cell Res. 239: 161–168, 1998.
- [72084] 10414. Cerretti, D. P.; DuBose, R. F.; Black, R. A.; Nelson, N.: Isolation of two novel metalloproteinase-disintegrin (ADAM) cDNAs that show testis-specific gene expression. Biochem. Biophys. Res. Commun. 263: 810–815, 1999.
- [72085] 10415. Xu, R.; Cai, J.; Xu, T.; Zhou, W.; Ying, B.; Deng, K.; Zhao, S.; Li, C.: Molecular cloning and mapping of a novel ADAM gene (ADAM29) to human chromosome 4. Genomics 62: 537–539, 1999.
- [72086] 10416. Jacobs, S.; Schilf, C.; Fliegert, F.; Koling, S.; Weber, Y.; Schurmann, A.; Joost, H.-G.: ADP-ribosylation factor (ARF)-like 4, 6, and 7 represent a subgroup of the ARF family characterized by rapid nucleotide exchange and a nuclear localization signal. FEBS Lett. 456: 384–388, 1999.
- [72087] 10417. Lako, M.; Lindsay, S.; Bullen, P.; Wilson, D. I.; Robson, S. C.; Strachan, T.: A novel mammalian Wnt gene, WNT8B, shows brain-restricted expression in early development, with sharply delimited expression boundaries in

the developing forebrain. Hum. Molec. Genet. 7:  
813–822,1998.

- [72088] 10418.Lako, M.; Strachan, T.; Curtis, A. R. J.; Lindsay, S.: Isolationand characterization of WNT8B, a novel human Wnt gene that maps to10q24. Genomics 35: 386–388, 1996.
- [72089] 10419.Toda, T.; Iida, A.; Miwa, T.; Nakamura, Y.; Imai, T.: Isolationand characterization of a novel gene encoding nuclear protein at alocus (D11S636) tightly linked to multiple endocrine neoplasia type1 (MEN1). Hum. Molec. Genet. 3: 465–470, 1994.
- [72090] 10420.Nusse, R.; Varmus, H. E.: Many tumors induced by the mouse mammarytumor virus contain a provirus integrated in the same region of thehost genome. Cell 31: 99–109, 1982.
- [72091] 10421.Scholler, J. K.; Kanner, S. B.: The human p167 gene encodes aunique structural protein that contains centrosomin A homology andassociates with a multicomponent complex. DNA Cell Biol. 16: 515–531,1997.
- [72092] 10422.Abdulkadir, S. A.; Magee, J. A.; Peters, T. J.; Kaleem, Z.; Naughton,C. K.; Humphrey, P. A.; Milbrandt, J.: Conditional loss of Nkx3.1in adult mice induces prostatic intraepithelial neoplasia. Molec.Cell. Biol. 22: 1495–1503,



2002.

- [72093] 10423.Bieberich, C. J.; Fujita, K.; He, W. W.; Jay, G.: Prostate-specific and androgen-dependent expression of a novel homeobox gene. *J. Biol.Chem.* 271: 31779–31782, 1996.
- [72094] 10424.Korkmaz, K. S.; Korkmaz, C. G; Ragnhildstveit, E.; Kizildag, S.;Pretlow, T. G.; Saatcioglu, F.: Full-length cDNA sequence and genomic organization of human NKX3A--alternative forms and regulation by both androgens and estrogens. *Gene* 260: 25–36, 2000.
- [72095] 10425.He, W. W.; Sciavolino, P. J.; Wing, J.; Augustus, M.; Hudson, P.;Meissner, P. S.; Curtis, R. T.; Shell, B. K.; Bostwick, D. G.; Tindall,D. J.; Gelmann, E. P.; Abate-Shen, C.; Carter, K. C.: A novel human prostate-specific, androgen-regulated homeobox gene (NKX3.1) that maps to 8p21, a region frequently deleted in prostate cancer. *Genomics* 43:69–77, 1997.
- [72096] 10426.Argyropoulos, G.; Brown, A. M.; Willi, S. M.; Zhu, J.; He, Y.;Reitman, M.; Geva, S. M.; Spruill, I.; Garvey, W. T.: Effects of mutations in the human uncoupling protein 3 gene on the respiratory quotient and fat oxidation in severe obesity and type 2 diabetes. *J.Clin. Invest.* 102: 1345–1351, 1998.

- [72097] 10427.Boss, O.; Giacobino, J.-P.; Muzzin, P.: Genomic structure of uncouplingprotein-3 (UCP3) and its assignment to chromosome 11q13. *Genomics* 47:425-426, 1998.
- [72098] 10428.Boss, O.; Samec, S.; Paoloni-Giacobino, A.; Rossier, C.; Dulloo,A.; Seydoux, J.; Muzzin, P.; Giacobino, J.-P.: Uncoupling protein-3:a new member of the mitochondrial carrier family with tissue-specificexpression. *FEBS Lett.* 408: 39-42, 1997.
- [72099] 10429.Brown, A. M.; Dolan, J. W.; Willi, S. M.; Garvey, W. T.; Argyropoulos,G.: Endogenous mutations in human uncoupling protein 3 alter itsfunctional properties. *FEBS Lett.* 464: 189-193, 1999.
- [72100] 10430.Bilaud, T.; Brun, C.; Ancelin, K.; Koering, C. E.; Laroche, T.;Gilson, E.: Telomeric localization of TRF2, a novel human teloboxprotein. *Nature Genet.* 17: 236-239, 1997.
- [72101] 10431.Broccoli, D.; Smogorzewska, A.; Chong, L.; de Lange, T.: Humantelomeres contain two distinct Myb-related proteins, TRF1 and TRF2. *NatureGenet.* 17: 231-235, 1997.
- [72102] 10432.Karlseder, J.; Smogorzewska, A.; de Lange, T.: Senescence inducedby altered telomere state, not telom-

ere loss. Science 295: 2446–2449,2002.

- [72103] 10433.Sakaguchi, A. Y.; Padalecki, S. S.; Mattern, V.; Rodriguez, A.;Leach, R. J.; McGill, J. R.; Chavez, M.; Giamb Bernardi, T. A.: Chromosomal sublocalization of the transcribed human telomere repeat binding factor 2 gene and comparative mapping in the mouse. Somat. Cell Molec. Genet. 24:157–163, 1998.
- [72104] 10434.van Steensel, B.; Smogorzewska, A.; de Lange, T.: TRF2 protects human telomeres from end-to-end fusions. Cell 92: 401–413, 1998.
- [72105] 10435.Parr, B. A.; McMahon, A. P.: Sexually dimorphic development of the mammalian reproductive tract requires Wnt-7a. Nature 395: 707–710,1998.
- [72106] 10436.Barron-Casella, E. A.; Torres, M. A.; Scherer, S. W.; Heng, H.H. Q.; Tsui, L.-C.; Casella, J. F.: Sequence analysis and chromosomal localization of human Cap Z: conserved residues within the actin-binding domain may link Cap Z to gelsolin/severin and profilin protein families. J.Biol. Chem. 270: 21472–21479, 1995.
- [72107] 10437.Hart, M. C.; Korshunova, Y. O.; Cooper, J. A.: Mapping of the mouse actin capping protein alpha subunit genes and pseudogenes. Genomics 39:264–270, 1997.
- [72108] 10438.Cooper, J. A.: Personal Communication. St. Louis,

Mo. 3/8/1999.

- [72109] 10439. Cardoso, C.; Mignon, C.; Hetet, G.; Grandchamps, B.; Fontes, M.; Colleaux, L.: The human EZH2 gene: genomic organisation and revised mapping in 7q35 within the critical region for malignant myeloid disorders. *Eur. J. Hum. Genet.* 8: 174–180, 2000.
- [72110] 10440. Chen, H.; Rossier, C.; Antonarakis, S. E.: Cloning of a human homolog of the *Drosophila* enhancer of zeste gene (EZH2) that maps to chromosome 21q22.2. *Genomics* 38: 30–37, 1996.
- [72111] 10441. Dohner, K.; Brown, J.; Hehmann, U.; Hetzel, C.; Stewart, J.; Lowther, G.; Scholl, C.; Frohling, S.; Cuneo, A.; Tsui, L. C.; Lichter, P.; Scherer, S. W.; Dohner, H.: Molecular cytogenetic characterization of a critical region in bands 7q35–q36 commonly deleted in malignant myeloid disorders. *Blood* 92: 4031–4035, 1998.
- [72112] 10442. Laible, G.; Haynes, A. R.; Lebersorger, A.; O'Carroll, D.; Mattei, M.-G.; Denny, P.; Brown, S. D. M.; Jenuwein, T.: The murine polycomb-group genes *Ezh1* and *Ezh2* map close to Hox gene clusters on mouse chromosomes 11 and 6. *Mammalian Genome* 10: 311–314, 1999.
- [72113] 10443. Varambally, S.; Dhanasekaran, S. M.; Zhou, M.; Barrette, T. R.; Kumar-Sinha, C.; Sanda, M. G.; Ghosh, D.;

Pienta, K. J.; Sewalt, R.G. A. B.; Otte, A. P.; Rubin, M. A.; Chinnaiyan, A. M.: The polycombgroup protein EZH2 is involved in progression of prostate cancer. *Nature* 419:624–629, 2002.

[72114] 10444.Morohoshi, F.; Arai, K.; Takahashi, E.; Tanigami, A.; Ohki, M.: Cloning and mapping of a human RBP56 gene encoding a putative RNA binding protein similar to FUS/TLS and EWS proteins. *Genomics* 38:51–57, 1996.

[72115] 10445.Batt, J.; Asa, S.; Fladd, C.; Rotin, D.: Pituitary, pancreatic and gut neuroendocrine defects in protein tyrosine phosphatase-sigma-deficient mice. *Molec. Endocr.* 16: 155–169, 2002.

[72116] 10446.Elchebly, M.; Wagner, J.; Kennedy, T. E.; Lanctot, C.; Michaliszyn, E.; Itie, A.; Drouin, J.; Tremblay, M. L.: Neuroendocrine dysplasia in mice lacking protein tyrosine phosphatase sigma. *Nature Genet.* 21:330–333, 1999.

[72117] 10447.Pulido, R.; Serra-Pages, C.; Tang, M.; Streuli, M.: The LAR/PTPdelta/PTP sigma subfamily of transmembrane protein-tyrosine-phosphatases: multiple human LAR, PTP delta, and PTP sigma isoforms are expressed in a tissue-specific manner and associate with the LAR-interacting protein LIP.1. *Proc. Nat. Acad. Sci.* 92: 11686–11690, 1995.

- [72118] 10448.Wagner, J.; Gordon, L. A.; Heng, H. H. Q.; Tremblay, M. L.; Olsen,A. S.: Physical mapping of receptor type protein tyrosine phosphatasesigma (PTPRS) to human chromosome 19p13.3. *Genomics* 38: 76–78, 1996.
- [72119] 10449.Wallace, M. J.; Batt, J.; Fladd, C. A.; Henderson, J. T.; Skarnes,W.; Rotin, D.: Neuronal defects and posterior pituitary hypoplasia in mice lacking the receptor tyrosine phosphatase PTP-sigma. *NatureGenet.* 21: 334–338, 1999.
- [72120] 10450.Bruyns, E.; Mincheva, A.; Bruyns, R. M.; Kirchgessner, H.; Weitz,S.; Lichter, P.; Meuer, S.; Schraven, B.: Sequence, genomic organization, and chromosomal localization of the human LPAP (PTPRCAP) and mouseCD45-AP/LSM-1 genes. *Genomics* 38: 79–83, 1996.
- [72121] 10451.Schraven, B.; Schoenhaut, D.; Bruyns, E.; Koretzky, G.; Eckerskorn,C.; Wallich, R.; Kirchgessner, H.; Sakorafas, P.; Labkovsky, B.; Ratnofsky,S.; Meuer, S.: LPAP, a novel 32-kDa phosphoprotein that interacts with CD45 in human lymphocytes. *J. Biol. Chem.* 269: 29102–29111, 1994.
- [72122] 10452.Spicer, A. P.; Augustine, M. L.; McDonald, J. A.: Molecular cloning and characterization of a putative mouse hyaluronan synthase. *J.Biol. Chem.* 271: 23400–23406,

1996.

- [72123] 10453.Skalhegg, B. S.; Huang, Y.; Su, T.; Idzerda, R. L.; McKnight, G.S.; Burton, K. A.: Mutation of the C-alpha subunit of PKA leads to growth retardation and sperm dysfunction. *Molec. Endocr.* 16: 630-639, 2002.
- [72124] 10454.Tasken, K.; Solberg, R.; Zhao, Y.; Hansson, V.; Jahnsen, T.; Siciliano, M. J.: The gene encoding the catalytic subunit C-alpha of cAMP-dependent protein kinase (locus PRKACA) localizes to human chromosome region 19p13.1. *Genomics* 36: 535-538, 1996.
- [72125] 10455.Kim, J.; Uyemura, K.; Van Dyke, M. K.; Legaspi, A. J.; Rea, T.H.; Shuai, K.; Modlin, R. L.: A role for IL-12 receptor expression and signal transduction in host defense in leprosy. *J. Immun.* 167:779-786, 2001.
- [72126] 10456.Morton, S. M.; Bocaccio, I.; Depetris, D.; Mattei, M.; Dessein, A.: Assignment of IL12RB2 to human chromosome 1p31.3-p31.2 between D1S230 and D1S198. *Cytogenet. Cell Genet.* 79: 282-283, 1997.
- [72127] 10457.McCright, B.; Brothman, A. R.; Virshup, D. M.: Assignment of human protein phosphatase 2A regulatory subunit genes B56-alpha, B56-beta, B56-gamma, B56-delta, and B56-epsilon (PPP2R5A--PPP2R5E), highly expressed in muscle and brain, to chromosome regions

1q41, 11q12, 3p21, 6p21.1, and 7p11.2-to-p12. *Genomics* 36: 168–170, 1996.

[72128] 10458. McCright, B.; Virshup, D. M.: Identification of a new family of protein phosphatase 2A regulatory subunits. *J. Biol. Chem.* 270:26123–26128, 1995.

[72129] 10459. Horikawa, S.; Tsukada, K.: Molecular cloning and developmental expression of a human kidney S-adenosylmethionine synthetase. *FEBS Lett.* 312: 37–41, 1992.

[72130] 10460. Mao, Z.; Liu, S.; Cai, J.; Huang, Z.-Z.; Lu, S. C.: Cloning and functional characterization of the 5-prime-flanking region of human methionine adenosyltransferase 2A gene. *Biochem. Biophys. Res. Commun.* 248:479–484, 1998.

[72131] 10461. Clark, J. A.; Bonner, T. I.; Kim, A. S.; Usdin, T. B.: Multiple regions of ligand discrimination revealed by analysis of chimeric parathyroid hormone 2 (PTH2) and PTH/PTH-related peptide (PTHrP) receptors. *Molec. Endocr.* 12: 193–206, 1998.

[72132] 10462. Usdin, T. B.; Gruber, C.; Bonner, T. I.: Identification and functional expression of a receptor selectively recognizing parathyroid hormone, the PTH2 receptor. *J. Biol. Chem.* 270: 15455–15458, 1995.



- [72133] 10463.Usdin, T. B.; Modi, W.; Bonner, T. I.: Assignment of the humanPTH2 receptor gene (PTHR2) to chromosome 2q33 by fluorescence in situhybridization. *Genomics* 37: 140–141, 1996.
- [72134] 10464.Faure, S.; Meyer, L.; Costagliola, D.; Vaneensberghe, C.; Genin,E.; Autran, B.; French ALT and IM-MUNOCO Study Groups; Delfraisay,J.–F.; SEROCO Study Group; McDermott, D. H.; Murphy, P. M.; Debre,P.; Theodorou, I.; Cambadiere, C.: Rapid progression to AIDS in HIV+individuals with a structural variant of the chemokine receptor CX(3)CR1. *Science* 287:2274–2277, 2000.
- [72135] 10465.Imai, T.; Hieshima, K.; Haskell, C.; Baba, M.; Nagira, M.; Nishimura,M.; Kakizaki, M.; Takagi, S.; Nomiyama, H.; Schall, T. J.; Yoshie,O.: Identification and molecular characterization of fractalkinereceptor CX3CR1, which mediates both leukocyte migration and adhesion. *Cell* 91:521–530, 1997.
- [72136] 10466.Moatti, D.; Faure, S.; Fumeron, F.; Amara, M. E. W.; Seknadji,P.; McDermott, D. H.; Debre, P.; Aumont, M. C.; Murphy, P. M.; deProst, D.; Combadiere, C.: Polymorphism in the fractalkine receptorCX3CR1 as a genetic risk factor for coronary artery disease. *Blood* 97:1925–1928, 2001.

- [72137] 10467. Raport, C. J.; Schweickart, V. L.; Eddy, R. L., Jr.; Shows, T. B.; Gray, P. W.: The orphan G-protein-coupled receptor-encoding gene V28 is closely related to genes for chemokine receptors and is expressed in lymphoid and neural tissues. *Gene* 163: 295–299, 1995.
- [72138] 10468. Tripp, R. A.; Jones, L. P.; Haynes, L. M.; Zheng, H.; Murphy, P. M.; Anderson, L. J.: CX3C chemokine mimicry by respiratory syncytial virus G glycoprotein. *Nature Immun.* 2: 732–738, 2001.
- [72139] 10469. Lafreniere, R. G.; Rochefort, D. L.; Chretien, N.; Neville, C. E.; Korneluk, R. G.; Zuo, L.; Wei, Y.; Lichter, J.; Rouleau, G. A.: Isolation and genomic structure of a human homolog of the yeast periodic tryptophan protein 2 (PWP2) gene mapping to 21q22.3. *Genome Res.* 6: 1216–1226, 1996.
- [72140] 10470. Lalioti, M. D.; Chen, H.; Rossier, C.; Shafaatian, R.; Reid, J. D.; Antonarakis, S. E.: Cloning the cDNA of human PWP2, which encodes a protein with WD repeats and maps to 21q22.3. *Genomics* 35: 321–327, 1996.
- [72141] 10471. Yamakawa, K.; Gao, D.-Q.; Korenberg, J. R.: A periodic tryptophan protein 2 gene homologue (PWP2H) in the candidate region of progressive myoclonus epilepsy on 21q22.3. *Cytogenet. Cell Genet.* 74: 140–145, 1996.

- [72142] 10472.Hanes, J.; von der Kammer, H.; Klaudiny, J.; Scheit, K. H.: Characterization by cDNA cloning of two new human protein kinases: evidence by sequence comparison of a new family of mammalian protein kinases. *J. Molec.Biol.* 244: 665–672, 1994.
- [72143] 10473.Talmadge, C. B.; Finkernagel, S.; Sumegi, J.; Sciorra, L.; Rabinow, L.: Chromosomal mapping of three human LAMMER protein-kinase-encoding genes. *Hum. Genet.* 103: 523–524, 1998.
- [72144] 10474.Andersen, G.; Poterszman, A.; Egly, J. M.; Moras, D.; Thierry, J.-C: The crystal structure of human cyclin H. *FEBS Lett.* 397:65–69, 1996.
- [72145] 10475.Eki, T.; Okumura, K.; Abe, M.; Kagotani, K.; Taguchi, H.; Murakami, Y.; Pan, Z.-Q.; Hanaoka, F.: Mapping of the human genes encoding cyclin H (CCNH) and the CDK-activating kinase (CAK) assembly factor MAT1 (MNAT1) to chromosome bands 5q13.3–q14 and 14q23, respectively. *Genomics* 47:115–120, 1998.
- [72146] 10476.Fisher, R. P.; Morgan, D. O.: A novel cyclin associates with MO15/CDK7 to form the CDK-activating kinase. *Cell* 78: 713–724, 1994.
- [72147] 10477.Shiekhattar, R.; Mermelstein, F.; Fisher, R. P.; Drapkin, R.; Dynlacht, B.; Wessling, H. C.; Morgan, D. O.; Rein-

berg: Cdk-activating kinase complex is a component of human transcription factor TFIIH. *Nature* 374:283–287, 1995.

- [72148] 10478. Alders, M.; Hodges, M.; Hadjantonakis, A.-K.; Postmus, J.; vanWijk, I.; Blik, J.; de Meulemeester, M.; Westerveld, A.; Guillemot, F.; Oudejans, C.; Little, P.; Mannens, M.: The human Achaete-Scute homologue 2 (ASCL2, HASH2) maps to chromosome 11p15.5, close to IGF2 and is expressed in extravillous trophoblasts. *Hum. Molec. Genet.* 6:859–867, 1997.
- [72149] 10479. Guillemot, F.; Nagy, A.; Auerbach, A.; Rossant, J.; Joyner, A.L.: Essential role of Mash-2 in extraembryonic development. *Nature* 371:333–336, 1994.
- [72150] 10480. Banga, S. S.; Ozer, H. L.; Park, S.-K.; Lee, S.-T.: Assignment of PTK7 encoding a receptor protein tyrosine kinase-like molecule to human chromosome 6p21.1–p12.2 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 76: 43–44, 1997.
- [72151] 10481. Lee, S.-T.; Strunk, K. M.; Spritz, R. A.: A survey of protein tyrosine kinase mRNAs expressed in normal human melanocytes. *Oncogene* 8:3403–3410, 1993.
- [72152] 10482. Mossie, K.; Jallal, B.; Alves, F.; Sures, I.; Plowman, G. D.; Ullrich, A.: Colon carcinoma kinase-4 defines a new

subclass of the receptortyrosine kinase family. *Oncogene* 11: 2179–2184, 1995.

[72153] 10483.Park, S.–K.; Lee, H.–S.; Lee, S.–T.: Characterization of the humanfull–length PTK7 cDNA encoding a receptor protein tyrosine kinase–likemolecule closely related to chick KLG. *J. Biochem.* 119: 235–239,1996.

[72154] 10484.Sotiropoulou, G.; Anisowicz, A.; Sager, R.: Identification, cloning,and characterization of cystatin M, a novel cysteine proteinase inhibitor,down–regulated in breast cancer. *J. Biol. Chem.* 272: 903–910, 1997.

[72155] 10485.Stenman, G.; Astrom, A.–K.; Roijer, E.; Sotiropoulou, G.; Zhang,M.; Sager, R.: Assignment of a novel cysteine proteinase inhibitor(CST6) to 11q13 by fluorescence in situ hybridization. *Cytogenet.Cell Genet.* 76: 45–46, 1997.

[72156] 10486.Kohler, M.; Ansieau, S.; Prehn, S.; Leutz, A.; Haller, H.; Hartmann,E.: Cloning of two novel human importin–alpha subunits and analysisof the expression pattern of the importin–alpha protein family. *FEBSLett.* 417: 104–108, 1997.

[72157] 10487.Moore, M. S.; Blobel, G.: The two steps of nuclear import, targetingto the nuclear envelope and translocation through the nuclear pore,require different cytosolic fac–

tors. Cell 69: 939–950, 1992.

- [72158] 10488. Takeda, S.; Fujiwara, T.; Shimizu, F.; Kawai, A.; Shinomiya, K.; Okuno, S.; Ozaki, K.; Katagiri, T.; Shimada, Y.; Nagata, M.; Watanabe, T.; Takaichi, A.; Kuga, Y.; Suzuki, M.; Hishigaki, H.; Takahashi, E.; Shin, S.; Nakamura, Y.; Hirai, Y.: Isolation and mapping of karyopherin  $\alpha$ -3 (KPNA3), a human gene that is highly homologous to genes encoding *Xenopus* importin, yeast SRP1 and human RCH1. Cytogenet. Cell Genet. 76:87–93, 1997.
- [72159] 10489. Zhang, C.-Y.; Baffy, G.; Perret, P.; Krauss, S.; Peroni, O.; Grujic, D.; Hagen, T.; Vidal-Puig, A.; Boss, O.; Kim, Y.-B.; Zheng, X. X.; Wheeler, M. B.; Shulman, G. I.; Chan, C. B.; Lowell, B. B.: Uncoupling protein-2 negatively regulates insulin secretion and is a major link between obesity, beta cell dysfunction, and type 2 diabetes. Cell 105:745–755, 2001.
- [72160] 10490. Edlund, A.; Johansson, T.; Leidvik, B.; Hansson, L.: Structure of the human kappa-casein gene. Gene 174: 65–69, 1996.
- [72161] 10491. Sprecher, C. A.; Morgenstern, K. A.; Mathewes, S.; Dahlen, J. R.; Schrader, S. K.; Foster, D. C.; Kisiel, W.: Molecular cloning, expression, and partial characterization of two novel members of the ovalbumin family of serine

proteinase inhibitors. J. Biol. Chem. 270:  
29854–29861,1995.

[72162] 10492.Chevalier, D.; Cauffiez, C.; Bernard, C.; Lo-Guidice, J.-M.; Allorge,D.; Fazio, F.; Ferrari, N.; Libersa, C.; Lhermitte, M.; D'Halluin,J.-C.; Broly, F.: Characterization of new mutations in the codingsequence and 5-prime-untranslated region of the human prostacyclin-synthase gene (CYP8A1). Hum. Genet. 108: 148–155, 2001.

[72163] 10493.Barrett, A. J.; Rawlings, N. D.: Evolutionary lines of cysteinepeptidases. Biol. Chem. 382: 727–733, 2001.

[72164] 10494.Fernandes-Alnemri, T.; Takahashi, A.; Armstrong, R.; Krebs, J.;Fritz, L.; Tomaselli, K. J.; Wang, L.; Yu, Z.; Croce, C. M.; Salveson,G.; Earnshaw, W. C.; Litwack, G.; Alnemri, E. S.: Mch3, a novel humanapoptotic cysteine protease highly related to CPP32. Cancer Res. 55:6045–6052, 1995.

[72165] 10495.Riedl, S. J.; Fuentes-Prior, P.; Renatus, M.; Kairies, N.; Krapp,S.; Huber, R.; Salvesen, G. S.; Bode, W.: Structural basis for theactivation of human procaspase-7. Proc. Nat. Acad. Sci. 98: 14790–14795,2001.

[72166] 10496.Park, W. S.; Lee, J. H.; Shin, M. S.; Park, J. Y.; Kim, H. S.;Lee, J. H.; Kim, Y. S.; Lee, S. N.; Xiao, W.; Park, C. H.;

Lee, S.H.; Yoo, N. J.; Lee, J. Y.: Inactivating mutations of the caspase-10 gene in gastric cancer. *Oncogene* 21: 2919-2925, 2002.

[72167] 10497. Shin, M. S.; Kim, H. S.; Kang, C. S.; Park, W. S.; Kim, S. Y.; Lee, S. N.; Lee, J. H.; Park, J. Y.; Jang, J. J.; Kim, C. W.; Kim, S. H.; Lee, J. Y.; Yoo, N. J.; Lee, S. H.: Inactivating mutations of CASP10 gene in non-Hodgkin lymphomas. *Blood* 99: 4094-4099, 2002.

[72168] 10498. Vincenz, C.; Dixit, V. M.: Fas-associated death domain protein interleukin-1-beta-converting enzyme 2 (FLICE2), an ICE/Ced-3 homologue, is proximally involved in CD95- and p55-mediated death signaling. *J. Biol. Chem.* 272: 6578-6583, 1997.

[72169] 10499. Wang, J.; Chun, H. J.; Wong, W.; Spencer, D. M.; Lenardo, M. J.: Caspase-10 is an initiator caspase in death receptor signaling. *Proc. Nat. Acad. Sci.* 98: 13884-13888, 2001.

[72170] 10500. Wang, J.; Zheng, L.; Lobito, A.; Chan, F. K.; Dale, J.; Sneller, M.; Yao, X.; Puck, J. M.; Straus, S. E.; Lenardo, M. J.: Inherited human caspase 10 mutations underlie defective lymphocyte and dendritic cell apoptosis in autoimmune lymphoproliferative syndrome type II. *Cell* 98: 47-58, 1999.



- [72171] 10501.Chun, H. J.; Zheng, L.; Ahmad, M.; Wang, J.; Speirs, C. K.; Siegel,R. M.; Dale, J. K.; Puck, J.; Davis, J.; Hall, C. G.; Skoda-Smith,S.; Atkinson, T. P.; Straus, S. E.; Lenardo, M. J.: Pleiotropic defects in lymphocyte activation caused by caspase-8 mutations lead to human immunodeficiency. *Nature* 419: 395-399, 2002.
- [72172] 10502.Grenet, J.; Teitz, T.; Wei, T.; Valentine, V.; Kidd, V. J.: Structure and chromosome localization of the human CASP8 gene. *Gene* 226: 225-232, 1999.
- [72173] 10503.Kischkel, F. C.; Kioschis, P.; Weitz, S.; Poustka, A.; Lichter,P.; Krammer, P. H.: Assignment of CASP8 to human chromosome band 2q33-q34 and Casp8 to the murine syntenic region on chromosome 1B-proximal C by in situ hybridization. *Cytogenet. Cell Genet.* 82: 95-96, 1998.
- [72174] 10504.Sanchez, I.; Xu, C.-J.; Juo, P.; Kakizaka, A.; Blenis, J.; Yuan,J.: Caspase-8 is required for cell death induced by expanded polyglutamine repeats. *Neuron* 22: 623-633, 1999.
- [72175] 10505.Berkovic, S. F.; Kennerson, M. L.; Howell, R. A.; Scheffer, I.E.; Hwang, P. A.; Nicholson, G. A.: Phenotypic expression of benign familial neonatal convulsions linked to chromosome 20. *Arch. Neurol.* 51:1125-1128, 1994.
- [72176] 10506.Varfolomeev, E. E.; Schuchmann, M.; Luria, V.; Chi-

annilkulchai,N.; Beckmann, J. S.; Mett, I. L.; Rebrikov, D.; Brodianski, V. M.;Kemper, O. C.; Kollet, O.; Lapidot, T.; Soffer, D.; Sobe, T.; Avraham,K. B.; Goncharov, T.; Holtmann, H.; Lonai, P.; Wallach, D.: Targetedisruption of the mouse Caspase 8 gene ablates cell death inductionby the TNF receptors, Fas/Apo1, and DR3 and is lethal prenatally. *Immunity* 9:267–276, 1998.

[72177] 10507.Kalchman, M. A.; Koide, H. B.; McCutcheon, K.; Graham, R. K.; Nichol,K.; Nishiyama, K.; Kazemi–Esfarjani, P.; Lynn, F. C.; Wellington,C.; Metzler, M.; Goldberg, Y. P.; Kanazawa, I.; Gietz, R. D.; Hayden,M. R.: HIP1, a human homologue of *S. cerevisiae* Slap2, interactswith membrane-associated huntingtin in the brain. *Nature Genet.* 16:44–53, 1997.

[72178] 10508.Wanker, E. E.; Rovira, C.; Scherzinger, E.; Hasenbank, R.; Walter,S.; Tait, D.; Colicelli, J.; Lehrach, H.: HIP–I: a huntingtin interactingprotein isolated by the yeast two–hybrid system. *Hum. Molec. Genet.* 6:487–495, 1997.

[72179] 10509.Eki, T; Okumura, K.; Shiratori, A.; Abe, M.; Nogami, M.; Taguchi,H.; Shibata, T.; Murakami, Y.; Hanaoka, F.: Assignment of the closesthuman homologue (DNA2L; KIAA0083) of the yeast Dna2 helicase genetochromosome band 10q21.3–q22.1 *Genomics* 37: 408–410, 1996.

- [72180] 10510.Nagase, T; Miyajima, N; Tanaka, A.; Sazuka, T.; Seki, N.; Sato,S.; Tabata, S.; Ishikawa, K.; Kawarabayashi, Y.; Kotani, H.; Nomura,N.: Prediction of the coding sequences of unidentified human genes.III. The coding sequences of 40 new genes (KIAA0081–KIAA0120) deducedby analysis of cDNA clones from human cell line KG–1. DNA Res. 2:37–43, 1995.
- [72181] 10511.Kim, J. W.; Lee, Y.; Lee, I. A.; Kang, H. B.; Choe, Y. K.; Choe,I. S.: Cloning and expression of human cDNA encoding Na(+),K(+)-ATPasegamma-subunit. Biochim. Biophys. Acta 1350: 133–135, 1997.
- [72182] 10512.Langkopf, A.; Hammarback, J. A.; Muller, R.; Vallee, R. B.; Garner,C. C.: Microtubule-associated proteins 1A and LC2: Two proteins encodedin one messenger RNA. J. Biol. Chem. 267: 16561–16566, 1992.
- [72183] 10513.Sun, X.–H.; Copeland, N. G.; Jenkins, N. A.; Baltimore, D.: Idproteins Id1 and Id2 selectively inhibit DNA binding by one classof helix–loop–helix proteins. Molec. Cell. Biol. 11: 5603–5611,1991.
- [72184] 10514.Kordeli, E.; Bennett, V.: Distinct ankyrin isoforms at neuroncell bodies and nodes of Ranvier resolved using erythrocyte ankyrin-deficientmice. J. Cell Biol. 114: 1243–1259, 1991.

- [72185] 10515.Reche, P. A.; Soumelis, V.; Gorman, D. M.; Clifford, T.; Liu, M.; Travis, M.; Zurawski, S. M.; Johnston, J.; Liu, Y.-J.; Spits, H.; de Waal Malefyt, R.; Kastelein, R. A.; Bazan, J. F.: Human thymicstromal lymphopoietin preferentially stimulates myeloid cells. *J.Immun.* 167: 336–343, 2001.
- [72186] 10516.Nagase, T.; Kikuno, R.; Nakayama, M.; Hirosawa, M.; Ohara, O.:Prediction of the coding sequences of unidentified human genes. XVIII.The complete sequences of 100 new cDNA clones from brain which codefor large proteins in vitro. *DNA Res.* 7: 273–281, 2000.
- [72187] 10517.Mukai, J.; Hachiya, T.; Shoji–Hoshino, S.; Kimura, M. T.; Nadano,D.; Suvanto, P.; Hanaoka, T.; Li, Y.; Irie, S.; Greene, L. A.; Sato,T.–A.: NADE, a p75NTR–associated cell death executor, is involvedin signal transduction mediated by the common neurotrophin receptorp75NTR. *J. Biol. Chem.* 275: 17566–17570, 2000.
- [72188] 10518.Rapp, G.; Freudenstein, J.; Klaudiny, J.; Mucha, J.; Wempe, F.;Zimmer, M.; Scheit, K. H.: Characterization of three abundant mRNAsfrom human ovarian granulosa cells. *DNA Cell Biol.* 9: 479–485, 1990.
- [72189] 10519.Scott, A. F.: Personal Communication. Baltimore, Md. 10/8/2001.
- [72190] 10520.Kurochkin, I. V.; Yonemitsu, N.; Funahashi, S.; No–

mura, H.: ALEX1, a novel human armadillo repeat protein that is expressed differentially in normal tissues and carcinomas. *Biochem. Biophys. Res. Commun.* 280:340–347, 2001.

- [72191] 10521. Nagase, T.; Ishikawa, K.; Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. IX. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro. *DNA Res.* 5: 31–39, 1998.
- [72192] 10522. Chuang, T.-H.; Ulevitch, R. J.: Cloning and characterization of a sub-family of human Toll-like receptors: hTLR7, hTLR8 and hTLR9. *Europ. Cytokine Netw.* 11: 372–378, 2000.
- [72193] 10523. Du, X.; Poltorak, A.; Wei, Y.; Beutler, B.: Three novel mammalian toll-like receptors: gene structure, expression, and evolution. *Europ. Cytokine Netw.* 11: 362–371, 2000.
- [72194] 10524. Kadowaki, N.; Ho, S.; Antonenko, S.; de Waal Malefyt, R.; Kastelein, R. A.; Bazan, F.; Liu, Y.-J.: Subsets of human dendritic cell precursors express different Toll-like receptors and respond to different microbial antigens. *J. Exp. Med.* 194: 863–869, 2001.
- [72195] 10525. Wirth, B.; Schmidt, T.; Hahnen, E.; Rudnik-Schoneborn, S.; Krawczak, M.; Muller-Myhsok, B.; Schon-

ling, J.; Zerres, K.: De novo rearrangements found in 2% of index patients with spinal muscular atrophy: mutational mechanisms, parental origin, mutation rate, and implications for genetic counseling. *Am. J. Hum. Genet.* 61: 1102–1111, 1997.

[72196] 10526. Frattini, A.; Orchard, P. J.; Sobacchi, C.; Giliani, S.; Abinun, M.; Mattsson, J. P.; Keeling, D. J.; Andersson, A.-K.; Wallbrandt, P.; Zecca, L.; Notarangelo, L. D.; Vezzoni, P.; Villa, A.: Defects in TCIRG1 subunit of the vacuolar proton pump are responsible for a subset of human autosomal recessive osteopetrosis. *Nature Genet.* 25:343–346, 2000.

[72197] 10527. Afzal, A. R.; Rajab, A.; Fenske, C. D.; Oldridge, M.; Elanko, N.; Ternes-Pereira, E.; Tuysuz, B.; Murday, V. A.; Patton, M. A.; Wilkie, A. O. M.; Jeffery, S.: Recessive Robinow syndrome, allelic to dominant brachydactyly type B, is caused by mutation of ROR2. *Nature Genet.* 25:419–422, 2000.

[72198] 10528. van Bokhoven, H.; Celli, J.; Kayserili, H.; van Beusekom, E.; Balci, S.; Brussel, W.; Skovby, F.; Kerr, B.; Percin, E. F.; Akarsu, N.; Brunner, H. G.: Mutation of the gene encoding the ROR2 tyrosine kinase causes autosomal recessive Robinow syndrome. *Nature Genet.* 25:423–426,

2000.

- [72199] 10529.den Dunnen, J. T.; Grootscholten, P. M.; Bakker, E.; Blonden, L. A. J.; Ginjaar, H. B.; Wapenaar, M. C.; van Paassen, H. M. B.; van Broeckhoven, C.; Pearson, P. L.; van Ommen, G. J. B.: Topography of the Duchenne muscular dystrophy (DMD) gene: FIGE and cDNA analysis of 194 cases reveals 115 deletions and 13 duplications. *Am. J. Hum. Genet.* 45: 835–847, 1989.
- [72200] 10530.Claudio, J. O.; Liew, C.–C.; Dempsey, A. A.; Cukerman, E.; Stewart, A. K.; Na, E.; Atkins, H. L.; Iscove, N. N.; Hawley, R. G.: Identification of sequence-tagged transcripts differentially expressed within the human hematopoietic hierarchy. *Genomics* 50: 44–52, 1998.
- [72201] 10531.Claudio, J. O.; Liew, C.–C.; Ma, J.; Heng, H. H. Q.; Stewart, A.K.; Hawley, R. G.: Cloning and expression analysis of a novel WD repeat gene, WDR3, mapping to 1p12–p13. *Genomics* 59: 85–89, 1999.
- [72202] 10532.Eng, B. H.; Guerinot, M. L.; Eide, D.; Saier, M. H., Jr.: Sequence analyses and phylogenetic characterization of the ZIP family of metal ion transport proteins. *J. Membr. Biol.* 166: 1–7, 1998.
- [72203] 10533.Lioumi, M.; Ferguson, C. A.; Sharpe, P. T.; Freeman, T.; Marenholz, I.; Mischke, D.; Heizmann, C.; Ragoussis, J.:

Isolation and characterization of human and mouse ZIRT1, a member of the IRT1 family of transporters, mapping within the epidermal differentiation complex. *Genomics* 62:272–280, 1999.

[72204] 10534. Lioumi, M.; Olavesen, M. G.; Nizetic, D.; Ragoussis, J.: High-resolution YAC fragmentation map of 1q21. *Genomics* 49: 200–208, 1998.

[72205] 10535. Leiper, J. M.; Santa Maria, J.; Chubb, A.; MacAllister, R. J.; Charles, I. G.; Whitley, G. S.; Vallance, P.: Identification of two human dimethylarginine dimethylaminohydrolases with distinct tissue distributions and homology with microbial arginine deiminases. *Biochem. J.* 343: 209–214, 1999.

[72206] 10536. Tran, C. T. L.; Fox, M. F.; Vallance, P.; Leiper, J. M.: Chromosomal localization, gene structure, and expression pattern of DDAH1: comparison with DDAH2 and implications for evolutionary origins. *Genomics* 68:101–105, 2000.

[72207] 10537. Vanhalst, K.; Kools, P.; Eynde, E. V.; van Roy, F.: The human and murine protocadherin-beta one-exon gene families show high evolutionary conservation, despite the difference in gene number. *FEBS Lett.* 495:120–125, 2001.



- [72208] 10538.Wu, Q.; Maniatis, T.: A striking organization of a large family of human neural cadherin like cell adhesion genes. *Cell* 97: 779–790,1999.
- [72209] 10539.Katayama, H.; Ota, T.; Morita, K.; Terada, Y.; Suzuki, F.; Katoh,O.; Tatsuka, M.: Human AIM-1: cDNA cloning and reduced expression during endomitosis in megakaryocyte–lineage cells. *Gene* 224: 1–7,1998.
- [72210] 10540.Kimura, M.; Matsuda, Y.; Yoshioka, T.; Sumi, N.; Okano, Y.: Identification and characterization of STK12/Aik2: a human gene related to aurora of *Drosophila* and yeast IPL1. *Cytogenet. Cell Genet.* 82: 147–152,1998.
- [72211] 10541.Tatsuka, M.; Katayama, H.; Ota, T.; Tanaka, T.; Odashima, S.; Suzuki,F.; Terada, Y.: Multinuclearity and increased ploidy caused by overexpression of the aurora– and Ipl1–like midbody–associated protein mitotic kinase in human cancer cells. *Cancer Res.* 58: 4811–4816, 1998.
- [72212] 10542.Chen, C. M.; Kraut, N.; Groudine, M.; Weintraub, H.: I–mf, a novel myogenic repressor, interacts with members of the MyoD family. *Cell* 86:731–741, 1996.
- [72213] 10543.Kraut, N.: The gene encoding I–mf (Mdfi) maps to human chromosome 6p21 and mouse chromosome 17. *Mammalian Genome* 8: 618–619, 1997.
- [72214] 10544.Kraut, N.; Snider, L.; Chen, C.–M. A.; Tapscott, S. J.;

Groudine, M.: Requirement of the mouse *l-mfa* gene for placental development and skeletal patterning. *EMBO J.* 17: 6276–6288, 1998.

[72215] 10545. Boynton, S.; Tully, T.: *Latheo*, a new gene involved in associative learning and memory in *Drosophila melanogaster*, identified from P-element mutagenesis. *Genetics* 131: 655–672, 1992.

[72216] 10546. Pinto, S.; Quintana, D. G.; Smith, P.; Mihalek, R. M.; Hou, Z.-H.; Boynton, S.; Jones, C. J.; Hendricks, M.; Velinzon, K.; Wohlschlegel, J. A.; Austin, R. J.; Lane, W. S.; Tully, T.; Dutta, A.: *Latheo* encodes a subunit of the origin recognition complex and disrupts neuronal proliferation and adult olfactory memory when mutant. *Neuron* 23:45–54, 1999.

[72217] 10547. Springer, J.; Nanda, I.; Hoehn, K.; Schmid, M.; Grummt, F.: Identification and chromosomal localization of murine *ORC3*, a new member of the mouse origin recognition complex. *Cytogenet. Cell Genet.* 87: 245–251, 1999.

[72218] 10548. Tugal, T.; Zou-Yang, X. H.; Gavin, K.; Pappin, D.; Canas, B.; Kobayashi, R.; Hunt, T.; Stillman, B.: The *Orc4p* and *Orc5p* subunits of the *Xenopus* and human origin recognition complex are related to *Orc1p* and *Cdc6p*. *J. Biol. Chem.* 273: 32421–32429, 1998.

- [72219] 10549. Butcher, S.; Arney, K. L.; Cook, G. P.: MAFA-L, an ITIM-containing receptor encoded by the human NK cell gene complex and expressed by basophils and NK cells. *Europ. J. Immun.* 28: 3755–3762, 1998.
- [72220] 10550. Hanke, T.; Corral, L.; Vance, R. E.; Raulet, D. H.: 2F1 antigen, the mouse homolog of the rat 'mast cell function-associated antigen', is a lectin-like type II transmembrane receptor expressed by natural killer cells. *Europ. J. Immun.* 28: 4409–4417, 1998.
- [72221] 10551. Lamers, M. B. A. C.; Lamont, A. G.; Williams, D. H.: Human MAFA has alternatively spliced variants. *Biochim. Biophys. Acta* 1399:209–212, 1998.
- [72222] 10552. Voehringer, D.; Kaufmann, M.; Pircher, H.: Genomic structure, alternative splicing, and physical mapping of the killer cell lectin-like receptor G1 gene (KLRG1), the mouse homologue of MAFA. *Immunogenetics* 52:206–211, 2001.
- [72223] 10553. Bahler, M.: Myosins on the move to signal transduction. *Curr. Opin. Cell Biol.* 8: 18–22, 1996.
- [72224] 10554. Gorman, S. W.; Haider, N. B.; Grieshammer, U.; Swiderski, R. E.; Kim E; Welch, J. W.; Searby, C.; Leng, S.; Carmi, R.; Sheffield, V. C.; Duhl, D. M.: The cloning and developmental expression of unconventional myosin IXA (MYO9A) a gene in the Bardet-Biedl syndrome (BBS4) re-

gionat chromosome 15q22–q23. *Genomics* 59: 150–160, 1999.

[72225] 10555. Dunham, I.; Shimizu, N.; Roe, B. A.; Chissoe, S.; Hunt, A. R.; Collins, J. E.; Bruskiewich, R.; Beare, D. M.; Clamp, M.; Smink, L. J.; Ainscough, R.; Almeida, J. P.; and 205 others: The DNA sequence of human chromosome 22. *Nature* 402: 489–495, 1999.

[72226] 10556. Kimura, T.; Ivell, R.; Rust, W.; Mizumoto, Y.; Ogita, K.; Kusui, C.; Matsumura, Y.; Azuma, C.; Murata, Y.: Molecular cloning of a human MafF homologue, which specifically binds to the oxytocin receptor gene in term myometrium. *Biochem. Biophys. Res. Commun.* 264: 86–92, 1999.

[72227] 10557. Hiki, K.; D'Andrea, R. J.; Furze, J.; Crawford, J.; Woollatt, E.; Sutherland, G. R.; Vadas, M. A.; Gamble, J. R.: Cloning, characterization, and chromosomal location of a novel human K(+)–Cl(–) cotransporter. *J. Biol. Chem.* 274: 10661–10667, 1999.

[72228] 10558. Mount, D. B.: Personal Communication. Nashville, Tenn. 3/28/2000.

[72229] 10559. Mount, D. B.; Mercado, A.; Song, L.; Xu, J.; George, A. L., Jr.; Delpire, E.; Gamba, G.: Cloning and characterization of KCC3 and KCC4, new members of the cation–

chloride cotransporter gene family. J.Biol. Chem. 274: 16355–16362, 1999.

[72230] 10560.Herzing, L. B. K.; Kim, S.-J.; Cook, E. H., Jr.; Ledbetter, D.H.: The human aminophospholipid-transporting ATPase gene ATP10C maps adjacent to UBE3A and exhibits similar imprinted expression. Am.J. Hum. Genet. 68: 1501–1505, 2001.

[72231] 10561.Dode, C.; Le Du, N.; Cuisset, L.; Letourneur, F.; Berthelot, J.-M.; Vaudour, G.; Meyrier, A.; Watts, R. A.; Scott, D. G. I.; Nicholls, A.; Granel, B.; Frances, C.; Garcier, F.; Edery, P.; Boulinguez, S.; Domergues, J.-P.; Delpech, M.; Grateau, G.: New mutations of CIAS1 that are responsible for Muckle-Wells syndrome and familial cold urticaria: a novel mutation underlies both syndromes. Am. J. Hum. Genet. 70:1498–1506, 2002.

[72232] 10562.Feldmann, J.; Prieur, A.-M.; Quartier, P.; Berquin, P.; Certain, S.; Cortis, E.; Teillac-Hamel, D.; Fischer, A.; de Saint Basile, G.: Chronic infantile neurological cutaneous and articular syndrome is caused by mutations in CIAS1, a gene highly expressed in polymorphonuclear cells and chondrocytes. Am. J. Hum. Genet. 71: 198–203, 2002.

[72233] 10563.Greeve, I.; Hermans-Borgmeyer, I.; Brellinger, C.; Kasper, D.; Gomez-Isla, T.; Behl, C.; Levkau, B.; Nitsch, R.

M.: The human DIMINUTO/DWARF1 homolog seladin-1 confers resistance to Alzheimer's disease-associated neurodegeneration and oxidative stress. *J. Neurosci.* 20: 7345–7352, 2000.

[72234] 10564. Sarkar, D.; Imai, T.; Kambe, F.; Shibata, A.; Ohmori, S.; Siddiq, A.; Hayasaka, S.; Funahashi, H.; Seo, H.: The human homolog of Diminuto/Dwarf1 gene (hDiminuto): a novel ACTH-responsive gene overexpressed in benign cortisol-producing adrenocortical adenomas. *J. Clin. Endocr. Metab.* 86: 5130–5137, 2001.

[72235] 10565. Weidenhammer, E. M.; Singh, M.; Ruiz-Noriega, M.; Woolford, J. L., Jr.: The PRP31 gene encodes a novel protein required for pre-mRNA splicing in *Saccharomyces cerevisiae*. *Nucleic Acids Res.* 24: 1164–1170, 1996.

[72236] 10566. Hattori, M.; Fujiyama, A.; Taylor, T. D.; Watanabe, H.; Yada, T.; Park, H.-S.; Toyoda, A.; Ishii, K.; Totoki, Y.; Choi, D.-K.; Groner, Y.; Soeda, E.; and 52 others: The DNA sequence of human chromosome 21. *Nature* 405: 311–319, 2000. Note: Erratum: *Nature*: 407: 110 only, 2000.

[72237] 10567. Wilcox, E. R.; Burton, Q. L.; Naz, S.; Riazuddin, S.; Smith, T. N.; Ploplis, B.; Belyatseva, I.; Ben-Yosef, T.; Liburd, N. A.; Morell, R. J.; Kachar, B.; Wu, D. K.; Griffith, A.

J.; Riazuddin, S.; Friedman, T. B.: Mutations in the gene encoding tight junction claudin-14 cause recessive deafness DFNB29. *Cell* 104: 165–172, 2001.

[72238] 10568. Dunlevy, J. R.; Berryhill, B. L.; Vergnes, J.-P.; SundarRaj, N.; Hassell, J. R.: Cloning, chromosomal localization, and characterization of cDNA from a novel gene, SH3BP4, expressed by human corneal fibroblasts. *Genomics* 62: 519–524, 1999.

[72239] 10569. Wong, W. T.; Schumacher, C.; Salcini, A. E.; Romano, A.; Castagnino, P.; Pelicci, P. G.; DiFiore, P. P.: A protein-binding domain, EH, identified in the receptor tyrosine kinase substrate Eps15 and conserved in evolution. *Proc. Nat. Acad. Sci.* 92: 9530–9534, 1995.

[72240] 10570. Yamadori, T.; Baba, Y.; Matsushita, M.; Hashimoto, S.; Kurosaki, M.; Kurosaki, T.; Kishimoto, T.; Tsukada, S.: Bruton's tyrosine kinase activity is negatively regulated by Sab, the Btk-SH3 domain-binding protein. *Proc. Nat. Acad. Sci.* 96: 6341–6346, 1999.

[72241] 10571. Engqvist-Goldstein, A. E. Y.; Kessels, M. M.; Chopra, V. S.; Hayden, M. R.; Drubin, D. G.: An actin-binding protein of the Sla2/Huntingtin-interacting protein 1 family is a novel component of clathrin-coated pits and vesicles. *J. Cell Biol.* 147: 1503–1518, 1999.

- [72242] 10572.Baba, Y.; Matsushita, M.; Matsuda, Y.; Inazawa, J.; Yamadori, T.; Hashimoto, S.; Kishimoto, T.; Tsukada, S.: Assignment of SH3BP5/Sh3bp5 encoding Sab, an SH3 domain-binding protein which preferentially associates with Bruton's tyrosine kinase, to human chromosome 1q43 and mouse chromosome 14B by in situ hybridization. *Cytogenet. Cell Genet.* 87:221–222, 1999.
- [72243] 10573.Matsushita, M.; Yamadori, T.; Kato, S.; Takemoto, Y.; Inazawa, J.; Baba, Y.; Hashimoto, S.; Sekine, S.; Arai, S.; Kunikata, T.; Kurimoto, M.; Kishimoto, T.; Tsukada, S.: Identification and characterization of a novel SH3-domain binding protein, Sab, which preferentially associates with Bruton's tyrosine kinase (Btk). *Biochem. Biophys. Res. Commun.* 245:337–343, 1998.
- [72244] 10574.Itoh, T.; Koshiba, S.; Kigawa, T.; Kikuchi, A.; Yokoyama, S.; Takenawa, T.: Role of the ENTH domain in phosphatidylinositol-4,5-bisphosphate binding and endocytosis. *Science* 291: 1047–1051, 2001.
- [72245] 10575.Seki, N.; Muramatsu, M.; Sugano, S.; Suzuki, Y.; Nakagawara, A.; Ohhira, M.; Hayashi, A.; Hori, T.; Saito, T.: Cloning, expression analysis, and chromosomal localization of HIP1R, an isolog of huntingtin interacting protein (HIP1). *J. Hum. Genet.* 43: 268–271, 1998.



- [72246] 10576. Marton, M. J.; Vazquez de Aldana, C. R.; Qiu, H.; Chakraborty, K.; Hinnebusch, A. G.: Evidence that GCN1 and GCN20, translational regulators of GCN4, function on elongating ribosomes in activation of eIF2- $\alpha$  kinase GCN2. *Molec. Cell. Biol.* 17: 4474–4489, 1997.
- [72247] 10577. Nash, S. R.; Giros, B.; Kingsmore, S. F.; Kim, K. M.; El-Mestikawy, S.; Dong, Q.; Fumagalli, F.; Seldin, M. F.; Caron, M. G.: Cloning, gene structure and genomic localization of an orphan transporter from mouse kidney with six alternatively-spliced isoforms. *Receptors Channels* 6:113–128, 1998.
- [72248] 10578. Scott, A. F.: Personal Communication. Baltimore, Md. 2/5/2001.
- [72249] 10579. Meguro, M.; Kashiwagi, A.; Mitsuya, K.; Nakao, M.; Kondo, I.; Saitoh, S.; Oshimura, M.: A novel maternally expressed gene, ATP10C, encodes a putative aminophospholipid translocase associated with Angelman syndrome. *Nature Genet.* 28: 19–20, 2001.
- [72250] 10580. Nakakura, E. K.; Watkins, D. N.; Schuebel, K. E.; Sriuranpong, V.; Borges, M. W.; Nelkin, B. D.; Ball, D. W.: Mammalian Scratch: a neural-specific Snail family transcriptional repressor. *Proc. Nat. Acad. Sci.* 98: 4010–4015, 2001.

- [72251] 10581.Scott, A. F.: Personal Communication. Baltimore, Md. 6/21/2001.
- [72252] 10582.Hoatlin, M. E.; Zhi, Y.; Ball, H.; Silvery, K.; Melnick, A.; Stone,S.; Arai, S.; Hawe, N.; Owen, G.; Zelent, A.; Licht, J. D.: A novelBTB/POZ transcriptional repressor protein interacts with the Fanconianemia group C protein and PLZF. Blood 94: 3737–3747, 1999.
- [72253] 10583.Lin, W.; Lai, C.–H.; Tang, C.–J. C.; Huang, C.–J.; Tang, T. K.: Identification and gene structure of a novel human PLZF–relatedtranscription factor gene, TZFP. Biochem. Biophys. Res. Commun. 264:789–795, 1999.
- [72254] 10584.Yokoyama–Kobayashi, M.; Sugano, S.; Kato, T.; Kato, S.: A signalsequence detection system using secreted protease activity as an indicator. Gene 163:193–196, 1995.
- [72255] 10585.Yokoyama–Kobayashi, M.; Yamaguchi, T.; Sekine, S.; Kato, S.: Selectionof cDNAs encoding putative type II membrane proteins on the cell surfacefrom a human full–length cDNA bank. Gene 228: 161–167, 1999.
- [72256] 10586.Olivier, E.; Soury, E.; Risler, J. L.; Smith, F.; Schneider, K.;Lochner, K.; Jouzeau, J. Y.; Fey, G. H.; Salier, J. P.: A novel setof hepatic mRNAs preferentially expressed during an acute inflammationin rat represents mostly intra–

cellular proteins. *Genomics* 57: 352–364,1999.

- [72257] 10587.Olivier, E.; Soury, E.; Ruminy, P.; Husson, A.; Parmentier, F.;Daveau, M.; Salier, J.–P.: Fetuin–B, a second member of the fetuinfamily in mammals. *Biochem. J.* 350: 589–597, 2000.
- [72258] 10588.Hampe, J.; Grebe, J.; Nikolaus, S.; Solberg, C.; Croucher, P. J.P.; Mascheretti, S.; Jahnsen, J.; Moum, B.; Klump, B.; Krawczak, M.;Mirza, M. M.; Foelsch, U. R.; Vatn, M.; Schreiber, S.: Associationof NOD2 (CARD 15) genotype with clinical course of Crohn's disease:a cohort study. *Lancet* 359: 1661–1665, 2002.
- [72259] 10589.Hugot, J.–P.; Chamaillard, M.; Zouali, H.; Lesage, S.; Cezard,J.–P.; Belaiche, J.; Almer, S.; Tysk, C.; O'Morain, C. A.; Gassull,M.; Binder, V.; Finkel, Y.; and 8 others: Association of NOD2 leucine–richrepeat variants with susceptibility to Crohn's disease. *Nature* 411:599–603, 2001.
- [72260] 10590.Murillo, L.; Crusius, J. B. A.; van Bodegraven, A. A.; Alizadeh,B. Z.; Pena, A. S.: CARD15 gene and the classification of Crohn'sdisease. *Immunogenetics* 54: 59–61, 2002.
- [72261] 10591.Ogura, Y.; Bonen, D. K.; Inohara, N.; Nicolae, D. L.; Chen, F.F.; Ramos, R.; Britton, H.; Moran, T.; Karaliuskas, R.; Duerr, R.H.; Achkar, J.–P.; Brant, S. R.; Bayless, T. M.;

Kirschner, B. S.; Hanauer, S. B.; Nunez, G.; Cho, J. H.: A frameshift mutation in Nod2 associated with susceptibility to Crohn's disease. *Nature* 411: 603–606, 2001.

[72262] 10592. Ogura, Y.; Inohara, N.; Benito, A.; Chen, F. F.; Yamaoka, S.; Nunez, G.: Nod2, a Nod1/Apaf-1 family member that is restricted to monocytes and activates NF- $\kappa$ B. *J. Biol. Chem.* 276: 4812–4818, 2001.

[72263] 10593. Lesage, S.; Zouali, H.; Cezard, J.-P.; EPWG-IBD Group; Colombel, J.-F.; EPIMAD Group; Belaiche, J.; GETAID Group; Almer, S.; Tysk, C.; O'Morain, C.; Gassull, M.; Binder, V.; Finkel, Y.; Modigliani, R.; Gower-Rousseau, C.; Macry, J.; Merlin, F.; Chamaillard, M.; Jannot, A.-S.; Thomas, G.; Hugot, J.-P.: CARD15/NOD2 mutational analysis and genotype-phenotype correlation in 612 patients with inflammatory bowel disease. *Am. J. Hum. Genet.* 70: 845–857, 2002.

[72264] 10594. van Heel, D. A.; McGovern, D. P. B.; Cardon, L. R.; Dechairo, B. M.; Lench, N. J.; Carey, A. H.; Jewell, D. P.: Fine mapping of the IBD1 locus did not identify Crohn disease-associated NOD2 variants: implications for complex disease genetics. *Am. J. Med. Genet.* 111: 253–259, 2002.

[72265] 10595. Vermeire, S.; Wild, G.; Kocher, K.; Cousineau, J.; Dufresne, L.; Bitton, A.; Langelier, D.; Pare, P.; Lapointe, G.;

Cohen, A.; Daly, M. J.; Rioux, J. D.: CARD15 genetic variation in a Quebec population: prevalence, genotype-phenotype relationship, and haplotype structure. *Am. J. Hum. Genet.* 71: 74–83, 2002.

[72266] 10596. Yamazaki, K.; Takazoe, M.; Tanaka, T.; Kazumori, T.; Nakamura, Y.: Absence of mutation in the NOD2/CARD15 gene among 483 Japanese patients with Crohn's disease. *J. Hum. Genet.* 47: 469–472, 2002.

[72267] 10597. Fliss, M. S.; Hinkle, P. M.; Bancroft, C.: Expression cloning and characterization of PREB (prolactin regulatory element binding), a novel WD motif DNA-binding protein with a capacity to regulate prolactin promoter activity. *Molec. Endocr.* 13: 644–657, 1999.

[72268] 10598. Taylor Clelland, C. L.; Levy, B.; McKie, J. M.; Duncan, A. M. V.; Hirschhorn, K.; Bancroft, C.: Cloning and characterization of human PREB; a gene that maps to a genomic region associated with trisomy 2p syndrome. *Mammalian Genome* 11: 675–681, 2000.

[72269] 10599. Li, B.; Oestreich, S.; de Lange, T.: Identification of human Rap1: implications for telomere evolution. *Cell* 101: 471–483, 2000.

[72270] 10600. Lieb, J. D.; Liu, X.; Botstein, D.; Brown, P. O.: Promoter-specific binding of Rap1 revealed by genome-wide

maps of protein–DNA association. *NatureGenet.* 28: 327–334, 2001.

- [72271] 10601.Honore, B.; Leffers, H.; Madsen, P.; Rasmussen, H. H.; Vandekerckhove,J.; Celis, J. E.: Molecular cloning and expression of a transformation–sensitivehuman protein containing the TPR motif and sharing identity to thestress–inducible yeast protein STI1. *J. Biol. Chem.* 267: 8485–8491,1992.
- [72272] 10602.Scheufler, C.; Brinker, A.; Bourenkov, G.; Pegoraro, S.; Moroder,L.; Bartunik, H.; Hartl, F. U.; Moarefi, I.: Structure of TPR domain–peptidecomplexes: critical elements in the assembly of the Hsp70–Hsp90 multichaperonemachine. *Cell* 101: 199–210, 2000.
- [72273] 10603.Chong, S. S.; Tanigami, A.; Roschke, A. V.; Ledbetter, D. H.:14–3–3–epsilon has no homology to LIS1 and lies telomeric to it onchromosome 17p13.3 outside the Miller–Dieker syndrome chromosome region. *GenomeRes.* 6: 735–741, 1996.
- [72274] 10604.Conklin, D. S.; Galaktionov, K.; Beach, D.: 14–3–3 proteins associatewith cdc25 phosphatases. *Proc. Nat. Acad. Sci.* 92: 7892–7896, 1995.
- [72275] 10605.Jin, D.–Y.; Lyu, M. S.; Kozak, C. A.; Jeang, K.–T.: Function of14–3–3 proteins. *Nature* 382: 308 only, 1996.

- [72276] 10606.Luk, S. C. W.; Garcia-Barcelo, M.; Tsui, S. K. W.; Fung, K. P.; Lee, C. Y.; Waye, M. M. Y.: Assignment of the human 14-3-3 epsilon isoform (YWHAE) to human chromosome 17p13 by in situ hybridization. *Cytogenet. Cell Genet.* 78: 105-106, 1997.
- [72277] 10607.Slantz-Kesler, K.; Moore, J. T.; Lombard, M.; Zhang, J.; Hollingsworth, R.; Weiner, M. P.: Identification of the human Mnk2 gene (MKNK2) through protein interaction with estrogen receptor beta. *Genomics* 69:63-71, 2000.
- [72278] 10608.Li, Y.; He, X.; Schembri-King, J.; Jakes, S.; Hayashi, J.: Cloning and characterization of human Lnk, an adaptor protein with pleckstrin homology and Src homology 2 domains that can inhibit T cell activation. *J. Immunol.* 164: 5199-5206, 2000.
- [72279] 10609.Takaki, S.; Sauer, K.; Iritani, B. M.; Chien, S.; Ebihara, Y.; Tsuji, K.; Takatsu, K.; Perlmutter, R. M.: Control of B cell production by the adaptor protein Lnk: definition of a conserved family of signal-modulating proteins. *Immunity* 13: 599-609, 2000.
- [72280] 10610.Fiscella, M.; Zhang, H.; Fan, S.; Sakaguchi, K.; Shen, S.; Mercer, W. E.; Vande Woude, G. F.; O'Connor, P. M.; Appella, E.: Wip1, a novel human protein phosphatase that is induced in response to ionizing radiation in a

p53-dependent manner. Proc. Nat. Acad. Sci. 94:  
6048–6053,1997.

[72281] 10611.Li, J.; Yang, Y.; Peng, Y.; Austin, R. J.; van Eyndhoven, W. G.;Nguyen, K. C. Q.; Gabriele, T.; McCurrach, M. E.; Marks, J. R.; Hoey,T.; Lowe, S. W.; Powers, S.: Oncogenic properties of PPM1D locatedwithin a breast cancer amplification epicenter at 17q23. Nature Genet. 31:133–134, 2002.

[72282] 10612.Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayasi, Y.;1. Fiscella, M.; Zhang, H.; Fan, S.; Sakaguchi, K.; Shen, S.; Mercer,W. E.; Vande Woude, G. F.; O'Connor, P. M.; Appella, E.: Wip1, a novel human protein phosphatase that is induced in response to ionizing radiation in a p53-dependent manner. Proc. Nat. Acad. Sci. 94: 6048–6053,1997.

[72283] 10613.Stover, C. M.; Schwaeble, W. J.; Lynch, N. J.; Thiel, S.; Speicher,M. R.: Assignment of the gene encoding mannan-binding lectin-associatedserine protease 2 (MASP2) to human chromosome 1p36.3–p36.2 by in situhybridization and somatic cell hybrid analysis. Cytogenet. Cell Genet. 84:148–149, 1999.

[72284] 10614.Stover, C. M.; Thiel, S.; Thelen, M.; Lynch, N. J.; Vorup-Jensen,T.; Jensenius, J. C.; Schwaeble, W. J.: Two



constituents of the initiation complex of the mannan-binding lectin activation pathway of complement are encoded by a single structural gene. *J. Immun.* 162: 3481–3490, 1999.

[72285] 10615. Takahashi, M.; Endo, Y.; Fujita, T.; Matsushita, M.: A truncated form of mannan-binding lectin-associated serine protease (MASP)-2 expressed by alternative polyadenylation is a component of the lectin complement pathway. *Int. Immun.* 11: 859–863, 1999.

[72286] 10616. Thiel, S.; Vorup-Jensen, T.; Stover, C. M.; Schwaeble, W.; Laursen, S. B.; Poulsen, K.; Willis, A. C.; Eggleton, P.; Hansen, S.; Holmskov, U.; Reid, K. B. M.; Jensenius, J. C.: A second serine protease associated with mannan-binding lectin that activates complement. *Nature* 386: 506–510, 1997.

[72287] 10617. Ferguson, K. M.; Kavran, J. M.; Sankaran, V. G.; Fournier, E.; Isakoff, S. J.; Skolnik, E. Y.; Lemmon, M. A.: Structural basis for discrimination of 3-phosphoinositides by pleckstrin homology domains. *Molec. Cell* 6: 373–384, 2000.

[72288] 10618. Franco, M.; Boretto, J.; Robineau, S.; Monier, S.; Goud, B.; Chardin, P.; Chavrier, P.: ARNO3, a Sec7-domain guanine nucleotide exchange factor for ADP ribosylation

factor 1, is involved in the control of Golgi structure and function. Proc. Nat. Acad. Sci. 95: 9926–9931, 1998.

- [72289] 10619. Klarlund, J. K.; Guilherme, A.; Holik, J. J.; Virbasius, J. V.; Chawla, A.; Czech, M. P.: Signaling by phosphoinositide-3,4,5-triphosphate through proteins containing pleckstrin and Sec7 homology domains. Science 275:1927–1930, 1997.
- [72290] 10620. Drewes, T.; Senkel, S.; Holewa, B.; Ryffel, G. U.: Human hepatocyte nuclear factor 4 isoforms are encoded by distinct and differentially expressed genes. Molec. Cell. Biol. 16: 925–931, 1996.
- [72291] 10621. Taraviras, S.; Mantamadiotis, T.; Dong-Si, T.; Mincheva, A.; Lichter, P.; Drewes, T.; Ryffel, G. U.; Monaghan, A. P.; Schutz, G.: Primary structure, chromosomal mapping, expression and transcriptional activity of murine hepatocyte nuclear factor 4- $\gamma$ . Biochim. Biophys. Acta 1490:21–32, 2000.
- [72292] 10622. Gauss, R.; Seifert, R.; Kaupp, U. B.: Molecular identification of a hyperpolarization-activated channel in sea urchin sperm. Nature 393:583–587, 1998.
- [72293] 10623. Seifert, R.; Scholten, A.; Gauss, R.; Mincheva, A.; Lichter, P.; Kaupp, U. B.: Molecular characterization of a slowly gating human hyperpolarization-activated channel

predominantly expressed in thalamus, heart, and testis.

Proc. Nat. Acad. Sci. 96: 9391–9396, 1999.

[72294] 10624. Aihara, Y.; Mashima, H.; Onda, H.; Hisano, S.; Kasuya, H.; Hori, T.; Yamada, S.; Tomura, H.; Yamada, Y.; Inoue, I.; Kojima, I.; Takeda, J.: Molecular cloning of a novel brain-type Na(+)-dependent inorganic phosphate cotransporter. J. Neurochem. 74: 2622–2625, 2000.

[72295] 10625. Bellocchio, E. E.; Reimer, R. J.; Fremeau, R. T., Jr.; Edwards, R. H.: Uptake of glutamate into synaptic vesicles by an inorganic phosphate transporter. Science 289: 957–960, 2000.

[72296] 10626. Ni, B.; Du, Y.; Wu, X.; DeHoff, B. S.; Rosteck, P. R., Jr.; Paul, S. M.: Molecular cloning, expression, and chromosomal localization of a human brain-specific Na(+)-dependent inorganic phosphate cotransporter. J. Neurochem. 66: 2227–2238, 1996.

[72297] 10627. Takamori, S.; Rhee, J. S.; Rosenmund, C.; Jahn, R.: Identification of a vesicular glutamate transporter that defines a glutamatergic phenotype in neurons. Nature 407: 189–194, 2000.

[72298] 10628. Blackwood, D.: P300, a state and a trait marker in schizophrenia. Lancet 355: 771–772, 2000.

[72299] 10629. Blackwood, D. H. R.; Fordyce, A.; Walker, M. T.; St.

Clair, D.M.; Porteous, D. J.; Muir, W. J.: Schizophrenia and affective disorders--cosegregation with a translocation at chromosome 1q42 that directly disrupts brain-expressed genes: clinical and P300 findings in a family. *Am. J. Hum. Genet.* 69:428–433, 2001.

[72300] 10630. Ekelund, J.; Hovatta, I.; Parker, A.; Paunio, T.; Varilo, T.; Martin, R.; Suhonen, J.; Ellonen, P.; Chan, G.; Sinheimer, J. S.; Sobel, E.; Juvonen, H.; Arajärvi, R.; Partonen, T.; Suvisaari, J.; Lonnqvist, J.; Meyer, J.; Peltonen, L.: Chromosome 1 loci in Finnish schizophrenia families. *Hum. Molec. Genet.* 10: 1611–1617, 2001.

[72301] 10631. Millar, J. K.; Wilson–Annan, J. C.; Anderson, S.; Christie, S.; Taylor, M. S.; Semple, C. A. M.; Devon, R. S.; St. Clair, D. M.; Muir, W. J.; Blackwood, D. H. R.; Porteous, D. J.: Disruption of two novel genes by a translocation cosegregating with schizophrenia. *Hum. Molec. Genet.* 9: 1415–1423, 2000.

[72302] 10632. St. Clair, D.; Blackwood, D.; Muir, W.; Carothers, A.; Walker, M.; Spowart, G.; Gosden, C.; Evans, H. J.: Association within a family of a balanced autosomal translocation with major mental illness. *Lancet* 336:13–16, 1990.

[72303] 10633. Bulfone, A.; Menguzzato, E.; Broccoli, V.; Marchitello, A.; Gattuso, C.; Mariani, M.; Consalez, G. G.; Mar-

tinez, S.; Ballabio, A.; Banfi, S.: Barhl1, a gene belonging to a new subfamily of mammalian homeobox genes, is expressed in migrating neurons of the CNS. Hum. Molec. Genet. 9: 1443–1452, 2000.

[72304] 10634. Alessi, D. R.; Deak, M.; Casamayor, A.; Caudwell, F. B.; Morrice, N.; Norman, D. G.; Gaffney, P.; Reese, C. B.; MacDougall, C. N.; Harbison, D.; Ashworth, A.; Bownes, M.: 3-phosphoinositide-dependent protein kinase-1 (PDK1): structural and functional homology with the Drosophila D-STPK61 kinase. Curr. Biol. 7: 776–789, 1997.

[72305] 10635. Bertin, J.; Nir, W.-J.; Fischer, C. M.; Tayber, O. V.; Errada, P. R.; Grant, J. R.; Keilty, J. J.; Gosselin, M. L.; Robinson, K. E.; Wong, G. H. W.; Glucksmann, M. A.; DiStefano, P. S.: Human CARD4 protein is a novel CED-4/Apaf-1 cell death family member that activates NF- $\kappa$ B. J. Biol. Chem. 274: 12955–12958, 1999.

[72306] 10636. Cohen, D. E.; Green, R. M.; Wu, M. K.; Beier, D. R.: Cloning, tissue-specific expression, gene structure and chromosomal localization of human phosphatidylcholine transfer protein. Biochim. Biophys. Acta 1447: 265–270, 1999.

[72307] 10637. van Helvoort, A.; de Brouwer, A.; Ottenhoff, R.; Brouwers, J. F. H. M.; Wijnholds, J.; Beijnen, J. H.; Rijneveld,

A.; van der Valk, M. A.; Majoor, D.; Voorhout, W.; Wirtz, K. W. A.; Elferink, R. P. J. O.; Borst, P.: Mice without phosphatidylcholine transfer protein have no defects in the secretion of phosphatidylcholine into bile or into lung airspaces. *Proc. Nat. Acad. Sci.* 96: 11501–11506, 1999.

[72308] 10638. de Rooij, J.; Zwartkruis, F. J. T.; Verheijen, M. H. G.; Cool, R. H.; Nijman, S. M. B.; Wittinghofer, A.; Bos, J. L.: Epac is a Rap1 guanine-nucleotide-exchange factor directly activated by cyclic AMP. *Nature* 396: 474–477, 1998.

[72309] 10639. Kawasaki, H.; Springett, G. M.; Mochizuki, N.; Toki, S.; Nakaya, M.; Matsuda, M.; Housman, D. E.; Graybiel, A. M.: A family of cAMP-binding proteins that directly activate Rap1. *Science* 282: 2275–2279, 1998.

[72310] 10640. Hahn, Y.; Lee, J.; Seong, C.; Yoon, J.; Chung, J. H.: Structural analysis of phylogenetically conserved J domain protein gene. *Biochim. Biophys. Acta* 1447: 325–333, 1999.

[72311] 10641. Hollopeter, G.; Jantzen, H.-M.; Vincent, D.; Li, G.; England, L.; Ramakrishnan, V.; Yang, R.-B.; Nurden, P.; Nurden, A.; Julius, D.; Conley, P. B.: Identification of the platelet ADP receptor targeted by antithrombotic drugs. *Nature* 409: 202–206, 2001.

[72312] 10642. Nurden, P.; Savi, P.; Heilmann, E.; Bihour, C.; Her-

bert, J.-M.;Maffrand, J.-P.; Nurden, A.: An inherited bleeding disorder linked to a defective interaction between ADP and its receptor on platelets:its influence on glycoprotein IIb-IIIa complex function. *J. Clin.Invest.* 95: 1612-1622, 1995.

[72313] 10643.Hill, K. E.; Dasouki, M.; Phillips, J. A., III; Burk, R. F.: Human selenoprotein P gene maps to 5q31. *Genomics* 36: 550-551, 1996.

[72314] 10644.Hill, K. E.; Lloyd, R. S.; Burk, R. F.: Conserved nucleotide sequences in the open reading frame and 3-prime untranslated region of selenoprotein P mRNA. *Proc. Nat. Acad. Sci.* 90: 537-541, 1993.

[72315] 10645.Keshan Disease Research Group of the Chinese Academy of Medical Sciences: Observations on effect of sodium selenite in prevention of Keshan disease. *Chinese Med. J.* 92: 471-476, 1979.

[72316] 10646.Eberhart, C. G.; Maines, J. Z.; Wasserman, S. A.: Meiotic cell cycle requirement for a fly homologue of human Deleted in Azoospermia. *Nature* 381:783-785, 1996.

[72317] 10647.Rugglu, M.; Speed, R.; Taggart, M.; McKay, S. J.; Kilanowski, F.;Saunders, P.; Derin, J.; Cooke, H. J.: The mouse Dazl gene encodes a cytoplasmic protein essential for gametogenesis. *Nature* 389: 73-77,1997.

- [72318] 10648. Shan, Z.; Hirschmann, P.; Seebacher, T.; Edelmann, A.; Jauch, A.; Morell, J.; Urbitsch, P.; Vogt, P. H.: A SPGY copy homologous to the mouse gene Dazla and the Drosophila gene boule is autosomal and expressed only in the human male gonad. Hum. Molec. Genet. 5: 2005–2011, 1996.
- [72319] 10649. Slee, R.; Grimes, B.; Speed, R. M.; Taggart, M.; Maguire, S. M.; Ross, A.; McGill, N. I.; Saunders, P. T. K.; Cooke, H. J.: A human DAZ transgene confers partial rescue of the mouse Daz1 null phenotype. Proc. Nat. Acad. Sci. 96: 8040–8045, 1999.
- [72320] 10650. Yen, P. H.; Chai, N. N.; Salido, E. C.: The human autosomal gene DAZLA: testis specificity and a candidate for male infertility. Hum. Molec. Genet. 5: 2013–2017, 1996.
- [72321] 10651. Akiyama, T. E.; Sakai, S.; Lambert, G.; Nicol, C. J.; Matsusue, K.; Pimprale, S.; Lee, Y.-H.; Ricote, M.; Glass, C. K.; Brewer, H. B., Jr.; Gonzalez, F. J.: Conditional disruption of the peroxisome proliferator-activated receptor gamma gene in mice results in lowered expression of ABCA1, ABCG1, and apoE in macrophages and reduced cholesterol efflux. Molec. Cell. Biol. 22: 2607–2619, 2002.
- [72322] 10652. Ardlie, K. G.; Lunetta, K. L.; Seielstad, M.: Testing for population subdivision and association in four case–



control studies. *Am. J. Hum. Genet.* 71: 304–311, 2002.

- [72323] 10653. Barak, Y.; Nelson, M. C.; Ong, E. S.; Jones, Y. Z.; Ruiz-Lozano, P.; Chien, K. R.; Koder, A.; Evans, R. M.: PPAR-gamma is required for placental, cardiac, and adipose tissue development. *Molec. Cell* 4:585–595, 1999.
- [72324] 10654. Barroso, I.; Gurnell, M.; Crowley, V. E. F.; Agostini, M.; Schwabe, J. W.; Soos, M. A.; Masien, G. L.; Williams, T. D. M.; Lewis, H.; Schafer, A. J.; Chatterjee, V. K. K.; O'Rahilly, S.: Dominant negative mutations in human PPAR-gamma associated with severe insulin resistance, diabetes mellitus and hypertension. *Nature* 402: 880–883, 1999.
- [72325] 10655. Beamer, B. A.; Yen, C.-J.; Andersen, R. E.; Muller, D.; Elahi, D.; Cheskin, L. J.; Andres, R.; Roth, J.; Shuldiner, A. R.: Association of the pro12ala variant in the peroxisome proliferator-activated receptor-gamma-2 gene with obesity in two Caucasian populations. *Diabetes* 47: 1806–1808, 1998.
- [72326] 10656. Beamer, B. A.; Negri, C.; Yen, C.-J.; Gavrilova, O.; Rumberger, J. M.; Durcan, M. J.; Yarnall, D. P.; Hawkins, A. L.; Griffin, C. A.; Burns, D. K.; Roth, J.; Reitman, M.; Shuldiner, A. R.: Chromosomal localization and partial genomic structure of the human peroxisome proliferator ac-

tivated receptor-gamma (hPPAR-gamma) gene.

Biochem.Biophys. Res. Commun. 233: 756-759, 1997.

- [72327] 10657.Chawla, A.; Boisvert, W. A.; Lee, C.-H.; Laffitte, B. A.; Barak,Y.; Joseph, S. B.; Liao, D.; Nagy, L.; Edwards, P. A.; Curtiss, L.K.; Evans, R. M.; Tontonoz, P.: A PPAR-gamma-LXR-ABCA1 pathway in macrophages is involved in cholesterol efflux and atherogenesis. Molec.Cell 7: 161-171, 2001.
- [72328] 10658.Deeb, S. S.; Fajas, L.; Nemoto, M.; Pihlajamaki, J.; Mykkanen,L.; Kuusisto, J.; Laakso, M.; Fujimoto, W.; Auwerx, J.: A pro12alasubstitution in PPAR-gamma-2 associated with decreased receptor activity, lower body mass index and improved insulin sensitivity. Nature Genet. 20:284-287, 1998.
- [72329] 10659.Elbrecht, A.; Chen, Y.; Cullinan, C. A.; Hayes, N.; Leibowitz,M. D.; Moller, D. E.; Berger, J.: Molecular cloning, expression and characterization of human peroxisome proliferator activated receptors gamma-1 and gamma-2. Biochem. Biophys. Res. Commun. 224: 431-437,1996.
- [72330] 10660.Fajas, L.; Auboeuf, D.; Raspe, E.; Schoonjans, K.; Lefebvre, A.M.; Saladin, R.; Najib, J.; Laville, M.; Fruchart, J.-C.; Deeb, S.; Vidal-Puig, A.; Flier, J.; Briggs, M. R.; Staels,

B.; Vidal, H.; Auwerx, J.: The organization, promoter analysis, and expression of the human PPAR-gamma gene. *J. Biol. Chem.* 272: 18779–18789, 1997.

- [72331] 10661. Gampe, R. T., Jr.; Montana, V. G.; Lambert, M. H.; Miller, A. B.; Bledsoe, R. K.; Milburn, M. V.; Kliewer, S. A.; Willson, T. M.; Xu, H. E.: Asymmetry in the PPAR-gamma/RXR-alpha crystal structure reveals the molecular basis of heterodimerization among nuclear receptors. *Molec. Cell* 5: 545–555, 2000.
- [72332] 10662. Chakraborti, A.; Lippman, D. L.; Loh, H. H.; Kozak, C. A.; Lee, N. M.: Genetic mapping of opioid binding protein gene(s) to mouse chromosome 9. *Mammalian Genome* 4: 179–182, 1993.
- [72333] 10663. Cho, T. M.; Hasegawa, J.-I.; Ge, B.-I.; Loh, H. H.: Purification to apparent homogeneity of a mu-type opioid receptor from rat brain. *Proc. Nat. Acad. Sci.* 83: 4138–4142, 1986.
- [72334] 10664. Shark, K. B.; Lee, N. M.: Cloning, sequencing and localization to chromosome 11 of a cDNA encoding a human opioid-binding cell adhesion molecule (OBCAM). *Gene* 155: 213–217, 1995.
- [72335] 10665. Burmeister, M.; Meyer, G. E.: The trefoil gene maps to mouse chromosome 17. *Mammalian Genome* 8:

223–224, 1997.

- [72336] 10666.Chinery, R.; Williamson, J.; Poulsom, R.: The gene encoding humanintestinal trefoil factor (TFF3) is located on chromosome 21q22.3clustered with other members of the trefoil peptide family. *Genomics* 32:281–284, 1996.
- [72337] 10667.Mashimo , H.; Wu, D.–C.; Podolsky, D. K.; Fishman, M. C.: Impaireddefense of intestinal mucosa in mice lacking intestinal trefoil factor. *Science* 274:262–265, 1996.
- [72338] 10668.Podolsky, D. K.; Lynch–Devaney, K.; Stow, J. L.; Oates, P.; Murgue,B.; DeBeaumont, M.; Sands, B. E.; Mahida, Y. R.: Identification ofhuman intestinal trefoil factor: goblet cell–specific expression ofa peptide targeted for apical secretion. *J. Biol. Chem.* 268: 6694–6702,1993.
- [72339] 10669.Probst, J. C.; Zetzsche, T.; Weber, M.; Theilemann, P.; Skutella,T.; Landgraf, R.; Jirikowski, G. F.: Human intestinal trefoil factoris expressed in human hypothalamus and pituitary: evidence for a novelneuropeptide. *FASEB J.* 10: 1518–1523, 1996.
- [72340] 10670.Schmitt, H.; Wundrack, I.; Beck, S.; Gott, P.; Welter, C.; Shizuya,H.; Simon, M. I.; Blin, N.: A third P–domain peptide gene (TFF3),human intestinal trefoil factor, maps to 21q22.3. *Cytogenet. CellGenet.* 72: 299–302, 1996.
- [72341] 10671.Taupin, D.; Wu, D.–C.; Jeon, W.–K.; Devaney, K.;

Wang, T. C.; Podolsky, D. K.: The trefoil gene family are coordinately expressed immediate-early genes: EGF receptor- and MAP kinase-dependent interregulation. *J. Clin. Invest.* 103: R31-R38, 1999.

[72342] 10672. Thim, L.: A new family of growth factor-like peptides: 'trefoil' disulphide loop structures as a common feature in breast cancer associated peptide (pS2), pancreatic spasmolytic polypeptide (PSP), and frog skin peptides (spasmolysins). *FEBS Lett.* 250: 85-90, 1989.

[72343] 10673. Sugimoto, Y.; Yatsunami, K.; Tsujimoto, M.; Khorana, H. G.; Ichikawa, A.: The amino acid sequence of a glutamic acid-rich protein from bovine retina as deduced from the cDNA sequence. *Proc. Nat. Acad. Sci.* 88: 3116-3119, 1991.

[72344] 10674. Marigo, V.; Roberts, D. J.; Lee, S. M. K.; Tsukurov, O.; Levi, T.; Gastier, J. M.; Epstein, D. J.; Gilbert, D. J.; Copeland, N. G.; Seidman, C. E.; Jenkins, N. A.; Seidman, J. G.; McMahon, A. P.; Tabin, C.: Cloning, expression, and chromosomal location of SHH and IHH: two human homologues of the *Drosophila* segment polarity gene hedgehog. *Genomics* 28:44-51, 1995.

[72345] 10675. McIntire, J. J.; Umetsu, S. E.; Akbari, O.; Potter, M.; Kuchroo, V. K.; Barsh, G. S.; Freeman, G. J.; Umetsu, D. T.;

DeKruyff, R. H.: Identification of Tapr (an airway hyperre-activity regulatory locus) and the linked Tim gene family. *Nature Immun.* 2: 1109–1116, 2001.

[72346] 10676. Hwang, B. J.; Ford, J. M.; Hanawalt, P. C.; Chu, G.: Expression of the p48 xeroderma pigmentosum gene is p53-dependent and is involved in global genomic repair. *Proc. Nat. Acad. Sci.* 96: 424–428, 1999.

[72347] 10677. Tahir, S. A.; Yang, G.; Ebara, S.; Timme, T. L.; Satoh, T.; Li, L.; Goltsov, A.; Ittmann, M.; Morrisett, J. D.; Thompson, T. C.: Secreted caveolin-1 stimulates cell survival/clonal growth and contributes to metastasis in androgen-insensitive prostate cancer. *Cancer Res.* 61: 3882–3885, 2001.

[72348] 10678. England, S. K.; Uebele, V. N.; Kodali, J.; Bennett, P. B.; Tamkun, M. M.: A novel K<sup>+</sup> channel beta-subunit (hKv-beta-1.3) is produced via alternative mRNA splicing. *J. Biol. Chem.* 270: 28531–28534, 1995.

[72349] 10679. Kratz, C. P.; Emerling, B. M.; Bonifas, J.; Wang, W.; Green, E. D.; Le Beau, M. M.; Shannon, K. M.: Genomic structure of the PIK3CG gene on chromosome band 7q22 and evaluation as a candidate myeloid tumor suppressor. *Blood* 99: 372–374, 2002.

[72350] 10680. Temple, I. K.; Gardner, R. J.; Robinson, D. O.;

Kibirige, M. S.; Ferguson, A. W.; Baum, J. D.; Barber, J. C. K.; James, R. S.; Shield, J. P. H.: Further evidence for an imprinted gene for neonatal diabetes localised to chromosome 6q22-q23. *Hum. Molec. Genet.* 5: 1117-1124, 1996.

[72351] 10681. Ahmad, W.; Li, S.; Chen, H.; Tuck-Muller, C. M.; Pittler, S. J.; Aronson, N. N., Jr.: Lysosomal chitobiase (CTB) and the G-protein gamma-5 subunit (GNG5) genes co-localize to human chromosome 1p22. *Cytogenet. Cell Genet.* 71: 44-46, 1995.

[72352] 10682. Fisher, K. J.; Aronson, N. N.: Characterization of the cDNA and genomic sequence of a G protein gamma subunit (gamma-5). *Molec. Cell Biol.* 12: 1585-1591, 1992.

[72353] 10683. Fiaschi, T.; Marzella, R.; Veggi, D.; Marzocchini, R.; Raugei, G.; Rocchi, M.; Ramponi, G.: Assignment of the human erythrocyte acylphosphatase gene (ACYP1) to chromosome band 14q24.3. *Cytogenet. Cell Genet.* 81: 235-236, 1998.

[72354] 10684. Fiaschi, T.; Raugei, G.; Marzocchini, R.; Chiarugi, P.; Cirri, P.; Ramponi, G.: Cloning and expression of the cDNA coding for the erythrocyte isoenzyme of human acylphosphatase. *FEBS Lett.* 367:145-148, 1995.

[72355] 10685. Goldowitz, D.; Smeyne, R. J.: Tune into the weaver channel. *Nature Genet.* 11: 107-109, 1995.

- [72356] 10686.Hess, E. J.: Identification of the weaver mouse mutation: the end of the beginning. *Neuron* 16: 1073–1076, 1996.
- [72357] 10687.Lane, P. W.: New mutation: Weaver, *wv*. *Mouse News Letter* 32–33, 1964.
- [72358] 10688.Lesage, F.; Duprat, F.; Fink, M.; Guillemare, E.; Coppola, T.; Lazdunski, M.; Hugnot, J.–P.: Cloning provides evidence for a family of inward rectifier and G–protein coupled K(+) channels in the brain. *FEBS Lett.* 353: 37–42, 1994.
- [72359] 10689.Rakic, P.; Sidman, R. L.: Sequence of developmental abnormalities leading to granule cell deficit in cerebellar cortex of weaver mutant mice. *J. Comp. Neurol.* 152: 103–132, 1973.
- [72360] 10690.Sakura, H.; Bond, C.; Warren–Perry, M.; Horsley, S.; Kearney, L.; Tucker, S.; Adelman, J.; Turner, R.; Ashcroft, F. M.: Characterization and variation of a human inwardly–rectifying K–channel gene (KCNJ6): a putative ATP–sensitive K–channel subunit. *FEBS Lett.* 367: 193–197, 1995.
- [72361] 10691.Tsaur, M.–L.; Menzel, S.; Lai, F.–P.; Espinosa, R., III; Concannon, P.; Spielman, R. S.; Hanis, C. L.; Cox, N. J.; Le Beau, M. M.; German, M. S.; Jan, L. Y.; Bell, G. I.; Stoffel, M.:



Isolation of a cDNA clone encoding a K(ATP) channel-like protein expressed in insulin-secreting cells, localization of the human gene to chromosome band 21q22.1 and linkage studies with NIDDM. *Diabetes* 44: 592–596, 1995.

[72362] 10692. Yasuda, K.; Sakura, H.; Mori, Y.; Iwamoto, K.; Shimokawa, K.; Kadowaki, H.; Hagura, R.; Akanuma, Y.; Adelman, J. P.; Yazaki, Y.; Ashcroft, F. M.; Kadowaki, T.: No evidence for mutations in a putative subunit of the beta-cell ATP-sensitive potassium channel (K-ATP channel) in Japanese NIDDM patients. *Biochem. Biophys. Res. Commun.* 211:1036–1040, 1995.

[72363] 10693. Efiok, B. J. S.; Chiorini, J. A.; Safer, B.: A key transcription factor for eukaryotic initiation factor-2-alpha is strongly homologous to developmental transcription factors and may link metabolic genes to cellular growth and development. *J. Biol. Chem.* 269: 18921–18930, 1994.

[72364] 10694. Gopalakrishnan, L.; Scarpulla, R. C.: Structure, expression, and chromosomal assignment of the human gene encoding nuclear respiratory factor 1. *J. Biol. Chem.* 270: 18019–18025, 1995.

[72365] 10695. Spelbrink, J. N.; Van den Bogert, C.: The pre-mRNA of nuclear respiratory factor 1, a regulator of mitochondrial biogenesis, is alternatively spliced in human tissues

and cell lines. Hum. Molec.Genet. 4: 1591–1596, 1995.

[72366] 10696.Tiranti, V.; Rossi, E.; Rocchi, M.; DiDonato, S.; Zufardi, O.;Zeviani, M.: The gene (NFE2L1) for human NRF–1, an activator involvedin nuclear–mitochondrial interactions, maps to 7q32. Genomics 27:555–557, 1995.

[72367] 10697.Ohshiro, T.; Yagami, T.; Zhang, C.; Matsuzaki, F.: Role of corticaltumour–suppressor proteins in asymmetric division of Drosophila neuroblast. Nature 408:593–596, 2000.

- [72368] 10698.Peng, C.-Y.; Manning, L.; Albertson, R.; Doe, C. Q.: The tumour-suppressorgenes lgl and dlg regulate basal protein targeting in Drosophila neuroblasts. *Nature* 408:596–600, 2000.
- [72369] 10699.Strand, D.; Unger, S.; Corvi, R.; Hartenstein, K.; Schenkel, H.;Kalmes, A.; Merdes, G.; Neumann, B.; Krieg-Schneider, F.; Coy, J.F.; Poustka, A.; Schwab, M.; Mechler B. M.: A human homologue ofthe Drosophila tumour suppressor gene l(2)gl maps to 17p11.2–12 andcodes for a cytoskeletal protein that associates with nonmuscle myosinII heavy chain. *Oncogene* 11: 291–301, 1995.
- [72370] 10700.Timchenko, L. T.; Miller, J. W.; Timchenko, N. A.; DeVore, D. R.;Datar, K. V.; Lin, L.; Roberts, R.; Caskey, C. T.; Swanson, M. S.: Identification of a (CUG)<sub>n</sub> triplet repeat RNA-binding protein andits expression in myotonic dystrophy. *Nucleic Acids Res.* 24: 4407–4414,1996.
- [72371] 10701.Timchenko, L. T.; Timchenko, N. A.; Caskey, C. T.; Roberts, R.: Novel proteins with binding specificity for DNA CTG repeats andRNA CUG repeats: implications for myotonic dystrophy. *Hum. Molec.Genet.* 5: 115–121, 1996.
- [72372] 10702.Wary, K. K.; Mariotti, A.; Zurzolo, C.; Giancotti, F. G.: A requirementfor caveolin-1 and associated kinase Fyn

in integrin signaling and anchorage-dependent cell growth. *Cell* 94: 625–634, 1998.

[72373] 10703. Yang, G.; Truong, L. D.; Timme, T. L.; Ren, C.; Wheeler, T. M.; Park, S. H.; Nasu, Y.; Bangma, C. H.; Kattan, M. W.; Scardino, P. T.; Thompson, T. C.: Elevated expression of caveolin is associated with prostate and breast cancer. *Clin. Cancer Res.* 4: 1873–1880, 1998.

[72374] 10704. Meij, I. C.; Koenderink, J. B.; van Bokhoven, H.; Assink, K. F. H.; Groenestege, W. T.; de Pont, J. J. H. H. M.; Bindels, R. J. M.; Monnens, L. A. H.; van den Heuvel, L. P. W. J.; Knoers, N. V. A. M.: Dominant isolated renal magnesium loss is caused by misrouting of the Na<sup>+</sup>,K<sup>+</sup>-ATPase gamma-subunit. *Nature Genet.* 26: 265–266, 2000. Note: Erratum. *Nature Genet.* 27: 125 only, 2001.

[72375] 10705. Sweadner, K. J.; Rael, E.: The FXYD gene family of small ion transport regulators or channels: cDNA sequence, protein signature sequence, and expression. *Genomics* 68: 41–56, 2000.

[72376] 10706. Sweadner, K. J.; Wetzel, R. K.; Arystarkhova, E.: Genomic organization of the human FXYD2 gene encoding the gamma subunit of the Na,K-ATPase. *Biochem. Biophys. Res. Commun.* 279: 196–201, 2000.

[72377] 10707. Blair, E.; Redwood, C.; Ashrafian, H.; Oliveira, M.;

Broxholme, J.; Kerr, B.; Salmon, A.; Ostman-Smith, I.; Watkins, H.: Mutations in the gamma-2 subunit of AMP-activated protein kinase cause familial hypertrophic cardiomyopathy: evidence for the central role of energy-compromise in disease pathogenesis. *Hum. Molec. Genet.* 10: 1215-1220, 2001.

[72378] 10708. Siderovski, D. P.; Heximer, S. P.; Forsdyke, D. R.: A human gene encoding a putative basic helix-loop-helix phosphoprotein whose mRNA increases rapidly in cycloheximide-treated blood mononuclear cells. *DNACell Biol.* 13: 125-147, 1994.

[72379] 10709. Hirai, H.; Roussel, M. F.; Kato, J.-Y.; Ashmun, R. A.; Sherr, C. J.: Novel INK4 proteins, p19 and p18, are specific inhibitors of cyclin D-dependent kinases CDK4 and CDK6. *Molec. Cell. Biol.* 15: 2672-2681, 1995.

[72380] 10710. Okuda, T.; Hirai, H.; Valentine, V. A.; Shurtleff, S. A.; Kidd, V. J.; Lahti, J. M.; Sherr, C. J.; Downing, J. R.: Molecular cloning, expression pattern, and chromosomal localization of human CDKN2D/INK4d, an inhibitor of cyclin D-dependent kinases. *Genomics* 29: 623-630, 1995.

[72381] 10711. Sago, H.; Kitagawa, M.; Obata, S.; Mori, N.; Taketani, S.; Rochelle, J. M.; Seldin, M. F.; Davidson, M.; St. John, T.; Suzuki, S. T.: Cloning, expression, and chromoso-

mal localization of a novel cadherin-related protein, protocadherin-3. *Genomics* 29: 631–640, 1995.

[72382] 10712. Fiorucci, S.; Mencarelli, A.; Palazzetti, B.; Distrutti, E.; Vergnolle, N.; Hollenberg, M. D.; Wallace, J. L.; Morelli, A.; Cirino, G.: Proteinase-activated receptor 2 is an anti-inflammatory signal for colonic lamina propria lymphocytes in a mouse model of colitis. *Proc. Nat. Acad. Sci.* 98:13936–13941, 2001.

[72383] 10713. Nystedt, S.; Emilsson, K.; Larsson, A.-K.; Strombeck, B.; Sundelin, J.: Molecular cloning and functional expression of the gene encoding the human proteinase activated receptor 2. *Europ. J. Biochem.* 232:84–89, 1995.

[72384] 10714. Nystedt, S.; Emilsson, K.; Wahlestedt, C.; Sundelin, J.: Molecular cloning of a potential proteinase-activated receptor. *Proc. Nat. Acad. Sci.* 91: 9208–9212, 1994.

[72385] 10715. Devlin, A. M.; Ling, E.; Peerson, J. M.; Fernando, S.; Clarke, R.; Smith, A. D.; Halsted, C. H.: Glutamate carboxypeptidase II: a polymorphism associated with lower levels of serum folate and hyperhomocysteinemia. *Hum. Molec. Genet.* 9: 2837–2844, 2000.

[72386] 10716. Israeli, R. S.; Powell, C. T.; Fair, W. R.; Heston, W. D. W.: Molecular cloning of a complementary DNA encoding a

prostate-specific membrane antigen. *Cancer Res.* 53: 227–230, 1993.

[72387] 10717. Leek, J.; Lench, N.; Maraj, B.; Bailey, A.; Carr, I. M.; Andersen, S.; Cross, J.; Whelan, P.; MacLennan, K. A.; Meredith, D. M.; Markham, A. F.: Prostate-specific membrane antigen: evidence for the existence of a second related human gene. *Brit. J. Cancer* 72: 583–588, 1995.

[72388] 10718. Maraj, B. H.; Leek, J. P.; Karayi, M.; Ali, M.; Lench, N. J.; Markham, A. F.: Detailed genetic mapping around a putative prostate-specific membrane antigen locus on human chromosome 11p11.2. *Cytogenet. Cell Genet.* 81: 3–9, 1998.

[72389] 10719. Ikemoto, M.; Arai, H.; Feng, D.; Tanaka, K.; Aoki, J.; Dohmae, N.; Takio, K.; Adachi, H.; Tsujimoto, M.; Inoue, K.: Identification of a PDZ-domain-containing protein that interacts with the scavenger receptor class B type I. *Proc. Nat. Acad. Sci.* 97: 6538–6543, 2000.

[72390] 10720. O'Keefe, D. S.; Su, S. L.; Bacich, D. J.; Horiguchi, Y.; Luo, Y.; Powell, C. T.; Zandvliet, D.; Russell, P. J.; Molloy, P. L.; Nowak, N. J.; Shows, T. B.; Mullins, C.; Vonder Haar, R. A.; Fair, W. R.; Heston, W. D. W.: Mapping, genomic organization and promoter analysis of the human prostate-specific membrane antigen gene. *Biochim. Biophys. Acta*

1443: 113–127, 1998.

- [72391] 10721.Rinker–Schaeffer, C. W.; Hawkins, A. L.; Su, S. L.; Israeli, R.S.; Griffin, C. A.; Isaacs, J. T.; Heston, W. D. W.: Localization and physical mapping of the prostate-specific membrane antigen (PSM) gene to human chromosome 11. *Genomics* 30: 105–108, 1995.
- [72392] 10722.Beltrame, J. F.; Sasayama, S.; Maseri, A.: Racial heterogeneity in coronary artery vasomotor reactivity: differences between Japanese and Caucasian patients. *J. Am. Coll. Cardiol.* 33: 1442–1452, 1999.
- [72393] 10723.Inagaki, N.; Inazawa, J.; Seino, S.: cDNA sequence, gene structure, and chromosomal localization of the human ATP-sensitive potassium channel, uK(ATP)–1, gene (KCNJ8). *Genomics* 30: 102–104, 1995.
- [72394] 10724.Inagaki, N.; Tsuura, Y.; Namba, N.; Masuda, K.; Gono, T.; Horie, M.; Seino, Y.; Mizuta, M.; Seino, S.: Cloning and functional characterization of a novel ATP-sensitive potassium channel ubiquitously expressed in rat tissues, including pancreatic islets, pituitary, skeletal muscle, and heart. *J. Biol. Chem.* 270: 5691–5694, 1995.
- [72395] 10725.MacAlpin, R. N.: Cardiac arrest and sudden unexpected death in variant angina: complications of coronary spasm that can occur in the absence of severe organic



coronary stenosis. *Am. Heart J.* 125:1011–1017, 1993.

[72396] 10726. Bartsch, J. W.; Mukai, H.; Takahashi, N.; Ronsiek, M.; Fuchs, S.; Jockusch, H.; Ono, Y.: The protein kinase N (PKN) gene PRKCL1/Prkcl1 maps to human chromosome 19p12–p13.1 and mouse chromosome 8 with close linkage to the myodystrophy (myd) mutation. *Genomics* 49: 129–132, 1998.

[72397] 10727. Mukai, H.; Ono, Y.: A novel protein kinase with leucine zipper-like sequences: its catalytic domain is highly homologous to that of protein kinase C. *Biochem. Biophys. Res. Commun.* 199: 897–904, 1994.

[72398] 10728. Palmer, R. H.; Ridden, J.; Parker, P. J.: Identification of multiple, novel, protein kinase C-related gene products. *FEBS Lett.* 356: 5–8, 1994.

[72399] 10729. Palmer, R. H.; Ridden, J.; Parker, P. J.: Cloning and expression patterns of two members of a novel protein-kinase-C-related kinase family. *Europ. J. Biochem.* 227: 344–351, 1995.

[72400] 10730. Yu, W.; Liu, J.; Morrice, N. A.; Wettenhall, R. E. H.: Isolation and characterization of a structural homologue of human PRK2 from rat liver. *J. Biol. Chem.* 272: 10030–10034, 1997.

[72401] 10731. Durkin, M. E.; Loechel, F.; Mattei, M.-G.; Gilpin, B.

J.; Albrechtsen, R.; Wewer, U. M.: Tissue-specific expression of the human laminin  $\alpha$ -5-chain, and mapping of the gene to human chromosome 20q13.2-13.3 and to distal mouse chromosome 2 near the locus for the ragged (Ra) mutation. FEBS Lett. 411: 296-300, 1997.

[72402] 10732. Miner, J. H.; Lewis, R. M.; Sanes, J. R.: Molecular cloning of a novel laminin chain,  $\alpha$ -5, and widespread expression in adult mouse tissues. J. Biol. Chem. 270: 28523-28526, 1995.

[72403] 10733. Hernandez, A.; Park, J. P.; Lyon, G. J.; Mohandas, T. K.; St. Germain, D. L.: Localization of the type 3 iodothyronine deiodinase (DIO3) gene to human chromosome 14q32 and mouse chromosome 12F1. Genomics 53:119-121, 1998.

[72404] 10734. Huang, S. A.; Tu, H. M.; Harney, J. W.; Venihaki, M.; Butte, A. J.; Kozakewich, H. P. W.; Fishman, S. J.; Larsen, P. R.: Severe hypothyroidism caused by type 3 iodothyronine deiodinase in infantile hemangiomas. New Eng. J. Med. 343: 185-189, 2000.

[72405] 10735. Salvatore, D.; Low, S. C.; Berry, M.; Maia, A. L.; Harney, J. W.; Croteau, W.; St. German, D. L.; Larsen, P. R.: Type 3 iodothyronine deiodinase: cloning, in vitro expression, and functional analysis of the placental selenoen-

zyme. J. Clin. Invest. 96: 2421–2430, 1995.

[72406] 10736.Kolodkin, A. L.; Matthes, D. J.; Goodman, C. S.: The semaphorins encode a family of transmembrane and secreted growth cone guidance molecules. Cell 75: 1389–1399, 1993.

[72407] 10737.Luo, Y.; Raible, D.; Raper, J. A.: Collapsin: a protein in brain that induces the collapse and paralysis of neuronal growth cones. Cell 75:217–227, 1993.

[72408] 10738.Marin, O.; Yaron, A.; Bagri, A.; Tessier-Lavigne, M.; Rubenstein, J. L. R.: Sorting of striatal and cortical interneurons regulated by semaphorin–neuropilin interactions. Science 293: 872–875, 2001.

[72409] 10739.Sekido, Y.; Bader, S.; Latif, F.; Chen, J.–Y.; Duh, F.–M.; Wei, M.–H.; Albanesi, J. P.; Lee, C.–C.; Lerman, M. I.; Minna, J. D.: Human semaphorins A(V) and IV reside in the 3p21.3 small cell lung cancer deletion region and demonstrate distinct expression patterns. Proc.Nat. Acad. Sci. 93: 4120–4125, 1996.

[72410] 10740.Xiang, R.–H.; Hensel, C. H.; Garcia, D. K.; Carlson, H. C.; Kok, K.; Daly, M. C.; Kerbacher, K.; van den Berg, A.; Veldhuis, P.; Buys, C. H. C. M.; Naylor S. L.: Isolation of the human semaphorin III/F gene (SEMA3F) at chromosome 3p21, a region deleted in lung cancer. Genomics

32:39–48, 1996.

- [72411] 10741.Ardehali, H.; Tiller, G. E.; Printz, R. L.; Mochizuki, H.; Prochazka,M.; Granner, D. K.: A novel (TA)<sub>n</sub> polymorphism in the hexokinase II gene: application to noninsulin-dependent diabetes mellitus in the Pima Indians. *Hum. Genet.* 97: 482–485, 1996.
- [72412] 10742.Echwald, S. M.; Bjorbaek, C.; Hansen, T.; Clausen, J. O.; Vestergaard,H.; Zierath, J. R.; Printz, R. L.; Granner, D. K.; Pedersen, O.:Identification of four amino acid substitutions in hexokinase II and studies of relationships to NIDDM, glucose effectiveness, and insulin sensitivity. *Diabetes* 44: 347–353, 1995.
- [72413] 10743.Heikkinen, S.; Suppola, S.; Malkki, M.; Deeb, S. S.; Janne, J.;Laakso, M.: Mouse hexokinase II gene: structure, cDNA, promoter analysis,and expression pattern. *Mammalian Genome* 11: 91–96, 2000.
- [72414] 10744.Laakso, M.; Malkki, M.; Deeb, S. S.: Amino acid substitutions in hexokinase II among patients with NIDDM. *Diabetes* 44: 330–334,1995.
- [72415] 10745.Lehto, M.; Xiang, K.; Stoffel, M.; Espinosa, R., III; Groop, L.C.; Le Beau, M. M.; Bell, G. I.: Human hexokinase II: localization of the polymorphic gene to chromosome 2. *Diabetologia* 36: 1299–1302,1993.

- [72416] 10746.Mathupala, S. P.; Heese, C.; Pedersen, P. L.: Glucose catabolism in cancer cells: the type II hexokinase promoter contains functionally active response elements for the tumor suppressor p53. *J. Biol. Chem.* 272:22776–22780, 1997.
- [72417] 10747.Mathupala, S. P.; Rempel, A.; Pedersen, P. L.: Glucose catabolism in cancer cells: isolation, sequence, and activity of the promoter for type II hexokinase. *J. Biol. Chem.* 270: 16918–16925, 1995.
- [72418] 10748.Vidal-Puig, A.; Printz, R. L.; Stratton, I. M.; Granner, D. K.; Moller, D. E.: Analysis of the hexokinase II gene in subjects with insulin resistance and NIDDM and detection of a gln142-to-his substitution. *Diabetes* 44:340–346, 1995.
- [72419] 10749.Luo, Y.; Roeder, R. G.: Cloning, functional characterization, and mechanism of action of the B-cell-specific transcriptional activator OCA-B. *Molec. Cell. Biol.* 15: 4115–4124, 1995.
- [72420] 10750.Staudt, L. M.; Lenardo, M. J.: Immunoglobulin gene transcription. *Ann.Rev. Immun.* 9: 373–398, 1991.
- [72421] 10751.Strubin, M.; Newell, J. W.; Matthias, P.: OBF-1, a novel B cell-specific coactivator that stimulates immunoglobulin promoter activity through association with

octamer-binding proteins. *Cell* 80: 497–506, 1995.

[72422] 10752.Hahm, K.; Kim, G.; Turck, C. W.; Smale, S. T.: Isolation of amurine gene encoding a nucleic acid-binding protein with homology to hnRNP K. *Nucleic Acids Res.* 21: 3894 only, 1993.

[72423] 10753.Leffers, H.; Dejgaard, K.; Celis, J. E.: Characterisation of two major cellular poly(rC)-binding human proteins, each containing three K-homologous (KH) domains. *Europ. J. Biochem.* 230: 447–453, 1995.

[72424] 10754.Bouchard, M. J.; Wang, L.-H.; Schneider, R. J.: Calcium signaling by HBx protein in hepatitis B virus DNA replication. *Science* 294:2376–2378, 2001.

[72425] 10755.Calalb, M. B.; Polte, T. R.; Hanks, S. K.: Tyrosine phosphorylation of focal adhesion kinase at sites in the catalytic domain regulates kinase activity: a role for Src family kinases. *Molec. Cell. Biol.* 15:954–963, 1995.

[72426] 10756.Ganem, D.: The X files—one step closer to closure. *Science* 294:2299–2300, 2001.

[72427] 10757.Herzog, H.; Nicholl, J.; Hort, Y. J.; Sutherland, G. R.; Shine, J.: Molecular cloning and assignment of FAK2, a novel human focal adhesion kinase, to 8p11.2–p22 by nonisotopic in situ hybridization. *Genomics* 32:484–486, 1996.

- [72428] 10758.Lev, S.; Moreno, H.; Martinez, R.; Canoll, P.; Peles, E.; Musacchio, J. M.; Plowman, G. D.; Rudy, B.; Schlessinger, J.: Protein tyrosine kinase PYK2 involved in  $\text{Ca}^{2+}$ -induced regulation of ion channel and MAP kinase functions. *Nature* 376: 737–745, 1995.
- [72429] 10759.Manser, E.; Leung, T.; Salihuddin, H.; Tan, L.; Lim, L.: A non-receptor tyrosine kinase that inhibits the GTPase activity of p21(cdc42). *Nature* 363:364–367, 1993.
- [72430] 10760.Matsuya, M.; Sasaki, H.; Aoto, H.; Mitaka, T.; Nagura, K.; Ohba, T.; Ishino, M.; Takahashi, S.; Suzuki, R.; Sasaki, T.: Cell adhesion kinase beta forms a complex with a new member, Hic-5, of proteins localized at focal adhesions. *J. Biol. Chem.* 273: 1003–1014, 1998.
- [72431] 10761.Aubry, F.; Mattei, M.-G.; Barque, J.-P.; Galibert, F.: Chromosomal localization and expression pattern of the RNase L inhibitor gene. *FEBS Lett.* 381: 135–139, 1996.
- [72432] 10762.Diriong, S.; Salehzada, T.; Bisbal, C.; Martinand, C.; Taviaux, S.: Localization of the ribonuclease L inhibitor gene (RNS4I), a new member of the interferon-regulated 2–5A pathway, to 4q31 by fluorescence in situ hybridization. *Genomics* 32: 488–490, 1996.
- [72433] 10763.Meyer, J.; Wirth, J.; Held, M.; Schempp, W.; Scherer, G.: SOX20, a new member of the SOX gene family, is lo-

cated on chromosome 17p13. *Cytogenet. Cell Genet.* 72: 246–249, 1996.

- [72434] 10764. Vujic, M.; Rajic, T.; Goodfellow, P. N.; Stevanovic, M.: cDNA characterization and high resolution mapping of the human SOX20 gene. *Mammalian Genome* 9: 1059–1061, 1998.
- [72435] 10765. Ide, H.; Saito–Ohara, F.; Ohnami, S.; Osada, Y.; Ikeuchi, T.; Yoshida, T.; Terada, M.: Assignment of the BMPR1A and BMPR1B genes to human chromosome 10q22.3 and 4q23–q24 by in situ hybridization and radiation hybrid mapping. *Cytogenet. Cell Genet.* 81: 285–286, 1998.
- [72436] 10766. Schuffenhauer, S.; Lichtner, P.; Peykar–Derakhshandeh, P.; Murken, J.; Haas, O. A.; Back, E.; Wolff, G.; Zabel, B.; Barisic, I.; Rauch, A.; Borochowitz, Z.; Dallapiccola, B.; Ross, M.; Meitinger, T.: Deletion mapping on chromosome 10p and definition of a critical region for the second DiGeorge syndrome locus (DGS2). *Europ. J. Hum. Genet.* 6: 213–225, 1998.
- [72437] 10767. Pizzuti, A.; Amati, F.; Calabrese, G.; Mari, A.; Colosimo, A.; Silani, V.; Giardino, L.; Ratti, A.; Penso, D.; Calza, L.; Palka, G.; Scarlato, G.; Novelli, G.; Dallapiccola, B.: cDNA characterization and chromosomal mapping of two



human homologs of the *Drosophila* dishevelled polarity-gene. *Hum. Molec. Genet.* 5: 953–958, 1996.

[72438] 10768.Semenov, M. V.; Snyder, M.: Human dishevelled genes constitute a DHR-containing multigene family. *Genomics* 42: 302–310, 1997.

[72439] 10769.Lee, J.; Hahn, Y.; Yun, J. H.; Mita, K.; Chung, J. H.: Characterization of JDP genes, an evolutionarily conserved J domain-only protein family, from human and moths. *Biochim. Biophys. Acta* 1491: 355–363, 2000.

[72440] 10770.Qiu, J.; Qian, Y.; Chen, V.; Guan, M.-X.; Shen, B.: Human exonuclease 1 functionally complements its yeast homologues in DNA recombination, RNA primer removal, and mutation avoidance. *J. Biol. Chem.* 274:17893–17900, 1999.

[72441] 10771.Schmutte, C.; Marinescu, R. C.; Sadoff, M. M.; Guerrette, S.; Overhauser, J.; Fishel, R.: Human exonuclease I interacts with the mismatch repair protein hMSH2. *Cancer Res.* 58: 4537–4542, 1998.

[72442] 10772.Tishkoff, D. X.; Amin, N. S.; Viars, C. S.; Arden, K. C.; Kolodner, R. D.: Identification of a human gene encoding a homologue of *Saccharomyces cerevisiae* EXO1, an exonuclease implicated in mismatch repair and recombination. *Cancer Res.* 58: 5027–5031, 1998.

- [72443] 10773.Wilson, D. M., III; Carney, J. P.; Coleman, M. A.; Adamson, A.W.; Christensen, M.; Lamerdin, J. E.: Hex1: a new human Rad2 nucleasefamily member with homology to yeast exonuclease 1. *Nucleic AcidsRes.* 26: 3762–3768, 1998.
- [72444] 10774.Carson–Walter, E. B.; Watkins, D. N.; Nanda, A.; Vogelstein, B.;Kinzler, K. W.; St. Croix, B.: Cell surface tumor endothelial markersare conserved in mice and humans. *Cancer Res.* 61: 6649–6655, 2001.
- [72445] 10775.Christian, S.; Ahorn, H.; Koehler, A.; Eisenhaber, F.; Rodi, H.–P.;Garin–Chesa, P.; Park, J. E.; Rettig, W. J.; Lenter, M. C.: Molecularcloning and characterization of endosialin, a C-type lectin–like cellsurface receptor of tumor endothelium. *J. Biol. Chem.* 276: 7408–7414,2001.
- [72446] 10776.St. Croix, B.; Rago, C.; Velculescu, V.; Traverso, G.; Romans,K. E.; Montgomery, E.; Lal, A.; Riggins, G. J.; Lengauer, C.; Vogelstein,B.; Kinzler, K. W.: Genes expressed in human tumor endothelium. *Science* 289:1197–1202, 2000.
- [72447] 10777.Gosling, J.; Dairaghi, D. J.; Wang, Y.; Hanley, M.; Talbot, D.;Miao, Z.; Schall, T. J.: Cutting edge: identification of a novelchemokine receptor that binds dendritic cell– and T cell–active chemokinesincluding ELC, SLC, and

TECK. J. Immun. 164: 2851–2856, 2000.

[72448] 10778.Cohn, R. D.; Campbell, K. P.: Molecular basis of muscular dystrophies. MuscleNerve 23: 1456–1471, 2000.

[72449] 10779.Gillespie, C. S.; Lee, M.; Fantes, J. F.; Brophy, P. J.: The gene encoding the Schwann cell protein periaxin localizes on mouse chromosome 7 (Prx). Genomics 41: 297–298, 1997.

[72450] 10780.Gillespie, C. S.; Sherman, D. L.; Fleetwood–Walker, S. M.; Cottrell, D. F.; Tait, S.; Garry, E. M.; Wallace, V. C. J.; Ure, J.; Griffiths, I. R.; Smith, A. Brophy, P. J.: Peripheral demyelination and neuropathic pain behavior in periaxin-deficient mice. Neuron 26: 523–531, 2000.

[72451] 10781.Schuler, G. D.: Sequence mapping by electronic PCR. Genome Res. 7:541–550, 1997.

[72452] 10782.Takashima, H.; Boerkoel, C. F.; De Jonghe, P.; Ceuterick, C.; Martin, J.–J.; Voit, T.; Schroder, J.–M.; Williams, A.; Brophy, P. J.; Timmerman, V.; Lupski, J. R.: Periaxin mutations cause a broad spectrum of demyelinating neuropathies. Ann. Neurol. 51: 709–715, 2002.

[72453] 10783.Liu, Q.–Y.; Wang, L. F.; Miao, S. Y.; Catterall, J. F.: Expression and characterization of a novel human sperm membrane protein. Biol.Reprod. 54: 323–330, 1996.

[72454] 10784.Miao, S.; Yan, Y.; Li, Y.; Bai, Y.; Wei, S.; Zong, C.;

Zhao, M.;Zong, S.; Wang, L.: cDNA encoding a human sperm membrane proteinBS-84. Prog. Natural Sci. 5: 119-122, 1995.

[72455] 10785.Wang, H.; Miao, S.; Chen, D.; Wang, L.; Koide, S. S.: Assignmentof chromosomal locus and evidence for alternatively spliced mRNAs of a human sperm membrane protein (hSMP-1). Biochim. Biophys. Acta 1447:119-124, 1999.

[72456] 10786.Horie, M.; Mitsumoto, Y.; Kyushiki, H.; Kanemoto, N.; Watanabe,A.; Taniguchi, Y.; Nishino, N.; Okamoto, T.; Kondo, M.; Mori, T.;Noguchi, K.; Nakamura, Y.; Takahashi, E.; Tanigami, A.: Identificationand characterization of TM-EFF2, a novel survival factor for hippocampaland mesencephalic neurons. Genomics 67: 146-152, 2000.

[72457] 10787.Uchida, T.; Wada, K.; Akamatsu, T.; Yonezawa, M.; Noguchi, H.;Mizoguchi, A.; Kasuga, M.; Sakamoto, C.: A novel epidermal growthfactor-like molecule containing two follistatin modules stimulatestyrosine phosphorylation of erbB-4 in MKN28 gastric cancer cells. Biochem.Biophys. Res. Commun. 266: 593-602, 1999.

[72458] 10788.Young, J.; Biden, K. G.; Simms, L. A.; Huggard, P.; Karamatic,R.; Eyre, H. J.; Sutherland, G. R.; Herath, N.; Barker, M.; Anderson,G. J.; Fitzpatrick, D. R.; Ramm, G. A.;

Jass, J. R.; Leggett, B. A.: HPP1: a transmembrane protein-encoding gene commonly methylated in colorectal polyps and cancers. *Proc. Nat. Acad. Sci.* 98: 265–270, 2001.

[72459] 10789. Blasina, A.; de Weyer, I. V.; Laus, M. C.; Luyten, W. H.; Parker, A. E.; McGowan, C. H.: A human homologue of the checkpoint kinase Cds1 directly inhibits Cdc25 phosphatase. *Curr. Biol.* 14: 1–10, 1999.

[72460] 10790. Brown, A. L.; Lee, C.-H.; Schwarz, J. K.; Mitiku, N.; Piwnicka-Worms, H.; Chung, J. H.: A human Cds1-related kinase that functions downstream of ATM protein in the cellular response to DNA damage. *Proc. Nat. Acad. Sci.* 96: 3745–3750, 1999.

[72461] 10791. Chaturvedi, P.; Eng, W. K.; Zhu, Y.; Mattern, M. R.; Mishra, R.; Hurle, M. R.; Zhang, X.; Annan, R. S.; Lu, Q.; Faucette, L. F.; Scott, G. F.; Li, X.; Carr, S. A.; Johnson, R. K.; Winkler, J. D.; Zhou, B. B.: Mammalian Chk2 is a downstream effector of the ATM-dependent DNA damage checkpoint pathway. *Oncogene* 18: 4047–4054, 1999.

[72462] 10792. Chehab, N. H.; Malikzay, A.; Appel, M.; Hazonetis, T. D.: Chk2/hCds1 functions as a DNA damage checkpoint in G-1 by stabilizing p53. *Genes Dev.* 14: 278–288, 2000.

[72463] 10793. Ino, Y.; Wahrer, D. C. R.; Bell, D. W.; Haber, D. A.;

Louis, D.N.: Mutation analysis of the hCHK2 gene in primary human malignant gliomas. (Letter) *Neurogenetics* 3: 45–46, 2000.

[72464] 10794. Lee, S. B.; Kim, S. H.; Bell, D. W.; Wahrer, D. C. R.; Schiripo, T. A.; Jorczak, M. M.; Sgroi, D. C.; Garber, J. E.; Li, F. P.; Nichols, K. E.; Varley, J. M.; Godwin, A. K.; Shannon, K. M.; Harlow, E.; Haber, D. A.: Destabilization of CHK2 by a missense mutation associated with Li–Fraumeni syndrome. *Cancer Res.* 61: 8062–8067, 2001.

[72465] 10795. Lopes, M.; Cotta–Ramusino, C.; Pelliccioli, A.; Liberi, G.; Plevani, P.; Muzi–Falconi, M. Newlon, C. S.; Foiani, M.: The DNA replication checkpoint response stabilizes stalled replication forks. *Nature* 412:557–561, 2001.

[72466] 10796. Matsuoka, S.; Huang, M.; Elledge, S. J.: Linkage of ATM to cell cycle regulation by the Chk2 protein kinase. *Science* 282: 1893–1897, 1998.

[72467] 10797. Meijers–Heijboer, H.; van den Ouweland, A.; Klijn, J.; Wasielewski, M.; de Snoo, A.; Oldenburg, R.; Hollestelle, A.; Houben, M.; Crepin, E.; van Veghel–Plandsoen, M.; Elstrodt, F.; van Duijn, C.; and 29 others: Low–penetrance susceptibility to breast cancer due to CHEK2\*1100delC in noncarriers of BRCA1 or BRCA2 mutations. *Nature Genet.* 31: 55–59, 2002.

- [72468] 10798. Miller, C. W.; Ikezoe, T.; Krug, U.; Hofmann, W.-K.; Tavor, S.; Vegesna, V.; Tsukasaki, K.; Takeuchi, S.; Koeffler, H. P.: Mutations of the CHK2 gene are found in some osteosarcomas, but are rare in breast, lung, and ovarian tumors. *Genes Chromosomes Cancer* 33: 17–21, 2002.
- [72469] 10799. Sogo, J. M.; Lopes, M.; Foiani, M.: Fork reversal and ssDNA accumulation at stalled replication forks owing to checkpoint defects. *Science* 297:599–602, 2002.
- [72470] 10800. Vahteristo, P.; Bartkova, J.; Eerola, H.; Syrjäkoski, K.; Ojala, S.; Kilpivaara, O.; Tamminen, A.; Kononen, J.; Aittomäki, K.; Heikkilä, P.; Holli, K.; Blomqvist, C.; Bartek, J.; Kallioniemi, O.-P.; Nevanlinna, H.: A CHEK2 genetic variant contributing to a substantial fraction of familial breast cancer. *Am. J. Hum. Genet.* 71: 432–438, 2002.
- [72471] 10801. Vahteristo, P.; Tamminen, A.; Karvinen, P.; Eerola, H.; Eklund, C.; Aaltonen, L. A.; Blomqvist, C.; Aittomäki, K.; Nevanlinna, H.: p53, CHK2, and CHK1 genes in Finnish families with Li–Fraumeni syndrome: further evidence of CHK2 in inherited cancer predisposition. *Cancer Res.* 61: 5718–5722, 2001.
- [72472] 10802. De, S. K.; Enders, G. C.; Andrews, G. K.: High levels of metallothionein messenger RNAs in male germ cells of the adult mouse. *Molec. Endocr.* 5:628–636, 1991.

- [72473] 10803.Salehi-Ashtiani, K.; Widrow, R. J.; Markert, C. L.; Goldberg, E.: Testis-specific expression of a metallothionein I-driven transgene correlates with undermethylation of the locus in testicular DNA. *Proc.Nat. Acad. Sci.* 90: 8886–8890, 1993.
- [72474] 10804.Sugihara, T.; Wadhwa, R.; Kaul, S. C.; Mitsui, Y.: A novel testis-specific metallothionein-like protein, tesmin, is an early marker of male germcell differentiation. *Genomics* 57: 130–136, 1999.
- [72475] 10805.Larrouy, D.; Vidal, H.; Andreelli, F.; Laville, M.; Langin, D.: Cloning and mRNA tissue distribution of human PPAR-gamma coactivator-1. *Int.J. Obesity* 23: 1327–1332, 1999.
- [72476] 10806.Lin, J.; Wu, H.; Tarr, P. T.; Zhang, C.-Y.; Wu, Z.; Boss, O.; Michael, L. F.; Puigserver, P.; Isotani, E.; Olson, E. N.; Lowell, B. B.; Bassel-Duby, R.; Spiegelman, B. M.: Transcriptional co-activator PGC-1-alpha drives the formation of slow-twitch muscle fibres. *Nature* 418: 797–801, 2002.
- [72477] 10807.Monsalve, M.; Wu, Z.; Adelmant, G.; Puigserver, P.; Fan, M.; Spiegelman, B. M.: Direct coupling of transcription and mRNA processing through the thermogenic coactivator PGC-1. *Molec. Cell* 6: 307–316, 2000.
- [72478] 10808.Puigserver, P.; Adelmant, G.; Wu, Z.; Fan, M.; Xu, J.;



O'Malley,B.; Spiegelman, B. M.: Activation of PPAR–gamma coactivator–1 throughtranscription factor docking. Science 286: 1368–1371, 1999.

- [72479] 10809.Puigserver, P.; Rhee, J.; Lin, J.; Wu, Z.; Yoon, J. C.; Zhang,C.–Y.; Krauss, S.; Mootha, V. K.; Lowell, B. B.; Spiegelman, B. M.: Cytokine stimulation of energy expenditure through p38 MAP kinaseactivation of PPAR–gamma coactivator–1. Molec. Cell 8: 971–982,2001.
- [72480] 10810.Puigserver, P.; Wu, Z.; Park, C. W.; Graves, R.; Wright, M.; Spiegelman,B. M.: A cold–inducible coactivator of nuclear receptors linked toadaptive thermogenesis. Cell 92: 829–839, 1998.
- [72481] 10811.Waite, L. L.; Person, E. C.; Zhou, Y.; Lim, K.–H.; Scanlan, T.S.; Taylor, R. N.: Placental peroxisome proliferator–activated receptor–gammaais up–regulated by pregnancy serum. J. Clin. Endocr. Metab. 85:3808–3814, 2000.
- [72482] 10812.Wu, Z.; Puigserver, P.; Andersson, U.; Zhang, C.; Adelmant, G.;Mootha, V.; Troy, A.; Cinti, S.; Lowell, B.; Scarpulla, R. C.; Spiegelman,B. M.: Mechanisms controlling mitochondrial biogenesis and respirationthrough the thermogenic coactivator PGC–1. Cell 98: 115–124, 1999.
- [72483] 10813.Yoon, J. C.; Puigserver, P.; Chen, G.; Donovan, J.; Wu, Z.; Rhee,J.; Adelmant, G.; Stafford, J.; Kahn, C. R.;

Granner, D. K.; Newgard, C. B.; Spiegelman, B. M.: Control of hepatic gluconeogenesis through the transcriptional coactivator PGC-1. *Nature* 413: 131–138, 2001.

[72484] 10814. Ellis, J. H.; Ashman, C.; Burden, M. N.; Kilpatrick, K. E.; Morse, M. A.; Hamblin, P. A.: GRID: a novel Grb-2-related adapter protein that interacts with the activated T cell costimulatory receptor CD28. *J. Immun.* 164: 5805–5814, 2000.

[72485] 10815. Qiu, M.; Hua, S.; Agrawal, M.; Li, G.; Cai, J.; Chan, E.; Zhou, H.; Luo, Y.; Liu, M.: Molecular cloning and expression of human Grap-2, a novel leukocyte-specific SH2- and SH3-containing adaptor-like protein that binds to Gab-1. *Biochem. Biophys. Res. Commun.* 253: 443–447, 1998.

[72486] 10816. Yoder, J.; Pham, C.; Iizuka, Y.-M.; Kanagawa, O.; Liu, S. K.; McGlade, J.; Cheng, A. M.: Requirement for the SLP-76 adaptor GADS in T cell development. *Science* 291: 1987–1991, 2001.

[72487] 10817. Granger, S. W.; Butrovich, K. D.; Houshmand, P.; Edwards, W. R.; Ware, C. F.: Genomic characterization of LIGHT reveals linkage to an immune response locus on chromosome 19p13.3 and distinct isoforms generated by alternate splicing or proteolysis. *J. Immun.* 167:

5122–5128,2001.

- [72488] 10818.Mauri, D. N.; Ebner, R.; Montgomery, R. I.; Kochel, K. D.; Cheung,T. C.; Yu, G. L.; Ruben, S.; Murphy, M.; Eisenberg, R. J.; Cohen,G. H.; Spear, P. G.; Ware, C. F.: LIGHT, a new member of the TNFsuperfamily, and lymphotoxin alpha are ligands for herpesvirus entrymediator. *Immunity* 8: 21–30, 1998.
- [72489] 10819.Harrop, J. A.; McDonnell, P. C.; Brigham–Burke, M.; Lyn, S. D.;Minton, J.; Tan, K. B.; Dede, K.; Spampanato, J.; Silverman, C.;Hensley, P.; DiPrinzio, R.; Emery, J. G.; Deen, K.; Eichman, C.; Chabot–Fletcher,M.; Truneh, A.; Young, P. R.: Herpesvirus entry mediator ligand (HVEM–L),a novel ligand for HVEM/TR2, stimulates proliferation of T cells andinhibits HT29 cell growth. . *J. Biol. Chem.* 273: 27548–27556, 1998.
- [72490] 10820.Hadjantonakis, A.–K.; Formstone, C. J.; Little, P. F. R.: mCelsr1is an evolutionarily conserved seven–pass transmembrane receptor andis expressed during mouse embryonic development. *Mech. Dev.* 78:91–95, 1998.
- [72491] 10821.Hadjantonakis, A.–K.; Sheward, W. J.; Harmar, A. J.; de Galan,L.; Hoovers, J. M. N.; Little, P. F. R.: Celsr1, a neural–specificgene encoding an unusual seven–pass transmembrane receptor, maps tomouse chromosome 15

and human chromosome 22qter. *Genomics* 45:  
97–104, 1997.

- [72492] 10822. Hewett–Emmett, D.; Tashian, R. E.: Functional diversity, conservation, and convergence in the evolution of the alpha–, beta–, and gamma–carbonic anhydrase gene families. *Molec. Phylogenet. Evol.* 5: 50–77, 1996.
- [72493] 10823. Hewett–Emmett, D.; Wiebauer, K.: Personal Communication. Houston, Texas 9/1999.
- [72494] 10824. Kleiderlein, J. J.; Nisson, P. E.; Jessee, J.; Li, W.–B.; Becker, K. G.; Derby, M. L.; Ross, C. A.; Margolis, R. L.: CCG repeats in cDNAs from human brain. *Hum. Genet.* 103: 666–673, 1998.
- [72495] 10825. Lovejoy, D. A.; Hewett–Emmett, D.; Porter, C. A.; Cepoi, D.; Sheffield, A.; Vale, W. W.; Tashian, R. E.: Evolutionarily conserved, 'acatalytic' carbonic anhydrase–related protein XI contains a sequence motif present in the neuropeptide sauvagine: the human CA–RP XI gene (CA11) is embedded between the secretor gene cluster and the DBP gene at 19q13.3. *Genomics* 54: 484–493, 1998.
- [72496] 10826. Bellingham, J.; Gregory–Evans, K.; Gregory–Evans, C. Y.: Sequence and tissue expression of a human novel carbonic anhydrase–related protein, CARP–2, mapping to chromosome 19q13.3. *Biochem. Biophys. Res. Comm.* 253:

364–367, 1998.

- [72497] 10827.Fujikawa–Adachi, K.; Nishimori, I.; Taguchi, T.; Yuri, K.; Onishi,S.: cDNA sequence, mRNA expression, and chromosomal localizationof human carbonic anhydrase–related protein, CA–RP XI. *Biochim. Biophys.Acta* 1431: 518–524, 1999.
- [72498] 10828.Lindsay, M. E.; Plafker, K.; Smith, A. E.; Clurman, B. E.; Macara,I. G.: Npap60/Nup50 is a tri–stable switch that stimulates importin–alpha:beta–mediatednuclear protein import. *Cell* 110: 349–360, 2002.
- [72499] 10829.Trichet, V.; Shkolny, D.; Dunham, I.; Beare, D.; Mc–Dermid, H. E.: Mapping and complex expression pattern of the human NPAP60L nucleoporingene. *Cytogenet. Cell Genet.* 85: 221–226, 1999.
- [72500] 10830.de Veer, M. J.; Sim, H.; Whisstock, J. C.; Devenish, R. J.; Ralph,S. J.: IFI60/ISG60/IFIT4, a new member of the human IFI54/IFIT2 familyof interferon–stimulated genes. *Genomics* 54: 267–277, 1998.
- [72501] 10831.Yu, M.; Tong, J.–H.; Mao, M.; Kan, L.–X.; Liu, M.–M.; Sun, Y.–W.;Fu, G.; Jing, Y.–K.; Yu, L.; Lepaslier, D.; Lanotte, M.; Wang, Z.–Y.;Chen, Z.; Waxman, S.; Wang, Y.–X.; Tan, J.–Z.; Chen, S.–J.: Cloningof a gene (RIG–G) associated with retinoic acid–induced differentiationof acute promye–

locytic leukemia cells and representing a new member of a family of interferon-stimulated genes. Proc. Nat. Acad. Sci. 94:7406–7411, 1997.

[72502] 10832. Zhu, H.; Cong, J.-P.; Shen, T.: Use of differential display analysis to assess the effect of human cytomegalovirus infection on the accumulation of cellular RNAs: induction of interferon-responsive RNAs. Proc. Nat. Acad. Sci. 94: 13985–13990, 1997.

[72503] 10833. Hirai, H.; Tanaka, K.; Takano, S.; Ichimasa, M.; Nakamura, M.; Nagata, K.: Cutting edge: agonistic effect of indomethacin on a prostaglandin D<sub>2</sub> receptor, CRTH2. J. Immun. 168: 981–985, 2002.

[72504] 10834. Nagata, K.; Tanaka, K.; Ogawa, K.; Kemmotsu, K.; Imai, Yoshie, O.; Abe, H.; Tada, K.; Nakamura, M.; Sugamura, K.; Takano, S.: Selective expression of a novel surface molecule by human Th2 cells in vivo. J. Immun. 162: 1278–1286, 1999.

[72505] 10835. Grundemann, D.; Schechinger, B.; Rappold, G. A.; Schomig, E.: Molecular identification of the corticosterone-sensitive extraneuronal catecholamine transporter. Nature Neurosci. 1: 349–351, 1998.

[72506] 10836. Boettger, T.; Hubner, C. A.; Maler, H.; Rust, M. B.; Beck, F. X.; Jentsch, T. J.: Deafness and renal tubular acido-

sis in mice lacking the K-Cl co-transporter Kcc4. *Nature* 416: 874–878, 2002.

[72507] 10837. Kotake, K.; Ozaki, N.; Mizuta, M.; Sekiya, S.; Inagaki, N.; Seino, S.: Noc2, a putative zinc finger protein involved in exocytosis in endocrine cells. *J. Biol. Chem.* 272: 29407–29410, 1997.

[72508] 10838. Smith, J. S.; Tachibana, I.; Allen, C.; Chiappa, S. A.; Lee, H. K.; McIver, B.; Jenkins, R. B.; Raffel, C.: Cloning of a human ortholog (RPH3AL) of (RNO) Rph3al from a candidate 17p13.3 medulloblastoma tumor suppressor locus. *Genomics* 59: 97–101, 1999.

[72509] 10839. Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Kawarabayashi, Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.: Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001–KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1. *DNA Res.* 1: 27–35, 1994.

[72510] 10840. Kimura, M.; Okano, Y.: Identification and assignment of the human NIMA-related protein kinase 7 gene (NEK7) to human chromosome 1q31.3. *Cytogenet. Cell Genet.* 94: 33–38, 2001.

[72511] 10841. Li, M. Z.; Yu, L.; Liu, Q.; Chu, J. Y.; Zhao, S. Y.: As-

signment of NEK6, a NIMA-related gene, to human chromosome 9q33.3–q34.11 by radiation hybrid mapping. Cytogenet. Cell Genet. 87: 271–272, 1999.

[72512] 10842. Favier, D.; Gonda, T. J.: Detection of proteins that bind to the leucine zipper motif of c-Myb. Oncogene 9: 305–311, 1994.

[72513] 10843. Keough, R.; Woollatt, E.; Crawford, J.; Sutherland, G. R.; Plummer, S.; Casey, G.; Gonda, T. J.: Molecular cloning and chromosomal mapping of the human homologue of MYB binding protein (P160) 1A (MYBBP1A) to 17p13.3. Genomics 62: 483–489, 1999.

[72514] 10844. Tavner, F. J.; Simpson, R.; Tashiro, S.; Favier, D.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Macmillan, E. M.; Lutwyche, J.; Keough, R. A.; Ishii, S.; Gonda, T. J.: Molecular cloning reveals that the p160 Myb-binding protein is a novel, predominantly nucleolar protein which may play a role in transactivation by Myb. Molec. Cell Biol. 18: 989–1002, 1998.

[72515] 10845. Gilbert, D. J.; Engel, H.; Wang, X.; Grzeschik, K.-H.; Copeland, N. G.; Jenkins, N. A.; Kilimann, M. W.: The neurobeachin gene (Nbea) identifies a new region of homology between mouse central chromosome 3 and human chromosome 13q13. Mamm. Genome 10: 1030–1031, 1999.



- [72516] 10846.Nagle, D. L.; Karim, M. A.; Woolf, E. A.; Holmgren, L.; Bork, P.; Misumi, D. J.; McGrail, S. H.; Dussault, B. J., Jr.; Perou, C. M.; Boissy, R. E.; Duyk, G. M.; Spritz, R. A.; Moore, K. J.: Identification and mutation analysis of the complete gene for Chediak–Higashi syndrome. *Nature–Genet.* 14: 307–311, 1996.
- [72517] 10847.Wang, X.; Herberg, F. W.; Laue, M. M.; Wullner, C.; Hu, B.; Petrasch–Parwez, E.; Kilimann, M, W.: Neu–robeachin: a protein kinase A–anchoring, beige/Chediak–Higashi protein homolog implicated in neuronal membrane traffic. *J. Neurosci.* 20: 8551–8565, 2000.
- [72518] 10848.Bunn, R. C.; Jensen, M. A.; Reed, B. C.: Protein interactions with the glucose transporter binding protein GLUT1CBP that provide a link between GLUT1 and the cytoskeleton. *Molec. Biol. Cell* 10:819–832, 1999.
- [72519] 10849.De Vries, L.; Lou, X.; Zhao, G.; Zheng, B.; Farquhar, M. G.: GIPC, a PDZ domain containing protein, interacts specifically with the Cterminus of RGS–GAIP. *Proc. Nat. Acad. Sci.* 95: 12340–12345, 1998.
- [72520] 10850.Von Kap–Herr, C.; Kandala, G.; Mann, S. S.; Hart, T. C.; Pettenati, M. J.; Setaluri, V.: Assignment of PDZ domain–containing protein GIPC gene (C19orf3) to human

chromosome band 19p13.1 by in situ hybridization and radiation hybrid mapping. *Cytogenet. Cell Genet.* 89: 234–235, 2000.

- [72521] 10851. Alessi, D. R.; James, S. R.; Downes, C. P.; Holmes, A. B.; Gaffney, P. R. J.; Reese, C. B.; Cohen, P.: Characterization of a 3-phosphoinositide-dependent protein kinase which phosphorylates and activates protein kinase B- $\alpha$ . *Curr. Biol.* 7: 261–269, 1997.
- [72522] 10852. Sandrini, F.; Farmakidis, C.; Kirschner, L. S.; Wu, S.-M.; Tullio-Pelet, A.; Lyonnet, S.; Metzger, D. L.; Boudony, C. J.; Tiosano, D.; Chan, W.-Y.; Stratakis, C. A.: Spectrum of mutations of the AAAS gene in Allgrove syndrome: lack of mutations in six kindreds with isolated resistance to corticotropin. *J. Clin. Endocr. Metab.* 86: 5433–5437, 2001.
- [72523] 10853. Ludwig, D.; Lorenz, J.; Dejana, E.; Bohlen, P.; Hicklin, D. J.; Witte, L.; Pytowski, B.: cDNA cloning, chromosomal mapping, and expression analysis of human VE-cadherin-2. *Mammalian Genome* 11: 1030–1033, 2000.
- [72524] 10854. Telo, P.; Breviario, F.; Huber, P.; Panzeri, C.; Dejana, E.: Identification of a novel cadherin (vascular endothelial cadherin-2) located at intercellular junctions in endothelial cells. *J. Biol. Chem.* 273: 17565–17572, 1998.

- [72525] 10855.Adams, J. C.; Seed, B.; Lawler, J.: Muskelin, a novel intracellularmediator of cell adhesive and cytoskeletal re-sponses to thrombospondin-1. EMBOJ. 17: 4964-4974, 1998.
- [72526] 10856.Adams, J. C.; Zhang, L.: cDNA cloning of human muskelin and localisationof the muskelin (MKLN1) gene to human chromosome 7q32 and mouse chromo-some6B1/B2 by physical mapping and FISH. Cytogenet. Cell Genet. 87:19-21, 1999.
- [72527] 10857.Moynihan, T. P.; Ardley, H. C.; Nuber, U.; Rose, S. A.; Jones,P. F.; Markham, A. F.; Scheffner, M.; Robinson, P. A.: The ubiquitin-conjugatingenzymes Ubch7 and Ubch8 interact with RING finger/IBR motif-containingdomains of HHARI and H7-AP1. J. Biol. Chem. 274: 30963-30968, 1999.
- [72528] 10858.Tan, N. G. S.; Ardley, H. C.; Rose, S. A.; Leek, J. P.; Markham,A. F.; Robinson, P. A.: Characterisation of the human and mouse orthologuesof the Drosophila ariadne gene. Cytogenet. Cell Genet. 90: 242-245,2000.
- [72529] 10859.Janssen, J. W. G.; Imoto, I.; Inoue, J.; Shimada, Y.; Ueda, M.;Imamura, M.; Bartram, C. R.; Inazawa, J.: MYEOV, a gene at 11q13,is coamplified with CCND1, but epige-netically inactivated in a subsetof esophageal squamous

cell carcinomas. J. Hum. Genet. 47: 460–464,2002.

- [72530] 10860.Janssen, J. W. G.; Vaandrager, J.–W.; Heuser, T.; Jauch, A.; Kluin,P. M.; Geelen, E.; Bergsagel, P. L.; Kuehl, W. M.; Drexler, H. G.;Otsuki, T.; Bartram, C. R.; Schuuring, E.: Concurrent activationof a novel putative transforming gene, myeov, and cyclin D1 in a subsetof multiple myeloma cell lines with t(11;14)(q13;q32). Blood 95:2691–2698, 2000.
- [72531] 10861.Di Cunto, F.; Calautti, E.; Hsiao, J.; Ong, L.; Topley, G.; Turco,E.; Dotto, G. P.: Citron Rho–interacting kinase, a novel tissue–specificser/thr kinase encompassing the Rho–Rac–binding protein citron. J.Biol. Chem. 273: 29706–29711, 1998.
- [72532] 10862.Di Cunto, F.; Imarisio, S.; Hirsch, E.; Broccoli, V.; Bulfone,A.; Migheli, A.; Atzori, C.; Turco, E.; Triolo, R.; Dotto, G. P.;Silengo, L.; Altruda, F.: Defective neurogenesis in citron kinaseknockout mice by altered cytokinesis and massive apoptosis. Neuron 28:115–127, 2000.
- [72533] 10863.Madaule, P.; Furuyashiki, T.; Reid, T.; Ishizaki, T.; Watanabe,G.; Morii, N.; Narumiya, S.: A novel partner for the GTP–bound formsof rho and rac. FEBS Lett. 377: 243–248, 1995.
- [72534] 10864.Ishizaki, J.; Suzuki, N.; Higashino, K.; Yokota, Y.;

Ono, T.; Kawamoto, K.; Fujii, N.; Arita, H.; Hanasaki, K.:  
Cloning and characterization of novel mouse and human  
secretory phospholipase A(2)s. *J. Biol. Chem.* 274:  
24973–24979, 1999.

[72535] 10865. Baudat, F.; Manova, K.; Yuen, J. P.; Jasin, M.;  
Keeney, S.: Chromosome synapsis defects and sexually di-  
morphic meiotic progression in mice lacking Spo11. *Molec.  
Cell* 6: 989–998, 2000.

[72536] 10866. Romanienko, P. J.; Camerini-Otero, R. D.: The  
mouse Spo11 gene is required for meiotic chromosome  
synapsis. *Molec. Cell* 6: 975–987, 2000.

[72537] 10867. Romanienko, P. J.; Camerini-Otero, R. D.: Cloning,  
characterization, and localization of mouse and human  
SPO11. *Genomics* 61: 156–169, 1999.

[72538] 10868. Shannon, M.; Richardson, L.; Christian, A.; Handel,  
M. A.; Thelen, M. P.: Differential gene expression of mam-  
malian SPO11/TOP6A homologs during meiosis. *FEBS Lett.*  
462: 329–334, 1999.

[72539] 10869. Lopez-Coviella, I.; Berse, B.; Krauss, R.; Thies, R.  
S.; Blusztajn, J. K.: Induction and maintenance of the neu-  
ronal cholinergic phenotype in the central nervous system  
by BMP-9. *Science* 289: 313–316, 2000.

[72540] 10870. Burn, T. C.; Connors, T. D.; Van Raay, T. J.; Dack-

owski, W. R.; Millholland, J. M.; Klinger, K. W.; Landes, G. M.: Generation of a transcriptional map for a 700-kb region surrounding the polycystic kidney disease type 1 (PKD1) and tuberous sclerosis type 2 (TSC2) disease genes on human chromosome 16p13.3. *Genome Res.* 6: 525–537, 1996.

[72541] 10871. Pullen, N.; Dennis, P. B.; Andjelkovic, M.; Dufner, A.; Kozma, S. C.; Hemmings, B. A.; Thomas, G.: Phosphorylation and activation of p70(s6k) by PDK1. *Science* 279: 707–710, 1998.

[72542] 10872. Stephens, L.; Anderson, K.; Stokoe, D.; Erdjument-Bromage, H.; Painter, G. F.; Holmes, A. B.; Gaffney, P. R. J.; Reese, C. B.; McCormick, F.; Tempst, P.; Coadwell, J.; Hawkins, P. T.: Protein kinase B kinases that mediate phosphatidylinositol 3,4,5-triphosphate-dependent activation of protein kinase B. *Science* 279: 710–714, 1998.

[72543] 10873. Uebele, V. N.; Lagrutta, A.; Wade, T.; Figueroa, D. J.; Liu, Y.; McKenna, E.; Austin, C. P.; Bennett, P. B.; Swanson, R.: Cloning and functional expression of two families of beta-subunits of the large conductance calcium-activated K(+) channel. *J. Biol. Chem.* 275: 23211–23218, 2000.

[72544] 10874. Kuroku, Y.; Soyama, A.; Fujita, E.; Urase, K.;

Tsukahara, T.; Momoi, T.: RA70 is a src kinase-associated protein expressed ubiquitously. *Biochem. Biophys. Res. Comm.* 252: 738–742, 1998.

- [72545] 10875. Liu, J.; Kang, H.; Raab, M.; da Silva, A. J.; Kraeft, S.-K.; Rudd, C. E.: FYB (FYN binding protein) serves as a binding partner for lymphoid protein and FYN kinase substrate SKAP55 and a SKAP55-related protein in T cells. *Proc. Nat. Acad. Sci.* 95: 8779–8784, 1998.
- [72546] 10876. Marie-Cardine, A.; Verhagen, A. M.; Eckerskorn, C.; Schraven, B.: SKAP-HOM, a novel adaptor protein homologous to the FYN-associated protein SKAP55. *FEBS Lett.* 435: 55–60, 1998.
- [72547] 10877. Chai, J.; Du, C.; Wu, J.-W.; Kyin, S.; Wang, X.; Shi, Y.: Structural and biochemical basis of apoptotic activation by Smac/DIABLO. *Nature* 406: 855–862, 2000.
- [72548] 10878. Du, C.; Fang, M.; Li, Y.; Li, L.; Wang, X.: Smac, a mitochondrial protein that promotes cytochrome c-dependent caspase activation by eliminating IAP inhibition. *Cell* 102: 33–42, 2000.
- [72549] 10879. Okada, H.; Suh, W.-K.; Jin, J.; Woo, M.; Du, C.; Elia, A.; Duncan, G. S.; Wakeham, A.; Itie, A.; Lowe, S. W.; Wang, X.; Mak, T. W.: Generation and characterization of Smac/DIABLO-deficient mice. *Molec. Cell. Biol.* 22: 3509–3517,

2002.

[72550] 10880.Scott, A. F.: Personal Communication. Baltimore, Md. 8/18/2000.

[72551] 10881.Verhagen, A. M.; Ekert, P. G.; Pakusch, M.; Silke, J.; Connolly,L. M.; Reid, G. E.; Moritz, R. L.; Simpson, R. J.; Vaux, D. L.: Identificationof DIABLO, a mammalian protein that promotes apoptosis by bindingto and antagonizing IAP proteins. Cell 102: 43–53, 2000.

[72552] 10882.Behrens, R.; Nolting, A.; Reimann, F.; Schwarz, M.; Waldschutz,R.; Pongs, O.: hKCNMB3 and hKCNMB4, cloning and characterizationof two members of the large–conductance calcium–activated potassiumchannel beta subunit family. FEBS Lett. 474: 99–106, 2000.

[72553] 10883.Brenner, R.; Jegla, T. J.; Wickenden, A.; Liu, Y.; Aldrich, R.W.: Cloning and functional characterization of novel large conductancecalcium–activated potassium channel beta subunits, hKCNMB3 and hKCNMB4. J.Biol. Chem. 275: 6453–6461, 2000.

[72554] 10884.Riazi, M. A.; Brinkman–Mills, P.; Johnson, A.; Naylor, S. L.; Minoshima,S.; Shimizu, N.; Baldini, A.; McDer–mid, H. E.: Identification ofa putative regulatory subunit of a calcium–activated potassium channelin the dup(3q) syndrome region and a related sequence on 22q11.2. Ge–



nomics 62:90–94, 1999.

- [72555] 10885.Hemmi, H.; Takeuchi, O.; Kawai, T.; Kaisho, T.; Sato, S.; Sanjo,H.; Matsumoto, M.; Hoshino, K.; Wagner, H.; Takeda, K.; Akira, S.: A toll–like receptor recognizes bacterial DNA. *Nature* 408: 740–745,2000.
- [72556] 10886.Leadbetter, E. A.; Rifkin, I. R.; Hohlbaum, A. M.; Beaudette, B.C.; Shlomchik, M. J.; Marshak–Rothstein, A.: Chromatin–IgG complexesactivate B cells by dual engagement of IgM and Toll–like receptors. *Nature* 416:603–607, 2002.
- [72557] 10887.Takeshita, F.; Leifer, C. A.; Gursel, I.; Ishii, K. J.; Takeshita,S.; Gursel, M.; Klinman, D. M.: Cutting edge: role of Toll–like receptor9 in CpG DNA–induced activation of human cells. *J. Immun.* 167: 3555–3558,2001.
- [72558] 10888.Verthelyi, D.; Ishii, K. J.; Gursel, M.; Takeshita, F.; Klinman,D. K.: Human peripheral blood cells differentially recognize andrespond to two distinct CpG motifs. *J. Immun.* 166: 2372–2377, 2001.
- [72559] 10889.Bagrodia, S.; Taylor, S. J.; Jordon, K. A.; Van Aelst, L.; Cerione,R. A.: A novel regulator of p21–activated kinases. *J. Biol. Chem.* 273:23633–23636, 1998.
- [72560] 10890.Oh, W. K.; Yoo, J. C.; Jo, D.; Song, Y. H.; Kim, M. G.; Park, D.: Cloning of a SH3 domain–containing proline–rich

protein, p85SPR, and its localization in focal adhesion.

Biochem. Biophys. Res. Commun. 235:794–798, 1997.

- [72561] 10891. Ehrmann, D. A.; Schwarz, P. E. H.; Hara, M.; Tang, X.; Horikawa, Y.; Imperial, J.; Bell, G. I.; Cox, N. J.: Relationship of calpain-10 genotype to phenotypic features of polycystic ovary syndrome. *J. Clin. Endocr. Metab.* 87: 1669–1673, 2002.
- [72562] 10892. Elbein, S. C.; Chu, W.; Ren, Q.; Hemphill, C.; Schay, J.; Cox, N. J.; Hanis, C. L.; Hasstedt, S. J.: Role of calpain-10 gene variants in familial type 2 diabetes in Caucasians. *J. Clin. Endocr. Metab.* 87:650–654, 2002.
- [72563] 10893. Fullerton, S. M.; Bartoszewicz, A.; Ybazeta, G.; Horikawa, Y.; Bell, G. I.; Kidd, K. K.; Cox, N. J.; Hudson, R. R.; Di Rienzo, A.: Geographic and haplotype structure of candidate type 2 diabetes-susceptibility variants at the calpain-10 locus. *Am. J. Hum. Genet.* 70: 1096–1106, 2002.
- [72564] 10894. Tsai, H.-J.; Sun, G.; Weeks, D. E.; Kaushal, R.; Wolujewicz, M.; McGarvey, S. T.; Tufa, J.; Viali, S.; Deka, R.: Type 2 diabetes and three calpain-10 gene polymorphisms in Samoans: no evidence of association. *Am. J. Hum. Genet.* 69: 1236–1244, 2001.
- [72565] 10895. Weiss, K. M.; Terwilliger, J. D.: How many diseases

does it take to map a gene with SNPs? *Nature Genet.* 26: 151–157, 2000.

[72566] 10896. Hanis, C. L.; Boerwinkle, E.; Chakraborty, R.; Ellsworth, D. L.; Concannon, P.; Stirling, B.; Morrison, V. A.; Wapelhorst, B.; Spielman, R. S.; Gogolin-Ewens, K. J.; Shephard, J. M.; Williams, S. R.; and 21 others: A genome-wide search for human non-insulin-dependent (type 2) diabetes genes reveals a major susceptibility locus on chromosome 2. *Nature Genet.* 13: 161–166, 1996.

[72567] 10897. Weber, A.; Wienker, T. F.; Jung, M.; Easton, D.; Dean, H. J.; Heinrichs, C.; Reis, A.; Clark, A. J. L.: Linkage of the gene for the triple A syndrome to chromosome 12q13 near the type II keratin gene cluster. *Hum. Molec. Genet.* 5: 2061–2066, 1996.

[72568] 10898. Bousquet, O.; Basseville, M.; Vila-Porcile, E.; Billette de Villemeur, T.; Hauw, J.-J.; Landrieu, P.; Portier, M.-M.: Aggregation of a subpopulation of vimentin filaments in cultured human skin fibroblasts derived from patients with giant axonal neuropathy. *Cell. Motil. Cytoskeleton* 33: 115–129, 1996.

[72569] 10899. Kehlenbaumer, G.; Young, P.; Oberwittler, C.; Hunermund, G.; Schirmacher, A.; Domschke, K.; Ringelstein, B.; Stogbauer, F.: Giant axonal neuropathy (GAN):

case report and two novel mutations in the gigaxonin gene. *Neurology* 58:1273–1276, 2002. Note: Erratum: *Neurology* 58: 1444, 2002.

[72570] 10900.Pena, S. D.: Giant axonal neuropathy: an inborn error of organization of intermediate filaments. *Muscle Nerve* 5: 166–172, 1982.

[72571] 10901.Prineas, J. W.; Ouvrier, R. A.; Wright, R. G.; Walsh, J. C.; McLeod, J. G.: Giant axonal neuropathy: a generalized disorder of cytoplasmic microfilament formation. *J. Neuropath. Exp. Neurol.* 35: 458–470, 1976.

[72572] 10902.Bowe, A. E.; Finnegan, R.; Jan de Beur, S. M.; Cho, J.; Levine, M. A.; Kumar, R.; Schiavi, S. C.: FGF-23 inhibits renal tubular phosphate transport and is a PHEX substrate. *Biochem. Biophys. Res. Commun.* 284:977–981, 2001.

[72573] 10903.Shimada, T.; Mizutani, S.; Muto, T.; Yoneya, T.; Hino, R.; Takeda, S.; Takeuchi, Y.; Fujita, T.; Fukumoto, S.; Yamashita, T.: Cloning and characterization of FGF23 as a causative factor of tumor-induced osteomalacia. *Proc. Nat. Acad. Sci.* 98: 6500–6505, 2001.

[72574] 10904.Strewler, G. J.: FGF23, hypophosphatemia, and rickets: has phosphorylation been found? (Commentary) *Proc. Nat. Acad. Sci.* 98: 5945–5946, 2001.

[72575] 10905.White, K. E.; Jonsson, K. B.; Carn, G.; Hampson, G.;

Spector, T.D.; Mannstadt, M.; Lorenz-Depiereux, B.; Miyauchi, A.; Yang, I. M.; Ljunggren, O.; Meitinger, T.; Strom, T. M.; Juppner, H.; Econs, M.J.: The autosomal dominant hypophosphatemic rickets (ADHR) gene is a secreted polypeptide overexpressed by tumors that cause phosphate wasting. *J. Clin. Endocr. Metab.* 86: 497–500, 2001.

[72576] 10906. Yamashita, T.; Yoshioka, M.; Itoh, N.: Identification of a novel fibroblast growth factor, FGF-23, preferentially expressed in the ventrolateral thalamic nucleus of the brain. *Biochem. Biophys. Res. Commun.* 277: 494–498, 2000.

[72577] 10907. Barker, R. L.; Gleich, G. J.; Pease, L. R.: Acidic precursor revealed in human eosinophil granule major basic protein cDNA. *J. Exp. Med.* 168: 1493–1498, 1988.

[72578] 10908. Li, M.-S.; Sun, L.; Satoh, T.; Fisher, L. M.; Spry, C. J. F.: Human eosinophil major basic protein, a mediator of allergic inflammation, is expressed by alternative splicing from two promoters. *Biochem. J.* 305: 921–927, 1995.

[72579] 10909. McGrogan, M.; Simonsen, C.; Scott, R.; Griffith, J.; Ellis, N.; Kennedy, J.; Campanelli, D.; Nathan, C.; Gabay, J.: Isolation of a complementary DNA clone encoding a precursor to human eosinophil major basic protein. *J. Exp.*

Med. 168: 2295–2308, 1988.

[72580] 10910. Plager, D. A.; Weiler, D. A.; Loegering, D. A.; Johnson, W. B.; Haley, L.; Eddy, R. L.; Shows, T. B.; Gleich, G. J.: Comparative structure, proximal promoter elements, and chromosome location of the human eosinophil major basic protein genes. *Genomics* 71: 271–281, 2001.

[72581] 10911. Wasmoen, T. L.; Bell, M. P.; Loegering, D. A.; Gleich, G. J.; Prendergast, F. G.; McKean, D. J.: Biochemical and amino acid sequence analysis of human eosinophil granule major basic protein. *J. Biol. Chem.* 263: 12559–12563, 1988.

[72582] 10912. Weller, P. F.; Ackerman, S. J.; Smith, J. A.: Eosinophil granule proteins: major basic protein is distinct from the smaller subunit of eosinophil peroxidase. *J. Leukoc. Biol.* 43: 1–4, 1988.

[72583] 10913. Yoshimatsu, K.; Ohya, Y.; Shikata, Y.; Seto, T.; Hasegawa, Y.; Tanaka, I.; Kawamura, T.; Kitoh, K.; Toyoshima, S.; Osawa, T.: Purification and cDNA cloning of a novel factor produced by a human T-cell hybridoma: sequence homology with animal lectins. *Molec. Immun.* 29: 537–546, 1992.

[72584] 10914. Frey, N.; Richardson, J. A.; Olson, E. N.: Calsarcins, a novel family of sarcomeric calcineurin-binding proteins.

Proc. Nat. Acad.Sci. 97: 14632–14637, 2000.

- [72585] 10915.Faulkner, G.; Pallavicini, A.; Comelli, A.; Salamon, M.; Bortoletto,G.; Ievolella, C.; Trevisan, S.; Kojic, S.; Dalla Vecchia, F.; Laveder,P.; Valle, G.; Lanfranchi, G.: FATZ, a filamin-, actinin-, and telethonin-binding protein of the Z-disc of skeletal muscle. J. Biol. Chem. 275: 41234–41242,2000.
- [72586] 10916.Takada, F.; Vander Woude, D. L.; Tong, H.-Q.; Thompson, T. G.;Watkins, S. C.; Kunkel, L. M.; Beggs, A. H.: Myozenin: an alpha-actinin-and gamma-filamin-binding protein of skeletal muscle Z lines. Proc.Nat. Acad. Sci. 98: 1595–1600, 2001.
- [72587] 10917.Scott, A. F.: Personal Communication. Baltimore, Md. 3/13/2001.
- [72588] 10918.Drane, P.; Barel, M.; Balbo, M.; Frade, R.: Identification ofRB18A, a 205 kDa new p53 regulatory protein which shares antigenicand functional properties with p53. Oncogene 15: 3013–3024, 1997.
- [72589] 10919.Mansharamani, M.; Hewetson, A.; Chilton, B. S.: Cloning and characterizationof an atypical type IV P-type ATPase that binds to the RING motifof RUSH transcription factors. J. Biol. Chem. 276: 3641–3649, 2001.
- [72590] 10920.Nakayama, Y.; Weissman, S. M.; Bothwell, A. L. M.:

BXMAS1 identifies a cluster of homologous genes differentially expressed in B cells. *Biochem. Biophys. Res. Commun.* 285: 830–837, 2001.

[72591] 10921. Desai, R.; Peretz, A.; Idelson, H.; Lazarovici, P.; Attali, B.:  $\text{Ca}^{2+}$ -activated  $\text{K}^{+}$  channels in human leukemic Jurkat T cells: molecular cloning, biochemical and functional characterization. *J. Biol. Chem.* 275: 39954–39963, 2000.

[72592] 10922. Maguchi, M.; Nishida, W.; Kohara, K.; Kuwano, A.; Kondo, I.; Hiwada, K.: Molecular cloning and gene mapping of human basic and acidic calponins. *Biochem. Biophys. Res. Commun.* 217: 238–244, 1995.

[72593] 10923. Miano, J. M.; Krahe, R.; Garcia, E.; Elliott, J. M.; Olson, E. N.: Expression, genomic structure and high resolution mapping to 19p13.2 of the human smooth muscle cell calponin gene. *Gene* 197: 215–224, 1997.

[72594] 10924. Miano, J. M.; Thomas, S.; Distèche, C. M.: Expression and chromosomal mapping of the mouse smooth muscle calponin gene. *Mammalian Genome* 12: 187–191, 2001.

[72595] 10925. Strasser, P.; Gimona, M.; Moessler, H.; Herzog, M.; Small, J. V.: Mammalian calponin: identification and expression of genetic variants. *FEBS Lett.* 330: 13–18, 1993.



- [72596] 10926.Duetsch, G.; Illig, T.; Loesgen, S.; Rohde, K.; Kloop, N.; Herbon,N.; Cohlke, H.; Altmueller, J.; Wjst, M.: STAT6 as an asthma candidategene: polymorphism–screening, association and haplotype analysis ina Caucasian sib–pair study. Hum. Molec. Genet. 11: 613–621, 2002.
- [72597] 10927.Wu, H.–K.; Heng, H. H. Q.; Shi, X.–M.; Forsdyke, D. R.; Tsui, L.–C.;Mak, T. W.; Minden, M. D.; Siderovski, D. P.: Differential expressionof a basic helix–loop–helix phosphoprotein gene, G0S8, in acute leukemiaand localization to human chromosome 1q31. Leukemia 9: 1291–1298,1995.
- [72598] 10928.Bogerd, H. P.; Fridell, R. A.; Madore, S.; Cullen, B. R.: Identificationof a novel cellular cofactor for the Rev/Rex class of retroviral regulatoryproteins. Cell 82: 485–494, 1995.
- [72599] 10929.Fritz, C. C.; Zapp, M. L.; Green, M. R.: A human nucleoporin–likeprotein that specifically interacts with HIV Rev. Nature 376: 530–533,1995.
- [72600] 10930.Jones, T.; Sheer, D.; Bevec, D.; Kappel, B.; Hauber, J.; Steinkasserer,A.: The human HIV–1 Rev binding–protein hRIP/Rab (HRB) maps to chromosome2q36. Genomics 40: 198–199, 1997.
- [72601] 10931.Kang–Decker, N.; Mantchev, G. T.; Juneja, S. C.;

McNiven, M. A.;van Deursen, J. M. A.: Lack of acrosome formation in Hrb-deficientmice. Science 294: 1531–1533, 2001.

[72602] 10932.Salcini, A. E.; Confalonieri, S.; Doria, M.; Santolini, E.; Tassi,E.; Minenkova, O.; Cesareni, G.; Pelicci, P. G.; Di Fiore, P. P.:Binding specificity and in vivo targets of the EH domain, a novelprotein–protein interaction module. Genes Dev. 11: 2239–2249, 1997.

[72603] 10933.Fish, K. J.; Cegielska, A.; Getman, M. E.; Landes, G. M.; Virshup,D. M.: Isolation and characterization of human casein kinase I–epsilon(CKI), a novel member of the CKI gene family. J. Biol. Chem. 270:14875–14883, 1995.

[72604] 10934.Kloss, B.; Price, J. L.; Saez, L.; Blau, J.; Rothenfluh, A.; Wesley,C. S.; Young, M. W.: The Drosophila clock gene double–time encodesa protein closely–related to human casein kinase I–epsilon. Cell 94:97–107, 1998.

[72605] 10935.Lowrey, P. L.; Shimomura, K.; Antoch, M. P.; Yamazaki, S.; Zamenides,P. D.; Ralph, M. R.; Menaker, M.; Takahashi, J. S.: Positional synteniccloning and functional characterization of the mammalian circadianmutation tau. Science 288: 483–491, 2000.

[72606] 10936.Graves, P. R.; Haas, D. W.; Hagedorn, C. H.; De–Paoli–Roach, A.A.; Roach, P. J.: Molecular cloning, expres–

sion, and characterization of a 49-kilodalton casein kinase I isoform from rat testis. *J. Biol. Chem.* 268: 6394–6401, 1993.

[72607] 10937. Kusuda, J.; Hidari, N.; Hirai, M.; Hashimoto, K.: Sequence analysis of the cDNA for the human casein kinase 1-delta (CSNK1D) gene and its chromosomal localization. *Genomics* 32: 140–143, 1996.

[72608] 10938. Kools, P. F. J.; Roebroek, A. J. M.; van de Velde, H. J. K.; Marynen, P.; Bullerdiek, J.; Van de Ven, W. J. M.: Regional mapping of the human NSP gene to chromosome region 14q21–q22 by fluorescence in situ hybridization analysis. *Cytogenet. Cell Genet.* 66: 48–50, 1994.

[72609] 10939. Roebroek, A. J. M.; Ayoubi, T. A. Y.; van de Velde, H. J. K.; Schoenmakers, E. F. P. M.; Pauli, I. G. L.; Van de Ven, W. J. M.: Genomic organization of the human NSP gene, prototype of a novel gene family encoding reticulons. *Genomics* 32: 191–199, 1996.

[72610] 10940. Roebroek, A. J. M.; van de Velde, H. J. K.; Van Bokhoven, A.; Broers, J. L. V.; Ramaekers, F. C. S.; Van de Ven, W. J. M.: Cloning and expression of alternative transcripts of a novel neuroendocrine-specific gene and identification of its 135-kDa translational product. *J. Biol. Chem.* 268: 13439–13447, 1993.

- [72611] 10941. Senden, N. H. M.; van de Velde, H. J. K.; Broers, J. L. V.; Timmer, E. D. J.; Roebroek, A. J. M.; van de Ven, W. J. M.; Ramaekers, F. C. S.: Cluster-10 lung-cancer antibodies recognize NSPs, novel neuro-endocrine proteins associated with membranes of the endoplasmic reticulum. *Int. J. Cancer* (Suppl. 8): 84–88, 1994.
- [72612] 10942. van de Velde, H. J. K.; Roebroek, A. J. M.; van Leeuwen, F. W.; Van de Ven, W. J. M.: Molecular analysis of expression in rat brain of NSP-A, a novel neuroendocrine-specific protein of the endoplasmic reticulum. *Molec. Brain Res.* 23: 81–92, 1994.
- [72613] 10943. van de Velde, H. J. K.; Senden, N. H. M.; Roskams, T. A. D.; Broers, J. L. V.; Ramaekers, F. C. S.; Roebroek, A. J. M.; Van de Ven, W. J. M.: NSP-encoded reticulons are neuroendocrine markers of a novel category in human lung cancer diagnosis. *Cancer Res.* 54: 4769–4776, 1994.
- [72614] 10944. Baron, B. W.; Anastasi, J.; Thirman, M. J.; Furukawa, Y.; Fears, S.; Kim, D. C.; Simone, F.; Birkenbach, M.; Montag, A.; Sadhu, A.; Zeleznik-Le, N.; McKeithan, T. W.: The human programmed cell death-2 (PDCD2) gene is a target of BCL6 repression: implications for a role of BCL6 in the down-regulation of apoptosis. *Proc. Nat. Acad. Sci.* 99:2860–2865, 2002.

- [72615] 10945.Cohen, J. J.; Duke, R. C.: Glucocorticoid activation of a calcium-dependent endonuclease in thymocyte nuclei leads to cell death. *J. Immun.* 132:38–42, 1984.
- [72616] 10946.Kawakami, T.; Furukawa, Y.; Sudo, K.; Saito, H.; Takami, S.; Takahashi, E.; Nakamura, Y.: Isolation and mapping of a human gene (PDCD2) that is highly homologous to Rp8, a rat gene associated with programmed cell death. *Cytogenet. Cell Genet.* 71: 41–43, 1995.
- [72617] 10947.Owens, G. P.; Cohen, J. J.: Identification of genes involved in programmed cell death. *Cancer Metastasis Rev.* 11: 149–156, 1992.
- [72618] 10948.Sellins, K. S.; Cohen, J. J.: Gene induction by gamma-irradiation leads to DNA fragmentation in lymphocytes. *J. Immun.* 139: 3199–3206, 1987.
- [72619] 10949.Wyllie, A. H.: Glucocorticoid-induced thymocyte apoptosis is associated with endogenous endonuclease activation. *Nature* 284: 555–556, 1980.
- [72620] 10950.Bullrich, F.; Druck, T.; Kunapuli, P.; Gomez, J.; Gripp, K. W.; Schlegelberger, B.; Lasota, J.; Aronson, M.; Cannizzaro, L. A.; Huebner, K.; Benovic, J. L.: Chromosomal mapping of the genes GPRK5 and GPRK6 encoding G protein-coupled receptor kinases GRK5 and GRK6. *Cytogenet. Cell Genet.* 70: 250–254, 1995.

- [72621] 10951.Haribabu, B.; Snyderman, R.: Identification of additional members of human G-protein-coupled receptor kinase multigene family. *Proc.Nat. Acad. Sci.* 90: 9398–9402, 1993.
- [72622] 10952.Miki, T.; Suzuki, M.; Shibasaki, T.; Uemura, H.; Sato, T.; Yamaguchi,K.; Koseki, H.; Iwanaga, T.; Nakaya, H.; Seino, S.: Mouse model of Prinzmetal angina by disruption of the inward rectifier Kir6.1. *NatureMed.* 8: 466–472, 2002.
- [72623] 10953.Prinzmetal, M.; Kenamer, R.; Merliss, R.; Wada, T.; Bor, N.:Angina pectoris. 1. A variant form of angina pectoris: preliminary report. *Am. J. Med.* 27: 375–388, 1959.
- [72624] 10954.Pristipino, C.; Beltrame, J. F.; Finocchiaro, M. L.; Hattori, R.;Fujita, M.; Mongiardo, R.; Cianflone, D.; Sanna, T.; Sasayama, S.;Maseri, A.: Major racial differences in coronary constrictor response between Japanese and Caucasians with recent myocardial infarction. *Circulation* 101:1102–1108, 2000.
- [72625] 10955.Cherel, M.; Sorel, M.; Apiou, F.; Lebeau, B.; Dubois, S.; Jacques,Y.; Minvielle, S.: The human interleukin–11 receptor alpha gene (IL11RA):genomic organization and chromosome mapping. *Genomics* 32: 49–53,1996.
- [72626] 10956.Cherel, M.; Sorel, M.; Lebeau, B.; Dubois, S.;

Moreau, J.-F.; Bataille, R.; Minvielle, S.; Jacques, Y.: Molecular cloning to two isoforms of a receptor for the human hematopoietic cytokine interleukin-11. *Blood* 86:2534-2540, 1995.

[72627] 10957. Neuhaus, H.; Bettenhausen, B.; Bilinski, P.; Simon-Chazottes, D.; Guenet, J.-L.; Gossler, A.: Etl2, a novel putative type-I cytokine receptor expressed during mouse embryogenesis at high levels in skin and cells with skeletogenic potential. *Dev. Biol.* 166: 531-542, 1994.

[72628] 10958. Van Leuven, F.; Stas, L.; Hilliker, C.; Miyake, Y.; Bilinski, P.; Gossler, A.: Molecular cloning and characterization of the human interleukin-11 receptor alpha-chain gene, IL11RA, located on chromosome 9p13. *Genomics* 31: 65-70, 1996.

[72629] 10959. Magrangeas, F.; Pitiot, G.; Dubois, S.; Bragado-Nilsson, E.; Cherel, M.; Jobert, S.; Lebeau, B.; Boisteau, O.; Lethe, B.; Mallet, J.; Jacques, Y.; Minvielle, S.: Cotranscription and intergenic splicing of human galactose-1-phosphate uridylyltransferase and interleukin-11 receptor alpha-chain genes generate a fusion mRNA in normal cells: implication for the production of multidomain proteins during evolution. *J. Biol. Chem.* 273: 16005-16010, 1998.

- [72630] 10960. Avraham, K. B.; Hasson, T.; Sobe, T.; Balsara, B.; Testa, J. R.; Skvorak, A. B.; Morton, C. C.; Copeland, N. G.; Jenkins, N. A.: Characterization of unconventional MYO6, the human homologue of the gene responsible for deafness in Snell's waltzer mice. *Hum. Molec. Genet.* 6: 1225–1231, 1997.
- [72631] 10961. Avraham, K. B.; Hasson, T.; Steel, K. P.; Kingsley, D. M.; Russell, L. B.; Mooseker, M. S.; Copeland, N. G.; Jenkins, N. A.: The mouse Snell's waltzer deafness gene encodes an unconventional myosin required for structural integrity of inner ear hair cells. *Nature Genet.* 11: 369–375, 1995.
- [72632] 10962. Hasson, T.; Mooseker, M. S.: Porcine myosin-VI: characterization of a new mammalian unconventional myosin. *J. Cell. Biol.* 127: 425–440, 1994.
- [72633] 10963. Hasson, T.; Skowron, J. F.; Gilbert, D. J.; Avraham, K. B.; Perry, W. L.; Bement, W. M.; Anderson, B. L.; Sherr, E. H.; Chen, Z.-Y.; Greene, L. A.; Ward, D. C.; Corey, D. P.; Mooseker, M. S.; Copeland, N. G.; Jenkins, N. A.: Mapping of unconventional myosins in mouse and human. *Genomics* 36: 431–439, 1996.
- [72634] 10964. Rock, R. S.; Rice, S. E.; Wells, A. L.; Purcell, T. J.; Spudich, J. A.; Sweeney, H. L.: Myosin VI is a processive



motor with a largestep size. Proc. Nat. Acad. Sci. 98: 13655–13659, 2001.

[72635] 10965.Wells, A. L.; Lin, A. W.; Chen, L.–Q.; Safer, D.; Cain, S. M.;Hasson, T.; Carragher, B. O.; Milligan, R. A.; Sweeney, H. L.: MyosinVI is an actin–based motor that moves backwards. Nature 401: 505–508,1999.

[72636] 10966.Melchionda, S.; Ahituv, N.; Bisceglia, L.; Sobe, T.; Glaser, F.;Rabionet, R.; Arbones, M. L.; Notarangelo, A.; Di Iorio, E.; Carella,M.; Zelante, L.; Estivill, X.; Avraham, K. B.; Gasparini, P.: MYO6,the human homologue of the gene responsible for deafness in Snell'sWaltzer mice, is mutated in autosomal dominant nonsyndromic hearingloss. Am. J. Hum. Genet. 69: 635–640, 2001.

[72637] 10967.Chittum, H. S.; Himeno, S.; Hill. K. E.; Burk, R. F.: Multipleforms of selenoprotein P in rat plasma. Arch. Biochem. Biophys. 325:124–128, 1996.

[72638] 10968.Bonne, S.; van Hengel, J.; van Roy, F.: Chromosomal mapping ofhuman armadillo genes belonging to the p120(ctn)/plakophilin subfamily. Genomics 51:452–454, 1998.

[72639] 10969.Drab, M.; Verkade, P.; Elger, M.; Kasper, M.; Lohn, M.; Lauterbach,B.; Menne, J.; Lindschau, C.; Mende, F.; Luft, F. C.; Schedl, A.;Haller, H.; Kurzchalia, T. V.: Loss of

caveolae, vascular dysfunction, and pulmonary defects in caveolin-1 gene-disrupted mice. *Science* 293:2449–2452, 2001.

[72640] 10970. Engelman, J. A.; Zhang, X.; Galbiati, F.; Volonte, D.; Sotgia, F.; Pestell, R. G.; Minetti, C.; Scherer, P. E.; Okamoto, T.; Lisanti, M. P.: Molecular genetics of the caveolin gene family: implications for human cancers, diabetes, Alzheimer disease, and muscular dystrophy. *Am. J. Hum. Genet.* 63: 1578–1587, 1998.

[72641] 10971. Engelman, J. A.; Zhang, X. L.; Galbiati, F.; Lisanti, M. P.: Chromosomal localization, genomic organization, and developmental expression of the murine caveolin gene family (Cav-1, -2, and -3): Cav-1 and Cav-2 genes map to a known tumor suppressor locus (6-A2/7q31). *FEBS Lett.* 429:330–336, 1998.

[72642] 10972. Engelman, J. A.; Zhang, X. L.; Lisanti, M. P.: Genes encoding human caveolin-1 and -2 are co-localized to the D7S522 locus (7q31.1), a known fragile site (FRA7G) that is frequently deleted in human cancers. *FEBS Lett.* 436: 403–410, 1998.

[72643] 10973. Feron, O.; Dessy, C.; Moniotte, S.; Desager, J.-P.; Balligand, J.-L.: Hypercholesterolemia decreases nitric oxide production by promoting the interaction of caveolin

and endothelial nitric oxidesynthase. J. Clin. Invest. 103: 897–905, 1999.

[72644] 10974.Giordano, S.; Ponzetto, C.; Di Renzo, M. F.; Cooper, C. S.; Comoglio,P. M.: Tyrosine kinase receptor indistinguishable from the c-metprotein. Nature 339: 155–156, 1989.

[72645] 10975.Glenney, J. R., Jr.: The sequence of human caveolin reveals identitywith VIP21, a component of transport vesicles. FEBS Lett. 314: 45–48,1992.

[72646] 10976.Kurzchalia, T. V.; Dupree, P.; Parton, R. G.; Kellner, R.; Virta,H.; Lehnert, M.; Simons, K.: VIP21, a 21–kD membrane protein is anintegral component of trans–Golgi–network–derived transport vesicles. J.Cell Biol. 118 1003–1014, 1992.

[72647] 10977.Scherer, P. E.; Okamoto, T.; Chun, M.; Nishimoto, I.; Lodish,H. F.; Lisanti, M. P.: Identification, sequence, and expression ofcaveolin–2 defines a caveolin gene family. Proc. Nat. Acad. Sci. 93:131–135, 1996.

[72648] 10978.Scherer, P. E.; Tang, Z.; Chun, M.; Sargiacomo, M.; Lodish, H.F.; Lisanti, M. P.: Caveolin isoforms differ in their N–terminalprotein sequence and subcellular distribution: identification andepitope mapping of an isoform–specific monoclonal antibody probe. J.Biol. Chem. 270

16395–16401, 1995.

- [72649] 10979. Razani, B.; Engelman, J. A.; Wang, X. B.; Schubert, W.; Zhang, X. L.; Marks, C. B.; Macaluso, F.; Russell, R. G.; Li, M.; Pestell, R. G.; Di Vizio, D.; Hou, H., Jr.; Kneitz, B.; Lagaud, G.; Christ, G. J.; Edelman, W.; Lisanti, M. P.: Caveolin-1 null mice are viable but show evidence of hyperproliferative and vascular abnormalities. *J. Biol. Chem.* 276: 38121–38138, 2001.
- [72650] 10980. Bell, D. R.; Plant, N. J.; Rider, C. G.; Na, L.; Brown, S.; Ateitalla, I.; Acharya, S. K.; Davies, M. H.; Elias, E.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Elcombe, C. R.: Species-specific induction of cytochrome P-450 4A RNAs: PCR cloning of partial guinea-pig, human and mouse CYP4A cDNAs. *Biochem. J.* 294: 173–180, 1993.
- [72651] 10981. Imaoka, S.; Ogawa, H.; Kimura, S.; Gonzalez, F. J.: Complete cDNA sequence and cDNA-directed expression of CYP4A11, a fatty acid omega-hydroxylase expressed in human kidney. *DNA Cell Biol.* 12: 893–899, 1993.
- [72652] 10982. Kawashima, H.; Kusunose, E.; Kikuta, Y.; Kinoshita, H.; Tanaka, S.; Yamamoto, S.; Kishimoto, T.; Kusunose, M.: Purification and cDNA cloning of human liver CYP4A fatty acid omega-hydroxylase. *J. Biochem.* 116: 74–80, 1994.
- [72653] 10983. Palmer, C. N. A.; Richardson, T. H.; Griffin, K. J.;

Hsu, M.-H.; Muerhoff, A. S.; Clark, J. E.; Johnson, E. F.: Characterization of a cDNA encoding a human kidney, cytochrome P-450 4A fatty acid omega-hydroxylase and the cognate enzyme expressed in *Escherichia coli*. *Biochim. Biophys. Acta* 1172: 161-166, 1993.

[72654] 10984. Muller, J.; Ory, S.; Copeland, T.; Piwnicka-Worms, H.; Morrison, D. K.: C-TAK1 regulates Ras signaling by phosphorylating the MAPK scaffold, KSR1. *Molec. Cell* 8: 983-993, 2001.

[72655] 10985. Rosenthal, R. L.: Haemorrhage in PTA (factor XI) deficiency. (Abstract) *Proc. 10th Int. Cong. Soc. Hemat., Stockholm*, 1964.

[72656] 10986. Laporte, J.; Hu, L. J.; Kretz, C.; Mandel, J.-L.; Kioschis, P.; Coy, J. F.; Klauck, S. M.; Poustka, A.; Dahl, N.: A gene mutated in X-linked myotubular myopathy defines a new putative tyrosine phosphatase family conserved in yeast. *Nature Genet.* 13: 175-182, 1996.

[72657] 10987. Rosenthal, R. L.; Dreskin, O. H.; Rosenthal, N.: Plasma thromboplastin antecedent (PTA) deficiency: clinical, coagulation, therapeutic and hereditary aspects of a new hemophilia-like disease. *Blood* 10: 120-131, 1955.

[72658] 10988. Seligsohn, U.: Factor XI (PTA) deficiency. In: Goodman, R. E.; Motulsky, A. G.: *Genetic Diseases Among*

Ashkenazi Jews. New York:Raven Press (pub.) 1979. Pp. 141–148.

[72659] 10989.Vinazzer, H.: Partieller familiaerer Faktor–XI–Mangel. Blut 15:263–267, 1967.

[72660] 10990.Wistinghausen, B.; Reischer, A.; Oddoux, C.; Ostrer, H.; Nardi,M.; Karpatkin, M.: Severe factor XI deficiency in an Arab familyassociated with a novel mutation in exon 11. Brit. J. Haemat. 99:575–577, 1997.

[72661] 10991.Zivelin, A.; Bauduer, F.; Ducout, L.; Peretz, H.; Rosenberg, N.;Yatuv, R.; Seligsohn, U.: Factor XI deficiency in French Basquesis caused predominantly by an ancestral cys38–to–arg mutation in thefactor XI gene. Blood 99: 2448–2454, 2002.

[72662] 10992.Hsueh, Y.–P.; Wang, T.–F.; Yang, F.–C.; Sheng, M.: Nuclear transcriptionand transcription regulation by the membrane–associated guanylatekinase CASK/LIN–2. Nature 404: 298–302, 2000.

[72663] 10993.De Smet, C.; Lurquin, C.; van der Bruggen, P.; De Plaen, E.; Brasseur,F.; Boon, T.: Sequence and expression pattern of the human MAGE2gene. Immunogenetics 39: 121–129, 1994.

[72664] 10994.Chambost, H.; Van Baren, N.; Brasseur, F.; Godelaine, D.; Xerri,L.; Landi, S. J.; Theate, I.; Plumas, J.; Spag–

noli, G. C.; Michel, G.; Coulie, P. G.; Olive, D.: Expression of gene MAGE-A4 in Reed-Sternberg cells. *Blood* 95: 3530-3533, 2000.

[72665] 10995. Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XI. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.* 5: 277-286, 1998.

[72666] 10996. Gospe, S. M., Jr.; Lazaro, R. P.; Lava, N. S.; Grootscholten, P. M.; Scott, M. O.; Fischbeck, K. H.: Familial X-linked myalgia and cramps: a nonprogressive myopathy associated with a deletion in the dystrophin gene. *Neurology* 39: 1277-1280, 1989.

[72667] 10997. Kingston, H. M.; Sarfarazi, M.; Thomas, N. S. T.; Harper, P. S.: Localisation of the Becker muscular dystrophy gene on the short arm of the X chromosome by linkage to cloned DNA sequences. *Hum. Genet.* 67: 6-17, 1984.

[72668] 10998. Kingston, H. M.; Thomas, N. S. T.; Pearson, P. L.; Sarfarazi, M.; Harper, P. S.: Genetic linkage between Becker muscular dystrophy and a polymorphic DNA sequence on the short arm of the X chromosome. *J. Med. Genet.* 20:

255–258, 1983.

- [72669] 10999.Reardon, W.; Middleton–Price, H. R.; Sandkuijl, L.; Phelps, P.; Bellman, S.; Luxon, L.; Pembrey, M. E.; Malcolm, S.: A multipedigree linkage study of X–linked deafness: linkage to Xq13–q21 and evidence for genetic heterogeneity. *Genomics* 11: 885–894, 1991.
- [72670] 11000.Cardoso, C.; Timsit, S.; Villard, L.; Khrestchatisky, M.; Fontes, M.; Colleaux, L.: Specific interaction between the XNP/ATR–X gene product and the SET domain of the human EZH2 protein. *Hum. Molec. Genet.* 7: 679–684, 1998.
- [72671] 11001.Bairoch, A.: Personal Communication. Geneva, Switzerland 5/13/1994.
- [72672] 11002.Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.–H.; Goldstein, L. S.: Axonal transport of amyloid precursor protein is mediated by direct binding to the kinesin light chain subunit of kinesin–I. *Neuron* 28:449–459, 2000.
- [72673] 11003.Luo, Y.; Bolon, B.; Kahn, S.; Bennett, B. D.; Babu–Khan, S.; Denis, P.; Fan, W.; Kha, H.; Zhang, J.; Gong, Y.; Martin, L.; Louis, J.–C.; Yan, Q.; Richards, W. G.; Citron, M.; Vassar, R.: Mice deficient in BACE1, the Alzheimer's beta–secretase, have normal phenotype and abolished beta–amyloid generation. *Nature Neurosci.* 4: 231–232, 2001.



- [72674] 11004.Schiemann, B.; Gommerman, J. L.; Vora, K.; Cachero, T. G.; Shulga-Morskaya,S.; Dobles, M.; Frew, E.; Scott, M. L.: An essential role for BAFFin the normal development of B cells through a BCMA-independent pathway. Science 2111-2114,2001.
- [72675] 11005.Okamoto, N.; Hori, S.; Akazawa, C.; Hayashi, Y.; Shigemoto, R.;Mizuno, N.; Nakanishi, S.: Molecular characterization of a new metabotropicglutamate receptor mGluR7 coupled to inhibitory cyclic AMP signaltransduction. J. Biol. Chem. 269: 1231-1236, 1994.
- [72676] 11006.Ango, F.; Prezeau, L.; Muller, T.; Tu, J. C.; Xiao, B.; Worley,P. F.; Pin, J. P.; Bockaert, J.; Fagni, L.: Agonist-independent activationof metabotropic glutamate receptors by the intracellular protein Homer. Nature 411:962-965, 2001.
- [72677] 11007.Offenberg, H. H.; Schalk, J. A. C.; Meuwissen, R. L. J.; van Aalderen,M.; Kester, H. A.; Dietrich, A. J. J.; Heyting, C.: SCP2: a majorprotein component of the axial elements of synaptonemal complexesof the rat. Nucleic Acids Res. 26: 2572-2579, 1998.
- [72678] 11008.Schalk, J. A. C.; Offenberg, H. H.; Peters, E.; Groot, N. P. B.;Hoovers, J. M. N.; Heyting, C.: Isolation and characterization ofthe human SCP2 cDNA and chromosomal

localization of the gene. *MammalianGenome* 10: 642–644, 1999.

- [72679] 11009.Komada, M.; McLean, D. J.; Griswold, M. D.; Russell, L. D.; Soriano,P.: E-MAP-115, encoding a microtubule-associated protein, is a retinoicacid-inducible gene required for spermatogenesis. *Genes Dev.* 14:1332–1342, 2000.
- [72680] 11010.Masson, D.; Kreis, T. E.: Identification and molecular characterizationof E-MAP-115, a novel microtubule-associated protein predominantlyexpressed in epithelial cells. *J. Cell Biol.* 123: 357–371, 1993.
- [72681] 11011.Kowalski, P. E.; Freeman, J. D.; Nelson, D. T.; Mager, D. L.:Genomic structure and evolution of a novel gene (PLA2L) with duplicatedphospholipase A2-like domains. *Genomics* 39: 38–46, 1997.
- [72682] 11012.Liu, M.; Parker, R. M. C.; Darby, K.; Eyre, H. J.; Copeland, N.G.; Crawford, J.; Gilbert, D. J.; Sutherland, G. R.; Jenkins, N. A.;Herzog, H.: GPR56, a novel secretin-like human G-protein-coupledreceptor gene. *Genomics* 55: 296–305, 1999.
- [72683] 11013.Zendman, A. J. W.; Cornelissen, I. M. H. A.; Weidle, U. H.; Ruiter,D. J.; van Muijen, G. N. P.: TM7XN1, a novel human EGF-TM7-like cDNA,detected with mRNA differen-

tial display using human melanoma celllines with different metastatic potential. FEBS Lett. 446: 292–298,1999.

- [72684] 11014.Nicolaou, M.; Song, Y.–Q.; Sato, C. A.; Orlacchio, A.; Kawarai,T.; Medeiros, H.; Liang, Y.; Sorbi, S.; Richard, E.; Rogaev, E. I.;Moliaka, Y.; Bruni, A. C.; Jorge, R.; Percy, M.; Duara, R.; Farrer,L. A.; St George–Hyslop, P.; Rogaeva, E. A.: Mutations in the openreading frame of the beta–site APP cleaving enzyme (BACE) locus arenot a common cause of Alzheimer's disease. Neurogenetics 3: 203–206,2001.
- [72685] 11015.Saunders, A. J.; Kim, T.–W.; Tanzi, R. E.: BACE maps to chromosome11 and a BACE homolog, BACE2, reside in the obligate Down syndromeregion of chromosome 21. Science 286: 1255a, 1999. Note: ElectronicPublication.
- [72686] 11016.Sinha, S.; Anderson, J. P.; Barbour, R.; Basi, G. S.; Caccavello,R.; Davis, D.; Doan, M.; Dovey, H. F.; Frigon, N.; Hong, J.; Jacobson–Croak,K.; Jewett, N.; and 15 others: Purification and cloning of amyloidprecursor protein beta–secretase from human brain. Nature 402: 537–540,1999.
- [72687] 11017.Vassar, R.; Bennett, B. D.; Babu–Khan, S.; Kahn, S.; Mendiaz,E. A.; Dents, P.; Taplow, D. B.; Ross, S.; Amara, P.; Loeloff,R.; Luo, Y.; Fisher, S.; and 12 others: Beta–secretase cleavageof Alzheimer's amyloid precursor protein by the transmembrane asparticprotease BACE. Sci–

ence 286: 735–741, 1999.

- [72688] 11018. Yan, R.; Bienkowski, M. J.; Shuck, M. E.; Miao, H.; Tory, M. C.; Pauley, A. M.; Brashler, J. R.; Stratman, N. C.; Mathews, W. R.; Buhl, A. E.; Carter, D. B.; Tomasselli, A. G.; Parodi, L. A.; Heinrikson, R. L.; Gurney, M. E.: Membrane-anchored aspartyl protease with Alzheimer's disease beta-secretase activity. *Nature* 402: 533–536, 1999.
- [72689] 11019. Jagla, K.; Dolle, P.; Mattei, M.-G.; Jagla, T.; Schuhbauer, B.; Dretzen, G.; Bellard, F.; Bellard, M.: Mouse Lbx1 and human LBX1 define a novel mammalian homeobox gene family related to the Drosophila ladybird genes. *Mech. Dev.* 53: 345–356, 1995.
- [72690] 11020. Mennerich, D.; Schafer, K.; Braun, T.: Pax-3 is necessary but not sufficient for lbx1 expression in myogenic precursor cells of the limb. *Mech. Dev.* 73: 147–158, 1998.
- [72691] 11021. Schafer, K.; Braun, T.: Early specification of limb muscle precursor cells by the homeobox gene Lbx1h. *Nature Genet.* 23: 213–216, 1999.
- [72692] 11022. Wilson, P. J.; McGlinn, E.; Marsh, A.; Evans, T.; Arnold, J.; Wright, K.; Biden, K.; Young, J.; Wainwright, B.; Wicking, C.; Chenevix-Trench, G.: Sequence variants of DLC1 in colorectal and ovarian tumours. *Hum. Mutat.* 15: 156–165, 2000.

- [72693] 11023.Yuan, B.-Z.; Miller, M. J.; Keck, C. L.; Zimonjic, D. B.; Thorgeirsson,S. S.; Popescu, N. C.: Cloning, characterization, and chromosomallocalization of a gene frequently deleted in human liver cancer (DLC-1)homologous to rat RhoGAP. *Cancer Res.* 58: 2196–2199, 1998.
- [72694] 11024.Yuan, B.-Z.; Yang, Y.; Keck–Waggoner, C. L.; Zimonjic, D. B.; Thorgeirsson,S. S.; Popescu, N. C.: Assignment and cloning of mouse Arhgap7 tochromosome 8A4–B2, a conserved syntenic region of human chromosome8p22–p21. *Cytogenet. Cell Genet.* 87: 189–190, 1999.
- [72695] 11025.Arnould, C.; Philippe, C.; Bourdon, V.; Gregoire, M. J.; Berger,R.; Jonveaux, P.: The signal transducer and activator of transcriptionSTAT5b gene is a new partner of retinoic acid receptor alpha in acutepromyelocytic–like leukaemia. *Hum. Molec. Genet.* 8: 1741–1749, 1999.
- [72696] 11026.Lin, J.-X.; Mietz, J.; Modi, W. S.; John, S.; Leonard, W. J.:Cloning of human Stat5B: reconstitution of interleukin–2–induced Stat5Aand Stat5B DNA binding activity in COS–7 cells. *J. Biol. Chem.* 271:10738–10744, 1996.
- [72697] 11027.Shinoura, N.; Shamraj, O. I.; Hugenholtz, H.; Zhu, J. G.; McBlack,P.; Warnick, R.; Tew, J. J.; Wani, M. A.; Menon, A. G.: Identificationand partial sequence of a cDNA that is

differentially expressed in human brain tumors. *Cancer Lett.* 89: 215–221, 1995.

[72698] 11028. Rumsby, G.; Cregeen, D. P.: Identification and expression of cDNA for human hydroxypyruvate/glyoxylate reductase. *Biochim. Biophys. Acta* 1446: 383–388, 1999.

[72699] 11029. Webster, K. E.; Ferree, P. M.; Holmes, R. P.; Cramer, S. D.: Identification of missense, nonsense, and deletion mutations in the GRHPR gene in patients with primary hyperoxaluria type II (PH2). *Hum. Genet.* 107:176–185, 2000.

[72700] 11030. Cremona, O.; Di Paolo, G.; Wenk, M. R.; Luthi, A.; Kim, W. T.; Takei, K.; Daniell, L.; Nemoto, Y.; Shears, S. B.; Flavell, R. A.; McCormick, D. A.; De Camilli, P.: Essential role of phosphoinositide metabolism in synaptic vesicle recycling. *Cell* 99: 179–188, 1999.

[72701] 11031. Cremona, O.; Nimmakayalu, M.; Haffner, C.; Bray-Ward, P.; Ward, D. C.; De Camilli, P.: Assignment of SYNJ1 to human chromosome 21q22.2 and Synj12 (sic) to the murine homologous region on chromosome 16C3–4 by in situ hybridization. *Cytogenet. Cell Genet.* 88: 89–90, 2000.

[72702] 11032. Haffner, C.; Takei, K.; Chen, H.; Ringstad, N.; Hudson, A.; Butler, M. H.; Salcini, A. E.; Di Fiore, P. P.; De

Camilli, P.: Synaptojanin1: localization on coated endocytic intermediates in nerve terminals and interaction of its 170 kDa isoform with Eps15. *FEBS Lett.* 419:175–180, 1997.

[72703] 11033. McPherson, P. S.; Garcia, E. P.; Slepnev, V. I.; David, C.; Zhang, X.; Grabs, D.; Sossin, W. S.; Bauerfeind, R.; Nemoto, Y.; De Camilli, P.: A presynaptic inositol-5-phosphatase. *Nature* 379: 353–357, 1996.

[72704] 11034. Saito, T.; Guan, F.; Papolos, D. F.; Lau, S.; Klein, M.; Fann, C. S. J.; Lachman, H. M.: Mutation analysis of SYNJ1: a possible candidate gene for chromosome 21q22-linked bipolar disorder. *Molec. Psychiat.* 6: 387–395, 2001.

[72705] 11035. Aita, V. M.; Liang, X. H.; Murty, V. V. V. S.; Pincus, D. L.; Yu, W.; Cayanis, E.; Kalachikov, S.; Gilliam, T. C.; Levine, B.: Cloning and genomic organization of beclin 1, a candidate tumor suppressor gene on chromosome 17q21. *Genomics* 59: 59–65, 1999.

[72706] 11036. Liang, X. H.; Jackson, S.; Seaman, M.; Brown, K.; Kempkes, B.; Hibshoosh, H.; Levine, B.: Induction of autophagy and inhibition of tumorigenesis by beclin 1. *Nature* 402: 672–676, 1999.

[72707] 11037. Liang, X. H.; Kleeman, L. K.; Jiang, H. H.; Gordon, G.; Goldman, J. E.; Berry, G.; Herman, B.; Levine, B.: Protec-

tion against fatal Sindbis virus encephalitis by beclin, a novel Bcl-2-interacting protein. *J. Virol.* 72: 8586–8596, 1998.

- [72708] 11038. Liang, X. H.; Yu, J.; Brown, K.; Levine, B.: Beclin 1 contains a leucine-rich nuclear export signal that is required for its autophagy and tumor suppressor function. *Cancer Res.* 61: 3443–3449, 2001.
- [72709] 11039. Hatamura, I.; Kanauchi, Y.; Takahara, M.; Fujiwara, M.; Muragaki, Y.; Ooshima, A.; Ogino, T.: A nonsense mutation in TRPS1 in a Japanese family with trichorhino-phalangeal syndrome type I. (Letter) *Clin. Genet.* 59: 366–367, 2001.
- [72710] 11040. Hilton, M. J.; Sawyer, J. M.; Gutierrez, L.; Hogart, A.; Kung, T. C.; Wells, D. E.: Analysis of novel and recurrent mutations responsible for the trichorhino-phalangeal syndromes. *J. Hum. Genet.* 47: 103–106, 2002.
- [72711] 11041. Patel, N.; Brinkman-Van der Linden, E. C. M.; Altmann, S. W.; Gish, K.; Balasubramanian, S.; Timans, J. C.; Peterson, D.; Bell, M. P.; Bazan, J. F.; Varki, A.; Kastelein, R. A.: OB-BP1/Siglec-6: a leptin- and sialic acid-binding protein of the immunoglobulin superfamily. *J. Biol. Chem.* 274: 22729–22738, 1999. Note: Erratum: *J. Biol. Chem.* 274: 28058 only, 1999.



- [72712] 11042. Miller, A. F.; Harvey, S. A. K.; Thies, R. S.; Olson, M. S.: Bonemorphogenetic protein-9: an autocrine/paracrine cytokine in the liver. *J. Biol. Chem.* 275: 17937-17945, 2000.
- [72713] 11043. Scott, A. F.: Personal Communication. Baltimore, Md. 7/13/2000.
- [72714] 11044. Lin, C.-Y.; Anders, J.; Johnson, M.; Sang, Q. A.; Dickson, R. B.: Molecular cloning of cDNA for matriptase, a matrix-degrading serineprotease with trypsin-like activity. *J. Biol. Chem.* 274: 18231-18236, 1999.
- [72715] 11045. Oberst, M.; Anders, J.; Xie, B.; Singh, B.; Ossandon, M.; Johnson, M.; Dickson, R. B.; Lin, C.-Y.: Matriptase and HAI-1 are expressed by normal and malignant epithelial cells in vitro and in vivo. *Am. J. Path.* 158: 1301-1311, 2001.
- [72716] 11046. Kandil, E.; Kohda, K.; Ishibashi, T.; Tanaka, K.; Kasahara, M.: PA28 subunits of the mouse proteasome: primary structures and chromosomal localization of the genes. *Immunogenetics* 46: 337-344, 1997.
- [72717] 11047. Nikaido, T.; Shimada, K.; Shibata, M.; Hata, M.; Sakamoto, M.; Takasaki, Y.; Sato, C.; Takahashi, T.; Nishida, Y.: Cloning and nucleotide sequence of cDNA for Ki antigen, a highly conserved nuclear protein detected

with sera from patients with systemic lupus erythematosus. Clin.Exp. Immun. 79: 209–214, 1990.

- [72718] 11048.Bednarek, A. K.; Keck–Waggoner, C. L.; Daniel, R. L.; Laflin, K.J.; Bergsagel, P. L.; Kiguchi, K.; Brenner, A. J.; Aldaz, C. M.:WWOX, the FRA16D gene, behaves as a suppressor of tumor growth. CancerRes. 61: 8068–8073, 2001.
- [72719] 11049.Bednarek, A. K.; Laflin, K. J.; Daniel, R. L.; Liao, Q.; Hawkins,K. A.; Aldaz, C. M.: WWOX, a novel WW domain-containing protein mappingto human chromosome 16q23.3–24.1, a region frequently affected inbreast cancer. Cancer Res. 60: 2140–2145, 2000.
- [72720] 11050.Chang, N.–S.; Pratt, N.; Heath, J.; Schultz, L.; Sleve, D.; Carey,G. B.; Zevotek, N.: Hyaluronidase induction of a WW domain-containingoxidoreductase that enhances tumor necrosis factor cytotoxicity. J.Biol. Chem. 276: 3361–3370, 2001.
- [72721] 11051.Krummel, K. A.; Denison, S. R.; Calhoun, E.; Phillips, L. A.; Smith,D. I.: The common fragile site FRA16D and its associated gene WWOXare highly conserved in the mouse at Fra8E1. Genes Chromosomes Cancer 34:154–167, 2002.
- [72722] 11052.Miyake, A.; Mochizuki, S.; Yokoi, H.; Kohda, M.; Fu–

ruichi, K.:New ether-a-go-go K(+) channel family members localized in human telencephalon. J.Biol. Chem. 274: 25018–25025, 1999.

[72723] 11053.Acampora, D.; Postiglione, M. P.; Avantaggiato, V.; Di Bonito,M.; Vaccarino, F. M.; Michaud, J.; Simeone, A.: Progressive impairmentof developing neuroendocrine cell lineages in the hypothalamus ofmice lacking the Orthopedia gene. Genes Dev. 13: 2787–2800, 1999.

[72724] 11054.Lin, X.; State, M. W.; Vaccarino, F. M.; Greally, J.; Hass, M.;Leckman, J. F.: Identification, chromosomal assignment, and expressionanalysis of the human homeodomain-containing gene Orthopedia (OTP). Genomics 60:96–104, 1999.

[72725] 11055.Gongora, C.; David, G.; Pintard, L.; Tissot, C.; Hua, T. D.; Dejean,A.; Mechti, N.: Molecular cloning of a new interferon-induced PMLnuclear body-associated protein. J. Biol. Chem. 272: 19457–19463,1997.

[72726] 11056.Mattei, M. G.; Tissot, C.; Gongora, C.; Mechti, N.: Assignmentof ISG20 encoding a new interferon-induced PML nuclear body-associatedprotein, to chromosome 15q26 by in situ hybridization. Cytogenet.Cell Genet. 79: 286–287, 1997.

[72727] 11057.Pentecost, B. T.: Expression and estrogen regula-

tion of the HEM45mRNA in human tumor lines and in the rat uterus. *J. Steroid Biochem.Molec. Biol.* 64: 25–33, 1998.

- [72728] 11058.Berchtold, S.; Muhl–Zurbes, P.; Heufler, C.; Winkelner, P.; Schuler,G.; Steinkasserer, A.: Cloning, recombinant expression and biochemicalcharacterization of the murine CD83 molecule which is specificallyupregulated during dendritic cell maturation. *FEBS Lett.* 461: 211–216,1999.
- [72729] 11059.Fujimoto, Y.; Tu, L.; Miller, A. S.; Bock, C.; Fujimoto, M.; Doyle,C.; Steeber, D. A.; Tedder, T. F.: CD83 expression influences CD4+T cell development in the thymus. *Cell* 108: 755–767, 2002.
- [72730] 11060.Kozlow, E. J.; Wilson, G. L.; Fox, C. H.; Kehrl, J. H.: SubtractivecDNA cloning of a novel member of the Ig gene superfamily expressedat high levels in activated B lymphocytes. *Blood* 81: 454–461, 1993.
- [72731] 11061.Olavesen, M.G.; Bentley, E.; Mason, R. V.; Stephens, R. J.; Ragoussis,J.: Fine mapping of 39 ESTs on human chromosome 6p23–p25. *Genomics* 46:303–306, 1997.
- [72732] 11062.Scholler, N.; Hayden–Ledbetter, M.; Hellstrom, K.–E.; Hellstrom,I.; Ledbetter, J. A.: CD83 is a sialic acid–binding Ig–like lectin(Siglec) adhesion receptor that binds

monocytes and a subset of activated CD8(+) T cells. *J. Immunol.* 166: 3865–3872, 2001.

- [72733] 11063. Schinkmann, K.; Blenis, J.: Cloning and characterization of a human STE20-like protein kinase with unusual cofactor requirements. *J. Biol. Chem.* 272: 28695–28703, 1997.
- [72734] 11064. Zhou, T.-H.; Ling, K.; Guo, J.; Zhou, H.; Wu, Y.-L.; Jing, Q.; Ma, L.; Pei, G.: Identification of a human brain-specific isoform of mammalian STE20-like kinase 3 that is regulated by cAMP-dependent protein kinase. *J. Biol. Chem.* 275: 2513–2519, 2000.
- [72735] 11065. St-Pierre, M. V.; Hagenbuch, B.; Ugele, B.; Meier, P. J.; Stallmach, T.: Characterization of an organic anion-transporting polypeptide (OATP-B) in human placenta. *J. Clin. Endocr. Metab.* 87: 1856–1863, 2002.
- [72736] 11066. Murthy, A.; Gonzalez-Agosti, C.; Cordero, E.; Pinney, D.; Candia, C.; Solomon, F.; Gusella, J.; Ramesh, V.: NHE-RF, a regulatory cofactor for Na(+)-H(+) exchange, is a common interactor for merlin and ERM (ERM) proteins. *J. Biol. Chem.* 273: 1273–1276, 1998.
- [72737] 11067. Reczek, D.; Berryman, M.; Bretscher, A.: Identification of EBP50: a PDZ-containing phosphoprotein that associates with members of the ezrin-radixin-moesin family.

J. Cell Biol. 139: 169–179, 1997.

[72738] 11068. Shenolikar, S.; Voltz, J. W.; Minkoff, C. M.; Wade, J. B.; Weinman, E. J.: Targeted disruption of the mouse NHERF-1 gene promotes internalization of proximal tubule sodium-phosphate cotransporter type IIa and renal phosphate wasting. Proc. Nat. Acad. Sci. 99: 11470–11475, 2002.

[72739] 11069. Boucher, C. A.; Winchester, C. L.; Hamilton, G. M.; Winter, A. D.; Johnson, K. J.; Bailey, M. E. S.: Structure, mapping and expression of the human gene encoding the homeodomain protein, SIX2. Gene 247: 145–151, 2000.

[72740] 11070. Hasan, S.; Stucki, M.; Hassa, P. O.; Imhof, R.; Gehrig, P.; Hunziker, P.; Hubscher, U.; Hottiger, M. O.: Regulation of human flap endonuclease-1 activity by acetylation through the transcriptional coactivator p300. Molec. Cell 7: 1221–1231, 2001.

[72741] 11071. Hiraoka, L. R.; Harrington, J. J.; Gerhard, D. S.; Lieber, M. R.; Hsieh, C.-L.: Sequence of human FEN-1, a structure-specific endonuclease, and chromosomal localization of the gene (FEN1) in mouse and human. Genomics 25: 220–225, 1995.

[72742] 11072. Hosfield, D. J.; Mol, C. D.; Shen, B.; Tainer, J. A.: Structure of the DNA repair and replication endonuclease

and exonuclease FEN-1:coupling DNA and PCNA binding to FEN-1 activity. Cell 95: 135–146,1998.

[72743] 11073.Kucherlapati, M.; Yang, K.; Kuraguchi, M.; Zhao, J.; Lia, M.; Heyer,J.; Kane, M. F.; Fan, K.; Russell, R.; Brown, A. M. C.; Kneitz, B.;Edelmann, W.; Kolodner, R. D.; Lipkin, M.; Kucherlapati, R.: Haploinsufficiencyof Flap endonuclease (Fen1) leads to rapid tumor progression. Proc.Nat. Acad. Sci. 99: 9924–9929, 2002.

[72744] 11074.Otto, C. J.; Almqvist, E.; Hayden, M. R.; Andrew, S. E.: The 'flap'endonuclease gene FEN1 is excluded as a candidate gene implicatedin the CAG repeat expansion underlying Huntington disease. Clin.Genet. 59: 122–127, 2001.

[72745] 11075.Spiro, C.; Pelletier, R.; Rolfsmeier, M. L.; Dixon, M. J.; Lahue,R. S.; Gupta, G.; Park, M. S.; Chen, X.; Mariappan, S. V. S.; McMurray,C. T.: Inhibition of FEN-1 processing by DNA secondary structureat trinucleotide repeats. Molec. Cell 4: 1079–1085, 1999.

[72746] 11076.Tishkoff, D. X.; Filosi, N.; Gaida, G. M.; Kolodner, R. D.: Anovel mutation avoidance mechanism dependent on S. cerevisiae RAD27is distinct from DNA mismatch repair. Cell 88: 253–263, 1997.

[72747] 11077.Vermeesch, J. R.; Mertens, G.; David, G.; Marynen,

P.: Assignment of the human glypican gene (GPC1) to 2q35–q37 by fluorescence in situ hybridization. *Genomics* 25: 327–329, 1995.

- [72748] 11078. Ono, Y.; Ohno, M.; Shimura, Y.: Identification of a putative RNA helicase (HRH1), a human homolog of yeast Prp22. *Molec. Cell. Biol.* 14:7611–7620, 1994.
- [72749] 11079. Fulop, V.; Bocskei, Z.; Polgar, L.: Prolyl oligopeptidase: an unusual beta-propeller domain regulates proteolysis. *Cell* 94: 161–170, 1998.
- [72750] 11080. Goossens, F. J.; Wauters, J. G.; Vanhoof, G. C.; Bossuyt, P. J.; Schatteman, K. A.; Loens, K.; Scharpe, S. L.: Subregional mapping of the human lymphocyte prolyl oligopeptidase gene (PREP) to human chromosome 6q22. *Cytogenet. Cell Genet.* 74: 99–101, 1996.
- [72751] 11081. Vanhoof, G.; Goossens, F.; Hendriks, L.; De Meester, I.; Hendriks, D.; Vriend, G.; Van Broeckhoven, C.; Scharpe, S.: Cloning and sequence analysis of the gene encoding human lymphocyte prolyl endopeptidase. *Gene* 149:363–366, 1994.
- [72752] 11082. Goldstein, L. A.; Gherzi, G.; Pineiro-Sanchez, M. L.; Salamone, M.; Yeh, Y.; Flessate, D.; Chen, W.-T.: Molecular cloning of seprase: a serine integral membrane protease from human melanoma. *Biochim. Biophys. Acta* 1361:



11–19, 1997.

- [72753] 11083. Mathew, S.; Scanlan, M. J.; Mohan Raj, B. K.; Murty, V. V. V. S.; Garin-Chesa, P.; Old, L. J.; Rettig, W. J.; Chaganti, R. S. K.: The gene for fibroblast activation protein alpha (FAP), a putative cell surface-bound serine protease expressed in cancer stroma and wound healing, maps to chromosome band 2q23. *Genomics* 25: 335–337, 1995.
- [72754] 11084. Pineiro-Sanchez, M. L.; Goldstein, L. A.; Dodt, J.; Howard, L.; Yeh, Y.; Tran, H.; Argraves, W. S.; Chen, W.-T.: Identification of the 170-kDa melanoma membrane-bound gelatinase (seprase) as a serine integral membrane protease. *J. Biol. Chem.* 272: 7595–7601, 1997.
- [72755] 11085. Scanlan, M. J.; Raj, B. K. M.; Calvo, B.; Garin-Chesa, P.; Sanz-Moncasi, M. P.; Healey, J. H.; Old, L. J.; Rettig, W. J.: Molecular cloning of fibroblast activation protein alpha, a member of the serine protease family selectively expressed in stromal fibroblasts of epithelial cancers. *Proc. Nat. Acad. Sci.* 91: 5657–5661, 1994.
- [72756] 11086. Okumura, K.; Nogami, M.; Taguchi, H.; Dean, F. B.; Chen, M.; Pan, Z.-Q.; Hurwitz, J.; Shiratori, A.; Murakami, Y.; Ozawa, K.; Eki, T.: Assignment of the 36.5-kDa (RFC5), 37-kDa (RFC4), 38-kDa (RFC3), and 40-kDa (RFC2) subunit genes of human replication factor C to chromosome–

bands 12q24.2–q24.3, 3q27, 13q12.3–q13, and 7q11.23. Genomics 25:274–278, 1995.

- [72757] 11087. Peoples, R.; Perez–Jurado, L.; Wang, Y.–K.; Kaplan, P.; Francke, U.: The gene for replication factor C subunit 2 (RFC2) is within the 7q11.23 Williams syndrome deletion. (Letter) Am. J. Hum. Genet. 58:1370–1373, 1996.
- [72758] 11088. Wang, Y.; Cortez, D.; Yazdi, P.; Neff, N.; Elledge, S. J.; Qin, J.: BASC, a super complex of BRCA1–associated proteins involved in the recognition and repair of aberrant DNA structures. Genes Dev. 14:927–939, 2000.
- [72759] 11089. Thim, L.; Woldike, H. F.; Nielsen, P. F.; Christensen, M.; Lynch–Devaney, K.; Podolsky, D. K.: Characterization of human and rat intestinal trefoil factor produced in yeast. Biochemistry 34: 4757–4764, 1995.
- [72760] 11090. Itoh, T.; Mori, T.; Ohkubo, H.; Yamaizumi, M.: A newly identified patient with clinical xeroderma pigmentosum phenotype has a non–sense mutation in the DDB2 gene and incomplete repair in (6–4) photoproducts. J. Invest. Derm. 113: 251–257, 1999.
- [72761] 11091. Acebron, A.; Aza–Blanc, P.; Rossi, D. L.; Lamas, L.; Santisteban, P.: Congenital human thyroglobulin defect due to low expression of the thyroid–specific transcription factor TTF–1. J. Clin. Invest. 96:781–785, 1995.

- [72762] 11092. Guazzi, S.; Price, M.; De Felice, M.; Damante, G.; Mattei, M.-G.; Di Lauro, R.: Thyroid nuclear factor 1 (TTF-1) contains a homeodomain and displays a novel DNA binding specificity. *EMBO J.* 9: 3631-3639, 1990.
- [72763] 11093. Ikeda, K.; Clark, J. C.; Shaw-White, J. R.; Stahlman, M. T.; Boutell, C. J.; Whitsett, J. A.: Gene structure and expression of human thyroid transcription factor-1 in respiratory epithelial cells. *J. Biol. Chem.* 270: 8108-8114, 1995.
- [72764] 11094. Kimura, S.; Hara, Y.; Pineau, T.; Fernandez-Salguero, P.; Fox, C. H.; Ward, J. M.; Gonzalez, F. J.: The T/ebp null mouse: thyroid-specific enhancer-binding protein is essential for the organogenesis of the thyroid, lung, ventral forebrain, and pituitary. *Genes Dev.* 10:60-69, 1996.
- [72765] 11095. Fernandes-Alnemri, T.; Armstrong, R. C.; Krebs, J.; Srinivasula, S. M.; Wang, L.; Bullrich, F.; Fritz, L. C.; Trapani, J. A.; Tomaselli, K. J.; Litwack, G.; Alnemri, E. S.: In vitro activation of CPP32 and Mch3 by Mch4, a novel human apoptotic cysteine protease containing two FADD-like domains. *Proc. Nat. Acad. Sci.* 93: 7464-7469, 1996.
- [72766] 11096. Fernandes-Alnemri, T.; Litwack, G.; Alnemri, E. S.: CPP32, a novel human apoptotic protein with homology to

Caenorhabditis elegans cell death protein Ced-3 and mammalian interleukin-1 beta-converting enzyme. J.Biol. Chem. 269: 30761-30764, 1994.

- [72767] 11097.Fernando, P.; Kelly, J. F.; Balazsi, K.; Slack, R. S.; Megeney, L. A.: Caspase 3 activity is required for skeletal muscle differentiation. Proc.Nat. Acad. Sci. 99: 11025-11030, 2002.
- [72768] 11098.Huang, Y.; Shin, N.-H.; Sun, Y.; Wang, K. K. W.: Molecular cloning and characterization of a novel caspase-3 variant that attenuates apoptosis induced by proteasome inhibition. Biochem. Biophys. Res. Commun. 283: 762-769, 2001.
- [72769] 11099.Kuida, K.; Zheng, T. S.; Na, S.; Kuan, C.; Yang, D.; Karasuyama, H.; Rakio, P.; Flavell, R. A.: Decreased apoptosis in the brain and premature lethality in CPP32-deficient mice. Nature 384: 368-372, 1996.
- [72770] 11100.Levkau, B.; Koyama, H.; Raines, E. W.; Clurman, B. E.; Herren, B.; Orth, K.; Roberts, J. M.; Ross, R.: Cleavage of p21(Cip1/Waf1) and p27(Kip1) mediates apoptosis in endothelial cells through activation of Cdk2: role of a caspase cascade. Molec. Cell 1: 553-563, 1998.
- [72771] 11101.Nasir, J.; Theilmann, J. L.; Chopra, V.; Jones, A. M.; Walker, D.; Rasper, D. M.; Vaillancourt, J. P.; Hewitt, J. E.;

Nicholson,D. W.; Hayden, M. R.: Localization of the cell death genes CPP32and Mch-2 to human chromosome 4q. Mammalian Genome 8: 56-59, 1997.

[72772] 11102.Tiso, N.; Pallavicini, A.; Muraro, T.; Zimbello, R.; Apolloni,E.; Valle, G.; Lanfranchi, G.; Danieli, G. A.: Chromosomal localizationof the human genes, CPP32, Mch2, Mch3, and Ich-1, involved in cellularapoptosis. Biochem. Biophys. Res. Commun. 225: 983-989, 1996.

[72773] 11103.Woo, M.; Hakem, R.; Soengas, M. S.; Duncan, G. S.; Shahinian,A.; Kagi, K.; Hakem, A.; McCurrach, M.; Khoo, W.; Kaufman, S. A.;Senaldi, G.; Howard, T.; Lowe, S. W.; Mak, T. W.: Essential contributionof caspase 3/CPP32 to apoptosis and its associated nuclear changes. GenesDev. 12: 806-819, 1998.

[72774] 11104.Nicholson, D. W.; Ali, A.; Thornberry, N. A.; Vaillancourt, J.P.; Ding, C. K.; Gallant, M.; Gareau, Y.; Griffin, P. R.; Labelle,M.; Lazebnik, Y. A.; Munday, N. A.; Raju, S. M.; Smulson, M. E.; Yamin,T.-T.; Yu, V. L.; Miller, D. K.: Identification and inhibition ofthe ICE/CED-3 protease necessary for mammalian apoptosis. Nature 376:37-43, 1995. MEDLINE UID: 95319529

[72775] 11105.Bergeron, L.; Perez, G. I.; Macdonald, G.; Shi, L.; Sun, Y.; Jurisicova,A.; Varmuza, S.; Latham, K. E.; Flaws, J.

A.; Salter, J. C. M.; Hara, H.; Moskowitz, M. A.; Li, E.; Greenberg, A.; Tilly, J. L.; Yuan, J.: Defects in regulation of apoptosis in caspase-2-deficient mice. *GenesDev.* 12: 1304-1314, 1998.

[72776] 11106. Garcia, C. K.; Goldstein, J. L.; Pathak, R. K.; Anderson, R. G.W.; Brown, M. S.: Molecular characterization of a membrane transporter for lactate, pyruvate, and other monocarboxylates: implications for the Cori cycle. *Cell* 76: 865-873, 1994.

[72777] 11107. Garcia, C. K.; Li, X.; Luna, J.; Francke, U.: cDNA cloning of the human monocarboxylate transporter 1 and chromosomal localization of the SLC16A1 locus to 1p13.2-p12. *Genomics* 23: 500-503, 1994.

[72778] 11108. Kim, C. M.; Goldstein, J. L.; Brown, M. S.: cDNA cloning of mev, a mutant protein that facilitates cellular uptake of mevalonate, and identification of the point mutation responsible for its gain of function. *J. Biol. Chem.* 267: 23113-23121, 1992.

[72779] 11109. Kingsmore, S. F.; Souryal, C. A.; Watson, M. L.; Patel, D. D.; Seldin, M. F.: Physical and genetic linkage of the genes encoding Ly-9 and CD48 on mouse and human chromosomes 1. *Immunogenetics* 42:59-62, 1995.

[72780] 11110. Kozak, C. A.; Davidson, W. F.; Morse, H. C., III: Ge-

netic and functional relationships of the retroviral and lymphocyte alloantigen loci on mouse chromosome 1. Immunogenetics 19: 163–168, 1984.

- [72781] 11111. Sandrin, M. S.; Gumley, T. P.; Henning, M. M.; Vaughan, H. A.; Gonez, L. J.; Trapani, J. A.; McKenzie, I. F. C.: Isolation and characterization of cDNA clones for mouse Ly-9. J. Immun. 149: 1636–1641, 1992.
- [72782] 11112. Leek, J. P.; Moynihan, T. P.; Anwar, R.; Bonthron, D. T.; Markham, A. F.; Lench, N. J.: Assignment of Indian hedgehog (IHH) to human chromosome bands 2q33–q35 by in situ hybridization. Cytogenet. Cell Genet. 76: 187–188, 1997.
- [72783] 11113. Porter, J. A.; Young, K. E.; Beachy, P. A.: Cholesterol modification of hedgehog signaling proteins in animal development. Science 274: 255–259, 1996.
- [72784] 11114. Zhang, X. M.; Ramalho-Santos, M.; McMahon, A. P.: Smoothed mutants reveal redundant roles for Shh and Ihh signaling including regulation of L/R asymmetry by the mouse node. Cell 105: 781–792, 2001.
- [72785] 11115. Geurts, J. M. W.; Schoenmakers, E. F. P. M.; Roijer, E.; Astrom, A.-K.; Stenman, G.; van de Ven, W. J. M.: Identification of NF1B as recurrent translocation partner gene of HMGIC in pleomorphic adenomas. Oncogene 16: 865–872,

1998.

- [72786] 11116.Fiedorek, F. T., Jr.; Kay, E. S.: Mapping of the insulin promoter factor 1 gene (Ip1) to distal mouse chromosome 5. *Genomics* 28:581–584, 1995.
- [72787] 11117.Hani, E. H.; Stoffers, D. A.; Chevre, J.–C.; Durand, E.; Stanojevic,V.; Dina, C.; Habener, J. F.; Froguel, P.: Defective mutations in the insulin promoter factor–1 (IPF–1) gene in late–onset type 2 diabetes mellitus. *J. Clin. Invest.* 104: R41–R48, 1999.
- [72788] 11118.Tang, J. Y.; Hwang, B. J.; Ford, J. M.; Hanawalt, P. C.; Chu, G.: Xeroderma pigmentosum p48 gene enhances global genomic repair and suppresses UV–induced mutagenesis. *Molec. Cell* 5: 737–744, 2000.
- [72789] 11119.Birmingham, J. R., Jr.; Arden, K. C.; Naumova, A. K.; Sapienza,C.; Viars, C. S.; Fu, X.–D.; Khotz, J.; Manley, J. L.; Rosenfeld,M. G.: Chromosomal localization of mouse and human genes encoding the splicing factors ASF/SF2 (SFRS1) and SC–35 (SFRS2). *Genomics* 29:70–79, 1995.
- [72790] 11120.Wang, J.; Takagaki, Y.; Manley, J. L.: Targeted disruption of an essential vertebrate gene: ASF/SF2 is required for cell viability. *GenesDev.* 10: 2588–2599, 1996.
- [72791] 11121.Hopfner, K.–P.; Karcher, A.; Craig, L.; Woo, T. T.; Carney, J.P.; Tainer, J. A.: Structural biochemistry and in–



teraction architecture of the DNA double-strand break repair Mre11 nuclease and Rad50-ATPase. *Cell* 105:473-485, 2001.

[72792] 11122. Stracker, T. H.; Carson, C. T.; Weitzman, M. D.: Adenovirus oncoproteins inactivate the Mre11-Rad50-NBS1 DNA repair complex. *Nature* 418:348-352, 2002.

[72793] 11123. Zhong, Q.; Chen, C.-F.; Li, S.; Chen, Y.; Wang, C.-C.; Xiao, J.; Chen, P.-L.; Sharp, Z. D.; Lee, W.-H.: Association of BRCA1 with the hRad50-hMre11-p95 complex and the DNA damage response. *Science* 285:747-750, 1999.

[72794] 11124. Zhu, X.-D.; Kuster, B.; Mann, M.; Petrini, J. H. J.; de Lange, T.: Cell-cycle-regulated association of RAD50/MRE11/NBS1 with TRF2 and human telomeres. *Nature Genet.* 25: 347-352, 2000.

[72795] 11125. Virbasius, C. A.; Virbasius, J. V.; Scarpulla, R. C.: NRF-1, an activator involved in nuclear-mitochondrial interactions, utilizes a new DNA-binding domain conserved in a family of developmental regulators. *Genes Dev.* 7: 2431-2445, 1993.

[72796] 11126. Choi, S. S.; Park, I.-C.; Yun, J. W.; Sung, Y. C.; Hong, S.-I.; Shin, H.-S.: A novel Bcl-2 related gene, Bfl-1,

- is overexpressed in stomach cancer and preferentially expressed in bone marrow. *Oncogene* 11:1693–1698, 1995.
- [72797] 11127. Choi, S. S.; Park, S. H.; Kim, U.-J.; Shin, H.-S.: Bfl-1, a Bcl-2-related gene, is the human homolog of the murine A1, and maps to chromosome 15q24.3. *Mammalian Genome* 8: 781–782, 1997.
- [72798] 11128. D'Sa-Eipper, C.; Subramanian, T.; Chinnadurai, G.: Bfl-1, a bcl-2 homologue, suppresses p53-induced apoptosis and exhibits potent cooperative transforming activity. *Cancer Res.* 56: 3879–3882, 1996.
- [72799] 11129. Lin, E. Y.; Kozak, C. A.; Orlofsky, A.; Prystowsky, M. B.: The bcl-2 family member, Bcl2a1, maps to mouse chromosome 9 and human chromosome 15. *Mammalian Genome* 8: 293–294, 1997.
- [72800] 11130. Lin, E. Y.; Orlofsky, A.; Berger, M. S.; Prystowsky, M. B.: Characterization of A1, a novel hemopoietic-specific early-response gene with sequence similarity to bcl-2. *J. Immun.* 151: 1979–1988, 1993.
- [72801] 11131. Lin, E. Y.; Orlofsky, A.; Wang, H.-G.; Reed, J. C.; Prystowsky, M. B.: A1, a Bcl-2 family member, prolongs cell survival and permits myeloid differentiation. *Blood* 87: 983–992, 1996.
- [72802] 11132. Xiang, Z.; Ahmed, A. A.; Moller, C.; Nakayama, K.;

Hatakeyama,S.; Nilsson, G.: Essential role of the prosurvival bcl-2 homologueA1 in mast cell survival after allergic activation. J. Exp. Med. 194:1561–1569, 2001.

- [72803] 11133.Bass, B. L.; Weintraub, H.: An unwinding activity that covalently modifies its double-stranded RNA substrate. Cell 55: 1089–1098,1988.
- [72804] 11134.Kim, U.; Wang, Y.; Sanford, T.; Zeng, Y.; Nishikura, K.: Molecular cloning of cDNA for double-stranded RNA adenosine deaminase, a candidate enzyme for nuclear RNA editing. Proc. Nat. Acad. Sci. 91: 11457–11461,1994.
- [72805] 11135.O'Connell, M. A.; Krause, S.; Higuchi, M.; Hsuan, J. J.; Totty,N. F.; Jenny, A.; Keller, W.: Cloning of cDNAs encoding mammalian double-stranded RNA-specific adenosine deaminase. Molec. Cell. Biol. 15:1389–1397, 1995.
- [72806] 11136.Patterson, J. B.; Samuel, C. E.: Expression and regulation by interferon of a double-stranded-RNA-specific adenosine deaminase from human cells: evidence for two forms of the deaminase. Molec. Cell.Biol. 15: 5376–5388, 1995.
- [72807] 11137.Wang, Q.; Khillan, J.; Gadue, P.; Nishikura, K.: Requirement of the RNA editing deaminase ADAR1 gene for embryonic erythropoiesis. Science 290:1765–1768, 2000.
- [72808] 11138.Wang, Y.; Zeng, Y.; Murray, J. M.; Nishikura, K.: Ge-

nomic organization and chromosomal location of the human dsRNA adenosine deaminase gene: the enzyme for glutamate-activated ion channel RNA editing. J.

Molec.Biol. 254: 184–195, 1995.

[72809] 11139. Weier, H.-U. G.; George, C. X.; Greulich, K. M.; Samuel, C. E.: The interferon-inducible, double-stranded RNA-specific adenosine deaminase gene (DSRAD) maps to human chromosome 1q21.1–21.2. Genomics 30: 372–375, 1995.

[72810] 11140. Weier, H.-U. G.; George, C. X.; Lersch, R. A.; Breitwieser, S.; Cheng, J.-F.; Samuel, C. E.: Assignment of the RNA-specific adenosine deaminase gene (Adar) to mouse chromosome 3F2 by in situ hybridization. Cytogenet. Cell Genet. 89: 214–215, 2000.

[72811] 11141. Kawagoe, H.; Soma, O.; Goji, J.; Nishimura, N.; Narita, M.; Inazawa, J.; Nakamura, H.; Sano, K.: Molecular cloning and chromosomal assignment of the human brain-type phosphodiesterase I/nucleotide pyrophosphatase gene (PDNP2). Genomics 30: 380–384, 1995.

[72812] 11142. Murata, J.; Lee, H. Y.; Clair, T.; Krutzsch, H. C.; Arestad, A. A.; Sobel, M. E.; Liotta, L. A.; Stracke, M. L.: cDNA cloning of human tumor motility-stimulating protein, autotaxin, reveals a homology with phosphodi-

esterases. J. Biol. Chem. 269: 30479–30484, 1994.

[72813] 11143.Narita, M.; Goji, J.; Nakamura, H.; Sano, K. Molecular cloning, expression, and localization of a brain-specific phosphodiesterase I/nucleotide (PD-I-alpha) from rat brain. J. Biol. Chem. 269: 28235–28242, 1994.

[72814] 11144.Piao, J.-H.; Matsuda, Y.; Nakamura, H.; Sano, K.: Assignment of Pdn2, the gene encoding phosphodiesterase I/nucleotide pyrophosphatase2, to mouse chromosome 15D2. Cytogenet. Cell Genet. 87: 172–174, 1999.

[72815] 11145.Tokumura, A.; Majima, E.; Kariya, Y.; Tominaga, K.; Kogure, K.; Yasuda, K.; Fukuzawa, K.: Identification of human plasma lysophospholipase D, a lysophosphatidic acid-producing enzyme, as autotaxin, a multifunctional phosphodiesterase. J. Biol. Chem. 277: 39436–39442, 2002.

[72816] 11146.Lin, Q.; Schwarz, J.; Bucana, C.; Olson, E. N.: Control of mouse cardiac morphogenesis and myogenesis by transcription factor MEF2C. Science 276:1404–1407, 1997.

[72817] 11147.Erickson, H. P.: A tenascin knockout with a phenotype. Nature Genet. 17: 5–7, 1997.

[72818] 11148.Akama, T. O.; Nakagawa, H.; Sugihara, K.; Narisawa, S.; Ohyama, C.; Nishimura, S.-I.; O'Brien, D. A.; Moremen, K. W.; Millan, J. L.; Fukuda, M. N.: Germ cell sur-

vival through carbohydrate-mediated interaction with Ser-toli cells. *Science* 295: 124–127, 2002.

- [72819] 11149. Bai, R.-Y.; Koester, C.; Ouyang, T.; Hahn, S. A.; Hammerschmidt, M.; Peschel, C.; Duyster, J.: SMIF, a Smad4-interacting protein that functions as a co-activator in TGF-beta signalling. *Nature Cell Biol.* 4:181–190, 2002.
- [72820] 11150. Akiyama, Y.; Tsubouchi, N.; Yuasa, Y.: Frequent somatic mutations of hMSH3 with reference to microsatellite instability in hereditary nonpolyposis colorectal cancer. *Biochem. Biophys. Res. Commun.* 236:248–252, 1997.
- [72821] 11151. Chittenden, T.; Harrington, E. A.; O'Connor, R.; Flemington, C.; Lutz, R. J.; Evan, G. I.; Guild, B. C.: Induction of apoptosis by the Bcl-2 homologue Bak. *Nature* 374: 733–736, 1995.
- [72822] 11152. Herberg, J. A.; Phillips, S.; Beck, S.; Jones, T.; Sheer, D.; Wu, J. J.; Prochazka, V.; Barr, P. J.; Kiefer, M. C.; Trowsdale, J.: Genomic structure and domain organisation of the human Bak gene. *Gene* 211:87–94, 1998.
- [72823] 11153. Kiefer, M. C.; Brauer, M. J.; Powers, V. C.; Wu, J. J.; Umansky, S. R.; Tomei, L. D.; Barr, P. J.: Modulation of apoptosis by the widely distributed Bcl-2 homologue Bak. *Nature* 374: 736–739, 1995.
- [72824] 11154. Ulrich, E.; Kauffmann-Zeh, A.; Hueber, A.-O.;

Williamson, J.; Chittenden, T.; Ma, A.; Evan, G.: Gene structure, cDNA sequence, and expression of murine Bak, a proapoptotic Bcl-2 family member. *Genomics* 44:195–200, 1997.

[72825] 11155. Higuchi, M.; Maas, S.; Single, F. N.; Hartner, J.; Rozov, A.; Burnashev, N.; Feldmeyer, D.; Sprengel, R.; Seeburg, P. H.: Point mutation in an AMPA receptor gene rescues lethality in mice deficient in the RNA-editing enzyme ADAR2. *Nature* 406: 78–81, 2000.

[72826] 11156. Lai, F.; Chen, C.-X.; Carter, K. C.; Nishikura, K.: Editing of glutamate receptor B subunit ion channel RNAs by four alternatively spliced DRADA2 double-stranded RNA adenosine deaminases. *Molec. Cell. Biol.* 17: 2413–2424, 1997.

[72827] 11157. Melcher, T.; Maas, S.; Herb, A.; Sprengel, R.; Seeburg, P. H.; Higuchi, M.: A mammalian RNA editing enzyme. *Nature* 379: 460–463, 1996.

[72828] 11158. Mittaz, L.; Scott, H. S.; Rossier, C.; Seeburg, P. H.; Higuchi, M.; Antonarakis, S. E.: Cloning of a human RNA editing deaminase (ADARB1) of glutamate receptors that maps to chromosome 21q22.3. *Genomics* 41:210–217, 1997.

[72829] 11159. O'Connell, M. A.; Gerber, A.; Keller, W.: Purification

of humandouble-stranded RNA-specific editase 1 (hRED1) involved in editingof brain glutamate receptor B pre-mRNA. J. Biol. Chem. 272: 473–478,1997.

- [72830] 11160.Villard, L.; Tassone, F.; Haymowicz, M.; Welborn, R.; Gardiner,K.: Map location, genomic organization and expression patterns ofthe human RED1 RNA editase. Somat. Cell Molec. Genet. 23: 135–145,1997.
- [72831] 11161.Yang, J.–H.; Sklar, P.; Axel, R.; Maniatis, T.: Purification andcharacterization of a human RNA adenosine deaminase for glutamatereceptor B pre-mRNA editing. Proc. Nat. Acad. Sci. 94: 4354–4359,1997.
- [72832] 11162.Lu, C.–M.; Han, J.; Rado, T. A.; Brown, G. B.: Differential expressionof two sodium channel subtypes in human brain. FEBS Letters 303:53–58, 1992.
- [72833] 11163.Zhao, Z.; Lee, C.–C.; Monckton, D. G.; Yazdani, A.; Coolbaugh,M. I.; Li, X.; Bailey, J.; Shen, Y.; Caskey, C. T.: Characterizationandgenomic mapping of genes and pseudogenes of a new human proteintyrosine phosphatase. Genomics 35: 172–181, 1996.
- [72834] 11164.Chopra, V. S.; Metzler, M.; Rasper, D. M.; Engqvist–Goldstein,A. E. Y.; Singaraja, R.; Gan, L.; Fichter, K. M.; McCutcheon, K.;Drubin, D.; Nicholson, D. W.; Hayden, M. R.: HIP12 is a non-proapoptoticmember of a gene family



including HIP1, an interacting protein with huntingtin.

Mammalian Genome 11: 1006–1015, 2000.

- [72835] 11165. Borrás, T.; Rowlette, L. L. S.; Tamm, E. R.; Gottanka, J.; Epstein, D. L.: Effects of elevated intraocular pressure on outflow facility and TIGR/MYOC expression in perfused human anterior segments. Invest. Ophthalmol. Vis. Sci. 43: 33–40, 2002.
- [72836] 11166. Cerretti, D. P.; Nelson, N.: Characterization of the genes from mouse LERK-3/Ephrin-A3 (Epl3), mouse LERK-4/Ephrin-A4 (Epl4), and human LERK-6/Ephrin-A2 (EPLG6): conservation of intron/exon structure. Genomics 47: 131–135, 1998.
- [72837] 11167. Ma, T.; Yang, B.; Kuo, W.-L.; Verkman, A. S.: cDNA cloning and gene structure of a novel water channel expressed exclusively in human kidney: evidence for a gene cluster of aquaporins at chromosome locus 12q13. Genomics 35: 543–550, 1996.
- [72838] 11168. Yasui, M.; Kwon, T.-H.; Knepper, M. A.; Nielsen, S.; Agre, P.: Aquaporin-6: an intracellular vesicle water channel protein in renal epithelia. Proc. Nat. Acad. Sci. 96: 5808–5813, 1999.
- [72839] 11169. Shan, X.; Bourdeau, A.; Rhoton, A.; Wells, D. E.; Cohen, E. H.; Landgraf, B. E.; Palfree, R. G. E.: Characteriza-

tion and mapping to human chromosome 8q24.3 of Ly-6-related gene 9804 encoding an apparent homologue of mouse TSA-1. *J. Immun.* 160: 197–208, 1998.

- [72840] 11170. Bova, G. S.; MacGrogan, D.; Levy, A.; Pin, S. S.; Bookstein, R.; Isaacs, W. B.: Physical mapping of chromosome 8p22 markers and their homozygous deletion in a metastatic prostate cancer. *Genomics* 35:46–54, 1996.
- [72841] 11171. An, S.; Yang, J.; Xia, M.; Goetzl, E. J.: Cloning and expression of the EP2 subtype of human receptors for prostaglandin E2. *Biochem. Biophys. Res. Comm.* 197: 263–270, 1993.
- [72842] 11172. Bastien, L.; Sawyer, N.; Grygorczyk, R.; Metters, K. M.; Adam, M.: Cloning, functional expression and characterization of the human prostaglandin E2 receptor EP2 subtype. *J. Biol. Chem.* 269: 11873–11877, 1994.
- [72843] 11173. Coleman, R. A.; Grix, S. P.; Head, S. A.; Louttit, J. B.; Mallett, A.; Sheldrick, R. L. G.: A novel inhibitory prostanoid receptor in piglet saphenous vein. *Prostaglandins* 47: 151–168, 1994.
- [72844] 11174. Foord, S. M.; Marks, B.; Stolz, M.; Bufflier, E.; Fraser, N. J.; Lee, M. G.: The structure of the prostaglandin EP4 receptor gene and related pseudogenes. *Genomics* 35: 182–188, 1996.

- [72845] 11175.Mori, K.; Tanaka, I.; Kotani, M.; Miyaoka, F.; Sando, T.; Muro,S.; Sasaki, Y.; Nakagawa, O.; Ogawa, Y.; Usui, T.; Ozaki, S.; Ichikawa,A.; Narumiya, S.; Nakao, K.: Gene expression of the human prostaglandinE receptor EP4 subtype: differential regulation in monocytoid andlymphoid lineage cells by phorbol ester. J. Molec. Med. 74: 333–336,1996.
- [72846] 11176.Regan, J. W.; Bailey, T. J.; Pepperl, D. J.; Pierce, K. L.; Bogardus,A. M.; Donello, J. E.; Fairbairn, C. E.; Kedzie, K. M.; Woodward,D. F.; Gil, D. W.: Cloning of a novel human prostaglandin receptorwith characteristics of the pharmacologically defined EP2 subtype. Molec.Pharm. 46: 213–220, 1994.
- [72847] 11177.Segi, E.; Sugimoto, Y.; Yamasaki, A.; Aze, Y.; Oida, H.; Nishimura,T.; Murata, T.; Matsuoka, T.; Ushikubi, F.; Hirose, M.; Tanaka, T.;Yoshida, N.; Narumiya, S.; Ichikawa, A.: Patent ductus arteriosusand neonatal death in prostaglandin receptor EP4–deficient mice. Biochem.Biophys. Res. Comm. 246: 7–12, 1998.
- [72848] 11178.Yoshida, K.; Oida, H.; Kobayashi, T.; Maruyama, T.; Tanaka, M.;Katayama, T.; Yamaguchi, K.; Segi, E.; Tsuboyama, T.; Matsushita,M.; Ito, K.; Ito, Y.; Sugimoto, Y.; Ushikubi, F.; Ohuchida, S.; Kondo,K.; Nakamura, T.; Naru–

miya, S.: Stimulation of bone formation and prevention of bone loss by prostaglandin E EP4 receptor activation. Proc.Nat. Acad. Sci. 99: 4580–4585, 2002.

- [72849] 11179.Jadayel, D. M.; Osborne, L. R.; Coignet, L. J. A.; Zani, V. J.;Tsui, L.–C.; Scherer, S. W.; Dyer, M. J. S.: The BCL7 gene family:deletion of BCL7B in Williams syndrome. Gene 224: 35–44, 1998.
- [72850] 11180.Arima, T.; Drewell, R. A.; Arney, K. L.; Inoue, J.; Makita, Y.;Hata, A.; Oshimura, M.; Wake, N.; Surani, M. A.: A conserved imprintingcontrol region at the HYMAI/ZAC domain is implicated in transientneonatal diabetes mellitus. Hum. Molec. Genet. 10: 1475–1483, 2001.
- [72851] 11181.Joung, I.; Strominger, J. L.; Shin, J.: Molecular cloning of aphosphotyrosine–independent ligand of the p56–lck SH2 domain. Proc.Nat. Acad. Sci. 93: 5991–5995, 1996.
- [72852] 11182.Laurin, N.; Brown, J. P.; Morissette, J.; Raymond, V.: Recurrentmutation of the gene encoding sequestosome 1 (SQSTM1/p62) in Pagetdisease of bone. Am. J. Hum. Genet. 70: 1582–1588, 2002.
- [72853] 11183.Park, I.; Chung, J.; Walsh, C. T.; Yun, Y.; Strominger, J. L.;Shin, J.: Phosphotyrosine–independent binding of a 62–kDa proteinto the src homology 2 (SH2) do–

main of p56-lck and its regulation by phosphorylation of ser-59 in the lck unique N-terminal region. Proc.Nat. Acad. Sci. 92: 12338–12342, 1995.

[72854] 11184.Vadlamudi, R. K.; Joung, I.; Strominger, J. L.; Shin, J.: p62, a phosphotyrosine-independent ligand of the SH2 domain of p56-lck, belongs to a new class of ubiquitin-binding proteins. J. Biol. Chem. 271:20235–20237, 1996.

[72855] 11185.Akbar, G. K. M.; Dasari, V. R.; Webb, T. E.; Ayyanathan, K.; Pillarisetti, K.; Sandhu, A. K.; Athwal, R. S.; Daniel, J. L.; Ashby, B.; Barnard, E. A.; Kunapuli, S. P.: Molecular cloning of a novel P2 purinoceptor from human erythroleukemia cells. J. Biol. Chem. 271: 18363–18367, 1996.

[72856] 11186.Chen, X.-S.; Sheller, J. R.; Johnson, E. N.; Funk, C. D.: Role of leukotrienes revealed by targeted disruption of 5-lipoxygenase gene. Nature 372: 179–182, 1994.

[72857] 11187.Devchand, P. R.; Keller, H.; Peters, J. M.; Vazquez, M.; Gonzalez, F. J.; Wahli, W.: The PPAR alpha-leukotriene B4 pathway to inflammation control. Nature 384: 39–43, 1996.

[72858] 11188.Kato, K.; Yokomizo, T.; Izumi, T.; Shimizu, T.: Cell-specific transcriptional regulation of human leukotriene B4 receptor gene. J. Exp. Med. 192: 413–420, 2000.

- [72859] 11189.Owman, C.; Nilsson, C.; Lolait, S. J.: Cloning of cDNA encoding a putative chemoattractant receptor. *Genomics* 37: 187–194, 1996.
- [72860] 11190.Samuelsson, B.; Dahlen, S. E.; Lindgren, J. A.; Rouzer, C. A.; Serhan, C. N.: Leukotrienes and lipoxins: structures, biosynthesis, and biological effects. *Science* 237: 1171–1176, 1987.
- [72861] 11191.Yokomizo, T.; Izumi, T.; Chang, K.; Takuwa, Y.; Shimizu, T.: A G-protein-coupled receptor for leukotriene B<sub>4</sub> that mediates chemotaxis. *Nature* 387:620–624, 1997.
- [72862] 11192.Kubo, Y.; Reuveny, E.; Slesinger, P. A.; Jan, Y. N.; Jan, L. Y.: Primary structure and functional expression of a rat G-protein-coupled muscarinic potassium channel. *Nature* 364: 802–806, 1993.
- [72863] 11193.Schoots, O.; Voskoglou, T.; Van Tol, H. H. M.: Genomic organization and promoter analysis of the human G-protein-coupled K<sup>+</sup> channel Kir3.1(KCNJ3/HGIRK1). *Genomics* 39: 279–288, 1997.
- [72864] 11194.Schoots, O.; Yue, K.-T.; MacDonald, J. F.; Hampson, D. R.; Nobrega, J. N.; Dixon, L. M.; Van Tol, H. H. M.: Cloning of a G protein-activated inwardly rectifying potassium channel from human cerebellum. *Molec. Brain Res.* 39: 23–30, 1996.

- [72865] 11195.Stoffel, M.; Espinosa, R., III; Powell, K. L.; Philipson, L. H.;Le Beau, M. M.; Bell, G. I.: Human G-protein-coupled inwardly rectifyingpotassium channel (GIRK1) gene (KCNJ3): localization to chromosome2 and identification of a simple tandem repeat polymorphism. Genomics 21:254–256, 1994.
- [72866] 11196.Phillips, H. A.; Scheffer, I. E.; Berkovic, S. F.; Hollway, G.E.; Sutherland, G. R.; Mulley, J. C.: Localization of a gene forautosomal dominant nocturnal frontal lobe epilepsy to chromosome 20q13.2. NatureGenet. 10: 117–118, 1995.
- [72867] 11197.Phillips, H. A.; Scheffer, I. E.; Crossland, K. M.; Bhatta, K.P.; Fish, D. R.; Marsden, C. D.; Howell, S. J. L.; Stephenson, J.B. P.; Tolmie, J.; Plazzi, G.; Eeg-Olofsson, O.; Singh, R.; Lopes-Cendes,I.; Andermann, E.; Andermann, F.; Berkovic, S. F.; Mulley, J. C.:Autosomal dominant nocturnal frontal-lobe epilepsy: genetic heterogeneityand evidence for a second locus at 15q24. Am. J. Hum. Genet. 63:1108–1116, 1998.
- [72868] 11198.D'Arcangelo, G.: Personal Communication. Nutley, N. J. 6/2/1995.
- [72869] 11199.D'Arcangelo, G.; Homayouni, R.; Keshvara, L.; Rice, D. S.; Sheldon,M.; Curran, T.: Reelin is a ligand for

lipoprotein receptors. *Neuron* 24:471–479, 1999.

- [72870] 11200.D'Arcangelo, G.; Miao, G. G.; Chen, S.-C.; Soares, H. D.; Morgan, J. I.; Curran, T.: A protein related to extracellular matrix proteins deleted in the mouse mutant reeler. *Nature* 374: 719–723, 1995.
- [72871] 11201.DeSilva, U.; D'Arcangelo, G.; Braden, V. V.; Chen, J.; Miao, G.G.; Curran, T.; Green, E. D.: The human reelin gene: isolation, sequencing, and mapping on chromosome 7. *Genome Res.* 7: 157–164, 1997.
- [72872] 11202.Green, M. C.: Catalog of mutant genes and polymorphic loci. In: Lyon, M. F.; Searle, A. G.: *Genetic Variants and Strains of the Laboratory Mouse*. Oxford: Oxford Univ. Press (pub.) (2nd ed.): 1989.
- [72873] 11203.Hack, I.; Bancila, M.; Loulier, K.; Carroll, P.; Cremer, H.: Reelin is a detachment signal in tangential chain-migration during postnatal neurogenesis. *Nature Neurosci.* 5: 939–945, 2002.
- [72874] 11204.Hirotsune, S.; Takahara, T.; Sasaki, N.; Hirose, K.; Yoshiki, A.; Ohashi, T.; Kusakabe, M.; Murakami, Y.; Muramatsu, M.; Watanabe, S.; Nakao, K.; Katsuki, M.; Hayashizaki, Y.: The reeler gene encodes a protein with an EGF-like motif expressed by pioneer neurons. *Nature-Genet.* 10: 77–83, 1995.



- [72875] 11205.Hong, S. E.; Shugart, Y. Y.; Huang, D. T.; Al Shahwan, S.; Grant,P. E.; Hourihane, J. O.; Martin, N. D. T.; Walsh, C. A.: Autosomalrecessive lissencephaly with cerebellar hypoplasia is associated withhuman RELN mutations. *Nature Genet.* 26: 93–96, 2000. Note: Erratum:*Nature Genet.* 27: 225 only, 2001.
- [72876] 11206.Hourihane, J. O.; Bennett, C. P.; Chaudhuri, R.; Robb, S. A.;Martin, N. D. T.: A sibship with a neuronal migration defect, cerebellarhypoplasia and congenital lymphedema. *Neuropediatrics* 24: 43–46,1993.
- [72877] 11207.Impagnatiello, F.; Guidotti, A. R.; Pesold, C.; Dwivedi, Y.; Caruncho,H.; Pisu, M. G.; Uzunov, D. P.; Smalheiser, N. R.; Davis, J. M.; Pandey,G. N.; Pappas, G. D.; Tueting, P.; Sharma, R. P.; Costa, E.: A decreaseof reelin expression as a putative vulnerability factor in schizophrenia. *Proc.Nat. Acad. Sci.* 95: 15718–15723, 1998.
- [72878] 11208.Magdaleno, S.; Keshvara, L.; Curran, T.: Rescue of ataxia andpreplate splitting by ectopic expression of reelin in reeler mice. *Neuron* 33:573–586, 2002.
- [72879] 11209.Royaux, I.; Lambert de Rouvroit, C.; D'Arcangelo, G.; Demirov,D.; Goffinet, A. M.: Genomic organization of the mouse reelin gene. *Genomics* 46:240–250, 1997.

- [72880] 11210.Yip, J. W.; Yip, Y. P. L.; Nakajima, K.; Capriotti, C.: Reelin controls position of autonomic neurons in the spinal cord. *Proc.Nat. Acad. Sci.* 97: 8612–8616, 2000.
- [72881] 11211.Cattaneo, M.; Lecchi, A.; Randi, A. M.; McGregor, J. L.; Mannucci, P. M.: Identification of a new congenital defect of platelet function characterized by severe impairment of platelet responses to adenosine diphosphate. *Blood* 80: 2787–2796, 1992.
- [72882] 11212.Pawson, T.; Gish, G. D.: SH2 and SH3 domains: from structure to function. *Cell* 71: 359–362, 1992.
- [72883] 11213.Welsh, M.; Mares, J.; Karlsson, T.; Lavergne, C.; Breant, B.; Claesson-Welsh, L.: Shb is a ubiquitously expressed Src homology 2 protein. *Oncogene* 9:19–27, 1994.
- [72884] 11214.Yulug, I. G.; Hillermann, R.; Fisher, E. M. C.: The SHB adaptor protein maps to human chromosome 9. *Genomics* 24: 615–617, 1994.
- [72885] 11215.Bansal-Pakala, P.; Jember, A. G.–H.; Croft, M.: Signaling through OX40 (CD134) breaks peripheral T-cell tolerance. *Nature Med.* 7:907–912, 2001.
- [72886] 11216.Birkeland, M. L.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.A.; Barclay, A. N.: Gene structure and chromosomal localization of the mouse homologue of rat OX40

protein. *Europ. J. Immun.* 25: 926–930, 1995.

[72887] 11217. Jember, A. G.-H.; Zuberi, R.; Liu, F.-T.; Croft, M.: Development of allergic inflammation in a murine model of asthma is dependent on the costimulatory receptor OX40. *J. Exp. Med.* 193: 387–392, 2001.

[72888] 11218. Latza, U.; Durkop, H.; Schnittger, S.; Ringeling, J.; Eitelbach, F.; Hummel, M.; Fonatsch, C.; Stein, H.: The human OX40 homolog: cDNA structure, expression and chromosomal assignment of the ACT35 antigen. *Europ. J. Immun.* 24: 677–683, 1994.

[72889] 11219. Anderson, D. W.; Probst, F. J.; Belyantseva, I. A.; Fridell, R. A.; Beyer, L.; Martin, D. M.; Wu, D.; Kachar, B.; Friedman, T. B.; Raphael, Y.; Camper, S. A.: The motor and tail regions of myosin XV are critical for normal structure and function of auditory and vestibular hair cells. *Hum. Molec. Genet.* 9: 1729–1738, 2000.

[72890] 11220. Probst, F. J.; Fridell, R. A.; Raphael, Y.; Saunders, T. L.; Wang, A.; Liang, Y.; Morell, R. J.; Touchman, J. W.; Lyons, R. H.; Noben-Trauth, K.; Friedman, T. B.; Camper, S. A.: Correction of deafness in shaker-2 mice by an unconventional myosin in a BAC transgene. *Science* 280: 1444–1447, 1998.

[72891] 11221. Wang, A.; Liang, Y.; Fridell, R. A.; Probst, F. J.;

Wilcox, E.R.; Touchman, J. W.; Morton, C. C.; Morell, R. J.; Noben-Trauth, K.; Camper, S. A.; Friedman, T. B.: Association of unconventional myosin MYO15 mutations with human nonsyndromic deafness DFNB3. *Science* 280:1447–1451, 1998.

[72892] 11222. Wilton, S. D.; Lim, L.; Dorosz, S. D.; Gunn, H. C.; Eyre, H. J.; Callen, D. F.; Laing, N. G.: Assignment of the human alpha-tropomyosin gene TPM4 to band 19p13.1 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 72: 294–296, 1996.

[72893] 11223. Lazzaro, M. A.; Picketts, D. J.: Cloning and characterization of the murine Imitation Switch (ISWI) genes: differential expression patterns suggest distinct developmental roles for Snf2h and Snf2l. *J. Neurochem.* 77: 1145–1156, 2001.

[72894] 11224. Okabe, I.; Bailey, L. C.; Attree, O.; Srinivasan, S.; Perkel, J.M.; Laurent, B. C.; Carlson, M.; Nelson, D. L.; Nussbaum, R. L.: Cloning of human and bovine homologs of SNF2/SWI2: a global activator of transcription in yeast *S. cerevisiae*. *Nucleic Acids Res.* 20:4649–4655, 1992.

[72895] 11225. Stahl, W. L.; Eakin, T. J.; Owens, J. W. M., Jr.; Breininger, J.F.; Filuk, P. E.; Anderson, W. R.: Plasma membrane Ca(2+)-ATPase isoforms: distribution of mRNAs in

rat brain by in situ hybridization. *Molec. Brain Res.* 16: 223–231, 1992.

[72896] 11226. De Plaen, E.; Arden, K.; Traversari, C.; Gaforio, J. J.; Szikora, J.-P.; De Smet, C.; Brasseur, F.; van der Bruggen, P.; Lethe, B.; Lurquin, C.; Brasseur, R.; Chomez, P.; De Backer, O.; Cavenee, W.; Boon, T.: Structure, chromosomal localization, and expression of 12 genes of the MAGE family. *Immunogenetics* 40: 360–369, 1994.

[72897] 11227. Rogner, U. C.; Wilke, K.; Steck, E.; Korn, B.; Poustka, A.: The melanoma antigen gene (MAGE) family is clustered in the chromosomal band Xq28. *Genomics* 29: 725–731, 1995.

[72898] 11228. Chakarova, C.; Wehnert, M. S.; Uhl, K.; Sakthivel, S.; Vosberg, H.-P.; van der Ven, P. F. M.; Furst, D. O.: Genomic structure and fine mapping of the two human filamin gene paralogues FLNB and FLNC and comparative analysis of the filamin gene family. *Hum. Genet.* 107: 597–611, 2000.

[72899] 11229. Fink, J. M.; Dobyns, W. B.; Guerrini, R.; Hirsch, B. A.: Identification of a duplication of Xq28 associated with bilateral periventricular nodular heterotopia. *Am. J. Hum. Genet.* 61: 379–387, 1997.

[72900] 11230. Fox, J. W.; Lamperti, E. D.; Eksioglu, Y. Z.; Hong, S. E.; Feng, Y.; Graham, D. A.; Scheffer, I. E.; Dobyns, W. B.;

Hirsch, B. A.;Radtke, R. A.; Berkovic, S. F.; Huttenlocher, P. R.; Walsh, C. A.: Mutations in filamin 1 prevent migration of cerebral cortical neuronsin human periventricular heterotopia. *Neuron* 21: 1315–1325, 1998.

[72901] 11231.Gariboldi, M.; Maestrini, E.; Canzian, F.; Manenti, G.; De Gregorio,L.; Rivella, S.; Chatterjee, A.; Herman, G. E.; Archidiacono, N.;Antonacci, R.; Pierotti, M. A.; Dragani, T. A.; Toniolo, D.: Comparativemapping of the actin-binding protein 280 genes in human and mouse. *Genomics* 21:428–430, 1994.

[72902] 11232.Bashir, R.; Britton, S.; Strachan, T.; Keers, S.; Vafiadaki, E.;Lako, M.; Richard, I.; Marchand, S.; Bourg, N.; Argov, Z.; Sadeh,M.; Mahjneh, I.; Marconi, G.; Passos-Bueno, M. R.; Moreira, E. S.;Zatz, M.; Beckmann, J. S.; Bushby, K.: A gene related to *Caenorhabditiselegans* spermatogenesis factor *fer-1* is mutated in limb-girdle muscular dystrophy type 2B. *Nature Genet.* 20: 37–42, 1998.

[72903] 11233.Liu, J.; Aoki, M.; Illa, I.; Wu, C.; Fardeau, M.; Angelini, C.;Serrano, C.; Urtizberea, J. A.; Hentati, F.; Ben Hamida, M.; Bohlega,S.; Culper, E. J.; Amato, A. A.; Bossie, K.; Oeltjen, J.; Bejaoui,K.; McKenna-Yasek, D.; Hosler, B. A.; Schurr, E.; Arahata, K.; deJong, P. J.; Brown, R. H., Jr.: Dysferlin, a novel skeletal musclegene, is mutated in

Miyoshi myopathy and limb girdle muscular dystrophy.

NatureGenet. 20: 31–36, 1998.

- [72904] 11234.Aghai, E.; Yaniv, I.; David, M.: Factor XI deficiency in an ArabMoslem family in Israel. Scand. J. Haemat. 32: 327–331, 1984.
- [72905] 11235.Asakai, R.; Chung, D. W.; Davie, E. W.; Seligsohn, U.: FactorXI deficiency in Ashkenazi Jews in Israel. New Eng. J. Med. 325:153–158, 1991.
- [72906] 11236.Asakai, R.; Chung, D. W.; Ratnoff, O. D.; Davie, E. W.: FactorXI (plasma thromboplastin antecedent) deficiency in Ashkenazi Jewsis a bleeding disorder that can result from three types of point mutations. Proc.Nat. Acad. Sci. 86: 7667–7671, 1989.
- [72907] 11237.Asakai, R.; Davie, E. W.; Chung, D. W.: Organization of the gene for human factor XI. Biochemistry 26: 7221–7228, 1987.
- [72908] 11238.Bauduer, F.; Dupreuilh, F.; Ducout, L.; Marti, B.: Factor XI deficiency in the French Basque country. Haemophilia 5: 187–190, 1999.
- [72909] 11239.Bertina, R. M.; Veltkamp, J. J.: A genetic variant of factor XI with decreased capacity for Ca ion binding. Brit. J. Haemat. 42:623–635, 1979.
- [72910] 11240.Biggs, R.; MacFarlane, R. G.: Human Blood Coagu–

lation and Its Disorders. Oxford: Blackwell (pub.) (3rd ed.): 1962.

[72911] 11241. Bolton-Maggs, P. H. B.: Factor XI deficiency. *Bailliere's Clin. Haemat.* 9: 355–368, 1996.

[72912] 11242. Braunstein, K. M.; Noyes, C. M.; Griffith, M. J.; Lundblad, R. L.; Roberts, H. R.: Characterization of the defect in activation of factor IX by human factor XIa. *J. Clin. Invest.* 68:1420–1426, 1981.

[72913] 11243. Buetow, K. H.; Shiang, R.; Yang, P.; Nakamura, Y.; Lathrop, G. M.; White, R.; Wasmuth, J. J.; Wood, S.; Berdahl, L. D.; Leysens, N. J.: A detailed multipoint map of human chromosome 4 provides evidence for linkage heterogeneity and position-specific recombination rates. *Am. J. Hum. Genet.* 48: 911–925, 1991.

[72914] 11244. Clarkson, K.; Rosenfeld, B.; Fair, J.; Klein, A.; Bell, W.: Factor XI deficiency acquired by liver transplantation. *Ann. Intern. Med.* 115:877–879, 1991.

[72915] 11245. Dodds, W. J.; Kull, J. E.: Canine factor XI (plasma thromboplastin antecedent) deficiency. *J. Lab. Clin. Med.* 78: 746–752, 1971.

[72916] 11246. Dzik, W. H.; Arkin, C. F.; Jenkins, R. L.: Transfer of congenital factor XI deficiency from a donor to a recipient by liver transplantation. (Letter) *New Eng. J. Med.* 316:



1217–1218, 1987.

[72917] 11247.Fujikawa, K.; Chung, D. W.; Hendrickson, L. E.; Davie, E. W.:Amino acid sequence of human factor XI, a blood coagulation factorwith four tandem repeats that are highly homologous with plasma prekallikrein. *Biochemistry* 25:2417–2424, 1986.

[72918] 11248.Imanaka, Y.; Lal, K.; Nishimura, T.; Bolton–Maggs, P. H. B.; Tuddenham,E. G. D.; McVey, J. H.: Identification of two novel mutations in non–Jewish factor XI deficiency. *Brit. J. Haemat.* 90: 916–920, 1995.

[72919] 11249.Kato, A.; Asakai, R.; Davie, E. W.; Aoki, N.: Factor XI gene(F11) is located on the distal end of the long arm of human chromosome4. *Cytogenet. Cell Genet.* 52: 77–78, 1989.

[72920] 11250.Litz, C. E.; Swaim, W. R.; Dalmaso, A. P.: Factor XI deficiency:genetic and clinical studies of a single kindred. *Am. J. Hemat.* 28:8–12, 1988.

[72921] 11251.Mannhalter, C.; Hellstern, P.; Deutsch, E.: Identification of a defective factor XI cross–reacting material in a factor XI–deficient patient. *Blood* 70: 31–37, 1987.

[72922] 11252.Meijers, J. C. M.; Davie, E. W.; Chung, D. W.: Expression of human blood coagulation factor XI: characterization of the defective factor XI type III deficiency. *Blood*

79: 1435–1440, 1992.

- [72923] 11253. Mitchell, M.; Cutler, J.; Thompson, S.; Moore, G.; Rees, E. J.; Smith, M.; Savidge, G.; Alhaq, A.: Heterozygous factor XI deficiency associated with three novel mutations. *Brit. J. Haemat.* 107: 763–765, 1999.
- [72924] 11254. Muir, W. A.; Ratnoff, O. D.: The prevalence of plasma thromboplastin antecedent (PTA factor XI) deficiency. *Blood* 44: 569–570, 1974.
- [72925] 11255. Niskanen, E. O.; Saito, H.; Cline, M. J.: Plasma thromboplastin antecedent (factor XI) deficiency in a black family. *Arch. Intern. Med.* 141: 936–941, 1981.
- [72926] 11256. Peretz, U.; Zivelin, A.; Usher, S.; Eichel, R.; Seligsohn, U.: Identification of a new mutation in the factor XI gene of an Ashkenazi-Jew with severe factor XI deficiency. (Abstract) *Blood* 82 (suppl. 1):66a only, 1993.
- [72927] 11257. Pugh, R. E.; McVey, J. H.; Tuddenham, E. G. D.; Hancock, J. F.: Six point mutations that cause factor XI deficiency. *Blood* 85:1509–1516, 1995.
- [72928] 11258. Ragni, M. V.; Sinha, D.; Seaman, F.; Lewis, J. H.; Spero, J. A.; Walsh, P. N.: Comparison of bleeding tendency, factor XI coagulant activity, and factor XI antigen in 25 factor XI-deficient kindreds. *Blood* 65:719–724, 1985.
- [72929] 11259. Rapaport, S. I.; Proctor, R. R.; Patch, M. J.; Yettra,

M.: The mode of inheritance of PTA deficiency: evidence for the existence of major PTA deficiency and minor PTA deficiency. *Blood* 18: 149–165, 1961.

[72930] 11260. Rimon, A.; Schiffman, S.; Feinstein, D. I.; Rapaport, S. I.: Factor–XI activity and factor–XI antigen in homozygous and heterozygous factor–XI deficiency. *Blood* 48: 165–174, 1976.

[72931] 11261. Foresti, V.; Ferrari, C.: Central hypothyroidism: isolated thyrotropin–releasing hormone deficiency or resistance of pituitary thyrotropes? (Letter) *J. Endocr. Invest.* 8: 577 only, 1985.

[72932] 11262. Katakami, H.; Kato, Y.; Inada, M.; Imura, H.: Hypothalamic hypothyroidism due to isolated thyrotropin–releasing hormone (TRH) deficiency. *J. Endocr. Invest.* 7: 231–233, 1984.

[72933] 11263. Niimi, H.; Inomata, H.; Sasaki, N.; Nakajima, H.: Congenital isolated thyrotrophin releasing hormone deficiency. *Arch. Dis. Child.* 57: 877–878, 1982.

[72934] 11264. Roller, M. L.; Camper, S. A.: Localization of the thyrotropin–releasing hormone gene, *Trh*, on mouse chromosome 6. *Mammalian Genome* 6: 443–444, 1995.

[72935] 11265. Yamada, M.; Radovick, S.; Wondisford, F. E.; Nakayama, Y.; Weintraub, B. D.; Wilber, J. F.: Cloning and

structure of human genomic DNA and hypothalamic cDNA encoding human preprothyrotropin-releasing hormone. *Molec. Endocr.* 4: 551–556, 1990.

- [72936] 11266. Yamada, M.; Satoh, T.; Monden, T.; Mori, M.: Assignment of the thyrotropin-releasing hormone gene (TRH) to human chromosome 3q13.3–q21 by in situ hybridization. *Cytogenet. Cell Genet.* 87: 275 only, 1999.
- [72937] 11267. Yamada, M.; Wondisford, F. E.; Radovick, S.; Nakayama, Y.; Weintraub, B. D.; Wilber, J. F.: Assignment of human preprothyrotropin-releasing hormone (TRH) gene to chromosome 3. *Somat. Cell Molec. Genet.* 17:97–100, 1991.
- [72938] 11268. McDermott, J. C.; Cardoso, M. C.; Yu, Y.-T.; Andres, V.; Leifer, D.; Krainc, D.; Lipton, S. A.; Nadal-Ginard, B.: hMEF2C gene encodes skeletal muscle- and brain-specific transcription factors. *Molec. Cell. Biol.* 13: 2564–2577, 1993.
- [72939] 11269. Anderson, G. J.; Murphy, T. L.; Cowley, L.; Evans, B. A.; Halliday, J. W.; McLaren, G. D.: Mapping the gene for sex-linked anemia: an inherited defect of intestinal iron absorption in the mouse. *Genomics* 48:34–39, 1998.
- [72940] 11270. Bannerman, R. M.: Genetic defects of iron transport. *Fed. Proc.* 35:2281–2285, 1976.

- [72941] 11271. Edwards, J. A.; Hoke, J. E.; Mattioli, M.; Reichlin, M.: Ferritin distribution and synthesis in sex-linked anemia. *J. Lab. Clin. Med.* 90:68–76, 1977.
- [72942] 11272. Falconer, D. S.; Isaacson, J. H.: The genetics of sex-linked anaemia in the mouse. *Genet. Res.* 3: 248–250, 1962.
- [72943] 11273. Grewal, M. S.: A sex-linked anaemia in the mouse. *Genet. Res.* 3:238–247, 1962.
- [72944] 11274. Lee, G. R.; Nacht, S.; Lukens, J. N.; Cartwright, G. E.: Iron metabolism in copper-deficient swine. *J. Clin. Invest.* 47: 2058–2069, 1968.
- [72945] 11275. Vulpe, C. D.; Kuo, Y.-M.; Murphy, T. L.; Cowley, L.; Askwith, C.; Libina, N.; Gitschier, J.; Anderson, G. J.: Hephaestin, a ceruloplasmin homologue implicated in intestinal iron transport, is defective in the sla mouse. *Nature Genet.* 21: 195–199, 1999.
- [72946] 11276. Karumanchi, S. A.; Jha, V.; Ramchandran, R.; Karihaloo, A.; Tsiokas, L.; Chan, B.; Dhanabal, M.; Hanai, J.; Venkataraman, G.; Shriver, Z.; Keiser, N.; Kalluri, R.; and 9 others: Cell surface glypicans are low-affinity endostatin receptors. *Molec. Cell* 7: 811–822, 2001.
- [72947] 11277. Veugelers, M.; Vermeesch, J.; Watanabe, K.; Yamaguchi, Y.; Marynen, P.; David, G.: GPC4, the gene for hu-

man K-glypican, flanks GPC3 onXq26: deletion of the GPC3-GPC4 gene cluster in one family with Simpson-Golabi-Behmelsyndrome. *Genomics* 53: 1-11, 1998.

- [72948] 11278.Emes, R. D.; Ponting, C. P.: A new sequence motif linking lissencephaly,Treacher Collins and oral-facial-digital type 1 syndromes, microtubuledynamics and cell migration. *Hum. Molec. Genet.* 10: 2813-2820, 2001.
- [72949] 11279.Copley, L. M.; Zhao, W. D.; Kopacz, K.; Herman, G. E.; Kioschis,P.; Poustka, A.; Taudien, S.; Platzer, M.: Exclusion of mutationsin the MTMR1 gene as a frequent cause of X-linked myotubular myopathy.(Letter) *Am. J. Med. Genet.* 107: 256-258, 2002.
- [72950] 11280.Kioschis, P.; Wiemann, S.; Heiss, N. S.; Francis, F.; Gotz, C.;Poustka, A.; Taudien, S.; Platzer, M.; Wiehe, T.; Beckmann, G.; Weber,J.; Nordsiek, G.; Rosenthal, A.: Genomic organization of a 225-kbregion in Xq28 containing the gene for X-linked myotubular myopathy(MTM1) and a related gene (MTMR1). *Genomics* 54: 256-266, 1998.
- [72951] 11281.Laporte, J.; Blondeau, F.; Buj-Bello, A.; Tentler, D.; Kretz, C.;Dahl, N.; Mandel, J.-L.: Characterization of the myotubularin dualspecificity phosphatase gene family from yeast to human. *Hum. Molec.Genet.* 7: 1703-1712, 1998.

- [72952] 11282.Gorlin, J. B.; Henske, E.; Warren, S. T.; Kunst, C. B.; D'Urso, M.; Palmieri, G.; Hartwig, J. H.; Bruns, G.; Kwiatkowski, D. J.: Actin-binding protein (ABP-280) filamin gene (FLN) maps telomeric to the color vision locus (R/GCP) and centromeric to G6PD in Xq28. *Genomics* 17:496-498, 1993.
- [72953] 11283.Jager, R. J.; Anvret, M.; Hall, K.; Scherer, G.: A human XY female with a frame shift mutation in the candidate testis-determining gene SRY. *Nature* 348: 452-454, 1990.
- [72954] 11284.Koopman, P.; Gubbay, J.; Vivian, N.; Goodfellow, P.; Lovell-Badge, R.: Male development of chromosomally female mice transgenic for Sry. *Nature* 351: 117-121, 1991.
- [72955] 11285.Koopman, P.; Munsterberg, A.; Capel, B.; Vivian, N.; Lovell-Badge, R.: Expression of a candidate sex-determining gene during mouse testis differentiation. *Nature* 348: 450-452, 1990.
- [72956] 11286.McElreavey, K. D.; Vilain, E.; Boucek, C.; Vidaud, M.; Jaubert, F.; Richaud, F.; Fellous, M.: XY sex reversal associated with a nonsense mutation in SRY. *Genomics* 13: 838-840, 1992.
- [72957] 11287.Page, D. C.; Mosher, R.; Simpson, E. M.; Fisher, E. M. C.; Mardon, G.; Pollack, J.; McGillivray, B.; de la

Chapelle, A.; Brown, L. G.: The sex-determining region of the human Y chromosome encodes a fingerprotein. *Cell* 51: 1091–1104, 1987.

[72958] 11288. Taylor, H.; Barter, R. H.; Jacobson, C. B.: Neoplasms of dysgenetic gonads. *Am. J. Obstet. Gynec.* 96: 816–823, 1966.

[72959] 11289. Vilain, E.; McElreavey, K.; Jaubert, F.; Raymond, J.-P.; Richaud, F.; Fellous, M.: Familial case with sequence variant in the testis-determining region associated with two sex phenotypes. *Am. J. Hum. Genet.* 50:1008–1011, 1992.

[72960] 11290. Gorlin, J. B.; Yamin, R.; Egan, S.; Stewart, M.; Stos-  
sel, T. P.; Kwiatkowski, D. J.; Hartwig, J. H.: Human en-  
dothelial actin-binding protein (ABP-280, nonmuscle fil-  
amin): a molecular leaf spring. *J. Cell Biol.* 111:  
1089–1105, 1990.

[72961] 11291. Huttenlocher, P. R.; Taravath, S.; Mojtahedi, S.:  
Periventricular heterotopia and epilepsy. *Neurology* 44:  
51–55, 1994.

[72962] 11292. Kunst, C. B.; Henske, E.; Hartwig, J. H.;  
Kwiatkowski, D. J.; D'Urso, M.; Bruns, G.; Warren, S. T.;  
Gorlin, J. B.: The dystrophin-like actin binding protein 280  
gene maps between DXS52 and G6PD overlapping the



Emery–Dreifuss muscular dystrophy locus. (Abstract) Am. J. Hum. Genet. 51: A21, 1992.

[72963] 11293. Maestrini, E.; Patrosso, C.; Mancini, M.; Rivella, S.; Rocchi, M.; Repetto, M.; Villa, A.; Frattini, A.; Zoppe, M.; Vezzoni, P.; Toniolo, D.: Mapping of two genes encoding isoforms of the actin-binding protein ABP-280, a dystrophin like protein, to Xq28 and to chromosome 7. Hum. Molec. Genet. 2: 761–766, 1993.

[72964] 11294. Maestrini, E.; Rivella, S.; Tribioli, C.; Purtilo, D.; Rocchi, M.; Archidiacono, N.; Toniolo, D.: Probes for CpG islands on the distal long arm of the human X chromosome are clustered in Xq24 and Xq28. Genomics 8: 664–670, 1990.

[72965] 11295. Patrosso, M. C.; Repetto, M.; Villa, A.; Milanesi, L.; Frattini, A.; Faranda, S.; Mancini, M.; Maestrini, E.; Toniolo, D.; Vezzoni, P.: The exon–intron organization of the human X-linked gene (FLN1) encoding actin-binding protein 280. Genomics 21: 71–76, 1994.

[72966] 11296. Sheen, V. L.; Dixon, P. H.; Fox, J. W.; Hong, S. E.; Kinton, L.; Sisodiya, S. M.; Duncan, J. S.; Dubeau, F.; Schaffer, I. E.; Schachter, S. C.; Wilner, A.; Henchy, R.; and 18 others: Mutations in the X-linked filamin 1 gene cause periventricular nodular heterotopia in males as well as in

females. Hum. Molec. Genet. 10: 1775–1783,2001.

[72967] 11297.Small, K.; Wagener, M.; Warren, S. T.: Isolation and characterization of the complete mouse emerin gene.

Mammalian Genome 8: 337–341,1997.

[72968] 11298.Vadlamudi, R. K.; Li, F.; Adam, L.; Nguyen, D.; Ohta, Y.; Stossel, T. P.; Kumar, R.: Filamin is essential in actin cytoskeletal assembly mediated by p21-activated kinase 1. Nature Cell Biol. 4: 681–690,2002.

[72969] 11299.Faranda, S.; Frattini, A.; Vezzoni, P.: The human genes encoding renin-binding protein and host cell factor are closely linked in Xq28 and transcribed in the same direction. Gene 155: 237–239, 1995.

[72970] 11300.Frattini, A.; Chatterjee, A.; Faranda, S.; Sacco, M. G.; Villa, A.; Herman, G. E.; Vezzoni, P.: The chromosome localization and the HCF repeats of the human host cell factor gene (HCFC1) are conserved in the mouse homologue. Genomics 32: 277–280, 1996.

[72971] 11301.Frattini, A.; Faranda, S.; Redolfi, E.; Zucchi, I.; Villa, A.; Patrosso, M. C.; Strina, D.; Susani, L.; Vezzoni, P.: Genomic organization of the human VP16 accessory protein, a housekeeping gene (HCFC1) mapping to Xq28. Genomics 23: 30–35, 1994.

[72972] 11302.Wilson, A. C.; LaMarco, K.; Peterson, M. G.; Herr,

W.: The VP16accessory protein HCF is a family of polypeptides processed from a large precursor protein. *Cell* 74: 115–125, 1993.

[72973] 11303. Wilson, A. C.; Parrish, J. E.; Massa, H. F.; Nelson, D. L.; Trask, B. J.; Herr, W.: The gene encoding the VP16–accessory protein HCF(HCFC1) resides in human Xq28 and is highly expressed in fetal tissues and the adult kidney. *Genomics* 25: 462–468, 1995.

[72974] 11304. Zoppe, M.; Frattini, A.; Faranda, S.; Vezzoni, P.: The complete sequence of the host cell factor 1 (HCFC1) gene and its promoter: a role for YY1 transcription factor in the regulation of its expression. *Genomics* 34: 85–91, 1996.

[72975] 11305. Maestrini, E.; Tamagnone, L.; Longati, P.; Cremona, O.; Gulisano, M.; Bione, S.; Tamanini, F.; Neel, B. G.; Toniolo, D.; Comoglio, P. M.: A family of transmembrane proteins with homology to the MET–hepatocyte growth factor receptor. *Proc. Nat. Acad. Sci.* 93: 674–678, 1996.

[72976] 11306. Tamagnone, L.; Artigiani, S.; Chen, H.; He, Z.; Ming, G.; Song, H.; Chedotal, A.; Winberg, M. L.; Goodman, C. S.; Poo, M.; Tessier–Lavigne, M.; Comoglio, P. M.: Plexins are a large family of receptors for transmembrane, secreted, and GPI–anchored semaphorins in vertebrates. *Cell* 99: 71–80, 1999.

- [72977] 11307.Becker, P. E.: Two new families of benign sex-linked recessivemuscular dystrophy. *Rev. Canad. Biol.* 21: 551–566, 1962.
- [72978] 11308.Becker, P. E.: Eine neue X-chromosomale Muskeldystrophie. *ActaPsychiat. Neurol. Scand.* 193: 427, 1955.
- [72979] 11309.Becker, P. E.: Neue Ergebnisse der Genetik der Muskeldystrophien. *ActaGenet. Statist. Med.* 7: 303–310, 1957.
- [72980] 11310.Bushby, K. M. D.; Cleghorn, N. J.; Curtis, A.; Haggerty, I. D.;Nicholson, L. V. B.; Johnson, M. A.; Harris, J. B.; Bhattacharya,S. S.: Identification of a mutation in the promoter region of thedystrophin gene in a patient with atypical Becker muscular dystrophy. *Hum.Genet.* 88: 195–199, 1991.
- [72981] 11311.Doriguzzi, C.; Palmucci, L.; Mongini, T.; Chiado-Piat, L.; Restagno,G.; Ferrone, M.: Exercise intolerance and recurrent myoglobinuriaas the only expression of Xp21 Becker type muscular dystrophy. *J.Neurol.* 240: 269–271, 1993.
- [72982] 11312.England, S. B.; Nicholson, L. V. B.; Johnson, M. A.; Forrest,S. M.; Love, D. R.; Zubrzycka-Gaarn, E. E.; Bulman, D. E.; Harris,J. B.; Davies, K. E.: Very mild muscular dystro–

phy associated with the deletion of 46% dystrophin. *Nature* 343: 180–182, 1990.

[72983] 11313. Derry, J. M. J.; Barnard, P. J.: Physical linkage of the A-raf-1, properdin, synapsin I, and TIMP genes on the human and mouse X chromosomes. *Genomics* 12:632–638, 1992.

[72984] 11314. Aronsson, F. C.; Magnusson, P.; Andersson, B.; Karsten, S. L.; Shibasaki, Y.; Lendon, C. L.; Goate, A. M.; Brookes, A. J: The NIK protein kinase and C17orf1 genes: chromosomal mapping, gene structures and mutational screening in frontotemporal dementia and parkinsonism-linked to chromosome 17. *Hum. Genet.* 103: 340–345, 1998.

[72985] 11315. Malinin, N. L.; Boldin, M. P.; Kovalenko, A. V.; Wallach, D.: MAP3K-related kinase involved in NF- $\kappa$ B induction by TNF, CD95 and IL-1. *Nature* 385: 540–544, 1997.

[72986] 11316. Shinkura, R.; Kitada, K.; Matsuda, F.; Tashiro, K.; Ikuta, K.; Suzuki, M.; Kogishi, K.; Serikawa, T.; Honjo, T.: Alymphoplasia is caused by a point mutation in the mouse gene encoding Nf- $\kappa$ B-inducing kinase. *Nature Genet.* 22: 74–77, 1999.

[72987] 11317. Smith, C.; Andreakos, E.; Crawley, J. B.; Brennan, F.

M.; Feldmann, M.; Foxwell, B. M. J.: NF-kappa-B-inducing kinase is dispensable for activation of NF-kappa-B in inflammatory settings but essential for lymphotoxin beta receptor activation of NF-kappa-B in primary human fibroblasts. *J. Immun.* 167: 5895–5903, 2001.

[72988] 11318. Yin, L.; Wu, L.; Wesche, H.; Arthur, C. D.; White, J. M.; Goeddel, D. V.; Schreiber, R. D.: Defective lymphotoxin-beta receptor-induced NF-kappa-B transcriptional activity in NIK-deficient mice. *Science* 291: 2162–2165, 2001.

[72989] 11319. Aronsson, F. C.; Magnusson, P.; Andersson, B.; Karsten, S. L.; Shibasaki, Y.; Lendon, C. L.; Goate, A. M.; Brookes, A. J.: The NIK protein kinase and C17orf1 genes: chromosomal mapping, gene structures and mutational screening in frontotemporal dementia and parkinsonism-linked to chromosome 17. *Hum. Genet.* 103: 340–345, 1998.

[72990] 11320. Yayoshi-Yamamoto, S.; Taniuchi, I.; Watanabe, T.: FRL, a novel formin-related protein, binds to Rac and regulates cell motility and survival of macrophages. *Molec. Cell. Biol.* 20: 6872–6881, 2000.

[72991] 11321. Gress, T. M.; Muller-Pillasch, F.; Geng, M.; Zimmerhackl, F.; Zehetner, G.; Friess, H.; Buchler, M.; Adler, G.;

Lehrach, H.: A pancreaticcancer-specific expression profile. *Oncogene* 13: 1819–1830, 1996.

[72992] 11322.Muller-Pillasch, F.; Wallrapp, C.; Lacher, U.; Friess, H.; Buchler,M.; Adler, G.; Gress, T. M.: Identification of a new tumour-associatedantigen TM4SF5 and its expression in human cancer. *Gene* 208: 25–30,1998.

[72993] 11323.Borggreffe, T.; Masat, L.; Wabl, M.; Riwar, B.; Cattoretti, G.;Jessberger, R.: Cellular, intracellular, and developmental expressionpatterns of murine SWAP-70. *Europ. J. Immun.* 29: 1812–1822, 1999.

[72994] 11324.Borggreffe, T.; Wabl, M.; Akhmedov, A. T.; Jessberger, R.: A B-cell-specificDNA recombination complex. *J. Biol. Chem.* 273: 17025–17035, 1998.

[72995] 11325.Masat, L.; Caldwell, J.; Armstrong, R.; Khoshnevisan, H.; Jessberger,R.; Herndier, B.; Wabl, M.; Ferrick, D.: Association of SWAP-70 withthe B cell antigen receptor complex. *Proc. Nat. Acad. Sci.* 97: 2180–2184,2000.

[72996] 11326.Masat, L.; Liddell, R. A.; Mock, B. A.; Kuo, W.-L.; Jessberger,R.; Wabl, M.; Morse, H. C., III: Mapping of the SWAP70 gene to mousechromosome 7 and human chromosome 11p15. *Immunogenetics* 51: 16–19,2000.

[72997] 11327.Shinohara, M.; Terada, Y.; Iwamatsu, A.; Shinohara, A.; Mochizuki,N.; Higuchi, M.; Gotoh, Y.; Ihara, S.; Nagata,

S.; Itoh, H.; Fukui,Y.; Jessberger, R.: SWAP-70 is a guanine-nucleotide-exchange factor that mediates signalling of membrane ruffling. *Nature* 416: 759-763,2002.

[72998] 11328.Miura, K.; Suzuki, K.; Tokino, T.; Isomura, M.; Inazawa, J.; Matsuno,S.; Nakamura, Y.: Detailed deletion mapping in squamous cell carcinomas of the esophagus narrows a region containing a putative tumor suppressor-gene to about 200 kilobases on distal chromosome 9q. *Cancer Res.* 56:1629-1634, 1996.

[72999] 11329.Nishiwaki, T.; Daigo, Y.; Kawasoe, T.; Nakamura, Y.: Isolation and mutational analysis of a novel human cDNA, DEC1 (deleted in esophageal cancer 1), derived from the tumor suppressor locus in 9q32. *Genes Chromosomes Cancer* 27: 169-176, 2000.

[73000] 11330.Nemoto, Y.; Yamamoto, T.; Takada, S.; Matsui, Y.; Obinata, M.:Antisense RNA of the latent period gene (MER5) inhibits the differentiation of murine erythroleukemia cells. *Gene* 91: 261-265, 1990.

[73001] 11331.Shih, S.-F.; Wu, Y.-H.; Hung, C.-H.; Yang, H.-Y.; Lin, J.-Y.:Abrin triggers cell death by inactivating a thiol-specific antioxidant protein. *J. Biol. Chem.* 276: 21870-21877, 2001.

[73002] 11332.Tsuji, K.; Copeland, N. G.; Jenkins, N. A.; Obinata,



M.: Mammalian antioxidant protein complements alkylhydroperoxide reductase (ahpC) mutation in *Escherichia coli*. *Biochem. J.* 307: 377–381, 1995.

- [73003] 11333. Wonsey, D. R.; Zeller, K. I.; Dang, C. V.: The c-Myc target gene PRDX3 is required for mitochondrial homeostasis and neoplastic transformation. *Proc. Nat. Acad. Sci.* 99: 6649–6654, 2002.
- [73004] 11334. Kamimoto, T.; Zama, T.; Aoki, R.; Muro, Y.; Hagiwara, M.: Identification of a novel kinesin-related protein, KRMP1, as a target for mitotic peptidyl-prolyl isomerase Pin1. *J. Biol. Chem.* 276: 37520–37528, 2001.
- [73005] 11335. Akagi, T.; Motegi, M.; Tamura, A.; Suzuki, R.; Hosokawa, Y.; Suzuki, H.; Ota, H.; Nakamura, S.; Morishima, Y.; Taniwaki, M.; Seto, M.: A novel gene, MALT1 at 18q21, is involved in t(11;18) (q21;q21) found in low-grade B-cell lymphoma of mucosa-associated lymphoid tissue. *Oncogene* 18: 5785–5794, 1999.
- [73006] 11336. Dierlamm, J.; Baens, M.; Wlodarska, I.; Stefanova-Ouzounova, M.; Hernandez, J. M.; Hossfeld, D. K.; De Wolf-Peeters, C.; Hagemeijer, A.; Van den Berghe, H.; Marynen, P.: The apoptosis inhibitor gene API2 and a novel 18q gene, MLT, are recurrently rearranged in the t(11;18)(q21;q21) associated with mucosa-associated

lymphoid tissue lymphomas. *Blood* 93: 3601–3609, 1999.

- [73007] 11337. Ott, G.; Katzenberger, T.; Greiner, A.; Kalla, J.; Rosenwald, A.; Heinrich, U.; Ott, M. M.; Muller-Hermelink, H. K.: The t(11;18)(q21;q21) chromosome translocation is a frequent and specific aberration in low-grade but not high-grade malignant non-Hodgkin's lymphomas of the mucosa-associated lymphoid tissue (MALT-) type. *Cancer Res.* 57:3944–3948, 1997.
- [73008] 11338. Sato, Y.; Akiyama, Y.; Tanizawa, T.; Shibata, T.; Saito, K.; Mori, S.; Kamiyama, R.; Yuasa, Y.: Molecular characterization of the genomic breakpoint junction in the t(11;18)(q21;q21) translocation of a gastric MALT lymphoma. *Biochem. Biophys. Res. Commun.* 280: 301–306, 2001.
- [73009] 11339. Valladeau, J.; Clair-Moninot, V.; Dezutter-Dambuyant, C.; Pin, J.-J.; Kissenpfennig, A.; Mattei, M.-G.; Ait-Yahia, S.; Bates, E. E. M.; Malissen, B.; Koch, F.; Fossiez, F.; Romani, N.; Lebecque, S.; Saeland, S.: Identification of mouse langerin/CD207 in Langerhans cells and some dendritic cells of lymphoid tissues. *J. Immun.* 168:782–792, 2002.
- [73010] 11340. Valladeau, J.; Ravel, O.; Dezutter-Dambuyant, C.; Moore, K.; Kleijmeer, M.; Liu, Y.; Duvert-Frances, V.; Vin-

cent, C.; Schmitt, D.; Davoust, J.; Caux, C.; Lebecque, S.; Saeland, S.: Langerin, a novel C-type lectin specific to Langerhans cells, is an endocytic receptor that induces the formation of Birbeck granules. *Immunity* 12: 71–81, 2000.

[73011] 11341. Reichenberger, E.; Tiziani, V.; Watanabe, S.; Park, L.; Ueki, Y.; Santanna, C.; Baur, S. T.; Shiang, R.; Grange, D. K.; Beighton, P.; Gardner, J.; Hamersma, H.; Sellars, S.; Ramesar, R.; Lidral, A. C.; Sommer, A.; Raposo do Amaral, C. M.; Gorlin, R. J.; Mulliken, J. B.; Olsen, B. R.: Autosomal dominant craniometaphyseal dysplasia is caused by mutations in the transmembrane protein ANK. *Am. J. Hum. Genet.* 68:1321–1326, 2001.

[73012] 11342. Ansel, K. M.; Harris, R. B. S.; Cyster, J. G.: CXCL13 is required for B1 cell homing, natural antibody production, and body cavity immunity. *Immunity* 16:67–76, 2002.

[73013] 11343. Gunn, M. D.; Ngo, V. N.; Ansel, K. M.; Ekland, E. H.; Cyster, J. G.; Williams, L. T.: A B-cell-homing chemokine made in lymphoid follicles activates Burkitt's lymphoma receptor-1. *Nature* 391: 799–803, 1998.

[73014] 11344. Legler, D. F.; Loetscher, M.; Roos, R. S.; Clark-Lewis, I.; Baggiolini, M.; Moser, B.: B cell-attracting chemokine 1, a human CXC chemokine expressed in lym-

phoid tissues, selectively attracts B lymphocytes via BLR1/CXCR5. *J. Exp. Med.* 187: 655–660, 1998.

[73015] 11345. Derst, C.; Engel, H.; Grzeschik, K.-H.; Daut, J.: Genomic structure and chromosome mapping of human and mouse RAMP genes. *Cytogenet. Cell Genet.* 90: 115–118, 2000.

[73016] 11346. Amler, L. C.; Bauer, A.; Corvi, R.; Dihlmann, S.; Praml, C.; Cavenee, W. K.; Schwab, M.; Hampton, G. M.: Identification and characterization of novel genes located at the t(1;15)(p36.2;q24) translocation breakpoint in the neuroblastoma cell line NGP. *Genomics* 64: 195–202, 2000.

[73017] 11347. Yanagisawa, H.; Bundo, M.; Miyashita, T.; Okamura-Oho, Y.; Tadokoro, K.; Tokunaga, K.; Yamada, M.: Protein binding of a DRPLA family through arginine–glutamic acid dipeptide repeats is enhanced by extended polyglutamine. *Hum. Molec. Genet.* 9: 1433–1442, 2000.

[73018] 11348. Oh, J.; Takahashi, R.; Kondo, S.; Mizoguchi, A.; Adachi, E.; Sasahara, R. M.; Nishimura, S.; Imamura, Y.; Kitayama, H.; Alexander, D. B.; Ide, C.; Horan, T. P.; Arakawa, T.; Yoshida, H.; Nishikawa, S.; Itoh, Y.; Seiki, M.; Itohara, S.; Takahashi, C.; Noda, M.: The membrane-anchored MMP inhibitor RECK is a key regulator of extracellular matrix

integrity and angiogenesis. *Cell* 107: 789–800, 2001.

- [73019] 11349. Takahashi, C.; Sheng, Z.; Horan, T. P.; Kitayama, H.; Maki, M.; Hitomi, K.; Kitaura, Y.; Takai, S.; Sasahara, R. M.; Horimoto, A.; Ikawa, Y.; Ratzkin, B. J.; Arakawa, T.; Noda, M.: Regulation of matrix metalloproteinase–9 and inhibition of tumor invasion by the membrane–anchored glycoprotein RECK. *Proc. Nat. Acad. Sci.* 95: 13221–13226, 1998.
- [73020] 11350. Nagase, T.; Ishikawa, I.; Nakajima, D.; Ohira, M.; Seki, N.; Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; O'Hara, O.: Prediction of the coding sequences of unidentified human genes. VII. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro. *DNA Res.* 4: 141–150, 1997.
- [73021] 11351. Koseki, T.; Inohara, N.; Chen, S.; Nunez, G.: ARC, an inhibitor of apoptosis expressed in skeletal muscle and heart that interacts selectively with caspases. *Proc. Nat. Acad. Sci.* 95: 5156–5160, 1998.
- [73022] 11352. Li, P.–F.; Li, J.; Muller, E.–C.; Otto, A.; Dietz, R.; von Harsdorf, R.: Phosphorylation by protein kinase CK2: a signaling switch for the caspase–inhibiting protein ARC. *Molec. Cell* 10: 247–258, 2002.
- [73023] 11353. Muto, A.; Hoshino, H.; Madisen, L.; Yanai, N.; Obi–

nata, M.; Karasuyama, H.; Hayashi, N.; Nakauchi, H.; Yamamoto, M.; Groudine, M.; Igarashi, K.: Identification of Bach2 as a B-cell-specific partner for small Maf proteins that negatively regulate the immunoglobulin heavy chain-gene 3-prime enhancer. EMBO J. 17: 5734–5743, 1998.

- [73024] 11354. Sasaki, S.; Ito, E.; Toki, T.; Maekawa, T.; Kanezaki, R.; Umenai, T.; Muto, A.; Nagai, H.; Kinoshita, T.; Yamamoto, M.; Inazawa, J.; Taketo, M. M.; Nakahata, T.; Igarashi, K.; Yokoyama, M.: Cloning and expression of human B cell-specific transcription factor BACH2 mapped to chromosome 6q15. Oncogene 19: 3739–3749, 2000.
- [73025] 11355. Gayle, M. A.; Slack, J. L.; Bonnert, T. P.; Renshaw, B. R.; Sonoda, G.; Taguchi, T.; Testa, J. R.; Dower, S. K.; Sims, J. E.: Cloning of a putative ligand for the T1/ST2 receptor. J. Biol. Chem. 271: 5784–5789, 1996.
- [73026] 11356. Shimaoka, T.; Kume, N.; Minami, M.; Hayashida, K.; Kataoka, H.; Kita, T.; Yonehara, S.: Molecular cloning of a novel scavenger receptor for oxidized low density lipoprotein, SR-PSOX, on macrophages. J. Biol. Chem. 275: 40663–40666, 2000.
- [73027] 11357. Wilbanks, A.; Zondlo, S. C.; Murphy, K.; Mak, S.; Soler, D.; Langdon, P.; Andrew, D. P.; Wu, L.; Briskin, M.: Expression cloning of the STRL33/BONZO/TYMSTR ligand

reveals elements of CC, CXC, CX3C chemokines. J.Immun. 166: 5145–5154, 2001.

- [73028] 11358.Brill, S.; Li, S.; Lyman, C. W.; Church, D. M.; Wasmuth, J. J.; Weissbach, L.; Bernards, A.; Snijders, A. J.: The Ras GTPase-activating-protein-related human protein IQ-GAP2 harbors a potential actin binding domain and interacts with calmodulin and Rho family GTPases. Molec. Cell. Biol. 16:4869–4878, 1996.
- [73029] 11359.Bitner-Glindzicz, M.; Turnpenny, P.; Hoglund, P.; Kaariainen, H.; Sankila, E.-M.; van der Maarel, S. M.; de Kok, Y. J. M.; Ropers, H.-H.; Cremers, F. P. M.; Pembrey, M.; Malcolm, S.: Further mutations in brain 4 (POU3F4) clarify the phenotype in the X-linked deafness, DFN3. Hum.Molec. Genet. 4: 1467–1469, 1995.
- [73030] 11360.de Kok, Y. J. M.; Cremers, C. W. R. J.; Ropers, H.-H.; Cremers, F. P. M.: The molecular basis of X-linked deafness type 3 (DFN3) in two sporadic cases: identification of a somatic mosaicism for a POU3F4 missense mutation. Hum. Mutat. 10: 207–211, 1997.
- [73031] 11361.de Kok, Y. J. M.; Merkx, G. F. M.; van der Maarel, S. M.; Huber, I.; Malcolm, S.; Ropers, H.-H.; Cremers, F. P. M.: A duplication/paracentric inversion associated with familial X-linked deafness (DFN3) suggests the presence of a reg-

ulatory element more than 400 kb upstream of the POU3F4 gene. Hum. Molec. Genet. 4: 2145–2150, 1995.

- [73032] 11362. de Kok, Y. J. M.; van der Maarel, S. M.; Bitner-Glindzicz, M.; Huber, I.; Monaco, A. P.; Malcolm, S.; Pembrey, M. E.; Ropers, H.-H.; Cremers, F. P. M.: Association between X-linked mixed deafness and mutations in the POU domain gene POU3F4. Science 267: 685–688, 1995.
- [73033] 11363. Douville, P. J.; Atanasoski, S.; Tobler, A.; Fontana, A.; Schwab, M. E.: The brain-specific POU-box gene Brn4 is a sex-linked transcription factor located on the human and mouse X chromosomes. Mammalian Genome 5: 180–182, 1994.
- [73034] 11364. Friedman, R. A.; Bykhovskaya, Y.; Tu, G.; Talbot, J. M.; Wilson, D. F.; Parnes, L. S.; Fischel-Ghodsian, N.: Molecular analysis of the POU3F4 gene in patients with clinical and radiographic evidence of X-linked mixed deafness with perilymphatic gusher. Ann. Otol. Rhinol. Laryng. 106: 320–325, 1997.
- [73035] 11365. Minowa, O.; Ikeda, K.; Sugitani, Y.; Oshima, T.; Nakai, S.; Katori, Y.; Suzuki, M.; Furukawa, M.; Kawase, T.; Zheng, Y.; Ogura, M.; Asada, Y.; Watanabe, K.; Yamanaka, H.; Gotoh, S.; Nishi-Takeshima, M.; Sugimoto, T.; Kikuchi, T.; Takasaka, T.; Noda, T.: Altered cochlear fibrocytes in a



mouse model of DFN3 nonsyndromic deafness. *Science* 285: 1408–1411, 1999.

[73036] 11366. Phelps, P. D.; Reardon, W.; Pembrey, M.; Bellman, S.; Luxon, L.: X-linked deafness, stapes gushers and a distinctive defect of the inner ear. *Neuroradiology* 33: 326–330, 1991.

[73037] 11367. Phippard, D.; Boyd, Y.; Reed, V.; Fisher, G.; Masson, W. K.; Evans, E. P.; Saunders, J. C.; Crenshaw, E. B., III: The sex-linked fidget mutation abolishes Brn4/Pou3f4 gene expression in the embryonic inner ear. *Hum. Molec. Genet.* 9: 79–85, 2000.

[73038] 11368. Brown, C. J.; Miller, A. P.; Carrel, L.; Rupert, J. L.; Davies, K. E.; Willard, H. F.: The DXS423E gene in Xp11.21 escapes X chromosome inactivation. *Hum. Molec. Genet.* 4: 251–255, 1995.

[73039] 11369. Sumara, I.; Vorlaufer, E.; Gieffers, C.; Peters, B. H.; Peters, J.-M.: Characterization of vertebrate cohesin complexes and their regulation in prophase. *J. Cell Biol.* 151: 749–761, 2000.

[73040] 11370. Lowe, D. G.; Dizhoor, A. M.; Liu, K.; Gu, Q.; Spencer, M.; Laura, R.; Lu, L.; Hurley, J. B.: Cloning and expression of a second photoreceptor-specific membrane retina guanylyl cyclase (RetGC), RetGC-2. *Proc. Nat.*

Acad.Sci. 92: 5535–5539, 1995.

[73041] 11371.Teague, P. W.; Aldred, M. A.; Jay, M.; Dempster, M.; Harrison,C.; Carothers, A. D.; Hardwick, L. J.; Evans, H. J.; Strain, L.; Brock,D. J. H.; Bunday, S.; Jay, B.; Bird, A. C.; Bhattacharya, S. S.; Wright,A. F.: Heterogeneity analysis in 40 X-linked retinitis pigmentosafamilies. Am. J. Hum. Genet. 55: 105–111, 1994.

[73042] 11372.Bodrug, S. E.; Ray, P. N.; Gonzalez, I. L.; Schmickel, R. D.;Sylvester, J. E.; Worton, R. G.: Molecular analysis of a constitutionalX-autosome translocation in a female with muscular dystrophy. Science 237:1620–1624, 1987.

[73043] 11373.Boyce, F. M.; Beggs, A. H.; Feener, C.; Kunkel, L. M.: Dystrophinis transcribed in brain from a distant upstream promoter. Proc. Nat.Acad. Sci. 88: 1276–1280, 1991.

[73044] 11374.Boyd, Y.; Buckle, V. J.: Cytogenetic heterogeneity of translocationsassociated with Duchenne muscular dystrophy. Clin. Genet. 29: 108–115,1986.

[73045] 11375.Bulman, D. E.; Gangopadhyay, S. B.; Bebchuck, K. G.; Worton, R.G.; Ray, P. N.: Point mutation in the human dystrophin gene: identificationthrough Western blot analysis. Genomics 10: 457–460, 1991.

[73046] 11376.Burke, J. F.; Mogg, A. E.: Suppression of a nonsense mutationin mammalian cells in vivo by the aminoglycoside

antibiotics G-418 and paromomycin. *Nucleic Acids Res.* 13: 6265–6272, 1985.

- [73047] 11377. Burnette, W. N.: 'Western blotting': electrophoretic transfer of proteins from sodium dodecyl sulfate–polyacrylamide gels to unmodified nitrocellulose and radiographic detection with antibody and radioiodinated–protein A. *Anal. Biochem.* 112: 195–203, 1981.
- [73048] 11378. Chamberlain, J. S.; Pearlman, J. A.; Muzny, D. M.; Gibbs, R. A.; Ranier, J. E.; Reeves, A. A.; Caskey, C. T.: Expression of the murine Duchenne muscular dystrophy gene in muscle and brain. *Science* 239:1416–1418, 1988.
- [73049] 11379. Chelly, J.; Concorde, J.-P.; Kaplan, J.-C.; Kahn, A.: Illegitimate transcription: transcription of any gene in any cell type. *Proc. Nat. Acad. Sci.* 86: 2617–2621, 1989.
- [73050] 11380. Chelly, J.; Gilgenkrantz, H.; Hugnot, J. P.; Hamard, G.; Lambert, M.; Recan, D.; Akli, S.; Cometto, M.; Kahn, A.; Kaplan, J. C.: Illegitimate transcription: application to the analysis of truncated transcripts of the dystrophin gene in nonmuscle cultured cells from Duchenne and Becker patients. *J. Clin. Invest.* 88: 1161–1166, 1991.
- [73051] 11381. Chelly, J.; Hamard, G.; Koulakoff, A.; Kaplan, J.-C.; Kahn, A.; Berwald-Netter, Y.: Dystrophin gene transcribed from different promoters in neuronal and glial cells. *Nature*

344: 64–65, 1990.

- [73052] 11382.Chelly, J.; Kaplan, J.-C.; Maire, P.; Gautron, S.; Kahn, A.:Transcription of the dystrophin gene in human muscle and non-muscle tissues. *Nature* 333: 858–860, 1988.
- [73053] 11383.Clemens, P. R.; Ward, P. A.; Caskey, C. T.; Bulman, D. E.; Fenwick, R. G.: Premature chain termination mutation causing Duchenne muscular dystrophy. *Neurology* 42: 1775–1782, 1992.
- [73054] 11384.Cooper, B. J.; Valentine, B. A.; Wilson, S.; Patterson, D. F.; Concannon, P. W.: Canine muscular dystrophy: confirmation of X-linked inheritance. *J. Hered.* 79: 405–408, 1988.
- [73055] 11385.Covone, A. E.; Lerone, M.; Romeo, G.: Genotype–phenotype correlation and germline mosaicism in DMD/BMD patients with deletions of the dystrophin gene. *Hum. Genet.* 87: 353–360, 1991.
- [73056] 11386.Cox, G. A.; Cole, N. M.; Matsumura, K.; Phelps, S. F.; Hauschka, S. D.; Campbell, K. P.; Faulkner, J. A.; Chamberlain, J. S.: Overexpression of dystrophin in transgenic mdx mice eliminates dystrophic symptoms without toxicity. *Nature* 364: 725–729, 1993.
- [73057] 11387.Cox, G. A.; Sunada, Y.; Campbell, K. P.; Chamber-

lain, J. S.:Dp71 can restore the dystrophin-associated glycoprotein complex in muscle but fails to prevent dystrophy. *Nature Genet.* 8: 333–339,1994.

[73058] 11388.Comi, G. P.; Ciafaloni, E.; de Silva, H. A. R.; Prella, A.; Bardoni,A.; Rigoletto, C.; Robotti, M.; Bresolin, N.; Moggio, M.; Fortunato,F.; Ciscato, P.; Turconi, A.; Rose, A. D.; Scarlato, G.: A G(+1)-to-Atransversion at the 5-prime splice site of intron 69 of the dystrophingene causing the absence of peripheral nerve Dp116 and severe clinical involvement in a DMD patient. *Hum. Molec. Genet.* 4: 2171–2174, 1995.

[73059] 11389.Crawford, G. E.; Lu, Q. L.; Partridge, T. A.; Chamberlain, J.S.: Suppression of revertant fibers in mdx mice by expression of a functional dystrophin. *Hum. Molec. Genet.* 10: 2745–2750, 2001.

[73060] 11390.Darras, B. T.; Blattner, P.; Harper, J. F.; Spiro, A. J.; Alter,S.; Francke, U.: Intragenic deletions in 21 Duchenne muscular dystrophy(DMD)/Becker muscular dystrophy (BMD) families studied with the dystrophin cDNA: location of breakpoints on HindIII and BglII exon-containing fragment maps, meiotic and mitotic origin of the mutations. *Am. J.Hum. Genet.* 43: 620–629, 1988.

[73061] 11391.Darras, B. T.; Francke, U.: Normal human genomic

restriction-fragment patterns and polymorphisms revealed by hybridization with the entire dystrophin cDNA. *Am. J. Hum. Genet.* 43: 612-619, 1988.

[73062] 11392. Darras, B. T.; Francke, U.: A partial deletion of the muscular dystrophy gene transmitted twice by an unaffected male. *Nature* 329:556-558, 1987.

[73063] 11393. Davies, K. E.; Smith, T. J.; Bunday, S.; Read, A. P.; Flint, T.; Bell, M.; Speer, A.: Mild and severe muscular dystrophy associated with deletions in Xp21 of the human X chromosome. *J. Med. Genet.* 25:9-13, 1988.

[73064] 11394. De Angelis, F. G.; Sthandier, O.; Berarducci, B.; Toso, S.; Galluzzi, G.; Ricci, E.; Cossu, G.; Bozzoni, I.: Chimeric snRNA molecules carrying antisense sequences against the splice junctions of exon 51 of the dystrophin pre-mRNA induce exon skipping and restoration of a dystrophin synthesis in delta-48-50 DMD cells. *Proc Nat. Acad. Sci.* 99: 9456-9461, 2002.

[73065] 11395. den Dunnen, J. T.; Bakker, E.; Klein Breteler, E. G.; Pearson, P. L.; van Ommen, G. J. B.: Direct mutation of more than 50% of the Duchenne muscular dystrophy mutations by field inversion gels. *Nature* 329:640-642, 1987.

[73066] 11396. Dickson, G.; Pizzey, J. A.; Elsom, V. E.; Love, D.;

Davies, K.E.; Walsh, F. S.: Distinct dystrophin mRNA species are expressed in embryonic and adult mouse skeletal muscle. FEBS Lett. 242: 47–52, 1988.

- [73067] 11397. Dominguez–Steglich, M.; Meng, G.; Bettecken, T.; Muller, C. R.; Schmid, M.: The dystrophin gene is autosomally located on a microchromosome in chicken. Genomics 8: 536–540, 1990.
- [73068] 11398. Doolittle, R. F.: Similar amino acid sequences: chance or common ancestry? Science 214: 149–159, 1981.
- [73069] 11399. Bassi, M. T.; Bergen, A. A. B.; Bitoun, P.; Charles, S. J.; Clementi, M.; Gosselin, R.; Hurst, J.; Lewis, R. A.; Lorenz, B.; Meitinger, T.; Messiaen, L.; Ramesar, R. S.; Ballabio, A.; Schiaffino, M. V.: Diverse prevalence of large deletions within the OA1 gene in ocular albinism type 1 patients from Europe and North America. Hum. Genet. 108: 51–54, 2001.
- [73070] 11400. Bassi, M. T.; Schiaffino, M. V.; Renieri, A.; De Nigris, F.; Galli, L.; Bruttini, M.; Gebbia, M.; Bergen, A. A. B.; Lewis, R. A.; Ballabio, A.: Cloning of the gene for ocular albinism type 1 from the distal short arm of the X chromosome. Nature Genet. 10: 13–19, 1995.
- [73071] 11401. Bergen, A. A. B.; Samanns, C.; Schuurman, E. J. M.; van Osch, L.; van Dorp, D. B.; Pinckers, A. J. L. G.; Bakker,

E.; Gal, A.; van Ommen, G. J. B.; Bleeker-Wagemakers, E. M.: Multipoint linkage analysis in X-linked ocular albinism of the Nettleship-Falls type. *Hum. Genet.* 88:162-166, 1991.

[73072] 11402. Bergen, A. A. B.; Schuurman, E. J. M.; van den Born, L. I.; Samanns, C.; van Dorp, D. B.; Pinckers, A. J. L. G.; Bakker, E.; van Ommen, G. J. B.; Gal, A.; Bleeker-Wagemakers, E. M.: Carrier detection in X-linked ocular albinism of the Nettleship-Falls type by DNA analysis. *Clin. Genet.* 41: 135-138, 1992.

[73073] 11403. Bergen, A. A. B.; Zijp, P.; Schuurman, E. J. M.; Bleeker-Wagemakers, E. M.; Apkarian, P.; van Ommen, G.-J. B.: Refinement of the localization of the X-linked ocular albinism gene. *Genomics* 16: 272-273, 1993.

[73074] 11404. Bouloux, P.-M. G.; Kirk, J.; Munroe, P.; Duke, V.; Meindl, A.; Hilson, A.; Grant, D.; Carter, N.; Betts, D.; Meitinger, T.; Besser, G. M.: Deletion analysis maps ocular albinism proximal to the steroid sulphatase locus. *Clin. Genet.* 43: 169-173, 1993.

[73075] 11405. Charles, S. J.; Green, J. S.; Moore, A. T.; Barton, D. E.; Yates, J. R. W.: Genetic mapping of X-linked ocular albinism: linkage analysis in a large Newfoundland kindred. *Genomics* 16: 259-261, 1993.



- [73076] 11406.Charles, S. J.; Moore, A. T.; Yates, J. R. W.: Genetic mapping of X linked ocular albinism: linkage analysis in British families. *J.Med. Genet.* 29: 552–554, 1992.
- [73077] 11407.Cooper, D. N.; Krawczak, M.: Mechanisms of insertional mutagenesis in human genes causing genetic disease. *Hum. Genet.* 87: 409–415, 1991.
- [73078] 11408.Creel, D.; O'Donnell, F. E., Jr.; Witkop, C. J., Jr.: Visual system anomalies in human ocular albinos. *Science* 201: 931–933, 1978.
- [73079] 11409.d'Addio, M.; Pizzigoni, A.; Bassi, M. T.; Baschiroto, C.; Valetti, C.; Incerti, B.; Clementi, M.; De Luca, M.; Bal-labio, A.; Schiaffino, M. V.: Defective intracellular transport and processing of OA1 is a major cause of ocular albinism type 1. *Hum. Molec. Genet.* 9: 3011–3018, 2000.
- [73080] 11410.Engelhard, C. F.: Eine Familie mit hereditaerem Nystagmus. *Z.Ges. Neurol. Psychiat.* 28: 319–338, 1915.
- [73081] 11411.Fialkow, P. J.; Giblett, E. R.; Motulsky, A. G.: Measurable linkage between ocular albinism and Xg. *Am. J. Hum. Genet.* 19: 63–69, 1967.
- [73082] 11412.Francois, J.; Deweer, J. P.: Albinisme oculaire lie au sexe et alterations caracteristiques du fond d'oeil chez les femmes heterozygotes. *Ophthalmologica* 126:209–221, 1953.

- [73083] 11413. Gillespie, F. D.: Ocular albinism with report of a family with female carriers. *Arch. Ophthalmol.* 66: 774–777, 1961.
- [73084] 11414. Incerti, B.; Cortese, K.; Pizzigoni, A.; Surace, E. M.; Varani, S.; Coppola, M.; Jeffery, G.; Seeliger, M.; Jaissle, G.; Bennett, D. C.; Marigo, V.; Schiaffino, M. V.; Tacchetti, C.; Ballabio, A.: Oa1 knock-out: new insights on the pathogenesis of ocular albinism type 1. *Hum. Molec. Genet.* 9: 2781–2788, 2000.
- [73085] 11415. Jaeger, C.; Jay, B.: X-linked ocular albinism: a family containing a manifesting heterozygote, and an affected male married to a female with autosomal recessive ocular albinism. *Hum. Genet.* 56: 299–304, 1981.
- [73086] 11416. Johnson, G. J.; Gillan, J. G.; Pearce, W. G.: Ocular albinism in Newfoundland. *Can. J. Ophthalmol.* 6: 237–248, 1971.
- [73087] 11417. Kidd, J. R.; Castiglione, C. M.; Davies, K. E.; Pakstis, A. J.; Gusella, J.; Sparkes, R. S.; Pearson, P.; Willard, H.; Kidd, K. K.: Mapping the locus for X-linked ocular albinism (OA). (Abstract) *Am. J. Hum. Genet.* 37: A161 only, 1985.
- [73088] 11418. Krawczak, M.; Reiss, J.; Cooper, D. N.: The mutational spectrum of single base-pair substitutions in mRNA

splice junctions of humangenes: causes and consequences. Hum. Genet. 90: 41–54, 1992.

[73089] 11419.Lein, J. N.; Stewart, C. T.; Moll, F. C.: Sex-linked hereditary nystagmus. Pediatrics 18: 214–217, 1956.

[73090] 11420.Lyon, M. F.: Sex chromatin and gene action in the mammalian X-chromosome. Am.J. Hum. Genet. 14: 135–148, 1962.

[73091] 11421.Meindl, A.; Hosenfeld, D.; Bruckl, W.; Schuffenhauer, S.; Jenderny, J.; Bacskulin, A.; Oppermann, H.-C.; Swensson, O.; Bouloux, P.; Meitinger, T.: Analysis of a terminal Xp22.3 deletion in a patient with six monogenic disorders: implications for the mapping of X linked ocular albinism. J. Med. Genet. 30: 838–842, 1993.

[73092] 11422.Negrelli, B. C.: L'albinisme oculaire lie au sexe dans le cadre du depistage des heterozygotes en ophtalmologie. J. Genet. Hum. 8:108 only, 1959.

[73093] 11423.Newton, J. M.; Orlow, S. J.; Barsh, G. S.: Isolation and characterization of a mouse homolog of the X-linked ocular albinism (OA1) gene. Genomics 37:219–225, 1996.

[73094] 11424.O'Donnell, F. E., Jr.; Green, W. R.; Fleishman, J. A.; Hambrick, G. W.: X-linked ocular albinism in Blacks: ocular albinism cum pigmento. Arch.Ophthal. 96: 1189–1192, 1978.

- [73095] 11425.O'Donnell, F. E., Jr.; Hambrick, G. W., Jr.; Green, W. R.; Iliff,W. J.; Stone, D. L.: X-linked ocular albinism: an oculocutaneousmacromelanosomal disorder. Arch. Ophthalmol. 94: 1883-1892, 1976.
- [73096] 11426.Oetting, W. S.: New insights into ocular albinism type 1 (OA1):mutations and polymorphisms of the OA1 gene. Hum. Mutat. 19: 85-92,2002.
- [73097] 11427.Pearce, W. G.; Johnson, G. J.; Gillan, J. G.: Nystagmus in afemale carrier of ocular albinism. J. Med. Genet. 9: 126-128, 1972.
- [73098] 11428.Pearce, W. G.; Johnson, G. J.; Sanger, R.: Ocular albinism andXg. (Letter) Lancet I: 1072 only, 1971.
- [73099] 11429.Pearce, W. G.; Sanger, R.: X mapping in man: evidence againstdirect measurable linkage between ocular albinism and deutan colourblindness. J. Med. Genet. 13: 319 only, 1976.
- [73100] 11430.Gaugitsch, H. W.; Prieschl, E. E.; Kalthoff, F.; Huber, N. E.;Baumruker, T.: A novel transiently expressed, integral membrane proteinlinked to cell activation: molecular cloning via the rapid degradationsignal AUUUA. J. Biol. Chem 267: 11267-11273, 1992.
- [73101] 11431.Kanai, Y.; Segawa, H.; Miyamoto, K.; Uchino, H.; Takeda, H.; Endou,H.: Expression cloning and characteri-

zation of a transporter for large neutral amino acids activated by the heavy chain of 4F2 antigen (CD98). J. Biol. Chem. 273: 23629–23632, 1998.

[73102] 11432. Takahara, K.; Kessler, E.; Biniaminov, L.; Brusel, M.; Eddy, R.L.; Jani-Sait, S.; Shows, T. B.; Greenspan, D. S.: Type I procollagen COOH-terminal proteinase enhancer protein: identification, primary structure, and chromosomal localization of the cognate human gene (PCOLCE). J. Biol. Chem. 269: 26280–26285, 1994.

[73103] 11433. Takahara, K.; Osborne, L.; Elliott, R. W.; Tsui, L.-C.; Scherer, S. W.; Greenspan, D. S.: Fine mapping of the human and mouse genes for the type I procollagen COOH-terminal proteinase enhancer protein. Genomics 31:253–256, 1996.

[73104] 11434. Amagai, M.; Wang, Y.; Minoshima, S.; Kawamura, K.; Green, K. J.; Nishikawa, T.; Shimizu, N.: Assignment of the human genes for desmocollin 3 (DSC3) and desmocollin 4 (DSC4) to chromosome 18q12. Genomics 25:330–332, 1995.

[73105] 11435. Kawamura, K.; Watanabe, K.; Suzuki, T.; Yamakawa, T.; Kamiyama, T.; Nakagawa, H.; Tsurufuji, S.: cDNA cloning and expression of a novel human desmocollin. J. Biol. Chem. 269: 26295–26302, 1994.

- [73106] 11436.King, I. A.; Sullivan, K. H.; Bennett, R., Jr.; Buxton, R. S.:The desmocollins of human foreskin epidermis: identification and chromosomal assignment of a third gene and expression patterns of the three isoforms. *J.Invest. Derm.* 105: 314–321, 1995.
- [73107] 11437.Warburg, O.: On the origin of cancer cells. *Science* 123: 309–314,1956.
- [73108] 11438.Yang, R.–B.; Foster, D. C.; Garbers, D. L.; Fulle, H.–J.: Twomembrane forms of guanylyl cyclase found in the eye. *Proc. Nat. Acad.Sci.* 92: 602–606, 1995.
- [73109] 11439.Dubrovsky, A. L.; Taratuto, A. L.; Sevelever, G.; Schultz, M.;Pegoraro, E.; Hoop, R. C.; Hoffman, E. P.: Duchenne muscular dystrophyand myotonic dystrophy in the same patient. *Am. J. Med. Genet.* 55:342–348, 1995.
- [73110] 11440.Cazzola, M.; Bergamaschi, G.: X-linked Wiskott–Aldrich syndromein a girl. (Letter) *New Eng. J. Med.* 338: 1850 only, 1998.
- [73111] 11441.Emery, A. E. H.: *Duchenne Muscular Dystrophy*. Oxford, UK: OxfordUniversity Press (pub.) (2nd ed.): 1993.
- [73112] 11442.Fabb, S. A.; Wells, D. J.; Serpente, P.; Dickson, G.: Adeno-associatedvirus vector gene transfer and sarcolemmal expression of a 144 kDamicro-dystrophin effectively restores the dystrophin-associated proteincom–

plex and inhibits myofibre degeneration in nude/mdx mice. *Hum.Molec. Genet.* 11: 733–741, 2002.

[73113] 11443.Feener, C. A.; Boyce, F. M.; Kunkel, L. M.: Rapid detection of CA polymorphisms in cloned DNA: application to the 5–prime region of the dystrophin gene. *Am. J. Hum. Genet.* 48: 621–627, 1991.

[73114] 11444.Ferlini, A.; Galie, N.; Merlini, L.; Sewry, C.; Branzi, A.; Muntoni, F.: A novel Alu–like element rearranged in the dystrophin gene causes a splicing mutation in a family with X–linked dilated cardiomyopathy. *Am.J. Hum. Genet.* 63: 436–446, 1998.

[73115] 11445.Finnegan, D. J.: Eukaryotic transposable elements and genome evolution. *Trends Genet.* 5: 103–107, 1989.

[73116] 11446.Forrest, S. M.; Cross, G. S.; Speer, A.; Gardner–Medwin, D.; Burn, J.; Davies, K. E.: Preferential deletion of exons in Duchenne and Becker muscular dystrophies. *Nature* 329: 638–640, 1987.

[73117] 11447.Francke, U.; Ochs, H. D.; de Martinville, B.; Giacalone, J.; Lindgren, V.; Distèche, C.; Pagon, R. A.; Hofker, M. H.; van Ommen, G.–J. B.; Pearson, P. L.; Wedgwood, R. J.: Minor Xp21 chromosome deletion in a male associated with expression of Duchenne muscular dystrophy, chronic granulomatous disease, retinitis pigmentosa, and McLeod

syndrome. Am.J. Hum. Genet. 37: 250–267, 1985.

- [73118] 11448.Furst, D.; Nave, R.; Osborn, M.; Weber, K.; Bardosi, A.; Archidiacono,N.; Ferro, M.; Romano, V.; Romeo, G.: Nebulin and titin expressionin Duchenne muscular dystrophy appears normal. FEBS Lett. 224: 49–53,1987.
- [73119] 11449.Giacalone, J. P.; Francke, U.: Common sequence motifs at therearrangement sites of a constitutional X/ autosome translocation andassociated deletion. Am. J. Hum. Genet. 50: 725–741, 1992.
- [73120] 11450.Gillard, E. F.; Chamberlain, J. S.; Murphy, E. G.; Duff, C. L.;Smith, B.; Burghes, A. H. M.; Thompson, M. W.; Sutherland, J.; Oss,I.; Bodrug, S. E.; Klamut, H. J.; Ray, P. N.; Worton, R. G.: Molecularand phenotypic analysis of patients with deletions within the deletion–richregion of the Duchenne muscular dystrophy (DMD) gene. Am. J. Hum.Genet. 45: 507–520, 1989.
- [73121] 11451.Ginjaar, I. B.; Kneppers, A. L. J.; Meulen, J.–D. M.; Anderson,L. V. B.; Bremmer–Bout, M.; van Deutekom, J. C. T.; Weegenaar, J.;den Dunnen, J. T.; Bakker, E.: Dystrophin nonsense mutation inducesdifferent levels of exon 29 skipping and leads to variable phenotypeswithin one BMD family. Europ. J. Hum. Genet. 8: 793–796, 2000.
- [73122] 11452.Greenberg, D. S.; Sunada, Y.; Campbell, K. P.;



Yaffe, D.; Nudel, U.: Exogenous Dp71 restores the levels of dystrophin associated proteins but does not alleviate muscle damage in mdx mice. *Nature Genet.* 8:340–344, 1994.

[73123] 11453. Gussoni, E.; Soneoka, Y.; Strickland, C. D.; Buzney, E. A.; Khan, M. K.; Flint, A. F.; Kunkel, L. M.; Mulligan, R. C.: Dystrophin expression in the mdx mouse restored by stem cell transplantation. *Nature* 401:390–394, 1999.

[73124] 11454. Hagiwara, Y.; Mizuno, Y.; Takemitsu, M.; Matsuzaki, T.; Nonaka, I.; Ozawa, E.: Dystrophin-positive muscle fibers following C2 myoblast transplantation into mdx nude mice. *Acta Neuropath.* 90: 592–600, 1995.

[73125] 11455. Hagiwara, Y.; Nishio, H.; Kitoh, Y.; Takeshima, Y.; Narita, N.; Wada, H.; Yokoyama, M.; Nakamura, H.; Matsuo, M.: A novel point mutation (G(–1) to T) in a 5-prime splice donor site of intron 13 of the dystrophin gene results in exon skipping and is responsible for Becker muscular dystrophy. *Am. J. Hum. Genet.* 54: 53–61, 1994.

[73126] 11456. Hammonds, R. G., Jr.: Protein sequence of DMD gene is related to actin-binding domain of alpha-actinin. (Letter) *Cell* 51: 1, 1987.

[73127] 11457. Harper, S. Q.; Hauser, M. A.; Dello Russo, C.; Duan, D.; Crawford, R. W.; Phelps, S. F.; Harper, H. A.; Robinson, A. S.; Engelhardt, J. F.; Brooks, S. V.; Chamberlain, J. S.:

Modular flexibility of dystrophin: implications for gene therapy of Duchenne muscular dystrophy. *Nature Med.* 8: 253–261, 2002.

- [73128] 11458. Hart, K. A.; Hodgson, S.; Walker, A.; Cole, C. G.; Johnson, L.; Dubowitz, V.; Bobrow, M.: DNA deletions in mild and severe Becker muscular dystrophy. *Hum. Genet.* 75: 281–285, 1987.
- [73129] 11459. Hodgson, S. V.; Abbs, S.; Clark, S.; Manzur, A.; Heckmatt, J. Z. H.; Dubowitz, V.; Bobrow, M.: Correlation of clinical and deletion data in Duchenne and Becker muscular dystrophy, with special reference to mental ability. *Neuromusc. Disord.* 2: 269–276, 1992.
- [73130] 11460. Hoffman, E. P.; Brown, R. H., Jr.; Kunkel, L. M.: The protein product of the Duchenne muscular dystrophy locus. *Cell* 51: 919–928, 1987.
- [73131] 11461. Hoffman, E. P.; Knudson, C. M.; Campbell, K. P.; Kunkel, L. M.: Subcellular fractionation of dystrophin to the triads of skeletal muscle. *Nature* 330: 754–758, 1987.
- [73132] 11462. Hoffman, E. P.; Monaco, A. P.; Feener, C. C.; Kunkel, L. M.: Conservation of the Duchenne muscular dystrophy gene in mice and humans. *Science* 238: 347–350, 1987.
- [73133] 11463. Hoop, R. C.; Russo, L. S.; Riconda, D. L.; Schwartz,

L. S.; Hoffman, E. P.: Restoration of half the normal dystrophin sequence in a double-deletion Duchenne muscular dystrophy family. *Am. J. Med. Genet.* 49: 323–327, 1994.

[73134] 11464. Hoffman, E. P.; Fischbeck, K. H.; Brown, R. H.; Johnson, M.; Medori, R.; Loike, J. D.; Harris, J. B.; Waterston, R.; Brooke, M.; Specht, L.; Kopsky, W.; Chamberlain, J.; Caskey, C. T.; Shapiro, F.; Kunkel, L. M.: Characterization of dystrophin in muscle-biopsy specimens from patients with Duchenne's or Becker's muscular dystrophy. *New Eng. J. Med.* 318: 1363–1368, 1988.

[73135] 11465. Howard, P. L.; Dally, G. Y.; Wong, M. H.; Ho, A.; Weleber, R. G.; Pillers, D.-A. M.; Ray, P. N.: Localization of dystrophin isoform Dp71 to the inner limiting membrane of the retina suggests a unique functional contribution of Dp71 in the retina. *Hum. Molec. Genet.* 7:1385–1391, 1998.

[73136] 11466. Hu, X.; Burghes, A. H. M.; Bulman, D. E.; Ray, P. N.; Worton, R. G.: Evidence for mutation by unequal sister chromatid exchange in the Duchenne muscular dystrophy gene. *Am. J. Hum. Genet.* 44:855–863, 1989.

[73137] 11467. Birnbaumer, M.: Vasopressin receptors. *TEM* 11: 406–410, 2000.

[73138] 11468. Kools, P.; Vanhalst, K.; Wan den Eynde, E.; van Roy,

F.: The humancadherin-10 gene: complete coding sequence, predominant expression in the brain and mapping on chromosome 5p13-13. FEBS Lett. 452:328-334, 1999.

[73139] 11469.Leder, S.; Weber, Y.; Altafaj, X.; Estivill, X.; Joost, H.-G.;Becker, W.: Cloning and characterization of DYRK1B, a novel member of the DYRK family of protein kinases.

Biochem. Biophys. Res. Commun. 254:474-479, 1999.

[73140] 11470.Lee, K.; Deng, X.; Friedman, E.: Mirk protein kinase is a mitogen-activated protein kinase substrate that mediates survival of colon cancer cells. Cancer Res. 60: 3631-3637, 2000.

[73141] 11471.Coyle, A. J.; Lehar, S.; Lloyd, C.; Tian, J.; Delaney, T.; Manning, S.; Nguyen, T.; Burwell, T.; Schneider, H.; Gonzalo, J. A.; Gosselin, M.; Owen, L. R.; Rudd, C. E.; Gutierrez-Ramos, J. C.: The CD28-related molecule ICOS is required for effective T cell-dependent immune responses. Immunity 13:95-105, 2000.

[73142] 11472.Dong, C.; Juedes, A. E.; Temann, U. A.; Shresta, S.; Allison, J.P.; Ruddle, N. H.; Flavell, R. A.: ICOS co-stimulatory receptor is essential for T-cell activation and function. Nature 409: 97-101, 2001.

[73143] 11473.Haimila, K. E.; Partanen, J. A.; Holopainen, P. M.: Genetic polymorphism of the human ICOS gene. Immuno-

genetics 53: 1028–1032, 2002.

- [73144] 11474.Hutloff, A.; Dittrich, A. M.; Beier, K. C.; Eljaschewitsch, B.;Kraft, R.; Anagnostopoulos, I.; Kroczeck, R. A.: ICOS is an inducibleT-cell co-stimulator structurally and functionally related to CD28. *Nature* 397:263–266, 1999.
- [73145] 11475.McAdam, A. J.; Greenwald, R. J.; Levin, M. A.; Chernova, T.; Malenkovich,N.; Ling, V.; Freeman, G. J.; Sharpe, A. H.: ICOS is critical forCD40-mediated antibody class switching. *Nature* 409: 102–105, 2001.
- [73146] 11476.Pack, S. D.; Pak, E.; Tanigami, A.; Ledbetter, D. H.; Fukuda, M.N.: Assignment of the bystin gene BYSL to human chromosome band 6p21.1by in situ hybridization. *Cytogenet. Cell Genet.* 83: 76–77, 1998.
- [73147] 11477.Medina, M.; Marinescu, R. C.; Overhauser, J.; Kosik, K. S.: Hemizygosityof delta-catenin (CTNND2) is associated with severe mental retardationin cri-du-chat syndrome. *Genomics* 63: 157–164, 2000.
- [73148] 11478.Paffenholz, R.; Franke, W. W.: Identification and localizationof a neurally expressed member of the plakoglobin/armadillo multigenefamily. *Differentiation* 61: 293–304, 1997.
- [73149] 11479.Zhou, J.; Liyanage, U.; Medina, M.; Ho, C.; Simmons, A. D.; Lovett.M.; Kosik, K. S.: Presenilin 1 interac-

tion in the brain with a novel member of the armadillo family. *Neuroreport* 8: 2085–2090, 1997.

- [73150] 11480. Hatzfeld, M.; Nachtsheim, C.: Cloning and characterization of a new armadillo family member, p0071, associated with the junctional plaque: evidence for a subfamily of closely related proteins. *J. Cell Sci.* 109: 2767–2778, 1996.
- [73151] 11481. Burger, J.; Fonknechten, N.; Hoeltzenbein, M.; Neumann, L.; Bratanoff, E.; Hazan, J.; Reis, A.: Hereditary spastic paraplegia caused by mutations in the SPG4 gene. *Europ. J. Hum. Genet.* 8: 771–776, 2000.
- [73152] 11482. Burger, J.; Metzke, H.; Paternotte, C.; Schilling, F.; Hazan, J.; Reis, A.: Autosomal dominant spastic paraplegia with anticipation maps to a 4-cM interval on chromosome 2p21–p24 in a large German family. *Hum. Genet.* 98: 371–375, 1996.
- [73153] 11483. Errico, A.; Ballabio, A.; Rugarli, E. I.: Spastin, the protein mutated in autosomal dominant hereditary spastic paraplegia, is involved in microtubule dynamics. *Hum. Molec. Genet.* 11: 153–163, 2002.
- [73154] 11484. Lindsey, J. C.; Lusher, M. E.; McDermott, C. J.; White, K. D.; Reid, E.; Rubinsztein, D. C.; Bashir, R.; Hazan, J.; Shaw, P. J.; Bushby, K. M. D.: Mutation analysis of the

spastin gene (SPG4) inpatients with hereditary spastic paraplegia. *J. Med. Genet.* 37:759–765, 2000.

[73155] 11485.Takahashi, N.; Tuiki, H.; Saya, H.; Kaibuchi, K.: Localization of the gene coding for ROCK II/Rho kinase on human chromosome 2p24. *Genomics* 55:235–237, 1999.

[73156] 11486.Gruber, A. D.; Pauli, B. U.: Clustering of the human CLCA gene family on the short arm of chromosome 1 (1p22–31). *Genome* 42: 1030–1032, 1999.

[73157] 11487.Kedra, D.; Pan, H.–Q.; Seroussi, E.; Fransson, I.; Guilbaud, C.; Collins, J. E.; Dunham, I.; Blennow, E.; Roe, B. A.; Piehl, F.; Dumanski, J. P.: Characterization of the human synaptogyrin gene family. *Hum.Genet.* 103: 131–141, 1998.

[73158] 11488.Hazan, J.; Davoine, C. S.; Mavel, D.; Fonknechten, N.; Paternotte, C.; Fizames, C.; Cruaud, C.; Samson, D.; Muselet, D.; Vega–Czarny, N.; Brice, A.; Gyapay, G.; Heilig, R.; Fontaine, B.; Weissenbach, J.: A fine integrated map of the SPG4 locus excludes an expanded CAG repeat in chromosome 2p–linked autosomal dominant spastic paraplegia. *Genomics* 60:309–319, 1999.

[73159] 11489.Hazan, J.; Fonknechten, N.; Mavel, D.; Paternotte, C.; Samson, D.; Artiguenave, F.; Davoine, C.–S.; Cruaud, C.; Durr, A.; Wincker, P.; Brottier, P.; Cattolico, L.; Barbe, V.;

Burgunder, J.-M.; Prud'homme, J.-F.; Brice, A.; Fontaine, B.; Heilig, R.; Weissenbach, J.: Spastin, a new AAA protein, is altered in the most frequent form of autosomal dominant spastic paraplegia. *Nature Genet.* 23: 296–303, 1999.

[73160] 11490. Bedford, F. K.; Ashworth, A.; Enver, T.; Wiedemann, L. M.: HEX: a novel homeobox gene expressed during haematopoiesis and conserved between mouse and human. *Nucleic Acids Res.* 21: 1245–1249, 1993.

[73161] 11491. Crompton, M. R.; Bartlett, T. J.; MacGregor, A. D.; Manfioletti, G.; Buratti, E.; Giancotti, V.; Goodwin, G. H.: Identification of a novel vertebrate homeobox gene expressed in haematopoietic cells. *Nucleic Acids Res.* 20: 5661–5667, 1992.

[73162] 11492. D'Elia, A. V.; Tell, G.; Russo, D.; Arturi, F.; Puglisi, F.; Manfioletti, G.; Gattei, V.; Mack, D. L.; Cataldi, P.; Filetti, S.; Di Loreto, C.; Damante, G.: Expression and localization of the homeodomain-containing protein HEX in human thyroid tumors. *J. Clin. Endocr. Metab.* 87: 1376–1383, 2002.

[73163] 11493. Hromas, R.; Radich, J.; Collins, S.: PCR cloning of an orphan homeobox gene (PRH) preferentially expressed in myeloid and liver cells. *Biochem. Biophys. Res. Commun.* 195: 976–983, 1993.



- [73164] 11494.Morgutti, M.; Demori, E.; Pecile, V.; Amoroso, A.; Rustighi, A.;Manfioletti, G.: Genomic organization and chromosome mapping of thehuman homeobox gene HHEX. *Cytogenet. Cell Genet.* 94: 30–32, 2001.
- [73165] 11495.Tanaka, T.; Inazu, T.; Yamada, K.; Myint, Z.; Keng, V. W; Inoue,Y.; Taniguchi, N.; Noguchi, T.: cDNA cloning and expression of rathomeobox gene, Hex, and functional characterization of the protein. *Biochem.J.* 339: 111–117, 1999.
- [73166] 11496.Miles, M. F.; Barhite, S.; Sganga, M.; Elliott, M.: Phosducin–likeprotein: an ethanol–responsive potential modulator of guanine nucleotide–bindingprotein function. *Proc. Nat. Acad. Sci.* 90: 10831–10835, 1993.
- [73167] 11497.Thibault, C.; Wang, J. F.; Charnas, R.; Mirel, D.; Barhite, S.;Miles, M. F.: Cloning and characterization of the rat and human phosducin–likeprotein genes: structure, expression and chromosomal localization. *Biochim.Biophys. Acta* 1444: 346–354, 1999.
- [73168] 11498.Pope, R. K.; Pestonjamas, K. N.; Smith, K. P.; Wulfkuhle, J. D.;Strassel, C. P.; Lawrence, J. B.; Luna, E. J.: Cloning, characterization,and chromosomal localization of human supervillin (SVIL). *Genomics* 52:342–351, 1998.
- [73169] 11499.Ting, H.–J.; Yeh, S.; Nishimura, K.; Chang, C.: Su–

pervillin associates with androgen receptor and modulates its transcriptional activity. Proc.Nat. Acad. Sci. 99: 661–666, 2002.

- [73170] 11500. Bianchi, V.; Robles, R.; Alberio, L.; Furlan, M.; Lammle, B.: Von Willebrand factor–cleaving protease (ADAMTS13) in thrombocytopenic disorders: a severely deficient activity is specific for thrombotic thrombocytopenic purpura. Blood 100: 710–713, 2002.
- [73171] 11501. Fujikawa, K.; Suzuki, H.; McMullen, B.; Chung, D.: Purification of human von Willebrand factor–cleaving protease and its identification as a new member of the metalloproteinase family. Blood 98: 1662–1666, 2001.
- [73172] 11502. Furlan, M.; Robles, R.; Solenthaler, M.; Wassmer, M.; Sandoz, P.; Lammle, B.: Deficient activity of von Willebrand factor–cleaving protease in chronic relapsing thrombotic thrombocytopenic purpura. Blood 89: 3097–3103, 1997.
- [73173] 11503. Gerritsen, H. E.; Robles, R.; Lammle, B.; Furlan, M.: Partial amino acid sequence of purified von Willebrand factor–cleaving protease. Blood 98: 1654–1661, 2001.
- [73174] 11504. Kokame, K.; Matsumoto, M.; Soejima, K.; Yagi, H.; Ishizashi, H.; Funato, M.; Tamai, H.; Konno, M.; Kamide, K.; Kawano, Y.; Miyata, T.; Fujimura, Y.: Mutations and com–

mon polymorphisms in ADAMTS13 gene responsible for von Willebrand factor-cleaving protease activity. *Proc. Nat. Acad. Sci.* 99: 11902–11907, 2002.

- [73175] 11505. Alexopoulou, L.; Holt, A. C.; Medzhitov, R.; and Flavell, R. A.: Recognition of double stranded RNA and activation of NF $\kappa$ B by Toll-like receptor 3. *Nature* 413: 732–738, 2001.
- [73176] 11506. Advani, R. J.; Bae, H.-R.; Bock, J. B.; Chao, D. S.; Doung, Y.-C.; Prekeris, R.; Yoo, J.-S.; Scheller, R. H.: Seven novel mammalian SNARE proteins localize to distinct membrane compartments. *J. Biol. Chem.* 273: 10317–10324, 1998.
- [73177] 11507. Bock, J. B.; Scheller, R. H.: A fusion of new ideas. *Nature* 387: 133–135, 1997.
- [73178] 11508. Bui, T. D.; Wong, S. H.; Lu, L.; Hong, W.: Endobrevin maps to chromosome 2 in human and chromosome 6 in mouse. *Genomics* 54: 579–580, 1998.
- [73179] 11509. Wong, S. H.; Zhang, T.; Xu, Y.; Subramaniam, V. N.; Griffiths, G.; Hong, W.: Endobrevin, a novel synaptobrevin/VAMP-like protein preferentially associated with the early endosome. *Molec. Biol. Cell* 9: 1549–1563, 1998.
- [73180] 11510. Hewett-Emmett, D.; Tashian, R. E.: Functional diversity, conservation, and convergence in the evolution of

the alpha-, beta- and gamma- carbonicanhydrase gene families. *Molec. Phylogenet. Evol.* 5: 50–77, 1996.

[73181] 11511.Nakagawa, Y.; Uemura, H.; Hirao, Y.; Yoshida, K.; Saga, S.; Yoshikawa,K.: Radiation hybrid mapping of the human MN/CA9 locus to chromosomeband 9p12–p13. *Genomics* 53: 118–119, 1998.

[73182] 11512.Opavsky, R.; Pastorekova, S.; Zelnik, V.; Gibadulinova, A.; Stanbridge,E. J.; Zavada, J.; Kettmann, R.; Pastorek, J.: Human MN/CA9 gene,a novel member of the carbonic anhydrase family: structure and exonto protein domain relationships. *Genomics* 33: 480–487, 1996.

[73183] 11513.Soilleux, E. J.; Barten, R.; Trowsdale, J.: Cutting edge: DC-SIGN;a related gene, DC-SIGNR; and CD23 form a cluster on 19p13. *J. Immun.* 165:2937–2942, 2000.

[73184] 11514.Matoskova, B.; Wong, W. T.; Nomura, N.; Robbins, K. C.; Di Fiore,P. P.: RN-tre specifically binds to the SH3 domain of eps8 with highaffinity and confers growth advantage to NIH3T3 upon carboxy-terminaltruncation. *Oncogene* 12: 2679–2688, 1996.

[73185] 11515.Matoskova, B.; Wong, W. T.; Seki, N.; Nagase, T.; Nomura, N.; Robbins,K. C.; Di Fiore, P. P.: RN-tre identifies a family of tre-relatedproteins displaying a novel potential protein binding domain. *Oncogene* 12:2563–2571, 1996.

- [73186] 11516.Lee, M. P.; Brandenburg, S.; Landes, G. M.; Adams, M.; Miller,G.; Feinberg, A. P.: Two novel genes in the center of the 11p15 imprinted domain escape genomic imprinting. *Hum. Molec. Genet.* 8: 683–690,1999.
- [73187] 11517.Nakamura, M.; Masuda, H.; Horii, J.; Kuma, K.; Yokoyama, N.; Ohba,T.; Nishitani, H.; Miyata, T.; Tanaka, M.; Nishimoto, T.: When overexpressed,a novel centrosomal protein, RanBPM, causes ectopic microtubule nucleations similar to gamma-tubulin. *J. Cell Biol.* 143: 1041–1052, 1998.
- [73188] 11518.Nishitani, H.; Hirose, E.; Uchimura, Y.; Nakamura, M.; Umeda,M.; Nishii, K.; Mori, N.; Nishimoto, T.: Full-sized RanBPM cDNA encodes a protein possessing a long stretch of proline and glutamine within the N-terminal region, comprising a large protein complex. *Gene* 272:25–33, 2001.
- [73189] 11519.Miyamoto, T.; Kanazawa, N.; Kato, S.; Kawakami, M.; Inoue, Y.;Kuhara, T.; Inoue, T.; Takeshita, K.; Tsujino, S.: Diagnosis of Japanese patients with HHH syndrome by molecular genetic analysis: a common mutation, R179X. *J. Hum. Genet.* 46: 260–262, 2001.
- [73190] 11520.Nakajima, M.; Ishii, S.; Mito, T.; Takeshita, K.; Takashima, S.;Takakura, H.; Inoue, I.; Saheki, T.; Akiyoshi,

H.; Ichihara, K.:Clinical, biochemical and ultrastructural study on the pathogenesis of hyperornithinemia–hyperammonemia–homocitrullinuria syndrome. *BrainDev.* 10: 181–185, 1988.

[73191] 11521.Tsujino, S.; Kanazawa, N.; Ohashi, T.; Eto, Y.; Saito, T.; Kira,J.; Yamada, T.: Three novel mutations (G27E, in-sAAC, R179X) in theORNT1 gene of Japanese patients with hyperornithinemia, hyperammonemia,and homocitrullinuria syndrome. *Ann. Neurol.* 47: 625–631, 2000.

[73192] 11522.Grand, R. J. A.; Milner, A. E.; Mustoe, T.; Johnson, G. D.; Owen,D.; Grant, M. L.; Gregory, C. D.: A novel protein expressed in mammalian cells undergoing apoptosis. *Exp. Cell Res.* 218: 439–451, 1995.

[73193] 11523.Twist, C. J.; Beier, D. R.; Disteché, C. M.; Edelhoff, S.; Tedder,T. F.: The mouse Cd83 gene: structure, domain organization, and chromosomal localization. *Immunogenetics* 48: 383–393, 1998.

[73194] 11524.Lee, S. B.; Rao, A. K.; Lee, K.–H.; Yang, X.; Bae, Y. S.; Rhee,S. G.: Decreased expression of phospholipase C–b2 isozyme in human platelets with impaired function. *Blood* 88: 1684–1691, 1996.

[73195] 11525.Mao, G. F.; Vaidyula, V. R.; Kunapuli, S. P.; Rao, A. K.: Lineage–specific defect in gene expression in human

platelet phospholipase C-beta-2 deficiency. *Blood* 99: 905-911, 2002.

[73196] 11526. Park, D.; Jhon, D.-Y.; Kriz, R.; Knopf, J.; Rhee, S. G.: Cloning, sequencing, expression, and Gq-independent activation of phospholipase C-beta-2. *J. Biol. Chem.* 267: 16048-16055, 1992.

[73197] 11527. Park, S. H.; Ryu, S. H.; Suh, P. G.; Kim, H.: Assignment of human PLCB2 encoding PLC-beta-2 to human chromosome 15q15 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 83: 48-49, 1998.

[73198] 11528. Rao, A. K.; Kowalska, M. A.; Disa, J.: Impaired cytoplasmic ionized calcium mobilization in inherited platelet secretion defects. *Blood* 74:664-672, 1989.

[73199] 11529. Yang, X.; Sun, L.; Ghosh, S.; Rao, A. K.: Human platelet signaling defect characterized by impaired production of 1,4,5 inositol triphosphate and phosphatidic acid, and diminished pleckstrin phosphorylation. Evidence for defective phospholipase C activation. *Blood* 88: 1676-1683, 1996.

[73200] 11530. Gillen, C. M.; Brill, S.; Payne, J. A.; Forbush, B., III: Molecular cloning and functional expression of the K-Cl cotransporter from rabbit, rat, and human: a new member of the cation-chloride cotransporter family. *J. Biol. Chem.*

271: 16237–16244, 1996.

- [73201] 11531.Larsen, F.; Solheim, J.; Kristensen, T.; Kolsto, A. B.; Prydz,H.: A tight cluster of five unrelated human genes on chromosome 16q22.1. *Hum.Molec. Genet.* 2: 1589–1595, 1993.
- [73202] 11532.Pellegrino, C. M.; Rybicki, A. C.; Musto, S.; Nagel, R. L.; Schwartz,R. S.: Molecular identification and expression of erythroid K:Clcotransporter in human and mouse erythroleukemic cells. *Blood CellsMolec. Dis.* 24: 31–40, 1998.
- [73203] 11533.Hammond, E. M.; Brunet, C. L.; Johnson, G. D.; Parkhill, J.; Milner,A. E.; Brady, G.; Gregory, C. D.; Grand, R. J. A.: Homology between a human apoptosis specific protein and the product of APG5, a gene involved in autophagy in yeast. *FEBS Lett.* 425: 391–395, 1998.
- [73204] 11534.Schmeiser, K.; Armstrong, S.; Hammond, E. M.; Grand, R. J. A.: Assignment of the yeast APG5 human homologue APG5L to chromosome band 6q21 by fluorescence in situ hybridisation. *Cytogenet. Cell Genet.* 87:213–214, 1999.
- [73205] 11535.Nakayama, M.; Nakajima, D.; Nagase, T.; Nomura, N.; Seki, N.; Ohara,O.: Identification of high-molecular-weight proteins with multiple EGF-like motifs by motif-



trap screening. *Genomics* 51: 27–34, 1998.

[73206] 11536.Wu, Q.; Maniatis, T.: Large exons encoding multiple ectodomains are a characteristic feature of protocadherin genes. *Proc. Nat. Acad.Sci.* 97: 3124–3129, 2000.

[73207] 11537.Falcon–Perez, J. M.; Starcevic, M.; Gautam, R.; Dell'Angelica, E. C.: BLOC–1, a novel complex containing the pallidin and muted proteins involved in the biogenesis of melanosomes and platelet–dense granules. *J. Biol. Chem.* 277: 28191–28199, 2002.

[73208] 11538.Huang, L.: Personal Communication. San Francisco, Calif. 2/3/2000.

[73209] 11539.Huang, L.; Kuo, Y.–M.; Gitschier, J.: The pallid gene encodes a novel, syntaxin 13–interacting protein involved in platelet storage pool deficiency. *Nature Genet.* 23: 329–332, 1999.

[73210] 11540.Paine–Saunders, S.; Viviano, B. L.; Saunders, S.: GPC6, a novel member of the glypican gene family, encodes a product structurally related to GPC4 and is colocalized with GPC5 on human chromosome 13. *Genomics* 57:455–458, 1999.

[73211] 11541.Veugelers, M.; De Cat, B.; Ceulemans, H.; Bruystens, A. M.; Coomans, C.; Durr, J.; Vermeesch, J.; Marynen, P.; David, G.: Glypican–6, a new member of the glypican

family of cell surface heparan sulfate proteoglycans. *J. Biol. Chem.* 274: 26968–26977, 1999.

- [73212] 11542. Takei, Y.; Sasaki, S.; Fujiwara, T.; Takahashi, E.; Muto, T.; Nakamura, Y.: Molecular cloning of a novel gene similar to myeloid antigen CD33 and its specific expression in placenta. *Cytogenet. Cell Genet.* 78:295–300, 1997.
- [73213] 11543. Jacobs, S.; Schurmann, A.; Becker, W.; Bockers, T. M.; Copeland, N. G.; Jenkins, N. A.; Joost, H.-G.: The mouse ADP-ribosylation factor-like 4 gene: two separate promoters direct specific transcription in tissues and testicular germ cell. *Biochem. J.* 335: 259–265, 1998.
- [73214] 11544. Bootcov, M. R.; Bauskin, A. R.; Valenzuela, S. M.; Moore, A. G.; Bansal, M.; He, X. Y.; Zhang, H. P.; Donnellan, M.; Mahler, S.; Pryor, K.; Walsh, B. J.; Nicholson, R. C.; Fairlie, W. D.; Por, S. B.; Robbins, J. M.; Breit, S. N.: MIC-1, a novel macrophage inhibitory cytokine, is a divergent member of the TGF- $\beta$  superfamily. *Proc. Nat. Acad. Sci.* 94: 11514–11519, 1997.
- [73215] 11545. Bottner, M.; Laaff, M.; Schechinger, B.; Rappold, G.; Unsicker, K.; Suter-Crazzolara, C.: Characterization of the rat, mouse, and human genes of growth/differentiation factor-15/macrophage inhibiting cytokine-1

(GDF-15/MIC-1). *Gene* 237: 105-111, 1999.

[73216] 11546. Brown, D. A.; Breit, S. N.; Buring, J.; Fairlie, W. D.; Bauskin, A. R.; Liu, T.; Ridker, P. M.: Concentration in plasma of macrophageinhibitory cytokine-1 and risk of cardiovascular events in women: a nested case-control study. *Lancet* 359: 2159-2163, 2002.

[73217] 11547. Fairlie, W. D.; Russell, P. K.; Wu, W. M.; Moore, A. G.; Zhang, H.-P.; Brown, P. K.; Bauskin, A. R.; Breit, S. N.: Epitope mapping of the transforming growth factor-beta superfamily protein, macrophageinhibitory cytokine-1 (MIC-1): identification of at least five distinct epitope specificities. *Biochemistry* 40: 65-73, 2001.

[73218] 11548. Hromas, R.; Hufford, M.; Sutton, J.; Xu, D.; Li, Y.; Lu, L.: PLAB, a novel placental bone morphogenetic protein. *Biochim. Biophys. Acta* 1354: 40-44, 1997.

[73219] 11549. Moore, A. G.; Brown, D. A.; Fairlie, W. D.; Bauskin, A. R.; Brown, P. K.; Munier, M. L. C.; Russell, P. K.; Salamonsen, L. A.; Wallace, E. M.; Breit, S. N.: The transforming growth factor-beta superfamily cytokine macrophage inhibitory cytokine-1 is present in high concentrations in the serum of pregnant women. *J. Clin. Endocr. Metab.* 85: 4781-4788, 2000.

[73220] 11550. Paralkar, V. M.; Vail, A. L.; Grasser, W. A.; Brown, T.

A.; Xu, H.; Vukicevic, S.; Ke, H. Z.; Qi, H.; Owen, T. A.; Thompson, D. D.: Cloning and characterization of a novel member of the transforming growth factor- $\beta$ /bone morphogenetic protein family. *J. Biol. Chem.* 273:13760–13767, 1998.

- [73221] 11551. Yokoyama–Kobayashi, M.; Saeki, M.; Sekine, S.; Kato, S.: Human cDNA encoding a novel TGF- $\beta$  super-family protein highly expressed in placenta. *J. Biochem.* 122: 622–626, 1997.
- [73222] 11552. Conklin, D. C.; Rixon, M. W.; Kuestner, R. E.; Maurer, M. F.; Whitmore, T. E.; Millar, R. P.: Cloning and gene expression of a novel human ribonucleoprotein. *Biochim. Biophys. Acta* 1492: 465–469, 2000.
- [73223] 11553. Kataoka, N.; Yong, J.; Kim, V. N.; Velazquez, F.; Perkinson, R. A.; Wang, F.; Dreyfuss, G.: Pre-mRNA splicing imprints mRNA in the nucleus with a novel RNA-binding protein that persists in the cytoplasm. *Molec. Cell* 6: 673–682, 2000.
- [73224] 11554. Salicioni, A. M.; Xi, M.; Vanderveer, L. A.; Balsara, B.; Testa, J. R.; Dunbrack, R. L., Jr.; Godwin, A. K.: Identification and structural analysis of human RBM8A and RBM8B: two highly conserved RNA-binding motif proteins that interact with OVCA1, a candidate tumor suppressor.

Genomics 69:54–62, 2000.

[73225] 11555.Zhao, X.-F.; Nowak, N. J.; Shows, T. B.; Aplan, P. D.: MAGOH interacts with a novel RNA-binding protein. Genomics 63: 145–148, 2000.

[73226] 11556.Fischle, W.; Emiliani, S.; Hendzel, M. J.; Nagase, T.; Nomura, N.; Voelter, W.; Verdin, E.: A new family of human histone deacetylases related to *Saccharomyces cerevisiae* HDA1p. J. Biol. Chem. 274: 11713–11720, 1999.

[73227] 11557.Pazin, M. J.; Kadonaga, J. T.: What's up and down with histone deacetylation and transcription? Cell 89: 325–328, 1997.

[73228] 11558.Zhou, L.-J.; Tedder, T. F.: Human blood dendritic cells selectively express CD83, a member of the immunoglobulin superfamily. J. Immun. 154:3821–3835, 1995.

[73229] 11559.Zhou, L. J.; Schwarting, R.; Smith, H. M.; Tedder, T. F.: A novel cell-surface molecule expressed by human interdigitating reticulum cells, Langerhans cells, and activated lymphocytes is a new member of the Ig superfamily. J. Immun. 149: 735–742, 1992.

[73230] 11560.Bost-Usinger, L.; Chen, R. J.; Hillman, D.; Park, P.; Burnside, B.: Multiple kinesin family members expressed in teleost retina and RPE include a novel C-terminal kinesin.

Exp. Eye Res. 64: 781–794,1997.

- [73231] 11561.Hoang, E. H.; Whitehead, J. L.; Dose, A. C.; Burnside, B.: Cloning of a novel C-terminal kinesin (KIFC3) that maps to human chromosome 16q13–q21 and thus is a candidate gene for Bardet–Biedl syndrome. Genomics 52:219–222, 1998.
- [73232] 11562.Norman, D. A. M.; Barton, P. J. R.: Isolation, sequence, and chromosomal localisation of the human I(kappa)BR gene. Ann. Hum. Genet. 64:15–23, 2000.
- [73233] 11563.Ray, P.; Zhang, D.–H.; Elias, J. A.; Ray, A.: Cloning of a differentially expressed I–kappa–B–related protein. J. Biol. Chem. 270: 10680–10685,1995.
- [73234] 11564.Uchiumi, T.; Hinoshita, E.; Haga, S.; Nakamura, T.; Tanaka, T.;Toh, S.; Furukawa, M.; Kawabe, T.; Wada, M.; Kagotani, K.; Okumura,K.; Kohno, K.; Akiyama, S.; Kuwano, M.: Isolation of a novel human canalicular multi-specific organic anion transporter, cMOAT2/MRP3, and its expression in cisplatin-resistant cancer cells with decreased ATP-dependent drug transport. Biochem. Biophys. Res. Commun. 252:103–110, 1998.
- [73235] 11565.Bruhn, S. L.; Pil, P. M.; Essigmann, J. M.; Housman, D. E.; Lippard,S. J.: Isolation and characterization of human cDNA clones encoding a high mobility group box pro–

tein that recognizes structural distortions to DNA caused by binding of the anticancer agent cisplatin. *Proc. Nat. Acad. Sci.* 89: 2307–2311, 1992.

- [73236] 11566. Orphanides, G.; LeRoy, G.; Chang, C.-H.; Luse, D. S.; Reinberg, D.: FACT, a factor that facilitates transcript elongation through nucleosomes. *Cell* 92: 105–116, 1998.
- [73237] 11567. Acker, J.; Wintzerith, M.; Vigneron, M.; Keding, C.: A 14.4 kDa acidic subunit of human RNA polymerase II with a putative leucine-zipper. *DNASEq.* 4: 329–331, 1994.
- [73238] 11568. Pusch, C.; Wang, Z.; Roe, B.; Blin, N.: Genomic structure of the RNA polymerase II small subunit (hRPB14.4) locus (POLRF) (sic) and mapping to 22q13.1 by sequence identity. *Genomics* 34: 440–442, 1996.
- [73239] 11569. Caira, F.; Antonson, P.; Pelto-Huikko, M.; Treuter, E.; Gustafsson, J.-A.: Cloning and characterization of RAP250, a novel nuclear receptor coactivator. *J. Biol. Chem.* 275: 5308–5317, 2000.
- [73240] 11570. Guan, X. Y.; Xu, J.; Anzick, S. L.; Zhang, H.; Trent, J. M.; Meltzer, P. S.: Hybrid selection of transcribed sequences from microdissected DNA: isolation of genes within amplified region at 20q11–q13.2 in breast cancer. *Cancer Res.* 56: 3446–3450, 1996.

- [73241] 11571.Ko, L.; Cardona, G. R.; Chin, W. W.: Thyroid hormone receptor-binding protein, an LXXLL motif-containing protein, functions as a general coactivator. *Proc. Nat. Acad. Sci.* 97: 6212–6217, 2000.
- [73242] 11572.Lee, S.-K.; Anzick, S. L.; Choi, J.-E.; Bubendorf, L.; Guan, X.-Y.; Jung, Y.-K.; Kallioniemi, O. P.; Kononen, J.; Trent, J. M.; Azorsa, D.; Jhun, B.-H.; Cheong, J. H.; Lee, Y. C.; Meltzer, P. S.; Lee, J.W.: A nuclear factor, ASC-2, as a cancer-amplified transcriptional coactivator essential for ligand-dependent transactivation by nuclear receptors in vivo. *J. Biol. Chem.* 274: 34283–34293, 1999.
- [73243] 11573.Mahajan, M. A.; Samuels, H. H: A new family of nuclear receptor coregulators that integrate nuclear receptor signaling through CREB-binding protein. *Molec. Cell. Biol.* 20: 5048–5063, 2000.
- [73244] 11574.Zhu, Y.; Kan, L.; Qi, C.; Kanwar, Y. S.; Yeldandi, A. V.; Rao, M. S.; Reddy, J. K.: Isolation and characterization of peroxisome proliferator-activated receptor (PPAR) interacting protein (PRIP) as a coactivator for PPAR. *J. Biol. Chem.* 275: 13510–13516, 2000.
- [73245] 11575.Glockner, G.; Scherer, S.; Schattevoy, R.; Borigt, A.; Weber, J.; Tsui, L.-C.; Rosenthal, A.: Large-scale sequencing of two regions in human chromosome 7q22:



analysis of 650 kb of genomic sequence around the EPO and CUTL1 loci reveals 17 genes. *Genome Res.* 8: 1060–1073, 1998.

[73246] 11576. Yokouchi, M.; Suzuki, R.; Masuhara, M.; Komiya, S.; Inoue, A.; Yoshimura, A.: Cloning and characterization of APS, an adaptor molecule containing PH and SH2 domains that is tyrosine phosphorylated upon B-cell receptor stimulation. *Oncogene* 15: 7–15, 1997.

[73247] 11577. Still, I. H.; Hamilton, M.; Vince, P.; Wolfman, A.; Cowell, J.K.: Cloning of TACC1, an embryonically expressed, potentially transforming coiled coil containing gene, from the 8p11 breast cancer amplicon. *Oncogene* 18:4032–4038, 1999.

[73248] 11578. Still, I. H.; Vince, P.; Cowell, J. K.: The third member of the transforming acidic coiled coil-containing gene family, TACC3, maps to 4p16, close to translocation breakpoints in multiple myeloma, and is upregulated in various cancer cell lines. *Genomics* 58: 165–170, 1999.

[73249] 11579. Burmester, T.; Weich, B.; Reinhardt, S.; Hankeln, T.: A vertebrate globin expressed in the brain. *Nature* 407: 520–523, 2000.

[73250] 11580. Moens, L.; Dewilde, S.: Globins in the brain. *Nature* 407: 461–462, 2000.

- [73251] 11581.Scott, A. F.: Personal Communication. Baltimore, Md. 9/28/2000.
- [73252] 11582.Vilches, C.; Gardiner, C. M.; Parham, P.: Gene structure and promoter variation of expressed and nonexpressed variants of the KIR2DL5 gene. *J.Immun.* 165: 6416–6421, 2000.
- [73253] 11583.Vilches, C.; Rajalingam, R.; Uhrberg, M.; Gardiner, C. M.; Young, N. T.; Parham, P.: KIR2DL5, a novel killer-cell receptor with a D0–D2 configuration of Ig-like domains. *J. Immun.* 164: 5797–5804, 2000.
- [73254] 11584.Bates, E. E. M.; Fournier, N.; Garcia, E.; Valladeau, J.; Durand, I.; Pin, J.-J.; Zurawski, S. M.; Patel, S.; Abrams, J. S.; Lebecque, S.; Garrone, P.; Saeland, S.: APCs express DCIR, a novel C-type lectin surface receptor containing an immunoreceptor tyrosine-based inhibitory motif. *J. Immun.* 163: 1973–1983, 1999.
- [73255] 11585.Huang, X.; Yuan, Z.; Chen, G.; Zhang, M.; Zhang, W.; Yu, Y.; Cao, X.: Cloning and characterization of a novel ITIM containing lectin-like immunoreceptor LLIR and its two transmembrane region deletion variants. *Biochem.Biophys. Res. Commun.* 281: 131–140, 2001.
- [73256] 11586.Seeger, M.; Kraft, R.; Ferrell, K.; Bech-Otschir, D.; Dumdey, R.; Schade, R.; Gordon, C.; Naumann, M.; Dubiel,

W.: A novel protein complex involved in signal transduction possessing similarities to 26S proteasome subunits. FASEB J. 12: 469–478, 1998.

- [73257] 11587. Ann, K.; Kowalchuk, J. A.; Loyet, K. M.; Martin, T. F. J.: Novel  $\text{Ca}^{2+}$ -binding protein (CAPS) related to UNC-31 required for  $\text{Ca}^{2+}$ -activated exocytosis. J. Biol. Chem. 272: 19637–19640, 1997.
- [73258] 11588. Berwin, B.; Floor, E.; Martin, T. F. J.: CAPS (mammalian UNC-31) protein localizes to membranes involved in dense-core vesicle exocytosis. Neuron 21: 137–145, 1998.
- [73259] 11589. Hirose, M.; Nagase, T.; Ishikawa, K.; Kikuno, R.; Nomura, N.; Ohara, O.: Characterization of cDNA clones selected by the GeneMark analysis from size-fractionated cDNA libraries from human brain. DNA Res. 6: 329–336, 1999.
- [73260] 11590. Walent, J. H.; Porter, B. W.; Martin, T. F. J.: A novel 145 kDa brain cytosolic protein reconstitutes  $\text{Ca}^{2+}$ -regulated secretion in permeable neuroendocrine cells. Cell 70: 765–775, 1992.
- [73261] 11591. Veldhuis, B.; Spruit, L.; Dauwerse, H. G.; Breuning, M. H.; Peters, D. J.: Genes homologous to the autosomal dominant polycystic kidney disease genes (PKD1 and

PKD2). *Europ. J. Hum. Genet.* 7: 860–872,1999.

- [73262] 11592.Hughes, J.; Ward, C. J.; Aspinwall, R.; Butler, R.; Harris, P.C.: Identification of a human homologue of the sea urchin receptor for egg jelly: a polycystic kidney disease-like protein. *Hum. Molec.Genet.* 8: 543–549, 1999.
- [73263] 11593.Curtis, B. M.; Scharnowske, S.; Watson, A. J.: Sequence and expression of a membrane-associated C-type lectin that exhibits CD4-independent binding of human immunodeficiency virus envelope glycoprotein gp120. *Proc.Nat. Acad. Sci.* 89: 8356–8360, 1992.
- [73264] 11594.Feinberg, H.; Mitchell, D. A.; Drickamer, K.; Weis, W. I.: Structural basis for selective recognition of oligosaccharides by DC-SIGN and DC-SIGNR. *Science* 294: 2163–2166, 2001.
- [73265] 11595.Geijtenbeek, T. B. H.; Kwon, D. S.; Torensma, R.; van Vliet, S.J.; van Duijnhoven, G. C. F.; Middel, J.; Cornelissen, I. L. M. H.A.; Nottet, H. S. L. M.; KewalRamani, V. N.; Littman, D. R.; Figdor, C. G.; van Kooyk, Y.: DC-SIGN, a dendritic cell-specific HIV-1-binding protein that enhances trans-infection of T cells. *Cell* 100: 587–597,2000.
- [73266] 11596.Geijtenbeek, T. B. H.; Torensma, R.; van Vliet, S. J.; van Duijnhoven, G. C. F.; Adema, G. J.; van Kooyk, Y.; Figdor, C. G.: Identification of DC-SIGN, a novel dendritic cell-

specific ICAM-3 receptor that supports primary immune responses. *Cell* 100: 575-585, 2000.

- [73267] 11597. Battini, J.-L.; Rasko, J. E. J.; Miller, A. D.: A human cell-surface receptor for xenotropic and polytropic murine leukemia viruses: possible role in G protein-coupled signal transduction. *Proc. Nat. Acad. Sci.* 96:1385-1390, 1999.
- [73268] 11598. Levy, J. A.: Xenotropism: the elusive viral receptor finally uncovered. *Proc. Nat. Acad. Sci.* 96: 802-804, 1999.
- [73269] 11599. Taylor, C. S.; Nouri, A.; Lee, C. G.; Kozak, C.; Kabat, D.: Cloning and characterization of a cell surface receptor for xenotropic and polytropic murine leukemia viruses. *Proc. Nat. Acad. Sci.* 96: 927-932, 1999.
- [73270] 11600. Yang, Y.-L.; Guo, L.; Xu, S.; Holland, C. A.; Kitamura, T.; Hunter, K.; Cunningham, J. M.: Receptors for polytropic and xenotropic mouse leukemia viruses encoded by a single gene at Rmc1. *Nature Genet.* 21:216-219, 1999.
- [73271] 11601. Aksoy, S.; Raftogianis, R.; Weinshilboum, R.: Human histamine N-methyltransferase gene: structural characterization and chromosomal localization. *Biochem. Biophys. Res. Commun.* 219: 548-554, 1996.
- [73272] 11602. Preuss, C. V.; Wood, T. C.; Szumlanski, C. L.; Raftogianis, R. B.; Otterness, D. M.; Girard, B.; Scott, M. C.;

Weinshilboum, R. M.: Human histamine N-methyltransferase pharmacogenetics: common genetic polymorphisms that alter activity. *Molec. Pharm.* 53: 708–717, 1998.

[73273] 11603. Price, R. A.; Scott, M. C.; Weinshilboum, R. M.: Genetic segregation analysis of red blood cell (RBC) histamine N-methyltransferase (HNMT) activity. *Genet. Epidemiol.* 10: 123–131, 1993.

[73274] 11604. Scott, M. C.; Van Loon, J. A.; Weinshilboum, R. M.: Pharmacogenetics of N-methylation: heritability of human erythrocyte histamine N-methyltransferase activity. *Clin. Pharm. Ther.* 43: 256–262, 1988.

[73275] 11605. Yamauchi, K.; Sekizawa, K.; Suzuki, H.; Nakazawa, H.; Ohkawara, Y.; Katayose, D.; Ohtsu, H.; Tamura, G.; Shibahara, S.; Takemura, M.; Maeyama, K.; Watanabe, T.; Sasaki, H.; Shirato, K.; Takishima, T.: Structure and function of human histamine N-methyltransferase: critical enzyme in histamine metabolism in airway. *Am. J. Physiol.* 267: L342–L349, 1994.

[73276] 11606. Yan, L.; Galinsky, R. E.; Bernstein, J. A.; Liggett, S. B.; Weinshilboum, R. M.: Histamine N-methyltransferase pharmacogenetics: association of a common functional polymorphism with asthma. *Pharmacogenetics*

10:261–266, 2000.

[73277] 11607. Bartoloni, L.; Blouin, J.-L. C.; Sainsbury, A. J.; Gos, A.; Morris, M. A.; Affara, N. A.; DeLozier-Blanchet, C. D.; Antonarakis, S. E.: Assignment of the human dynein heavy chain gene DNAH17L to human chromosome 17p12 by in situ hybridization and radiation hybrid mapping. *Cytogenet. Cell Genet.* 84: 188–189, 1999.

[73278] 11608. Bartoloni, L.; Blouin, J.-L.; Maiti, A. K.; Sainsbury, A.; Rossier, C.; Gehrig, C.; She, J.-X.; Marron, M. P.; Lander, E. S.; Meeks, M.; Chung, E.; Armengot, M.; Jorissen, M.; Scott, H. S.; DeLozier-Blanchet, C. D.; Gardiner, R. M.; Antonarakis, S. E.: Axonemal beta heavy chain dynein DNAH9: cDNA sequence, genomic structure, and investigation of its role in primary ciliary dyskinesia. *Genomics* 72: 21–33, 2001.

[73279] 11609. Maiti, A. K.; Mattei, M.-G.; Jorissen, M.; Volz, A.; Zeigler, A.; Bouvagnet, P.: Identification, tissue specific expression, and chromosomal localisation of several human dynein heavy chain genes. *Europ. J. Hum. Genet.* 8: 923–932, 2000.

[73280] 11610. Milisav, I.; Jones, M. H.; Affara, N. A.: Characterization of a novel human dynein-related gene that is specifically expressed in testis. *Mammalian Genome* 7: 667–672,

1996.

- [73281] 11611.Hart, M. J.; Callow, M. G.; Souza, B.; Polakis, P.: IQ-GAP1, acalmodulin-binding protein with a rasGAP-related domain, is a potentialeffector for cdc42Hs. EMBO J. 15: 2997-3005, 1996.
- [73282] 11612.Sugimoto, N.; Imoto, I.; Fukuda, Y.; Kurihara, N.; Kuroda, S.;Tanigami, A.; Kaibuchi, K.; Kamiyama, R.; Inazawa, J.: IQGAP1, anegative regulator of cell-cell adhesion, is upregulated by gene amplificationat 15q26 in gastric cancer cell lines HSC39 and 40A. J. Hum. Genet. 46:21-25, 2001.
- [73283] 11613.Weissbach, L.; Settleman, J.; Kalady, M. F.; Snijders, A. J.; Murthy,A. E.; Yan, Y.-X.; Bernards, A.: Identification of a human RasGAP-relatedprotein containing calmodulin-binding motifs. J. Biol. Chem. 269:20517-20521, 1994.
- [73284] 11614.Harrington, J. J.; Lieber, M. R.: Functional domains within FEN-1and RAD2 define a family of structure-specific endonucleases: implicationsfor nucleotide excision repair. Genes Dev. 8: 1344-1355, 1994.
- [73285] 11615.Kato, H.; Torigoe, T.: Radioimmunoassay for tumor antigen of humancervical squamous cell carcinoma. Cancer 40: 1621-1628, 1977.



- [73286] 11616.Kuwano, A.; Kondo, I.; Kishi, F.; Suminami, Y.; Kato, H.: Assignment of the squamous cell carcinoma antigen locus (SCC) to 18q21 by insitu hybridization. *Genomics* 30: 626, 1995.
- [73287] 11617.Suminami, Y.; Kishi, F.; Sekiguchi, K.; Kato, H.: Squamous cell carcinoma antigen is a new member of the serine protease inhibitors. *Biochem.Biophys. Res. Commun.* 181: 51–58, 1991.
- [73288] 11618.Barnes, R. C.; Worrall, D. M.: Identification of a novel human serpin gene; cloning sequencing and expression of leupin. *FEBS Lett.* 373:61–65, 1995.
- [73289] 11619.Schick, C.; Kamachi, Y.; Bartuski, A. J.; Cataltepe, S.; Schechter, N. M.; Pemberton, P. A.; Silverman, G. A.: Squamous cell carcinoma antigen 2 is a novel serpin that inhibits the chymotrypsin-like proteinases cathepsin G and mast cell chymase. *J. Biol. Chem.* 272: 1849–1855, 1997.
- [73290] 11620.Dahl, M. R.; Thiel, S.; Matsushita, M.; Fujita, T.; Willis, A.C.; Christensen, T.; Vorup-Jensen, T.; Jensenius, J. C.: MASP-3 and its association with distinct complexes of the mannan-binding lectin complement activation pathway. *Immunity* 15: 127–135, 2001.
- [73291] 11621.Endo, Y.; Sato, T.; Matsushita, M.; Fujita, T.: Exon structure of the gene encoding the human mannose-bind-

ing protein-associated serine protease light chain: comparison with complement C1r and C1s genes. *Int. Immun.* 8: 1355–1358, 1996.

[73292] 11622. Matsushita, M.; Thiel, S.; Jensenius, J. C.; Terai, I.; Fujita, T.: Proteolytic activities of two types of mannose-binding lectin-associated serine protease. *J. Immun.* 165: 2637–2642, 2000.

[73293] 11623. Sato, T.; Endo, Y.; Matsushita, M.; Fujita, T.: Molecular characterization of a novel serine protease involved in activation of the complement system by mannose-binding protein. *Int. Immun.* 6: 665–669, 1994.

[73294] 11624. Takada, F.; Seki, N.; Matsuda, Y.; Takayama, Y.; Kawakami, M.: Localization of the genes for the 100-kDa complement-activating component of C3a-reactive factor (CRARF and Crarf) to human 3q27–q28 and mouse 16B2–B3. *Genomics* 25: 757–759, 1995.

[73295] 11625. Takada, F.; Takayama, Y.; Hatsuse, H.; Kawakami, M.: A new member of the C1s family of complement proteins found in a bactericidal factor, C3a-reactive factor, in human serum. *Biochem. Biophys. Res. Commun.* 196: 1003–1009, 1993.

[73296] 11626. Takayama, Y.; Takada, F.; Nowatari, M.; Kawakami, M.; Matsuura, N.: Gene structure of the P100 serine-

protease component of the human Ra-reactive factor.

Molec. Immun. 36: 505–514, 1999.

[73297] 11627.Dessen, A.; Tang, J.; Schmidt, H.; Stahl, M.; Clark, J. D.; Seehra, J.; Somers, W. S.: Crystal structure of human cytosolic phospholipase A(2) reveals a novel topology and catalytic mechanism. Cell 97:349–360, 1999.

[73298] 11628.Hansen, L.; Urioste, S.; Petersen, H. V.; Jensen, J. N.; Eiberg, H.; Barbetti, F.; Serup, P.; Hansen, T.; Pedersen, O.: Missense mutations in the human insulin promoter factor-1 gene and their relation to maturity-onset diabetes of the young and late-onset type 2 diabetes mellitus in Caucasians. J. Clin. Endocr. Metab. 85: 1323–1326, 2000.

[73299] 11629.Johansson, K. A.; Grapin-Botton, A.: Development and diseases of the pancreas. Clin. Genet. 62: 14–23, 2002.

[73300] 11630.Jonsson, J.; Carlsson, L.; Edlund, T.; Edlund, H.: Insulin-promoter-factor 1 is required for pancreas development in mice. Nature 371: 606–609, 1994.

[73301] 11631.Leonard, J.; Peers, B.; Johnson, T.; Ferreri, I.; Lee, K.; Montminy, M. R.: Characterization of somatostatin transactivating factor-1, a novel homeobox factor that stimulates somatostatin expression in pancreatic islet cells. Molec. Endocr. 7: 1275–1283, 1993.

- [73302] 11632. Macfarlane, W. M.; Frayling, T. M.; Ellard, S.; Evans, J. C.; Allen, L. I. S.; Bulman, M. P.; Ayres, S.; Shepherd, M.; Clark, P.; Millward, A.; Demaine, A.; Wilkin, T.; Docherty, K.; Hattersley, A. T.: Missense mutations in the insulin promoter factor-1 gene predispose to type 2 diabetes. *J. Clin. Invest.* 104: R33–R39, 1999.
- [73303] 11633. Miller, C. P.; McGehee, R. E., Jr.; Habener, J. F.: IDX-1: a new homeodomain transcription factor expressed in rat pancreatic islets and duodenum that transactivates the somatostatin gene. *EMBO J.* 13: 1145–1156, 1994.
- [73304] 11634. Ohlsson, H.; Karlsson, K.; Edlund, T.: IPF1, a homeodomain-containing transactivator of the insulin gene. *EMBO J.* 12: 4251–4259, 1993.
- [73305] 11635. Sharma, S.; Jhala, U. S.; Johnson, T.; Ferreri, K.; Leonard, J.; Montminy, M.: Hormonal regulation of an islet-specific enhancer in the pancreatic homeobox gene STF-1. *Molec. Cell. Biol.* 17: 2598–2604, 1997.
- [73306] 11636. Stoffel, M.; Stein, R.; Wright, C. V. E.; Espinosa, R., III; LeBeau, M. M.; Bell, G. I.: Localization of human homeodomain transcription factor insulin promoter factor 1 (IPF1) to chromosome band 13q12.1. *Genomics* 28: 125–126, 1995.
- [73307] 11637. Stoffers, D. A.; Ferrer, J.; Clarke, W. L.; Habener, J.

F.: Early-onset type-II diabetes mellitus (MODY4) linked to IPF1. (Letter) *Nature Genet.* 17: 138–141, 1997.

[73308] 11638. Stoffers, D. A.; Stanojevic, V.; Habener, J. F.: Insulin promoter factor-1 gene mutation linked to early-onset type 2 diabetes mellitus directs expression of a dominant negative isoform. *J. Clin. Invest.* 102:232–241, 1998.

[73309] 11639. Watada, H.; Kajimoto, Y.; Kaneto, H.; Matsuoka, T.; Fujitani, Y.; Miyazaki, J.; Yamasaki, Y.: Involvement of the homeodomain-containing transcription factor PDX-1 in islet amyloid polypeptide gene transcription. *Biochem. Biophys. Res. Commun.* 229: 746–751, 1996.

[73310] 11640. Wright, N. M.; Metzger, D. L.; Borowitz, S. M.; Clarke, W. L.: Permanent neonatal diabetes mellitus and pancreatic exocrine insufficiency resulting from congenital pancreatic agenesis. *Am. J. Dis. Child.* 147:607–609, 1993.

[73311] 11641. Ashford, M. L. J.; Bond, C. T.; Blair, T. A.; Adelman, J. P.: Cloning and functional expression of a rat heart KATP channel. *Nature* 370:456–459, 1994.

[73312] 11642. Bond, C. T.; Pessia, M.; Xia, X.-M.; Lagrutta, A.; Kavanaugh, M. P.; Adelman, J. P.: Cloning and expression of a family of inward rectifier potassium channels. *Receptors Channels* 2: 183–191, 1994.

- [73313] 11643. Tucker, S. J.; James, M. R.; Adelman, J. P.: Assignment of K(ATP)-1, the cardiac ATP-sensitive potassium channel gene (KCNJ5), to human chromosome 11q24. *Genomics* 28: 127-128, 1995.
- [73314] 11644. Wickman, K.; Seldin, M. F.; Gendler, S. J.; Clapham, D. E.: Partial structure, chromosome localization, and expression of the mouse *Girk4* gene. *Genomics* 40: 395-401, 1997.
- [73315] 11645. Jenkins, N. A.: Personal Communication. Frederick, Md. 5/12/1997.
- [73316] 11646. Khurana, T. S.; Engle, E. C.; Bennett, R. R.; Silverman, G. A.; Selig, S.; Bruns, G. A. P.; Kunkel, L. M.: (CA) repeat polymorphism in the chromosome 18 encoded dystrophin-like protein. *Hum. Molec. Genet.* 3: 841 only, 1994.
- [73317] 11647. Eisenberg, I.; Avidan, N.; Potikha, T.; Hochner, H.; Chen, M.; Olender, T.; Barash, M.; Shemesh, M.; Sadeh, M.; Grabov-Nardini, G.; Shmlevich, I.; Friedmann, A.; Karpati, G.; Bradley, W. G.; Baumbach, L.; Lancet, D.; Ben Asher, E.; Beckmann, J. S.; Argov, Z.; Mitrani-Rosenbaum, S.: The UDP-N-acetylglucosamine 2-epimerase/N-acetylmannosamine kinase gene is mutated in recessive hereditary inclusion body myopathy.

NatureGenet. 29: 83–87, 2001.

- [73318] 11648.Metzinger, L.; Blake, D. J.; Squier, M. V.; Anderson, L. V. B.;Deconinck, A. E.; Nawrotzki, R.; Hilton–Jones, D.; Davies, K. E.:Dystrobrevin deficiency at the sarcolemma of patients with muscular dystrophy. Hum. Molec. Genet. 6: 1185–1191, 1997.
- [73319] 11649.Sadoulet–Puccio, H. M.; Feener, C. A.; Schaid, D. J.; Thibodeau,S. N.; Michels, V. V.; Kunkel, L. M.: The genomic organization of human dystrobrevin. Neurogenetics 1: 37–42, 1997.
- [73320] 11650.Sadoulet–Puccio, H. M.; Khurana, T. S.; Cohen, J. B.; Kunkel, L.M.: Cloning and characterization of the human homologue of a dystrophinrelated phosphoprotein found at the Torpedo electric organ post–synaptic membrane. Hum. Molec. Genet. 5: 489–496, 1996.
- [73321] 11651.Yoshida, M.; Hama, H.; Ishikawa–Sakurai, M.; Ima–mura, M.; Mizuno,Y.; Araishi, K.; Wakabayashi–Takai, E.; Noguchi, S.; Sasaoka, T.;Ozawa, E.: Biochemical evidence for association of dystrobrevin with the sarcoglycan–sar–cospan complex as a basis for understanding sarcogly–canopathy. Hum.Molec. Genet. 9: 1033–1040, 2000.
- [73322] 11652.Bauer, W. R.; Hayes, J. J.; White, J. H.; Wolffe, A. P.: Nucleosome structural changes due to acetylation. J.

Molec. Biol. 236: 685–690,1994.

- [73323] 11653.Furukawa, Y.; Kawakami, T.; Sudo, K.; Inazawa, J.; Matsumine, A.;Akiyama, T.; Nakamura, Y.: Isolation and mapping of a human gene(RPD3L1) that is homologous to RPD3, a transcription factor in *Saccharomyces cerevisiae*. Cytogenet. Cell Genet. 73: 130–133, 1996.
- [73324] 11654.Lee, D. Y.; Hayes, J. J.; Pruss, D.; Wolffe, A. P.: A positive role for histone acetylation in transcription factor access to nucleosomal DNA. Cell 72: 73–84, 1993.
- [73325] 11655.Luo, J.; Su, F.; Chen, D.; Shiloh, A.; Gu, W.: Deacetylation of p53 modulates its effect on cell growth and apoptosis. Nature 408:377–381, 2000.
- [73326] 11656.Magnaghi-Jaulin, L.; Groisman, R.; Naguibneva, I.; Robin, P.; Lorain, S.; Le Villain, J. P.; Troalen, F.; Trouche, D.; Harel-Bellan, A.: Retinoblastoma protein represses transcription by recruiting a histone deacetylase. Nature 391: 601–605, 1998.
- [73327] 11657.Taunton, J.; Hassig, C. A.; Schreiber, S. L.: A mammalian histone deacetylase related to the yeast transcriptional regulator Rpd3p. Science 272:408–411, 1996.
- [73328] 11658.Vidal, M.; Gaber, R. F.: RPD3 encodes a second factor required to achieve maximum positive and negative transcriptional states in *Saccharomyces cerevisiae*. Molec.



Cell. Biol. 11: 6317–6327, 1991.

- [73329] 11659.Vidal, M.; Strich, R.; Esposito, R. E.; Gaber, R. F.: RPD1 (SIN3/UME4)is required for maximal activation and repression of diverse yeastgenes. Molec. Cell. Biol. 11: 6306–6316, 1991.
- [73330] 11660.Wade, P. A.: Transcriptional control at regulatory checkpointsby histone deacetylases: molecular connections between cancer andchromatin. Hum. Molec. Genet. 10: 693–698, 2001.
- [73331] 11661.Wolffe, A. P.: Histone deacetylase: a regulator of transcription. Science 272:371–372, 1996.
- [73332] 11662.Zhang, Y.; Sun, Z.–W.; Iratni, R.; Erdjument–Bromage, H.; Tempst,P.; Hampsey, M.; Reinberg, D.: SAP30, a novel protein conserved betweenhuman and yeast, is a component of a histone deacetylase complex. Molec.Cell 1: 1021–1031, 1998.
- [73333] 11663.Chappell, T. G.; Welch, W. J.; Schlossman, D. M.; Palter, K. B.;Schlesinger, M. J.; Rothman, J. E.: Uncoating ATPase is a memberof the 70 kilodalton family of stress proteins. Cell 45: 3–13, 1986.
- [73334] 11664.Dworniczak, B.; Mirault, M.–E.: Structure and expression of ahuman gene coding for a 71 kd heat shock 'cognate' protein. NucleicAcids Res. 15: 5181–5197,

1987.

- [73335] 11665.Tavaria, M.; Gabriele, T.; Anderson, R. L.; Mirault, M.-E.; Baker,E.; Sutherland, G.; Kola, I.: Localization of the gene encoding thehuman heat shock cognate protein, HSP73, to chromosome 11. *Genomics* 29:266–268, 1995.
- [73336] 11666.Ungewickell, E.: The 70-kd mammalian heat shock proteins are structurallyand functionally related to the un-coating protein that releases clathrintriskelia from coated vesicles. *EMBO J.* 4: 3385–3391, 1985.
- [73337] 11667.Camoretti–Mercado, B.; Forsythe, S. M.; LeBeau, M. M.; Espinosa,R., III; Vieira, J. E.; Halayko, A. J.; Willadsen, S.; Kurtz, B.;Ober, C.; Evans, G. A.; Thweatt, R.; Shapiro, S.; Niu, Q.; Qin, Y.;Padrid, P. A.; Solway, J.: Expression and cytogenetic localizationof the human SM22 gene (TAGLN). *Genomics* 49: 452–457, 1998.
- [73338] 11668.Lawson, D.; Harrison, M.; Shapland, C.: Fibroblast transgelinand smooth muscle SM22–alpha are the same protein, the expressionof which is down–regulated in many cell lines. *Cell Motil. Cytoskeleton* 38:250–257, 1997.
- [73339] 11669.Lees–Miller, J. P.; Heeley, D. H.; Smillie, L. B.; Kay, C. M.:Isolation and characterization of an abundant and novel 22–kDa protein(SM22) from chicken gizzard smooth

muscle. J. Biol. Chem. 262: 2988–2993, 1987.

[73340] 11670. Nishida, W.; Kitami, Y.; Abe, M.; Hiwada, K.: Gene cloning and nucleotide sequence of SM22- $\alpha$  from the chicken gizzard smooth muscle. Biochem. Int. 23: 663–668, 1991.

[73341] 11671. Shapland, C.; Hsuan, J. J.; Totty, N. F.; Lawson, D.: Purification and properties of transgelin: a transformation and shape change sensitive actin-gelling protein. J. Cell Biol. 121: 1065–1073, 1993.

[73342] 11672. Shapland, C.; Lowings, P.; Lawson, D.: Identification of new actin-associated polypeptides that are modified by viral transformation and changes in cell shape. J. Cell Biol. 107: 153–161, 1988.

[73343] 11673. Solway, J.; Seltzer, J.; Samaha, F. F.; Kim, S.; Alger, L. E.; Niu, Q.; Morrissey, E. E.; Ip, H. S.; Parmacek, M. S.: Structure and expression of a smooth muscle cell-specific gene, SM22- $\alpha$ . J. Biol. Chem. 270: 13460–13469, 1995.

[73344] 11674. Stanier, P.; Abu-Hayyeh, S.; Murdoch, J. N.; Eddleston, J.; Copp, A. J.: Paralogous Sm22- $\alpha$  (Tagln) genes map to mouse chromosomes 1 and 9: further evidence for a paralogous relationship. Genomics 51: 144–147, 1998.

[73345] 11675. Thweatt, R.; Lumpkin, C. K., Jr.; Goldstein, S.: A

novel gene encoding a smooth muscle protein is overexpressed in senescent human fibroblasts. *Biochem. Biophys. Res. Commun.* 187: 1–7, 1992.

[73346] 11676. Tamanini, F.; Bontekoe, C.; Bakker, C. E.; van Unen, L.; Anar, B.; Willemsen, R.; Yoshida, M.; Galjaard, H.; Oostra, B. A.; Hoogeveen, A. T.: Different targets for the fragile X-related proteins revealed by their distinct nuclear localizations. *Hum. Molec. Genet.* 8: 863–869, 1999.

[73347] 11677. Tamanini, F.; Kirkpatrick, L. L.; Schonkeren, J.; van Unen, L.; Bontekoe, C.; Bakker, C.; Nelson, D. L.; Galjaard, H.; Oostra, B. A.; Hoogeveen, A. T.: The fragile X-related proteins FXR1P and FXR2P contain a functional nucleolar-targeting signal equivalent to the HIV-1 regulatory proteins. *Hum. Molec. Genet.* 9: 1487–1493, 2000.

[73348] 11678. Radice, P.; Pensotti, V.; Jones, C.; Perry, H.; Pierotti, M. A.; Tunnacliffe, A.: The human archain gene, ARCN1, has highly conserved homologs in rice and *Drosophila*. *Genomics* 26: 101–106, 1995.

[73349] 11679. Tunnacliffe, A.; van de Vrugt, H.; Pensotti, V.; Radice, P.: The coatamer protein delta-COP, encoded by the archain gene, is conserved across diverse eukaryotes. *Mammalian Genome* 7: 784–786, 1996.

[73350] 11680. Hawtin, S. R.; Wesley, V. J.; Parslow, R. A.; Simms,

J.; Miles, A.; McEwan, K.; Wheatley, M.: A single residue (arg46) located within the N-terminus of the V(1a) vasopressin receptor is critical for binding vasopressin but not peptide or nonpeptide antagonists. *Molec. Endocr.* 16:600–609, 2002.

[73351] 11681. Morel, A.; O'Carroll, A.-M.; Brownstein, M. J.; Lolait, S. J.: Molecular cloning and expression of a rat V1a arginine vasopressin receptor. *Nature* 356: 523–526, 1992.

[73352] 11682. Thibonnier, M.; Auzan, C.; Madhun, Z.; Wilkins, P.; Berti-Mattera, L.; Clauser, E.: Molecular cloning, sequencing, and functional expression of a cDNA encoding the human V1a vasopressin receptor. *J. Biol. Chem.* 269:3304–3310, 1994.

[73353] 11683. Thibonnier, M.; Graves, M. K.; Wagner, M. S.; Auzan, C.; Clauser, E.; Willard, H. F.: Structure, sequence, expression, and chromosomal localization of the human V(1a) vasopressin receptor gene. *Genomics* 31:327–334, 1996.

[73354] 11684. Young, L. J.; Nilsen, R.; Waymire, K. G.; MacGregor, G. R.; Insel, T. R.: Increased affiliative response to vasopressin in mice expressing the V(1A) receptor from a monogamous vole. *Nature* 400: 766–768, 1999.

[73355] 11685. Langer, L. O., Jr.; Cervenka, J.; Camargo, M.: A se-

vere autosomalrecessive acromesomelic dysplasia, the Hunter–Thompson type, and comparison with the Grebe type. *Hum. Genet.* 81: 323–328, 1989.

[73356] 11686.Robin, N. H.: Personal Communication. Cleveland, Ohio 11/3/1997.

[73357] 11687.Runner, M. N.: Linkage of brachypodism: a new member of linkagegroup V of the house mouse. *J. Hered.* 50: 81–84, 1959.

[73358] 11688.Thomas, J. T.; Lin, K.; Nandedkar, M.; Camargo, M.; Cervenka,J.; Luyten, F. P.: A human chondrodysplasia due to a mutation in a TGF–beta superfamily member. *Nature Genet.* 12: 315–317, 1996.

[73359] 11689.Tsumaki, N.; Tanaka, K.; Arikawa–Hirasawa, E.; Nakase, T.; Kimura,T.; Thomas, J. T.; Ochi, T.; Luyten, F. P.; Yamada, Y.: Role of CDMP–1 in skeletal morphogenesis: promotion of mesenchymal cell recruitment and chondrocyte differentiation. *J. Cell Biol.* 144: 161–173, 1999.

[73360] 11690.Huh, T.–L.; Kim, Y.–O.; Oh, I.–U.; Song, B. J.; Inazawa, J.: Assignment of the human mitochondrial NAD(+)-specific isocitrate dehydrogenase alpha subunit (IDH3A) gene to 15q25.1–q25.2 by in situ hybridization. *Genomics* 31:295–296, 1996.

[73361] 11691.Kim, Y.–O.; Oh, I.–U.; Park, H.–S.; Jeng, J.; Song, B.

J.; Huh, T.-L.: Characterization of a cDNA clone for human NAD(+)-specific isocitrate dehydrogenase alpha-subunit and structural comparison with its isoenzymes from different species. *Biochem. J.* 308: 63–68, 1995.

[73362] 11692. Amann, J.; Kidd, V. J.; Lahti, J. M.: Characterization of putative human homologues of the yeast chromosome transmission fidelity gene, CHL1. *J. Biol. Chem.* 272: 3823–3832, 1997.

[73363] 11693. Amann, J.; Valentine, M.; Kidd, V. J.; Lahti, J. M.: Localization of Chl1-related helicase genes to human chromosome regions 12p11 and 12p13: similarity between parts of these genes and conserved human telomeric-associated DNA. *Genomics* 32: 260–265, 1996.

[73364] 11694. Frank, S.; Werner, S.: The human homologue of the yeast CHL1 gene is a novel keratinocyte growth factor-regulated gene. *J. Biol. Chem.* 271: 24337–24340, 1996.

[73365] 11695. Barnes, L. D.; Garrison, P. N.; Siprashvili, Z.; Guranowski, A.; Robinson, A. K.; Ingram, S. W.; Croce, C. M.; Ohta, M.; Huebner, K.: Fhit, a putative tumor suppressor in humans, is a dinucleoside 5-prime,5-tripleprime-P(1),P(3)-triphosphate hydrolase. *Biochemistry* 35: 11529–11535, 1996.

[73366] 11696. Bernar, J.; Funderburk, S. J.; Sparkes, R. S.: The in-

ducible fragile site on chromosome 3. (Letter) Hum. Genet. 66: 373 only, 1984.

- [73367] 11697 Jin, C.-J.; Miners, J. O.; Lillywhite, K. J.; Mackenzie, P. I.: cDNA cloning and expression of two new members of the human liver UDP-glucuronosyltransferase 2B subfamily. Biochem. Biophys. Res. Commun. 194: 496–503, 1993.
- [73368] 11698 Krasnewich, D.; Kozak, C. A.; Nebert, D. W.; Mackenzie, P. I.: Localization of UDP glucuronosyltransferase gene(s) on mouse chromosome
- [73369] 11699 Somat. Cell Molec. Genet. 13: 179–182, 1987.5. Monaghan, G.; Clarke, D. J.; Povey, S.; See, C. G.; Boxer, M.; Burchell, B.: Isolation of a human YAC contig encompassing a cluster of UGT2 genes and its regional localization to chromosome 4q13. Genomics 23: 496–499, 1994.
- [73370] 11700 Monaghan, G.; Povey, S.; Burchell, B.; Boxer, M.: Localization of a bile acid UDP-glucuronosyltransferase gene (UGT2B) to chromosome 4 using the polymerase chain reaction. Genomics 13: 908–909, 1992.
- [73371] 11701 Riedy, M.; Wang, J. Y.; Miller, A. P.; Buckler, A.; Hall, J.; Guida, M.: Genomic organization of the UGT2b gene cluster on human chromosome 4q13. Pharmacogenetics 10: 251–260, 2000.
- [73372] 11702 Buc-Caron, M. H.; Condamine, H.; Kerjaschki, D.:



Rat Heymann nephritis antigen is closely related to brushin, a glycoprotein present in early mouse embryo epithelia. *Ann. Inst. Pasteur/Immun.* 138: 707–722, 1987.

[73373] 11703. Chatelet, F.; Brianti, E.; Ronco, P.; Roland, J.; Verroust, P.: Ultrastructural localization by monoclonal antibodies of brush border antigens expressed by glomeruli: II. Extrarenal distribution. *Am. J. Path.* 122: 512–519, 1986.

[73374] 11704. Chowdhary, B. P.; Lundgren, S.; Johansson, M.; Hjalm, G.; Akerstrom, G.; Gustavsson, I.; Rask, L.: In situ hybridization mapping of a 500-kDa calcium-sensing protein gene (LRP2) to human chromosome region 2q31–q32.1 and porcine chromosome region 15q22–q24. *Cytogenet. Cell Genet.* 71: 120–123, 1995.

[73375] 11705. Farquhar, M. G.: The unfolding story of megalin (gp330): now recognized as a drug receptor (Editorial) *J. Clin. Invest.* 96: 1184 only, 1995.

[73376] 11706. Farquhar, M. G.; Saito, A.; Kerjaschki, D.; Orlando, R. A.: The Heymann nephritis antigenic complex: megalin (gp330) and RAP. *J. Am. Soc. Nephrol.* 6: 35–47, 1995.

[73377] 11707. Hjalm, G.; Murray, E.; Crumley, G.; Harazim, W.; Lundgren, S.; Onyango, I.; Bo, E. K.; Larsson, M.; Juhlin, C.; Hellman, P.; Davis, H.; Akerstrom, G.; Rask, L.; Morse, B.:

Cloning and sequencing of human gp330, a Ca(2+)-binding receptor with potential intracellular signaling properties. *Europ. J. Biochem.* 239: 132–137, 1996.

- [73378] 11708. Arinami, T.; Kondo, I.; Hamaguchi, H.; Nakajima, S.: Multifocal meningiomas in a patient with a constitutional ring chromosome 22. *J. Med. Genet.* 23: 178–180, 1986.
- [73379] 11709. Petrella, R.; Levine, S.; Wilmot, P. L.; Ashar, K. D.; Casamassima, A. C.; Shapiro, L. R.: Multiple meningiomas in a patient with constitutional ring chromosome 22. *Am. J. Med. Genet.* 47: 184–186, 1993.
- [73380] 11710. Peyrard, M.; Fransson, I.; Xie, Y.-G.; Han, F.-Y.; Rutledge, M. H.; Swahn, S.; Collins, J. E.; Dunham, I.; Collins, V. P.; Dumanski, J. P.: Characterization of a new member of the human beta-adaptin gene family from chromosome 22q12, a candidate meningioma gene. *Hum. Molec. Genet.* 3: 1393–1399, 1994.
- [73381] 11711. Peyrard, M.; Pan, H.-Q.; Kedra, D.; Fransson, I.; Swahn, S.; Hartman, K.; Clifton, S. W.; Roe, B. A.; Dumanski, J. P.: Structure of the promoter and genomic organization of the human beta-prime-adaptin gene (BAM22) from chromosome 22q12. *Genomics* 36: 112–117, 1996.
- [73382] 11712. Zankl, H.; Zang, K. D.: Correlations between clinical and cytogenetical data in 180 human meningiomas. *Can-*

cer Genet. Cytogenet. 1: 351–356,1980.

- [73383] 11713.Bai, C.; Sen, P.; Hofmann, K.; Ma, L.; Goebel, M.; Harper, J. W.; Elledge, S. J.: SKP1 connects cell cycle regulators to the ubiquitinproteolysis machinery through a novel motif, the F-box. Cell 86:263–274, 1996.
- [73384] 11714.Kraus, B.; Pohlschmidt, M.; Leung, A. L. S.; Germino, G. G.; Snarey,A.; Schneider, M. C.; Reeders, S. T.; Frischauf, A.–M.: A novel cyclin gene (CCNF) in the region of the polycystic kidney disease gene (PKD1). Genomics 24:27–33, 1994.
- [73385] 11715.Obermayr, F.; Sutherland, H. F.; Kraus, B.; Frischauf, A.–M.: Mouse cyclin F maps to a conserved linkage group on mouse chromosome17. Mammalian Genome 6: 149–150, 1995.
- [73386] 11716.Donaldson, S. H.; Hirsh, A.; Li, D. C.; Holloway, G.; Chao, J.;Boucher, R. C.; Gabriel, S. E.: Regulation of the epithelial sodiumchannel by serine proteases in human airways. J. Biol. Chem. 277:8338–8345, 2002.

- [73387] 11717. Harvey, K. F.; Dinudom, A.; Cook, D. I.; Kumar, S.: The Nedd4-like protein KIAA0439 is a potential regulator of the epithelial sodium channel. *J. Biol. Chem.* 276: 8597–8601, 2001.
- [73388] 11718. Arriza, J. L.; Kavanaugh, M. P.; Fairman, W. A.; Wu, Y.-N.; Murdoch, G. H.; North, R. A.; Amara, S. G.: Cloning and expression of a human neutral amino acid transporter with structural similarity to the glutamate transporter gene family. *J. Biol. Chem.* 268: 15329–15332, 1993.
- [73389] 11719. Hofmann, K.; Duker, M.; Fink, T.; Lichter, P.; Stoffel, W.: Human neutral amino acid transporter ASCT1: structure of the gene (SLC1A4) and localization to chromosome 2p13–p15. *Genomics* 24: 20–26, 1994.
- [73390] 11720. Shafqat, S.; Tamarappoo, B. K.; Kilberg, M. S.; Purnam, R. S.; McNamara, J. O.; Guadano-Ferraz, A.; Freneau, R. T., Jr.: Cloning and expression of a novel Na(+)-dependent neutral amino acid transporter structurally related to mammalian Na(+)/glutamate cotransporters. *J. Biol. Chem.* 268: 15351–15355, 1993.
- [73391] 11721. Zerangue, N.; Kavanaugh, M. P.: ASCT-1 is a neutral amino acid exchanger with chloride channel activity. *J. Biol. Chem.* 271: 27991–27994, 1996.
- [73392] 11722. Albertin, G.; Rossi, G. P.; Majone, F.; Tiso, N.; Mat-

tara, A.;Danieli, G. A.; Pessina, A. C.; Palu, G.: Fine mapping of the humanendothelin-converting enzyme gene by fluorescent in situ hybridizationand radiation hybrids. Biochem. Biophys. Res. Commun. 221: 682–687,1996.

[73393] 11723.Sharp, J. D.; Pickard, R. T.; Chiou, X. G.; Manetta, J. V.; Kovacevic,S.; Miller, J. R.; Varshavsky, A. D.; Roberts, E. F.; Strifler, B.A.; Brems, D. N.; Kramer, R. M.: Serine 228 is essential for catalyticactivities of 85–kDa cytosolic phospholipase A2. J. Biol. Chem. 269:23250–23254, 1994.

[73394] 11724.Tamimi, R.; Steingrimsson, E.; Copeland, N. G.; Dyer–Montgomery,K.; Lee, J. E.; Hernandez, R.; Jenkins, N. A.; Tapscott, S. J.: TheNEUROD gene maps to human chromosome 2q32 and mouse chromosome 2. Genomics 34:418–421, 1996.

[73395] 11725.Newton, J. S.; Deed, R. W.; Mitchell, E. L. D.; Murphy, J. J.;Norton, J. D.: A B cell specific immediate early human gene is locatedon chromosome band 1q31 and encodes an alpha helical basic phosphoprotein. Biochim.Biophys. Acta 1216: 314–316, 1993.

[73396] 11726.Sierra, D. A.; Gilbert, D. J.; Householder, D.; Grishin, N. V.;Yu, K.; Ukidwe, P.; Barker, S. A.; He, W.; Wensel, T. G.; Otero, G.;Brown, G.; Copeland, N. G.; Jenkins, N. A.; Wilkie, T. M.: Evolutionof the regulators of G–

protein signaling multigene family in mouse and human.  
Genomics 79: 177–185, 2002.

- [73397] 11727. Hofstra, R. M. W.; Valdenaire, O.; Arch, E.; Osinga, J.; Kroes, H.; Loffler, B.-M.; Hamosh, A.; Meijers, C.; Buys, C. H. C. M.: A loss-of-function mutation in the endothelin-converting enzyme 1 (ECE-1) associated with Hirschsprung disease, cardiac defects, and autonomic dysfunction. (Letter) Am. J. Hum. Genet. 64: 304–308, 1999.
- [73398] 11728. Matsuoka, R.; Sawamura, T.; Yamada, K.; Yoshida, M.; Furutani, Y.; Ikura, T.; Shiraki, T.; Hoshikawa, H.; Shimada, K.; Tanzawa, K.; Masaki, T.: Human endothelin converting enzyme gene (ECE1) mapped to chromosomal region 1p36.1. Cytogenet. Cell Genet. 72: 322–324, 1996.
- [73399] 11729. Schmidt, M.; Kroger, B.; Jacob, E.; Seulberger, H.; Subkowski, T.; Otter, R.; Meyer, T.; Schmalzing, G.; Hillen, H.: Molecular characterization of human and bovine endothelin converting enzyme (ECE-1). FEBS Lett. 356: 238–243, 1994.
- [73400] 11730. Shimada, K.; Matsushita, Y.; Wakabayashi, K.; Takahashi, M.; Matsubara, A.; Iijima, Y.; Tanzawa, K.: Cloning and functional expression of human endothelin-converting enzyme cDNA. Biochem. Biophys. Res. Commun. 207: 807–812, 1995.

- [73401] 11731.Valdenaire, O.; Rohrbacher, E.; Mattei, M.-G.: Organization of the gene encoding the human endothelin-converting enzyme (ECE-1). *J.Biol. Chem.* 270: 29794-29798, 1995.
- [73402] 11732.Yorimitsu, K.; Moroi, K.; Inagaki, N.; Saito, T.; Masuda, Y.; Masaki, T.; Seino, S.; Kimura, S.: Cloning and sequencing of a human endothelin converting enzyme in renal adenocarcinoma (ACHN) cells producing endothelin-2. *Biochem. Biophys. Res. Commun.* 208: 721-727, 1995.
- [73403] 11733.Ivey-Hoyle, M.; Conroy, R.; Huber, H. E.; Goodhart, P. J.; Oliff, A.; Heimbrook, D. C.: Cloning and characterization of E2F-2, a novel protein with the biochemical properties of transcription factor E2F. *Molec.Cell. Biol.* 13: 7802-7812, 1993.
- [73404] 11734.Lees, J. A.; Saito, M.; Vidal, M.; Valentine, M.; Look, T.; Harlow, E.; Dyson, N.; Helin, K.: The retinoblastoma protein binds to a family of E2F transcription factors. *Molec. Cell. Biol.* 13: 7813-7825, 1993.
- [73405] 11735.Cloud, J. E.; Rogers, C.; Reza, T. L.; Ziebold, U.; Stone, J. R.; Picard, M. H.; Caron, A. M.; Bronson, R. T.; Lees, J. A.: Mutant mouse models reveal the relative roles of E2F1 and E2F3 in vivo. *Molec.Cell. Biol.* 22: 2663-2672, 2002.

- [73406] 11736.He, Y.; Armanious, M. K.; Thomas, M. J.; Cress, W. D.: Identification of E2F-3B, an alternative form of E2F-3 lacking a conserved N-terminal region. *Oncogene* 19: 3422-3433, 2000.
- [73407] 11737.Sharp, J. D.; White, D. L.; Chiou, X. G.; Goodson, T.; Gamboa, G. C.; McClure, D.; Burgett, S.; Hoskins, J.; Skatrud, P. L.; Sportsman, J. R.; Becker, G. W.; Kang, L. H.; Roberts, E. F.; Kramer, R. M.: Molecular cloning and expression of human Ca(2+)-sensitive cytosolic phospholipase A2. *J. Biol. Chem.* 266: 14850-14853, 1991.
- [73408] 11738.Sheridan, A. M.; Force, T.; Yoon, H.-J.; O'Leary, E.; Choukroun, G.; Taheri, M. R.; Bonventre, J. V.: PLIP, a novel splice variant of Tip60, interacts with group IV cytosolic phospholipase A2, induces apoptosis, and potentiates prostaglandin production. *Molec. Cell. Biol.* 21: 4470-4481, 2001.
- [73409] 11739.Skorecki, K. L.: Personal Communication. Toronto, Canada 4/19/1995.
- [73410] 11740.Tay, A.; Simon, J. S.; Squire, J.; Hamel, K.; Jacob, H. J.; Skorecki, K.: Cytosolic phospholipase A2 gene in human and rat: chromosomal localization and polymorphic markers. *Genomics* 26: 138-141, 1995.
- [73411] 11741.Halford, M. M.; Armes, J.; Buchert, M.; Meskenaite,



V.; Grail,D.; Hibbs, M. L.; Wilks, A. F.; Farlie, P. G.; Newgreen, D. F.; Hovens,C. M.; Stacker, S. A.: Ryk-deficient mice exhibit craniofacial defectsassociated with perturbed Eph receptor crosstalk. *Nature Genet.* 25:414–418, 2000.

[73412] 11742.Nakamura, S.; Stock, D. W.; Wydner, K. L.; Bollekens, J. A.; Takeshita,K.; Nagai, B. M.; Chiba, S.; Kitamura, T.; Freeland, T. M.; Zhao,Z.; Minowada, J.; Lawrence, J. B.; Weiss, K. M.; Ruddle, F. H.: Genomicanalysis of a new mammalian distal-less gene: *Dlx7*. *Genomics* 38:314–324, 1996.

[73413] 11743.Singh, J.; Itahana, Y.; Parrinello, S.; Murata, K.; Desprez, P.–Y.: Molecular cloning and characterization of a zinc finger proteininvolved in Id–1–stimulated mammary epithelial cell growth. *J. Biol.Chem.* 276: 11852–11858, 2001.

[73414] 11744.Kumar, S.; Kinoshita, M.; Noda, M.; Copeland, N. G.; Jenkins, N.A.: Induction of apoptosis by the mouse *Nedd2* gene, which encodesa protein similar to the product of the *Caenorhabditis elegans* celldeath gene *ced–3* and the mammalian IL–1–beta–converting enzyme. *GenesDev.* 8: 1613–1626, 1994.

[73415] 11745.Bollag, R. J.; Siegfried, Z.; Cebra–Thomas, J. A.; Garvey, N.;Davison, E. M.; Silver, L. M.: An ancient family

of embryonically expressed mouse genes sharing a conserved protein motif with the Tlocus. *Nature Genet.* 7: 383–389, 1994.

[73416] 11746. Campbell, C.; Goodrich, K.; Casey, G.; Beatty, B.: Cloning and mapping of a human gene (TBX2) sharing a highly conserved protein motif with the *Drosophila omb* gene. *Genomics* 28: 255–260, 1995.

[73417] 11747. Campbell, C. E.; Casey, G.; Goodrich, K.: Genomic structure of TBX2 indicates conservation with distantly related T-box genes. *Mammalian Genome* 9: 70–73, 1998.

[73418] 11748. Jacobs, J. J. L.; Keblusek, P.; Robanus-Maandag, E.; Kristel, P.; Lingbeek, M.; Nederlof, P. M.; van Welsem, T.; van de Vijver, M. J.; Koh, E. Y.; Daley, G. Q.; van Lohuizen, M.: Senescence bypass screen identifies TBX2, which represses Cdkn2a(p19ARF) and is amplified in a subset of human breast cancers. *Nature Genet.* 26: 291–299, 2000.

[73419] 11749. Law, D. J.; Gebhur, T.; Garvey, N.; Agulnik, S. I.; Silver, L.M.: Identification, characterization, and localization to chromosome 17q21–22 of the human TBX2 homolog, member of a conserved developmental gene family. *Mammalian Genome* 6: 793–797, 1995.

[73420] 11750. Pflugfelder, G. O.; Roth, H.; Poeck, B.: A homology domain shared between *Drosophila optomotor-blind* and

mouse Brachyury is involved in DNA binding. *Biochem. Biophys. Res. Commun.* 186: 918–925, 1992.

- [73421] 11751. Sowden, J. C.; Holt, J. K. L.; Meins, M.; Smith, H. K.; Bhattacharya, S. S.: Expression of *Drosophila* omb-related T-box genes in the developing human and mouse neural retina. *Invest. Ophthalmol. Vis. Sci.* 42: 3095–3102, 2001.
- [73422] 11752. Walter, L.; Dirks, B.; Rothermel, E.; Heyens, M.; Szpirer, C.; Levan, G.; Gunther, E.: A novel, conserved gene of the rat that is developmentally regulated in the testis. *Mammalian Genome* 5: 216–221, 1994.
- [73423] 11753. Walter, L.; Marynen, P.; Szpirer, J.; Levan, G.; Gunther, E.: Identification of a novel conserved human gene, TEGT. *Genomics* 28:301–304, 1995.
- [73424] 11754. Xu, Q.; Reed, J. C.: Bax inhibitor-1, a mammalian apoptosis suppressor identified by functional screening in yeast. *Molec. Cell* 1: 337–346, 1998.
- [73425] 11755. Hsu, Y.-C.; Perin, M. S.: Human neuronal pentraxin II (NPTX2): conservation, genomic structure, and chromosomal localization. *Genomics* 28:220–227, 1995.
- [73426] 11756. Schlimgen, A. K.; Helms, J. A.; Vogel, H.; Perin, M. S.: Neuronal pentraxin, a secreted protein with homology to acute phase proteins of the immune system. *Neuron* 14: 519–526, 1995.

- [73427] 11757.Kumar, S.; White, D. L.; Takai, S.; Turczynowicz, S.; Juttner,C. A.; Hughes, T. P.: Apoptosis regulatory gene NEDD2 maps to human chromosome segment 7q34–35, a region frequently affected in haematological neoplasms. Hum. Genet. 95: 641–644, 1995.
- [73428] 11758.Lassus, P.; Opitz–Araya, X.; Lazebnik, Y.: Requirement for caspase–2 in stress–induced apoptosis before mitochondrial permeabilization. Science 297:1352–1354, 2002.
- [73429] 11759.Aksoy, I. A.; Callen, D. F.; Apostolou, S.; Her, C.; Weinshilboum,R. M.: Thermolabile phenol sulfotransferase gene (STM) localization to human chromosome 16p11.2. Genomics 23: 275–277, 1994.
- [73430] 11760.Aksoy, I. A.; Weinshilboum, R. M.: Human thermolabile phenol sulfotransferase gene (STM): molecular cloning and structural characterization. Biochem.Biophys. Res. Commun. 208: 786–795, 1995.
- [73431] 11761.Wood, T. C.; Aksoy, I. A.; Aksoy, S.; Weinshilboum, R. M.: Human liver thermolabile phenol sulfotransferase: cDNA cloning, expression and characterization. Biochem. Biophys. Res. Commun. 198: 1119–1127, 1994.
- [73432] 11762.Zhu, X.; Veronese, M. E.; Bernard, C. C. A.; Sansom, L. N.; McManus,M. E.: Identification of two human brain

aryl sulfotransferase cDNAs. Biochem.Biophys. Res. Commun. 195: 120–127, 1993.

[73433] 11763.Fuhlbrigge, R. C.; Kieffer, J. D.; Armerding, D.; Kupper, T. S.: Cutaneous lymphocyte antigen is a specialized form of PSGL-1 expressed on skin-homing T cells. Nature 389: 978–981, 1997.

[73434] 11764.Herron, M. J.; Nelson, C. M.; Larson, J.; Snapp, K. R.; Kansas, G. S.; Goodman, J. L.: Intracellular parasitism by the human granulocyte ehrlichiosis bacterium through the P-selectin ligand, PSGL-1. Science 288:1653–1656, 2000.

[73435] 11765.Veldman, G. M.; Bean, K. M.; Cumming, D. A.; Eddy, R. L.; Sait, S. N. J.; Shows, T. B.: Genomic organization and chromosomal localization of the gene encoding human P-selectin glycoprotein ligand. J. Biol.Chem. 270: 16470–16475, 1995.

[73436] 11766.Yang, J.; Galipeau, J.; Kozak, C. A.; Furie, B. C.; Furie, B.: Mouse P-selectin glycoprotein ligand-1: molecular cloning, chromosomal localization, and expression of a functional P-selectin receptor. Blood 87:4176–4186, 1996.

[73437] 11767.Brown, C. B.; Boyer, A. S.; Runyan, R. B.; Barnett, J. V.: Requirement of type III TGF-beta receptor for endocardial cell transformation in the heart. Science 283:

2080–2082, 1999.

- [73438] 11768. Lewis, K. A.; Gray, P. C.; Blount, A. L.; MacConell, L. A.; Wiater, E.; Bilezikjian, L. M.; Vale, W.: Betaglycan binds inhibin and can mediate functional antagonism of activin signalling. *Nature* 404:411–414, 2000.
- [73439] 11769. Hu, Y.-F.; Luscher, B.; Admon, A.; Mermod, N.; Tjian, R.: Transcription factor AP-4 contains multiple dimerization domains that regulate dimer specificity. *Genes Dev.* 4: 1741–1752, 1990.
- [73440] 11770. Mermod, N.; Williams, T. J.; Tjian, R.: Enhancer binding factors AP-4 and AP-1 act in concert to activate SV40 late transcription in vitro. *Nature* 332: 557–561, 1988.
- [73441] 11771. Carr, C. S.; Sharp, P. A.: A helix–loop–helix protein related to the immunoglobulin E box-binding proteins. *Molec. Cell. Biol.* 10:4384–4388, 1990.
- [73442] 11772. Bonner, C. A.; Loftus, S. K.; Wasmuth, J. J.: Isolation, characterization, and precise physical localization of human CDX1, a caudal-type homeobox gene. *Genomics* 28: 206–211, 1995.
- [73443] 11773. Subramanian, V.; Meyer, B. I.; Gruss, P.: Disruption of the murine homeobox gene *Cdx1* affects axial skeletal identities by altering the mesodermal expression domains

of Hox genes. *Cell* 83: 641–653, 1995.

- [73444] 11774.Henske, E. P.; Short, M. P.; Jozwiak, S.; Bovey, C. M.; Ramlakhan, S.; Haines, J. L.; Kwiatkowski, D. J.: Identification of VAV2 on 9q34 and its exclusion as the tuberous sclerosis gene TSC1. *Ann. Hum. Genet.* 59: 25–37, 1995.
- [73445] 11775.Schwientek, T.; Nomoto, M.; Levery, S. B.; et al: Control of O-glycan branch formation. *J. Biol. Chem.* 274: 4504–4512, 1999.
- [73446] 11776.Walczak, H.; Degli-Esposti, M. A.; Johnson, R. S.; Smolak, P. J.; Waugh, J. Y.; Boiani, N.; Timour, M. S.; Gerhart, M. J.; Schooley, K. A.; Smith, C. A.; Goodwin, R. G.; Rauch, C. T.: TRAIL-R2: a novel apoptosis-mediating receptor for TRAIL. *EMBO J.* 16: 5386–5397, 1997.
- [73447] 11777.Wu, G. S.; Burns, T. F.; McDonald, E. R., III; Jiang, W.; Meng, R.; Krantz, I. D.; Kao, G.; Gan, D.-D.; Zhou, J.-Y.; Muschel, R.; Hamilton, S. R.; Spinner, N. B.; Markowitz, S.; Wu, G.; El-Deiry, W. S.: KILLER/DR5 is a DNA damage-inducible p53-regulated death receptor gene. *Nature Genet.* 17: 141–143, 1997.
- [73448] 11778.Degli-Esposti, M. A.; Smolak, P. J.; Walczak, H.; Waugh, J.; Huang, C.-P.; DuBose, R. F.; Goodwin, R. G.; Smith, C. A.: Cloning and characterization of TRAIL-R3, a novel member of the emerging TRAIL receptor family.

J.Exp. Med. 186: 1165–1170, 1997.

- [73449] 11779.Kanaar, R.; Troelstra, C.; Swagemakers, S. M. A.; Essers, J.; Smit, B.; Franssen, J.–H.; Pastink, A.; Bezzubova, O. Y.; Buerstedde, J.–M.; Clever, B.; Heyer, W.–D.; Hoeijmakers, J. H. J.: Human and mouse homologs of the *Saccharomyces cerevisiae* RAD54 DNA repair gene: evidence for functional conservation. *Curr. Biol.* 6: 828–838, 1996.
- [73450] 11780.Matsuda, M.; Miyagawa, K.; Takahashi, M.; Fukuda, T.; Kataoka, T.; Asahara, T.; Inui, H.; Watatani, M.; Yasutomi, M.; Kamada, N.; Dohi, K.; Kamiya, K.: Mutations in the RAD54 recombination gene in primary cancers. *Oncogene* 18: 3427–3430, 1999.
- [73451] 11781.Tucker, J. E.; Winkfein, R. J.; Cooper, C. B.; Schnetkamp, P. P.: cDNA cloning of the human retinal rod Na–Ca + K exchanger: comparison with a revised bovine sequence. *Invest. Ophthalm. Vis. Sci.* 39: 435–440, 1998.
- [73452] 11782.Tucker, J. E.; Winkfein, R. J.; Murthy, S. K.; Friedman, J. S.; Walter, M. A.; Demetrick, D. J.; Schnetkamp, P. P. M.: Chromosomal localization and genomic organization of the human retinal rod Na–Ca+K exchanger. *Hum. Genet.* 103: 411–414, 1998.
- [73453] 11783.Dennis, J. A.; Healy, P. J.; Beaudet, A. L.; O'Brien, W. E.: Molecular definition of bovine argininosuccinate syn–



thetase deficiency. Proc.Nat. Acad. Sci. 86: 7947–7951, 1989.

[73454] 11784.McCarrey, J. R.; Riggs, A. D.: Determinator–inhibitor pairs as a mechanism for threshold setting in development: a possible function for pseudogenes. Proc. Nat. Acad. Sci. 83: 679–683, 1986.

[73455] 11785.Su, T.–S.; Nussbaum, R. L.; Airhart, S.; Ledbetter, D. H.; Mohandas, T.; O'Brien, W. E.; Beaudet, A. L.: Human chromosomal assignments for 14 argininosuccinate synthetase pseudogenes: cloned DNAs as reagents for cytogenetic analysis. Am. J. Hum. Genet. 36: 954–964, 1984.

[73456] 11786.Hara, Y.; Wakamori, M.; Ishii, M.; Maeno, E.; Nishida, M.; Yoshida, T.; Yamada, H.; Shimizu, S.; Mori, E.; Kudoh, J.; Shimizu, S.; Kurose, H.; Okada, Y.; Imoto, K.; Mori, Y.: LTRPC2 Ca(2+)–permeable channel activated by changes in redox status confers susceptibility to cell death. Molec. Cell 9: 163–173, 2002.

[73457] 11787.Harteneck, C.; Plant, T. D.; Schultz, G.: From worm to man: three subfamilies of TRP channels. Trends Neurosci. 23: 159–166, 2000.

[73458] 11788.Kudoh, J.; Nagamine, K.; Asakawa, S.; Abe, I.; Kawasaki, K.; Maeda, H.; Tsujimoto, S.; Minoshima, S.; Ito, F.; Shimizu, N.: Localization of 16 exons to a 450–kb re–

gion involved in the autoimmune polyglandular disease type I (APECED) on human chromosome 21q22.3. DNA Res. 4:45–52, 1997.

- [73459] 11789. Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.; Asakawa, S.; Ito, F.; Shimizu, N.: Molecular cloning of a novel putative Ca(2+) channel protein (TRPC7) highly expressed in brain. Genomics 54:124–131, 1998.
- [73460] 11790. Perraud, A.–L.; Fleig, A.; Dunn, C. A.; Bagley, L. A.; Launay, P.; Schmitz, C.; Stokes, A. J.; Zhu, Q.; Bessman, M. J.; Penner, R.; Kinet, J.–P.; Scharenberg, A. M.: ADP–ribose gating of the calcium–permeable LTRPC2 channel revealed by Nudix motif homology. Nature 411: 595–599, 2001.
- [73461] 11791. Sano, Y.; Inamura, K.; Miyake, A.; Mochizuki, S.; Yokoi, H.; Matsushime, H.; Furuichi, K.: Immunocyte Ca(2+) influx system mediated by LTRPC2. Science 293:1327–1330, 2001.
- [73462] 11792. Koyama, N.; Ishibashi, K.; Kuwahara, M.; Inase, N.; Ichioka, M.; Sasaki, S.; Marumo, F.: Cloning and functional expression of human aquaporin 8 cDNA and analysis of its gene. Genomics 54: 169–172, 1998.
- [73463] 11793. Viggiano, L.; Rocchi, M.; Svelto, M.; Calamita, G.: Assignment of the aquaporin–8 water channel gene (AQP8) to human chromosome 16p12. Cytogenet. Cell Genet. 84:

208–210, 1999.

- [73464] 11794.Graler, M. H.; Bernhardt, G.; Lipp, M.: EDG6, a novel G–protein–coupled receptor related to receptors for bioactive lysophospholipids, is specifically expressed in lymphoid tissue. *Genomics* 53: 164–169, 1998.
- [73465] 11795.Jedlicka, A. E.; Taylor, E. W.; Meyers, D. A.; Liu, Z.; Levitt, R. C.: Localization of the highly polymorphic locus D19S120 to 19p13.3 by linkage. *Cytogenet. Cell. Genet.* 65: 140 only, 1994.
- [73466] 11796.Koegl, M.; Hoppe, T.; Schlenker, S.; Ulrich, H. D.; Mayer, T. U.; Jentsch, S.: A novel ubiquitination factor, E4, is involved in multiubiquitin chain assembly. *Cell* 96: 635–644, 1999.
- [73467] 11797.Nonaka, S.; Tanaka, Y.; Okada, Y.; Takeda, S.; Harada, A.; Kanai, Y.; Kido, M.; Hirokawa, N.: Randomization of left–right asymmetry due to loss of nodal cilia generating leftward flow of extraembryonic fluid in mice lacking KIF3B motor protein. *Cell* 95: 829–837, 1998.
- [73468] 11798.Yamazaki, H.; Nakata, T.; Okada, Y.; Hirokawa, N.: KIF3A/B: a heterodimeric kinesin superfamily protein that works as a microtubule plus end–directed motor for membrane organelle transport. *J. Cell Biol.* 130: 1387–1399, 1995.

- [73469] 11799. Allikmets, R.; Schriml, L. M.; Hutchinson, A.; Romano-Spica, V.; Dean, M.: A human placenta-specific ATP-binding cassette gene (ABCP) on chromosome 4q22 that is involved in multidrug resistance. *Cancer Res.* 58: 5337–5339, 1998.
- [73470] 11800. Bailey-Dell, K. J.; Hassel, B.; Doyle, L. A.; Ross, D. D.: Promoter characterization and genomic organization of the human breast cancer resistance protein (ATP-binding cassette transporter G2) gene. *Biochim. Biophys. Acta* 1520: 234–241, 2001.
- [73471] 11801. Doyle, L. A.; Yang, W.; Abruzzo, L. V.; Krogmann, T.; Gao, Y.; Rishi, A. K.; Ross, D. D.: A multidrug resistance transporter from human MCF-7 breast cancer cells. *Proc. Nat. Acad. Sci.* 95: 15665–15670, 1998.
- [73472] 11802. Suzuki, N.; Zara, J.; Sato, T.; Ong, E.; Bakhiet, N.; Oshima, R. G.; Watson, K. L.; Fukuda, M. N.: A cytoplasmic protein, bystin, interacts with trophinin, tastin, and cytokeratin and may be involved in trophinin-mediated cell adhesion between trophoblast and endometrial epithelial cells. *Proc. Nat. Acad. Sci.* 95: 5027–5032, 1998.
- [73473] 11803. Cerutti, A.; Schaffer, A.; Goodwin, R. G.; Shah, S.; Zan, H.; Ely, S.; Casali, P.: Engagement of CD153 (CD30 ligand) by CD30-positive T cells inhibits class switch DNA

recombination and antibody production in human IgD-positive IgM-positive B cells. *J. Immun.* 165: 786–794, 2000.

- [73474] 11804. Croager, E. J.; Abraham, L. J.: Characterisation of the human CD30 ligand gene structure. *Biochim. Biophys. Acta* 1353: 231–235, 1997.
- [73475] 11805. Smith, C. A.; Gruss, H.-J.; Davis, T.; Anderson, D.; Farrah, T.; Baker, E.; Sutherland, G. R.; Brannan, C. I.; Copeland, N. G.; Jenkins, N. A.; Grabstein, K. H.; Gliniak, B.; and 9 others: CD30 antigen, a marker for Hodgkin's lymphoma, is a receptor whose ligand defines an emerging family of cytokines with homology to TNF. *Cell* 73: 1349–1360, 1993.
- [73476] 11806. Hurskainen, T. L.; Hirohata, S.; Seldin, M. F.; Apte, S. S.: ADAM-TS5, ADAM-TS6, and ADAM-TS7, novel members of a new family of zinc metalloproteases: general features and genomic distribution of the ADAM-TS family. *J. Biol. Chem.* 274: 25555–25563, 1999.
- [73477] 11807. Tang, B. L.; Hong, W.: ADAMTS: a novel family of proteases with an ADAM protease domain and thrombospondin 1 repeats. *FEBS Lett.* 445: 223–225, 1999.
- [73478] 11808. Tortorella, M. D.; Burn, T. C.; Pratta, M. A.; Abbaszade, I.; Hollis, J. M.; Liu, R.; Rosenfeld, S. A.; Copeland,

R. A.; Decicco, C. P.; Wynn, R.; Rockwell, A.; Yang, F.; and 16 others: Purification and cloning of aggrecanase-1: a member of the ADAMTS family of proteins. *Science* 284:1664–1666, 1999.

[73479] 11809. Higashide, T.; Inana, G.: Characterization of the gene for HRG4(UNC119), a novel photoreceptor synaptic protein homologous to unc-119. *Genomics* 57:446–450, 1999.

[73480] 11810. Higashide, T.; McLaren, M. J.; Inana, G.: Localization of HRG4, a photoreceptor protein homologous to unc-119, in ribbon synapse. *Invest. Ophthalmol. Vis. Sci.* 39: 690–698, 1998.

[73481] 11811. Higashide, T.; Murakami, A.; McLaren, M. J.; Inana, G.: Cloning of the cDNA for a novel photoreceptor protein. *J. Biol. Chem.* 271:1797–1804, 1996.

[73482] 11812. Swanson, D. A.; Chang, J. T.; Campochiaro, P. A.; Zack, D. J.; Valle, D.: Mammalian orthologs of *C. elegans* unc-119 highly expressed in photoreceptors. *Invest. Ophthalmol. Vis. Sci.* 39: 2085–2094, 1998.

[73483] 11813. Gundelfinger, E.: Personal Communication. Magdeburg, FRG. 1/8/1999.

[73484] 11814. Hashida, H.; Goto, J.; Zhao, N.; Takahashi, N.; Hirai, M.; Kanazawa, I.; Sakaki, Y.: Cloning and mapping of

ZNF231, a novel brain-specific gene encoding neuronal double zinc finger protein whose expression is enhanced in a neurodegenerative disorder, multiple system atrophy (MSA). *Genomics* 54: 50–58, 1998.

[73485] 11815. Santorelli, F. M.; Patrono, C.; Fortini, D.; Tessa, A.; Comanducci, G.; Bertini, E.; Pierallini, A.; Amabile, G. A.; Casali, C.: Intrafamilial variability in hereditary spastic paraplegia associated with an SPG4 gene mutation. *Neurology* 55: 702–705, 2000.

[73486] 11816. Sauter, S.; Mitterski, B.; Klimpe, S.; Bonsch, D.; Schols, L.; Visbeck, A.; Papke, T.; Hopf, H. C.; Engel, W.; Deufel, T.; Epplen, J. T.; Neesen, J.: Mutation analysis of the spastin gene (SPG4) in patients in Germany with autosomal dominant hereditary spastic paraplegia. *Hum. Mutat.* 20: 127–132, 2002.

[73487] 11817. Svenson, I. K.; Ashley-Koch, A. E.; Pericak-Vance, M. A.; Marchuk, D. A.: A second leaky splice-site mutation in the spastin gene. (Letter) *Am. J. Hum. Genet.* 69: 1407–1409, 2001.

[73488] 11818. Flannery, C. R.; Hughes, C. E.; Schumacher, B. L.; Tudor, D.; Aydelotte, M. B.; Kuettner, K. E.; Caterson, B.: Articular cartilage superficial zone protein (SZP) is homologous to megakaryocyte stimulating factor precursor and is

a multifunctional proteoglycan with potential growth-promoting, cytoprotective, and lubricating properties in cartilage metabolism. *Biochem. Biophys. Res. Commun.* 254: 535–541, 1999.

[73489] 11819. Ikegawa, S.; Sano, M.; Koshizuka, Y.; Nakamura, Y.: Isolation, characterization and mapping of the mouse and human PRG4 (proteoglycan4) genes. *Cytogenet. Cell Genet.* 90: 291–297, 2000.

[73490] 11820. Merberg, D. M.; Fitz, L. J.; Temple, P.; Giannotti, J.; Murtha, P.; Fitzgerald, M.; Scaltreto, H.; Kelleher, K.; Preissner, K.; Kriz, R.; Jacobs, K.; Turner, K.: In: Preissner, K. T.; Rosenblatt, S.; Kost, C.; Wegerhoff, J.; Mosher, D. F. (eds.): *Biology of Vitronectins and Their Receptors*. Elsevier Science, B.V. 1993. Pp. 45–53.

[73491] 11821. Schumacher, B. L.; Block, J. A.; Schmid, T. M.; Ayselotte, M. B.; Kuettner, K. E.: A novel proteoglycan synthesized and secreted by chondrocytes of the superficial zone of articular cartilage. *Arch. Biochem. Biophys.* 15: 144–152, 1994.

[73492] 11822. Pennica, D.; Swanson, T. A.; Welsh, J. W.; Roy, M. A.; Lawrence, D. A.; Lee, J.; Brush, J.; Taneyhill, L. A.; Deuel, B.; Lew, M.; Watanabe, C.; Cohen, R. L.; Melhem, M. F.; Finley, G. G.; Quirke, P.; Goddard, A. D.; Hillan, K. J.;



Gurney, A. L.; Botstein, D.; Levine, A. J.: WISP genes are members of the connective tissue growth factor family that are up-regulated in Wnt-1-transformed cells and aberrantly expressed in human colon tumors. *Proc. Nat. Acad. Sci.* 95: 14717–14722, 1998.

[73493] 11823. Tanaka, S.; Sugimachi, K.; Saeki, H.; Kinoshita, J.; Ohga, T.; Shimada, M.; Maehara, Y.; Sugimachi, K.: A novel variant of WISP1 lacking a von Willebrand type C module overexpressed in scirrhous gastric carcinoma. *Oncogene* 20: 5525–5532, 2001.

[73494] 11824. Gee, S.; Krauss, S. W.; Miller, E.; Aoyagi, K.; Arenas, J.; Conboy, J. G.: Cloning of mDEAH9, a putative RNA helicase and mammalian homologue of *Saccharomyces cerevisiae* splicing factor Prp43. *Proc. Nat. Acad. Sci.* 94: 11803–11807, 1997.

[73495] 11825. Imamura, O.; Sugawara, M.; Furuichi, Y.: Cloning and characterization of a putative human RNA helicase gene of the DEAH-box protein family. *Biochem. Biophys. Res. Commun.* 240: 335–340, 1997.

[73496] 11826. Sano, K.; Tanihara, H.; Heimark, R. L.; Obata, S.; Davidson, M.; St. John, T.; Taketani, S.; Suzuki, S.: Protocadherins: a large family of cadherin-related molecules in central nervous system. *EMBO J.* 12: 2249–2256, 1993.

- [73497] 11827.Eisenblatter, T.; Galla, H.-J.: A new multidrug resistance protein at the blood-brain barrier. *Biochem. Biophys. Res. Commun.* 293:1273–1278, 2002.
- [73498] 11828.Miyake, K.; Mickley, L.; Litman, T.; Zhan, Z.; Robey, R.; Cristensen, B.; Brangi, M.; Greenberger, L.; Dean, M.; Fojo, T.; Bates, S. E.: Molecular cloning of cDNAs which are highly overexpressed in mitoxantrone-resistant cells: demonstration of homology to ABC transport genes. *Cancer Res.* 59:8–13, 1999.
- [73499] 11829.Ozvegy, C.; Litman, T.; Szakacs, G.; Nagy, Z.; Bates, S.; Varadi, A.; Sarkadi, B.: Functional characterization of the human multidrug transporter, ABCG2, expressed in insect cells. *Biochem. Biophys. Res. Commun.* 285: 111–117, 2001.
- [73500] 11830.Dominguez, O.; Ashhab, Y.; Sabater, L.; Belloso, E.; Caro, P.; Pujol-Borrell, R.: Cloning of ARE-containing genes by AU-motif-directed display. *Genomics* 54: 278–286, 1998.
- [73501] 11831.Kostrub, C. F.; Knudsen, K.; Subramani, S.; Enoch, T.: Hus1p, a conserved fission yeast checkpoint protein, interacts with Rad1p and is phosphorylated in response to DNA damage. *EMBO J.* 17: 2055–2066, 1998.
- [73502] 11832.Lieberman, H. B.; Hopkins, K. M.; Nass, M.; Demet-

rick, D.; Davey, S.: A human homolog of the *Schizosaccharomyces pombe* rad9+ checkpoint control gene. *Proc. Nat. Acad. Sci.* 93: 13890–13895, 1996.

[73503] 11833. Katashima, R.; Iwahana, H.; Fujimura, M.; Yamaoka, T.; Ishizuka, T.; Tatibana, M.; Itakura, M.: Molecular cloning of a human cDNA for the 41-kDa phosphoribosylpyrophosphate synthetase-associated protein. *Biochim. Biophys. Acta* 1396: 245–250, 1998.

[73504] 11834. Katashima, R.; Iwahana, H.; Fujimura, M.; Yamaoka, T.; Itakura, M.: Assignment of the human phosphoribosylpyrophosphate synthetase-associated protein 41 gene (PRPSAP2) to 17p11.2–p12. *Genomics* 54: 180–181, 1998.

[73505] 11835. Ando, A.; Kikuti, Y. Y.; Kawata, H.; Okamoto, N.; Imai, T.; Eki, T.; Yokoyama, K.; Soeda, E.; Ikemura, T.; Abe, K.; Inoko, H.: Cloning of a new kinesin-related gene located at the centromeric end of the human MHC region. *Immunogenetics* 39: 194–200, 1994.

[73506] 11836. Endow, S. A.; Higuchi, H.: A mutant of the motor protein kinesin that moves in both directions on microtubules. *Nature* 406: 913–916, 2000.

[73507] 11837. Hoyt, M. A.; He, L.; Totis, L.; Saunders, W. S.: Loss of function of *Saccharomyces cerevisiae* kinesin-related CIN8 and KIP1 is suppressed by KAR3 motor domain mu-

tations. *Genetics* 135: 35–44, 1993.

[73508] 11838. Janatipour, M.; Naumov, Y.; Ando, A.; Sugimura, K.; Okamoto, N.; Tsuji, K.; Abe, K.; Inoko, H.: Search for MHC-associated genes in human: five new genes centromeric to HLA-DP with yet unknown functions. *Immunogenetics* 35:272–278, 1992.

[73509] 11839. Koike, N.; Hida, A.; Numano, R.; Hirose, M.; Sakaki, Y.; Tei, H.: Identification of the mammalian homologues of the *Drosophila* timeless gene, timeless1. *FEBS Lett.* 441: 427–431, 1998.

[73510] 11840. Sangoram, A. M.; Saez, L.; Antoch, M. P.; Gekakis, N.; Staknis, D.; Whiteley, A.; Fruechte, E. M.; Vitaterna, M. H.; Shimomura, K.; King, D. P.; Young, M. W.; Weitz, C. J.; Takahashi, J. S.: Mammalian circadian autoregulatory loop: a timeless ortholog and mPer1 interact and negatively regulate CLOCK-BMAL1-induced transcription. *Neuron* 21:1101–1113, 1998.

[73511] 11841. Zylka, M. J.; Shearman, L. P.; Levine, J. D.; Jin, X.; Weaver, D. R.; Reppert, S. M.: Molecular analysis of mammalian timeless. *Neuron* 21:1115–1122, 1998.

[73512] 11842. Ardley, H. C.; Rose, S. A.; Tan, N.; Leek, J. P.; Markham, A. F.; Robinson, P. A.: Genomic organization of the human ubiquitin-conjugating enzyme gene, UBE2L6 on

chromosome 11q12. Cytogenet. Cell Genet. 89:137–140, 2000.

[73513] 11843.Kumar, S.; Kao, W. H.; Howley, P. M.: Physical interaction between specific E2 and Hect E3 enzymes determines functional cooperativity. J.Biol. Chem. 272: 13548–13554, 1997.

[73514] 11844.tom Dieck, S.; Sanmarti-Vila, L.; Langnaese, K.; Richter, K.; Kindler, S.; Soyke, A.; Wex, H.; Smalla, K.-H.; Kampf, U.; Franzer, J.-T.; Stumm, M.; Garner, C. C.; Gundelfinger, E. D.: Bassoon, a novel zinc-finger CAG/glutamine-repeat protein selectively localized at the active zone of presynaptic nerve terminals. J. Cell Biol. 142: 499–509, 1998.

[73515] 11845.Winter, c.; tom Dieck, S.; Boeckers, T. M.; Bockmann, J.; Kampf, U.; Sanmarti-Vila, L.; Langnaese, K.; Altmann, W.; Stumm, M.; Soyke, A.; Wieacker, P.; Garner, C. C.; Gundelfinger, E. D.: The presynaptic cytomatrix protein Bassoon: sequence and chromosomal localization of the human BSN gene. Genomics 57: 389–397, 1999.

[73516] 11846.Bui, T. D.; Levy, E. R.; Subramaniam, V. N.; Lowe, S. L.; Hong, W.: cDNA characterization and chromosomal mapping of human Golgi SNARE GS27 and GS28 to chromosome 17. Genomics 57: 285–288, 1999.

- [73517] 11847.Lowe, S. L.; Peter, F.; Subramaniam, V. N.; Wong, S. H.; Hong,W.: A SNARE involved in protein transport through the Golgi apparatus. *Nature* 389:881–884, 1997.
- [73518] 11848.Nagahama, M.; Orci, L.; Ravazzola, M.; Amherdt, M.; Lacomis, L.;Tempst, P.; Rothman, J. E.; Sollner, T. H.: A v-SNARE implicatedin intra-Golgi transport. *J. Cell Biol.* 133: 507–516, 1996.
- [73519] 11849.Subramaniam, V. N.; Peter, F.; Philp, R.; Wong, S. H.; Hong, W.: GS28, a 28-kilodalton Golgi SNARE that participates in ER-Golgitransport. *Science* 272: 1161–1163, 1996.
- [73520] 11850.Hay, J. C.; Chao, D. S.; Kuo, C. S.; Scheller, R. H.: Proteininteractions regulating vesicle transport between the endoplasmicreticulum and Golgi apparatus in mammalian cells. *Cell* 89: 149–158,1997.
- [73521] 11851.Li, J.; Ding, S.-F.; Habib, N. A.; Fermor, B. F.; Wood, C. B.;Gilmour, R. S.: Partial characterization of a cDNA for human stearoyl-CoA desaturase and changes in its mRNA expression in some normal and malignant tissues. *Int. J. Cancer* 57: 348–352, 1994.
- [73522] 11852.Ntambi, J. M.; Miyazaki, M.; Stoehr, J. P.; Lan, H.; Kendzierski,C. M.; Yandell, B. S.; Song, Y.; Cohen, P.; Friedman, J. M.; Attie,A. D.: Loss of stearoyl-CoA desat-

urase-1 function protects mice against adiposity. *Proc. Nat. Acad. Sci.* 99: 11482–11486, 2002.

- [73523] 11853. Thiede, M. A.; Ozols, J.; Strittmatter, P.: Construction and sequence of cDNA for rat liver stearyl coenzyme A desaturase. *J. Biol. Chem.* 261:13230–13235, 1986.
- [73524] 11854. Zhang, L.; Ge, L.; Parimoo, S.; Stenn, K.; Prouty, S. M.: Human stearyl-CoA desaturase: alternative transcripts generated from a single gene by usage of tandem polyadenylation sites. *Biochem. J.* 340:255–264, 1999.
- [73525] 11855. Zheng, Y.; Eilertsen, K. J.; Ge, L.; Zhang, L.; Sundberg, J. P.; Prouty, S. M.; Stenn, K. S.; Parimoo, S.: *Scd1* is expressed in sebaceous glands and is disrupted in the *asebia* mouse. (Letter) *Nature Genet.* 23:268–270, 1999.
- [73526] 11856. Duh, F.-M.; Latif, F.; Weng, Y.; Geil, L.; Modi, W.; Stackhouse, T.; Matsumura, F.; Duan, D. R.; Linehan, W. M.; Lerman, M. I.; Gnarr, J. R.: cDNA cloning and expression of the human homolog of the sea urchin fascin and *Drosophila* singed genes which encodes an actin-bundling protein. *DNA Cell Biol.* 13: 821–827, 1994.
- [73527] 11857. Mosialos, G.; Birkenbach, M.; Ayehunie, S.; Matsumura, F.; Pinkus, G. S.; Kieff, E.; Langhoff, E.: Circulating human dendritic cells differentially express high levels of a 55-kd actin-bundling protein. *Am. J. Path.* 148: 593–600,

1996.

- [73528] 11858. Ono, S.; Yamakita, Y.; Yamashiro, S.; Matsudaira, P. T.; Gnarr, J. R.; Obinata, T.; Matsumura, F.: Identification of an actin binding region and a protein kinase C phosphorylation site on human fascin. *J. Biol. Chem.* 272: 2527–2533, 1997.
- [73529] 11859. Pinkus, G. S.; Pinkus, J. L.; Langhoff, E.; Matsumura, F.; Yamashiro, S.; Mosialos, G.; Said, J. W.: Fascin, a sensitive new marker for Reed–Sternberg cells of Hodgkin's disease: evidence for a dendritic or B cell derivation? *Am. J. Path.* 150: 543–562, 1997.
- [73530] 11860. Sonderbye, L.; Magerstadt, R.; Blatman, R. N.; Prefer, F. I.; Langhoff, E.: Selective expression of human fascin (p55) by dendritic leukocytes. *Adv. Exp. Med. Biol.* 471: 41–46, 1997.
- [73531] 11861. Yamakita, Y.; Ono, S.; Matsumura, F.; Yamashiro, S.: Phosphorylation of human fascin inhibits its actin binding and bundling activities. *J. Biol. Chem.* 271: 12632–12638, 1996.
- [73532] 11862. Yamashiro–Matsumura, S.; Matsumura, F.: Intracellular localization of the 55–kD actin–bundling protein in cultured cells: spatial relationships with actin, alpha-actinin, tropomyosin, and fimbrin. *J. Cell Biol.* 103:631–640,



1986.

- [73533] 11863.Yamashiro–Matsumura, S.; Matsumura, F.: Purification and characterization of an F–actin–bundling 55–kilodalton protein from HeLa cells. *J.Biol. Chem.* 260: 5087–5097, 1985.
- [73534] 11864.Carapeti, M.; Aguiar, R. C. T.; Chase, A.; Goldman, J. M.; Cross, N. C. P.: Assignment of the steroid receptor coactivator–1 (SRC–1) gene to human chromosome band 2p23. *Genomics* 52: 242–244, 1998.
- [73535] 11865.Hayashi, Y.; Ohmori, S.; Ito, T.; Seo, H.: A splicing variant of steroid receptor coactivator–1 (SRC–1E): the major isoform of SRC–1 to mediate thyroid hormone action. *Biochem. Biophys. Res. Commun.* 236:83–87, 1997.
- [73536] 11866.Jenster, G.; Spencer, T. E.; Burcin, M. M.; Tsai, S. Y.; Tsai, M.–J.; O'Malley, B. W.: Steroid receptor induction of gene transcription: a two–step model. *Proc. Nat. Acad. Sci.* 94: 7879–7884, 1997.
- [73537] 11867.Liu, Z.; Wong, J.; Tsai, S. Y.; Tsai, M. J.; O'Malley, B. W.: Steroid receptor coactivator–1 (SRC–1) enhances ligand–dependent and receptor–dependent cell–free transcription of chromatin. *Proc. Nat. Acad. Sci.* 96: 9485–9490, 1999.
- [73538] 11868.Onate, S. A.; Tsai, S. Y.; Tsai, M.–J.; O'Malley, B. W.:

Sequence and characterization of a coactivator for the steroid hormone receptor superfamily. *Science* 270: 1354–1357, 1995.

- [73539] 11869. Takeshita, A.; Yen, P. M.; Misiti, S.; Cardona, G. R.; Liu, Y.; Chin, W. W.: Molecular cloning and properties of a full-length putative thyroid hormone receptor coactivator. *Endocrinology* 137: 3594–3597, 1996.
- [73540] 11870. Masson, J.-Y.; Stasiak, A. Z.; Stasiak, A.; Benson, F. E.; West, S. C.: Complex formation by the human RAD51C and XRCC3 recombination repair proteins. *Proc. Nat. Acad. Sci.* 98: 8440–8446, 2001.
- [73541] 11871. Selfors, L. M.; Schutzman, J. L.; Borland, C. Z.; Stern, M. J.: Soc-2 encodes a leucine-rich repeat protein implicated in fibroblast growth factor receptor signaling. *Proc. Nat. Acad. Sci.* 95: 6903–6908, 1998.
- [73542] 11872. Sieburth, D. S.; Sun, Q.; Han, M.: SUR-8, a conserved Ras-binding protein with leucine-rich repeats, positively regulates Ras-mediated signaling in *C. elegans*. *Cell* 94: 119–130, 1998.
- [73543] 11873. Gibbs, P. E. M.; McGregor, W. G.; Maher, V. M.; Nis-son, P.; Lawrence, C. W.: A human homolog of the *Saccharomyces cerevisiae* REV3 gene, which encodes the catalytic subunit of DNA polymerase zeta. *Proc. Nat. Acad. Sci.* 95:

6876–6880, 1998.

- [73544] 11874. Morelli, C.; Mungall, A. J.; Negrini, M.; Barbanti-Brodano, G.; Croce, C. M.: Alternative splicing, genomic structure, and fine chromosomal localization of REV3L. *Cytogenet. Cell Genet.* 83: 18–20, 1998.
- [73545] 11875. Xiao, W.; Lechler, T.; Chow, B. L.; Fontanie, T.; Augustus, M.; Carter, K. C.; Wei, Y.-F.: Identification, chromosomal mapping and tissue-specific expression of hREV3 encoding a putative human DNA polymerase zeta. *Carcinogenesis* 19: 945–949, 1998.
- [73546] 11876. Kahn, M. L.; Zheng, Y.-W.; Huang, W.; Bigornia, V.; Zeng, D.; Moff, S.; Farese, R. V., Jr.; Tam, C.; Coughlin, S. R.: A dual thrombin receptor system for platelet activation. *Nature* 394: 690–694, 1998.
- [73547] 11877. Xu, W.-F.; Andersen, H.; Whitmore, T. E.; Presnell, S. R.; Yee, D. P.; Ching, A.; Gilbert, T.; Davie, E. W.; Foster, D. C.: Cloning and characterization of human protease-activated receptor 4. *Proc. Nat. Acad. Sci.* 95: 6642–6646, 1998.
- [73548] 11878. Santoro, B.; Grant, S. G. N.; Bartsch, D.; Kandel, E. R.: Interactive cloning with the SH3 domain of N-src identifies a new brain specific ion channel protein, with homology to Eag and cyclic nucleotide-gated channels. *Proc. Nat.*

Acad. Sci. 94: 14815–14820, 1997.

- [73549] 11879.Santoro, B.; Liu, D. T.; Yao, H.; Bartsch, D.; Kandel, E. R.; Siegelbaum, S. A.; Tibbs, G. R.: Identification of a gene encoding a hyperpolarization-activated pacemaker channel of brain. *Cell* 93: 717–729, 1998.
- [73550] 11880.Stevens, D. R.; Seifert, R.; Bufe, B.; Muller, F.; Kremmer, E.; Gauss, R.; Meyerhof, W.; Kaupp, U. B.; Lindemann, B.: Hyperpolarization-activated channels HCN1 and HCN4 mediate responses to sour stimuli. *Nature* 413:631–635, 2001.
- [73551] 11881.Wainger, B. J.; DeGennaro, M.; Santoro, B.; Siegelbaum, S. A.; Tibbs, G. R.: Molecular mechanism of cAMP modulation of HCN pacemaker channels. *Nature* 411: 805–810, 2001.
- [73552] 11882.Ludwig, A.; Zong, X.; Stieber, J.; Hullin, R.; Hofmann, F.; Biel, M.: Two pacemaker channels from human heart with profoundly different activation kinetics. *EMBO J.* 18: 2323–2329, 1999.
- [73553] 11883.Casari, G.; De Fusco, M.; Ciarmatori, S.; Zeviani, M.; Mora, M.; Fernandez, P.; De Michele, G.; Filla, A.; Coccozza, S.; Marconi, R.; Durr, A.; Fontaine, B.; Ballabio, A.: Spastic paraplegia and OXPHOS impairment caused by mutations in paraplegin, a nuclear-encoded mitochondrial metallo-

protease. Cell 93: 973–983, 1998.

- [73554] 11884. De Michele, G.; De Fusco, M.; Cavalcanti, F.; Filla, A.; Marconi, R.; Volpe, G.; Monticelli, A.; Ballabio, A.; Casari, G.; Coccozza, S.: A new locus for autosomal recessive hereditary spastic paraplegia maps to chromosome 16q24.3. Am. J. Hum. Genet. 63: 135–139, 1998.
- [73555] 11885. Settasatian, C.; Whitmore, S. A.; Crawford, J.; Bilton, R. L.; Cleton-Jansen, A.-M.; Sutherland, G. R.; Callen, D. F.: Genomic structure and expression analysis of the spastic paraplegia gene, SPG7. Hum. Genet. 105: 139–144, 1999.
- [73556] 11886. de Lecea, L.; Criado, J. R.; Prospero-Garcia, O.; Gautvik, K. M.; Schweitzer, P.; Danielson, P. E.; Dunlop, C. L. M.; Siggins, G. R.; Henriksen, S. J.; Sutcliffe, J. G.: A cortical neuropeptide with neuronal depressant and sleep-modulating properties. Nature 381: 242–245, 1996.
- [73557] 11887. Kerjaschki, D.; Farquhar, M. G.: Immunocytochemical localization of the Heymann nephritis antigen (GP330) in glomerular epithelial cells of normal Lewis rats. J. Exp. Med. 157: 667–686, 1983.
- [73558] 11888. Korenberg, J. R.; Argraves, K. M.; Chen, X.-N.; Tran, H.; Strickland, D.; Argraves, W. S.: Chromosomal localization of human genes for the LDL receptor family

member glycoprotein 330 (LRP2) and its associated protein RAP (LRPAP1). *Genomics* 22: 88–93, 1994.

[73559] 11889. Kounnas, M. Z.; Loukinova, E. B.; Stefansson, S.; Harmony, J. A.K.; Brewer, B. H.; Strickland, D. K.; Argraves, W. S.: Identification of glycoprotein 330 as an endocytic receptor for apolipoprotein J/clusterin. *J. Biol. Chem.* 270: 13070–13075, 1995.

[73560] 11890. Leheste, J.-R.; Rolinski, B.; Vorum, H.; Hilpert, J.; Nykjaer, A.; Jacobsen, C.; Aucouturier, P.; Moskaug, J. O.; Otto, A.; Christensen, E. I.; Willnow, T. E.: Megalin knock-out mice as an animal model of low molecular weight proteinuria. *Am. J. Path.* 155: 1361–1370, 1999.

[73561] 11891. Marino, M.; Chiovato, L.; Friedlander, J. A.; Latrofa, F.; Pinchera, A.; McCluskey, R. T.: Serum antibodies against megalin (GP330) in patients with autoimmune thyroiditis. *J. Clin. Endocr. Metab.* 84: 2468–2474, 1999.

[73562] 11892. Moestrup, S. K.; Cui, S.; Vorum, H.; Bregengard, C.; Bjorn, S.E.; Norris, K.; Gliemann, J.; Christensen, E. I.: Evidence that epithelial glycoprotein 330/megalin mediates uptake of polybasic drugs. *J. Clin. Invest.* 96: 1404–1413, 1995.

[73563] 11893. Nykjaer, A.; Dragun, D.; Walther, D.; Vorum, H.; Jacobsen, C.; Herz, J.; Melsen, F.; Christensen, E. I.; Willnow,

T. E.: An endocytic pathway essential for renal uptake and activation of the steroid 25-(OH)vitamin D3. *Cell* 96: 507–515, 1999.

- [73564] 11894. Raychowdhury, R.; Niles, J. L.; McCluskey, R. T.; Smith, J. A.: Autoimmune target in Heymann nephritis is a glycoprotein with homology to the LDL receptor. *Science* 244: 1163–1165, 1989.
- [73565] 11895. Saito, A.; Pietromonaco, S.; Loo, A. K.-C.; Farquhar, M. G.: Complete cloning and sequencing of rat gp330/`megalin,' a distinctive member of the low density lipoprotein receptor gene family. *Proc. Nat. Acad. Sci.* 91: 9725–9729, 1994.
- [73566] 11896. Schmitz, C.; Hilpert, J.; Jacobsen, C.; Boensch, C.; Christensen, E. I.; Luft, F. C.; Willnow, T. E.: Megalin deficiency offers protection from renal aminoglycoside accumulation. *J. Biol. Chem.* 277: 618–622, 2002.
- [73567] 11897. Willnow, T. E.; Hilpert, J.; Armstrong, S. A.; Rohlmann, A.; Hammer, R. E.; Burns, D. K.; Herz, J.: Defective forebrain development in mice lacking gp330/megalin. *Proc. Nat. Acad. Sci.* 93: 8460–8464, 1996.
- [73568] 11898. Gostout, B.; Liu, Q.; Sommer, S. S.: 'Cryptic' repeating tripletsof purines and pyrimidines (cRRY(i)) are fre-

quent and polymorphic:analysis of coding cRRY(i) in the proopiomelanocortin (POMC) and TATA-bindingprotein (TBP) genes. *Am. J. Hum. Genet.* 52: 1182–1190, 1993.

[73569] 11899.Hobbs, N. K.; Bondareva, A. A.; Barnett, S.; Capecchi, M. R.; Schmidt,E. E.: Removing the vertebrate-specific TBP N terminus disrupts placentalbeta-2M-dependent interactions with the maternal immune system. *Cell* 110:43–54, 2002.

[73570] 11900.Imbert, G.; Trottier, Y.; Beckmann, J.; Mandel, J. L.: The gene for the TATA binding protein (TBP) that contains a highly polymorphicprotein coding CAG repeat maps to 6q27. *Genomics* 21: 667–668, 1994.

[73571] 11901.Jones, A. L.; Middle, F.; Guy, C.; Spurlock, G.; Cairns, N. J.; McGuffin, P.; Craddock, N.; Owen, M.; O'Donovan, M. C.: No evidence for expanded polyglutamine sequences in bipolar disorder and schizophrenia. *Molec.Psychiat.* 2: 478–482, 1997.

[73572] 11902.Kao, C. C.; Lieberman, P. M.; Schmidt, M. C.; Zhou, Q.; Pei, R.;Berk, A. J.: Cloning of a transcriptionally active human TATA bindingfactor. *Science* 248: 1646–1650, 1990.

[73573] 11903.Koide, R.; Kobayashi, S.; Shimohata, T.; Ikeuchi, T.; Maruyama,M.; Saito, M.; Yamada, M.; Takahashi, H.; Tsuji,



S.: A neurological disease caused by an expanded CAG trinucleotide repeat in the TATA-binding protein gene: a new polyglutamine disease? Hum. Molec. Genet. 8:2047–2053, 1999.

[73574] 11904. Nakamura, K.; Jeong, S.-Y.; Uchihara, T.; Anno, M.; Nagashima, K.; Nagashima, T.; Ikeda, S.; Tsuji, S.; Kanazawa, I.: SCA17, a novel autosomal dominant cerebellar ataxia caused by an expanded polyglutamine in TATA-binding protein. Hum. Molec. Genet. 10: 1441–1448, 2001.

[73575] 11905. Nikolov, D. B.; Hu, S.-H.; Lin, J.; Gasch, A.; Hoffmann, A.; Horikoshi, M.; Chua, N.-H.; Roeder, R. G.; Burley, S. K.: Crystal structure of TFIID TATA-box binding protein. Nature 360: 40–46, 1992.

[73576] 11906. Peterson, M. G.; Tanese, N.; Pugh, B. F.; Tjian, R.: Functional domains and upstream activation properties of cloned human TATA binding protein. Science 248: 1625–1630, 1990.

[73577] 11907. Polymeropoulos, M. H.; Rath, D. S.; Xiao, H.; Merrill, C. R.: Trinucleotide repeat polymorphism at the human transcription factor IID gene. Nucleic Acids Res. 19: 4307 only, 1991.

[73578] 11908. Rosen, D. R.; Trofatter, J. A.; Brown, R. H., Jr.: Map-

ping of the human TATA-binding protein gene (TBP) to chromosome 6qter. *Cytogenet. Cell Genet.* 69: 279–280, 1995.

- [73579] 11909. Rubinsztein, D. C.; Leggo, J.; Crow, T. J.; DeLisi, L. E.; Walsh, C.; Jain, S.; Paykel, E. S.: Analysis of polyglutamine-coding repeats in the TATA-binding protein in different human populations and in patients with schizophrenia and bipolar affective disorder. *Am. J. Med. Genet.* 67: 495–498, 1996.
- [73580] 11910. Saito, F.; Yamamoto, T.; Horikoshi, M.; Ikeuchi, T.: Direct mapping of the human TATA box-binding protein (TBP) gene to 6q27 by fluorescence in situ hybridization. *Jpn. J. Hum. Genet.* 39: 421–425, 1994.
- [73581] 11911. Serrano, M.; Hannon, G. J.; Beach, D.: A new regulatory motif in cell-cycle control causing specific inhibition of cyclin D/CDK4. *Nature* 366: 704–707, 1993.
- [73582] 11912. Testa, J. R.; Zhou, J.; Bell, D. W.; Yen, T. J.: Chromosomal localization of the genes encoding the kinetochore proteins CENPE and CENPF to human chromosomes 4q24–q25 and 1q32–q41, respectively, by fluorescence in situ hybridization. *Genomics* 23: 691–693, 1994.
- [73583] 11913. Pennica, D.; Swanson, T. A.; Shaw, K. J.; Kuang, W.-J.; Gray, C. L.; Beatty, B. G.; Wood, W. I.: Human car-

diotrophin-1: protein and gene structure, biological and binding activities, and chromosomal localization. Cytokine 8: 183-189, 1996.

[73584] 11914. Fukudome, K.; Esmon, C. T.: Identification, cloning, and regulation of a novel endothelial cell protein C/activated protein C receptor. J. Biol. Chem. 269: 26486-26491, 1994.

[73585] 11915. Hayashi, T.; Nakamura, H.; Okada, A.; Takebayashi, S.; Wakita, T.; Yuasa, H.; Okumura, K.; Suzuki, K.: Organization and chromosomal localization of the human endothelial protein C receptor gene. Gene 238: 367-373, 1999.

[73586] 11916. Mincheva, A.; Rothbarth, K.; Werner, D.; Lichter, P.: Assignment of the gene encoding centrosome-associated protein CCD41 to mouse chromosome 2H. Mammalian Genome 6: 444 only, 1995.

[73587] 11917. Rothbarth, K.; Dabaghian, A. R. H.; Stammer, H.; Werner, D.: One single mRNA encodes the centrosomal protein CCD41 and the endothelial cell protein C receptor (EPCR). FEBS Lett. 458: 77-80, 1999.

[73588] 11918. Rothbarth, K.; Petzelt, C.; Lu, X.; Todorov, I. T.; Joswig, G.; Pepperkok, R.; Ansorge, W.; Werner, D.: cDNA-derived molecular characteristics and antibodies to a new

centrosome-associated and G2/M phase-prevalent protein. *J. Cell Sci.* 104: 19–30, 1993.

[73589] 11919. Simmonds, R. E.; Lane, D. A.: Structural and functional implications of the intron/exon organization of the human endothelial cell protein C/activated protein C receptor (EPCR) gene: comparison with the structure of CD1/major histocompatibility complex alpha-1 and alpha-2 domains. *Blood* 94:632–641, 1999.

[73590] 11920. Ye, X.; Fukudome, K.; Tsuneyoshi, N.; Satoh, T.; Tokunaga, O.; Sugawara, K.; Mizokami, H.; Kimoto, M.: The endothelial cell protein C receptor (EPCR) functions as a primary receptor for protein C activation on endothelial cells in arteries, veins, and capillaries. *Biochem. Biophys. Res. Commun.* 259: 671–677, 1999.

[73591] 11921. Purrello, M.; Pietro, C. D.; Mirabile, E.; Rapisarda, A.; Rimini, R.; Tine, A.; Pavone, L.; Motta, S.; Grzeschik, K.-H.; Sichel, G.: Physical mapping at 6q27 of the locus for the TATA box-binding protein, the DNA-binding subunit of TFIID and a component of SL1 and TFIIB, strongly suggests that it is single copy in the human genome. *Genomics* 22:94–100, 1994.

[73592] 11922. Candia, A. F.; Hu, J.; Crosby, J.; Lalley, P. A.; Noden, D.; Nadeau, J. H.; Wright, C. V. E.: Mox-1 and Mox-2 de-

fine a novel homeobox gene subfamily and are differentially expressed during early mesodermal patterning in mouse embryos. *Development* 116: 1123–1136, 1992.

[73593] 11923. Gorski, D. H.; LePage, D. F.; Patel, C. V.; Copeland, N. G.; Jenkins, N. A.; Walsh, K.: Molecular cloning of a diverged homeobox gene that is rapidly down-regulated during the G0/G1 transition in vascular smooth muscle cells. *Molec. Cell Biol.* 13: 3722–3733, 1993.

[73594] 11924. Grigoriou, M.; Kastrinaki, M.-C.; Modi, W. S.; Theodorakis, K.; Mankoo, B.; Pachnis, V.; Karagogeos, D.: Isolation of the human MOX2 homeobox gene and localization to chromosome 7p22.1–p21.3. *Genomics* 26:550–555, 1995.

[73595] 11925. LePage, D. F.; Altomare, D. A.; Testa, J. R.; Walsh, K.: Molecular cloning and localization of the human GAX gene to 7p21. *Genomics* 24:535–540, 1994.

[73596] 11926. Mankoo, B. S.; Collins, N. S.; Ashby, P.; Grigorieva, E.; Pevny, L. H.; Candia, A.; Wright, C. V. E.; Rigby, P. W. J.; Pachnis, V.: Mox2 is a component of the genetic hierarchy controlling limb muscle development. *Nature* 400: 69–73, 1999.

[73597] 11927. Lindor, N. M.; Furuichi, Y.; Kitao, S.; Shimamoto, A.; Arndt, C.; Jalal, S.: Rothmund–Thomson syndrome due to

RECQ4 helicase mutations: report and clinical and molecular comparisons with Bloom syndrome and Werner syndrome. *Am. J. Med. Genet.* 90: 223–228, 2000.

[73598] 11928. Puranam, K. L.; Blackshear, P. J.: Cloning and characterization of RECQL, a potential human homologue of the *Escherichia coli* DNA helicase RecQ. *J. Biol. Chem.* 269: 29838–29845, 1994.

[73599] 11929. Puranam, K. L.; Kennington, E.; J.-Sait, S. N.; Shows, T. B.; Rochelle, J. M.; Seldin, M. F.; Blackshear, P. J.: Chromosomal localization of the gene encoding the human DNA helicase RECQL and its mouse homologue. *Genomics* 26: 595–598, 1995.

[73600] 11930. Gollner, H.; Bouwman, P.; Mangold, M.; Karis, A.; Braun, H.; Rohner, I.; Del Rey, A.; Besedovsky, H.-O.; Meinhardt, A.; van den Broek, M.; Cutforth, T.; Grosveld, F.; Philipsen, S.; Suske, G.: Complex phenotype of mice homozygous for a null mutation in the Sp4 transcription factor gene. *Genes Cells* 6: 689–697, 2001.

[73601] 11931. Hagen, G.; Dennig, J.; Preiss, M.; Beato, M.; Suske, G.: Functional analyses of the transcription factor Sp4 reveal properties distinct from Sp1 and Sp3. *J. Biol. Chem.* 270: 24989–24994, 1995.

[73602] 11932. Hagen, G.; Muller, S.; Beato, M.; Suske, G.: Cloning

by recognitionsite screening of two novel GT box binding proteins: a family of Sp1related genes. *Nucleic Acids Res.* 20: 5519–5525, 1992.

- [73603] 11933.Kalff–Suske, M.; Kunz, J.; Grzeschik, K.–H.; Suske, G.: HumanSp4 transcription factor gene (SP4) maps to chromosome 7p15. *Genomics* 26:631–633, 1995.
- [73604] 11934.Nguyen–Tran, V. T. B.; Kubalak, S. W.; Minamisawa, S.; Fiset, C.;Wollert, K. C.; Brown, A. B.; Ruiz–Lozano, P.; Barrere–Lemaire, S.;Kondo, R.; Norman, L. W.; Gourdie, R. G.; Rahme, M. M.; Feld, G. K.;Clark, R. B.; Giles, W. R.; Chien, K. R.: A novel genetic pathwayfor sudden cardiac death via defects in the transition between ventricularand conduction system cell lineages. *Cell* 102: 671–682, 2000.
- [73605] 11935.Lamour, V.; Lecluse, Y.; Desmaze, C.; Spector, M.; Bodescot, M.;Aurias, A.; Osley, M. A.; Lipinski, M.: A human homolog of the *S.cerevisiae* HIR1 and HIR2 transcriptional repressors cloned from theDiGeorge syndrome critical region. *Hum. Molec. Genet.* 4: 791–799,1995.
- [73606] 11936.Llevadot, R.; Scambler, P.; Estivill, X.; Pritchard, M.: Genomicorganization of TUPLE1/HIRA: a gene implicated in DiGeorge syndrome. *MammalianGenome* 7: 911–914, 1996.

- [73607] 11937.Lorain, S.; Quivy, J.-P.; Monier-Gavelle, F.; Scamps, C.; Lecluse, Y.; Almouzni, G.; Lipinski, M.: Core histones and HIRIP3, a novel histone-binding protein, directly interact with WD repeat protein HIRA. *Molec. Cell. Biol.* 18: 5546–5556, 1998.
- [73608] 11938.Magnaghi, P.; Roberts, C.; Lorain, S.; Lipinski, M.; Scambler, P. J.: HIRA, a mammalian homologue of *Saccharomyces cerevisiae* transcriptional co-repressors, interacts with Pax3. *Nature Genet.* 20: 74–77, 1998.
- [73609] 11939.Mattei, M.-G.; Halford, S.; Scambler, P. J.: Mapping of the *TUPLE1* gene to mouse chromosome 16A-B1. *Genomics* 23: 717–718, 1994.
- [73610] 11940.Roberts, C.; Daw, S. C. M.; Halford, S.; Scambler, P. J.: Cloning and developmental expression analysis of chick Hira (*Chira*), a candidate gene for DiGeorge syndrome. *Hum. Molec. Genet.* 6: 237–245, 1997.
- [73611] 11941.Wilming, L. G.; Snoeren, C. A. S.; van Rijswijk, A.; Grosveld, F.; Meijers, C.: The murine homologue of HIRA, a DiGeorge syndrome candidate gene, is expressed in embryonic structures affected in human CATCH22 patients. *Hum. Molec. Genet.* 6: 247–258, 1997.
- [73612] 11942.Kim, I.-G.; Gorman, J. J.; Park, S.-C.; Chung, S.-I.; Steinert, P. M.: The deduced sequence of the novel pro-



transglutaminase E (TGase3) of human and mouse. J. Biol. Chem. 268: 12682–12690, 1993.

- [73613] 11943. Kim, I.-G.; Lee, S.-C.; Lee, J.-H.; Yang, J.-M.; Chung, S.-I.; Steinert, P. M.: Structure and organization of the human transglutaminase3 gene: evolutionary relationship to the transglutaminase family. J. Invest. Derm. 103: 137–142, 1994.
- [73614] 11944. Marchese, A.; Docherty, J. M.; Nguyen, T.; Heiber, M.; Cheng, R.; Heng, H. H. Q.; Tsui, L.-C.; Shi, X.; George, S. R.; O'Dowd, B. F.: Cloning of human genes encoding novel G protein-coupled receptors. Genomics 23:609–618, 1994.
- [73615] 11945. Homey, B.; Alenius, H.; Muller, A.; Soto, H.; Bowman, E. P.; Yuan, W.; McEvoy, L.; Lauerma, A. I.; Assmann, T.; Bunemann, E.; Lehto, M.; Wolff, H.; Yen, D.; Marxhausen, H.; To, W.; Sedgwick, J.; Ruzicka, T.; Lehmann, P.; Zlotnik, A.: CCL27–CCR10 interactions regulate T cell-mediated skin inflammation. Nature Med. 8: 157–165, 2002.
- [73616] 11946. Homey, B.; Wang, W.; Soto, H.; Buchanan, M. E.; Wiesenborn, A.; Catron, D.; Muller, A.; McClanahan, T. K.; Dieu-Nosjean, M.-C.; Orozco, R.; Ruzicka, T.; Lehmann, P.; Oldham, E.; Zlotnik, A.: Cutting edge: the orphan

chemokine receptor G protein-coupled receptor-2 (GPR-2,CCR10) binds the skin-associated chemokine CCL27 (CTACK/ALP/ILC). J.Immun. 164: 3465-3470, 2000.

[73617] 11947.Jarmin, D. I.; Rits, M.; Bota, D.; Gerard, N. P.; Graham, G. J.;Clark-Lewis, I.; Gerard, C.: Cutting edge: identification of the orphan receptor G-protein-coupled receptor 2 as CCR10, a specific receptor for the chemokine ESKine. J. Immun. 164: 3460-3464, 2000.

[73618] 11948.Wang, W.; Soto, H.; Oldham, E. R.; Buchanan, M. E.; Homey, B.;Catron, D.; Jenkins, N.; Copeland, N. G.; Gilbert, D. J.; Nguyen,N.; Abrams, J.; Kershenovich, D.; Smith, K.; McClanahan, T.; Vicari,A. P.; Zlotnik, A.: Identification of a novel chemokine (CCL28),which binds CCR10 (GPR2). J. Biol. Chem. 275: 22313-22323, 2000.

[73619] 11949.Akao, Y.; Matsuda, Y.: Identification and chromosome mapping of the mouse homologue of the human gene (DDX6) that encodes a putative RNA helicase of the DEAD box protein family. Cytogenet. Cell Genet. 75:38-44, 1996.

[73620] 11950.Akao, Y.; Seto, M.; Takahashi, T.; Kubonishi, I.; Miyoshi, I.;Nakazawa, S.; Tsujimoto, Y.; Croce, C. M.; Ueda, R.: Molecular cloning of the chromosomal breakpoint of a

B-cell lymphoma with the t(11;14)(q23;q32) chromosome translocation. *Cancer Res.* 51: 1574–1576, 1991.

- [73621] 11951. Akao, Y.; Seto, M.; Yamamoto, K.; Iida, S.; Nakazawa, S.; Inazawa, J.; Abe, T.; Takahashi, T.; Ueda, R.: The RCK gene associated with t(11;14) translocation is distinct from the MLL/ALL-1 gene with t(4;11) and t(11;19) translocations. *Cancer Res.* 52: 6083–6087, 1992.
- [73622] 11952. Akao, Y.; Tsujimoto, Y.; Finan, J.; Nowell, P. C.; Croce, C. M.: Molecular characterization of a t(11;14)(q23;q32) chromosome translocation in a B-cell lymphoma. *Cancer Res.* 50: 4856–4859, 1990.
- [73623] 11953. Lu, D.; Yunis, J. J.: Cloning, expression and localization of an RNA helicase gene from a human lymphoid cell line with chromosomal breakpoint 11q23.3. *Nucleic Acids Res.* 20: 1967–1972, 1992.
- [73624] 11954. Seto, M.; Yamamoto, K.; Takahashi, T.; Ueda, R.: Cloning and expression of a murine cDNA homologous to the human RCK/P54, a lymphoma-linked chromosomal translocation junction gene on 11q23. *Gene* 166: 293–296, 1995.
- [73625] 11955. Tunnacliffe, A.; Perry, H.; Radice, P.; Budarf, M. L.; Emanuel, B. S.: A panel of sequence tagged sites for chromosome band 11q23. *Genomics* 17: 744–747, 1993.

- [73626] 11956.Hallier, E.; Jager, R.; Deutschmann, S.; Bolt, H. M.; Peter, H.: Glutathione conjugation and cytochrome P-450 metabolism of methylchloride in vitro. *Toxicol. in Vitro* 4: 513-517, 1990.
- [73627] 11957.Pemle, S.; Schroeder, K. R.; Spencer, S. R.; Meyer, D. J.; Hallier,E.; Bolt, H. M.; Ketterer, B.; Taylor, J. B.: Human glutathione S-transferase theta (GSTT1): cDNA cloning and the characterization of a genetic polymorphism. *Biochem. J.* 300: 271-276, 1994.
- [73628] 11958.Peter, H.; Deutschmann, S.; Reichel, C.; Hallier, E.: Metabolism of methyl chloride by human erythrocytes. *Arch. Toxicol.* 63: 351-355, 1989.
- [73629] 11959.Schroder, K. R.; Wiebel, F. A.; Reich, S.; Dannappel, D.; Bolt,H. M.; Hallier, E.: Glutathione S-transferase (GST) theta polymorphism influences background SCE rate. *Arch. Toxicol.* 69: 505-507, 1995.
- [73630] 11960.Webb, G.; Vaska, V.; Coggan, M.; Board, P.: Chromosomal localization of the gene for the human theta class glutathione transferase (GSTT1). *Genomics* 33:121-123, 1996.
- [73631] 11961.Wiebel, F. A.; Dommermuth, A.; Thier, R.: The hereditary transmission of the glutathione transferase hGSTT1-1 conjugator phenotype in a large family. *Phar-*

macogenetics 9: 251–256, 1999.

- [73632] 11962.Dumon, K. R.; Ishii, H.; Fong, L. Y. Y.; Zanesi, N.; Fidanza, V.; Mancini, R.; Vecchione, A.; Baffa, R.; Trapasso, F.; During, M. J.; Huebner, K.; Croce, C. M.: FHIT gene therapy prevents tumor development in Fhit-deficient mice. Proc. Nat. Acad. Sci. 98: 3346–3351, 2001.
- [73633] 11963.Ikeda, A.; Zheng, Q. Y.; Zuberi, A. R.; Johnson, K. R.; Naggert, J. K.; Nishina, P. M.: Microtubule-associated protein 1A is a modifier of tubby hearing (moth1). Nature Genet. 30: 401–405, 2002.
- [73634] 11964.Roy, S. K.; Hu, J.; Meng, Q.; Xia, Y.; Shapiro, P. S.; Reddy, S.P. M.; Platanius, L. C.; Lindner, D. J.; Johnson, P. F.; Pritchard, C.; Pages, G.; Pouyssegur, J.; Kalvakolanu, D. V.: MEKK1 plays a critical role in activating the transcription factor C/EBP-beta-dependent gene expression in response to IFN-gamma. Proc. Nat. Acad. Sci. 99:7945–7950, 2002.
- [73635] 11965.Chou, D.; Miyashita, T.; Mohrenweiser, H. W.; Ueki, K.; Kastury, K.; Druck, T.; von Deimling, A.; Huebner, K.; Reed, J. C.; Louis, D. N.: The BAX gene maps to the glioma candidate region at 19q13.3, but is not altered in human gliomas. Cancer Genet. Cytogenet. 88:136–140, 1996.
- [73636] 11966.Deckwerth, T. L.; Elliott, J. L.; Knudson, C. M.;

Johnson, E. M., Jr.; Snider, W. D.; Korsmeyer, S. J.: BAX is required for neuronal death after trophic factor deprivation and during development. *Neuron* 17:401–411, 1996.

[73637] 11967. Knudson, C. M.; Tung, K. S. K.; Tourtellotte, W. G.; Brown, G. A. J.; Korsmeyer, S. J.: Bax-deficient mice with lymphoid hyperplasia and male germ cell death. *Science* 270: 96–99, 1995.

[73638] 11968. LeBlanc, H.; Lawrence, D.; Varfolomeev, E.; Totpal, K.; Morlan, J.; Schow, P.; Fong, S.; Schwall, R.; Sinicropi, D.; Ashkenazi, A.: Tumor-cell resistance to death receptor-induced apoptosis through mutational inactivation of the proapoptotic Bcl-2 homolog Bax. *Nature Med.* 8: 274–281, 2002.

[73639] 11969. Lindsten, T.; Ross, A. J.; King, A.; Zong, W.-X.; Rathmell, J. C.; Shiels, H. A.; Ulrich, E.; Waymire, K. G.; Mahar, P.; Frauwirth, K.; Chen, Y.; Wei, M.; and 9 others: The combined functions of proapoptotic Bcl-2 family members Bak and Bax are essential for normal development of multiple tissues. *Molec. Cell* 6: 1389–1399, 2000.

[73640] 11970. Marzo, I.; Brenner, C.; Zamzami, N.; Jurgensmeier, J. M.; Susin, S. A.; Vieira, H. L. A.; Prevost, M.-C.; Xie, Z.; Matsuyama, S.; Reed, J. C.; Kroemer, G.: Bax and adenine nucleotide translocator cooperate in the mitochondrial

control of apoptosis. *Science* 281: 2027–2031,1998.

- [73641] 11971.Matikainen, T.; Perez, G. I.; Jurisicova, A.; Pru, J. K.; Schlezinger, J. J.; Ryu, H.-Y.; Laine, J.; Sakai, T.; Korsmeyer, S. J.; Casper, R. F.; Sherr, D. H.; Tilly, J. L.: Aromatic hydrocarbon receptor-driven Bax gene expression is required for premature ovarian failure caused by biohazardous environmental chemicals. *Nature Genet.* 28: 355–360,2001.
- [73642] 11972.Matsuda, Y.; Kusano, H.; Tsujimoto, Y.: Chromosomal assignment of the Bcl2-related genes, Bcl2l and Bax, in the mouse and rat. *Cytogenet. Cell Genet.* 74: 107–110, 1996.
- [73643] 11973.Meijerink, J. P. P.; Mensink, E. J. B. M.; Wang, K.; Sedlak, T.W.; Sloetjes, A. W.; de Witte, T.; Waksman, G.; Korsmeyer, S. J.: Hematopoietic malignancies demonstrate loss-of-function mutations of BAX. *Blood* 91: 2991–2997, 1998.
- [73644] 11974.Miyashita, T.; Reed, J. C.: Tumor suppressor p53 is a direct transcriptional activator of the human bax gene. *Cell* 80: 293–299,1995.
- [73645] 11975.Oltvai, Z. N.; Millman, C. L.; Korsmeyer, S. J.: Bcl-2 heterodimers in vivo with a conserved homolog, Bax, that accelerates programmed cell death. *Cell* 74: 609–619, 1993.

- [73646] 11976.Perez, G. I.; Robles, R.; Knudson, C. M.; Flaws, J. A.; Korsmeyer,S. J.; Tilly, J. L.: Prolongation of ovarian lifespan into advancedchronological age by Bax–deficiency. *Nature Genet.* 21: 200–203,1999.
- [73647] 11977.Rampino, N.; Yamamoto, H.; Ionov, Y.; Li, Y.; Sawai, H.; Reed,J. C.; Perucho, M.: Somatic frameshift mutations in the BAX genein colon cancers of the microsatellite mu–tator phenotype. *Science* 275:967–969, 1997.
- [73648] 11978.Suzuki, M.; Youle, R. J.; Tjandra, N.: Structure of Bax: coregulationof dimer formation and intracellular lo–calization. *Cell* 103: 645–654,2000.
- [73649] 11979.Wei, M. C.; Zong, W.–X.; Cheng, E. H.–Y.; Lindsten, T.; Panoutsakopoulou,V.; Ross, A. J.; Roth, K. A.; MacGre–gor, G. R.; Thompson, C. B.; Korsmeyer,S. J.: Proapoptotic BAX or BAK: a requisite gateway to mitochondrial dysfunc–tion and death. *Science* 292: 727–730, 2001.
- [73650] 11980.Zhang, L.; Yu, J.; Park, B. H.; Kinzler, K. W.; Vogel–stein, B.: Role of BAX in the apoptotic response to anti–cancer agents. *Science* 290:989–992, 2000.
- [73651] 11981.Dasari, V. R.; Sandhu, A. K.; Mills, D. C. B.; Athwal, R. S.; Kunapuli,S. P.: Mapping of the P2U purinergic recep–tor gene to human chromosome11q13.5–14.1. *Somat. Cell Molec. Genet.* 22: 75–79, 1996.



- [73652] 11982.Katzur, A. C.; Koshimizu, T.-A.; Tomic, M.; Schultze-Mosgau, A.;Ortmann, O.; Stojilkovic, S. S.: Expression and responsiveness ofP2Y2 receptors in human endometrial cancer cell lines. J. Clin. Endocr.Metab. 84: 4085–4091, 1999.
- [73653] 11983.Parr, C. E.; Sullivan, D. M.; Paradiso, A. M.; Lazarowski, E. R.;Burch, L. H.; Olsen, J. C.; Erb, L.; Weisman, G. A.; Boucher, R. C.;Turner, J. T.: Cloning and expression of a human P(2U) nucleotidereceptor, a target for cystic fibrosis pharmacotherapy. Proc. Nat.Acad. Sci. 91: 3275–3279, 1994.
- [73654] 11984.Somers, G. R.; Hammet, F.; Woollatt, E.; Richards, R. I.; Southey,M. C.; Venter, D. J.: Chromosomal localization of the human P2Y(6)purinoceptor gene and phylogenetic analysis of the P2y purinoceptorfamily. Genomics 44: 127–130, 1997.
- [73655] 11985.Tai, C.-J.; Kang, S. K.; Cheng, K. W.; Choi, K.-C.; Nathwani, P.S.; Leung, P. C. K.: Expression and regulation of P2U–purinergicreceptor in human granulosa–luteal cells. J. Clin. Endocr. Metab. 85:1591–1597, 2000.
- [73656] 11986.Santamarina-Fojo, S.; Peterson, K.; Knapper, C.; Qiu, Y.; Freeman,L.; Cheng, J.-F.; Osorio, J.; Remaley, A.; Yang, X.-P.; Haudenschild,C.; Prades, C.; Chimini, G.;

Blackmon, E.; Francois, T.; Duverger, N.; Rubin, E. M.;  
Rosier, M.; Deneffe, P.; Fredrickson, D. S.; Brewer, H. B., Jr.:  
Complete genomic sequence of the human ABCA1 gene:  
analysis of the human and mouse ATP-binding cassette A  
promoter. *Proc. Nat. Acad. Sci.* 97: 7987–7992, 2000.  
Note: Erratum: *Proc. Nat. Acad. Sci.* 99: 1098 only, 2002.

[73657] 11987. Szakacs, G.; Langmann, T.; Ozvegy, C.; Orso, E.;  
Schmitz, G.; Varadi, A.; Sarkadi, B.: Characterization of the  
ATPase cycle of human ABCA1: implications for its function  
as a regulator rather than an active transporter. *Biochem.  
Biophys. Res. Commun.* 288: 1258–1264, 2001.

[73658] 11988. Utech, M.; Hobbel, G.; Rust, S.; Reinecke, H.; Ass-  
mann, G.; Walter, M.: Accumulation of RhoA, RhoB, RhoG,  
and Rac1 in fibroblasts from Tangier disease subjects sug-  
gests a regulatory role of Rho family proteins in choles-  
terol efflux. *Biochem. Biophys. Res. Commun.*  
280: 229–236, 2001.

[73659] 11989. Zhao, L.-X.; Zhou, C.-J.; Tanaka, A.; Nakata, M.;  
Hirabayashi, T.; Amachi, T.; Shioda, S.; Ueda, K.; Inagaki,  
N.: Cloning, characterization and tissue distribution of the  
rat ATP-binding cassette (ABC) transporter ABC2/ABCA2.  
*Biochem J.* 350: 865–872, 2000.

[73660] 11990. Zwarts, K. Y.; Clee, S. M.; Zwinderman, A. H.; En-

gert, J. C.; Singaraja, R.; Loubser, O.; James, E.; Roomp, K.; Hudson, T. J.; Jukema, J. W.; Kastelein, J. J. P.; Hayden, M. R.: ABCA1 regulatory variants influence coronary artery disease independent of effects on plasma lipid levels. Clin. Genet. 61: 115–125, 2002.

[73661] 11991. Fears, S.; Mathieu, C.; Zeleznik-Le, N.; Huang, S.; Rowley, J. D.; Nucifora, G.: Intergenic splicing of MDS1 and EVI1 occurs in normal tissues as well as in myeloid leukemia and produces a new member of the PR domain family. Proc. Nat. Acad. Sci. 93: 1642–1647, 1996.

[73662] 11992. Mochizuki, N.; Shimizu, S.; Nagasawa, T.; Tanaka, H.; Taniwaki, M.; Yokota, J.; Morishita, K.: A novel gene, MEL1, mapped to 1p36.3 is highly homologous to the MDS1/EVI1 gene and is transcriptionally activated in t(1;3)(p36;q21)–positive leukemia cells. Blood 96:3209–3214, 2000.

[73663] 11993. Ing, Y. L.; Leung, I. W. L.; Heng, H. H. Q.; Tsui, L.-C.; Lassam, N. J.: MLK-3: identification of a widely-expressed protein kinase bearing an SH3 domain and a leucine zipper–basic region domain. Oncogene 9:1745–1750, 1994.

[73664] 11994. Bernard, O. A.; Mauchauffe, M.; Mecucci, C.; Van Den Berghe, H.; Berger, R.: A novel gene, AF-1p, fused to

HRX in t(1;11)(p32;q23), is not related to AF-4, AF-9 nor ENL. *Oncogene* 9: 1039–1045, 1994.

[73665] 11995. Fazioli, F.; Minichiello, L.; Matoska, V.; Castagnino, P.; Miki, T.; Wong, W. T.; Di Fiore, P. P.: Eps8, a substrate for the epidermal growth factor receptor kinase, enhances EGF-dependent mitogenic signals. *EMBO J.* 12: 3799–3808, 1993.

[73666] 11996. Fazioli, F.; Minichiello, L.; Matoskova, B.; Wong, W. T.; Di Fiore, P. P.: eps 15, a novel tyrosine kinase substrate, exhibits transforming activity. *Molec. Cell. Biol.* 13: 5814–5828, 1993.

[73667] 11997. Fazioli, F.; Wong, W. T.; Ullrich, S. J.; Sakaguchi, K.; Appella, E.; Di Fiore, P. P.: The ezrin-like family of tyrosine kinase substrates: receptor-specific pattern of tyrosine phosphorylation and relationship to malignant transformation. *Oncogene* 8: 1335–1345, 1993.

[73668] 11998. Wong, W. T.; Kraus, M. H.; Carlomagno, F.; Zelano, A.; Druck, T.; Croce, C. M.; Huebner, K.; Di Fiore, P. P.: The human eps15 gene, encoding a tyrosine kinase substrate, is conserved in evolution and maps to 1p31–p32. *Oncogene* 9: 1591–1597, 1994.

[73669] 11999. Kozu, T.; Henrich, B.; Schafer, K. P.: Structure and expression of the gene (HNRPA2B1) encoding the human

hnRNP protein A2/B1. Genomics 25:365–371, 1995.

- [73670] 12000.Kordeli, E.; Lambert, S.; Bennett, V.: Ankyrin-G: a new ankyringene with neural-specific isoforms localized at the axonal initial segment and node of Ranvier. J. Biol. Chem. 270: 2352–2359, 1995.
- [73671] 12001.Hsu, L. C.; Chang, W.-C.; Yoshida, A.: Human aldehyde dehydrogenase genes, ALDH7 and ALDH8: genomic organization and gene structure comparison. Gene 189:89–94, 1997.
- [73672] 12002.Hsu, L. C.; Chang, W.-C.; Yoshida, A.: Cloning of a cDNA encoding human ALDH7, a new member of the aldehyde dehydrogenase family. Gene 151:285–289, 1994.
- [73673] 12003.Aruga, J.; Yokota, N.; Hashimoto, M.; Furuichi, T.; Fukuda, M.; Mikoshiba, K.: A novel zinc finger protein, Zic, is involved in neurogenesis, especially in the cell lineage of cerebellar granule cells. J. Neurochem. 63:1880–1890, 1994.
- [73674] 12004.Salero, E.; Perez-Sen, R.; Aruga, J.; Gimenez, C.; Zafra, F.: Transcription factors Zic1 and Zic2 bind and transactivate the apolipoprotein E gene promoter. J. Biol. Chem. 276: 1881–1888, 2001.
- [73675] 12005.Yokota, N.; Aruga, J.; Takai, S.; Yamada, K.; Hamazaki, M.; Iwase, T.; Sugimura, H.; Mikoshiba, K.: Pre-

dominant expression of humanZic in cerebellar granule cell lineage and medulloblastoma. CancerRes. 56: 377–383, 1996.

- [73676] 12006.Chen, N. N.; Chang, C.–F.; Gallia, G. L.; Kerr, D. A.; Johnson,E. M.; Krachmarov, C. P.; Barr, S. M.; Frisque, R. J.; Bollag, B.;Khalili, K.: Cooperative action of cellular proteins YB–1 and Pur–alphawith the tumor antigen of the human JC polyomavirus determines theirinteraction with the viral lytic control element. Proc. Nat. Acad.Sci. 92: 1087–1091, 1995.
- [73677] 12007.Gallia, G. L.; Johnson, E. M.; Khalili, K.: Pur–alpha: a multifunctionalsingle–stranded DNA– and RNA–binding protein. Nucleic Acids Res. 28:3197–3205, 2000.
- [73678] 12008.Kennedy, G. C.; Rutter, W. J.: Pur–1, a zinc–finger protein thatbinds to purine–rich sequences, transactivates an insulin promoterin heterologous cells. Proc. Nat. Acad. Sci. 89: 11498–11502, 1992.
- [73679] 12009.Ma, Z.–W.; Pejovic, T.; Najfeld, V.; Ward, D. C.; Johnson, E. M.: Localization of PURA, the gene encoding the sequence–specific single–stranded–DNA–bindingprotein Pur–alpha, to chromosome band 5q31. Cytogenet. Cell Genet. 71:64–67, 1995.
- [73680] 12010.Agerberth, B.; Gunne, H.; Odeberg, J.; Kogner, P.;

Boman, H. G.;Gudmundsson, G. H.: FALL-39, a putative human peptide antibiotic,is cysteine-free and expressed in bone marrow and testis. Proc. Nat.Acad. Sci. 92: 195-199, 1995.

[73681] 12011.Frohm, M.; Agerberth, B.; Ahangari, G.; Stahle-Backdahl, M.; Liden,S.; Wigzell, H.; Gudmundsson, G. H.: The expression of the gene codingfor the antibacterial peptide LL-37 is induced in human keratinocytesduring inflammatory disorders. J. Biol. Chem. 272: 15258-15263,1997.

[73682] 12012.Gudmundsson, G. H.; Agerberth, B.; Odeberg, J.; Bergman, T.; Olsson,B.; Salcedo, R.: The human gene FALL39 and processing of the cathelinprecursor to the an-tibacterial peptide LL-37 in granulocytes. Europ.J. Biochem. 238: 325-332, 1996.

[73683] 12013.Kirikae, T.; Hirata, M.; Yamasu, H.; Kirikae, F.; Tamura, H.; Kayama,F.; Nakatsuka, K.; Yokochi, T.; Nakano, M.: Protective effects ofa human 18-kilodalton cationic antimicrobial protein (CAP18)-derivedpeptide against murine endotoxemia. Infect. Immun. 66: 1861-1868,1998.

[73684] 12014.Matsuzono, Y.; Kinoshita, N.; Tamura, S.; Shi-mozawa, N.; Hamasaki,M.; Ghaedi, K.; Wanders, R. J. A.;

Suzuki, Y.; Kondo, N.; Fujiki, Y.: Human PEX19: cDNA cloning by functional complementation, mutation analysis in a patient with Zellweger syndrome, and potential role in peroxisomal membrane assembly. *Proc. Nat. Acad. Sci.* 96: 2116–2121, 1999.

[73685] 12015. Kinoshita, N.; Ghaedi, K.; Shimozawa, N.; Wanders, R. J. A.; Matsuzono, Y.; Imanaka, T.; Okumoto, K.; Suzuki, Y.; Kondo, N.; Fujiki, Y.: Newly identified Chinese hamster ovary cell mutants are defective in biogenesis of peroxisomal membrane vesicles (peroxisomal ghosts), representing a novel complementation group in mammals *J. Biol. Chem.* 273:24122–24130, 1998.

[73686] 12016. Chen, K.-S.; Gunaratne, P. H.; Hoheisel, J. D.; Young, I. G.; Gabor Miklos, G. L.; Greenberg, F.; Shaffer, L. G.; Campbell, H. D.; Lupski, J. R.: The human homologue of the *Drosophila melanogaster* flightless-I gene (*flil*) maps within the Smith–Magenis microdeletion critical region in 17p11.2. *Am. J. Hum. Genet.* 56: 175–182, 1995.

[73687] 12017. Heisterkamp, N.; Kaartinen, V.; van Soest, S.; Bokoch, G. M.; Groffen, J.: Human ABR encodes a protein with GAP–rac activity and homology to the DBL nucleotide exchange factor domain. *J. Biol. Chem.* 268:16903–16906, 1993.



- [73688] 12018.Heisterkamp, N.; Morris, C.; Groffen, J.: ABR, an active BCR-related gene. *Nucleic Acids Res.* 17: 8821–8831, 1989.
- [73689] 12019.McDonald, J. D.; Daneshvar, L.; Willert, J. R.; Matsuura, K.; Waldman, F.; Cogen, P. H.: Physical mapping of chromosome 17p13.3 in the region of a putative tumor suppressor gene important in medulloblastoma. *Genomics* 23:229–232, 1994.
- [73690] 12020.Shimomura, H.; Sanke, T.; Hanabusa, T.; Tsunoda, K.; Furuta, H.; Nanjo, K.: Nonsense mutation of islet-1 gene (Q310X) found in a type 2 diabetic patient with a strong family history. *Diabetes* 49: 1597–1600, 2000.
- [73691] 12021.Tanizawa, Y.; Riggs, A. C.; Dagogo-Jack, S.; Vaxillaire, M.; Froguel, P.; Liu, L.; Donis-Keller, H.; Permutt, M. A.: Isolation of the human LIM/homeodomain gene islet-1 and identification of a simple sequence repeat 1. *Diabetes* 43: 935–941, 1994.
- [73692] 12022.Kleiman, F. E.; Manley, J. L.: Functional interaction of BRCA1-associated BARD1 with polyadenylation factor CstF-50. *Science* 285: 1576–1579, 1999.
- [73693] 12023.Ishibashi, T.; Bottaro, D. P.; Chan, A.; Miki, T.; Aaronson, S.A.: Expression cloning of a human dual-specificity phosphatase. *Proc. Nat. Acad. Sci.* 89:

12170–12174, 1992.

[73694] 12024.Kamb, A.; Futreal, P. A.; Rosenthal, J.; Cochran, C.; Harshman,K. D.; Liu, Q.; Phelps, R. S.; Tavitigian, S. V.; Tran, T.; Hussey,C.; Bell, R.; Miki, Y.; Swensen, J.; Hobbs, M. R.; Marks, J.; Bennett,L. M.; Barrett, J. C.; Wiseman, R. W.; Shattuck–Eidens, D.: Localization of the VHR phosphatase gene and its analysis as a candidate for BRCA1. Genomics 23:163–167, 1994.

[73695] 12025.Corti, O.; Finocchiaro, G.; Rossi, E.; Zuffardi, O.; DiDonato,S.: Molecular cloning of cDNAs encoding human carnitine acetyltransferase and mapping of the corresponding gene to chromosome 9q34.1. Genomics 23:94–99, 1994.

[73696] 12026.Kalaria, R. N.; Harik, S. I.: Carnitine acetyltransferase activity in the human brain and its microvessels is decreased in Alzheimer's disease. Ann. Neurol. 32: 583–586, 1992.

[73697] 12027.van der Leij, F. R.; Huijkman, N. C. A.; Boomsma, C.; Kuipers,J. R. G.; Bartelds, B.: Genomics of the human carnitine acyltransferase genes. Molec. Genet. Metab. 71: 139–153, 2000.

[73698] 12028.Couch, F. J.; Rommens, J. M.; Neuhausen, S. L.; Belanger, C.; Dumont,M.; Abel, K.; Bell, R.; Berry, S.; Bogden,

R.; Cannon-Albright, L.; Farid, L.; Frye, C.; and 30 others: Generation of an integrated transcription map of the BRCA2 region on chromosome 13q12-q13. *Genomics* 36:86-99, 1996.

[73699] 12029. Nelms, K.; Snow, A. J.; Noben-Trauth, K.: Dok1 encoding p62(dok) maps to mouse chromosome 6 and human chromosome 2 in a region of translocation in chronic lymphocytic leukemia. *Genomics* 53: 243-245, 1998.

[73700] 12030. Wisniewski, D.; Strife, A.; Wojchiewicz, D.; Lambek, C.; Clarkson, B.: A 62-kilodalton tyrosine phosphoprotein constitutively present in primary chronic phase chronic myelogenous leukemia enriched lineage negative blast populations. *Leukemia* 8: 688-693, 1994.

[73701] 12031. Gengyo-Ando, K.; Kitayama, H.; Mukaida, M.; Ikawa, Y.: A murine neural-specific homolog corrects cholinergic defects in *Caenorhabditis elegans* unc-18 mutants. *J. Neurosci.* 16: 6695-6702, 1996.

[73702] 12032. Pevsner, J.; Hsu, S.-C.; Scheller, R. H.: n-Sec1: a neural-specific syntaxin-binding protein. *Proc. Nat. Acad. Sci.* 91: 1445-1449, 1994.

[73703] 12033. Swanson, D. A.; Steel, J. M.; Valle, D.: Identification and characterization of the human ortholog of rat STXBP1, a protein implicated in vesicle trafficking and neurotrans-

mitter release. *Genomics* 48: 373–376,1998.

[73704] 12034.Verhage, M.; Mala, A. S.; Plomp, J. J.; Brussaard, A. B.; Heeroma,J. H.; Vermeer, H.; Toonen, R. F.; Hammer, R. E.; van den Berg, T.K.; Missler, M.; Geuze, H. J.; Sudhof, T. C.: Synaptic assembly ofthe brain in the absence of neurotransmitter secretion. *Science* 287:864–869, 2000.

[73705] 12035.Cox, P. R.; Zoghbi, H. Y.: Sequencing, expression analysis, andmapping of three unique human tropomodulin genes and their mouse orthologs. *Genomics* 63:97–107, 2000.

[73706] 12036.Watakabe, A.; Kobayashi, R.; Helfman, D. M.: N-tropomodulin: anovel isoform of tropomodulin identified as the major binding proteinto brain tropomyosin. *J. Cell Sci.* 109: 2299–2310, 1996.

[73707] 12037.Galvin, K. M.; Donovan, M. J.; Lynch, C. A.; Meyer, R. I.; Paul,R. J.; Lorenz, J. N.; Fairchild–Huntress, V.; Dixon, K. L.; Dunmore,J. H.; Gimbrone, M. A., Jr.; Falb, D.; Huszar, D.: A role for Smad6in development and homeostasis of the cardiovascular system. *NatureGenet.* 24: 171–174, 2000.

[73708] 12038.Topper, J. N.; Cai, J.; Qiu, Y.; Anderson, K. R.; Xu, Y.–Y.; Deeds,J. D.; Feeley, R.; Gimeno, C. J.; Woolf, E. A.; Tayber, O.; Mays,G. G.; Sampson, B. A.; Schoen, F. J.; Gim–

brone, M. A., Jr.; Falb, D.: Vascular MADs: two novel MAD-related genes selectively inducible by flow in human vascular endothelium. *Proc. Nat. Acad. Sci.* 94:9314–9319, 1997.

[73709] 12039. Festing, M. F. W. (ed.): Inbred strains in biomedical research. New York: Oxford University Press, 1979. p. 255.

[73710] 12040. Illa, I.; Serrano-Munuera, C.; Gallardo, E.; Lasa, A.; Rojas-Garcia, R.; Palmer, J.; Gallano, P.; Baiget, M.; Matsuda, C.; Brown, R. H.: Distal anterior compartment myopathy: a dysferlin mutation causing a new muscular dystrophy phenotype. *Ann. Neurol.* 49: 130–134, 2001.

[73711] 12041. Illarioshkin, S. N.; Ivanova-Smolenskaya, I. A.; Greenberg, C. R.; Nylen, E.; Sukhorukov, V. S.; Poleshchuk, V. V.; Markova, E. D.; Wrogemann, K.: Identical dysferlin mutation in limb-girdle muscular dystrophy type 2B and distal myopathy. *Neurology* 55: 1931–1933, 2000.

[73712] 12042. Liu, J.; Wu, C.; Bossie, K.; Bejaoui, K.; Hosler, B. A.; Gingrich, J. C.; Ben Hamida, M.; Hentati, F.; Schurr, E.; de Jong, P. J.; Brown, R. H., Jr.: Generation of 3-Mb PAC contig spanning the Miyoshi myopathy/limb-girdle muscular dystrophy (MM/LGMD2B) locus on chromosome 2p13. *Genomics* 49:23–29, 1998.

- [73713] 12043.Mahjneh, I.; Vannelli, G.; Bushby, K.; Marconi, G. P.: A largeinbred Palestinian family with two forms of muscular dystrophy. Neuromusc.Disord. 2: 277–283, 1992.
- [73714] 12044.Weiler, T.; Bashir, R.; Anderson, L. V. B.; Davison, K.; Moss,J. A.; Britton, S.; Nylen, E.; Keers, S.; Vafiadaki, E.; Greenberg,C. R.; Bushby, K. M. D.; Wrogemann, K.: Identical mutation in patientswith limb girdle muscular dystrophy type 2B or Miyoshi myopathy suggestsa role for modifier gene(s). Hum. Molec. Genet. 8: 871–877, 1999.
- [73715] 12045.Pennica, D.; Arce, V.; Swanson, T. A.; Vejsada, R.; Pollock, R.A.; Armanini, M.; Dudley, K.; Phillips, H. S.; Rosenthal, A.; Kato,A. C.; Henderson, C. E.: Cardiotrophin-1, a cytokine present in embryonicmuscle, supports long-term survival of spinal motoneurons. Neuron 17:63–74, 1996.
- [73716] 12046.Pennica, D.; King, K. L.; Shaw, K. J.; Luis, E.; Rullamas, J.;Luoh, S.–M.; Darbonne, W. C.; Knutzon, D. S.; Yen, R.; Chien, K. R.;Baker, J. B.; Wood, W. I.: Expression cloning of cardiotrophin 1,a cytokine that induces cardiac myocyte hypertrophy. Proc. Nat. Acad.Sci. 92: 1142–1146, 1995.
- [73717] 12047.Ghosh, A.: Learning more about NMDA receptor regulation. Science 295:449–451, 2002.

- [73718] 12048.Grunwald, I. C.; Korte, M.; Wolfer, D.; Wilkinson, G. A.; Unsicker,K.; Lipp, H.-P.; Bonhoeffer, T.; Klein, R.: Kinase-independent requirement of EphB2 receptors in hippocampal synaptic plasticity. *Neuron* 32:1027–1040, 2001.
- [73719] 12049.Henderson, J. T.; Georgiou, J.; Jia, Z.; Robertson, J.; Elowe,S.; Roder, J. C.; Pawson, T.: The receptor tyrosine kinase EphB2 regulates NMDA-dependent synaptic function. *Neuron* 32: 1041–1056,2001.
- [73720] 12050.Himanen, J.-P.; Rajashankar, K. R.; Lackmann, M.; Cowan, C. A.;Henkemeyer, M.; Nikolov, D. B.: Crystal structure of an Eph receptor–ephrin complex. *Nature* 414: 933–938, 2001.
- [73721] 12051.Takasu, M. A.; Dalva, M. B.; Zigmond, R. E.; Greenberg, M. E.: Modulation of NMDA receptor-dependent calcium influx and gene expression through EphB receptors. *Science* 295: 491–495, 2002.
- [73722] 12052.Doudney, K.; Murdoch, J. N.; Paternotte, C.; Bentley, L.; Gregory,S.; Copp, A. J.; Stanier, P.: Comparative physical and transcript maps of approximately 1 Mb around loop–tail, a gene for severe neuraltube defects on distal mouse chromosome 1 and human chromosome 1q22–q23. *Genomics* 72:180–192, 2001.

- [73723] 12053.Kibar, Z.; Vogan, K. J.; Groulx, N.; Justice, M. J.; Underhill, D. A.; Gros, P.: Ltap, a mammalian homolog of *Drosophila Strabismus/VanGogh*, is altered in the mouse neural tube mutant loop-tail. *Nature Genet.* 28: 251–255, 2001.
- [73724] 12054.Mullick, A.; Trasler, D.; Gros, P.: High-resolution linkage map in the vicinity of the Lp locus. *Genomics* 26: 479–488, 1995.
- [73725] 12055.Murdoch, J. N.; Doudney, K.; Paternotte, C.; Copp, A. J.; Stanier, P.: Severe neural tube defects in the loop-tail mouse result from mutation of Lpp1, a novel gene involved in floor plate specification. *Hum. Molec. Genet.* 10: 2593–2601, 2001.
- [73726] 12056.Stanier, P.; Henson, J. N.; Eddleston, J.; Moore, G. E.; Copp, A. J.: Genetic basis of neural tube defects: the mouse gene loop-tail maps to a region of chromosome 1 syntenic with human 1q21–q23. *Genomics* 26:473–478, 1995.
- [73727] 12057.Yu, J. X.; Chao, L.; Chao, J.: Prostasin is a novel human serine proteinase from seminal fluid: purification, tissue distribution, and localization in prostate gland. *J. Biol. Chem.* 269: 18843–18848, 1994.
- [73728] 12058.Yu, J. X.; Chao, L.; Chao, J.: Molecular cloning, tis-



sue-specific expression, and cellular localization of human prostatic mRNA. J. Biol. Chem. 270: 13483–13489, 1995.

- [73729] 12059. Yu, J. X.; Chao, L.; Ward, D. C.; Chao, J.: Structure and chromosomal localization of the human prostatic (PRSS8) gene. Genomics 32: 334–340, 1996.
- [73730] 12060. Arber, S.; Halder, G.; Caroni, P.: Muscle LIM protein, a novel essential regulator of myogenesis, promotes myogenic differentiation. Cell 79: 221–231, 1994.
- [73731] 12061. Fung, Y. W.; Wang, R. X.; Heng, H. H. Q.; Liew, C. C.: Mapping of a human LIM protein (CLP) to human chromosome 11p15.1 by fluorescence in situ hybridization. Genomics 28: 602–603, 1995.
- [73732] 12062. Becker-Andre, M.; Andre, E.; DeLamarter, J. F.: Identification of nuclear receptor mRNAs by RT-PCR amplification of conserved zinc-finger motif sequences. Biochem. Biophys. Res. Commun. 194: 1371–1379, 1993.
- [73733] 12063. Carlberg, C.; Hooft van Huijsduijnen, R.; Staple, J. K.; DeLamarter, J. F.; Becker-Andre, M.: RZR's, a new family of retinoid-related orphan receptors that function as both monomers and homodimers. Molec. Endocr. 8: 757–770, 1994.
- [73734] 12064. Giguere, V.; Beatty, B.; Squire, J.; Copeland, N. G.; Jenkins, N. A.: The orphan nuclear receptor ROR- $\alpha$

(RORA) maps to a conserved region of homology on human chromosome 15q21–q22 and mouse chromosome 9. *Genomics* 28: 596–598, 1995.

- [73735] 12065. Giguere, V.; Tini, M.; Flock, G.; Ong, E.; Evans, R. M.; Otulakowski, G.: Isoform-specific amino-terminal domains dictate DNA-binding properties of ROR- $\alpha$ , a novel family of orphan hormone nuclear receptors. *Genes Dev.* 8: 538–553, 1994.
- [73736] 12066. Hamilton, B. A.; Frankel, W. N.; Kerrebrock, A. W.; Hawkins, T. L.; FitzHugh, W.; Kusumi, K.; Russell, L. B.; Mueller, K. L.; vanBerkel, V.; Birren, B. W.; Kruglyak, L.; Lander, E. S.: Disruption of the nuclear hormone receptor ROR- $\alpha$  in staggerer mice. *Nature* 379: 736–739, 1996.
- [73737] 12067. Matysiak-Scholze, U.; Nehls, M.: The structural integrity of ROR- $\alpha$  isoforms is mutated in staggerer mice: cerebellar coexpression of ROR- $\alpha$ -1 and ROR- $\alpha$ -4. *Genomics* 43: 78–84, 1997.
- [73738] 12068. Meyer, T.; Kneissel, M.; Mariani, J.; Fournier, B.: In vitro and in vivo evidence for orphan nuclear receptor ROR- $\alpha$  function in bone metabolism. *Proc. Nat. Acad. Sci.* 97: 9197–9202, 2000.
- [73739] 12069. Ueda, H. R.; Chen, W.; Adachi, A.; Wakamatsu, H.; Hayashi, S.; Takasugi, T.; Nagano, M.; Nakahama, K.;

Suzuki, Y.; Sugano, S.; Iino, M.; Shigeyoshi, Y.; Hashimoto, S.: A transcription factor response element for gene expression during circadian night. *Nature* 418:534–539, 2002.

[73740] 12070. Zanjani, H. S.; Herrup, K.; Guastavino, J.-M.; Delhay-Bouchaud, N.; Mariani, J.: Developmental studies of the inferior olivary nucleus in staggerer mutant mice. *Develop. Brain Res.* 82: 18–28, 1994.

[73741] 12071. Masternak, K.; Barras, E.; Zufferey, M.; Conrad, B.; Corthals, G.; Aebersold, R.; Sanchez, J.-C.; Hochstrasser, D. F.; Mach, B.; Reith, W.: A gene encoding a novel RFX-associated transactivator is mutated in the majority of MHC class II deficiency patients. *Nature Genet.* 20: 273–277, 1998.

[73742] 12072. Prange, C. K.; Pennacchio, L. A.; Lieuallen, K.; Fan, W.; Lennon, G. G.: Characterization of the human neurocan gene, CSPG3. *Gene* 221:199–205, 1998.

[73743] 12073. Rauch, U.; Grimpe, B.; Kulbe, G.; Arnold-Ammer, I.; Beier, D. R.; Fassler, R.: Structure and chromosomal localization of the mouse neurocan gene. *Genomics* 28: 405–410, 1995.

[73744] 12074. Rauch, U.; Karthikeyan, L.; Maurel, P.; Margolis, R. U.; Margolis, R. K.: Cloning and primary structure of neu-

rocan, a developmentally regulated, aggregating chondroitin sulfate proteoglycan of brain. *J. Biol. Chem.* 267: 19536–19547, 1992.

[73745] 12075. Clement, S.; Krause, U.; Desmedt, F.; Tanti, J.-F.; Behrends, J.; Pesesse, X.; Sasaki, T.; Penninger, J.; Doherty, M.; Malaisse, W.; Dumont, J. E.; Le Marchand-Brustel, Y.; Erneux, C.; Hue, L.; Schurmans, S.: The lipid phosphatase SHIP2 controls insulin sensitivity. *Nature* 409:92–97, 2001.

[73746] 12076. Habib, T.; Hejna, J. A.; Moses, R. E.; Decker, S. J.: Growth factors and insulin stimulate tyrosine phosphorylation of the 51C/SHIP2 protein. *J. Biol. Chem.* 273: 18605–18609, 1998.

[73747] 12077. Hejna, J. A.; Saito, H.; Merkens, L. S.; Tittle, T. V.; Jakobs, P. M.; Whitney, M. A.; Grompe, M.; Friedberg, A. S.; Moses, R. E.: Cloning and characterization of a human cDNA (INPPL1) sharing homology with inositol polyphosphate phosphatases. *Genomics* 29: 285–287, 1995.

[73748] 12078. Pesesse, X.; Deleu, S.; De Smedt, F.; Drayer, L.; Erneux, C.: Identification of a second SH2-domain-containing protein closely related to the phosphatidylinositol polyphosphate 5-phosphatase SHIP. *Biochem. Biophys. Res. Commun.* 239: 697–700, 1997.

- [73749] 12079.Schurmans, S.; Carrio, R.; Behrends, J.; Pouillon, V.; Merino,J.; Clement, S.: The mouse SHIP2 (Inpp1) gene: complementary DNA,genomic structure, promoter analysis, and gene expression in the embryoand adult mouse. *Genomics* 62: 260–271, 1999.
- [73750] 12080.Drummond, J. T.; Genschel, J.; Wolf, E.; Modrich, P.: DHFR/MSH3amplification in methotrexate–resistant cells alters the hMutS–alpha/hMutS–betaratio and reduces the efficiency of base–base mismatch repair. *Proc.Nat. Acad. Sci.* 94: 10144–10149, 1997.
- [73751] 12081.Fujii, H.; Shimada, T.: Isolation and characteriza–tion of cDNAclones derived from the divergently tran–scribed gene in the regionupstream from the human dihy–drofolate reductase gene. *J. Biol. Chem.* 264:10057–10064, 1989.
- [73752] 12082.Inokuchi, K.; Ikejima, M.; Watanabe, A.; Nakajima, E.; Orimo, H.;Nomura, T.; Shimada, T.: Loss of expression of the human MSH3 genein hematological malignancies. *Biochem. Biophys. Res. Commun.* 214:171–179, 1995.
- [73753] 12083.Linton, J. P.; Yen, J. Y.–J.; Selby, E.; Chen, Z.; Chin–sky, J.M.; Liu, K.; Kellems, R. E.; Crouse, G. F.: Dual bidi–rectional promotersat the mouse dhfr locus: cloning and characterization of two mRNAclasses of the divergently

transcribed Rep-1 gene. *Molec. Cell. Biol.* 9:3058–3072, 1989.

[73754] 12084.Marra, G.; Iaccarino, I.; Lettieri, T.; Roscilli, G.; Delmastro, P.; Jiricny, J.: Mismatch repair deficiency associated with overexpression of the MSH3 gene. *Proc. Nat. Acad. Sci.* 95: 8568–8573, 1998.

[73755] 12085.Nakajima, E.; Orimo, H.; Ikejima, M.; Shimada, T.: Nine-bp repeat polymorphism in exon 1 of the hMSH3 gene. *Jpn. J. Hum. Genet.* 40:343–345, 1995.

[73756] 12086.Risinger, J. I.; Umar, A.; Boyd, J.; Berchuck, A.; Kunkel, T. A.; Barrett, J. C.: Mutation of MSH3 in endometrial cancer and evidence for its functional role in heteroduplex repair. *Nature Genet.* 14:102–109, 1996.

[73757] 12087.Smith, M. L.; Mitchell, P. J.; Crouse, G. F.: Analysis of the mouse Dhfr/Rep-3 major promoter region by using linker-scanning and internal deletion mutations and DNase I footprinting. *Molec. Cell. Biol.* 10: 6003–6012, 1990.

[73758] 12088.Watanabe, A.; Ikejima, M.; Suzuki, N.; Shimada, T.: Genomic organization and expression of the human MSH3 gene. *Genomics* 31: 311–318, 1996.

[73759] 12089.Yin, J.; Kong, D.; Wang, S.; Zou, T.–T. Souza, R. F.; Smolinski, K. N.; Lynch, P. M.; Hamilton, S. R.; Sugimura, H.; Powell, S. M.; Young, J.; Abraham, J. M.; Meltzer, S. J.:

Mutation of hMSH3 and hMSH6 mismatch repair genes in genetically unstable human colorectal and gastric carcinomas. *Hum. Mutat.* 10: 474–478, 1997.

[73760] 12090. Chan, A. M.-L.; McGovern, E. S.; Catalano, G.; Fleming, T. P.; Miki, T.: Expression cDNA cloning of a novel oncogene with sequence similarity to regulators of small GTP-binding proteins. *Oncogene* 9: 1057–1063, 1994.

[73761] 12091. Miki, T.; Fleming, T. P.; Crescenzi, M.; Molloy, C. J.; Blam, S. B.; Reynolds, S. H.; Aaronson, S. A.: Development of a highly efficient expression cDNA cloning system: application to oncogene isolation. *Proc. Nat. Acad. Sci.* 88: 5167–5171, 1991.

[73762] 12092. Takai, S.; Chan, A. M.-L.; Yamada, K.; Miki, T.: Assignment of the human TIM proto-oncogene to 7q33–q35. *Cancer Genet. Cytogenet.* 83:87–89, 1995.

[73763] 12093. Burger, B.; Uhlhaas, S.; Mangold, E.; Propping, P.; Friedl, W.; Jenne, D.; Dockter, G.; Back, W.: Novel de novo mutation of MADH4/SMAD4 in a patient with juvenile polyposis. (Letter) *Am. J. Med. Genet.* 110:289–291, 2002.

[73764] 12094. Kurima, K.; Peters, L. M.; Yang, Y.; Riazuddin, S.; Ahmed, Z. M.; Naz, S.; Arnaud, D.; Drury, S.; Mo, J.; Makishima, T.; Ghosh, M.; Menon, P. S. N.; and 13 others: Dominant and recessive deafness caused by mutations of a

novel gene, TMC1, required for cochlear hair-cell function. Nature Genet. 30: 277–284, 2002.

[73765] 12095. Bryant, P. J.; Huettner, B.; Held, L. I., Jr.; Ryerse, J.; Szidonya, J.: Mutations at the fat locus interfere with cell proliferation control and epithelial morphogenesis in Drosophila. Dev. Biol. 129:541–554, 1988.

[73766] 12096. Dunne, J.; Hanby, A. M.; Poulson, R.; Jones, T. A.; Sheer, D.; Chin, W. G.; Da, S. M.; Zhao, Q.; Beverley, P. C. L.; Owen, M. J.: Molecular cloning and tissue expression of FAT, the human homologue of the Drosophila fat gene that is located on chromosome 4q34–q35 and encodes a putative adhesion molecule. Genomics 30: 207–223, 1995.

[73767] 12097. Hortsch, M.; Goodman, C. S.: Cell and substrate adhesion molecules in Drosophila. Ann. Rev. Cell. Biol. 7: 505–557, 1991.

[73768] 12098. Mahoney, P. A.; Weber, U.; Onofrechuk, P.; Biessmann, H.; Bryant, P. J.; Goodman, C. S.: The fat tumor suppressor gene in Drosophila encodes a novel member of the cadherin gene superfamily. Cell 67:853–868, 1991.

[73769] 12099. Crowe, P. D.; VanArsdale, T. L.; Walter, B. N.; Ware, C. F.; Hession, C.; Ehrenfels, B.; Browning, J. L.; Din, W. S.; Goodwin, R. G.; Smith, C. A.: A lymphotoxin-beta-specific receptor. Science 264: 707–710, 1994.



- [73770] 12100.Nakamura, T.; Tashiro, K.; Nazarea, M.; Nakano, T.; Sasayama, S.;Honjo, T.: The murine lymphotoxin-beta receptor cDNA: isolation bythe signal sequence trap and chromosomal mapping. *Genomics* 30: 312-319,1995.
- [73771] 12101.Tashiro, K.; Tada, H.; Heilker, R.; Shirozu, M.; Nakano, T.; Honjo,T.: Signal sequence trap: a cloning strategy for secreted proteinsand type I membrane proteins. *Science* 261: 600-603, 1993.
- [73772] 12102.George, A.; Sabsay, B.; Simonian, P. A.; Veis, A.: Characterizationof a novel dentin matrix acidic phosphoprotein: implications for inductionof biomineralization . *J. Biol. Chem.* 268: 12624-12630, 1993.
- [73773] 12103.Hirst, K. L.; Simmons, D.; Feng, J.; Aplin, H.; Dixon, M. J.; MacDougall,M.: Elucidation of the sequence and the genomic organization of thehuman dentin matrix acidic phosphoprotein 1 (DMP1) gene: exclusionof the locus from a causative role in the pathogenesis of dentinogenesisimperfecta type II. *Genomics* 42: 38-45, 1997.
- [73774] 12104.MacDougall, M.; DuPont, B. R.; Simmons, D.; Leach, R. J.: Assignmentof DMP1 to human chromosome 4 band q21 by in situ hybridization. *Cytogenet.Cell. Genet.* 74: 189 only, 1996.
- [73775] 12105.Lu, Z.; Xu, S.; Joazeiro, C.; Cobb, M. H.; Hunter, T.:

The PHDdomain of MEKK1 acts as an E3 ubiquitin ligase and mediates ubiquitinationand degradation of ERK1/2. Molec. Cell 9: 945–956, 2002.

- [73776] 12106.Bohm, H.; Benndorf, R.; Gaestel, M.; Gross, B.; Nurnberg, P.; Kraft,R.; Otto, A.; Bielka, H.: The growth–related protein P23 of the Ehrlichascites tumor: translational control, cloning and primary structure. Biochem.Int. 19: 277–286, 1989.
- [73777] 12107.Barnard, R. C.; Pascall, J. C.; Brown, K. D.; McKay, I. A.; Williams,N. S.; Bustin, S. A.: Coding sequence of ERF–1, the human homologueof Tis11b/cMG1, members of the Tis11 family of early response genes. NucleicAcids Res. 21: 3580 only, 1993.
- [73778] 12108.Bustin, S. A.; Xiao–Feng, N.; Barnard, R. C.; Kumar, V.; Pascall,J. C.; Brown, K. D.; Leigh, I. M.; Williams, N. S.; McKay, I. A.:Cloning and characterisation of ERF1, a human member of the Tis11family of early–response genes. DNA Cell Biol. 13: 449–459, 1994.
- [73779] 12109.Maclean, K. N.; See, C. G.; McKay, I. A.; Bustin, S. A.: The humanimmediate early gene BRF1 maps to chromosome 14q22–q24. Genomics 30:89–90, 1995.
- [73780] 12110.Ning, Z.–Q.; Norton, J. D.; Li, J.; Murphy, J. J.: Distinct mechanismsfor rescue from apoptosis in Ramos hu–

man B cells by signaling through CD40 and interleukin-4 receptor: a role for inhibition of an early response gene, Berg 36. *Europ. J. Immun.* 26: 2356–2363, 1996.

[73781] 12111. Chihade, J. W.; Brown, J. R.; Schimmel, P. R.; Ribas de Pouplana, L.: Origin of mitochondria in relation to evolutionary history of eukaryotic alanyl-tRNA synthetase.

*Proc. Nat. Acad. Sci.* 97: 12153–12157, 2000.

[73782] 12112. Maas, S.; Kim, Y.-G.; Rich, A.: Genomic clustering of tRNA-specific adenosine deaminase ADAT1 and two tRNA synthetases. *Mammalian Genome* 12:387–393, 2001.

[73783] 12113. Derynck, R.; Gelbart, W. M.; Harland, R. M.; Heldin, C.-H.; Kern, S. E.; Massague, J.; Melton, D. A.; Mlodzik, M.; Padgett, R. W.; Roberts, A. B.; Smith, J.; Thomsen, G. H.; Vogelstein, B.; Wang, X.-F.: Nomenclature: vertebrate mediators of TGF-beta family signals. (Letter) *Cell* 87:173 only, 1996.

[73784] 12114. Friedl, W.; Kruse, R.; Uhlhaas, S.; Stolte, M.; Scharmann, B.; Keller, K. M.; Jungck, M.; Stern, M.; Loff, S.; Back, W.; Propping, P.; Jenne, D. E.: Frequent 4-bp deletion in exon 9 of the SMAD4/MADH4 gene in familial juvenile polyposis patients. *Genes Chromosomes Cancer* 25:403–406, 1999.

- [73785] 12115.Hahn, S. A.; Schutte, M.; Hoque, T. M. S.; Moskaluk, C. A.; daCosta, L. T.; Rozenblum, E.; Weinstein, C. L.; Fischer, A.; Yeo, C.J.; Hruban, R. H.; Kern, S. E.: DPC4, a candidate tumor suppressorgene at human chromosome 18q21.1. *Science* 271: 350–354, 1996.
- [73786] 12116.Howe, J. R.; Shellnut, J.; Wagner, B.; Ringold, J. C.; Sayed, M. G.; Ahmed, A. F.; Lynch, P. M.; Amos, C. I.; Sistonen, P.; Aaltonen, L. A.: Common deletion of SMAD4 in juvenile polyposis is a mutational hotspot. *Am. J. Hum. Genet.* 70: 1357–1362, 2002.
- [73787] 12117.Inman, G. J.; Nicolas, F. J.; Hill, C. S.: Nucleocytoplasmic shuttling of Smads 2, 3, and 4 permits sensing of TGF-beta receptor activity. *Molec. Cell* 10: 283–294, 2002.
- [73788] 12118.Kim, S. K.; Fan, Y.; Papadimitrakopoulou, V.; Clayman, G.; Hittelman, W. N.; Hong, W. K.; Lotan, R.; Mao, L.: DPC4, a candidate tumor suppressorgene, is altered infrequently in head and neck squamous cell carcinoma. *Cancer Res.* 56: 2519–2521, 1996.
- [73789] 12119.Kinzler, K. W.; Vogelstein, B.: Landscaping the cancer terrain. *Science* 280:1036–1037, 1998.
- [73790] 12120.MacGrogan, D.; et al.; et al: Comparative mutational analysis of DPC4 (Smad4) in prostatic and colorectal carcinomas. *Oncogene* 15:1111–1114, 1997.

- [73791] 12121.Roth, S.; Johansson, M.; Loukola, A.; Peltomaki, P.; Jarvinen,H.; Mecklin, J.-P.; Aaltonen, L. A.: Mutation analysis of SMAD2,SMAD3, and SMAD4 genes in hereditary non-polyposis colorectal cancer. J.Med. Genet. 37: 298-300, 2000.
- [73792] 12122.Schutte, M.; Hruban, R. H.; Hedrick, L.; Cho, K. R.; Nadasdy,G. M.; Weinstein, C. L.; Bova, G. S.; Isaacs, W. B.; Cairns, P.; Nawroz,H.; Sidransky, D.; Casero, R. A., Jr.; Meltzer, P. S.; Hahn, S. A.;Kern, S. E.: DPC4 gene in various tumor types. Cancer Res. 56:2527-2530, 1996.
- [73793] 12123.Shioda, T.; Lechleider, R. J.; Dunwoodie, S. L.; Li, H.; Yahata,T.; de Caestecker, M. P.; Fenner, M. H.; Roberts, A. B.; Isselbacher,K. J.: Transcriptional activating activity of Smad4: roles of SMADhetero-oligomerization and enhancement by an associating transactivator. Proc.Nat. Acad. Sci. 95: 9785-9790, 1998.
- [73794] 12124.Sirard, C.; de la Pompa, J. L.; Elia, A.; Itie, A.; Mirtos, C.;Cheung, A.; Hahn, S.; Wakeham, A.; Schwartz, L.; Kern, S. E.; Rossant,J.; Mak, T. W.: The tumor suppressor gene Smad4/Dpc4 is requiredfor gastrulation and later for anterior development of the mouse embryo. GenesDev. 12: 107-119, 1998.
- [73795] 12125.Tagaki, Y.; et al.; et al: Somatic alterations of the

DPC4 gene in human colorectal cancers in vivo. *Gastroenterology* 111: 1369–1372, 1996.

[73796] 12126. Takaku, K.; Oshima, M.; Miyoshi, H.; Matsui, M.; Seldin, M. F.; Taketo, M. M.: Intestinal tumorigenesis in compound mutant mice of both Dpc4 (Smad4) and Apc genes. *Cell* 92: 645–656, 1998.

[73797] 12127. Thiagalingam, S.; Lengauer, C.; Leach, F. S.; Schutte, M.; Hahn, S. A.; Overhauser, J.; Willson, J. K. V.; Markowitz, S.; Hamilton, S. R.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Evaluation of candidate tumour suppressor genes on chromosome 18 in colorectal cancers. *Nature-Genet.* 13: 343–346, 1996.

[73798] 12128. Zawel, L.; Dai, J. L.; Buckhaults, P.; Zhou, S.; Kinzler, K. W.; Vogelstein, B.; Kern, S. E.: Human Smad3 and Smad4 are sequence-specific transcription activators. *Molec. Cell* 1: 611–617, 1998.

[73799] 12129. Zhou, S.; Buckhaults, P.; Zawel, L.; Bunz, F.; Riggs, G.; LeDai, J.; Kern, S. E.; Kinzler, K. W.; Vogelstein, B.: Targeted deletion of Smad4 shows it is required for transforming growth factor beta and activin signaling in colorectal cancer cells. *Proc. Nat. Acad. Sci.* 95: 2412–2416, 1998.

[73800] 12130. Abbs, S.; Roberts, R. G.; Mathew, C. G.; Bentley, D.

R.; Bobrow, M.: Accurate assessment of intragenic recombination frequency within the Duchenne muscular dystrophy gene. *Genomics* 7: 602–606, 1990.

[73801] 12131. Ahn, A. H.; Kunkel, L. M.: The structural and functional diversity of dystrophin. *Nature Genet.* 3: 283–291, 1993.

[73802] 12132. Alwine, J. C.; Kemp, D. J.; Stark, G. R.: Method for detection of specific RNAs in agarose gels by transfer to diazobenzyloxymethyl-paper and by hybridization with DNA probes. *Proc. Nat. Acad. Sci.* 74: 5350–5354, 1977.

[73803] 12133. Angelini, C.; Beggs, A. H.; Hoffman, E. P.; Fanin, M.; Kunkel, L. M.: Enormous dystrophin in a patient with Becker muscular dystrophy. *Neurology* 40: 808–812, 1990.

[73804] 12134. Badorff, C.; Berkely, N.; Mehrotra, S.; Talhouk, J. W.; Rhoads, R. E.; Knowlton, K. U.: Enteroviral protease 2A directly cleaves dystrophin and is inhibited by a dystrophin-based substrate analogue. *J. Biol. Chem.* 275: 11191–11197, 2000.

[73805] 12135. Badorff, C.; Lee, G.-H.; Lamphear, B. J.; Martone, M. E.; Campbell, K. P.; Rhoads, R. E.; Knowlton, K. U.: Enteroviral protease 2A cleaves dystrophin: evidence of cytoskeletal disruption in an acquired cardiomyopathy. *Nature Med.* 5: 320–326, 1999.

- [73806] 12136. Bakker, E.; Pearson, P. L.: Mutation of the Duchenne muscular dystrophy gene associated with meiotic recombination. (Letter) Clin. Genet. 30: 347–349, 1986.
- [73807] 12137. Bakker, E.; Hofker, M. H.; Goor, N.; Mandel, J. L.; Wrogemann, K.; Davies, K. E.; Kunkel, L. M.; Willard, H. F.; Fenton, W. A.; Sandkuyl, L.; Majoor-Krakauer, D.; van Essen, A. J.; Jahoda, M. G. J.; Sachs, E. S.; van Ommen, G. J. B.; Pearson, P. L.: Prenatal diagnosis and carrier detection of Duchenne muscular dystrophy with closely linked RFLPs. Lancet I: 655–658, 1985.
- [73808] 12138. Bakker, E.; Van Broeckhoven, C.; Bonten, E. J.; van de Vooren, M. J.; Veenema, H.; Van Hul, W.; Van Ommen, G. J. B.; Vandenberghe, A.; Pearson, P. L.: Germline mosaicism and Duchenne muscular dystrophy mutations. Nature 329: 554–556, 1987.
- [73809] 12139. Bar, S.; Barnea, E.; Levy, Z.; Neuman, S.; Yaffe, D.; Nudel, U.: A novel product of the Duchenne muscular dystrophy gene which greatly differs from the known isoforms in its structure and tissue distribution. Biochem. J. 272: 557–560, 1990.
- [73810] 12140. Barbieri, A. M.; Soriani, N.; Tubiello, G. M.; Ferrari, M.; Carrera, P.: A nonsense mutation (gln-673-term) in



exon 17 of the human dystrophingene detected by heteroduplex analysis. Hum. Genet. 96: 343–344,1995.

[73811] 12141.Bartlett, R. J.; Pericak–Vance, M. A.; Koh, J.; Yamaoka, L. H.;Chen, J. C.; Hung, W.–Y.; Speer, M. C.; Wapenaar, M. C.; Van Ommen,G. J. B.; Bakker, E.; Pearson, P. L.; Kandt, R. S.; Siddique, T.;Gilbert, J. R.; Lee, J. E.; Sirotkin–Roses, M. J.; Roses, A. D.:Duchenne muscular dystrophy: high frequency of deletions. Neurology 38:1–4, 1988.

[73812] 12142.Barton–Davis, E. R.; Cordier, L.; Shoturma, D. I.; Leland, S.E.; Sweeney, H. L.: Aminoglycoside antibiotics restore dystrophinfunction to skeletal muscles of mdx mice. J. Clin. Invest. 104:375–381, 1999.

[73813] 12143.Bastianutto, C.; Bestard, J. A.; Lahnakoski, K.; Broere, D.; DeVisser, M.; Zaccolo, M.; Pozzan, T.; Ferlini, A.; Muntoni, F.; Patarnello,T.; Klamut, H. J.: Dystrophin muscle enhancer 1 is implicated inthe activation of non-muscle isoforms in the skeletal muscle of patientswith X-linked dilated cardiomyopathy. Hum. Molec. Genet. 10: 2627–2635,2001.

[73814] 12144.Baumbach, L. L.; Chamberlain, J. S.; Ward, P. A.; Farwell, N.J.; Caskey, C. T.: Molecular and clinical correlation of deletionleading to Duchenne and Becker muscular dystrophies. Neurology 39:465–474, 1989.

- [73815] 12145.Baumbach, L. L.; Ward, P. A.; Fenwick, R.; Caskey, C. T.: Analysis of mutations at the Duchenne muscular dystrophy locus provides no evidence for illegitimate recombination in deletion formation. (Abstract) Am.J. Hum. Genet. 45 (suppl.): A173, 1989.
- [73816] 12146.Beggs, A. H.; Koenig, M.; Boyce, F. M.; Kunkel, L. M.: Detection of 98% of DMD/BMD gene deletions by polymerase chain reaction. Hum.Genet. 86: 45–48, 1990.
- [73817] 12147.Berko, B. A.; Swift, M.: X-linked dilated cardiomyopathy. NewEng. J. Med. 316: 1186–1191, 1987.
- [73818] 12148.Bettecken, T.; Muller, C. R.: Identification of a 220-kb insertion into the Duchenne gene in a family with an atypical course of muscular dystrophy. Genomics 4: 592–596, 1989.
- [73819] 12149.Bies, R. D.: X-linked dilated cardiomyopathy. (Letter) New Eng.J. Med. 330: 368–369, 1994.
- [73820] 12150.Bies, R. D.; Caskey, C. T.; Fenwick, R.: An intact cysteine-rich domain is required for dystrophin function. J. Clin. Invest. 90:666–672, 1992.
- [73821] 12151.Bittner, R. E.; Streubel, B.; Shorny, S.; Schaden, G.; Voit, T.; Hoyer, H.: Coisogenic all-plus-one immunization: a model for identifying missing proteins in null-mutant conditions. Antibodies to dystrophin in mdx mouse after

transplantation of muscle from normal coisogenic donor.  
Neuropediatrics 25: 176–182, 1994.

- [73822] 12152. Ryu, S.; Zhou, S.; Ladurner, A. G.; Tjian, R.: The transcriptional cofactor complex CRSP is required for activity of the enhancer-binding protein Sp1. Nature 397: 446–450, 1999.
- [73823] 12153. Mohapatra, B.; Verma, S.; Shankar, S.; Suri, A.: Molecular cloning of human testis mRNA specifically expressed in haploid germ cells, having structural homology with the A-kinase anchoring proteins. Biochem. Biophys. Res. Commun. 244: 540–545, 1998.
- [73824] 12154. Turner, R. M. O.; Johnson, L. R.; Haig-Ladewig, L.; Gerton, G. L.; Moss, S. B.: An X-linked gene encodes a major human sperm fibrous sheath protein, hAKAP82: genomic organization, protein kinase A-R11 binding, and distribution of the precursor in the sperm tail. J. Biol. Chem. 273: 32135–32141, 1998.
- [73825] 12155. Dever, T. E.; Wei, C.-L.; Benkowski, L. A.; Browning, K.; Merrick, W. C.; Hershey, J. W. B.: Determination of the amino acid sequence of rabbit, human, and wheat germ protein synthesis factor eIF-4C by cloning and chemical sequencing. J. Biol. Chem. 269: 3212–3218, 1994.
- [73826] 12156. Lee, S.-H.; Kim, W.-H.; Kim, H.-K.; Woo, K.-M.;

Nam, H.-S.; Kim, H.-S.; Kim, J.-G.; Cho, M.-H.: Altered expression of the fragile histidine triad gene in primary gastric adenocarcinomas. *Biochem. Biophys. Res. Commun.* 284: 850–855, 2001.

[73827] 12157. Markkanen, A.; Heinonen, K.; Knuutila, S.; de la Chapelle, A.: Methotrexate-induced increase in gap formation in human chromosome band 3p14. *Hereditas* 96: 317–319, 1982.

[73828] 12158. Markkanen, A.; Knuutila, S.; de la Chapelle, A.: Inducible fragile site on chromosome 3. (Letter) *Hum. Genet.* 65: 217 only, 1983.

[73829] 12159. Anderson, D. M.; Kumaki, S.; Ahdieh, M.; Bertles, J.; Tometsko, M.; Loomis, A.; Giri, J.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Shapiro, D. N.; Morris, S. W.; Park, L. S.; Cosman, D.: Functional characterization of the human interleukin-15 receptor alpha chain and close linkage of IL15RA and IL2RA genes. *J. Biol. Chem.* 270: 29862–29869, 1995.

[73830] 12160. Giri, J. G.; Kumaki, S.; Ahdieh, M.; Friend, D. J.; Loomis, A.; Shanebeck, K.; DuBose, R.; Cosman, D.; Park, L. S.; Anderson, D. M.: Identification and cloning of a novel IL-15 binding protein that is structurally related to the alpha of the IL-2 receptor. *EMBO J.* 14:3654–3663, 1995.

- [73831] 12161.Yasunaga, S.; Grati, M.; Cohen-Salmon, M.; El-Amraoui, A.; Mustapha, M.; Salem, N.; El-Zir, E.; Loiselet, J.; Petit, C.: A mutation in OTOF, encoding otoferlin, a FER-1-like protein, causes DFNB9, a nonsyndromic form of deafness. *Nature Genet.* 21: 363-369, 1999.
- [73832] 12162.Berry, A.; Scott, H. S.; Kudoh, J.; Talior, I.; Korostishevsky, M.; Wattenhofer, M.; Guipponi, M.; Barras, C.; Rossier, C.; Shibuya, K.; Wang, J.; Kawasaki, K.; Asakawa, S.; Minoshima, S.; Shimizu, N.; Antonarakis, S.; Bonne-Tamir, B.: Refined localization of autosomal recessive nonsyndromic deafness DFNB10 locus using 34 novel microsatellite markers, genomic structure, and exclusion of six known genes in the region. *Genomics* 68: 22-29, 2000.
- [73833] 12163.Good, P. J.; Chen, Q.; Warner, S. J.; Herring, D. C.: A family of human RNA-binding proteins related to the *Drosophila* Bruno transcriptional regulator. *J. Biol. Chem.* 275: 28583-28592, 2000.
- [73834] 12164.Roberts, R.; Timchenko, N. A.; Miller, J. W.; Reddy, S.; Caskey, C. T.; Swanson, M. S.; Timchenko, L. T.: Altered phosphorylation and intracellular distribution of a (CUG)<sub>n</sub> triplet repeat RNA-binding protein in patients with myotonic dystrophy and in myotonin protein kinase knockout mice. *Proc. Nat. Acad. Sci.* 94: 13221-13226, 1997.

- [73835] 12165.Li, Z.; Jiang, H.; Xie, W.; Zhang, Z.; Smrcka, A. V.; Wu, D.:Roles of PLC-beta-2 and -beta-3 and PI3K-gamma in chemoattractant-mediated signal transduction. *Science* 287: 1046-1049, 2000.
- [73836] 12166.Fowler, K. J.; Saffery, R.; Irvine, D. V.; Trowell, H. E.; Choo, K. H. A.: Mouse centromere protein F (Cenpf) gene maps to the distal region of chromosome 1 by interspecific backcross analysis. *Cytogenet. Cell Genet.* 82: 180-181, 1998.
- [73837] 12167.Liao, H.; Winkfein, R. J.; Mack, G.; Rattner, J. B.; Yen, T. J.: CENP-F is a protein of the nuclear matrix that assembles onto kinetochores at late G2 and is rapidly degraded after mitosis. *J. Cell Biol.* 130:507-518, 1995.
- [73838] 12168.Rattner, J. B.; Rao, A.; Fritzler, M. J.; Valencia, D. W.; Yen, T. J.: CENP-F is a ca 400 kDa kinetochore protein that exhibits a cell-cycle dependent localization. *Cell Motil. Cytoskeleton* 26:214-226, 1993.
- [73839] 12169.Panagopoulos, I.; Mencinger, M.; Dietrich, C. U.; Bjerkehagen, B.; Saeter, G.; Mertens, F.; Mandahl, N.; Heim, S.: Fusion of the RBP56 and CHN genes in extraskeletal myxoid chondrosarcomas with translocation (9;17)(q22;q11). *Oncogene* 18: 7594-7598, 1999.
- [73840] 12170.Gassmann, M.; Casagrande, F.; Orioli, D.; Simon,

H.; Lai, C.; Klein, R.; Lemke, G.: Aberrant neural and cardiac development in mice lacking the ErbB4 neuregulin receptor. *Nature* 378: 390–394, 1995.

[73841] 12171. Golding, J. P.; Trainor, P.; Krumlauf, R.; Gassmann, M.: Defects in pathfinding by cranial neural crest cells in mice lacking the neuregulin receptor ErbB4. *Nature Cell Biol.* 2: 103–109, 2000.

[73842] 12172. Zimonjic, D. B.; Alimandi, M.; Miki, T.; Popescu, N. C.; Kraus, M. H.: Localization of the human HER4/erbB-4 gene to chromosome 2. *Oncogene* 10:1235–1237, 1995.

[73843] 12173. Adibi, S. A.: The oligopeptide transporter (Pept-1) in human intestine: biology and function. *Gastroenterology* 113: 332–340, 1997.

[73844] 12174. Fei, Y. -J.; Kanai, Y.; Nussberger, S.; Ganapathy, V.; Leibach, F. H.; Romero, M. F.; Singh, S. K.; Boron, W. F.; Hediger, M. A.: Expression cloning of a mammalian proton-coupled oligopeptide transporter. *Nature* 368:563–566, 1994.

[73845] 12175. Liang, R.; Fei, Y. -J.; Prasad, P. D.; Ramamoorthy, S.; Han, H.; Yang-Feng, T. L.; Hediger, M. A.; Ganapathy, V.; Leibach, F. H.: Human intestinal H(+)/peptide cotransporter: cloning, functional expression, and chromosomal localization. *J. Biol. Chem.* 270: 6456–6463, 1995.

- [73846] 12176.Heiber, M.; Docherty, J. M.; Shah, G.; Nguyen, T.; Cheng, R.; Heng,H. H. Q.; Marchese, A.; Tsui, L.-C.; Shi, X.; George, S. R.; O'Dowd,B. F.: Isolation of three novel human genes encoding G protein-coupledreceptors. *DNA Cell Biol.* 14: 25-35, 1995.
- [73847] 12177.Mahadevan, M. S.; Baird, S.; Bailly, J. E.; Shutler, G. G.; Sabourin,L. A.; Tsilfidis, C.; Neville, C. E.; Narang, M.; Korneluk, R. G.: Isolation of a novel G protein-coupled receptor (GPR4) localizedto chromosome 19q13.3. *Genomics* 30: 84-88, 1995.
- [73848] 12178.Ye, R. D.; Prossnitz, E. R.; Zou, A.; Cochrane, C. G.: Characterizationof a human cDNA that encodes a functional receptor for platelet activatingfactor. *Biochem. Biophys. Res. Commun.* 180: 105-111, 1991.
- [73849] 12179.Hossain, A.; Saunders, G. F.: The human sex-determining geneSRY is a direct target of WT1. *J. Biol. Chem.* 276: 16817-16823,2001.
- [73850] 12180.Iida, T.; Nakahori, Y.; Komaki, R.; Mori, E.; Hayashi, N.; Tsutsumi,O.; Taketani, Y.; Nakagome, Y.: A novel mis-sense mutation in theHMG box of the SRY gene in a patient with XY sex reversal. *Hum. Molec.Genet.* 3: 1437-1438, 1994.
- [73851] 12181.Jager, R. J.; Harley, V. R.; Pfeiffer, R. A.; Goodfellow,



P. N.;Scherer, G.: A familial mutation in the testis-determining gene SRYshared by both sexes. Hum. Genet. 90: 350–355, 1992.

[73852] 12182.Kawakura, K.; Miyake, Y.–I.; Murakami, R.–K.; Kondoh, S.; Hirata,T.–I.; Kaneda, Y.: Deletion of the SRY region on the Y chromosomedetected in bovine gonadal hypoplasia (XY female) by PCR. Cytogenet.Cell Genet. 72: 183–184, 1996.

[73853] 12183.Lahr, G.; Maxson, S. C.; Mayer, A.; Just, W.; Pilgrim, C.; Reisert,I.: Transcription of the Y chromosomal gene, Sry, in adult mousebrain. Molec. Brain Res. 33: 179–182, 1995.

[73854] 12184.Li, B.; Zhang, W.; Chan, G.; Jancso–Radek, A.; Liu, S.; Weiss,M. A.: Human sex reversal due to impaired nuclear localization ofSRY: a clinical correlation. J. Biol. Chem. 276: 46480–46484, 2001.

[73855] 12185.Lovell–Badge, R.; Robertson, E.: XY female mice resulting froma heritable mutation in the primary testis-determining gene, Tdy. Development 109:635–646, 1990.

[73856] 12186.Margarit, E.; Coll, M. D.; Oliva, R.; Gomez, D.; Soler, A.; Ballesta,F.: SRY gene transferred to the long arm of the X chromosome in aY–positive XX true hermaphrodite. Am. J. Med. Genet. 90: 25–28,2000.

- [73857] 12187.Mittwoch, U.: Sex determination and sex reversal: genotype, phenotype,dogma and semantics. Hum. Genet. 89: 467–479, 1992.
- [73858] 12188.Muller, J.; Schwartz, M.; Skakkebaek, N. E.: Analysis of the sex–determining region of the Y–chromosome (SRY) in sex reversed patients:point–mutation in SRY causing sex–reversion in a 46,XY female. J.Clin. Endocr. Metab. 75: 331–333, 1992.
- [73859] 12189.Peters, R.; King, C.–Y.; Ukiyama, E.; Falsafi, S.; Donahoe, P.K.; Weiss, M. A.: An SRY mutation causing human sex reversal resolves a general mechanism of structure–specific DNA recognition: application to the four–way DNA junction. Biochemistry 34: 4569–4576, 1995.
- [73860] 12190.Schaffler, A.; Barth, N.; Winkler, K.; Zietz, B.; Rummeler, P.;Knuchel, R.; Scholmerich, J.; Palitzsch, K.–D.: Identification of a new missense mutation (gly95glu) in a highly conserved codon within the high–mobility group box of the sex–determining region Y gene:report on a 46,XY female with gonadal dysgenesis and yolk–sac tumor. J.Clin. Endocr. Metab. 85: 2287–2292, 2000.
- [73861] 12191.Schmitt–Ney, M.; Thiele, H.; Kaltwasser, P.; Bardoni, B.; Cisternino,M.; Scherer, G.: Two novel SRY missense mutations reducing DNA binding identified in XY females

and their mosaic fathers. *Am. J. Hum. Genet.* 56:862–869, 1995.

[73862] 12192.Sinclair, A. H.; Berta, P.; Palmer, M. S.; Hawkins, R.; Griffiths,B. L.; Smith, M.; Foster, J. W.; Frischauf, A.–M.; Lovell–Badge, R.;Goodfellow, P. N.: A gene from the human sex–determining region encodesa protein with homology to a conserved DNA–binding motif.. *Nature* 346:240–245, 1990.

[73863] 12193.Su, H.; Lau, Y.–F. C.: Identification of the transcriptionalunit, structural organization, and promoter sequence of the humansex–determining region Y (SRY) gene, using a reverse genetic approach. *Am.J. Hum. Genet.* 52: 24–38, 1993.

[73864] 12194.Tucker, P. K.; Lundrigan, B. L.: Rapid evolution of the sex determininglocus in Old World mice and rats. *Nature* 364: 715–717, 1993.

[73865] 12195.Uehara, S.; Funato, T.; Yaegashi, N.; Suzuki, H.; Sato, J.; Sasaki,T.; Yajima, A.: SRY mutation and tumor formation on the gonads ofXY pure gonadal dysgenesis patients. *Cancer Genet. Cytogenet.* 113:78–84, 1999.

[73866] 12196.Uehara, S.; Hashiyada, M.; Sato, K.; Nata, M.; Funato, T.; Okamura,K.: Complete XY gonadal dysgenesis and aspects of the SRY genotypeand gonadal tumor for–

mation. J. Hum. Genet. 47: 279–284, 2002.

- [73867] 12197.Veitia, R.; Ion, A.; Barbaux, S.; Jobling, M. A.; Souleyreau, N.; Ennis, K.; Ostrer, H.; Tosi, M.; Meo, T.; Chibani, J.; Fellous, M.; McElreavey, K.: Mutations and sequence variants in the testis-determining region of the Y chromosome in individuals with a 46,XY female phenotype. Hum.Genet. 99: 648–652, 1997.
- [73868] 12198.Werner, M. H.; Huth, J. R.; Gronenborn, A. M.; Clore, G. M.:Molecular basis of human 46X,Y (sic) sex reversal revealed from the three-dimensional solution structure of the human SRY–DNA complex. Cell 81:705–714, 1995.
- [73869] 12199.Whitfield, L. S.; Lovell–Badge, R.; Goodfellow, P. N.: Rapid sequence evolution of the mammalian sex-determining gene SRY. Nature 364:713–715, 1993.
- [73870] 12200.Zeng, Y.; Ren, Z.; Zhang, M.; Huang, Y.; Zeng, F.; Huang, S.:A new de novo mutation (A113T) in HMG box of the SRY gene leads to XY gonadal dysgenesis. J. Med. Genet. 30: 655–657, 1993.
- [73871] 12201.Zenteno, J. C.; Lopez, M.; Vera, C.; Mendez, J. P.; Kofman–Alfaro, S.: Two SRY–negative XX male brothers without genital ambiguity. Hum.Genet. 100: 606–610, 1997.

- [73872] 12202.Henkemeyer, M.; Orioli, D.; Henderson, J. T.; Saxton, T. M.; Roder,J.; Pawson, T.; Klein, R.: Nuk controls pathfinding of commissuralaxons in the mammalian central nervous system. *Cell* 86: 35–46, 1996.
- [73873] 12203.Ikegaki, N.; Tang, X. X.; Liu, X.–G.; Biegel, J. A.; Allen, C.;Yoshioka, A.; Sulman, E. P.; Brodeur, G. M.; Pleasure, D. E.: Molecularcharacterization and chromosomal localization of DRT (EPHT3): a developmentallyregulated human protein–tyrosine kinase gene of the EPH family. *Hum.Molec. Genet.* 4: 2033–2045, 1995.
- [73874] 12204.Saito, T.; Seki, N.; Matsuda, Y.; Kitahara, M.; Murata, M.; Kanda,N.; Nomura, N.; Yamamoto, T.; Hori, T.: Identification of the humanERK gene as a putative receptor tyrosine kinase and its chromosomallocalization to 1p36.1: a comparative mapping of human, mouse, andrat chromosomes. *Genomics* 26: 382–384, 1995.
- [73875] 12205.Wybenga–Groot, L. E.; Baskin, B.; Ong, S. H.; Tong, J.; Pawson,T.; Sicheri, F.: Structural basis for autoinhibition of the EphB2receptor tyrosine kinase by the unphosphorylated juxtamembrane region. *Cell* 106:745–757, 2001.
- [73876] 12206.Johnson, D. F.; Hamon, M.; Fischel–Ghodsian, N.: Characterizationof the human mitochondrial ribosomal S12 gene. *Genomics* 52: 363–368,1998.

- [73877] 12207. Shah, Z. H.; Migliosi, V.; Miller, S. C. M.; Wang, A.; Friedman, T. B.; Jacobs, H. T.: Chromosomal locations of three human nucleargenes (RPSM12, TUFM, and AFG3L1) specifying putative components of the mitochondrial gene expression apparatus. *Genomics* 48: 384–388, 1998.
- [73878] 12208. Honore, B.; Leffers, H.; Madsen, P.; Celis, J. E.: Interferon- $\gamma$  up-regulates a unique set of proteins in human keratinocytes: molecular cloning and expression of the cDNA encoding the RGD-sequence-containing protein IGUP I5111. *Europ. J. Biochem.* 218: 421–430, 1993.
- [73879] 12209. Preckel, T.; Fung-Leung, W.-P.; Cai, Z.; Vitiello, A.; Salter-Cid, L.; Winqvist, O.; Wolfe, T. G.; Von Herrath, M.; Angulo, A.; Ghazal, P.; Lee, J.-D.; Fourie, A. M.; Wu, Y.; Pang, J.; Ngo, K.; Peterson, P. A.; Fruh, K.; Yang, Y.: Impaired immunoproteasome assembly and immune responses in PA28  $-/-$  mice. *Science* 286: 2162–2165, 1999.
- [73880] 12210. Realini, C.; Dubiel, W.; Pratt, G.; Ferrell, K.; Reichsteiner, M.: Molecular cloning and expression of a gamma-interferon-inducible activator of the multicatalytic protease. *J. Biol. Chem.* 269: 20727–20732, 1994.
- [73881] 12211. Trachtulec, Z.; Forejt, J.: Synteny of orthologous genes conserved in mammals, snake, fly, nematode, and

fission yeast. Mammalian Genome 12:227–231, 2001.

[73882] 12212.Reif, K.; Ekland, E. H.; Ohl, L.; Nakano, H.; Lipp, M.; Forster,R.; Cyster, J. G.: Balanced responsiveness to chemoattractants from adjacent zones determines B-cell position. Nature 416: 94–99, 2002.

[73883] 12213.Zhu, K.; Baudhuin, L. M.; Hong, G.; Williams, F. S.; Cristina,K. L.; Kabarowski, J. H. S.; Witte, O. N.; Xu, Y.: Sphingosylphosphorylcholine and lysophosphatidylcholine are ligands for the G protein-coupled receptor GPR4. J. Biol. Chem. 276: 41325–41335, 2001.

[73884] 12214.Vinik, B. S.; Kay, E. S.; Fiedorek, F. T., Jr.: Mapping of the MEK kinase gene (Mekk) to mouse chromosome 13 and human chromosome 5. Mammalian Genome 6: 782–783, 1995.

[73885] 12215.Yujiri, T.; Sather, S.; Fanger, G. R.; Johnson, G. L.: Role of MEKK1 in cell survival and activation of JNK and ERK pathways defined by targeted gene disruption. Science 282: 1911–1914, 1998.

[73886] 12216.Arriza, J. L.; Weinberger, C.; Cerelli, G.; Glaser, T. M.; Handelin, B. L.; Housman, D. E.; Evans, R. M.: Cloning of human mineralocorticoid receptor complementary DNA: structural and functional kinship with the glucocorticoid receptor. Science 237: 268–275, 1987.

- [73887] 12217.Berger, S.; Bleich, M.; Schmid, W.; Cole, T. J.; Peters, J.; Watanabe,H.; Kriz, W.; Warth, R.; Greger, R.; Schutz, G.: Mineralocorticoidreceptor knockout mice: pathophysiology of Na<sup>+</sup> metabolism. Proc.Nat. Acad. Sci. 95: 9424–9429, 1998.
- [73888] 12218.Fan, Y.–S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.M.; Nowak, N. J.; Shows, T. B.: The human mineralocorticoid receptorgene (MLR) is located on chromosome 4 at q31.2. Cytogenet. Cell Genet. 52:83–84, 1989.
- [73889] 12219.Geller, D. S.; Farhi, A.; Pinkerton, N.; Fradley, M.; Moritz, M.;Spitzer, A.; Meinke, G.; Tsai, F. T. F.; Sigler, P. B.; Lifton, R.P.: Activating mineralocorticoid receptor mutation in hypertensionexacerbated by pregnancy. Science 289: 119–123, 2000.
- [73890] 12220.Geller, D. S.; Rodriguez–Soriano, J.; Vallo Boado, A.; Schifter,S.; Bayer, M.; Chang, S. S.; Lifton, R. P.: Mutations in the mineralocorticoidreceptor gene cause autosomal dominant pseudohypoaldosteronism typel. Nature Genet. 19: 279–281, 1998.
- [73891] 12221.Hellal–Levy, C.; Fagart, J.; Souque, A.; Wurtz, J.–M.; Moras, D.;Rafestin–Oblin, M.–E.: Crucial role of the H11–H12 loop in stabilizingthe active conformation of the human mineralocorticoid receptor. Molec.Endocr. 14:



1210–1221, 2000.

- [73892] 12222. Le Menuet, D.; Isnard, R.; Bichara, M.; Viengchareun, S.; Muffat-Joly, M.; Walker, F.; Zennaro, M.-C.; Lombes, M.: Alteration of cardiac and renal functions in transgenic mice overexpressing human mineralocorticoid receptor. *J. Biol. Chem.* 276: 38911–38920, 2001.
- [73893] 12223. Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor, J. M.: Regional chromosomal assignment of the human mineralocorticoid receptor gene to 4q31.1. *Hum. Genet.* 85: 130–132, 1990.
- [73894] 12224. Morrison, N.; Harrap, S. B.; Arriza, J. L.; Boyd, E.; Connor, J. M.: Regional chromosomal assignment of the human mineralocorticoid receptor gene to 4q31.1. (Abstract) *Cytogenet. Cell Genet.* 51: 1048, 1989.
- [73895] 12225. Veenstra, G. J. C.; Weeks, D. L.; Wolffe, A. P.: Distinct roles for TBP and TBP-like factor in early embryonic gene transcription in *Xenopus*. *Science* 290: 2312–2314, 2000.
- [73896] 12226. Zuhlke, C.; Hellenbroich, Y.; Dalski, A.; Kononowa, N.; Hagenah, J.; Vieregge, P.; Riess, O.; Klein, C.; Schwinger, E.: Different types of repeat expansion in the TATA-binding protein gene are associated with a new form

of inherited ataxia. *Europ. J. Hum. Genet.* 9:  
160–164, 2001.

[73897] 12227. Charest, A.; Wagner, J.; Muise, E. S.; Heng, H. H. Q.; Tremblay, M. L.: Structure of the murine MPTP-PEST gene: genomic organization and chromosomal mapping. *Genomics* 28: 501–507, 1995.

[73898] 12228. Takekawa, M.; Itoh, F.; Hinoda, Y.; Adachi, M.; Ariyama, T.; Inazawa, J.; Imai, K.; Yachi, A.: Chromosomal localization of the protein tyrosine phosphatase G1 gene and characterization of the aberrant transcripts in human colon cancer cells. *FEBS Lett.* 339: 222–228, 1994.

[73899] 12229. Takekawa, M.; Itoh, F.; Hinoda, Y.; Arimura, Y.; Toyota, M.; Sekiya, M.; Adachi, M.; Imai, K.; Yachi, A.: Cloning and characterization of a human cDNA encoding a novel putative cytoplasmic protein-tyrosine-phosphatase. *Biochem. Biophys. Res. Commun.* 189: 1223–1230, 1992.

[73900] 12230. Yang, Q.; Co, D.; Sommercorn, J.; Tonks, N. K.: Cloning and expression of PTP-PEST: a novel, human, non-transmembrane protein tyrosine phosphatase. *J. Biol. Chem.* 268: 6622–6628, 1993.

[73901] 12231. Coopman, P. J. P.; Do, M. T. H.; Barth, M.; Bowden, E. T.; Hayes, A. J.; Basyuk, E.; Blancato, J. K.; Vezza, P. R.; McLeskey, S. W.; Mangeat, P. H.; Mueller, S. C.: The Syk ty-

rosine kinase suppresses malignant growth of human breast cancer cells. *Nature* 406: 742–747, 2000.

[73902] 12232. Inatome, R.; Yanagi, S.; Takano, T.; Yamamura, H.: A critical role for Syk in endothelial cell proliferation and migration. *Biochem. Biophys. Res. Commun.* 286: 195–199, 2001.

[73903] 12233. Mocsai, A.; Zhou, M.; Meng, F.; Tybulewicz, V. L.; Lowell, C. A.: Syk is required for integrin signaling in neutrophils. *Immunity* 16:547–558, 2002.

[73904] 12234. Muller, B.; Cooper, L.; Terhorst, C.: Molecular cloning of the human homologue to the pig protein-tyrosine kinase syk. *Immunogenetics* 39:359–362, 1994.

[73905] 12235. Taniguchi, T.; Kobayashi, T.; Kondo, J.; Takahashi, K.; Nakamura, H.; Suzuki, J.; Nagai, K.; Yamada, T.; Nakamura, S.; Yamamura, H.: Molecular cloning of a porcine gene syk that encodes a 72-kDa protein-tyrosine kinase showing high susceptibility to proteolysis. *J. Biol. Chem.* 266:15790–15796, 1991.

[73906] 12236. Zioncheck, T. F.; Harrison, M. L.; Isaacson, C. C.; Geahlen, R. L.: Generation of an active protein-tyrosine kinase from lymphocytes by proteolysis. *J. Biol. Chem.* 263: 19195–19202, 1988.

[73907] 12237. Satre, M. A.; Zgombic-Knight, M.; Duester, G.: The

complete structure of human class IV alcohol dehydrogenase (retinol dehydrogenase) determined from the ADH7 gene. J. Biol. Chem. 269: 15606–15612, 1994.

[73908] 12238. Yokoyama, H.; Baraona, E.; Lieber, C. S.: Molecular cloning and chromosomal localization of the ADH7 gene encoding human class IV (sigma) ADH. Genomics 31: 243–245, 1996.

[73909] 12239. Zgombic-Knight, M.; Foglio, M. H.; Duester, G.: Genomic structure and expression of the ADH7 gene encoding human class IV alcohol dehydrogenase, the form most efficient for retinol metabolism in vitro. J. Biol. Chem. 270: 4305–4311, 1995.

[73910] 12240. Bashir, M. M.; Abrams, W. R.; Tucker, T.; Sellinger, B.; Budarf, M.; Emanuel, B.; Rosenbloom, J.: Molecular cloning and characterization of the bovine and human tuftelin genes. Connect. Tissue Res. 39: 13–24, 1998.

[73911] 12241. Deutsch, D.: Structure and function of enamel gene products. Anat. Rec. 224: 189–210, 1989.

[73912] 12242. Beguin, P.; Nagashima, K.; Gono, T.; Shibasaki, T.; Takahashi, K.; Kashima, Y.; Ozaki, N.; Geering, K.; Iwanaga, T.; Seino, S.: Regulation of  $\text{Ca}^{2+}$  channel expression at the cell surface by the small G-protein kir/Gem. Nature 411: 701–706, 2001.

- [73913] 12243. Maguire, J.; Santoro, T.; Jensen, P.; Siebenlist, U.; Yewdell, J.; Kelly, K.: GEM: an induced, immediate early protein belonging to the Ras family. *Science* 265: 241–244, 1994.
- [73914] 12244. Santoro, T.; Maguire, J.; McBride, O. W.; Avraham, K. B.; Copeland, N. G.; Jenkins, N. A.; Kelly, K.: Chromosomal organization and transcriptional regulation of human GEM and localization of the human and mouse Gem loci encoding an inducible Ras-like protein. *Genomics* 30: 558–564, 1995.
- [73915] 12245. Inoue, I.; Taniuchi, I.; Kitamura, D.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Watanabe, T.: Characteristics of the mouse genomic histamine H1 receptor gene. *Genomics* 36: 178–181, 1996.
- [73916] 12246. Le Coniat, M.; Traiffort, E.; Ruat, M.; Arrang, J.-M.; Berger, R.: Chromosomal localization of the human histamine H1-receptor gene. *Hum. Genet.* 94: 186–188, 1994.
- [73917] 12247. Ma, R. Z.; Gao, J.; Meeker, N. D.; Fillmore, P. D.; Tung, K. S. K.; Watanabe, T.; Zachary, J. F.; Offner, H.; Blankenhorn, E. P.; Teuscher, C.: Identification of Bphs, an autoimmune disease locus, as histamine receptor H-1. *Science* 297: 620–623, 2002.

- [73918] 12248.Yamashita, M.; Fukui, H.; Sugama, K.; Horio, Y.; Ito, S.; Mizuguchi,H.; Wada, H.: Expression cloning of a cDNA encoding the bovine histamineH1 receptor. *Proc. Nat. Acad. Sci.* 88: 11515–11519, 1991.
- [73919] 12249.Bauer, S.; Groh, V.; Wu, J.; Steinle, A.; Phillips, J. H.; Lanier,L. L.; Spies, T.: Activation of NK cells and T cells by NKG2D, areceptor for stress–inducible MICA. *Science* 285: 727–729, 1999.
- [73920] 12250.Diefenbach, A.; Jensen, E. R.; Jamieson, A. M.; Raulet, D. H.:Rae1 and H60 ligands of the NKG2D receptor stimulate tumour immunity. *Nature* 413:165–171, 2001.
- [73921] 12251.Girardi, M.; Oppenheim, D. E.; Steele, C. R.; Lewis, J. M.; Glusac,E.; Filler, R.; Hobby, P.; Sutton, B.; Tigelaar, R. E.; Hayday, A.C.: Regulation of cutaneous malignancy by gamma–delta T cells. *Science* 294:605–609, 2001.
- [73922] 12252.Groh, V.; Rhinehart, R.; Randolph–Habecker, J.; Topp, M. S.; Riddell,S. R.; Spies, T.: Costimulation of CD8–alpha–beta T cells by NKG2Dvia engagement by MIC induced on virus–infected cells. *Nature Immun.* 2:255–260, 2001.
- [73923] 12253.Groh, V.; Steinle, A.; Bauer, S.; Spies, T.: Recogni– tion of stress–inducedMHC molecules by intestinal epithe– lial gamma–delta T cells. *Science* 279:1737–1740, 1998.

- [73924] 12254.Groh, V.; Wu, J.; Yee, C.; Spies, T.: Tumour-derived soluble MICligands impair expression of NKG2D and T-cell activation. *Nature* 419:734–738, 2002.
- [73925] 12255.Li, P.; Morris, D. L.; Willcox, B. E.; Steinle, A.; Spies, T.;Strong, R. K.: Complex structure of the activating immunoreceptorNKG2D and its MHC class I-like ligand MICA. *Nature Immun.* 2: 443–451,2001.
- [73926] 12256.Jepsen, K. J.; Wu, F.; Peragallo, J. H.; Paul, J.; Roberts, L.;Ezura, Y.; Oldberg, A.; Birk, D. E.; Chakravarti, S.: A syndromeof joint laxity and impaired tendon integrity in lumican- and fibromodulin-deficientmice. *J. Biol. Chem.* 277: 35532–35540, 2002.
- [73927] 12257.Sztrolovics, R.; Chen, X.-N.; Grover, J.; Roughley, P. J.; Korenberg,J. R.: Localization of the human fibromodulin gene (FMOD) to chromosome1q32 and completion of the cDNA sequence. *Genomics* 23: 715–717,1994.
- [73928] 12258.Dalton, S.; Treisman, R.: Characterization of SAP-1, a proteinrecruited by serum response factor to the c-fos serum response element. *Cell* 68:597–612, 1992.
- [73929] 12259.Mo, Y.; Vaessen, B.; Johnston, K.; Marmorstein, R.: Structuresof SAP-1 bound to DNA targets from the E74 and c-fos promoters: insightsinto DNA sequence discrimination by Ets proteins. *Molec. Cell* 2:201–212, 1998.

- [73930] 12260.Shipley, J.; Sheer, D.; Dalton, S.; Treisman, R.; Patel, K.: Mapping of the human SAP1 (SRF accessory protein 1) gene and SAP2, a gene encoding a related protein, to chromosomal bands 1q32 and 12q23, respectively. *Genomics* 23:710–711, 1994.
- [73931] 12261.Giovane, A.; Pintzas, A.; Maira, S.–M.; Sobieszczuk, P.; Wasylyk, B.: Net, a new ets transcription factor that is activated by Ras. *Genes Dev.* 8: 1502–1513, 1994.
- [73932] 12262.Lopez, M.; Oettgen, P.; Akbarali, Y.; Dendorfer, U.; Libermann, T. A.: ERP, a new member of the ets transcription factor/oncoprotein family: cloning, characterization, and differential expression during B-lymphocyte development. *Molec. Cell. Biol.* 14: 3292–3309, 1994.
- [73933] 12263.Kelner, G. S.; Kennedy, J.; Bacon, K. B.; Kleyensteuber, S.; Largaespada, D. A.; Jenkins, N. A.; Copeland, N. G.; Bazan, J. F.; Moore, K. W.; Schall, T. J.; Zlotnik, A.: Lymphotactin: a cytokine that represents a new class of chemokine. *Science* 266: 1395–1399, 1994.
- [73934] 12264.Kennedy, J.; Kelner, G. S.; Kleyensteuber, S.; Schall, T. J.; Weiss, M. C.; Yssel, H.; Schneider, P. V.; Cocks, B. G.; Bacon, K. B.; Zlotnik, A.: Molecular cloning and functional characterization of human lymphotactin. *J. Immun.* 155: 203–209, 1995.



- [73935] 12265.Muller, S.; Dorner, B.; Korthauer, U.; Mages, H. W.; D'Apuzzo,M.; Senger, G.; KroczeK, R. A.: Cloning of ATAC, an activation–induced,chemokine–related molecule exclusively expressed in CD8+ T lymphocytes. *Europ.J. Immun.* 25: 1744–1748, 1995.
- [73936] 12266.Yoshida, T.; Imai, T.; Kakizaki, M.; Nishimura, M.; Takagi, S.;Yoshie, O.: Identification of single C motif–1/lymphotactin receptorXCR1. *J. Biol. Chem.* 273: 16551–16554, 1998.
- [73937] 12267.Yoshida, T.; Imai, T.; Kakizaki, M.; Nishimura, M.; Yoshie, O.: Molecular cloning of a novel C or gamma type chemokine, SCM–1. *FEBSLett.* 360: 155–159, 1995.
- [73938] 12268.Yoshida, T.; Imai, T.; Takagi, S.; Nishimura, M.; Ishikawa, I.;Yaoi, T.; Yoshie, O.: Structure and expression of two highly relatedgenes encoding SCM–1/human lymphotactin. *FEBS Lett.* 395: 82–88,1996.
- [73939] 12269.Ikeuchi, T.; Asaka, T.; Saito, M.; Tanaka, H.; Higuchi, S.; Tanaka,K.; Saida, K.; Uyama, E.; Mizusawa, H.; Fukuhara, N.; Nonaka, I.;Takamori, M.; Tsuji, S.: Gene locus for autosomal recessive distalmyopathy with rimmed vacuoles maps to chromosome 9. *Ann. Neurol.* 41:432–437, 1997.
- [73940] 12270.Erickson, J. D.; Varoqui, H.; Schafer, M. K.–H.; Modi,

W.; Diebler, M.-F.; Weihe, E.; Rand, J.; Eiden, L. E.; Bonner, T. I.; Usdin, T.B.: Functional identification of a vesicular acetylcholine transporter and its expression from a 'cholinergic' gene locus. *J. Biol. Chem.* 269:21929–21932, 1994.

[73941] 12271. Tanahashi, H.; Tabira, T.: Alzheimer's disease-associated presenilin2 interacts with DRAL, an LIM-domain protein. *Hum. Molec. Genet.* 9:2281–2289, 2000.

[73942] 12272. Erkinheimo, T.-L.; Saukkonen, K.; Narko, K.; Jalkanen, J.; Ylikorkala, O.; Ristimäki, A.: Expression of cyclooxygenase-2 and prostanoid receptors by human myometrium. *J. Clin. Endocr. Metab.* 85: 3468–3475, 2000.

[73943] 12273. Hla, T.; Neilson, K.: Human cyclooxygenase-2 cDNA. *Proc. Nat. Acad. Sci.* 89: 7384–7388, 1992.

[73944] 12274. Jones, D. A.; Carlton, D. P.; McIntyre, T. M.; Zimmerman, G. A.; Prescott, S. M.: Molecular cloning of human prostaglandin endoperoxidesynthase type II and demonstration of expression in response to cytokines. *J. Biol. Chem.* 268: 9049–9054, 1993.

[73945] 12275. Kraemer, S. A.; Meade, E. A.; DeWitt, D. L.: Prostaglandin endoperoxidesynthase gene structure: identification of the transcriptional start site and 5-prime-flanking regulatory sequences. *Arch. Biochem. Biophys.* 293:391–400, 1992.

- [73946] 12276.Lassus, P.; Wolff, H.; Andersson, S.: Cyclooxygenase-2 in human perinatal lung. *Pediat. Res.* 47: 602–605, 2000.
- [73947] 12277.Lim, H.; Paria, B. C.; Das, S. K.; Dinchuk, J. E.; Langenbach, R.; Trzaskos, J. M.; Dey, S. K.: Multiple female reproductive failures in cyclooxygenase 2-deficient mice. *Cell* 91: 197–208, 1997.
- [73948] 12278.Jacquemin, P.; Durviaux, S. M.; Jensen, J.; Godfraind, C.; Gradwohl, G.; Guillemot, F.; Madsen, O. D.; Carmeliet, P.; Dewerchin, M.; Collen, D.; Rousseau, G. G.; Lemaigre, F. P.: Transcription factor hepatocyte nuclear factor 6 regulates pancreatic endocrine cell differentiation and controls expression of the proendocrine gene *ngn3*. *Molec. Cell Biol.* 20: 4445–4454, 2000.
- [73949] 12279.Jacquemin, P.; Lannoy, V. J.; Rousseau, G. G.; Lemaigre, F. P.: OC-2, a novel mammalian member of the ONECUT class of homeodomain transcription factors whose function in liver partially overlaps with that of hepatocyte nuclear factor-6. *J. Biol. Chem.* 274: 2665–2671, 1999.
- [73950] 12280.Lemaigre, F. P.; Durviaux, S. M.; Truong, O.; Lannoy, V. J.; Hsuan, J. J.; Rousseau, G. G.: Hepatocyte nuclear factor 6, a transcription factor that contains a novel type of

homeodomain and a single cutdomain. Proc. Nat. Acad. Sci. 93: 9460–9464, 1996.

- [73951] 12281. Pierreux, C. E.; Stafford, J.; Demonte, D.; Scott, D. K.; Vandenhoute, J.; O'Brien, R. M.; Granner, D. K.; Rousseau, G. G.; Lemaigre, F. P.: Antigluco-corticoid activity of hepatocyte nuclear factor-6. Proc. Nat. Acad. Sci. 96: 8961–8966, 1999.
- [73952] 12282. Samadani, U.; Costa, R. H.: The transcriptional activator hepatocyte nuclear factor 6 regulates liver gene expression. Molec. Cell. Biol. 16: 6273–6284, 1996.
- [73953] 12283. Vaisse, C.; Kim, J.; Espinosa, R., III; Le Beau, M. M.; Stoffel, M.: Pancreatic islet expression studies and polymorphic DNA markers in the genes encoding hepatocyte nuclear factor-3- $\alpha$ , -3- $\beta$ , -4- $\gamma$ , and -6. Diabetes 46: 1364–1367, 1997.
- [73954] 12284. Iacobazzi, V.; Palmieri, F.; Runswick, M. J.; Walker, J. E.: Sequences of the human and bovine genes for the mitochondrial 2-oxoglutarate carrier. DNA Seq. 3: 79–88, 1992.
- [73955] 12285. Piccininni, S.; Iacobazzi, V.; Lauria, G.; Rocchi, M.; Palmieri, F.: Assignment of the oxoglutarate carrier gene (SLC20A4) to human chromosome 17p13.3. Cytogenet. Cell Genet. 83: 256–257, 1998.

- [73956] 12286.Abbott, G. W.; Sesti, F.; Splawski, I.; Buck, M. E.; Lehmann, M.H.; Timothy, K. W.; Keating, M. T.; Goldstein, S. A. N.: MiRP1 forms I(Kr) potassium channels with HERG and is associated with cardiac arrhythmia. *Cell* 97: 175–187, 1999.
- [73957] 12287.Schroeder, B. C.; Waldegger, S.; Fehr, S.; Bleich, M.; Warth, R.; Greger, R.; Jentsch, T. J.: A constitutively open potassium channel formed by KCNQ1 and KCNE3. *Nature* 403: 196–199, 2000.
- [73958] 12288.Harvey, T. J.; Hooper, J. D.; Myers, S. A.; Stephenson, S.-A.; Ashworth, L. K.; Clements, J. A.: Tissue-specific expression patterns and fine mapping of the human kallikrein (KLK) locus on proximal 19q13.4. *J.Biol. Chem.* 275: 37397–37406, 2000.
- [73959] 12289.Chaudhry, F. A.; Reimer, R. J.; Krizaj, D.; Barber, D.; Storm-Mathisen, J.; Copenhagen, D. R.; Edwards, R. H.: Molecular analysis of system N suggests novel physiological roles in nitrogen metabolism and synaptic transmission. *Cell* 99: 769–780, 1999.
- [73960] 12290.Yamada, K.; Nishida, K.; Hibi, M.; Hirano, T.; Matsuda, Y.: Comparative FISH mapping of Gab1 and Gab2 genes in human, mouse and rat. *Cytogenet. Cell Genet.* 94: 39–42, 2001.

- [73961] 12291.Afonina, E.; Stauber, R.; Pavlakis, G. N.: The human poly(A)-bindingprotein 1 shuttles between the nucleus and the cytoplasm. *J. Biol.Chem.* 273: 13015–13021, 1998.
- [73962] 12292.Deo, R. C.; Bonanno, J. B.; Sonenberg, N.; Burley, S. K.: Recognitionof polyadenylate RNA by the poly(A)-binding protein. *Cell* 98: 835–845,1999.
- [73963] 12293.Gorlach, M.; Burd, C. G.; Dreyfuss, G.: The mRNA poly(A)-bindingprotein: localization, abundance, and RNA-binding specificity. *Exp.Cell Res.* 211: 400–407, 1994.
- [73964] 12294.Grange, T.; Martins de Sa, C.; Oddos, J.; Pictet, R.: Human mRNApolyadenylate binding protein: evolutionary conservation of a nucleicacid binding motif. *Nucleic Acids Res.* 15: 4771–4787, 1987.
- [73965] 12295.Danpure, C. J.; Purdue, P. E.; Fryer, P.; Griffiths, S.; Allsop,J.; Lumb, M. J.; Guttridge, K. M.; Jennings, P. R.; Scheinman, J.I.; Mauer, S. M.; Davidson, N. O.: Enzymological and mutational analysisof a complex primary hyperoxaluria type I phenotype involving alanine:glyoxylateaminotransferase peroxisome-to-mitochondrion mistargeting and intraperoxisomalaggregation. *Am. J. Hum. Genet.* 53: 417–432, 1993.

- [73966] 12296.Mori, M.; Oda, T.; Nishiyama, K.; Serikawa, T.; Yamada, J.; Ichiyama,A.: A single serine:pyruvate aminotransferase gene on rat chromosome9q34–q36. *Genomics* 13: 686–689, 1992.
- [73967] 12297.Purdue, P. E.; Allsop, J.; Isaya, G.; Rosenberg, L. E.; Danpure,C. J.: Mistargeting of peroxisomal L-alanine:glyoxylate aminotransferaseto mitochondria in primary hyperoxaluria patients depends upon activationof a cryptic mitochondrial targeting sequence by a point mutation. *Proc.Nat. Acad. Sci.* 88: 10900–10904, 1991.
- [73968] 12298.Purdue, P. E.; Lumb, M. J.; Allsop, J.; Minatogawa, Y.; Danpure,C. J.: A glycine–to–glutamate substitution abolishes alanine:glyoxylateaminotransferase catalytic activity in a subset of patients with primaryhyperoxaluria type 1. *Genomics* 13: 215–218, 1992.
- [73969] 12299.Purdue, P. E.; Takada, Y.; Danpure, C. J.: Identification ofmutations associated with peroxisome–to–mitochondrion mistargetingof alanine/glyoxylate aminotransferase in primary hyperoxaluria type1. *J. Cell Biol.* 111: 2341–2351, 1990.
- [73970] 12300.Takada, Y.; Kaneko, N.; Esumi, H.; Purdue, P. E.; Danpure, C.J.: Human peroxisomal L-alanine:glyoxylate aminotransferase: evolutionaryloss of a mitochondrial tar–

geting signal by point mutation of the initiation codon.

Biochem. J. 268: 517–520, 1990.

- [73971] 12301.von Schnakenburg, C.; Rumsby, G.: Primary hyperoxaluria type 1: a cluster of new mutations in exon 7 of the AGXT gene. J. Med. Genet. 34: 489–492, 1997.
- [73972] 12302.Kobayashi, H.; Hino, M.; Shimodahira, M.; Iwakura, T.; Ishihara, T.; Ikekubo, K.; Ogawa, Y.; Nakao, K.; Kura-hachi, H.: Missense mutation of TRPS1 in a family of tricho-rhino-phalangeal syndrome type III. Am. J. Med. Genet. 107: 26–29, 2002.
- [73973] 12303.Lipkin, S. M.; Moens, P. B.; Wang, V.; Lenzi, M.; Shanmugarajah, D.; Gilgeous, A.; Thomas, J.; Cheng, J.; Touchman, J. W.; Green, E.D.; Schwartzberg, P.; Collins, F. S.; Cohen, P. E.: Meiotic arrest and aneuploidy in MLH3-deficient mice. Nature Genet. 31: 385–390, 2002.
- [73974] 12304.Lipkin, S. M.; Wang, V.; Jacoby, R.; Banerjee-Basu, S.; Baxevanis, A. D.; Lynch, H. T.; Elliott, R. M.; Collins, F. S.: MLH3: a DNA mismatch repair gene associated with mammalian microsatellite instability. Nature Genet. 24: 27–35, 2000.
- [73975] 12305.Lipkin, S. M.; Wang, V.; Stoler, D. L.; Anderson, G. R.; Kirsch, I.; Hadley, D.; Lynch, H. T.; Collins, F. S.: Germline and somatic mutation analyses in the DNA mis-



match repair gene MLH3: evidence for somatic mutation in colorectal cancers. *Hum. Mutat.* 17: 389–396, 2001.

[73976] 12306. Wu, Y.; Berends, M. J. W.; Sijmons, R. H.; Mensink, R. G. J.; Verlinde, E.; Kooi, K. A.; van der Sluis, T.; Kempinga, C.; van der Zee, A. G. J.; Hollema, H.; Buys, C. H. C. M.; Kleibeuker, J. H.; Hofstra, R. M. W.: A role for MLH3 in hereditary nonpolyposis colorectal cancer. *Nature Genet.* 29: 137–138, 2001.

[73977] 12307. Sahara, S.; Aoto, M.; Eguchi, Y.; Imamoto, N.; Yoneda, Y.; Tsujimoto, Y.: Acinus is a caspase-3-activated protein required for apoptotic chromatin condensation. *Nature* 401: 168–173, 1999.

[73978] 12308. Hata, M.; Ohtsuka, K.: Characterization of HSE sequences in human Hsp40 gene: structural and promoter analysis. *Biochim. Biophys. Acta* 1397: 43–55, 1998.

[73979] 12309. Hata, M.; Okumura, K.; Seto, M.; Ohtsuka, K.: Genomic cloning of a human heat shock protein 40 (Hsp40) gene (HSPF1) and its chromosomal localization to 19p13.2. *Genomics* 38: 446–449, 1996.

[73980] 12310. Ohtsuka, K.: Cloning of a cDNA for heat-shock protein hsp40, a human homologue of bacterial DnaJ. *Biochem. Biophys. Res. Commun.* 197: 235–240, 1993.

[73981] 12311. Janitz, K.; Wild, A.; Beck, S.; Savasta, S.; Beluffi, G.;

Ziegler,A.; Volz, A.: Genomic organization of the HSET locus and the possible association of HLA-linked genes with immotile cilia syndrome (ICS). Immunogenetics 49:644–652, 1999.

[73982] 12312.Muravenko, O. V.; Gizatullin, R. Z.; Protopopov, A. I.; Kashuba,V. I.; Zabarovsky, E. R.; Zelenin, A. V.: Assignment of CDK5R2 coding for the cyclin-dependent kinase 5, regulatory subunit 2 (NCK5A1 protein) to human chromosome band 2q35 by fluorescent in situ hybridization. Cytogenet.Cell Genet. 89: 160–161, 2000.

[73983] 12313.Nilden, F.; Backstrom, A.; Bark, C.: Molecular cloning and characterisation of a mouse gene encoding an isoform of the neuronal cyclin-dependent kinase 5 (CDK5) activator. Biochim. Biophys. Acta 1398: 371–376,1998.

[73984] 12314.Tang, D.; Yeung, J.; Lee, K-Y.; Matsushita, M.; Matsui, H.; Tomizawa,K.; Hatase, O.; Wang, J. H.: An isoform of the neuronal cyclin-dependent kinase 5 (Cdk5) activator. J. Biol. Chem. 270: 26897–26903, 1995.

[73985] 12315.DuPont, B. R.; Hu, C.-C.; Reveles, X.; Simmer, J. P.: Assignment of serine protease 17 (PRSS17) to human chromosome bands 19q13.3–q13.4 by in situ hybridization. Cytogenet. Cell Genet. 86: 212–213, 1999.

[73986] 12316.Nelson, P. S.; Gan, L.; Ferguson, C.; Moss, P.; Geli-

nas, R.; Hood, L.; Wang, K.: Molecular cloning and characterization of prostase, an androgen-regulated serine protease with prostate-restricted expression. *Proc. Nat. Acad. Sci.* 96: 3114–3119, 1999.

[73987] 12317. Hallas, C.; Pekarsky, Y.; Itoyama, T.; Varnum, J.; Bichi, R.; Rothstein, J. L.; Croce, C. M.: Genomic analysis of human and mouse TCL1 loci reveals a complex of tightly clustered genes. *Proc. Nat. Acad. Sci.* 96: 14418–14423, 1999.

[73988] 12318. Pekarsky, Y.; Hallas, C.; Isobe, M.; Russo, G.; Croce, C. M.: Abnormalities at 14q32.1 in T cell malignancies involve two oncogenes. *Proc. Nat. Acad. Sci.* 96: 2949–2951, 1999.

[73989] 12319. Koga, Y.; Fabrizi, G. M.; Mita, S.; Arnaudo, E.; Lomax, M. I.; Aqua, M. S.; Grossman, L. I.; Schon, E. A.: Sequence of a cDNA specifying subunit VIIc of human cytochrome c oxidase. *Nucleic Acids Res.* 18: 684 only, 1990.

[73990] 12320. Chaudhary, P. M.; Ferguson, C.; Nguyen, V.; Nguyen, O.; Massa, H. F.; Eby, M.; Jasmin, A.; Trask, B. J.; Hood, L.; Nelson, P. S.: Cloning and characterization of two Toll/interleukin-1 receptor-like genes TIL3 and TIL4: evidence for a multi-gene receptor family in humans.

Blood 91: 4020–4027, 1998.

- [73991] 12321.Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.; Grandy, D.;Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an 11q22.3–9p22translocation with affective disorder: proximity of the dopamine D2receptor gene relative to the translocation breakpoint. (Abstract) Am.J. Hum. Genet. 45 (suppl.): A220 only, 1989.
- [73992] 12322.Abuladze, N.; Lee, I.; Newman, D.; Hwang, J.; Boorer, K.; Pushkin,A.; Kurtz, I.: Molecular cloning, chromosomal localization, tissue distribution, and functional expression of the human pancreatic sodiumbicarbonate cotransporter. J. Biol. Chem. 273: 17689–17695, 1998.
- [73993] 12323.Burnham, C. E.; Amlal, H.; Wang, Z.; Shull, G. E.; Soleimani, M.: Cloning and functional expression of a human kidney  $\text{Na}^+:\text{HCO}_3^-$  cotransporter. J.Biol. Chem. 272: 19111–19114, 1997.
- [73994] 12324.Choi, I.; Romero, M. F.; Khandoudi, N.; Bril, A.; Boron, W. F.: Cloning and characterization of a human electrogenic  $\text{Na}(+)\text{--HCO}(3-)$  cotransporter isoform (hhNBC). Am. J. Physiol. 276: C576–C584, 1999.
- [73995] 12325.Igarashi, T.; Inatomi, J.; Sekine, T.; Cha, S. H.; Kanai, Y.; Kunimi,M.; Tsukamoto, K.; Satoh, H.; Shimadzu, M.; Tozawa, F.; Mori, T.;Shiobara, M.; Seki, G.; Endou, H.:

Mutations in SLC4A4 cause permanent isolated proximal renal tubular acidosis with ocular abnormalities. (Letter) Nature Genet. 23: 264–265, 1999.

- [73996] 12326. Romero, M. F.; Boron, W. F.: Electrogenic Na(+)/HCO(3-) cotransporters: cloning and physiology. Annu. Rev. Physiol. 61: 699–723, 1999.
- [73997] 12327. Soleimani, M.; Burnham, C. E.: Physiologic and molecular aspects of the Na(+):HCO(3-) cotransporter in health and disease processes. Kidney Int. 57: 371–384, 2000.
- [73998] 12328. Usui, T.; et al.; et al.: Pflugers Arch. 438: 458–462, 1999.
- [73999] 12329. Zhou, Y.-D.; Barnard, M.; Tian, H.; Li, X.; Ring, H. Z.; Francke, U.; Shelton, J.; Richardson, J.; Russell, D. W.; McKnight, S. L.: Molecular characterization of two mammalian bHLH-PAS domain proteins selectively expressed in the central nervous system. Proc. Nat. Acad. Sci. 94: 713–718, 1997.
- [74000] 12330. Garcia, J. A.; Zhang, D.; Estill, S. J.; Michnoff, C.; Rutter, J.; Reick, M.; Scott, K.; Diaz-Arrastia, R.; McKnight, S. L.: Impaired cued and contextual memory in NPAS2-deficient mice. Science 288: 2226–2230, 2000.
- [74001] 12331. Spencer, E.; Jiang, J.; Chen, Z. J.: Signal-induced

ubiquitination of I-kappa-B-alpha by the F-box protein Slimb/beta-TrCP. *Genes Dev.* 13:284-294, 1999.

- [74002] 12332. Yaron, A.; Hatzubai, A.; Davis, M.; Lavon, I.; Amit, S.; Manning, A. M.; Andersen, J. S.; Mann, M.; Mercurio, F.; Ben-Neriah, Y.: Identification of the receptor component of the I-kappa-B-alpha-ubiquitin ligase. *Nature* 396:590-594, 1998.
- [74003] 12333. Lucas, C. A.; Rughani, A.; Hoh, J. F. Y.: Expression of extraocular myosin heavy chain in rabbit laryngeal muscle. *J. Muscle Res. Cell Motil.* 16: 368-378, 1995.
- [74004] 12334. Wieczorek, D. F.; Periasamy, M.; Butler-Browne, G. S.; Whalen, R. G.; Nadal-Ginard, B.: Co-expression of multiple myosin heavy chain genes, in addition to a tissue-specific one, in extraocular musculature. *J. Cell Biol.* 101: 618-629, 1985.
- [74005] 12335. Winters, L. M.; Briggs, M. M.; Schachat, F.: The human extraocular muscle myosin heavy chain gene (MYH13) maps to the cluster of fast and developmental myosin genes on chromosome 17. *Genomics* 54: 188-189, 1998.
- [74006] 12336. Briskin, C.; Heineman, A.; Chavarria, T.; Elenbaas, B.; Tan, J.; Dey, S. K.; McMahon, J. A.; McMahon, A. P.; Weinberg, R. A.: Essential function of Wnt-4 in mammary gland development downstream of progesterone signaling.

Genes Dev. 14: 650–654, 2000.

[74007] 12337.Jordan, B. K.; Mohammed, M.; Ching, S. T.; Delot, E.; Chen, X.-N.;Dewing, P.; Swain, A.; Rao, P. N.; Elejalde, B. R.; Vilain, E.: Up-regulationof WNT-4 signaling and dosage-sensitive sex reversal in humans. Am.J. Hum. Genet. 68: 1102–1109, 2001.

[74008] 12338.Stark, K.; Vainio, S.; Vassileva, G.; McMahon, A. P.: Epithelialtransformation of metanephric mesenchyme in the developing kidneyregulated by Wnt-4. Nature 372: 679–683, 1994.

[74009] 12339.Vainio, S.; Heikkila, M.; Kispert, A.; Chin, N.; McMahon, A. P.: Female development in mammals is regulated by Wnt-4 signalling. Nature 397:405–409, 1999.

[74010] 12340.Vilain, A.; Apiou, F.; Dutrillaux, B.; Malfoy, B.: Assignmentof candidate DNA methyltransferase gene (DNMT2) to human chromosomeband 10p15.1 by in situ hybridization. Cytogenet. Cell Genet. 82:120 only, 1998.

[74011] 12341.Yoder, J. A.; Bestor, T. H.: A candidate mammalian DNA methyltransferaserelated to pmt1p of fission yeast. Hum. Molec. Genet. 7: 279–284,1998.

[74012] 12342.Avraham, K. B.; Cho, B. C.; Gilbert, D.; Fujii, H.; Okamoto, K.;Shimazaki, T.; Ito, T.; Shoji, H.; Wakamatsu, Y.; Kondoh, H.; Takahashi,N.; Muramatsu, M.; Hamada, H.;

Copeland, N. G.; Jenkins, N. A.: Murine chromosomal location of four class III POU transcription factors. *Genomics* 18:131–133, 1993.

[74013] 12343. Egan, S.; Herbrick, J.-A.; Tsui, L.-C.; Cohen, B.; Flock, G.; Beatty, B.; Scherer, S. W.: Mapping of the human lunatic fringe (LFNG) gene to 7p22 and manic fringe (MFNG) to 22q12. *Genomics* 54: 576–577, 1998.

[74014] 12344. Johnston, S. H.; Rauskolb, C.; Wilson, R.; Prabhakaran, B.; Irvine, K. D.; Vogt, T. F.: A family of mammalian Fringe genes implicated in boundary determination and the Notch pathway. *Development* 124:2245–2254, 1997.

[74015] 12345. Moran, J. L.; Johnston, S. H.; Rauskolb, C.; Bhalerao, J.; Bowcock, A. M.; Vogt, T. F.: Genomic structure, mapping, and expression analysis of the mammalian lunatic, manic, and radical fringe genes. *Mammalian Genome* 10: 535–541, 1999.

[74016] 12346. Van Tine, B. A.; Knops, J.; Shaw, G. M.; May, W. A.: Assignment of human MFNG, manic fringe *Drosophila* homolog, to 22q13.1 using tyramide fluorescence in situ hybridization (T-FISH). *Cytogenet. Cell Genet.* 87:132–133, 1999.

[74017] 12347. Laufer, E.; Dahn, R.; Orozco, O. E.; Yeo, C.-Y.;



Pisenti, J.; Henrique,D.; Abbott, U. K.; Fallon, J. F.; Tabin, C.: Expression of Radicalfringe in limb–bud ectoderm regulates apical ectodermal ridge formation. *Nature* 386:366–373, 1997.

[74018] 12348.Rodriguez–Esteban, C.; Schwabe, J. W. R.; De La Pena, J.; Foy,B.; Eshelman, B.; Izpisua Belmonte, J. C.: Radical fringe positionsthe apical ectodermal ridge at the dorsoventral boundary of the vertebratelimb. *Nature* 386: 360–366, 1997.

[74019] 12349.Hofmann, Y.; Becker, J.; Wright, F.; Avner, E. D.; Mrug, M.; Guay–Woodford,L. M.; Somlo, S.; Zerres, K.; Germino, G. G.; Onuchic, L. F.: Genomicstructure of the gene for the human P1 protein (MCM3) and its exclusionas a candidate for autosomal recessive polycystic kidney disease. *Europ.J. Hum. Genet.* 8: 163–166, 2000.

[74020] 12350.Starborg, M.; Brundell, E.; Gell, K.; Larsson, C.; White, I.; Daneholt,B.; Hoog, C.: A murine replication protein accumulates temporarilyin the heterochromatic regions of nuclei prior to initiation of DNAreplication. *J. Cell Sci.* 108: 927–934, 1995.

[74021] 12351.Budde, S. M. S.; van den Heuvel, L. P. W. J.; Janssen, A. J.; Smeets,R. J. P.; Buskens, C. A. F.; DeMeirleir, L.; Van Coster, R.; Baethmann,M.; Voit, T.; Trijbels, J. M. F.;

Smeitink, J. A. M.: Combined enzymatic complex I and III deficiency associated with mutations in the nuclear-encoded NDUF54 gene. *Biochem. Biophys. Res. Commun.* 275: 63–68, 2000.

[74022] 12352. Papa, S.; Sardanelli, A. M.; Cocco, T.; Speranza, F.; Scacco, S.C.; Technikova-Dobrova, Z.: The nuclear encoded-18 kDa (IP) AQDQ subunit of bovine heart complex I is phosphorylated by the mitochondrial cAMP-dependent protein kinase. *FEBS Lett.* 379: 299–301, 1996.

[74023] 12353. Petruzzella, V.; Vergari, R.; Puzziferri, I.; Boffoli, D.; Lamantea, E.; Zeviani, M.; Papa, S.: A nonsense mutation in the NDUF54 gene encoding the 18 kDa (AQDQ) subunit of complex I abolishes assembly and activity of the complex in a patient with Leigh-like syndrome. *Hum. Molec. Genet.* 10: 529–535, 2001.

[74024] 12354. van den Heuvel, L.; Ruitenbeek, W.; Smeets, R.; Gelman-Kohan, Z.; Elpeleg, O.; Loeffen, J.; Trijbels, F.; Mariman, E.; de Bruijn, D.; Smeitink, J.: Demonstration of a new pathogenic mutation in human complex I deficiency: a 5-bp duplication in the nuclear gene encoding the 18-kD (AQDQ) subunit. *Am. J. Hum. Genet.* 62: 262–268, 1998.

[74025] 12355. Walker, J. E.; Arizmendi, J. M.; Dupuis, A.; Fearnley, I. M.; Finel, M.; Medd, S. M.; Pilkington, S. J.; Runswick, M.

J.; Skehel, J. M.: Sequences of 20 subunits of NADH:ubiquinone oxidoreductase from bovine heart mitochondria: application of a novel strategy for sequencing proteins using the polymerase chain reaction. *J. Molec. Biol.* 226:1051–1072, 1992.

- [74026] 12356. Chicheportiche, Y.; Bourdon, P. R.; Xu, H.; Hsu, Y.-M.; Scott, H.; Hession, C.; Garcia, I.; Browning, J. L.: TWEAK, a new secreted ligand in the tumor necrosis factor family that weakly induces apoptosis. *J. Biol. Chem.* 272: 32401–32410, 1997.
- [74027] 12357. Kaptein, A.; Jansen, M.; Dilaver, G.; Kitson, J.; Dash, L.; Wang, E.; Owen, M. J.; Bodmer, J.-L.; Tschopp, J.; Farrow, S. N.: Studies on the interaction between TWEAK and the death receptor WSL-1/TRAMP(DR3). *FEBS Lett.* 485: 135–141, 2000.
- [74028] 12358. Marsters, S. A.; Sheridan, J. P.; Pitti, R. M.; Brush, J.; Goddard, A.; Ashkenazi, A.: Identification of a ligand for the death-domain-containing receptor Apo3. *Curr. Biol.* 8: 525–528, 1998.
- [74029] 12359. Schneider, P.; Schwenzer, R.; Haas, E.; Muhlenbeck, F.; Schubert, G.; Scheurich, P.; Tschopp, J.; Wajant, H.: TWEAK can induce cell death via endogenous TNF and TNF receptor 1. *Europ. J. Immun.* 29:1785–1792, 1999.

- [74030] 12360.Wiley, S. R.; Cassiano, L.; Lofton, T.; Davis-Smith, T.; Winkles,J. A.; Lindner, V.; Liu, H.; Daniel, T. O.; Smith, C. A.; Fanslow,W. C.: A novel TNF receptor family member binds TWEAK and is implicatedin angiogenesis. *Immunity* 15: 837–846, 2001.
- [74031] 12361.Hoey, T.; Sun, Y.–L.; Williamson, K.; Xu, X.: Isolation of twonew members of the NF-AT gene family and functional characterizationof the NF-AT proteins. *Immunity* 2: 461–472, 1995.
- [74032] 12362.Hoey, T.; Sun, Y. L.; Williamson, K.; Xu, X.: Isolation of twonew members of the NF-AT gene family and functional characterizationof the NF-AT proteins. *Immunity* 2: 461–472, 1995.
- [74033] 12363.Arany, Z.; Newsome, D.; Oldread, E.; Livingston, D. M.; Eckner,R.: A family of transcriptional adaptor proteins targeted by theE1A oncoprotein. *Nature* 374: 81–84, 1995.
- [74034] 12364.de Lecea, L.; Ruiz-Lozano, P.; Danielson, P. E.; Peelle-Kirley,J.; Foye, P. E.; Frankel, W. N.; Sutcliffe, J. G.: Cloning, mRNAexpression, and chromosomal mapping of mouse and human preprocortistatin. *Genomics* 42:499–506, 1997.
- [74035] 12365.Ejeskar, K.; Abel, F.; Sjoberg, R.–M.; Backstrom, J.;

Kogner, P.;Martinsson, T.: Fine mapping of the human preprocortistatin gene(CORT) to neuroblastoma consensus deletion region 1p36.3–p36.2, but absence of mutations in primary tumors. Cytogenet. Cell Genet. 89:62–66, 2000.

[74036] 12366.Fukusumi, S.; Kitada, C.; Takekawa, S.; Kizawa, H.; Sakamoto, J.;Miyamoto, M.; Hinuma, S.; Kitano, K.; Fujino, M.: Identification and characterization of a novel human cortistatin–like peptide. Biochem.Biophys. Res. Commun. 232: 157–163, 1997.

[74037] 12367.Dong, C.; Zhu, S.; Wang, T.; Yoon, W.; Li, Z.; Alvarez, R. J.;ten Dijke, P.; White, B.; Wigley, F. M.; Goldschmidt–Clermont, P.J.: Deficient Smad7 expression: a putative molecular defect in scleroderma. Proc.Nat. Acad. Sci. 99: 3908–3913, 2002.

[74038] 12368.Kavsak, P.; Rasmussen, R. K.; Causing, C. G.; Bonni, S.; Zhu, H.;Thomsen, G. H.; Wrana, J. L.: Smad7 binds to Smurf2 to form an E3ubiquitin ligase that targets the TGF–beta receptor for degradation. Molec.Cell 6: 1365–1375, 2000.

[74039] 12369.Lallemand, F.; Mazars, A.; Prunier, C.; Bertrand, F.; Kornprost,M.; Gallea, S.; Roman–Roman, S.; Cherqui, G.; Atfi, A.: Smad7 inhibits the survival nuclear factor kappa–B and potentiates apoptosis in epithelial cells. Oncogene 20:

879–884, 2001.

- [74040] 12370.Roijer, E.; Moren, A.; ten Dijke, P.; Stenman, G.: Assignment of the Smad7 gene (MADH7) to human chromosome 18q21.1 by fluorescence in situ hybridization. *Cytogenet. Cell Genet.* 81: 189–190, 1998.
- [74041] 12371.Cravatt, B. F.; Giang, D. K.; Mayfield, S. P.; Boger, D. L.; Lerner, R. A.; Gilula, N. B.: Molecular characterization of an enzyme that degrades neuromodulatory fatty-acid amides. *Nature* 384: 83–87, 1996.
- [74042] 12372.Giang, D. K.; Cravatt, B. F.: Molecular characterization of human and mouse fatty acid amide hydrolases. *Proc. Nat. Acad. Sci.* 94:2238–2242, 1997.
- [74043] 12373.Sipe, J. C.; Chiang, K.; Gerber, A. L.; Beutler, E.; Cravatt, B.F.: A missense mutation in human fatty acid amide hydrolase associated with problem drug use. *Proc. Nat. Acad. Sci.* 99: 8394–8399, 2002.
- [74044] 12374.Wan, M.; Cravatt, B. F.; Ring, H. Z.; Zhang, X.; Francke, U.: Conserved chromosomal location and genomic structure of human and mouse fatty-acid amide hydrolase genes and evaluation of clasper as a candidate neurological mutation. *Genomics* 54: 408–414, 1998.
- [74045] 12375.Falany, C. N.; Johnson, M. R.; Barnes, S.; Diasio, R. B.: Glycine and taurine conjugation of bile acids by a single

enzyme: molecularcloning and expression of a human liver bile acid CoA:amino acid N-acyltransferase. J.Biol. Chem. 269: 19375-19379, 1994.

[74046] 12376.Johnson, M. R.; Barnes, S.; Kwakye, J. B.; Diasio, R. B.: Purificationand characterization of bile acid-CoA:amino acid N-acyltransferasefrom human liver. J. Biol. Chem. 266: 10227-10233, 1991.

[74047] 12377.Baier, M.; Bannert, N.; Werner, A.; Lang, K.; Kurth, R.: Molecularcloning, sequence, expression, and processing of the interleukin 16precursor. Proc. Nat. Acad. Sci. 94: 5273-5277, 1997.

[74048] 12378.Roh, M. H.; Makarova, O.; Liu, C.-J.; Shin, K.; Lee, S.; Laurinec,S.; Goyal, M.; Wiggins, R.; Margolis, B.: The Maguk protein, Pals1,functions as an adapter, linking mammalian homologues of Crumbs andDiscs Lost. J. Cell Biol. 157: 161-172, 2002.

[74049] 12379.Bannert, N.; Baier, M.; Werner, A.; Kurth, R.: Interleukin-16or not? Nature 381: 30 only, 1996.

[74050] 12380.Bannert, N.; Kurth, R.; Baier, M.: The gene encoding mouse interleukin-16consists of seven exons and maps to Chromosome 7 D2-D3. Immunogenetics 49:704-706, 1999.

[74051] 12381.Cruikshank, W.; Center, D. M.: Modulation of lym-

phocyte migration by human lymphokines. II. Purification of a lymphotactic factor (LCF). *J. Immun.* 128: 2569–2574, 1982.

[74052] 12382. Cruikshank, W. W.; Center, D. M.; Nisar, N.; Wu, M.; Natke, B.; Theodore, A. C.; Kornfeld, H.: Molecular and functional analysis of a lymphocyte chemoattractant factor: association of biologic function with CD4 expression. *Proc. Nat. Acad. Sci.* 91: 5109–5113, 1994.

[74053] 12383. Drwinga, H. L.; Toji, L. H.; Kim, C. H.; Greene, A. E.; Mulivor, R. A.: NIGMS human/rodent somatic cell hybrid mapping panels 1 and 2. *Genomics* 16: 311–314, 1993.

[74054] 12384. Hudson, T. J.; Stein, L. D.; Gerety, S. S.; Ma, J.; Castle, A. B.; Silva, J.; Slonim, D. K.; Baptista, R.; Kruglyak, L.; Xu, S. H.; Hu, X.; Colbert, A. M. E.; and 39 others: An STS-based map of the human genome. *Science* 270: 1945–1954, 1995.

[74055] 12385. Keane, J.; Nicoll, J.; Kim, S.; Wu, D. M. H.; Cruikshank, W. W.; Brazer, W.; Natke, B.; Zhang, Y.; Center, D. M.; Kornfeld, H.: Conservation of structure and function between human and murine IL-16. *J. Immun.* 160: 5945–5954, 1998.

[74056] 12386. Scheuerpflug, C. G.; Lichter, P.; Debatin, K.-M.; Mincheva, A.: Assignment of TRADD to human chromo-



some band 16q22 by in situ hybridization. *Cytogenet. Cell Genet.* 92: 347–348, 2001.

[74057] 12387. Lennon–Dumenil, A.–M.; Barbouche, M.–R.; Védrenne, J.; Prod'Homme, T.; Bejaoui, M.; Ghariani, S.; Charon, D.; Fellous, M.; Dellagi, K.; Alcaide–Loridan, C.: Uncoordinated HLA–D gene expression in a RFXANK–defective patient with MHC class II deficiency. *J. Immun.* 166:5681–5687, 2001.

[74058] 12388. Lisowska–Groszpiere, B.; Fondaneche, M. C.; Rols, M. P.; Griscelli, C.; Fischer, A.: Two complementation groups account for most cases of inherited MHC class II deficiency. *Hum. Molec. Genet.* 3: 953–958, 1994.

[74059] 12389. Boll, W.; Wagner, P.; Mantei, N.: Structure of the chromosomal gene and cDNAs coding for lactase–phlorizin hydrolase in humans with adult–type hypolactasia or persistence of lactase. *Am. J. Hum. Genet.* 48:889–902, 1991.

[74060] 12390. Wiszniewski, W.; Fondaneche, M.–C.; Lambert, N.; Masternak, K.; Picard, C.; Notarangelo, L.; Schwartz, K.; Bal, J.; Reith, W.; Alcaide, C.; de Saint Basile, G.; Fischer, A.; Lisowska–Groszpiere, B.: Founder effect for a 26–bp deletion in the RFXANK gene in North African major histocompatibility complex class II–deficient patients belonging to complementation group B. *Immunogenetics* 51: 261–267,

2000.

- [74061] 12391. Harvey, C. B.; Fox, M. F.; Jeggo, P. A.; Mantei, N.; Povey, S.; Swallow, D. M.: Regional localization of the lactase-phlorizin hydrolase gene, LCT, to chromosome 2q21. *Ann. Hum. Genet.* 57: 179–185, 1993.
- [74062] 12392. Kruse, T. A.; Bolund, L.; Byskov, A.; Sjostrom, H.; Noren, O.; Mantei, N.; Semenza, G.: Mapping of the human lactase-phlorizin hydrolase gene to chromosome 2. (Abstract) *Cytogenet. Cell Genet.* 51: 1026, 1989.
- [74063] 12393. Kruse, T. A.; Bolund, L.; Grzeschik, K.-H.; Ropers, H. H.; Sjostrom, H.; Noren, O.; Mantei, N.; Semenza, G.: The human lactase-phlorizin hydrolase gene is located on chromosome 2. *FEBS Lett.* 240: 123–126, 1988.
- [74064] 12394. Mantei, N.; Villa, M.; Enzler, T.; Wacker, H.; Boll, W.; James, O.; Hunziker, W.; Semenza, G.: Complete primary structure of human and rabbit lactase-phlorizin hydrolase: implications for biosynthesis, membrane anchoring and evolution of the enzyme. *EMBO J.* 7: 2705–2713, 1988.
- [74065] 12395. Stec, I.; Wright, T. J.; van Ommen, G.-J. B.; de Boer, P. A. J.; van Haeringen, A.; Moorman, A. F. M.; Altherr, M. R.; den Dunnen, J. T.: WHSC1, a 90 kb SET domain-containing gene, expressed in early development and homologous to a *Drosophila* dysmorphia gene maps in the Wolf-

Hirschhorn syndrome critical region and is fused to IgH int(4;14) multiple myeloma. Hum. Molec. Genet. 7: 1071–1082, 1998.

[74066] 12396.Ame, J.–C.; Apiou, F.; Jacobson, E. L.; Jacobson, M. K.: Assignment of the poly(ADP–ribose) glycohydrolase gene (PARG) to human chromosome 10q11.23 and mouse chromosome 14B by in situ hybridization. Cytogenet. Cell Genet. 85: 269–270, 1999.

[74067] 12397.Lin, W.; Ame, J.–C.; Aboul–Ela, N.; Jacobson, E. L.; Jacobson, M. K.: Isolation and characterization of the cDNA encoding bovine poly(ADP–ribose) glycohydrolase. J. Biol. Chem. 272: 11895–11901, 1997.

[74068] 12398.Brown, S. D.; Twells, R. C. J.; Hey, P. J.; Cox, R. D.; Levy, E.R.; Soderman, A. R.; Metzker, M. L.; Caskey, C. T.; Todd, J. A.; Hess, J. F.: Isolation and characterization of LRP6, a novel member of the low density lipoprotein receptor gene family. Biochem. Biophys. Res. Commun. 248: 879–888, 1998.

[74069] 12399.Mao, B.; Wu, W.; Li, Y.; Hoppe, D.; Stannek, P.; Glinka, A.; Niehrs, C.: LDL–receptor–related protein 6 is a receptor for Dickkopf proteins. Nature 411:321–325, 2001.

[74070] 12400.Pinson, K. I.; Brennan, J.; Monkley, S.; Avery, B. J.;

Skarnes,W. C.: An LDL-receptor-related protein mediates Wnt signalling in mice. *Nature* 407: 535–538, 2000.

[74071] 12401.Tamai, K.; Semenov, M.; Kato, Y.; Spokony, R.; Liu, C.; Katsuyama,Y.; Hess, F.; Saint-Jeannet, J.-P.; He, X.: LDL-receptor-related proteins in Wnt signal transduction. *Nature* 407: 530–535, 2000.

[74072] 12402.Wehrli, M.; Dougan, S. T.; Caldwell, K.; O'Keefe, L.; Schwartz,S.; Vaizel-Ohayon, D.; Schejter, E.; Tomlinson, A.; DiNardo, S.:Arrow encodes an LDL-receptor-related protein essential for Wingless signalling. *Nature* 407: 527–530, 2000. Note: Erratum: *Nature* 410:847 only, 2001.

[74073] 12403.Vinkemeier, U.; Obermann, W.; Weber, K.; Furst, D. O.: The globular head domain of titin extends into the center of the sarcomeric M band:cDNA cloning, epitope mapping and immunoelectron microscopy of two titin-associated proteins. *J. Cell Sci.* 106: 319–330, 1993.

[74074] 12404.van der Ven, P. F. M.; Speel, E. J. M.; Albrechts, J. C. M.; Ramaekers,F. C. S.; Hopman, A. H. N.; Furst, D. O.: Assignment of the human gene for endosarcomeric cytoskeletal M-protein (MYOM2) to 8p23.3. *Genomics* 55:253–255, 1999.

[74075] 12405.Black, R. A.; Rauch, C. T.; Kozlosky, C. J.; Peschon,

J. J.; Slack, J. L.; Wolfson, M. F.; Castner, B. J.; Stocking, K. L.; Reddy, P.; Srinivasan, S.; Nelson, N.; Boiani, N.; Schooley, K. A.; Gerhart, M.; Davis, R.; Fitzner, J. N.; Johnson, R. S.; Paxton, R. J.; March, C. J.; Cerretti, D. P.: A metalloproteinase disintegrin that releases tumour-necrosis factor-alpha from cells. *Nature* 385: 729-733, 1997.

[74076] 12406. Hirohata, S.; Seldin, M. F.; Apte, S. S.: Chromosomal assignment of two ADAM genes, TACE (ADAM17) and MLTNB (ADAM19), to human chromosomes 2 and 5, respectively, and of MltNb to mouse chromosome 11. *Genomics* 54: 178-179, 1998.

[74077] 12407. Moss, M. L.; Jin, S.-L. C.; Milla, M. E.; Burkhart, W.; Carter, H. L.; Chen, W.-J.; Clay, W. C.; Didsbury, J. R.; Hasler, D.; Hoffman, C. R.; Kost, T. A.; Lambert, M. H.; and 13 others: Cloning of a disintegrin metalloproteinase that processes precursor tumour-necrosis factor-alpha. *Nature* 385: 733-736, 1997.

[74078] 12408. Patel, I. R.; Attur, M. G.; Patel, R. N.; Stuchin, S. A.; Abagyan, R. A.; Abramson, S. B.; Amin, A. R.: TNF-alpha convertase enzyme from human arthritis-affected cartilage: isolation of cDNA by differential display, expression of the active enzyme, and regulation of TNF-alpha. *J. Immun.* 160: 4570-4579, 1998.

- [74079] 12409. Peschon, J. J.; Slack, J. L.; Reddy, P.; Stocking, K. L.; Sunnarborg, S. W.; Lee, D. C.; Russell, W. E.; Castner, B. J.; Johnson, R. S.; Fitzner, J. N.; Boyce, R. W.; Nelson, N.; Kozlosky, C. J.; Wolfson, M. F.; Rauch, C. T.; Cerretti, D. P.; Paxton, R. J.; March, C. J.; Black, R. A.: An essential role for ectodomain shedding in mammalian development. *Science* 282: 1281–1284, 1998.
- [74080] 12410. Yamazaki, K.; Mizui, Y.; Sagane, K.; Tanaka, I.: Genetic mapping of mouse tumor necrosis factor- $\alpha$  converting enzyme (Tace) to chromosome 12. *Genomics* 49: 336–337, 1998.
- [74081] 12411. Inoue, D.; Reid, M.; Lum, L.; Kratzschmar, J.; Weskamp, G.; Myung, Y. M.; Baron, R.; Blobel, C. P.: Cloning and initial characterization of mouse meltrin beta and analysis of the expression of four metalloprotease-disintegrins in bone cells. *J. Biol. Chem.* 273: 4180–4187, 1998.
- [74082] 12412. Iyer, N. V.; Leung, S. W.; Semenza, G. L.: The human hypoxia-inducible factor 1- $\alpha$  gene: HIF1A structure and evolutionary conservation. *Genomics* 52: 159–165, 1998.
- [74083] 12413. Kline, D. D.; Peng, Y.-J.; Manalo, D. J.; Semenza, G. L.; Prabhakar, N. R.: Defective carotid body function and

impaired ventilatory response to chronic hypoxia in mice partially deficient for hypoxia-inducible factor 1- $\alpha$ .  
Proc. Nat. Acad. Sci. 99: 821–826, 2002.

- [74084] 12414. Lando, D.; Peet, D. J.; Whelan, D. A.; Gorman, J. J.; Whitelaw, M. L.: Asparagine hydroxylation of the HIF trans-activation domain: a hypoxic switch. Science 295: 858–861, 2002.
- [74085] 12415. Marti, H. H.; Katschinski, D. M.; Wagner, K. F.; Schaffer, L.; Stier, B.; Wenger, R. H.: Isoform-specific expression of hypoxia-inducible factor-1- $\alpha$  during the late stages of mouse spermiogenesis. Molec. Endocr. 16: 234–243, 2002.
- [74086] 12416. Maxwell, P. H.; Wiesener, M. S.; Chang, G.-W.; Clifford, S. C.; Vaux, E. C.; Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.; Maher, E. R.; Ratcliffe, P. J.: The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. Nature 399: 271–275, 1999.
- [74087] 12417. Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W. G., Jr.; Pavletich, N. P.: Structure of an HIF-1- $\alpha$ -pVHL complex: hydroxyproline recognition in signaling. Science 296: 1886–1889, 2002.
- [74088] 12418. Semenza, G. L.: HIF-1 and human disease: one

highly involved factor. *Genes Dev.* 14: 1983–1991, 2000.

- [74089] 12419. Semenza, G. L.; Rue, E. A.; Iyer, N. V.; Pang, M. G.; Kearns, W. G.: Assignment of the hypoxia-inducible factor 1- $\alpha$  gene to a region of conserved synteny on mouse chromosome 12 and human chromosome 14q. *Genomics* 34: 437–439, 1996.
- [74090] 12420. Sutter, C. H.; Laughner, E.; Semenza, G. L.: Hypoxia-inducible factor 1- $\alpha$  protein expression is controlled by oxygen-regulated ubiquitination that is disrupted by deletions and missense mutations. *Proc. Nat. Acad. Sci.* 97: 4748–4753, 2000.
- [74091] 12421. Wenger, R. H.; Rolfs, A.; Kvietikova, I.; Spielmann, P.; Zimmermann, D. R.; Gassmann, M.: The mouse gene for hypoxia-inducible factor-1- $\alpha$ —genomic organization, expression and characterization of an alternative first exon and 5-prime flanking sequence. *Europ. J. Biochem.* 246: 155–165, 1997.
- [74092] 12422. Wenger, R. H.; Rolfs, A.; Marti, H. H.; Guenet, J.-L.; Gassmann, M.: Nucleotide sequence, chromosomal assignment and mRNA expression of mouse hypoxia-inducible factor-1- $\alpha$ . *Biochem. Biophys. Res. Commun.* 223: 54–59, 1996.
- [74093] 12423. Wenger, R. H.; Rolfs, A.; Spielmann, P.; Zimmer-



mann, D. R.; Gassmann, M.: Mouse hypoxia-inducible factor-1-alpha is encoded by two different mRNA isoforms: expression from a tissue-specific and a housekeeping-type promoter. *Blood* 91: 3471-3480, 1998.

[74094] 12424. Tian, H.; Hammer, R. E.; Matsumoto, A. M.; Russell, D. W.; McKnight, S. L.: The hypoxia-responsive transcription factor EPAS1 is essential for catecholamine homeostasis and protection against heart failure during embryonic development. *Genes Dev.* 12: 3320-3324, 1998.

[74095] 12425. Tian, H.; McKnight, S. L.; Russell, D. W.: Endothelial PAS domain protein 1 (EPAS1), a transcription factor selectively expressed in endothelial cells. *Genes Dev.* 11: 72-82, 1997.

[74096] 12426. Hovanessian, A. G.; Laurent, A. G.; Chebath, J.; Galabru, J.; Robert, N.; Svab, J.: Identification of 69-kd and 100-kd forms of 2-5A synthetase in interferon-treated human cells by specific monoclonal antibodies. *EMBO J.* 6: 1273-1280, 1987.

[74097] 12427. Marie, I.; Galabru, J.; Svab, J.; Hovanessian, A. G.: Preparation and characterization of polyclonal antibodies specific for the 69 and 100 k-dalton forms of human 2-5A synthetase. *Biochem. Biophys. Res. Commun.* 160: 580-587, 1989.

- [74098] 12428. Marie, I.; Hovanessian, A. G.: The 69-kDa 2-5A synthetase is composed of two homologous and adjacent functional domains. *J. Biol. Chem.* 267:9933-9939, 1992.
- [74099] 12429. Kools, P.; Van Imschoot, G.; van Roy, F.: Characterization of three novel human cadherin genes (CDH7, CDH19, and CDH20) clustered on chromosome 18q22-q23 and with high homology to chicken cadherin-7. *Genomics* 68:283-295, 2000.
- [74100] 12430. Dantzig, A. H.; Hoskins, J.; Tabas, L. B.; Bright, S.; Shepard, R. L.; Jenkins, I. L.; Duckworth, D. C.; Sportsman, J. R.; Mackensen, D.; Rosteck, P. R., Jr.; Skatrud, P. L.: Association of intestinal peptide transport with a protein related to the cadherin superfamily. *Science* 264:430-433, 1994.
- [74101] 12431. Shibata, T.; Shimoyama, Y.; Gotoh, M.; Hirohashi, S.: Identification of human cadherin-14, a novel neurally specific type II cadherin, by protein interaction cloning. *J. Biol. Chem.* 272: 5236-5240, 1997.
- [74102] 12432. Barabino, S. M. L.; Hubner, W.; Jenny, A.; Minvielle-Sebastia, L.; Keller, W.: The 30 kDa subunit of mammalian cleavage and polyadenylation specificity factor and its yeast homolog are RNA binding zinc finger proteins. *Genes Dev.* 11: 1703-1716, 1997.

- [74103] 12433.Nemeroff, M. E.; Barabino, S. M. L.; Li, Y.; Keller, W.; Krug,R. M.: Influenza virus NS1 protein interacts with the cellular 30kDa subunit of CPSF and inhibits 3-prime end formation of cellularpre-mRNAs. *Molec. Cell* 1: 991-1000, 1998.
- [74104] 12434.Hsu, D. R.; Economides, A. N.; Wang, X.; Eimon, P. M.; Harland,R. M.: The *Xenopus* dorsalizing factor gremlin identifies a novelfamily of secreted proteins that antago-nize BMP activities. *Molec.Cell* 1: 673-683, 1998.
- [74105] 12435.Topol, L. Z.; Modi, W. S.; Koochekpour, S.; Blair, D. G.: DRM-Gremlin(CKTSF1B1) maps to human chromosome 15 and is highly expressed inadult and fetal brain. *Cyto-genet. Cell Genet.* 89: 79-84, 2000.
- [74106] 12436.Beaconsfield, P.; Rainsbury, R.; Kalton, G.: Glucose-6-phosphatedehydrogenase deficiency and the incidence of cancer. *Oncologia* 19:11-19, 1965.
- [74107] 12437.Ben-Bassat, J.; Ben-Ishay, D.: Hereditary hemolytic anemia associatedwith glucose-6-phosphate dehydroge-nase deficiency (Mediterranean type). *IsraelJ. Med. Sci.* 5: 1053-1059, 1969.
- [74108] 12438.Benabadji, M.; Merad, F.; Benmoussa, M.; Tra-buchet, G.; Junien,C.; Dreyfus, J. C.; Kaplan, J. C.: Hetero-geneity of glucose-6-phosphatedehydrogenase deficiency

in Algeria. Hum. Genet. 40: 177–184, 1978.

- [74109] 12439. Benohr, H. C.; Klumpp, F.; Waller, H. D.: Glucose-6-phosphat-Dehydrogenase Typ Schwaben. Dtsch. Med. Wschr. 96: 1029–1032, 1971.
- [74110] 12440. Benohr, H. C.; Waller, H. D.: Eigenschaften der Glucose-6-p-dehydrogenase, Typ Tübingen. Klin. Wschr. 48: 71–74, 1970.
- [74111] 12441. Benohr, H. C.; Waller, H. D.; Arnold, H.; Blume, K. G.; Lohr, G. W.: Glucose-6-P-Dehydrogenase Typ Bordensee (eine neue Enzymvariante). Klin. Wschr. 49: 1058–1062, 1971.
- [74112] 12442. Beutler, E.: Glucose 6-phosphate dehydrogenase deficiency, a new Indian variant, G6PD Jammu. In: Sen, N. N.; Basu, A. K.: Trends in Haematology. Calcutta: Chatterjea Memorial Committee (pub.) 1975. Pp. 279–283.
- [74113] 12443. Beutler, E.: Personal Communication. La Jolla, Calif. 11/12/1990.
- [74114] 12444. Beutler, E.: G6PD deficiency. Blood 84: 3613–3636, 1994.
- [74115] 12445. Beutler, E.: Glucose-6-phosphate dehydrogenase deficiency. New Eng. J. Med. 324: 169–174, 1991.
- [74116] 12446. Beutler, E.: Selectivity of proteases as a basis for tissue distribution of enzymes in hereditary deficiencies.

Proc. Nat. Acad. Sci. 80:3767–3768, 1983.

- [74117] 12447. Beutler, E.: Glucose-6-phosphate dehydrogenase deficiency. In: Wintrobe, M. M.: Red Cell Metabolism in Hemolytic Anemia. New York: Plenum Press (pub.) 1978.
- [74118] 12448. Beutler, E.: The hemolytic effect of primaquine and related compounds: a review. Blood 14: 103–139, 1959.
- [74119] 12449. Beutler, E.; Grooms, A. M.; Morgan, S. K.; Trinidad, F.: Chronic severe hemolytic anemia due to G-6-PD Charleston: a new deficiency variant. J. Pediatr. 80: 1005–1009, 1972.
- [74120] 12450. Beutler, E.; Hartman, K.; Gelbart, T.; Forman, L.: G-6-PD Walter Reed: possible insight into 'structural' NADP in G-6-PD. Am. J. Hemat. 23: 25–30, 1986.
- [74121] 12451. Beutler, E.; Keller, J. W.; Matsumoto, F.: A new glucose 6-phosphate dehydrogenase (G6PD) variant associated with nonspherocytic hemolytic anemia: G6PD Atlanta. I.R.C.S. 4: 479, 1976.
- [74122] 12452. Beutler, E.; Kuhl, W.: The NT 1311 polymorphism of G6PD: G6PD Mediterranean mutation may have originated independently in Europe and Asia. Am. J. Hum. Genet. 47: 1008–1012, 1990.
- [74123] 12453. Beutler, E.; Kuhl, W.: Linkage between a PvuII restriction fragment length polymorphism and G6PD A-

(202A/376G): evidence for a single origin of the common G6PD A- mutation. Hum. Genet. 85: 9–11, 1990.

- [74124] 12454. Beutler, E.; Kuhl, W.; Gelbart, T.; Forman, L.: DNA sequence abnormalities of human glucose-6-phosphate dehydrogenase variants. J. Biol. Chem. 266: 4145–4150, 1991.
- [74125] 12455. Beutler, E.; Kuhl, W.; Ramirez, E.; Lisker, R.: Some Mexican glucose-6-phosphate dehydrogenase variants revisited. Hum. Genet. 86: 371–374, 1991.
- [74126] 12456. Beutler, E.; Kuhl, W.; Saenz, G. F.; Rodriguez, W.: Mutation analysis of glucose-6-phosphate dehydrogenase (G6PD) variants in Costa Rica. Hum. Genet. 87: 462–464, 1991.
- [74127] 12457. Beutler, E.; Mathai, C. K.; Smith, J. E.: Biochemical variants of glucose-6-phosphate dehydrogenase giving rise to congenital nonspherocytic hemolytic disease. Blood 31: 131–150, 1968.
- [74128] 12458. Beutler, E.; Matsumoto, F.: A new glucose 6-phosphate dehydrogenase variant: G6PD (-) Los Angeles. I.R.C.S. 5: 89, 1977.
- [74129] 12459. Beutler, E.; Matsumoto, F.; Daiber, A.: Nonspherocytic hemolytic anemia due to G-6-PD Panama. I.R.C.S. 2: 1389, 1974.

- [74130] 12460.Beutler, E.; Rosen, R.: Nonspherocytic congenital hemolytic anemia due to a new G-6-PD variant: G-6-PD Alhambra. *Pediatrics* 45: 230-235, 1970.
- [74131] 12461.Beutler, E.; Westwood, B.; Prchal, J. T.; Vaca, G.; Bartsocas, C. S.; Baronciani, L.: New glucose-6-phosphate dehydrogenase mutations from various ethnic groups. *Blood* 80: 255-256, 1992.
- [74132] 12462.Beutler, E.; Yoshida, A.: Human glucose-6-phosphate dehydrogenase variants: a supplementary tabulation. *Ann. Hum. Genet.* 37: 151-156, 1973.
- [74133] 12463.Boivin, P.; Galand, C.: Nouvelles variantes de la glucose-6-phosphate dehydrogenase erythrocytaire. *Rev. Franc. Etud. Clin. Biol.* 13:30-39, 1968.
- [74134] 12464.Botha, M. C.; Dern, R. J.; Mitchell, M.; West, C.; Beutler, E.: G6PD Capetown, a variant of glucose-6-phosphate dehydrogenase. *Am.J. Hum. Genet.* 21: 547-551, 1969.
- [74135] 12465.Boyer, S. H.; Graham, J. B.: Linkage between the X chromosomal loci for glucose-6-phosphate dehydrogenase electrophoretic variation and hemophilia A. *Am. J. Hum. Genet.* 17: 320-324, 1965.
- [74136] 12466.Boyer, S. H.; Porter, I. H.; Weilbaecher, R. G.: Electrophoretic heterogeneity of glucose-6-phosphate dehy-

drogenase and its relationship to enzyme deficiency in man. Proc. Nat. Acad. Sci. 48: 1868–1876, 1962.

- [74137] 12467. Busch, D.; Bote, K.: Glucose-6-phosphate-dehydrogenase-Defect in Deutschland. II. Eigenschaften des Enzyms (Typ Freiburg). Klin. Wschr. 48: 74–78, 1970.
- [74138] 12468. Calabro, V.; Giacobbe, A.; Vallone, D.; Montanaro, V.; Cascone, A.; Filosa, S.; Battistuzzi, G.: Genetic heterogeneity at the glucose-6-phosphate dehydrogenase locus in southern Italy: a study on a population from the Matera district. Hum. Genet. 86: 49–53, 1990.
- [74139] 12469. Cappadoro, M.; Giribaldi, G.; O'Brien, E.; Turrini, F.; Mannu, F.; Ulliers, D.; Simula, G.; Luzzatto, L.; Arese, P.: Early phagocytosis of glucose-6-phosphate dehydrogenase (G6PD)-deficient erythrocytes parasitized by Plasmodium falciparum may explain malaria protection in G6PD deficiency. Blood 92: 2527–2534, 1998.
- [74140] 12470. Cappellini, M. D.; Sampietro, M.; Toniolo, D.; Carandina, G.; Pittalis, S.; Martinez di Montemuros, F.; Tavazzi, D.; Fiorelli, G.: Biochemical and molecular characterization of a new sporadic glucose-6-phosphate dehydrogenase variant described in Italy: G6PD Modena. Brit. J. Haemat. 87: 209–211, 1994.



- [74141] 12471.Carandina, G.; Moretto, E.; Zecchi, G.; Conighi, C.: Glucose6-phosphate dehydrogenase Ferrara. A new variant of G6PD identified in Northern Italy. *Acta Haemat.* 56: 116-122, 1976.
- [74142] 12472.Carson, P. E.; Flanagan, C. L.; Ickes, C. E.; Alving, A. S.: Enzymatic deficiency in primaquine-sensitive erythrocytes. *Science* 124:484-485, 1956.
- [74143] 12473.Poy, F.; Yaffe, M. B.; Sayos, J.; Saxena, K.; Morra, M.; Sumegi, J.; Cantley, L. C.; Terhorst, C.; Eck, M. J.: Crystal structures of the XLP protein SAP reveal a class of SH2 domains with extended, phosphotyrosine-independent sequence recognition. *Molec. Cell* 4:555-561, 1999.
- [74144] 12474.Provisor, A. J.; Iaccone, J. J.; Chilcote, R. R.; Neiburger, R.G.; Crussi, F. G.; Baehner, R. L.: Acquired agammaglobulinemia after a life-threatening illness with clinical and laboratory features of infectious mononucleosis in three related male children. *New Eng.J. Med.* 293: 62-65, 1975.
- [74145] 12475.Purtilo, D. T.: Pathogenesis and phenotypes of an X-linked recessive lymphoproliferative syndrome. *Lancet* II: 882-885, 1976.
- [74146] 12476.Purtilo, D. T.: X-linked lymphoproliferative syndrome: an immunodeficiency disorder with acquired

agammaglobulinemia, fatal infectious mononucleosis, or malignant lymphoma. Arch. Path. Lab. Med. 105: 119–121, 1981.

- [74147] 12477. Purtilo, D. T.; Bhawan, J.; Hutt, L. M.; De Nicola, L.; Szymanski, I.; Yang, J. P. S.; Boto, W.; Naier, R.; Thorley-Lawson, D.: Epstein-Barrvirus in the X-linked recessive lymphoproliferative syndrome. Lancet I: 798–801, 1978.
- [74148] 12478. Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.: Fatal infectious mononucleosis in familial lymphohistiocytosis. (Letter) New Eng. J. Med. 201: 736 only, 1974.
- [74149] 12479. Purtilo, D. T.; Cassel, C. K.; Yang, J. P. S.; Harper, R.; Stephenson, S. R.; Landing, B. H.; Vewter, G. F.: X-linked recessive progressive combined variable immunodeficiency (Duncan's disease). Lancet I: 935–941, 1975.
- [74150] 12480. Purtilo, D. T.; DeFlorio, D., Jr.; Hutt, L. M.; Bhawan, J.; Yang, J. P. S.; Otto, R. L.; Edwards, W.: Variable phenotypic expression of an X-linked recessive lymphoproliferative syndrome. New Eng. J. Med. 297: 1077–1081, 1977.
- [74151] 12481. Purtilo, D. T.; Grierson, H. L.: Methods of detection of new families with X-linked lymphoproliferative disease. Cancer Genet. Cytogenet. 51: 143–153, 1991.
- [74152] 12482. Purtilo, D. T.; Sakamoto, K.; Barnabei, V.; Seeley, J.; Bechtold, T.; Rogers, G.; Yetz, J.; Harada, S.; the XLP col-

laborators: Epstein-Barrvirus-induced diseases in boys with the X-linked lymphoproliferative syndrome (XLP): update on studies of the registry. *Am. J. Med.* 73:49-56, 1982.

[74153] 12483. Purtilo, D. T.; Yang, J. P. S.; Allegra, S.; DeFlorio, D.; Hutt, L. M.; Soltani, M.; Vawter, G. F.: Hematopathology and pathogenesis of the X-linked recessive lymphoproliferative syndrome. *Am. J. Med.* 62:225-233, 1977.

[74154] 12484. Sanger, W. G.; Grierson, H. L.; Skare, J.; Wyandt, H.; Pirruccello, S.; Fordyce, R.; Purtilo, D. T.: Partial Xq25 deletion in a family with the X-linked lymphoproliferative disease (XLP). *Cancer Genet. Cytogenet.* 47: 163-169, 1990.

[74155] 12485. Sayos, J.; Wu, C.; Morra, M.; Wang, N.; Zhang, X.; Allen, D.; van Schaik, S.; Notarangelo, L.; Gehat, R.; Roncarolo, M. G.; Oettgen, H.; De Vries, J. E.; Aversall, G.; Terhorst, C.: The X-linked lymphoproliferative-disease gene product SAP regulates signals induced through the co-receptor SLAM. *Nature* 395: 462-469, 1998.

[74156] 12486. Scher, I.: The CBA/N mouse strain: an experimental model illustrating the influence of the X-chromosome on immunity. *Adv. Immun.* 33: 1-71, 1982.

[74157] 12487. Schuster, V.; Kreth, H. W.: X-linked lymphoproliferative disease. In: Ochs, H. D.; Smith, C. I. E.; Puck, J. M.

(eds.): Primary Immunodeficiency Diseases: A Molecular and Genetic Approach. New York: Oxford University Press 1999. Pp. 222–232.

- [74158] 12488. Seemayer, T. A.; Gross, T. G.; Egeler, R. M.; Pirruccello, S. J.; Davis, D. J.; Kelly, C. M.; Okano, M.; Lanyi, A.; Sumegi, J.: X-linked lymphoproliferative disease: twenty-five years after the discovery. *Pediat. Res.* 38: 471–478, 1995.
- [74159] 12489. Skare, J.; Grierson, H.; Wyandt, H.; Sanger, W.; Milunsky, J.; Purtilo, D.; Sullivan, J.; Milunsky, A.: Genetics of the X-linked lymphoproliferative syndrome. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A161 only, 1989.
- [74160] 12490. Skare, J.; Madan, S.; Glaser, J.; Purtilo, D.; Nitowsky, H.; Pulijaal, V.; Milunsky, A.: First prenatal diagnosis of X-linked lymphoproliferative disease. *Am. J. Med. Genet.* 44: 79–81, 1992.
- [74161] 12491. Skare, J.; Milunsky, A.; Byron, K.; Sullivan, J.: The mutation causing X-linked lymphoproliferative syndrome lies in Xq26. (Abstract) *Am. J. Hum. Genet.* 41: A185 only, 1987.
- [74162] 12492. Skare, J.; Wu, B.-L.; Madan, S.; Pulijaal, V.; Purtilo, D.; Haber, D.; Nelson, D.; Sylla, B.; Grierson, H.; Nitowsky, H.; Glaser, J.; Wissink, J.; White, B.; Holden, J.; Housman,

D.; Lenoir, G.; Wyandt, H.; Milunsky, A.: Characterization of three overlapping deletions causing X-linked lymphoproliferative disease. *Genomics* 16: 254–255, 1993.

- [74163] 12493. Skare, J. C.; Grierson, H. L.; Sullivan, J. L.; Nussbaum, R. L.; Purtilo, D. T.; Sylla, B. S.; Lenoir, G. M.; Reilly, D. S.; White, B. N.; Milunsky, A.: Linkage analysis of seven kindreds with the X-linked lymphoproliferative syndrome (XLP) confirms that the XLP locus is near DXS42 and DXS37. *Hum. Genet.* 82: 354–358, 1989.
- [74164] 12494. Skare, J. C.; Milunsky, A.; Byron, K. S.; Sullivan, J. L.: Mapping the X-linked lymphoproliferative syndrome. *Proc. Nat. Acad. Sci.* 84: 2015–2018, 1987.
- [74165] 12495. Skare, J. C.; Sullivan, J. L.; Milunsky, A.: Mapping the mutation causing the X-linked lymphoproliferative syndrome in relation to restriction fragment length polymorphisms on Xq. *Hum. Genet.* 82: 349–353, 1989.
- [74166] 12496. Steinherz, R.; Levy, Y.; Litwin, A.; Nitzan, M.; Friedman, E.; Levin, S.: X-linked lymphoproliferative syndrome: a new kindred with variable phenotypic expression. *Am. J. Dis. Child.* 139: 191–193, 1985.
- [74167] 12497. Nehlin, J. O.; Hara, E.; Kuo, W.-L.; Collins, C.; Campisi, J.: Genomic organization, sequence, and chromosomal localization of the human helix-loop-helix *Id1* gene.

Biochem. Biophys. Res. Commun. 231:628–634, 1997.

[74168] 12498. Loftin, C. D.; Trivedi, D. B.; Tiano, H. F.; Clark, J. A.; Lee, C. A.; Epstein, J. A.; Morham, S. G.; Breyer, M. D.; Nguyen, M.; Hawkins, B. M.; Goulet, J. L.; Smithies, O.; Koller, B. H.; Langenbach, R.: Failure of ductus arteriosus closure and remodeling in neonatal mice deficient in cyclooxygenase-1 and cyclooxygenase-2. Proc. Nat. Acad. Sci. 98: 1059–1064, 2001.

[74169] 12499. Macchia, L.; Di Paola, R.; Guerrese, M.-C.; Chiechi, L. M.; Tursi, A.; Caiaffa, M. F.; Haeggstrom, J. Z.: Expression of prostaglandin endoperoxide H synthase 1 and 2 in human placenta at term. Biochem. Biophys. Res. Commun. 233: 496–501, 1997.

[74170] 12500. Morham, S. G.; Langenbach, R.; Loftin, C. D.; Tiano, H. F.; Vouloumanos, N.; Jennette, J. C.; Mahler, J. F.; Kluckman, K. D.; Ledford, A.; Lee, C. A.; Smithies, O.: Prostaglandin synthase 2 gene disruption causes severe renal pathology in the mouse. Cell 83: 473–482, 1995.

[74171] 12501. Neufang, G.; Furstenberger, G.; Heidt, M.; Marks, F.; Muller-Decker, K.: Abnormal differentiation of epidermis in transgenic mice constitutively expressing cyclooxygenase-2 in skin. Proc. Nat. Acad. Sci. 98: 7629–7634, 2001.

- [74172] 12502.O'Banion, M. K.; Sadowski, H. B.; Winn, V.; Young, D. A.: A serum-and glucocorticoid-regulated 4-kilobase mRNA encodes a cyclooxygenase-related protein. *J. Biol. Chem.* 266: 23261-23267, 1991.
- [74173] 12503.O'Banion, M. K.; Winn, V. D.; Young, D. A.: cDNA cloning and functional activity of a glucocorticoid-regulated inflammatory cyclooxygenase. *Proc.Nat. Acad. Sci.* 89: 4888-4892, 1992.
- [74174] 12504.Oshima, M.; Dinchuk, J. E.; Kargman, S. L.; Oshima, H.; Hancock, B.; Kwong, E.; Trzaskos, J. M.; Evans, J. F.; Taketo, M. M.: Suppression of intestinal polyposis in *Apc(delta-716)* knockout mice by inhibition of cyclooxygenase 2 (COX-2). *Cell* 87: 803-809, 1996.
- [74175] 12505.Salmenkivi, K.; Haglund, C.; Ristimäki, A.; Arola, J.; Heikkilä, P.: Increased expression of cyclooxygenase-2 in malignant pheochromocytomas. *J.Clin. Endocr. Metab.* 86: 5615-5619, 2001.
- [74176] 12506.Tay, A.; Squire, J. A.; Goldberg, H.; Skorecki, K.: Assignment of the human prostaglandin-endoperoxide synthase 2 (PTGS2) gene to 1q25 by fluorescence in situ hybridization. *Genomics* 23: 718-719, 1994.
- [74177] 12507.Swan, S. K.; Rudy, D. W.; Lasseter, K. C.; Ryan, C. F.; Buechel, K. L.; Lambrecht, L. J.; Pinto, M. B.; Dilzer, S.

C.; Obrda, O.; Sundblad, K. J.; Gumbs, C. P.; Ebel, D. L.; Quan, H.; Larson, P. J.; Schwartz, J. I.; Musliner, T. A.; Gertz, B. J.; Brater, D. C.; Yao, S.-L.: Effect of cyclooxygenase-2 inhibition on renal function in elderly persons receiving a low-salt diet: a randomized controlled trial. *Ann. Intern. Med.* 133: 1-9, 2000.

- [74178] 12508. Tazawa, R.; Xu, X.-M.; Wu, K. K.; Wang, L.-H.: Characterization of the genomic structure, chromosomal location and promoter of human prostaglandin H synthase-2 gene. *Biochem. Biophys. Res. Commun.* 203:190-199, 1994.
- [74179] 12509. Tsujii, M.; DuBois, R. N.: Alterations in cellular adhesion and apoptosis in epithelial cells overexpressing prostaglandin endoperoxide synthase 2. *Cell* 83: 493-501, 1995.
- [74180] 12510. Zhou, X.-L.; Lei, Z. M.; Rao, C. V.: Treatment of human endometrial gland epithelial cells with chorionic gonadotropin/luteinizing hormone increases the expression of the cyclooxygenase-2 gene. *J. Clin. Endocr. Metab.* 84: 3364-3377, 1999.
- [74181] 12511. Maho, A.; Bensimon, A.; Vassart, G.; Parmentier, M.: Mapping of the CCXCR1, CX3CR1, CCBP2 and CCR9 genes to the CCR cluster within the 3p21.3 region of the



human genome. *Cytogenet. Cell Genet.* 87:265–268, 1999.

- [74182] 12512. Anderson, D. M.; Johnson, L.; Glaccum, M. B.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Kirstein, M. N.; Shapiro, D. N.; Morris, S. W.; Grabstein, K.; Cosman, D.: Chromosomal assignment and genomic structure of IL15. *Genomics* 25: 701–706, 1995.
- [74183] 12513. Grabstein, K. H.; Eisenman, J.; Shanebeck, K.; Rauch, C.; Srinivasan, S.; Fung, V.; Beers, C.; Richardson, J.; Schoenborn, M. A.; Ahdieh, M.; Johnson, L.; Alderson, M. R.; Watson, J. D.; Anderson, D. M.; Giri, J. G.: Cloning of a T cell growth factor that interacts with the beta chain of the interleukin-2 receptor. *Science* 264: 965–968, 1994.
- [74184] 12514. Roberts, A. I.; Lee, L.; Schwarz, E.; Groh, V.; Spies, T.; Ebert, E. C.; Jabri, B.: Cutting edge: NKG2D receptors induced by IL-15 costimulate CD28-negative effector CTL in the tissue microenvironment. *J. Immun.* 167: 5527–5530, 2001.
- [74185] 12515. Robinson, P.; Okhuysen, P. C.; Chappell, C. L.; Lewis, D. E.; Shahab, I.; Lahoti, S.; White, A. C., Jr.: Expression of IL-15 and IL-4 in IFN-gamma-independent control of experimental human *Cryptosporidium parvum* infection. *Cytokine* 15: 39–46, 2001.

- [74186] 12516.Chen, X.; Vinkemeier, U.; Zhao, Y.; Jeruzalmi, D.; Darnell, J.E., Jr.; Kuriyan, J.: Crystal structure of a tyrosine phosphorylatedSTAT-1 dimer bound to DNA. *Cell* 93: 827–839, 1998.
- [74187] 12517.Copeland, N. G.; Gilbert, D. J.; Schindler, C.; Zhong, Z.; Wen,Z.; Darnell, J. E., Jr.; Mui, A. L.–F.; Miyajima, A.; Quelle, F. W.;Ihle, J. N.; Jenkins, N. A.: Distribution of the mammalian Stat genefamily in mouse chromosomes. *Genomics* 29: 225–228, 1995.
- [74188] 12518.Darnell, J. E., Jr.; Kerr, I. M.; Stark, G. M.: Jak–STAT pathwaysand transcriptional activation in response to IFNs and other extracellularsignaling proteins. *Science* 264: 1415–1421, 1994.
- [74189] 12519.Durbin, J. E.; Hackenmiller, R.; Simon, M. C.; Levy, D. E.: Targetedisruption of the mouse Stat1 gene results in compromised innate immunityto viral disease. *Cell* 84: 443–450, 1996.
- [74190] 12520.Haddad, B.; Pabon–Pena, C. R.; Young, H.; Sun, W. H.: Assignmentof STAT1 to human chromosome 2q32 by FISH and radiation hybrids. *Cytogenet.Cell Genet.* 83: 58–59, 1998.
- [74191] 12521.Ihle, J. N.: STATs: signal transducers and activators of transcription. *Cell* 84:331–334, 1996.

- [74192] 12522.Meraz, M. A.; White, J. M.; Sheehan, K. C. F.; Bach, E. A.; Rodig, S. J.; Dighe, A. S.; Kaplan, D. H.; Riley, J. K.; Greenlund, A. C. Campbell, D.; Carver-Moore, K.; DuBois, R. N.; Clark, R.; Aguet, M.; Schreiber, R. D.: Targeted disruption of the Stat1 gene in mice reveals unexpected physiologic specificity in the JAK-STAT signaling pathway. *Cell* 84:431-442, 1996.
- [74193] 12523.Mowen, K. A.; Tang, J.; Zhu, W.; Schurter, B. T.; Shuai, K.; Herschman, H. R.; David, M.: Arginine methylation of STAT1 modulates IFN- $\alpha$ - $\beta$ -induced transcription. *Cell* 104: 731-741, 2001.
- [74194] 12524.Ramana, C. V.; Chatterjee-Kishore, M.; Nguyen, H.; Stark, G. R.: Complex roles of Stat1 in regulating gene expression. *Oncogene* 19:2619-2627, 2000.
- [74195] 12525.Yamamoto, K.; Kobayashi, H.; Arai, A.; Miura, O.; Hirosawa, S.; Miyasaka, N.: cDNA cloning, expression and chromosome mapping of the human STAT4 gene: both STAT4 and STAT1 genes are mapped to 2q32.2-q32.3. *Cytogenet. Cell Genet.* 77: 207-210, 1997.
- [74196] 12526.Ware, R. E.; Howard, T. A.; Kamitani, T.; Chang, H.-M.; Yeh, E.T. H.; Seldin, M. F.: Chromosomal assignment of genes involved in glycosylphosphatidylinositol anchor biosynthesis: implications for the pathogenesis of

paroxysmal nocturnal hemoglobinuria. Blood  
83:3753–3757, 1994.

- [74197] 12527.Watanabe, R.; Inoue, N.; Westfall, B.; Taron, C. H.; Orlean, P.;Takeda, J.; Kinoshita, T.: The first step of glycosylphosphatidylinositolbiosynthesis is mediated by a complex of PIG–A, PIG–H, PIG–C and GPI1. EMBOJ. 17: 877–885, 1998.
- [74198] 12528.Watanabe, R.; Kinoshita, T.; Masaki, R.; Yamamoto, A.; Takeda,J.; Inoue, N.: PIG–A and PIG–H, which participate in glycosylphosphatidylinositolanchor biosynthesis, form a protein complex in the endoplasmic reticulum. J.Biol. Chem. 271: 26868–26875, 1996.
- [74199] 12529.Chernajovsky, Y.; Brown, A.; Clark, J.: Human kinesin light (beta)chain gene: DNA sequence and functional characterization of its promoterand first exon. DNA Cell Biol. 15: 965–974, 1996.
- [74200] 12530.Goedert, M.; Marsh, S.; Carter, N.: Localization of the humankinesin light chain gene (KNS2) to chromosome 14q32.3 by fluorescencein situ hybridization. Genomics 32: 173–175, 1996.
- [74201] 12531.Kamal, A.; Stokin, G. B.; Yang, Z.; Xia, C.; Goldstein, L. S.:Axonal transport of amyloid precursor protein is mediated by directbinding to the kinesin light chain subunit

of kinesin-I. *Neuron* 28:449–459, 2000.

- [74202] 12532. Butzow, R.; Huhtala, M.-L.; Bohn, H.; Virtanen, I.; Seppala, M.: Purification and characterization of placental protein 5. *Biochem. Biophys. Res. Commun.* 150: 483–490, 1988.
- [74203] 12533. Kisiel, W.; Sprecher, C. A.; Foster, D. C.: Evidence that a second human tissue factor pathway inhibitor (TFPI-2) and human placental protein 5 are equivalent. (Letter) *Blood* 84: 4384–4385, 1994.
- [74204] 12534. Miyagi, Y.; Yasumitsu, H.; Eki, T.; Miyata, S.; Koshikawa, N.; Hirahara, F.; Aoki, I.; Misugi, K.; Miyazaki, K.: Assignment of the human PP5/TFPI-2 gene to 7q22 by FISH and PCR-based human/rodent cell hybrid mapping panel analysis. *Genomics* 35: 267–268, 1996.
- [74205] 12535. Miyagi, Y.; Yasumitsu, H.; Mizushima, H.; Koshikawa, N.; Matsuda, Y.; Itoh, H.; Hori, T.-A.; Aoki, I.; Misugi, K.; Miyazaki, K.: Cloning of the cDNA encoding mouse PP5/TFPI-2 and mapping of the gene to chromosome 6. *DNA and Cell Biology* 15: 947–954, 1996.
- [74206] 12536. Siiteri, J. E.; Koistinen, R.; Salem, H. T.; Bohn, H.; Seppala, M.: Placental protein 5 is related to blood coagulation and fibrinolytic systems. *Life Sci.* 30: 1885–1891, 1982.

- [74207] 12537.Sprecher, C. A.; Kisiel, W.; Mathewes, S.; Foster, D. C.: Molecularcloning, expression, and partial characterization of a second humantissue-factor-pathway inhibitor. Proc. Nat. Acad. Sci. 91: 3353–3357,1994.
- [74208] 12538.Maglott, D. R.; Durkin, A. S.; Lane, S. A.; Callen, D. F.; Feldblyum,T. V.; Nierman, W. C.: The gene for membrane protein E16 (D16S469E)maps to human chromosome 16q24.3 and is expressed in human brain,thymus, and retina. Genomics 23: 303–304, 1994.
- [74209] 12539.Kastury, K.; Druck, T.; Huebner, K.; Barletta, C.; Acampora, D.;Simeone, A.; Faiella, A.; Boncinelli, E.: Chromosome locations ofhuman EMX and OTX genes. Genomics 22: 41–45, 1994.
- [74210] 12540.Simeone, A.; Acampora, D.; Gulisano, M.; Stornaiuolo, A.; Boncinelli,E.: Nested expression domains of four homeobox genes in developingrostral brain. Nature 358: 687–690, 1992.
- [74211] 12541.Bishop, K. M.; Goudreau, G.; O'Leary, D. D. M.: Regulation ofarea identity in the mammalian neocortex by Emx2 and Pax6. Science 288:344–349, 2000.
- [74212] 12542.Boncinelli, E.; Gulisano, M.; Spada, F.; Broccoli, V.: Emx andOtx gene expression in the developing mouse brain. Ciba Found. Symp. 193:100–116, 1995.

- [74213] 12543.Bosetti, A.; Faiella, A.; Boncinelli, E.; Consalez, G. G.: Linkagemapping of Emx2 to mouse chromosome 19. Mammalian Genome 8: 71–72,1997.
- [74214] 12544.Noonan, F. C.; Mutch, D. G.; Mallon, M. A.; Goodfellow, P. J.:Characterization of the homeodomain gene EMX2: sequence conservation,expression analysis, and a search for mutations in endometrial cancers. Genomics 76:37–44, 2001.
- [74215] 12545.Acampora, D.; Mazan, S.; Avantaggiato, V.; Barone, P.; Tuorto,F.; Lallemand, Y.; Brulet, P.; Simeone, A.: Epilepsy and brain abnormalitiesin mice lacking the Otx1 gene. Nature Genet. 14: 218–222, 1996.
- [74216] 12546.Boncinelli, E.; Gulisano, M.; Broccoli, V.: Emx and Otx homeoboxgenes in the developing mouse brain. J. Neurobiol. 24: 1356–1366,1993.
- [74217] 12547.Frantz, G. D.; Weimann, J. M.; Levin, M. E.; McConnell, S. K.:Otx1 and Otx2 define layers and regions in developing cerebral cortexand cerebellum. J. Neurosci. 14: 5725–5740, 1994.
- [74218] 12548.Avraham, S.; Jiang, S.; Ota, S.; Fu, Y.; Deng, B.; Dowler, L. L.;White, R. A.; Avraham, H.: Structural and functional studies of theintracellular tyrosine kinase MATK gene and its translated product. J.Biol. Chem. 270:

1833–1842, 1995.

- [74219] 12549. Bennett, B. D.; Cowley, S.; Jiang, S.; London, R.; Deng, B.; Grabarek, J.; Groopman, J. E.; Goeddel, D. V.; Avraham, H.: Identification and characterization of a novel tyrosine kinase from megakaryocytes. *J. Biol. Chem.* 269: 1068–1074, 1994.
- [74220] 12550. Klages, S.; Adam, D.; Class, K.; Fargnoli, J.; Bolen, J. B.; Penhallow, R. C.: Ctk: a protein-tyrosine kinase related to Csk that defines an enzyme family. *Proc. Nat. Acad. Sci.* 91: 2597–2601, 1994.
- [74221] 12551. Sakano, S.; Iwama, A.; Inazawa, J.; Ariyama, T.; Ohno, M.; Suda, T.: Molecular cloning of a novel non-receptor tyrosine kinase, HYL (hematopoietic consensus tyrosine-lacking kinase). *Oncogene* 9: 1155–1161, 1994.
- [74222] 12552. Zrihan-Licht, S.; Lim, J.; Keydar, I.; Sliwkowski, M. X.; Groopman, J. E.; Avraham, H.: Association of Csk-homologous kinase (CHK) (formerly MATK) with HER-2/ErbB-2 in breast cancer cells. *J. Biol. Chem.* 272: 1856–1863, 1997.
- [74223] 12553. Shimizu, S.; Narita, M.; Tsujimoto, Y.: Bcl-2 family proteins regulate the release of apoptogenic cytochrome c by the mitochondrial channel VDAC. *Nature* 399: 483–487, 1999.



- [74224] 12554.Nichols, A. F.: Personal Communication. Berkeley, Calif. 10/4/1995.
- [74225] 12555.Campbell, H. D.; Schimansky, T.; Claudianos, C.; Ozsarac, N.; Kasprzak,A. B.; Cotsell, J. N.; Young, I. G.; de Couet, H. G.; Gabor Miklos,G. L.: The *Drosophila melanogaster* flightless-I gene involved ingastrulation and muscle degeneration encodes gelsolin-like and leucine-richrepeat domains and is conserved in *Caenorhabditis elegans* and humans. *Proc.Nat. Acad. Sci.* 90: 11386–11390, 1993.
- [74226] 12556.Harmon, D. L.; Gardner–Medwin, D.; Stirling, J. L.: Two new mutationsin a late infantile Tay–Sachs patient are both in exon 1 of the beta–hexosaminidasealpha subunit gene. *J. Med. Genet.* 30: 123–128, 1993.
- [74227] 12557.Buckle, V.; Mondello, C.; Darling, S.; Craig, I. W.; Goodfellow,P. N.: Homologous expressed genes in the human sex chromosome pairingregion. *Nature* 317: 739–741, 1985.
- [74228] 12558.Slaugenhaupt, S. A.; Roca, A. L.; Liebert, C. B.; Altherr, M. R.;Gusella, J. F.; Reppert, S. M.: Mapping of the gene for the Mel1a–melatoninreceptor to human chromosome 4 (MTNR1A) and mouse chromosome 8 (Mtnr1a). *Genomics* 27:355–357, 1995.

- [74229] 12559.Hattori, Y.; Yamashiro, Y.; Ohba, Y.; Miyaji, T.; Morishita, M.;Yamamoto, K.; Yamamoto, K.; Narai, S.; Kimura, A.: A new beta-thalassemiamutation (initiation codon ATG-to-GTG) found in the Japanese population. Hemoglobin 15:317-325, 1991.
- [74230] 12560.Holmes, L. B.: Norrie's disease--an X-linked syndrome of retinalmalformation, mental retardation and deafness. New Eng. J. Med. 284:367-368, 1971.
- [74231] 12561.Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Doi, N.; Nakagawa,M.; Ozawa, M.; Kuroda, N.: Novel mutation at the initiation codonin the Norrie disease gene in two Japanese families. Hum. Genet. 95:105-108, 1995.
- [74232] 12562.Isashiki, Y.; Ohba, N.; Yanagita, T.; Hokita, N.; Hotta, Y.; Hayakawa,M.; Fujiki, K.; Tanabe, U.: Mutations in the Norrie disease gene:a new mutation in a Japanese family. (Letter) Brit. J. Ophthal. 79:703-708, 1995.
- [74233] 12563.Johnson, K.; Mintz-Hittner, H. A.; Conley, Y. P.; Ferrell, R.E.: X-linked exudative vitreoretinopathy caused by an arginine toleucine substitution (R121L) in the Norrie disease protein. Clin.Genet. 50: 113-115, 1996.
- [74234] 12564.Johnston, S. S.; Hanna, J. E.; Nevin, N. C.; Bryars, J. H.: Norrie'sdisease. Birth Defects Orig. Art. Ser. 18(6): 729-738, 1982.

- [74235] 12565.Katayama, S.; Wohlfeld, M.; Golbus, M. S.: First demonstration of recombination between the gene for Norrie disease and probe L1.28. *Am.J. Med. Genet.* 30: 967–970, 1988.
- [74236] 12566.Kivlin, J. D.; Sanborn, G. E.; Wright, E.; Cannon, L.; Carey, J.: Further linkage data on Norrie disease. *Am. J. Med. Genet.* 26:733–736, 1987.
- [74237] 12567.Lindsay, S.; Thiselton, D. L.; Bateman, J. B.; Ngo, J. T.; Sparkes, R. S.; Coleman, M.; Davies, K. E.; Bhattacharya, S. S.: Localisation of the gene for Norrie disease to be-  
tween DXS7 and DXS426 on Xp. *Hum.Genet.* 88: 349–350, 1992.
- [74238] 12568.Meindl, A.; Berger, W.; Meitinger, T.; van de Pol, D.; Achatz, H.; Dorner, C.; Haasemann, M.; Hellebrand, H.; Gal, A.; Cremers, F.; Ropers, H.–H.: Norrie disease is caused by mutations in an extracellular protein resembling C-  
terminal globular domain of mucins. *Nature Genet.* 2:139–143, 1992.
- [74239] 12569.Meindl, A.; Lorenz, B.; Achatz, H.; Hellebrand, H.; Schmitz–Valckenberg, P.; Meitinger, T.: Missense muta-  
tions in the NDP gene in patients with a less severe course of Norrie disease. *Hum. Molec. Genet.* 4:489–490, 1995.
- [74240] 12570.Meitinger, T.; Meindl, A.; Bork, P.; Rost, B.; Sander,

C.; Haasemann,M.; Murken, J.: Molecular modeling of the Norrie disease proteinpredicts a cystine knot growth factor tertiary structure. *NatureGenet.* 5: 376–380, 1993.

[74241] 12571.Moreira-Filho, C. A.; Neustein, I.: A presumptive new variantof Norrie's disease. *J. Med. Genet.* 16: 125–128, 1979.

[74242] 12572.Nance, W. E.; Hara, S.; Hansen, A.; Elliott, J.; Lewis, M.; Chown,B.: Genetic linkage studies in a Negro kindred with Norrie's disease. *Am.J. Hum. Genet.* 21: 423–429, 1969.

[74243] 12573.Ngo, J.; Spence, M. A.; Cortessis, V.; Bateman, J. B.; Sparkes,R. S.: Duplicate report crossing over in Norrie disease family. (Letter) *Am.J. Med. Genet.* 33: 286, 1989.

[74244] 12574.Ngo, J. T.; Bateman, J. B.; Cortessis, V.; Sparkes, R. S.; Mohandas,T.; Inana, G.; Spence, M. A.: Norrie disease: linkage analysis usinga 4.2–kb RFLP detected by a human ornithine aminotransferase cDNAprobe. *Genomics* 4: 539–545, 1989.

[74245] 12575.Ngo, J. T.; Spence, M. A.; Cortessis, V.; Sparkes, R. S.; Bateman,J. B.: Recombinational event between Norrie disease and DXS7 loci. *Clin.Genet.* 34: 43–47, 1988.

[74246] 12576.Norrie, G.: Nogle Blindhedsaarsager: en oversigt. *Hospitalstidende* 76:141–147, 1933.

- [74247] 12577.Ohba, N.; Yamashita, T.: Primary vitreoretinal dysplasia resemblingNorrie's disease in a female: association with X autosome chromosomaltranslocation. *Brit. J. Ophthal.* 70: 64–71, 1986.
- [74248] 12578.Phillips, C. I.; Newton, M.; Duvall, J.; Holloway, S.; Levy, A.M.: Probably Norrie's disease due to mutation: two sporadic sibshipsof two males each, a necropsy of one case, and, given Norrie's disease,a calculation of the gene mutation frequency. *Brit. J. Ophthal.* 70:305–313, 1986.
- [74249] 12579.Rehm, H. L.; Gutierrez–Espeleta, G. A.; Garcia, R.; Jimenez, G.;Khetarpal, U.; Priest, J. M.; Sims, K. B.; Keats, B. J. B.; Morton,C. C.: Norrie disease gene mutation in a large Costa Rican kindredwith a novel phenotype including venous insufficiency. *Hum. Mutat.*9: 402–408, 1997.
- [74250] 12580.Schuback, D. E.; Chen, Z. Y.; Craig, I. W.; Breakefield, X. O.;Sims, K. B.: Mutations in the Norrie disease gene. *Hum. Mutat.* 5:285–292, 1995.
- [74251] 12581.Shastry, B. S.: Identification of a recurrent missense mutationin the Norrie disease gene associated with a simplex case of exudativevitreoretinopathy. *Biochem. Biophys. Res. Commun.* 246: 35–38, 1998.
- [74252] 12582.Shastry, B. S.; Hejtmancik, J. F.; Plager, D. A.; Hartzler, M.K.; Trese, M. T.: Linkage and candidate gene

analysis in X-linked familial exudative vitreoretinopathy.

Genomics 27: 341–344, 1995.

- [74253] 12583. Shastry, B. S.; Hejtmancik, J. F.; Trese, M. T.: Identification of novel missense mutations in the Norrie disease gene associated with one X-linked and four sporadic cases of familial exudative vitreoretinopathy. Hum. Mutat. 9: 396–401, 1997.
- [74254] 12584. Sims, K. B.; Lebo, R. V.; Benson, G.; Shalish, C.; Schuback, D.; Chen, Z. Y.; Bruns, G.; Craig, I. W.; Golbus, M. S.; Breakefield, X. O.: The Norrie disease gene maps to a 150 kb region on chromosome Xp11.3. Hum. Molec. Genet. 1: 83–89, 1992.
- [74255] 12585. Sims, K. B.; Ozelius, L.; Corey, T.; Rinehart, W. B.; Liberfarb, R.; Haines, J.; Chen, W. J.; Norio, R.; Sankila, E.; de la Chapelle, A.; Murphy, D. L.; Gusella, J.; Breakefield, X. O.: Norrie disease gene is distinct from the monoamine oxidase genes. Am. J. Hum. Genet. 45: 424–434, 1989.
- [74256] 12586. Taylor, P. J.; Coates, T.; Newhouse, M. L.: Episkopi blindness: hereditary blindness in a Greek Cypriot family. Brit. J. Ophthalmol. 43: 340–344, 1959.
- [74257] 12587. Torrente, I.; Mangino, M.; Gennarelli, M.; Novelli, G.; Giannotti, A.; Vadala, P.; Dallapiccola, B.: Two new missense mutations (A105T and C110G) in the norrin gene in

two Italian families with Norrie disease and familial exudative vitreoretinopathy. (Letter) *Am. J. Med. Genet.* 72: 242–244, 1997.

[74258] 12588. Burgoyne, P. S.: Genetic homology and crossing over in the X and Y chromosomes of mammals. *Hum. Genet.* 61: 85–90, 1982.

[74259] 12589. Cooke, H. J.; Brown, W. R. A.; Rappold, G. A.: Hypervariable telomeric sequences from the human sex chromosomes are pseudoautosomal. *Nature* 317:687–692, 1985.

[74260] 12590. Darling, S. M.; Banting, G. S.; Pym, B.; Wolfe, J.; Goodfellow, P. N.: Cloning an expressed gene shared by the human sex chromosomes. *Proc. Nat. Acad. Sci.* 83: 135–139, 1986.

[74261] 12591. Dracopoli, N. C.; Rettig, W. J.; Albino, A. P.; Esposito, D.; Archidiacono, N.; Rocchi, M.; Siniscalco, M.; Old, L. J.: Genes controlling gp25/30 cell-surface molecules map to chromosomes X and Y and escape X-inactivation. *Am. J. Hum. Genet.* 37: 199–207, 1985.

[74262] 12592. Gelin, C.; Aubrit, F.; Phalipon, A.; Raynal, B.; Cole, S.; Kaczorek, M.; Bernard, A.: The E2 antigen, a 32 kD glycoprotein involved in T-cell adhesion processes, is the MIC2 gene product. *EMBO J.* 8:3253–3259, 1989.

- [74263] 12593. Geller, R. L.; Shapiro, L. J.; Mohandas, T. K.: Fine mapping of the distal short arm of the human X chromosome using X/Y translocations. *Am. J. Hum. Genet.* 38: 884–890, 1986.
- [74264] 12594. Goodfellow, P.; Banting, G.; Sheer, D.; Ropers, H. H.; Caine, A.; Ferguson-Smith, M. A.; Povey, S.; Voss, R.: Genetic evidence that a Y-linked gene in man is homologous to a gene on the X chromosome. *Nature* 302: 346–349, 1983.
- [74265] 12595. Goodfellow, P.; Pym, B.; Mohandas, T.; Shapiro, L. J.: The cell surface antigen locus, MIC2X, escapes X-inactivation. *Am. J. Hum. Genet.* 36: 777–782, 1984.
- [74266] 12596. Goodfellow, P. J.; Darling, S. M.; Thomas, N. S.; Goodfellow, P. N.: A pseudoautosomal gene in man. *Science* 234: 740–743, 1986.
- [74267] 12597. Goodfellow, P. N.; Tippet, P.: A human quantitative polymorphism related to Xg blood groups. *Nature* 289: 404–405, 1981.
- [74268] 12598. Levy, R.; Dilley, J.; Fox, R. I.; Warnke, R.: A human thymus-leukemia antigen defined by hybridoma monoclonal antibodies. *Proc. Nat. Acad. Sci.* 76: 6552–6556, 1979.
- [74269] 12599. Pettersen, R. D.; Bernard, G.; Olafsen, M. K.;



Pourtein, M.; Lie, S. O.: CD99 signals caspase-independent T cell death. *J. Immun.* 166:4931–4942, 2001.

- [74270] 12600. Ropers, H. H.; Zimmer, J.; Strobl, G.; Goodfellow, P.: The MIC2X(12E7) locus maps distally from STS on Xp. (Abstract) *Cytogenet. Cell Genet.* 40: 736 only, 1985.
- [74271] 12601. Simmler, M.-C.; Rouyer, F.; Vergnaud, G.; Nystrom-Lahti, M.; Ngo, K. Y.; de la Chapelle, A.; Weissenbach, J.: Pseudoautosomal DNA sequences in the pairing region of the human sex chromosomes. *Nature* 317:692–697, 1985.
- [74272] 12602. Tippett, P.; Shaw, M.-A.; Green, C. A.; Daniels, G. L.: The 12E7 red cell quantitative polymorphism: control by the Y-borne locus, Yg. *Ann. Hum. Genet.* 50: 339–347, 1986.
- [74273] 12603. Janz, R.; Sudhof, T. C.; Hammer, R. E.; Unni, V.; Siegelbaum, S. A.; Bolshakov, V. Y.: Essential roles in synaptic plasticity for synaptogyrin I and synaptophysin I. *Neuron* 24: 687–700, 1999.
- [74274] 12604. McMahon, H. T.; Bolshakov, V. Y.; Janz, R.; Hammer, R. E.; Siegelbaum, S. A.; Sudhof, T. C.: Synaptophysin, a major synaptic vesicle protein, is not essential for neurotransmitter release. *Proc. Nat. Acad. Sci.* 93:4760–4764, 1996.

- [74275] 12605. Brown, C. J.; Sekiguchi, T.; Nishimoto, T.; Willard, H. F.: Regional localization of CCG1 gene which complements hamster cell cycle mutation BN462 to Xq11-Xq13. *Somat. Cell Molec. Genet.* 15: 93-96, 1989.
- [74276] 12606. Derry, J. M. J.; Barnard, P. J.: Localization of the Ccg1 gene on the mouse X chromosome. (Abstract) *Cytogenet. Cell Genet.* 51:988, 1989.
- [74277] 12607. Dikstein, R.; Ruppert, S.; Tjian, R.: TAFII250 is a bipartite protein kinase that phosphorylates the basal transcription factor RAP74. *Cell* 84:781-790, 1996.
- [74278] 12608. Giles, R. E.; Ruddle, F. H.: X-linkage of a human genetic locus that corrects the DNA synthesis lesion in tsC1AGOH mouse cells. *Genetics* 93:975-996, 1979.
- [74279] 12609. Hisatake, K.; Hasegawa, S.; Takada, R.; Nakatani, Y.; Horikoshi, M.; Roeder, R. G.: The p250 subunit of native TATA box-binding factor TFIID is the cell-cycle regulatory protein CCG1. *Nature* 362: 179-181, 1993.
- [74280] 12610. Jacobson, R. H.; Ladurner, A. G.; King, D. S.; Tjian, R.: Structure and function of a human TAFII250 double bromodomain module. *Science* 288:1422-1425, 2000.
- [74281] 12611. Jha, K. K.; Ozer, H. L.: Genetic studies with a mutant mouse cell, ts-2 Balb-3T3, with a temperature-sensitive defect in DNA synthesis. (Abstract) *Genetics* 86:

s32–s33, 1977.

- [74282] 12612.Jha, K. K.; Siniscalco, M.; Ozer, H. L.: Temperature-sensitivemutants of Balb–3T3 cells. III. Hybrids between ts2 and other mousemutant cells affected in DNA synthesis and correction of ts2 defectby human X chromosome. *Somat. Cell Genet.* 6: 603–614, 1980.
- [74283] 12613.Almind, K.; Ahlgren, M. G.; Hansen, T.; Urhammer, S. A.; Clausen,J. O.; Pedersen, O.: Discovery of a Met300Val variant in Shc andstudies of its relationship to birth weight and length, impaired insulinsecretion, insulin resistance, and type 2 diabetes mellitus. *J. Clin.Endocr. Metab.* 84: 2241–2244, 1999.
- [74284] 12614.Harun, R. B.; Smith, K. K.; Leek, J. P.; Markham, A. F.; Norris,A.; Morrison, J. F. J.: Characterization of human SHC p66 cDNA andits processed pseudogene mapping to Xq12–q13.1. *Genomics* 42: 349–352,1997.
- [74285] 12615.McGlade, J.; Cheng, A.; Pelicci, G.; Pelicci, P. G.; Pawson, T.: Shc proteins are phosphorylated and regulated by the v–src and v–fpsprotein–tyrosine–kinases. *Proc. Nat. Acad. Sci.* 89: 8869–8873, 1992.
- [74286] 12616.Migliaccio, E.; Giorgio, M.; Mele, S.; Pelicci, G.; Reboldi, P.;Pandolfi, P. P.; Lanfranccone, L.; Pelicci, P. G.: The p66(shc) adaptorprotein controls oxidative stress re–

sponse and life span in mammals. *Nature* 402:309–313, 1999.

[74287] 12617. Nemoto, S.; Finkel, T.: Redox regulation of fork-head proteins through a p66shc-dependent signaling pathway. *Science* 295: 2450–2452, 2002.

[74288] 12618. Pelicci, G.; Lanfrancone, L.; Grignani, F.; McGlade, J.; Cavallo, F.; Forni, G.; Nicoletti, I.; Grignani, F.; Pawson, T.; Pelicci, P.G.: A novel transforming protein (SHC) with an SH2 domain is implicated in mitogenic signal transduction. *Cell* 70: 93–104, 1992.

[74289] 12619. Yulug, I. G.; Egan, S. E.; See, C. G.; Fisher, E. M. C.: A human SHC-related sequence maps to chromosome 17, the SHC gene maps to chromosome 1. *Hum. Genet.* 96: 245–248, 1995.

[74290] 12620. Zhang, L.; Camerini, V.; Bender, T. P.; Ravichandran, K. S.: A nonredundant role for the adapter protein Shc in thymic T cell development. *Nature Immun.* 3: 749–755, 2002.

[74291] 12621. Abramovitz, M.; Boie, Y.; Nguyen, T.; Rushmore, T. H.; Bayne, M.A.; Metters, K. M.; Slipetz, D. M.; Grygorczyk, R.: Cloning and expression of a cDNA for the human prostanoid FP receptor. *J. Biol. Chem.* 269:2632–2636, 1994.

- [74292] 12622.Betz, R.; Lagercrantz, J.; Kedra, D.; Dumanski, J. P.; Nordenskjold,A.: Genomic structure, 5-prime flanking sequences, and precise localizationin 1p31.1 of the human prostaglandin F receptor gene. *Biochem. Biophys.Res. Commun.* 254: 413–416, 1999.
- [74293] 12623.Sugimoto, Y.; Yamasaki, A.; Segi, E.; Tsuboi, K.; Aze, Y.; Nishimura,T.; Oida, H.; Yoshida, N.; Tanaka, T.; Katsuyama, M.; Hasumoto, K.;Murata, T.; Hirata, M.; Ushikubi, F.; Negishi, M.; Ichikawa, A.; Narumiya,S.: Failure of parturition in mice lacking the prostaglandin F receptor. *Science* 277:681–683, 1997.
- [74294] 12624.Geppert, M.; Khvotchev, M.; Krasnoperov, V.; Goda, Y.; Missler,M.; Hammer, R. E.; Ichtchenko, K.; Petrenko, A. G.; Sudhof, T. C.: Neurexin I-alpha is a major alpha-latrotoxin receptor that cooperatesin alpha-latrotoxin action. *J. Biol. Chem.* 273: 1705–1710, 1998.
- [74295] 12625.Ichtchenko, K.; Hata, Y.; Nguyen, T.; Ullrich, B.; Missler, M.;Moomaw, C.; Sudhof, T. C.: Neuroligin 1: a splice site-specific ligandfor beta-neurexins. *Cell* 81: 435–443, 1995.
- [74296] 12626.Kleiderlein, J. J.; Nisson, P. E.; Jessee, J.; Li, W.-B.; Becker,K. G.; Derby, M. L.; Ross, C. A.; Margolis, R. L.: CCG repeats incDNAs from human brain. *Hum. Genet.* 103:

666–673, 1998. Note: Erratum:Hum. Genet. 104: 113 only, 1999.

- [74297] 12627.Apte, S. S.; Mattei, M.–G.; Olsen, B. R.: Mapping of human BAXgene to chromosome 19q13.3–q13.4 and isolation of a novel alternativelyspliced transcript, BAX–delta. Genomics 26: 592–594, 1995.
- [74298] 12628.Niyonsaba, F.; Iwabuchi, K.; Someya, A.; Hirata, M.; Matsuda, H.;Ogawa, H.; Nagaoka, I.: A cathelicidin family of human antibacterialpeptide LL–37 induces mast cell chemotaxis. Immunology 106: 20–26,2002.
- [74299] 12629.Margolis, R.: Personal Communication. Baltimore, Md. 3/29/2000.
- [74300] 12630.Missler, M.; Sudhof, T. C.: Neurexins: three genes and 1001 products. TrendsGenet. 14: 20–26, 1998.
- [74301] 12631.Ikeshima, H.; Imai, S.; Shimoda, K.; Hata, J.; Takano, T.: Expressionof a MADS box gene, MEF2D, in neurons of the mouse central nervoussystem: implication of its binary function in myogenic and neurogeniccell lineages. Neurosci. Lett. 200: 117–120, 1995.
- [74302] 12632.Delhase, M.; Hayakawa, M.; Chen, Y.; Karin, M.: Positive and negativeregulation of I–kappa–B kinase activity through IKK–beta subunit phosphorylation. Science 284:309–312, 1999.

- [74303] 12633.DiDonato, J. A.; Hayakawa, M.; Rothwarf, D. M.; Zandi, E.; Karin,M.: A cytokine-responsive IkappaB kinase that activates the transcriptionfactor NF-kappaB. *Nature* 388: 548–554, 1997.
- [74304] 12634.Mercurio, F.; Zhu, H.; Murray, B. W.; Shevchenko, A.; Bennett,B. L.; Li, J.; Young, D. B.; Barbosa, M.; Mann, M.; Manning, A.; Rao,A.: IKK-1 and IKK-2: cytokine-activated I-kappa-B kinases essentialfor NF-kappa-B activation. *Science* 278: 860–866, 1997.
- [74305] 12635.Matsushima, A.; Kaisho, T.; Rennert, P. D.; Nakano, H.; Kurosawa,K.; Uchida, D.; Takeda, K.; Akira, S.; Matsumoto, M.: Essential roleof nuclear factor (NF)-kappa-B-inducing kinase and inhibitor of kappa-B(I-kappa-B) kinase alpha in NF-kappa-B activation through lymphotoxinbeta receptor, but not through tumor necrosis factor receptor I. *J.Exp. Med.* 193: 631–636, 2001.
- [74306] 12636.Brzezinski, A.: Melatonin in humans. *New Eng. J. Med.* 336: 186–195,1997.
- [74307] 12637.Liu, R.-Y.; Zhou, J.-N.; van Heerikhuize, J.; Hofman, M. A.; Swaab,D. F.: Decreased melatonin levels in post-mortem cerebrospinal fluidin relation to aging, Alzheimer's disease, and apolipoprotein E-ep-

silon-4/4genotype. J. Clin. Endocr. Metab. 84: 323–327, 1999.

- [74308] 12638.Nelson, C. S.; Ikeda, M.; Gompf, H. S.; Robinson, M. L.; Fuchs, N. K.; Yoshioka, T.; Neve, K. A.; Allen, C. N.: Regulation of melatonin<sub>1a</sub> receptor signaling and trafficking by asparagine-124. Molec. Endocr. 15:1306–1317, 2001.
- [74309] 12639.Reppert, S. M.; Weaver, D. R.: Melatonin madness. Cell 83: 1059–1062, 1995.
- [74310] 12640.Reppert, S. M.; Weaver, D. R.; Ebisawa, T.: Cloning and characterization of a mammalian melatonin receptor that mediates reproductive and circadian responses. Neuron 13: 1177–1185, 1994.
- [74311] 12641.Muller-Pillasch, F.; Zimmerhackl, F.; Lacher, U.; Schultz, N.; Hameister, H.; Varga, G.; Friess, H.; Buchler, M.; Adler, G.; Gress, T. M.: Cloning of novel transcripts of the human guanine-nucleotide-exchange factor Mss4: in situ chromosomal mapping and expression in pancreatic-cancer. Genomics 46: 389–396, 1997.
- [74312] 12642.Yu, H.; Schreiber, S. L.: Cloning, Zn(2+) binding, and structural characterization of the guanine nucleotide exchange factor human Mss4. Biochemistry 34:9103–9110, 1995.



- [74313] 12643.Yabe, D.; Nakamura, T.; Kanazawa, N.; Tashiro, K.; Honjo, T.:Calumenin, a  $\text{Ca}(2+)$ –binding protein retained in the endoplasmic reticulumwith a novel carboxyl–terminal sequence, HDEF. J. Biol. Chem. 272:18232–18239, 1997.
- [74314] 12644.Yabe, D.; Taniwaki, M.; Nakamura, T.; Kanazawa, N.; Tashiro, K.;Honjo, T.: Human calumenin gene (CALU): cDNA isolation and chromosomal mapping to 7q32. Genomics 49: 331–333, 1998.
- [74315] 12645.Ito, T.; Yang, M.; May, W. S.: RAX, a cellular activator for double–strandedRNA–dependent protein kinase during stress signaling. J. Biol. Chem. 274:15427–15432, 1999.
- [74316] 12646.Patel, R. C.; Sen, G. C.: PACT, a protein activator of the interferon–inducedprotein kinase, PKR. EMBO J. 17: 4379–4390, 1998.
- [74317] 12647.Scott, A. F.: Personal Communication. Baltimore, Md. 2001.
- [74318] 12648.Shearman, L. P.; Zylka, M. J.; Weaver, D. R.; Kolakowski, L. F., Jr.; Reppert, S. M.: Two period homologs: circadian expression and photic regulation in the suprachiasmatic nuclei. Neuron 19: 1261–1269, 1997.
- [74319] 12649.Toh, K. L.; Jones, C. R.; He, Y.; Eide, E. J.; Hinz, W. A.; Virshup, D. M.; Ptacek, L. J.; Fu, Y.–H.: An hPer2 phos–

phorylation site mutation in familial advanced sleep phase syndrome. *Science* 291: 1040–1043, 2001.

- [74320] 12650. Engelender, S.; Wanner, T.; Kleiderlein, J. J.; Wakabayashi, K.; Tsuji, S.; Takahashi, H.; Ashworth, R.; Margolis, R. L.; Ross, C. A.: Organization of the human synphilin-1 gene, a candidate for Parkinson's disease. *Mammalian Genome* 11: 763–766, 2000.
- [74321] 12651. Chien, W.; Pei, L.: A novel binding factor facilitates nuclear translocation and transcriptional activation function of the pituitary tumor-transforming gene product. *J. Biol. Chem.* 275: 19422–19427, 2000.
- [74322] 12652. Yaspo, M.-L.; Aaltonen, J.; Horelli-Kuitunen, N.; Peltonen, L.; Lehrach, H.: Cloning of a novel human putative type Ia integral membrane protein mapping to 21q22.3. *Genomics* 49: 133–136, 1998.
- [74323] 12653. Yaspo, M.-L.; Gellen, L.; Mott, R.; Korn, B.; Nizetic, D.; Poustka, A. M.; Lehrach, H.: Model for a transcript map of human chromosome 21: isolation of new coding sequences from exon and enriched cDNA libraries. *Hum. Molec. Genet.* 4: 1291–1304, 1995.
- [74324] 12654. Ghiso, J. A.; Holton, J.; Miravalle, L.; Calero, M.; Lashley, T.; Vidal, R.; Houlden, H.; Wood, N.; Neubert, T. A.; Rostagno, A.; Plant, G.; Revesz, T.; Frangione, B.: Sys-

temic amyloid deposits in familial British dementia. *J. Biol. Chem.* 276: 43909–43914, 2001.

[74325] 12655. Stromgren, E.; Dalby, A.; Dalby, M.; Ranheim, B.: *Acta Neurol. Scand.* 46 (suppl. 43): 97–98, 1970.

[74326] 12656. Gandhi, R.; Elble, R. C.; Gruber, A. D.; Schreur, K. D.; Ji, H.-L.; Fuller, C. M.; Pauli, B. U.: Molecular and functional characterization of a calcium-sensitive chloride channel from mouse lung. *J. Biol. Chem.* 273: 32096–32101, 1998.

[74327] 12657. Gruber, A. D.; Elble, R. C.; Ji, H.-L.; Schreur, K. D.; Fuller, C. M.; Pauli, B. U.: Genomic cloning, molecular characterization, and functional analysis of human CLCA1, the first human member of the family of  $\text{Ca}^{2+}$ -activated  $\text{Cl}^-$  channel proteins. *Genomics* 54:200–214, 1998.

[74328] 12658. Gruber, A. D.; Schreur, K. D.; Ji, H.-L.; Fuller, C. M.; Pauli, B. U.: Molecular cloning and transmembrane structure of hCLCA2 from human lung, trachea, and mammary gland. *Am. J. Physiol.* 276: C1261–C1270, 1999.

[74329] 12659. Liao, C.; Wang, X. Y.; Wei, H. Q.; Li, S. Q.; Merghoub, T.; Pandolfi, P. P.; Wolgemuth, D. J.: Altered myelopoiesis and the development of acute myeloid leukemia in transgenic mice overexpressing cyclin A1. *Proc. Nat. Acad. Sci.* 98: 6853–6858, 2001.

- [74330] 12660.Liu, D.; Matzuk, M. M.; Sung, W. K.; Guo, Q.; Wang, P.; Wolgemuth, D. J.: Cyclin A1 is required for meiosis in the male mouse. *NatureGenet.* 20: 377–380, 1998.
- [74331] 12661.Muller, C.; Yang, R.; Beck–von–Peccoz, L.; Idos, G.; Verbeek, W.;Koeffler, H. P.: Cloning of the cyclin A1 genomic structure and characterization of the promoter region: GC boxes are essential for cell cycle–regulatedtranscription of the cyclin A1 gene. *J. Biol. Chem.* 274: 11220–11228, 1999.
- [74332] 12662.Albertsen, H. M.; Smith, S. A.; Mazoyer, S.; Fujimoto, E.; Stevens, J.; Williams, B.; Rodriguez, P.; Cropp, C. S.; Slijepcevic, P.; Carlson, M.; Robertson, M.; Bradley, P.; and 9 others: A physical map and candidate genes in the BRCA1 region on chromosome 17q12–21. *NatureGenet.* 7: 472–479, 1994.
- [74333] 12663.Han, H.–J.; Sudo, K.; Inazawa, J.; Nakamura, Y.: Isolation and mapping of a human gene (RABL) encoding a small GTP–binding protein homologous to the Ras–related RAB gene. *Cytogenet. Cell Genet.* 73:137–139, 1996.
- [74334] 12664.Csoka, A. B.; Scherer, S. W.; Stern, R.: Expression analysis of six paralogous human hyaluronidase genes clustered on chromosomes 3p21 and 7q31. *Genomics* 60: 356–361, 1999.

- [74335] 12665.Heckel, D.; Comtesse, N.; Brass, N.; Blin, N.; Zang, K. D.; Meese,E.: Novel immunogenic antigen homologous to hyaluronidase in meningioma. *Hum.Molec. Genet.* 7: 1859–1872, 1998.
- [74336] 12666.Dolganov, G. M.; Maser, R. S.; Novikov, A.; Tosto, L.; Chong, S.;Bressan, D. A.; Petrini, J. H. J.: Human Rad50 is physically associatedwith human Mre11: identification of a conserved multiprotein compleximplicated in recombinational DNA repair. *Molec. Cell Biol.* 16:4832–4841, 1996.
- [74337] 12667.Hopfner, K.–P.; Craig, L.; Moncalian, G.; Zinkel, R. A.; Usui,T.; Owen, B. A. L.; Karcher, A.; Henderson, B.; Bodmer, J.–L.; McMurray,C. T.; Carney, J. P.; Petrini, J. H. J.; Tainer, J. A.: The Rad50zinc–hook is a structure joining Mre11 complexes in DNA recombinationand repair. *Nature* 418: 562–566, 2002.
- [74338] 12668.Luo, G.; Yao, M. S.; Bender, C. F.; Mills, M.; Bladl, A. R.; Bradley,A.; Petrini, J. H. J.: Disruption of mRad50 causes embryonic stemcell lethality, abnormal embryonic development, and sensitivity toionizing radiation. *Proc. Nat. Acad. Sci.* 96: 7376–7381, 1999.
- [74339] 12669.Trujillo, K. M.; Yuan, S.–S. F.; Lee, E. Y.–H. P.; Sung, P.: Nucleaseactivities in a complex of human recombina–

tion and DNA repair factors Rad50, Mre11, and p95. *J. Biol. Chem.* 273: 21447–21450, 1998.

[74340] 12670. Clark, J.; Lu, Y. J.; Sidhar, S. K.; Parker, C.; Gill, S.; Smedley, D.; Hamoudi, R.; Linehan, W.; Shipley, J.; Cooper, C.: Fusion of splicing factor genes PSF and NonO (p54nrb) to the TFE3 gene in papillary renal cell carcinoma. *Oncogene* 15: 2233–2239, 1997.

[74341] 12671. Heimann, P.; El Housni, H.; Ogur, G.; Weterman, M. A. J.; Petty, E. M.; Vassart, G.: Fusion of a novel gene, RCC17, to the TFE3 gene in t(X;17)(p11.2;q25.3)-bearing papillary renal cell carcinomas. *Cancer Res.* 61: 4130–4135, 2001.

[74342] 12672. Henthorn, P. S.; Stewart, C. C.; Kadesch, T.; Puck, J. M.: The gene encoding human TFE3, a transcription factor that binds the immunoglobulin heavy-chain enhancer, maps to Xp11.22. *Genomics* 11: 374–378, 1991.

[74343] 12673. Joyama, S.; Ueda, T.; Shimizu, K.; Kudawara, I.; Mano, M.; Funai, H.; Takemura, K.; Yoshikawa, H.: Chromosome rearrangement at 17q25 and Xp11.2 in alveolar soft-part sarcoma: a case report and review of the literature. *Cancer* 86: 1246–1250, 1999.

[74344] 12674. Ladanyi, M.; Lui, M. Y.; Antonescu, C. R.; Krause-Boehm, A.; Meindl, A.; Argani, P.; Healey, J. H.; Ueda, T.;

Yoshikawa, H.; Meloni-Ehrig, A.; Sorensen, P. H. B.; Mertens, F.; Mandahl, N.; van den Berghe, H.; Sciort, R.; Dal Cin, P.; Bridge, J.: The der(17)t(X;17)(p11;q25) of human alveolar soft part sarcoma fuses the TFE3 transcription factor gene to ASPL, a novel gene at 17q25. *Oncogene* 20: 48–57, 2001.

[74345] 12675. Macchi, P.; Notarangelo, L.; Giliani, S.; Strina, D.; Repetto, M.; Sacco, M. G.; Vezzoni, P.; Villa, A.: The genomic organization of the human transcription factor 3 (TFE3) gene. *Genomics* 28: 491–494, 1995.

[74346] 12676. Shipley, J. M.; Birdsall, S.; Clark, J.; Crew, J.; Gill, S.; Linehan, M.; Gnarr, J.; Gisher, S.; Craig, I. W.; Cooper, C. S.: Mapping the X chromosome breakpoint in two papillary renal cell carcinoma cell lines with a t(X;1)(p11.2;q21.2) and the first report of a female case. *Cytogenet. Cell. Genet.* 71: 280–284, 1995.

[74347] 12677. Sidhar, S. K.; Clark, J.; Gill, S.; Hamoudi, R.; Crew, A. J.; Gwilliam, R.; Ross, M.; Linehan, W. M.; Birdsall, S.; Shipley, J.; Cooper, C. S.: The t(X;1)(p11.2;q21.2) translocation in papillary renal cell carcinoma fuses a novel gene PRCC to the TFE3 transcription factor gene. *Hum. Molec. Genet.* 5: 1333–1338, 1996.

[74348] 12678. Larola, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.:

Control of mRNA decay by heat shock-ubiquitin-proteasome pathway. *Science* 284:499-502, 1999.

- [74349] 12679. Deutsch, D.; Palmon, A.; Dafni, L.; Fisher, L.; Ter-  
mine, J. D.; Young, M.: Cloning, sequencing, and charac-  
terization of tuftelin: a novel acidic enamel protein. *Con-  
nect. Tissue Res.* 27: 121 only, 1992.
- [74350] 12680. Deiss, L. P.; Feinstein, E.; Berissi, H.; Cohen, O.;  
Kimchi, A.: Identification of a novel serine/threonine ki-  
nase and a novel 15-kD protein as potential mediators of  
the gamma interferon-induced cell death. *Genes Dev.* 9:  
15-30, 1995.
- [74351] 12681. Feinstein, E.; Druck, T.; Kastury, K.; Berissi, H.;  
Goodart, S.A.; Overhauser, J.; Kimchi, A.; Huebner, K.: As-  
signment of DAP1 and DAPK: genes that positively mediate  
programmed cell death triggered by IFN-gamma--to chro-  
mosome regions 5p12.2 (sic) and 9q34.1, respectively.  
*Genomics* 29:305-307, 1995.
- [74352] 12682. Fink, T. M.; Vaesen, M.; Kratzin, H. D.; Lichter, P.;  
Zimmer, M.: Localization of the gene encoding the puta-  
tive human HLA class II associated protein (PHAPI) to chro-  
mosome 15q22.3-q23 by fluorescence in situ hybridiza-  
tion. *Genomics* 29: 309-310, 1995.
- [74353] 12683. Li, M.; Makkinje, A.; Damuni, Z.: Molecular identifi-



cation of  $\text{p1}^{\text{L}}(PP2A)$ , a novel potent heat-stable inhibitor protein of protein phosphatase 2A. *Biochemistry* 35: 6998–7002, 1996.

[74354] 12684. Matilla, A.; Koshy, B. T.; Cummings, C. J.; Isobe, T.; Orr, H. T.; Zoghbi, H. Y.: The cerebellar leucine-rich acidic nuclear protein interacts with ataxin-1. *Nature* 389: 974–978, 1997. Note: Erratum: *Nature* 391: 818 only, 1998.

[74355] 12685. Sandberg, A. A.; Morgan, R.; McCallister, J. A.; Kaiser-McCaw, B.; Hecht, F.: Acute myeloblastic leukemia (AML) with  $t(6;9)(p23;q34)$ : a specific subgroup of AML? *Cancer Genet. Cytogenet.* 10: 139–142, 1983.

[74356] 12686. Vaesen, M.; Barnikol-Watanabe, S.; Gotz, H.; Awni, L. A.; Cole, T.; Zimmermann, B.; Kratzin, H. D.; Hilschmann, N.: Purification and characterization of two putative HLA class II associated proteins: PHAPI and PHAPII. *Biol. Chem. Hoppe-Seyler* 375: 113–126, 1994.

[74357] 12687. von Lindern, M.; van Baal, S.; Wiegant, J.; Raap, A.; Hagemeijer, A.; Grosveld, G.: Can, a putative oncogene associated with myeloid leukemogenesis may be activated by fusion of its 3-prime half to different genes: characterization of the set gene. *Molec. Cell Biol.* 12: 3346–3355, 1992.

- [74358] 12688.Hughes, K. A.; Hurlstone, A. F. L.; Tobias, E. S.; McFarlane, R.; Black, D. M.: Absence of ST7 mutations in tumor-derived cell lines and tumors. *Nature Genet.* 29: 380–381, 2001.
- [74359] 12689.Thomas, N. A.; Choong, D. Y. H.; Jokubaitis, V. J.; Neville, P.J.; Campbell, I. G.: Mutation of the ST7 tumor suppressor gene on 7q31.1 is rare in breast, ovarian and colorectal cancers. *Nature Genet.* 29: 379–380, 2001.
- [74360] 12690.Zenklusen, J. C.; Conti, C. J.; Green, E. D.: Mutational and functional analyses reveal that ST7 is a highly conserved tumor-suppressor gene on human chromosome 7q31. *Nature Genet.* 27: 392–398, 2001.
- [74361] 12691.Zenklusen, J. C.; Rodriguez, L. V.; LaCava, M.; Wang, Z.; Goldstein, L. S.; Conti, C. J.: Novel susceptibility locus for mouse hepatomas: evidence for a conserved tumor suppressor gene. *Genome Res.* 6: 1070–1076, 1996.
- [74362] 12692.Zenklusen, J. C.; Weitzel, J. N.; Ball, H. G.; Conti, C. J.: Allelic loss at 7q31.1 in human primary ovarian carcinomas suggests the existence of a tumor suppressor gene. *Oncogene* 11: 359–363, 1995.
- [74363] 12693.Lee, P. L.; Gelbart, T.; West, C.; Adams, M.; Blackstone, R.; Beutler, E.: Three genes encoding zinc finger proteins on human chromosome 6p21.3: members of a

new subclass of the Kruppel gene family containing the conserved SCAN box domain. *Genomics* 43: 191–201, 1997.

[74364] 12694. Tirosvoutis, K. N.; Divane, A.; Jones, M.; Affara, N. A.: Characterization of a novel zinc finger gene (ZNF165) mapping to 6p21 that is expressed specifically in testis. *Genomics* 28: 485–490, 1995.

[74365] 12695. Deutsch, D.; Palmon, A.; Fisher, L. W.; Kolodny, N.; Termine, J. D.; Young, M. F.: Sequencing of bovine enamelin (tuftelin), a novel acidic enamel protein. *J. Biol. Chem.* 266: 16021–16028, 1991.

[74366] 12696. Deutsch, D.; Palmon, A.; Young, M. F.; Selig, S.; Kearns, W. G.; Fisher, L. W.: Mapping of the human tuftelin (TUFT1) gene to chromosome 1 by fluorescence in situ hybridization. (Abstract) *Mammalian Genome* 5: 461–462, 1994.

[74367] 12697. MacDougall, M.; Simmons, D.; Dodds, A.; Knight, C.; Luan, X.; Zeichner-David, M.; Zhang, C.; Ryu, O. H.; Qian, Q.; Simmer, J. P.; Hu, C.-C.: Cloning, characterization, and tissue expression pattern of mouse tuftelin cDNA. *J. Dent. Res.* 77: 1970–1978, 1998.

[74368] 12698. Dickeson, S. K.; Helmkamp, G. M., Jr.; Yarbrough, L. R.: Sequence of a human cDNA encoding phosphatidyli-

nositol transfer protein and occurrence of a related sequence in widely divergent eukaryotes. *Gene* 142:301–305, 1994.

- [74369] 12699. Dickeson, S. K.; Lim, C. N.; Schuyler, G. T.; Dalton, T. P.; Helmkamp, G. M., Jr.; Yarbrough, L. R.: Isolation and sequence of cDNA clones encoding rat phosphatidylinositol transfer protein. *J. Biol. Chem.* 264:16557–16564, 1989.
- [74370] 12700. Hay, J. C.; Martin, T. F. J.: Phosphatidylinositol transfer protein required for ATP-dependent priming of  $\text{Ca}^{2+}$ -activated secretion. *Nature* 366:572–575, 1993.
- [74371] 12701. Vihtelic, T. S.; Goebel, M.; Milligan, S.; O'Tousa, J. E.; Hyde, D. R.: Localization of *Drosophila* retinal degeneration B, a membrane-associated phosphatidylinositol transfer protein. *J. Cell Biol.* 122: 1013–1022, 1993.
- [74372] 12702. Wirtz, K. W. A.: Phospholipid transfer proteins. *Annu. Rev. Biochem.* 60:73–99, 1991.
- [74373] 12703. Fitzgibbon, J.; Pilz, A.; Gayther, S.; Appukuttan, B.; Dulai, K. S.; Delhanty, J. D. A.; Helmkamp, G. M., Jr.; Yarbrough, L. R.; Hunt, D. M.: Localization of the gene encoding human phosphatidylinositol transfer protein (PITPN) to 17p13.3: a gene showing homology to the *Drosophila* retinal degeneration B gene (*rdgB*). *Cytogenet. Cell Genet.*

67:205–207, 1994.

- [74374] 12704. Hamilton, B. A.; Smith, D. J.; Mueller, K. L.; Kerrebrock, A. W.; Bronson, R. T.; van Berkel, V.; Daly, M. J.; Kruglyak, L.; Reeve, M. P.; Nemhauser, J. L.; Hawkins, T. L.; Rubin, E. M.; Lander, E. S.: The vibrator mutation causes neurodegeneration via reduced expression of PITP- $\alpha$ : positional complementation cloning and extragenic suppression. *Neuron* 18:711–722, 1997.
- [74375] 12705. Hara, T.; Yamauchi, M.; Takahashi, E.; Hoshino, M.; Aoki, K.; Ayusawa, D.; Kawakita, M.: The UDP-galactose translocator gene is mapped to band Xp11.23–p11.22 containing the Wiskott–Aldrich syndrome locus. *Somatic Cell Molec. Genet.* 19: 571–575, 1993.
- [74376] 12706. Ishida, N.; Miura, N.; Yoshioka, S.; Kawakita, M.: Molecular cloning and characterization of a novel isoform of the human UDP-galactose transporter, and of related complementary DNAs belonging to the nucleotide-sugar transporter gene family. *J. Biochem.* 120: 1074–1078, 1996.
- [74377] 12707. Miura, N.; Ishida, N.; Hoshino, M.; Yamauchi, M.; Hara, T.; Ayusawa, D.; Kawakita, M.: Human UDP-galactose translocator: molecular cloning of a complementary DNA that complements the genetic defect of a mutant cell line

deficient in UDP–galactose translocator. J. Biochem. 120:236–241, 1996.

- [74378] 12708.Lund, A.; Udd, B.; Juvonen, V.; Andersen, P. M.; Cederquist, K.; Davis, M.; Gellera, C.; Kolmel, C.; Ronnevi, L.–O.; Sperfeld, A.–D.; Sorensen, S.–A.; Tranebjaerg, L.; Van Maldergem, L.; Watanabe, M.; Weber, M.; Yeung, L.; Savontaus, M.–L.: Multiple founder effects in spinal and bulbar muscular atrophy (SBMA, Kennedy disease) around the world. Europ. J. Hum. Genet. 9: 431–436, 2001.
- [74379] 12709.Edelhoff, S.; Ayer, D. E.; Zervos, A. S.; Steingrims-son, E.; Jenkins, N. A.; Copeland, N. G.; Eisenman, R. N.; Brent, R.; Disteche, C. M.: Mapping of two genes encoding members of a distinct subfamily of MAX interacting proteins: MAD to human chromosome 2 and mouse chromosome 6, and MXI1 to chromosome 10 and mouse chromosome 19. Oncogene 9:665–668, 1994.
- [74380] 12710.Ferguson, G. D.; Anagnostaras, S. G.; Silva, A. J.; Herschman, H. R.: Deficits in memory and motor performance in synaptotagmin IV mutant mice. Proc. Nat. Acad. Sci. 97: 5598–5603, 2000.
- [74381] 12711.Ferguson, G. D.; Chen, X.–N.; Korenberg, J. R.; Herschman, H. R.: The human synaptotagmin IV gene defines

an evolutionary break point between syntenic mouse and human chromosome regions but retains ligand inducibility and tissue specificity. *J. Biol. Chem.* 275: 36920–36926, 2000.

[74382] 12712. Holterhus, P.-M.; Wiebel, J.; Sinnecker, G. H. G.; Bruggenwirth, H. T.; Sippell, W. G.; Brinkmann, A. O.; Kruse, K.; Hiort, O.: Clinical and molecular spectrum of somatic mosaicism in androgen insensitivity syndrome. *Pediatr. Res.* 46: 684–690, 1999.

[74383] 12713. Foresta, C.; Bertella, A.; Moro, E.; Roverato, A.; Merico, M.; Ferlin, A.: Sertoli cell function in infertile patients with and without microdeletions of the azoospermia factors on the Y chromosome long arm. *J. Clin. Endocr. Metab.* 86: 2414–2419, 2001.

[74384] 12714. Kobayashi, K.; Mizuno, K.; Hida, A.; Komaki, R.; Tomita, K.; Matsushita, I.; Namiki, M.; Iwamoto, T.; Tamura, S.; Minowada, S.; Nakahori, Y.; Nakagome, Y.: PCR analysis of the Y chromosome long arm in azoospermic patients: evidence for a second locus required for spermatogenesis. *Hum. Molec. Genet.* 3: 1965–1967, 1994.

[74385] 12715. Agulnik, A. I.; Zharkikh, A.; Boettger-Tong, H.; Bourgeron, T.; McElreavey, K.; Bishop, C. E.: Evolution of the DAZ gene family suggests that Y-linked DAZ plays lit-

tle, or a limited, role in spermatogenesisbut underlines a recent African origin for human populations. Hum.Molec. Genet. 7: 1371–1377, 1998.

[74386] 12716.Anonymous: A missing piece on the Y. (Editorial) Nature Genet. 10:367–368, 1995.

[74387] 12717.Cooke, H. J.; Lee, M.; Kerr, S.; Ruggiu, M.: A murine homologueof the human DAZ gene is autosomal and expressed only in male andfemale gonads. Hum. Molec. Genet. 5: 513–516, 1996.

[74388] 12718.Foresta, C.; Moro, E.; Garolla, A.; Onisto, M.; Ferlin, A.: Ychromosome microdeletions in cryptorchidism and idiopathic infertility. J.Clin. Endocr. Metab. 84: 3660–3665, 1999.

[74389] 12719.Makova, K. D.; Li, W.–H.: Strong male–driven evolution of DNAsequences in humans and apes. Nature 416: 624–626, 2002.

[74390] 12720.Menke, D. B.; Mutter, G. L.; Page, D. C.: Expression of DAZ, anazoospermia factor candidate, in human spermatogonia. (Letter) Am.J. Hum. Genet. 60: 237–241, 1997.

[74391] 12721.Moro, E.; Ferlin, A.; Yen, P. H.; Franchi, P. G.; Palka, G.; Foresta,C.: Male infertility caused by a de novo partial deletion of theDAZ cluster on the Y chromosome. J. Clin.



Endocr. Metab. 85: 4069–4073,2000.

- [74392] 12722.Reijo, R.; Lee, T.-Y.; Salo, P.; Alagappan, R.; Brown, L. G.;Rosenberg, M.; Rozen, S.; Jaffe, T.; Straus, D.; Hovatta, O.; de laChapelle, A.; Silber, S.; Page, D. C.: Diverse spermatogenic defects in humans caused by Y chromosome deletions encompassing a novel RNA-binding protein gene. Nature Genet. 10: 383–393, 1995.
- [74393] 12723.Reijo, R.; Seligman, J.; Dinulos, M. B.; Jaffe, T.; Brown, L.G.; Disteche, C. M.; Page, D. C.: Mouse autosomal homolog of DAZ,a candidate male sterility gene in humans, is expressed in male germ cells before and after puberty. Genomics 35: 346–352, 1996.
- [74394] 12724.Saxena, R.; Brown, L. G.; Hawkins, T.; Alagappan, R. K.; Skaletsky,H.; Reeve, M. P.; Reijo, R.; Rozen, S.; Dinulos, M. B.; Disteche,C. M.; Page, D. C.: The DAZ gene cluster on the human Y chromosome arose from an autosomal gene that was transposed, repeatedly amplified and pruned. Nature Genet. 14: 292–299, 1996.
- [74395] 12725.Najmabadi, H.; Huang, V.; Yen, P.; Subbarao, M. N.; Bhasin, D;Banaag, L.; Naseeruddin, S.; de Kretser, D. M.; Baker, H. W. G.; McLachlan,R. I.; Loveland, K. A.; Bhasin, S.: Substantial prevalence of microdeletions of the Y-chromosome in infertile men with idiopathic azoospermia

and oligozoospermia detected using a sequence-tagged site-based mapping strategy. *J. Clin. Endocr. Metab.* 81: 1347–1352, 1996.

[74396] 12726. Wang, G.; Sawai, N.; Kotliarova, S.; Kanazawa, I.; Nukina, N.: Ataxin-3, the MJD1 gene product, interacts with the two human homologs of yeast DNA repair protein RAD23, HHR23A and HHR23B. *Hum. Molec. Genet.* 9: 1795–1803, 2000.

[74397] 12727. Ng, J. M. Y.; Vrieling, H.; Sugawara, K.; Ooms, M. P.; Grootegeed, J. A.; Vreeburg, J. T. M.; Visser, P.; Beems, R. B.; Gorgels, T. G. M. F.; Hanaoka, F.; Hoeijmakers, J. H. J.; van der Horst, G. T. J.: Developmental defects and male sterility in mice lacking the ubiquitin-like DNA repair gene mHR23B. *Molec. Cell. Biol.* 22: 1233–1245, 2002.

[74398] 12728. Volker, M.; Mone, M. J.; Karmakar, P.; van Hoffen, A.; Schul, W.; Vermeulen, W.; Hoeijmakers, J. H. J.; van Driel, R.; van Zeeland, A. A.; Mullenders, L. H. F.: Sequential assembly of the nucleotide excision repair factors in vivo. *Molec. Cell* 8: 213–224, 2001.

[74399] 12729. Abramson, J. S.; Mills, E. L.; Sawyer, M. K.; Regelman, W. R.; Nelson, J. D.; Quie, P. G.: Recurrent infections and delayed separation of the umbilical cord in an infant with abnormal phagocytic cell locomotion and oxidative

response during particle phagocytosis. J. Pediat.  
99:887–894, 1981.

- [74400] 12730.Allende, L. M.; Hernandez, M.; Corell, A.; Garcia-Perez, M. A.;Varela, P.; Moreno, A.; Caragol, I.; Garcia-Martin, F.; Guillen-Perales,J.; Olive, T.; Espanol, T.; Arnaiz-Villena, A.: A novel CD18 genomicdeletion in a patient with severe leucocyte adhesion deficiency: apossible CD2/lymphocyte function-associated antigen-1 functional associationin humans. Immunology 99: 440–450, 2000.
- [74401] 12731.Anderson, D. C.; Springer, T. A.: Leukocyte adhesion deficiency:an inherited defect in the Mac-1, LFA-1, and p150,95 glycoproteins. Annu.Rev. Med. 38: 175–194, 1987.
- [74402] 12732.Arnaout, M. A.; Dana, N.; Gupta, S. K.; Tenen, D. G.; Fathallah,D. M.: Point mutations impairing cell surface expression of the commonbeta subunit (CD18) in a patient with leukocyte adhesion molecule(Leu-CAM) deficiency. J. Clin. Invest. 85: 977–981, 1990.
- [74403] 12733.Arnaout, M. A.; Pitt, J.; Cohen, H. J.; Melamed, J.; Rosen, F.S.; Colten, H. R.: Deficiency of a granulocyte-membrane glycoprotein(gp150) in a boy with recurrent bacterial infections. New Eng. J.Med. 306: 693–699, 1982.
- [74404] 12734.Arnaout, M. A.; Spits, H.; Terhorst, C.; Pitt, J.; Todd,

R. F.,III: Deficiency of a leukocyte surface glycoprotein (LFA-1) in two patients with Mo1 deficiency: effects of cell activation on Mo1/LFA-1 surface expression in normal and deficient leukocytes. J. Clin. Invest. 74:1291-1300, 1984.

[74405] 12735. Anderson, D. C.; Schmalstieg, F. C.; Finegold, M. J.; Hughes, B.J.; Rothlein, R.; Miller, L. J.; Kohl, S.; Tosi, M. F.; Jacobs, R.L.; Waldrop, T. C.; Goldman, A. S.; Shearer, W. T.; Springer, T. A.: The severe and moderate phenotypes of heritable Mac-1, LFA-1 deficiency: their quantitative definition and relation to leukocyte dysfunction and clinical features. J. Infect. Dis. 152: 668-689, 1985.

- [74406] 12736. Breviario, F.; d'Aniello, E. M.; Golay, J.; Peri, G.; Bottazzi, B.; Bairoch, A.; Saccone, S.; Marzella, R.; Predazzi, V.; Rocchi, M.; Della Valle, G.; Dejana, E.; Mantovani, A.; Introna, M.: Interleukin-1-inducible genes in endothelial cells: cloning of a new gene related to C-reactive protein and serum amyloid P component. *J. Biol. Chem.* 267: 22190-22197, 1992.
- [74407] 12737. Lee, T. H.; Lee, G. W.; Ziff, E. B.; Vilcek, J.: Isolation and characterization of eight tumor necrosis factor-induced gene sequences from human fibroblasts. *Molec. Cell. Biol.* 10: 1982-1988, 1990.
- [74408] 12738. Pallini, R.; Guazzi, G. C.; Cannella, C.; Cacace, M. G.: Cloning and sequence analysis of the human liver rhodanese: comparison with the bovine and chicken enzymes. *Biochem. Biophys. Res. Commun.* 180: 887-893, 1991.
- [74409] 12739. Greco, A.; Mariani, C.; Miranda, C.; Lupas, A.; Pagliardini, S.; Pomati, M.; Pierotti, M. A.: The DNA rearrangement that generates the TRK-T3 oncogene involves a novel gene on chromosome 3 whose product has a potential coiled-coil domain. *Molec. Cell Biol.* 15: 6118-6127, 1995.
- [74410] 12740. Mencinger, M.; Panagopoulos, I.; Andreasson, P.; Lassen, C.; Mitelman, F.; Aman, P.: Characterization and

chromosomal mapping of the human TFG gene involved in thyroid carcinoma. *Genomics* 41: 327–331, 1997.

- [74411] 12741. Kramps, T.; Peter, O.; Brunner, E.; Nellen, D.; Froesch, B.; Chatterjee, S.; Murone, M.; Zullig, S.; Basler, K.: Wnt/Wingless signaling requires BCL9/legless-mediated recruitment of pygopus to the nuclear beta-catenin-TCF complex. *Cell* 109: 47–60, 2002.
- [74412] 12742. Willis, T. G.; Zalcberg, I. R.; Coignet, L. J. A.; Wlodarska, M.; Stul, D. M.; Jadayel, D. M.; Bastard, C.; Treleaven, J. G.; Catovsky, D.; Silva, M. L. M.; Dyer, M. J. S.: Molecular cloning of translocation t(1;14)(q21;q32) defines a novel gene (BCL9) at chromosome 1q21. *Blood* 91:1873–1881, 1998.
- [74413] 12743. Kim, D.-H.; Iijima, H.; Goto, K.; Sakai, J.; Ishii, H.; Kim, H.-J.; Suzuki, H.; Kondo, H.; Saeki, S.; Yamamoto, T.: Human apolipoprotein E receptor 2: a novel lipoprotein receptor of the low density lipoprotein receptor family predominantly expressed in brain. *J. Biol. Chem.* 271:8373–8380, 1996.
- [74414] 12744. Kim, D.-H.; Magoori, K.; Inoue, T. R.; Mao, C. C.; Kim, H.-J.; Suzuki, H.; Fujita, T.; Endo, Y.; Saeki, S.; Yamamoto, T. T.: Exon/intron organization, chromosome localization, alternative splicing, and transcription units of

the human apolipoprotein E receptor 2 gene. J. Biol. Chem. 272:8498–8504, 1997.

- [74415] 12745. Davidson, R. G.; Nitowsky, H. M.; Childs, B.: Demonstration of two populations of cells in the human female heterozygous for glucose-6-phosphate dehydrogenase variants. Proc. Nat. Acad. Sci. 50: 481–485, 1963.
- [74416] 12746. De Flora, A.; Morelli, A.; Benatti, U.; Giuntini, P.; Ferraris, A. M.; Galiano, S.; Ravazzolo, R.; Gaetani, G. F.: G6PD Napoli and Ferrara II: two new glucose-6-phosphate dehydrogenase variants having similar characteristics but different intracellular lability and specific activity. Brit. J. Haemat. 48: 417–423, 1981.
- [74417] 12747. Dern, R. J.; McCurdy, P. R.; Yoshida, A.: A new structural variant of glucose-6-phosphate dehydrogenase with a high production rate (G6PD Hektoen). J. Lab. Clin. Med. 73: 283–290, 1969.
- [74418] 12748. De Vita, G.; Alcalay, M.; Sampietro, M.; Cappelini, M. D.; Fiorelli, G.; Toniolo, D.: Two point mutations are responsible for G6PD polymorphism in Sardinia. Am. J. Hum. Genet. 44: 233–240, 1989.
- [74419] 12749. Du, C.; Xu, Y.; Hua, X.; Wu, Q.; Liu, L.; Wu, M.: Studies on erythrocyte glucose-6-phosphate dehydroge-

nase variants in Chinese.III: Gd(-) Miaozi-Baisha. Acta Genet. Sinica 11(2): 153-158, 1984.

- [74420] 12750.Du, C.-S.; Hua, X.-Y.; Wu, Q.-L.; Li, C.-Q.; Zheng, J.-F.; Li,H.-L.: Studies on erythrocyte glucose-6-phosphate dehydrogenase variants in Chinese. IV. Gd(-) Gaohe associated with paroxysmal nocturnal hemoglobinuria. Chinese J. Pathophysiol. 1: 12-15, 1985.
- [74421] 12751.Du, C. S.; Xu, Y. K.; Hua, X. Y.; Wu, Q. L.; Liu, L. B.: Glucose-6-phosphatedehydrogenase variants and their frequency in Guangdong, China. Hum.Genet. 80: 385-388, 1988.
- [74422] 12752.Eber, S. W.; Gahr, M.; Schroter, W.: Glucose-6-phosphate dehydrogenase(G6PD) Iserlohn and G6PD Regensburg: two new severe enzyme defects in German families. Blut 51: 109-115, 1985.
- [74423] 12753.Echard, G.; Gillois, M.: G6PD--alpha-GAL-PGK--HPRT synteny in the rabbit, *Oryctolagus cuniculus*. (Abstract) Cytogenet. Cell Genet. 25:148-149, 1979.
- [74424] 12754.Elizondo, J.; Saenz, G. F.; Paez, C. A.; Ramon, M.; Garcia, M.;Gutierrez, A.; Estrada, M.: G6PD Puerto Limon: a new deficient variant of glucose-6-phosphate dehydrogenase associated with congenital nonsphero-



cytichemolytic anemia. Hum. Genet. 62: 110–112, 1982.

[74425] 12755.Engstrom, P. F.; Beutler, E.: G-6-PD Tripler: a unique variant associated with chronic hemolytic disease. Blood 36: 10–13, 1970.

[74426] 12756.Epstein, C. J.: Mammalian oocytes: X-chromosome activity. Science 163:1078–1079, 1969.

[74427] 12757.Ermakov, N. V.; Chernyak, N. B.; Tokarev, Y. N.: Properties of new variant of glucose-6-phosphate dehydrogenase (Regar variant): glucose metabolism in erythrocytes containing abnormal enzyme. Biokhimiia 48:577–583, 1983.

[74428] 12758.Estrada, M.; Garcia, M.; Gutierrez, A.; Quintero, I.; Gonzalez, R.: G6PD Varadero. Vox Sang. 43: 102–104, 1982.

[74429] 12759.Fairbanks, V. F.; Nepo, A. G.; Beutler, E.; Dickson, E. R.; Honig, G.: Glucose-6-phosphate dehydrogenase variants: reexamination of G6PD Chicago and Cornell and a new variant (G6PD Pea Ridge) resembling G6PD Chicago. Blood 55: 216–220, 1980.

[74430] 12760.Faust, C. J.; Levinson, B.; Gitschier, J.; Herman, G. E.: Extension of the physical map in the region of the mouse X chromosome homologous to human Xq28 and identification of an exception to conserved linkage. Ge-

nomics 13:1289–1295, 1992.

[74431] 12761.Feldman, R.; Gromisch, D. S.; Luhby, A. L.; Beutler, E.: Congenital nonspherocytic hemolytic anemia due to glucose-6-phosphate dehydrogenase East Harlem: a new deficient variant. J. Pediat. 90: 89–91, 1977.

[74432] 12762.Fernandez, M.; Fairbanks, V. F.: Glucose-6-phosphate dehydrogenase deficiency in the Philippines: report of a new variant--G6PD Panay. Mayo Clin. Proc. 43: 645–660, 1968.

[74433] 12763.Ferraris, A. M.; Broccia, G.; Meloni, T.; Forteleoni, G.; Gaetani, G. F.: Glucose-6-phosphate dehydrogenase deficiency and incidence of hematologic malignancy. Am. J. Hum. Genet. 42: 516–520, 1988.100. Filosa, E.: Personal Communication. La Jolla, Calif. 1989.101. Filosa, S.; Calabro, V.; Vallone, D.; Poggi, V.; Mason, P.; Pagnini, D.; Alfinito, F.; Rotoli, B.; Martini, G.; Luzzatto, L.; Battistuzzi, G.: Molecular basis of chronic non-spherocytic haemolytic anaemia: a new G6PD variant (393arg-to-his) with abnormal K(m) G6P and marked in vivo instability. Brit. J. Haemat. 80: 111–116, 1992.102. Filosa, S.; Giacometti, N.; Wangwei, C.; De Mattia, D.; Pagnini, D.; Alfinito, F.; Schettini, F.; Luzzatto, L.; Martini, G.: Somatic-cell selection is a major determinant of the blood-cell phe-

notype in heterozygotes for glucose-6-phosphate dehydrogenase mutations causing severe enzyme deficiency. Am. J. Hum. Genet. 59: 887-895, 1996.103. Fiorelli, G.; Manoussakis, C.; Sampietro, M.; Pittalis, S.; Guglielmino, C. R.; Cappellini, M. D.: Different polymorphic variants of glucose-6-phosphate dehydrogenase (G6PD) in Italy. Ann. Hum. Genet. 53: 229-236, 1989.104. Fite, E.; Morell, F.; Zuazu, J.; Julia, A.; Morera, J.: Leucocyte glucose-6-phosphate dehydrogenase deficiency and necrotizing pneumonia. Europ. J. Resp. Dis. 64: 150-154, 1983.105. Francke, U.; Bakay, B.; Connor, J. D.; Coldwell, J. G.; Nyhan, W. L.: Linkage relationships of X-linked enzymes glucose-6-phosphate dehydrogenase and hypoxanthine guanine phosphoribosyltransferase. Am. J. Hum. Genet. 26: 512-522, 1974.106. Friedman, M. J.; Trager, W.: The biochemistry of resistance to malaria. Sci. Am. 244(3): 154-164, 1981.107. Frigerio, R.; Sole, G.; Olla, N.; Lovicu, M.; Passiu, G.; Carcassi, U.: Cagliari II: a new G-6-PD variant. Haematologica 72: 241-243, 1987.108. Fujii, H.; Miwa, S.; Takegawa, S.; Takahashi, K.; Hirono, A.; Takizawa, T.; Morisaki, T.; Kanno, H.; Taguchi, T.; Okamura, J.: Gd(-) Gifu and Gd(-) Fukuoka: two new variants of glucose-6-phosphate dehydrogenase found in Japan.

Hum. Genet. 66: 276–278, 1984.109. Fujii, H.; Miwa, S.; Tani, K.; Takegawa, S.; Fujinami, N.; Takahashi, K.; Nakayama, S.; Konno, M.; Sato, T.: Glucose 6–phosphate dehydrogenase variants: a unique variant (G6PD Kobe) showed an extremely increased affinity for galactose 6–phosphate and a new variant (G6PD Sapporo) resembling G6PD Pea Ridge. Hum. Genet. 58: 405–407, 1981.110. Gaetani, G. F.; Galiano, S.; Melani, C.; Miglino, M.; Forni, G. L.; Napoli, G.; Perrone, L.; Ferraris, A. M.: A new glucose–6–phosphate dehydrogenase variant with congenital nonspherocytic hemolytic anemia (G6PD Genova): biochemical characterization and mosaicism expression in the heterozygote. Hum. Genet. 84: 337–340, 1990.111. Gahr, M.; Bornhalm, D.; Schroeter, W.: Biochemische Eigenschaften einer neuen Variante des Glucose–6–phosphate dehydrogenase (G6PD) Mangels mit Favis-mus: G6PD Bielefeld. Klin. Wschr. 55: 379–384, 1977.112. Gahr, M.; Bornhalm, D.; Schroeter, W.: Haemolytic anemia due to glucose 6–phosphate dehydrogenase (G6PD) deficiency: demonstration of two new biochemical variants, G6PD Hamm and G6PD Tarsus. Brit. J. Haemat. 33: 363–370, 1976.113. Gahr, M.; Schroeter, W.: Glucose 6–phosphate dehydrogenase (G6PD) Hamburg, a new vari-

ant with chronic nonspherocytic hemolytic anemia. *Euro. J. Clin. Invest.* 4: 187–191, 1974.114. Gahr, M.; Schroeter, W.; Sturzenegger, M.; Bornhalm, D.; Marti, H. R.: Glucose 6-phosphate dehydrogenase (G6PD) deficiency in Switzerland. *Helv. Paediat. Acta* 31: 159–166, 1976.115. Geerdink, R. A.; Horst, R.; Staal, G. E.: An Iraqi Jewish family with a new red cell glucose 6-phosphate dehydrogenase variant (Gd-Bagdad) and kernicterus. *Israel J. Med. Sci.* 9: 1040–1043, 1973.116. Gellin, J.; Benne, F.; Renard, C.; Vaiman, M.; Hors-Cayla, M. C.; Gillois, M.: Pig gene mapping: synteny, attempt to assign the histocompatibility complex (SLA). (Abstract) *Cytogenet. Cell Genet.* 25:159, 1979.117. Gomez-Gallego, F.; Garrido-Pertierra, A.; Bautista, J. M.: Structural defects underlying protein dysfunction in human glucose-6-phosphate dehydrogenase A- deficiency. *J. Biol. Chem.* 275: 9256–9262, 2000.118. Gonzalez, R.; Estrada, M.; Garcia, M.; Gutierrez, A.: G6PD Ciudad de la Habana: a new slow variant with deficiency found in a Cuban family. *Hum. Genet.* 55: 133–135, 1980.119. Gonzalez, R.; Wade, M.; Estrada, M.; Svach, E.; Colombo, B.: G6PD Pinar del Rio: a new variant discovered in a Cuban family. *Biochem. Genet.* 15: 909–913, 1977.120. Goss, S. J.; Harris, H.: Gene transfer by means

of cell fusion. I. Statistical mapping of the human X-chromosome by analysis of radiation-induced gene segregation. *J. Cell Sci.* 25: 17–37, 1977. 121. Gourdin, D.; Vergnes, H.; Bouloux, C.; Ruffie, J.; Gherardi, M.: Polymorphism of erythrocyte G6PD in the baboon. *Am. J. Phys. Anthropol.* 37: 281–288, 1972. 122. Gray, G. R.; Stamatoyanopoulos, G.; Naiman, S. C.; Kliman, M. R.; Klebanoff, S. J.; Austin, T.; Yoshida, A.; Robinson, G. C. G.: Neutrophil dysfunction, chronic granulomatous disease, and non-spherocytic hemolytic anemia caused by complete deficiency of glucose-6-phosphate dehydrogenase. *Lancet* II: 530–534, 1973. 123. Grossman, A.; Ramanathan, K.; Justice, P.; Gordon, J.; Shahidi, N. T.; Hsia, D. Y. Y.: Congenital nonspherocytic hemolytic anemia associated with erythrocyte G-6-PD deficiency in a Negro family. *Pediatrics* 37: 624–629, 1966. 124. Gutierrez, A.; Garcia, M.; Estrada, M.; Quintero, I.; Gonzalez, R.: Glucose-6-phosphate dehydrogenase (G6PD) Guantnamo and G6PD Caujeri: two new glucose-6-phosphate dehydrogenase-deficient variants found in Cuba. *Biochem. Genet.* 25: 231–238, 1987. 125. Hall, K.; Schreeder, M. T.; Prchal, J. T.: G6PD Huntsville: a new glucose-6-phosphate dehydrogenase associated with chronic hemolytic anemia. *Hum.*

Genet. 79: 90–91, 1988.126. Harkonen, M.; Vuopio, P.: Red cell glucose–6–phosphate dehydrogenase deficiency in Finland. Ann. Clin. Res. 6: 187–197, 1974.127. Harley, J. D.; Agar, N. S.; Yoshida, A.: Glucose 6–phosphate dehydrogenase variant Gd(+) Alexandra associated with neonatal jaundice and Gd(–) Camperdown in a young man with lamellar cataracts. J. Lab. Clin. Med. 91: 295–300, 1978.128. Helge, H.; Borner, K.: Kongenitale nichtsphaerozytare haemolytische Anaemie, Katarakt und Glucose–6–phosphat–dehydrogenase–mangel. Dtsch. Med. Wschr. 91: 1584–1589, 1966.129. Hirono, A.; Beutler, E.: Alternative splicing of human glucose–6–phosphate dehydrogenase messenger RNA in different tissues. J. Clin. Invest. 83: 343–346, 1989.130. Hirono, A.; Beutler, E.: Molecular cloning and nucleotide sequence of cDNA for human glucose–6–phosphate dehydrogenase variant A(–). Proc. Nat. Acad. Sci. 85: 3951–3954, 1988.131. Hirono, A.; Fujii, H.; Shima, M.; Miwa, S.: G6PD Nara: a new class 1 glucose–6–phosphate dehydrogenase variant with an eight amino acid deletion. Blood 82: 3250–3252, 1998.132. Hirono, A.; Kawate, K.; Honda, A.; Fujii, H.; Miwa, S.: A single mutation 202G–A in the human glucose–6–phosphate dehydrogenase gene (G6PD) can

cause acute hemolysis by itself. (Letter) Blood 99: 1498only, 2002.133. Hirono, A.; Kuhl, W.; Gelbart, T.; Forman, L.; Fairbanks, V.F.; Beutler, E.: Identification of the binding domain for NADP(+)of human glucose-6-phosphate dehydrogenase by sequence analysis of mutants. Proc. Nat. Acad. Sci. 86: 10015-10017, 1989.134. Hitzeroth, H. W.; Bender, K.: Age-dependency of somatic selection in South African Negro G-6-PD heterozygotes. Hum. Genet. 58: 338-343, 1981.135. Honig, G. R.; Habacon, E.; Vida, L. N.; Matsumoto, F.; Beutler, E.: Three new variants of glucose-6-phosphate dehydrogenase associated with chronic nonspherocytic hemolytic anemia: G6PD Lincoln Park, G6PD Arlington Heights, and G6PD West Town. Am. J. Hemat. 6: 353-360, 1979.136. Hook, E. B.; Stamatoyannopoulos, G.; Yoshida, A.; Motulsky, A.G.: Glucose-6-phosphate dehydrogenase Madrona: a slow electrophoretic glucose-6-phosphate dehydrogenase variant with kinetic characteristics similar to those of normal type. J. Lab. Clin. Med. 72: 404-409, 1968.137. Hors-Cayla, M. C.; Heuertz, S.; Van Cong, N.; Benne, F.: Cattle gene mapping by somatic cell hybridization. (Abstract) Cytogenet. Cell Genet. 25: 165-166, 1979.138. Howell, E. B.; Nelson, A. J.; Jones, O. W.: A new G-6-PD variant associ-



ated with chronic non-spherocytic haemolytic anaemia in a Negrofamilly. J. Med. Genet. 9: 160–164, 1972.139.

Hutz, M. H.; Yoshida, A.; Salzano, F. M.: Three rare G-6-PDvariants from Porto Alegre, Brazil. Hum. Genet. 39: 191–197, 1977.140. Ishwad, C. S.; Naik, S. N.: A new glucose-6-phosphate dehydrogenasevariant (G-6-PD Kalyan) found in a Koli family. Hum. Genet. 66:171–175, 1984.141. Johnson, G. J.; Kaplan, M. E.; Beutler, E.: G6PD Long Prairie:a new mutant exhibiting normal sensitivity to inhibition by NADPHand accompanied by nonspherocytic hemolytic anemia. Blood 49: 247–251,1977.142. Johnston, P. G.; VandeBerg, J. L.; Sharman, G. B.: Inheritanceof erythrocyte glucose 6-phosphate dehydrogenase in the red-neckedwallaby, *Macropus rufogriseus* (Desmarest) consistent with paternalX inactivation. Biochem. Genet. 13: 235–242, 1975.143. Junien, C.; Kaplan, J.-C.; Meienhofer, M. C.; Maigret, P.; Sender,A.: G6PD Baudelocque: a new unstable variant characterized in culturedfibroblasts. Enzyme 18: 48–59, 1974.144. Kaeda, J. S.; Chhotray, G. P.; Ranjit, M. R.; Bautista, J. M.;Reddy, P. H.; Stevens, D.; Naidu, J. M.; Britt, R. P.; Vulliamy, T.J.; Luzzatto, L.; Mason, P. J.: A new glucose-6-phosphate dehydrogenasevariant, G6PD Orissa (44 ala-gly), is the major polymorphic vari-

antin tribal populations in India. Am. J. Hum. Genet. 57:  
 1335–1341, 1995. 145. Kageoka, T.; Satoh, C.; Goriki, K.;  
 Fujita, M.; Neriishi, S.; Yamamura, K.; Kaneko, J.; Masunari,  
 N.: Electrophoretic variants of blood proteins in Japanese.  
 IV. Prevalence and enzymologic characteristics of G6PD  
 variants in Hiroshima and Nagasaki. Hum. Genet. 70:  
 101–108, 1985. 146. Kahn, A.; Bernard, J.-F.; Cottreau, D.;  
 Mazie, J.; Boivin, P.: Gd(–) Abrami, a deficient G6PD variant  
 with hemizygous expression in blood cells of a woman  
 with primary myelofibrosis. Humangenetik 30:41–46,  
 1975. 147. Kahn, A.; Boivin, P.; Hakim, J.; Lagneau, J.: Het-  
 erogeneity of glucose-6-phosphate dehydrogenase ery-  
 throcyte deficiency in the black race: kinetic study  
 and description of two new variants Gd(–) Dakar et  
 Gd(–) Mali. Nouv. Rev. Franc. Hemat. 11:741–758,  
 1971. 148. Kahn, A.; Boivin, P.; Lagneau, J.: Phenotypes de  
 la glucose-6-phosphate dehydrogenase erythrocytaire  
 dans la race noire. Humangenetik 18:261–270, 1973. 149.  
 Kahn, A.; Boulard, M.; Hakim, J.; Schaison, G.; Boivin, P.;  
 Bernard, J.: Anémie hémolytique congénitale non sphéro-  
 cytaire par déficit en glucose 6-phosphate dehydrogenase  
 erythrocytaire: description de deux nouveaux variants:  
 Gd(–) Saint Louis (Paris) et Gd(–) Hayem. Nouv. Rev. Franc.

Hemat. 14: 587–600, 1974.150. Kahn, A.; Dao, C.; Cottreau, D.; Bilski-Pasquier, G.: 'Gd(-)Hotel Dieu': a new G-6PD variant with chronic hemolysis in a Negropatient from Senegal. Hum. Genet. 39: 353–358, 1977.151. Kahn, A.; Exters, A.; Habedank, M.: Gd(-) Aachen, a new variant of deficient glucose-6-phosphate dehydrogenase. Humangenetik 32:171–180, 1976.152. Kahn, A.; Hakim, J.; Cottreau, D.; Boivin, P.: Gd (-) Matam. An African glucose 6-phosphate dehydrogenase variant with enzyme deficiency. Biochemical and immunological properties in various hemopoietic tissues. Clin. Chim. Acta 59: 183–190, 1975.153. Kahn, A.; North, M. L.; Cottreau, D.; Giron, G.; Lang, J. M.: G6PD Vientiane: a new glucose-6-phosphate dehydrogenase variant with increased stability. Hum. Genet. 43: 85–89, 1978.154. Kahn, A.; North, M. L.; Messer, J.; Boivin, P.: G-6PD 'Ankara': a new G-6PD variant with deficiency found in a Turkish family. Humangenetik 27:247–250, 1975.155. Kanno, H.; Huang, I.-Y.; Kan, Y. W.; Yoshida, A.: Two structural genes on different chromosomes are required for encoding the major subunit of human red cell glucose-6-phosphate dehydrogenase. Cell 58:595–606, 1989.156. Kanno, H.; Takano, T.; Fujii, H.; Tani, K.; Morisaki, T.; Hirono, A.; Kumakawa, T.; Ogura, H.;

Takahashi, K.; Tsutsumi, H.; Miwa, S.: A new glucose-6-phosphate variant (G6PD Iwate) associated with congenital non-spherocytic hemolytic anemia. *Acta Haemat. Jpn.* 51: 715-719, 1987. 157.

Kaplan, J. C.; Hanzlickova-Leroux, A.; Nicholas, A. M.; Rosa, R.; Weiler, C.; Lepercq, G.: A new glucose-6-phosphate dehydrogenase variant (G6PD Port-Royal). *Enzyme* 12: 25-32, 1971. 158.

Kaplan, J. C.; Rosa, R.; Seringe, P.; Hoeffel, J. C.: Le polymorphisme genetique de la glucose-6-phosphate deshydrogenase erythrocytaire chez l'homme. II. Etude d'une nouvelle variete a activite diminuee: le type 'Kabyle.'. *Enzym. Biol. Clin.* 8: 332-340, 1967. 159.

Kaplan, M.; Hammerman, C.; Vreman, H. J.; Stevenson, D. K.; Beutler, E.: Acute hemolysis and severe neonatal hyperbilirubinemia in glucose-6-phosphate dehydrogenase deficient heterozygotes. *J. Pediat.* 139: 137-140, 2001. 160.

Kaplan, M.; Renbaum, P.; Levy-Lahad, E.; Hammerman, C.; Lahad, A.; Beutler, E.: Gilbert syndrome and glucose-6-phosphate dehydrogenase deficiency: a dose-dependent genetic interaction crucial to neonatal hyperbilirubinemia. *Proc. Nat. Acad. Sci.* 94: 12128-12132, 1997. 161.

Kappas, A.; Drummond, G. S.; Valaes, T.: A single dose of Sn-mesoporphyrin prevents development of severe hyperbilirubinemia in glucose-

6-phosphatedehydrogenase-deficient newborns. *Pediatrics* 108: 25–30, 2001.162. Karadsheh, N. S.; Awidi, A. S.; Tarawneh, M. S.: Two new glucose-6phosphate dehydrogenase (G6PD) variants associated with hemolytic anemia: G6PD Amman-1 and G6PD Amman-2. *Am. J. Hemat.* 22: 185–192, 1986.163. Kay, A. C.; Kuhl, W.; Prchal, J.; Beutler, E.: The origin of glucose-6-phosphate-dehydrogenase (G6PD) polymorphisms in African-Americans. *Am. J. Hum. Genet.* 50: 394–398, 1992.164. Kirkman, H. N.; Kidson, C.; Kennedy, M.: Variants of human glucose-6-phosphatedehydrogenase. Studies of samples from New Guinea. In: Beutler, E.: *Hereditary Disorders of Erythrocyte Metabolism*. New York: Grune and Stratton (pub.) 1968. Pp. 126–145.165. Kirkman, H. N.; Lie-Injo, L. E.: Variants of glucose 6-phosphatedehydrogenase in Indonesia. *Nature* 221: 959, 1969.166. Kirkman, H. N.; McCurdy, P. R.; Naiman, J. L.: Functionally abnormal glucose-6-phosphate dehydrogenases. *Cold Spring Harbor Symp. Quant. Biol.* 29: 391–398, 1964.167. Kirkman, H. N.; Ramot, B.; Lee, J. T.: Altered aggregational properties in a genetic variant of human glucose-6-phosphate dehydrogenase. *Biochem. Genet.* 3: 137–150, 1969.168. Kirkman, H. N.;

Riley, H. D., Jr.: Congenital nonspherocytic hemolytic anemia. *Am. J. Dis. Child.* 102: 313–320, 1961.169. Kirkman, H. N.; Rosenthal, I. M.; Simon, E. B.; Carson, P. E.; Brinson, A. G.: 'Chicago I' variant of glucose-6-phosphate dehydrogenase in congenital hemolytic disease. *J. Lab. Clin. Med.* 63: 715–725, 1964.170. Kirkman, H. N.; Schettini, E.; Pickard, B. M.: Mediterranean variant of glucose-6-phosphate dehydrogenase. *J. Lab. Clin. Med.* 63: 726–735, 1964.171. Kirkman, H. N.; Simon, E. R.; Pickard, B. M.: Seattle variant of glucose-6-phosphate dehydrogenase. *J. Lab. Clin. Med.* 66: 834–840, 1965.172. Kissin, C.; Cotte, J.: Etude d'un variant de glucose-6-phosphate dehydrogenase: I B type Constantine. *Enzyme* 11: 277–284, 1970.173. Kitao, T.; Ito, K.; Hattori, K.; Matsuki, T.; Yoneyama, Y.: G6PD Kanazawa: a new variant of glucose-6-phosphate dehydrogenase associated with congenital nonspherocytic hemolytic anemia. *Acta Haemat.* 68: 131–135, 1982.174. Kojima, H.: Congenital nonspherocytic hemolytic disease (CNHD) due to a G-6-PD variant: G-6-PD Kyoto. *Acta Haemat. Jpn.* 35: 32–38, 1972.175. Koliakos, G.; Kalomenopoulou, M.; Grammatikos, P.; Dimitriadou, A.; Kouzi-Koliakos, K.; Zacharaki, R.; Skaragas, G.; Kokka, A.; Trakatellis, A.: A

new glucose 6-phosphate dehydrogenase variant (G6PD Thessaloniki) in a patient with idiopathic myelofibrosis. Hum. Hered. 39: 141–149, 1989. 176. Krasnopol'skaya, K. D.; Bochkov, N. P.: Genetic heterogeneity of hereditary enzymopathies. Vestn. Akad. Med. Nauk. SSSR 9: 56–64, 1982. 177. Krasnopol'skaya, K. D.; Shatskaya, T. L.; Filippov, I. K.; Annenkov, G. A.; Zakharova, T. V.; Mekhtiev, N. K.; Movsum-Zade, K. M.: Genetic heterogeneity of G6PD deficiency: study of mutant alleles in Sheki district of Azerbaijan. Genetika 13: 1455–1461, 1977. 178. Kumakawa, T.; Suzuki, S.; Fujii, H.; Miwa, S.: Frequency of glucose 6-phosphate dehydrogenase (G6PD) deficiency in Tokyo and a new variant: G6PD Musashino. Acta Haemat. Jpn. 50: 25–28, 1987. 179. Kurdi-Haidar, B.; Mason, P. J.; Berrebi, A.; Ankrah-Badu, G.; Al-Ali, A.; Oppenheim, A.; Luzzatto, L.: Origin and spread of the glucose-6-phosphate dehydrogenase variant (G6PD-Mediterranean) in the Middle East. Am. J. Hum. Genet. 47: 1013–1019, 1990. 180. Kwiatkowska, J.; Kacprzak-Bergman, I.: New erythrocyte glucose-6-phosphate dehydrogenase variant. Acta Haemat. 46: 188–192, 1971. 181. Kwok, C. J.; Martin, A. C. R.; Au, S. W. N.; Lam, V. M. S.: G6PDdb, an integrated database of

glucose-6-phosphate dehydrogenase(G6PD) mutations.

Hum. Mutat. 19: 217-224, 2002.182. Lee, K. T.; Thomas,

W. A.; Janakidevi, K.; Kroms, M.; Reiner, J. M.; Borg, K. Y.:

Mosaicism in female hybrid hares heterozygous for glu-

cose-6-phosphate dehydrogenase (G-6-PD). I. General

properties of a hybrid hare model with special reference to

atherogenesis. Exp. Molec. Path. 34: 191-201, 1981.183.

Lenzerini, L.; Meera Khan, P.; Filippi, G.; Rattazzi, M.

C.; Rat, A. K.: Characterization of glucose-6-phosphate

dehydrogenase variants. I. Occurrence of a G6PD Seattle-

like variant in Sardinia and its interaction with G6PD

Mediterranean variant. Am. J. Hum. Genet. 21: 142-153,

1969.184. Lisker, R.; Linares, C.; Motulsky, A. G.: Glu-

cose-6-phosphate dehydrogenase Mexico, a new variant

with enzyme deficiency, abnormal mobility and absence of

hemolysis. J. Lab. Clin. Med. 29: 788-793, 1972.185.

Lisker, R.; Perez-Briceno, R.; Beutler, E.: A new glucose-

6-phosphate dehydrogenase variant, Gd(-) Tepic, charac-

terized by moderate enzyme deficiency and mild episodes

of hemolytic anemia. Hum. Genet. 69:19-21, 1985.186.

Lisker, R.; Perez-Briceno, R.; Rave, V.; Yoshida, A.: Glu-

cose-6-phosphate dehydrogenase Gd(-) Distrito Federal:

nueva variante asociada a deficiencia enzimática moderada



y anemia hemolitica ocasional. Rev. Invest. Clin. 33:209–211, 1981.187. Lisker, R.; Perez Briceno, R.; Aguil-  
lar, L.; Yoshida, A.: A variant glucose-6-phosphate dehy-  
drogenase Gd(-) Chiapas associated with moderate en-  
zyme deficiency and occasional hemolytic anemia. Hum.  
Genet. 43:81–84, 1978.188. Lisker, R.; Perez Briceno, R.;  
Zavala, C.; Navarrette, J. I.; Wessels, M.; Yoshida, A.: A glu-  
cose 6-phosphate dehydrogenase Gd(-)Castilla variant  
characterized by mild deficiency associated with drug-in-  
duced hemolytic anemia. J. Lab. Clin. Med. 90: 754–759,  
1977.189. Liu, Y.; Phelan, J.; Go, R. C. P.; Prchal, J. F.; Pr-  
chal, J.T.: Rapid determination of clonality by detection of  
two closely-linked X chromosome exonic polymorphisms  
using allele-specific PCR. J. Clin. Invest. 99: 1984–1990,  
1997.190. Long, W. K.; Kirkman, H. N.; Sutton, H. H.: Elec-  
trophoretically slow variants of glucose-6-phosphate de-  
hydrogenase from red cells of Negroes. J. Lab. Clin. Med.  
65: 81–87, 1965.191. Longo, L.; Vanegas, O. C.; Patel, M.;  
Rosti, V.; Li, H.; Waka, J.; Merghoub, T.; Pandolfi, P. P.; No-  
taro, R.; Manova, K.; Luzzatto, L.: Maternally transmitted  
severe glucose 6-phosphate dehydrogenase deficiency is  
an embryonic lethal. EMBO J. 21: 4229–4239, 2002.192.  
Luzzatto, L.: Genetic heterogeneity and pathophysiology

of G6PD deficiency. *Brit. J. Haemat.* 28: 151–156, 1974.

193. Luzzatto, L.: Personal Communication. London, England 4/1990.

194. Luzzatto, L.; Afolayam, A.: Enzyme properties of different types of human erythrocyte glucose-6-phosphate dehydrogenase with characterization of two new genetic variants. *J. Clin. Invest.* 47: 1833–1842, 1968.

195. Luzzatto, L.; Martini, G.: X-Linked Wiskott-Aldrich syndrome in a girl. (Letter) *New Eng. J. Med.* 338: 1850–1851, 1998.

196. Luzzatto, L.; Usanga, E. A.; Bien-zle, U.; Esan, G. F. J.; Fusuan, F. A.: Imbalance in X-chromosome expression: evidence for a human X-linked gene affecting growth of hemopoietic cells. *Science* 205:1418–1420, 1979.

197. Luzzatto, L.; Usanga, E. A.; Reddy, S.: Glucose-6-phosphate dehydrogenase deficient red cells: resistance to infection by malarial parasites. *Science* 164: 839–842, 1969.

198. MacDonald, D.; Town, M.; Mason, P.; Vulliamy, T.; Luzzatto, L.; Goff, D. K.: Deficiency in red blood cells. (Letter) *Nature* 350:115, 1991.

199. Maeda, M.; Constantoulakis, P.; Chen, C.-S.; Stamatoyannopoulos, G.; Yoshida, A.: Molecular abnormalities of a human glucose-6-phosphate dehydrogenase variant associated with undetectable enzyme activity and immunologically cross-reacting material. *Am. J. Hum. Genet.*

51:386–395, 1992.200. Mallouh, A. A.; Abu-Osba, Y. K.: Bacterial infections in children with glucose-6-phosphate dehydrogenase deficiency. *J. Pediat.* 111:850–852, 1987.201. Mamlok, R. J.; Mamlok, V.; Mills, G. C.; Daeschner, C. W., III; Schmalstieg, F. C.; Anderson, D. C.: Glucose-6-phosphate dehydrogenase deficiency, neutrophil dysfunction and *Chromobacterium violaceum* sepsis. *J. Pediat.* 111: 852–854, 1987.202. Mamlok, R. J.; Mills, G. C.; Goldblum, R. M.; Daeschner, C. W.: Glucose-6-phosphate dehydrogenase Beaumont: a new variant with severe enzyme deficiency and chronic nonspherocytic hemolytic anemia. *Enzyme* 34:15–21, 1985.203. Mandelli, F.; Amadori, S.; DeLaurenzi, A.; Kahn, A.; Isacchi, G.; Papa, G.: Glucose-6-phosphate dehydrogenase Velletri: a new variant with reduced activity in a patient with congenital non-spherocytic hemolytic anemia. *Acta Haemat.* 57: 121–126, 1977.204. Marks, P. A.; Banks, J.; Gross, R.: Genetic heterogeneity of glucose-6-phosphate dehydrogenase deficiency. *Nature* 194: 454–456, 1962.205. Martin, S. K.; Miller, L. H.; Alling, D.; Okoye, V. C.; Esan, G. J. F.; Osunkoya, B. O.; Deane, M.: Severe malaria and glucose-6-phosphate-dehydrogenase deficiency: a reappraisal of the malaria-G6PD hypothesis. *Lancet* I:524–526,

1979.206. Martini, G.; Toniolo, D.; Vulliamy, T.; Luzzatto, L.; Dono, R.; Viglietto, G.; Paonessa, G.; D'Urso, M.; Persico, M. G.: Structural analysis of the X-linked gene encoding human glucose 6-phosphate dehydrogenase. *EMBOJ.* 5: 1849–1855, 1986.207. Mason, P. J.: New insights into G6PD deficiency. (Annotation) *Brit.J. Haemat.* 94: 585–591, 1996.208. Mason, P. J.; Stevens, D. J.; Luzzatto, L.; Brenner, S.; Aparicio, S.: Genomic structure and sequence of the *Fugu rubripes* glucose-6-phosphate dehydrogenase gene (G6PD). *Genomics* 26: 587–591, 1995.209. McCann, S. R.; Smithwick, A. M.; Temperley, I. J.; Tipton, K.: G6PD (Dublin): chronic non-spherocytic haemolytic anaemia resulting from glucose-6-phosphate dehydrogenase deficiency in an Irish kindred. *J. Med. Genet.* 17: 191–193, 1980.210. McCurdy, P. R.: Use of genetic linkage for the detection of female carriers of hemophilia. *New Eng. J. Med.* 285: 218–219, 1971.211. McCurdy, P. R.; Blackwell, R. Q.; Todd, D.; Tso, S. C.; Tuchinda, S.: Further studies on glucose-6-phosphate dehydrogenase deficiency in Chinese subjects. *J. Lab. Clin. Med.* 75: 788–797, 1970.212. McCurdy, P. R.; Kamel, K.; Selim, O.: Heterogeneity of red cell glucose 6-phosphate dehydrogenase (G6PD) deficiency in

Egypt. J. Lab. Clin. Med. 84: 673–680, 1974.213. McCurdy, P. R.; Kirkman, H. N.; Naiman, J. L.; Jim, R. T. S.; Pickard, B. M.: A Chinese variant of glucose-6-phosphate dehydrogenase. J. Lab. Clin. Med. 67: 374–385, 1966.214. McCurdy, P. R.; Mahmood, L.: Red cell glucose-6-phosphate dehydrogenase deficiency in Pakistan. J. Lab. Clin. Med. 76: 943–948, 1970.215. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.: Variants of glucose-6-phosphate dehydrogenase (G-6-PD) associated with G-6-PD deficiency in Puerto Ricans. J. Lab. Clin. Med. 82: 432–437, 1973.216. McCurdy, P. R.; Maldonado, N. I.; Dillon, D. E.; Conrad, M. E.: Variants of glucose-6-phosphate dehydrogenase (G-6-PD) associated with G-6-PD deficiency in Puerto Ricans. J. Lab. Clin. Med. 82: 432–437, 1973.217. Meloni, T.; Carta, F.; Forteleoni, G.; Carta, A.; Ena, F.; Meloni, G. F.: Glucose 6-phosphate dehydrogenase deficiency and cataract of patients in northern Sardinia. Am. J. Ophthalmol. 110: 661–664, 1990.218. Mentzer, W. C., Jr.; Warner, R.; Addiego, J.; Smith, B.; Walter, T.: G6PD San Francisco: a new variant of glucose-6-phosphate dehydrogenase associated with congenital nonspherocytic hemolytic anemia. Blood 55: 195–198, 1980.219. Mesbah-Namin, S. A.; Sanati, M. H.; Mowjoodi, A.; Mason, P. J.; Vulliamy, T. J.;

Noori-Daloui, M. R.: Three major glucose-6-phosphatedehydrogenase-deficient polymorphic variants identified in Mazandaran state of Iran. *Brit. J. Haemat.* 117: 763–764, 2002.220. Miller, D. R.; Wollman, M. R.: A new variant of glucose 6-phosphatedehydrogenase deficiency hereditary hemolytic anemia, G6PD Cornell:erythrocyte, leukocyte and platelet studies. *Blood* 44: 277–284, 1974.221. Mills, G. C.; Alperin, J. B.; Trimmer, K. B.: Studies on variant glucose-6-phosphate dehydrogenase: G6PD Fort Worth. *Biochem. Med.* 13:264–275, 1975.222. Milner, G.; Delamore, I. W.; Yoshida, A.: G-6-PD Manchester: a new variant associated with chronic nonspherocytic hemolytic anemia. *Blood* 43:271–276, 1974.223. Miwa, S.; Fujii, H.: Molecular basis of erythroenzymopathies associated with hereditary hemolytic anemia: tabulation of mutant enzymes. *Am. J. Hemat.* 51: 122–132, 1996.224. Miwa, S.; Fujii, H.; Nakashima, K.; Miura, Y.; Yamada, K.; Hagiwara, T.; Fukuda, M.: Three new electrophoretically normal glucose-6-phosphatedehydrogenase variants associated with congenital nonspherocytic hemolytic anemia found in Japan: G6PD Ogikubo, Yokohama, and Akita. *Hum. Genet.* 45:11–17, 1978.225. Miwa, S.; Fujii, H.; Nakatsuji, T.;

Ishida, Y.; Oda, E.; Kaneto, A.; Motokawa, M.; Ariga, Y.; Fukuchi, S.; Sasai, S.; Hiraoka, K.; Kashii, H.; Kodama, T.: Four new electrophoretically slow-moving glucose-6-phosphate dehydrogenase variants associated with congenital nonspherocytic hemolytic anemia found in Japan: Gd(-) Kurume, Gd(-) Fukushima, Gd(-) Yamaguchi, and Gd(-) Wakayama. *Am. J. Hemat.* 5:131-138, 1978.226. Miwa, S.; Nakashima, K.; Ono, J.; Fujii, H.; Suzuki, E.: Three glucose 6-phosphate dehydrogenase variants found in Japan. *Hum. Genet.* 36:327-334, 1977.227. Miwa, S.; Ono, J.; Nakashima, K.; Abe, S.; Kageoka, T.; Shinohara, K.; Isobe, J.; Yamaguchi, H.: Two new glucose 6-phosphate dehydrogenase variants associated with congenital nonspherocytic hemolytic anemia found in Japan: Gd(-) Tokushima and Gd(-) Tokyo. *Am. J. Hemat.* 1:433-442, 1976.228. Modiano, G.; Battistuzzi, G.; Esan, G. J. F.; Testa, U.; Luzzatto, L.: Genetic heterogeneity of 'normal' human erythrocyte glucose-6-phosphate dehydrogenase: an isoelectrophoretic polymorphism. *Proc. Nat. Acad. Sci.* 76: 852-856, 1979.229. Mohrenweiser, H. W.; Neel, J. V.: Frequency of thermostability variants: estimation of total 'rare' variant frequency in human populations. *Proc. Nat. Acad. Sci.* 78: 5729-5733,

1981.230. Morelli, A.; Benatti, U.; Guida, L.; De Flora, A.: G6PD Cagliari: a new low activity glucose 6-phosphate dehydrogenase variant characterized by enhanced intracellular lability. Hum. Genet. 66: 62-65, 1984.231. Morisaki, T.; Fujii, H.; Takegawa, S.; Tani, K.; Hirono, A.; Takizawa, T.; Takahashi, K.; Shinogi, M.; Teshirogi, T.; Miwa, S.: G6PD Sendagi: a new glucose-6-phosphate dehydrogenase variant associated with congenital hemolytic anemia. Hum. Genet. 65: 214-215, 1983.232. Nafa, K.; Reghis, A.; Osmani, N.; Baghli, L.; Benabadji, M.; Kaplan, J.-C.; Villamy, T. J.; Luzzatto, L.: G6PD Aures: a new mutation (48 ile-to-thr) causing mild G6PD deficiency is associated with favism. Hum. Molec. Genet. 2: 81-82, 1993.233. Nagel, R. L.; Ranney, H. M.: Genetic epidemiology of structural mutations of the beta-globin gene. Semin. Hemat. 27: 342-359, 1990.234. Nakai, T.; Yoshida, A.: G6PD Heian. A glucose-6-phosphate dehydrogenase variant associated with hemolytic anemia found in Japan. Clin. Chim. Acta 51: 199-203, 1974.235. Nakashima, K.; Ono, J.; Abe, S.; Miwa, S.; Yoshida, A.: G6PD Ube: a glucose 6-phosphate dehydrogenase variant found in four unrelated Japanese families. Am. J. Hum. Genet. 29: 24-30, 1977.236. Nakatsuji, T.; Miwa, S.: Incidence and charac-



teristics of glucose-6-phosphatedehydrogenase variants in Japan. Hum. Genet. 51: 297-305, 1979.237. Nance, W. E.: Turner's syndrome, twinning, and an unusual variant of glucose-6-phosphate dehydrogenase. Am. J. Hum. Genet. 16: 380-392, 1964.238. Necheles, T. F.; Snyder, L. M.; Strauss, W.: Glucose-6-phosphatedehydrogenase Boston. A new variant associated with congenital nonspherocytic hemolytic disease. Humangenetik 13: 218-221, 1971.239. Niazi, G.; Adeyokunu, A.; Westwood, B.; Beutler, E.: G6PD Aures: a rare mutant of G6PD in Saudi Arabia: molecular and clinical presentations. Saudi Med. J. 17: 311-314, 1996.240. Ninfali, P.; Baronciani, L.; Bardoni, A.; Bresolin, N.: Muscle expression of glucose-6-phosphate dehydrogenase deficiency in different variants. Clin. Genet. 48: 232-237, 1995.241. Notaro, R.; Afolayan, A.; Luzzatto, L.: Human mutations in glucose-6-phosphate dehydrogenase reflect evolutionary history. FASEB J. 14: 485-494, 2000.242. Nowicki, L.; Strobel, S.; Martin, H.; Koschwitz, U.: Ueber eine neue erythrocytäre glucose-6-phosphatdehydrogenase Variante, Typ Frankfurt. Klin. Wschr. 52: 478-484, 1974.243. Nsouly, G. M.; Prchal, J. T.: Characterization of a new G6PD variant and its associated oxidative damage. Clin. Res. 29: 829, 1981.244.

O'Brien, S. J.: The extent and character of biochemical genetic variation in the domestic cat. *J. Hered.* 71: 2–8, 1980.245. Ogura, H.; Morisaki, T.; Tani, K.; Kanno, H.; Tsutsumi, H.; Takahashi, K.; Miyamori, T.; Fujii, H.; Miwa, S.: A new glucose-6-phosphatedehydrogenase variant (G6PD Tsukui) associated with congenital hemolytic anemia. *Hum. Genet.* 78: 369–371, 1988.246. Ohno, S.: *Sex Chromosomes and Sex-linked Genes*. Berlin and New York: Springer (pub.) 1967.247. Orzalesi, N.; Sorcinelli, R.; Guiso, G.: Increased incidence of cataracts in male subjects deficient in glucose-6-phosphate dehydrogenase. *Arch. Ophthalmol.* 99: 69–70, 1981.248. Othieno-Obel, A.: East African variant of glucose-6-phosphatedehydrogenase. *East Afr. Med. J.* 49: 230–234, 1972.249. Pai, G. S.; Sprenkle, J. A.; Do, T. T.; Marenzi, C. E.; Migeon, B. R.: Localization of loci for hypoxanthine phosphoribosyltransferase and glucose-6-phosphate dehydrogenase and biochemical evidence of nonrandom X chromosome expression from studies of a human X-autosome translocation. *Proc. Nat. Acad. Sci.* 77: 2810–2813, 1980.250. Panich, V.: G6PD variants in Lao-tians. *Hum. Hered.* 24: 285–290, 1974.251. Panich, V.: G6PD Intanon, a new glucose 6-phosphate dehydrogenase-

sevariant. Humangenetik 21: 203–205, 1974.252. Panich, V.: G6PD characterization in Thailand. Genetics 74(suppl.): s208 only, 1973.253. Panich, V.: Glucose–6–phosphate dehydrogenase in Thailand. Hum.Genet. 53: 227–228, 1980.254. Panich, V.; Bumrungtrakul, P.; Jitjai, C.; Kamolmatayakul, S.; Khoprasert, B.; Klaisuvan, C.; Kongmuang, U.; Maneechai, P.; Pornpatkul, M.; Ruengrairatanaroje, P.; Surapruk, P.; Viriyayudhakorn, S.: Glucose–6–phosphatedehydrogenase deficiency in South Vietnamese. Hum. Hered. 30: 361–364, 1980.255. Panich, V.; Na–Nakorn, S.: G6PD variants in Thailand. J. Med.Assoc. Thai. 63: 537–543, 1980.256. Panich, V.; Sungnate, T.: Characterization of glucose 6–phosphatedehydrogenase in Thailand: the occurrence of 6 variants among 50 G6PDdeficient Thai. Humangenetik 18: 39–46, 1973.257. Panich, V.; Sungnate, T.; Na–Nakorn, S.: Acute intravascularhemolysis and renal failure in a new glucose 6–phosphate dehydrogenasevariant: G6PD Siriraj. J. Med. Assoc. Thai. 55: 726–731, 1972.258. Panich, V.; Sungnate, T.; Wasi, P.; Na–Nakorn, S.: G–6–PD Mahidol: the most common glucose–6–phosphate dehydrogenase variant in Thailand. J. Med. Assoc. Thai. 55: 576–585, 1972.259. Pawlak, A. L.; Mazurkiewicz, C. A.; Ordynski, J.;

Ruzynkova,D.; Horst, A.: G6PD Poznan, variant with severe enzyme deficiency. *Humangenetik* 28:163–165, 1975.260. Pawlak, A. L.; Zagorski, Z.; Ruzynkova, D.; Horst, A.: Polish variant of glucose–6–phosphate dehydrogenase (G–6–PD Lublin). *Humangenetik* 10:340–343, 1970.261. Pekrun, A.; Eber, S. W.; Schroter, W.: G6PD Avenches and G6PDMoosburg: biochemical and erythrocyte membrane characterization. *Blut* 58:11–14, 1989.262. Perroni, L.; Tassara, P.; Baldi, M.; Reali, R.; Scartezzini,P.: G6PD variants detected in Genoa area.In: Weatherall, D. J.; Fiorelli,G.; Gorini, S.: *Advances in Red Blood Cell Biology*. New York: RavenPress (pub.) 1982. Pp. 409–416.263. Persico, M. G.; Toniolo, D.; Nobile, C.; D'Urso, M.; Luzzatto,L.: cDNA sequences of human glucose 6–phosphate dehydrogenase cloned in pBR322. *Nature* 294: 778–780, 1981.264. Persico, M. G.; Viglietto, G.; Martini, G.; Toniolo, D.; Paonessa,G.; Moscatelli, C.; Dono, R.; Vulliamy, T.; Luzzatto, L.; D'Urso,M.: Isolation of human glucose–6–phosphate dehydrogenase (G6PD) cDNA-clones: primary structure of the protein and unusual 5–prime non–coding region. *Nucleic Acids Res.* 14: 2511–2522, 1986.265. Picat, C.; Etiemble, J.; Boivin, P.; Le Prise, P.–Y.: Gd(–)Rennes, a new deficient variant of glu–

cose-6-phosphate dehydrogenase associated with congenital nonspherocytic hemolytic anemia found in France. Hum. Genet. 55: 125-127, 1980.266. Pinto, P. V. C.; Newton, W. A., Jr.; Richardson, K. E.: Evidence for four types of erythrocyte glucose-6-phosphate dehydrogenase from G-6-PD deficient human subjects. J. Clin. Invest. 45: 823-831, 1966.267. Poggi, V.: Personal Communication. London, England 1989.268. Poon, M.-C.; Hall, K.; Scott, C. W.; Prchal, J. T.: G6PD Viangchan: a new glucose 6-phosphate dehydrogenase variant from Laos. Hum. Genet. 78:98-99, 1988.269. Porter, I. H.; Boyer, S. H.; Watson-Williams, E. J.; Adam, A.; Szeinberg, A.; Siniscalco, M.: Variation of glucose-6-phosphate dehydrogenase in different populations. Lancet I: 895-899, 1964.270. Porter, I. H.; Schulze, J.; McKusick, V. A.: Genetical linkage between the loci for glucose-6-phosphate dehydrogenase deficiency and colour-blindness in American Negroes. Ann. Hum. Genet. 26: 107-122, 1962.271. Prchal, J.; Moreno, H.; Conrad, M.; Vitek, A.: G-6-PD Dothan: a new variant associated with chronic hemolytic anemia. I.R.C.S. 7:348, 1979.272. Prchal, J. T.: Personal Communication. Birmingham, Ala. 1985.273. Prchal, J. T.; Crist, W. M.; Malluh, A.; Vitek, A.; Tauxe, W.N.; Carroll, A. J.: A new

glucose-6-phosphate dehydrogenase deficient variant in a patient with Chediak-Higashi syndrome. *Blood* 56: 476-480, 1980.274. Prchal, J. T.; Hall, K.; Csepregy, M.; Lilly, M.; Berkow, R.; Scott, C. W.: Two apparent glucose-6-phosphate dehydrogenase variants in normal XY males: G6PD Alabama. *Am. J. Med.* 84: 517-523, 1988.275. Pretsch, W.; Charles, D. J.; Merkle, S.: X-linked glucose-6-phosphate-dehydrogenase deficiency in *Mus musculus*. *Biochem. Genet.* 26: 89-103, 1988.276. Puck, J. M.; Willard, H. F.: X inactivation in females with X-linked disease. *New Eng. J. Med.* 338: 325-327, 1998.277. Ramot, B.; Ben-Bassat, I.; Shchory, M.: New glucose-6-phosphate dehydrogenase variants observed in Israel and their association with congenital nonspherocytic hemolytic disease. *J. Lab. Clin. Med.* 74: 895-901, 1969.278. Ramot, B.; Brok, F.: A new glucose-6-phosphate dehydrogenase mutant (Tel-Hashomer mutant). *Ann. Hum. Genet.* 28: 167-172, 1964.279. Rattazzi, M. C.; Corash, L. M.; Van Zzanen, G. E.; Jaffe, E. R.; Piomelli, S.: G6PD deficiency and chronic hemolysis: four new mutants--relationships between clinical syndrome and enzyme kinetics. *Blood* 38: 205-218, 1971.280. Rattazzi, M. C.; Lenzerini, L.; Meera Khan, P.; Luzzatto, L.:

Characterization of glucose-6-phosphate dehydrogenase variants.II. G6PD Kephallonia, G6PD Attica, and G6PD 'Seattle-like' found inGreece. Am. J. Hum. Genet. 21: 154-167, 1969.281. Ravindranath, Y.; Beutler, E.: Two new variants of glucose-6-phosphatedehydrogenase associated with hereditary non-spherocytic hemolyticanemia: G6PD Wayne and G6PD Huron. Am. J. Hemat. 24: 357-363, 1987.282. Reys, L.; Manso, C.; Stamatoyannopoulos, G.: Genetic studieson Southeastern Bantu of Mozambique. I. Variants of glucose-6-phosphatedehydrogenase. Am. J. Hum. Genet. 22: 203-215, 1970.283. Rinaldi, A.; Filippi, G.; Siniscalco, M.: Variability of redcell phenotypes between and within individuals in an unbiased sampleof 77 heterozygotes for G6PD deficiency in Sardinia. Am. J. Hum.Genet. 28: 496-505, 1976.284. Rosenstrauss, M.; Chasin, L. A.: Isolation of mammalian cellmutants deficient in glucose-6-phosphate dehydrogenase activity: linkage to hypoxanthine phosphoribosyl transferase. Proc. Nat. Acad. Sci. 72:493-497, 1975.285. Roth, E. F., Jr.; Raventos-Suarez, C.; Rinaldi, A.; Nagel, R.L.: Glucose-6-phosphate dehydrogenase deficiency inhibits in vitro growth of Plasmodium falciparum. Proc. Nat. Acad. Sci. 80: 298-299,1983.286. Roychoudhury, A. K.; Nei, M.: Hu-

man Polymorphic Genes: WorldDistribution. New York:  
 Oxford Univ. Press (pub.) 1988.287. Ruwando, C.; Khea, S.  
 C.; Snow, R. W.; Yates, S. N. R.; Kwiatkoweld,D.; Gupta, S.;  
 Warn, P.; Alisopp, G. E. M.; Gilbert, S. C.; Peschu,N.; New-  
 bold, C. I.; Greenwood, S. M.; Marsh, K.; Hill, A. V.  
 S.:Natural selection of hemi- and heterozygotes for G6PD  
 deficiency inAfrica by resistance to severe malaria. Nature  
 376: 246–249, 1995.288. Saenz, G. F.; Chaves, M.;  
 Berrantes, A.; Elizondo, J.; Montero,A. G.; Yoshida, A.: A  
 glucose–6–phosphate dehydrogenase variant,Gd(–) Santa-  
 maria found in Costa Rica. Acta Haemat. 72: 37–40,  
 1984.289. Saidi, N.; Hors–Cayla, M. C.; Van Cong, N.;  
 Benne, F.: Sheepline mapping by somatic cell hybridiza-  
 tion. (Abstract) Cytogenet.Cell Genet. 25: 200, 1979.290.  
 Samuel, A. P. W.; Saha, N.; Omer, A.; Hoffbrand, A. V.:  
 Quantitativeexpression of G6PD activity of different phe-  
 notypes of G6PD and haemoglobin in a Sudanese popula-  
 tion. Hum. Hered. 31: 110–115, 1981.291. Sansone, G.;  
 Perroni, L.; Testa, U.; Marenzi, C.; Luzzatto, L.: New genetic  
 variants of glucose 6–phosphate dehydrogenase (G6PD)in  
 Italy. Ann. Hum. Genet. 45: 97–104, 1981.292. Sansone,  
 G.; Perroni, L.; Yoshida, A.: Glucose–6–phosphate dehy-  
 drogenasevariants from Italian subjects associated with



severe neonatal jaundice. Brit.J. Haemat. 31: 159–165, 1975.293. Sansone, G.; Perroni, L.; Yoshida, A.; Dave, V.: A new glucose6–phosphate dehydrogenase variant (Gd Trinacria) in two unrelated families of Sicilian ancestry. Ital. J. Biochem. 26: 44–50, 1977.294. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Annenkov, G. A.: A description of new mutant forms of erythrocyte glucose–6–phosphate dehydrogenase isolated at the territory of the Soviet Union. Genetika 11:116–122, 1975.295. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Idelson, L. J.: The new form of glucose 6–phosphate dehydrogenase (G6PD 'Kaluga') from erythrocytes of a patient with chronic non–spherocytic hemolytic anemia. Vopr.Med. Khim. 22: 764–768, 1976.296. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Idelson, L. J.: Mutant forms of erythrocyte glucose 6–phosphate dehydrogenase in Ashkenazi: description of two new variants, G6PD Kirovograd and G6PD Zhitomir. Humangenetik 33:175–178, 1976.297. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Tzoneva, M.; Mavrudieva, M.; Toncheva, D.: Variants of erythrocyte glucose–6–phosphate dehydrogenase (G6PD) in Bulgarian populations. Hum. Genet. 54: 115–117, 1980.298. Shatskaya, T. L.; Krasnopol'skaya, K. D.; Zakharova, T. V.: Regularities of

distribution of Gd-alleles in Azerbaijan. II. Identification of G6PD mutant forms. *Genetika* 16: 2217–2225, 1980.299. Shows, T. B.; Brown, J. A.: Human X-linked genes regionally mapped utilizing X-autosome translocations and somatic cell hybrids. *Proc. Nat. Acad. Sci.* 72: 2125–2129, 1975.300. Shows, T. B.; Brown, J. A.; Chapman, V. M.: Comparative genemapping of HPRT, G6PD and PGK in man, mouse, and Muntjac deer. *Birth Defects Orig. Art. Ser. XII(7)*: 436–439, 1976.301. Shows, T. B.; Tashian, R. E.; Brewer, G. J.: Erythrocyte glucose-6-phosphate dehydrogenase in Caucasians: new inherited variant. *Science* 145:1056–1057, 1964.302. Sidi, Y.; Aderka, D.; Brok-Simoni, F.; Benjamin, D.; Ramot, B.; Pinkhas, J.: Viral hepatitis with extreme hyperbilirubinemia, massive hemolysis and encephalopathy in a patient with a new G6PD variant. *Israel J. Med. Sci.* 16: 130–133, 1980.303. Siegel, N. H.; Beutler, E.: Hemolytic anemia caused by G-6-PD Carswell, a new variant. *Ann. Intern. Med.* 75: 437–439, 1971.304. Smith, J. E.; Ryer, K.; Wallace, L.: Glucose-6-phosphate dehydrogenase deficiency in a dog. *Enzyme* 21: 379–382, 1976.305. Snyder, L. M.; Necheles, T. F.; Reddy, W. J.: G-6-PD Worcester: a new variant, associated with X-linked optic atrophy. *Am. J.*

Med. 49:125–132, 1970.306. Stamatoyannopoulos, G.; Kotsakis, P.; Voigtlander, V.; Akrivakis, A.; Motulsky, A. G.: Electrophoretic diversity of glucose–6–phosphatedehydrogenase among Greeks. Am. J. Hum. Genet. 22: 587–596, 1970.307. Stamatoyannopoulos, G.; Voigtlander, V.; Akrivakis, A.: Thessaly variant of glucose–6–phosphate dehydrogenase. Humangenetik 9: 23–25, 1970.308. Stamatoyannopoulos, G.; Voigtlander, V.; Kotsakis, P.; Akrivakis, A.: Genetic diversity of the 'Mediterranean' glucose–6–phosphatedehydrogenase deficiency phenotype. J. Clin. Invest. 50: 1253–1261, 1971.309. Stamatoyannopoulos, G.; Yoshida, A.; Bacopoulos, C.; Motulsky, A. G.: Athens variant of glucose–6–phosphate dehydrogenase. Science 157:831–833, 1967.310. Stevens, D. J.; Wanachiwanawin, W.; Mason, P. J.; Vulliamy, T.J.; Luzzatto, L.: G6PD Canton a common deficient variant in SouthEast Asia caused by a 459 arg–to–leu mutation. Nucleic Acids Res. 18:7190, 1990.311. Stocco dos Santos, R. C.; Barretto, O. C. O.; Nonoyama, K.; Castro, N. H. C.; Ferraz, O. P.; Walter–Moura, J.; Vescio, C. C. S.; Becak, W.: X–linked syndrome: mental retardation, hip luxation, and G6PD variant (Gd(+)) Butantan. Am. J. Med. Genet. 39: 133–136, 1991.312. Stockham, S. L.;

Harvey, J. W.; Kinden, D. A.: Equine glucose-6-phosphatedehydrogenase deficiency. *Vet. Path.* 31: 518-527, 1994.313. Streiff, F.; Vigneron, C.: Anemie hemolytique chronique par deficit en glucose 6-phosphate deshydrogenase dans une famille d'origineLorraine. *Nouv. Rev. Franc. Hemat.* 11: 279-290, 1971.314. Takahashi, K.; Fujii, H.; Takegawa, S.; Tani, K.; Hirono, A.;Takizawa, T.; Kawakatsu, T.; Miwa, S.: A new glucose-6-phosphatedehydrogenase variant (G6PD Nagano) associated with congenital hemolyticanemia. *Hum. Genet.* 62: 368-370, 1982.315. Takizawa, T.; Fujii, H.; Takegawa, S.; Takahashi, K.; Hirono,A.; Morisaki, T.; Kanno, H.; Oka, R.; Yoshioka, H.; Miwa, S.: A uniqueelectrophoretic slow-moving glucose 6-phosphate dehydrogenase variant(G6PD Asahikawa) with a markedly acidic pH optimum. *Hum. Genet.* 68:70-72, 1984.316. Takizawa, T.; Huang, I.-Y.; Ikuta, T.; Yoshida, A.: Human glucose-6-phosphatedehydrogenase: primary structure and cDNA cloning. *Proc. Nat. Acad.Sci.* 83: 4157-4161, 1986.317. Takizawa, T.; Yoneyama, Y.; Miwa, S.; Yoshida, A.: A single nucleotide base transition is the basis of the common human glucose-6-phosphatedehydrogenase variant A(+). *Genomics* 1: 228-231, 1987.318. Takizawa, T.; Yoshida,

A.: Molecular abnormality of the common glucose-6-phosphate dehydrogenase variant, G6PD A(+), and restriction-fragment-length polymorphism. (Abstract) Am. J. Hum. Genet. 41: A241, 1987.319. Talalak, P.; Beutler, E.: G-6-PD Bangkok: a new variant found in congenital non-spherocytic hemolytic disease (CNHD). Blood 33:772-776, 1969.320. Tanaka, K. R.; Beutler, E.: Hereditary hemolytic anemia due to glucose-6-phosphate dehydrogenase Torrance: a new variant. J. Lab. Clin. Med. 73: 657-667, 1969.321. Tang, T. K.; Huang, C.-S.; Huang, M. J.; Tam, K.-B.; Yeh, C.-H.; Tang, C.-J. C.: Diverse point mutations result in glucose-6-phosphate dehydrogenase (G6PD) polymorphism in Taiwan. Blood 79: 2135-2140, 1992.322. Testa, U.; Meloni, T.; Lania, A.; Battistuzzi, G.; Cutillo, S.; Luzzatto, L.: Genetic heterogeneity of glucose-6-phosphate dehydrogenase deficiency in Sardinia. Hum. Genet. 56: 99-105, 1980.323. Thigpen, J. T.; Steinberg, M. H.; Beutler, E.; Gillespie, G.T., Jr.; Dreiling, B. J.; Morrison, F. S.: Glucose-6-phosphate dehydrogenase Jackson, a new variant associated with hemolytic anemia. Acta Haemat. 51:310-314, 1974.324. Tishkoff, S. A.; Varkonyi, R.; Cahinhinan, N.; Abbes, S.; Argyropoulos, G.; Destro-Bisol, G.; Drousiotou, A.; Dangerfield, B.; Lefranc,

G.;Loiselet, J.; Piro, A.; Stoneking, M.; Tagarelli, A.; Tagarelli, G.;Touma, E. H.; Williams, S. M.; Clark, A. G.: Haplotype diversity and linkage disequilibrium at human G6PD: recent origin of alleles that confer malarial resistance. *Science* 293: 455–462, 2001.325. Tokarev, Y. N.; Chernyak, N. B.; Batischev, A. I.; Lanzina, N.V.; Alexeyev, G. A.: Etude des propriétés électrophorétiques et cinétiques de la glucose-6-phosphate déshydrogénase (Gd) d'érythrocytes dans les déficits héréditaires de l'enzyme: description d'une nouvelle variante de glucose-6-phosphate déshydrogénase: la Gd Kremenchug. *Nouv.Rev. Franc. Hemat.* 20: 557–564, 1978.326. Toncheva, D.: Variants of glucose-6-phosphate dehydrogenase in a Vietnamese population. *Hum. Hered.* 36: 348–351, 1986.327. Toncheva, D.; Tzoneva, M.: Genetic polymorphism of G6PD in a Bulgarian population. *Hum. Genet.* 67: 340–342, 1984.328. Toniolo, D.; Martini, G.; Migeon, B. R.; Dono, R.: Expression of the G6PD locus on the human X chromosome is associated with demethylation of three CpG islands within 100 kb of DNA. *EMBO J.* 7: 401–406, 1988.329. Town, M.; Athanasiou-Metaxa, M.; Luzzatto, L.: Intragenic interspecific complementation of glucose 6-phosphate dehydrogenase in human-ham-

stercell hybrids. *Somat. Cell Molec. Genet.* 16: 97–108, 1990.330. Town, M.; Bautista, J. M.; Mason, P. J.; Luzzatto, L.: Both mutations in G6PD A- are necessary to produce the G6PD deficient phenotype. *Hum. Molec. Genet.* 1: 171–174, 1992.331. Usanga, E. A.; Bienzle, U.; Cancedda, K.; Fasuan, F. A.; Ajayi, O.; Luzzatto, L.: Genetic variants of human erythrocyte glucose 6-phosphate dehydrogenase: new variants in West Africa characterized by column chromatography. *Ann. Hum. Genet.* 40: 279–286, 1977.332. Vaca, G.; Ibarra, B.; Garcia Cruz, D.; Medina, C.; Romero, F.; Cantu, J. M.; Beutler, E.: G-6-PD Jalisco and G-6-PD Morelia: two new Mexican variants. *Hum. Genet.* 71: 82–85, 1985.333. Vaca, G.; Ibarra, B.; Romero, F.; Olivares, N.; Cantu, J. M.; Beutler, E.: G-6-PD Guadalajara: a new mutant associated with chronic nonspherocytic hemolytic anemia. *Hum. Genet.* 61: 175–176, 1982.334. Ventura, A.; Panizon, F.; Soranzo, M. R.; Veneziano, G.; Sansone, G.; Testa, U.; Luzzatto, L.: Congenital dyserythropoietic anaemia type II associated with a new type of G6PD deficiency (G6PD Gabrovizza). *Acta Haemat.* 71: 227–234, 1984.335. Vergnes, H.; Gherardi, M.; Bouloux, C.: Erythrocyte glucose-6-phosphate dehydrogenase in the Niokolonko (Malinke of the Niokolo) of the Eastern Senegal.

gal: identification of a slow variant with normal activity (Tacoma-like). Hum.Hered. 25: 80-87, 1975.336.

Vergnes, H.; Gherardi, M.; Quilici, J. C.; Yoshida, A.; Giaccardy, R.: G6PD Luz-Saint-Sauveur: a new variant with abnormal electrophoretic mobility, mild enzyme deficiency and absence of hemolytic disorders. I.R.C.S. 7:14,

1973.337. Vergnes, H.; Gherardi, M.; Yoshida, A.: G6PD Lozere and Trinacria-like: segregation of two non-

hemolytic variants in a French family. Hum.Genet. 34:

293-298, 1976.338. Vergnes, H.; Ribet, A.; Bommelaer, G.; Amadiou, J.; Brun, H.: GD(-) Muret and GD(-)

Colomiers, two new variants of glucose-

6-phosphatedehydrogenase associated with favism. Hum.

Genet. 57: 332-334, 1981.339. Vergnes, H.; Yoshida, A.;

Gourdin, D.; Gherardi, M.; Bierme, R.; Ruffie, J.: Glucose

6-phosphate dehydrogenase Toulouse: a new variant with

marked instability and severe deficiency discovered in a

family of Mediterranean ancestry. Acta Haemat. 51:

240-249, 1974.340. Viglietto, G.; Montanaro, V.; Calabro,

V.; Vallone, D.; D'Urso, M.; Persico, M. G.; Battistuzzi, G.:

Common glucose-6-phosphate dehydrogenase(G6PD)

variants from the Italian population: biochemical and

molecular characterization. Ann. Hum. Genet. 54: 1-15,



1990.341. Vives-Corróns, J.-L.; Kuhl, W.; Pujades, M. A.; Beutler, E.:Molecular genetics of the glucose-6-phosphate dehydrogenase (G6PD)Mediterranean variant and description of a new G6PD mutant, G6PD Andalus(1361A). Am.J. Hum. Genet. 47: 575-579, 1990.342. Vives-Corróns, J. L.; Feliu, E.; Pujades, M. A.; Cardellach,F.; Rozman, C.; Carreras, A.; Jou, J. M.; Vallespi, M. T.; Zuazu,F. J.: Severe glucose-6-phosphate dehydrogenase (G6PD) deficiencyassociated with chronic hemolytic anemia, granulocyte dysfunction,and increased susceptibility to infections: description of a new molecularvariant (G6PD Barcelona). Blood 59: 428-434, 1982.343. Vives-Corróns, J. L.; Pujades, A.: Heterogeneity of 'Mediterranean type' glucose-6-phosphate dehydrogenase (G6PD) deficiency in Spainand description of two new variants associated with favism. Hum.Genet. 60: 216-221, 1982.344. Vives-Corróns, J. L.; Pujades, A.; Curia, M. D.: Caracterizacimolecular de la glucosa-6-fosfato deshidrogenasa (G6PD) en 24 casosde deficit enzimatico y descripcion de una nueva variante (G6PD-Betica). Sangre 25:1049-1064, 1980.345. Vives-Corróns, J. L.; Pujades, M. A.; Petit, J.; Colomer, D.;Corbella, M.; Aguilar i Bascompte, J. L.; Merino, A.: Chronic nonspherocyctic hemolytic anemia

(CNSHA) and glucose 6 phosphate dehydrogenase (G6PD) deficiency in a patient with familial amyloidotic polyneuropathy (FAP): molecular study of a new variant (G6PD Clinic) with markedly acidic pH optimum. Hum. Genet. 81: 161–164, 1989.346. Vulliamy, T.; Beutler, E.; Luzzatto, L.: Variants of glucose-6-phosphate dehydrogenase are due to missense mutations spread throughout the coding region of the gene. Hum. Mutat. 2: 159–167, 1993.347. Vulliamy, T.; Mason, P.; Luzzatto, L.: The molecular basis of glucose-6-phosphate dehydrogenase deficiency. Trends Genet. 8: 138–143, 1992.348. Vulliamy, T.; Rovira, A.; Yusoff, N.; Colomer, D.; Luzzatto, L.; Vives-Corrons, J.-L.: Independent origin of single and double mutations in the human glucose 6-phosphate dehydrogenase gene. Hum. Mutat. 8: 311–318, 1996.349. Vulliamy, T. J.: Personal Communication. London, England 1989.350. Vulliamy, T. J.; D'Urso, M.; Battistuzzi, G.; Estrada, M.; Foulkes, N. S.; Martini, G.; Calabro, V.; Poggi, V.; Giordano, R.; Town, M.; Luzzatto, L.; Persico, M. G.: Diverse point mutations in the human glucose-6-phosphate dehydrogenase gene cause enzyme deficiency and mild or severe hemolytic anemia. Proc. Nat. Acad. Sci. 85: 5171–5175, 1988.351. Vulliamy, T. J.;

Kaeda, J. S.; Ait-Chafa, D.; Mangerini, R.; Roper, D.; Barbot, J.; Mehta, A. B.; Athanassiou-Metaxa, M.; Luzzatto, L.; Mason, P. J.: Clinical and haematological consequences of recurrent G6PD mutations and a single new mutation causing chronic nonspherocytic haemolytic anaemia. *Brit. J. Haemat.* 101: 670–675, 1998.352. Vulliamy, T. J.; Othman, A.; Town, M.; Nathwani, A.; Falusi, A. G.; Mason, P. J.; Luzzatto, L.: Polymorphic sites in the African population detected by sequence analysis of the glucose-6-phosphate dehydrogenase gene outline the evolution of the variants A and A-. *Proc. Nat. Acad. Sci.* 88: 8568–8571, 1991.353. Vulliamy, T. J.; Wanachiwanawin, W.; Mason, P. J.; Luzzatto, L.: G6PD Mahidol, a common deficient variant in South East Asia is caused by a (163)glycine-to-serine mutation. *Nucleic Acids Res.* 17:5868, 1989.354. Vuopio, P.; Harkonen, M.; Helske, T.; Naeveri, H.: Red cell glucose-6-phosphate dehydrogenase deficiency in Finland: characterization of a new variant with severe enzyme deficiency. *Scand. J. Haemat.* 15:145–152, 1975.355. Vuopio, P.; Harkonen, M.; Johnsson, P.; Nuutinen, M.: Red cell glucose-phosphate dehydrogenase deficiency in Finland. *Ann. Clin. Res.* 5: 168–178, 1973.356. Waitz, R.; Boivin, P.; Oberling, F.; Casenave, J. P.; North, M.

L.; Mayer, S.: Variante Gd(-) Strasbourg de la glucose-6-phosphatedehydrogenase. *Nouv. Rev. Franc. Hemat.* 10: 312-314, 1970.357. Wang, Y. M.; Patterson, J. H.; Van Eys, J.: The potential use of xylitol in glucose-6-phosphate dehydrogenase deficiency anemia. *J. Clin. Invest.* 50: 1421-1428, 1971.358. Weimer, T. A.; Salzano, F. M.; Hutz, M. H.: Erythrocyte isozymes and hemoglobin types in a southern Brazilian population. *J. Hum. Evol.* 10: 319-328, 1981.359. Weimer, T. A.; Schuler, L.; Beutler, E.; Salzano, F. M.: Gd(+)Laguna, a new rare glucose-6-phosphate dehydrogenase variant from Brazil. *Hum. Genet.* 65: 402-404, 1984.360. Weinreich, J.; Busch, D.; Gottstein, U.; Schaefer, J.; Rohr, J.: Ueber zwei neue Faelle von hereditaerer nicht-sphaerocytaerer haemolytischer Anaemie bei glucose-6-phosphat-dehydrogenase-Defekt in einer Nord Deutschen Familie. *Klin. Wschr.* 46: 146-149, 1968.361. Welch, S. G.; McGregor, I. A.; Williams, K.: A new variant of human erythrocyte G6PD occurring at a high frequency amongst the population of two villages in The Gambia, West Africa. *Hum. Genet.* 40: 305-309, 1978.362. Westring, D. W.; Pisciotto, A. V.: Anemia, cataracts, and seizures in patients with glucose-6-phosphate dehydroge-

nase deficiency. Arch.Intern. Med. 118: 385–390, 1966.363. WHO: Nomenclature of glucose–6–phosphate dehydrogenase in man. Bull. WHO 36: 319–322, 1967. Note: See Also: Canad. Med. Assoc.J. 97: 422–424, 1967.364. WHO: Scientific group on the standardization of procedures for the study of glucose–6–phosphate dehydrogenase. WHO Techn. Rep.(pub.) Ser. No. 366: 1967.365. Wilson, W. W.: Congenital hemolytic anemia due to a deficiency of glucose 6–phosphate dehydrogenase. Rocky Mt. Med. J. 73: 160–162, 1976.366. Witt, I.; Yoshioka, S.: Biochemical characterization of a glucose–6–phosphate dehydrogenase variant with favism: G–6–PD Zaehringen. Klin. Wschr. 50:205–209, 1972.367. Wong, P. W. K.; Shih, L.–Y.; Hsia, D. Y. Y.: Characterization of glucose–6–phosphate dehydrogenase among Chinese. Nature 208:1323–1324, 1965.368. Yermakov, N.; Tokarev, J.; Chernjak, N.; Schoenian, G.; Grieger, M.; Guckler, G.; Jacobasch, G.; Mahmudova, M.; Bahramov, S.: New stable mutant Gd(–) variants: G6PD Tashkent and G6PD Nucus: molecular basis of hereditary enzyme deficiency. Acta Biol. Med. Ger. 40:559–562, 1981.369. Yoshida, A.: Personal Communication. Duarte, Calif. 2/26/1996.370. Yoshida, A.: Amino acid substitution (histidine to tyrosine) in a glu–

cose-6-phosphatate dehydrogenase variant (G6PD Hek-  
toen) associated with over-production. J. Mol. Biol. 52:  
483-490, 1970.371. Yoshida, A.: A single amino acid  
substitution (asparagine to aspartic acid) between normal  
(B plus) and the common Negro variant (A plus) of human  
glucose-6-phosphate dehydrogenase. Proc. Nat. Acad. Sci.  
57: 835-840, 1967.372. Yoshida, A.: Human glucose-  
6-phosphate dehydrogenase: purification and characteri-  
zation of Negro type variant (A+) and comparison with-  
normal enzyme (B+). Biochem. Genet. 1: 81-99,  
1967.373. Yoshida, A.; Baur, E. W.; Motulsky, A. G.: A  
Philippino glucose-6-phosphate dehydrogenase variant  
(G6PD Union) with enzyme deficiency and altered substrate  
specificity. Blood 35: 506-513, 1970.374. Yoshida, A.;  
Beutler, E.: Human glucose-6-phosphate dehydrogenase-  
variants: a supplementary tabulation. Ann. Hum. Genet.  
41: 347-355, 1978.375. Yoshida, A.; Beutler, E.; Motulsky,  
A. G.: Table of human glucose-  
6-phosphate dehydrogenase variants. Bull. WHO 45:  
243-253, 1971.376. Yoshida, A.; Stamatoyannopoulos,  
G.; Motulsky, A. G.: Negro variant of glucose-6-phosphate  
dehydrogenase deficiency (A-) in man. Science  
155: 97-99, 1967.377. Yoshida, A.; Takizawa, T.: The

same extra FokI cleavage site exists in glucose-6-phosphate dehydrogenase variants A(+) and A(-). Am.J. Hum. Genet. 43: 131-133, 1988.378. Yoshida, A.; Takizawa, T.; Prchal, J. T.: RFLP of the X chromosome-linked glucose-6-phosphate dehydrogenase locus in blacks. Am. J. Hum. Genet. 42:872-876, 1988.379. Zinkham, W. H.: A deficiency of glucose-6-phosphate dehydrogenase activity in lens from individuals with primaquine-sensitive erythrocytes. Bull. Johns Hopkins Hosp. 109: 206-216, 1961.380. Zuo, L.; Chen, E.; Du, C. S.; Chang, C. N.; Chiu, D. T. Y.: Genetic study of Chinese G6PD variants by direct PCR sequencing. (Abstract) Blood 76(suppl. 1): 51A, 1990.

[74434] 12764. Campuzano, V.; Montermini, L.; Molto, M. D.; Piane, L.; Cossee, M.; Cavalcanti, F.; Monros, E.; Rodius, F.; Duclos, F.; Monticelli, A.; Zara, F.; Canizares, J.; Koutnikova, H.; Bidichandani, S. I.; Gellera, C.; Brice, A.; Trouillas, P.; De Michele, G.; Filla, A.; De Frutos, R.; Palau, F.; Patel, P. I.; Di Donato, S.; Mandel, J.-L.; Coccozza, S.; Koenig, M.; Pandolfo, M.: Friedreich's ataxia: autosomal recessive disease caused by an intronic GAA triplet repeat expansion. Science 271: 1423-1427, 1996.

[74435] 12765. Ilyin, G. P.; Riialand, M.; Pigeon, C.; Guguen-Guil-

louzo, C.:cDNA cloning and expression analysis of new members of the mammalianF-box protein family. Genomics 67: 40–47, 2000.

- [74436] 12766.Inase, N.; Fushimi, K.; Ishibashi, K.; Uchida, S.; Ichioka, M.;Sasaki, S.; Marumo, F.: Isolation of human aquaporin 3 gene. J.Biol. Chem. 270: 17913–17916, 1995.
- [74437] 12767.Ishibashi, K.; Sasaki, S.; Saito, F.; Ikeuchi, T.; Marumo, F.:Structure and chromosomal localization of a human water channel (AQP3)gene. Genomics 27: 352–354, 1995. Note: Erratum: Genomics 30: 633only, 1995.
- [74438] 12768.Ma, T.; Hara, M.; Sougrat, R.; Verbavatz, J.–M.; Verkman, A. S.: Impaired stratum corneum hydration in mice lacking epidermal waterchannel aquaporin–3. J. Biol. Chem. 277: 17147–17153, 2002.
- [74439] 12769.Ishibashi, K.; Sasaki, S.; Fushimi, K.; Uchida, S.; Kuwahara, M.;Saito, H.; Furukawa, T.; Nakajima, K.; Yamaguchi, Y.; Gojobori, T.;Marumo, F.: Molecular cloning and expression of a member of the aquaporinfamily with permeability to glycerol and urea in addition to waterexpressed at the basolateral membrane of kidney collecting duct cells. Proc.Nat. Acad. Sci. 91: 6269–6273, 1994.



- [74440] 12770.Ma, T.; Song, Y.; Yang, B.; Gillespie, A.; Carlson, E. J.; Epstein,C. J.; Verkman, A. S.: Nephrogenic diabetes insipidus in mice lackingaquaporin-3 water channels. *Proc. Nat. Acad. Sci.* 97: 4386–4391,2000.
- [74441] 12771.Mulders, S. M.; Olde Weghuis, D.; van Boxtel, J. A. F.; Geurtsvan Kessel, A.; Echevarria, M.; van Os, C. H.; Deen, P. M. T.: Localizationof the human gene for aquaporin 3 (AQP3) to chromosome 9, region p21–p12,using fluorescent in situ hybridization. *Cytogenet. Cell Genet.* 72:303–305, 1996.
- [74442] 12772.Sougrat, R.; Morand, M.; Gondran, C.; Barre, P.; Gobin, R.; Bonte,F.; Dumas, M.; Verbavatz, J.–M.: Functional expression of AQP3 inhuman skin epidermis and reconstructed epidermis. *J. Invest. Derm.* 118:678–685, 2002.
- [74443] 12773.Andersson, P.; McGuire, J.; Rubio, C.; Gradin, K.; Whitelaw, M.L.; Pettersson, S.; Hanberg, A.; Poellinger, L.: A constitutivelyactive dioxin/aryl hydrocarbon receptor induces stomach tumors. *Proc.Nat. Acad. Sci.* 99: 9990–9995, 2002.
- [74444] 12774.Ema, M.; Matsushita, N.; Sogawa, K.; Ariyama, T.; Inazawa, J.;Nemoto, T.; Ota, M.; Oshimura, M.; Fujii–Kuriyama, Y.: Human arylhydrocarbonreceptor: functional

expression and chromosomal assignment to 7p21.

J.Biochem. 116: 845–851, 1994.

- [74445] 12775. Le Beau, M. M.; Carver, L. A.; Espinosa, R., III; Schmidt, J. V.; Bradfield, C. A.: Chromosomal localization of the human AHR locus encoding the structural gene for the Ah receptor to 7p21–p15. Cytogenet. Cell Genet. 66: 172–176, 1994.
- [74446] 12776. Micka, J.; Milatovich, A.; Menon, A.; Grabowski, G. A.; Puga, A.; Nebert, D. W.: Human Ah receptor (AHR) gene: localization to 7p15 and suggestive correlation of polymorphism with CYP1A1 inducibility. Pharmacogenetics 7:95–101, 1997.
- [74447] 12777. Shimizu, Y.; Nakatsuru, Y.; Ichinose, M.; Takahashi, Y.; Kume, H.; Mimura, J.; Fujii–Kuriyama, Y.; Ishikawa, T.: Benzo[a]pyrenecarcinogenicity is lost in mice lacking the aryl hydrocarbon receptor. Proc. Nat. Acad. Sci. 97: 779–782, 2000.
- [74448] 12778. Adams, M. D.; Kelley, J. M.; Gocayne, J. D.; Dubnick, M.; Polymeropoulos, M. H.; Xiao, H.; Merril, C. R.; Wu, A.; Olde, B.; Moreno, R. F.; Kerlavage, A. R.; McCombie, W. R.; Venter, J. C.: Complementary DNA sequencing: expressed sequence tags and human genome project. Science 252: 1651–1656, 1991.

- [74449] 12779. Polvi, A.; Armstrong, E.; Lai, C.; Lemke, G.; Huebner, K.; Spritz, R. A.; Guida, L. C.; Nicholls, R. D.; Alitalo, K.: The human TYRO3 gene and pseudogene are located in chromosome 15q14–q25. *Gene* 134:289–293, 1993.
- [74450] 12780. Chen, P.; Hao, W.; Rife, L.; Wang, X. P.; Shen, D.; Chen, J.; Ogden, T.; Van Boemel, G. B.; Wu, L.; Yang, M.; Fong, H. K. W.: A photic visual cycle of rhodopsin regeneration is dependent on Rgr. *Nature Genet.* 28: 256–260, 2001.
- [74451] 12781. Chen, X.-N.; Korenberg, J. R.; Jiang, M.; Shen, D.; Fong, H. K. W.: Localization of the human RGR opsin gene to chromosome 10q23. *Hum. Genet.* 97: 720–722, 1996.
- [74452] 12782. Jiang, M.; Pandey, S.; Fong, H. K. W.: An opsin homologue in the retina and pigment epithelium. *Invest. Ophthalmol. Visual Sci.* 34:3669–3678, 1993.
- [74453] 12783. Morimura, H.; Saindelle-Ribeaudeau, F.; Berson, E. L.; Dryja, T. P.: Mutations in RGR, encoding a light-sensitive opsin homologue, in patients with retinitis pigmentosa. (Letter) *Nature Genet.* 23:393–394, 1999.
- [74454] 12784. Shen, D.; Jiang, M.; Hao, W.; Tao, L.; Salazar, M.; Fong, H. K. W.: A human opsin-related gene that encodes a retinaldehyde-binding protein. *Biochemistry* 33: 13117–13125, 1994.

- [74455] 12785.Asano, H.; Ishida, A.; Hasegawa, M.; Ono, T.; Yoshida, M. C.; Taniguchi,M.; Kanno, M.: The mouse Mel-18 'RING-finger' gene: genomic organization,promoter analysis and chromosomal assignment. DNA Seq. 3: 369-377,1993.
- [74456] 12786.Ishida, A.; Asano, H.; Hasegawa, M.; Koseki, H.; Ono, T.; Yoshida,M. C.; Taniguchi, M.; Kanno, M.: Cloning and chromosome mapping ofthe human Mel-18 gene which encodes a DNA-binding protein with a new'RING-finger' motif. Gene 129: 249-255, 1993.
- [74457] 12787.Tagawa, M.; Sakamoto, T.; Shigemoto, K.; Matsubara, H.; Tamura,Y.; Ito, T.; Nakamura, I.; Okitsu, A.; Imai, K.; Taniguchi, M.: Expressionof novel DNA-binding protein with zinc finger structure in varioustumor cells. J. Biol. Chem. 265: 20021-20026, 1990.
- [74458] 12788.Hara, E.; Yamaguchi, T.; Nojima, H.; Ide, T.; Campisi, J.; Okayama,H.; Oda, K.: Id-related genes encoding helix-loop-helix proteinsare required for G1 progression and are repressed in senescent humanfibroblasts. J. Biol. Chem. 269: 2139-2145, 1994.
- [74459] 12789.Mathew, S.; Chen, W.; Murty, V. V. V. S.; Benezra, R.; Chaganti,R. S. K.: Chromosomal assignment of human ID1 and ID2 genes. Genomics 30:385-387, 1995.

- [74460] 12790.Chen, M. X.; McPartlin, A. E.; Brown, L.; Chen, Y. H.; Barker,H. M.; Cohen, P. T. W.: A novel human protein serine/threonine phosphatase,which possesses four tetratri-copeptide repeat motifs and localizesto the nucleus. EMBO J. 13: 4278–4290, 1994.
- [74461] 12791.Xu, X.; Lagercrantz, J.; Zickert, P.; Bajalica–Lagercrantz, S.;Zetterberg, A.: Chromosomal localization and 5–prime sequence ofthe human protein serine/threonine phosphatase 5–prime gene. Biochem.Biophys. Res. Commun. 218: 514–517, 1996.
- [74462] 12792.Yong, W. H.; Ueki, K.; Chou, D.; Reeves, S. A.; von Deimling, A.;Gusella, J. F.; Mohrenweiser, H. W.; Buckler, A. J.; Louis, D. N.: Cloning of a highly conserved human protein serine–threonine phosphatasegene from the glioma candidate region on chromosome 19q13.3. Genomics 29:533–536, 1995.
- [74463] 12793.Breitbart, R. E.; Liang, C.; Smoot, L. B.; Laheru, D. A.; Mahdavi,V.; Nadal–Ginard, B.: A fourth human MEF2 transcription factor, hMEF2D,is an early marker of the myogenic lineage. Development 118: 1095–1106,1993.
- [74464] 12794.Hobson, G. M.; Krahe, R.; Garcia, E.; Siciliano, M. J.; Funanage,V. L.: Regional chromosomal assignments for four members of the MADSdomain transcription enhancer

factor 2 (MEF2) gene family to human chromosomes 15q26, 19p12, 5q14, and 1q12–q23. *Genomics* 29: 704–711, 1995.

- [74465] 12795. Mao, Z.; Bonni, A.; Xia, F.; Nadal-Vicans, M.; Greenberg, M. E.: Neuronal activity-dependent cell survival mediated by transcription factor MEF2. *Science* 286: 785–790, 1999.
- [74466] 12796. Martin, J. F.; Miano, J. M.; Hustad, C. M.; Copeland, N. G.; Jenkins, N. A.; Olson, E. N.: A Mef2 gene that generates a muscle-specific isoform via alternative mRNA splicing. *Molec. Cell. Biol.* 14: 1647–1656, 1994.
- [74467] 12797. Pollock, R.; Treisman, R.: Human SRF-related proteins: DNA-binding properties and potential regulatory targets. *Genes Dev.* 5: 2327–2341, 1991.
- [74468] 12798. Yu, Y.-T.; Breitbart, R. E.; Smoot, L. B.; Lee, Y.; Mahdavi, V.; Nadal-Ginard, B.: Human myocyte-specific enhancer factor 2 comprises a group of tissue-restricted MADS box transcription factors. *Genes Dev.* 6: 1783–1798, 1992.
- [74469] 12799. Youn, H.-D.; Sun, L.; Prywes, R.; Liu, J. O.: Apoptosis of T cells mediated by  $\text{Ca}^{2+}$ -induced release of the transcription factor MEF2. *Science* 286: 790–793, 1999.
- [74470] 12800. Krainc, D.; Haas, M.; Ward, D. C.; Lipton, S. A.;

Bruns, G.; Leifer, D.: Assignment of human myocyte-specific enhancer binding factor 2C (hMEF2C) to human chromosome 5q14 and evidence that MEF2C is evolutionarily conserved. *Genomics* 29: 809–811, 1995.

[74471] 12801. Leifer, D.; Krainc, D.; Yu, Y.-T.; McDermott, J.; Breitbart, R.E.; Heng, J.; Neve, R. L.; Kosofsky, B.; Nadal-Ginard, B.; Lipton, S. A.: MEF2C, a MADS/MEF2-family transcription factor expressed in a laminar distribution in cerebral cortex. *Proc. Nat. Acad. Sci.* 90:1546–1550, 1993.

[74472] 12802. Greengard, P.; Valtorta, F.; Czernik, A. J.; Benfenati, F.: Synaptic vesicle phosphoproteins and regulation of synaptic function. *Science* 259:780–785, 1993.

[74473] 12803. Li, L.; Chin, L.-S.; Greengard, P.; Copeland, N. G.; Gilbert, D.J.; Jenkins, N. A.: Localization of the synapsin II (SYN2) gene to human chromosome 3 and mouse chromosome 6. *Genomics* 28: 365–366, 1995.

[74474] 12804. McCright, B.; Rivers, A. M.; Audlin, S.; Virshup, D. M.: The B56 family of protein phosphatase 2A (PP2A) regulatory subunits encodes differentiation-induced phosphoproteins that target PP2A to both nucleus and cytoplasm. *J. Biol. Chem.* 271: 22081–22089, 1996.

[74475] 12805. Van Hoof, C.; Aly, M. S.; Garcia, A.; Cayla, X.; Cas-

siman, J. J.;Merlevede, W.; Goris, J.: Structure and chromosomal localization of the human gene of the phosphotyrosyl phosphatase activator (PTPA) of protein phosphatase 2A. *Genomics* 28: 261–272, 1995.

[74476] 12806.Andre, E.; Becker–Andre, M.: Expression of an N-terminally truncated form of human focal adhesion kinase in brain. *Biochem. Biophys. Res. Commun.* 190: 140–147, 1993.

[74477] 12807.Fiedorek, F. T., Jr.; Kay, E. S.: Mapping of the focal adhesion kinase (Fadk) gene to mouse chromosome 15 and human chromosome 8. *Mammalian Genome* 6: 123–126, 1995.

[74478] 12808.Polte, T. R.; Hanks, S. K.: Interaction between focal adhesion kinase and Crk-associated tyrosine kinase substrate p130-Cas. *Proc. Nat. Acad. Sci.* 92: 10678–10682, 1995.

[74479] 12809.Schaller, M. D.; Borgman, C.; Cobb, B. S.; Vines, R. R.; Reynolds, A. B.; Parsons, J. T.: pp125(FAK), a structurally distinctive protein-tyrosine kinase associated with focal adhesions. *Proc. Nat. Acad. Sci.* 89:5192–5196, 1992.

[74480] 12810.Arnemann, J.; Epplen, J. T.; Cooke, H. J.; Sauermann, U.; Engel, W.; Schmidtke, J.: A human Y-



chromosomal DNA sequence expressed in testicular tissue. Nucleic Acids Res. 15: 8713–8724, 1987.

- [74481] 12811. Arnemann, J.; Jakubiczka, S.; Thuring, S.; Schmidtke, J.: Cloning and sequence analysis of a human Y-chromosome-derived, testicular cDNA, TSPY. Genomics 11: 108–114, 1991.
- [74482] 12812. Guttenbach, M.; Muller, U.; Schmid, M.: A human moderately repeated Y-specific DNA sequence is evolutionarily conserved in the Y chromosome of the great apes. Genomics 13: 363–367, 1992.
- [74483] 12813. Jakubiczka, S.; Schnieders, F.; Schmidtke, J.: A bovine homologue of the human TSPY gene. Genomics 17: 732–735, 1993.
- [74484] 12814. Lau, Y.-F. C.: Sex chromosome genetics '99: gonadoblastoma, testicular and prostate cancers, and the TSPY gene. Am. J. Hum. Genet. 64:921–927, 1999.
- [74485] 12815. Manz, E.; Schnieders, F.; Brechlin, A. M.; Schmidtke, J.: TSPY-related sequences represent a micro-heterogeneous gene family organized as constitutive elements in DYZ5 tandem repeat units on the human Y chromosome. Genomics 17:726–731, 1993.
- [74486] 12816. Mazeyrat, S.; Mitchell, M. J.: Rodent Y chromosome TSPY gene is functional in rat and non-functional in

mouse. Hum. Molec. Genet. 7:557–562, 1998.

[74487] 12817.Schnieders, F.; Dork, T.; Arnemann, J.; Vogel, T.; Werner, M.;Schmidtke, J.: Testis–specific protein, Y–encoded (TSPY) expressionin testicular tissues. Hum. Molec. Genet. 5: 1801–1807, 1996.

[74488] 12818.Vogel, T.; Boettger–Tong, H.; Nanda, I.; Dechend, F.; Agulnik,A. I.; Bishop, C. E.; Schmid, M.; Schmidtke, J.: A murine TSPY. ChromosomeRes. 6: 35–40, 1998.

[74489] 12819.Zhang, J. S.; Yang–Feng, T. L.; Muller, U.; Mohandas, T. K.; deJong, P. J.; Lau, Y.–F. C.: Molecular isolation and characterizationof an expressed gene from the human Y–chromosome. Hum. Molec. Genet. 1:717–726, 1992.

[74490] 12820.Donnelly, P.; Tavare, S.; Balding, D. J.; Griffiths, R. C.: Estimatingthe age of the common ancestor of men from the ZFY intron.(Letter) Science 272:1357–1359, 1996.

[74491] 12821.Dorit, R. L.; Akashi, H.; Gilbert, W.: Absence of polymorphismat the ZFY locus on the human Y chromosome. Science 268: 1183–1185,1995.

[74492] 12822.Dorit, R. L.; Akashi, H.; Gilbert, W.: Estimating the age of thecommon ancestor of men from the ZFY intron.(Letter) Science 272:1361–1362, 1996.

[74493] 12823.Fu, Y.–X.; Li, W.–H.: Estimating the age of the com–

mon ancestor of men from the ZFY intron. (Letter) *Science* 272: 1356–1357, 1996.

- [74494] 12824. Mahaffey, C. L.; Bayleran, J. K.; Yeh, G. Y.; Lee, T. C.; Page, D. C.; Simpson, E. M.: Intron/exon structure confirms that mouse *Zfy1* and *Zfy2* are members of the ZFY gene family. *Genomics* 41: 123–127, 1997.
- [74495] 12825. Palmer, M. S.; Berta, P.; Sinclair, A. H.; Pym, B.; Goodfellow, P. N.: Comparison of human ZFY and ZFX transcripts. *Proc. Nat. Acad. Sci.* 87: 1681–1685, 1990.
- [74496] 12826. Rogers, J.; Samollow, P. B.; Comuzzie, A. G.: Estimating the age of the common ancestor of men from the ZFY intron. (Letter) *Science* 272: 1360–1361, 1996.
- [74497] 12827. Sinclair, A. H.; Foster, J. W.; Spencer, J. A.; Page, D. C.; Palmer, M.; Goodfellow, P. N.; Graves, J. A. M.: Sequences homologous to ZFY, a candidate human sex-determining gene, are autosomal in marsupials. *Nature* 336: 780–783, 1988.
- [74498] 12828. Slattery, J. P.; Sanner-Wachter, L.; O'Brien, S. J.: Novel gene conversion between X–Y homologues located in the nonrecombining region of the Y chromosome in Felidae (mammalia). *Proc. Nat. Acad. Sci.* 97: 5307–5312, 2000.
- [74499] 12829. Weiss, G.; von Haeseler: Estimating the age of the

common ancestor of men from the ZFY intron. (Letter) *Science* 272: 1359–1360, 1996.

- [74500] 12830. Zambrowicz, B. P.; Findley, S. D.; Simpson, E. M.; Page, D. C.; Palmiter, R. D.: Characterization of the murine *Zfy1* and *Zfy2* promoters. *Genomics* 24:406–408, 1994.
- [74501] 12831. Mach, B.; Steimle, V.; Martinez–Soria, E.; Reith, W.: Regulation of MHC class II genes: lessons from a disease. *Annu. Rev. Immun.* 14:301–331, 1996.
- [74502] 12832. Reith, W.: Personal Communication. Geneva, Switzerland 5/30/1997.
- [74503] 12833. Scholl, T.; Mahanta, S. K.; Strominger, J. L.: Specific complex formation between the type II bare lymphocyte syndrome–associated transactivators CIITA and RFX5. *Proc. Nat. Acad. Sci.* 94: 6330–6334, 1997.
- [74504] 12834. Emery, P.; Durand, B.; Mach, B.; Reith, W.: RFX proteins, a novel family of DNA binding proteins conserved in the eukaryotic kingdom. *Nucleic Acids Res.* 24: 803–807, 1996.
- [74505] 12835. Pugliati, L.; Reith, W.; Fey, S.; Mach, B.: Mapping the RF–X gene, encoding a DNA–binding protein controlling HLA class II gene expression, to 19p13. (Abstract) *Cytogenet. Cell Genet.* 51: 1061 only, 1989.
- [74506] 12836. Mizuta, M.; Inagaki, N.; Nemoto, Y.; Matsukura, S.;

Takahashi,M.; Seino, S.: Synaptotagmin III is a novel iso-form of rat synaptotagmin expressed in endocrine and neuronal cells. J. Biol. Chem. 269: 11675–11678,1994.

[74507] 12837.Li, X.-J.; Wang, D.-Y.; Zhu, Y.; Guo, R.-J.; Wang, X.-D.; Lubomir,K.; Mukai, K.; Sasaki, H.; Yoshida, H.; Oka, T.; Machinami, R.; Shinmura,K.; Tanaka, M.; Sugimura, H.: Mxi1 mutations in human neurofibrosarcomas. Jpn.J. Cancer Res. 90: 740–746, 1999.

[74508] 12838.Prochownik, E. V.; Grove, L. E.; Deubler, D.; Zhu, X. L.; Stephenson,R. A.; Rohr, L. R.; Yin, X.; Brothman, A. R.: Commonly occurring loss and mutation of the MXI1 gene in prostate cancer. Genes Chromosomes Cancer 22: 295–304, 1998.

[74509] 12839.Schreiber–Agus, N.; Meng, Y.; Hoang, T.; Hou, H., Jr.; Chen, K.;Greenberg, R.; Cordon–Cardo, C.; Lee, H.–W.; DePinho, R. A.: Role of Mxi1 in ageing organ systems and the regulation of normal and neoplastic growth. Nature 393: 483–487, 1998.

[74510] 12840.Shapiro, D. N.; Valentine, V.; Eagle, L.; Yin, X.; Morris, S. W.;Prochownik, E. V.: Assignment of the human MAD and MXI1 genes to chromosomes 2p12–p13 and 10q24–q25. Genomics 23: 282–285, 1994.

[74511] 12841.Wechsler, D. S.; Hawkins, A. L.; Li, X.; Jabs, E. W.;

Griffin,C. A.; Dang, C. V.: Localization of the human Mxi1 transcriptionfactor gene (MXI1) to chromosome 10q24–q25. *Genomics* 21: 669–672,1994.

[74512] 12842.Wechsler, D. S.; Shelly, C. A.; Dang, C. V.: Genomic organizationof human MXI1, a putative tumor suppressor gene. *Genomics* 32: 466–470,1996.

[74513] 12843.Zervos, A. S.; Gyuris, J.; Brent, R.: Mxi1, a protein that specificallyinteracts with Max to bind Myc–Max recognition sites. *Cell* 72: 223–232,1993. Note: Erratum: *Cell* 79: 389 only, 1994.

[74514] 12844.Boie, Y.; Rushmore, T. H.; Darmon–Goodwin, A.; Grygorczyk, R.;Slipetz, D. M.; Metters, K. M.; Abramovitz, M.: Cloning and expressionof a cDNA for the human prostanoid IP receptor. *J. Biol. Chem.* 269:12173–12178, 1994.

[74515] 12845.Ishikawa, T.; Tamai, Y.; Rochelle, J. M.; Hirata, M.; Namba, T.;Sugimoto, Y.; Ichikawa, A.; Narumiya, S.; Taketo, M. M.; Seldin, M.F.: Mapping of the genes encoding mouse prostaglandin D, E, and Fand prostacyclin receptors. *Genomics* 32: 285–288, 1996.

[74516] 12846.Murata, T.; Ushikubi, F.; Matsuoka, T.; Hirata, M.; Yamasaki, A.;Sugimoto, Y.; Ichikawa, A.; Aze, Y.; Tanaka, T.; Yoshida, N.; Ueno,A.; Oh–ishi, S.; Narumiya, S.: Altered

pain perception and inflammatory response in mice lacking prostacyclin receptor. *Nature* 388: 678–682, 1997.

- [74517] 12847. Ogawa, Y.; Tanaka, I.; Inoue, M.; Yoshitake, Y.; Isse, N.; Nakagawa, O.; Usui, T.; Itoh, H.; Yoshimasa, T.; Narumiya, S.; Nakao, K.: Structural organization and chromosomal assignment of the human prostacyclin receptor gene. *Genomics* 27: 142–148, 1995.
- [74518] 12848. Hoffman, I.; Balling, R.: Chromosomal localization of the murine cadherin-11. *Mammalian Genome* 6: 304 only, 1995.
- [74519] 12849. Okazaki, M.; Takeshita, S.; Kawai, S.; Kikuno, R.; Tsujimura, A.; Kudo, A.; Amann, E.: Molecular cloning and characterization of OB-cadherin, a new member of cadherin family expressed in osteoblasts. *J. Biol. Chem.* 269: 12092–12098, 1994.
- [74520] 12850. Tanihara, H.; Sano, K.; Heimark, R. L.; St. John, T.; Suzuki, S.: Cloning of five human cadherins clarifies characteristic features of cadherin extracellular domain and provides further evidence for two structurally different types of cadherin. *Cell Adhes. Commun.* 2: 15–26, 1994.
- [74521] 12851. Christie, P. T.; Curley, A.; Nesbit, M. A.; Chapman, C.; Genet, S.; Harper, P. S.; Keeling, S. L.; Wilkie, A. O. M.; Winter, R. M.; Thakker, R. V.: Mutational analysis in X-

linked spondyloepiphysealdysplasia tarda. J. Clin. Endocr. Metab. 86: 3233–3236, 2001.

[74522] 12852.Fiedler, J.; Bittner, M.; Puhl, W.; Brenner, R. E.: Mutations in the X-linked spondyloepiphyseal dysplasia tarda (SEDL) coding sequence are not a common cause of early primary osteoarthritis in men. (Letter) Clin.Genet. 62: 94–95, 2002.

[74523] 12853.Gecz, J.; Hillman, M. A.; Gedeon, A. K.; Cox, T. C.; Baker, E.; Mulley, J. C.: Gene structure and expression study of the SEDL gene for spondyloepiphyseal dysplasia tarda. Genomics 69: 242–251, 2000.

[74524] 12854.Gedeon, A. K.; Colley, A.; Jamieson, R.; Thompson, E. M.; Rogers, J.; Sillence, D.; Tiller, G. E.; Mulley, J. C.; Gecz, J.: Identification of the gene (SEDL) causing X-linked spondyloepiphyseal dysplasia tarda. Nature Genet. 22: 400–404, 1999.

[74525] 12855.Gedeon, A. K.; Tiller, G. E.; Le Merrer, M.; Heuertz, S.; Tranebjaerg, L.; Chitayat, D.; Robertson, S.; Glass, I. A.; Savarirayan, R.; Cole, W. G.; Rimoin, D. L.; Kousseff, B. G.; Ohashi, H.; Zabel, B.; Munnich, A.; Gecz, J.; Mulley, J. C.: The molecular basis of X-linked spondyloepiphysealdysplasia tarda. Am. J. Hum. Genet. 68: 1386–1397, 2001.

[74526] 12856.Grunebaum, E.; Arpaia, E.; MacKenzie, J. J.; Fitz-



patrick, J.; Ray, P. N.; Roifman, C. M.: A missense mutation in the SEDL gene results in delayed onset of X linked spondyloepiphyseal dysplasia in a large pedigree. (Letter) J. Med. Genet. 38: 409–411, 2001.

[74527] 12857. Mumm, S.; Christie, P. T.; Finnegan, P.; Jones, J.; Dixon, P. H.; Pannett, A. A. J.; Harding, B.; Gottesman, G. S.; Thakker, R. V.; Whyte, M. P.: A five-base pair deletion in the sedlin gene causes spondyloepiphyseal dysplasia tarda in a six-generation Arkansas kindred. J. Clin. Endocr. Metab. 85: 3343–3347, 2000.

[74528] 12858. Mumm, S.; Zhang, X.; Vacca, M.; D'Esposito, M.; Whyte, M. P.: The sedlin gene for spondyloepiphyseal dysplasia tarda escapes X-inactivation and contains a non-canonical splice site. Gene 273: 285–293, 2001.

[74529] 12859. Takahashi, T.; Takahashi, I.; Tsuchida, S.; Oyama, K.; Komatsu, M.; Saito, H.; Takada, G.: An SEDL gene mutation in a Japanese kindred of X-linked spondyloepiphyseal dysplasia tarda. (Letter) Clin. Genet. 61: 319–320, 2002.

[74530] 12860. Tiller, G. E.; Hannig, V. L.; Dozier, D.; Carrel, L.; Trevarthen, K. C.; Wilcox, W. R.; Mundlos, S.; Haines, J. L.; Gedeon, A. K.; Gecz, J.: A recurrent RNA-splicing mutation in the SEDL gene causes X-linked spondyloepiphyseal dysplasia tarda. Am. J. Hum. Genet. 68: 1398–1407, 2001.

- [74531] 12861. Whyte, M. P.; Gottesman, G. S.; Eddy, M. C.; McAlister, W. H.: X-linked recessive spondyloepiphyseal dysplasia tarda: clinical and radiographic evolution in a 6-generation kindred and review of the literature. *Medicine* 78: 9-25, 1999.
- [74532] 12862. Becker, M. A.; Heidler, S. A.; Bell, G. I.; Seino, S.; Le Beau, M. M.; Westbrook, C. A.; Neuman, W.; Shapiro, L. J.; Mohandas, T. K.; Roessler, B. J.; Palella, T. D.: Cloning of cDNAs for human phosphoribosylpyrophosphatesynthetases 1 and 2 and X chromosome localization of PRPS1 and PRPS2 genes. *Genomics* 8: 555-561, 1990.
- [74533] 12863. Buchner, G.; Montini, E.; Andolfi, G.; Quaderi, N.; Cainarca, S.; Messali, S.; Bassi, M. T.; Ballabio, A.; Meroni, G.; Franco, B.: MID2, a homologue of the Opitz syndrome gene MID1: similarities in subcellular localization and differences in expression during development. *Hum. Molec. Genet.* 8: 1397-1407, 1999.
- [74534] 12864. Braverman, N.; Lin, P.; Moebius, F. F.; Obie, C.; Moser, A.; Glossmann, H.; Wilcox, W. R.; Rimoïn, D. L.; Smith, M.; Kratz, L.; Kelley, R. I.; Valle, D.: Mutations in the gene encoding 3-beta-hydroxysteroid-delta(8),delta(7)-isomerase cause X-linked dominant Conradi-Hunermann syndrome. *Nature*

Genet. 22:291–294, 1999.

- [74535] 12865.Cho, S. Y.; Kim, J. H.; Paik, Y. K.: Cholesterol biosynthesis from lanosterol: differential inhibition of sterol delta 8–isomerase and other lanosterol–converting enzymes by tamoxifen. *Molec. Cells* 8:233–239, 1998.
- [74536] 12866.Clayton, P. T.; Kalter, D. C.; Atherton, D. J.; Besley, G. T.; Broadhead, D. M.: Peroxisomal enzyme deficiency in X–linked dominant Conradi–Hunermann syndrome. *J. Inherit. Metab. Dis.* 12: 358–360, 1989.
- [74537] 12867.Derry, J. M. J.; Gormally, E.; Means, G. D.; Zhao, W.; Meindl, A.; Kelley, R. I.; Boyd, Y.; Herman, G. E.: Mutations in a delta(8)–delta(7)sterol isomerase in the tattered mouse and X–linked dominant chondrodysplasia punctata. *Nature Genet.* 22: 286–290, 1999.
- [74538] 12868.Grange, D. K.; Kratz, L. E.; Braverman, N. E.; Kelley, R. I.: CHILD syndrome caused by deficiency of 3–beta–hydroxysteroid–delta–8,delta–7–isomerase. *Am.J. Med. Genet.* 90: 328–335, 2000.
- [74539] 12869.Hanner, M.; Moebius, F. F.; Weber, F.; Grabner, M.; Striessnig, J.; Glossmann, H.: Phenylalkylamine Ca(2+) antagonist binding protein: molecular cloning, tissue distribution, and heterologous expression. *J. Biol. Chem.* 270: 7551–7557, 1995.

- [74540] 12870.Has, C.; Bruckner-Tuderman, L.; Muller, D.; Floeth, M.; Folkers, E.; Donnai, D.; Traupe, H.: The Conradi-Hunermann-Happle syndrome (CDPX2) and emopamil binding protein: novel mutations, and somatic and gonadal mosaicism. *Hum. Molec. Genet.* 9: 1951-1955, 2000.
- [74541] 12871.Holmes, R. D.; Wilson, G. N.; Hajra, A. K.: Peroxisomal enzyme deficiency in the Conradi-Hunerman (sic) form of chondrodysplasia punctata. *New Eng. J. Med.* (Letter) 316: 1608 only, 1987.
- [74542] 12872.Ikegawa, S.; Ohashi, H.; Ogata, T.; Honda, A.; Tsukahara, M.; Kubo, T.; Kimizuka, M.; Shimode, M.; Hasegawa, T.; Nishimura, G.; Nakamura, Y.: Novel and recurrent EBP mutations in X-linked dominant chondrodysplasia punctata. *Am. J. Med. Genet.* 94: 300-305, 2000.
- [74543] 12873.Kelley, R. I.; Wilcox, W. G.; Smith, M.; Kratz, L. E.; Moser, A.; Rimoin, D. S.: Abnormal sterol metabolism in patients with Conradi-Hunermann-Happle syndrome and sporadic lethal chondrodysplasia punctata. *Am. J. Med. Genet.* 83: 213-219, 1999.
- [74544] 12874.Aksoy, S.; Brandriff, B. F.; Ward, A.; Little, P. F. R.; Weinshilboum, R. M.: Human nicotinamide N-methyltransferase gene: molecular cloning, structural characterization and chromosomal localization. *Genomics*

29:555–561, 1995.

- [74545] 12875.Aksoy, S.; Szumlanski, C. L.; Weinshilboum, R. M.: Human livernicotinamide N–methyltransferase: cDNA cloning, expression, and biochemical characterization. *J. Biol. Chem.* 269: 14835–14840, 1994.
- [74546] 12876.Bennett, B. D.; Wang, Z.; Kuang, W.–J.; Wang, A.; Groopman, J.E.; Goeddel, D. V.; Scadden, D. T.: Cloning and characterization of HTK, a novel transmembrane tyrosine kinase of the EPH subfamily. *J. Biol. Chem.* 269: 14211–14218, 1994.
- [74547] 12877.Berclaz, G.; Andres, A.–C.; Albrecht, D.; Dreher, E.; Ziemiecki, A.; Gusterson, B. A.; Crompton, M. R.: Expression of the receptor protein tyrosine kinase myk–1/htk in normal and malignant mammary epithelium. *Biochem. Biophys. Res. Commun.* 226: 869–875, 1996.
- [74548] 12878.Gerety, S. S.; Wang, H. U.; Chen, Z.–F.; Anderson, D. J.: Symmetrical mutant phenotypes of the receptor EphB4 and its specific transmembrane ligand ephrin–B2 in cardiovascular development. *Molec. Cell* 4: 403–414, 1999.
- [74549] 12879.Moynihan, T. P.; Ardley, H. C.; Leek, J. P.; Thompson, J.; Brindle, N. S.; Markham, A. F.; Robinson, P. A.: Characterization of a human ubiquitin–conjugating en–

zyme gene UBE2L3. Mammalian Genome 7:  
520–525,1996.

- [74550] 12880.Flanagan, J. R.; Becker, K. G.; Ennist, D. L.; Gleason, S. L.; Driggers, P. H.; Levi, B.-Z.; Appella, E.; Ozato, K.: Cloning of a negative transcription factor that binds to the upstream conserved region of Moloney murine leukemia virus. *Molec. Cell. Biol.* 12:38–44, 1992.
- [74551] 12881.Hariharan, N.; Kelley, D. E.; Perry, R. P.: Delta, a transcription factor that binds to downstream elements in several polymerase II promoters, is a functionally versatile zinc finger protein. *Proc.Nat. Acad. Sci.* 88: 9799–9803, 1991.
- [74552] 12882.Oei, S. L.; Shi, Y.: Transcription factor Yin Yang 1 stimulates poly(ADP-ribosyl)ation and DNA repair. *Biochem. Biophys. Res. Commun.* 284:450–454, 2001.
- [74553] 12883.Park, K.; Atchison, M. L.: Isolation of a candidate repressor/activator, NF-E1 (YY-1, delta), that binds to the immunoglobulin kappa 3-prime enhancer and the immunoglobulin heavy-chain micro-E1 site.. *Proc.Nat. Acad. Sci.* 88: 9804–9808, 1991.
- [74554] 12884.Shi, Y.; Seto, E.; Chang, L.-S.; Shenk, T.: Transcriptional repression by YY1, a human GLI-Kruppel-related protein, and relief of repression by adenovirus E1A protein.

Cell 67: 377–388, 1991.

- [74555] 12885. Yao, Y.-L.; Dupont, B. R.; Ghosh, S.; Fang, Y.; Leach, R. J.; Seto, E.: Cloning, chromosomal localization and promoter analysis of the human transcription factor YY1. *Nucleic Acids Res.* 26: 3776–3783, 1998.
- [74556] 12886. Zhu, W.; Lossie, A. C.; Camper, S. A.; Gumucio, D. L.: Chromosomal localization of the transcription factor YY1 in the mouse and human. *Mammalian Genome* 5: 234–236, 1994.
- [74557] 12887. Hakimi, M.-A.; Bochar, D. A.; Schmiesing, J. A.; Dong, Y.; Barak, O. G.; Speicher, D. W.; Yokomori, K.; Shiekhata, R.: A chromatin remodelling complex that loads cohesin onto human chromosomes. *Nature* 418:994–998, 2002.
- [74558] 12888. Muchardt, C.; Reyes, J. C.; Bourachot, B.; Leguoy, E.; Yaniv, M.: The hbrm and BRG-1 proteins, components of the human SNF/SWI complex, are phosphorylated and excluded from the condensed chromosomes during mitosis. *EMBO J.* 15: 3394–3402, 1996.
- [74559] 12889. Muchardt, C.; Yaniv, M.: A human homologue of *Saccharomyces cerevisiae* SNF2/SWI2 and *Drosophila* brm genes potentiates transcriptional activation by the glucocorticoid receptor. *EMBO J.* 12: 4279–4290, 1993.

- [74560] 12890. Muchardt, C.; Yaniv, M.; Mattei, M.-G.: Assignment of HBRM, the human homolog of *S. cerevisiae* SNF2/SWI2 and *Drosophila* brm genes, to chromosome region 9p23-p24, by in situ hybridization. *Mammalian Genome* 5: 241-243, 1994.
- [74561] 12891. Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Kawarabayashi, Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.: Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40 new genes (KIAA0001-KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line KG-1. *DNA Res.* 1: 27-35, 1994.
- [74562] 12892. Kikuno, R.; Nagase, T.; Ishikawa, K.; Hirose, M.; Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. XIV. The complete sequences of 100 new cDNA clones from brain which code for large proteins in vitro. *DNA Res.* 6: 197-205, 1999.
- [74563] 12893. Polsky, D.; Young, A. Z.; Busam, K. J.; Alani, R. M.: The transcriptional repressor of p16/Ink4a, Id1, is up-regulated in early melanomas. *Cancer Res.* 61: 6008-6011, 2001.
- [74564] 12894. Walker, J. L.; Dixon, J.; Fenton, C. R.; Hungerford, D. W.;



J.; Lynch, S. A.; Stenhouses, S. A. R.; Christian, A.; Craig, I. W.: Two new mutations in exon 3 of the NDP gene: S73X and S101F associated with severe and less severe ocular phenotype, respectively. *Hum. Mutat.* 9:53–56, 1997.

[74565] 12895. Lee, D. K.; Horikoshi, M.; Roeder, R. G.: Interaction of TFIID in the minor groove of the TATA element. *Cell* 67: 1241–1250, 1991.

[74566] 12896. Wilson, R. B.; Kiledjian, M.; Shen, C.-P.; Benezra, R.; Zwollo, P.; Dymecki, S. M.; Desiderio, S. V.; Kadesch, T.: Repression of immunoglobulin enhancers by the helix–loop–helix protein Id: implications for B–lymphoid–cell development. *Molec. Cell. Biol.* 11: 6185–6191, 1991.

[74567] 12897. Battaglia, G.; Princivale, A.; Forti, F.; Lizier, C.; Zeviani, M.: Expression of the SMN gene, the spinal muscular atrophy determining gene, in the mammalian central nervous system. *Hum. Molec. Genet.* 6:1961–1971, 1997.

[74568] 12898. Burglen, L.; Lefebvre, S.; Clermont, O.; Burlet, P.; Viollet, L.; Cruaud, C.; Munnich, A.; Melki, J.: Structure and organization of the human survival motor neurone (SMN) gene. *Genomics* 32: 479–482, 1996.

[74569] 12899. Burglen, L.; Spiegel, R.; Ignatius, J.; Cobben, J. M.; Landrieu, P.; Lefebvre, S.; Munnich, A.; Melki, J.: SMN gene deletion in variant of infantile spinal muscular atrophy.

(Letter) Lancet 346: 316–317,1995.

- [74570] 12900.Bussaglia, E.; Clermont, O.; Tizzano, E.; Lefebvre, S.; Burglen,L.; Cruaud, C.; Urtizberea, J. A.; Colomer, J.; Munnich, A.; Baiget,M.; Melki, J.: A frame–shift deletion in the survival motor neurongene in Spanish spinal muscular atrophy patients. Nature Genet. 11:335–337, 1995.
- [74571] 12901.Callebaut, I.; Mornon, J. P.: The human EBNA–2 coactivator p100:multidomain organization and relationship to the staphylococcal nucleasefold and to the tudor protein involved in Drosophila melanogasterdevelopment. Biochem. J. 321: 125–132, 1997.
- [74572] 12902.Campbell, L.; Daniels, R. J.; Dubowitz, V.; Davies, K. E.: Maternalmosaicism for a second mutational event in a type I spinal muscularatrophy family. Am. J. Hum. Genet. 63: 37–44, 1998.
- [74573] 12903.Campbell, L.; Hunter, K. M. D.; Mohaghegh, P.; Tinsley, J. M.;Brasch, M. A.; Davies, K. E.: Direct interaction of Smn with dp103,a putative RNA helicase: a role for Smn in transcription regulation? Hum.Molec. Genet. 9: 1093–1100, 2000.
- [74574] 12904.Chen, K.–L.; Wang, Y. L.; Rennert, H.; Joshi, I.; Mills, J. K.;Leonard, D. G. B.; Wilson, R. B.: Duplications and de novo deletionsof the SMNt gene demonstrated by fluores–

cence-based carrier testing for spinal muscular atrophy.

Am. J. Med. Genet. 85: 463–469, 1999.

[74575] 12905. Warburg, M.: Norrie's disease: a new hereditary bilateral pseudotumor of the retina. Acta Ophthalmol. (Copenh) 39: 757–772, 1961.

[74576] 12906. Warburg, M.: Norrie's disease (atrofia bulborum hereditaria). Acta Ophthalmol. 41: 134–146, 1963.

[74577] 12907. Warburg, M.: Norrie's disease, a congenital progressive oculo-acoustico-cerebral degeneration. Acta Ophthalmol. 89 (suppl.): 1–147, 1966.

[74578] 12908. Warburg, M.; Hauge, M.; Sanger, R.: Norrie's disease and the Xg blood group system: linkage data. Acta Genet. Statist. Med. 15: 103–115, 1965.

[74579] 12909. Whitnall, S. E.; Norman, R. M.: Microphthalmia and the visual pathways: a case associated with blindness and imbecility, and sex-linked. Brit. J. Ophthalmol. 24: 229–244, 1940.

[74580] 12910. Wilson, W. M. G.: Congenital blindness (pseudoglioma) occurring as a sex-linked developmental anomaly. Canad. Med. Assoc. J. 60: 580–584, 1949.

[74581] 12911. Wolff, G.; Mayerova, A.; Wienker, T. F.; Atalianis, P.; Ioannou, P.; Warburg, M.: Clinical reinvestigation and linkage analysis in the family with Episkopi blindness (Norrie

disease). *J. Med. Genet.* 29:816–819, 1992.

[74582] 12912.Wong, F.; Goldberg, M. F.; Hao, Y.: Identification of a nonsensemutation at codon 128 of the Norrie's disease gene in a male infant. *Arch.Ophthal.* 111: 1553–1557, 1993.

[74583] 12913.Woodruff, G.; Newbury–Ecob, R.; Plaha, D. S.; Young, I. D.: Manifestingheterozygosity in Norrie's disease?. *Brit. J. Ophthal.* 77: 813–814,1993.

[74584] 12914.Zhu, D.; Antonarakis, S. E.; Schmeckpeper, B. J.; Diergaarde,P. J.; Greb, A. E.; Maumenee, I. H.: Microdeletion in the X–chromosomeand prenatal diagnosis in a family with Norrie disease. *Am. J. Med.Genet.* 33: 485–488, 1989.

[74585] 12915.Avner, P.; Bucan, M.; Arnaud, D.; Lehrach, H.; Rapp, U.: A–rafoncogene localizes on mouse X chromosome to region some 10–17 centimorgansproximal to hypoxanthine phosphoribosyltransferase gene. *Somat. CellMolec. Genet.* 13: 267–272, 1987.

[74586] 12916.Beck, T. W.; Huleihel, M.; Gunnell, M.; Bonner, T. I.; Rapp, U.R.: The complete coding sequence of the human A–raf–1 oncogene andtransforming activity of a human A–raf carrying retrovirus. *NucleicAcids Res.* 15: 595–609, 1987.

- [74587] 12917.Huebner, K.; ar-Rushdi, A.; Griffin, C. A.; Isobe, M.; Kozak, C.;Emanuel, B. S.; Nagarajan, L.; Cleveland, J. L.; Bonner, T. I.; Goldsborough,M. D.; Croce, C. M.; Rapp, U.: Actively transcribed genes in the raf oncogene group, located on the X chromosome in mouse and human. Proc.Nat. Acad. Sci. 83: 3934–3938, 1986.
- [74588] 12918.Lee, J.–E.; Beck, T. W.; Brennscheidt, U.; DeGennaro, L. J.; Rapp,U. R.: The complete sequence and promoter activity of the human A-raf-1gene (ARAF1). Genomics 20: 43–55, 1994.
- [74589] 12919.Mark, G. E.; Seeley, T. W.; Shows, T. B.; Mountz, J. D.: Pks,a raf-related sequence in humans. Proc. Nat. Acad. Sci. 83: 6312–6316,1986.
- [74590] 12920.Popescu, N. C.; Mark, G. E.: Localization of the pKs gene, a rafrelated sequence on human chromosomes X and 7. Oncogene 4: 517–519,1989.
- [74591] 12921.Dowdy, S. F.; Fasching, C. L.; Araujo, D.; Lai, K.–M.; Livanos,E.; Weissman, B. E.; Stanbridge, E. J.: Suppression of tumorigenicityin Wilms' tumor by the p15.5–p14 region of chromosome 11. Science 254:293–295, 1991.
- [74592] 12922.Dowdy, S. F.; Lai, K.–M.; Weissman, B. E.; Matsui, Y.; Hogan, B.L. M.; Stanbridge, E. J.: The isolation and characterization of a novel cDNA demonstrating an altered

mRNA level in nontumorigenic Wilms' microcell hybrid cells. *Nucleic Acids Res.* 19: 5763–5769, 1991.

- [74593] 12923. Farmer, A. A.; Loftus, T. M.; Mills, A. A.; Sato, K. Y.; Neill, J. D.; Tron, T.; Yang, M.; Trumpower, B. L.; Stanbridge, E. J.: Extreme evolutionary conservation of QM, a novel c-Jun associated transcription factor. *Hum. Molec. Genet.* 3: 723–728, 1994.
- [74594] 12924. Kaneko, K.; Kobayashi, H.; Onodera, O.; Miyatake, T.; Tsuji, S.: Genomic organization of a cDNA (QM) demonstrating an altered mRNA level in nontumorigenic Wilms' microcell hybrid cells and its localization to Xq28. *Hum. Molec. Genet.* 1: 529–533, 1992.
- [74595] 12925. Korn, B.; Sedlacek, Z.; Manca, A.; Kioschis, P.; Konnecki, D.; Lehrach, H.; Poustka, A.: A strategy for the selection of transcribed sequences in the Xq28 region. *Hum. Molec. Genet.* 1: 235–242, 1992.
- [74596] 12926. van den Ouweland, A. M. W.; Verdijk, M.; Mannens, M. M. A. M.; van Oost, B. A.: The QM gene is X-linked and therefore not involved in suppression of tumorigenesis in Wilms' tumor. *Hum. Molec. Genet.* 90:144–146, 1992.
- [74597] 12927. Weissman, B. E.; Saxon, P. J.; Pasquale, S. R.; Jones, G. R.; Geiser, A. G.; Stanbridge, E. J.: Introduction of a normal human chromosome 11 into a Wilms' tumor cell line

controls its tumorigenic expression. Science  
236:175–180, 1987.

- [74598] 12928.Nishimoto, T.; Sekiguchi, T.; Kai, R.; Yamashita, K.; Takahashi,T.; Sekiguchi, M.: Large-scale selection and analysis of temperature-sensitive mutants for cell reproduction from BHK cells. Somat. Cell Genet. 8:811–812, 1982.
- [74599] 12929.O'Brien, T.; Tjian, R.: Functional analysis of the human TAFII250N-terminal kinase domain. Molec. Cell 1: 905–911, 1998.
- [74600] 12930.Ruppert, S.; Wang, E. H.; Tjian, R.: Cloning and expression of human TAF(II)250: a TBP-associated factor implicated in cell-cycle regulation. Nature 362: 175–179, 1993.
- [74601] 12931.Schwartz, H. E.; Holmes, S.; Meiss, H. K.: Assignment of temperature-sensitive mutations of BHK cells to the X-chromosome. (Abstract) J. Cell Biol. 75:393A, 1977.
- [74602] 12932.Schwartz, H. E.; Moser, G. C.; Holmes, S.; Meiss, H. K.: Assignment of temperature-sensitive mutations of BHK cells to the X-chromosome. Somat.Cell Genet. 5: 217–224, 1979.
- [74603] 12933.Sekiguchi, T.; Miyata, T.; Nishimoto, T.: Molecular cloning of the cDNA of human X chromosomal gene

(CCG1) which complements the temperature-sensitive G(1) mutants, tsBN462 and ts13, of the BHK cell line. EMBO J. 7: 1683–1687, 1988.

[74604] 12934. Sekiguchi, T.; Nohiro, Y.; Nakamura, Y.; Hisamoto, N.; Nishimoto, T.: The human CCG1 gene, essential for progression of the G1 phase, encodes a 210-kilodalton nuclear DNA-binding protein. Molec. Cell Biol. 11: 3317–3325, 1991.

[74605] 12935. Sekiguchi, T.; Yoshida, M. C.; Sekiguchi, M.; Nishimoto, T.: Isolation of a human X chromosome-linked gene essential for progression from G1 to S phase of the cell cycle. Exp. Cell Res. 169: 395–407, 1987.

[74606] 12936. Simchen, G.: Cell cycle mutants. Annu. Rev. Genet. 12: 161–191, 1978.

[74607] 12937. Slater, M. L.; Ozer, H. L.: Temperature-sensitive mutants of Balb-3T3 cells: description of a mutant affected in cellular and polyomavirus DNA synthesis. Cell 7: 289–295, 1976.

[74608] 12938. Starr, D. B.; Hawley, D. K.: TFIID binds in the minor groove of the TATA box. Cell 67: 1231–1240, 1991.

[74609] 12939. Abel, A.; Walcott, J.; Woods, J.; Duda, J.; Merry, D. E.: Expression of expanded repeat androgen receptor produces neurologic disease in transgenic mice. Hum. Molec.



Genet. 10: 107–116, 2001.

- [74610] 12940. Bailey, C. K.; Andriola, I. F. M.; Kampinga, H. H.; Merry, D. E.: Molecular chaperones enhance the degradation of expanded polyglutamine repeat androgen receptor in a cellular model of spinal and bulbar muscular atrophy. Hum. Molec. Genet. 11: 515–523, 2002.
- [74611] 12941. Batch, J. A.; Williams, D. M.; Davies, H. R.; Brown, B. D.; Evans, B. A. J.; Hughes, I. A.; Patterson, M. N.: Androgen receptor gene mutations identified by SSCP in fourteen subjects with androgen insensitivity syndrome. Hum. Molec. Genet. 1: 497–503, 1992.
- [74612] 12942. Beitel, L. K.; Kazemi-Esfarjani, P.; Kaufman, M.; Lumbroso, R.; DiGeorge, A. M.; Killinger, D. W.; Trifiro, M. A.; Pinsky, L.: Substitution of arginine–839 by cysteine or histidine in the androgen receptor causes different receptor phenotypes in cultured cells and coordinated degrees of clinical androgen resistance. J. Clin. Invest. 94: 546–554, 1994.
- [74613] 12943. Belsham, D. D.; Pereira, F.; Greenberg, C. R.; Liao, S.; Wrogemann, K.: Leu676-to-pro mutation of the androgen receptor causes complete androgen insensitivity syndrome in a large Hutterite kindred. Hum. Mutat. 5: 28–33, 1995.

- [74614] 12944.Bevan, C. L.; Brown, B. B.; Davies, H. R.; Evans, B. A. J.; Hughes,I. A.; Patterson, M. N.: Functional analysis of six androgen receptormutations identified in patients with partial androgen insensitivitysyndrome. Hum. Molec. Genet. 5: 265–273, 1996.
- [74615] 12945.Biancalana, V.; Serville, F.; Pommier, J.; Julien, J.; Hanauer,A.; Mandel, J. L.: Moderate instability of the trinucleotide repeatin spinobulbar muscular atrophy. Hum. Molec. Genet. 1: 255–258,1992.
- [74616] 12946.Boehmer, A. L. M.; Brinkmann, A. O.; Niermeijer, M. F.; Bakker,L.; Halley, D. J. J.; Drop, S. L. S.: Germ–line and somatic mosaicismin the androgen insensitivity syndrome: implications for genetic counseling.(Letter) Am. J. Hum. Genet. 60: 1003–1006, 1997.
- [74617] 12947.Boehmer, A. L. M.; Brinkmann, A. O.; Nijman, R. M.; Verleun–Mooijman,M. C. T.; de Ruiter, P.; Niermeijer, M. F.; Drop, S. L. S.: Phenotypicvariation in a family with partial androgen insensitivity syndromeexplained by differences in 5–alpha dihydrotestosterone availability. J.Clin. Endocr. Metab. 86: 1240–1246, 2001.
- [74618] 12948.Bouvattier, C.; Carel, J.–C.; Lecoindre, C.; David, A.; Sultan,C.; Bertrand, A.–M., Morel, Y.; Chaussain, J.–L.: Postnatal changesof T, LH, and FSH in 46,XY infants with

mutations in the AR gene. J.Clin. Endocr. Metab. 87: 29–32, 2002.

- [74619] 12949. Brown, C. J.; Goss, S. J.; Lubahn, D. B.; Joseph, D. R.; Wilson, E. M.; French, F. S.; Willard, H. F.: Androgen receptor locus on the human X chromosome: regional localization to Xq11–12 and description of a DNA polymorphism. Am. J. Hum. Genet. 44: 264–269, 1989.
- [74620] 12950. Brown, T. R.; Lubahn, D. B.; Wilson, E. M.; Joseph, D. R.; French, F. S.; Migeon, C. J.: Deletion of the steroid-binding domain of the human androgen receptor gene in one family with complete androgen insensitivity syndrome: evidence for further genetic heterogeneity in this syndrome. Proc. Nat. Acad. Sci. 85: 8151–8155, 1988.
- [74621] 12951. Bruggenwirth, H. T.; Boehmer, A. L. M.; Ramnarain, S.; Verleun-Mooijman, M. C. T.; Satijn, D. P. E.; Trapman, J.; Grootegoed, J. A.; Brinkmann, A. O.: Molecular analysis of the androgen-receptor gene in a family with receptor-positive partial androgen insensitivity: an unusual type of intronic mutation. Am. J. Hum. Genet. 61: 1067–1077, 1997.
- [74622] 12952. Buchanan, G.; Yang, M.; Harris, J. M.; Nahm, H. S.; Han, G.; Moore, N.; Bentel, J. M.; Matusik, R. J.; Horsfall, D. J.; Marshall, V. R.; Greenberg, N. M.; Tilley, W. D.: Muta-

tions at the boundary of the hinge and ligand binding domain of the androgen receptor confer increased transactivation function. *Molec. Endocr.* 15: 46–56, 2001.

- [74623] 12953. Butler, R.; Leigh, P. N.; McPhaul, M. J.; Gallo, J.-M.: Truncated forms of the androgen receptor are associated with polyglutamine expansion in X-linked spinal and bulbar muscular atrophy. *Hum. Molec. Genet.* 7:121–127, 1998.
- [74624] 12954. Muller, G.; Schempp, W.: Mapping the human ZFX locus to Xp21.3 by in situ hybridization. *Hum. Genet.* 82: 82–84, 1989.
- [74625] 12955. Palmer, M. S.; Sinclair, A. H.; Berta, P.; Ellis, N. A.; Goodfellow, P. N.; Abbas, N. E.; Fellous, M.: Genetic evidence that ZFY is not the testis-determining factor. *Nature* 342: 937–939, 1989.
- [74626] 12956. Masutani, C.; Sugasawa, K.; Yanagisawa, J.; Sonoyama, T.; Ui, M.; Enomoto, T.; Takio, K.; Tanaka, K.; van der Spek, P. J.; Bootsma, D.; Hoeijmakers, J. H. J.; Hanaoka, F.: Purification and cloning of a nucleotide excision repair complex involving the xeroderma pigmentosum group C protein and a human homologue of yeast RAD23. *EMBO J.* 13:1831–1843, 1994.
- [74627] 12957. Coover, D. D.; Le, T. T.; McAndrew, P. E.; Strass-

wimmer, J.; Crawford, T. O.; Mendell, J. R.; Coulson, S. E.; Androphy, E. J.; Prior, T. W.; Burghes, A. H. M.: The survival motor neuron protein in spinal muscular atrophy. *Hum. Molec. Genet.* 6: 1205–1214, 1997.

[74628] 12958. Campbell, H. D.; Fountain, S.; McLennan, I. S.; Berven, L. A.; Crouch, M. F.; Davy, D. A.; Hooper, J. A.; Waterford, K.; Chen, K.-S.; Lupski, J. R.; Ledermann, B.; Young, I. G.; Matthaei, K. I.: Fliih, a gelsolin-related cytoskeletal regulator essential for early mammalian embryonic development. *Molec. Cell. Biol.* 22: 3518–3526, 2002.

[74629] 12959. Campbell, H. D.; Fountain, S.; Young, I. G.; Claudianos, C.; Hoheisel, J. D.; Chen, K.-S.; Lupski, J. R.: Genomic structure, evolution, and expression of human FLII, a gelsolin and leucine-rich-repeat family member: overlap with LLGL. *Genomics* 42: 46–54, 1997.

[74630] 12960. Connelly, M. A.; Zhang, H.; Kieleczawa, J.; Anderson, C. W.: The promoters for human DNA-PK(cs) (PRKDC) and MCM4: divergently transcribed genes located at chromosome 8 band q11. *Genomics* 47: 71–83, 1998.

[74631] 12961. Aksoy, I. A.; Wood, T. C.; Weinshilboum, R.: Human liver estrogensulfotransferase: identification by cDNA cloning and expression. *Biochem. Biophys. Res. Commun.*

200: 1621–1629, 1994.

- [74632] 12962. Bernier, F.; Leblanc, G.; Labrie, F.; Luu-The, V.: Structure of human estrogen and aryl sulfotransferase gene: two mRNA species issued from a single gene. *J. Biol. Chem.* 269: 28200–28205, 1994.
- [74633] 12963. Her, C.; Aksoy, I. A.; Kimura, S.; Brandriff, B. F.; Wasmuth, J. J.; Weinshilboum, R. M.: Human estrogen sulfotransferase gene (STE): cloning, structure, and chromosomal localization. *Genomics* 29: 16–23, 1995.
- [74634] 12964. Weinshilboum, R.: Personal Communication. Rochester, Minn. 10/9/1995.
- [74635] 12965. Weinshilboum, R. M.; Otterness, D. M.; Aksoy, I. A.; Wood, T. C.; Her, C.; Raftogianis, R. B.: Sulfotransferase molecular biology: cDNAs and genes. *FASEB J.* 11: 3–14, 1997.
- [74636] 12966. Bartley, T. D.; Bogenberger, J.; Hunt, P.; Li, Y.-S.; Lu, H. S.; Martin, F.; Chang, M.-S.; Samal, B.; Nichol, J. L.; Swift, S.; Johnson, M. J.; Hsu, R.-Y.; and 41 others: Identification and cloning of a megakaryocyte growth and development factor that is a ligand for the cytokine receptor Mpl. *Cell* 77: 1117–1124, 1994.
- [74637] 12967. Chang, M.; McNinch, J.; Basu, R.; Shutter, J.; Hsu, R.; Perkins, C.; Mar, V.; Suggs, S.; Welcher, A.; Li, L.; Lu, H.;

Bartley, T.; Hunt, P.; Martin, F.; Samal, B.; Bogenberger, J.: Cloning and characterization of the human megakaryocyte growth and development factor (MGDF) gene. *J. Biol. Chem.* 270: 511–514, 1995.

[74638] 12968. Chang, M.-S.; Hsu, R.-Y.; McNinch, J.; Copeland, N. G.; Jenkins, N. A.: The gene for murine megakaryocyte growth and development factor (thrombopoietin, Thpo) is located on mouse chromosome 16. *Genomics* 26:636–637, 1995.

[74639] 12969. de Sauvage, F. J.; Hass, P. E.; Spencer, S. D.; Malloy, B. E.; Gurney, A. L.; Spencer, S. A.; Darbonne, W. C.; Henzel, W. J.; Wong, S. C.; Kuang, W.-J.; Oles, K. J.; Hultgren, B.; Solberg, L. A., Jr.; Goeddel, D. V.; Eaton, D. L.: Stimulation of megakaryocytopoiesis and thrombopoiesis by the c-Mpl ligand. *Nature* 369: 533–538, 1994.

[74640] 12970. Farese, A. M.; Hunt, P.; Grab, L. B.; MacVittie, T. J.: Combined administration of recombinant human megakaryocyte growth and development factor and granulocyte colony-stimulating factor enhances multilineage hematopoietic reconstitution in nonhuman primates after radiation-induced marrow aplasia. *J. Clin. Invest.* 97: 2145–2151, 1996.

[74641] 12971. Foster, D. C.; Sprecher, C. A.; Grant, F. J.; Kramer, J.

M.; Kuijper, J. L.; Holly, R. D.; Whitmore, T. E.; Heipel, M. D.; Bell, L. A.; Ching, A. F. T.; McGrane, V.; Hart, C.; O'Hara, P. J.; Lok, S.: Human thrombopoietin: gene structure, cDNA sequence, expression, and chromosomal localization. *Proc. Nat. Acad. Sci.* 91: 13023–13027, 1994.

[74642] 12972. Ghilardi, N.; Wiestner, A.; Kikuchi, M.; Ohsaka, A.; Skoda, R. C.: Hereditary thrombocythaemia in a Japanese family is caused by a novel point mutation in the thrombopoietin gene. *Brit. J. Haemat.* 107:310–316, 1999.

[74643] 12973. Kaushansky, K.; Lok, S.; Holly, R. D.; Broudy, V. C.; Lin, N.; Bailey, M. C.; Forstrom, J. W.; Buddle, M. M.; Oort, P. J.; Hagen, F. S.; Roth, G. J.; Papayannopoulou, T.; Foster, D. C.: Promotion of megakaryocyte progenitor expansion and differentiation by the c-Mpl ligand thrombopoietin. *Nature* 369: 568–571, 1994.

[74644] 12974. Kondo, T.; Okabe, M.; Sanada, M.; Kurosawa, M.; Suzuki, S.; Kobayashi, M.; Hosokawa, M.; Asaka, M.: Familial essential thrombocythemia associated with one-base deletion in the 5'-prime-untranslated region of the thrombopoietin gene. *Blood* 92: 1091–1096, 1998.

[74645] 12975. Li, J.; Yang, C.; Xia, Y.; Bertino, A.; Glaspy, J.; Roberts, M.; Kuter, D. J.: Thrombocytopenia caused by the development of antibodies to thrombopoietin. *Blood* 98:



3241–3248, 2001.

- [74646] 12976.Lok, S.; Kaushansky, K.; Holly, R. D.; Kuijper, J. L.; Lofton–Day,C. E.; Oort, P. J.; Grant, F. J.; Heipel, M. D.; Burkhead, S. K.;Kramer, J. M.; Bell, L. A.; Sprecher, C. A.; Blumberg, H.; Johnson,R.; Prunkard, D.; Ching, A. F. T.; Mathewes, S. L.; Bailey, M. C.;Forstrom, J. W.; Buddle, M. M.; Osborn, S. G.; Evans, S. J.; Sheppard,P. O.; Presnell, S. R.; O'Hara, P. J.; Hagen, F. S.; Roth, G. J.;Foster, D. C.: Cloning and expression of murine thrombopoietin cD–NAand stimulation of platelet production in vivo. *Nature* 369: 565–568,1994.
- [74647] 12977.Metcalf, D.: Thrombopoietin: at last. *Nature* 369: 519–520,1994.
- [74648] 12978.Methia, N.; Louache, F.; Vainchenker, W.; Wendling, F.: Oligodeoxynucleotidesantisense to the proto–oncogene c–mpl specifically inhibit in vitromegakaryocy–topoiesis. *Blood* 82: 1395–1401, 1993.
- [74649] 12979.Ratajczak, M. Z.; Ratajczak, J.; Marlicz, W.; Pletcher, C. H.,Jr.; Machalinski, B.; Moore, J.; Hung, H.; Gewirtz, A. M.: Recombinanthuman thrombopoietin (TPO) stimulates erythropoiesis by inhibitingerythroid progenitor cell apoptosis. *Brit. J. Haemat.* 98: 8–17,1997.
- [74650] 12980.Sohma, Y.; Akahori, H.; Seki, N.; Hori, T.; Ogami,

K.; Kato, T.; Shimada, Y.; Kawamura, K.; Miyazaki, H.:  
Molecular cloning and chromosomal localization of the human thrombopoietin gene. FEBS Lett. 353: 57–61, 1994.

[74651] 12981. Wendling, F.; Maraskovsky, E.; Debili, N.; Florindo, C.; Teepe, M.; Titeux, M.; Methia, N.; Breton-Gorius, J.; Cosman, D.; Vainchenker, W.: c-Mpl ligand is a humoral regulator of megakaryocytopoiesis. Nature 369: 571–574, 1994.

[74652] 12982. Dualan, R.; Brody, T.; Keeney, S.; Nichols, A. F.; Admon, A.; Linn, S.: Chromosomal localization and cDNA cloning of the genes (DDB1 and DDB2) for the p127 and p48 subunits of a human damage-specific DNA binding protein. Genomics 29: 62–69, 1995.

[74653] 12983. Koken, M. H. M.; Reynolds, P.; Jaspers-Dekker, I.; Prakash, L.; Prakash, S.; Bootsma, D.; Hoeijmakers, J. H. J.: Structural and functional conservation of two human homologs of the yeast DNA repair gene RAD6. Proc. Nat. Acad. Sci. 88: 8865–8869, 1991.

[74654] 12984. Knoll, A.; Schunkert, H.; Reichwald, K.; Danser, A. H. J.; Bauer, D.; Platzer, M.; Stein, G.; Rosenthal, A.: Human renin binding protein: complete genomic sequence and association of an intronic T/C polymorphism with the prorenin level in males. Hum. Molec. Genet. 6:

1527–1534,1997.

- [74655] 12985.Laan, M.; Paabo, S.: Mapping genes by drift-generated linkage disequilibrium. (Letter) Am. J. Hum. Genet. 63: 654–656, 1998.
- [74656] 12986.Takahashi, S.; Inoue, H.; Miyake, Y.: The human gene for renin-binding protein. J. Biol. Chem. 267: 13007–13013, 1992.
- [74657] 12987.Terwilliger, J. D.; Zollner, S.; Laan, M.; Paabo, S.: Mapping genes through the use of linkage disequilibrium generated by genetic drift: 'drift mapping' in small populations with no demographic expansion. Hum. Hered. 48: 138–154, 1998.
- [74658] 12988.van den Ouweland, A. M. W.; Verdijk, M.; Kiochis, P.; Poustka, A.; van Oost, B. A.: The human renin-binding protein gene (RENBP) maps in Xq28. Genomics 21: 279–281, 1994.
- [74659] 12989.Warren, S. T.; Knight, S. J. L.; Peters, J. F.; Stayton, C. L.; Consalez, G. G.; Zhang, F.: Isolation of the human chromosomal band Xq28 within somatic cell hybrids by fragile X site breakage. Proc. Nat. Acad. Sci. 87: 3856–3860, 1990.
- [74660] 12990.Calvo, R. M.; Asuncion, M.; Sancho, J.; San Millan, J. L.; Escobar-Morreale, H. F.: The role of the CAG repeat

polymorphism in the androgen receptorgene and of skewed X-chromosome inactivation, in the pathogenesis of hirsutism. *J. Clin. Endocr. Metab.* 85: 1735–1740, 2000.

[74661] 12991. Gao, X.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Gridley, T.: Assignment of the murine Notch2 and Notch3 genes to chromosomes 3 and 17. *Genomics* 49: 160–161, 1998.

[74662] 12992. Larsson, C.; Lardelli, M.; White, I.; Lendahl, U.: The human NOTCH1, 2, and 3 genes are located at chromosome positions 9q34, 1p13–p11, and 19p13.2–p13.1 in regions of neoplasia-associated translocation. *Genomics* 24:253–258, 1994.

[74663] 12993. Dichgans, M.; Herzog, J.; Gasser, T.: NOTCH3 mutation involving three cysteine residues in a family with typical CADASIL. *Neurology* 57:1714–1717, 2001.

[74664] 12994. Joutel, A.; Corpechot, C.; Ducros, A.; Vahedi, K.; Chabriat, H.; Mouton, P.; Alamowitch, S.; Domenga, V.; Cecillion, M.; Marechal, E.; Maciazek, J.; Vayssiere, C.; Cruaud, C.; Cabanis, E.-A.; Ruchoux, M. M.; Weissenbach, J.; Bach, J. F.; Boussier, M. G.; Tournier-Lasserre, E.: Notch3 mutations in CADASIL, a hereditary adult-onset condition causing stroke and dementia. *Nature* 383: 707–710, 1996.

- [74665] 12995.Joutel, A.; Dodick, D. D.; Parisi, J. E.; Cecillon, M.; Tournier-Lasserre, E.; Bousser, M. G.: De novo mutation in the Notch3 gene causing CADASIL. *Ann.Neurol.* 47: 388–391, 2000.
- [74666] 12996.Joutel, A.; Vahedi, K.; Corpechot, C.; Troesch, A.; Chabriat, H.; Vayssiere, C.; Cruaud, C.; Maciazek, J.; Weissenbach, J.; Bousser, M.-G.; Bach, J.-F.; Tournier-Lasserre, E.: Strong clustering and stereotyped nature of Notch3 mutations in CADASIL patients. *Lancet* 350:1511–1515, 1997.
- [74667] 12997.Rebay, I.; Fehon, R. G.; Artavanis-Tsakonas, S.: Specific truncations of Drosophila Notch define dominant activated and dominant negative forms of the receptor. *Cell* 74: 319–329, 1993.
- [74668] 12998.Deed, R. W.; Hirose, T.; Mitchell, E. L. D.; Santibanez-Koref, M. F.; Norton, J. D.: Structural organisation and chromosomal mapping of the human Id-3 gene. *Gene* 151: 309–314, 1994.
- [74669] 12999.Ellmeier, W.; Aguzzi, A.; Kleiner, E.; Kurzbaue, R.; Weith, A.: Mutually exclusive expression of a helix-loop-helix gene and N-myc in human neuroblastomas and in normal development. *EMBO J.* 11: 2563–2571, 1992.
- [74670] 13000.Kee, B. L.; Rivera, R. R.; Murre, C.: Id3 inhibits B

lymphocyte progenitor growth and survival in response to TGF- $\beta$ . *Nature Immun.* 2:242–247, 2001.

- [74671] 13001. Lyden, D.; Young, A. Z.; Zagzag, D.; Yan, W.; Gerald, W.; O'Reilly, R.; Bader, B. L.; Hynes, R. O.; Zhuang, Y.; Manova, K.; Benezra, R.: Id1 and Id3 are required for neurogenesis, angiogenesis and vascularization of tumour xenografts. *Nature* 401: 670–677, 1999.
- [74672] 13002. Pan, L.; Sato, S.; Frederick, J. P.; Sun, X.-H.; Zhuang, Y.: Impaired immune responses and B-cell proliferation in mice lacking the Id3 gene. *Molec. Cell. Biol.* 19: 5969–5980, 1999.
- [74673] 13003. Yeh, K.; Lim, R. W.: Genomic organization and promoter analysis of the murine Id3 gene. *Gene* 254: 163–171, 2000.
- [74674] 13004. Braun, A.; Kammerer, S.; Weissenhorn, W.; Weiss, E. H.; Cleve, H.: Sequence of a putative human housekeeping gene (HK33) localized on chromosome 1. *Gene* 146: 291–295, 1994.
- [74675] 13005. Gotte, K.; Girzalsky, W.; Linkert, M.; Baumgart, E.; Kammerer, S.; Kunau, W.-H.; Erdmann, R.: Pex19p, a farnesylated protein essential for peroxisome biogenesis. *Molec. Cell. Biol.* 18: 616–628, 1998.
- [74676] 13006. James, G. L.; Goldstein, J. L.; Pathak, R. K.; Ander-

son, R. G.W.; Brown, M. S.: PxF, a prenylated protein of peroxisomes J. Biol.Chem. 269: 14182–14190, 1994.

- [74677] 13007.Kammerer, S.; Arnold, N.; Gutensohn, W.; Mewes, H.–W.; Kunau, W.–H.;Hofler, G.; Roscher, A. A.; Braun, A.: Genomic organization and molecular characterization of a gene encoding HsPXF, a human peroxisomal farnesylated–protein. Genomics 45: 200–210, 1997.
- [74678] 13008.Caskey, C. T.; Pizzuti, A.; Fu, Y.–H.; Fenwick, R. G., Jr.; Nelson,D. L.: Triplet repeat mutations in human disease. Science 256:784–789, 1992.
- [74679] 13009.Chang, B.; Zheng, S. L.; Hawkins, G. A.; Isaacs, S. D.; Wiley,K. E.; Turner, A.; Carpten, J. D.; Bleecker, E. R.; Walsh, P. C.;Trent, J. M.; Meyers, D. A.; Isaacs, W. B.; Xu, J.: Polymorphic GGC repeats in the androgen receptor gene are associated with hereditary and sporadic prostate cancer risk. Hum. Genet. 110: 122–129, 2002.
- [74680] 13010.Chang, C.; Kokontis, J.; Liao, S.: Molecular cloning of human and rat complementary DNA encoding androgen receptors. Science 240:324–326, 1988.
- [74681] 13011.Choong, C. S.; Kemppainen, J. A.; Zhou, Z.–X.; Wilson, E. M.:Reduced androgen receptor gene expression with first exon CAG repeat expansion. Molec. Endocr. 10: 1527–1535, 1996.

- [74682] 13012.Choong, C. S.; Quigley, C. A.; French, F. S.; Wilson, E. M.:A novel missense mutation in the amino-terminal domain of the humanandrogen receptor gene in a family with partial androgen insensitivitysyndrome causes reduced efficiency of protein translation. J. Clin.Invest. 98: 1423–1431, 1996.
- [74683] 13013.Chu, J.; Zhang, R.; Zhao, Z.; Zou, W.; Han, Y.; Qi, Q.; Zhang,H.; Wang, J.–C.; Tao, S.; Liu, X.; Luo, Z.: Male fertility is compatiblewith an Arg840Cys substitution in the AR in a large Chinese familywith divergent phenotypes of AR insensitivity syndrome. J. Clin.Endocr. Metab. 87: 347–351, 2002.
- [74684] 13014.Coetzee, G. A.; Ross, R. K.: Re: Prostate cancer and the androgenreceptor. (Letter) J. Nat. Cancer Inst. 86: 872–873, 1994.
- [74685] 13015.Correa–Cerro, L.; Wohr, G.; Haussler, J.; Berthon, P.; Drelon,E.; Mangin, P.; Fournier, G.; Cussenot, O.; Kraus, P.; Just, W.; Paiss,T.; Cantu, J. M.; Vogel, W.: (CAG)nCAA and GGN repeats in the humanandrogen receptor gene are not associated with prostate cancer ina French–German population. Europ. J. Hum. Genet. 7: 347–362, 1999.
- [74686] 13016.Dowsing, A. T.; Yong, E. L.; Clark, M.; McLachlan, R. I.; de Kretser,D. M.; Trounson, A. O.: Linkage between



male infertility and trinucleotide repeat expansion in the androgen-receptor gene. *Lancet* 354: 640–643, 1999.

[74687] 13017. Elo, J. P.; Kvist, L.; Leinonen, K.; Isomaa, V.; Henttu, P.; Lukkarinen, O.; Vihko, P.: Mutated human androgen receptor gene detected in a prostatic cancer patient is also activated by estradiol. *J. Clin. Endocr. Metab.* 80: 3494–3500, 1995.

[74688] 13018. Gaddipati, J. P.; McLeod, D. G.; Heidenberg, H. B.; Sesterhenn, I. A.; Finger, M. J.; Moul, J. W.; Srivastava, S.: Frequent detection of codon 877 mutation in the androgen receptor gene in advanced prostate cancers. *Cancer Res.* 54: 2861–2864, 1994.

[74689] 13019. Gehring, U.; Tomkins, G. M.: Characterization of a hormone receptor defect in the androgen-insensitivity mutant. *Cell* 3: 59–64, 1974.

[74690] 13020. Gingrich, J. R.; Barrios, R. J.; Kattan, M. W.; Nahm, H. S.; Finegold, M. J.; Greenberg, N. M.: Androgen-independent prostate cancer progression in the TRAMP model. *Cancer Res.* 57: 4687–4691, 1997.

[74691] 13021. Gingrich, J. R.; Greenberg, N. M.: A transgenic mouse prostate cancer model. *Toxicol. Path.* 24: 502–504, 1996.

[74692] 13022. Giovannucci, E.; Stampfer, M. J.; Krithivas, K.;

Brown, M.; Brufsky, A.; Talcott, J.; Hennekens, C. H.; Kantoff, P. W.: The CAG repeat within the androgen receptor gene and its relationship to prostate cancer. *Proc. Nat. Acad. Sci.* 94: 3320–3323, 1997.

[74693] 13023. Gottlieb, B.; Beitel, L. K.; Trifiro, M. A.: Variable expressivity and mutation databases: the androgen receptor gene mutations database. *Hum. Mutat.* 17: 382–388, 2001.

[74694] 13024. Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Pinsky, L.: The androgen receptor gene mutations database. *Nucleic Acids Res.* 25: 158–162, 1997.

[74695] 13025. Gottlieb, B.; Trifiro, M.; Lumbroso, R.; Vasiliou, D. M.; Pinsky, L.: The androgen receptor gene mutations database. *Nucleic Acids Res.* 24: 151–154, 1996.

[74696] 13026. Griffin, J. E.: Androgen resistance—the clinical and molecular spectrum. *New Eng. J. Med.* 326: 611–618, 1992.

[74697] 13027. Grino, P. B.; Isidro-Gutierrez, R. F.; Griffin, J. E.; Wilson, J. D.: Androgen resistance associated with a qualitative abnormality of the androgen receptor and responsive to high dose androgen therapy. *J. Clin. Endocr. Metab.* 68: 578–584, 1989.

[74698] 13028. Hardy, D. O.; Scher, H. I.; Bogenreider, T.; Sabba-

tini, P.; Zhang,Z.-F.; Nanus, D. M.; Catterall, J. F.: Androgen receptor CAG repeatlengths in prostate cancer: correlation with age of onset. J. Clin.Endocr. Metab. 81: 4400-4405, 1996.

[74699] 13029.Hellwinkel, O. J.-C.; Holterhus, P.-M.; Struve, D.; Marschke,C.; Homburg, N.; Hiort, O.: A unique exonic splicing mutation inthe human androgen receptor gene indicates a physiologic relevanceof regular androgen receptor transcript variants. J. Clin. Endocr.Metab. 86: 2569-2575, 2001.

[74700] 13030.Hickey, T.; Chandy, A.; Norman, R. J.: The androgen receptorCAG repeat polymorphism and X-chromosome inactivation in AustralianCaucasian women with infertility related to polycystic ovary syndrome. J.Clin. Endocr. Metab. 87: 161-165, 2002.

[74701] 13031.Hiort, O.; Sinnecker, G. H. G.; Holterhus, P.-M.; Nitsche, E.M.; Kruse, K.: Inherited and de novo androgen receptor gene mutations:investigation of single-case families. J. Pediat. 132: 939-943,1998.

[74702] 13032.Holterhus, P.-M.; Bruggenwirth, H. T.; Hiort, O.; Kleinkauf-Houcken,A.; Kruse, K.; Sinnecker, G. H. G.; Brinkmann, A. O.: Mosaicism dueto a somatic mutation of the androgen receptor gene determines phenotypein an-

drogen insensitivity syndrome. J. Clin. Endocr. Metab. 82:3584–3589, 1997.

- [74703] 13033. Shibuya, H.; Yamaguchi, K.; Shirakabe, K.; Tonegawa, A.; Gotoh, Y.; Ueno, N.; Irie, K.; Nishida, E.; Matsumoto, K.: TAB1: an activator of the TAK1 MAPKKK in TGF-beta signal transduction. Science 272:1179–1182, 1996.
- [74704] 13034. Boyd, J. M.; Subramanian, T.; Schaeper, U.; La Regina, M.; Bayley, S.; Chinnadurai, G.: A region in the C-terminus of adenovirus 2/5E1a protein is required for association with a cellular phosphoprotein and important for the negative modulation of T24-ras mediated transformation, tumorigenesis and metastasis. EMBO J. 469–478, 1993.
- [74705] 13035. Furusawa, T.; Moribe, H.; Kondoh, H.; Higashi, Y.: Identification of CtBP1 and CtBP2 as corepressors of zinc finger-homeodomain factor delta-EF1. Molec. Cell. Biol. 19: 8581–8590, 1999.
- [74706] 13036. Katsanis, N.; Fisher, E. M. C.: A novel C-terminal binding protein (CTBP2) is closely related to CTBP1, an adenovirus E1A-binding protein, and maps to human chromosome 21q21.3. Genomics 47: 294–299, 1998.
- [74707] 13037. Schaeper, U.; Boyd, J. M.; Verma, S.; Uhlmann, E.;

Subramanian,T.; Chinnadurai, G.: Molecular cloning and characterization of acellular phosphoprotein that interacts with a conserved C-terminaldomain of adenovirus E1A involved in negative modulation of oncogenictransformation. Proc. Nat. Acad. Sci. 92: 10467–10471, 1995.

[74708] 13038.Sewalt, R. G. A. B.; Gunster, M. J.; van der Vlag, J.; Satijn,D. P. E.; Otte, A. P.: C-terminal binding protein is a transcriptionalrepressor that interacts with a specific class of vertebrate polycombproteins. Molec. Cell. Biol. 19: 777–787, 1999.

[74709] 13039.Zhang, Q.; Piston, D. W.; Goodman, R. H.: Regulation of corepressorfunction by nuclear NADH. Science 295: 1895–1897, 2002.

[74710] 13040.Watanabe, T.; Inoue, S.; Hiroi, H.; Orimo, A.; Kawashima, H.; Muramatsu,M.: Isolation of estrogen–responsive genes with a CpG island library. Molec.Cell. Biol. 18: 442–449, 1998.

[74711] 13041.Adra, C. N.; Kobayashi, H.; Rowley, J. D.; Lim, B.: Assignmentof the human GDID4 gene, a GDP/GTP–exchange regulator, to chromosome12p12.3. Genomics 24: 188–190, 1994.

[74712] 13042.Leffers, H.; Nielsen, M. S.; Andersen, A. H.; Honore, B.; Madsen,P.; Vandekerckhove, J.; Celis, J. E.: Indentifica–

tion of two human Rho GDP dissociation inhibitor proteins whose overexpression leads to disruption of the actin cytoskeleton. *Exp. Cell Res.* 209: 165–174, 1993.

- [74713] 13043. Lelias, J.-M.; Adra, C. N.; Wulf, G. M.; Guillemot, J.-C.; Khagad, M.; Caput, D.; Lim, B.: cDNA cloning of a human mRNA preferentially expressed in hematopoietic cells and with homology to a GDP-dissociation inhibitor for the rho GTP-binding proteins. *Proc. Nat. Acad. Sci.* 90:1479–1483, 1993.
- [74714] 13044. Scherle, P.; Behrens, T.; Staudt, L. M.: Ly-GDI, a GDP-dissociation inhibitor of the RhoA GTP-binding protein, is expressed preferentially in lymphocytes. *Proc. Nat. Acad. Sci.* 90: 7568–7572, 1993.
- [74715] 13045. Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of coding sequences of unidentified human genes. X. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro. *DNAREs.* 5: 169–176, 1998.
- [74716] 13046. Waldegger, S.; Erdel, M.; Nagl, U. O.; Barth, P.; Raber, G.; Steuer, S.; Utermann, G.; Paulmichl, M.; Lang, F.: Genomic organization and chromosomal localization of the human SGK protein kinase gene. *Genomics* 51:299–302,

1998.

- [74717] 13047. Webster, M. K.; Goya, L.; Ge, Y.; Maiyar, A. C.; Firestone, G.L.: Characterization of sgk, a novel member of the serine/threonine protein kinase gene family which is transcriptionally induced by glucocorticoids and serum. *Molec. Cell. Biol.* 13: 2031–2040, 1993.
- [74718] 13048. Aoki, K.; Ishida, R.; Kasai, M.: Isolation and characterization of a cDNA encoding a translin-like protein, TRAX. *FEBS Lett.* 401:109–112, 1997.
- [74719] 13049. Meng, G.; Aoki, K.; Tokura, K.; Nakahara, K.; Inazawa, J.; Kasai, M.: Genomic structure and chromosomal localization of the gene encoding TRAX, a translin-associated factor X. *J. Hum. Genet.* 45: 305–308, 2000.
- [74720] 13050. Eudy, J. D.; Yao, S.; Weston, M. D.; Ma-Edmonds, M.; Talmadge, C. B.; Cheng, J. J.; Kimberling, W. J.; Sumegi, J.: Isolation of a gene encoding a novel member of the nuclear receptor superfamily from the critical region of Usher syndrome type IIa at 1q41. *Genomics* 50:382–384, 1998.
- [74721] 13051. Greschik, H.; Wurtz, J.-M.; Sanglier, S.; Bourguet, W.; van Dorsselaer, A.; Moras, D.; Renaud, J.-P.: Structural and functional evidence for ligand-independent transcriptional activation by the estrogen-related receptor 3. *Molec. Cell* 9: 303–313, 2002.

- [74722] 13052.Matsuyoshi, N.; Imamura, S.: Multiple cadherins are expressed in human fibroblasts. *Biochem. Biophys. Res. Commun.* 235: 355–358,1997.
- [74723] 13053.Heron, L.; Virsolvy, A.; Peyrolier, K.; Gribble, F. M.; Le Cam,A.; Ashcroft, F. M.; Bataille, D.: Human alpha-endosulfine, a possible regulator of sulfonylurea-sensitive K(ATP) channel: molecular cloning,expression and biological properties. *Proc. Nat. Acad. Sci.* 95:8387–8391, 1998.
- [74724] 13054.Scott, A. F.: Personal Communication. Baltimore, Md. 1/11/2001.
- [74725] 13055.Bertilsson, G.; Heidrich, J.; Svensson, K.; Asman, M.; Jendeberg,L.; Sydow–Backman, M.; Ohlsson, R.; Postlind, H.; Blomquist, P.; Berkenstam,A.: Identification of a human nuclear receptor defines a new signaling pathway for CYP3A induction. *Proc. Nat. Acad. Sci.* 95: 12208–12213,1998.
- [74726] 13056.Blumberg, B.; Sabbagh, W., Jr.; Juguilon, H.; Bolado, J., Jr.;van Meter, C. M.; Ong, E. S.; Evans, R. M.: SXR, a novel steroid and xenobiotic–sensing nuclear receptor. *Genes Dev.* 12: 3195–3205,1998.
- [74727] 13057.Watkins, R. E.; Wisely, G. B.; Moore, L. B.; Collins, J. L.; Lambert,M. H.; Williams, S. P.; Willson, T. M.; Kliewer, S. A.; Redinbo, M.R.: The human nuclear xenobiotic receptor



PXR: structural determinantsof directed promiscuity. *Science* 292: 2329–2333, 2001.

- [74728] 13058.Grenet, J.; Valentine, V.; Kitson, J.; Li, H.; Farrow, S. N.; Kidd,V. J.: Duplication of the DR3 gene on human chromosome 1p36 and itsdeletion in human neuroblas-toma. *Genomics* 49: 385–393, 1998.
- [74729] 13059.Adato, A.; Raskin, L.; Petit, C.; Bonne–Tamir, B.: Deafness heterogeneityin a Druze isolate from the Middle East: novel OTOF and PDS mutations,low prevalence of GJB2 35delG mutation and indication for a new DFNBlo-cus. *Europ. J. Hum. Genet.* 8: 437–442, 2000.
- [74730] 13060.Yasunaga, S.; Grati, M.; Chardenoux, S.; Smith, T. N.; Friedman,T. B.; Lalwani, A. K.; Wilcox, E. R.; Petit, C.: OTOF encodes multiplelong and short isoforms: genetic evidence that the long ones underlierecessive deafness DFNB9. *Am. J. Hum. Genet.* 67: 591–600, 2000.
- [74731] 13061.Yasunaga, S.; Petit, C.: Physical map of the region surroundingthe OTOFERLIN locus on chromosome 2p22–p23. *Genomics* 66: 110–112,2000.
- [74732] 13062.Niederreither, K.; Subbarayan, V.; Dolle, P.; Cham-bon, P.: Embryonicretinoic acid synthesis is essential for early mouse post–implantationdevelopment. *Nature Genet.* 21: 444–448, 1999.

- [74733] 13063. Ono, Y.; Fukuhara, N.; Yoshie, O.: TAL1 and LIM-only protein synergistically induce retinaldehyde dehydrogenase 2 expression in T-cell acute lymphoblastic leukemia by acting as cofactors for GATA3. *Molec. Cell. Biol.* 18: 6939–6950, 1998.
- [74734] 13064. Perlmann, T.: Retinoid metabolism: a balancing act. *Nature Genet.* 31:7–9, 2002. Note: Erratum: *Nature Genet.* 31: 221 only, 2002.
- [74735] 13065. Wang, X.; Penzes, P.; Napoli, J. L.: Cloning of a cDNA encoding an aldehyde dehydrogenase and its expression in *Escherichia coli*: recognition of retinal as substrate. *J. Biol. Chem.* 271: 16288–16293, 1996.
- [74736] 13066. Zhao, D.; McCaffery, P.; Ivins, K. J.; Neve, R. L.; Hogan, P.; Chin, W. W.; Drager, U. C.: Molecular identification of a major retinoic-acid-synthesizing enzyme, a retinaldehyde-specific dehydrogenase. *Europ. J. Biochem.* 240:15–22, 1996.
- [74737] 13067. McGwire, G. B.; Tan, F.; Michel, B.; Rehli, M.; Skidgel, R. A.: Identification of a membrane-bound carboxypeptidase as the mammalian homolog of duck gp180, a hepatitis B virus-binding protein. *Life Sci.* 60: 715–724, 1997.
- [74738] 13068. Riley, D. A.; Tan, F.; Miletich, D. J.; Skidgel, R. A.:

Chromosomal localization of the genes for human carboxypeptidase D (CPD) and the active 50-kilodalton subunit of human carboxypeptidase N (CPN1). *Genomics* 50:105–108, 1998.

[74739] 13069. Tan, F.; Rehli, M.; Krause, S. W.; Skidgel, R. A.: Sequence of human carboxypeptidase D reveals it to be a member of the regulatory carboxypeptidase family with three tandem active site domains. *Biochem. J.* 327: 81–87, 1997.

[74740] 13070. Gebhard, W.; Schube, M.; Eulitz, M.: cDNA cloning and complete primary structure of the small, active subunit of human carboxypeptidase N (kininase 1). *Eur. J. Biochem.* 178: 603–607, 1989.

[74741] 13071. Skidgel, R. A.; Bennett, C. D.; Schilling, J. W.; Tan, F.; Weerasinghe, D. K.; Erdos, E. G.: Amino acid sequence of the N-terminus and selected tryptic peptides of the active subunit of human plasma carboxypeptidase N: comparison with other carboxypeptidases. *Biochem. Biophys. Res. Commun.* 154: 1323–1329, 1988.

[74742] 13072. Rajadhyaksha, A.; Riviere, M.; Van Vooren, P.; Szpirer, J.; Szpirer, C.; Babin, J.; Bina, M.: Assignment of AR1, transcription factor 20 (TCF20), to human chromosome 22q13.3 with somatic cell hybrids and in situ hy-

bridization. *Cytogenet. Cell Genet.* 81: 176–177, 1998.

- [74743] 13073.Sanz, L.; Moscat, J.; Diaz-Meco, M. T.: Molecular characterization of a novel transcription factor that controls stromelysin expression. *Molec.Cell. Biol.* 15: 3164–3170, 1995.
- [74744] 13074.Ding, H.; Descheemaeker, K.; Marynen, P.; Nelles, L.; Carvalho, T.; Carmo-Fonseca, M.; Collen, D.; Belayew, A.: Characterization of a helicase-like transcription factor involved in the expression of the human plasminogen activator inhibitor-1 gene. *DNA Cell Biol.* 15:429–442, 1996.
- [74745] 13075.Lin, Y.; Sheridan, P. L.; Jones, K. A.; Evans, G. A.: The HIP116SNF2/SWI2-related transcription factor gene (SNF2L3) is located on human chromosome 3q25.1–q26.1 *Genomics* 27: 381–382, 1995.
- [74746] 13076.Moinova, H. R.; Chen, W.-D.; Shen, L.; Smiraglia, D.; Olechnowicz, J.; Ravi, L.; Kasturi, L.; Myeroff, L.; Plass, C.; Parsons, R.; Minna, J.; Willson, J. K. V.; Green, S. B.; Issa, J.-P.; Markowitz, S. D.: HLTF gene silencing in human colon cancer. *Proc. Nat. Acad. Sci.* 99:4562–4567, 2002.
- [74747] 13077.Sheridan, P. L.; Schorpp, M.; Voz, M. L.; Jones, K. A.: Cloning of an SNF2/SWI2-related protein that binds specifically to the SPH motifs of the SV40 enhancer and to the HIV-1 promoter. *J. Biol. Chem.* 270:4575–4587, 1995.

- [74748] 13078.Ambros, P. F.; Schmid, J.; Rumpler, S.; Binder, B. R.; de Martin,R.: Localization of the human I-kappa-B kinase-beta (IKBKB) to chromosome8p11.2 by fluorescence in situ hybridization and radiation hybridmapping. *Genomics* 54: 575-576, 1998.
- [74749] 13079.Pasparakis, M.; Courtois, G.; Hafner, M.; Schmidt-Supprian, M.;Nenci, A.; Toksoy, A.; Krampert, M.; Goe-beler, M.; Gillitzer, R.;Israel, A.; Krieg, T.; Rajewsky, K.; Haase, I.: TNF-mediated inflammatory skin disease in mice with epidermis-specific deletion of IKK2. *Nature* 417:861-866, 2002.
- [74750] 13080.Rossi, A.; Kapahi, P.; Natoli, G.; Takahashi, T.; Chen, Y.; Karin,M.; Santoro, M. G.: Anti-inflammatory cyclopentenone prostaglandinsare direct inhibitors of I-kappa-B kinase. *Nature* 403: 103-108,2000.
- [74751] 13081.Shindo, M.; Nakano, H.; Sakon, S.; Yagita, H.; Mi-hara, M.; Okumura,K.: Assignment of I-kappa-B kinase beta (IKBKB) to human chromosomeband 8p12-p11 by in situ hybridization. *Cytogenet. Cell Genet.* 82:32-33, 1998.
- [74752] 13082.Tang, G.; Yang, J.; Minemoto, Y.; Lin, A.: Blocking caspase-3-mediatedproteolysis of IKK-beta suppresses TNF-alpha-induced apoptosis. *Molec.Cell* 8: 1005-1016, 2001.

- [74753] 13083.Woronicz, J. D.; Gao, X.; Cao, Z.; Rothe, M.; Goeddel, D. V.:IkappaB kinase-beta: NF-kappa-B activation and complex formation withIkappaB kinase-alpha and NIK. Science 278: 866-869, 1997.
- [74754] 13084.Yin, M.-J.; Yamamoto, Y.; Gaynor, R. B.: The anti-inflammatoryagents aspirin and salicylate inhibit the activity of I-kappa-B kinase-beta. Nature 396:77-80, 1998.
- [74755] 13085.Zandi, E.; Rothwarf, D. M.; Delhase, M.; Hayakawa, M.; Karin,M.: The IkappaB kinase complex (IKK) contains two kinase subunits,IKKalpha and IKKbeta, necessary for IkappaB phosphorylation and NF-kappa-Bactivation. Cell 91: 243-252, 1997.
- [74756] 13086.Aoki, M.; Hamada, F.; Sugimoto, T.; Sumida, S.; Akiyama, T.; Toyoshima,K.: The human cot proto-oncogene encodes two protein serine/threoninekinases with different transforming activities by alternative initiationof translation. J. Biol. Chem. 268: 22723-22732, 1993.
- [74757] 13087.Kitson, J.; Raven, T.; Jiang, Y.-P.; Goeddel, D. V.; Giles, K.M.; Pun, K.-T.; Grinham, C. J.; Brown, R.; Farrow, S. N.: A death-domain-containingreceptor that mediates apoptosis. Nature 384: 372-375, 1996.
- [74758] 13088.Marsters, S. A.; Sheridan, J. P.; Donahue, C. J.; Pitti,

R. M.; Gray, C. L.; Goddard, A. D.; Bauer, K. D.; Ashkenazi, A.: Apo-3, a new member of the tumor necrosis factor receptor family, contains a death domain and activates apoptosis and NF-kappa-B. *Curr. Biol.* 6:1669-1676, 1996.

[74759] 13089. Screaton, G. R.; Xu, X.-N.; Olsen, A. L.; Cowper, A. E.; Tan, R.; McMichael, A. J.; Bell, J. I.: LARD: a new lymphoid-specific death domain containing receptor regulated by alternative pre-mRNA splicing. *Proc. Nat. Acad. Sci.* 94: 4615-4619, 1997.

[74760] 13090. Wang, E. C. Y.; Thern, A.; Denzel, A.; Kitson, J.; Farrow, S. N.; Owen, M. J.: DR3 regulates negative selection during thymocyte development. *Molec. Cell. Biol.* 21: 3451-3461, 2001.

[74761] 13091. Bataille, V.; Hiles, R.; Bishop, J. A. N.: Retinoblastoma, melanoma and the atypical mole syndrome. *Brit. J. Derm.* 132: 134-138, 1995.

[74762] 13092. Chilosì, M.; Doglioni, C.; Yan, Z.; Lestani, M.; Menestrina, F.; Sorio, C.; Benedetti, A.; Vinante, F.; Pizzolo, G.; Inghirami, G.: Differential expression of cyclin-dependent kinase 6 in cortical thymocytes and T-cell lymphoblastic lymphoma/leukemia. *Am. J. Pathol.* 152:209-217, 1998.

[74763] 13093. Costello, J. F.; Plass, C.; Arap, W.; Chapman, V. M.;

Held, W.A.; Berger, M. S.; Su Huang, H. J.; Cavenee, W. K.: Cyclin-dependent kinase 6 (cdk6) amplification in human gliomas identified using two-dimensional separation of genomic DNA. *Cancer Res.* 57: 1250–1254, 1997.

[74764] 13094. Guan, K.-L.; Jenkins, C. W.; Li, Y.; Nichols, M. A.; Wu, X.; O'Keefe, C. L.; Matera, A. G.; Xiong, Y.: Growth suppression by p18, a p16(INK4/MTS1)- and p14(INK4B/MTS2)-related CDK6 inhibitor, correlates with wild-type pRb function. *Genes Dev.* 8: 2939–2952, 1994.

[74765] 13095. Lien, H.-C.; Lin, C.-W.; Huang, P.-H.; Chang, M.-L.; Hsu, S.-M.: Expression of cyclin-dependent kinase 6 (cdk6) and frequent loss of CD44 in nasal-nasopharyngeal NK/T-cell lymphomas: comparison with CD56-negative peripheral T-cell lymphomas. *Lab. Invest.* 80: 893–900, 2000.

[74766] 13096. Meyerson, M.; Harlow, E.: Identification of G1 kinase activity for cdk6, a novel cyclin D partner. *Molec. Cell. Biol.* 14: 2077–2086, 1994.

[74767] 13097. Shennan, M. G.; Badin, A.-C.; Walsh, S.; Summers, A.; From, L.; McKenzie, M.; Goldstein, A. M.; Tucker, M. A.; Hogg, D.; Lassam, N.: Lack of germline CDK6 mutations in familial melanoma. *Oncogene* 19: 1849–1852, 2000.

[74768] 13098. Traboulsi, E. I.; Zimmerman, L. E.; Manz, H. J.: Cu-



taneous malignant melanoma in survivors of heritable retinoblastoma. Arch. Ophthalmol. 106:1059–1061, 1988.

- [74769] 13099. Blais, A.; Labrie, Y.; Pouliot, F.; Lachance, Y.; Labrie, C.: Structure of the gene encoding the human cyclin-dependent kinase inhibitor p18 and mutational analysis in breast cancer. Biochem. Biophys. Res. Commun. 247: 146–153, 1998.
- [74770] 13100. Lapointe, J.; Lachance, Y.; Labrie, Y.; Labrie, C.: A p18 mutant defective in CDK6 binding in human breast cancer cells. Cancer Res. 56:4586–4589, 1996.
- [74771] 13101. Bontemps, Y.; Maquart, F.-X.; , Wegrowski, Y.: Human UDP-glucose dehydrogenase gene: complete cloning and transcription start mapping. Biochem. Biophys. Res. Commun. 275: 981–985, 2000.
- [74772] 13102. Hempel, J.; Perozich, J.; Romovacek, H.; Hinich, A.; Kuo, I.; Feingold, D. S.: UDP-glucose dehydrogenase from bovine liver: primary structure and relationship to other dehydrogenases. Protein Sci. 3: 1074–1080, 1994.
- [74773] 13103. Marcu, O.; Stathakis, D. G.; Marsh, J. L.: Assignment of the UGDH locus encoding UDP-glucose dehydrogenase to human chromosome band 4p15.1 by radiation hybrid mapping. Cytogenet. Cell Genet. 86: 244–245, 1999.

- [74774] 13104.Spicer, A. P.; Kaback, L. A.; Smith, T. J.; Seldin, M. F.: Molecularcloning and characterization of the human and mouse UDP-glucose dehydrogenasegenes. J. Biol. Chem. 273: 25117–25124, 1998.
- [74775] 13105.Walsh, E. C.; Stainier, D. Y. R.: UDP-glucose dehydrogenase requiredfor cardiac valve formation in zebrafish. Science 293: 1670–1674,2001.
- [74776] 13106.Bernard, M.; Sanseau, P.; Henry, C.; Couturier, A.; Prigent, C.: Cloning of STK13, a third human protein kinase related to DrosophilaAurora and budding yeast Ipl1 that maps on chromosome 19q13.3–ter. Genomics 53:406–409, 1998.
- [74777] 13107.Tseng, T.–C.; Chen, S.–H.; Hsu, Y.–P. P.; Tang, T. K.: Proteinkinase profile of sperm and eggs: cloning and characterization oftwo novel testis-specific protein kinases (AIE1, AIE2) related toyeast and fly chromosome segregation regulators. DNA Cell Biol. 17:823–833, 1998.
- [74778] 13108.Becker, W.; Weber, Y.; Wetzol, K.; Eirimbter, K.; Tejedor, F. J.;Joost, H.–G.: Sequence characteristics, subcellular localization,and substrate specificity of DYRK-related kinases, a novel familyof dual specificity protein kinases. J. Biol. Chem. 273: 25893–25902,1998.
- [74779] 13109.Hsu, H.; Xiong, J.; Goeddel, D. V.: The TNF receptor

1-associated protein TRADD signals cell death and NF- $\kappa$ -B activation. *Cell* 81:495–504, 1995.

[74780] 13110. Pan, M.-G.; Xiong, J.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Goeddel, D. V.: Sequence, genomic organization, and chromosomal localization of the mouse TRADD gene. *J. Inflamm.* 46: 168–175, 1996.

[74781] 13111. Bao, J.; Zervos, A. S.: Isolation and characterization of Nmi, a novel partner of Myc proteins. *Oncogene* 12: 2171–2176, 1996.

[74782] 13112. Zhu, M.; John, S.; Berg, M.; Leonard, W. J.: Functional association of Nmi with Stat5 and Stat1 in IL-2- and IFN- $\gamma$ -mediated signaling. *Cell* 96:121–130, 1999.

[74783] 13113. Phillips, N. J.; Zeigler, M. R.; Deaven, L. L.: A cDNA from the ovarian cancer critical region of deletion on chromosome 17p13.3. *Cancer Lett.* 102: 85–90, 1996.

[74784] 13114. Schultz, D. C.; Vanderveer, L.; Berman, D. B.; Hamilton, T. C.; Wong, A. J.; Godwin, A. K.: Identification of two candidate tumor suppressor genes on chromosome 17p13.3. *Cancer Res.* 56: 1997–2002, 1996.

[74785] 13115. Kirchhausen, T.; Davis, A. C.; Frucht, S.; O'Brine Greco, B.; Payne, G. S.; Tubb, B.: AP17 and AP19, the mammalian small chains of the clathrin-associated protein complexes show homology to Yap17p, their putative ho-

molog in yeast. J. Biol. Chem. 266: 11153–11157, 1991.

[74786] 13116. Peyrard, M.; Parveneh, S.; Lagercrantz, S.; Ekman, M.; Fransson, I.; Sahlen, S.; Dumanski, J. P.: Cloning, expression pattern, and chromosomal assignment to 16q23 of the human gamma-adaptin gene (ADTG). Genomics 50:275–280, 1998.

[74787] 13117. Takatsu, H.; Sakurai, M.; Shin, H.-W.; Murakami, K.; Nakayama, K.: Identification and characterization of novel clathrin adaptor-related proteins. J. Biol. Chem. 273: 24693–24700, 1998.

[74788] 13118. Albarosa, R.; Colombo, B. M.; Roz, L.; Magnani, I.; Pollo, B.; Cirenei, N.; Giani, C.; Conti, A. M.; DiDonato, S.; Finocchiaro, G.: Deletion mapping of gliomas suggests the presence of two small regions for candidate tumor-suppressor genes in a 17-cM interval on chromosome 10q. Am. J. Hum. Genet. 58: 1260–1267, 1996.

[74789] 13119. Maier, D.; Zhang, Z.; Taylor, E.; Hamou, M.-F.; Gratzl, O.; VanMeir, E. G.; Scott, R. J.; Merlo, A.: Somatic deletion mapping on chromosome 10 and sequence analysis of PTEN/MMAC1 point to the 10q25–26 region as the primary target in low-grade and high-grade gliomas. Oncogene 16:3331–3335, 1998.

[74790] 13120. Rasheed, B. K.; McLendon, R. E.; Friedman, H. S.;

Friedman, A.H.; Fuchs, H. E.; Bigner, D. D.; Bigner, S. H.: Chromosome 10 deletion mapping in human gliomas: a common deletion region in 10q25. *Oncogene* 10:2243–2246, 1995.

[74791] 13121. Andre, E.; Conquet, F.; Steinmayr, M.; Stratton, S. C.; Porciatti, V.; Becker–Andre, M.: Disruption of retinoid-related orphan receptor beta changes circadian behavior, causes retinal degeneration and leads to vacillans phenotype in mice. *EMBO J.* 17: 3867–3877, 1998.

[74792] 13122. Paravicini, G.; Steinmayr, M.; Andre, E.; Becker–Andre, M.: The metastasis suppressor candidate nucleotide diphosphate kinase NM23 specifically interacts with members of the ROR/RZR nuclear orphan receptor subfamily. *Biochem. Biophys. Res. Commun.* 227: 82–87, 1996.

[74793] 13123. Sirlin, J. L.: Vacillans, a neurological mutant in the house mouse linked with brown. *J. Genet.* 54: 42–48, 1956.

[74794] 13124. Vidal–Puig, A.; Solanes, G.; Grujic, D.; Flier, J. S.; Lowell, B. B.: UCP3: an uncoupling protein homologue expressed preferentially and abundantly in skeletal muscle and brown adipose tissue. *Biochem. Biophys. Res. Commun.* 235: 79–82, 1997.

[74795] 13125. Vidal–Puig, A. J.; Grujic, D.; Zhang, C.–Y.; Hagen,

T.; Boss, O.; Ido, Y.; Szczepanik, A.; Wade, J.; Mootha, V.; Cortright, R.; Muoio, D. M.; Lowell, B. B.: Energy metabolism in uncoupling protein3 gene knockout mice. *J. Biol. Chem.* 275: 16258–16266, 2000.

[74796] 13126. Lovering, R.; Hanson, I. M.; Borden, K. L. B.; Martin, S.; O'Reilly, N. J.; Evan, G. I.; Rahman, D.; Pappin, D. J. C.; Trowsdale, J.; Freemont, P. S.: Identification and preliminary characterization of a protein motif related to the zinc finger. *Proc. Nat. Acad. Sci.* 90: 2112–2116, 1993.

[74797] 13127. Bourdi, M.; Demady, D.; Martin, J. L.; Jabbour, S. K.; Martin, B. M.; George, J. W.; Pohl, L. R.: cDNA cloning and baculovirus expression of the human liver endoplasmic reticulum P58: characterization as a protein disulfide isomerase isoform, but not as a protease or acarnitine acyltransferase. *Arch. Biochem. Biophys.* 323: 397–403, 1995.

[74798] 13128. Briquet-Laugier, V.; Xia, Y.-R.; Rooke, K.; Mehra-bian, M.; Lysis, A. J.; Doolittle, M. H.: Mapping of three members of the mouse protein disulfide isomerase family. *Mammalian Genome* 9: 176–177, 1998.

[74799] 13129. Hirano, N.; Shibasaki, F.; Sakai, R.; Tanaka, T.; Nishida, J.; Yazaki, Y.; Takenawa, T.; Hirai, H.: Molecular cloning of the human glucose-regulated protein ERp57/GRP58, a thiol-dependent reductase: identification

of its secretory form and inducible expression by the oncogenic transformation. *Europ. J. Biochem.* 234: 336–342, 1995.

[74800] 13130. Koivunen, P.; Helaakoski, T.; Annunen, P.; Veijola, J.; Räsänen, S.; Pihlajaniemi, T.; Kivirikko, K. I.: ERp60 does not substitute for protein disulphide isomerase as the beta-subunit of prolyl 4-hydroxylase. *Biochem. J.* 316: 599–605, 1996.

[74801] 13131. Koivunen, P.; Horelli-Kuitunen, N.; Helaakoski, T.; Karvonen, P.; Jaakkola, M.; Palotie, A.; Kivirikko, K. I.: Structures of the human gene for the protein disulfide isomerase-related polypeptide ERp60 and a processed gene and assignment of these genes to 15q15 and 1q21. *Genomics* 42:397–404, 1997.

[74802] 13132. Oliver, J. D.; van der Wal, F. J.; Bulleid, N. J.; High, S.: Interaction of the thiol-dependent reductase ERp57 with nascent glycoproteins. *Science* 275:86–88, 1997.

[74803] 13133. Didsbury, J.; Weber, R. F.; Bokoch, G. M.; Evans, T.; Snyderman, R.: rac, a novel ras-related family of proteins that are botulinum toxin substrates. *J. Biol. Chem.* 264: 16378–16382, 1989.

[74804] 13134. Palmiter, R. D.; Cole, T. B.; Quaife, C. J.; Findley, S. D.: ZnT3, a putative transporter of zinc into synaptic vesi-

cles. *Proc. Nat. Acad. Sci.* 93: 14934–14939, 1996.

[74805] 13135. Palmiter, R. D.; Findley, S. D.: Cloning and functional characterization of a mammalian zinc transporter that confers resistance to zinc. *EMBO J.* 14: 639–649, 1995.

[74806] 13136. Vallee, B. L.; Falchuk, K. H.: The biochemical basis of zinc physiology. *Physiol. Rev.* 73: 79–118, 1993.

[74807] 13137. Berthelsen, J.; Viggiano, L.; Schulz, H.; Ferretti, E.; Consalez, G. G.; Rocchi, M.; Blasi, F.: PKNOX1, a gene encoding PREP1, a new regulator of Pbx activity, maps on human chromosome 21q22.3 and murine chromosome 17B/C. *Genomics* 47: 323–324, 1998.

[74808] 13138. Chen, H.; Rossier, C.; Nakamura, Y.; Lynn, A.; Chakravarti, A.; Antonarakis, S. E.: Cloning of a novel homeobox-containing gene, PKNOX1, and mapping to human chromosome 21q22.3. *Genomics* 41: 193–200, 1997.

[74809] 13139. Morita, K.; Furuse, M.; Fujimoto, K.; Tsukita, S.: Claudin multigene family encoding four-transmembrane domain protein components of tight junction strands. *Proc. Nat. Acad. Sci.* 96: 511–516, 1999.

[74810] 13140. Chiang, P.-W.; Fogel, E.; Jackson, C. L.; Lieuallen, K.; Lennon, G.; Qu, X.; Wang, S.-Q.; Kurnit, D. M.: Isolation, sequencing, and mapping of the human homologue of the



yeast transcription factor, SPT5. *Genomics* 38: 421–424, 1996.

[74811] 13141. Chiang, P.-W.; Stubbs, L.; Zhang, L.; Kurnit, D. M.: Isolation of murine SPT5 homologue: completion of the isolation and characterization of human and murine homologues of yeast chromatin structural protein complex SPT4, SPT5, and SPT6. *Genomics* 47: 426–428, 1998.

[74812] 13142. Grover, J.; Chen, X.-N.; Korenberg, J. R.; Roughley, P. J.: The structure and chromosome location of the human chondroadherin gene (CHAD). *Genomics* 45: 379–385, 1997.

[74813] 13143. Larsson, T.; Sommarin, Y.; Paulsson, M.; Antonsson, P.; Hedbom, E.; Wendel, M.; Heinegard, D.: Cartilage matrix proteins: a basic 36-kDa protein with a restricted distribution to cartilage and bone. *J. Biol. Chem.* 266: 20428–20433, 1991.

[74814] 13144. Tribioli, C.; Frasch, M.; Lufkin, T.: Bapx1: an evolutionary conserved homologue of the *Drosophila* bagpipe homeobox gene is expressed in planar mesoderm and the embryonic skeleton. *Mech. Dev.* 65: 145–162, 1997.

[74815] 13145. Tribioli, C.; Lufkin, T.: Molecular cloning, chromosomal mapping and developmental expression of BAPX1, a novel human homeobox-containing gene homologous to

*Drosophila bagpipe*. *Gene* 203: 225–233, 1997.

- [74816] 13146. Yoshiura, K.-I.; Murray, J. C.: Sequence and chromosomal assignment of human BAPX1, a bagpipe-related gene, to 4p16.1: a candidate gene for skeletal dysplasia. *Genomics* 45: 425–428, 1997.
- [74817] 13147. de Coo, R. F. M.; Buddiger, P.; Smeets, H. J. M.; van Oost, B. A.: Molecular cloning and characterization of the human mitochondrial NADH: oxidoreductase 10-kDa gene (NDUFV3). *Genomics* 45: 434–437, 1997.
- [74818] 13148. Fritzler, M. J.; Lung, C.-C.; Hamel, J. C.; Griffith, K. J.; Chan, E. K. L.: Molecular characterization of golgin-245, a novel Golgi complex protein containing a granin signature. *J. Biol. Chem.* 270: 31262–31268, 1995.
- [74819] 13149. Chen, L.-S. K.; Lo, C. F.; Numann, R.; Cuddy, M.: Characterization of the human and rat phospholemman (PLM) cDNAs and localization of the human PLM gene to chromosome 19q13.1. *Genomics* 41: 435–443, 1997.
- [74820] 13150. Bianchi, M. C.; Tosetti, M.; Fornai, F.; Alessandri, M. G.; Cipriani, P.; De Vito, G.; Canapicchi, R.: Reversible brain creatine deficiency in two sisters with normal blood creatine level. *Ann. Neurol.* 47: 511–513, 2000.
- [74821] 13151. Humm, A.; Fritsche, E.; Mann, K.; Gohl, M.; Huber, R.: Recombinant expression and isolation of human L-

arginine:glycine amidinotransferase and identification of its active-site cysteine residue. *Biochem.J.* 322: 771–776, 1997.

[74822] 13152. Humm, A.; Huber, R.; Mann, K.: The amino acid sequences of human and pig L-arginine:glycine amidinotransferase. *FEBS Lett.* 339: 101–107, 1994.

[74823] 13153. Item, C. B.; Stockler-Ipsiroglu, S.; Stromberger, C.; Muhl, A.; Alessandri, M. G.; Bianchi, M. C.; Tosetti, M.; Fornai, F.; Cioni, G.: Arginine:glycine amidinotransferase deficiency: the third inborn error of creatine metabolism in humans. *Am. J. Hum. Genet.* 69: 1127–1133, 2001.

[74824] 13154. Yasuda, H.; Shima, N.; Nakagawa, N.; Yamaguchi, K.; Kinosaki, M.; Mochizuki, S.; Tomoyasu, A.; Yano, K.; Goto, M.; Murakami, A.; Tsuda, E.; Morinaga, T.; Higashio, K.; Udagawa, N.; Takahashi, N.; Suda, T.: Osteoclast differentiation factor is a ligand for osteoprotegerin/osteoclastogenesis-inhibitory factor and is identical to TRANCE/RANKL. *Proc. Nat. Acad. Sci.* 95: 3597–3602, 1998.

[74825] 13155. Bey, F.; Silva Pereira, I.; Coux, O.; Viegas-Pequignot, E.; Recillas-Targa, F.; Nothwang, H. G.; Dutrillaux, B.; Scherrer, K.: The prosomal RNA-binding protein p27K is a member of the alpha-type human prosomal gene family. *Molec. Gen. Genet.* 237: 193–205, 1993.

- [74826] 13156. Heard, D. J.; Norby, P. L.; Holloway, J.; Vissing, H.: Human ERR-gamma, a third member of the estrogen receptor-related receptor (ERR) subfamily of orphan nuclear receptors: tissue-specific isoforms are expressed during development in the adult. *Molec. Endocr.* 14: 382-392, 2000.
- [74827] 13157. Hong, H.; Yang, L.; Stallcup, M. R.: Hormone-independent transcriptional activation and coactivator binding by novel orphan nuclear receptor ERR3. *J. Biol. Chem.* 274: 22618-22626, 1999.
- [74828] 13158. Fisher, D. A.; Smith, J. F.; Pillar, J. S.; St. Denis, S. H.; Cheng, J. B.: Isolation and characterization of PDE9A, a novel human cGMP-specific phosphodiesterase. *J. Biol. Chem.* 273: 15559-15564, 1998.
- [74829] 13159. Guipponi, M.; Scott, H. S.; Kudoh, J.; Kawasaki, K.; Shibuya, K.; Shintani, A.; Asakawa, S.; Chen, H.; Lalioti, M. D.; Rossier, C.; Minoshima, S.; Shimizu, N.; Antonarakis, S. E.: Identification and characterization of a novel cyclic nucleotide phosphodiesterase gene (PDE9A) that maps to 21q22.3: alternative splicing of mRNA transcripts, genomic structure and sequence. *Hum. Genet.* 103: 386-392, 1998.
- [74830] 13160. Soderling, S. H.; Bayuga, S. J.; Beavo, J. A.: Identifi-

cation and characterization of a novel family of cyclic nucleotide phosphodiesterases. *J. Biol. Chem.* 273: 15553–15558, 1998.

[74831] 13161. Ishibashi, K.; Kuwahara, M.; Gu, Y.; Kageyama, Y.; Tohsaka, A.; Suzuki, F.; Marumo, F.; Sasaki, S.: Cloning and functional expression of a new water channel abundantly expressed in the testis permeable to water, glycerol, and urea. *J. Biol. Chem.* 272: 20782–20786, 1997.

[74832] 13162. Ishibashi, K.; Yamauchi, K.; Kageyama, Y.; Saito-Ohara, F.; Ikeuchi, T.; Marumo, F.; Sasaki, S.: Molecular characterization of human aquaporin-7 gene and its chromosomal mapping. *Biochim. Biophys. Acta* 1399: 62–66, 1998.

[74833] 13163. Kuriyama, H.; Kawamoto, S.; Ishida, N.; Ohno, I.; Mita, S.; Matsuzawa, Y.; Matsubara, K.; Okubo, K.: Molecular cloning and expression of a novel human aquaporin from adipose tissue with glycerol permeability. *Biochem. Biophys. Res. Commun.* 241: 53–58, 1997.

[74834] 13164. Billin, A. N.; Eilers, A. L.; Queva, C.; Ayer, D. E.: Mlx, a novel Max-like BHLHZip protein that interacts with the Max network of transcription factors. *J. Biol. Chem.* 274: 36344–36350, 1999.

[74835] 13165. Bjerknes, M.; Cheng, H.: TCFL4: a gene at 17q21.1

encoding a putative basic helix-loop-helix leucine-zipper transcription factor. *Gene* 181:7-11, 1996.

- [74836] 13166. Cairo, S.; Merla, G.; Urbinati, F.; Ballabio, A.; Raymond, A.: WBSCR14, a gene mapping to the Williams-Beuren syndrome deleted region, is a new member of the Mlx transcription factor network. *Hum. Molec. Genet.* 10: 617-627, 2001.
- [74837] 13167. Fukuoka, S.-I.; Freedman, S. D.; Scheele, G. A.: A single gene encodes membrane-bound and free forms of GP-2, the major glycoprotein in pancreatic secretory (zymogen) granule membranes. *Proc. Nat. Acad. Sci.* 88: 2898-2902, 1991.
- [74838] 13168. Fukuoka, S.-I.; Freedman, S. D.; Yu, H.; Sukhatme, V. P.; Scheele, G. A.: GP-2/THP gene family encodes self-binding glycosylphosphatidylinositol-anchored proteins in apical secretory compartments of pancreas and kidney. *Proc. Nat. Acad. Sci.* 89: 1189-1193, 1992.
- [74839] 13169. Fukuoka, S.-I.; Suzuki, M.; Okabayashi, K.; Takahashi, E.: Assignment of pancreatic zymogen granule membrane protein GP2 (GP2) to human chromosome band 9q21.11 to q21.2 by in situ hybridization. *Cytogenet. Cell Genet.* 79: 231-232, 1997.
- [74840] 13170. Kooy, J.; Toh, B.-H.; Pettitt, J. M.; Erlich, R.; Glee-

son, P. A.: Human autoantibodies as reagents to conserved Golgi components: characterization of a peripheral, 230-kDa compartment-specific Golgi protein. *J. Biol. Chem.* 267: 20255–20263, 1992.

[74841] 13171. Snow, B. E.; Antonio, L.; Suggs, S.; Gutstein, H. B.; Siderovski, D. P.: Molecular cloning and expression analysis of rat Rgs12 and Rgs14. *Biochem. Biophys. Res. Commun.* 233: 770–777, 1997.

[74842] 13172. Kimple, R. J.; Kimple, M. E.; Betts, L.; Sondek, J.; Siderovski, D. P.: Structural determinants for GoLoco-induced inhibition of nucleotide release by G- $\alpha$  subunits. *Nature* 416: 878–881, 2002.

[74843] 13173. Takesono, A.; et al; et al: Receptor-independent activators of heterotrimeric G-protein signalling pathways. *J. Biol. Chem.* 274: 33202–33205, 1999.

[74844] 13174. Traver, S.; Bidot, C.; Spassky, N.; Baltauss, T.; de Tand, M.-F.; Thomas, J.-L.; Zalc, B.; Janoueix-Lerosey, I.; de Gunzburg, J.: RGS14 is a novel Rap effector that preferentially regulates the GTPase activity of G- $\alpha$ -0. *Biochem. J.* 350: 19–29, 2000.

[74845] 13175. Buckbinder, L.; Velasco-Miguel, S.; Chen, Y.; Xu, N.; Talbott, R.; Gelbert, L.; Gao, J.; Seizinger, B. R.; Gutkind, J. S.; Kley, N.: The p53 tumor suppressor targets a novel

regulator of G proteinsignaling. Proc. Nat. Acad. Sci. 94: 7868–7872, 1997.

- [74846] 13176.Chen, C.-K.; Wieland, T.; Simon, M. I.: RGS-r, a retinal specificRGS protein, binds an intermediate conformation of transducin andenhances recycling. Proc. Nat. Acad. Sci. 93: 12885–12889, 1996.
- [74847] 13177.Snow, B. E.; Antonio, L.; Suggs, S.; Siderovski, D. P.: Cloningof a retinally abundant regulator of G-protein signaling (RGS-r/RGS16):genomic structure and chromosomal localization of the human gene. Gene 206:247–253, 1998.
- [74848] 13178.Zheng, B.; Chen, D.; Farquhar, M. G.: MIR16, a putative membraneglycerophosphodiester phosphodiesterase, interacts with RGS16. Proc.Nat. Acad. Sci. 97: 3999–4004, 2000.
- [74849] 13179.LeBoeuf, R. C.; Caldwell, M.; Guo, Y.; Metz, C.; Davitz, M. A.;Olson, L. K.; Deeg, M. A.: Mouse glycosylphosphatidylinositol-specificphospholipase D (Gpld1) characterization. Mammalian Genome 9: 710–714,1998.
- [74850] 13180.Scallon, B. J.; Fung, W.-J. C.; Tsang, T. C.; Li, S.; Kado-Fong,H.; Huang, K.-S.; Kochan, J. P.: Primary structure and functionalactivity of a phosphatidylinositol-glycan-specific phospholipase D. Science 252:446–448,



1991.

- [74851] 13181.Schofield, J. N.; Rademacher, T. W.: Structure and expression of the human glycosylphosphatidylinositol phospholipase D1 (GPLD1) gene. *Biochim. Biophys. Acta* 1494: 189–194, 2000.
- [74852] 13182.Tsang, T. C.; Fung, W.-J.; Levine, J.; Metz, C. N.; Davitz, M.A.; Burns, D. K.; Huang, K.-S.; Kochan, J. P.: Isolation and expression of two human glycosylphosphatidylinositol phospholipase D (GPI-PLD) cDNAs. (Abstract) *FASEB J. (supp.)* 6: A1922 only, 1992.
- [74853] 13183.Schmitz, F.; Konigstorfer, A.; Sudhof, T. C.: RIBEYE, a component of synaptic ribbons: a protein's journey through evolution provides insight into synaptic ribbon function. *Neuron* 28: 857–872, 2000.
- [74854] 13184.Bergelson, J. M.; Cunningham, J. A.; Droguett, G.; Kurt-Jones, E. A.; Krithivas, A.; Hong, J. S.; Horwitz, M. S.; Crowell, R. L.; Finberg, R. W.: Isolation of a common receptor for coxsackie B viruses and adenoviruses 2 and 5. *Science* 275: 1320–1323, 1997.
- [74855] 13185.Bowles, K. R.; Gibson, J.; Wu, J.; Shaffer, L. G.; Towbin, J. A.; Bowles, N. E.: Genomic organization and chromosomal localization of the human Cocksackievirus B-adenovirus receptor gene. *Hum. Genet.* 105:354–359,

1999.

- [74856] 13186. Carson, S. D.; Chapman, N. N.; Tracy, S. M.: Purification of the putative coxsackievirus B receptor from HeLa cells. *Biochem. Biophys. Res. Commun.* 233: 325–328, 1997.
- [74857] 13187. Griffin, L. D.; Kearney, D.; Ni, J.; Jaffe, R.; Fricker, F. J.; Webber, S.; Demmler, G.; Gelb, B. D.; Towbin, J. A.: Analysis of formalin-fixed and frozen myocardial autopsy samples for viral genome in childhood myocarditis and dilated cardiomyopathy with endocardial fibroelastosis using polymerase chain reaction (PCR). *Cardiovasc. Path.* 4: 3–11, 1995.
- [74858] 13188. Martin, A. B.; Webber, S.; Fricker, F. J.; Jaffe, R.; Demmler, G.; Kearney, D.; Zhang, Y.-H.; Bodurtha, J.; Gelb, B.; Ni, J.; Bricker, J. T.; Towbin, J. A.: Acute myocarditis: rapid diagnosis by PCR in children. *Circulation* 90: 330–339, 1994.
- [74859] 13189. Pauschinger, M.; Bowles, N. E.; Fuentes-Garcia, F. J.; Pham, V.; Kuhl, U.; Schwimmbeck, P. L.; Schultheiss, H.-P.; Towbin, J. A.: Detection of adenoviral genome in the myocardium of adult patients with idiopathic left ventricular dysfunction. *Circulation* 99: 1348–1354, 1999.
- [74860] 13190. Tomko, R. P.; Xu, R.; Philipson, L.: HCAR and

MCAR: the human and mouse receptors for subgroup C adenoviruses and group B coxsackieviruses. *Proc. Nat. Acad. Sci.* 94: 3352–3356, 1997.

[74861] 13191. Margolis, R. L.; Abraham, M. R.; Gatchell, S. B.; Li, S.-H.; Kidwai, A. S.; Breschel, T. S.; Stine, O. C.; Callahan, C.; McInnis, M. G.; Ross, C. A.: cDNAs with long CAG trinucleotide repeats from human brain. *Hum. Genet.* 100: 114–122, 1997.

[74862] 13192. Dale, M.; Hammond, D. W.; Cox, A.; Nicklin, M. J. H.: The human gene encoding the interleukin-1 receptor accessory protein (IL1RAP) maps to chromosome 3q28 by fluorescence in situ hybridization and radiation hybrid mapping. *Genomics* 47: 325–326, 1998.

[74863] 13193. Huang, J.; Gao, X.; Li, S.; Cao, Z.: Recruitment of IRAK to the interleukin 1 receptor complex requires interleukin 1 receptor accessory protein. *Proc. Nat. Acad. Sci.* 94: 12829–12832, 1997.

[74864] 13194. Wesche, H.; Korrherr, C.; Kracht, M.; Falk, W.; Resch, K.; Martin, M. U.: The interleukin-1 receptor accessory protein (IL-1RAcP) is essential for IL-1-induced activation of interleukin-1 receptor-associated kinase (IRAK) and stress-activated protein kinases (SAP kinases). *J. Biol. Chem.* 272: 7727–7731, 1997.

- [74865] 13195.Saha, P.; Chen, J.; Thome, K. C.; Lawlis, S. J.; Hou, Z.-H.; Hendricks,M.; Parvin, J. D.; Dutta, A.: Human CDC6/Cdc18 associates with Orc1and cyclin-cdk and is selectively eliminated from the nucleus at theonset of S phase. *Molec. Cell. Biol.* 18: 2758–2767, 1998.
- [74866] 13196.Williams, R. S.; Shohet, R. V.; Stillman, B.: A human proteinrelated to yeast Cdc6p. *Proc. Nat. Acad. Sci.* 94: 142–147, 1997.
- [74867] 13197.Yan, Z.; DeGregori, J.; Shohet, R.; Leone, G.; Stillman, B.; Nevins,J. R.; Williams, R. S.: Cdc6 is regulated by E2F and is essentialfor DNA replication in mammalian cells. *Proc. Nat. Acad. Sci.* 95:3603–3608, 1998.
- [74868] 13198.Brais, B.; Bouchard, J.-P.; Xie, Y.-G.; Rochefort, D. L.; Chretien,N.; Tome, F. M. S.; Lafreniere, R. G.; Rommens, J. M.; Uyama, E.;Nohira, O.; Blumen, S.; Korczyn, A. D.; Heutink, P.; Mathieu, J.;Duranceau, A.; Codere, F.; Fardeau, M.; Rouleau, G. A.: Short GCGexpansions in the PABP2 gene cause oculopharyngeal muscular dystrophy. *NatureGenet.* 18: 164–167, 1998. Note: Erratum: *Nature Genet.* 19: 404 only,1998.
- [74869] 13199.Calado, A.; Tome, F. M. S.; Brais, B.; Rouleau, G. A.; Kuhn, U.;Wahle, E.; Carmo-Fonseca, M.: Nuclear inclusions in oculopharyngealmuscular dystrophy consist of poly(A)

binding protein 2 aggregates which sequester poly(A) RNA. Hum. Molec. Genet. 9: 2321–2328, 2000.

[74870] 13200. Fan, X.; Dion, P.; Laganier, J.; Brais, B.; Rouleau, G. A.: Oligomerization of polyalanine expanded PABPN1 facilitates nuclear protein aggregation that is associated with cell death. Hum. Molec. Genet. 10: 2341–2351, 2001.

[74871] 13201. Ikeda, S.; Shiva, N.; Ikeda, A.; Smith, R. S.; Nusinowitz, S.; Yan, G.; Lin, T. R.; Chu, S.; Heckenlively, J. R.; North, M. A.; Naggert, J. K.; Nishina, P. M.; Duyao, M. P.: Retinal degeneration but not obesity is observed in null mutants of the tubby-like protein 1 gene. Hum. Molec. Genet. 9: 155–163, 2000.

[74872] 13202. North, M. A.; Naggert, J. K.; Yan, Y.; Noben-Trauth, K.; Nishina, P. M.: Molecular characterization of TUB, TULP1, and TULP2, members of the novel tubby gene family and their possible relation to ocular diseases. Proc. Nat. Acad. Sci. 94: 3128–3133, 1997.

[74873] 13203. Paloma, E.; Hjelmqvist, L.; Bayes, M.; Garcia-Sandoval, B.; Ayuso, C.; Balcells, S.; Gonzalez-Duarte, R.: Novel mutations in the TULP1 gene causing autosomal recessive retinitis pigmentosa. Invest. Ophthalmol. Vis. Sci. 41: 656–659, 2000.

[74874] 13204. Aoki, N.; Ishii, T.; Ohira, S.; Yamaguchi, Y.; Negi,

M.; Adachi,T.; Nakamura, R.; Matsuda, T.: Stage specific expression of milkfat globule membrane glycoproteins in mouse mammary gland: comparison of MFG-E8, butyrophilin, and CD36 with a major milk protein, beta-casein. *Biochim.Biophys. Acta* 1334: 182–190, 1997.

[74875] 13205.Collins, C.; Nehlin, J. O.; Stubbs, J. D.; Kowbel, D.; Kuo, W.-L.; Parry, G.: Mapping of a newly discovered human gene homologous to the apoptosis associated-murine mammary protein, MFG-E8, to chromosome 15q25. *Genomics* 39: 117–118, 1997.

[74876] 13206.Haggqvist, B.; Naslund, J.; Sletten, K.; Westermarck, G. T.; Mucchiano, G.; Tjernberg, L. O.; Nordstedt, C.; Engstrom, U.; Westermarck, P.: Medin: an integral fragment of aortic smooth muscle cell-produced lectin adherin forms the most common human amyloid. *Proc. Nat. Acad.Sci.* 96: 8669–8674, 1999.

[74877] 13207.Hanayama, R.; Tanaka, M.; Miwa, K.; Shinohara, A.; Iwamatsu, A.; Nagata, S.: Identification of a factor that links apoptotic cells to phagocytes. *Nature* 417: 182–187, 2002.

[74878] 13208.Larocca, D.; Peterson, J. A.; Urrea, R.; Kuniyoshi, J.; Bistrain, A. M.; Ceriani, R. L.: A Mr 46,000 human milk fat globule protein that is highly expressed in human breast

tumors contains factor VIII-like domains. *Cancer Res.* 51: 4994–4998, 1991.

[74879] 13209. Stubbs, J. D.; Lekutis, C.; Singer, K. L.; Bui, A.; Yuzuki, D.; Srinivasan, U.; Parry, G.: cDNA cloning of a mouse mammary epithelial cell surface protein reveals the existence of epidermal growth factor-like domains linked to factor VIII-like sequences. *Proc. Nat. Acad. Sci.* 87:8417–8421, 1990.

[74880] 13210. Zhang, J.; Kuehl, P.; Green, E. D.; Touchman, J. W.; Watkins, P. B.; Daly, A.; Hall, S. D.; Maurel, P.; Relling, M.; Brimer, C.; Yasuda, K.; Wrighton, S. A.; Hancock, M.; Kim, R. B.; Strom, S.; Thummel, K.; Russell, C. G.; Hudson, J. R., Jr.; Schuetz, E. G.; Boguski, M. S.: The human pregnane X receptor: genomic structure and identification and functional characterization of natural allelic variants. *Pharmacogenetics* 11:555–572, 2001.

[74881] 13211. Su, L.-K.; Qi, Y.: Characterization of human MAPRE genes and their proteins. *Genomics* 71: 143–149, 2001.

[74882] 13212. Arai, T.; Akiyama, Y.; Okabe, S.; Ando, M.; Endo, M.; Yuasa, Y.: Genomic structure of the human Smad3 gene and its infrequent alterations in colorectal cancers. *Cancer Lett.* 122: 157–163, 1998.

[74883] 13213. Zhang, Y.; Feng, X.-H.; Wu, R.-Y.; Derynck, R.: Re-

ceptor-associated Mad homologues synergize as effectors of the TGF-beta response. *Nature* 383:168-172, 1996.

[74884] 13214. Zhu, Y.; Richardson, J. A.; Parada, L. F.; Graff, J. M.: Smad3 mutant mice develop metastatic colorectal cancer. *Cell* 94: 703-714, 1998.

[74885] 13215. Bruno, E.; Horrigan, S. K.; Van Den Berg, D.; Rozler, E.; Fitting, P. R.; Moss, S. T.; Westbrook, C.; Hoffman, R.: The Smad5 gene is involved in the intracellular signaling pathways that mediate the inhibitory effects of transforming growth factor-beta on human hematopoiesis. *Blood* 91:1917-1923, 1998.

[74886] 13216. Gemma, A.; Hagiwara, K.; Vincent, F.; Ke, Y.; Hancock, A. R.; Nagashima, M.; Bennett, W. P.; Harris, C. C.: hSmad5 gene, a human hSmad family member: its full length cDNA, genomic structure, promoter region and mutation analysis in human tumors. *Oncogene* 16: 951-956, 1998.

[74887] 13217. Cole, A. M.; Kim, Y.-H.; Tahk, S.; Hong, T.; Weis, P.; Waring, A. J.; Ganz, T.: Calcitermin, a novel antimicrobial peptide isolated from human airway secretions. *FEBS Lett.* 504: 5-10, 2001.

[74888] 13218. Guignard, F.; Mauel, J.; Markert, M.: Identification and characterization of a novel human neutrophil protein



related to the S100 family. *Biochem.J.* 309: 395–401, 1995.

[74889] 13219.Ilg, E. C.; Troxler, H.; Burgisser, D. M.; Kuster, T.; Markert, M.; Guignard, F.; Hunziker, P.; Birchler, N.; Heizmann, C. W.: Aminoacid sequence determination of human S100A12 (P6, Calgranulin C, CGRP, CAAF1) by tandem mass spectrometry. *Biochem. Biophys. Res. Commun.* 225:146–150, 1996.

[74890] 13220.Marti, T.; Erttmann, K. D.; Gallin, M. Y.: Host-parasite interaction in human onchocerciasis: identification and sequence analysis of a novel human calgranulin. *Biochem. Biophys. Res. Commun.* 221: 454–458, 1996.

[74891] 13221.Wicki, R.; Marenholz, I.; Mischke, D.; Schafer, B. W.; Heizmann, C. W.: Characterization of the human S100A12 (calgranulin C, p6, CAAF1, CGRP) gene, a new member of the S100 gene cluster on chromosome 1q21. *Cell Calcium* 20: 459–464, 1996.

[74892] 13222.Tan, K. B.; Harrop, J.; Reddy, M.; Young, P.; Terrett, J.; Emery, J.; Moore, G.; Truneh, A.: Characterization of a novel TNF-like ligand and recently described TNF ligand and TNF receptor superfamily genes and their constitutive and inducible expression in hematopoietic and non-hematopoietic cells. *Gene* 204: 35–46, 1997.

- [74893] 13223. Bonini, J. A.; Martin, S. K.; Dralyuk, F.; Roe, M. W.; Philipson, L. H.; Steiner, D. F.: Cloning, expression, and chromosomal mapping of a novel human CC-chemokine receptor (CCR10) that displays high-affinity binding for MCP-1 and MCP-3. *DNA Cell Biol.* 16: 1249–1256, 1997.
- [74894] 13224. Nibbs, R. J. B.; Wylie, S. M.; Yang, J.; Landau, N. R.; Graham, G. J.: Cloning and characterization of a novel promiscuous human beta-chemokine receptor D6. *J. Biol. Chem.* 272: 32078–32083, 1997.
- [74895] 13225. Zieger, B.; Tran, H.; Hainmann, I.; Wunderle, D.; Zgaga-Griesz, A.; Blaser, S.; Ware, J.: Characterization and expression analysis of two human septin genes, PNUTL1 and PNUTL2. *Gene* 261: 197–203, 2000.
- [74896] 13226. Brandt, S.; Jentsch, T. J.: ClC-6 and ClC-7 are two novel broadly expressed members of the ClC chloride channel family. *FEBS Lett.* 377: 15–20, 1995.
- [74897] 13227. Eggermont, J.; Buyse, G.; Voets, T.; Tytgat, J.; De Smedt, H.; Droogmans, G.: Alternative splicing of ClC-6 (a member of the ClC chloride-channel family) transcripts generates three truncated isoforms one of which, ClC-6c, is kidney-specific. *Biochem. J.* 325: 269–276, 1997.
- [74898] 13228. Cleiren, E.; Benichou, O.; Van Hul, E.; Gram, J.; Bollerslav, J.; Singer, F. R.; Beaverson, K.; Aledo, A.; Whyte,

M. P.; Yoneyama, T.; deVernejou, M.-C.; Van Hul, W.: Albers-Schonberg disease (autosomal dominant osteopetrosis, type II) results from mutations in the CLCN7 chloride channel gene. *Hum. Molec. Genet.* 10: 2861–2867, 2001.

[74899] 13229. Hedblom, E.; Kirkness, E. F.: A novel class of GABA-A receptor subunit in tissues of the reproductive system. *J. Biol. Chem.* 272:15346–15350, 1997.

[74900] 13230. Caloca, M. J.; Garcia-Bermejo, M. L.; Blumberg, P. M.; Lewin, N.E.; Kremmer, E.; Mischak, H.; Wang, S.; Nacro, K.; Bienfait, B.; Marquez, V. E.; Kazanietz, M. G.: Beta-2-chimaerin is a novel target for diacylglycerol: binding properties and changes in subcellular localization mediated by ligand binding to its C1 domain. *Proc. Nat. Acad. Sci.* 96: 11854–11859, 1999.

[74901] 13231. Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Cerebellar beta-2-chimaerin, a GTPase-activating protein for p21 Ras-related Rac is specifically expressed in granule cells and has a unique N-terminal SH2 domain. *J. Biol. Chem.* 269: 12888–12892, 1994.

[74902] 13232. Leung, T.; How, B.-E.; Manser, E.; Lim, L.: Germ cell beta-chimaerin, a new GTPase-activating protein for p21rac, is specifically expressed during the acrosomal assembly stage in rat testis. *J. Biol. Chem.* 268:3813–3816,

1993.

- [74903] 13233.Yuan, S.; Miller, D. W.; Barnett, G. H.; Hahn, J. F.; Williams,B. R.: Identification and characterization of human beta 2-chimaerin:association with malignant transformation in astrocytoma. *CancerRes.* 55: 3456–3461, 1995.
- [74904] 13234.Chung, E.; Hanukoglu, A.; Rees, M.; Thompson, R.; Dillon, M.; Hanukoglu,I.; Bistritzer, T.; Kuhnle, U.; Seckl, J.; Gardiner, R. M.: Exclusionof the locus for autosomal recessive pseudohypoaldosteronism type1 from the mineralocorticoid receptor gene region on human chromosome4q by linkage analysis. *J. Clin. Endocr. Metab.* 80: 3341–3345, 1995.
- [74905] 13235.Nagase, T.; Ishikawa, K.; Suyama, M.; Kikuno, R.; Hirosawa, M.;Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Predictionof the coding sequences of unidentified human genes. XIII. The complete sequences of 100 new cDNA clones from brain which code for large proteinsin vitro. *DNA Res.* 6: 63–70, 1999.
- [74906] 13236.Philibert, R. A.; Winfield, S. L.; Sandhu, H. K.; Martin, B. M.;Ginns, E. I.: The structure and expression of the human neuroligin–3gene. *Gene* 246: 303–310, 2000.
- [74907] 13237.Arico, M.; Imashuku, S.; Clementi, R.; Hibi, S.; Ter–

amura, T.; Danesino, C.; Haber, D. A.; Nichols, K. E.: Hemophagocytic lymphohistiocytosis due to germline mutations in SH2D1A, the X-linked lymphoproliferative disease gene. *Blood* 97: 1131–1133, 2001.

[74908] 13238. Arkwright, P. D.; Makin, G.; Will, A. M.; Ayres, M.; Gokhale, D. A.; Fergusson, W. D.; Taylor, G. M.: X linked lymphoproliferative disease in a United Kingdom family. *Arch. Dis. Child.* 79: 52–55, 1998.

[74909] 13239. Bar, R. S.; DeLor, C. J.; Clausen, K. P.; Hurtubise, P.; Henle, W.; Hewetson, J. F.: Fatal infectious mononucleosis in a family. *New Eng. J. Med.* 290: 363–367, 1974.

[74910] 13240. Benoit, L.; Wang, X.; Pabst, H. F.; Dutz, J.; Tan, R.: Cutting edge: defective NK cell activation in X-linked lymphoproliferative disease. *J. Immun.* 165: 3549–3553, 2000.

[74911] 13241. Brandau, O.; Schuster, V.; Weiss, M.; Hellebrand, H.; Fink, F. M.; Kreczy, A.; Friedrich, W.; Strahm, B.; Niemeyer, C.; Belohradsky, B. H.; Meindl, A.: Epstein–Barr virus–negative boys with non–Hodgkin lymphoma are mutated in the SH2D1A gene, as are patients with X-linked lymphoproliferative disease (XLP). *Hum. Molec. Genet.* 8: 2407–2413, 1999.

[74912] 13242. Coffey, A. J.; Brooksbank, R. A.; Brandau, O.;

Oohashi, T.; Howell, G. R.; Bye, J. M.; Cahn, A. P.; Durham, J.; Heath, P.; Wray, P.; Pavitt, R.; Wilkinson, J.; and 31 others: Host response to EBV infection in X-linked lymphoproliferative disease results from mutations in an SH2-domain encoding gene. *Nature Genet.* 20: 129–135, 1998.

[74913] 13243. Czar, M. J.; Kersh, E. N.; Mijares, L. A.; Lanier, G.; Lewis, J.; Yap, G.; Chen, A.; Sher, A.; Duckett, C. S.; Ahmed, R.; Schwartzberg, P. L.: Altered lymphocyte responses and cytokine production in mice deficient in the X-linked lymphoproliferative disease gene SH2D1A/DSHP/SAP. *Proc. Nat. Acad. Sci.* 98: 7449–7454, 2001.

[74914] 13244. Dutz, J. P.; Benoit, L.; Wang, X.; Demetrick, D. J.; Junker, A.; de Sa, D.; Tan, R.: Lymphocytic vasculitis in X-linked lymphoproliferative disease. *Blood* 97: 95–100, 2001.

[74915] 13245. Grierson, H. L.; Skare, J.; Church, J.; Silberman, T.; Davis, J. R.; Kobrinsky, N.; McGregor, R.; Israels, S.; McCarty, J.; Andrews, L. G.; Blecha, T.; Erdman, S.; Obringer, A.; Scharnhorst, D.; Purtilo, D. T.: Evaluation of families wherein a single male manifests a phenotype of X-linked lymphoproliferative disease (XLP). *Am. J. Med. Genet.* 47: 458–463, 1993.

- [74916] 13246.Hambleton, G.; Cottom, D. G.: Familial lymphoma. Proc. Roy.Soc. Med. 62: 1095 only, 1969.
- [74917] 13247.Hamilton, J. K.; Paquin, L. A.; Sullivan, J. L.; Maurer, H. S.;Cruzi, F. G.; Provisor, A. J.; Steuber, C. P.; Hawkins, E.; Yawn,D.; Cornet, J.; Clausen, K.; Finkelstein, G. Z.; Landing, B.; Grunnet,M.; Purtilo, D. T.: X-linked lympho-proliferative syndrome registryreport. J. Pediat. 96: 669-673, 1980.
- [74918] 13248.Harris, A.; Docherty, Z.: X-linked lymphoprolifera-tive disease:a karyotype analysis. Cytogenet. Cell Genet. 47: 92-94, 1988.
- [74919] 13249.Harris, A.; Lenoir, G. M.; Lankester, S. A.: X-linked lymphoproliferativedisease: linkage studies using DNA probes. Clin. Genet. 33: 162-168,1988.
- [74920] 13250.Hayoz, D.; Lenoir, G. M.; Nicole, A.; Pugin, P.; Regamey, C.:X-linked lymphoproliferative syndrome: identification of a large familyin Switzerland. Am. J. Med. 84: 529-534, 1988.
- [74921] 13251.Klein, G.; Klein, E.: Sinking surveillance's flagship. Nature 395:441-445, 1998.
- [74922] 13252.Levine, P. H.; Kamaraju, L. S.; Connelly, R. R.; Be-rard, C. W.;Dorfman, R. F.; Magrath, I.; Easton, J. M.: The American Burkitt'sLymphoma Registry: eight years' expe-

rience. Cancer 49: 1016–1022,1982.

[74923] 13253.Loeffel, S.; Chang, C.–H.; Heyn, R.; Harada, S.; Lipscomb, H.;Sinangil, F.; Volsky, D. J.; McClain, K.; Ochs, H.; Purtilo, D. T.: Necrotizing lymphoid vasculitis in X–linked lymphoproliferativesyndrome. Arch. Path. Lab. Med. 109: 546–550, 1985.

[74924] 13254.Lyon, M. F.; Loutit, J. F.: X–linked factor in acquired immunodeficiencysyndrome?. (Letter) Lancet I: 768 only, 1983.

[74925] 13255.Mulley, J. C.; Turner, A. M.; Gedeon, A. K.; Berdoukas, V. A.;Huang, T. H. M.; Ledbetter, D. H.; Grier–son, H.; Purtilo, D. T.:X–linked lymphoproliferative disease: prenatal detection of an unaffectedhistocompatible male. Clin. Genet. 42: 76–79, 1992.

[74926] 13256.Eva, A.; Aaronson, S. A.: Isolation of a new human oncogene froma diffuse B–cell lymphoma. Nature 316: 273–275, 1985.

[74927] 13257.Anderson, S. R.; Warburg, M.: Norrie's disease. Arch. Ophthal. 66:614–618, 1961.

[74928] 13258.Bergen, A. A. B.; Wapenaar, M. C.; Schuurman, E. J. M.; Diergaarde,P. J.; Lerach, H.; Monaco, A. P.; Bakker, E.; Bleeker–Wagemakers,E. M.; van Ommen, G. J. B.: Detection of a new submicroscopic Norriedisease deletion interval



with a novel DNA probe isolated by differentialAlu PCR fingerprint cloning. Cytogenet. Cell Genet. 62: 231–235,1993.

[74929] 13259.Berger, W.; Meindl, A.; van de Pol, T. J. R.; Cremers, F. P. M.;Ropers, H. H.; Doerner, C.; Monaco, A.; Bergen, A. A. B.; Lebo, R.;Warburg, M.; Zergollern, L.; Lorenz, B.; Gal, A.; Bleeker–Wagemakers,E. M.; Meitinger, T.: Isolation of a candidate gene for Norrie diseaseby positional cloning. Nature Genet. 1: 199–203, 1992.

[74930] 13260.Berger, W.; van de Pol, D.; Bachner, D.; Oerlemans, F.; Winkens,H.; Hameister, H.; Wieringa, B.; Hendriks, W.; Ropers, H.–H.: An animal model for Norrie disease (ND): gene targeting of the mouseND gene. Hum. Molec. Genet. 5: 51–59, 1996.

[74931] 13261.Eva, A.; Vecchio, G.; Rao, C. D.; Tronick, S. R.; Aaronson, S.A.: The predicted DBL oncogene product defines a distinct class of transforming proteins. Proc. Nat. Acad. Sci. 85: 2061–2065, 1988.

[74932] 13262.Galland, F.; Stefanova, M.; Lafage, M.; Birnbaum, D.: Localization of the 5–prime end of the MCF2 oncogene to human chromosome 15q15–q23.Cytogenet. Cell Genet. 60: 114–116, 1992.

[74933] 13263.Grant, S. G.; Mattei, M.–G.; Galland, F.; Stephenson,

D. A.; Keitz, B. T.; Birnbaum, D.; Chapman, V. M.: Localization of the mouse MCF-2(Dbl) protooncogene within a conserved linkage group on the mouse X chromosome. *Cytogenet. Cell Genet.* 54: 175–181, 1990.

[74934] 13264. Nguyen, C.; Pontarotti, P.; Birnbaum, D.; Chimini, G.; Rey, J.A.; Mattei, J.-F.; Jordan, B. R.: Large scale physical mapping in the q27 region of the human X chromosome: the coagulation factor IX gene and the mcf.2 transforming sequence are separated by at most 270 kilobase pairs and are surrounded by several 'HTF islands.' *EMBO J.* 6: 3285–3289, 1987.

[74935] 13265. Nguyen, C.; Poustka, A.-M.; Djabali, M.; Roux, D.; Mattei, J.-F.; Lehrach, H.; Jordan, B. R.: Large-scale mapping and chromosome jumping in the q27 region of the human X chromosome. *Genomics* 5: 298–303, 1989.

[74936] 13266. Noguchi, T.; Mattei, M.-G.; Oberle, I.; Planche, J.; Imbert, J.; Pelassy, C.; Birg, F.; Birnbaum, D.: Localization of the mcf.2 transforming sequence to the X chromosome. *EMBO J.* 6: 1301–1307, 1987.

[74937] 13267. Ron, D.; Tronick, S. R.; Aaronson, S. A.; Eva, A.: Molecular cloning and characterization of the human DBL proto-oncogene: evidence that its overexpression is sufficient to transform NIH/3T3 cells. *EMBO J.* 7: 2465–2473,

1988.

- [74938] 13268.Srivastava, S. K.; Wheelock, R. H. P.; Aaronson, S. A.; Eva, A.: Identification of the protein encoded by the human diffuse B-cell lymphoma (dbl) oncogene. *Proc. Nat. Acad. Sci.* 83: 8868–8872, 1986.
- [74939] 13269.Tronick, S. R.; McBride, O. W.; Popescu, N. C.; Eva, A.: Chromosomal localization of DBL oncogene sequences. *Genomics* 5: 546–553, 1989.
- [74940] 13270.Giovane, A.; Sobieszczuk, P.; Mignon, C.; Mattei, M.–G.; Wasylyk, B.: Locations of the ets subfamily members net, elk1, and sap1 (ELK3, ELK1, and ELK4) on three homologous regions of the mouse and human genomes. *Genomics* 29: 769–772, 1995.
- [74941] 13271.Janz, M.; Lehmann, U.; Olde Weghuis, D.; de Leeuw, B.; Geurts van Kessel, A.; Gilgenkrantz, S.; Hipskind, R. A.; Nordheim, A.: Refined mapping of the human Ets-related gene Elk-1 to Xp11.2–p11.4, distal to the OATL1 region. *Hum. Genet.* 94: 442–444, 1994.
- [74942] 13272.Rao, V. N.; Huebner, K.; Isobe, M.; ar-Rushdi, A.; Croce, C. M.; Reddy, E. S. P.: Elk, tissue-specific ets-related genes on chromosomes X and 14 near translocation breakpoints. *Science* 244: 66–70, 1989.
- [74943] 13273.Tamai, Y.; Taketo, M.; Nozaki, M.; Seldin, M. F.:

Mouse Elk oncogene maps to chromosome X and a novel Elk oncogene (Elk3) maps to chromosome 10. Genomics 26: 414–416, 1995.

- [74944] 13274. Yamauchi, T.; Toko, M.; Suga, M.; Hatakeyama, T.; Isobe, M.: Structural organization of the human Elk1 gene and its processed pseudogene Elk2. DNA Res. 6: 21–27, 1999.
- [74945] 13275. Bermingham, J. R., Jr.; Scherer, S. S.; O'Connell, S.; Arroyo, E.; Kalla, K. A.; Powell, F. L.; Rosenfeld, M. G.: Tst-1/Oct-6/SCIP regulates a unique step in peripheral myelination and is required for normal respiration. Genes Dev. 10: 1751–1762, 1996.
- [74946] 13276. Jaegle, M.; Mandemakers, W.; Broos, L.; Zwart, R.; Karis, A.; Visser, P.; Grosveld, F.; Meijer, D.: The POU factor Oct-6 and Schwann cell differentiation. Science 273: 507–510, 1996.
- [74947] 13277. Sumiyama, K.; Washio-Watanabe, K.; Ono, T.; Yoshida, M. C.; Hayakawa, T.; Ueda, S.: Human class III POU genes, POU3F1 and POU3F3, map to chromosomes 1p34.1 and 3p14.1. Mammalian Genome 9: 180–181, 1998.
- [74948] 13278. Costache, M.; Apoil, P.-A.; Cailleau, A.; Elmgren, A.; Larson, G.; Henry, S.; Blancher, A.; Iordachescu, D.;

Oriol, R.; Mollicone, R.: Evolution of fucosyltransferase genes in vertebrates. *J. Biol. Chem.* 272: 29721–29728, 1997.

[74949] 13279. Yanagidani, S.; Uozumi, N.; Ihara, Y.; Miyoshi, E.; Yamaguchi, N.; Taniguchi, N.: Purification and cDNA cloning of GDP-L-Fuc:N-acetyl-beta-D-glucosaminide:alpha-1-6fucosyltransferase (alpha-1-6 FucT) from human gastric cancer MKN45 cells. *J. Biochem.* 121: 626–632, 1997.

[74950] 13280. Bekri, S.; Adelaide, J.; Merscher, S.; Grosgeorge, J.; Caroli-Bosc, F.; Perucca-Lostanlen, D.; Kelley, P. M.; Pebusque, M.-J.; Theillet, C.; Birnbaum, D.; Gaudray, P.: Detailed map of a region commonly amplified at 11q13–q14 in human breast carcinoma. *Cytogenet. Cell Genet.* 79: 125–131, 1997.

[74951] 13281. Goodwin, R. G.; Alderson, M. R.; Smith, C. A.; Armitage, R. J.; VandenBos, T.; Jerzy, R.; Tough, T. W.; Schoenborn, M. A.; Davis-Smith, T.; Hennen, K.; Falk, B.; Cosman, D.; Baker, E.; Sutherland, G. R.; Grabstein, K. H.; Farrah, T.; Giri, J. G.; Beckmann, M. P.: Molecular and biological characterization of a ligand for CD27 defines a new family of cytokines with homology to tumor necrosis

factor. Cell 73:447–456, 1993.

- [74952] 13282. Wright, R. M.; Vaitaitis, G. M.; Wilson, C. M.; Repine, T. B.; Terada, L. S.; Repine, J. E.: cDNA cloning, characterization, and tissue-specific expression of human xanthine dehydrogenase/xanthine oxidase. Proc. Nat. Acad. Sci. 90: 10690–10694, 1993.
- [74953] 13283. Kawabata, M.; Saeki, K.: Multiple alternative transcripts of the human homologue of the mouse TRAD/R51H3/RAD51D gene, a member of the RecA/RAD51 gene family. Biochem. Biophys. Res. Commun. 257: 156–162, 1999.
- [74954] 13284. Pittman, D. L.; Weinberg, L. R.; Schimenti, J. C.: Identification, characterization, and genetic mapping of Rad51d, a new mouse and human RAD51/RecA-related gene. Genomics 49: 103–111, 1998.
- [74955] 13285. Bell, B.; Scheer, E.; Tora, L.: Identification of hTAFII80–delta links apoptotic signaling pathways to transcription factor TFIID function. Molec. Cell 8: 591–600, 2001.
- [74956] 13286. Hisatake, K.; Ohta, T.; Takada, R.; Guermah, M.; Horikoshi, M.; Nakatani, Y.; Roeder, R. G.: Evolutionary conservation of human TATA-binding–polypeptide-associated factors TAFII31 and TAFII80

and interactions of TAFII80 with otherTAFs and with general transcription factors. Proc. Nat. Acad. Sci. 92:8195–8199, 1995.

- [74957] 13287.Weinzierl, R. O. J.; Ruppert, S.; Dynlacht, B. D.; Tanese, N.;Tjian, R.: Cloning and expression of Drosophila TAFII60 and humanTAFII70 reveal conserved interactions with other subunits of TFIID. EMBOJ. 12: 5303–5309, 1993.
- [74958] 13288.Tsai, K. J.; Chen, S. K.; Ma, Y. L.; Hsu, W. L.; Lee, E. H. Y.: sgk, a primary glucocorticoid–induced gene, facilitates memory consolidationof spatial learning in rats. Proc. Nat. Acad. Sci. 99: 3990–3995,2002.
- [74959] 13289.Waldegger, S.; Barth, P.; Raber, G.; Lang, F.: Cloning and characterizationof a putative human serine/threonine protein kinase transcriptionallymodified during anisotonic and isotonic alterations of cell volume. Proc.Nat. Acad. Sci. 94: 4440–4445, 1997.
- [74960] 13290.Mackay, D. J. G.; Coupe, A.–M.; Shield, J. P. H.; Storr, J. N.P.; Temple, I. K.; Robinson, D. O.: Relaxation of imprinted expressionof ZAC and HYMAI in a patient with transient neonatal diabetes mellitus. Hum.Genet. 110: 139–144, 2002.
- [74961] 13291.Spengler, D.; Villalba, M.; Hoffmann, A.; Pantaloni,

C.; Houssami, S.; Bockaert, J.; Journot, L.: Regulation of apoptosis and cell cycle arrest by Zac1, a novel zinc finger protein expressed in the pituitary gland and the brain. EMBO J. 16: 2814–2825, 1997.

[74962] 13292. Temple, I. K.; James, R. S.; Crolla, J. A.; Sitch, F. L.; Jacobs, P. A.; Howell, W. M.; Betts, P.; Baum, J. D.; Shield, J. P. H.: An imprinted gene(s) for diabetes? (Letter) Nature Genet. 9: 110–112, 1995.

[74963] 13293. Varrault, A.; Ciani, E.; Apiou, F.; Bilanges, B.; Hoffmann, A.; Pantaloni, C.; Bockaert, J.; Spengler, D.; Journot, L.: hZAC encodes a zinc finger protein with antiproliferative properties and maps to a chromosomal region frequently lost in cancer. Proc. Nat. Acad. Sci. 95: 8835–8840, 1998.

[74964] 13294. Jiang, Y.-W.; Veschambre, P.; Erdjument-Bromage, H.; Tempst, P.; Conaway, J. W.; Conaway, R. C.; Kornberg, R. D.: Mammalian mediator of transcriptional regulation and its possible role as an end-point of signal transduction pathways. Proc. Nat. Acad. Sci. 95: 8538–8543, 1998.

[74965] 13295. Adams, M. D.; Soares, M. B.; Kerlavage, A. R.; Fields, C.; Venter, J. C.: Rapid cDNA sequencing (expressed sequence tags) from a directionally cloned human infant brain cDNA library. Nature Genet. 4: 373–380, 1993.



- [74966] 13296.Dai, K.-S.; Liew, C.-C.: A novel human striated muscle RING zincfinger protein, SMRZ, interacts with SMT3b via its RING domain. *J.Biol. Chem.* 276: 23992–23999, 2001.
- [74967] 13297.Abdollahi, A.; Godwin, A. K.; Miller, P. D.; Getts, L. A.; Schultz,D. C.; Taguchi, T.; Testa, J. R.; Hamilton, T. C.: Identificationof a gene containing zinc–finger motifs based on lost expression inmalignantly transformed rat ovarian surface epithelial cells. *CancerRes.* 57: 2029–2034, 1997.
- [74968] 13298.Abdollahi, A.; Roberts, D.; Godwin, A. K.; Schultz, D. C.; Sonoda,G.; Testa, J. R.; Hamilton, T. C.: Identifica–tion of a zinc–fingergene at 6q25: a chromosomal region implicated in development of manysolid tumors. *Onco–gene* 14: 1973–1979, 1997.
- [74969] 13299.Kamiya, M.; Judson, H.; Okazaki, Y.; Kusakabe, M.; Muramatsu, M.;Takada, S.; Takagi, N.; Arima, T.; Wake, N.; Kamimura, K.; Satomura,K.; Hermann, R.; Bonthron, D. T.; Hayashizaki, Y.: The cell cyclecontrol gene ZAC/PLAGL1 is imprinted: a strong candidate gene fortransient neonatal diabetes. *Hum. Molec. Genet.* 9: 453–460, 2000.
- [74970] 13300.Ide, H.; Katoh, M.; Sasaki, H.; Yoshida, T.; Aoki, K.; Nawa, Y.;Osada, Y.; Sugimura, T.; Terada, M.: Cloning of

human bone morphogenetic protein type 1B receptor (BMPR-1B) and its expression in prostate cancer in comparison with other BMPRs. *Oncogene* 14: 1377-1382, 1997.

[74971] 13301. ten Dijke, P.; Yamashita, H.; Ichijo, H.; Franzen, P.; Laiho, M.; Miyazono, K.; Heldin, C.-H.: Characterization of type I receptors for transforming growth factor-beta and activin. *Science* 264: 101-104, 1994.

[74972] 13302. Blixt, A.; Mahlapuu, M.; Bjursell, C.; Darnfors, C.; Johannesson, T.; Enerback, S.; Carlsson, P.: The two-exon gene of the human forkhead transcription factor FREAC-2 (FKHL6) is located at 6p25.3. *Genomics* 53:387-390, 1998.

[74973] 13303. Best, J. L.; Presky, D. H.; Swerlick, R. A.; Burns, D. K.; Chu, W.: Cloning of a full-length cDNA sequence encoding a cdc2-related protein kinase from human endothelial cells. *Biochem. Biophys. Res. Commun.* 208: 562-568, 1995.

[74974] 13304. Garriga, J.; Mayol, X.; Grana, X.: The CDC2-related kinase PITALRE is the catalytic subunit of active multimeric protein complexes. *Biochem. J.* 319: 293-298, 1996.

[74975] 13305. Grana, X.; De Luca, A.; Sang, N.; Fu, Y.; Claudio, P. P.; Rosenblatt, J.; Morgan, D. O.; Giordano, A.: PITALRE, a

nuclear CDC2-related protein kinase that phosphorylates the retinoblastoma protein in vitro. *Proc. Nat. Acad. Sci.* 91: 3834–3838, 1994.

[74976] 13306. Liu, H.; Rice, A. P.: Genomic organization and characterization of promoter function of the human CDK9 gene. *Gene* 252: 51–59, 2000.

[74977] 13307. Triepels, R.; van den Heuvel, L.; Loeffen, J.; Smeets, R.; Trijbels, F.; Smeitink, J.: The nuclear-encoded human NADH:ubiquinone oxidoreductase NDUF8 subunit: cDNA cloning, chromosomal localization, tissue distribution, and mutation detection in complex-I-deficient patients. *Hum. Genet.* 103:557–563, 1998.

[74978] 13308. Bai, C.; Connolly, B.; Metzker, M. L.; Hilliard, C. A.; Liu, X.; Sandig, V.; Soderman, A.; Galloway, S. M.; Liu, Q.; Austin, C. P.; Caskey, C. T.: Overexpression of M68/DCR3 in human gastrointestinal tract tumors independent of gene amplification and its location in a four-gene cluster. *Proc. Nat. Acad. Sci.* 97: 1230–1235, 2000.

[74979] 13309. Pitti, R. M.; Marsters, S. A.; Lawrence, D. A.; Roy, M.; Kischkel, F. C.; Dowd, P.; Huang, A.; Donahue, C. J.; Sherwood, S. W.; Baldwin, D. T.; Godowski, P. J.; Wood, W. I.; Gurney, A. L.; Hillan, K. J.; Cohen, R. L.; Goddard, A. D.; Botstein, D.; Ashkenazi, A.: Genomic amplification of a de-

coy receptor for Fas ligand in lung and coloncancer. Nature 396: 699–703, 1998.

- [74980] 13310. Ringstad, N.; Nemoto, Y.; De Camilli, P.: The SH3p4/Sh3p8/SH3p13 protein family: binding partners for synaptojanin and dynamin via a Grb2-like Src homology 3 domain. Proc. Nat. Acad. Sci. 94: 8569–8574, 1997.
- [74981] 13311. Sittler, A.; Walter, S.; Wedemeyer, N.; Hasenbank, R.; Scherzinger, E.; Eickhoff, H.; Bates, G. P.; Lehrach, H.; Wanker, E. E.: SH3GL3 associates with the huntington exon 1 protein and promotes the formation of polyglutamine-containing protein aggregates. Molec. Cell 2: 427–436, 1998.
- [74982] 13312. Sparks, A. B.; Hoffman, N. G.; McConnell, S. J.; Fowlkes, D. M.; Kay, B. K.: Cloning of ligand targets: systematic isolation of SH3 domain-containing proteins. Nature Biotech. 14: 741–744, 1996.
- [74983] 13313. Chinnaiyan, A. M.; O'Rourke, K.; Yu, G.-L.; Lyons, R. H.; Garg, M.; Duan, D. R.; Xing, L.; Gentz, R.; Ni, J.; Dixit, V. M.: Signal transduction by DR3, a death domain-containing receptor related to TNFR-1 and CD95. Science 274: 990–992, 1996.
- [74984] 13314. Fracchiolla, N. S.; Colombo, G.; Finelli, P.; Maiolo, A. T.; Neri, A.: EHT, a new member of the MTG8/ETO gene family, maps on 20q11 region and is deleted in acute

myeloid leukemias. (Letter) Blood 92:3481–3484, 1998.

- [74985] 13315. Kitabayashi, I.; Ida, K.; Morohoshi, F.; Yokoyama, A.; Mitsuhashi, N.; Shimizu, K.; Nomura, N.; Hayashi, Y.; Ohki, M.: The AML1–MTG8 leukemic fusion protein forms a complex with a novel member of the MTG8 (ETO/CDR) family, MTGR1. Molec. Cell. Biol. 18: 846–858, 1998.
- [74986] 13316. Nelson, H.; Mandiyan, S.; Noumi, T.; Moriyama, Y.; Miedel, M. C.; Nelson, N.: Molecular cloning of cDNA encoding the C subunit of H(+)-ATPase from bovine chromaffin granules. J. Biol. Chem. 265: 20390–20393, 1990.
- [74987] 13317. van Hille, B.; Vanek, M.; Richener, H.; Green, J. R.; Bilbe, G.: Cloning and tissue distribution of subunits C, D, and E of the human vacuolar H(+)-ATPase. Biochem. Biophys. Res. Commun. 197: 15–21, 1993.
- [74988] 13318. Fujiwara, T.; Suzuki, M.; Tanigami, A.; Ikenoue, T.; Omata, M.; Chiba, T.; Tanaka, K.: The BTRC gene, encoding a human F-box/WD40-repeat protein, maps to chromosome 10q24–q25. Genomics 58: 104–105, 1999.
- [74989] 13319. Margottin, F.; Bour, S. P.; Durand, H.; Selig, L.; Benichou, S.; Richard, V.; Thomas, D.; Strebel, K.; Benarous, R.: A novel human WD protein, h-beta TrCp, that interacts with HIV-1 Vpu connects CD4 to the ER degradation pathway through an F-box motif. Molec. Cell

1:565–574, 1998.

- [74990] 13320.Fakruddin, J. M.; Chaganti, R. S. K.; Murty, V. V. V. S.: Lackof BCL10 mutations in germ cell tumors and B cell lymphomas. *Cell* 97:683–688, 1999.
- [74991] 13321.Isaacson, P. G.; Spencer, J.: The biology of low grade MALT lymphoma. *J.Clin. Path.* 48: 395–397, 1995.
- [74992] 13322.Ruland, J.; Duncan, G. S.; Elia, A.; del Barco Barantes, I.; Nguyen,L.; Plyte, S.; Millar, D. G.; Bouchard, D.; Wakeham, A.; Ohashi, P.S.; Mak, T. W.: Bcl10 is a positive regulator of antigen receptor–inducedactivation of NF–kappa–B and neural tube closure. *Cell* 104: 33–42,2001.
- [74993] 13323.Thome, M.; Martinon, F.; Hofmann, K.; Rubio, V.; Steiner, V.; Schneider,P.; Mattmann, C.; Tschopp, J.: Equine herpesvirus–2 E10 gene product,but not its cellular homologue, activates NF–kappa–B transcriptionfactor and c–Jun N–terminal kinase. *J. Biol. Chem.* 274: 9962–9968,1999.
- [74994] 13324.Willis, T. G.; Jadayel, D. M.; Du, M.–Q.; Peng, H.; Perry, A. R.;Abdul–Rauf, M.; Price, H.; Karran, L.; Majeko–dunmi, O.; Wlodarska,I.; Pan, L.; Crook, T.; Hamoudi, R.; Isaacson, P. G.; Dyer, M. J.S.: Bcl10 is involved in t(1;14)(p22;q32) of MALT B cell lymphomaand mutated in multiple tumor types *Cell* 96: 35–45, 1999.

- [74995] 13325. Wotherspoon, A. C.; Pan, L. X.; Diss, T. C.; Isaacson, P. G.: Cytogenetic study of B-cell lymphoma of mucosa-associated lymphoid tissue. *Cancer Genet. Cytogenet.* 58: 35–38, 1992.
- [74996] 13326. Anderson, P.; Nagler-Anderson, C.; O'Brien, C.; Levine, H.; Watkins, S.; Slayter, H. S.; Blue, M. L.; Schlossman, S. F.: A monoclonal antibody reactive with a 15-kDa cytoplasmic granule-associated protein defines a subpopulation of CD8<sup>+</sup> T lymphocytes. *J. Immun.* 144: 574–582, 1990.
- [74997] 13327. Forch, P.; Puig, O.; Kedersha, N.; Martinez, C.; Granneman, S.; Seraphin, B.; Anderson, P.; Valcarcel, J.: The apoptosis-promoting factor TIA-1 is a regulator of alternative pre-mRNA splicing. *Molec. Cell* 6: 1089–1098, 2000.
- [74998] 13328. Kawakami, A.; Tian, Q.; Streuli, M.; Poe, M.; Edelhoff, S.; Distech, C. M.; Anderson, P.: Intron-exon organization and chromosomal localization of the human TIA-1 gene. *J. Immun.* 152: 4937–4945, 1994.
- [74999] 13329. Tian, Q.; Streuli, M.; Saito, H.; Schlossman, S. F.; Anderson, P.: A polyadenylate binding protein localized to the granules of cytolytic lymphocytes induces DNA fragmentation in target cells. *Cell* 67: 629–639, 1991.

- [75000] 13330.Zheng, B.; Albrecht, U.; Kaasik, K.; Sage, M.; Lu, W.; Vaishnav,S.; Li, Q.; Sun, Z. S.; Eichele, G.; Bradley, A.; Lee, C. C.: Nonredundant roles of the mPer1 and mPer2 genes in the mammalian circadian clock. *Cell* 105:683–694, 2001.
- [75001] 13331.Cheng, J.–F.; Boyartchuk, V.; Zhu, Y.: Isolation and mapping of human chromosome 21 cDNA: progress in constructing a chromosome 21 expression map. *Genomics* 23: 75–84, 1994.
- [75002] 13332.Imamura, Y.; Kubota, R.; Wang, Y.; Asakawa, S.; Kudoh, J.; Mashima,Y.; Oguchi, Y.; Shimizu, N.: Human retina–specific amine oxidase(RAO): cDNA cloning, tissue expression, and chromosomal mapping. *Genomics* 40:277–283, 1997.
- [75003] 13333.Imamura, Y.; Noda, S.; Mashima, Y.; Kudoh, J.; Oguchi, Y.; Shimizu,N.: Human retina–specific amine oxidase: genomic structure of the gene (AOC2), alternatively spliced variant, and mRNA expression in retina. *Genomics* 51: 293–298, 1998.
- [75004] 13334.Sirotkin, H.; O'Donnell, H.; DasGupta, R.; Halford, S.; St. Jore,B.; Puech, A.; Parimoo, S.; Morrow, B.; Skoultschi, A.; Weissman, S.M.; Scambler, P.; Kucherlapati, R.: Identification of a new human catenin gene family member



(ARVCF) from the region deleted in velo-cardio-facialsyndrome. Genomics 41: 75–83, 1997.

[75005] 13335.Andrews, N. C.: Mining copper transport genes. Proc. Nat. Acad.Sci. 98: 6543–6545, 2001.

[75006] 13336.Boultwood, J.; Strickson, A. J.; Jabs, E. W.; Cheng, J.-F.; Fidler,C.; Wainscoat, J. S.: Physical mapping of the human ATX1 homologue(HAH1) to the critical region of the 5q- syndrome within 5q32, and immediately adjacent to the SPARC gene. Hum. Genet. 106: 127–129,2000.

[75007] 13337.Hamza, I.; Faisst, A.; Prohaska, J.; Chen, J.; Gruss, P.; Gitlin,J. D.: The metallochaperone Atox1 plays a critical role in perinatalcopper homeostasis. Proc. Nat. Acad. Sci. 98: 6848–6852, 2001.

[75008] 13338.Klomp, L. W. J.; Lin, S.-J.; Yuan, D. S.; Klausner, R. D.; Culotta,V. C.; Gitlin, J. D.: Identification and functional expression ofHAH1, a novel human gene involved in copper homeostasis. J. Biol.Chem. 272: 9221–9226, 1997.

[75009] 13339.Castro, A. M.; Snyder, L. M.: G6PD San Jose: a new variant characterizedby NADPH inhibition studies. Humangenetik 21: 361–363, 1974.

[75010] 13340.Cayanis, E.; Gomperts, E. D.; Balinsky, D.; Disler, P.; Myers,A.: G6PD Hillbrow: a new variant of glucose-6-phosphate dehydrogenaseassociated with drug-in-

duced haemolytic anaemia. Brit. J. Haemat. 30:343–350, 1975.

[75011] 13341.Cayanis, E.; Lane, A. B.; Jenkins, T.; Nurse, G. T.; Balinsky,D.: Glucose–6–phosphate dehydrogenase Por–bandar: a new slow variantwith slightly reduced activity in a South African family of Indiandescent. Biochem. Genet. 15: 765–773, 1977.

[75012] 13342.Cederbaum, A. I.; Beutler, E.: Nonspherocytic hemolytic anemiadue to G6PD Grand Prairie. I.R.C.S. 3: 579, 1975.

[75013] 13343.Chan, T. K.; Todd, D.: Characteristics and distribution of glucose–6–phosphatedehydrogenase–deficient variants in South China. Am. J. Hum. Genet. 24:475–484, 1972.

[75014] 13344.Chan, T. K.; Todd, D.; Lai, M. C. S.: Glucose 6–phosphate dehydrogenase:identity of erythrocyte and leukocyte enzyme with report of a newvariant in Chinese. Biochem. Genet. 6: 119–124, 1972.

[75015] 13345.Chang, J.–G.; Chiou, S.–S.; Perng, L.–I.; Chen, T.–C.; Liu, T.–C.;Lee, L.–S.; Chen, P.–H.; Tang, T. K.: Molecular characterizationof glucose–6–phosphate dehydrogenase (G6PD) deficiency by naturaland amplification created restriction sites: five mutations accountfor most G6PD defi–

ciency cases in Taiwan. *Blood* 80: 1079–1082, 1992.

- [75016] 13346. Chen, E. Y.; Cheng, A.; Lee, A.; Kuang, W.-J.; Hillier, L.; Green, P.; Schlessinger, D.; Ciccodicola, A.; D'Urso, M.: Sequence of human glucose-6-phosphate dehydrogenase cloned in plasmids and a yeast artificial chromosome. *Genomics* 10: 792–800, 1991.
- [75017] 13347. Chen, H.-L.; Huang, M.-J.; Huang, C.-S.; Tang, T. K.: G6PD NanKang(517 T-to-C; 173 phe-to-leu): a new Chinese G6PD variant associated with neonatal jaundice. *Hum. Hered.* 46: 201–204, 1996.
- [75018] 13348. Chernyak, N. B.; Batischev, A. I.; Lanzina, N. V.; Tokarev, Y. N.; Alexeyev, G. A.: Electrophoretic and kinetic properties of glucose-6-phosphate dehydrogenase from erythrocytes or patients with hemolytic anemia, related to deficiency of the enzyme activity. *Vopr. Med. Khim.* 23:166–171, 1977.
- [75019] 13349. Childs, B.; Zinkham, W.; Browne, E. A.; Kimbro, E. L.; Torbert, J. V.: A genetic study of a defect in glutathione metabolism of the erythrocytes. *Bull. Johns Hopkins Hosp.* 102: 21–37, 1958.
- [75020] 13350. Chiu, D. T. Y.; Zuo, L.; Chao, L.; Chen, E.; Louie, E.; Lubin, B.; Liu, T. Z.; Du, C. S.: Molecular characterization of glucose-6-phosphate dehydrogenase (G6PD) deficiency in

patients of Chinese descent and identification of new base substitutions in the human G6PD gene. *Blood* 81:2150–2154, 1993.

[75021] 13351. Chockkalingam, K.; Board, P. G.: Further evidence for heterogeneity of glucose-6-phosphate dehydrogenase deficiency in Papua New Guinea. *Hum. Genet.* 56: 209–212, 1980.

[75022] 13352. Chockkalingam, K.; Board, P. G.; Breguet, G.: Glucose-6-phosphate dehydrogenase variants of Bali Island (Indonesia). *Hum. Genet.* 60:60–62, 1982.

[75023] 13353. Chockkalingam, K.; Board, P. G.; Nurse, G. T.: Glucose-6-phosphate dehydrogenase deficiency in Papua New Guinea: the description of 13 new variants. *Hum. Genet.* 60: 189–192, 1982.

[75024] 13354. Chuanshu, D.; Yankang, X.; Lin, W. Q. R.; Xiaoyun, H.: Studies on erythrocyte glucose-6-phosphate dehydrogenase variants in Chinese. I. Gd(-) Lizu-Baisha. *Acta Acad. Med. Zhong Shan* 2: 649–658, 1981.

[75025] 13355. Cocco, P.; Todde, P.; Fornera, S.; Manca, M. B.; Manca, P. Sias, A. R.: Mortality in a cohort of men expressing the glucose-6-phosphate dehydrogenase deficiency. *Blood* 91: 706–709, 1998.

[75026] 13356. Cohn, J.; Carter, N.; Warburg, M.: Glucose-

6-phosphate dehydrogenase deficiency in a native Danish family: a new variant. *Scand. J. Haemat.* 23:403–406, 1979.

[75027] 13357.Colonna–Romano, S.; Iolascon, A.; Lippo, S.; Pinto, L.; Cutillo, S.; Battistuzzi, G.: Genetic heterogeneity at the glucose–6-phosphate dehydrogenase locus in southern Italy: a study on the population of Naples. *Hum. Genet.* 69: 228–232, 1985.

[75028] 13358.Cooper, D. W.; Johnston, P. G.; Murtagh, C. E.; Sharman, G. B.; Vandeberg, J. L.; Poole, W. E.: Sex-linked isozymes and sex-chromosome evolution and inactivation in kangaroos. In: Markert, C. L.: *Isozymes. Developmental Biology*. New York: Academic Press (pub.) III: 1975. Pp. 559–573.

[75029] 13359.Cooper, M. R.; Dechatelet, L. R.; McCall, C. E.; Lavia, M. F.; Spurr, C. L.; Baehner, R. L.: Complete deficiency of leukocyte glucose–6-phosphate dehydrogenase with defective bactericidal activity. *J. Clin. Invest.* 51:769–778, 1972.

[75030] 13360.Corash, L.; Spielberg, S.; Bartsocas, C.; Boxer, L.; Steinherz, R.; Sheetz, M.; Egan, M.; Schlesselman, J.; Schulman, J. D.: Reduced chronic hemolysis during high-dose vitamin E administration in Mediterranean-type glu-

cose-6-phosphate dehydrogenase deficiency. New Eng. J. Med. 303:416-420, 1980.

[75031] 13361. Corcoran, C. M.; Calabro, V.; Tamagnini, G.; Town, M.; Haidar, B.; Vulliamy, T. J.; Mason, P. J.; Luzzatto, L.: Molecular heterogeneity underlying the G6PD Mediterranean phenotype. Hum. Genet. 88: 688-690, 1992.

[75032] 13362. Costa, E.; Cabeda, J. M.; Vieira, E.; Pinto, R.; Pereira, S. A.; Ferraz, L.; Santos, R.; Barbot, J.: Glucose-6-phosphate dehydrogenase Aveiro: a de novo mutation associated with chronic nonspherocytic hemolytic anemia. Blood 95: 1499-1501, 2000.

[75033] 13363. Crookston, J. H.; Yoshida, A.; Lin, M.; Booser, D. J.: G6PD Toronto. Biochem. Genet. 8: 259-265, 1973.

[75034] 13364. Csepregy, M.; Hall, M. K.; Berkow, R. L.; Jackson, S.; Prchal, J. T.: Characterization of a new G6PD variant: G6PD Titusville. Am. J. Med. Sci. 297: 114-117, 1989.

[75035] 13365. Csepregy, M.; Yeilding, A.; Lilly, M.; Hall, K.; Scott, C. W.; Prchal, J. T.: Characterization of a new glucose-6-phosphate dehydrogenase variant: G6PD Central City. Am. J. Hemat. 28: 61-62, 1988.

[75036] 13366. D'Urso, M.; Luzzatto, L.; Perroni, L.; Ciccodicola, A.; Gentile, G.; Peluso, I.; Persico, M. G.; Pizzella, T.; Toniolo, D.; Vulliamy, T. J.: An extensive search for RFLP in the hu-

man glucose-6-phosphatedehydrogenase locus has revealed a silent mutation in the coding sequence. *Am.J. Hum. Genet.* 42: 735–741, 1988.

- [75037] 13367. Basse, F.; Stout, J. G.; Sims, P. J.; Wiedmer, T.: Isolation of an erythrocyte membrane protein that mediates  $\text{Ca}^{2+}$ -dependent transbilayer movement of phospholipid. *J. Biol. Chem.* 271: 17205–17210, 1996.
- [75038] 13368. Garrett, R. M.; Johnson, J. L.; Graf, T. N.; Feigenbaum, A.; Rajagopalan, K. V.: Human sulfite oxidase R160Q: identification of the mutation in a sulfite oxidase-deficient patient and expression and characterization of the mutant enzyme. *Proc. Nat. Acad. Sci.* 95: 6394–6398, 1998.
- [75039] 13369. Kisker, C.; Schindelin, H.; Pacheco, A.; Wehbi, W. A.; Garrett, R. M.; Rajagopalan, K. V.; Enemark, J. H.; Rees, D. C.: Molecular basis of sulfite oxidase deficiency from the structure of sulfite oxidase. *Cell* 91: 973–983, 1997.
- [75040] 13370. Corral, J.; Forster, A.; Thompson, S.; Lampert, F.; Kaneko, Y.; Slater, R.; Kroes, W. G.; van der Schoot, C. E.; Ludwig, W.-D.; Karpas, A.; Pocock, C.; Cotter, F.; Rabbitts, T. H.: Acute leukemias of different lineages have similar MLL gene fusions encoding related chimeric proteins resulting from chromosomal translocation. *Proc. Nat. Acad.*

Sci. 90:8538–8542, 1993.

- [75041] 13371. Parry, P.; Wei, Y.; Evans, G.: Cloning and characterization of the t(X;11) breakpoint from a leukemic cell line identify a new member of the forkhead gene family. *Genes Chromosomes Cancer* 11: 79–84, 1994.
- [75042] 13372. Peters, U.; Haberhausen, G.; Kostrzewa, M.; Nolte, D.; Muller, U.: AFX1 and p54(nrb): fine mapping, genomic structure, and exclusion as candidate genes of X-linked dystonia parkinsonism. *Hum. Genet.* 100:569–572, 1997.
- [75043] 13373. Bohme, B.; VandenBos, T.; Cerretti, D. P.; Park, L. S.; Holtrich, U.; Rubsamen-Waigmann, H.; Strebhardt, K.: Cell-cell adhesion mediated by binding of membrane-anchored ligand LERK-2 to the EPH-related receptor human embryonal kinase 2 promotes tyrosine kinase activity. *J. Biol. Chem.* 271: 24747–24752, 1996.
- [75044] 13374. Fletcher, F. A.; Huebner, K.; Shaffer, L. G.; Fairweather, N. D.; Monaco, A. P.; Muller, U.; Druck, T.; Simoneaux, D. K.; Chelly, J.; Belmont, J. W.; Beckmann, M. P.; Lyman, S. D.: Assignment of the gene (EPLG2) encoding a high-affinity binding protein for the receptor tyrosine kinase Elk to a 200-kilobase pair region in human chromosome Xq12. *Genomics* 25: 334–335, 1995.
- [75045] 13375. Fletcher, F. A.; Renshaw, B.; Hollingsworth, T.;



Baum, P.; Lyman, S. D.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Davison, B. L.: Genomic organization and chromosomal localization of mouse *Eplg2*, a gene encoding a binding protein for the receptor tyrosine kinase *Elk*. *Genomics* 24: 127–132, 1994.

[75046] 13376. Palmer, A.; Zimmer, M.; Erdmann, K. S.; Eulenburg, V.; Porthin, A.; Heumann, R.; Deutsch, U.; Klein, R.: EphrinB phosphorylation and reverse signaling: regulation by Src kinases and PTP-BL phosphatase. *Molec. Cell* 9: 725–737, 2002.

[75047] 13377. Bourke, E.; Frindt, G.; Flynn, P.; Schreiner, G. E.: Primary hyperoxaluria with normal alpha-ketoglutarate:glyoxylate carboxylase activity: treatment with isocarboxazid. *Ann. Intern. Med.* 76: 279–284, 1972.

[75048] 13378. Danpure, C. J.: Personal Communication. Middlesex, England 6/16/1988.

[75049] 13379. Danpure, C. J.: Primary hyperoxaluria type 1 and peroxisome-to-mitochondrion mistargeting of alanine:glyoxylate aminotransferase. *Biochimie* 75: 309–315, 1993.

[75050] 13380. Danpure, C. J.; Jennings, P. R.: Peroxisomal alanine:glyoxylate aminotransferase deficiency in primary hyperoxaluria type I. *FEBS Lett.* 201: 20–24, 1986.

- [75051] 13381.Danpure, C. J.; Jennings, P. R.; Watts, R. W. E.: Enzymological diagnosis of primary hyperoxaluria type 1 by measurement of hepatic alanine:glyoxylate aminotransferase activity. *Lancet* I: 289–291, 1987.
- [75052] 13382.Brunelli, S.; Faiella, A.; Capra, V.; Nigro, V.; Simone, A.; Cama, A.; Boncinelli, E.: Germline mutation in the homeobox gene EMX2 in patients with severe schizencephaly. *Nature Genet.* 12: 94–96, 1996.
- [75053] 13383.Rulten, S.; Thorpe, J.; Kay, J.: Identification of eukaryotic parvulin homologues: a new subfamily of peptidylprolyl cis–trans isomerases. *Biochem. Biophys. Res. Commun.* 259: 557–562, 1999.
- [75054] 13384.Barton, D. E.; Yang–Feng, T. L.; Francke, U.: The human tyrosine aminotransferase gene mapped to the long arm of chromosome 16 (region 16q22–q24) by somatic cell hybrid analysis and in situ hybridization. *Hum. Genet.* 72: 221–224, 1986.
- [75055] 13385.Beinfang, D. C.; Kuwabara, T.; Pueschel, S. M.: The Richner–Hanhart syndrome: report of a case with associated tyrosinemia. *Arch. Ophthalmol.* 94:1133–1137, 1976.
- [75056] 13386.Bohnert, A.; Anton–Lamprecht, I.: Richner–Hanhart's syndrome: ultrastructural abnormalities of epidermal keratinization indicating a causal relationship to high in–

tracellular tyrosine levels. J. Invest.Derm. 79: 68–74, 1982.

[75057] 13387.Buist, N.: Phenylketonuria and related problems.In: Nyhan, W.L.: Amino Acid Metabolism and Genetic Variation. New York: McGraw–Hill(pub.) 1967. Pp. 117 only.

[75058] 13388.Chitayat, D.; Balbul, A.; Hani, V.; Mamer, O. A.; Clow, C.; Scriver,C. R.: Hereditary tyrosinaemia type II in a consanguineous Ashkenazijewish family: intrafamilial variation in phenotype; absence of parentalphenotype effects on the fetus. J. Inherit. Metab. Dis. 15: 198–203,1992.

[75059] 13389.Crovato, F.; Desirello, G.; Gatti, R.; Babbini, N.; Rebora, A.: Richner–Hanhart syndrome spares a plantar autograft. Arch. Derm. 121:539–540, 1985.

[75060] 13390.Fellman, J. H.; Vanbellinghen, P. J.; Jones, R. T.; Koler, R. D.: Soluble and mitochondrial forms of tyrosine aminotransferase: relationship to human tyrosinemia. Biochemistry 8: 615–622, 1969.

[75061] 13391.Garibaldi, L. R.; Siliato, F.; De Martini, I.; Scarsi, M. R.; Romano,C.: Oculocutaneous tyrosinosis: report of two cases in the same family. Helv.Paediat. Acta 32: 173–180, 1977.

[75062] 13392.Goldsmith, L. A.; Kang, E. S.; Bienfang, D. C.; Jim–

bow, K.; Gerald, P. S.; Baden, H. P.: Tyrosinemia with plan-  
tar and palmar keratosis and keratitis. J. Pediat. 83:  
798–805, 1973.

[75063] 13393. Hanhart, E.: Neue Sonderformen von Keratosis  
palmo-plantaris, u.a. eine regelmässig-dominante mit  
systematisierten Lipomen, ferner 2 einfach-rezessive mit  
Schwachsinn und z.T. mit Hornhautveränderungen des  
Auges (Ektodermatosyndrom). Dermatologica 94:  
286–308, 1947.

[75064] 13394. Hunziker, N.: Richner-Hanhart syndrome and ty-  
rosinemia type II. Dermatologica 160:180–189, 1980.

[75065] 13395. Kennaway, N. G.; Buist, N. R. M.: Metabolic studies  
in a patient with hepatic cytosol tyrosine aminotransferase  
deficiency. Pediat. Res. 5: 287–297, 1971.

[75066] 13396. Muller, G.; Scherer, G.; Zentgraf, H.; Ruppert, S.;  
Herrmann, B.; Lehrach, H.; Schutz, G.: Isolation, characteri-  
zation and chromosomal mapping of the mouse tyrosine  
aminotransferase gene. J. Molec. Biol. 184:367–373,  
1985.

[75067] 13397. Natt, E.; Kao, F.-T.; Rettenmeier, R.; Scherer, G.:  
Assignment of the human tyrosine aminotransferase gene  
to chromosome 16. Hum. Genet. 72: 225–228, 1986.

[75068] 13398. Huang, L.; Gitschier, J.: A novel gene involved in

zinc transportis deficient in the lethal milk mouse. *Nature Genet.* 17: 292–297,1997.

[75069] 13399.McKusick, V. A.: *On the X Chromosome of Man.* Washington: Am.Inst. Biol. Sci. (pub.) 1964.

[75070] 13400.Ichinose, H.; Ohye, T.; Suzuki, T.; Sumi-Ichinose, C.; Nomura,T.; Hagino, Y.; Nagatsu, T.: Molecular cloning of the human Nurr1gene: characterization of the human gene and cDNAs. *Gene* 230: 233–239,1999.

[75071] 13401.Law, S. W.; Conneely, O. M.; DeMayo, F. J.; O'Malley, B. W.: Identificationof a new brain-specific transcription factor, NURR1. *Molec. Endocr.* 6:2129–2135, 1992.

[75072] 13402.Mages, H. W.; Rilke, O.; Bravo, R.; Senger, G.; Kroczeck, R. A.: NOT, a human immediate-early response gene closely related to thesteroid/thyroid hormone receptor NAK1/TR3. *Molec. Endocr.* 8: 1583–1591,1994.

[75073] 13403.McEvoy, A. N.; Murphy, E. A.; Ponnio, T.; Conneely, O. M.; Bresnihan,B.; FitzGerald, O.; Murphy, E. P.: Activation of nuclear orphan receptorNURR1 transcription by NF-kappaB and cyclic adenosine 5-prime-monophosphateresponse element-binding protein in rheumatoid arthritis synovialtissue. *J. Immun.* 168: 2979–2987, 2002.

- [75074] 13404.Okabe, T.; Takayanagi, R.; Imasaki, K.; Haji, M.; Nawata, H.; Watanabe,T.: cDNA cloning of a NGFI-B/nur77-related transcription factor from an apoptotic human T cell line. *J. Immun.* 154: 3871–3879, 1995.
- [75075] 13405.Torii, T.; Kwarai, T.; Nakamura, S.; Kawakami, H.: Organization of the human orphan nuclear receptor Nurr1 gene. *Gene* 230: 225–232,1999.
- [75076] 13406.Xu, P.-Y.; Liang, R.; Jankovic, J.; Hunter, C.; Zeng, Y.-X.; Ashizawa,T.; Lai, D.; Le, W.-D.: Association of homozygous 7048G7049 variant in the intron six of Nurr1 gene with Parkinson's disease. *Neurology* 58:881–884, 2002.
- [75077] 13407.Hanahan, D.: Signaling vascular morphogenesis and maintenance. *Science* 277:48–50, 1997.
- [75078] 13408.Wong, S. M. E.; Lowe, A. W.: Sequence of the cDNA encoding humanGP-2, the major membrane protein in the secretory granule of the exocrine pancreas. *Gene* 171: 311–312, 1996.
- [75079] 13409.Yamamura, T.; Hitomi, J.; Nagasaki, K.; Suzuki, M.; Takahashi,E.; Saito, S.; Tsukada, T.; Yamaguchi, K.: Human CAAF1 gene--molecular cloning, gene structure, and chromosome mapping. *Biochem. Biophys. Res. Commun.* 221: 356–360, 1996.

- [75080] 13410.Miyoshi, J.; Higashi, T.; Mukai, H.; Ohuchi, T.; Kakunaga, T.:Structure and transforming potential of the human cot oncogene encodinga putative protein kinase. Molec. Cell. Biol. 11: 4088–4096, 1991.
- [75081] 13411.Ohara, R.; Miyoshi, J.; Aoki, M.; Toyoshima, K.: The murine cotproto–oncogene: genome structure and tissue–specific expression. Jpn.J. Cancer Res. 84: 518–525, 1993.
- [75082] 13412.Gould, D. B.; Walter, M. A.: Cloning, characteriza–tion, localization,and mutational screening of the human BARX1 gene. Genomics 68: 336–342,2000.
- [75083] 13413.Tissier–Seta, J. P.; Mucchielli, M. L.; Mark, M.; Mattei, M. G.;Goridis, C.; Brunet, J. F.: Barx1, a new mouse homeodomain transcriptionfactor expressed in cranio–fa–cial ectomesenchyme and the stomach. Mech.Dev. 51: 3–15, 1995.
- [75084] 13414.Ishihara, H.; Shibasaki, Y.; Kizuki, N.; Wada, T.; Yazaki, Y.;Asano, T.; Oka, Y.: Type I phosphatidylinositol–4–phosphate 5–kinases:cloning of the third isoform and deletion/substitution analysis ofmembers of this novel lipid kinase family. J. Biol. Chem. 273: 8741–8748,1998.
- [75085] 13415.Besset, S.; Vincourt, J.–B.; Amalric, F.; Girard, J.–P.: Nuclearlocalization of PAPS synthetase 1: a sulfate activa–tion pathway inthe nucleus of eukaryotic cells. FASEB J. 14:

345–354, 2000.

- [75086] 13416. Faiyaz ul Haque, M.; King, L. M.; Krakow, D.; Cantor, R. M.; Rusiniak, M. E.; Swank, R. T.; Superti-Furga, A.; Haque, S.; Abbas, H.; Ahmad, W.; Ahmad, M.; Cohn, D. H.: Mutations in orthologous genes in human spondyloepimetaphyseal dysplasia and the brachymorphic mouse. *Nature Genet.* 20: 157–162, 1998.
- [75087] 13417. Geller, D. H.; Henry, J. G.; Belch, J.; Schwartz, N. B.: Co-purification and characterization of ATP-sulfurylase and adenosine-5-prime-phosphosulfate kinase from rat chondrosarcoma. *J. Biol. Chem.* 262: 7374–7382, 1987.
- [75088] 13418. Girard, J.-P.; Baekkevold, E. S.; Amalric, F.: Sulfation in high endothelial venules: cloning and expression of the human PAPS synthetase. *FASEBJ.* 12: 603–612, 1998.
- [75089] 13419. Lyle, S.; Stanczak, J.; Ng, K.; Schwartz, N. B.: Rat chondrosarcoma ATP sulfurylase and adenosine 5-prime-phosphosulfate kinase reside on a single bifunctional protein. *Biochemistry* 33: 5920–5925, 1994.
- [75090] 13420. Venkatachalam, K. V.; Akita, H.; Strott, C. A.: Molecular cloning, expression, and characterization of human bifunctional 3-prime-phosphoadenosine 5-prime-phosphosulfate synthase and its functional domains. *J. Biol. Chem.* 273:



19311–19320, 1998.

- [75091] 13421. Park, Y. C.; Ye, H.; Hsia, C.; Segal, D.; Rich, R. L.; Liou, H.-C.; Myszka, D. G.; Wu, H.: A novel mechanism of TRAF signaling revealed by structural and functional analyses of the TRADD–TRAF2 interaction. *Cell* 101:777–787, 2000.
- [75092] 13422. Asao, H.; Sasaki, Y.; Arita, T.; Tanaka, N.; Endo, K.; Kasai, H.; Takeshita, T.; Endo, Y.; Fujita, T.; Sugamura, K.: Hrs is associated with STAM, a signal–transducing adaptor molecule: its suppressive effect on cytokine–induced cell growth. *J. Biol. Chem.* 272: 32785–32791, 1997.
- [75093] 13423. Takeshita, T.; Arita, T.; Asao, H.; Tanaka, N.; Higuchi, M.; Kuroda, H.; Kaneko, K.; Munakata, H.; Endo, Y.; Fujita, T.; Sugamura, K.: Cloning of a novel signal–transducing adaptor molecule containing an SH3 domain and ITAM. *Biochem. Biophys. Res. Commun.* 225: 1035–1039, 1996.
- [75094] 13424. Everett, R. D.; Meredith, M.; Orr, A.; Cross, A.; Kathoria, M.; Parkinson, J.: A novel ubiquitin–specific protease is dynamically associated with the PML nuclear domain and binds to a herpesvirus regulatory protein. *EMBO J.* 16: 1519–1530, 1997. Note: Corrected republication of *EMBO J.*: 16: 566–577, 1997.

- [75095] 13425.Li, M.; Chen, D.; Shiloh, A.; Luo, J.; Nikolaev, A. Y.; Qin, J.;Gu, W.: Deubiquitination of p53 by HAUSP is an important pathwayfor p53 stabilization. *Nature* 416: 648–653, 2002.
- [75096] 13426.Robinson, P. A.; Lomonte, P.; Leek, J. P.; Markham, A. F.; Everett,R. D.: Assignment of herpesvirus–associated ubiquitin–specific proteasegene HAUSP to human chromosome band 16p13.3 by in situ hybridization. *Cytogenet.Cell Genet.* 83: 100 only, 1998.
- [75097] 13427.Zhou, G.; Bao, Z. Q.; Dixon, J. E.: Components of a new humanprotein kinase signal transduction pathway. *J. Biol. Chem.* 270:12665–12669, 1995.
- [75098] 13428.Lee, J.–D.; Ulevitch, R. J.; Han, J.: Primary structure of BMK1:a new mammalian MAP kinase. *Biochem. Biophys. Res. Commun.* 213:715–724, 1995.
- [75099] 13429.Purandare, S. M.; Lee, J.–D.; Patel, P. I.: Assignment of bigMAP kinase (PRKM7) to human chromosome 17 band p11.2 with somaticcell hybrids. *Cytogenet. Cell Genet.* 83: 258–259, 1998.
- [75100] 13430.Regan, C. P.; Li, W.; Boucher, D. M.; Spatz, S.; Su, M. S.; Kuida,K.: Erk5 null mice display multiple extraembryonic vascular and embryoniccardiovascular defects. *Proc. Nat. Acad. Sci.* 99: 9248–9253, 2002.

- [75101] 13431. Barlow, G. M.; Chen, X.-N.; Shi, Z. Y.; Lyons, G. E.; Kurnit, D. M.; Celle, L.; Spinner, N. B.; Zackai, E.; Pettenati, M. J.; Van Riper, A. J.; Vekemans, M. J.; Mjaatvedt, C. H.; Korenberg, J. R.: Down syndrome congenital heart disease: a narrowed region and a candidate gene. *Genet. Med.* 3: 91–101, 2001.
- [75102] 13432. Schmucker, D.; Clemens, J. C.; Shu, H.; Worby, C. A.; Xiao, J.; Muda, M.; Dixon, J. E.; Zipursky, S. L.: *Drosophila* Dscam is an axon guidance receptor exhibiting extraordinary molecular diversity. *Cell* 101:671–684, 2000.
- [75103] 13433. Yamakawa, K.; Huo, Y.-K.; Haendel, M. A.; Hubert, R.; Chen, X.-N.; Lyons, G. E.; Korenberg, J. R.: DSCAM: a novel member of the immunoglobulin superfamily maps in a Down syndrome region and is involved in the development of the nervous system. *Hum. Molec. Genet.* 7: 227–237, 1998.
- [75104] 13434. Zhu, J.; Petersen, S.; Tessarollo, L.; Nussenzweig, A.: Targeted disruption of the Nijmegen breakage syndrome gene NBS1 leads to early embryonic lethality in mice. *Curr. Biol.* 11: 105–109, 2001.
- [75105] 13435. Cooper, P. R.; Smilnich, N. J.; Day, C. D.; Nowak, N. J.; Reid, L. H.; Pearsall, R. S.; Reece, M.; Prawitt, D.; Lan-

ders, J.; Housman, D. E.; Winterpacht, A.; Zabel, B. U.; Pelletier, J.; Weissman, B. E.; Shows, T. B.; Higgins, M. J.: Divergently transcribed overlapping genes expressed in liver and kidney and located in the 11p15.5 imprinted domain. *Genomics* 49: 38–51, 1998.

- [75106] 13436. Kershaw, D. B.; Beck, S. G.; Wharram, B. L.; Wiggins, J. E.; Goyal, M.; Thomas, P. E.; Wiggins, R. C.: Molecular cloning and characterization of human podocalyxin-like protein: orthologous relationship to rabbit PCLP1 and rat podocalyxin. *J. Biol. Chem.* 272: 15708–15714, 1997.
- [75107] 13437. Kershaw, D. B.; Wiggins, J. E.; Wharram, B. L.; Wiggins, R. C.: Assignment of the human podocalyxin-like protein (PODXL) gene to 7q32–q33. *Genomics* 45: 239–240, 1997.
- [75108] 13438. Chan, K. K.; Tsui, S. K. W.; Lee, S. M. Y.; Luk, S. C. W.; Liew, C. C.; Fung, K. P.; Waye, M. M. Y.; Lee, C. Y.: Molecular cloning and characterization of FHL2, a novel LIM domain protein preferentially expressed in human heart. *Gene* 210: 345–350, 1998.
- [75109] 13439. Genini, M.; Schwalbe, P.; Scholl, F. A.; Remppis, A.; Mattei, M.-G.; Schafer, B. W.: Subtractive cloning and characterization of DRAL, a novel LIM-domain protein down-regulated in rhabdomyosarcoma. *DNACell Biol.* 16:

433–442, 1997.

- [75110] 13440.Claverie–Martin, F.; Wang, M.; Cohen, S. N.: ARD–1 cDNA from humancells encodes a site–specific single–strand endoribonuclease thatfunctionally resembles Escherichia coli RNase E. *J. Biol. Chem.* 272:13823–13828, 1997.
- [75111] 13441.Van Eynde, A.; Wera, S.; Beullens, M.; Torrekens, S.; Van Leuven,F.; Stalmans, W.; Bollen, M.: Molecular cloning of NIPP–1, a nuclearinhibitor of protein phosphatase–1, reveals homology with polypeptidesinvolved in RNA processing. *J. Biol. Chem.* 270: 28068–28074, 1995.
- [75112] 13442.Tekin, M.; Dogu, F.; Tacyildiz, N.; Akar, E.; Ikin–ciogullari,A.; Ogur, G.; Yavuz, G.; Babacan, E.; Akar, N.: 657del5 mutationin the NBS1 gene is associated with Nijmegen breakage syndrome ina Turkish family. *Clin. Genet.* 62: 84–88, 2002.
- [75113] 13443.Varon, R.; Reis, A.; Henze, G.; Einsiedel, H. G.; Sperling, K.;Seeger, K.: Mutations in the Nijmegen break–age syndrome gene (NBS1)in childhood acute lymphoblastic leukemia (ALL). *Cancer Res.* 61:3570–3572, 2001.
- [75114] 13444.Wilda, M.; Demuth, I.; Concannon, P.; Sperling, K.; Hameister,H.: Expression pattern of the Nijmegen break–age syndrome gene, Nbs1,during murine development.

Hum. Molec. Genet. 9: 1739–1744, 2000.

- [75115] 13445.Wu, X.; Ranganathan, V.; Weisman, D. S.; Heine, W. F.; Ciccone, D. N.; O'Neill, T. B.; Crick, K. E.; Pierce, K. A.; Lane, W. S.; Rathbun, G.; Livingston, D. M.; Weaver, D. T.: ATM phosphorylation of Nijmegen breakage syndrome protein is required in a DNA damage response. Nature 405:477–482, 2000.
- [75116] 13446.Zhao, S.; Weng, Y.-C.; Yuan, S.-S. F.; Lin, Y.-T.; Hsu, H.-C.; Lin, S.-C. J.; Gerbino, E.; Song, M.; Zdzienicka, M. Z.; Gatti, R. A.; Shay, J. W.; Ziv, Y.; Shiloh, Y.; Lee, E. Y.-H. P.: Functional link between ataxia-telangiectasia and Nijmegen breakage syndrome gene products. Nature 405: 473–477, 2000.
- [75117] 13447.Mankodi, A.; Urbinati, C. R.; Yuan, Q.-P.; Moxley, R. T.; Sansone, V.; Krym, M.; Henderson, D.; Schalling, M.; Swanson, M. S.; Thornton, C. A.: Muscleblind localizes to nuclear foci of aberrant RNA in myotonic dystrophy types 1 and 2. Hum. Molec. Genet. 10: 2165–2170, 2001.
- [75118] 13448.Hittner, H. M.; Kretzer, F. L.; Antoszyk, J. H.; Ferrell, R. E.; Mehta, R. S.: Variable expressivity of autosomal dominant anterior segment mesenchymal dysgenesis in six generations. Am. J. Ophthalmol. 93:57–70, 1982.
- [75119] 13449.Semina, E. V.; Ferrell, R. E.; Mintz-Hittner, H. A.;

Bitoun, P.; Alward, W. L. M.; Reiter, R. S.; Funkhauser, C.; Daack-Hirsch, S.; Murray, J. C.: A novel homeobox gene PITX3 is mutated in families with autosomal-dominant cataracts and ASMD. *Nature Genet.* 19: 167–170, 1998.

[75120] 13450. Semina, E. V.; Murray, J. C.; Reiter, R.; Hrstka, R. F.; Graw, J.: Deletion in the promoter region and altered expression of Pitx3 homeobox gene in aphakia mice. *Hum. Molec. Genet.* 9: 1575–1585, 2000.

[75121] 13451. Semina, E. V.; Reiter, R. S.; Murray, J. C.: Isolation of a new homeobox gene belonging to the Pitx/Rieg family: expression during lens development and mapping to the aphakia region on mouse chromosome 19. *Hum. Molec. Genet.* 6: 2109–2116, 1997.

[75122] 13452. Kent, J.; Lee, M.; Schedl, A.; Boyle, S.; Fantes, J.; Powell, M.; Rushmere, N.; Abbott, C.; van Heyningen, V.; Bickmore, W. A.: The reticulocalbin gene maps to the WAGR region in human and to the Smalley-Harwell deletion in mouse. *Genomics* 42: 260–267, 1997.

[75123] 13453. Ozawa, M.: Cloning of a human homologue of mouse reticulocalbin reveals conservation of structural domains in the novel endoplasmic reticulum resident  $\text{Ca}^{2+}$ -binding protein with multiple EF-hand motifs. *J. Biochem.* 117: 1113–1119, 1995.

- [75124] 13454.Ozawa, M.; Muramatsu, T.: Reticulocalbin, a novel endoplasmicreticulum resident  $\text{Ca}^{2+}$ -binding protein with multiple EF-hand motifsand a carboxyl-terminal HDEL sequence. J. Biol. Chem. 268: 699-705,1993.
- [75125] 13455.Luo, G.; Leroy, E.; Kozak, C. A.; Polymeropoulos, M. H.; Horowitz,R.: Mapping of the gene (NRAP) encoding N-RAP in the mouse and humangenomes. Genomics 45: 229-232, 1997.
- [75126] 13456.Luo, G.; Zhang, J. Q.; Nguyen, T.-P.; Herrera, A. H.; Paterson,B.; Horowitz, R.: Complete cDNA sequence and tissue localizationof N-RAP, a novel nebulin-related protein of striated muscle. CellMotil. Cytoskeleton 38: 75-90, 1997.
- [75127] 13457.Ichikawa, S.; Ozawa, K.; Hirabayashi, Y.: Assignment of a UDP-glucose:ceramideglucosyltransferase gene (Ugcg) to mouse chromosome band 4B3 by insitu hybridization. Cytogenet. Cell Genet. 83: 14-15, 1998.
- [75128] 13458.Ichikawa, S.; Ozawa, K.; Hirabayashi, Y.: Assignment of a UDP-glucose:ceramideglucosyltransferase gene (UGCG) to human chromosome band 9q31 by insitu hybridization. Cytogenet. Cell Genet. 79: 233-234, 1997.
- [75129] 13459.Ichikawa, S.; Sakiyama, H.; Suzuki, G.; Hidari, K. I.-P. J.; Hirabayashi,Y.: Expression cloning of a cDNA for



human ceramide glucosyltransferase that catalyzes the first glycosylation step of glycosphingolipid synthesis. Proc.Nat. Acad. Sci. 93: 4638–4643, 1996.

- [75130] 13460.Watanabe, R.; Wu, K.; Paul, P.; Marks, D. L.; Kobayashi, T.; Pittelkow, M. R.; Pagano, R. E.: Up-regulation of glucosylceramide synthase expression and activity during human keratinocyte differentiation. J.Biol. Chem. 273: 9651–9655, 1998.
- [75131] 13461.Strahm, B.; Rittweiler, K.; Duffner, U.; Brandau, O.; Orłowska-Volk, M.; Karajannis, M. A.; zur Stadt, U.; Tie-mann, M.; Reiter, A.; Brandis, M.; Meindl, A.; Niemeyer, C. M.: Recurrent B-cell non-Hodgkin's lymphoma in two brothers with X-linked lymphoproliferative disease with-out evidence for Epstein-Barr virus infection. Brit. J. Haemat. 108:377–382, 2000.
- [75132] 13462.Sylla, B. S.; Murphy, K.; Cahir-McFarland, E.; Lane, W. S.; Mosialos, G.; Kieff, E.: The X-linked lymphoproliferative syndrome gene product SH2D1A associates with p62(dok) (Dok1) and activates NF-kappa-beta. Proc.Nat. Acad. Sci. 97: 7470–7475, 2000.
- [75133] 13463.Sylla, B. S.; Wang, Q.; Hayoz, D.; Lathrop, G. M.; Lenoir, G.M.: Multipoint linkage mapping of the Xq25–q26 region in a family affected by the X-linked lymphoprolifer-

ative syndrome. Clin. Genet. 36:359–462, 1989.

- [75134] 13464. Tangye, S. G.; Lazetic, S.; Woollatt, E.; Sutherland, G. R.; Lanier, L. L.; Phillips, J. H.: Cutting edge: human 2B4, an activating NK cell receptor, recruits the protein tyrosine phosphatase SHP-2 and the adaptor signaling protein SAP. J. Immun. 162: 6981–6985, 1999.
- [75135] 13465. Tangye, S. G.; Phillips, J. H.; Lanier, L. L.; Nichols, K. E.: Cutting edge: functional requirement for SAP in 2B4-mediated activation of human natural killer cells as revealed by the X-linked lymphoproliferative syndrome. J. Immun. 165: 2932–2936, 2000.
- [75136] 13466. Thorley-Lawson, D. A.; Schooley, R. T.; Bhan, A. K.; Nadler, L. M.: Epstein-Barr virus superinduces a new human B cell differentiation antigen (B-LAST 1) expressed on transformed lymphoblasts. Cell 30:415–425, 1982.
- [75137] 13467. Vowels, M. R.; Lam-Po-Tang, R.; Berdoukas, V.; Ford, D.; Thierry, D.; Purtilo, D.; Gluckman, E.: Correction of X-linked lymphoproliferative disease by transplantation of cord-blood stem cells. New Eng. J. Med. 329: 1623–1625, 1993.
- [75138] 13468. Williams, L. L.; Rooney, C. M.; Conley, M. E.; Brenner, M. K.; Krance, R. A.; Heslop, H. E.: Correction of Duncan's syndrome by allogeneic bone marrow transplanta-

tion. *Lancet* 342: 587–588, 1993.

- [75139] 13469. Wu, C.; Nguyen, K. B.; Pien, G. C.; Wang, N.; Gullo, C.; Duncan, H.; Sosa, M. R.; Edwards, M. J.; Borrow, P.; Satoskar, A. R.; Sharpe, A. H.; Biron, C. A.; Terhorst, C.: SAP controls T cell response to virus and terminal differentiation of T(H)2 cells. *Nature Immun.* 2:410–414, 2001.
- [75140] 13470. Wyandt, H. E.; Skare, J. C.; Grierson, H. L.; Purtilo, D. T.; Milunsky, A.: Detection of a chromosomal deletion of Xq25 in an affected male with X-linked lymphoproliferative disease. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A108 only, 1989.
- [75141] 13471. Yin, L.; Ferrand, V.; Lavoue, M.-F.; Hayoz, D.; Philippe, N.; Souillet, G.; Seri, M.; Giacchino, R.; Castagnola, E.; Hodgson, S.; Sylla, B. S.; Romeo, G.: SH2D1A mutation analysis for diagnosis of XLP in typical and atypical patients. *Hum. Genet.* 105: 501–505, 1999.
- [75142] 13472. Yin, L.; Tocco, T.; Pauly, S.; Lenoir, G. M.; Romeo, G.: Absence of SH2D1A point mutation in 62 Burkitts lymphoma cell lines. *Am. J. Hum. Genet.* 65 (suppl. 1868): A331 only, 1999.
- [75143] 13473. Laxminarayan, K. M.; Chan, B. K.; Tetaz, T.; Bird, P. I.; Mitchell, C. A.: Characterization of a cDNA encoding the 43-kDa membrane-associated inositol-polyphosphate

5-phosphatase. J. Biol. Chem. 269: 17305–17310,1994.

[75144] 13474.Fink, J. K.; Jones, S. M.; Esposito, C.; Wilkowski, J.: Humanmicrotubule-associated protein 1a (MAP1A) gene: genomic organization,cDNA sequence, and developmen- tal-and tissue-specific expression. Genomics 35:577–585, 1996.

[75145] 13475.Asker, C.; Steinitz, M.; Andersson, K.; Sumegi, J.; Klein, G.;Ingvarsson, S.: Nucleotide sequence of the rat Bmyc gene.. Oncogene 4:1523–1527, 1989.

[75146] 13476.Ingvarsson, S.; Sundaresan, S.; Jin, P.; Francke, U.; Asker, C.;Sumegi, J.; Klein, G.; Sejersen, T.: Chromosome localization andexpression pattern of Lmyc and Bmyc in murine embryonal carcinomacells. Oncogene 3: 679–685, 1988.

[75147] 13477.Kojima, T.; Inazawa, J.; Takamatsu, J.; Rosenberg, R. D.; Saito,H.: Human ryudocan core protein: molecular cloning and characterizationof the cDNA, and chromoso- mal localization of the gene. Biochem. Biophys.Res. Com- mun. 190: 814–822, 1993.

[75148] 13478.Kojima, T.; Shworak, N. W.; Rosenberg, R. D.: Molecular cloningand expression of two distinct cDNA- encoding heparan sulfate proteoglycancore proteins from a rat endothelial cell line. J. Biol. Chem. 267:4870–4877,

1992.

- [75149] 13479.Yu, H.; Humphries, D. E.; Watkins, M.; Karlinsky, J. B.: Molecularcloning of the human ryudocan promoter. Biochem. Biophys. Res. Commun. 212:1139–1144, 1995.
- [75150] 13480.McBride, O. W.; Yi, H. F.; Srivastava, M.: The human cytochrome b561 gene (CYB561) is located at 17q11–qter. Genomics 21: 662–663,1994.
- [75151] 13481.Srivastava, M.: Genomic structure and expression of the humangene encoding cytochrome b(561), an integral protein of the cromaffingranule membrane. J. Biol. Chem. 270: 22714–22720, 1995.
- [75152] 13482.Albarosa, R.; DiDonato, S.; Finocchiaro, G.: Redefinition of thecoding sequence of the MXI1 gene and identification of a polymorphicrepeat in the 3–prime non–coding region that allows the detectionof loss of heterozygosity of chromosome 10q25 in glioblastomas. Hum.Genet. 95: 709–711, 1995.
- [75153] 13483.Mitchell, C. A.; Speed, C. J.; Nicholl, J.; Sutherland, G. R.:Chromosomal mapping of the gene (INPP5A) encoding the 43–kDa membrane–associatedinositol polyphosphate 5–phosphatase to 10q26.3 by fluorescence insitu hybridization. Genomics 31: 139–140, 1996.
- [75154] 13484.Sugasawa, K.; Ng, J. M. Y.; Masutani, C.; Iwai, S.;

van der Spek, P. J.; Eker, A. P. M.; Hanaoka, F.; Bootsma, D.; Hoeijmakers, J. H. J.: Xeroderma pigmentosum group C protein complex is the initiator of global genome nucleotide excision repair. *Molec. Cell* 2: 223–232, 1998.

[75155] 13485. van der Spek, P. J.; Smit, E. M. E.; Beverloo, H. B.; Sugasawa, K.; Masutani, C.; Hanaoka, F.; Hoeijmakers, J. H. J.; Hagemmeijer, A.: Chromosomal localization of three repair genes: the xeroderma pigmentosum group C gene and two human homologs of yeast RAD23. *Genomics* 23:651–658, 1994.

[75156] 13486. van der Spek, P. J.; Visser, C. E.; Hanaoka, F.; Smit, B.; Hagemmeijer, A.; Bootsma, D.; Hoeijmakers, J. H. J.: Cloning, comparative mapping, and RNA expression of the mouse homologues of the *Saccharomyces cerevisiae* nucleotide excision repair gene RAD23. *Genomics* 31: 20–27, 1996.

[75157] 13487. Netik, A.; Forss-Petter, S.; Holzinger, A.; Molzer, B.; Unterrainer, G.; Berger, J.: Adrenoleukodystrophy-related protein can compensate functionally for adrenoleukodystrophy protein deficiency (X-ALD): implications for therapy. *Hum. Molec. Genet.* 8: 907–913, 1999.

[75158] 13488. Rajantie, I.; Ekman, N.; Iljin, K.; Arighi, E.; Gunji, Y.; Kaukonen, J.; Palotie, A.; Dewerchin, M.; Carmeliet, P.; Ali-

talo, K.: Bmx tyrosinekinase has a redundant function downstream of angiopoietin and vascular endothelial growth factor receptors in arterial endothelium.

Molec.Cell. Biol. 21: 4647–4655, 2001.

- [75159] 13489. Tamagnone, L.; Lahtinen, I.; Mustonen, T.; Virtaneva, K.; Francis, F.; Muscatelli, F.; Alitalo, R.; Edvard Smith, C. I.; Larsson, C.; Alitalo, K.: BMX, a novel nonreceptor tyrosine kinase gene of the BTK/ITK/TEC/TKK family located in chromosome Xp22.2. *Oncogene* 9:3683–3688, 1994.
- [75160] 13490. Dinulos, M. B.; Bassi, M. T.; Rugarli, E. I.; Chapman, V.; Ballabio, A.; Distech, C. M.: A new region of conservation is defined between human and mouse X chromosomes. *Genomics* 35: 244–247, 1996.
- [75161] 13491. Schiaffino, M. V.; Bassi, M. T.; Rugarli, E. I.; Renieri, A.; Galli, L.; Ballabio, A.: Cloning of a human homologue of the *Xenopus laevis* APX gene from the ocular albinism type 1 critical region. *Hum. Molec. Genet.* 4: 373–382, 1995.
- [75162] 13492. Bachner, D.; Sedlacek, Z.; Korn, B.; Hameister, H.; Poustka, A.: Expression patterns of two human genes coding for different rabGDP-dissociation inhibitors (GDIs), extremely conserved proteins involved in cellular transport. *Hum. Molec. Genet.* 4: 701–708, 1995.

- [75163] 13493.Black, G. C. M.; Perveen, R.; Bonshek, R.; Cahill, M.; Clayton-Smith, J.; Lloyd, I. C.; McLeod, D.: Coats' disease of the retina (unilateral retinal telangiectasis) caused by somatic mutation in the NDP gene: a role for norrin in retinal angiogenesis. *Hum. Molec. Genet.* 8:2031–2035, 1999.
- [75164] 13494.Mao, M.; Fu, G.; Wu, J.-S.; Zhang, Q.-H.; Zhou, J.; Kan, L.-X.; Huang, Q.-H.; He, K.-L.; Gu, B.-W.; Han, Z.-G.; Shen, Y.; Gu, J.; Yu, Y.-P.; Xu, S.-H.; Wang, Y.-X.; Chen, S.-J.; Chen, Z.: Identification of genes expressed in human CD34+ hematopoietic stem/progenitor cells by expressed sequence tags and efficient full-length cDNA cloning. *Proc. Nat. Acad. Sci.* 95: 8175–8180, 1998.
- [75165] 13495.Chen, Y.-T.; Gure, A. O.; Tsang, S.; Stockert, E.; Jager, E.; Knuth, A.; Old, L. J.: Identification of multiple cancer/testis antigens by allogeneic antibody screening of a melanoma cell line library. *Proc. Nat. Acad. Sci.* 95: 6919–6923, 1998.
- [75166] 13496.Lucas, S.; De Smet, C.; Arden, K. C.; Viars, C. S.; Lethe, B.; Lurquin, C.; Boon, T.: Identification of a new MAGE gene with tumor-specific expression by representational difference analysis. *Cancer Res.* 58:743–752, 1998.
- [75167] 13497.Pearce, W. G.; Sanger, R.; Race, R. R.: Ocular al-



binism and Xg. Lancet I:1282–1283, 1968.

- [75168] 13498.Rosenberg, T.; Schwartz, M.: X-linked ocular albinism: prevalence and mutations--a national study. Europ. J. Hum. Genet. 6: 570–577, 1998.
- [75169] 13499.Schiaffino, M. V.; Bassi, M. T.; Galli, L.; Renieri, A.; Bruttini, M.; De Nigris, F.; Bergen, A. A. B.; Charles, S. J.; Yates, J. R.W., Meindl, A.; Lewis, R. A.; King, R. A.; Ballabio, A.: Analysis of the OA1 gene reveals mutations in only one-third of patients with X-linked ocular albinism. Hum. Molec. Genet. 4: 2319–2325, 1995.
- [75170] 13500.Schiaffino, M. V.; d'Addio, M.; Alloni, A.; Baschirotto, C.; Valetti, C.; Cortese, K.; Puri, C.; Bassi, M. T.; Colla, C.; De Luca, M.; Tacchetti, C.; Ballabio, A.: Ocular albinism: evidence for a defect in an intracellular signal transduction system. Nature Genet. 23: 108–112, 1999.
- [75171] 13501.Schnur, R. E.; Gao, M.; Wick, P. A.; Keller, M.; Benke, P. J.; Edwards, M. J.; Grix, A. W.; Hockey, A.; Jung, J. H.; Kidd, K. K.; Kistenmacher, M.; Levin, A. V.; and 11 others: OA1 mutations and deletions in X-linked ocular albinism. Am. J. Hum. Genet. 62: 800–809, 1998.
- [75172] 13502.Schnur, R. E.; Trask, B. J.; van den Engh, G.; Punnett, H. H.; Kistenmacher, M.; Tomeo, M. A.; Naidu, R. E.; Nussbaum, R. L.: An Xp22 microdeletion associated with

ocular albinism and ichthyosis:approximation of break-points and estimation of deletion size by usingcloned DNA probes and flow cytometry. Am. J. Hum. Genet. 45: 706–720,1989.

- [75173] 13503.Schnur, R. E.; Wick, P. A.; Bailey, C.; Rebbeck, T.; Weleber,R. G.; Wagstaff, J.; Grix, A. W.; Pagon, R. A.; Hockey, A.; Edwards,M. J.: Phenotypic variability in X-linked ocular albinism: relationship to linkage genotypes. Am. J. Hum. Genet. 55: 484–496, 1994.
- [75174] 13504.Vogt, A.: Die Iris: Albinismus solum bulbi. Atlas Spalt-Lampen-Mikroskopie(pub.) (3rd ed.): 1942.
- [75175] 13505.Waardenburg, P. J.; Van den Bosch, J.: X-chromosomal ocular albinism in Dutch family. Ann. Hum. Genet. 21: 101–122, 1956.
- [75176] 13506.Zhang, Y.; McMahon, R.; Charles, S. J.; Green, J. S.; Moore, A.T.; Barton, D. E.; Yates, J. R. W.: Genetic mapping of the Kallmannsyndrome and X linked ocular albinism gene loci. J. Med. Genet. 30:923–925, 1993.
- [75177] 13507.Shapira, E.; Ben-Yoseph, Y.; Eyal, G.; Russell, A.: Enzymaticallyinactive red cell carbonic anhydrase B in a family with renal tubularacidosis. J. Clin. Invest. 53: 59–63, 1974.
- [75178] 13508.Cianfriglia, M.; Miggiano, V. C.; Meo, T.; Muller, H.

J.; Muller,E.; Battistuzzi, G.: Evidence for syntenic between the rabbit genes coding for HPRT, PGK and G6PD in mouse-rabbit somatic cell hybrids.(Abstract) Cytogenet. Cell Genet. 25: 142, 1979.

[75179] 13509.Holterhus, P. M.; Sinnecker, G. H. G.; Wollmann, H. A.; Struve,D.; Homburg, N.; Kruse, K.; Hiort, O.: Expression of two functionally different androgen receptors in a patient with androgen insensitivity. Europ.J. Pediat. 158: 702-706, 1999.

[75180] 13510.Shimmin, L. C.; Chang, B.-H.; Li, W.-H.: Male-driven evolution of DNA sequences. Nature 362: 745-747, 1993.

[75181] 13511.Warburg, M.: Personal Communication. Copenhagen, Denmark 1966.

[75182] 13512.Imasaki, K.; Okabe, T.; Murakami, H.; Tanaka, Y.; Haji, M.; Takayanagi,R.; Nawata, H.: Androgen insensitivity syndrome due to new mutations in the DNA-binding domain of the androgen receptor. Molec. Cell.Endocr. 120: 15-24, 1996.

[75183] 13513.Jakubiczka, S.; Nedel, S.; Werder, E. A.; Schleiermacher, E.;Theile, U.; Wolff, G.; Wieacker, P.: Mutations of the androgen receptor gene in patients with complete androgen insensitivity. Hum. Mutat. 9:57-61, 1997.

- [75184] 13514.Kang, H.-Y.; Yeh, S.; Fujimoto, N.; Chang, C.: Cloning and characterization of human prostate coactivator ARA54, a novel protein that associates with the androgen receptor. *J. Biol. Chem.* 274: 8570–8576, 1999.
- [75185] 13515.Kazemi-Esfarjani, P.; Beitel, L. K.; Trifiro, M.; Kaufman, M.; Rennie, P.; Sheppard, P.; Matusik, R.; Pinsky, L.: Substitution of valine-865 by methionine or leucine in the human androgen receptor causes complete or partial androgen insensitivity, respectively with distinct androgen receptor phenotypes. *Molec. Endocr.* 7: 37–46, 1993.
- [75186] 13516.Kittles, R. A.; Young, D.; Weinrich, S.; Hudson, J.; Argyropoulos, G.; Ukoli, F.; Adams-Campbell, L.; Dunston, G. M.: Extent of linkage disequilibrium between the androgen receptor gene CAG and GGC repeats in human populations: implications for prostate cancer risk. *Hum. Genet.* 109: 253–261, 2001.
- [75187] 13517.Klocker, H.; Kaspar, F.; Eberle, J.; Uberreiter, S.; Radmayr, C.; Bartsch, G.: Point mutation in the DNA binding domain of the androgen receptor in two families with Reifenshtein syndrome. *Am. J. Med. Genet.* 50: 1318–1327, 1992.
- [75188] 13518.Knoke, I.; Allera, A.; Wieacker, P.: Significance of the CAG repeat length in the androgen receptor gene (AR)

for the transactivation function of an M780I mutant AR.

Hum. Genet. 104: 257–261, 1999.

- [75189] 13519. Kobayashi, Y.; Miwa, S.; Merry, D. E.; Kume, A.; Mei, L.; Doyu, M.; Sobue, G.: Caspase-3 cleaves the expanded androgen receptor protein of spinal and bulbar muscular atrophy in a polyglutamine repeat length-dependent manner. Biochem. Biophys. Res. Commun. 252: 145–150, 1998.
- [75190] 13520. Koivisto, P. A.; Schleutker, J.; Helin, H.; Ehren-van Eekelen, C.; Kallioniemi, O.-P.; Trapman, J.: Androgen receptor gene alterations and chromosomal gains and losses in prostate carcinomas appearing during finasteride treatment for benign prostatic hyperplasia. Clin. Cancer Res. 5: 3578–3582, 1999.
- [75191] 13521. Kooy, R. F.; Reyniers, E.; Storm, K.; Vits, L.; van Velzen, D.; de Ruiter, P. E.; Brinkmann, A. O.; de Paepe, A.; Willems, P. J.: CAG repeat contraction in the androgen receptor gene in three brothers with mental retardation. Am. J. Med. Genet. 85: 209–213, 1999.
- [75192] 13522. La Spada, A.; Fischbeck, K. H.: Androgen receptor gene defect in X-linked spinal and bulbar muscular atrophy. (Abstract) Am. J. Hum. Genet. 49 (suppl.): 20, 1991.
- [75193] 13523. Lewis, M.; Kaita, H.; Giblett, E. R.; Anderson, J.;

Philipps,S.; Steinberg, A. G.; McAlpine, P. J.: Multiplicity of genetic polymorphisms of blood in the Schmiedeleut Hutterites. *Am. J. Med. Genet.* 22:477–485, 1985.

- [75194] 13524.Lim, H. N.; Chen, H.; McBride, S.; Dunning, A. M.; Nixon, R. M.;Hughes, I. A.; Hawkins, J. R.: Longer polyglutamine tracts in the androgen receptor are associated with moderate to severe undermasculinized genitalia in XY males. *Hum. Molec. Genet.* 9: 829–834, 2000.
- [75195] 13525.Lobaccaro, J.–M.; Lumbroso, S.; Belon, C.; Galtier–Dereure, F.;Bringer, J.; Lesimple, T.; Namer, M.; Cutuli, B. F.; Pujol, H.; Sultan,C.: Androgen receptor gene mutation in male breast cancer. *Hum.Molec. Genet.* 2: 1799–1802, 1993.
- [75196] 13526.Lubahn, D. B.; Brown, T. R.; Simental, J. A.; Higgs, H. N.; Migeon,C. J.; Wilson, E. M.; French, F. S.: Sequence of the intron/exon junctions of the coding region of the human androgen receptor gene and identification of a point mutation in a family with complete androgen insensitivity. *Proc. Nat. Acad. Sci.* 86: 9534–9538, 1989.
- [75197] 13527.Lubahn, D. B.; Joseph, D. R.; Sar, M.; Tan, J.; Higgs, H. N.;Larson, R. E.; French, F. S.; Wilson, E. M.: The human androgen receptor:complementary deoxyribonucleic acid cloning, sequence analysis and gene expression in

prostate. *Molec. Endocr.* 2: 1265–1275, 1988.

[75198] 13528. Lubahn, D. B.; Joseph, D. R.; Sullivan, P. M.; Willard, H. F.; French, F. S.; Wilson, E. M.: Cloning of human androgen receptor complementary DNA and localization to the X chromosome. *Science* 240:327–330, 1988.

[75199] 13529. Lumbroso, S.; Lobaccaro, J. M.; Georget, V.; Leger, J.; Poujol, N.; Terouanne, B.; Evain-Brion, D.; Czernichow, P.; Sultan, C.: A novel substitution (leu707-to-arg) in exon 4 of the androgen receptor gene causes complete androgen resistance. *J. Clin. Endocr. Metab.* 81:1984–1988, 1996.

[75200] 13530. Stuppia, L.; Calabrese, G.; Franchi, P. G.; Mingarelli, R.; Gatta, V.; Palka, G.; Dallapiccola, B.: Widening of a Y-chromosome interval-6 deletion transmitted from a father to his infertile son accounts for an oligozoospermia critical region distal to the RBM1 and DAZ genes. (Letter) *Am. J. Hum. Genet.* 59: 1393–1395, 1996.

[75201] 13531. Brown, G. M.; Furlong, R. A.; Sargent, C. A.; Erickson, R. P.; Longepied, G.; Mitchell, M.; Jones, M. H.; Hargreave, T. B.; Cooke, H. J.; Affara, N. A.: Characterisation of the coding sequence and fine mapping of the human DF-FRY gene and comparative expression analysis and mapping to the Sxr-b interval of the mouse Y chromosome of

theDffry gene. Hum. Molec. Genet. 7: 97–107, 1998.

[75202] 13532.Ferlin, A.; Moro, E.; Garolla, A.; Foresta, C.: Human male infertilityand Y chromosome deletions: role of the AZF–candidate genes DAZ, RBMand DFFRY. Hum. Reprod. 14: 1710–1716, 1999.

[75203] 13533.Foresta, C.; Ferlin, A.; Moro, E.: Deletion and expression analysisof AZFa genes on the human Y chromosome revealed a major role forDBY in male infertility. Hum. Molec. Genet. 9: 1161–1169, 2000.

[75204] 13534.Sargent, C. A.; Boucher, C. A.; Kirsch, S.; Brown, G.; Weiss,B.; Trundley, A.; Burgoyne, P.; Saut, N.; Durand, C.; Levy, N.; Terriou,P.; Hargreave, T.; Cooke, H.; Mitchell, M.; Rappold, G. A.; Affara,N. A.: The critical region of overlap defining the AZFa male infertilityinterval of proximal Yq contains three transcribed sequences. J.Med. Genet. 36: 670–677, 1999.

[75205] 13535.Shen, P.; Wang, F.; Underhill, P. A.; Franco, C.; Yang, W.–H.;Roxas, A.; Sung, R.; Lin, A. A.; Hyman, R. W.; Vollrath, D.; Davis,R. W.; Cavalli–Sforza, L. L.; Oefner, P. J.: Population genetic implicationsfrom sequence variation in four Y chromosome genes. Proc. Nat. Acad.Sci. 97: 7354–7359, 2000.

[75206] 13536.Sun, C.; Skaletsky, H.; Birren, B.; Devon, K.; Tang,



Z.; Silber,S.; Oates, R.; Page, D. C.: An azoospermic man with a de novo pointmutation in the Y–chromosomal gene USP9Y. *Nature Genet.* 23: 429–432,1999.

[75207] 13537.Thomson, R.; Pritchard, J. K.; Shen, P.; Oefner, P. J.; Feldman,M. W.: Recent common ancestry of human Y chromosomes: evidence fromDNA sequence data. *Proc. Nat. Acad. Sci.* 97: 7360–7365, 2000.

[75208] 13538.Schiebel, K.; Winkelmann, M.; Mertz, A.; Xu, X.; Page, D. C.; Weil,D.; Petit, C.; Rappold,, G. A.: Abnormal XY interchange between anovel isolated protein kinase gene, PRKY, and its homologue, PRKX,accounts for one third of all (Y+)XX males and (Y–)XY females. *Hum.Molec. Genet.* 6: 1985–1989, 1997.

[75209] 13539.Greenfield, A.; Scott, D.; Pennisi, D.; Ehrmann, I.; Ellis, P.;Cooper, L.; Simpson, E.; Koopman, P.: An H–YDb epitope is encodedby a novel mouse Y chromosome gene. *Nature Genet.* 14: 474–478, 1996.

[75210] 13540.Scott, D.; Addey, C.; Ellis, P; James, E.; Mitchell, M. J.; Saut,N.; Jurcevic, S.; Simpson, E.: Dendritic cells permit identificationof genes encoding MHC class II–restricted epitopes of transplantationantigens. *Immunity* 12: 711–720, 2000.

[75211] 13541.Wu, G.; Chai, J.; Suber, T. L.; Wu, J.–W.; Du, C.;

Wang, X.; Shi,Y.: Structural basis of IAP recognition by Smac/DIABLO. *Nature* 408:1008–1012, 2000.

[75212] 13542.Macke, J. P.; Hu, N.; Hu, S.; Bailey, M.; King, V. L.; Brown,T.; Hamer, D.; Nathans, J.: Sequence variation in the androgen receptorgene is not a common determinant of male sexual orientation. *Am.J. Hum. Genet.* 53: 844–852, 1993.

[75213] 13543.Yang, R.–B.; Fulle, H.–J.; Garbers, D. L.: Chromosomal localizationand genomic organization of genes encoding guanylyl cyclase receptorsexpressed in olfactory sensory neurons and retina. *Genomics* 31:367–372, 1996.

[75214] 13544.Nakamura, H.; Izumoto, Y.; Kambe, H.; Kuroda, T.; Mori, T.; Kawamura,K.; Yamamoto, H.; Kishimoto, T.: Molecular cloning of complementaryDNA for a novel human hepatoma–derived growth factor: its homologywith high mobility group–1 protein. *J. Biol. Chem.* 269: 25143–25149,1994.

[75215] 13545.Wanschura, S.; Schoenmakers, E. F. P. M.; Huysmans, C.; Bartnitzke,S.; Van de Ven, W. J. M.; Bullerdiek, J.: Mapping of the gene encodingthe human hepatoma–derived growth factor (HDGF) with homology to thehigh–mobility group (HMG)–1 protein to Xq25. *Genomics* 32: 298–300,1996.

- [75216] 13546.Baens, M.; Marynen, P.: A human homologue (BICD1) of the Drosophilabicaudal-D gene. Genomics 45: 601–606, 1997.
- [75217] 13547.Bullock, S. L.; Ish-Horowicz, D.: Conserved signals and machineryfor RNA transport in Drosophila oogenesis and embryogenesis. Nature 414:611–616, 2001.
- [75218] 13548.McMurtrie, E. B.; Barbosa, M. D. F. S.; Zerial, M.; Kingsmore,S. F.: Rab17 and Rab18, small GTPases with specificity for polarizedepithelial cells: genetic mapping in the mouse. Genomics 45: 623–625,1997.
- [75219] 13549.Schafer, U.; Seibold, S.; Schneider, A.; Neugebauer, E.: Isolationand characterisation of the human rab18 gene after stimulation ofendothelial cells with histamine. FEBS Lett. 466: 148–154, 2000.
- [75220] 13550.Doupnik, C. A.; Davidson, N.; Lester, H. A.: The inward rectifierpotassium channel family. Curr. Opin. Neurobiol. 5: 268–277, 1995.
- [75221] 13551.Tada, Y.; Horio, Y.; Takumi, T.; Terayama, M.; Tsuji, L.; Copeland,N. G.; Jenkins, N. A.; Kurachi, Y.: Assignment of the glial inwardlyrectifying potassium channel K(AB)–2/Kir4.1 (Kcnj10) gene to the distalregion of mouse chromosome 1. Genomics 45: 629–630, 1997.
- [75222] 13552.Takumi, T.; Ishii, T.; Horio, Y.; Morishige, K.–I.;

Takahashi,N.; Yamada, M.; Yamashita, T.; Kiyama, H.;  
Sohmiya, K.; Nakanishi,S.; Kurachi, Y.: A novel ATP-  
dependent inward rectifier potassiumchannel expressed  
predominantly in glial cells. J. Biol. Chem.  
270:16339–16436, 1995.

[75223] 13553.Schenker, T.; Trueb, B.: Assignment of the gene for  
a developmentallyregulated GTP-binding protein (DRG2)  
to human chromosome bands 17p13–p12by in situ hy-  
bridization. Cytogenet. Cell Genet. 79: 274–275, 1997.

[75224] 13554.Heikkinen, J.; Risteli, M.; Wang, C.; Latvala, J.;  
Rossi, M.; Valtavaara,M.; Myllyla, R.: Lysyl hydroxylase 3 is  
a multifunctional proteinpossessing collagen glucosyl-  
transferase activity. J. Biol. Chem. 275:36158–36163,  
2000.

[75225] 13555.Passoja, K.; Rautavuoma, K.; Ala-Kokko, L.; Koso-  
nen, T.; Kivirikko,K. I.: Cloning and characterization of a  
third human lysyl hydroxylaseisoform. Proc. Nat. Acad.  
Sci. 95: 10482–10486, 1998.

[75226] 13556.Valtavaara, M.; Szpirer, C.; Szpirer, J.; Myllyla, R.:  
Primarystructure, tissue distribution, and chromosomal  
localization of anovel isoform of lysyl hydroxylase (lysyl  
hydroxylase 3). J. Biol.Chem. 273: 12881–12886, 1998.

[75227] 13557.Rohan, P. J.; Davis, P.; Moskaluk, C. A.; Kearns, M.;

Krutzsch,H.; Siebenlist, U.; Kelly, K.: PAC-1: a mitogen-induced nuclear proteintyrosine phosphatase. Science 259: 1763-1766, 1993.

- [75228] 13558.Ward, Y.; Gupta, S.; Jensen, P.; Wartmann, M.; Davis, R. J.; Kelly,K.: Control of MAP kinase activation by the mitogen-induced threonine/tyrosinephosphatase PAC1. Nature 367: 651-654, 1994.
- [75229] 13559.Yi, H.; Morton, C. C.; Weremowicz, S.; McBride, O. W.; Kelly, K.: Genomic organization and chromosomal localization of the DUSP2 gene,encoding a MAP kinase phosphatase, to human 2p11.2-q11. Genomics 28:92-96, 1995.
- [75230] 13560.Ishibashi, T.; Bottaro, D. P.; Michieli, P.; Kelley, C. A.; Aaronson,S. A.: A novel dual specificity phosphatase induced by serum stimulationand heat shock. J. Biol. Chem. 269: 29897-29902, 1994.
- [75231] 13561.Kwak, S. P.; Dixon, J. E.: Multiple dual specificity protein tyrosinephosphatases are expressed and regulated differentially in liver celllines. J. Biol. Chem. 270: 1156-1160, 1995.
- [75232] 13562.Kovalenko, O. V.; Golub, E. I.; Bray-Ward, P.; Ward, D. C.; Radding,C. M.: A novel nucleic acid-binding protein that interacts with humanRad51 recombinase. Nucleic

Acids Res. 25: 4946–4953, 1997.

- [75233] 13563. Mizuta, R.; LaSalle, J. M.; Cheng, H.-L.; Shinohara, A.; Ogawa, H.; Copeland, N.; Jenkins, N. A.; Lalande, M.; Alt, F. W.: RAB22 and RAB163/mouse BRCA2: proteins that specifically interact with the RAD51 protein. Proc. Nat. Acad. Sci. 94: 6927–6932, 1997.
- [75234] 13564. Blasius, R.; Weber, R. G.; Lichter, P.; Ogilvie, A.: A novel orphan G protein-coupled receptor primarily expressed in the brain is localized on human chromosomal band 2q21. J. Neurochem. 70: 1357–1365, 1998.
- [75235] 13565. Raport, C. J.; Schweickart, V. L.; Chantry, D.; Eddy, R. L., Jr.; Shows, T. B.; Godiska, R.; Gray, P. W.: New members of the chemokine receptor gene family. J. Leukoc. Biol. 59: 18–23, 1996.
- [75236] 13566. Shindo, M.; Nakano, H.; Kuroyanagi, H.; Shirasawa, T.; Mihara, M.; Gilbert, D. J.; Jenkins, N. A.; Copeland, N. G.; Yagita, H.; Okumura, K.: cDNA cloning, expression, subcellular localization, and chromosomal assignment of mammalian aurora homologues, aurora-related kinase (ARK)1 and 2. Biochem. Biophys. Res. Commun. 244: 285–292, 1998.
- [75237] 13567. Adachi, H.; Tsujimoto, M.; Hattori, M.; Arai, H.; Inoue, K.: cDNA cloning of human cytosolic platelet-acti-

vating factor acetylhydrolase gamma-subunit and its mRNA expression in human tissues. *Biochem. Biophys. Res. Commun.* 214: 180–187, 1995.

[75238] 13568. Presky, D. H.; Yang, H.; Minetti, L. J.; Chua, A. O.; Nabavi, N.; Wu, C.-Y.; Gately, M. K.; Gubler, U.: A functional interleukin 12 receptor complex is composed of two beta-type cytokine receptor subunits. *Proc. Nat. Acad. Sci.* 93: 14002–14007, 1996.

[75239] 13569. Yamamoto, K.; Kobayashi, H.; Miura, O.; Hirosawa, S.; Miyasaka, N.: Assignment of IL12RB1 and IL12RB2, interleukin-12 receptor beta-1 and beta-2 chains, to human chromosome 19 band p13.1 and chromosome 1 band p31.2, respectively, by in situ hybridization. *Cytogenet. Cell Genet.* 77: 257–258, 1997.

[75240] 13570. Lafreniere, R. G.; Rochefort, D. L.; Kibar, Z.; Fon, E. A.; Han, F.-Y.; Cochius, J.; Kang, X.; Baird, S.; Korneluk, R. G.; Andermann, E.; Rommens, J. M.; Rouleau, G. A.: Isolation and characterization of GT335, a novel human gene conserved in *Escherichia coli* and mapping to 21q22.3. *Genomics* 38: 264–272, 1996.

[75241] 13571. Nagamine, K.; Kudoh, J.; Minoshima, S.; Kawasaki, K.; Asakawa, S.; Ito, F.; Shimizu, N.: Isolation of cDNA for a novel human protein KNP-I that is homologous to the E.

coli SCRP-27A protein from the autoimmune polyglandular disease type I (APECED) region of chromosome 21q22.3. Biochem. Biophys. Res. Commun. 225: 608–616, 1996.

- [75242] 13572.Scott, H. S.; Chen, H.; Rossier, C.; Lalioti, M. D.; Antonarakis, S. E.: Isolation of a human gene (HES1) with homology to an Escherichia coli and a zebrafish protein that maps to chromosome 21q22.3. Hum.Genet. 99: 616–623, 1997.
- [75243] 13573.Bernier-Villamor, V.; Sampson, D. A.; Matunis, M. J.; Lima, C.D.: Structural basis for E2-mediated SUMO conjugation revealed by a complex between ubiquitin-conjugating enzyme Ubc9 and RanGAP1. Cell 108:345–356, 2002.
- [75244] 13574.Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association of FHIT (fragile histidine triad), a candidate tumour suppressor gene, with the ubiquitin-conjugating enzyme hUBC9. Biochem. J. 352: 443–448, 2000.
- [75245] 13575.Tachibana, M.; Iwata, N.; Watanabe, A.; Nobukuni, Y.; Ploplis, B.; Kajigaya, S.: Assignment of the gene for a ubiquitin-conjugating enzyme (UBE2I) to human chromosome band 16p13.3 by in situ hybridization. Cytogenet.Cell Genet. 75: 222–223, 1996.
- [75246] 13576.Wang, Z.-Y.; Qiu, Q.-Q.; Seufert, W.; Taguchi, T.;



Testa, J. R.;Whitmore, S. A.; Callen, D. F.; Welsh, D.; Shenk, T.; Deuel, T. F.: Molecular cloning of the cDNA and chromosome localization of the gene for human ubiquitin-conjugating enzyme 9. J. Biol. Chem. 271:24811–24816, 1996.

[75247] 13577.Yasugi, T.; Howley, P. M.: Identification of the structural and functional human homolog of the yeast ubiquitin conjugating enzyme UBC9. Nucleic Acids Res. 24: 2005–2010, 1996.

[75248] 13578.Bord, S.; Horner, A.; Beavan, S.; Compston, J.: Estrogen receptors alpha and beta are differentially expressed in developing human bone. J.Clin. Endocr. Metab. 86: 2309–2314, 2001.

[75249] 13579.Watanabe, T. K.; Fujiwara, T.; Kawai, A.; Shimizu, F.; Takami, S.; Hirano, H.; Okuno, S.; Ozaki, K.; Takeda, S.; Shimada, Y.; Nagata, M.; Takaichi, A.; Takahashi, E.; Nakamura, Y.; Shin, S.: Cloning, expression, and mapping of UBE2I, a novel gene encoding a human homologue of yeast ubiquitin-conjugating enzymes which are critical for regulating the cell cycle. Cytogenet. Cell Genet. 72: 86–89, 1996.

[75250] 13580.Chaidarun, S. S.; Swearingen, B.; Alexander, J. M.: Differential expression of estrogen receptor-beta (ER-beta)

in human pituitary tumors: functional interactions with ER-  
alpha and a tumor-specific splice variant. J. Clin. Endocr.  
Metab. 83: 3308–3315, 1998.

[75251] 13581. Chu, S.; Mamers, P.; Burger, H. G.; Fuller, P. J.: Es-  
trogen receptor isoform gene expression in ovarian stro-  
mal and epithelial tumors. J. Clin. Endocr. Metab. 85:  
1200–1205, 2000.

[75252] 13582. Enmark, E.; Peltö-Huikko, M.; Grandien, K.; Lager-  
crantz, S.; Lagercrantz, J.; Fried, G.; Nordenskjöld, M.;  
Gustafsson, J.-A.: Human estrogen receptor beta-gene  
structure, chromosomal localization, and expression pat-  
tern. J. Clin. Endocr. Metab. 82: 4258–4265, 1997.

[75253] 13583. Esmaili, B.; Harvey, J. T.; Hewlett, B.: Immunohis-  
tochemical evidence for estrogen receptors in meibomian  
glands. Ophthalmology 107: 180–184, 2000.

[75254] 13584. Forsell, C.; Enmark, E.; Axelman, K.; Blomberg, M.;  
Wahlund, L.-O.; Gustafsson, J.-A.; Lannfelt, L.: Investiga-  
tions of a CA repeat in the oestrogen receptor beta gene in  
patients with Alzheimer's disease. Europ. J. Hum. Genet. 9:  
802–804, 2001.

[75255] 13585. Korach, K. S.; Couse, J. F.; Curtis, S. W.; Washburn,  
T. F.; Lindzey, J.; Kimbro, K. S.; Eddy, E. M.; Migliaccio, S.;  
Snedeker, S. M.; Lubahn, D. B.; Schomberg, D. W.; Smith, E.

P.: Estrogen receptor gene disruption:molecular characterization and experimental and clinical phenotypes. Recent Prog. Horm. Res. 51: 159–188, 1996.

[75256] 13586.Krege, J. H.; Hodgin, J. B.; Couse, J. F.; Enmark, E.; Warner, M.; Mahler, J. F.; Sar, M.; Korach, K. S.; Gustafsson, J.-A.; Smithies, O.: Generation and reproductive phenotypes of mice lacking estrogenreceptor beta. Proc. Nat. Acad. Sci. 95: 15677–15682, 1998.

[75257] 13587.Krezel, W.; Dupont, S.; Krust, A.; Chambon, P.; Chapman, P. F.: Increased anxiety and synaptic plasticity in estrogen receptor beta-deficientmice. Proc. Nat. Acad. Sci. 98: 12278–12282, 2001.

[75258] 13588.Kuiper, G. G. J. M.; Enmark, E.; Peltö-Huikko, M.; Nilsson, S.; Gustafsson, J.-A.: Cloning of a novel estrogen receptor expressed in rat prostate and ovary. Proc. Nat. Acad. Sci. 93: 5925–5930, 1996.

[75259] 13589.Mosialos, G.; Birkenbach, M.; Yalamanchili, R.; VanArsdale, T.; Ware, C.; Kieff, E.: The Epstein–Barr virus transforming protein LMP1 engages signaling proteins for the tumor necrosis factor receptor family. Cell 80: 389–399, 1995.

[75260] 13590.Rothe, M.; Wong, S. C.; Henzel, W. J.; Goeddel, D. V.: A novel family of putative signal transducers associated

with the cytoplasmic domain of the 75 kDa tumor necrosis factor receptor. *Cell* 78: 681–692, 1994.

[75261] 13591. Siemienski, K.; Peters, N.; Scheurich, P.; Wajant, H.: Organization of the human tumour necrosis factor receptor-associated factor 1 (TRAF1) gene and mapping to chromosome 9q33–34. *Gene* 195: 35–39, 1997.

[75262] 13592. Tsitsikov, E. N.; Laouini, D.; Dunn, I. F.; Sannikova, T. Y.; Davidson, L.; Alt, F. W.; Geha, R. S.: TRAF1 is a negative regulator of TNF signaling: enhanced TNF signaling in TRAF1-deficient mice. *Immunity* 15: 647–657, 2001.

[75263] 13593. Jacquemin, P.; Hwang, J.-J.; Martial, J. A.; Dolle, P.; Davidson, I.: A novel family of developmentally regulated mammalian transcription factors containing the TEA/ATTS DNA binding domain. *J. Biol. Chem.* 271: 21775–21785, 1996.

[75264] 13594. Bejjani, B. A.; Lewis, R. A.; Tomey, K. F.; Anderson, K. L.; Dueker, D. K.; Jabak, M.; Astle, W. F.; Otterud, B.; Leppert, M.; Lupski, J. R.: Mutations in CYP1B1, the gene for cytochrome P4501B1, are the predominant cause of primary congenital glaucoma in Saudi Arabia. *Am. J. Hum. Genet.* 62: 325–333, 1998.

[75265] 13595. Bejjani, B. A.; Stockton, D. W.; Lewis, R. A.; Tomey, K. F.; Dueker, D. K.; Jabak, M.; Astle, W. F.; Lupski, J. R.:

Multiple CYP1B1 mutations and incomplete penetrance in an inbred population segregating primary congenital glaucoma suggest frequent de novo events and a dominant-modifier locus. *Hum. Molec. Genet.* 9: 367–374, 2000.

[75266] 13596. Lin, J.; Arnold, H. B.; Della-Fera, M. A.; Azain, M. J.; Hartzell, D. L.; Baile, C. A.: Myostatin knockout in mice increases myogenesis and decreases adipogenesis. *Biochem. Biophys. Res. Commun.* 291:701–706, 2002.

[75267] 13597. McPherron, A. C.; Lawler, A. M.; Lee, S.-J.: Regulation of skeletal muscle mass in mice by a new TGF- $\beta$  superfamily member. *Nature* 387:83–90, 1997.

[75268] 13598. Robins, P.; Lindahl, T.: DNA ligase IV from HeLa cell nuclei. *J. Biol. Chem.* 271: 24257–24261, 1996.

[75269] 13599. Roldan-Arjona, T.; Wei, Y.-F.; Carter, K. C.; Klungland, A.; Anselmino, C.; Wang, R.-P.; Augustus, M.; Lindahl, T.: Molecular cloning and functional expression of a human cDNA encoding the antimutator enzyme 8-hydroxyguanine-DNA glycosylase. *Proc. Nat. Acad. Sci.* 94: 8016–8020, 1997.

[75270] 13600. Guo, Z.; Turner, C.; Castle, D.: Relocation of the t-SNARE SNAP-23 from lamellipodia-like cell surface projections regulates compound exocytosis in mast cells. *Cell* 94: 537–548, 1998.

- [75271] 13601.Lazo, P. A.; Nadal, M.; Ferrer, M.; Area, E.; Hernandez-Torres,J.; Nabokina, S. M.; Mollinedo, F.; Estivill, X.: Genomic organization,chromosomal localization, alternative splicing, and isoforms of thehuman synaptosome-associated protein-23 gene implicated in vesicle-membranefusion processes. Hum. Genet. 108: 211-215, 2001.
- [75272] 13602.Mollinedo, F.; Lazo, P. A.: Identification of two isoforms ofthe vesicle-membrane fusion protein SNAP-23 in human neutrophils andHL-60 cells. Biochem. Biophys. Res. Commun. 231: 808-812, 1997.
- [75273] 13603.Ravichandran, V.; Chawla, A.; Roche, P. A.: Identification ofa novel syntaxin- and synaptobrevin/VAMP-binding protein, SNAP-23,expressed in non-neuronal tissues. J. Biol. Chem. 271: 13300-13303,1996.
- [75274] 13604.Shukla, A.; Corydon, T. J.; Nielsen, S.; Hoffmann, H. J.; Dahl,R.: Identification of three new splice variants of the SNARE proteinSNAP-23. Biochem. Biophys. Res. Commun. 285: 320-327, 2001.
- [75275] 13605.Mugita, N.; Kumura, Y.; Ogawa, M.; Saya, H.; Nakao, M.: Identificationof a novel, tissue-specific calpain htra-3; a human homologue of theCaenorhabditis elegans sex determination gene. Biochem. Biophys.Res. Commun.

239: 845–850, 1997.

- [75276] 13606.Sharpless, N. E.; Ferguson, D. O.; O'Hagan, R. C.; Castrillon,D. H.; Lee, C.; Farazi, P. A.; Alson, S.; Fleming, J.; Morton, C.C.; Frank, K.; Chin, L.; Alt, F. W.; DePinho, R. A.: Impaired nonhomologousend–joining provokes soft tissue sarcomas harboring chromosomal translocations,amplifications, and deletions. *Molec. Cell* 8: 1187–1196, 2001.
- [75277] 13607.Bohme, B.; Holtrich, U.; Wolf, G.; Luzius, H.; Grzeschik, K.–H.;Strebhardt, K.; Rubsamen–Waigmann, H.: PCR mediated detection ofa new human receptor–tyrosine–kinase, HEK 2. *Oncogene* 8: 2857–2862,1993.
- [75278] 13608.Ruiz, J. C.; Conlon, F. L.; Robertson, E. J.: Identification ofnovel protein kinases expressed in the myocardium of the developingmouse heart. *Mech. Dev.* 48: 153–164, 1994.
- [75279] 13609.Ashbourne, K. J.; Byth, B. C.; Meijers, J. C. M.; Cox, D. W.:Polymorphism of the protein C inhibitor (PCI) gene on chromosome 14. *Hum.Molec. Genet.* 2: 92 only, 1993.
- [75280] 13610.Sadler, J. E.: Combined factors V and VIII deficiency climbs ontothe map. (Editorial) *J. Clin. Invest.* 99: 555–556, 1997.
- [75281] 13611.Yasuda, T.; Nadano, D.; Iida, R.; Tanaka, Y.;

Nakanaga, M.; Kishi, K.: Discovery of a genetic polymorphism of human plasma protein C inhibitor (PCI): genetic survey utilizing isoelectric focusing followed by immunoblotting, immunological and biochemical characterization. *Hum. Genet.* 89: 265–269, 1992.

[75282] 13612. Eki, T.; Okumura, K.; Amin, A.; Ishiai, M.; Abe, M.; Nogami, M.; Taguchi, H.; Hurwitz, J.; Murakami, Y.; Hanaoka, F.: Mapping of the human homologue (ORC1L) of the yeast origin recognition complex subunit 1 gene to chromosome band 1p32. *Genomics* 36: 559–561, 1996.

[75283] 13613. Mendez, J.; Zou-Yang, H.; Kim, S.-Y.; Hidaka, M.; Tansey, W. P.; Stillman, B.: Human origin recognition complex large subunit is degraded by ubiquitin-mediated proteolysis after initiation of DNA replication. *Molec. Cell* 9: 481–491, 2002.

[75284] 13614. Bui, T. D.; Rankin, J.; Smith, K.; Huguet, E. L.; Ruben, S.; Strachan, T.; Harris, A. L.; Lindsay, S.: A novel human Wnt gene, WNT10B, maps to 12q13 and is expressed in human breast carcinomas. *Oncogene* 14: 1249–1253, 1997.

[75285] 13615. Hardiman, G.; Kastelein, R. A.; Bazan, J. F.: Isolation, characterization and chromosomal localization of human WNT10B. *Cytogenet. Cell Genet.* 77: 278–282, 1997.



- [75286] 13616.Ross, S. E.; Hemati, N.; Longo, K. A.; Bennett, C. N.; Lucas, P.C.; Erickson, R. L.; MacDougald, O. A.: Inhibition of adipogenesis by Wnt signaling. *Science* 289: 950–953, 2000.
- [75287] 13617.Meyerhardt, J. A.; Look, A. T.; Bigner, S. H.; Fearon, E. R.: Identification and characterization of neogenin, a DCC-related gene. *Oncogene* 14:1129–1136, 1997.
- [75288] 13618.Vielmetter, J.; Chen, X.-N.; Miskevich, F.; Lane, R. P.; Yamakawa, K.; Korenberg, J. R.; Dreyer, W. J.: Molecular characterization of human neogenin, a DCC-related protein, and the mapping of its gene (NEO1) to chromosomal position 15q22.3–q23. *Genomics* 41: 414–421, 1997.
- [75289] 13619.Vielmetter, J.; Kayyem, J. F.; Roman, J. M.; Dreyer, W. J.: Neogenin, an avian cell surface protein expressed during terminal neuronal differentiation, is closely related to the human tumor suppressor molecule deleted in colorectal cancer. *J. Cell Biol.* 127: 2009–2020, 1994.
- [75290] 13620.Rosenquist, T. A.; Zharkov, D. O.; Grollman, A. P.: Cloning and characterization of a mammalian 8-oxoguanine DNA glycosylase. *Proc. Nat. Acad. Sci.* 94: 7429–7434, 1997.
- [75291] 13621.Tani, M.; Shinmura, K.; Kohno, T.; Shiroishi, T.; Wakama, S.; Kim, S.-R.; Nohmi, T.; Kasai, H.; Takenoshita,

S.; Nagamachi, Y.; Yokota, J.: Genomic structure and chromosomal localization of the mouse Ogg1 gene that is involved in the repair of 8-hydroxyguanine in DNA damage. *Mammalian Genome* 9: 32–37, 1998.

- [75292] 13622. Lim, H. N.; Hawkins, J. R.; Hughes, I. A.: Genetic evidence to exclude the androgen receptor co-factor, ARA70 (NCOA4) as a candidate gene for the causation of undermasculinised genitalia. (Letter) *Clin. Genet.* 59: 284–286, 2001.
- [75293] 13623. Santoro, M.; Dathan, N. A.; Berlingieri, M. T.; Bongarzone, I.; Paulin, C.; Grieco, M.; Pierotti, M. A.; Vecchio, G.; Fusco, A.: Molecular characterization of RET/PTC3: a novel rearranged version of the RET proto-oncogene in a human thyroid papillary carcinoma. *Oncogene* 9: 509–516, 1994.
- [75294] 13624. Yeh, S.; Chang, C.: Cloning and characterization of a specific coactivator, ARA-70, for the androgen receptor in human prostate cells. *Proc. Nat. Acad. Sci.* 93: 5517–5521, 1996.
- [75295] 13625. Donghi, R.; Sozzi, G.; Pierotti, M. A.; Biunno, I.; Miozzo, M.; Fusco, A.; Grieco, M.; Santoro, M.; Vecchio, G.; Spurr, N. K.; Della Porta, G.: The oncogene associated with human papillary thyroid carcinoma (PTC) is assigned to

chromosome 10 q11–q12 in the same region as multiple endocrine neoplasia type 2A (MEN2A). *Oncogene* 4: 521–523, 1989.

- [75296] 13626. Nikiforova, M. N.; Stringer, J. R.; Blough, R.; Medvedovic, M.; Fagin, J. A.; Nikiforov, Y. E.: Proximity of chromosomal loci that participate in radiation-induced rearrangements in human cells. *Science* 290:138–141, 2000.
- [75297] 13627. Sozzi, G.; Pierotti, M. A.; Miozzo, M.; Donghi, R.; Radice, P.; De Benedetti, V.; Grieco, M.; Santoro, M.; Fusco, A.; Vecchio, G.; Mathew, C. G. P.; Ponder, B. A. J.; Spurr, N. K.: Refined localization to contiguous regions on chromosome 10q of the two genes (H4 and RET) that form the oncogenic sequence PTC. *Oncogene* 6: 339–342, 1991.
- [75298] 13628. Tong, Q.; Li, Y.; Smanik, P. A.; Fithian, L. J.; Xing, S.; Mazzaferri, E. L.; Jhiang, S. M.: Characterization of the promoter region and oligomerization domain of H4 (D10S170), a gene frequently rearranged with the ret proto-oncogene. *Oncogene* 10: 1781–1787, 1995.
- [75299] 13629. Araki, T.; Milbrandt, J.: Ninjurin, a novel adhesion molecule, is induced by nerve injury and promotes axonal growth. *Neuron* 17:353–361, 1996.
- [75300] 13630. Araki, T.; Zimonjic, D. B.; Popescu, N. C.; Milbrandt, J.: Mechanism of homophilic binding mediated by ninjurin,

a novel widely expressed adhesion molecule. *J. Biol. Chem.* 272: 21373–21380, 1997.

- [75301] 13631. Chadwick, B. P.; Heath, S. K.; Williamson, J.; Obermayr, F.; Patel, L.; Sheer, D.; Frischauf, A.-M.: The human homologue of the *ninjurin* gene maps to the candidate region of hereditary sensory neuropathy type I (HSN1). *Genomics* 47: 58–63, 1998.
- [75302] 13632. Mandich, P.; Bellone, E.; Di Maria, E.; Pigullo, S.; Pizzuti, A.; Schenone, A.; Soriani, S.; Varese, A.; Windebank, A. J.; Ajmar, F.: Exclusion of the *ninjurin* gene as a candidate for hereditary sensory neuropathies type I and type II. *Am. J. Med. Genet.* 83: 409–410, 1999.
- [75303] 13633. McAllister, G.; Whiting, P.; Hammond, E. A.; Knowles, M. R.; Atack, J. R.; Bailey, F. J.; Maigetter, R.; Ragan, C. I.: cDNA cloning of human and rat brain myo-inositol monophosphatase: expression and characterization of the human recombinant enzyme. *Biochem. J.* 284: 749–754, 1992.
- [75304] 13634. Sjøholt, G.; Molven, A.; Lovlie, R.; Wilcox, A.; Sikela, J. M.; Steen, V. M.: Genomic structure and chromosomal localization of a human myo-inositol monophosphatase gene (IMPA). *Genomics* 45: 113–122, 1997.
- [75305] 13635. Steen, V. M.; Gulbrandsen, A. K.; Eiken, H. G.; Berle,

J. O.:Lack of genetic variation in the coding region of the myo-inositolmonophosphatase gene in lithium-treated patients with manic depressive illness. *Pharmacogenetics* 6: 113–116, 1996.

[75306] 13636.McDonald, M. T.; Flejter, W.; Sheldon, S.; Putzi, M. J.; Gorski, J. L.: XY sex reversal and gonadal dysgenesis due to 9p24 monosomy. *Am.J. Med. Genet.* 73: 321–326, 1997.

[75307] 13637.Muroya, K.; Okuyama, T.; Goishi, K.; Ogiso, Y.; Fukuda, S.; Kameyama, J.; Sato, H.; Suzuki, Y.; Terasaki, H.; Gomyo, H.; Wakui, K.; Fukushima, Y.; Ogata, T.: Sex-determining gene(s) on distal 9p: clinical and molecular studies in six cases. *J. Clin. Endocr. Metab.* 85: 3094–3100, 2000.

[75308] 13638.Raymond, C. S.; Shamu, C. E.; Shen, M. M.; Seifert, K. J.; Hirsch, B.; Hodgkin, J.; Zarkower, D.: Evidence for evolutionary conservation of sex-determining genes. *Nature* 391: 691–695, 1998.

[75309] 13639.Shan, Z.; Zabel, B.; Trautmann, U.; Hillig, U.; Ottolenghi, C.; Wang, Y.; Haaf, T.: FISH mapping of the sex-reversal region on human chromosome 9p in two XY females and in primates. *Europ. J. Hum. Genet.* 8:167–173, 2000.

- [75310] 13640. Shen, M. M.; Hodgkin, J.: mab-3, a gene required for sex-specific yolk protein expression and a male-specific lineage in *C. elegans*. *Cell* 54:1019–1031, 1988.
- [75311] 13641. Smith, C. A.; McClive, P. J.; Western, P. S.; Reed, K. J.; Sinclair, A. H.: Conservation of a sex-determining gene. (Letter) *Nature* 402:601–602, 1999.
- [75312] 13642. Veitia, R.; Nunes, M.; Brauner, R.; Doco-Fenzy, M.; Joanny-Flinois, O.; Jaubert, F.; Lortat-Jacob, S.; Fellous, M.; McElreavey, K.: Deletion of distal 9p associated with 46,XY male to female sex reversal: definition of the break-points at 9p23.3–p24.1. *Genomics* 41: 271–274, 1997.
- [75313] 13643. Takekawa, M.; Posas, F.; Saito, H.: A human homolog of the yeast Ssk2/Ssk22 MAP kinase kinase kinases, MTK1, mediates stress-induced activation of the p38 and JNK pathways. *EMBO J.* 16: 4973–4982, 1997.
- [75314] 13644. Chapman, D. L.; Agulnik, I.; Hancock, S.; Silver, L. M.; Papaioannou, V. E.: Tbx6, a mouse T-box gene implicated in paraxial mesoderm formation at gastrulation. *Dev. Biol.* 180: 534–542, 1996.
- [75315] 13645. Chapman, D. L.; Papaioannou, V. E.: Three neural tubes in mouse embryos with mutations in the T-box gene Tbx6. *Nature* 391: 695–697, 1998.
- [75316] 13646. Papapetrou, C.; Putt, W.; Fox, M.; Edwards, Y. H.:

The human TBX6 gene: cloning and assignment to chromosome 16p11.2. *Genomics* 55:238–241, 1999.

[75317] 13647.Spicer, A. P.; Olson, J. S.; McDonald, J. A.: Molecular cloning and characterization of a cDNA encoding the third putative mammalian hyaluronan synthase. *J. Biol. Chem.* 272: 8957–8961, 1997.

[75318] 13648.Bagri, A.; Marin, O.; Plump, A. S.; Mak, J.; Pleasure, S. J.; Rubenstein, J. L. R.; Tessier-Lavigne, M.: Slit proteins prevent midline crossing and determine the dorsoventral position of major axonal pathways in the mammalian fore-brain. *Neuron* 33: 233–248, 2002.

[75319] 13649.Dallol, A.; Forgacs, E.; Martinez, A.; Sekido, Y.; Walker, R.; Kishida, T.; Rabbitts, P.; Maher, E. R.; Minna, J. D.; Latif, F.: Tumour specific promoter region methylation of the human homologue of the *Drosophila* Roundabout gene DUTT1 (ROBO1) in human cancers. *Oncogene* 21:3020–3028, 2002.

[75320] 13650.Kidd, T.; Brose, K.; Mitchell, K. J.; Fetter, R. D.; Tessier-Lavigne, M.; Goodman, C. S.; Tear, G.: Roundabout controls axon crossing of the CNS midline and defines a novel subfamily of evolutionarily conserved guidance receptors. *Cell* 92: 205–215, 1998.

[75321] 13651.Sundaresan, V.; Chung, G.; Heppell-Parton, A.;

Xiong, J.; Grundy, C.; Roberts, I.; James, L.; Cahn, A.; Bench, A.; Douglas, J.; Minna, J.; Sekido, Y.; Lerman, M.; Latif, F.; Bergh, J.; Li, H.; Lowe, N.; Ogilvie, D.; Rabbitts, P.: Homozygous deletions at 3p12 in breast and lung cancer. *Oncogene* 17: 1723–1729, 1998.

[75322] 13652. Zallen, J. A.; Yi, B. A.; Bargmann, C. I.: The conserved immunoglobulin superfamily member SAX-3/Robo directs multiple aspects of axon guidance in *C. elegans*. *Cell* 92: 217–227, 1998.

[75323] 13653. Fischer, U.; Heckel, D.; Michel, A.; Janka, M.; Hulsebos, T.; Meese, E.: Cloning of a novel transcription factor-like gene amplified in human glioma including astrocytoma grade I. *Hum. Molec. Genet.* 6: 1817–1822, 1997.

[75324] 13654. Gracia, E.; Fischer, U.; Elkahoul, A.; Trent, J. M.; Meese, E.; Meltzer, P. S.: Isolation of genes amplified in human cancers by microdissection mediated cDNA capture. *Hum. Molec. Genet.* 5: 595–600, 1996.

[75325] 13655. Gerard, M.; Hernandez, L.; Wevrick, R.; Stewart, C. L.: Disruption of the mouse *necln* gene results in early post-natal lethality. *Nature Genet.* 23: 199–202, 1999.

[75326] 13656. Hurst, L. D.; McVean, G.; Moore, T.: Imprinted genes have few and small introns. (Letter) *Nature Genet.* 12: 234–237, 1996.



- [75327] 13657.Jay, P.; Rougeulle, C.; Massacrier, A.; Moncla, A.; Mattei, M.-G.;Malzac, P.; Roeckel, N.; Taviaux, S.; Lefranc, J.-L. B.; Cau, P.;Berta, P.; Lalande, M.; Muscatelli, F.: The human necdin gene, NDN,is maternally imprinted and located in the Prader-Willi syndrome chromosomalregion. *Nature Genet.* 17: 357-360, 1997.
- [75328] 13658.MacDonald, H. R.; Wevrick, R.: The necdin gene is deleted in Prader-Willis syndrome and is imprinted in human and mouse. *Hum. Molec. Genet.* 6:1873-1878, 1997.
- [75329] 13659.Maruyama, K.; Usami, M.; Aizawa, T.; Yoshikawa, K.: A novel brain-specificmRNA encoding nuclear protein (necdin) expressed in neurally differentiatedembryonal carcinoma cells. *Biochem. Biophys. Res. Comm.* 178: 291-296,1991.
- [75330] 13660.Muscatelli, F.; Abrous, D. N.; Massacrier, A.; Boc-caccio, I.; LeMoal, M.; Cau, P.; Cremer, H.: Disruption of the mouse Necdin generesults in hypothalamic and behavioral alterations reminiscent ofthe human Prader-Willi syndrome. *Hum. Molec. Genet.* 9: 3101-3110,2000.
- [75331] 13661.Nakada, Y.; Taniura, H.; Uetsuki, T.; Inazawa, J.; Yoshikawa, K.: The human chromosomal gene for necdin, a neuronal growth suppressor,in the Prader-Willi syndrome deletion region. *Gene* 213: 65-72, 1998.

- [75332] 13662.Nicholls, R. D.: Incriminating gene suspects, Prader-Willi style. *NatureGenet.* 23: 132-134, 1999.
- [75333] 13663.Tsai, T.-F.; Armstrong, D.; Beaudet, A. L.: Necdin-deficient micedo not show lethality or the obesity and infertility of Prader-Willis syndrome. (Letter) *Nature Genet.* 22: 15-16, 1999.
- [75334] 13664.Watrin, F.; Roeckel, N.; Lacroix, L.; Mignon, C.; Mattei, M.-G.; Disteche, C.; Muscatelli, F.: The mouse necdin gene is expressed from the paternal allele only and lies in the 7C region of the mouse chromosome 7, a region of conserved synteny to the human Prader-Willi syndrome region. *Europ. J. Hum. Genet.* 5: 324-332, 1997.
- [75335] 13665.Delmas, V.; Stokes, D. G.; Perry, R. P.: A mammalian DNA-binding protein that contains a chromodomain and an SNF2/SWI2-like helicase domain. *Proc. Nat. Acad. Sci.* 90: 2414-2418, 1993.
- [75336] 13666.Woodage, T.; Basrai, M. A.; Baxevanis, A. D.; Hieter, P.; Collins, F. S.: Characterization of the CHD family of proteins. *Proc. Nat. Acad. Sci.* 94: 11472-11477, 1997.
- [75337] 13667.Ge, Q.; Nilasena, D. S.; O'Brien, C. A.; Frank, M. B.; Targoff, I. N.: Molecular analysis of a major antigenic region of the 240-kD protein of Mi-2 autoantigen. *J. Clin. Invest.* 96: 1730-1737, 1995.

- [75338] 13668.Seelig, H. P.; Renz, M.; Targoff, I. N.; Ge, Q.; Frank, M. B.:Two forms of the major antigenic protein of the dermatomyositis-specificMi-2 autoantigen. (Letter) Arthritis Rheum. 39: 1769–1771, 1996.
- [75339] 13669.Hu, G.; Chung, Y.-L.; Glover, T.; Valentine, V.; Look, A T.; Fearon,E. R.: Characterization of human homologs of the Drosophila sevenin absentia (sina) gene. Genomics 46: 103–111, 1997.
- [75340] 13670.Liu, J.; Stevens, J.; Rote, C. A.; Yost, H. J.; Hu, Y.; Neufeld,K. L.; White, R. L.; Matsunami, N.: Siah-1 mediates a novel beta-catenindegradation pathway linking p53 to the adenomatous polyposis coliprotein. Molec. Cell 7: 927–936, 2001.
- [75341] 13671.Nemani, M.; Linares-Cruz, G.; Bruzzoni-Giovanelli, H.; Roperch,J.-P.; Tuynder, M.; Bougueleret, L.; Cherif, D.; Medhioub, M.; Pasturaud,P.; Alvaro, V.; der Sarkissan, H.; Cazes, L.; Le Paslier, D.; Le Gall,I.; Israeli, D.; Dausset, J.; Sigaux, F.; Chumakov, I.; Oren, M.;Calvo, F.; Amson, R. B.; Cohen, D.; Telerman, A.: Activation of thehuman homologue of the Drosophila SINA gene in apoptosis and tumorsuppression. Proc. Nat. Acad. Sci. 93: 9039–9042, 1996.
- [75342] 13672.Ozkaynak, E.; Schnegelsberg, P. N. J.; Jin, D. F.;

Clifford, G.M.; Warren, F. D.; Drier, E. A.; Oppermann, H.: Osteogenic protein-2: a new member of the transforming growth factor-beta superfamily expressed in early embryogenesis. *J. Biol. Chem.* 267: 25220-25227, 1992.

[75343] 13673. Jang, W.; Weber, J. S.; Harkins, E. B.; Meisler, M. H.: Localization of the rhotekin gene RTKN on the physical maps of mouse chromosome 6 and human chromosome 2p13 and exclusion as a candidate for mnd2 and LGMD2B. *Genomics* 40: 506-507, 1997.

[75344] 13674. Reid, T.; Furuyashiki, T.; Ishizaki, T.; Watanabe, G.; Watanabe, N.; Fujisawa, K.; Morii, N.; Madaule, P.; Narumiya, S.: Rhotekin, a new putative target for Rho bearing homology to a serine/threonine kinase, PKN, and rhophilin in the Rho-binding domain. *J. Biol. Chem.* 271: 13556-13560, 1996.

[75345] 13675. Janssen, U.; Davis, E. M.; Le Beau, M. M.; Stoffel, W.: Human mitochondrial enoyl-CoA hydratase gene (ECHS1): structural organization and assignment to chromosome 10q26.2-q26.3. *Genomics* 40: 470-475, 1997.

[75346] 13676. Kanazawa, M.; Ohtake, A.; Abe, H.; Yamamoto, S.; Satoh, Y.; Takayanagi, M.; Niimi, H.; Mori, M.; Hashimoto, T.: Molecular cloning and sequence analysis of the cDNA for human mitochondrial short-chain enoyl-

CoA hydratase. Enzyme Protein 47: 9–13, 1993.

- [75347] 13677.Bingle, C. D.; Gowan, S.: Molecular cloning of the forkhead transcriptionfactor HNF-3-alpha from a human pulmonary adenocarcinoma cell line. Biochim.Biophys. Acta 1307: 17–20, 1996.
- [75348] 13678.Hannigan, G. E.; Bayani, J.; Weksberg, R.; Beatty, B.; Pandita,A.; Dedhar, S.; Squire, J.: Mapping of the gene encoding the integrin-linkedkinase, ILK, to human chromosome 11p15.5–p15.4. Genomics 42: 177–179,1997.
- [75349] 13679.Hannigan, G. E.; Leung–Hagesteijn, C.; Fitz–Gibbon, L.; Coppolino,M. G.; Radeva, G.; Filmus, J.; Bell, J. C.; Dedhar, S.: Regulationof cell adhesion and anchorage–dependent growth by a new beta–1–integrin–linkedprotein kinase. Nature 379: 91–96, 1996.
- [75350] 13680.Omeis, I. A.; Hsu, Y.–C.; Perin, M. S.: Mouse and human neuronalpentraxin 1 (NPXT1): conservation, genomic structure, and chromosomallocalization. Genomics 36: 543–545, 1996.
- [75351] 13681.Babic, A. M.; Kireeva, M. L.; Kolesnikova, T. V.; Lau, L. F.:CYR61, a product of a growth factor–inducible immediate early gene,promotes angiogenesis and tumor growth. Proc. Nat. Acad. Sci. 95:6355–6360, 1998.

- [75352] 13682.Jay, P.; Berge-Lefranc, J. L.; Marsollier, C.; Mejean, C.; Taviaux,S.; Berta, P.: The human growth factor-inducible immediate earlygene, CYR61, maps to chromosome 1p. *Oncogene* 14: 1753–1757, 1997.
- [75353] 13683.Kireeva, M. L.; Lam, S. C.–T.; Lau, L. F.: Adhesion of human umbilicalvein endothelial cells to the immediate-early gene product Cyr61 ismediated through integrin alpha(V)beta(3). *J. Biol. Chem.* 273: 3090–3096,1998.
- [75354] 13684.Martinerie, C.; Viegas-Pequignot, E.; Nguyen, V. C.; Perbal, B.: Chromosomal mapping and expression of the human cyr61 gene in tumourcells from the nervous system. *J. Clin. Path.* 50: 310–316, 1997.
- [75355] 13685.Sampath, D.; Zhu, Y.; Winneker, R. C.; Zhang, Z.: Aberrant expressionof Cyr61, a member of the CCN (CTGF/Cyr61/Cef10/NOVH) family, anddysregulation by 17-beta-estradiol and basic fibroblast growth factorin human uterine leiomyomas. *J. Clin. Endocr. Metab.* 86: 1707–1715,2001.
- [75356] 13686.Fujino, T.; Kang, M.–J.; Suzuki, H.; Iijima, H.; Yamamoto, T.:Molecular characterization and expression of rat acyl-CoA synthetase3. *J. Biol. Chem.* 271: 16748–16752, 1996.
- [75357] 13687.Minekura, H.; Fujino, T.; Kang, M.–J.; Fujita, T.;

Endo, Y.; Yamamoto, T. T.: Human acyl-coenzyme A synthetase 3 cDNA and localization of its gene (ACS3) to chromosome band 2q34-q35. *Genomics* 42: 180-181, 1997.

[75358] 13688. Kubota, N.; Terauchi, Y.; Miki, H.; Tamemoto, H.; Yamauchi, T.; Komeda, K.; Satoh, S.; Nakano, R.; Ishii, C.; Sugiyama, T.; Eto, K.; Tsubamoto, Y.; and 17 others: PPAR-gamma mediates high-fat diet-induced adipocyte hypertrophy and insulin resistance. *Molec. Cell* 4: 597-609, 1999.

[75359] 13689. Lapsys, N. M.; Kriketos, A. D.; Lim-Fraser, M.; Poynten, A. M.; Lowy, A.; Furler, S. M.; Chisholm, D. J.; Cooney, G. J.: Expression of genes involved in lipid metabolism correlate with peroxisome proliferator-activated receptor gamma expression in human skeletal muscle. *J. Clin. Endocr. Metab.* 85: 4293-4297, 2000.

[75360] 13690. Lehmann, J. M.; Moore, L. B.; Smith-Oliver, T. A.; Wilkison, W. O.; Willson, T. M.; Kliewer, S. A.: An antidiabetic thiazolidinedione is a high affinity ligand for peroxisome proliferator-activated receptor gamma (PPAR gamma). *J. Biol. Chem.* 270: 12953-12956, 1995.

[75361] 13691. Lowell, B. B.: PPAR-gamma: an essential regulator of adipogenesis and modulator of fat cell function. *Cell* 99: 239-242, 1999.

- [75362] 13692.Martin, G.; Schoonjans, K.; Staels, B.; Auwerx, J.: PPAR-gammaactivators improve glucose homeostasis by stimulating fatty acid uptakein the adipocytes. *Atherosclerosis* 137: S75-S80, 1998.
- [75363] 13693.Meirhaeghe, A.; Fajas, L.; Helbecque, N.; Cottel, D.; Lebel, P.;Dallongeville, J.; Deeb, S.; Auwerx, J.; Amouyel, P.: A genetic polymorphismof the peroxisome prolifera-tor-activated receptor gamma gene influencesplasma lep-tin levels in obese tumors. *Hum. Molec. Genet.* 7: 435-440,1998.
- [75364] 13694.Miles, P. D. G.; Barak, Y.; He, W.; Evans, R. M.; Olefsky, J.M.: Improved insulin-sensitivity in mice heterozygous for PPAR-gammadeficiency. *J. Clin. Invest.* 105: 287-292, 2000.
- [75365] 13695.Mueller, E.; Sarraf, P.; Tontonoz, P.; Evans, R. M.; Martin, K.J.; Zhang, M.; Fletcher, C.; Singer, S.; Spiegelman, B. M.: Terminaldifferentiation of human breast cancer through PPAR-gamma. *Molec.Cell.* 1: 465-470, 1998.
- [75366] 13696.Mueller, E.; Smith, M.; Sarraf, P.; Kroll, T.; Aiyer, A.; Kaufman,D. S.; Oh, W.; Demetri, G.; Figg, W. D.; Zhou, X.-P.; Eng, C.; Spiegelman,B. M.; Kantoff, P. W.: Effects of ligand activation of peroxisomeproliferator-activated receptor gamma in human prostate cancer. *Proc.Nat. Acad.*



Sci. 97: 10990–10995, 2000.

- [75367] 13697. Nagy, L.; Tontonoz, P.; Alvarez, J. G. A.; Chen, H.; Evans, R.M.: Oxidized LDL regulates macrophage gene expression through ligand activation of PPAR- $\gamma$ . *Cell* 93: 229–240, 1998.
- [75368] 13698. Oh, E. Y.; Min, K. M.; Chung, J. H.; Min, Y.-K.; Lee, M.-S.; Kim, K.-W.; Lee, M.-K.: Significance of pro12ala mutation in peroxisome proliferator-activated receptor- $\gamma$ 2 in Korean diabetic and obese subjects. *J. Clin. Endocr. Metab.* 85: 1801–1804, 2000.
- [75369] 13699. Pritchard, J. K.; Rosenberg, N. A.: Use of unlinked genetic markers to detect population stratification in association studies. *Am. J. Hum. Genet.* 65: 220–228, 1999.
- [75370] 13700. Ristow, M.; Muller-Wieland, D.; Pfeiffer, A.; Krone, W.; Kahn, C. R.: Obesity associated with a mutation in a genetic regulator of adipocyte differentiation. *New Eng. J. Med.* 339: 953–959, 1998.
- [75371] 13701. Rocchi, S.; Picard, F.; Vamecq, J.; Gelman, L.; Potier, N.; Zeyer, D.; Dubuquoy, L.; Bac, P.; Champy, M.-F.; Plunket, K. D.; Leesnitzer, L. M.; Blanchard, S. G.; Desreumaux, P.; Moras, D.; Renaud, J.-P.; Auwerx, J.: A unique PPAR- $\gamma$  ligand with potent insulin-sensitizing yet weak adipogenic activity. *Molec. Cell* 8: 737–747, 2001.

- [75372] 13702.Rosen, E. D.; Sarraf, P.; Troy, A. E.; Bradwin, G.; Moore, K.; Milstone, D. S.; Spiegelman, B. M.; Mortensen, R. M.: PPAR-gamma is required for the differentiation of adipose tissue in vivo and in vitro. *Molec. Cell* 4: 611–617, 1999.
- [75373] 13703.Ricote, M.; Huang, J.; Fajas, L.; Li, A.; Welch, J.; Najib, J.; Witztum, J. L.; Auwerx, J.; Palinski, W.; Glass, C. K.: Expression of the peroxisome proliferator-activated receptor gamma (PPAR-gamma) in human atherosclerosis and regulation in macrophages by colony stimulating factors and oxidized low density lipoprotein. *Proc. Nat. Acad. Sci.* 95:7614–7619, 1998.
- [75374] 13704.Savage, D. B.; Agostini, M.; Barroso, I.; Gurnell, M.; Luan, J.; Meirhaeghe, A.; Harding, A.-H.; Ihrke, G.; Rajanayagam, O.; Soos, M. A.; George, S.; Berger, D.; and 9 others: Digenic inheritance of severe insulin resistance in a human pedigree. *Nature Genet.* 31:379–384, 2002. Note: Erratum: *Nature Genet.* 32: 211 only, 2002.
- [75375] 13705.Smith, W. M.; Zhou, X.-P.; Kurose, K.; Gao, X.; Latif, F.; Kroll, T.; Sugano, K.; Cannistra, S. A.; Clinton, S. K.; Maher, E. R.; Prior, T. W.; Eng, C.: Opposite association of two PPARG variants with cancer: overrepresentation of H449H in endometrial carcinoma cases and underrepresenta-

tion of P12A in renal cell carcinoma cases. Hum. Genet. 109: 146–151, 2001.

- [75376] 13706. Tarrade, A.; Schoonjans, K.; Pavan, L.; Auwerx, J.; Rochette-Egly, C.; Evain-Brion, D.; Fournier, T.: PPAR-gamma/RXR-alpha heterodimers control human trophoblast invasion. J. Clin. Endocr. Metab. 86: 5017–5024, 2001.
- [75377] 13707. Tontonoz, P.; Hu, E.; Devine, J.; Beale, E. G.; Spiegelman, B. M.: PPAR gamma 2 regulates adipose expression of the phosphoenolpyruvate carboxykinase gene. Molec. Cell. Biol. 15: 351–357, 1995.
- [75378] 13708. Tontonoz, P.; Hu, E.; Graves, R. A.; Budavari, A. I.; Spiegelman, B. M.: mPPAR gamma 2: tissue-specific regulator of an adipocyte enhancer. Genes Dev. 8: 1224–1234, 1994.
- [75379] 13709. Tontonoz, P.; Nagy, L.; Alvarez, J. G. A.; Thomazy, V. A.; Evans, R. M.: PPAR-gamma promotes monocyte/macrophage differentiation and uptake of oxidized LDL. Cell 93: 241–252, 1998.
- [75380] 13710. Wang, M.: Isolation and characterization of a human gene encoding a single-strand-specific endoribonuclease. Ph.D. Thesis: Stanford Univ. , 1995.
- [75381] 13711. Wang, M.; Cohen, S. N.: ard-1: a human gene that

reverses the effects of temperature-sensitive and deletion mutations in the *Escherichia coli* *rne* gene and encodes an activity producing RNase E-like cleavages. *Proc. Nat. Acad. Sci.* 91: 10591–10595, 1994.

[75382] 13712. Beck, S.; Hanson, I.; Kelly, A.; Pappin, D. J.; Trowsdale, J.: A homologue of the *Drosophila* female sterile homeotic (*fsh*) gene in the class II region of the human MHC. *DNA Sequence* 2: 203–210, 1992.

[75383] 13713. Denis, G. V.; Green, M. R.: A novel, mitogen-activated nuclear kinase is related to a *Drosophila* developmental regulator. *Genes Dev.* 10: 261–271, 1996.

[75384] 13714. Okamoto, N.; Ando, A.; Kawai, J.; Yoshiwara, T.; Tsuji, K.; Inoko, H.: Orientation of HLA-DNA gene and identification of a CpG island-associated gene adjacent to DNA in human major histocompatibility complex class II region. *Hum. Immun.* 32: 221–228, 1991.

[75385] 13715. Thorpe, K. L.; Abdulla, S.; Kaufman, J.; Trowsdale, J.; Beck, S.: Phylogeny and structure of the RING3 gene. *Immunogenetics* 44: 391–396, 1996.

[75386] 13716. Cushman, L. J.; Camper, S. A.: Molecular basis of pituitary dysfunction in mouse and human. *Mammalian Genome* 12: 485–494, 2001.

[75387] 13717. Kozak, M.: Interpreting cDNA sequences: some in-

sights from studies on translation. *Mammalian Genome* 7: 563–574, 1996.

- [75388] 13718. Pellegrini-Bouiller, I.; Manrique, C.; Gunz, G.; Grino, M.; Zamora, A. J.; Figarella-Branger, D.; Grisoli, F.; Jaquet, P.; Enjalbert, A.: Expression of the members of the Ptx family of transcription factors in human pituitary adenomas. *J. Clin. Endocr. Metab.* 84:2212–2220, 1999.
- [75389] 13719. Brinke, A.; Green, P. M.; Giannelli, F.: Characterization of the gene (VBP1) and transcript for the von Hippel-Lindau binding protein and isolation of the highly conserved murine homologue. *Genomics* 45:105–112, 1997.
- [75390] 13720. Brinke, A.; Tagliavacca, L.; Naylor, J.; Green, P.; Giannelli, F.: Two chimaeric transcription units result from an inversion breaking intron 1 of the factor VIII gene and a region reportedly affected by reciprocal translocations in T-cell leukaemia. *Hum. Molec. Genet.* 5: 1945–1951, 1996.
- [75391] 13721. Clifford, S. C.; Walsh, S.; Hewson, K.; Green, E. K.; Brinke, A.; Green, P. M.; Giannelli, F.; Eng, C.; Maher, E. R.: Genomic organization and chromosomal localization of the human CUL2 gene and the role of von Hippel-Lindau tumor suppressor-binding protein (CUL2 and VBP1) mutation and loss in renal-cell carcinoma development. *Genes*

ChromosomesCancer 26: 20–28, 1999.

- [75392] 13722.Hemberger, M.; Himmelbauer, H.; Neumann, H. P. H.; Plate, K. H.;Schwarzkopf, G.; Fundele, R.: Expression of the von Hippel–Lindau–bindingprotein–1 (Vbp1) in fetal and adult mouse tissues. Hum. Molec. Genet. 8:229–236, 1999.
- [75393] 13723.Tsuchiya, H.; Iseda, T.; Hino, O.: Identification of a novel protein(VBP–1) binding to the von Hippel–Lindau (VHL) tumor suppressor geneproduct. Cancer Res. 56: 2881–2885, 1996.
- [75394] 13724.Corbaz, A.; ten Hove, T.; Herren, S.; Graber, P.; Schwartsburd,B.; Belzer, I.; Harrison, J.; Plitz, T.; Kosco–Vilbois, M. H.; Kim,S.–H.; Dinarello, C. A.; Novick, D.; van Deventer, S.; Chvatchko,Y.: IL–18–binding protein expression by endothelial cells and macrophages up–regulated during active Crohn's disease. J. Immun. 168: 3608–3616,2002.
- [75395] 13725.Turner, N.; Mason, P. J.; Brown, R.; Fox, M.; Povey, S.; Rees,A.; Pusey, C. D.: Molecular cloning of the human Goodpasture antigendemonstrates it to be the alpha–3 chain of type IV collagen. J. Clin.Invest. 89: 592–601, 1992.
- [75396] 13726.Hoffmann, M. M.; Jacob, S.; Luft, D.; Schmulling,

R.-M.; Rett, K.; Marz, W.; Haring, H.-U.; Matthaei, S.: Type I hyperlipoproteinemia due to a novel loss of function mutation of lipoprotein lipase, cys239-trp, associated with recurrent severe pancreatitis. *J. Clin. Endocr. Metab.* 85:4795-4798, 2000.

[75397] 13727. Holt, L. E., Jr.; Aylward, F. X.; Timbers, H. G.: Idiopathic familial lipemia. *Bull. Johns Hopkins Hosp.* 64: 279-314, 1939.

[75398] 13728. Ishimura-Oka, K.; Faustinella, F.; Kihara, S.; Smith, L. C.; Oka, K.; Chan, L.: A missense mutation (trp86-to-arg) in exon 3 of the lipoprotein lipase gene: a cause of familial chylomicronemia. *Am. J. Hum. Genet.* 50: 1275-1280, 1992.

[75399] 13729. Kastelein, J. J. P.; Groenemeyer, B. E.; Hallman, D. M.; Henderson, H.; Reymer, P. W. A.; Gagne, S. E.; Jansen, H.; Seidell, J. C.; Kromhout, D.; Jukema, J. W.; Bruschke, A. V. G.; Boerwinkle, E.; Hayden, M. R.; The Regress Study Group: The asn9 variant of lipoprotein lipase is associated with the -93G promoter mutation and an increased risk of coronary artery disease. *Clin. Genet.* 53: 27-33, 1998.

[75400] 13730. Kirchgessner, T. G.; Chuat, J.-C.; Heinzmann, C.; Etienne, J.; Guilhot, S.; Svenson, K.; Ameis, D.; Pilon, C.; d'Auriol, L.; Andalibi, A.; Schotz, M. C.; Galibert, F.; Lusi,

A. J.: Organization of the human lipoprotein lipase gene and evolution of the lipase gene family. *Proc. Nat. Acad. Sci.* 86: 9647–9651, 1989.

[75401] 13731. Kirchgessner, T. G.; Svenson, K. L.; Lusis, A. J.; Schotz, M.C.: The sequence of cDNA encoding lipoprotein lipase: a member of a lipase gene family. *J. Biol. Chem.* 262: 8463–8466, 1987.

[75402] 13732. Kobayashi, J.; Nishida, T.; Ameis, D.; Stahnke, G.; Schotz, M.C.; Hashimoto, H.; Fukamachi, I.; Shirai, K.; Saito, Y.; Yoshida, S.: A heterozygous mutation (the codon for ser477 to a stop codon) in lipoprotein lipase contributes to a defect in lipid interface recognition in a case with type I hyperlipidemia. *Biochem. Biophys. Res. Commun.* 182:70–77, 1992.

[75403] 13733. Kobayashi, J.; Sasaki, N.; Tashiro, J.; Inadera, H.; Saito, Y.; Yoshida, S.: A missense mutation (ala334-to-thr) in exon 7 of the lipoprotein lipase gene in a case with type I hyperlipidemia. *Biochem. Biophys. Res. Commun.* 191: 1046–1054, 1993.

[75404] 13734. Langlois, S.; Deeb, S.; Brunzell, J.; Kastelein, J. J.; Hayden, M. R.: A unique insertion accounts for a significant proportion of the mutations in the lipoprotein lipase (LPL) gene. (Abstract) *Am. J. Hum. Genet.* 43: A191, 1988.



- [75405] 13735.Langlois, S.; Deeb, S.; Brunzell, J. D.; Kastelein, J. J.; Hayden,M. R.: A major insertion accounts for a significant proportion of mutations underlying human lipoprotein lipase deficiency. *Proc. Nat.Acad. Sci.* 86: 948–952, 1989.
- [75406] 13736.Levak–Frank, S.; Radner, H.; Walsh, A.; Stollberger, R.; Knipping,G.; Hoefler, G.; Sattler, W.; Weinstock, P. H.; Breslow, J. L.; Zechner,R.: Muscle–specific overexpression of lipoprotein lipase causes a severe myopathy characterized by proliferation of mitochondria and peroxisomes in transgenic mice. *J. Clin. Invest.* 96: 976–986, 1995.
- [75407] 13737.Li, S.; Oka, K.; Galton, D.; Stocks, J.: Pvu–II RFLP at the human lipoprotein lipase (LPL) gene locus. *Nucleic Acids Res.* 16: 2358,1988.
- [75408] 13738.Lo, J. Y.; Smith, L. C.; Chan, L.: Lipoprotein lipase: role of intramolecular disulfide bonds in enzyme catalysis. *Biochem. Biophys.Res. Commun.* 206: 266–271, 1995.
- [75409] 13739.Ma, Y.; Henderson, H. E.; Julien, P.; Roederer, G.; Brunzell,J.; Hayden, M. R.: A missense mutation (pro–to–leu207) in the human lipoprotein lipase gene is the major cause of type I hyperlipoproteinemia in French Canadians. (Series) *Miami Short Reports. Advances in Gene Technology: The Molecular Biology of Human Genetic Disease.* New York: IRL Press (pub.) 1: 1991. Pp. 34 only.

- [75410] 13740.Ma, Y.; Henderson, H. E.; Ven Murthy, M. R.; Roederer, G.; Monsalve, M. V.; Clarke, L. A.; Normand, T.; Julien, P.; Gagne, C.; Lambert, M.; Davignon, J.; Lupien, P. J.; Brunzell, J.; Hayden, M. R.: A mutation in the human lipoprotein lipase gene as the most common cause of familial chylomicronemia in French Canadians. *New Eng. J. Med.* 324: 1761–1766, 1991.
- [75411] 13741.Ma, Y.; Liu, M.-S.; Chitayat, D.; Bruin, T.; Beisiegel, U.; Benlian, P.; Foubert, L.; De Gennes, J. L.; Funke, H.; Forsythe, I.; Blauchman, S.; Papanikolaou, M.; Erkelens, D. W.; Kastelein, J.; Brunzell, J.D.; Hayden, M. R.: Recurrent missense mutations at the first and second base of codon arg243 in human lipoprotein lipase in patients of different ancestries. *Hum. Mutat.* 3: 52–58, 1994.
- [75412] 13742.Ma, Y.; Liu, M.-S.; Ginzinger, D.; Frohlich, J.; Brunzell, J.D.; Hayden, M. R.: Gene–environment interaction in the conversion of a mild–to–severe phenotype in a patient homozygous for a ser172–to–cys mutation in the lipoprotein lipase gene. *J. Clin. Invest.* 91: 1953–1958, 1993.
- [75413] 13743.Ma, Y.; Wilson, B. I.; Bijvoet, S.; Henderson, H. E.; Cramb, E.; Roederer, G.; Ven Murthy, M. R.; Julien, P.; Bakker, H. D.; Kastelein, J. J. P.; Brunzell, J. D.; Hayden, M. R.: A missense mutation (asp250–to–asn) in exon 6 of the

human lipoprotein lipase gene causes chylomicronemia in patients of different ancestries. *Genomics* 13: 649–653, 1992.

- [75414] 13744. Mattei, M. G.; Etienne, J.; Chuat, J. C.; Nguyen, V. C.; Brault, D.; Bernheim, A.; Galibert, F.: Assignment of the human lipoprotein lipase (LPL) gene to chromosome band 8p22. *Cytogenet. Cell Genet.* 63:45–46, 1993.
- [75415] 13745. Miesenbock, G.; Holzl, B.; Foger, B.; Brandstatter, E.; Paulweber, B.; Sandhofer, F.; Patsch, J. R.: Heterozygous lipoprotein lipase deficiency due to a missense mutation as the cause of impaired triglyceride tolerance with multiple lipoprotein abnormalities. *J. Clin. Invest.* 91:448–455, 1993.
- [75416] 13746. Nevin, D. N.; Brunzell, J. D.; Deeb, S. S.: The LPL gene in individuals with familial combined hyperlipidemia and decreased LPL activity. *Arteriosclerosis Thromb.* 14: 869–873, 1994.
- [75417] 13747. Nevin, N. C.; Slack, J.: Hyperlipidaemic xanthomatosis II: mode of inheritance in 55 families with essential hyperlipidaemia and xanthomatosis. *J. Med. Genet.* 5: 9–28, 1968.
- [75418] 13748. Nickerson, D. A.; Taylor, S. L.; Weiss, K. M.; Clark, A. G.; Hutchinson, R. G.; Stengard, J.; Salomaa, V.; Varti-

ainen, E.; Boerwinkle, E.; Sing, C. F.: DNA sequence diversity in a 9.7-kb region of the human lipoprotein lipase gene. *Nature Genet.* 19: 233–240, 1998.

[75419] 13749. Lahn, B. T.; Page, D. C.: Retroposition of autosomal mRNA yielded testis-specific gene family on human Y chromosome. *Nature Genet.* 21: 429–433, 1999.

[75420] 13750. Yen, P. H.: A long-range restriction map of deletion interval 6 of the human Y chromosome: a region frequently deleted in azoospermic males. *Genomics* 54: 5–12, 1998.

[75421] 13751. MacLean, H. E.; Chu, S.; Warne, G. L.; Zajac, J. D.: Related individuals with different androgen receptor gene deletions. *J. Clin. Invest.* 91: 1123–1128, 1993.

[75422] 13752. Madgar, I.; Green, L.; Kent-First, M.; Weissenberg, R.; Gershoni-Baruch, R.; Goldman, B.; Friedman, E.: Genotyping of Israeli infertile men with idiopathic oligozoospermia. *Clin. Genet.* 62: 203–207, 2002.

[75423] 13753. Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Griffin, J. E.; Wilson, J. D.; McPhaul, M. J.: Definition of the human androgen receptor gene structure permits the identification of mutations that cause androgen resistance: premature termination of the receptor protein at amino acid residue 588 causes complete androgen resistance. *Molec. Endocr.* 4: 1105–1116, 1990.

[75424] 13754. Marcelli, M.; Tilley, W. D.; Wilson, C. M.; Wilson, J. D.; Griffin, J. E.; McPhaul, M. J.: A single nucleotide substitution introduces a premature termination codon into the androgen receptor gene of a patient with receptor-negative androgen resistance. *J. Clin. Invest.* 85:1522–1528, 1990.

- [75425] 13755. Marcelli, M.; Tilley, W. D.; Zoppi, S.; Griffin, J. E.; Wilson, J. D.; McPhaul, M. J.: Androgen resistance associated with a mutation of the androgen receptor at amino acid 772 (arg-to-cys) results from a combination of decreased messenger ribonucleic acid levels and impairment of receptor function. *J. Clin. Endocr.* 73: 318–325, 1991.
- [75426] 13756. McCampbell, A.; Taylor, J. P.; Taye, A. A.; Robitschek, J.; Li, M.; Walcott, J.; Merry, D.; Chai, Y.; Paulson, H.; Sobue, G.; Fischbeck, K. H.: CREB-binding protein sequestration by expanded polyglutamine. *Hum. Molec. Genet.* 9: 2197–2202, 2000.
- [75427] 13757. McPhaul, M. J.; Griffin, J. E.: Male pseudohermaphroditism caused by mutations of the human androgen receptor. *J. Clin. Endocr. Metab.* 84: 3435–3441, 1999.
- [75428] 13758. McPhaul, M. J.; Marcelli, M.; Tilley, W. D.; Griffin, J. E.; Isidro-Gutierrez, R. F.; Wilson, J. D.: Molecular basis of androgen resistance in a family with a qualitative abnormality of the androgen receptor and responsive to high-dose androgen therapy. *J. Clin. Invest.* 87: 1413–1421, 1991.
- [75429] 13759. McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Griffin, J. E.;

Wilson, J. D.: Genetic basis of endocrine disease 4: the spectrum of mutations in the androgen receptor gene that causes androgen resistance. *J. Clin. Endocr. Metab.* 76: 17–23, 1993.

- [75430] 13760. McPhaul, M. J.; Marcelli, M.; Zoppi, S.; Wilson, C. M.; Griffin, J. E.; Wilson, J. D.: Mutations in the ligand-binding domain of the androgen receptor gene cluster in two regions of the gene. *J. Clin. Invest.* 90: 2097–2101, 1992.
- [75431] 13761. McPhaul, M. J.; Schweikert, H.-U.; Allman, D. R.: Assessment of androgen receptor function in genital skin fibroblasts using a recombinant adenovirus to deliver an androgen-responsive reporter gene. *J. Clin. Endocr. Metab.* 82: 1944–1948, 1997.
- [75432] 13762. Mifsud, A.; Ramirez, S.; Yong, E. L.: Androgen receptor gene CAG trinucleotide repeats in anovulatory infertility and polycystic ovaries. *J. Clin. Endocr. Metab.* 85: 3484–3488, 2000.
- [75433] 13763. Migeon, B. R.; Brown, T. R.; Axelman, J.; Migeon, C. J.: Studies of the locus for androgen receptor: localization on the human X and evidence for homology with the Tfm locus in the mouse. *Proc. Nat. Acad. Sci.* 78: 6339–6343, 1981.

- [75434] 13764.Mongan, N. P.; Jaaskelainen, J.; Green, K.; Schwabe, J. W.; Shimura,N.; Dattani, M.; Hughes, I. A.: Two de novo mutations in the AR gene cause the complete androgen insensitivity syndrome in a pair of monozygotic twins. *J. Clin. Endocr. Metab.* 87: 1057–1061, 2002.
- [75435] 13765.Mononen, N.; Syrjakoski, K.; Matikainen, M.; Tam-mela, T. L. J.;Schleutker, J.; Kallioniemi, O.–P.; Trapman, J.; Koivisto, P. A.:Two percent of Finnish prostate cancer patients have a germ–line mutation in the hormone–binding domain of the androgen receptor gene. *Cancer Res.* 60: 6479–6481, 2000.
- [75436] 13766.Murono, K.; Mendonca, B. B.; Arnhold, I. J. P.; Rigon, A. C. M.M.; Migeon, C. J.; Brown, T. R.: Human androgen insensitivity due to point mutations encoding amino acid substitutions in the androgen receptor steroid–binding domain. *Hum. Mutat.* 6: 152–162, 1995.
- [75437] 13767.Nakao, R.; Haji, M.; Yanase, T.; Ogo, A.; Takayanagi, R.; Katsube,T.; Fukumaki, Y.; Nawata, H.: A single amino acid substitution (met786–to–val) in the steroid–binding domain of human androgen receptor leads to complete androgen insensitivity syndrome. *J. Clin. Endocr. Metab.* 74:1152–1157, 1992.
- [75438] 13768.Newmark, J. R.; Hardy, D. O.; Tonb, D. C.; Carter, B.



S.; Epstein, J. I.; Isaacs, W. B.; Brown, T. R.; Barrack, E. R.: Androgen receptor gene mutations in human prostate cancer. *Proc. Nat. Acad. Sci.* 89:6319–6323, 1992.

[75439] 13769. Nguyen, D.; Steinberg, S. V.; Rouault, E.; Chagnon, S.; Gottlieb, B.; Pinsky, L.; Trifiro, M.; Mader, S.: A G577R mutation in the human AR P box results in selective decreases in DNA binding and in partial androgen insensitivity syndrome. *Molec. Endocr.* 15: 1790–1802, 2001.

[75440] 13770. Ohno, S.: The Y-linked antigen locus and the X-linked Tfm locus as major regulatory genes of the mammalian sex determining mechanism. *J. Steroid Biochem.* 8: 585–592, 1977. 100. Ohno, S.: Simplicity of mammalian regulatory systems inferred by single gene determination of sex phenotypes. *Nature* 234: 134–137, 1971. 101. Ong, Y. C.; Wong, H. B.; Adaikan, G.; Yong, E. L.: Directed pharmacological therapy of ambiguous genitalia due to an androgen receptor gene mutation. (Letter) *Lancet* 354: 1444–1445, 1999. 102. Patterson, M. N.; Hughes, I. A.; Gottlieb, B.; Pinsky, L.: The androgen receptor gene mutations database. *Nucleic Acids Res.* 22:3560–3562, 1994. 103. Pinsky, L.; Kaufman, M.; Killinger, D. W.; Burko, B.; Shatz, D.; Volpe, R.: Human minimal androgen insensitivity with normal dihydrotestosterone-binding capacity in

cultured genital skin fibroblasts: evidence for an androgen-selective qualitative abnormality of the receptor. *Am. J. Hum. Genet.* 36:965–978, 1984.104. Pinsky, L.; Kaufman, M.; Levitsky, L. L.: Partial androgen resistance due to a distinctive qualitative defect of the androgen receptor. *Am. J. Med. Genet.* 27: 459–466, 1987.105. Pinsky, L.; Kaufman, M.; Summitt, R. L.: Congenital androgen insensitivity due to a qualitatively abnormal androgen receptor. *Am. J. Med. Genet.* 10: 91–99, 1981.106. Pinsky, L.; Trifiro, M.; Sebbaghian, N.; Kaufman, M.; Chang, C.; Trapman, J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris, C. J.; Brown, C. J.; Willard, H. F.; Sergovich, F.: A deletional alteration of the androgen receptor (AR) gene in a sporadic patient with complete androgen insensitivity (CAI) who is mentally retarded. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A212, 1989.107. Prior, L.; Bordet, S.; Trifiro, M. A.; Mhatre, A.; Kaufman, M.; Pinsky, L.; Wrogeman, K.; Belsham, D. D.; Pereira, F.; Greenberg, C.; Trapman, J.; Brinkman, A. O.; Chang, C.; Liao, S.: Replacement of arginine 773 by cysteine or histidine in the human androgen receptor causes complete androgen insensitivity with different receptor phenotypes. *Am. J. Hum. Genet.* 51: 143–155, 1992.108. Quigley, C. A.; Friedman, K. J.; Johnson, A.; Lafreniere,

R.G.; Silverman, L. M.; Lubahn, D. B.; Brown, T. R.; Wilson, E. M.; Willard, H. F.; French, F. S.: Complete deletion of the androgenreceptor gene: definition of the null phenotype of the androgen insensitivitysyndrome and determination of carrier status. *J. Clin. Endocr. Metab.* 74:927–933, 1992.109. Ris–Stalpers, C.; Kuiper, G. G. J. M.; Faber, P. W.; Schweikert, H. U.; van Rooij, H. C. J.; Zegers, N. D.; Hodgins, M. B.; Degenhart, H. J.; Trapman, J.; Brinkmann, A. O.: Aberrant splicing of androgenreceptor mRNA results in synthesis of a nonfunctional receptor protein in a patient with androgen insensitivity. *Proc. Nat. Acad. Sci.* 87:7866–7870, 1990.110. Rodien, P.; Mebarki, F.; Mowszowicz, I.; Chaussain, J.–L.; Young, J.; Morel, Y.; Schaison, G.: Different phenotypes in a family with androgen insensitivity caused by the same M780I point mutation in the androgen receptor gene. *J. Clin. Endocr. Metab.* 81: 2994–2998, 1996.111. Sai, T.; Seino, S.; Chang, C.; Trifiro, M.; Pinsky, L.; Mhatre, A.; Kaufman, M.; Lambert, B.; Trapman, J.; Brinkmann, A. O.; Rosenfield, R. L.; Liao, S.: An exonic point mutation of the androgen receptorgene in a family with complete androgen insensitivity. *Am. J. Hum. Genet.* 46: 1095–1100, 1990.112. Sammarco, I.; Grimaldi, P.; Rossi, P.; Cappa, M.; Moretti, C.; Frajese, G.;

Geremia, R.: Novel point mutation in the splice donorsite of exon–intron junction 6 of the androgen receptor gene in a patient with partial androgen insensitivity syndrome. *J. Clin. Endocr. Metab.* 85: 3256–3261, 2000.113. Schoenberg, M. P.; Hakimi, J. M.; Wang, S.; Bova, G. S.; Epstein, J. I.; Fischbeck, K. H.; Isaacs, W. B.; Walsh, P. C.; Barrack, E. R.: Microsatellite mutation (CAG (24–to–18)) in the androgen receptor gene in human prostate cancer *Biochem. Biophys. Res. Commun.* 198:74–80, 1994.114. Shang, Y.; Myers, M.; Brown, M.: Formation of the androgen receptor transcription complex. *Molec. Cell* 9: 601–610, 2002.115. Simeoni, S.; Mancini, M. A.; Stenoien, D. L.; Marcelli, M.; Weigel, N. L.; Zanisi, M.; Martini, L.; Poletti, A.: Motoneuronal cell death is not correlated with aggregate formation of androgen receptors containing an elongated polyglutamine tract. *Hum. Molec. Genet.* 9: 133–144, 2000.116. Sullivan, D. A.; Sullivan, B. D.; Ullman, M. D.; Rocha, E. M.; Krenzer, K. L.; Cermak, J. M.; Toda, I.; Doane, M. G.; Evans, J. E.; Wickham, L. A.: Androgen influence on the meibomian gland. *Invest. Ophthalm. Vis. Sci.* 41: 3732–3742, 2000.117. Sultan, C.; Lumbroso, S.; Poujol, N.; Belon, C.; Boudon, C.; Lobaccaro, J.–M.: Mutations of androgen receptor gene in androgen insensitivity syn–

dromes. J. Steroid Biochem. Molec. Biol. 46: 519–530, 1993.118. Sutherland, R. W.; Wiener, J. S.; Hicks, J. P.; Marcelli, M.; Gonzales, E. T.; Roth, D. R.; Lamb, D. J.: Androgen receptor gene mutations are rarely associated with isolated penile hypospadias. J. Urol. 156: 828–831, 1996.119. Taplin, M.-E.; Bubley, G. J.; Shuster, T. D.; Frantz, M. E.; Spooner, A. E.; Ogata, G. K.; Keer, H. N.; Balk, S. P.: Mutation of the androgen-receptor gene in metastatic androgen-independent prostate cancer. New Eng. J. Med. 332: 1393–1398, 1995.120. Tilley, W. D.; Marcelli, M.; Wilson, J. D.; McPhaul, M. J.: Characterization and expression of a cDNA encoding the human androgen-receptor. Proc. Nat. Acad. Sci. 86: 327–331, 1989.121. Trifiro, M.; Prior, L.; Pinsky, L.; Kaufman, M.; Chang, C.; Trapman, J.; Brinkmann, A. O.; Kuiper, G. G. J. M.; Ris, C.: A single transition at an exonic CpG site apparently abolishes androgen receptor (AR)-binding activity in a family with complete androgen insensitivity (CAI). (Abstract) Am. J. Hum. Genet. 45 (suppl.): A225, 1989.122. Trifiro, M.; Prior, R. L.; Sabbaghian, N.; Pinsky, L.; Kaufman, M.; Nylén, E. G.; Belsham, D. D.; Greenberg, C. R.; Wrogermann, K.: Amber mutation creates a diagnostic Mael site in the androgen receptor gene of a family with complete

androgen insensitivity. *Am. J. Med. Genet.* 40: 493–499, 1991.123. Tut, T. G.; Ghadessy, F. J.; Trifiro, M. A.; Pinsky, L.; Yong, E. L.: Long polyglutamine tracts in the androgen receptor are associated with reduced trans-activation, impaired sperm production, and male infertility. *J. Clin. Endocr. Metab.* 82: 3777–3782, 1997.124. Visakorpi, T.; Hyytinen, E.; Koivisto, P.; Tanner, M.; Keinänen, R.; Palmberg, C.; Palotie, A.; Tammela, T.; Isola, J.; Kallioniemi, O.-P.: In vivo amplification of the androgen receptor gene and progression of human prostate cancer. *Nature Genet.* 9: 401–406, 1995.125. Von Eckardstein, S.; Syska, A.; Gromoll, J.; Kamischke, A.; Simoni, M.; Nieschlag, E.: Inverse correlation between sperm concentration and number of androgen receptor CAG repeats in normal men. *J. Clin. Endocr. Metab.* 86: 2585–2590, 2001.126. Wang, Q.; Ghadessy, F. J.; Trounson, A.; de Kretser, D.; McLachlan, R.; Ng, S. C.; Yong, E. L.: Azoospermia associated with a mutation in the ligand-binding domain of an androgen receptor displaying normal ligand binding, but defective trans-activation. *J. Clin. Endocr. Metab.* 83: 4303–4309, 1998.127. Weidemann, W.; Peters, B.; Romalo, G.; Spindler, K.-D.; Schweikert, H.-U.: Response to androgen treatment in a patient with partial androgen insensitivity

and a mutation in the deoxyribonucleic acid-binding domain of the androgen receptor. *J. Clin. Endocr. Metab.* 83: 1173–1176, 1998.128. Welch, W. J.; Diamond, M. I.: Glucocorticoid modulation of androgen receptor nuclear aggregation and cellular toxicity is associated with distinct forms of soluble expanded polyglutamine protein. *Hum. Molec. Genet.* 10: 3063–3074, 2001.129. Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.; Landen, M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Polymorphisms of the androgen receptor gene and the estrogen receptor beta gene are associated with androgen levels in women. *J. Clin. Endocr. Metab.* 86:2562–2568, 2001.130. Wieacker, P.; Breckwoldt, M.; Gal, A.: Testicular feminization: diagnosis and search for closely linked restriction fragment length polymorphism. *Dis. Markers* 3: 213–218, 1985.131. Wieacker, P.; Griffin, J. E.; Wienker, T.; Lopez, J. M.; Wilson, J. D.; Breckwoldt, M.: Linkage analysis with RFLPs in families with androgen resistance syndromes: evidence for close linkage between the androgen receptor locus and the DXS1 segment. *Hum. Genet.* 76:248–252, 1987.132. Wilson, C. M.; McPhaul, M. J.: A and B forms of the androgen receptor are present in human genital skin fibroblasts. *Proc. Nat. Acad. Sci.* 91: 1234–1238,

1994.133. Wilson, J. D.: The promiscuous receptor: prostate cancer comes of age. (Editorial) *New Eng. J. Med.* 332: 1440–1441, 1995.134. Wilson, J. D.; Carlson, B. R.; Weaver, D. D.; Kovacs, W. J.; Griffin, J. E.: Endocrine and genetic characterization of cousins with male pseudohermaphroditism: evidence that the Lubs phenotype can result from a mutation that alters the structure of the androgen receptor. *Clin. Genet.* 26: 363–370, 1984.135. Wooster, R.; Mangion, J.; Eeles, R.; Smith, S.; Dowsett, M.; Averill, D.; Barrett-Lee, P.; Easton, D. F.; Ponder, B. A. J.; Stratton, M. R.: A germline mutation in the androgen receptor gene in two brothers with breast cancer and Reifenshtein syndrome. *Nature Genet.* 2: 132–134, 1992.136. Zhang, L.; Leeflang, E. P.; Yu, J.; Arnheim, N.: Studying human mutations by sperm typing: instability of CAG trinucleotide repeats in the human androgen receptor gene. *Nature Genet.* 7: 531–535, 1994.137. Zhu, Y.-S.; Cai, L.-Q.; Cordero, J. J.; Canovatchel, W. J.; Katz, M. D.; Imperato-McGinley, J.: A novel mutation in the CAG triplet region of exon 1 of androgen receptor gene causes complete androgen insensitivity syndrome in a large kindred. *J. Clin. Endocr. Metab.* 84: 1590–1594, 1999.138. Zoppi, S.; Wilson, C. M.; Harbison, M. D.; Griffin, J. E.; Wilson, J. D.;



McPhaul, M. J.; Marcelli, M.: Complete testicular feminization caused by an amino-terminal truncation of the androgen receptor with downstream initiation. *J. Clin. Invest.* 91: 1105–1112, 1993.

[75441] 13771. Konig, A.; Happle, R.; Fink-Puches, R.; Soyer, H. P.; Bornholdt, D.; Engel, H.; Grzeschik, K.-H.: A novel missense mutation of NSDHL in an unusual case of CHILD syndrome showing bilateral, almost symmetric involvement. *J. Am. Acad. Derm.* 46: 594–596, 2002.

[75442] 13772. Labit-Le Bouteiller, C.; Jamme, M. F.; David, M.; Silve, S.; Lanau, C.; Dhers, C.; Picard, C.; Rahier, A.; Taton, M.; Loison, G.; Caput, D.; Ferrara, P.; Lupker, J.: Antiproliferative effects of SR31747A in animal cell lines are mediated by inhibition of cholesterol biosynthesis at the sterol isomerase step. *Europ. J. Biochem.* 256: 342–349, 1998.

[75443] 13773. Liu, X. Y.; Dangel, A. W.; Kelley, R. I.; Zhao, W.; Denny, P.; Botcherby, M.; Cattanach, B.; Peters, J.; Hunsicker, P. R.; Mallon, A.-M.; Strivens, M. A.; Bate, R.; Miller, W.; Rhodes, M.; Brown, S. D. M.; Herman, G. E.: The gene mutated in bare patches and striated mice encodes a novel 3-beta-hydroxysteroid dehydrogenase. *Nature Genet.* 22: 182–187, 1999.

[75444] 13774. Schindelhauer, D.; Hellebrand, H.; Grimm, L.;

Bader, I.; Meitinger, T.; Wehnert, M.; Ross, M.; Meindl, A.: Long-range map of a 3.5-Mb region in Xp11.23-22 with a sequence-ready map from a 1.1-Mb gene-rich interval. *Genome Res.* 6: 1056-1069, 1996.

- [75445] 13775. Silve, S.; Dupuy, P. H.; Labit-Lebouteiller, C.; Kaghad, M.; Chalon, P.; Rahier, A.; Taton, M.; Lupker, J.; Shire, D.; Loison, G.: Emopamil-binding protein, a mammalian protein that binds a series of structurally diverse neuroprotective agents, exhibits delta(8)-delta(7) sterol isomerase activity in yeast. *J. Biol. Chem.* 271: 22434-22440, 1996.
- [75446] 13776. Traupe, H.; Muller, D.; Atherton, D.; Kalter, D. C.; Cremers, F. P. M.; van Oost, B. A.; Ropers, H.-H.: Exclusion mapping of the X-linked dominant chondrodysplasia punctata/ichthyosis/cataract/short stature (Happle) syndrome: possible involvement of an unstable pre-mutation. *Hum. Genet.* 89: 659-665, 1992.
- [75447] 13777. Jin, H.; Gardner, R. J.; Viswesvariah, R.; Muntoni, F.; Roberts, R. G.: Two novel members of the interleukin-1 receptor gene family, one deleted in Xp22.1-Xp21.3 mental retardation. *Europ. J. Hum. Genet.* 8: 87-94, 2000.
- [75448] 13778. Biamonti, G.; Ruggiu, M.; Saccone, S.; Della Valle, G.; Riva, S.: Two homologous genes, originated by dupli-

cation, encode the humanhnRNP proteins A2 and A1. *Nucleic Acids Res.* 22: 1996–2002, 1994.

- [75449] 13779.Stoddart, K. L.; Jermak, C.; Nagaraja, R.; Schlessinger, D.; Bech-Hansen, N. T.: Physical map covering a 2 Mb region in human Xp11.3 distalto DX6849. *Gene* 227: 111–116, 1999.
- [75450] 13780.Swanson, D. A.; Freund, C. L.; Ploder, L.; McInnes, R. R.; Valle, D.: A ubiquitin C-terminal hydrolase gene on the proximal short arm of the X chromosome: implications for X-linked retinal disorders. *Hum.Molec. Genet.* 5: 533–538, 1996.
- [75451] 13781.Narayanan, V.; Olinsky, S.; Dahle, E.; Naidu, S.; Zoghbi, H. Y.: Mutation analysis of the M6b gene in patients with Rett syndrome. *Am.J. Med. Genet.* 78: 165–168, 1998.
- [75452] 13782.Olinsky, S.; Loop, B. T.; DeKosky, A.; Ripepi, B.; Weng, W.; Cummins, J.; Wenger, S. L.; Yan, Y.; Lagenaur, C.; Narayanan, V.: Chromosomal mapping of the human M6 genes. *Genomics* 33: 532–536, 1996.
- [75453] 13783.Yan, Y.; Lagenaur, C.; Narayanan, V.: Molecular cloning of M6: identification of a PLP/DM20 gene family. *Neuron* 11: 423–431, 1993.
- [75454] 13784.Ciccodicola, A.; D'Esposito, M.; Esposito, T.; Gian-

francesco, F.;Migliaccio, C.; Miano, M. G.; Matarazzo, M. R.; Vacca, M.; Franze,A.; Cuccurese, M.; Cocchia, M.; Curci, A.: Differentially regulated and evolved genes in the fully sequenced Xq/Yq pseudoautosomal region. Hum.Molec. Genet. 9: 395–401, 2000.

[75455] 13785.D'Esposito, M.; Ciccodicola, A.; Gianfrancesco, F.; Esposito, T.;Flagiello, L.; Mazzarella, R.; Schlessinger, D.; D'Urso, M.: A synaptobrevin–like gene in the Xq28 pseudoautosomal region undergoes X inactivation. Nature–Genet. 13: 227–229, 1996.

[75456] 13786.Muller, D. J.; Schulze, T. G.; Jahnes, E.; Cichon, S.; Krauss,H.; Kesper, K.; Held, T.; Maier, W.; Propping, P.; Nothen, M. M.;Rietschel, M.: Association between a polymorphism in the pseudoautosomalX–linked gene SYBL1 and bipolar affective disorder. Am. J. Med. Genet.(Neuropsychiat. Genet.) 114: 74–78, 2002.

[75457] 13787.Saito, T.; Parsia, S.; Papolos, D. F.; Lachman, H. M.: Analysis of the pseudoautosomal X–linked gene SYBL1 in bipolar affective disorder:description of a new candidate allele for psychiatric disorders. Am.J. Med. Genet. (Neuropsychiat. Genet.) 96: 317–323, 2000.

[75458] 13788.Schaefer, L.; Ballabio, A.; Zoghbi, H. Y.: Cloning and characterization of a putative human holocytochrome

c-type synthetase gene (HCCS) isolated from the critical region for microphthalmia with linear skin defects (MLS). Genomics 34: 166–172, 1996.

[75459] 13789. Van den Veyver, I. B.; Subramanian, S.; Zoghbi, H. Y.: Genomic structure of a human holocytochrome c-type synthetase gene in Xp22.3 and mutation analysis in patients with Rett syndrome. Am. J. Med. Genet. 78: 179–181, 1998.

[75460] 13790. Andersson, M.; Page, D. C.; Pettay, D.; Subrt, I.; Turleau, C.; de Grouchy, J.; de la Chapelle, A.: Y; autosome translocations and mosaicism in the aetiology of 45,X maleness: assignment of fertility factor to distal Yq11. Hum. Genet. 79: 2–7, 1988.

[75461] 13791. Chandley, A. C.; Gosden, J. R.; Hargreave, T. B.; Spowart, G.; Speed, R. M.; McBeath, S.: Deleted Yq in the sterile son of a man with a satellited Y chromosome (Yqs). J. Med. Genet. 26: 145–153, 1989.

[75462] 13792. Franco, B.; Meroni, G.; Parenti, G.; Levilliers, J.; Bernard, L.; Gebbia, M.; Cox, L.; Maroteaux, P.; Sheffield, L.; Rappold, G. A.; Andria, G.; Petit, C.; Ballabio, A.: A cluster of sulfatase genes on Xp22.3: mutations in chondrodysplasia punctata (CDPX) and implications for warfarin embryopathy. Cell 81: 1–20, 1995.

- [75463] 13793.Meroni, G.; Franco, B.; Archidiacono, N.; Messali, S.; Andolfi,G.; Rocchi, M.; Ballabio, A.: Characterization of a cluster of sulfatasegenes on Xp22.3 suggests gene duplications in an ancestral pseudoautosomalregion. Hum. Molec. Genet. 5: 423–431, 1996.
- [75464] 13794.Lee, C. C.; Yazdani, A.; Wehnert, M.; Zhao, Z. Y.; Lindsay, E.A.; Bailey, J.; Coolbaugh, M. I.; Couch, L.; Xiong, M.; Chinault,A. C.; Baldini, A.; Caskey, C. T.: Isolation of chromosome–specificgenes by reciprocal probing of arrayed cDNA and cosmid libraries. Hum.Molec. Genet. 4: 1373–1380, 1995.
- [75465] 13795.Patzak, D.; Zhuchenko, O.; Lee, C.–C.; Wehnert, M.: Identification,mapping, and genomic structure of a novel X–chromosomal human gene(SMPX) encoding a small muscular protein. Hum. Genet. 105: 506–512,1999.
- [75466] 13796.Lahn, B. T.; Page, D. C.: A human sex–chromosomal gene familyexpressed in male germ cells and encoding variably charged proteins. Hum.Molec. Genet. 9: 311–319, 2000.
- [75467] 13797.Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayasi, Y.;Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura, N.: Predictionof the coding sequences of unidentified human genes. VI. The codingsequences of

80 new genes (KIAA0201–KIAA0280) deduced by analysis of cDNA clones from cell line KG–1 and brain. DNA Res. 3: 321–329, 1996. Note: Supplement: DNA Res. 3: 341–354, 1996.

- [75468] 13798. Numata, M.; Petrecca, K.; Lake, N.; Orlowski, J.: Identification of a mitochondrial Na<sup>+</sup>/H<sup>+</sup> exchanger. J. Biol. Chem. 273: 6951–6959, 1998.
- [75469] 13799. Buchner, G.; Orfanelli, U.; Quaderi, N.; Bassi, M. T.; Andolfi, G.; Ballabio, A.; Franco, B.: Identification of a new EGF-repeat-containing gene from human Xp22: a candidate for developmental disorders. Genomics 65:16–23, 2000.
- [75470] 13800. Yeung, G.; Mulero, J. J.; Berntsen, R. P.; Loeb, D. B.; Drmanac, R.; Ford, J. E.: Cloning of a novel epidermal growth factor repeat-containing gene EGFL6: expressed in tumor and fetal tissues. Genomics 62:304–307, 1999.
- [75471] 13801. Asao, H.; Okuyama, C.; Kumaki, S.; Ishii, N.; Tsuchiya, S.; Foster, D.; Sugamura, K.: Cutting edge: the common gamma-chain is an indispensable subunit of the IL–21 receptor complex. J. Immun. 167: 1–5, 2001.
- [75472] 13802. Lankes, W.; Griesmacher, A.; Grunwald, J.; Schwartz-Albiez, R.; Keller, R.: A heparin-binding protein involved in inhibition of smooth-muscle cell proliferation.

Biochem. J. 251: 831–842, 1988.

- [75473] 13803.Lankes, W. T.; Furthmayr, H.: Moesin: a member of the protein4.1–talín–ezrin family of proteins. Proc. Nat. Acad. Sci. 88: 8297–8301,1991.
- [75474] 13804.Shcherbina, A.; Bretscher, A.; Rosen, F. S.; Kenney, D. M.; Remold–O'Donnell,E.: The cytoskeletal linker protein moesin: decreased levels in Wiskott–Aldrichsyndrome platelets and identification of a cleavage pathway in normalplatelets. Brit. J. Haemat. 106: 216–223, 1999.
- [75475] 13805.Wilgenbus, K. K.; Hsieh, C.–L.; Lankes, W. T.; Milatovich, A.;Francke, U.; Furthmayr, H.: Structure and localization on the X chromosomeof the gene coding for the human filopodial protein moesin (MSN). Genomics 19:326–333, 1994.
- [75476] 13806.Deng, L.; Wang, C.; Spencer, E.; Yang, L.; Braun, A.; You, J.;Slaughter, C.; Pickart, C.; Chen, Z. J.: Activation of the I–kappa–Bcomplex by TRAF6 requires a dimeric ubiquitin–conjugating enzyme complexand a unique polyubiquitin chain. Cell 103: 351–361, 2000.
- [75477] 13807.Wang, C.; Deng, L.; Hong, M.; Akkaraju, G. R.; Inoue, J.; Chen,Z. J.: TAK1 is a ubiquitin–dependent kinase of MKK and IKK. Nature 412:346–351, 2001.
- [75478] 13808.Wong, B. R.; Besser, D.; Kim, N.; Arron, J. R.; Volo–



godskaia, M.; Hanafusa, H.; Choi, Y.: TRANCE, a TNF family member, activates Akt/PKB through a signaling complex involving TRAF6 and c-Src. *Molec. Cell* 4:1041–1049, 1999.

[75479] 13809. Nakano, H.; Oshima, H.; Chung, W.; Williams–Abbott, L.; Ware, C.F.; Yagita, H.; Okumura, K.: TRAF5, an activator of NF–kappaB and putative signal transducer for the lymphotoxin–beta receptor. *J. Biol. Chem.* 271: 14661–14664, 1996.

[75480] 13810. Nakano, H.; Sakon, S.; Koseki, H.; Takemori, T.; Tada, K.; Matsumoto, M.; Munechika, E.; Sakai, T.; Shirasawa, T.; Akiba, H.; Kobata, T.; Santee, S. M.; Ware, C. F.; Renner, P. D.; Taniguchi, M.; Yagita, H.; Okumura, K.: Targeted disruption of Traf5 gene causes defects in CD40– and CD27–mediated lymphocyte activation. *Proc. Nat. Acad. Sci.* 96: 9803–9808, 1999.

[75481] 13811. Nakano, H.; Shindo, M.; Yamada, K.; Yoshida, M. C.; Santee, S.M.; Ware, C. F.; Jenkins, N. A.; Gilbert, D. J.; Yagita, H.; Copeland, N. G.; Okumura, K.: Human TNF receptor–associated factor 5 (TRAF5): cDNA cloning, expression and assignment of the TRAF5 gene to chromosome 1q32. *Genomics* 42: 26–32, 1997.

[75482] 13812. Yost, C.; Farr, G. H., III; Pierce, S. B.; Ferkey, D. M.;

Chen, M. M.; Kimelman, D.: GBP, an inhibitor of GSK-3, is implicated in *Xenopus* development and oncogenesis. *Cell* 93: 1031–1041, 1998.

[75483] 13813. De Baere, E.; Speleman, F.; Van Roy, N.; De Paepe, A.; Messiaen, L.: Assignment of SHOX2 (alias OG12X and SHOT) to human chromosome bands 3q25–q26.1 by in situ hybridization. *Cytogenet. Cell Genet.* 82:228–229, 1998.

[75484] 13814. Fernandez-Valle, C.; Tang, Y.; Ricard, J.; Rodenas-Ruano, A.; Taylor, A.; Hackler, E.; Biggerstaff, J.; Iacovelli, J.: Paxillin binds schwannomin and regulates its density-dependent localization and effect on cell morphology. *Nature Genet.* 31: 354–362, 2002.

[75485] 13815. Glenney, J. R., Jr.; Zokas, L.: Novel tyrosine kinase substrates from Rous sarcoma virus-transformed cells are present in the membraneskeleton. *J. Cell Biol.* 108: 2401–2408, 1989.

[75486] 13816. Mazaki, Y.; Hashimoto, S.; Sabe, H.: Monocyte cells and cancer cells express novel paxillin isoforms with different binding properties to focal adhesion proteins. *J. Biol. Chem.* 272: 7437–7444, 1997.

[75487] 13817. Salgia, R.; Li, J.-L.; Lo, S. H.; Brunkhorst, B.; Kansas, G. S.; Sobhany, E. S.; Sun, Y.; Pisick, E.; Hallek, M.; Ernst,

T.; Tantravahi,R.; Chen, L. B.; Griffin, J. D.: Molecular cloning of human paxillin,a focal adhesion protein phosphorylated by P210(BCR/ABL). J. Biol.Chem. 270: 5039–5047, 1995.

- [75488] 13818.Turner, C. E.; Glenney, J. R., Jr.; Burridge, K.: Paxillin: anew vinculin–binding protein present in focal adhesions. J. CellBiol. 111: 1059–1068, 1990.
- [75489] 13819.Nguyen, V. T.; Kiss, T.; Michels, A. A.; Bensaude, O.: 7SK smallnuclear RNA binds to and inhibits the activity of CDK9/cyclin T complexes. Nature 414:322–325, 2001.
- [75490] 13820.Peng, J.; Zhu, Y.; Milton, J. T.; Price, D. H.: Identificationof multiple cyclin subunits of human P–TEFb. Genes Dev. 12: 755–762,1998.
- [75491] 13821.Yang, Z.; Zhu, Q.; Luo, K.; Zhou, Q.: The 7SK small nuclear RNAinhibits the CDK9/cyclin T1 kinase to control transcription. Nature 414:317–322, 2001.
- [75492] 13822.Adachi, H.; Tsujimoto, M.; Hattori, M.; Arai, H.; Inoue, K.: Differentialtissue distribution of the beta– and gamma–subunits of human cytosolicplatelet–activating factor acetylhydrolase (isoform 1). Biochem.Biophys. Res. Commun. 233: 10–13, 1997.
- [75493] 13823.Moro, F.; Arrigo, G.; Fogli, A.; Bernard, L.; Carozzo, R.: Thebeta and gamma subunits of the human

platelet-activating factor acetylhydrolase isoform Ib (PAFAH1B2 and PAFAH1B3) map to chromosome 11q23 and 19q13.1, respectively. *Genomics* 51: 157–159, 1998.

- [75494] 13824. Daigo, Y.; Isomura, M.; Nishiwaki, T.; Tamari, M.; Ishikawa, S.; Kai, M.; Murata, Y.; Takeuchi, K.; Yamane, Y.; Hayashi, R.; Minami, M.; Fujino, M. A.; Hojo, Y.; Uchiyama, I.; Takagi, T.; Nakamura, Y.: Characterization of a 1200-kb genomic segment of chromosome 3p22–p21.3. *DNAREs.* 6: 37–44, 1999.
- [75495] 13825. Erlich, R.; Gleeson, P. A.; Campbell, P.; Dietzsch, E.; Toh, B.–H.: Molecular characterization of trans-Golgi p230: a human peripheral membrane protein encoded by a gene on chromosome 6p12–22 contains extensive coiled-coil alpha-helical domains and a granin motif. *J. Biol. Chem.* 271: 8328–8337, 1996.
- [75496] 13826. Barr, F. A.; Nakamura, N.; Warren, G.: Mapping the interaction between GRASP65 and GM130, components of a protein complex involved in the stacking of Golgi cisternae. *EMBO J.* 17: 3258–3268, 1998.
- [75497] 13827. Boguski, M. S.; Lowe, T. M. J.; Tolstocher, C. M.: dbEST: database for 'expressed sequence tags.' (Letter) *Nature Genet.* 4: 332–333, 1993.

- [75498] 13828.Izon, D. J.; Aster, J. C.; He, Y.; Weng, A.; Karnell, F. G.; Patriub,V.; Xu, L.; Bakkour, S.; Rodriguez, C.; Allman, D.; Pear, W. S.:Deltex1 redirects lymphoid progenitors to the B cell lineage by antagonizingNotch1. *Immunity* 16: 231–243, 2002.
- [75499] 13829.Lee, L.; Dowhanick–Morrissette, J.; Katz, A.; Jukofsky, L.; Krantz,I. D.: Chromosomal localization, genomic characterization, and mappingto the Noonan syndrome critical region of the human Deltex (DTX1)gene. *Hum. Genet.* 107: 577–581, 2000.
- [75500] 13830.Matsuno, K.; Eastman, D.; Mitsiades, T.; Quinn, A. M.; Carcanciu,M. L.; Ordentlich, P.; Kadesch, T.; Artavanis–Tsakonas, S.: Humandeltex is a conserved regulator of Notch signalling. *Nature Genet.* 19:74–78, 1998.
- [75501] 13831.Chen, J. J.; Reid, C. E.; Band, V.; Androphy, E. J.: Interactionof papillomavirus E6 oncoproteins with a putative calcium–bindingprotein. *Science* 269: 529–531, 1995.
- [75502] 13832.Imai, T.; Matsuda, K.; Shimojima, T.; Hashimoto, T.; Masuhiro,Y.; Kitamoto, T.; Sugita, A.; Suzuki, K.; Matsumoto, H.; Masushige,S.; Nogi, Y.; Muramatsu, M.; Handa, H.; Kato, S.: ERC–55, a bindingprotein for the papilloma virus E6 oncoprotein, specifically interactswith vitamin D receptor among nuclear receptors. *Biochem. Bio-*

phys.Res. Commun. 233: 765–769, 1997.

[75503] 13833.Wang, J. Y.; Zhen, D. K.; Bianchi, D. W.; Androphy, E. J.; Chen,J. J.: Assignment of the gene for ERC–55 (RCN2) to human chromosomeband 15q22.33–q24.1 by in situ hybridization. Cytogenet. Cell Genet. 79:60–61, 1997.

[75504] 13834.Weis, K.; Griffiths, G.; Lamond, A. I.: The endoplasmic reticulumcalcium–binding protein of 55 kDa is a novel EF–hand protein retainedin the endoplasmic reticulum by a carboxyl–terminal His–Asp–Glu–Leumotif. J. Biol. Chem. 269: 19142–19150, 1994.

[75505] 13835.Dempsey, C. E.; Sakurai, H.; Sugita, T.; Guesdon, F.: Alternativesplicing and gene structure of the transforming growth factor beta–activatedkinase 1. Biochim. Biophys. Acta 1517: 46–52, 2000.

[75506] 13836.Kondo, M.; Osada, H.; Uchida, K.; Yanagisawa, K.; Masuda, A.; Takagi,K.; Takahashi, T.; Takahashi, T.: Molecular cloning of human TAK1and its mutational analysis in human lung cancer. Int. J. Cancer 75:559–563, 1998.

[75507] 13837.Sakurai, H.; Shigemori, N.; Hasegawa, K.; Sugita, T.: TGF–beta–activatedkinase 1 stimulates NF–kappa–B activation by an NF–kappa–B–inducingkinase–independent mechanism. Biochem. Biophys. Res. Commun.

243:545–549, 1998.

- [75508] 13838.Yamaguchi, K.; Shirakabe, K.; Shibuya, H.; Irie, K.; Oishi, I.;Ueno, N.; Taniguchi, T.; Nishida, E.; Matsumoto, K.: Identificationof a member of the MAPKKK family as a potential mediator of TGF–betasignal transduction. Science 270: 2008–2011, 1995.
- [75509] 13839.Chan, A. M.–L.; Chedid, M.; McGovern, E. S.; Popescu, N. C.; Miki,T.; Aaronson, S. A.: Expression cDNA cloning of a serine kinase transforminggene. Oncogene 8: 1329–1333, 1993.
- [75510] 13840.Frank, K. M.; Sekiguchi, J. M.; Seidl, K. J.; Swat, W.; Rathbun,G. A.; Cheng, H.–L.; Davidson, L.; Kangaloo, L.; Alt, F. W.: Lateembryonic lethality and impaired V(D)J recombination in mice lackingDNA ligase IV. Nature 396: 173–177, 1998.
- [75511] 13841.Frank, K. M.; Sharpless, N. E.; Gao, Y.; Sekiguchi, J. M.; Ferguson,D. O.; Zhu, C.; Manis, J. P.; Horner, J.; DePinho, R. A.; Alt, F.W.: DNA ligase IV deficiency in mice leads to defective neurogenesisand embryonic lethality via the p53 pathway. Molec. Cell 5: 993–1002,2000.
- [75512] 13842.Grawunder, U.; Zimmer, D.; Fugmann, S.; Schwarz, K.; Lieber, M.R.: DNA ligase IV is essential for V(D)J recombination and DNA double–strandbreak repair in human

precursor lymphocytes. *Molec. Cell* 2: 477–484, 1998.

[75513] 13843. O'Driscoll, M.; Cerosaletti, K. M.; Girard, P.-M.; Dai, Y.; Stumm, M.; Kysela, B.; Hirsch, B.; Gennery, A.; Palmer, S. E.; Seidel, J.; Gatti, R. A.; Varon, R.; Oettinger, M. A.; Neitzel, H.; Jeggo, P. A.; Concannon, P.: DNA ligase IV mutations identified in patients exhibiting developmental delay and immunodeficiency. *Molec. Cell* 8: 1175–1185, 2001.

[75514] 13844. Riballo, E.; Critchlow, S. E.; Teo, S.-H.; Doherty, A. J.; Priestley, A.; Broughton, B.; Kysela, B.; Beamish, H.; Plowman, N.; Arlett, C. F.; Lehmann, A. R.; Jackson, S. P.; Jeggo, P. A.: Identification of a defect in DNA ligase IV in a radiosensitive leukaemia patient. *Curr. Biol.* 9: 699–702, 1999.

[75515] 13845. Riballo, E.; Doherty, A. J.; Dai, Y.; Stiff, T.; Oettinger, M. A.; Jeggo, P. A.; Kysela, B.: Cellular and biochemical impact of a mutation in DNA ligase IV conferring clinical radiosensitivity. *J. Biol. Chem.* 276: 31124–31132, 2001.

[75516] 13846. Giachino, C.; Lantelme, E.; Lanzetti, L.; Saccone, S.; Della Valle, G.; Migone, N.: A novel SH3-containing human gene family preferentially expressed in the central nervous system. *Genomics* 41: 427–434, 1997.



- [75517] 13847.Furlan, M.; Robles, R.; Galbusera, M.; Remuzzi, G.; Kyrle, P. A.; Brenner, B.; Krause, M.; Scharer, I.; Aumann, V.; Mittler, U.; Solenthaler, M.; Lammle, B.: Von Willebrand factor-cleaving protease in thrombotic thrombocytopenic purpura and the hemolytic-uremic syndrome. *New Eng. J. Med.* 339: 1578–1584, 1998.
- [75518] 13848.Furlan, M.; Robles, R.; Lammle, B.: Partial purification and characterization of a protease from human plasma cleaving von Willebrand factor to fragments produced by in vivo proteolysis. *Blood* 87: 4223–4234, 1996.
- [75519] 13849.Nichols, A. F.; Ong, P.; Linn, S.: Mutations specific to the xeroderma pigmentosum group E Ddb- phenotype. *J. Biol. Chem.* 271: 24317–24320, 1996.
- [75520] 13850.Shisheva, A.; Sudhof, T. C.; Czech, M. P.: Cloning, characterization, and expression of a novel GDP dissociation inhibitor isoform from skeletal muscle. *Molec. Cell. Biol.* 14: 3459–3468, 1994.
- [75521] 13851.Grieff, M.; Whyte, M. P.; Thakker, R. V.; Mazzarella, R.: Sequence analysis of 139 kb in Xp22.1 containing spermine synthase and the 5-prime region of PEX. *Genomics* 44: 227–231, 1997.
- [75522] 13852.Korhonen, V. P.; Halmekyto, M.; Kauppinen, L.; Myohanen, S.; Wahlfors, J.; Keinonen, T.; Hyvonen, T.; Alho-

nen, L.; Eloranta, T.; Janne, J.: Molecular cloning of a cDNA encoding human spermine synthase. *DNACell Biol.* 14: 841–847, 1995.

[75523] 13853.Banfi, S.; Borsani, G.; Bulfone, A.; Ballabio, A.: *Drosophila*–related expressed sequences. *Hum. Molec. Genet.* 6: 1745–1753, 1997.

[75524] 13854.Castrillon, D. H.; Wasserman, S. A.: Diaphanous is required for cytokinesis in *Drosophila* and shares domains of similarity with the products of the limb deformity gene. *Development* 120: 3367–3377, 1994.

[75525] 13855.Philippe, C.; Cremers, F. P. M.; Chery, M.; Bach, I.; Abbadi, N.; Ropers, H. H.; Gilgenkrantz, S.: Physical mapping of DNA markers in the q13–q22 region of the human X chromosome. *Genomics* 17: 147–152, 1993.

[75526] 13856.Bione, S.; Sala, C.; Manzini, C.; Arrigo, G.; Zuffardi, O.; Banfi, S.; Borsani, G.; Jonveaux, P.; Philippe, C.; Zucotti, M.; Ballabio, A.; Toniolo, D.: A human homologue of the *Drosophila melanogaster* diaphanous gene is disrupted in a patient with premature ovarian failure: evidence for conserved function in oogenesis and implications for human sterility. *Am. J. Hum. Genet.* 62: 533–541, 1998.

[75527] 13857.Sala, C.; Arrigo, G.; Torri, G.; Martinazzi, F.; Riva, P.; Larizza, L.; Philippe, C.; Jonveaux, P.; Sloan, F.; Labella,

T.; Toniolo, D.: Eleven X chromosome breakpoints associated with premature ovarian failure (POF) map to a 15-Mb YAC contig spanning Xq21. *Genomics* 40:123–131, 1997.

[75528] 13858. Hanaoka, F.; Tandai, M.; Miyazawa, H.; Murakami, Y.; Hori, T.; Yamada, M.: Assignment of human DNA polymerase alpha gene (POLA) to the X chromosome. (Abstract) *Cytogenet. Cell Genet.* 40: 647 only, 1985.

[75529] 13859. Miyazawa, H.; Tandai, M.; Hanaoka, F.; Yamada, M.; Hori, T.; Shimizu, K.; Sekiguchi, M.: Identification of a DNA segment containing the human DNA polymerase alpha gene. *Biochem. Biophys. Res. Commun.* 139:637–643, 1986.

[75530] 13860. Wang, T. S.-F.; Pearson, B. E.; Suomalainen, H. A.; Mohandas, T.; Shapiro, L. J.; Schroder, J.; Korn, D.: Assignment of the gene for human DNA polymerase alpha to the X chromosome. *Proc. Nat. Acad. Sci.* 82: 5270–5274, 1985.

[75531] 13861. Watson, J. M.; Spencer, J. A.; Riggs, A. D.; Graves, J. A. M.: Sex chromosome evolution: platypus gene mapping suggests that part of the human X chromosome was originally autosomal. *Proc. Nat. Acad. Sci.* 88: 11256–11260, 1991.

[75532] 13862. Lachner, M.; O'Carroll, D.; Rea, S.; Mechtler, K.;

Jenuwein, T.: Methylation of histone H3 lysine 9 creates a binding site for HP1 proteins. *Nature* 410: 116–120, 2001.

- [75533] 13863. Melcher, M.; Schmid, M.; Aagaard, L.; Selenko, P.; Laible, G.; Jenuwein, T.: Structure–function analysis of SUV39H1 reveals a dominant role in heterochromatin organization, chromosome segregation, and mitotic progression. *Molec. Cell Biol.* 20: 3728–3841, 2000.
- [75534] 13864. Nielsen, S. J.; Schneider, R.; Bauer, U.–M.; Bannister, A. J.; Morrison, A.; O'Carroll, D.; Firestein, R.; Cleary, M.; Jenuwein, T.; Herrera, R. E.; Kouzarides, T.: Rb targets histone H3 methylation and HP1 to promoters. *Nature* 412: 561–565, 2001.
- [75535] 13865. Peters, A. H. F. M.; O'Carroll, D.; Scherthan, H.; Mechtler, K.; Sauer, S.; Schofer, C.; Weipoltshammer, K.; Pagani, M.; Lachner, M.; Kohlmaier, A.; Opravil, S.; Doyle, M.; Sibilia, M.; Jenuwein, T.: Loss of the Suv39h histone methyltransferases impairs mammalian heterochromatin and genome stability. *Cell* 107: 323–337, 2001.
- [75536] 13866. Rea, S.; Eisenhaber, F.; O'Carroll, D.; Strahl, B. D.; Sun, Z.–W.; Schmid, M.; Opravil, S.; Mechtler, K.; Ponting, C. P.; Allis, C. D.; Jenuwein, T.: Regulation of chromatin structure by site–specific histone H3 methyltransferases. *Nature* 406: 593–599, 2000.

- [75537] 13867.Scott, A. F.: Personal Communication. Baltimore, Md. 8/7/2000.
- [75538] 13868.Haltiwanger, R. S.; Blomberg, M. A.; Hart, G. W.: Glycosylationof nuclear and cytoplasmic proteins: purification and characterizationof a uridine diphospho-N-acetylglucosamine:polypeptide beta-N-acetylglucosaminyltransferase. J.Biol. Chem. 267: 9005-9013, 1992.
- [75539] 13869.Kreppel, L. K.; Blomberg, M. A.; Hart, G. W.: Dynamic glycosylationof nuclear and cytosolic proteins: cloning and characterization ofa unique O-GlcNAc transferase with multiple tetratricopeptide repeats. J.Biol. Chem. 272: 9308-9315, 1997.
- [75540] 13870.Lubas, W. A.; Frank, D. W.; Krause, M.; Hanover, J. A.: O-linkedGlcNAc transferase is a conserved nucleocytoplasmic protein containingtetratricopeptide repeats. J. Biol. Chem. 272: 9316-9324, 1997.
- [75541] 13871.Shafi, R.; Iyer, S. P. N.; Ellies, L. G.; O'Donnell, N.; Marek,K. W.; Chui, D.; Hart, G. W.; Marth, J. D.: The O-GlcNAc transferasegene resides on the X chromosome and is essential for embryonic stemcell viability and mouse ontogeny. Proc. Nat. Acad. Sci. 97: 5735-5739,2000.
- [75542] 13872.Yang, X.; Zhang, F.; Kudlow, J. E.: Recruitment of

O-GlcNAc transferase to promoters by corepressor mSin3A: coupling protein O-GlcNAcylation to transcriptional repression. *Cell* 110: 69–80, 2002.

- [75543] 13873. Furuta, S.; Kobayashi, A.; Miyazawa, S.; Hashimoto, T.: Cloning and expression of cDNA for a newly identified isozyme of bovine liver 3-hydroxyacyl-CoA dehydrogenase and its import into mitochondria. *Biochim. Biophys. Acta* 1350: 317–324, 1997.
- [75544] 13874. Hansis, C.; Jahner, D.; Spiess, A.-N.; Boettcher, K.; Ivell, R.: The gene for the Alzheimer-associated beta-amyloid-binding protein (ERAB) is differentially expressed in the testicular Leydig cells of the azoospermic by w/w(v) mouse. *Europ. J. Biochem.* 258: 53–60, 1998.
- [75545] 13875. He, X.-Y.; Schulz, H.; Yang, S.-Y.: A human brain L-3-hydroxyacyl-coenzyme A dehydrogenase is identical to an amyloid beta-peptide-binding protein involved in Alzheimer's disease. *J. Biol. Chem.* 273: 10741–10746, 1998.
- [75546] 13876. Yan, S. D.; Fu, J.; Soto, C.; Chen, X.; Zhu, H.; Al-Mohanna, F.; Collison, K.; Zhu, A.; Stern, E.; Saido, T.; Tohyama, M.; Ogawa, S.; Roher, A.; Stern, D.: An intracellular protein that binds amyloid-beta peptide and mediates neurotoxicity in Alzheimer's disease. *Nature*

389:689–695, 1997.

- [75547] 13877. Chamberlin, M. E.; Ubagai, T.; Mudd, S. H.; Wilson, W. G.; Leonard, J. V.; Chou, J. Y.: Demyelination of the brain is associated with methionine adenosyltransferase I/III deficiency. *J. Clin. Invest.* 98:1021–1027, 1996.
- [75548] 13878. Nishiyama, K.; Funai, T.; Katafuchi, R.; Hattori, F.; Onoyama, K.; Ichiyama, A.: Primary hyperoxaluria type I due to a point mutation of T to C in the coding region of the serine:pyruvate aminotransferase gene. *Biochem. Biophys. Res. Commun.* 176: 1093–1099, 1991.
- [75549] 13879. Pirulli, D.; Puzzer, D.; Ferri, L.; Crovella, S.; Amoroso, A.; Ferrettini, C.; Marangella, M.; Mazzola, G.; Florian, F.: Molecular analysis of hyperoxaluria type 1 in Italian patients reveals eight new mutations in the alanine:glyoxylate aminotransferase gene. *Hum. Genet.* 104: 523–525, 1999.
- [75550] 13880. Watts, R. W. E.; Calne, R. Y.; Rolles, K.; Danpure, C. J.; Morgan, S. H.; Mansell, M. A.; Williams, R.; Purkiss, P.: Successful treatment of primary hyperoxaluria type I by combined hepatic and renal transplantation. *Lancet* II:474–475, 1987.
- [75551] 13881. Cramer, S. D.; Ferree, P. M.; Lin, K.; Milliner, D. S.; Holmes, R. P.: The gene encoding hydroxypyruvate reduc-

tase (GRHPR) is mutated in patients with primary hyperoxaluria type II. Hum. Molec. Genet. 8:2063–2069, 1999.

- [75552] 13882. Natt, E.; Kida, K.; Odievre, M.; Di Rocco, M.; Scherer, G.: Point mutations in the tyrosine aminotransferase gene in tyrosinemia type II. Proc. Nat. Acad. Sci. 89: 9297–9301, 1992.
- [75553] 13883. Johnson, W. G.: Metabolic interference and the +/- heterozygote: a hypothetical form of simple inheritance which is neither dominant nor recessive. Am. J. Hum. Genet. 32: 374–386, 1980.
- [75554] 13884. Brenner, V.; Nyakatura, G.; Rosenthal, A.; Platzer, M.: Genomic organization of two novel genes on human Xq28: compact head to head arrangement of IDH-gamma and TRAP-delta is conserved in rat and mouse. Genomics 44:8–14, 1997.
- [75555] 13885. Achen, M. G.; Jeltsch, M.; Kukk, E.; Makinen, T.; Vitali, A.; Wilks, A. F.; Alitalo, K.; Stacker, S. A.: Vascular endothelial growth factor D (VEGF-D) is a ligand for the tyrosine kinases VEGF receptor 2 (Flk1) and VEGF receptor 3 (Flt4). Proc. Nat. Acad. Sci. 95: 548–553, 1998.
- [75556] 13886. Stacker, S. A.; Caesar, C.; Baldwin, M. E.; Thornton, G. E.; Williams, R. A.; Prevo, R.; Jackson, D. G.; Nishikawa, S.; Kubo, H.; Achen, M. G.: VEGF-D promotes the



metastatic spread of tumor cells via the lymphatics. *Nature Med.* 7: 186–191, 2001.

- [75557] 13887. Yamada, Y.; Nezu, J.; Shimane, M.; Hirata, Y.: Molecular cloning of a novel vascular endothelial growth factor, VEGF-D. *Genomics* 42:483–488, 1997.
- [75558] 13888. Davies, P. A.; Hanna, M. C.; Hales, T. G.; Kirkness, E. F.: Insensitivity to anaesthetic agents conferred by a class of GABA(A) receptor subunit. *Nature* 385:820–823, 1997.
- [75559] 13889. Sinkkonen, S. T.; Hanna, M. C.; Kirkness, E. F.; Korppe, E. R.: GABA-A receptor epsilon and theta subunits display unusual structural variation between species and are enriched in the rat locus ceruleus. *J. Neurosci.* 20: 3588–3595, 2000.
- [75560] 13890. Wilke, K.; Gaul, R.; Klauck, S. M.; Poustka, A.: A gene in human chromosome band Xq28 (GABRE) defines a putative new subunit class of the GABA(A) neurotransmitter receptor. *Genomics* 45: 1–10, 1997.
- [75561] 13891. Gecz, J.; Baker, E.; Donnelly, A.; Ming, J. E.; McDonald-McGinn, D. M.; Spinner, N. B.; Zackai, E. H.; Sutherland, G. R.; Mulley, J. C.: Fibroblast growth factor homologous factor 2 (FHF2): gene structure, expression and mapping to the Borjeson-Forssman-Lehmann syndrome region in

Xq26 delineated by a duplication breakpoint in a BFLS-like patient. Hum.Genet. 104: 56–63, 1999.

- [75562] 13892.Lovec, H.; Hartung, H.; Verdier, A.–S.; Mattei, M.–G.; Birnbaum,D.; Goldfarb, M.; Coulier, F.: Assignment of FGF13 to human chromosomeband Xq21 by in situ hybridization. Cytogenet. Cell Genet. 76: 183–184,1997.
- [75563] 13893.Smallwood, P. M.; Munoz–Sanjuan, I.; Tong, P.; Macke, J. P.; Hendry,S. H. C.; Gilbert, D. J.; Copeland, N. G.; Jenkins, N. A.; Nathans,J.: Fibroblast growth factor (FGF) homologous factors: new membersof the FGF family implicated in nervous system development. Proc.Nat. Acad. Sci. 93: 9850–9857, 1996.
- [75564] 13894.Jones, M. H.; Furlong, R. A.; Burkin, H.; Chalmers, I. J.; Brown,G. M.; Khwaja, O.; Affara, N. A.: The Drosophila developmental genefat facets has a human homologue in Xp11.4 which escapes X–inactivationand has related sequences on Yq11.2. Hum. Molec. Genet. 5: 1695–1701,1996.
- [75565] 13895.Stoffers, D. A.; Zinkin, N. T.; Stanojevic, V.; Clarke, W. L.;Habener, J. F.: Pancreatic agenesis attributable to a single nucleotidedeletion in the human IPF1 gene coding sequence. Nature Genet. 15:106–110, 1997.
- [75566] 13896.Theophilos, M. B.; Cox, D. W.; Mercer, J. F. B.: The

toxic milkmouse is a murine model of Wilson disease.

Hum. Molec. Genet. 5:1619–1624, 1996.

- [75567] 13897.Natt, E.; Westphal, E.–M.; Toth–Fejel, S. E.; Magenis, R. E.; Buist, N. R. M.; Rettenmeier, R.; Scherer, G.: Inherited and de novo deletion of the tyrosine aminotransferase gene locus at 16q22.1–q22.3 in a patient with tyrosinemia type II. Hum. Genet. 77: 352–358, 1987.
- [75568] 13898.Pelet, B.; Antener, I.; Faggioni, R.; Spahr, A.; Gautier, E.: Tyrosinemia without liver or renal damage with plantar and palmar keratosis and keratitis (hypertyrosinemia type II). Helv. Paediat. Acta 34: 177–183, 1979.
- [75569] 13899.Rehak, A.; Selim, M. M.; Yadav, G.: Richner–Hanhart syndrome (tyrosinaemia–II) (report of four cases without ocular involvement). Brit. J. Derm. 104: 469–475, 1981.
- [75570] 13900.Rettenmeier, R.; Natt, E.; Zentgraf, H.; Scherer, G.: Isolation and characterization of the human tyrosine aminotransferase gene. Nucleic Acids Res. 18: 3853–3861, 1990.
- [75571] 13901.Richner, H.: Hornhautaffektion bei Keratoma palmare et plantare hereditarium. Klin. Mbl. Augenheilk. 100: 580–588, 1938.
- [75572] 13902.Schmid, W.; Muller, G.; Schutz, G.; Gluecksohn–

Waelsch, S.: Deletions near the albino locus on chromosome 7 of the mouse affect the level of tyrosine aminotransferase mRNA. *Proc. Nat. Acad. Sci.* 82: 2866–2869, 1985.

[75573] 13903. Tallab, T.M.: Richner–Hanhart syndrome: importance of early diagnosis and early intervention. *J. Am. Acad. Dermatol.* 35: 857–859, 1996.

[75574] 13904. Ventura, G.; Biasini, G.; Petrozzi, M.: Cheratomia palmoplantare dissipatum associato a lesioni corneali in due fratelli. *Boll. Oculist.* 44: 497–510, 1965.

[75575] 13905. Waardenburg, P. J.; Franceschetti, A.; Klein, D.: *Genetics and Ophthalmology*. Springfield, Ill.: Charles C Thomas (pub.) 1: 1961. Pp. 515–517.

[75576] 13906. Westphal, E.–M.; Natt, E.; Grimm, T.; Odievre, M.; Scherer, G.: The human tyrosine aminotransferase gene: characterization of restriction fragment length polymorphisms and haplotype analysis in a family with tyrosinemia type II. *Hum. Genet.* 79: 260–264, 1988.

[75577] 13907. Perheentupa, J.: Autoimmune polyendocrinopathy–candidosis–ectodermal dystrophy (APECED). In: Eriksson, A. W.; Forsius, H. R.; Nevanlinna, H. R.; Workman, P. L.; Norio, R. K.: *Population Structure and Genetic Disorders*. New York: Academic Press (pub.) 1980. Pp. 583–587.

- [75578] 13908.Prader, A.: Personal Communication. Zurich, Switzerland 1972.
- [75579] 13909.Ramsay, C.; Winqvist, O.; Puhakka, L.; Halonen, M.; Moro, A.;Kampe, O.; Eskelin, P.; Pelto-Huikko, M.; Peltonen, L.: Aire deficientmice develop multiple features of APECED phenotype and show alteredimmune response. Hum. Molec. Genet. 11: 397-409, 2002.
- [75580] 13910.Rinderle, C.; Christensen, H.-M.; Schweiger, S.; Lehrach, H.;Yaspo, M.-L.: AIRE encodes a nuclear protein co-localizing with cytoskeletal filaments: altered sub-cellular distribution of mutants lacking PHDzinc fingers. Hum. Molec. Genet. 8: 277-290, 1999.
- [75581] 13911.Rosatelli, M. C.; Meloni, A.; Meloni, A.; Devoto, M.; Cao, A.;Scott, H. S.; Peterson, P.; Heino, M.; Krohn, K. J.; Nagamine, K.;Kudoh, J.; Shimizu, N.; Antonarakis, S. E.: A common mutation inSardinian autoimmune polyendocrinopathy-candidiasis-ectodermal dystrophypatients. Hum. Genet. 103: 428-434, 1998.
- [75582] 13912.Saenger, P.: Primary hypoaldosteronism due to zona glomerulosadefect. (Letter) New Eng. J. Med. 310: 1394 only, 1984.
- [75583] 13913.Saenger, P.; Levine, L. S.; Irvine, W. J.; Gottesdiener, K.; Rauh,W.; Sonino, N.; Chow, D.; New, M. I.: Progressive

adrenal failure in polyglandular autoimmune disease. *J. Clin. Endocr. Metab.* 54:863–868, 1982.

[75584] 13914.Scott, H. S.; Heino, M.; Peterson, P.; Mittaz, L.; Lalioti, M.D.; Betterle, C.; Cohen, A.; Seri, M.; Lerone, M.; Romeo, G.; Collin, P.; Salo, M.; and 9 others: Common mutations in autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy patients of different origins. *Molec. Endocr.* 12: 1112–1119, 1998.

[75585] 13915.Shapiro, M. S.; Zamir, R.; Weiss, E.; Radnay, J.; Shenkman, L.: The polyglandular deficiency syndrome: a new variant in Persian Jews. *J. Endocr. Invest.* 10: 1–7, 1987.

[75586] 13916.Spinner, M. W.; Blizzard, R. M.; Childs, B.: Clinical and genetic heterogeneity in idiopathic Addison's disease and hypoparathyroidism. *J. Clin. Endocr.* 28: 795–804, 1968.

[75587] 13917.Sweetnam, W. P.: Juvenile familial endocrinopathy. *Lancet* I:463–465, 1966.

[75588] 13918.Wang, C.–Y.; Davoodi–Semiromi, A.; Huang, W.; Connor, E.; Shi, J.–D.; She, J.–X.: Characterization of mutations in patients with autoimmune polyglandular syndrome type 1 (APS1). *Hum. Genet.* 103:681–685, 1998.

[75589] 13919.Whitaker, J. A.; Landing, B. H.; Esselborn, V. M.;

Williams, R.R.: Syndrome of familial juvenile hypoadrenocorticism, hypoparathyroidism and superficial moniliasis. J. Clin. Endocr. 16: 1374–1387, 1956.

- [75590] 13920. Wirfalt, A.: Genetic heterogeneity in autoimmune polyglandular failure. Acta Med. Scand. 210: 7–13, 1981.
- [75591] 13921. Zlotogora, J.; Shapiro, M. S.: Polyglandular autoimmune syndrometype I among Iranian Jews. J. Med. Genet. 29: 824–826, 1992.
- [75592] 13922. Hediger, M. A.: New view at C. Nature Med. 8: 445–446, 2002.
- [75593] 13923. Sotiriou, S.; Gispert, S.; Cheng, J.; Wang, Y.; Chen, A.; Hoogstraten-Miller, S.; Miller, G. F.; Kwon, O.; Levine, M.; Guttentag, S. H.; Nussbaum, R. L.: Ascorbic-acid transporter Slc23a1 is essential for vitamin C transport into the brain and for perinatal survival. Nature Med. 8: 514–517, 2002.
- [75594] 13924. Nauseef, W. M.; Brigham, S.; Cogley, M.: Hereditary myeloperoxidase deficiency due to a missense mutation of arginine 569 to tryptophan. J. Biol. Chem. 269: 1212–1216, 1994.
- [75595] 13925. Busard, B. L. S. M.; Renier, W. O.; Gabreels, F. J. M.; Jaspar, H. H. J.; van Haelst, U. J. G.; Slooff, J. L.: Lafora's disease: comparison of inclusion bodies in skin and in

brain. Arch. Neurol. 43:296–299, 1986.

[75596] 13926. Busard, H. L. S. M.; Gabreels–Festen, A. A. W. M.; Renier, W. O.; Gabreels, F. J. M.; Stadhouders, A. M.: Axilla skin biopsy: a reliable test for the diagnosis of Lafora's disease. Ann. Neurol. 21: 599–601, 1987.

[75597] 13927. Fluharty, A. L.; Porter, M. T.; Hirsh, G. A.; Pevida, E.; Kihara, H.: Metachromasia in fibroblasts from a patient with Lafora's disease. (Letter) Lancet II: 109–110, 1970.

[75598] 13928. Ganesh, S.; Agarwala, K. L.; Ueda, K.; Akagi, T.; Shoda, K.; Usui, T.; Hashikawa, T.; Osada, H.; Delgado–Escueta, A. V.; Yamakawa, K.: Laforin, defective in the progressive myoclonus epilepsy of Lafora type, is a dual–specificity phosphatase associated with polyribosomes. Hum. Molec. Genet. 9: 2251–2261, 2000.

[75599] 13929. Gomez–Garre, P.; Sanz, Y.; Rodriguez de Cordoba, S.; Serratosa, J. M.: Mutational spectrum of the EPM2A gene in progressive myoclonic epilepsy of Lafora: high degree of allelic heterogeneity and prevalence of deletions. Europ. J. Hum. Genet. 8: 946–954, 2000.

[75600] 13930. Harriman, D. G. F.; Millar, J. H. D.: Progressive familial myoclonic epilepsy in 3 families: its clinical features and pathological basis. Brain 78:325–349, 1955.

[75601] 13931. Janeway, R.; Ravens, J. R.; Pearce, L. A.; Odor, D. L.;



Suzuki,K.: Progressive myoclonus epilepsy with Lafora inclusion bodies.I. Clinical, genetic, histopathologic and biochemical aspects. Arch.Neurol. 16: 565–582, 1967.

[75602] 13932.Lehesjoki, A.–E.; Koskiniemi, M.; Pandolfo, M.; Antonelli, A.;Kyllerman, M.; Wahlstrom, J.; Nergardh, A.; Burmeister, M.; Sistonen,P.; Norio, R.; de la Chapelle, A.: Linkage studies in progressivemyoclonus epilepsy: Unverricht–Lundborg and Lafora's diseases. Neurology 42:1545–1550, 1992.

[75603] 13933.Maddox, L. O.; Descartes, M.; Collins, J.; Keating, J.; Rosenfeld,S.; Palmer, C.; Carroll, A. J.; Kuzniecky, R.: Identification of a recombination event narrowing the Lafora disease gene region. J.Med. Genet. 34: 590–591, 1997.

[75604] 13934.Amadou, C.; Ribouchon, M. T.; Mattei, M. G.; Jenkins, N. A.; Gilbert,D. J.; Copeland, N. G.; Avoustin, P.; Pontarotti, P.: Localization of new genes and markers to the distal part of the human major histocompatibility complex (MHC) region and comparison with the mouse: new insights into the evolution of mammalian genomes. Genomics 26: 9–20, 1995.

[75605] 13935.Solinas–Toldo, S.; Lengauer, C.; Fries, R.: Comparative genome map of human and cattle. Genomics 27: 489–496, 1995.

- [75606] 13936.Olives, B.; Martial, S.; Mattei, M.-G.; Matassi, G.; Rousselet,G.; Ripoche, P.; Cartron, J.-P.; Bailly, P.: Molecular characterization of a new urea transporter in the human kidney. *FEBS Lett.* 386: 156–160,1996.
- [75607] 13937.Ranade, K.; Wu, K.-W.; Hwu, C.-M.; Ting, C.-T.; Pei, D.; Pesich,R.; Hebert, J.; Chen, Y.-D. I.; Pratt, R.; Olshen, R.; Masaki, K.;Risch, N.; Cox, D. R.; Botstein, D.: Genetic variation in the human urea transporter-2 is associated with variation in blood pressure. *Hum.Molec. Genet.* 10: 2157–2164, 2001.
- [75608] 13938.Ansel, K. M.; Ngo, V. N.; Hyman, P. L.; Luther, S. A.; Forster,R.; Sedgwick, J. D.; Browning, J. L.; Lipp, M.; Cyster, J. G.: A chemokine-driven positive feedback loop organizes lymphoid follicles. *Nature* 406:309–314, 2000.
- [75609] 13939.Galko, M. J.; Tessier-Lavigne, M.: Function of an axonal chemoattractant modulated by metalloprotease activity. *Science* 289: 1365–1367, 2000.
- [75610] 13940.Stein, E.; Tessier-Lavigne, M.: Hierarchical organization of guidance receptors: silencing of netrin attraction by Slit through a Robo/DCC receptor complex. *Science* 291: 1928–1938, 2001.
- [75611] 13941.Stein, E.; Zou, Y.; Poo, M.; Tessier-Lavigne, M.: Binding of DCC by netrin-1 to mediate axon guidance in-

dependent of adenosineA2B receptor activation. Science 291: 1976–1982, 2001.

[75612] 13942.Lubahn, D. B.; Moyer, J. S.; Golding, T. S.; Couse, J. F.; Korach,K. S.; Smithies, O.: Alteration of reproductive function but not prenatal sexual development after insertional disruption of the mouseestrogen receptor gene.

Proc. Nat. Acad. Sci. 90: 11162–11166, 1993.

[75613] 13943.Hanna, I. H.; Dawling, S.; Roodi, N.; Guengerich, F. P.; Parl,F. F.: Cytochrome P450 1B1 (CYP1B1) pharmacogenetics: associationof polymorphisms with functional differences in estrogen hydroxylationactivity. Cancer Res. 60: 3440–3444, 2000.

[75614] 13944.Moore, J. T.; McKee, D. D.; Slentz–Kesler, K.; Moore, L. B.; Jones,S. A.; Horne, E. L.; Su, J.–L.; Kliewer, S. A.; Lehmann, J. M.; Willson,T. M.: Cloning and characterization of human estrogen receptor betaisoforms.

Biochem. Biophys. Res. Commun. 247: 75–78, 1998.

[75615] 13945.Mosselman, S.; Polman, J.; Dijkema, R.: ER–beta: identificationand characterization of a novel human estrogen receptor. FEBS Lett. 392:49–53, 1996.

[75616] 13946.Ogawa, S.; Emi, M.; Shiraki, M.; Hosoi, T.; Ouchi, Y.; Inoue,S.: Association of estrogen receptor beta (ESR2) gene polymorphismwith blood pressure. J. Hum. Genet.

45: 327–330, 2000.

- [75617] 13947.Ogawa, S.; Inoue, S.; Watanabe, T.; Orimo, A.; Hosoi, T.; Ouchi, Y.; Muramatsu, M.: Molecular cloning and characterization of human estrogen receptor beta-cx: a potential inhibitor of estrogen action in human. *Nucleic Acids Res.* 26: 3505–3512, 1998.
- [75618] 13948.Osterlund, M. K.; Gustafsson, J. A.; Keller, E.; Hurd, Y. L.: Estrogen receptor beta (ER-beta) messenger ribonucleic acid (mRNA) expression within the human forebrain: distinct distribution pattern to ER-alpha mRNA. *J. Clin. Endocr. Metab.* 85: 3840–3846, 2000.
- [75619] 13949.Pasquali, D.; Rossi, V.; Esposito, D.; Abbondanza, C.; Puca, G.A.; Bellastella, A.; Sinisi, A. A.: Loss of estrogen receptor beta expression in malignant human prostate cells in primary cultures and in prostate cancer tissues. *J. Clin. Endocr. Metab.* 86: 2051–2055, 2001.
- [75620] 13950.Petersen, D. N.; Tkalec, G. T.; Koza-Taylor, P. H.; Turi, T.G.; Brown, T. A.: Identification of estrogen receptor beta-2, a functional variant of estrogen receptor beta expressed in normal rat tissues. *Endocrinology* 139:1082–1092, 1998.
- [75621] 13951.Rissman, E. F.; Heck, A. L.; Leonard, J. E.; Shupnik, M. A.; Gustafsson, J.-A.: Disruption of estrogen receptor

beta gene impairs spatial learning in female mice. Proc. Nat. Acad. Sci. 99: 3996–4001, 2002.

- [75622] 13952. Rosenkranz, K.; Hinney, A.; Ziegler, A.; Hermann, H.; Fichter, M.; Mayer, H.; Siegfried, W.; Young, J. K.; Remschmidt, H.; Hebebrand, J.: Systematic mutation screening of the estrogen receptor beta gene in probands of different weight extremes: identification of several genetic variants. J. Clin. Endocr. Metab. 83: 4524–4527, 1998.
- [75623] 13953. Speirs, V.; Adams, I. P.; Walton, D. S.; Atkin, S. L.: Identification of wild-type and exon 5 deletion variants of estrogen receptor beta in normal human mammary gland. J. Clin. Endocr. Metab. 85: 1601–1605, 2000.
- [75624] 13954. Tsukamoto, K.; Inoue, S.; Hosoi, T.; Orimo, H.; Emi, M.: Isolation and radiation hybrid mapping of dinucleotide repeat polymorphism at the human estrogen receptor beta locus. J. Hum. Genet. 43: 73–74, 1998.
- [75625] 13955. Weihua, Z.; Saji, S.; Makinen, S.; Cheng, G.; Jensen, E. V.; Warner, M.; Gustafsson, J.-A.: Estrogen receptor (ER)-beta, a modulator of ER-alpha in the uterus. Proc. Nat. Acad. Sci. 97: 5936–5941, 2000.
- [75626] 13956. Westberg, L.; Baghaei, F.; Rosmond, R.; Hellstrand, M.; Landen, M.; Jansson, M.; Holm, G.; Bjorntorp, P.; Eriksson, E.: Polymorphisms of the androgen receptor gene and

the estrogen receptor gene are associated with androgen levels in women. *J. Clin. Endocr. Metab.* 86: 2562–2568, 2001.

- [75627] 13957. Zhu, Y.; Bian, Z.; Lu, P.; Karas, R. H.; Bao, L.; Cox, D.; Hodgins, J.; Shaul, P. W.; Thoren, P.; Smithies, O.; Gustafsson, J.-A.; Mendelsohn, M. E.: Abnormal vascular function and hypertension in mice deficient in estrogen receptor beta. *Science* 295: 505–508, 2002.
- [75628] 13958. DesGroseillers, L.; Lemieux, N.: Localization of a human double-stranded RNA-binding protein gene (STAU) to band 20q13.1 by fluorescence in situ hybridization. *Genomics* 36: 527–529, 1996.
- [75629] 13959. Marion, R. M.; Fortes, P.; Beloso, A.; Dotti, C.; Ortin, J.: A human sequence homologue of stau6 is an RNA-binding protein that is associated with polysomes and localizes to the rough endoplasmic reticulum. *Molec. Cell. Biol.* 19: 2212–2219, 1999.
- [75630] 13960. Wickham, L.; Duchaine, T.; Luo, M.; Nabi, I. R.; DesGroseillers, L.: Mammalian stau6 is a double-stranded-RNA- and tubulin-binding protein which localizes to the rough endoplasmic reticulum. *Molec. Cell. Biol.* 19: 2220–2230, 1999.
- [75631] 13961. Lee, J. E.; Hollenberg, S. M.; Snider, L.; Turner, D.

L.; Lipnick, N.; Weintraub, H.: Conversion of *Xenopus* ectoderm into neurons by NeuroD, a basic helix-loop-helix protein. *Science* 268: 836–844, 1995.

- [75632] 13962. Liu, M.; Pleasure, S. J.; Collins, A. E.; Noebels, J. L.; Naya, F. J.; Tsai, M.-J.; Lowenstein, D. H.: Loss of BETA2/NeuroD leads to malformation of the dentate gyrus and epilepsy. *Proc. Nat. Acad. Sci.* 97: 865–870, 2000.
- [75633] 13963. Malecki, M. T.; Jhala, U. S.; Antonellis, A.; Fields, L.; Doria, A.; Orban, T.; Saad, M.; Warram, J. H.; Montminy, M.; Krolewski, A. S.: Mutations in NEUROD1 are associated with the development of type 2 diabetes mellitus. *Nature Genet.* 23: 323–328, 1999.
- [75634] 13964. Naya, F. J.; Huang, H.-P.; Qiu, Y.; Mutoh, H.; DeMayo, F. J.; Leiter, A. B.; Tsai, M.-J.: Diabetes, defective pancreatic morphogenesis, and abnormal enteroendocrine differentiation in BETA2/neurod-deficient mice. *Genes Dev.* 11: 2323–2334, 1997.
- [75635] 13965. Naya, F. J.; Stellrecht, C. M.; Tsai, M. J.: Tissue-specific regulation of the insulin gene by a novel basic helix-loop-helix transcription factor. *Genes Dev.* 9: 1009–1019, 1995.
- [75636] 13966. Plasilova, M.; Stoilov, I.; Sarfarazi, M.; Kadasi, L.; Ferakova, E.; Ferak, V.: Identification of a single ancestral

CYP1B1 mutation in Slovak Gypsies (Roms) affected with primary congenital glaucoma. *J. Med. Genet.* 36: 290–294, 1999.

[75637] 13967. Schwartzman, M. L.; Balazy, M.; Masferrer, J.; Abraham, N. G.; McGiff, J. C.; Murphy, R. C.:

12(R)-hydroxyicosatetraenoic acid: a cytochrome

P450-dependent arachidonate metabolite that inhibits

Na<sup>+</sup>, K<sup>+</sup>-ATPase in the cornea. *Proc. Nat. Acad. Sci.* 84:

8125–8129, 1987.

[75638] 13968. Stoilov, I.; Akarsu, A. N.; Alozie, I.; Child, A.; Barsoum-Homsy, M.; Turacli, M. E.; Or, M.; Lewis, R. A.;

Ozdemir, N.; Brice, G.; Aktan, S. G.; Chevrette, L.; Coca-

Prados, M.; Sarfarazi, M.: Sequence analysis and homology

modeling suggest that primary congenital glaucoma on

2p21 results from mutations disrupting either the hinge

region or the conserved core structures of cytochrome

P4501B1. *Am. J. Hum. Genet.* 62: 573–584, 1998.

[75639] 13969. Stoilov, I.; Akarsu, A. N.; Sarfarazi, M.: Identifica-

tion of three different truncating mutations in cytochrome

P4501B1 (CYP1B1) as the principal cause of primary con-

genital glaucoma (buphthalmos) in families linked to the

GLC3A locus on chromosome 2p21. *Hum. Molec. Genet.*

6:641–647, 1997.



- [75640] 13970.Sutter, T. R.; Tang, Y. M.; Hayes, C. L.; Wo, Y.-Y. P.; Jabs, E.W.; Li, X.; Yin, H.; Cody, C. W.; Greenlee, W. F.: Complete cDNAsequence of a human dioxin-inducible mRNA identifies a new gene subfamilyof cytochrome P450 that maps to chromosome 2. J. Biol. Chem. 269:13092–13099, 1994.
- [75641] 13971.Tang, Y. M.; Wo, Y.-Y. P.; Stewart, J.; Hawkins, A. L.; Griffin,C. A.; Sutter, T. R.; Greenlee, W. F.: Isolation and characterizationof the human cytochrome P450 CYP1B1 gene. J. Biol. Chem. 271: 28324–28330,1996.
- [75642] 13972.Vincent, A.; Billingsley, G.; Priston, M.; Williams–Lyn, D.; Sutherland,J.; Glaser, T.; Oliver, E.; Walter, M. A.; Heathcote, G.; Levin, A.;Heon, E.: Phenotypic heterogeneity of CYP1B1: mutations in a patientwith Peters' anomaly. J. Med. Genet. 38: 324–326, 2001.
- [75643] 13973.Chen, H. T.; Bhandoola, A.; Difilippantonio, M. J.; Zhu, J.; Brown,M. J.; Tai, X.; Rogakou, E. P.; Brotz, T. M.; Bonner, W. M.; Ried,T.; Nussenzweig, A.: Response to RAG-mediated V(D)J cleavage by NBS1and gamma-H2AX. Science 290: 1962–1964, 2000.
- [75644] 13974.Petersen, S.; Casellas, R.; Reina–San–Martin, B.; Chen, H. T.;Difilippantonio, M. J.; Wilson, P. C.; Hanitsch, L.; Celeste, A.;Muramatsu, M.; Pilch, D. R.; Redon, C.; Ried,

T.; Bonner, W. M.; Honjo, T.; Nussenzweig, M. C.; Nussenzweig, A.: AID is required to initiate Nbs1/gamma-H2AX focus formation and mutations at sites of class switching. *Nature* 414:660–665, 2001.

- [75645] 13975. Morasso, M. I.; Yonescu, R.; Griffin, C. A.; Sargent, T. D.: Localization of human DLX8 to chromosome 17q21.3–q22 by fluorescence in situ hybridization. *Mammalian Genome* 8: 302–303, 1997.
- [75646] 13976. Quinn, L. M.; Johnson, B. V.; Nicholl, J.; Sutherland, G. R.; Kalionis, B.: Isolation and identification of homeobox genes from the human placenta including a novel member of the Distal-less family, DLX4. *Gene* 187:55–61, 1997.
- [75647] 13977. Boddy, M. N.; Howe, K.; Etkin, L. D.; Solomon, E.; Freemont, P. S.: PIC 1, a novel ubiquitin-like protein which interacts with the PML component of a multiprotein complex that is disrupted in acute promyelocytic leukaemia. *Oncogene* 13: 971–982, 1996.
- [75648] 13978. Desterro, J. M. P.; Rodriguez, M. S.; Hay, R. T.: SUMO-1 modification of I-kappa-B-alpha inhibits NF-kappa-B activation. *Molec. Cell* 2:233–239, 1998.
- [75649] 13979. Lapenta, V.; Chiurazzi, P.; van der Spek, P.; Pizzuti, A.; Hanaoka, F.; Brahe, C.: SMT3A, a human homologue of the *S. cerevisiae* SMT3 gene, maps to chromosome 21qter

and defines a novel gene family. *Genomics* 40:362–366, 1997.

- [75650] 13980.Mao, Y.; Sun, M.; Desai, S. D.; Liu, L. F.: SUMO–1 conjugation to topoisomerase I: a possible repair response to topoisomerase–mediated DNA damage. *Proc. Nat. Acad. Sci.* 97: 4046–4051, 2000.
- [75651] 13981.Okura, T.; Gong, L.; Kamitani, T.; Wada, T.; Okura, I.; Wei, C.–F.; Chang, H.–M.; Yeh, E. T. H.: Protection against Fas/APO–1– and tumor necrosis factor–mediated cell death by a novel protein, sentrin. *J. Immun.* 157: 4277–4281, 1996.
- [75652] 13982.Shen, Z.; Pardington–Purtymun, P. E.; Comeaux, J. C.; Moyzis, R.K.; Chen, D. J.: UBL1, a human ubiquitin–like protein associating with human RAD51/RAD52 proteins. *Genomics* 36: 271–279, 1996.
- [75653] 13983.Bengtsson, E.; Neame, P. J.; Heinegard, D.; Sommarin, Y.: The primary structure of a basic leucine–rich repeat protein, PRELP, found in connective tissues. *J. Biol. Chem.* 270: 25639–25644, 1995.
- [75654] 13984.Grover, J.; Chen, X.–N.; Korenberg, J. R.; Recklies, A. D.; Roughley, P. J.: The gene organization, chromosome location, and expression of a 55–kDa matrix protein (PRELP) of human articular cartilage. *Genomics*

38:109–117, 1996.

- [75655] 13985. Bigg, H. F.; Shi, Y. E.; Liu, Y. E.; Steffensen, B.; Overall, C.M.: Specific, high affinity binding of tissue inhibitor of metalloproteinases–4 (TIMP–4) to the COOH-terminal hemopexin-like domain of human gelatinase A. *J. Biol. Chem.* 272: 15496–15500, 1997.
- [75656] 13986. Greene, J.; Wang, M.; Liu, Y. E.; Raymond, L. A.; Rosen, C.; Shi, Y. E.: Molecular cloning and characterization of human tissue inhibitor of metalloproteinase 4. *J. Biol. Chem.* 271: 30375–30380, 1996.
- [75657] 13987. Leco, K. J.; Apte, S. S.; Taniguchi, G. T.; Hawkes, S. P.; Khokha, R.; Schultz, G. A.; Edwards, D. R.: Murine tissue inhibitor of metalloproteinases–4 (Timp–4): cDNA isolation and expression in adult mouse tissues. *FEBS Lett.* 401: 213–217, 1997.
- [75658] 13988. Olson, T. M.; Hirohata, S.; Ye, J.; Leco, K.; Seldin, M. F.; Apte, S. S.: Cloning of the human tissue inhibitor of metalloproteinase–4 gene (TIMP4) and localization of the TIMP4 and Timp4 gene to human chromosome 3p25 and mouse chromosome 6, respectively. *Genomics* 51:148–151, 1998.
- [75659] 13989. Wang, M.; Liu, Y. E.; Greene, J.; Sheng, S.; Fuchs, A.; Rosen, E. M.; Shi, Y. E.: Inhibition of tumor growth and

metastasis of human breast cancer cells transfected with tissue inhibitor of metalloproteinase4. *Oncogene* 14: 2767–2774, 1997.

[75660] 13990.Hsu, L. C.; Chang, W.-C.: Sequencing and expression of the human ALDH8 encoding a new member of the aldehyde dehydrogenase family. *Gene* 174:319–322, 1996.

[75661] 13991.Hsu, L. C.; Chang, W.-C.; Lin, S. W.; Yoshida, A.: Cloning and characterization of genes encoding four additional human aldehyde dehydrogenase isozymes. *Adv. Exp. Med. Biol.* 372: 159–168, 1995.

[75662] 13992.Rawson, R. B.; Zelenski, N. G.; Nijhawan, D.; Ye, J.; Sakai, J.; Hasan, M. T.; Chang, T. Y.; Brown, M. S.; Goldstein, J. L.: Complementation cloning of S2P, a gene encoding a putative metalloprotease required for intramembrane cleavage of SREBPs. *Molec. Cell* 1: 47–57, 1997.

[75663] 13993.Ye, J.; Rawson, R. B.; Komuro, R.; Chen, X.; Dave, U. P.; Prywes, R.; Brown, M. S.; Goldstein, J. L.: ER stress induces cleavage of membrane-bound ATF6 by the same proteases that process SREBPs. *Molec. Cell* 6: 1355–1364, 2000.

[75664] 13994.Zelenski, N. G.; Rawson, R. B.; Brown, M. S.; Goldstein, J. L.: Membrane topology of S2P, a protein required

for intramembraneous cleavage of sterol regulatory element-binding proteins. *J. Biol. Chem.* 274: 21973–21980, 1999.

- [75665] 13995. Kim, V. N.; Kataoka, N.; Dreyfuss, G.: Role of the nonsense-mediated decay factor hUpf3 in the splicing-dependent exon-exon junction complex. *Science* 293:1832–1836, 2001.
- [75666] 13996. Lykke-Andersen, J.; Shu, M.-D.; Steitz, J. A.: Human Upf protein targets an mRNA for nonsense-mediated decay when bound downstream of a termination codon. *Cell* 103: 1121–1131, 2000.
- [75667] 13997. Scott, A. F.: Personal Communication. Baltimore, Md. 2/9/2001.
- [75668] 13998. Serin, G.; Gersappe, A.; Black, J. D.; Aronoff, R.; Maquat, L.E.: Identification and characterization of human orthologues to *Saccharomyces cerevisiae* Upf2 protein and Upf3 protein (*Caenorhabditis elegans* SMG-4). *Molec. Cell. Biol.* 21: 209–223, 2001.
- [75669] 13999. de Leeuw, B.; Balemans, M.; Geurts van Kessel, A.: A novel Kruppel-associated box containing the SSX gene (SSX3) on the human X chromosome is not implicated in t(X;18)-positive synovial sarcomas. *Cytogenet. Cell Genet.* 73: 179–183, 1996.

- [75670] 14000.Gure, A. O.; Tureci, O.; Sahin, U.; Tsang, S.; Scanlan, M. J.; Jager, E.; Knuth, A.; Pfreundschuh, M.; Old, L. J.; Chen, Y.-T.:SSX: a multigene family with several members transcribed in normal testis and human cancer. *Int. J. Cancer* 72: 965–971, 1997.
- [75671] 14001.Kawai, A.; Woodruff, J.; Healey, J. H.; Brennan, M. F.; Antonescu, C. R.; Ladanyi, M.: SYT–SSX gene fusion as a determinant of morphology and prognosis in synovial sarcoma. *New Eng. J. Med.* 338: 153–160, 1998.
- [75672] 14002.Skytting, B.; Nilsson, G.; Brodin, B.; Xie, Y.; Lundeberg, J.; Uhlen, M.; Larsson, O.: A novel fusion gene, SYT–SSX4, in synovial sarcoma. (Letter) *J. Nat. Cancer Inst.* 91: 974–975, 1999.
- [75673] 14003.Yang, K.; Lui, W.-O.; Xie, Y.; Zhang, A.; Skytting, B.; Mandahl, N.; Larsson, C.; Larsson, O.: Co-existence of SYT–SSX1 and SYT–SSX2 fusions in synovial sarcomas. *Oncogene* 21: 4181–4190, 2002.
- [75674] 14004.Piccini, M.; Vitelli, F.; Seri, M.; Galletta, L. J. V.; Moran, O.; Bulfone, A.; Banfi, S.; Pober, B.; Renieri, A.: KCNE1-like gene is deleted in AMME contiguous gene syndrome: identification and characterization of the human and mouse homologs. *Genomics* 60: 251–257, 1999.
- [75675] 14005.Koda, T.; Kakinuma, M.: Molecular cloning of a

cDNA encoding a novel small GTP-binding protein. FEBS Lett. 328: 21–24, 1993.

[75676] 14006.Scott, A. F.: Personal Communication. Baltimore, Md. 5/14/2001.

[75677] 14007.Zheng, J. Y.; Koda, T.; Arimura, Y.; Kishi, M.; Kakinuma, M.:Structure and expression of the mouse S10 gene. Biochim. Biophys.Acta 1351: 47–50, 1997.

[75678] 14008.Tipnis, S. R.; Hooper, N. M.; Hyde, R.; Karran, E.; Christie, G.;Turner, A. J.: A human homolog of an-  
giotensin-converting enzyme:cloning and functional ex-  
pression as a captopril-insensitive carboxypeptidase.  
J.Biol. Chem. 275: 33238–33243, 2000.

[75679] 14009.Berta, P.; Hawkins, J. R.; Sinclair, A. H.; Taylor, A.; Griffiths,B. L.; Goodfellow, P. N.; Fellous, M.: Genetic evi-  
dence equatingSRY and the testis-determining factor. Na-  
ture 348: 448–450, 1990.

[75680] 14010.Disteche, C. M.; Casanova, M.; Saal, H.; Friedman, C.; Sybert,V.; Graham, J.; Thuline, H.; Page, D.; Fellous, M.: Small deletionsof the short arm of the Y chromosome in 46,XY females. Proc. Nat.Acad. Sci. 83: 7841–7844, 1986.

[75681] 14011.De Plaen, E.; De Backer, O.; Arnaud, D.; Bonjean, B.; Chomez, P.;Martelange, V.; Avner, P.; Baldacci, P.; Babinet, C.; Hwang, S.-Y.;Knowles, B.; Boon, T.: A new family of



mouse genes homologous to the human MAGE genes. *Genomics* 55: 176–184, 1999.

- [75682] 14012. Andersson, M.; Page, D. C.; de la Chapelle, A.: Chromosome Y-specific DNA is transferred to the short arm of X chromosome in human XX males. *Science* 233:786–788, 1986.
- [75683] 14013. Burgoyne, P. S.; Levy, E. R.; McLaren, A.: Spermatogenic failure in male mice lacking H-Y antigen. *Nature* 320: 170–172, 1986.
- [75684] 14014. Ferguson-Smith, M. A.: X-Y chromosomal interchange in the aetiology of true hermaphroditism and of XX Klinefelter's syndrome. *Lancet* II:475–476, 1966.
- [75685] 14015. Simpson, E.; Chandler, P.; Goulmy, E.; Distèche, C. M.; Ferguson-Smith, M. A.; Page, D. C.: Separation of the genetic loci for the H-Y antigen and for testis determination on human Y chromosome. *Nature* 326:876–878, 1987.
- [75686] 14016. Tiepolo, L.; Zuffardi, O.: Localization of factors controlling spermatogenesis in the nonfluorescent portion of the human Y chromosome long arm. *Hum. Genet.* 34: 119–124, 1976.
- [75687] 14017. Aalfs, C. M.; van den Berg, H.; Barth, P. G.; Hennekam, R. C. M.: The Hoyerlaal-Hreidarsson syndrome: the

fourth case of a separate entity with prenatal growth retardation, progressive pancytopenia and cerebellar hypoplasia. *Europ. J. Pediat.* 154: 304–308, 1995.

- [75688] 14018.Devriendt, K.; Matthijs, G.; Legius, E.; Schollen, E.; Blockmans, D.; van Geet, C.; Degreef, H.; Cassiman, J.-J.; Fryns, J.-P.: Skewed X-chromosome inactivation in female carriers of dyskeratosis congenita. *Am.J. Hum. Genet.* 60: 581–587, 1997.
- [75689] 14019.Hassock, S.; Vetrie, D.; Giannelli, F.: Mapping and characterization of the X-linked dyskeratosis congenita (DKC) gene. *Genomics* 55:21–27, 1999.
- [75690] 14020.Heiss, N. S.; Knight, S. W.; Vulliamy, T. J.; Klauck, S. M.; Wiemann, S.; Mason, P. J.; Poustka, A.; Dokal, I.: X-linked dyskeratosis congenita is caused by mutations in a highly conserved gene with putative nucleolar functions. *Nature Genet.* 19: 32–38, 1998.
- [75691] 14021.Heiss, N. S.; Megarbane, A.; Klauck, S. M.; Kreuz, F. R.; Makhoul, E.; Majewski, F.; Poustka, A.: One novel and two recurrent missense DKC1 mutations in patients with dyskeratosis congenita (DKC). *Genet.Counsel.* 12: 129–136, 2001.
- [75692] 14022.Abedinia, M.; Layfield, R.; Jones, S. M.; Nixon, P. F.; Mattick, J. S.: Nucleotide and predicted amino acid se–

quence of a cDNA clone encoding part of human transketolase. *Biochem. Biophys. Res. Commun.* 183:1159–1166, 1992.

[75693] 14023. Coy, J. F.; Dubel, S.; Kioschis, P.; Thomas, K.; Micklem, G.; Delius, H.; Poustka, A.: Molecular cloning of tissue-specific transcripts of a transketolase-related gene: implications for the evolution of new vertebrate genes. *Genomics* 32: 309–316, 1996.

[75694] 14024. McCool, B. A.; Plonk, S. G.; Martin, P. R.; Singleton, C. K.: Cloning of human transketolase cDNAs and comparison of the nucleotide sequence of the coding region in Wernicke-Korsakoff and non-Wernicke-Korsakoff individuals. *J. Biol. Chem.* 268: 1397–1404, 1993.

[75695] 14025. van der Maarel, S. M.; Scholten, I. H. J. M.; Huber, I.; Philippe, C.; Suijkerbuijk, R. F.; Gilgenkrantz, S.; Kere, J.; Cremers, F. P. M.; Ropers, H.-H.: Cloning and characterization of DXS6673E, a candidate gene for X-linked mental retardation in Xq13.1. *Hum. Molec. Genet.* 5:887–897, 1996.

[75696] 14026. Jacobi, F. K.; Broghammer, M.; Pesch, K.; Zrenner, E.; Berger, W.; Meindl, A.; Pusch, C. M.: Physical mapping and exclusion of GPR34 as the causative gene for congeni-

tal stationary night blindness type1. Hum. Genet. 107: 89–91, 2000.

- [75697] 14027. Marchese, A.; Sawzdargo, M.; Nguyen, T.; Cheng, R.; Heng, H. H.Q.; Nowak, T.; Im, D-S.; Lynch, K. R.; George, S. R.; O'Dowd, B. F.: Discovery of three novel orphan G-protein-coupled receptors. Genomics 56:12–21, 1999.
- [75698] 14028. Schoneberg, T.; Schulz, A.; Grosse, R.; Schade, R.; Henklein, P.; Schultz, G.; Gudermann, T.: A novel subgroup of class I G-protein-coupled receptors. Biochim. Biophys. Acta 1446: 57–70, 1999.
- [75699] 14029. Sanchis, D.; Fleury, C.; Chomiki, N.; Goubern, M.; Huang, Q.; Neverova, M.; Gregoire, F.; Easlick, J.; Raimbault, S.; Levi-Meyrueis, C.; Miroux, B.; Collins, S.; Seldin, M.; Richard, D.; Warden, C.; Bouillaud, F.; Ricquier, D.: BMCP1, a novel mitochondrial carrier with high expression in the central nervous system of humans and rodents, and respiration uncoupling activity in recombinant yeast. J. Biol. Chem. 273: 34611–34615, 1998.
- [75700] 14030. Blanco, P.; Sargent, C. A.; Boucher, C. A.; Mitchell, M.; Affara, N. A.: Conservation of PCDHX in mammals; expression of X/Y genes predominantly in brain. Mammalian Genome 11: 906–914, 2000.

- [75701] 14031.Ciccodicola, A.; D'Esposito, M.; Esposito, T.; Gianfrancesco, F.;Migliaccio, C.; Miano, M. G.; Matarazzo, M. R.; Vacca, M.; Franze,A.; Cuccurese, M.; Cocchia, M.; Curci, A.; and 9 others: Differentiallyregulated and evolved genes in the fully sequenced Xq/Yq pseudoautosomalregion. *Hum. Molec. Genet.* 9: 395–401, 2000.
- [75702] 14032.Mumm, S.; Molini, B.; Terrell, J.; Srivastava, A.; Schlessinger,D.: Evolutionary features of the 4–Mb Xq21.3 XY homology region revealedby a map at 60–kb resolution. *Genome Res.* 7: 307–314, 1997.
- [75703] 14033.Schwartz, A.; Chan, D. C.; Brown, L. G.; Alagappan, R.; Pettay,D.; Distech, C.; McGillivray, B.; de la Chapelle, A.; Page, D. C.: Reconstructing hominid Y evolution: X–homologous block, createdby X–Y transposition, was disrupted by Yp inversion through LINE–LINErecombination. *Hum. Molec. Genet.* 7: 1–11, 1998.
- [75704] 14034.Tilford, C. A.; Kuroda–Kawaguchi, T.; Skaletsky, H.; Rozen, S.;Brown, L. G.; Rosenberg, M.; McPherson, J. D.; Wylie, K.; Sekhon,M.; Kucaba, T. A.; Waterston, R. H.; Page, D. C.: A physical mapof the human Y chromosome. *Nature* 409: 943–945, 2001.
- [75705] 14035.Roberts, R. G.; Kendall, E.; Vetrie, D.; Bobrow, M.: Sequenceand chromosomal location of a human homo–

logue of LRPR1, an FSH primary response gene. Genomics 37: 122–124, 1996.

[75706] 14036.Slegtenhorst–Eegdeman, K. E.; Post, M.; Baarends, W. M.; Themmen, A. P. N.; Grootegeed, J. A.: Regulation of gene expression in Sertolicells by follicle–stimulating hormone (FSH): cloning and characterization of LRPR1, a primary response gene encoding a leucine–rich protein. Molec.Cell. Endocr. 108: 115–124, 1995.

[75707] 14037.Batch, J. A.; Evans, B. A. J.; Hughes, I. A.; Patterson, M. N.: Mutations of the androgen receptor gene identified in perineal hypospadias. J.Med. Genet. 30: 198–201, 1993.

[75708] 14038.Borkhardt, A.; Repp, R.; Haas, O. A.; Leis, T.; Harbott, J.; Kreuder, J.; Hammermann, J.; Henn, T.; Lampert, F.: Cloning and characterization of AFX, the gene that fuses to MLL in acute leukemias with a t(X;11)(q13;q23). Oncogene 14:195–202, 1997.

[75709] 14039.Chatterjee, A.; Faust, C. J.; Herman, G. E.: Genetic and physical mapping of the biglycan gene on the mouse X chromosome. Mammalian Genome 4: 33–36, 1993.

[75710] 14040.Das, S.; Metzenberg, A.; Pai, G. S.; Gitschier, J.: Mutational analysis of the biglycan gene excludes it as a candidate for X–linked dominant chondrodysplasia punct-

tata, dyskeratosis congenita, and incontinentia pigmenti.  
(Letter) Am. J. Hum. Genet. 54: 922–925, 1994.

- [75711] 14041. Fisher, L. W.; Heegaard, A.-M.; Vetter, U.; Vogel, W.; Just, W.; Termine, J. D.; Young, M. F.: Human biglycan gene: putative promoter, intron–exon junctions, and chromosomal localization. J. Biol. Chem. 266:14371–14377, 1991.
- [75712] 14042. Fisher, L. W.; Termine, J. D.; Young, M. F.: Deduced–protein sequence of bone small proteoglycan I (biglycan) shows homology with proteoglycan II (decorin) and several nonconnective tissue proteins in a variety of species. J. Biol. Chem. 264: 4571–4576, 1989.
- [75713] 14043. Geerkens, C.; Vetter, U.; Just, W.; Fedarko, N. S.; Fisher, L. W.; Young, M. F.; Termine, J. D.; Gehron Robey, P.; Wöhrle, D.; Vogel, W.: The X–chromosomal human biglycan gene BGN is subject to X inactivation but is transcribed like an X–Y homologous gene. Hum. Genet. 96:44–52, 1995.
- [75714] 14044. McBride, O. W.; Fisher, L. W.; Young, M. F.: Localization of PGI (biglycan, BGN) and PGII (decorin, DCN, PG-40) genes on human chromosomes Xq13–qter and 12q, respectively. Genomics 6: 219–225, 1990.
- [75715] 14045. Traupe, H.; van den Ouweland, A. M. W.; van Oost,

B. A.; Vogel, W.; Vetter, U.; Warren, S. T.; Rocchi, M.; Darlison, M. G.; Ropers, H.-H.: Fine mapping of the human biglycan (BGN) gene within the Xq28 region employing a hybrid cell panel. *Genomics* 13: 481–483, 1992.

[75716] 14046. Wegrowski, Y.; Pillarisetti, J.; Danielson, K. G.; Suzuki, S.; Iozzo, R. V.: The murine biglycan: complete cDNA cloning, genomic organization, promoter function, and expression. *Genomics* 30: 8–17, 1995.

[75717] 14047. Xu, T.; Bianco, P.; Fisher, L. W.; Longenecker, G.; Smith, E.; Goldstein, S.; Bonadio, J.; Boskey, A.; Heegaard, A.-M.; Sommer, B.; Satomura, K.; Dominguez, P.; Zhao, C.; Kulkarni, A. B.; Robey, P. G.; Young, M. F.: Targeted disruption of the biglycan gene leads to an osteoporosis-like phenotype in mice. *Nature Genet.* 20: 78–82, 1998.

[75718] 14048. Yoshida, K.; Sugano, S.: Identification of a novel protocadherin gene (PCDH11) on the human XY homology region in Xq21.3. *Genomics* 62: 540–543, 1999.

[75719] 14049. de la Chapelle, A.; Sankila, E.-M.; Lindlof, M.; Aula, P.; Norio, R.: Norrie disease caused by a gene deletion allowing carrier detection and prenatal diagnosis. *Clin. Genet.* 28: 317–320, 1985.

[75720] 14050. Sims, K. B.; de la Chapelle, A.; Norio, R.; Sankila, E.-M.; Hsu, Y.-P. P.; Rinehart, W. B.; Corey, T. J.; Ozelius,



L.; Powell, J. F.; Bruns, G.; Gusella, J. F.; Murphy, D. L.; Breakefield, X. O.: Monoamine oxidase deficiency in males with an X chromosome deletion. *Neuron* 2:1069–1076, 1989.

[75721] 14051. Grino, P. B.; Griffin, J. E.; Cushard, W. G., Jr.; Wilson, J. D.: A mutation of the androgen receptor associated with partial androgen resistance, familial gynecomastia, and fertility. *J. Clin. Endocr. Metab.* 66: 754–761, 1988.

[75722] 14052. Holterhus, P.-M.; Sinnecker, G. H. G.; Hiort, O.: Phenotypic diversity and testosterone-induced normalization of mutant L712F androgen receptor function in a kindred with androgen insensitivity. *J. Clin. Endocr. Metab.* 85: 3245–3250, 2000.

[75723] 14053. Hughes, I. A.; Evans, B. A. J.: Complete androgen insensitivity syndrome characterized by increased concentration of a normal androgen receptor in genital skin fibroblasts. *J. Clin. Endocr. Metab.* 63:309–315, 1986.

[75724] 14054. Imperato-McGinley, J.; Ip, N. Y.; Gautier, T.; Neuweiler, J.; Gruenspan, H.; Liao, S.; Chang, C.; Balazs, I.: DNA linkage analysis and studies of the androgen receptor gene in a large kindred with complete androgen insensitivity. *Am. J. Med. Genet.* 36: 104–108, 1990.

[75725] 14055. Jukier, L.; Kaufman, M.; Pinsky, L.; Peterson, R. E.:

Partial androgen resistance associated with secondary 5- $\alpha$ -reductase deficiency: identification of a novel qualitative androgen receptor defect and clinical implications. *J. Clin. Endocr. Metab.* 59: 679–688, 1984.

[75726] 14056. Kaufman, M.; Pinsky, L.; Bowin, A.; Au, M. W. S.: Familial external genital ambiguity due to a transformation defect of androgen-receptor complexes that is expressed with 5- $\alpha$ -dihydrotestosterone and the synthetic androgen methyltrienolone. *Am. J. Med. Genet.* 18: 493–507, 1984.

[75727] 14057. Kaufman, M.; Straisfeld, C.; Pinsky, L.: Male pseudohermaphroditism presumably due to target organ unresponsiveness to androgens: deficient 5- $\alpha$ -dihydrotestosterone binding in cultured skin fibroblasts. *J. Clin. Invest.* 58: 345–350, 1976.

[75728] 14058. Keenan, B. S.; Meyer, W. J., III; Hadjian, A. J.; Jones, H. W.; Migeon, C. J.: Syndrome of androgen insensitivity in man: absence of 5- $\alpha$ -dihydrotestosterone binding protein in skin fibroblasts. *J. Clin. Endocr.* 38: 1143–1146, 1974.

[75729] 14059. Liao, S.; Witte, D.: Autoimmune anti-androgen-receptor antibodies in human serum. *Proc. Nat. Acad. Sci.* 82: 8345–8348, 1985.

- [75730] 14060.Lin, S.-Y.; Ohno, S.: The binding of androgen receptor to DNA and RNA. *Biochim. Biophys. Acta* 654: 181–186, 1981.
- [75731] 14061.Aradhya, S.; Bardaro, T.; Galgoczy, P.; Yamagata, T.; Esposito, T.; Patlan, H.; Ciccodicola, A.; Munnich, A.; Kenwrick, S.; Platzner, M.; D'Urso, M.; Nelson, D. L.: Multiple pathogenic and benign genomic rearrangements occur at a 35 kb duplication involving the NEMO and LAGE2 genes. *Hum. Molec. Genet.* 10: 2557–2567, 2001.
- [75732] 14062.Aradhya, S.; Courtois, G.; Rajkovic, A.; Lewis, R. A.; Levy, M.; Israel, A.; Nelson, D. L.: Atypical forms ofcontinentia pigmenti in male individuals result from mutations of a cytosine tract in exon 10 of NEMO (IKK- $\gamma$ ). *Am. J. Hum. Genet.* 68: 765–771, 2001.
- [75733] 14063.Aradhya, S.; Woffendin, H.; Jakins, T.; Bardaro, T.; Esposito, T.; Smahi, A.; Shaw, C.; Levy, M.; Munnich, A.; D'Urso, M.; Lewis, R. A.; Kenwrick, S.; Nelson, D. L.: A recurrent deletion in the ubiquitously expressed NEMO (IKK- $\gamma$ ) gene accounts for the vast majority ofcontinentia pigmenti mutations. *Hum. Molec. Genet.* 10: 2171–2179, 2001.
- [75734] 14064.Doffinger, R.; Smahi, A.; Bessia, C.; Geissmann, F.; Feinberg, J.; Durandy, A.; Bodemer, C.; Kenwrick, S.;

Dupuis–Girod, S.; Blanche, S.; Wood, P.; Rabia, S. H.; and 16 others: X–linked anhidrotic ectodermal dysplasia with immunodeficiency is caused by impaired NF–kappa–B signaling. *Nature Genet.* 27: 277–285, 2001.

[75735] 14065. Jain, A.; Ma, C. A.; Liu, S.; Brown, M.; Cohen, J.; Strober, W.: Specific missense mutations in NEMO result in hyper–IgM syndrome with hypohydrotic (sic) ectodermal dysplasia. *Nature Immun.* 2: 223–228, 2001.

[75736] 14066. Jin, D. Y.; Jeang, K. T.: Isolation of full–length cDNA and chromosomal localization of human NF–kappaB modulator NEMO to Xq28. *J. Biomed. Sci.* 6: 115–120, 1999.

[75737] 14067. Kosaki, K.; Shimasaki, N.; Fukushima, H.; Hara, M.; Ogata, T.; Matsuo, N.: Female patient showing hypohydrotic ectodermal dysplasia and immunodeficiency (HED–ID). (Letter) *Am. J. Hum. Genet.* 69: 664–665, 2001.

[75738] 14068. Li, Q.; Van Antwerp, D.; Mercurio, F.; Lee, K.–F.; Verma, I. M.: Severe liver degeneration in mice lacking the I–kappa–B kinase 2 gene. *Science* 284: 321–325, 1999.

[75739] 14069. Li, Y.; Kang, J.; Friedman, J.; Tarassishin, L.; Ye, J.; Kovalenko, A.; Wallach, D.; Horwitz, M. S.: Identification of a cell protein (FIP–3) as a modulator of NF–kappa–B activity and as a target of an adenovirus inhibitor of tumor necrosis factor alpha–induced apoptosis. *Proc. Nat. Acad. Sci.*

96: 1042–1047, 1999.

- [75740] 14070. Makris, C.; Godfrey, V. L.; Krahn–Senftleben, G.; Takahashi, T.; Roberts, J. L.; Schwarz, T.; Feng, L.; Johnson, R. S.; Karin, M.: Female mice heterozygous for IKK–gamma/NEMO deficiencies develop a dermatopathy similar to the human X–linked disorder incontinentia pigmenti. *Molec. Cell* 5: 969–979, 2000.
- [75741] 14071. May, M. J.; D'Acquisto, F.; Madge, L. A.; Glockner, J.; Pober, J. S.; Ghosh, S.: Selective inhibition of NF–kappa–B activation by a peptide that blocks the interaction of NEMO with the I–kappa–B kinase complex. *Science* 289: 1550–1554, 2000.
- [75742] 14072. Roberts, J. L.; Morrow, B.; Vega–Rich, C.; Salafia, C. M.; Nitowsky, H. M.: Incontinentia pigmenti in a newborn male infant with DNA confirmation. *Am. J. Med. Genet.* 75: 159–163, 1998.
- [75743] 14073. Rothwarf, D. M.; Zandi, E.; Natoli, G.; Karin, M.: IKK–gamma is an essential regulatory subunit of the I–kappa–B kinase complex. *Nature* 395: 297–300, 1998.
- [75744] 14074. Rudolph, D.; Yeh, W.–C.; Wakeham, A.; Rudolph, B.; Nallainathan, D.; Potter, J.; Elia, A. J.; Mak, T. W.: Severe liver degeneration and lack of NF–kappa–B activation in NEMO/IKK–gamma–deficient mice. *Genes Dev.* 14:

854–862, 2000.

- [75745] 14075. Schmidt–Supprian, M.; Bloch, W.; Courtois, G.; Ad-dicks, K.; Israel, A.; Rajewsky, K.; Pasparakis, M.: NEMO/IKK–gamma–deficient mice model incontinentia pigmenti. *Molec. Cell* 5: 981–992, 2000.
- [75746] 14076. The International Incontinentia Pigmenti Consortium: Genomic rearrangement in NEMO impairs NF–kappa–B activation and is a cause of incontinentia pigmenti. *Nature* 405: 466–472, 2000.
- [75747] 14077. The International IP Consortium: Survival of male patients with incontinentia pigmenti carrying a lethal mutation can be explained by somatic mosaicism or Klinefelter syndrome. *Am. J. Hum. Genet.* 69: 1210–1217, 2001.
- [75748] 14078. Yamaoka, S.; Courtois, G.; Bessia, C.; Whiteside, S. T.; Weil, R.; Agou, F.; Kirk, H. E.; Kay, R. J.; Israel, A.: Complementation cloning of NEMO, a component of the I–kappa–B kinase complex essential for NF–kappa–B activation. *Cell* 93: 1231–1240, 1998.
- [75749] 14079. Zonana, J.; Elder, M. E.; Schneider, L. C.; Orlow, S. J.; Moss, C.; Golabi, M.; Shapira, S. K.; Farndon, P. A.; Wara, D. W.; Emmal, S. A.; Ferguson, B. M.: A novel X–linked disorder of immune deficiency and hypohidrotic ectodermal dysplasia is allelic to incontinentia pigmenti and due to

mutations in IKK-gamma (NEMO). Am. J. Hum. Genet. 67:1555–1562, 2000.

- [75750] 14080. Bauer, M. F.; Gempel, K.; Reichert, A. S.; Rappold, G. A.; Lichtner, P.; Gerbitz, K.-D.; Neupert, W.; Brunner, M.; Hofmann, S.: Genetic and structural characterization of the human mitochondrial inner membrane translocase. J. Molec. Biol. 289: 69–82, 1999.
- [75751] 14081. Moller, D. E.; Xia, C. H.; Tang, W.; Zhu, A. X.; Jakubowski, M.: Human rsk isoforms: cloning and characterization of tissue-specific expression. Am. J. Physiol. 266: C351–C359, 1994.
- [75752] 14082. Zhao, Y.; Bjorbaek, C.; Weremowicz, S.; Morton, C. C.; Moller, D. E.: RSK3 encodes a novel pp90rsk isoform with a unique N-terminal sequence: growth factor-stimulated kinase function and nuclear translocation. Molec. Cell. Biol. 15: 4353–4363, 1995.
- [75753] 14083. Au, H. C.; Seo, B. B.; Matsuno-Yagi, A.; Yagi, T.; Scheffler, I. E.: The NDUFA1 gene product (MWFE protein) is essential for activity of complex I in mammalian mitochondria. Proc. Nat. Acad. Sci. 96:4354–4359, 1999.
- [75754] 14084. Zhuchenko, O.; Wehnert, M.; Bailey, J.; Sun, Z. S.; Lee, C. C.: Isolation, mapping, and genomic structure of an X-linked gene for a subunit of human mitochondrial

complex I. Genomics 37: 281–288,1996.

[75755] 14085.Liu, Z.; Sun, C.; Olejniczak, E. T.; Meadows, R. P.; Betz, S. F.; Oost, T.; Herrmann, J.; Wu, J. C.; Fesik, S. W.: Structural basis for binding of Smac/DIABLO to the XIAP BIR3 domain. Nature 408:1004–1008, 2000.

[75756] 14086.Srinivasula, S. M.; Hegde, R.; Saleh, A.; Datta, P.; Shiozaki, E.; Chai, J.; Lee, R.-A.; Robbins, P. D.; Fernandes-Alnemri, T.; Shi, Y.; Alnemri, E. S.: A conserved XIAP–interaction motif in caspase–9 and Smac/DIABLO regulates caspase activity and apoptosis. Nature 410:112–116, 2001.

[75757] 14087.Bomont, P.; Cavalier, L.; Blondeau, F.; Ben Hamida, C.; Belal, S.; Tazir, M.; Demir, E.; Topaloglu, H.; Korinthenberg, R.; Tuysuz, B.; Landrieu, P.; Hentati, F.; Koenig, M.: The gene encoding gigaxonin, a new member of the cytoskeletal BTB/kelch repeat family, is mutated in giant axonal neuropathy. Nature Genet. 26: 370–374, 2000.

[75758] 14088.Treiber–Held, S.; Budjarjo–Welim, H.; Riemann, D.; Richter, J.; Kretzschmar, H. A.; Hanefeld, F.: Giant axonal neuropathy: a generalized disorder of intermediate filaments with longitudinal grooves in the hair. Neuropediatrics 25: 89–93, 1994.

[75759] 14089.Lieuallen, K.; Pennacchio, L. A.; Park, M.; Myers, R.



M.; Lennon, G. G.: Cystatin B-deficient mice have increased expression of apoptosis and glial activation genes. *Hum. Molec. Genet.* 10: 1867–1871, 2001.

- [75760] 14090. Mazarib, A.; Xiong, L.; Neufeld, M. Y.; Birnbaum, M.; Korczyn, A. D.; Pandolfo, M.; Berkovic, S. F.: Unverricht-Lundborg disease in a five-generation Arab family: instability of dodecamer repeats. *Neurology* 57:1050–1054, 2001.
- [75761] 14091. Pennacchio, L. A.; Bouley, D. M.; Higgins, K. M.; Scott, M. P.; Noebels, J. L.; Myers, R. M.: Progressive ataxia, myoclonic epilepsy and cerebellar apoptosis in cystatin B-deficient mice. *Nature Genet.* 20:251–258, 1998.
- [75762] 14092. Virtaneva, K.; D'Amato, E.; Miao, J.; Koskiniemi, M.; Norio, R.; Avanzini, G.; Franceschetti, S.; Michelucci, R.; Tassinari, C. A.; Omer, S.; Pennacchio, L. A.; Myers, R. M.; Dieguez-Lucena, J. L.; Krahe, R.; de la Chapelle, A.; Lehesjoki, A.-E.: Unstable minisatellite expansion causing recessively inherited myoclonus epilepsy, EPM1. *Nature-Genet.* 15: 393–396, 1997.
- [75763] 14093. Cannizzaro, L. A.; Chen, Y. Q.; Rafi, M. A.; Wenger, D. A.: Regional mapping of the human galactocerebrosidase gene (GALC) to 14q31 by in situ hybridization. *Cytogenet. Cell Genet.* 66: 244–245, 1994.

- [75764] 14094.De Gasperi, R.; Sosa, M. A. G.; Sartorato, E. L.; Battistini, S.; MacFarlane, H.; Gusella, J. F.; Krivit, W.; Kolodny, E. H.: Molecular heterogeneity of late-onset forms of globoid-cell leukodystrophy. *Am.J. Hum. Genet.* 59: 1233–1242, 1996.
- [75765] 14095.Duchen, L. W.; Eicher, E. M.; Jacobs, J. M.; Scaravilli, F.; Teixeira, F.: Hereditary leucodystrophy in the mouse: the new mutant twitcher. *Brain* 103:695–710, 1980.
- [75766] 14096.Minassian, B. A.; Lee, J. R.; Herbrick, J.-A.; Huizenga, J.; Soder, S.; Mungall, A. J.; Dunham, I.; Gardner, R.; Fong, C. G.; Carpenter, S.; Jardim, L.; Satishchandra, P.; Andermann, E.; Snead, O. C., III; Lopes-Cendes, I.; Tsui, L.-C.; Delgado-Escueta, A. V.; Rouleau, G.A.; Scherer, S. W.: Mutations in a gene encoding a novel protein tyrosine phosphatase cause progressive myoclonus epilepsy. *Nature Genet.* 20: 171–174, 1998.
- [75767] 14097.Norio, R.; Koskiniemi, M.: Progressive myoclonus epilepsy: genetic and nosological aspects with special reference to 107 Finnish patients. *Clin.Genet.* 15: 382–398, 1979.
- [75768] 14098.Ortiz-Hidalgo, C.: The man behind Lafora's bodies. *Am. J. Surg.Path.* 10: 358–361, 1986.
- [75769] 14099.Sainz, J.; Minassian, B. A.; Serratosa, J. M.; Gee, M.

N.; Sakamoto, L. M.; Iranmanesh, R.; Bohlega, S.; Baumann, R. J.; Ryan, S.; Sparkes, R. S.; Delgado-Escueta, A. V.:

Lafora progressive myoclonus epilepsy: narrowing the chromosome 6q24 locus by recombinations and homozygosity. (Letter) *Am. J. Hum. Genet.* 61: 1205–1209, 1997.

[75770] 14100. Sarlin, M. B.; Klopfer, H. W.; Mickle, W. A.; Heath, R. G.: The detection of carriers in hereditary myoclonic epilepsy. *Acta Genet. Med. Gemellol.* 9: 466–471, 1960.

[75771] 14101. Schwarz, G. A.; Yanoff, M.: Lafora's disease, distinct clinico-pathologic form of Unverricht's syndrome. *Arch. Neurol.* 12: 172–188, 1965.

[75772] 14102. Serratosa, J. M.; Delgado-Escueta, A. V.; Posada, I.; Shih, S.; Drury, I.; Berciano, J.; Zabala, J. A.; Antunez, M. C.; Sparkes, R. S.: The gene for progressive myoclonus epilepsy of the Lafora type maps to chromosome 6q. *Hum. Molec. Genet.* 4: 1657–1663, 1995.

[75773] 14103. Serratosa, J. M.; Gomez-Garre, P.; Gallardo, M. E.; Anta, B.; Beltran-Valero de Bernabe, D.; Lindhout, D.; Augustijn, P. B.; Tassinari, C. A.; Michelucci, R.; Malafosse, A.; Topcu, M.; Grid, D.; Dravet, C.; Berkovic, S. F.; Rodriguez de Cordoba, S.: A novel protein tyrosine phosphatase gene is mutated in progressive myoclonus epilepsy of the Lafora type (EPM2). *Hum. Molec. Genet.* 8: 345–352, 1999.

- [75774] 14104.Yanoff, M.; Schwarz, G. A.: Lafora's disease: a distinct geneticallydetermined form of Unverricht's syndrome. Genet. Hum. 14: 235–244,1965.
- [75775] 14105.Yokoi, S.; Austin, J.; Witmer, F.; Sakai, M.: Studies in myoclonusepilepsy (Lafora body forms). I. Isolation and preliminary characterizationof Lafora bodies in two cases. Arch. Neurol. 19: 15–33, 1968.
- [75776] 14106.Antonarakis, S.: Personal Communication. Geneva, Switzerland 4/8/1997.
- [75777] 14107.Lafreniere, R. G.; Rochefort, D. L.; Chretien, N.; Rommens, J.M.; Cochius, J. I.; Kalviainen, R.; Nousiainen, U.; Patry, G.; Farrell,K.; Soderfeldt, B.; Federico, A.; Hale, B. R.; Cossio, O. H.; Sorensen,T.; Pouliot, M. A.; Kmiec, T.; Uldall, P.; Janszky, J.; Pranzatelli,M. R.; Andermann, F.; Andermann, E.; Rouleau, G. A.: Unstable insertionof the 5–prime flanking region of the cystatin B gene is the mostcommon mutation in progressive myoclonus epilepsy type 1, EPM1. NatureGenet. 15: 298–302, 1997.
- [75778] 14108.Lalioi, M. D.; Mirotsou, M.; Buresi, C.; Peitsch, M. C.; Rossier,C.; Ouazzani, R.; Baldy–Moulinier, M.; Bottani, A.; Malafosse, A.;Antonarakis, S. E.: Identification of mutations in cystatin B, thegene responsible for the Unverricht–Lundborg type of progressive myoclonusepilepsy

(EPM1). Am. J. Hum. Genet. 60: 342–351, 1997.

- [75779] 14109.Lalioi, M. D.; Scott, H. S.; Buresi, C.; Rossier, C.; Bottani,A.; Morris, M. A.; Malafosse, A.; Antonarakis, S. E.: Dodecamer repeatexpansion in cystatin B gene in progressive myoclonus epilepsy. Nature 386:847–851, 1997.
- [75780] 14110.Carmel, R.: Gastric juice in congenital pernicious anemia containsno immunoreactive intrinsic factor molecule: study of three kindredswith variable ages at presentation, including a patient first diagnosedin adulthood. Am. J. Hum. Genet. 35: 67–77, 1983.
- [75781] 14111.Lyerla, T. A.; Konola, J. T.; Skiba, M. C.; Raghavan, S.: Galactocerebrosidaseactivity in somatic cell hybrids derived from twitcher mouse/controlhuman fibroblasts is associated with human chromosome 17. Am. J.Hum. Genet. 44: 198–207, 1989.
- [75782] 14112.Sakai, N.; Inui, K.; Fujii, N.; Fukushima, H.; Nishimoto, J.;Yanagihara,I.; Isegawa, Y.; Iwamatsu, A.; Okada, S.: Krabbe disease:isolation and characterization of a full-length cDNA for human galactocerebrosidase. Biochem.Biophys. Res. Commun. 198: 485–491, 1994.
- [75783] 14113.Sweet, H.: Twitcher (twi) is on chromosome 12. Mouse Newsletter 75:30, 1986.
- [75784] 14114.Roume, J.; Genin, E.; Cormier–Daire, V.; Ma, H. W.;

Mehaye, B.;Attie, T.; Razavi-Encha, F.; Fallet-Bianco, C.; Buenerd, A.; Clerget-Darpoux,F.; Munnich, A.; Le Merrer, M.: A gene for Meckel syndrome maps to chromosome 11q13. Am. J. Hum. Genet. 63: 1095-1101, 1998.

[75785] 14115.Fernandes, M.; Poirier, C.; Lespinasse, F.; Carle, G. F.: Themouse homologs of human GIF, DDB1, and CFL1 genes are located on chromosome19. Mammalian Genome 9: 339-342, 1998.

[75786] 14116.Hewitt, J. E.; Gordon, M. M.; Taggart, R. T.; Mohandas, T. K.;Alpers, D. H.: Human gastric intrinsic factor: characterization ofcDNA and genomic clones and localization to human chromosome 11. Genomics 10:432-440, 1991.

[75787] 14117.Katz, M.; Lee, S. K.; Cooper, B. A.: Vitamin B(12) malabsorptiondue to biologically inert intrinsic factor. New Eng. J. Med. 287:425-429, 1972.

[75788] 14118.Katz, M.; Mehlman, C. S.; Allen, R. H.: Isolation and characterizationof an abnormal human intrinsic factor. J. Clin. Invest. 53: 1274-1283,1974.

[75789] 14119.Levine, J. S.; Yang, Y.; Ducos, R.; Rosenberg, A. J.; Catron, P.G.; Podell, E. R.; Allen, R. H.: Juvenile pernicious anemia (JPA)due to an abnormal intrinsic factor (IF) that is markedly susceptible to acid and proteolysis. (Abstract)

Gastroenterology 80: 1210 only,1981.

- [75790] 14120.McNicholl, B.; Egan, B.: Congenital pernicious anemia: effectson growth, brain, and absorption of B12. Pediatrics 42: 149–156,1968.
- [75791] 14121.Remacha, A. F.; Sambeat, M. A.; Barcelo, M. J.; Mones, J.; Garcia–Die,J.; Gimferrer, E.: Congenital intrinsic factor deficiency in a Spanishpatient. Ann. Hemat. 64: 202–204, 1992.
- [75792] 14122.Yang, Y. M.; Ducos, R.; Rosenberg, A. J.; Catrou, P. G.; Levine,J. S.; Podell, E. R.; Allen, R. H.: Cobalamin malabsorption in threesiblings due to an abnormal intrinsic factor that is markedly susceptible to acid and proteolysis. J. Clin. Invest. 76: 2057–2065, 1985.
- [75793] 14123.Carney, J. P.; Maser, R. S.; Olivares, H.; Davis, E. M.; Le Beau,M.; Yates, J. R., III; Hays, L.; Morgan, W. F.; Petrini, J. H. J.: The hMre11/hRad50 protein complex and Nijmegen breakage syndrome:linkage of double–strand break repair to the cellular DNA damage response. Cell 93:477–486, 1998.
- [75794] 14124.Mailly, F.; Palmen, J.; Muller, D. P. R.; Gibbs, T.; Lloyd, J.;Brunzell, J.; Durrington, P.; Mitropoulos, K.; Betteridge, J.; Watts,G.; Lithell, H.; Angelico, F.; Humphries, S. E.; Talmud, P. J.: Familialipoprotein lipase (LPL) deficiency:

a catalogue of LPL gene mutations identified in 20 patients from the UK, Sweden, and Italy. *Hum. Mutat.* 10:465–473, 1997.

[75795] 14125. Normand, T.; Bergeron, J.; Fernandez–Margallo, T.; Bharucha, A.; Ven Murthy, M. R.; Julien, P.; Gagne, C.; Dionne, C.; De Braekeleer, M.; Ma, R.; Hayden, M. R.; Lupien, P. J.: Geographic distribution and genealogy of mutation 207 of the lipoprotein lipase gene in the French Canadian population of Quebec. *Hum. Genet.* 89: 671–675, 1992.

[75796] 14126. Monsalve, M. V.; Henderson, H.; Roederer, G.; Julien, P.; Deeb, S.; Kastelein, J. J. P.; Peritz, L.; Devlin, R.; Bruin, T.; Murthy, M. R. V.; Gagne, C.; Davignon, J.; Lupien, P. J.; Brunzell, J. D.; Hayden, M. R.: A missense mutation at codon 188 of the human lipoprotein lipase gene is a frequent cause of lipoprotein lipase deficiency in persons of different ancestries. *J. Clin. Invest.* 86: 728–734, 1990.

[75797] 14127. Cobben, J. M.; van der Steege, G.; Grootscholten, P.; de Visser, M.; Scheffer, H.; Buys, C. H. C. M.: Deletions of the survival motor neuron gene in unaffected siblings of patients with spinal muscular atrophy. *Am. J. Hum. Genet.* 57: 805–808, 1995.

[75798] 14128. Hahnen, E.; Schonling, J.; Rudnik–Schoneborn, S.;



Zerres, K.;Wirth, B.: Hybrid survival motor neuron genes in patients with autosomalrecessive spinal muscular atrophy: new insights into molecular mechanismsresponsible for the disease. *Am. J. Hum. Genet.* 59: 1057–1065, 1996.

[75799] 14129.Lefebvre, S.; Burglen, L.; Reboullet, S.; Clermont, O.; Burlet,P.; Viollet, L.; Benichou, B.; Cruaud, C.; Mil-lasseau, P.; Zeviani,M.; Le Paslier, D.; Frezal, J.; Cohen, D.; Weissenbach, J.; Munnich,A.; Melki, J.: Identification and characterization of a spinal muscularatrophy–determining gene. *Cell* 80: 155–165, 1995.

[75800] 14130.Matthijs, G.; Schollen, E.; Legius, E.; Devriendt, K.; Goemans,N.; Kayserili, H.; Apak, M. Y.; Cassiman, J.–J.: Unusual molecularfindings in autosomal recessive spinal muscular atrophy. *J. Med.Genet.* 33: 469–474, 1996.

[75801] 14131.Handschug, K.; Sperling, S.; Yoon, S.–J. K.; Hennig, S.; Clark,A. J. L.; Huebner, A.: Triple A syndrome is caused by mutations inAAAS, a new WD–repeat protein gene. *Hum. Molec. Genet.* 10: 283–290,2001.

[75802] 14132.Kaplan, J.; Gerber, S.; Larget–Piet, D.; Rozet, J.–M.; Dollfus,H.; Dufier, J.–L.; Odent, S.; Postel–Vinay, A.; Janin, N.; Briard,M.–L.; Frezal, J.; Munnich, A.: A gene for Star–gardt's disease (fundusflavimaculatus) maps to the short arm of chromosome 1. *Nature Genet.* 5:308–311, 1993.

- [75803] 14133.Coppa, G. V.; Giorgi, P. L.; Felici, L.; Gabrielli, O.; Donti,E.; Bernasconi, S.; Kresse, H.; Paschke, E.; Mastropaolo, C.: Clinicalheterogeneity in Sanfilippo disease (mucopolysaccharidosis III) typeD: presentation of two new cases. *Europ. J. Pediat.* 140: 130–133,1983.
- [75804] 14134.Dhoot, G. K.; Gustafsson, M. K.; Ai, X.; Sun, W.; Standiford, D.M.; Emerson, C. P., Jr.: Regulation of Wnt signaling and embryo patterningby an extracellular sulfatase. *Science* 293: 1663–1666, 2001.
- [75805] 14135.Freeman, C.; Clements, P. R.; Hopwood, J. J.: Human liver N–acetylglucosamine–6–sulphatesulphatase: purification and characterization. *Biochem. J.* 246:347–354, 1987.
- [75806] 14136.Freeman, C.; Hopwood, J. J.: Human liver N–acetylglucosamine–6–sulphatesulphatase: catalytic properties. *Biochem. J.* 246: 355–365, 1987.
- [75807] 14137.Gatti, R.; Borrone, C.; Durand, P.; De Virgiliis, S.; Sanna, G.;Cao, A.; von Figura, K.; Kresse, H.; Paschke, E.: Sanfilippo typeD disease: clinical findings in two patients with a new variant ofmucopolysaccharidosis III. *Europ. J. Pediat.* 138: 168–171, 1982.
- [75808] 14138.Kaplan, P.; Wolfe, L. S.: Sanfilippo syndrome type D. *J. Pediat.* 110:267–271, 1987.

- [75809] 14139.Kresse, H.; Paschke, E.; von Figura, K.; Gilberg, W.; Fuchs, W.: Sanfilippo disease type D: deficiency of N-acetylglucosamine-6-sulfatesulfatase required for heparan sulfate degradation. *Proc. Nat. Acad.Sci.* 77: 6822-6826, 1980.
- [75810] 14140.Robertson, D. A.; Callen, D. F.; Baker, E. G.; Morris, C. P.; Hopwood,J. J.: Chromosomal localization of the gene for human glucosamine-6-sulphatase to 12q14. *Hum. Genet.* 79: 175-178, 1988.
- [75811] 14141.Robertson, D. A.; Freeman, C.; Nelson, P. V.; Morris, C. P.; Hopwood,J. J.: Human glucosamine-6-sulfatase cDNA reveals homology with steroidsulfatase. *Biochem. Biophys. Res. Commun.* 157: 218-224, 1988.
- [75812] 14142.Siciliano, L.; Fiumara, A.; Pavone, L.; Freeman, C.; Robertson,D.; Morris, C. P.; Hopwood, J. J.; Di Natale, P.; Musumeci, S.; Horwitz,A. L.: Sanfilippo syndrome type D in two adolescent sisters. *J.Med. Genet.* 28: 402-405, 1991.
- [75813] 14143.Thompson, J. N.; Jones, M. Z.; Dawson, G.; Huffman, P. S.: N-acetylglucosamine-6-sulphatase deficiency in a Nubian goat: a model of Sanfilippo syndrome type D (mucopolysaccharidosis IIID). *J. Inherit. Metab. Dis.* 15:760-768, 1992.
- [75814] 14144.Faiella, A.; Brunelli, S.; Granata, T.; D'Incerti, L.;

Cardini,R.; Lenti, C.; Battaglia, G.; Boncinelli, E.: A number of schizencephalypatients including 2 brothers are heterozygous for germline mutations in the homeobox gene EMX2. *Europ. J. Hum. Genet.* 5: 186–190, 1997.

[75815] 14145.Bespalova, I. N.; Adkins, S.; Pranzatelli, M.; Burmeister, M.:Novel cystatin B mutation and diagnostic PCR assay in an Unverricht–Lundborgprogressive myoclonus epilepsy patient. *Am. J. Med. Genet.* 74: 467–471,1997.

[75816] 14146.Jarvinen, M.; Rinne, A.: Human spleen cysteine proteinase inhibitor:purification, fractionation into isoelectric variants and some propertiesof the variants. *Biochim. Biophys. Acta* 708: 210–217, 1982.

[75817] 14147.Jerala, R.; Trstenjak, M.; Lenarcic, B.; Turk, V.: Cloning a syntheticgene for human stefin B and its expression in *E. coli*. *FEBS Lett.* 239:41–44, 1988.

[75818] 14148.Lalioti, M. D.; Scott, H. S.; Antonarakis, S. E.: What is expandedin progressive myoclonus epilepsy? (Letter) *Nature Genet.* 17: 17only, 1997.

[75819] 14149.Lalioti, M. D.; Scott, H. S.; Genton, P.; Grid, D.; Ouazzani, R.;M'Rabet, A.; Ibrahim, S.; Gouider, R.; Dravet, C.; Chkili, T.; Bottani,A.; Buresi, C.; Malafosse, A.; Antonarakis, S. E.: A PCR amplificationmethod reveals insta-

bility of the dodecamer repeat in progressive myoclonus epilepsy (EPM1) and no correlation between the size of the repeat and age at onset. *Am. J. Hum. Genet.* 62: 842–847, 1998.

- [75820] 14150. Pennacchio, L. A.; Lehesjoki, A.-E.; Stone, N. E.; Willour, V. L.; Virtaneva, K.; Miao, J.; D'Amato, E.; Ramirez, L.; Faham, M.; Koskiniemi, M.; Warrington, J. A.; Norio, R.; de la Chapelle, A.; Cox, D. R.; Myers, R. M.: Mutations in the gene encoding cystatin B in progressive myoclonus epilepsy (EPM1). *Science* 271: 1731–1733, 1996.
- [75821] 14151. Pennacchio, L. A.; Myers, R. M.: Isolation and characterization of the mouse cystatin B gene. *Genome Res.* 6: 1103–1109, 1996.
- [75822] 14152. Pranzatelli, M. R.; Tate, E.; Huang, Y.; Haas, R. H.; Bodensteiner, J.; Ashwal, S.; Franz, D.: Neuropharmacology of progressive myoclonic epilepsy: response to 5-hydroxy-L-tryptophan. *Epilepsia* 36: 783–791, 1995.
- [75823] 14153. Turk, V.; Bode, W.: The cystatins: protein inhibitors of cysteine proteinases. *FEBS Lett.* 285: 213–219, 1991.
- [75824] 14154. Chang, S. C.; Hoang, B.; Thomas, J. T.; Vukicevic, S.; Luyten, F. P.; Ryba, N. J. P.; Kozak, C. A.; Reddi, A. H.; Moos, M., Jr.: Cartilage-derived morphogenetic proteins: new members of the transforming growth factor- $\beta$  su-

perfamily predominantly expressed in long bones during human embryonic development. J. Biol. Chem. 269: 28227–28234, 1994.

- [75825] 14155. Faiyaz–Ul–Haque, M.; Ahmad, W.; Wahab, A.; Haque, S.; Azim, A.C.; Zaidi, S. H. E.; Teebi, A. S.; Ahmad, M.; Cohn, D. H.; Siddique, T.; Tsui, L.–C.: Frameshift mutation in the cartilage–derived morphogenetic protein 1 (CDMP1) gene and severe acromesomelic chondrodysplasia resembling Grebe–type chondrodysplasia. Am. J. Med. Genet. 111: 31–37, 2002.
- [75826] 14156. Rousseau–Merck, M.–F.; Duro, D.; Berger, R.; Thiesen, H. J.: Chromosomal localization of two KOX zinc finger genes on chromosome bands 7q21–q22. Ann. Genet. 38: 81–84, 1995.
- [75827] 14157. Tommerup, N.; Vissing, H.: Isolation and fine mapping of 16 novel human zinc finger–encoding cDNAs identify putative candidate genes for developmental and malignant disorders. Genomics 27: 259–264, 1995.
- [75828] 14158. Canman, C. E.; Radany, E. H.; Parsels, L. A.; Davis, M. A.; Lawrence, T. S.; Maybaum, J.: Induction of resistance to fluorodeoxyuridine cytotoxicity and DNA damage in human tumor cells by expression of *Escherichia coli* deoxyuridine triphosphatase. Cancer Res. 54:

2296–2298,1994.

- [75829] 14159.Canman, C. E.; Tang, H. Y.; Normolle, D. P.; Lawrence, T. S.; Maybaum,J.: Variations in patterns of DNA damage induced in human colorectal tumor cells by 5–fluorodeoxyuridine: implications for mechanisms of resistance and cytotoxicity. *Proc. Nat. Acad. Sci.* 89: 10474–10478,1992.
- [75830] 14160.Chu, R.; Lin, Y.; Rao, M. S.; Reddy, J. K.: Cloning and identification of rat deoxyuridine triphosphatase as an inhibitor of peroxisome proliferator–activated receptor alpha. *J. Biol. Chem.* 271: 27670–27676, 1996.
- [75831] 14161.Cohen, D.; Heng, H. H. Q.; Shi, X.–M.; McIntosh, E. M.; Tsui, L.–C.; Pearlman, R. E.: Assignment of the human dUTPase gene (DUT) to chromosome 15q15–q21.1 by fluorescence in situ hybridization. *Genomics* 40:213–215, 1997.
- [75832] 14162.el–Hajj, H. H.; Zhang, H.; Weiss, B.: Lethality of a dut (deoxyuridine triphosphatase) mutation in *Escherichia coli*. *J. Bacteriol.* 170:1069–1075, 1988.
- [75833] 14163.Ladner, R. D.; Caradonna, S. J.: The human dUTPase gene encodes both nuclear and mitochondrial isoforms: differential expression of the isoforms and characterization of a cDNA encoding the mitochondrial species. *J.*

Biol. Chem. 272: 19072–19080, 1997.

- [75834] 14164.Ladner, R. D.; McNulty, D. E.; Carr, S. A.; Roberts, G. D.; Caradonna, S. J.: Characterization of distinct nuclear and mitochondrial forms of human deoxyuridine triphosphate nucleotidohydrolase. J. Biol. Chem. 271: 7745–7751, 1996.
- [75835] 14165.Lindahl, T.: DNA repair enzymes. Annu. Rev. Biochem. 51: 61–87, 1982.
- [75836] 14166.McIntosh, E. M.; Ager, D. D.; Gadsden, M. H.; Haynes, R. H.: Human dUTP pyrophosphatase: cDNA sequence and potential biological importance of the enzyme. Proc. Nat. Acad. Sci. 89: 8020–8024, 1992. Note: Erratum: Proc. Nat. Acad. Sci.: 90: 4328 only, 1993.
- [75837] 14167.Boring, L.; Gosling, J.; Cleary, M.; Charo, I. F.: Decreased lesion formation in CCR2  $-/-$  mice reveals a role for chemokines in the initiation of atherosclerosis. Nature 394: 894–897, 1998.
- [75838] 14168.Charo, I. F.; Myers, S. J.; Herman, A.; Franci, C.; Connolly, A.J.; Coughlin, S. R.: Molecular cloning and functional expression of two monocyte chemoattractant protein 1 receptors reveals alternative splicing of the carboxyl-terminal tails. Proc. Nat. Acad. Sci. 91: 2752–2756, 1994.



- [75839] 14169.Barrett, T. E.; Savva, R.; Panayotou, G.; Barlow, T.; Brown, T.;Jiricny, J.; Pearl, L. H.: Crystal structure of a G:T/U mismatch–specificDNA glycosylase: mismatch recognition by complementary–strand interactions. *Cell* 92:117–129, 1998.
- [75840] 14170.De Gregorio, L.; Gallinari, P.; Gariboldi, M.; Ma–nenti, G.; Pierotti,M. A.; Jiricny, J.; Dragani, T. A.: Genetic mapping of thymine DNAglycosylase (Tdg) gene and of one pseudogene in the mouse. *MammalianGenome* 7: 909–910, 1996.
- [75841] 14171.Lindahl, T.: DNA repair enzymes. *Ann. Rev. Biochem.* 51: 61–87,1982.
- [75842] 14172.Neddermann, P.; Gallinari, P.; Lettieri, T.; Schmid, D.; Truong,O.; Hsuan, J. J.; Wiebauer, K.; Jiricny, J.: Cloning and expressionof human G/T mismatch–specific thymine–DNA glycosylase. *J. Biol.Chem.* 271: 12767–12774, 1996.
- [75843] 14173.Sard, L.; Tornielli, S.; Gallinari, P.; Minoletti, F.; Jir–icny,J.; Lettieri, T.; Pierotti, M. A.; Sozzi, G.; Radice, P.: Chromosomallocalizations and molecular analysis of TDG gene–related sequences. *Genomics* 44:222–226, 1997.
- [75844] 14174.DiMarco, S. P.; Glover, T. W.; Miller, D. E.; Reines, D.; Warren,S. T.: Transcription elongation factor SII (TCEA) maps to human chromosome3p22–p21.3. *Genomics* 36:

185–188, 1996.

- [75845] 14175. Park, H.; Baek, K.; Jeon, C.; Agarwal, K.; Yoo, O.: Characterization of the gene encoding the human transcriptional elongation factor TFIIIS. *Gene* 139:263–267, 1994.
- [75846] 14176. Reines, D.: Nascent RNA cleavage by transcription elongation complexes. In: Conaway, R. C.; Conaway, J. W. (eds.): *Transcription: Mechanisms and Regulation*. New York: Raven Press 1994. Pp. 263–278.
- [75847] 14177. Thomas, M. J.; Platas, A. A.; Hawley, D. K.: Transcriptional fidelity and proofreading by RNA polymerase II. *Cell* 93: 627–637, 1998.
- [75848] 14178. Chang, C.; da Silva, S. L.; Ideta, R.; Lee, Y.; Yeh, S.; Burbach, J. P. H.: Human and rat TR4 orphan receptors specify a subclass of the steroid receptor superfamily. *Proc. Nat. Acad. Sci.* 91: 6040–6044, 1994.
- [75849] 14179. Chang, C.; Kokontis, J.; Acakpo-Satchivi, L.; Liao, S.; Takeda, H.; Chang, Y.: Molecular cloning of new human TR2 receptors: a class of steroid receptor with multiple ligand-binding domains. *Biochem. Biophys. Res. Commun.* 165: 735–741, 1989.
- [75850] 14180. Hirose, T.; Fujimoto, W.; Yamaai, T.; Kim, K. H.; Matsuura, H.; Jetten, A. M.: TAK1: Molecular cloning and

characterization of a new member of the nuclear receptor superfamily. *Molec. Endocr.* 1667–1680, 1994.

[75851] 14181. Yoshikawa, T.; DuPont, B. R.; Leach, R. J.; Detera-Wadleigh, S.D.: New variants of the human and rat nuclear hormone receptor, TR4: expression and chromosomal localization of the human gene. *Genomics* 35:361–366, 1996.

[75852] 14182. Applequist, S. E.; Selg, M.; Raman, C.; Jack, H.-M.: Cloning and characterization of HUPF1, a human homolog of the *Saccharomyces cerevisiae* nonsense mRNA-reducing UPF1 protein. *Nucleic Acids Res.* 25: 814–821, 1997.

[75853] 14183. Medghalchi, S. M.; Frischmeyer, P. A.; Mendell, J. T.; Kelly, A.G.; Lawler, A. M.; Dietz, H. C.: Rent1, a trans-effector of nonsense-mediated mRNA decay, is essential for mammalian embryonic viability. *Hum. Molec. Genet.* 10: 99–105, 2001.

[75854] 14184. Mendell, J. T.; ap Rhys, C. M. J.; Dietz, H. C.: Separable roles for rent1/hUpf1 in altered splicing and decay of nonsense transcripts. *Science* 298:419–371, 2002.

[75855] 14185. Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Dietz, H. C.: Cloning and characterization of a human regulator of nonsense transcript stability. (Abstract) *Am. J. Hum. Genet.* 59 (suppl.): A32 only, 1996.

- [75856] 14186.Perlick, H. A.; Medghalchi, S. M.; Spencer, F. A.; Kendzior, R.J., Jr.; Dietz, H. C.: Mammalian orthologues of a yeast regulator of nonsense transcript stability. *Proc. Nat. Acad. Sci.* 93: 10928–10932, 1996.
- [75857] 14187.Sun, X.; Perlick, H. A.; Dietz, H. C.; Maquat, L. E.: A mutated human homologue to yeast Upf1 protein has a dominant-negative effect on the decay of nonsense-containing mRNAs in mammalian cells. *Proc. Nat. Acad. Sci.* 95: 10009–10014, 1998.
- [75858] 14188.Eng, C.; Myers, S. M.; Kogon, M. D.; Sanicola, M.; Hession, C.; Cate, R. L.; Mulligan, L. M.: Genomic structure and chromosomal localization of the human GDNFR- $\alpha$  gene. *Oncogene* 16: 597–601, 1998.
- [75859] 14189.GFR- $\alpha$  Nomenclature Committee: Nomenclature of GPI-linked receptors for the GDNF ligand family. *Neuron* 19: 485 only, 1997.
- [75860] 14190.Gorodinsky, A.; Zimonjic, D. B.; Popescu, N. C.; Milbrandt, J.: Assignment of the GDNF family receptor  $\alpha$ -1 (GFRA1) to human chromosome band 10q26 by in situ hybridization. *Cytogenet. Cell Genet.* 78:289–290, 1997.
- [75861] 14191.Jing, S.; Wen, D.; Yu, Y.; Holst, P. L.; Luo, Y.; Fang, M.; Tamir, R.; Antonio, L.; Hu, Z.; Cupples, R.; Louis, J.-C.;

Hu, S.; Altroch, B. W.; Fox, G. M.: GDNF-induced activation of the Ret protein tyrosine kinase is mediated by GDNFR- $\alpha$ , a novel receptor for GDNF. *Cell* 85:1113–1124, 1996.

[75862] 14192. Paratcha, G.; Ledda, F.; Baars, L.; Culpier, M.; Besset, V.; Anders, J.; Scott, R.; Ibanez, C. F.: Released GFR- $\alpha$ -1 potentiates downstream signaling, neuronal survival, and differentiation via a novel mechanism of recruitment of c-Ret to lipid rafts. *Neuron* 29: 171–184, 2001.

[75863] 14193. Puliti, A.; Cinti, R.; Seri, M.; Ceccherini, I.; Romeo, G.: Assignment of mouse Gfra1, the homologue of a new human HSCR candidate gene, to the telomeric region of mouse chromosome 19. *Cytogenet. Cell Genet.* 78:291–294, 1997.

[75864] 14194. Shefelbine, S. E.; Khorana, S.; Schultz, P. N.; Huang, E.; Thobe, N.; Hu, Z. J.; Fox, G. M.; Jing, S.; Cote, G. J.; Gagel, R. F.: Mutational analysis of the GDNF/RET-GDNFR- $\alpha$  signaling complex in a kindred with vesicoureteral reflux. *Hum. Genet.* 102: 474–478, 1998.

[75865] 14195. Chen, H.; Chrast, R.; Rossier, C.; Gos, A.; Antonarakis, S. E.; Kudoh, J.; Yamaki, A.; Shindoh, N.; Maeda, H.; Minoshima, S.; Shimizu, N.: Single-minded and Down syndrome? (Letter) *Nature Genet.* 10:9–10, 1995.

[75866] 14196. Chrast, R.; Scott, H. S.; Madani, R.; Huber, L.;

Wolfer, D. P.;Prinz, M.; Aguzzi, A.; Lipp, H.-P.; Antonarakis, S. E.: Mice trisomicfor a bacterial artificial chromosome with the single-minded 2 gene(Sim2) show phenotypes similar to some of those present in the partialtrisomy 16 mouse models of Down syndrome. Hum. Molec. Genet. 9:1853–1864, 2000.

[75867] 14197.Dahmane, N.; Charron, G.; Lopes, C.; Yaspo, M.-L.; Maunoury, C.;Decorte, L.; Sinet, P.-M.; Bloch, B.; Delabar, J.-M.: Down syndrome–criticalregion contains a gene homologous to Drosophila sim expressed duringrat and human central nervous system development. Proc. Nat. Acad.Sci. 92: 9191–9195, 1995.

[75868] 14198.Ema, M.; Ikegami, S.; Hosoya, T.; Mimura, J.; Ohtani, H.; Nakao,K.; Inokuchi, K.; Katsuki, M.; Fujii–Kuriyama, Y.: Mild impairmentof learning and memory in mice overexpressing the mSim2 gene locatedon chromosome 16: an animal model of Down's syndrome. Hum. Molec.Genet. 8: 1409–1415, 1999.

[75869] 14199.Moffett, P.; Dayo, M.; Reece, M.; McCormick, M. K.; Pelletier,J.: Characterization of msim, a murine homologue of the Drosophilasim transcription factor. Genomics 35: 144–155, 1996.

[75870] 14200.Muenke, M.; Bone, L. J.; Mitchell, H. F.; Hart, I.;

Walton, K.;Hall-Johnson, K.; Ippel, E. F.; Dietz-Band, J.; Kvaloy, K.; Fan, C.-M.;Tessier-Lavigne, M.; Patterson, D.: Physical mapping of the holoprosencephalycritical region in 21q22.3, exclusion of SIM2 as a candidate gene for holoprosencephaly, and mapping of SIM2 to a region of chromosome 21 important for Down syndrome. *Am. J. Hum. Genet.* 57: 1074–1079,1995.

[75871] 14201.Yamaki, A.; Noda, S.; Kudoh, J.; Shindoh, N.; Maeda, H.; Minoshima,S.; Kawasaki, K.; Shimizu, Y.; Shimizu, N.: The mammalian single-minded(SIM) gene: mouse cDNA structure and diencephalic expression indicate a candidate gene for Down syndrome. *Genomics* 35: 136–143, 1996.

[75872] 14202.de Winter, J. P.; Leveille, F.; van Berkel, C. G. M.; Rooimans,M. A.; van der Weel, L.; Steltenpool, J.; Demuth, I.; Morgan, N. V.;Alon, N.; Bosnoyan-Collins, L.; Lightfoot, J.; Leegwater, P. A.; Waisfisz,Q.; Komatsu, K.; Arwert, F.; Pronk, J. C.; Mathew, C. G.; Digweed,M.; Buchwald, M.; Joenje, H.: Isolation of a cDNA representing the Fanconi anemia complementation group E gene. *Am. J. Hum. Genet.* 67:1306–1308, 2000. Note: Erratum: *Am. J. Hum. Genet.* 67: 1365 only,2000.

[75873] 14203.Sosa-Pineda, B.; Wigle, J. T.; Oliver, G.: Hepatocyte

migration during liver development requires Prox1. *Nature Genet.* 25: 254–255, 2000.

[75874] 14204. Wigle, J. T.; Chowdhury, K.; Gruss, P.; Oliver, G.: Prox1 function is crucial for mouse lens-fibre elongation. *Nature Genet.* 21: 318–322, 1999.

[75875] 14205. Wigle, J. T.; Oliver, G.: Prox1 function is required for the development of the murine lymphatic system. *Cell* 98: 769–778, 1999.

[75876] 14206. Zinovieva, R. D.; Duncan, M. K.; Johnson, T. R.; Torres, R.; Polymeropoulos, M. H.; Tomarev, S. I.: Structure and chromosomal localization of the human homeobox gene Prox 1. *Genomics* 35: 517–522, 1996.

[75877] 14207. Ladenburger, E. M.; Fackelmayer, F. O.; Hameister, H.; Knippers, R.: MCM4 and PRKDC, human genes encoding proteins MCM4 and DNA-PKcs, are close neighbours located on chromosome 8q12–q13. *Cytogenet. Cell Genet.* 77: 268–270, 1997.

[75878] 14208. Ma, Y.; Pannicke, U.; Schwarz, K.; Lieber, M. R.: Hairpin opening and overhang processing by an Artemis/DNA-dependent protein kinase complex in nonhomologous end joining and V(D)J recombination. *Cell* 108: 781–794, 2002.

[75879] 14209. Satoh, T.; Tsuruga, H.; Yabuta, N.; Ishidate, M., Jr.;



Nojima,H.: Assignment of the human CDC21 (MCM4) gene to chromosome 8q11.2. *Genomics* 46:525–526, 1997.

[75880] 14210.Joenje, H.; Lo Ten Foe, J. R.; Oostra, A. B.; van Berkel, C. G.M.; Rooimans, M. A.; Schroeder–Kurth, T.; Wegner, R.–D.; Gille, J.J. P.; Buchwald, M.; Arwert, F.: Classification of Fanconi anemiapartients by complementation analysis: evidence for a fifth geneticsubtype. *Blood* 86: 2156–2160, 1995.

[75881] 14211.Byk, T.; Dobransky, T.; Cifuentes–Diaz, C.; Sobel, A.: Identificationand molecular characterization of Unc–33–like phosphoprotein (Ulip),a putative mammalian homolog of the axonal guidance–associated unc–33gene product. *J. Neurosci.* 16: 688–701, 1996.

[75882] 14212.Choe, H.; Farzan, M.; Sun, Y.; Sullivan, N.; Rollins, B.; Ponath,P. D.; Wu, L.; Mackay, C. R.; LaRosa, G.; Newman, W.; Gerard, N.;Gerard, C.; Sodroski, J.: The beta–chemokine receptors CCR3 and CCR5facilitate infection by primary HIV–1 isolates. *Cell* 85: 1135–1148,1996.

[75883] 14213.Diriong, S.; Lory, P.; Williams, M. E.; Ellis, S. B.; Harpold,M. M.; Taviaux, S.: Chromosomal localization of the human genes foralpha–1A, alpha–1B, and alpha–1E voltage–dependent Ca(2+) channelsubunits. *Genomics* 30: 605–609, 1995.

- [75884] 14214.Ben-Porath, I.; Benvenisty, N.: Characterization of a tumor-associated gene, a member of a novel family of genes encoding membrane glycoproteins. *Gene* 183:69–75, 1996.
- [75885] 14215.Hong, Y.; Ohishi, K.; Inoue, N.; Endo, Y.; Fujita, T.; Takeda, J.; Kinoshita, T.: Structures and chromosomal localizations of the glycosylphosphatidylinositol synthesis gene PIGC and its pseudogene PIGCP1. *Genomics* 44: 347–349, 1997.
- [75886] 14216.Gaetano, C; Matsuo, T.; Thiele, C. J.: Identification and characterization of a retinoic acid-regulated human homologue of the unc-33-like phosphoprotein gene (hUlip) from neuroblastoma cells. *J. Biol. Chem.* 272: 12195–12201, 1997.
- [75887] 14217.Matsuo, T.; Stauffer, J. K.; Walker, R. L.; Meltzer, P.; Thiele, C. J.: Structure and promoter analysis of the human unc-33-like phosphoprotein gene: E-box required for maximal expression in neuroblastoma and myoblasts. *J. Biol. Chem.* 275: 16560–16568, 2000.
- [75888] 14218.Behm, F. G.; Smith, F. O.; Raimondi, S. C.; Pui, C.-H.; Bernstein, I. D.: Human homologue of the rat chondroitin sulfate proteoglycan, NG2, detected by monoclonal antibody 7.1, identifies childhood acute lymphoblastic

leukemias with t(4;11)(q21;q23) or t(11;19)(q23;p13) and MLL gene rearrangements. Blood 87: 1134–1139, 1996.

[75889] 14219. Pluschke, G.; Vanek, M.; Evans, A.; Dittmar, T.; Schmid, P.; Itin, P.; Filardo, E. J.; Reisfeld, R. A.: Molecular cloning of a human melanoma-associated chondroitin sulfate proteoglycan. Proc. Nat. Acad. Sci. 93: 9710–9715, 1996.

[75890] 14220. Rettig, W. J.; Dracopoli, N. C.; Goetzger, T. A.; Spengler, B. A.; Biedler, J. L.; Oettgen, H. F.; Old, L. J.: Somatic cell genetic analysis of human cell surface antigens: chromosomal assignments and regulation of expression in rodent–human hybrid cells. Proc. Nat. Acad. Sci. 81: 6437–6441, 1984.

[75891] 14221. Rettig, W. J.; Real, F. X.; Spengler, B. A.; Biedler, J. L.; Old, L. J.: Human melanoma proteoglycan: expression in hybrids controlled by intrinsic and extrinsic signals. Science 231: 1281–1284, 1986.

[75892] 14222. Smith, F. O.; Rauch, C.; Williams, D. E.; March, C. J.; Arthur, D.; Hilden, J.; Lampkin, B. C.; Buckley, J. D.; Buckley, C. V.; Woods, W. G.; Dinndorf, P. A.; Sorensen, P.; Kersey, J.; Hammond, D.; Bernstein, I. D.: The human homologue of rat NG2, a chondroitin sulfate proteoglycan, is not expressed on the cell surface of normal hematopoietic

cells but is expressed by acute myeloid leukemia blasts from poor-prognosis patients with abnormalities of chromosome band 11q23. *Blood* 87:1123–1133, 1996.

[75893] 14223. Gipp, J. J.; Bailey, H. H.; Mulcahy, R. T.: Cloning and sequencing of the cDNA for the light subunit of human liver gamma-glutamylcysteine synthetase and relative mRNA levels for heavy and light subunits in human normal tissues. *Biochem. Biophys. Res. Commun.* 206: 584–589, 1995.

[75894] 14224. Huang, C.-S.; Anderson, M. E.; Meister, A.: Amino acid sequence and function of the light subunit of rat kidney gamma-glutamylcysteine synthetase. *J. Biol. Chem.* 268: 20578–20583, 1993.

[75895] 14225. Sierra-Rivera, E.; Dasouki, M.; Summar, M. L.; Krishnamani, M. R. S.; Meredith, M.; Rao, P. N.; Phillips, J. A., III; Freeman, M. L.: Assignment of the human gene (GLCLR) that encodes the regulatory subunit of gamma-glutamylcysteine synthetase to chromosome 1p21. *Cytogenet. Cell Genet.* 72: 252–254, 1996.

[75896] 14226. Tsuchiya, K.; Mulcahy, R. T.; Reid, L. L.; Distech, C. M.; Kavanagh, T. J.: Mapping of the glutamate-cysteine ligase catalytic subunit gene (GLCLC) to human chromosome 6p12 and mouse chromosome 9D-E and of the reg-

ulatory subunit gene (GLCLR) to human chromosome 1p21-p22 and mouse chromosome 3H1-3. Genomics 30: 630-632, 1995.

- [75897] 14227. Rozet, J.-M.; Gerber, S.; Perrault, I.; Calvas, P.; Souied, E.; Chatelin, S.; Viegas-Pequignot, E.; Molina-Gomez, D.; Munnich, A.; Kaplan, J.: Structure and refinement of the physical mapping of the gamma-glutamylcysteine ligase regulatory subunit (GLCLR) gene to chromosome 1p22.1 within the critically deleted region of human malignant mesothelioma. Cytogenet. Cell Genet. 82: 91-94, 1998.
- [75898] 14228. Ellis, J. A.; Luzio, J. P.: Identification and characterization of a novel protein (p137) which transcytoses bidirectionally in Caco-2 cells. J. Biol. Chem. 270: 20717-20723, 1995.
- [75899] 14229. Gessler, M.; Klamt, B.; Tsaoussidou, S.; Ellis, J. A.; Luzio, J. P.: The gene encoding the GPI-anchored membrane protein p137(GPI)(M11S1) maps to human chromosome 11p13 and is highly conserved in the mouse. Genomics 32: 169-170, 1996.
- [75900] 14230. Inoue, N.; Watanabe, R.; Takeda, J.; Kinoshita, T.: PIG-C, one of the three human genes involved in the first step of glycosylphosphatidylinositol biosynthesis is a ho-

mologue of *Saccharomyces cerevisiae* GPI2.

Biochem.Biophys. Res. Commun. 226: 193–199, 1996.

[75901] 14231. Ring, H. Z.; Vameghi-Meyers, V.; Wang, W.; Crabtree, G. R.; Francke, U.: Five SWI/SNF-related, matrix-associated, actin-dependent regulator of chromatin (SMARC) genes are dispersed in the human genome. Genomics 51:140–143, 1998.

[75902] 14232. Wang, W.; Xue, Y.; Zhou, S.; Kuo, A.; Cairns, B. R.; Crabtree, G. R.: Diversity and specialization of mammalian SWI/SNF complexes. Genes Dev. 10: 2117–2130, 1996.

[75903] 14233. Kim, B.-T.; Kitagawa, H.; Tamura, J.; Saito, T.; Kusche-Gullberg, M.; Lindahl, U.; Sugahara, K.: Human tumor suppressor EXT gene family members EXTL1 and EXTL3 encode alpha-1,4-N-acetylglucosaminyl transferases that likely are involved in heparan sulfate/heparin biosynthesis. Proc. Nat. Acad. Sci. 98: 7176–7181, 2001.

[75904] 14234. Wise, C. A.; Clines, G. A.; Massa, H.; Trask, B. J.; Lovett, M.: Identification and localization of the gene for EXTL, a third member of the multiple exostoses gene family. Genome Res. 7: 10–16, 1997.

[75905] 14235. Moskow, J. J.; Bullrich, F.; Huebner, K.; Daar, I. O.; Buchberg, A. M.: Meis1, a PBX1-related homeobox gene

involved in myeloid leukemia in BXH-2 mice. *Molec. Cell. Biol.* 15: 5434–5443, 1995.

[75906] 14236. Steelman, S.; Moskow, J. J.; Muzynski, K.; North, C.; Druck, T.; Montgomery, J. C.; Huebner, K.; Daar, I. O.; Buchberg, A. M.: Identification of a conserved family of Meis1-related homeobox genes. *Genome Res.* 7:142–156, 1997.

[75907] 14237. Thorsteinsdottir, U.; Kroon, E.; Jerome, L.; Blasi, F.; Sauvageau, G.: Defining roles for HOX and MEIS1 genes in induction of acute myeloid leukemia. *Molec. Cell. Biol.* 21: 224–234, 2001.

[75908] 14238. Capdevila, J.; Tsukui, T.; Esteban, C. R.; Zappavigna, V.; Belmonte, J. C. I.: Control of vertebrate limb outgrowth by the proximal factor Meis2 and distal antagonism of BMPs by Gremlin. *Molec. Cell* 4: 839–849, 1999.

[75909] 14239. Nakamura, T.; Jenkins, N. A.; Copeland, N. G.: Identification of a new family of Pbx-related homeobox genes. *Oncogene* 13: 2235–2242, 1996.

[75910] 14240. Smith, J. E.; Afonja, O.; Yee, H. T.; Inghirami, G.; Takeshita, K.: Chromosomal mapping to 15q14 and expression analysis of the human MEIS2 homeobox gene. *Mammalian Genome* 8: 951–952, 1997.

[75911] 14241. Kobayashi, S.; Uemura, H.; Kohda, T.; Nagai, T.;

Chinen, Y.; Naritomi, K.; Kinoshita, E.; Ohashi, H.; Imaizumi, K.; Tsukahara, M.; Sugio, Y.; Tonoki, H.; Kishino, T.; Tanaka, T.; Yamada, M.; Tsutsumi, O.; Niikawa, N.; Kaneko-Ishino, T.; Ishino, F.: No evidence of PEG1/MEST gene mutations in Silver-Russell syndrome patients. *Am. J. Med. Genet.* 104:225–231, 2001.

[75912] 14242. Mizuno, K.; Hasegawa, K.; Katagiri, T.; Ogimoto, M.; Ichikawa, T.; Yakura, H.: MPTP-delta, a putative murine homolog of HPTP-delta, is expressed in specialized regions of the brain and in the B-cell lineage. *Molec. Cell. Biol.* 13: 5513–5523, 1993.

[75913] 14243. Uetani, N.; Kato, K.; Ogura, H.; Mizuno, K.; Kawano, K.; Mikoshiba, K.; Yakura, H.; Asano, M.; Iwakura, Y.: Impaired learning with enhanced hippocampal long-term potentiation in PTP-delta-deficient mice. *EMBO J.* 19: 2775–2785, 2000.

[75914] 14244. Heighway, J.; Betticher, D. C.; Hoban, P. R.; Altermatt, H. J.; Cowen, R.: Coamplification in tumors of KRAS2, type 2 inositol 1,4,5 triphosphate receptor gene, and a novel human gene, KRAG. *Genomics* 35:207–214, 1996.

[75915] 14245. Scott, A. F.; Elizaga, A.; Morrell, J.; Bergen, A.; Penno, M. B.: Characterization of a gene coamplified with Ki-ras in Y1 murine adrenal carcinoma cells that codes for



a putative membrane protein. *Genomics* 20:227–230, 1994.

[75916] 14246. Monte, D.; Baert, J. L.; Defossez, P. A.; de Launoit, Y.; Stehelin, D.: Molecular cloning and characterization of human ERM, a new member of the Ets family closely related to mouse PEA3 and ER81 transcription factors. *Oncogene* 9: 1397–1406, 1994.

[75917] 14247. Monte, D.; Coutte, L.; Dewitte, F.; Defossez, P.-A.; Le Coniat, M.; Stehelin, D.; Berger, R.; de Launoit, Y.: Genomic organization of the human ERM (ETV5) gene, a PEA3 group member of ETS transcription factors. *Genomics* 35: 236–240, 1996.

[75918] 14248. Protopopova, M. V.; Vorobieva, N. V.; Protopopov, A. I.; Gizatullin, R. Z.; Kashuba, V. I.; Klein, G.; Zabarovsky, E. R.; Graphodatsky, A. S.: Assignment of the ERM gene (ETV5) coding for the ets-related protein to human chromosome band 3q28 by in situ hybridization. *Cytogenet. Cell Genet.* 74: 220 only, 1996.

[75919] 14249. Bosher, J. M.; Williams, T.; Hurst, H. C.: The developmentally regulated transcription factor AP-2 is involved in c-erbB-2 overexpression in human mammary carcinoma. *Proc. Nat. Acad. Sci.* 92: 744–747, 1995.

[75920] 14250. Combadiere, C.; Ahuja, S. K.; Murphy, P. M.:

Cloning, chromosomal localization, and RNA expression of a human beta chemokine receptor-like gene. *DNA Cell Biol.* 14: 673–680, 1995.

- [75921] 14251. Combadiere, C.; Ahuja, S. K.; Van Damme, J.; Tiffany, H. L.; Gao, J.-L.; Murphy, P. M.: Monocyte chemoattractant protein-3 is a functional ligand for CC chemokine receptors 1 and 2. *J. Biol. Chem.* 270:29671–29675, 1995.
- [75922] 14252. Doranz, B. J.; Rucker, J.; Yi, Y.; Smyth, R. J.; Samson, M.; Peiper, S. C.; Parmentier, M.; Collman, R. G.; Doms, R. W.: A dual-tropic primary HIV-1 isolate that uses fusin and the beta-chemokine receptors CKR-5, CKR-3, and CKR-2b as fusion cofactors. *Cell* 85: 1149–1158, 1996.
- [75923] 14253. Mummidi, S.; Ahuja, S. S.; Gonzalez, E.; Anderson, S. A.; Santiago, E. N.; Stephan, K. T.; Craig, F. E.; O'Connell, P.; Tryon, V.; Clark, R. A.; Dolan, M. J.; Ahuja, S. K.: Genealogy of the CCR5 locus and chemokine system gene variants associated with altered rates of HIV-1 disease progression. *Nature Med.* 4: 786–793, 1998.
- [75924] 14254. Peters, W.; Dupuis, M.; Charo, I. F.: A mechanism for the impaired IFN-gamma production in C-C chemokine receptor 2 (CCR2) knockout mice: role of CCR2 in linking

the innate and adaptive immune responses. *J.Immun.* 165: 7072–7077, 2000.

- [75925] 14255.Peters, W.; Scott, H. M.; Chambers, H. F.; Flynn J. L.; Charo,I. F.; Ernst, J. D.: Chemokine receptor 2 serves an early and essential role in resistance to *Mycobacterium tuberculosis*. *Proc. Nat. Acad.Sci.* 98: 7958–7963, 2001.
- [75926] 14256.Samson, M.; Labbe, O.; Mollereau, C.; Vassart, G.; Parmentier,M.: Molecular cloning and functional expression of a new human CC–chemokine receptor gene. *Biochemistry* 35: 3362–3367, 1996.
- [75927] 14257.Samson, M.; Soularue, P.; Vassart, G.; Parmentier, M.: The genes encoding the human CC–chemokine receptors CC–CCR1 to CC–CCR5 (CMKBR1–CMKBR5) are clustered in the p21.3–p24 region of chromosome 3. *Genomics* 36:522–526, 1996.
- [75928] 14258.Sanders, S. K.; Crean, S. M.; Boxer, P. A.; Kellner, D.; LaRosa,G. J.; Hunt, S. W., III.: Functional differences between monocyte chemotactic protein–1 receptor A and monocyte chemotactic protein–1 receptor B expressed in a Jurkat T cell. *J. Immun.* 165: 4877–4883, 2000.
- [75929] 14259.Smith, M. W.; Dean, M.; Carrington, M.; Winkler, C.; Huttley,G. A.; Lomb, D. A.; Goedert, J. J.; O'Brien, T. R.; Jacobson, L. P.; Kaslow, R.; Buchbinder, S.; Vittinghoff, E.;

Vlahov, D.; Hoots, K.; Hilgartner, M. W.; Hemophilia Growth and Development Study (HGDS); Multicenter AIDS Cohort Study (MACS); Multicenter Hemophilia Cohort Study (MHCS); San Francisco City Cohort (SFCC); ALIVE Study; O'Brien, S. J.: Contrasting genetic influence of CCR2 and CCR5 variant on HIV-1 infection and disease progression. *Science* 277: 959-965, 1997.

[75930] 14260. Wong, L.-M.; Myers, S. J.; Tsou, C.-L.; Gosling, J.; Arai, H.; Charo, I. F.: Organization and differential expression of the human monocyte chemoattractant protein 1 receptor gene: evidence for the role of the carboxyl-terminal tail in receptor trafficking. *J. Biol. Chem.* 272: 1038-1045, 1997.

[75931] 14261. Donald, L. J.; Wang, H. S.; Hamerton, J. L.: Assignment of the gene for cystathionase (CYS) to human chromosome 16. (Abstract) *Cytogenet. Cell Genet.* 32: 268 only, 1982.

[75932] 14262. Frimpter, G. W.: Cystathioninuria: nature of the defect. *Science* 149: 1095-1096, 1965.

[75933] 14263. Frimpter, G. W.: Cystathioninuria, sulfite oxidase deficiency, and  $\beta$ -mercaptolactate-cysteine disulfiduria. In: Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: *The Metabolic Basis of Inherited Disease*. New York:

McGraw-Hill (pub.) (3rd ed.): 1972. Pp. 413-425.

- [75934] 14264.Frimpter, G. W.; Haymovitz, A.; Horwith, M.: Cystathioninuria. NewEng. J. Med. 268: 333-339, 1963.
- [75935] 14265.Harris, H.; Penrose, L. S.; Thomas, D. H. H.: Cystathioninuria. Ann.Hum. Genet. 23: 442-453, 1959.
- [75936] 14266.Lu, Y.; O'Dowd, B. F.; Orrego, H.; Israel, Y.: Cloning and nucleotidesequence of human liver cDNA encoding for cystathionine gamma-lyase. Biochem.Biophys. Res. Commun. 189: 749-758, 1992.
- [75937] 14267.Lyon, I. C. T.; Procopis, P. G.; Turner, B.: Cystathioninuriain a well baby population. Acta Paediat. Scand. 60: 324-328, 1971.
- [75938] 14268.Mongeau, J.-G.; Hilgartner, M.; Worthen, H. G.; Frimpter, G. W.: Cystathioninuria: study of an infant with normal mentality, thrombocytopenia,and renal calculi. J. Pediat. 69: 1113-1120, 1967.
- [75939] 14269.Pascal, T. A.; Gaull, G. E.; Beratis, N. G.; Gillam, B. M.; Tallan,H. H.: Cystathionase deficiency: evidence for genetic heterogeneityin primary cystathioninuria. Pediat. Res. 12: 125-133, 1978.
- [75940] 14270.Perry, T. L.; Hardwick, D. F.; Hansen, S.; Love, D. L.; Israels,S.: Cystathioninuria in two healthy siblings. New Eng. J. Med. 278:590-592, 1968.

- [75941] 14271.Schneiderman, L. J.: Latent cystathioninuria. *J. Med. Genet.* 4:260–263, 1967.
- [75942] 14272.Scott, C. R.; Dassell, S. W.; Clark, S. H.; Chiang-Teng, C.; Swedberg,K. R.: Cystathioninemia: a benign genetic condition. *J. Pediat.* 76:571–577, 1970.
- [75943] 14273.Shaw, K. N. F.; Lieberman, E.; Koch, R.; Donnell, G. N.: Cystathioninuria. *Am.J. Dis. Child.* 113: 119–128, 1967.
- [75944] 14274.Tada, K.; Yoshida, T.; Yokoyama, Y.; Sato, T.; Nakagawa, H.; Arakawa,T.: Cystathioninuria not associated with vitamin B6 dependency: aprobably new type of cystathioninuria. *Tohoku J. Exp. Med.* 95: 235–242,1968.
- [75945] 14275.Whelan, D. T.; Scriver, C. R.: Cystathioninuria and renal iminoglycinuria in a pedigree: a perspective on counseling. *New Eng. J. Med.* 278:924–927, 1968.
- [75946] 14276.Joenje, H.: Fanconi anaemia complementation groups in Germanyand The Netherlands. *Hum. Genet.* 97: 280–282, 1996.
- [75947] 14277.Joenje, H.; Oostra, A. B.; Wijker, M.; di Summa, F. M.; van Berkel,C. G. M.; Rooimans, M. A.; Ebell, W.; van Weel, M.; Pronk, J. C.;Buchwald, M.; Arwert, F.: Evidence for at least eight Fanconi anemiagenes. *Am. J. Hum. Genet.* 61: 940–944, 1997.

- [75948] 14278.Bongiovanni, A. M.: Disorders of adrenocortical steroid biogenesis.In:Stanbury, J. B.; Wyngaarden, J. B.; Fredrickson, D. S.: The MetabolicBasis of Inherited Disease. New York: McGraw-Hill (pub.) (3rded.): 1972. Pp. 857-885.
- [75949] 14279.Bongiovanni, A. M.; Root, A. W.: The adrenogenital syndrome. NewEng. J. Med. 268: 1283-1289; 1342-1351; 1391-1399, 1963.
- [75950] 14280.Boudreaux, D.; Waisman, J.; Skinner, D. G.; Low, R.: Giant adrenalmyelolipoma and testicular interstitial cell tumor in a man with congenital21-hydroxylase deficiency. Am. J. Surg. Path. 3: 109-123, 1979.
- [75951] 14281.Carroll, M. C.; Campbell, R. D.; Porter, R. R.: Mapping of steroid21-hydroxylase genes adjacent to complement component C4 genes inHLA, the major histocompatibility complex in man. Proc. Nat. Acad.Sci. 82: 521-525, 1985.
- [75952] 14282.Chaplin, D. D.; Galbraith, L. J.; Seidman, J. G.; White, P. C.;Parker, K. L.: Nucleotide sequence analysis of murine 21-hydroxylasegene: mutations affecting gene expression. Proc. Nat. Acad. Sci. 83:9601-9605, 1986.
- [75953] 14283.Charmandari, E.; Hindmarsh, P. C.; Johnston, A.; Brook, C. G.D.: Congenital adrenal hyperplasia due to

21-hydroxylase deficiency:alterations in cortisol pharmacokinetics at puberty. J. Clin. Endocr.Metab. 86: 2701-2708, 2001.

[75954] 14284.Childs, B.; Grumbach, M. M.; Van Wyk, J. J.: Virilizing adrenalhyperplasia: genetic and hormonal studies. J. Clin. Invest. 35:213-222, 1956.

[75955] 14285.Chiou, S.-H.; Hu, M.-C.; Chung, B.: A missense mutation at ile172-to-asnor arg356-to-trp causes steroid 21-hydroxylase deficiency. J. Biol.Chem. 265: 3549-3552, 1990.

[75956] 14286.Chrousos, G. P.; Loriaux, D. L.; Mann, D. L.; Cutler, G. B., Jr.: Late-onset 21-hydroxylase deficiency mimicking idiopathic hirsutismor polycystic ovarian disease: an allelic variant of congenital virilizingadrenal hyperplasia with a milder enzymatic defect. Ann. Intern.Med. 96: 143-148, 1982.

[75957] 14287.Collier, S.; Tassabehji, M.; Strachan, T.: A de novo pathologicalpoint mutation at the 21-hydroxylase locus: implications for geneconversion in the human genome. Nature Genet. 3: 260-265, 1993.

[75958] 14288.Cooper, D. Y.; Levin, S.; Narasimhulu, S.; Rosenthal, O.; Estabrook,R. W.: Photochemical action spectrum of the terminal oxidase of mixedfunction oxidase sys-



tems. Science 147: 400–402, 1965.

[75959] 14289.Cornean, R. E.; Hindmarsh, P. C.; Brook, C. G. D.: Obesity in 21-hydroxylase deficient patients. Arch. Dis. Child. 78: 261–263, 1998.

[75960] 14290.Couillin, P.; Kottler–Missonnier, M. L.; Grisard, M. C.; Hors, J.; Feingold, J.; Boue, J.; Boue, A.: HLA–A, B, C, DR alleles in congenital adrenal hyperplasia. Hum. Genet. 53: 389–392, 1980.

[75961] 14291.Cutfield, R. G.; Bateman, J. M.; Odell, W. D.: Infertility caused by bilateral testicular masses secondary to congenital adrenal hyperplasia (21-hydroxylase deficiency). Fertil. Steril. 40: 809–814, 1983.

[75962] 14292.Cutler, G. B., Jr.; Laue, L.: Congenital adrenal hyperplasia due to 21-hydroxylase deficiency. New Eng. J. Med. 323: 1806–1813, 1990.

[75963] 14293.Dacou–Voutetakis, C.; Dracopoulou, M.: High incidence of molecular defects of the CYP21 gene in patients with premature adrenarche. J. Clin. Endocr. Metab. 84: 1570–1574, 1999.

[75964] 14294.Day, D. J.; Speiser, P. W.; Schulze, E.; Bettendorf, M.; Fitness, J.; Barany, F.; White, P. C.: Identification of non-amplifying CYP21 genes when using PCR-based diagnosis of 21-hydroxylase deficiency in congenital adrenal

hyperplasia (CAH) affected pedigrees. Hum.Molec. Genet. 5: 2039–2048, 1996.

[75965] 14295.Deneux, C.; Tardy, V.; Dib, A.; Mornet, E.; Billaud, L.; Charron,D.; Morel, Y.; Kuttenn, F.: Phenotype–genotype correlation in 56women with nonclassical congenital adrenal hyperplasia due to 21–hydroxylasedeficiency. J. Clin. Endocr. Metab. 86: 207–213, 2001.

[75966] 14296.Donohoue, P. A.; Jospe, N.; Migeon, C. J.; Van Dop, C.: Two distinctareas of unequal crossingover within the steroid 21–hydroxylase genesproduce absence of CYP21B. Genomics 5: 397–406, 1989.

[75967] 14297.Dracopoulou–Vabouli, M.; Maniati–Christidi, M.; Dacou–Voutetakis,C.: The spectrum of molecular defects of the CYP21 gene in the Hellenicpopulation: variable concordance between genotype and phenotype inthe different forms of congenital adrenal hyperplasia. J. Clin. Endocr.Metab. 86: 2845–2848, 2001.

[75968] 14298.Duck, S. C.: Malignancy associated with congenital adrenal hyperplasia. J.Pediat. 99: 423–424, 1981.

[75969] 14299.Dumic, M.; Brkljadic, L.; Mardesic, D.; Plavsic, V.; Lukenda,M.; Kastelan, A.: 'Cryptic' form of congenital adrenal hyperplasiadue to 21–hydroxylase deficiency in the Yugoslav population. ActaEndocr. 109: 386–392,

1985.

- [75970] 14300. Dupont, B.; Smithwick, E. M.; Oberfield, S. E.; Lee, T. D.; Levine, L. S.: Close genetic linkage between HLA and congenital adrenal hyperplasia (21-hydroxylase deficiency). *Lancet* II: 1309–1312, 1977.
- [75971] 14301. Ezquieta, B.; Oliver, A.; Gracia, R.; Gancedo, P. G.: Analysis of steroid 21-hydroxylase gene mutations in the Spanish population. *Hum. Genet.* 96: 198–204, 1995.
- [75972] 14302. Ferenczi, A.; Garami, M.; Kiss, E.; Pek, M.; Sasvari-Szekely, M.; Barta, C.; Staub, M.; Solyom, J.; Fekete, G.: Screening for mutations of 21-hydroxylase gene in Hungarian patients with congenital adrenal hyperplasia. *J. Clin. Endocr. Metab.* 84: 2369–2372, 1999.
- [75973] 14303. Fitness, J.; Dixit, N.; Webster, D.; Torresani, T.; Pergolizzi, R.; Speiser, P. W.; Day, D. J.: Genotyping of CYP21, linked chromosome 6p markers, and a sex-specific gene in neonatal screening for congenital adrenal hyperplasia. *J. Clin. Endocr. Metab.* 84: 960–966, 1999.
- [75974] 14304. Fleischnick, E.; Awdeh, Z. L.; Raum, D.; Granados, J.; Alosco, S. M.; Crigler, J. F., Jr.; Gerald, P. S.; Giles, C. M.; Yunis, E. J.; Alper, C. A.: Extended MHC haplotypes in 21-hydroxylase-deficiency congenital adrenal hyperplasia: shared genotypes in unrelated patients. *Lancet* I: 152–156,

1983.

- [75975] 14305. Flori, J.; Tongio, M. M.; Kurtz, F.; Mayer, S.; Juif, J. G.: Liaison entre le gene responsable de l'hyperplasie sur-renalienne par deficit en 21 hydroxylase et les marqueurs du bras court du chromosome 6. *J. Genet. Hum.* 28: 185-193, 1980.
- [75976] 14306. Forest, M. G.; Betuel, H.; David, M.: Prenatal treatment in congenital adrenal hyperplasia due to 21-hydroxylase deficiency: up-date 88 of the French multicentric study. *Endocr. Res.* 15: 277-301, 1989.
- [75977] 14307. Galal, O. M.; Rudd, B. T.; Drayer, N. M.: Evaluation of deficiency of 21-hydroxylation in patients with congenital adrenal hyperplasia. *Arch. Dis. Child.* 43: 410-414, 1968.
- [75978] 14308. Ghanem, N.; Lobaccaro, J. M.; Buresi, C.; Abbal, M.; Halaby, G.; Sultan, C.; Lefranc, G.: Defective, deleted or converted CYP21B gene and negative association with a rare restriction fragment length polymorphism allele of the factor B gene in congenital adrenal hyperplasia. *Hum. Genet.* 86: 117-125, 1990.
- [75979] 14309. Rassool, F. V.; Le Beau, M. M.; Shen, M.-L.; Neilly, M. E.; Espinosa, R., III; Ong, S. T.; Boldog, F.; Drabkin, H.; McCarroll, R.; McKeithan, T. W.: Direct cloning of DNA se-

quences from the common fragile site region at chromosome band 3p14.2. *Genomics* 35: 109–117, 1996.

[75980] 14310. Rudduck, C.; Franzen, G.: A new heritable fragile site on human chromosome 3. *Hereditas* 98: 297–299, 1983.

[75981] 14311. Shi, Y.; Zou, M.; Farid, N. R.; Paterson, M. C.: Association of FHIT (fragile histidine triad), a candidate tumour suppressor gene, with the ubiquitin–conjugating enzyme hUBC9. *Biochem. J.* 352: 443–448, 2000.

[75982] 14312. Shiraishi, T.; Druck, T.; Mimori, K.; Flomenberg, J.; Berk, L.; Alder, H.; Miller, W.; Huebner, K.; Croce, C. M.: Sequence conservation at human and mouse orthologous common fragile regions, FRA3B/FHIT and Fra14A2/Fhit. *Proc. Nat. Acad. Sci.* 98: 5722–5727, 2001.

[75983] 14313. Sozzi, G.; Veronese, M. L.; Negrini, M.; Baffa, R.; Cotticelli, M. G.; Inoue, H.; Torielli, S.; Pilotti, S.; De Gregorio, L.; Pastorino, U.; Pierotti, M. A.; Ohta, M.; Huebner, K.; Croce, C. M.: The FHIT gene at 3p14.2 is abnormal in lung cancer. *Cell* 85: 17–26, 1996.

[75984] 14314. Stein, C. K.; Glover, T. W.; Palmer, J. L.; Glisson, B. S.: Direct correlation between FRA3B expression and cigarette smoking. *Genes Chromosomes Cancer* 34: 333–340, 2002.

- [75985] 14315. Virgilio, L.; Shuster, M.; Gollin, S. M.; Veronese, M. L.; Ohta, M.; Huebner, K.; Croce, C. M.: FHIT gene alterations in head and neck squamous cell carcinomas. *Proc. Nat. Acad. Sci.* 93: 9770–9775, 1996.
- [75986] 14316. Wegner, R.-D.: A new inducible fragile site on chromosome 3 (p14.2) in human lymphocytes. *Hum. Genet.* 63: 297–298, 1983.
- [75987] 14317. Wegner, R.-D.: Reply to the letter of A. Markkanen, S. Knuutila, and A. de la Chapelle. (Letter) *Hum. Genet.* 65: 218 only, 1983.
- [75988] 14318. Zanesi, N.; Fidanza, V.; Fong, L. Y.; Mancini, R.; Druck, T.; Valtieri, M.; Rudiger, T.; McCue, P. A.; Croce, C. M.; Huebner, K.: The tumor spectrum in FHIT-deficient mice. *Proc. Nat. Acad. Sci.* 98: 10250–10255, 2001.
- [75989] 14319. Nagase, T.; Seki, N.; Ishikawa, K.; Ohira, M.; Kawarabayashi, Y.; Ohara, O.; Tanaka, A.; Kotani, H.; Miyajima, N.; Nomura, N.: Prediction of the coding sequences of unidentified human genes. VI. The coding sequences of 80 new genes (KIAA0201–KIAA0280) deduced by analysis of cDNA clones from cell line KG-1 and brain. *DNA Res.* 3: 321–329, 1996.
- [75990] 14320. Palfree, R. G. E.; Sadro, L. C.; Solomon, S.: The gene encoding the human corticostatin HP-4 precursor contains

a recent 86-base duplication and is located on chromosome 8. *Molec. Endocr.* 7: 199–205, 1993.

- [75991] 14321. Singh, A.; Bateman, A.; Zhu, Q. Z.; Shimasaki, S.; Esch, F.; Solomon, S.: Structure of the novel human granulocyte peptide with anti-ACTH activity. *Biochem. Biophys. Res. Commun.* 155: 524–529, 1988.
- [75992] 14322. Lee, E.; Marcucci, M.; Daniell, L.; Pypaert, M.; Weisz, O. A.; Ochoa, G.-C.; Farsad, K.; Wenk, M. R.; De Camilli, P.: Amphiphysin2 (Bin1) and T-tubule biogenesis in muscle. *Science* 297: 1193–1196, 2002.
- [75993] 14323. Negorev, D.; Riethman, H.; Wechsler-Reya, R.; Sakamuro, D.; Prendergast, G. C.; Simon, D.: The Bin1 gene localizes to human chromosome 2q14 by PCR analysis of somatic cell hybrids and fluorescence in situ hybridization. *Genomics* 33: 329–331, 1996.
- [75994] 14324. Sakamuro, D.; Elliott, K. J.; Wechsler-Reya, R.; Prendergast, G. C.: BIN1 is a novel MYC-interacting protein with features of a tumour suppressor. *Nature Genet.* 14: 69–77, 1996.
- [75995] 14325. Ishizuka, T.; Ahmad, I.; Kita, K.; Sonoda, T.; Ishijima, S.; Sawa, K.; Suzuki, N.; Tatibana, M.: The human phosphoribosylpyrophosphate synthetase-associated protein 39 gene (PRPSAP1) is located in the chromosome re-

gion 17q24–q25. *Genomics* 33: 332–334, 1996.

[75996] 14326. Ishizuka, T.; Kita, K.; Sonoda, T.; Ishijima, S.; Sawa, K.; Suzuki, N.; Tatibana, M.: Cloning and sequencing of human complementary DNA for the phosphoribosylpyrophosphate synthetase-associated protein 39. *Biochim. Biophys. Acta* 1306: 27–30, 1996.

[75997] 14327. Kita, K.; Ishizuka, T.; Ishijima, S.; Sonoda, T.; Tatibana, M.: A novel 39-kDa phosphoribosylpyrophosphate synthetase-associated protein of rat liver: cloning, high sequence similarity to the catalytic subunits, and a negative regulatory role. *J. Biol. Chem.* 269: 8334–8340, 1994.

[75998] 14328. Endo, Y.; Sato, Y.; Matsushita, M.; Fujita, T.: Cloning and characterization of the human lectin P35 gene and its related gene. *Genomics* 36: 515–521, 1996.

[75999] 14329. Wayne, S.; Robertson, N. G.; DeClau, F.; Chen, N.; Verhoeven, K.; Prasad, S.; Tranebjarg, L.; Morton, C. C.; Ryan, A. F.; Van Camp, G.; Smith, R. J. H.: Mutations in the transcriptional activator EYA4 cause late-onset deafness at the DFNA10 locus. *Hum. Molec. Genet.* 10: 195–200, 2001.

[76000] 14330. Kanai, Y.; Miura, K.; Uehara, T.; Amagai, M.; Takeda, O.; Tanuma, S.; Kurosawa, Y.: Natural occurrence of Nuc in the sera of autoimmune-prone MRL/lpr mice.



Biochem. Biophys. Res. Commun. 196: 729–736, 1993.

[76001] 14331.Miura, K.; Hirai, M.; Kanai, Y.; Kurosawa, Y.: Organization of the human gene for nucleobindin (NUC) and its chromosomal assignment to 19q13.2–q13.4. Genomics 34: 181–186, 1996.

[76002] 14332.Miura, K.; Titani, K.; Kurosawa, Y.; Kanai, Y.: Molecular cloning of nucleobindin, a novel DNA-binding protein that contains both a signal peptide and a leucine zipper structure. Biochem. Biophys. Res. Commun. 187: 375–380, 1992.

[76003] 14333.Connelly, M. A.; Grady, R. C.; Mushinski, J. F.; Marcu, K. B.: PANG, a gene encoding a neuronal glycoprotein, is ectopically activated by intracisternal A-type particle long terminal repeats in murine plasmacytomas. Proc. Nat. Acad. Sci. 91: 1337–1341, 1994.

[76004] 14334.Mock, B. A.; Connelly, M. A.; McBride, O. W.; Kozak, C. A.; Marcu, K. B.: Plasmacytoma-associated neuronal glycoprotein, Pang, maps to mouse chromosome 6 and human chromosome 3. Genomics 34: 226–228, 1996.

[76005] 14335.Noben-Trauth, K.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N.A.; Sonoda, G.; Testa, J. R.; Klempnauer, K.-H.: Mybl2 (Bmyb) maps to mouse chromosome 2 and human chromosome 20q13.1. Genomics 35: 610–612, 1996.

- [76006] 14336.Ando, A.; Kikuti, Y. Y.; Shigenari, A.; Kawata, H.; Okamoto, N.; Shiina, T.; Chen, L.; Ikemura, T.; Abe, K.; Kimura, M.; Inoko, H.: cDNA cloning of the human homologues of the mouse Ke4 and Ke6 genes at the centromeric end of the human MHC region. *Genomics* 35: 600–602, 1996.
- [76007] 14337.Aziz, N.; Maxwell, M. M.; Brenner, B. M.: Coordinate regulation of 11-beta-HSD and Ke6 gene in cpk mouse: implications for steroid metabolic defect in PKD. *Am. J. Physiol.* 267: F791–F797, 1994.
- [76008] 14338.Aziz, N.; Maxwell, M. M.; St.-Jacques, B.; Brenner, B. M.: Downregulation of Ke 6, a novel gene encoded within the major histocompatibility complex, in murine polycystic kidney disease. *Molec. Cell. Biol.* 13:1847–1853, 1993.
- [76009] 14339.Kikuti, Y. Y.; Tamiya, G.; Ando, A.; Chen, L.; Kimura, M.; Ferreira, E.; Tsuji, K.; Trowsdale, J.; Inoko, H.: Physical mapping 220 kb centromeric of the human MHC and DNA sequence analysis of the 43-kb segment including the RING1, HKE6, and HKE4 genes. *Genomics* 42:422–435, 1997.
- [76010] 14340.Aziz, N.; Anderson, E.; Lee, G. Y.; Woo, D. D. L.: Arrested testis development in the cpk mouse may be the

result of abnormal steroid metabolism. *Molec. Cell. Endocr.* 171: 83–88, 2001.

[76011] 14341. Fomitcheva, J.; Baker, M. E.; Anderson, E.; Lee, G. Y.; Aziz, N.: Characterization of Ke 6, a new 17-beta-hydroxysteroid dehydrogenase, and its expression in gonadal tissues. *J. Biol. Chem.* 273: 22664–22671, 1998.

[76012] 14342. Cheung, A. H.; Stewart, R. J.; Marsden, P. A.: Endothelial Tie2/Tek ligands angiopoietin-1 (ANGPT1) and angiopoietin-2 (ANGPT2): regional localization of the human genes to 8q22.3–q23 and 8p23. *Genomics* 48:389–391, 1998.

[76013] 14343. Davis, S.; Aldrich, T. H.; Jones, P. F.; Acheson, A.; Compton, D. L.; Jain, V.; Ryan, T. E.; Bruno, J.; Radziejewski, C.; Maisonpierre, P. C.; Yancopoulos, G. D.: Isolation of angiopoietin-1, a ligand for the TIE2 receptor, by secretion-trap expression cloning. *Cell* 87:1161–1169, 1996.

[76014] 14344. Folkman, J.; D'Amore, P. A.: Blood vessel formation: what is its molecular basis? *Cell* 87: 1153–1155, 1996.

[76015] 14345. Marziliano, N.; Crovella, S.; Audero, E.; Pecile, V.; Bussolino, F.; Amoroso, A.; Garagna, S.: Genetic mapping of the mouse homologue of the human angiopoietin-1 gene (Agpt) to mouse chromosome 9E2 by in situ hy-

bridization. Cytogenet. Cell Genet. 87: 199–200, 1999.

- [76016] 14346.Sato, T. N.; Tozawa, Y.; Deutsch, U.; Wolburg–Buchholz, K.; Fujiwara,Y.; Gendron–Maguire, M.; Gridley, T.; Wolburg, H.; Risau, W.; Qin,Y.: Distinct roles of the receptor tyrosine kinases Tie–1 and Tie–2in blood vessel formation. Nature 376: 70–73, 1995.
- [76017] 14347.Suri, C.; Jones, P. F.; Patan, S.; Bartunkova, S.; Maisonpierre,P. C.; Davis, S.; Sato, T. N.; Yancopoulos, G. D.: Requisite roleof angiopoietin–1, a ligand for the TIE2 receptor, during embryonicangiogenesis. Cell 87: 1171–1180, 1996.
- [76018] 14348.Suri, C.; McClain, J.; Thurston, G.; McDonald, D. M.; Zhou, H.;Oldmixon, E. H.; Sato, T. N.; Yancopoulos, G. D.: Increased vascularizationin mice overexpressing angiopoietin–1. Science 282: 468–471, 1998.
- [76019] 14349.Valenzuela, D. M.; Griffiths, J. A.; Rojas, J.; Aldrich, T. H.;Jones, P. F.; Zhou, H.; McClain, J.; Copeland, N. G.; Gilbert, D.J.; Jenkins, N. A.; Huang, T.; Papadopoulos, N.; Maisonpierre, P.C.; Davis, S.; Yancopoulos, G. D.: Angiopoietins 3 and 4: diverginggene counterparts in mice and humans. Proc. Nat. Acad. Sci. 96:1904–1909, 1999.
- [76020] 14350.Chestkov, A. V.; Baka, I. D.; Kost, M. V.; Georgiev, G. P.; Buchman,V. L.: The d4 gene family in the human

genome. Genomics 36: 174–177,1996.

- [76021] 14351.Gabig, T. G.; Crean, C. D.; Klenk, A.; Long, H.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Quincey, D.; Parente, F.; Lespinasse, F.; Carle, G. F.; Gaudray, P.; and 13 others: Expression and chromosomal localization of the Requiem gene. Mammalian Genome 9: 660–665, 1998.
- [76022] 14352.Gabig, T. G.; Mantel, P. L.; Rosli, R.; Crean, C. D.: Requiem: a novel zinc finger gene essential for apoptosis in myeloid cells. J. Biol. Chem. 269: 29515–29519, 1994.
- [76023] 14353.Smith, J. E., Jr.; Bollekens, J. A.; Inghirami, G.; Takeshita, K.: Cloning and mapping of the MEIS1 gene, the human homolog of murine leukemogenic gene. Genomics 43: 99–103, 1997.
- [76024] 14354.Duperat, V. G.; Jacquelin, B.; Boisseau, P.; Arveiler, B.; Nurden, A. T.: Protease-activated receptor genes are clustered on 5q13. Blood 92:25–31, 1998.
- [76025] 14355.Ishihara, H.; Connolly, A. J.; Zeng, D.; Kahn, M. L.; Zheng, Y. W.; Timmons, C.; Tram, T.; Coughlin, S. R.: Protease-activated receptor3 is a second thrombin receptor in humans. Nature 386: 502–506, 1997.
- [76026] 14356.Kahn, M. L.; Zheng, Y. W.; Huang, W.; Bigornia, V.; Zeng, D.; Moff, S.; Farese, R. V., Jr.; Tam, C.; Coughlin, S. R.: A dual thrombin receptor system for platelet activation.

Nature 394: 690–694, 1998.

[76027] 14357.Nakanishi–Matsui, M.; Zheng, Y.–W.; Sulciner, D. J.; Weiss, E.J.; Ludeman, M. J.; Coughlin, S. R.: PAR3 is a co–factor for PAR4activation by thrombin. Nature 404: 609–613, 2000.

[76028] 14358.Sambrano, G. R.; Weiss, E. J.; Zheng, Y.–W.; Huang, W.; Coughlin,S. R.: Role of thrombin signalling in platelets in haemostasis andthrombosis. Nature 413: 74–78, 2001.

[76029] 14359.Schmidt, V.; Wainer, J.; Bahou, W.: Identification and characterizationof the human proteolytically activated receptor–3 (PAR–3) gene withinthe PAR gene cluster by positional mapping. (Abstract) Blood (suppl.1) 90: 284a only, 1997.

[76030] 14360.Paulweber, B.; Wiebusch, H.; Miesenboeck, G.; Funke, H.; Assmann,G.; Hoelzl, B.; Sippl, M. J.; Friedl, W.; Patsch, J. R.; Sandhofer,F.: Molecular basis of lipoprotein lipase deficiency in two Austrianfamilies with type I hyper–lipoproteinemia. Atherosclerosis 86: 239–250,1991.

[76031] 14361.Abe, M.; Suzuki, T.; Deguchi, T.: An improved method for genotypingof N–acetyltransferase polymorphism by polymerase chain reaction. Jpn.J. Hum. Genet. 38: 163–168, 1993.

[76032] 14362.Azad Khan, A. K.; Nurazzaman, M.; Truelove, S. C.:

The effect of the acetylator phenotype on the metabolism of sulphasalazine in man. *J. Med. Genet.* 20: 30–36, 1983.

- [76033] 14363. Bell, D. A.; Taylor, J. A.; Butler, M. A.; Stephens, E. A.; Wiest, J.; Brubaker, L. H.; Kadlubar, F. F.; Lucier, G. W.: Genotype/phenotype discordance for human arylamine N-acetyltransferase (NAT2) reveals a new slow-acetylator allele common in African-Americans. *Carcinogenesis* 14:1689–1692, 1993.
- [76034] 14364. Blum, M.; Grant, D.; Demierre, A.; Meyer, U. A.: N-acetylation pharmacogenetics: a gene deletion causes absence of arylamine N-acetyltransferase in liver of slow acetylator rabbits. *Proc. Nat. Acad. Sci.* 86: 9554–9557, 1989.
- [76035] 14365. Burrows, A. W.; Hockaday, T. D. R.; Mann, J. I.; Taylor, J. G.: Diabetic dimorphism according to acetylator status. *Brit. Med. J.* 1: 208–210, 1978.
- [76036] 14366. Cascorbi, I.; Brockmoller, J.; Bauer, S.; Reum, T.; Roots, I.: NAT2\*12A (803A–G) codes for rapid arylamine N-acetylation in humans. *Pharmacogenetics* 6:257–259, 1996.
- [76037] 14367. Cascorbi, I.; Drakoulis, N.; Brockmoller, J.; Maurer, A.; Sperling, K.; Roots, I.: Arylamine N-acetyltransferase (NAT2) mutations and their allelic linkage in unrelated

Caucasian individuals: correlation with phenotypic activity.  
Am. J. Hum. Genet. 57: 581–592, 1995.

- [76038] 14368. Delomenie, C.; Sica, L.; Grant, D. M.; Krishnamoorthy, R.; Dupret, J.-M.: Genotyping of the polymorphic N-acetyltransferase (NAT2\*) gene locus in two native African populations. Pharmacogenetics 6: 177–185, 1996.
- [76039] 14369. Evans, D. A. P.: N-acetyltransferase. Pharm. Therap. 42: 157–234, 1989.
- [76040] 14370. Evans, D. A. P.; Manley, K. A.; McKusick, V. A.: Genetic control of isoniazid metabolism in man. Brit. Med. J. 2: 485–491, 1960.
- [76041] 14371. Evans, D. A. P.; White, T. A.: Human acetylation polymorphism. J. Lab. Clin. Med. 63: 394–403, 1964.
- [76042] 14372. Frymoyer, J. W.; Jacox, R. F.: Studies of genetically controlled sulfadiazine acetylation in rabbit livers: possible identification of the heterozygous trait. J. Lab. Clin. Med. 62: 905–909, 1963.
- [76043] 14373. Glowinski, I. B.; Weber, W. W.: Biochemical characterization of genetically variant aromatic amine N-acetyltransferases in A/J and C57BL/6J mice. J. Biol. Chem. 257: 1431–1437, 1982.
- [76044] 14374. Glowinski, I. B.; Weber, W. W.: Genetic regulation of aromatic amine N-acetylation in inbred mice. J. Biol. Chem.



257: 1424–1430, 1982.

- [76045] 14375. Grant, D. M.; Morike, K.; Eichelbaum, M.; Meyer, U. A.: Acetylation pharmacogenetics: the slow acetylator phenotype is caused by decreased or absent arylamine N-acetyltransferase in human liver. *J. Clin. Invest.* 85: 968–972, 1990.
- [76046] 14376. Grant, D. M.; Tang, B. K.; Kalow, W.: A simple test for acetylator phenotype using caffeine. *Brit. J. Clin. Pharm.* 17: 459–464, 1984.
- [76047] 14377. Hoo, J. J.; Hussein, L.; Goedde, H. W.: A simplified micromethod for the determination of the acetylator phenotype. *J. Clin. Chem. Clin. Biochem.* 15: 329–331, 1977.
- [76048] 14378. Ilett, K. F.; David, B. M.; Detchon, P.; Castleden, W. M.; Kwa, R.: Acetylation phenotype in colorectal carcinoma. *Cancer Res.* 47: 1466–1469, 1987.
- [76049] 14379. Iselius, L.; Evans, D. A. P.: Formal genetics of isoniazid metabolism in man. *Clin. Pharmacokinetics* 8: 541–544, 1983.
- [76050] 14380. Kukongviriyapan, V.; Lulitanond, V.; Areejitranusorn, C.; Kongyingyose, B.; Laupattarakasem, P.: N-acetyltransferase polymorphism in Thailand. *Hum. Hered.* 34: 246–249, 1984.
- [76051] 14381. Kutt, H.; Brennan, R.; Dehejia, H.; Verebely, K.:

Diphenylhydantoinintoxication: a complication of isoni-  
azid therapy. *Am. Rev. Resp.Dis.* 101: 377–384, 1970.

[76052] 14382.Lang, N. P.; Chu, D. Z. J.; Hunter C. F.; Kendall, D.  
C.; Flammang,T. J.; Kadlubar, F. F.: Role of aromatic amine  
acetyltransferasein human colorectal cancer. *Arch. Surg.*  
121: 1259–1261, 1986.

[76053] 14383.Lee, E. J. D.; Zhao, B.; Seow–Choen, F.: Relationship  
betweenpolymorphism of N–acetyltransferase gene and  
susceptibility to colorectalcarcinoma in a Chinese popula-  
tion. *Pharmacogenetics* 8: 513–517,1998.

[76054] 14384.Lin, H. J.; Han, C.–Y.; Lin, B. K.; Hardy, S.: Ethnic  
distributionof slow acetylator mutations in the polymor-  
phic N–acetyltransferase(NAT2) gene. *Pharmacogenetics*  
4: 125–134, 1994.

[76055] 14385.Lin, H. J.; Han, C.–Y.; Lin, B. K.; Hardy, S.: Slow  
acetylatormutations in the human polymorphic N–  
acetyltransferase gene in 786Asians, blacks, Hispanics,  
and whites: application to metabolic epidemiology. *Am.J.*  
*Hum. Genet.* 52: 827–834, 1993.

[76056] 14386.Mashimo, M.; Suzuki, T.; Abe, M.; Deguchi, T.:  
Molecular genotypingof N–acetylation polymorphism to  
predict phenotype. *Hum. Genet.* 90:139–143, 1992.

[76057] 14387.Mattano, S. S.; Weber, W. W.: Kinetics of arylamine

N-acetyltransferase in tissues from rapid and slow acetylators mice. *Carcinogenesis* 8:133–137, 1987.

- [76058] 14388. McLaren, E. H.; Burden, A. C.; Moorhead, P. J.: Acetylator phenotype in diabetic neuropathy. *Brit. Med. J.* 2: 291–293, 1977.
- [76059] 14389. McQueen, C. A.; Maslansky, C. J.; Glowinski, I. B.; Crescenzi, S. B.; Weber, W. W.; Williams, G. M.: Relationship between the genetically determined acetylator phenotype and DNA damage induced by hydralazine and 2-aminofluorene in cultured rabbit hepatocytes. *Proc. Nat. Acad. Sci.* 79: 1269–1272, 1982.
- [76060] 14390. Giralt, M.; Park, E. A.; Gurney, A. L.; Liu, J.; Hakimi, P.; Hanson, R. W.: Identification of a thyroid hormone response element in the phosphoenolpyruvate carboxykinase (GTP) gene: evidence for synergistic interaction between thyroid hormone and cAMP cis-regulatory elements. *J. Biol. Chem.* 266: 21991–21996, 1991.
- [76061] 14391. Pilz, A. J.; Willer, E.; Povey, S.; Abbott, C. M.: The genes coding for phosphoenolpyruvate carboxykinase-1 (PCK1) and neuronal nicotinic acetylcholine receptor alpha-4 subunit (CHRNA4) map to human chromosome 20, extending the known region of homology with mouse chromosome 2. *Ann. Hum. Genet.* 56: 289–293, 1992.

- [76062] 14392.Olswang, Y.; Cohen, H.; Papo, O.; Cassuto, H.; Croniger, C. M.;Hakimi, P.; Tilghman, S. M.; Hanson, R. W.; Reshef, L.: A mutationin the peroxisome proliferator–activated receptor gamma–binding sitein the gene for the cytosolic form of phosphoenolpyruvate carboxykinasere–duces adipose tissue size and fat content in mice. *Proc. Nat. Acad.Sci.* 99: 625–630, 2002.
- [76063] 14393.Stoffel, M.; Xiang, K.; Espinosa, R., III; Cox, N. J.; Le Beau,M. M.; Bell, G. I.: cDNA sequence and localization of polymorphichuman cytosolic phosphoenolpyruvate carboxykinase gene (PCK1) to chromosome20, band q13.31: PCK1 is not tightly linked to maturity–onset diabetesof the young. *Hum. Molec. Genet.* 2: 1–4, 1993.
- [76064] 14394.Ting, C.–N.; Burgess, D. L.; Chamberlain, J. S.; Keith, T. P.;Falls, K.; Meisler, M. H.: Phosphoenolpyruvate carboxykinase (GTP):characterization of the human PCK1 gene and localization distal toMODY on chromosome 20. *Genomics* 16: 698–706, 1993.
- [76065] 14395.Vidnes, J.; Sovik, O.: Gluconeogenesis in infancy and childhood.III. Deficiency of the extramitochondrial form of hepatic phosphoenolpyruvatecarboxykinase in a case of persistent neonatal hypoglycaemia. *ActaPaediat. Scand.* 65: 301–312, 1976.

- [76066] 14396.Yu, H.; Thun, R.; Chandrasekharappa, S.; Trent, J. M.; Zhang, J.; Meisler, M. H.: Human PCK1 encoding phosphoenolpyruvate carboxykinase is located on chromosome 20q13.2. *Genomics* 15: 219–221, 1993.
- [76067] 14397.Pepe, G.; Chimienti, G.; Resta, F.; Di Perna, V.; Taricone, C.; Lovecchio, M.; Colacicco, A. M.; Capurso, A.: A new Italian case of lipoprotein lipase deficiency: a leu365-to-val change resulting in loss of enzyme activity. *Biochem. Biophys. Res. Commun.* 199:570–576, 1994.
- [76068] 14398.Reymer, P. W. A.; Groenemeyer, B. E.; Gagne, E.; Miao, L.; Appelman, E. E. G.; Seidel, J. C.; Kromhout, D.; Bijvoet, S. M.; van de Oever, K.; Bruin, T.; Hayden, M. R.; Kastelein, J. J. P.: A frequently occurring mutation in the lipoprotein lipase gene (Asn291Ser) contributes to the expression of familial combined hyperlipidemia. *Hum. Molec. Genet.* 4:1543–1549, 1995.
- [76069] 14399.Samuels, M. E.; Forbey, K. C.; Reid, J. E.; Abkevich, V.; Bulka, K.; Wardell, B. R.; Bowen, B. R.; Hopkins, P. N.; Hunt, S. C.; Ballinger, D. G.; Skolnick, M. H.; Wagner, S.: Identification of a common variant in the lipoprotein lipase gene in a large Utah kindred ascertained for coronary heart disease: the –93G/D9N variant predisposes to low HDL-C/high triglycerides. *Clin. Genet.* 59: 88–98, 2001.

- [76070] 14400.Schreibman, P. H.; Arons, D. L.; Saudek, C. D.; Arky, R. A.:Abnormal lipoprotein lipase in familial exogenous hypertriglyceridemia. J.Clin. Invest. 52: 2074–2082, 1973.
- [76071] 14401.Sprecher, D. L.; Kobayashi, J.; Rymaszewski, M.; Goldberg, I.J.; Harris, B. V.; Bellet, P. S.; Ameis, D.; Yunker, R. L.; Black,D. M.; Stein, E. A.; Schotz, M. C.; Wiginton, D. A.: Trp64–to–nonsensemutation in the lipoprotein lipase gene. J. Lipid Res. 33: 859–866,1992.
- [76072] 14402.Sternowsky, H. J.; Gaertner, U.; Stahnkel, N.; Kaukel, E.: Juvenilefamilial hypertriglyceridemia and growth retardation: clinical andbiochemical observations in three siblings. Europ. J. Pediat. 125:59–70, 1977.
- [76073] 14403.Takagi, A.; Ikeda, Y.; Tsutsumi, Z.; Shoji, T.; Yamamoto, A.:Molecular studies on primary lipoprotein lipase (LPL) deficiency:one base deletion (G–916) in exon 5 of LPL gene causes no detectableLPL protein due to the absence of LPL mRNA transcript. J. Clin. Invest. 89:581–591, 1992.
- [76074] 14404.Weinstock, P. H.; Bisgaier, C. L.; Aalto–Setala, K.; Radner, H.;Ramakrishnan, R.; Levak–Frank, S.; Essenburg, A. D.; Zechner, R.;Breslow, J. L.: Severe hypertriglyceridemia, reduced high densitylipoprotein, and neonatal

death in lipoprotein lipase knockout mice:mild hypertriglyceridemia with impaired very low density lipoprotein clearance in heterozygotes. J. Clin. Invest. 96: 2555–2568, 1995.

[76075] 14405.Wessler, S.; Avioli, L. A.: Familial hyperlipoproteinemia. J.A.M.A. 207:929–937, 1969.

[76076] 14406.Wilson, D. E.; Edwards, C. Q.; Chan, I.–F.: Phenotypic heterogeneity in the extended pedigree of a proband with lipoprotein lipase deficiency. Metabolism 32:1107–1114, 1983.

[76077] 14407.Wilson, D. E.; Emi, M.; Iverius, P.–H.; Hata, A.; Wu, L. L.; Hillas,E.; Williams, R. R.; Lalouel, J.–M.: Phenotypic expression of heterozygous lipoprotein lipase deficiency in the extended pedigree of a proband homozygous for a missense mutation. J. Clin. Invest. 86: 735–750, 1990.

[76078] 14408.Wilson, D. E.; Hata, A.; Kwong, L. K.; Lingam, A.; Shuhua, J.; Ridinger, D. N.; Yeager, C.; Kaltenborn, K. C.; Iverius, P.–H.; Lalouel, J.–M.: Mutations in exon 3 of the lipoprotein lipase gene segregating in a family with hypertriglyceridemia, pancreatitis, and non–insulin–dependent diabetes. J. Clin. Invest. 92: 203–211, 1993.

[76079] 14409.Wion, K. L.; Kirchgessner, T. G.; Lusi, A. J.; Schotz,

M. C.;Lawn, R. M.: Human lipoprotein lipase complementary DNA sequence. *Science* 235:1638–1641, 1987.

[76080] 14410.Wittrup, H. H.; Tybjaerg-Hansen, A.; Abildgaard, S.; Steffensen,R.; Schnohr, P.; Nordestgaard, B. G.: A common substitution (asn291ser)in lipoprotein lipase is associated with increased risk of ischemicheart disease. *J. Clin. Invest.* 99: 1606–1613, 1997.

[76081] 14411.Wittrup, H. H.; Tybjaerg-Hansen, A.; Steffensen, R.; Deeb, S.S.; Brunzell, J. D.; Jensen, G.; Nordestgaard, B. G.: Mutations inthe lipoprotein lipase gene associated with ischemic heart diseasein men: the Copenhagen City Heart Study. *Arterioscler. Thromb. Vasc.Biol.* 19: 1535–1540, 1999.

[76082] 14412.Wood, S.; Schertzer, M.; Hayden, M.; Ma, Y.: Support for foundereffect for two lipoprotein lipase (LPL) gene mutations in French Canadiansby analysis of GT microsatellites flanking the LPL gene. *Hum. Genet.* 91:312–316, 1993.

[76083] 14413.McQueen, C. A.; Weber, W. W.: Characterization of human lymphocyteN-acetyltransferase and its relationship to the isoniazid acetylatorpolymorphism. *Biochem. Genet.* 18: 889–904, 1980.

[76084] 14414.Nhachi, C. F. B.: Polymorphic acetylation of sul-



phamethazine in a Zimbabwe population. *J. Med. Genet.* 25: 29–31, 1988.

[76085] 14415. Ohsako, S.; Deguchi, T.: Cloning and expression of cDNAs for polymorphic and monomorphic arylamine N-acetyltransferases from human liver. *J. Biol. Chem.* 265: 4630–4634, 1990.

[76086] 14416. Parker, J. M.: Human variability in the metabolism of sulfamethazine. *Hum. Hered.* 19: 402–409, 1969.

[76087] 14417. Penketh, R. J. A.; Gibney, S. F. A.; Nurse, G. T.; Hopkinson, D. A.: Acetylator phenotypes in Papua New Guinea. *J. Med. Genet.* 20: 37–40, 1983.

[76088] 14418. Roberts–Thomson, I. C.; Ryan, P.; Khoo, K. K.; Hart, W. J.; McMichael, A. J.; Butler, R. N.: Diet, acetylator phenotype, and risk of colorectal neoplasia. *Lancet* 347: 1372–1374, 1996.

[76089] 14419. Roots, I.; Drakoulis, N.; Brockmoller, J.; Janicke, I.; Cuprunov, M.; Ritter, J.: Hydroxylation and acetylation phenotypes as genetic risk factors in certain malignancies. In: Kato, W.; Estabrook, R. W.; Cayen, M. N. (eds.): *Xenobiotic Metabolism and Disposition*. London: Taylor and Francis 1989. Pp. 499–506.

[76090] 14420. Schloot, W.; Goedde, H. W.: Studies on the polymorphism of isoniazid (INH) acetylation in rhesus monkeys

(*Macaca mulatta*). *Acta Genet. Statist. Med.* 18: 394–398, 1968.

- [76091] 14421. Schnakenberg, E.; Ehlers, C.; Feyerabend, W.; Werdin, R.; Hubotter, R.; Dreikorn, K.; Schloot, W.: Genotyping of the polymorphic N-acetyltransferase (NAT2) and loss of heterozygosity in bladder cancer patients. *Clin. Genet.* 53: 396–402, 1998.
- [76092] 14422. Sonnhag, C.; Karlsson, E.; Hed, J.: Procainamide-induced lupus erythematosus-like syndrome in relation to acetylator phenotype and plasma levels of procainamide. *Acta Med. Scand.* 206: 245–251, 1979.
- [76093] 14423. Sunahara, S.; Urano, M.; Ogawa, M.: Genetical and geographic studies on isoniazid inactivation. *Science* 134: 1530–1531, 1961.
- [76094] 14424. Tannen, R. H.; Weber, W. W.: Inheritance of acetylator phenotype in mice. *J. Pharm. Exp. Therap.* 213: 480–484, 1980.
- [76095] 14425. Timbrell, J. A.; Wright, J. M.; Baillie, T. A.: Monoacetylhydrazine as a metabolite of isoniazid in man. *Clin. Pharm. Therap.* 22: 602–609, 1977.
- [76096] 14426. Vansant, J.; Woosley, R. L.; John, J. T.; Sergent, J. S.: Normal distribution of acetylation phenotypes in systemic lupus erythematosus. *Arthritis Rheum.* 21: 192–195, 1978.

- [76097] 14427.Weber, W. W.: The Acetylators Genes and Drug Response. New York:Oxford Univ. Press (pub.) 1987.
- [76098] 14428.Woosley, R. L.; Drayer, D. E.; Reidenberg, M. M.; Nies, A. S.;Carr, K.; Oates, J. A.: Effect of acetylator phenotype on the rate at which procainamide induces antinuclear antibodies and the lupus syndrome. New Eng. J. Med. 298: 1157–1159, 1978.
- [76099] 14429.Pace, P.; Johnson, M.; Tan, W. M.; Mosedale, G.; Sng, C.; Hoatlin, M.; de Winter, J.; Joenje, H.; Gergely, F.; Patel, K. J.: FANCE: the link between Fanconi anaemia complex assembly and activity. EMBO J. 21: 3414–3423, 2002.
- [76100] 14430.Hodge, S. E.; Anderson, C. E.; Neiswanger, K.; Field, L. L.; Spence, M. A.; Sparkes, R. S.; Sparkes, M. C.; Crist, M.; Terasaki, P. I.; Rimoim, D. L.; Rotter, J. I.: Close genetic linkage between diabetes mellitus and Kidd blood group. Lancet II: 893–895, 1981.
- [76101] 14431.Faiyaz-Ul-Haque, M.; Ahmad, W.; Zaidi, S. H. E.; Haque, S.; Teebi, A. S.; Ahmad, M.; Cohn, D. H.; Tsui, L. C.: Mutation in the cartilage-derived morphogenetic protein-1 (CDMP1) gene in a kindred affected with fibular hypoplasia and complex brachydactyly (DuPan syndrome). Clin. Genet. 61:454–458, 2002.
- [76102] 14432.Touchman, J. W.; Anikster, Y.; Dietrich, N. L.;

Braden Maduro, V. V.; McDowell, G.; Shotelersuk, V.; Bouffard, G. G.; Beckstrom-Sternberg, S. M.; Gahl, W. A.; Green, E. D.: The genomic region encompassing the nephropathic cystinosis gene (CTNS): complete sequencing of a 200-kb segment and discovery of a novel gene within the common cystinosis-causing deletion. *Genome Res.* 10: 165–173, 2000.

[76103] 14433. Shapiro, M. B.; Senapathy, P.: RNA splice junctions of different classes of eukaryotes: sequence statistics and functional implications in gene expression. *Nucleic Acids Res.* 15: 7155–7174, 1987.

[76104] 14434. Jobard, F.; Lefevre, C.; Karaduman, A.; Blanchet-Baron, C.; Emre, S.; Weissenbach, J.; Ozguc, M.; Lathrop, M.; Prud'homme, J.-F.; Fischer, J.: Lipoxigenase-3 (ALOXE3) and 12(R)-lipoxigenase (ALOX12B) are mutated in non-bullous congenital ichthyosiform erythroderma (NCIE) linked to chromosome 17p13.1. *Hum. Molec. Genet.* 11: 107–113, 2002.

[76105] 14435. Varon, R.; Vissinga, C.; Platzer, M.; Cerosaletti, K. M.; Chrzanowska, K. H.; Saar, K.; Beckmann, G.; Seemanova, E.; Cooper, P. R.; Nowak, N. J.; Stumm, M.; Weemaes, C. M. R.; Gatti, R. A.; Wilson, R. K.; Digweed, M.; Rosenthal, A.; Sperling, K.; Concannon, P.; Reis, A.: Nibrin,

a novel DNA double-strand break repair protein, is mutated in Nijmegen breakage syndrome. *Cell* 93: 467–476, 1998.

[76106] 14436. ten Hove, T.; Corbaz, A.; Amitai, H.; Aloni, S.; Belzer, I.; Graber, P.; Driltenburg, P.; van Deventer, S. J.; Chvatchko, Y.; Te Velde, A. A.: Blockade of endogenous IL-18 ameliorates TNBS-induced colitis by decreasing local TNF- $\alpha$  production in mice. *Gastroenterology* 121:1372–1379, 2001.

[76107] 14437. Ishikawa, K.; Nagase, T.; Suyama, M.; Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Prediction of the coding sequences of unidentified human genes. X. The complete sequences of 100 new cDNA clones from brain which can code for large proteins in vitro. *DNAREs.* 5: 169–176, 1998.

[76108] 14438. Kipreos, E. T.; Lander, L. E.; Wing, J. P.; He, W. W.; Hedgecock, E. M.: *cul-1* is required for cell cycle exit in *C. elegans* and identifies a novel gene family. *Cell* 85: 829–839, 1996.

[76109] 14439. Rasooly, R. S.: Personal Communication. Baltimore, Md. 9/29/1998.

[76110] 14440. Wang, P. J.; McCarrey, J. R.; Yang, F.; Page, D. C.: An abundance of X-linked genes expressed in spermatogenesis.

gonia. *Nature Genet.* 27:422–426, 2001.

- [76111] 14441. Herold, A.; Suyama, M.; Rodrigues, J. P.; Braun, I. C.; Kutay, U.; Carmo-Fonseca, M.; Bork, P.; Izaurralde, E.: TAP (NXF1) belongs to a multigene family of putative RNA export factors with a conserved modular architecture. *Molec. Cell. Biol.* 20: 8996–9008, 2000.
- [76112] 14442. Yang, J.; Bogerd, H. P.; Wang, P. J.; Page, D. C.; Cullen, B. R.: Two closely related human nuclear export factors utilize entirely distinct export pathways. *Molec. Cell* 8: 397–406, 2001.
- [76113] 14443. Ikeda, M.; Ishida, O.; Hinoi, T.; Kishida, S.; Kikuchi, A.: Identification and characterization of a novel protein interacting with Ral-binding protein 1, a putative effector protein of Ral. *J. Biol. Chem.* 273:814–821, 1998.
- [76114] 14444. Koshiba, S.; Kigawa, T.; Iwahara, J.; Kikuchi, A.; Yokoyama, S.: Solution structure of the Eps15 homology domain of a human POB1 (partner of RalBP1). *FEBS Lett.* 442: 138–142, 1999.
- [76115] 14445. Crackower, M. A.; Sarao, R.; Oudit, G. Y.; Yagil, C.; Kozieradzki, I.; Scanga, S. E.; Oliveira-dos-Santos, A. J.; da Costa, J.; Zhang, L.; Pei, Y.; Scholey, J.; Ferrario, C. M.; Manoukian, A. S.; Chappell, M. C.; Backx, P. H.; Yagil, Y.; Penninger, J. M.: Angiotensin-converting enzyme 2 is an

essential regulator of heart function. *Nature* 417:822–828, 2002.

- [76116] 14446.Scanlan, M. J.; Gordan, J. D.; Williamson, B.; Stockert, E.; Bander, N. H.; Jongeneel, V.; Gure, A. O.; Jager, D.; Jager, E.; Knuth, A.; Chen, Y.-T.; Old, L. J.: Antigens recognized by autologous antibody in patients with renal-cell carcinoma. *Int. J. Cancer* 83: 456–464, 1999.
- [76117] 14447.Yan, Z.; Fedorov, S. A.; Mumby, M. C.; Williams, R. S.: PR48, a novel regulatory subunit of protein phosphatase 2A, interacts with Cdc6 and modulates DNA replication in human cells. *Molec. Cell. Biol.* 20:1021–1029, 2000.
- [76118] 14448.Gitzelmann, R.: Hereditary galactokinase deficiency, a newly recognized cause of juvenile cataracts. *Pediatr. Res.* 1: 14–23, 1967.
- [76119] 14449.Kalaydjieva, L.; Perez-Lezaun, A.; Angelicheva, D.; Onengut, S.; Dye, D.; Bosshard, N. U.; Jordanova, A.; Savov, A.; Yanakiev, P.; Kremensky, I.; Radeva, B.; Hallmayer, J.; Markov, A.; Nedkova, V.; Tournev, I.; Aneva, L.; Gitzelmann, R.: A founder mutation in the GK1 gene is responsible for galactokinase deficiency in Roma (Gypsies). *Am. J. Hum. Genet.* 65: 1299–1307, 1999.
- [76120] 14450.Stratakis, C. A.; Lin, J.-P.; Pras, E.; Rennert, O. M.;

Bourdony, C. J.; Chan, W.-Y.: Segregation of Allgrove (triple-A) syndrome in Puerto Rican kindreds with chromosome 12 (12q13) polymorphic markers. *Proc. Assoc. Am. Phys.* 109: 478–482, 1997.

[76121] 14451. Tullio-Pelet, A.; Salomon, R.; Hadj-Rabia, S.; Mugnier, C.; deLaet, M.-H.; Chaouachi, B.; Bakiri, F.; Brottier, P.; Cattolico, L.; Penet, C.; Begeot, M.; Naville, D.; Nicolino, M.; Chaussain, J.-L.; Weissenbach, J.; Munnich, A.; Lyonnet, S.: Mutant WD-repeat protein in triple-A syndrome. *Nature Genet.* 26: 332–335, 2000.

[76122] 14452. Camacho, J. A.; Obie, C.; Biery, B.; Goodman, B. K.; Hu, C.-A.; Almashanu, S.; Steel, G.; Casey, R.; Lambert, M.; Mitchell, G. A.; Valle, D.: Hyperornithinaemia-hyperammonaemia-homocitrullinuria syndrome is caused by mutations in a gene encoding a mitochondrial ornithine transporter. *Nature Genet.* 22: 151–158, 1999.

[76123] 14453. Darnell, J. E., Jr.: Reflections on STAT3, STAT5, and STAT6 as fat STATs. *Proc. Nat. Acad. Sci.* 93: 6221–6224, 1996.

[76124] 14454. Ghilardi, N.; Ziegler, S.; Wiestner, A.; Stoffel, R.; Heim, M. H.; Skoda, R. C.: Defective STAT signaling by the leptin receptor in diabetic mice. *Proc. Nat. Acad. Sci.* 93: 6231–6235, 1996.



- [76125] 14455. Mohandas, T. K.; Chen, X.-N.; Rowe, L. B.; Birkenmeier, E. H.; Fanning, A. S.; Anderson, J. M.; Korenberg, J. R.: Localization of the tight junction protein gene TJP1 to human chromosome 15q13, distal to the Prader-Willi/Angelman region, and to mouse chromosome 7. *Genomics* 30:594–597, 1995.
- [76126] 14456. Willott, E.; Balda, M. S.; Fanning, A. S.; Jameson, B.; Van Itallie, C.; Anderson, J. M.: The tight junction protein ZO-1 is homologous to the *Drosophila* discs-large tumor suppressor protein of septate junctions. *Proc. Nat. Acad. Sci.* 90: 7834–7838, 1993.
- [76127] 14457. Blannar, M. A.; Crossley, P. H.; Peters, K. G.; Steingrimsson, E.; Copeland, N. G.; Jenkins, N. A.; Martin, G. R.; Rutter, W. J.: Meso1, a basic helix-loop-helix protein involved in mammalian presomitic mesoderm development. *Proc. Nat. Acad. Sci.* 92: 5870–5874, 1995.
- [76128] 14458. Burgess, R.; Rawls, A.; Brown, D.; Bradley, A.; Olson, E. N.: Requirement of the paraxis gene for somite formation and musculoskeletal patterning. *Nature* 384: 570–573, 1996.
- [76129] 14459. Hidai, H.; Quertermous, E. E.; Espinosa, R., III; Le Beau, M. M.; Quertermous, T.: Genomic organization and chromosomal localization of the gene TCF15 encoding the

early mesodermal basic helix-loop-helix factor bHLH-EC2. Genomics 30: 598-601, 1995.

[76130] 14460. Quertermous, E. E.; Hidai, H.; Blannar, M. A.; Quertermous, T.: Cloning and characterization of a basic helix-loop-helix protein expressed in early mesoderm and the developing somites. Proc. Nat. Acad. Sci. 91:7066-7070, 1994.

[76131] 14461. Quertermous, T.: Personal Communication. Nashville, Tenn. 1/22/1996.

[76132] 14462. Clevidence, D. E.; Overdier, D. G.; Peterson, R. S.; Porcella, A.; Ye, H.; Paulson, K. E.; Costa, R. H.: Members of the HNF-3/forkhead family of transcription factors exhibit distinct cellular expression patterns in lung and regulate the surfactant protein B promoter. Dev. Biol. 166: 195-209, 1994.

[76133] 14463. Hellqvist, M.; Mahlapuu, M.; Blixt, A.; Enerback, S.; Carlsson, P.: The human forkhead protein FREAC-2 contains two functionally redundant activation domains and interacts with TBP and TFIIB. J. Biol. Chem. 273: 23335-23343, 1998.

[76134] 14464. Hellqvist, M.; Mahlapuu, M.; Samuelsson, L.; Enerback, S.; Carlsson, P.: Differential activation of lung-specific genes by two forkhead proteins, FREAC-1 and

FREAC-2. J. Biol. Chem. 271: 4482-4490, 1996.

- [76135] 14465.Larsson, C.; Hellqvist, M.; Pierrou, S.; White, I.; Enerback, S.; Carlsson, P.: Chromosomal localization of six human forkhead genes, freac-1 (FKHL5), -3 (FKHL7), -4 (FKHL8), -5 (FKHL9), -6 (FKHL10), and -8 (FKHL12). Genomics 30: 464-469, 1995.
- [76136] 14466.Pierrou, S.; Hellqvist, M.; Samuelsson, L.; Enerback, S.; Carlsson, P.: Cloning and characterization of seven human forkhead proteins: binding site specificity and DNA bending. EMBO J. 13: 5002-5012, 1994.
- [76137] 14467.Ernstsson, S.; Pierrou, S.; Hulander, M.; Cederberg, A.; Hellqvist, M.; Carlsson, P.; Enerback, S.: Characterization of the human forkhead gene FREAC-4. J. Biol. Chem. 271: 21094-21099, 1996.
- [76138] 14468.Hulander, M.; Wurst, W.; Carlsson, P.; Enerback, S.: The winged helix transcription factor Fkh10 is required for normal development of the inner ear. Nature Genet. 20: 374-376, 1998.
- [76139] 14469.De Smaele, E.; Zazzeroni, F.; Papa, S.; Nguyen, D. U.; Jin, R.; Jones, J.; Cong, R.; Franzoso, G.: Induction of gadd45-beta by NF-kappa-B downregulates pro-apoptotic JNK signaling. Nature 414: 308-313, 2001.
- [76140] 14470.Dong, C.; Yang, D. D.; Tournier, C.; Whitmarsh, A.

J.; Xu, J.; Davis, R. J.; Flavell, R. A.: JNK is required for effector T-cell function but not for T-cell activation. *Nature* 405: 91–94, 2000.

- [76141] 14471. Gupta, S.; Barrett, T.; Whitmarsh, A. J.; Cavanagh, J.; Sluss, H. K.; Derijard, B.; Davis, R. J.: Selective interaction of JNK protein kinase isoforms with transcription factors. *EMBO J.* 15: 2760–2770, 1996.
- [76142] 14472. Tournier, C.; Hess, P.; Yang, D. D.; Xu, J.; Turner, T. K.; Nimnual, A.; Bar-Sagi, D.; Jones, S. N.; Flavell, R. A.; Davis, R. J.: Requirement of JNK for stress-induced activation of the cytochrome c-mediated death pathway. *Science* 288: 870–874, 2000.
- [76143] 14473. Daugherty, B. L.; Springer, M. S.: The beta-chemokine receptor genes CCR1 (CMKBR1), CCR2 (CMKBR2), and CCR3 (CMKBR3) cluster within 285 kb on human chromosome 3p21. *Genomics* 41: 294–295, 1997.
- [76144] 14474. Ayyanathan, K.; Naylor, S. L.; Kunapuli, S. P.: Structural characterization and fine chromosomal mapping of the human P2Y(1) purinergic receptor gene (P2RY1). *Somat. Cell Molec. Genet.* 22: 419–424, 1996.
- [76145] 14475. Ayyanathan, K.; Webbs, T. E.; Sandhu, A. K.; Athwal, R. S.; Barnard, E. A.; Kunapuli, S. P.: Cloning and chromosomal localization of the human P2Y1 purinoceptor.

Biochem. Biophys. Res. Commun. 218: 783–788,1996.

- [76146] 14476. Leon, C.; Hechler, B.; Freund, M.; Eckly, A.; Vial, C.; Ohlmann, P.; Dierich, A.; LeMeur, M.; Cazenave, J.-P.; Gachet, C.: Defective platelet aggregation and increased resistance to thrombosis in purinergic P2Y<sub>1</sub> receptor-null mice. *J. Clin. Invest.* 104: 1731–1737, 1999.
- [76147] 14477. Leon, C.; Hechler, B.; Vial, C.; Leray, C.; Cazenave, J.-P.; Gachet, C.: The P2Y<sub>1</sub> receptor is an ADP receptor antagonized by ATP and expressed in platelets and megakaryoblastic cells. *FEBS Letts.* 403:26–30, 1997.
- [76148] 14478. Leon, C.; Vial, C.; Cazenave, J.-P.; Gachet, C.: Cloning and sequencing of a human cDNA encoding endothelial P2Y<sub>1</sub> purinoceptor. *Gene* 171:295–297, 1996.
- [76149] 14479. Matsuda, C.; Hayashi, Y. K.; Ogawa, M.; Aoki, M.; Murayama, K.; Nishino, I.; Nonaka, I.; Arahata, K.; Brown, R. H., Jr.: The sarcolemmal proteins dysferlin and caveolin-3 interact in skeletal muscle. *Hum. Molec. Genet.* 10: 1761–1766, 2001.
- [76150] 14480. Furlong, R. A.; Zhou, C. Y.; Ferguson-Smith, M. A.; Affara, N. A.: Characterization of a kinesin-related gene ATSV, within the tuberous sclerosis locus (TSC1) candidate region on chromosome 9q34. *Genomics* 33:421–429, 1996.

- [76151] 14481.Keller, M. P.; Seifried, B. A.; Rabin, B. A.; Chance, P. F.: Mapping of the kinesin-related gene ATSV to chromosome 2q37. *Hum. Genet.* 104:254–256, 1999.
- [76152] 14482.Kikkawa, M.; Okada, Y.; Hirokawa, N.: 15-angstrom resolution model of the monomeric kinesin motor, KIF1A. *Cell* 100: 241–252, 2000.
- [76153] 14483.Kikkawa, M.; Sablin, E. P.; Okada, Y.; Yajima, H.; Fletterick, R. J.; Hirokawa, N.: Switch-based mechanism of kinesin motors. *Nature* 411:439–445, 2001.
- [76154] 14484.Klopfenstein, D. R.; Tomishige, M.; Stuurman, N.; Vale, R. D.: Role of phosphatidylinositol(4,5)bisphosphate organization in membrane transport by the Unc104 kinesin motor. *Cell* 109: 347–358, 2002.
- [76155] 14485.Okada, Y.; Yamazaki, H.; Sekine-Aizawa, Y.; Hirokawa, N.: The neuron-specific kinesin superfamily protein KIF1A is a unique monomeric motor for anterograde axonal transport of synaptic vesicle precursors. *Cell* 81:769–780, 1995.
- [76156] 14486.Yonekawa, Y.; Harada, A.; Okada, Y.; Funakoshi, T.; Kanai, Y.; Takei, Y.; Terada, S.; Noda, T.; Hirokawa, N.: Defect in synaptic vesicle precursor transport and neuronal cell death in KIF1A motor protein-deficient mice. *J. Cell Biol.* 141: 431–441, 1998.

- [76157] 14487.Kikuta, Y.; Sogawa, K.; Haniu, M.; Kinosaki, M.; Kusunose, E.;Nojima, Y.; Yamamoto, S.; Ichihara, K.; Kusunose, M.; Fujii-Kuriyama,Y.: A novel species of cytochrome P-450 (P-450ib) specific for the small intestine of rabbits: cDNA cloning and its expression in COS cells. *J. Biol. Chem.* 266: 17821–17825, 1991.
- [76158] 14488.Ma, J.; Ramachandran, S.; Fiedorek, F. T., Jr.; Zeldin, D. C.: Mapping of the CYP2J cytochrome P450 genes to human chromosome 1 and mouse chromosome 4. *Genomics* 49: 152–155, 1998.
- [76159] 14489.Wu, S.; Moomaw, C. R.; Tomer, K. B.; Falck, J. R.; Zeldin, D. C.: Molecular cloning and expression of CYP2J2, a human cytochrome P450 arachidonic acid epoxygenase highly expressed in heart. *J. Biol.Chem.* 271: 3460–3468, 1996.
- [76160] 14490.Chiang, P.-W.; Wang, S.; Smithivas, P.; Song, W.-J.; Ramamoorthy,S.; Hillman, J.; Puett, S.; Van Keuren, M. L.; Crombez, E.; Kumar,A.; Glover, T. W.; Miller, D. E.; Tsai, C.-H.; Blackburn, C. C.; Chen,X.-N.; Sun, Z.; Cheng, J.-F.; Korenberg, J. R.; Kurnit, D. M.: Identification and analysis of the human and murine putative chromatin structure-regulator SUPT6H and Supt6h. *Genomics* 34: 328–333, 1996.

- [76161] 14491. Segre, J. A.; Nemhauser, J. L.; Taylor, B. A.; Nadeau, J. H.; Lander, E. S.: Positional cloning of the nude locus: genetic, physical and transcription maps of the region and mutations in the mouse and rat. *Genomics* 28:549–559, 1995.
- [76162] 14492. Han, J.; Knops, J. F.; Longshore, J. W.; King, P. H.: Localization of human elav-like neuronal protein 1 (Hel-N1) on chromosome 9p21 by chromosome microdissection polymerase chain reaction and fluorescence in situ hybridization. *Genomics* 36: 189–191, 1996.
- [76163] 14493. King, P. H.: Hel-N2: a novel isoform of Hel-N1 which is conserved in rat neural tissue and produced in early embryogenesis. *Gene* 151:261–265, 1994.
- [76164] 14494. King, P. H.; Levine, T. D.; Fremeau, R. T., Jr.; Keene, J. D.: Mammalian homologs of *Drosophila* ELAV localized to a neuronal subset can bind in vitro to the 3' UTR of mRNA encoding the *Id* transcriptional repressor. *J. Neurosci.* 14: 1943–52, 1994.
- [76165] 14495. Abel, K. J.; Brody, L. C.; Valdes, J. M.; Erdos, M. R.; McKinley, D. R.; Castilla, L. H.; Merajver, S. D.; Couch, F. J.; Friedman, L. S.; Ostermeyer, E. A.; Lynch, E. D.; King, M.-C.; Welcsh, P. L.; Osborne-Lawrence, S.; Spillman, M.; Bowcock, A. M.; Collins, F. S.; Weber, B. L.: Characteriza-



tion of EZH1, a human homolog of Drosophila enhancer of zeste near BRCA1. *Genomics* 37:161–171, 1996.

- [76166] 14496. Akazawa, K.; Yamane, S.; Shiota, H.; Naito, E.: A case of retinoblastoma associated with Rieger's anomaly and 13q deletion. *Jpn. J. Ophthalmol.* 25:321–325, 1981.
- [76167] 14497. Stathacopoulos, R. A.; Bateman, J. B.; Sparkes, R. S.; Hepler, R. S.: The Rieger syndrome and a chromosome 13 deletion. *J. Pediatr. Ophthalmol. Strabismus* 24: 198–203, 1987.
- [76168] 14498. Ferbus, D.; Le Chalony, C.; Prosperi, M.–T.; Muleris, M.; Vincent–Salomon, A.; Goubin, G.: Identification, nuclear localization, and binding activities of OZF, a human protein solely composed of zinc finger motifs. *Europ. J. Biochem.* 236: 991–995, 1996.
- [76169] 14499. Le Chalony, C.; Apiou, F.; Pibouin, L.; Dutrillaux, B.; Goubin, G.: Constitutive amplification of a zinc finger protein gene in cattle. *DNACell Biol.* 15: 83–88, 1996.
- [76170] 14500. Le Chalony, C.; Prosperi, M.–T.; Haluza, R.; Apiou, F.; Dutrillaux, B.; Goubin, G.: The OZF gene encodes a protein consisting essentially of zinc–finger motifs. *J. Molec. Biol.* 236: 399–404, 1994.
- [76171] 14501. Dell'Angelica, E. C.; Ohno, H.; Ooi, C. E.; Rabinovich, E.; Roche, K. W.; Bonifacino, J. S.: AP–3: an adap–

tor-like protein complex with ubiquitous expression.

EMBO J. 16: 917–928, 1997.

- [76172] 14502. Watanabe, T. K.; Shimizu, F.; Nagata, M.; Takaichi, A.; Fujiwara, T.; Nakamura, Y.; Takahashi, E.; Hirai, Y.: Cloning, expression pattern and mapping to 12p13.2–p13.1 of CLAPS3, a gene encoding a novel clathrin-adaptor small chain. Cytogenet. Cell Genet. 73: 214–217, 1996.
- [76173] 14503. Angrist, M.; Wells, D. E.; Chakravarti, A.; Pandey, A.: Chromosomal localization of the mouse Src-like adapter protein (Slap) gene and its putative human homolog SLA. Genomics 30: 623–625, 1995.
- [76174] 14504. Holland, S. J.; Liao, X. C.; Mendenhall, M. K.; Zhou, X.; Pardo, J.; Chu, P.; Spencer, C.; Fu, A.; Sheng, N.; Yu, P.; Pali, E.; Nagin, A.; and 14 others: Functional cloning of Src-like adapter protein-2 (SLAP-2), a novel inhibitor of antigen receptor signaling. J. Exp. Med. 194: 1263–1276, 2001.
- [76175] 14505. Kratchmarova, I.; Sosinowski, T.; Weiss, A.; Witter, K.; Vincenz, C.; Pandey, A.: Characterization of promoter region and genomic structure of the murine and human genes encoding Src like adapter protein. Gene 262:267–273, 2001.

- [76176] 14506.Meijerink, P. H. S.; Yanakiev, P.; Zorn, I.; Grierson, A. J.; Bikker,H.; Dye, D.; Kalaydjieva, L.; Baas, F.: The gene for the human Src-likeadaptor protein (hSLAP) is located within the 64-kb intron of the thyroglobulin gene. *Europ. J. Biochem.* 254: 297–303, 1998.
- [76177] 14507.Sosinowski, T.; Pandey, A.; Dixit, V. M.; Weiss, A.: Src-likeadaptor protein (SLAP) is a negative regulator of T cell receptorsignaling. *J. Exp. Med.* 191: 463–474, 2000.
- [76178] 14508.Bailey, S. M.; Cornforth, M. N.; Kurimasa, A.; Chen, D. J.; Goodwin,E. H.: Strand-specific postreplicative processing of mammalian telomeres. *Science* 293:2462–2465, 2001.
- [76179] 14509.Maniatis, T.: A ubiquitin ligase complex essential for the NF-kappa-B,Wnt/Wingless, and Hedgehog signaling pathways. *Genes Dev.* 13: 505–510,1999.
- [76180] 14510.Winston, J. T.; Strack, P.; Beer-Romero, P.; Chu, C. Y.; Elledge,S. J.; Harper, J. W.: The SCF(beta-TRCP)-ubiquitin ligase complexassociates specifically with phosphorylated destruction motifs in I-kappa-B-alpha and beta-catenin and stimulates I-kappa-B-alpha ubiquitinationin vitro. *Genes Dev.* 13: 270–283, 1999.
- [76181] 14511.Ahouse, J. J.; Hagerman, C. L.; Mittal, P.; Gilbert, D.

J.; Copeland, N. G.; Jenkins, N. A.; Simister, N. E.: Mouse MHC class I-like Fc receptor encoded outside the MHC. *J. Immun.* 151: 6076–6088, 1993.

[76182] 14512. Junghans, R. P.; Anderson, C. L.: The protection receptor for IgG catabolism is the beta-2-microglobulin-containing neonatal intestinal transport receptor. *Proc. Nat. Acad. Sci.* 93: 5512–5516, 1996.

[76183] 14513. Junghans, R. P.; Ebralidze, A.; Tiwari, B.: Does (CUG)<sub>n</sub> repeat in DMPK mRNA 'paint' chromosome 19 to suppress distant genes to create the diverse phenotype of myotonic dystrophy? A new hypothesis of long-range cis autosomal inactivation. *Neurogenetics* 3: 59–67, 2001.

[76184] 14514. Kandil, E.; Egashira, M.; Miyosi, O.; Niikawa, N.; Ishibashi, T.; Kasahara, M.: The human gene encoding the heavy chain of the major histocompatibility complex class I-like Fc receptor (FCGRT) maps to 19q13.3. *Cytogenet. Cell Genet.* 73: 97–98, 1996.

[76185] 14515. Story, C. M.; Mikulska, J. E.; Simister, N. E.: A major histocompatibility complex class I-like Fc receptor cloned from human placenta: possible role in transfer of immunoglobulin G from mother to fetus. *J. Exp. Med.* 180: 2377–2381, 1994.

[76186] 14516. Waldmann, T. A.; Strober, W.: Metabolism of im-

munoglobulins. *Prog.Allergy* 13: 1–110, 1969.

- [76187] 14517.Hart, T. C.; Price, J. A.; Bobby, P. L.; Pettenati, M. J.; Shashi,V.; Von Kap Herr, C.; Van Dyke, T. E.: Cytogenetic assignment andphysical mapping of the human DGKE gene to chromosome 17q22. *Genomics* 56:233–235, 1999.
- [76188] 14518.Tang, W.; Bunting, M.; Zimmerman, G. A.; McIntyre, T. M.; Prescott,S. M.: Molecular cloning of a novel human diacylglycerol kinase highlyselective for arachidonate–containing substrates. *J. Biol. Chem.* 271:10237–10241, 1996.
- [76189] 14519.Beddow, A. L.; Richards, S. A.; Orem, N. R.; Macara, I. G.: TheRan/TC4 GTPase–binding domain: identification by expression cloningand characterization of a conserved sequence motif. *Proc. Nat. Acad.Sci.* 92: 3328–3332, 1995.
- [76190] 14520.Fauser, S.; Aslanukov, A.; Roepman, R.; Ferreira, P. A.: Genomicorganization, expression, and localization of murine Ran–binding protein2 (RanBP2) gene. *Mammalian Genome* 12: 406–415, 2001.
- [76191] 14521.Krebber, H.; Bastians, H.; Hoheisel, J.; Lichter, P.; Ponstingl,H.; Joos, S.: Localization of the gene encoding the Ran–binding proteinRanBP2 to human chromosome

2q11–q13 by fluorescence in situ hybridization. *Genomics* 43:247–248, 1997.

[76192] 14522.Pichler, A.; Gast, A.; Seeler, J. S.; Dejean, A.; Melchior, F.: The nucleoporin RanBP2 has SUMO1 E3 ligase activity. *Cell* 108:109–120, 2002.

[76193] 14523.Wu, J.; Matunis, M. J.; Kraemer, D.; Blobel, G.; Coutavas, E.:Nup358, a cytoplasmically exposed nucleoporin with peptide repeats,Ran–GTP binding sites, zinc fingers, a cyclophilin A homologous domain,and a leucine–rich region. *J. Biol. Chem.* 270: 14209–14213, 1995.

[76194] 14524.Yokoyama, N.; Hayashi, N.; Seki, T.; Pante, N.; Ohba, T.; Nishii,K.; Kuma, K.; Hayashida, T.; Miyata, T.; Abei, U.; Fukui, M.; Nishimoto,T.: A giant nucleopore protein that binds Ran/TC4. *Nature* 376:184–188, 1995.

[76195] 14525.Dhar, S. K.; Yoshida, K.; Machida, Y.; Khaira, P.; Chaudhuri, B.;Wohlschlegel, J. A.; Leffak, M.; Yates, J.; Dutta, A.: Replicationfrom oriP of Epstein–Barr virus requires human ORC and is inhibitedby geminin. *Cell* 106: 287–296, 2001.

[76196] 14526.Gavin, K. A.; Hidaka, M.; Stillman, B.: Conserved initiator proteinsin eukaryotes. *Science* 270: 1667–1671, 1995.

- [76197] 14527.Ohtani, K.; DeGregori, J.; Leone, G.; Herendeen, D. R.; Kelly,T. J.; Nevins, J. R.: Expression of the HsOrc1 gene, a human ORC1homolog, is regulated by cell proliferation via the E2F transcriptionfactor. *Molec. Cell. Biol.* 16: 6977–6984, 1996.
- [76198] 14528.Takahara, K.; Bong, M.; Brevard, R.; Eddy, R. L.; Ha-ley, L. L.;Sait, S. J.; Shows, T. B.; Hoffman, G. G.; Greenspan, D. S.: Mouseand human homologues of the yeast origin of replication recognitioncomplex subunit ORC2 and chromosomal localization of the cognate hu-mangene ORC2L. *Genomics* 31: 119–122, 1996.
- [76199] 14529.Chang, A. C.–M.; Janosi, J.; Hulsbeek, M.; de Jong, D.; Jeffrey,K. J.; Noble, J. R.; Reddel, R. R.: A novel human cDNA highly homologousto the fish hormone stanniocalcin. *Molec. Cell. Endocr.* 112: 241–247,1995.
- [76200] 14530.Chang, A. C.–M.; Jeffrey, K. J.; Tokutake, Y.; Shi-mamoto, A.; Neumann,A. A.; Dunham, M. A.; Cha, J.; Sug-awara, M.; Furuichi, Y.; Reddel,R. R.: Human stanniocalcin (STC): genomic structure, chromosomallocalization, and the presence of CAG trinucleotide repeats. *Genomics* 47:393–398, 1998.
- [76201] 14531.Jellinek, D. A.; Chang, A. C.; Larsen, M. R.; Wang, X.; Robinson,P. J.; Reddel, R. R.: Stanniocalcin 1 and 2 are

secreted as phosphoproteins from human fibrosarcoma cells. *Biochem. J.* 350: 453–461, 2000.

- [76202] 14532. Olsen, H. S.; Cepeda, M. A.; Zhang, Q.-Q.; Rosen, C. A.; Vozzolo, B. L.; Wagner, G. F.: Human stanniocalcin: a possible hormonal regulator of mineral metabolism. *Proc. Nat. Acad. Sci.* 93: 1792–1796, 1996.
- [76203] 14533. Varghese, R.; Wong, C. K. C.; Deol, H.; Wagner, G. F.; DiMattia, G. E.: Comparative analysis of mammalian stanniocalcin genes. *Endocrinology* 139: 4714–4725, 1998.
- [76204] 14534. Wagner, G. F.; Guiraudon, C. C.; Milliken, C.; Copp, D. H.: Immunological and biological evidence for a stanniocalcin-like hormone in human kidney. *Proc. Nat. Acad. Sci.* 92: 1871–1875, 1995.
- [76205] 14535. Bartsch, O.; Wagner, A.; Hinkel, G. K.; Lichtner, P.; Murken, J.; Schuffenhauer, S.: No evidence for chromosomal microdeletions at the second DiGeorge syndrome locus on 10p near D10S585. (Letter) *Am. J. Med. Genet.* 83: 425–426, 1999.
- [76206] 14536. Daw, S. C. M.; Taylor, C.; Kraman, M.; Call, K.; Mao, J.; Schuffenhauer, S.; Meitinger, T.; Lipson, T.; Goodship, J.; Scambler, P.: A common region of 10p deleted in DiGeorge and velocardiofacial syndromes. *Nature Genet.* 13:



458–461, 1996.

- [76207] 14537.Lichtner, P.; Konig, R.; Hasegawa, T.; Van Esch, H.; Meitinger,T.; Schuffenhauer, S.: An HDR (hypoparathyroidism, deafness, renal dysplasia) syndrome locus maps distal to the DiGeorge syndrome region on 10p13/14. *J. Med. Genet.* 37: 33–37, 2000.
- [76208] 14538.Sorensen, O. E.; Follin, P.; Johnsen, A. H.; Calafat, J.; Tjabringa,G. S.; Hiemstra, P. S.; Borregaard, N.: Human cathelicidin, hCAP–18,is processed to the antimicrobial peptide LL–37 by extracellular cleavage with proteinase 3. *Blood* 97: 3951–3959, 2001.
- [76209] 14539.Zanetti, M.; Del Sal, G.; Storici, P.; Schneider, C.; Romeo, D.: The cDNA of the neutrophil antibiotic Bac5 predicts a pro–sequence homologous to a cysteine proteinase inhibitor that is common to other neutrophil antibiotics. *J. Biol. Chem.* 268: 522–526, 1993.
- [76210] 14540.Chen, C.–Y.; Gherzi, R.; Ong, S.–E.; Chan, E. L.; Rajmakers, R.;Prujn, G. J. M.; Stoecklin, G.; Moroni, C.; Mann, M.; Karin, M.:AU binding proteins recruit the exosome to degrade ARE–containing mRNAs. *Cell* 107: 451–464, 2001.
- [76211] 14541.Dangel, A. W.; Shen, L.; Mendoza, A. R.; Wu, L.–c.; Yu, C. Y.:Human helicase gene SKI2W in the HLA class III

region exhibits striking structural similarities to the yeast antiviral gene SKI2 and to the human gene KIAA0052: emergence of a new gene family. *Nucleic Acids Res.* 23: 2120–2126, 1995.

[76212] 14542. Lee, S.-G.; Lee, I.; Park, S. H.; Kang, C.; Song, K.: Identification and characterization of a human cDNA homologous to yeast SKI2. *Genomics* 25:660–666, 1995.

[76213] 14543. Shen, L.; Wu, L. C.; Sanlioglu, S.; Chen, R.; Mendoza, A. R.; Dangel, A. W.; Carroll, M. C.; Zipf, W. B.; Yu, C. Y.: Structure and genetics of the partially duplicated gene RP located immediately upstream of the complement C4A and the C4B genes in the HLA class III region: molecular cloning, exon–intron structure, composite retroposon, and breakpoint of gene duplication. *J. Biol. Chem.* 269: 8466–8476, 1994.

[76214] 14544. Yang, Z.; Shen, L.; Dangel, A. W.; Wu, L.-C.; Yu, C. Y.: Four ubiquitously expressed genes, RD (D6S45)–SKI2W (SKIV2L)–DOM3Z–RP1 (D6S60E), are present between complement component genes factor B and C4 in the class III region of the HLA. *Genomics* 53: 338–347, 1998.

[76215] 14545. Doyle, K.; Zhang, Y.; Baer, R.; Bina, M.: Distinguishable patterns of protein–DNA interactions involving complexes of basic helix–loop–helix proteins. *J. Biol. Chem.*

269: 12099–12105, 1994.

- [76216] 14546. Hu, J.-S.; Olson, E. N.; Kingston, R. E.: HEB, a helix-loop-helix protein related to E2A and ITF2 that can modulate the DNA-binding ability of myogenic regulatory factors. *Molec. Cell. Biol.* 12: 1031–1042, 1992.
- [76217] 14547. Sawada, S.; Littman, D. R.: A heterodimer of HEB and an E12-related protein interacts with the CD4 enhancer and regulates its activity in T-cell lines. *Molec. Cell. Biol.* 13: 5620–5628, 1993.
- [76218] 14548. Zhang, Y.; Babin, J.; Feldhaus, A. L.; Singh, H.; Sharp, P. A.; Bina, M.: HTF4: a new human helix-loop-helix protein. *Nucleic Acids Res.* 19: 4555 only, 1991.
- [76219] 14549. Zhang, Y.; Bina, M.: The nucleotide sequence of the human transcription factor HTF4a cDNA. *DNA Sequence* 2: 397–403, 1992.
- [76220] 14550. Zhang, Y.; Flejter, W. L.; Barcroft, C. L.; Riviere, M.; Szpirer, J.; Szpirer, C.; Bina, M.: Localization of the human HTF4 transcription factor 4 gene (TCF12) to chromosome 15q21. *Cytogenet. Cell Genet.* 68:235–238, 1995.
- [76221] 14551. Dubrulle, J.; McGrew, M. J.; Pourquie, O.: FGF signaling controls somite boundary position and regulates segmentation clock control of spatiotemporal Hox gene activation. *Cell* 106: 219–232, 2001.

- [76222] 14552.Fukuchi-Shimogori, T.; Grove, E. A.: Neocortex patterning by the secreted signaling molecule FGF8. *Science* 294: 1071–1074, 2001.
- [76223] 14553.Gemel, J.; Gorry, M.; Ehrlich, G. D.; MacArthur, C. A.: Structure and sequence of human FGF8. *Genomics* 35: 253–257, 1996.
- [76224] 14554.Rowen, L.; Young, J.; Birditt, B.; Kaur, A.; Madan, A.; Philipps, D. L.; Qin, S.; Minx, P.; Wilson, R. K.; Hood, L.; Graveley, B. R.: Analysis of the human neurexin genes: alternative splicing and the generation of protein diversity. *Genomics* 79: 587–597, 2002.
- [76225] 14555.Tabuchi, K.; Sudhof, T. C.: Structure and evolution of neurexin genes: insight into the mechanism of alternative splicing. *Genomics* 79:849–859, 2002.
- [76226] 14556.Ullrich, B.; Ushkaryov, Y. A.; Sudhof, T. C.: Cartography of neurexins: more than 1000 isoforms generated by alternative splicing and expressed in distinct subsets of neurons. *Neuron* 14: 497–507, 1995.
- [76227] 14557.Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T. C.: Neurexins: synaptic cell surface proteins related to the  $\alpha$ -latrotoxin receptor and laminin. *Science* 257: 50–56, 1992.
- [76228] 14558.Bergman, L.; Silins, G.; Grimmond, S.; Hummerich,

H.; Stewart,C.; Little, P.; Hayward, N.: A 500-kb sequence-ready cosmid contigand transcript map of the MEN1 region on 11q13. *Genomics* 55: 49–56,1999.

- [76229] 14559.Ushkaryov, Y. A.; Petrenko, A. G.; Geppert, M.; Sudhof, T. C.:Neurexins: synaptic cell surface proteins related to the alpha-latrotoxinreceptor and laminin. *Science* 257: 50–56, 1992.
- [76230] 14560.Scheiffele, P.; Fan, J.; Choh, J.; Fetter, R.; Serafini, T.:Neuroigin expressed in nonneuronal cells triggers presynaptic developmentin contacting axons. *Cell* 101: 657–669, 2000.
- [76231] 14561.Abderrahmani, A.; Steinmann, M.; Plaisance, V.; Niederhauser, G.;Haefliger, J.–A.; Mooser, V.; Bonny, C.; Nicod, P.; Waeber, G.: Thetranscriptional repressor REST determines the cell-specific expressionof the human MAPK8IP1 gene encoding IB1 (JIP–1). *Molec. Cell. Biol.* 21:7256–7627, 2001.
- [76232] 14562.Chen, Z.–F.; Paquette, A. J.; Anderson, D. J.: NRSF/ REST is requiredin vivo for repression of multiple neuronal target genes during embryogenesis. *NatureGenet.* 20: 136–142, 1998.
- [76233] 14563.Palm, K.; Metsis, M.; Timmusk, T.: Neuron-specific splicing of zinc finger transcription factor REST/NRSF/XBR

is frequent in neuroblastomas and conserved in human, mouse and rat. *Molec. Brain Res.* 72: 30–39, 1999.

- [76234] 14564. Schoenherr, C. J.; Anderson, D. J.: The neuron-restrictive silencer factor (NRSF): a coordinate repressor of multiple neuron-specific genes. *Science* 267: 1360–1363, 1995.
- [76235] 14565. Thiel, G.; Lietz, M.; Cramer, M.: Biological activity and modulator structure of RE-1-silencing transcription factor (REST), a repressor of neuronal genes. *J. Biol. Chem.* 273: 26891–26899, 1998.
- [76236] 14566. Cavalc, Y.; Popielarz, M.; Fuchs, J.-P.; Gattoni, R.; Stevenin, J.: Characterization and cloning of the human splicing factor 9G8: a novel 35 kDa factor of the serine/arginine protein family. *EMBO J.* 13: 2639–2649, 1994.
- [76237] 14567. Popielarz, M.; Cavalc, Y.; Mattei, M.-G.; Gattoni, R.; Stevenin, J.: The gene encoding human splicing factor 9G8: structure, chromosomal localization, and expression of alternatively processed transcripts. *J. Biol. Chem.* 270: 17830–17835, 1995.
- [76238] 14568. Chiang, C.-M.; Roeder, R. G.: Cloning of an intrinsic human TFIID subunit that interacts with multiple transcriptional activators. *Science* 267: 531–536, 1995.
- [76239] 14569. Wu, Q.; Zhang, T.; Cheng, J.-F.; Kim, Y.; Grimwood,

J.; Schmutz, J.; Dickson, M.; Noonan, J. P.; Zhang, M. Q.; Myers, R. M.; Maniatis, T.: Comparative DNA sequence analysis of mouse and human protocadherin gene clusters. *Genome Res.* 11: 389–404, 2001.

[76240] 14570. Aoki, K.; Inazawa, J.; Takahashi, T.; Nakahara, K.; Kasai, M.: Genomic structure and chromosomal localization of the gene encoding translin, a recombination hotspot binding protein. *Genomics* 43:237–241, 1997.

[76241] 14571. Aoki, K.; Nakahara, K.; Ikegawa, C.; Seto, M.; Takahashi, T.; Minowada, J.; Strominger, J. L.; Maziarz, R. T.; Kasai, M.: Nuclear proteins binding to a novel target sequence within the recombination hotspot regions of Bcl-2 and the immunoglobulin D(H) gene family. *Oncogene* 9:1109–1115, 1994.

[76242] 14572. Blair, I. P.; Gibson, R. R.; Bennett, C. L.; Chance, P. F.: Search for genes involved in Joubert syndrome: evidence that one or more major loci are yet to be identified and exclusion of candidate genes EN1, EN2, FGF8, and BARHL1. *Am. J. Med. Genet.* 107: 190–196, 2002.

[76243] 14573. Berge, K. E.; Tian, H.; Graf, G. A.; Yu, L.; Grishin, N. V.; Schultz, J.; Kwiterovich, P.; Shan, B.; Barnes, R.; Hobbs, H. H.: Accumulation of dietary cholesterol in sitosterolemia caused by mutations in adjacent ABC transporters. *Science*

290: 1771–1775, 2000.

- [76244] 14574. Bodzioch, M.; Orso, E.; Klucken, J.; Langmann, T.; Bottcher, A.; Diederich, W.; Drobnik, W.; Barlage, S.; Buchler, C.; Porsch-Ozcurumez, M.; Kaminski, W. E.; Hahmann, H. W.; Oette, K.; Rothe, G.; Aslanidis, C.; Lackner, K. J.; Schmitz, G.: The gene encoding ATP-binding cassette-transporter 1 is mutated in Tangier disease. *Nature Genet.* 22: 347–351, 1999.
- [76245] 14575. Brooks-Wilson, A.; Marcil, M.; Clee, S. M.; Zhang, L.-H.; Roomp, K.; van Dam, M.; Yu, L.; Brewer, C.; Collins, J. A.; Molhuizen, H. O. F.; Loubser, O.; Ouelette, B. F. F.; and 14 others: Mutations in ABC1 in Tangier disease and familial high-density lipoprotein deficiency. *Nature Genet.* 22: 336–345, 1999.
- [76246] 14576. Asamoah, A.; Wilson, A. F.; Elston, R. C.; Dalferes, E., Jr.; Berenson, G. S.: Segregation and linkage analyses of dopamine-beta-hydroxylase activity in a six-generation pedigree. *Am. J. Med. Genet.* 27: 613–621, 1987.
- [76247] 14577. Biaggioni, I.; Goldstein, D. S.; Atkinson, T.; Robertson, D.: Dopamine-beta-hydroxylase deficiency in humans. *Neurology* 40: 370–373, 1990.
- [76248] 14578. Biaggioni, I.; Robertson, D.: Endogenous restoration of noradrenaline by precursor therapy in dopamine-



beta-hydroxylase deficiency. Lancet II:1170–1172, 1987.

[76249] 14579.Craig, S. P.; Buckle, V. J.; Lamouroux, A.; Mallet, J.; Craig, I. W.: Localization of the human dopamine beta hydroxylase (DBH) gene to chromosome 9q34. Cytogenet. Cell Genet. 48: 48–50, 1988.

[76250] 14580.Dunnette, J.; Weinshilboum, R.: Human serum dopamine beta-hydroxylase: correlation of enzymatic activity with immunoreactive protein in genetically defined samples. Am. J. Hum. Genet. 28: 155–166, 1976.

[76251] 14581.Dunnette, J.; Weinshilboum, R.: Inheritance of low immunoreactive human plasma dopamine-beta-hydroxylase: radioimmunoassay studies. J. Clin. Invest. 60: 1080–1087, 1977.

[76252] 14582.Elston, R. C.; Namboodiri, K. K.; Hames, C. G.: Segregation and linkage analysis of dopamine-beta-hydroxylase activity. Hum. Hered. 29:284–292, 1979.

[76253] 14583.Gershon, E. S.; Goldin, L. R.: Segregation and linkage studies of plasma dopamine-beta-hydroxylase (DBH), erythrocyte catechol-O-methyltransferase (COMT) and platelet monoamine oxidase (MAO): possible linkage between the ABO locus and a gene controlling DBH activity. (Abstract) Am. J. Hum. Genet. 33: 136A only, 1981.

- [76254] 14584. Goldin, L. R.; Gershon, E. S.; Lake, C. R.; Murphy, D. L.; McGinniss, M.; Sparkes, R. S.: Segregation and linkage studies of plasma dopamine-beta-hydroxylase (DBH), erythrocyte catechol-O-methyltransferase (COMT), and platelet monoamine oxidase (MAO): possible linkage between the ABO locus and a gene controlling DBH activity. *Am. J. Hum. Genet.* 34: 250-262, 1982.
- [76255] 14585. Joh, T. H.; Baetge, E. E.; Reis, D. J.: Evidence for the existence of a single gene or linked genes coding for catecholamine biosynthetic enzymes. *Trans. Assoc. Am. Phys.* 96: 38-43, 1983.
- [76256] 14586. Joh, T. H.; Baetge, E. E.; Ross, M. E.; Albert, V. R.; Moon, H. M.; Reis, D. J.: Existence of catecholamine biosynthetic enzyme gene family. (Abstract) *Clin. Res.* 31: 528 only, 1983.
- [76257] 14587. Joh, T. H.; Baetge, E. E.; Ross, M. E.; Reis, D. J.: Biochemistry and molecular biology of catecholamine neurons: a single gene or gene family hypothesis. *Clin. Exp. Hypertension* 6A: 11-21, 1984.
- [76258] 14588. Kobayashi, K.; Kurosawa, Y.; Fujita, K.; Nagatsu, T.: Human dopamine beta-hydroxylase gene: two mRNA types having different 3-prime-terminal regions are produced through alternative polyadenylation. *Nucleic Acids Res.* 17:

1089–1102, 1989.

- [76259] 14589.Lamouroux, A.; Vigny, A.; Faucon Biguet, N.; Darmon, M. C.; Franck,R.; Henry, J.-P.; Mallet, J.: The primary structure of human dopamine-beta-hydroxylase:insights into the relationship between the soluble and the membrane-boundforms of the enzyme. *EMBO J.* 6: 3931–3937, 1987.
- [76260] 14590.Lea, R. A.; Dohy, A.; Jordan, K.; Quinlan, S.; Brimage, P. J.;Griffiths, L. R.: Evidence for allelic association of the dopaminebeta-hydroxylase gene (DBH) with susceptibility to typical migraine. *Neurogenetics* 3:35–40, 2000.
- [76261] 14591.Mathias, C. J.; Bannister, R. B.; Cortelli, P.; Heslop, K.; Polak,J. M.; Raimbach, S.; Springall, D. R.; Watson, L.: Clinical, autonomicand therapeutic observations in two siblings with postural hypotensionand sympathetic failure due to an inability to synthesize noradrenalinefrom dopamine because of a deficiency of dopamine beta hydroxylase. *Quart.J. Med.* 75: 617–633, 1990.
- [76262] 14592.McKinney, E. F.; Walton, R. T.; Yudkin, P.; Fuller, A.; Haldar,N. A.; Mant, D.; Murphy, M.; Welsh, K. I.; Marshall, S. E.: Associationbetween polymorphisms in dopamine metabolic enzymes and tobacco consumptionin smokers. *Pharmacogenetics* 10: 483–491, 2000.

- [76263] 14593.O'Malley, K. L.; Maun, A.; Raese, J.; Barchas, J. D.; Kedes,L.: Genes for catecholamine biosynthesis: cloning by expression and identification of the cDNA for rat dopamine beta-hydroxylase. *Proc.Nat. Acad. Sci.* 80: 2161-2165, 1983.
- [76264] 14594.Ogihara, T.; Nugent, C. A., Jr.; Shen, S.-W.; Goldfein, S.: Serum dopamine-beta-hydroxylase activity in parents and children. *J. Lab.Clin. Med.* 85: 566-573, 1975.
- [76265] 14595.Perry, S. E.; Summar, M. L.; Phillips, J. A., III; Robertson,D.: Linkage analysis of the human dopamine beta-hydroxylase gene. *Genomics* 10:493-495, 1991.
- [76266] 14596.Robertson, D.; Haile, V.; Perry, S. E.; Robertson, R. M.; Phillips,J. A., III; Biaggioni, I.: Dopamine beta-hydroxylase deficiency:a genetic disorder of cardiovascular regulation. *Hypertension* 18:1-8, 1991.
- [76267] 14597.Ross, S. B.; Wetterberg, L.; Myrhed, M.: Genetic control of plasma dopamine-beta-hydroxylase. *Life Sci.* 12: 529-532, 1973.
- [76268] 14598.Schanberg, S. M.; Stone, R. A.; Kirshner, N.; Gunnells, J. C.;Robinson, R. R.: Plasma dopamine beta-hydroxylase: a possible aid in the study and evaluation of hypertension. *Science* 183: 523-525,1974.
- [76269] 14599.Marlar, R. A.; Griffin, J. H.: Deficiency of protein C

inhibitorin combined factor V/VIII deficiency disease. J. Clin. Invest. 66:1186–1189, 1980.

- [76270] 14600.Meijers, J. C.; Chung, D. W.: Organization of the gene codingfor human protein C inhibitor (plasminogen activator inhibitor-3):assignment of the gene to chromosome 14. J. Biol. Chem. 266: 15028–15034,1991.
- [76271] 14601.Nichols, W. C.; Seligsohn, U.; Zivelin, A.; Terry, V. H.; Arnold,N. D.; Siemieniak, D. R.; Kaufman, R. J.; Ginsburg, D.: Linkage ofcombined factors V and VIII deficiency to chromosome 18q by homozygositymapping. J. Clin. Invest. 99: 596–601, 1997.
- [76272] 14602.Suzuki, K.; Deyashiki, Y.; Nishioka, J.; Kurachi, K.; Akira, M.;Yamamoto, S.; Hashimoto, S.: Characterization of a cDNA for humanprotein C inhibitor: a new member of the plasma serine protease inhibitorsuperfamily. J. Biol. Chem. 262: 611–616, 1987.
- [76273] 14603.Cathelineau, L.; Dinh, D. P.; Briand, P.; Kamoun, P.: Studieson complementation in argininosuccinate synthetase and argininosuccinatelyase deficiencies in human fibroblasts. Hum. Genet. 57: 282–284,1981.
- [76274] 14604.Kromberg, J. G. R.; Castle, D.; Zwane, E. M.; Jenkins, T.: Albinismand skin cancer in southern Africa. Clin. Genet. 36: 43–52, 1989.

- [76275] 14605.El Kahloun, A.; Chauvel, B.; Mauvieux, V.; Dorval, I.; Jouanolle, A.-M.; Gicquel, I.; Le Gall, J.-Y.; David, V.: Localization of seven new genes around the HLA-A locus. *Hum. Molec. Genet.* 2: 55–60, 1993.
- [76276] 14606.Beattie, E. C.; Stellwagen, D.; Morishita, W.; Bresnahan, J. C.; Ha, B. K.; Von Zastrow, M.; Beattie, M. S.; Malenka, R. C.: Control of synaptic strength by glial TNF- $\alpha$ . *Science* 295: 2282–2285, 2002.
- [76277] 14607.Beutler, B.; Krochin, N.; Milsark, I. W.; Luedke, C.; Cerami, A.: Control of cachectin (tumor necrosis factor) synthesis: mechanisms of endotoxin resistance. *Science* 232: 977–980, 1986.
- [76278] 14608.Brenner, D. A.; O'Hara, M.; Angel, P.; Chojkier, M.; Karin, M.: Prolonged activation of JUN and collagenase genes by tumour necrosis factor- $\alpha$ . *Nature* 337: 661–663, 1989.
- [76279] 14609.Broudy, V. C.; Kaushansky, K.; Segal, G. M.; Harlan, J. M.; Adamson, J. W.: Tumor necrosis factor type  $\alpha$  stimulates human endothelial cells to produce granulocyte/macrophage colony-stimulating factor. *Proc. Nat. Acad. Sci.* 83: 7467–7471, 1986.
- [76280] 14610.Bruce, A. J.; Boling, W.; Kindy, M. S.; Peschon, J.; Kraemer, P. J.; Carpenter, M. K.; Holtsberg, F. W.; Mattson,

M. P.: Altered neuronal and microglial responses to excitotoxic and ischemic brain injury in mice lacking TNF receptors. *Nature Med.* 2: 788–794, 1996.

[76281] 14611. Cabrera, M.; Shaw, M. A.; Sharples, C.; Williams, H.; Castes, M.; Convit, J.; Blackwell, J. M.: Polymorphism in tumor necrosis factor genes associated with mucocutaneous leishmaniasis. *J. Exp. Med.* 182:1259–1264, 1995.

[76282] 14612. Conway, D. J.; Holland, M. J.; Bailey, R. L.; Campbell, A. E.; Mahdi, O. S.; Jennings, R.; Mbeni, E.; Mabey, D. C.: Scarring trachoma is associated with polymorphism in the tumor necrosis factor alpha (TNF-alpha) gene promoter and with elevated TNF-alpha levels in tear fluid. *Infect. Immun.* 65: 1003–1006, 1997.

[76283] 14613. Davis, J. M.; Narachi, M. A.; Alton, N. K.; Arakawa, T.: Structure of human tumor necrosis factor alpha derived from recombinant DNA. *Biochemistry* 26:1322–1326, 1987.

[76284] 14614. Escobar-Morreale, H. F.; Calvo, R. M.; Sancho, J.; San Millan, J. L.: TNF-alpha and hyperandrogenism: a clinical, biochemical, and molecular genetic study. *J. Clin. Endocrin. Metab.* 86: 3761–3767, 2001.

[76285] 14615. Gorman, J. D.; Sack, K. E.; Davis, J. C., Jr.: Treatment of ankylosing spondylitis by inhibition of tumor

necrosis factor- $\alpha$ . *New Eng.J. Med.* 346: 1349–1356, 2002.

- [76286] 14616.Herrmann, S.-M.; Ricard, S.; Nicaud, V.; Mallet, C.; Arveiler,D.; Evans, A.; Ruidavets, J.-B.; Luc, G.; Bara, L.; Parra, H.-J.;Poirier, O.; Cambien, F.: Polymorphisms of the tumour necrosis factor- $\alpha$  gene, coronary heart disease and obesity. *Europ. J. Clin. Invest.* 28:59–66, 1998.
- [76287] 14617.Inoko, H.; Trowsdale, J.: Linkage of TNF genes to the HLA-B locus. *NucleicAcids Res.* 15: 8957–8962, 1987.
- [76288] 14618.Knight, J. C.; Udalova, I.; Hill, A. V. S.; Greenwood, B. M.;Peshu, N.; Marsh, K.; Kwiatkowski, D.: A polymorphism that affectsOCT-1 binding to the TNF promoter region is associated with severemalaria. *Nature Genet.* 22: 145–150, 1999.
- [76289] 14619.Koss, K.; Satsangi, J.; Fanning, G. C.; Welsh, K. I.; Jewell,D. P.: Cytokine (TNF- $\alpha$ , LT- $\alpha$ , and IL-10) polymorphisms ininflammatory bowel diseases and normal controls: differential effectson production and allele frequencies. *Genes Immun.* 1: 185–190, 2000.
- [76290] 14620.Marino, M. W.; Dunn, A.; Grail, D.; Inglese, M.; Noguchi, Y.;Richards, E.; Jungbluth, A.; Wada, H.; Moore, M.; Williamson, B.;Basu, S.; Old, L. J.: Characterization of tumor necrosis factor-deficientmice. *Proc. Nat. Acad. Sci.*



94: 8093–8098, 1997.

[76291] 14621. McCusker, S. M.; Curran, M. D.; Dynan, K. B.; McCullagh, C. D.; Urquhart, D. D.; Middleton, D.; Patterson, C. C.; McIlroy, S. P.; Passmore, A. P.: Association between polymorphism in regulatory region of gene encoding tumour necrosis factor- $\alpha$  and risk of Alzheimer's disease and vascular dementia: a case-control study. *Lancet* 357:436–439, 2001.

[76292] 14622. Remaley, A. T.; Rust, S.; Rosier, M.; Knapper, C.; Naudin, L.; Broccardo, C.; Peterson, K. M.; Koch, C.; Arnould, I.; Prades, C.; Duverger, N.; Funke, H.; Assman, G.; Dinger, M.; Dean, M.; Chimini, G.; Santamarina-Fojo, S.; Fredrickson, D. S.; Deneffe, P. Brewer, H. B., Jr.: Human ATP-binding cassette transporter 1 (ABC1): genomic organization and identification of the genetic defect in the original Tangier disease kindred. *Proc. Nat. Acad. Sci.* 96: 12685–12690, 1999.

[76293] 14623. Rust, S.; Rosier, M.; Funke, H.; Real, J.; Amoura, Z.; Piette, J.-C.; Deleuze, J.-F.; Brewer, H. B.; Duverger, N.; Deneffe, P.; Assmann, G.: Tangier disease is caused by mutations in the gene encoding ATP-binding cassette transporter 1. *Nature Genet.* 22: 352–355, 1999.

[76294] 14624. Young, S. G.; Fielding, C. J.: The ABCs of cholest-

terol efflux. *NatureGenet.* 22: 316–318, 1999.

[76295] 14625.Buchwald, M.: Complementation groups: one or more per gene? *NatureGenet.* 11: 228–230, 1995.

[76296] 14626.Stambolian, D.; Ai, Y.; Sidjanin, D.; Nesburn, K.; Sathe, G.;Rosenberg, M.; Bergsma, D. J.: Cloning of the galactokinase cDNAand identification of mutations in two families with cataracts. *NatureGenet.* 10: 307–312, 1995.

[76297] 14627.Coppola, G.; De Michele, G.; Cavalcanti, F.; Pianese, L.; Perretti,A.; Santoro, L.; Vita, G.; Toscano, A.; Amboni, M.; Grimaldi, G.;Salvatore, E.; Caruso, G.; Filla, A.: Why do some Friedreich's ataxiapatients retain tendon reflexes? A clinical, neurophysiological andmolecular study. *J. Neurol.* 246: 353–357, 1999.

[76298] 14628.Delatycki, M. B.; Knight, M.; Koenig, M.; Cossee, M.; Williamson,R.; Forrest, S. M.: G130V, a common FRDA point mutation, appearsto have arisen from a common founder. *Hum. Genet.* 105: 343–346,1999.

[76299] 14629.Delatycki, M. B.; Paris, D. B. B. P.; Gardner, R. J. M.; Nicholson,G. A.; Nassif, N.; Storey, E.; MacMillan, J. C.; Collins, V.; Williamson,R.; Forrest, S. M.: Clinical and genetic study of Friedreich ataxiain an Australian population. *Am. J. Med. Genet.* 87: 168–174, 1999.

[76300] 14630.Durr, A.; Cossee, M.; Agid, Y.; Campuzano, V.;

Mignard, C.; Penet, C.; Mandel, J.-L.; Brice, A.; Koenig, M.: Clinical and genetic abnormalities in patients with Friedreich's ataxia. *New Eng. J. Med.* 335: 1169–1175, 1996.

[76301] 14631. Filla, A.; De Michele, G.; Cavalcanti, F.; Pianese, L.; Monticelli, A.; Campanella, G.; Coccozza, S.: The relationship between trinucleotide (GAA) repeat length and clinical features in Friedreich ataxia. *Am. J. Hum. Genet.* 59: 554–560, 1996.

[76302] 14632. Fujita, R.; Hanauer, A.; Vincent, A.; Mandel, J.-L.; Koenig, M.: Physical mapping of two loci (D9S5 and D9S15) tightly linked to Friedreich ataxia locus (FRDA) and identification of nearby CpG islands by pulse-field gel electrophoresis. *Genomics* 10: 915–920, 1991.

[76303] 14633. Gray, J. V.; Johnson, K. J.: Waiting for frataxin. *Nature Genet.* 16: 323–325, 1997.

[76304] 14634. Victoria, T.; Rafi, M. A.; Wenger, D. A.: Cloning of the canine GALC cDNA and identification of the mutation causing globoid cell leukodystrophy in West Highland White and Cairn terriers. *Genomics* 33: 457–462, 1996.

[76305] 14635. Wenger, D. A.; Rafi, M. A.; Luzi, P.: Molecular genetics of Krabbe disease (globoid cell leukodystrophy): diagnostic and clinical implications. *Hum. Mutat.* 10: 268–279, 1997.

- [76306] 14636.Zlotogora, J.; Chakraborty, S.; Knowlton, R. G.; Wenger, D. A.: Krabbe disease locus mapped to chromosome 14 by genetic linkage. *Am.J. Hum. Genet.* 47: 37–44, 1990.
- [76307] 14637.Zlotogora, J.; Regev, R.; Zeigler, M.; Iancu, T. C.; Bach, G.: Krabbe disease: increased incidence in a highly inbred community. *Am.J. Med. Genet.* 21: 765–770, 1985.
- [76308] 14638.Rushton, A. R.; Dawson, G.: Genetic linkage studies of the humanglycosphingolipid beta-galactosidases. *Biochem. Genet.* 15: 1071–1082,1977.
- [76309] 14639.Balciunaite, G.; Keller, M. P.; Balciunaite, E.; Piali, L.; Zuklys,S.; Mathieu, Y. D.; Gill, J.; Boyd, R.; Sussman, D. J.; Hollander,G. A.: Wnt glycoproteins regulate the expression of FoxN1, the genedefective in nude mice. *Nature Immun.* 15Oct, 2002. Note: AdvanceElectronic Publication.
- [76310] 14640.Frank, J.; Pignata, C.; Panteleyev, A. A.; Prowse, D. M.; Baden,H.; Weiner, L.; Gaetaniello, L.; Ahmad, W.; Pozzi, N.; Caerhalmi–Friedman,P. B.; Aita, V. M.; Uyttendaele, H.; Gordon, D.; Ott, J.; Brissette,J. L.; Christiano, A. M.: Exposing the human nude phenotype. *Nature* 398:473–474, 1999.
- [76311] 14641.Nehls, M.; Pfeifer, D.; Schorpp, M.; Hedrich, H.; Boehm, T.: Newmember of the winged–helix protein fam–

ily disrupted in mouse and rat nude mutations. *Nature* 372: 103–107, 1994.

- [76312] 14642. Pignata, C.; Fiore, M.; Guzzetta, V.; Castaldo, A.; Sebastio, G.; Porta, F.; Guarino, A.: Congenital alopecia and nail dystrophy associated with severe functional T-cell immunodeficiency in two sibs. *Am. J. Med. Genet.* 65: 167–170, 1996.
- [76313] 14643. Schorpp, M.; Hofmann, M.; Dear, T. N.; Boehm, T.: Characterization of mouse and human nude genes. *Immunogenetics* 46: 509–515, 1997.
- [76314] 14644. Segre, J. A.; Nemhauser, J. L.; Taylor, B. A.; Nadeau, J. H.; Lander, E. S.: Positional cloning of the nude locus: genetic, physical, and transcription maps of the region and mutations in the mouse and rat. *Genomics* 28: 549–559, 1995.
- [76315] 14645. Delpire, E.; Lu, J.; England, R.; Dull, C.; Thorne, T.: Deafness and imbalance associated with inactivation of the secretory Na–K–2Cl co-transporter. *Nature Genet.* 22: 192–195, 1999.
- [76316] 14646. Evans, R. L.; Park, K.; Turner, R. J.; Watson, G. E.; Nguyen, H.-V.; Dennett, M. R.; Hand, A. R.; Flagella, M.; Shull, G. E.; Melvin, J. E.: Severe impairment of salivation in Na<sup>+</sup>/K<sup>+</sup>/2Cl<sup>–</sup> cotransporter (NKCC1)-deficient mice. *J.*

Biol. Chem. 275: 26720–26726, 2000.

- [76317] 14647. Payne, J. A.; Xu, J.-C.; Haas, M.; Lytle, C. Y.; Ward, D.; Forbush, B., III: Primary structure, functional expression, and chromosomal localization of the bumetanide-sensitive Na-K-Cl cotransporter in human colon. *J. Biol. Chem.* 270: 17977–17985, 1995.
- [76318] 14648. Quaggin, S. E.; Payne, J. A.; Forbush, B., III; Igarashi, P.: Localization of the renal Na-K-Cl cotransporter gene (Slc12a1) on mouse chromosome 2. *Mammalian Genome* 6: 557–561, 1995.
- [76319] 14649. Xu, J.-C.; Lytle, C.; Zhu, T. T.; Payne, J. A.; Benz, E., Jr.; Forbush, B., III: Molecular cloning and functional expression of the bumetanide-sensitive Na-K-Cl cotransporter. *Proc. Nat. Acad. Sci.* 91: 2201–2205, 1994.
- [76320] 14650. Daniels, S. E.; Bhattacharya, S.; James, A.; Leaves, N. I.; Young, A.; Hill, M. R.; Faux, J. A.; Ryan, G. F.; le Souef, P. N.; Lathrop, G. M.; Musk, A. W.; Cookson, W. O. C. M.: A genome-wide search for quantitative trait loci underlying asthma. *Nature* 383: 247–250, 1996.
- [76321] 14651. MacDonald, S. M.; Paznekas, W. A.; Jabs, E. W.: Chromosomal localization of tumor protein, translationally-controlled 1 (TPT1) encoding the human histamine releasing factor (HRF) to 13q12–q14. *Cytogenet. Cell Genet.*

(in-press), 1999.

- [76322] 14652. MacDonald, S. M.; Rafnar, T.; Langdon, J.; Lichtenstein, L. M.: Molecular identification of an IgE-dependent histamine-releasing factor. *Science* 269: 688–690, 1995.
- [76323] 14653. Zhang, K.; Saxon, A.; Max, E. E.: Two unusual forms of human immunoglobulin E encoded by alternative RNA splicing of the epsilon heavy chain membrane exons. *J. Exp. Med.* 176: 233–243, 1992.
- [76324] 14654. Simister, N. E.; Mostov, K. E.: An Fc receptor structurally related to MHC class I antigens. *Nature* 337: 184–187, 1989.
- [76325] 14655. Sedlacek, Z.; Konecki, D. S.; Korn, B.; Klauck, S. M.; Poustka, A.: Evolutionary conservation and genomic organization of XAP-4, an Xq28 located gene coding for a human rab GDP-dissociation inhibitor (GDI). *Mammalian Genome* 5: 633–639, 1995.
- [76326] 14656. Sedlacek, Z.; Munstermann, E.; Mincheva, A.; Lichter, P.; Poustka, A.: The human rab GDI beta gene with long retroposon-rich introns maps to 10p15 and its pseudogene to 7p11–p13. *Mammalian Genome* 9: 78–80, 1998.
- [76327] 14657. Desseyn, J.-L.; Aubert, J.-P.; van Seuningen, I.; Porchet, N.; Laine, A.: Genomic organization of the

3-prime region of the human mucin gene MUC5B. J. Biol. Chem. 272: 16873–16883, 1997.

- [76328] 14658. Desseyn, J.-L.; Guyonnet-Duperat, V.; Porchet, N.; Aubert, J.-P.; Laine, A.: Human mucin gene MUC5B, the 10.7-kb large central exon encodes various alternate subdomains resulting in a super-repeat: structural evidence for a 11p15.5 gene family. J. Biol. Chem. 272:3168–3178, 1997.
- [76329] 14659. Gipson, I. K.; Moccia, R.; Spurr-Michaud, S.; Argueso, P.; Gargiulo, A. R.; Hill, J. A., III; Offner, G. D.; Keutmann, H. T.: The amount of MUC5B in cervical mucus peaks at midcycle. J. Clin. Endocr. Metab. 86:594–600, 2001.
- [76330] 14660. Keates, A. C.; Nunes, D. P.; Afdhal, N. H.; Troxler, R. F.; Offner, G. D.: Molecular cloning of a major human gall bladder mucin: complete C-terminal sequence and genomic organization of MUC5B. Biochem. J. 324:295–303, 1997.
- [76331] 14661. Troxler, R. F.; Offner, G. D.; Zhang, F.; Iontcheva, I.; Oppenheim, F. G.: Molecular cloning of a novel high molecular weight mucin (MG1) from human sublingual gland. Biochem. Biophys. Res. Commun. 217:1112–1119, 1995.



- [76332] 14662.Birck, C.; Poch, O.; Romier, C.; Ruff, M.; Mengus, G.; Lavigne,A.-C.; Davidson, I.; Moras, D.: Human TAFII28 and TAFII18 interactthrough a histone fold encoded by atypical evolutionary conservedmotifs also found in the SPT3 family. *Cell* 94: 239–249, 1998.
- [76333] 14663.Mengus, G.; May, M.; Jacq, X.; Staub, A.; Tora, L.; Chambon, P.;Davidson, I.: Cloning and characterization of hTAFII18, hTAFII20and hTAFII28: three subunits of the human transcription factor TFIID. *EMBOJ.* 14: 1520–1531, 1995.
- [76334] 14664.Waisfisz, Q.; Saar, K.; Morgan, N. V.; Altay, C.; Leegwater, P.A.; de Winter, J. P.; Komatsu, K.; Evans, G. R.; Wegner, R.-D.; Reis,A.; Joenje, H.; Arwert, F.; Mathew, C. G.; Pronk, J. C.; Digweed,M.: The Fanconi anemia group E gene, FANCE, maps to chromosome 6p. *Am.J. Hum. Genet.* 64: 1400–1405, 1999.
- [76335] 14665.Wegner, R.-D.; Henrichs, I.; Joenje, H.; Schroeder-Kurth, T.:Fanconi anemia complementation group E: clinical and cytogenetic dataof the first patient. *Clin. Genet.* 50: 479–482, 1996.
- [76336] 14666.Green, P.; Lipman, D.; Hillier, L.; Waterston, R.; States, D.;Claverie, J.-M.: Ancient conserved regions in new gene sequencesand the protein databases. *Science*

259: 1711–1716, 1993.

- [76337] 14667.Schwartz, F.; Eisenman, R.; Knoll, J.; Gessler, M.; Bruns, G.:cDNA sequence, genomic organization, and evolutionary conservation of a novel gene from the WAGR region. *Genomics* 29: 526–532, 1995.
- [76338] 14668.Schwartz, F.; Neve, R.; Eisenman, R.; Gessler, M.; Bruns, G.:A WAGR region gene between PAX–6 and FSHB expressed in fetal brain. *Hum.Genet.* 94: 658–664, 1994.
- [76339] 14669.Krishnan, B. R.; Jamry, I.; Chaplin, D. D.: Feature mapping of the HLA class I region: localization of the POU5F1 and TCF19 genes. *Genomics* 30:53–58, 1995.
- [76340] 14670.Ku, D.–H.; Chang, C.; Koniecki, J.; Cannizzaro, L. A.; Boghosian–Sell,L.; Alder, H.; Baserga, R.: A new growth–regulated complementaryDNA with the sequence of a putative trans–activating factor. *CellGrowth Differ.* 2: 179–186, 1991.
- [76341] 14671.Joseph, R. E.; Walker, J.; Norris, F. A.: Assignment of the inositolpolyphosphate 4–phosphatase type I gene (INPP4A) to human chromosomeband 2q11.2 by in situ hybridization. *Cytogenet. Cell Genet.* 87:276–277, 1999.
- [76342] 14672.Norris, F. A.; Auethavekiat, V.; Majerus, P. W.: The isolationand characterization of cDNA encoding human and rat brain inositolpolyphosphate 4–phosphatase. *J.*

Biol. Chem. 270: 16128–16133, 1995.

- [76343] 14673. van den Maagdenberg, A. M. J. M.; van den Hurk, H. H.; Olde Weghuis, D.; Wieringa, B.; Geurts van Kessel, A.; Hendriks, W. J. A. J.: Assignment of the human protein tyrosine phosphatase epsilon (PTPRE) gene to chromosome 10q26 by fluorescence in situ hybridization. *Genomics* 30:128–129, 1995.
- [76344] 14674. Kim, D. S.; Jung, H.-H.; Park, S.-H.; Chin, H.: Isolation and characterization of the 5-prime-upstream region of the human N-type calcium channel alpha-1B subunit gene: chromosomal localization and promoter analysis. *J. Biol. Chem.* 272: 5098–5104, 1997.
- [76345] 14675. Williams, M. E.; Brust, P. F.; Feldman, D. H.; Patthi, S.; Simerson, S.; Maroufi, A.; McCue, A. F.; Velicelebi, G.; Ellis, S. B.; Harpold, M. M.: Structure and functional expression of an omega-conotoxin-sensitive human N-type calcium channel. *Science* 257: 389–395, 1992.
- [76346] 14676. Azim, A. C.; Knoll, J. H. M.; Marfatia, S. M.; Peel, D. J.; Bryant, P. J.; Chishti, A. H.: DLG1: chromosome location of the closest human homologue of the *Drosophila* discs large tumor suppressor gene. *Genomics* 30:613–616, 1995.
- [76347] 14677. Burgess, D. L.; Rafael, J. A.; Meisler, M. H.; Cham-

berlain, J.S.: Dlgh1, a mouse homolog of the Drosophila discs-large gene, is located on chromosome 16. Mammalian Genome 7: 623–624, 1996.

[76348] 14678. Hanada, T.; Lin, L.; Chandy, K. G.; Oh, S. S.; Chishti, A. H.: Human homologue of the Drosophila discs large tumor suppressor binds to p56lck tyrosine kinase and Shaker type Kv1.3 potassium channel in T lymphocytes. J. Biol. Chem. 272: 26899–26904, 1997.

[76349] 14679. Lue, R. A.; Marfatia, S. M.; Branton, D.; Chishti, A. H.: Cloning and characterization of hdlg: the human homologue of the Drosophila discs large tumor suppressor binds to protein 4.1. Proc. Nat. Acad. Sci. 91: 9818–9822, 1994.

[76350] 14680. Mori, K.; Iwao, K.; Miyoshi, Y.; Nakagawara, A.; Kofu, K.; Akiyama, T.; Arita, N.; Hayakawa, T.; Nakamura, Y.: Identification of brain-specific splicing variants of the hDLG1 gene and altered splicing in neuroblastoma cell lines. J. Hum. Genet. 43: 123–127, 1998.

[76351] 14681. Peters, L. L.; Ciciotte, S. L.; Lin, L.; Chishti, A. H.: The mouse homolog of the Drosophila discs large tumor suppressor gene maps to chromosome 16. Mammalian Genome 7: 619–620, 1996.

[76352] 14682. Bertaux, F.; Sharp, A. H.; Ross, C. A.; Lehrach, H.;

Bates, G.P.; Wanker, E.: HAP1–huntingtin interactions do not contribute to the molecular pathology in Huntington's disease transgenic mice. *FEBS Lett.* 426: 229–232, 1998.

[76353] 14683. Engelender, S.; Sharp, A. H.; Colomer, V.; Tokito, M. K.; Lanahan, A.; Worley, P.; Holzbaun, E. L. F.; Ross, C. A.: Huntingtin–associated protein 1 (HAP1) interacts with the p150(Glued) subunit of dynactin. *Hum. Molec. Genet.* 6: 2205–2212, 1997.

[76354] 14684. Li, S.–H.; Hosseini, S. H.; Gutekunst, C.–A.; Hersch, S. M.; Ferrante, R. J.; Li, X.–J.: A human HAP1 homologue: cloning, expression, and interaction with huntingtin. *J. Biol. Chem.* 273: 19220–19227, 1998.

[76355] 14685. Li, X.–J.; Li, S.–H.; Sharp, A. H.; Nucifora, F. C., Jr.; Schilling, G.; Lanahan, A.; Worley, P.; Snyder, S. H.; Ross, C. A.: A huntingtin–associated protein enriched in brain with implications for pathology. *Nature* 378: 398–402, 1995.

[76356] 14686. Nasir, J.; Duan, K.; Nichol, K.; Engelender, S.; Ashworth, R.; Colomer, V.; Thomas, S.; Distèche, C. M.; Hayden, M. R.; Ross, C. A.: Gene structure and map location of the murine homolog of the Huntington–associated protein, Hap1. *Mammalian Genome* 9: 565–570, 1998.

[76357] 14687. Nasir, J.; Lafuente, M.–J.; Duan, K.; Colomer, V.; Engelender, S.; Ingersoll, R.; Margolis, R. L.; Ross, C. A.;

Hayden, M. R.: Human huntingtin-associated protein (HAP-1) gene: genomic organisation and an intragenic polymorphism. *Gene* 254: 181–187, 2000.

[76358] 14688. Broccoli, D.; Chong, L.; Oelmann, S.; Fernald, A. A.; Marziliano, N.; van Steensel, B.; Kipling, D.; Le Beau, M. M.; de Lange, T.: Comparison of the human and mouse genes encoding the telomeric protein, TRF1: chromosomal localization, expression and conserved protein domains. *Hum. Molec. Genet.* 6: 69–76, 1997.

[76359] 14689. Chong, L.; van Steensel, B.; Broccoli, D.; Erdjument-Bromage, H.; Hanish, J.; Tempst, P.; de Lange, T.: A human telomeric protein. *Science* 270: 1663–1667, 1995.

[76360] 14690. Fairall, L.; Chapman, L.; Moss, H.; de Lange, T.; Rhodes, D.: Structure of the TRFH dimerization domain of the human telomeric proteins TRF1 and TRF2. *Molec. Cell* 8: 351–361, 2001.

[76361] 14691. Griffith, J. D.; Comeau, L.; Rosenfield, S.; Stansel, R. M.; Bianchi, A.; Moss, H.; de Lange, T.: Mammalian telomeres end in a large duplex loop. *Cell* 97: 503–514, 1999.

[76362] 14692. Kim, S.; Kaminker, P.; Campisi, J.: TIN2, a new regulator of telomere length in human cells. *Nature Genet.* 23: 405–412, 1999.

[76363] 14693. Marcand, S.; Gilson, E.; Shore, D.: A protein–

counting mechanism for telomere length regulation in yeast. *Science* 275: 986–990, 1997.

- [76364] 14694. Okabe, J.; Eguchi, A.; Masago, A.; Hayakawa, T.; Nakanishi, M.: TRF1 is a critical trans-acting factor required for de novo telomere formation in human cells. *Hum. Molec. Genet.* 9: 2639–2650, 2000.
- [76365] 14695. Young, A. C.; Chavez, M.; Giambernardi, T. A.; Mattern, V.; McGill, J. R.; Harris, J. M.; Sarosdy, M. F.; Patel, P.; Sakaguchi, A. Y.: Organization and expression of human telomere repeat binding factor genes. *Somat. Cell Molec. Genet.* 23: 275–286, 1997.
- [76366] 14696. Zakian, V. A.: Telomeres: beginning to understand the end. *Science* 270: 1601–1607, 1995.
- [76367] 14697. Brodsky, G.; Otterson, G. A.; Parry, B. B.; Hart, I.; Patterson, D.; Kaye, F. J.: Localization of STCH to human chromosome 21q11.1. *Genomics* 30: 627–628, 1995.
- [76368] 14698. Otterson, G. A.; Flynn, G. C.; Kratzke, R. A.; Coxon, A.; Johnston, P. G.; Kaye, F. J.: Stch encodes the 'ATPase core' of a microsomal stress 70 protein. *EMBO J.* 13: 1216–1225, 1994.
- [76369] 14699. Reeves, R. H.; Rue, E.; Yu, J.; Kao, F.-T.: Stch maps to mouse chromosome 16, extending the conserved synteny with human chromosome 21. *Genomics* 49: 156–157,

1998.

- [76370] 14700.Nielsen, P. J.; Rochelle, J. M.; Seldin, M. F.: The functional genes for protein synthesis initiation factor 4A1 and 4A2 map to mouse chromosomes 11 and 16. *Mammalian Genome* 4: 185–186, 1993.
- [76371] 14701.Nielsen, P. J.; Trachsel, H.: The mouse protein synthesis initiation factor 4A gene family includes two related functional genes which are differentially expressed. *EMBO J.* 7: 2097–2105, 1988.
- [76372] 14702.Sudo, K.; Takahashi, E.; Nakamura, Y.: Isolation and mapping of the human EIF4A2 gene homologous to the murine protein synthesis initiation factor 4A-II gene Eif4a2. *Cytogenet. Cell Genet.* 71: 385–388, 1995.
- [76373] 14703.Gladyshev, V. N.; Jeang, K.-T.; Stadtman, T. C.: Selenocysteine, identified as the penultimate C-terminal residue in human T-cell thioredoxin reductase, corresponds to TGA in the human placental gene. *Proc. Nat. Acad. Sci.* 93: 6146–6151, 1996.
- [76374] 14704.Zhang, K.; Lindsberg, P. J.; Tatlisumak, T.; Kaste, M.; Olsen, H. S.; Andersson, L. C.: Stanniocalcin: a molecular guard of neurons during cerebral ischemia. *Proc. Nat. Acad. Sci.* 97: 3637–3642, 2000.
- [76375] 14705.Gillett, G. T.; Fox, M. F.; Rowe, P. S. N.; Casimir, C.



M.; Povey,S.: Mapping of human non-muscle type cofilin (CFL1) to chromosome11q13 and muscle-type cofilin (CFL2) to chromosome 14. *Ann. Hum.Genet.* 60: 201–211, 1996.

[76376] 14706.Ogawa, K.; Tashima, M.; Yumoto, Y.; Okuda, T.; Sawada, H.; Okuma,M.; Maruyama, Y.: Coding sequence of human placenta cofilin cDNA. *Nuc.Acids Res.* 18: 7169 only, 1990.

[76377] 14707.Ono, S.; Minami, N.; Abe, H.; Obinata, T.: Characterization of a novel cofilin isoform that is predominantly expressed in mammalian skeletal muscle. *J. Biol. Chem.* 269: 15280–15286, 1994.

[76378] 14708.Georgakopoulos, T.; Thireos, G.: Two distinct yeast transcriptional activators require the function of the GCN5 protein to promote normal levels of transcription. *EMBO J.* 11: 4145–4152, 1992.

[76379] 14709.Inoue, M.; Isomura, M.; Ikegawa, S.; Fujiwara, T.; Shin, S.; Moriya,H.; Nakamura, Y.: Isolation and characterization of a human cDNA clone (GCN5L1) homologous to GCN5, a yeast transcription factor. *Cytogenet.Cell Genet.* 73: 134–136, 1996.

[76380] 14710.Emahazion, T.; Beskow, A.; Gyllenstein, U.; Brookes, A. J.: Intron based radiation hybrid mapping of 15 complex

I genes of the humanelectron transport chain. Cytogenet. Cell Genet. 82: 115–119, 1998.

- [76381] 14711.Gu, J. Z.; Lin, X.; Wells, D. E.: The human B22 subunit of theNADH–ubiquinone oxidoreductase maps to the region of chromosome 8involved in branchio–oto–renal syndrome. Genomics 35: 6–10, 1996.
- [76382] 14712.Lin, X.; Wells, D. E.; Kimberling, W. J.; Kumar, S.: Human NDUFB9gene: genomic organization and a possible candidate gene associatedwith deafness disorder mapped to chromosome 8q13. Hum. Hered. 49:75–80, 1999.
- [76383] 14713.Wilkinson, K. D.; Tashayev, V. L.; O'Connor, L. B.; Larsen, C.N.; Kasperek, E.; Pickart, C. M.: Metabolism of the polyubiquitin degradation signal: structure, mechanism, and role of isopeptidaseT. Biochemistry 34: 14535–14546, 1995.
- [76384] 14714.Byrd, P. J.; Cooper, P. R.; Stankovic, T.; Kullar, H. S.; Watts,G. D. J.; Robinson, P. J.; Taylor, M. R.: A gene transcribed fromthe bidirectional ATM promoter coding for a serine rich protein: aminoacid sequence, structure and expression studies. Hum. Molec. Genet. 5:1785–1791, 1996.
- [76385] 14715.Chen, X.; Yang, L.; Udar, N.; Liang, T.; Uhrhammer,

N.; Xu, S.; Bay, J.-O.; Wang, Z.; Dandekar, S.; Chiplunkar, S.; Klisak, I.; Telatar, M.; Yang, H.; Concannon, P.; Gatti, R. A.: CAND3: a ubiquitously expressed gene immediately adjacent and in opposite transcriptional orientation to the ATM gene at 11q23.1. *Mammalian Genome* 8: 129–133, 1997.

[76386] 14716. Imai, T.: Personal Communication. Chiba, Japan 9/12/1996.

[76387] 14717. Imai, T.; Sugawara, T.; Nishiyama, A.; Shimada, R.; Ohki, R.; Seki, N.; Sagara, M.; Ito, H.; Yamauchi, M.; Hori, T.: The structure and organization of the human NPAT gene. *Genomics* 42: 388–392, 1997.

[76388] 14718. Imai, T.; Yamauchi, M.; Seki, N.; Sugawara, T.; Saito, T.; Matsuda, Y.; Ito, H.; Nagase, T.; Nomura, N.; Hori, T.: Identification and characterization of a new gene physically linked to the ATM gene. *Genome Res.* 6: 439–447, 1996.

[76389] 14719. Borowsky, B.; Hoffman, B. J.: Analysis of a gene encoding two glycine transporter variants reveals alternative promoter usage and a novel gene structure. *J. Biol. Chem.* 273: 29077–29085, 1998.

[76390] 14720. Borowsky, B.; Mezey, E.; Hoffman, B. J.: Two glycine transporter variants with distinct localization in the CNS

and peripheral tissues are encoded by a common gene.

Neuron 10: 851–863, 1993.

[76391] 14721. Jones, E. M. C.; Fernald, A.; Bell, G. I.; Le Beau, M. M.: Assignment of SLC6A9 to human chromosome band 1p33 by in situ hybridization. Cytogenet. Cell Genet. 71: 211, 1995.

[76392] 14722. Kim, K.-M.; Kingsmore, S. F.; Han, H.; Yang-Feng, T. L.; Godinot, N.; Seldin, M. F.; Caron, M. G.; Giros, B.: Cloning of the human glycine transporter type 1: molecular and pharmacological characterization of novel isoform variants and chromosomal localization of the gene in the human and mouse genomes. Molec. Pharmacol. 45: 608–617, 1994.

[76393] 14723. Freeman, G. J.; Gribben, J. G.; Boussiotis, V. A.; Ng, J. W.; Restivo, V. A.; Lombard, L. A.; Gray, G. S.; Nadler, L. M. Cloning of B7-2: a CTLA-4 counter-receptor that costimulates human T cell proliferation. Science 262: 909–911, 1993.

[76394] 14724. Jeannin, P.; Magistrelli, G.; Aubry, J.-P.; Caron, G.; Gauchat, J.-F.; Renno, T.; Herbault, N.; Goetsch, L.; Blaecke, A.; Dietrich, P.-Y.; Bonnefoy, J.-Y.; Delneste, Y.: Soluble CD86 is a costimulatory molecule for human T lymphocytes. Immunity 13: 303–312, 2000.

- [76395] 14725.Jellis, C. L.; Wang, S. S.; Rennert, P.; Borriello, F.; Sharpe,A. H.; Green, N. R.; Gray, G. S.: Genomic organization of the genecoding for the costimulatory human B-lymphocyte antigen B7-2 (CD86). Immunogenetics 42:85-89, 1995.
- [76396] 14726.Arai, Y.; Hosoda, F.; Kobayashi, H.; Arai, K.; Hayashi, Y.; Kamada,N.; Kaneko, Y.; Ohki, M.: The inv(11)(p15q22) chromosome translocationof de novo and therapy-related myeloid malignancies results in fusionof the nucleoporin gene, NUP98, with the putative RNA heli-case gene,DDX10. Blood 89: 3936-3944, 1997.
- [76397] 14727.Dash, A. B.; Williams, I. R.; Kutok, J. L.; Tomasson, M. H.; Anastasiadou,E.; Lindahl, K.; Li, S.; Van Etten, R. A.; Borrow, J.; Housman, D.;Druker, B.; Gilliland, D. G.: A murine model of CML blast crisisinduced by cooperation between BCR/ABL and NUP98/HOXA9. Proc. Nat.Acad. Sci. 99: 7622-7627, 2002.
- [76398] 14728.Enninga, J.; Levy, D. E.; Blobel, G.; Fontoura, B. M. A.: Roleof nucleoporin induction in releasing an mRNA nuclear export block. Science 295:1523-1525, 2002.
- [76399] 14729.Fontoura, B. M.; Blobel, G.; Matunis, M. J.: A con-served biogenesispathway for nucleoporins: proteolytic processing of a 186-kilodaltonprecursor generates Nup98

and the novel nucleoporin, Nup96. *J. Cell Biol.* 144: 1097–1112, 1999.

[76400] 14730. Hodel, A. E.; Hodel, M. R.; Griffis, E. R.; Hennig, K. A.; Ratner, G. A.; Xu, S.; Powers, M. A.: The three-dimensional structure of the autoproteolytic, nuclear pore-targeting domain of the human nucleoporin Nup98. *Molec. Cell* 10: 347–358, 2002.

[76401] 14731. Jaju, R. J.; Fidler, C.; Haas, O. A.; Strickson, A. J.; Watkins, F.; Clark, K.; Cross, N. C. P.; Cheng, J.-F.; Aplan, P. D.; Kearney, L.; Boulton, J.; Wainscoat, J. S.: A novel gene, NSD1, is fused to NUP98 in the t(5;11)(q35;p15.5) in de novo childhood acute myeloid leukemia. *Blood* 98: 1264–1267, 2001.

[76402] 14732. Jaju, R. J.; Haas, C. A.; Neat, M.; et al: A new recurrent translocation, t(5;11)(q35p15.5), associated with del(5q) in childhood acute myeloid leukemia. *Blood* 94: 773–780, 1999.

[76403] 14733. Mizuno, T.; Kaibuchi, K.; Yamamoto, T.; Kawamura, M.; Sakoda, T.; Fujioka, H.; Matsuura, Y.; Takai, Y.: A stimulatory GDP/GTP exchange protein for smg p21 is active on the post-translationally processed form of c-Ki-ras p21 and rhoA p21. *Proc. Nat. Acad. Sci.* 88: 6442–6446, 1991.

[76404] 14734. Nakamura, T.; Largaespada, D. A.; Lee, M. P.; John-

son, L. A.; Ohyashiki, K.; Toyama, K.; Chen, S. J.; Willman, C. L.; Chen, I.-M.; Feinberg, A. P.; Jenkins, N. A.; Copeland, N. G.; Shaughnessy, J.D., Jr.: Fusion of the nucleoporin gene NUP98 to HOXA9 by the chromosometranslocation t(7;11)(p15;p15) in human myeloid leukaemia. *Nature-Genet.* 12: 154–158, 1996.

[76405] 14735. Radu, A.; Moore, M. S.; Blobel, G.: The peptide repeat domain of nucleoporin Nup98 functions as a docking site in transport across the nuclear pore complex. *Cell* 81: 215–222, 1995.

[76406] 14736. Rosati, R.; La Starza, R.; Veronese, A.; Aventin, A.; Schwienbacher, C.; Vallespi, T.; Negrini, M.; Martelli, M. F.; Mecucci, C.: NUP98 is fused to the NSD3 gene in acute myeloid leukemia associated with t(8;11)(p11.2;p15). *Blood* 99: 3857–3860, 2002.

[76407] 14737. von Kobbe, C.; van Deursen, J. M. A.; Rodrigues, J. P.; Sitterlin, D.; Bachi, A.; Wu, X.; Wilm, M.; Carmo-Fonseca, M.; Izaurralde, E.: Vesicular stomatitis virus matrix protein inhibits host cell gene expression by targeting the nucleoporin Nup98. *Molec. Cell* 6: 1243–1252, 2000.

[76408] 14738. Cheng, Y.; Austin, S. C.; Rocca, B.; Koller, B. H.; Coffman, T.M.; Grosser, T.; Lawson, J. A.; FitzGerald, G. A.: Role of prostacyclin in the cardiovascular response to

thromboxane A<sub>2</sub>. Science 296: 539–541,2002.

- [76409] 14739.Fuse, I.; Mito, M.; Hattori, A.; Higuchi, W.; Shibata, A.; Ushikubi,F.; Okuma, M.; Yahata, K.: Defective signal transduction inducedby thromboxane A<sub>2</sub> in a patient with a mild bleeding disorder: impairedphospholipase C activation despite normal phospholipase A<sub>2</sub> activation. Blood 81:994–1000, 1993.
- [76410] 14740.Hirata, M.; Hayashi, Y.; Ushikubi, F.; Yokota, Y.; Kageyama, R.;Nakanishi, S.; Narumiya, S.: Cloning and expression of cDNA for a human thromboxane A<sub>2</sub> receptor. Nature 349: 617–620, 1991.
- [76411] 14741.Hirata, T.; Kakizuka, A.; Ushikubi, F.; Fuse, I.; Okuma, M.; Narumiya,S.: Arg60–to–leu mutation of the human thromboxane A<sub>2</sub> receptor ina dominantly inherited bleeding disorder. J. Clin. Invest. 94: 1662–1667,1994.
- [76412] 14742.Watanabe, O.; Maruyama, I.; Arimura, K.; Kitajima, I.; Arimura,H.; Hanatani, M.; Matsuo, K.; Arisato, T.; Osame, M.: Overproductionof vascular endothelial growth factor/vascular permeability factoris causative in Crow–Fukase (POEMS) syndrome. Muscle Nerve 21: 1390–1397,1998.
- [76413] 14743.Wei, M.–H.; Popescu, N. C.; Lerman, M. I.; Merrill, M. J.; Zimonjic,D. B.: Localization of the human vascular



endothelial growth factor gene, VEGF, at chromosome 6p12. Hum. Genet. 97: 794–797, 1996.

[76414] 14744. Wong, A. K.; Alfert, M.; Castrillon, D. H.; Shen, Q.; Holash, J.; Yancopoulos, G. D.; Chin, L.: Excessive tumor-elaborated VEGF and its neutralization define a lethal paraneoplastic syndrome. Proc. Nat. Acad. Sci. 98: 7481–7486, 2001.

[76415] 14745. Wulff, C.; Wilson, H.; Lague, P.; Duncan, W. C.; Armstrong, D. G.; Fraser, H. M.: Angiogenesis in the human corpus luteum: localization and changes in angiopoietins, Tie-2, and vascular endothelial growth factor messenger ribonucleic acid. J. Clin. Endocr. Metab. 85: 4302–4309, 2000.

[76416] 14746. Ylikorkala, A.; Rossi, D. J.; Korsisaari, N.; Luukko, K.; Alitalo, K.; Henkemeyer, M.; Makela, T. P.: Vascular abnormalities and deregulation of VEGF in Lkb1-deficient mice. Science 293: 1323–1326, 2001.

[76417] 14747. Bodner, M.; Fridkin, M.; Gozes, I.: Coding sequences for vasoactive intestinal peptide and PHM-27 peptide are located on two adjacent exons in the human genome. Proc. Nat. Acad. Sci. 82: 3548–3551, 1985.

[76418] 14748. Delgado, M.; Abad, C.; Martinez, C.; Leceta, J.; Gomariz, R. P.: Vasoactive intestinal peptide prevents experi-

mental arthritis by downregulating both autoimmune and inflammatory components of the disease. *Nature Med.* 7: 563–568, 2001.

- [76419] 14749. Gotoh, E.; Yamagami, T.; Yamamoto, H.; Okamoto, H.: Chromosomal assignment of human VIP/PHM-27 gene to 6q26–q27 region by spot blot hybridization and in situ hybridization. *Biochem. Int.* 17: 555–562, 1988.
- [76420] 14750. Gozes, I.; Avidor, R.; Yahav, Y.; Katznelson, D.; Croce, C. M.; Huebner, K.: The gene encoding vasoactive intestinal peptide is located on human chromosome 6p21–6pter. *Hum. Genet.* 75: 41–44, 1987.
- [76421] 14751. Gozes, I.; Nakai, H.; Byers, M.; Avidor, R.; Weinstein, Y.; Shani, Y.; Shows, T. B.: Sequential expression in the nervous system of C-MYB and VIP genes, located in human chromosomal region 6q24. *Somat. Cell Molec. Genet.* 13: 305–313, 1987.
- [76422] 14752. Heinz-Erian, P.; Dey, R. D.; Flux, M.; Said, S. I.: Deficient vasoactive intestinal peptide innervation in sweat glands of cystic fibrosis patients. *Science* 229: 1407–1408, 1985.
- [76423] 14753. Itoh, N.; Obata, K.; Yanaihara, N.; Okamoto, H.: Human preprovasoactive intestinal polypeptide contains a novel PHI-27-like peptide, PHM-27. *Nature* 304: 547–549,

1983.

- [76424] 14754.Linder, S.; Barkhem, T.; Norberg, A.; Persson, H.; Schalling, M.;Hokfelt, T.; Magnusson, G.: Structure and expression of the geneencoding the vasoactive intestinal peptide precursor. *Proc. Nat.Acad. Sci.* 84: 605–609, 1987.
- [76425] 14755.Omary, M. B.; Kagnoff, M. F.: Identification of nuclear receptorsfor VIP on a human colonic adenocarcinoma cell line. *Science* 238:1578–1581, 1987.
- [76426] 14756.Couvineau, A.; Rouyer–Fessard, C.; Darmoul, D.; Maoret, J.–J.;Carrero, I.; Ogier–Denis, E.; Laburthe, M.: Human intestinal VIPreceptor: cloning and functional expression of two cDNA encoding proteinswith different N–terminal domains. *Biochem. Biophys. Res. Commun.* 200:769–776, 1994.
- [76427] 14757.Hashimoto, H.; Nishino, A.; Shintani, N.; Hagihara, N.; Copeland,N. G.; Jenkins, N. A.; Yamamoto, K.; Matsuda, T.; Ishihara, T.; Nagata,S.; Baba, A.: Genomic organization and chromosomal location of themouse vasoactive intestinal polypeptide 1 (VPAC–1) receptor. *Genomics* 58:90–93, 1999.
- [76428] 14758.Sreedharan, S. P.; Huang, J.–X.; Cheung, M.–C.; Goetzl, E. J.:Structure, expression, and chromosomal lo–

- calization of the type I human vasoactive intestinal peptide receptor gene. *Proc. Nat. Acad. Sci.* 92: 2939–2943, 1995.
- [76429] 14759. Sreedharan, S. P.; Patel, D. R.; Huang, J.-X.; Goetzl, E. J.: Cloning and functional expression of a human neuroendocrine vasoactive intestinal peptide receptor. *Biochem. Biophys. Res. Commun.* 193:546–553, 1993.
- [76430] 14760. Sreedharan, S. P.; Robichon, A.; Peterson, K. E.; Goetzl, E. J.: Cloning and expression of the human vasoactive intestinal peptide receptor. *Proc. Nat. Acad. Sci.* 88: 4986–4990, 1991.
- [76431] 14761. Ueno, S.; Maruki, Y.; Nakamura, M.; Tomemori, Y.; Kamae, K.; Tanabe, H.; Yamashita, Y.; Matsuda, S.; Kaneko, S.; Sano, A.: The gene encoding a newly discovered protein, chorein, is mutated in chorea-acanthocytosis. *Nature Genet.* 28: 121–122, 2001.
- [76432] 14762. Hirata, T.; Ushikubi, F.; Kakizuka, A.; Okuma, M.; Narumiya, S.: Two thromboxane A<sub>2</sub> receptor isoforms in human platelets: opposite coupling to adenylyl cyclase with different sensitivity to arg60-to-leu mutation. *J. Clin. Invest.* 97: 949–956, 1996.
- [76433] 14763. Nusing, R. M.; Hirata, M.; Kakizuka, A.; Eki, T.; Ozawa, K.; Narumiya, S.: Characterization and chromosomal mapping of the human thromboxane A<sub>2</sub> receptor

gene. J. Biol. Chem. 268: 25253–25259, 1993.

[76434] 14764.Schwengel, D. A.; Nouri, N.; Meyers, D. A.; Levitt, R. C.: Linkagemapping of the human thromboxane A2 receptor (TBXA2R) to chromosome19p13.3 using transcribed 3–prime untranslated DNA sequence polymorphisms. Genomics 18:212–215, 1993.

[76435] 14765.Thomas, D. W.; Mannon, R. B.; Mannon, P. J.; Latour, A.; Oliver, J. A.; Hoffman, M.; Smithies, O.; Koller, B. H.; Coffman, T. M.:Coagulation defects and altered hemodynamic responses in mice lackingreceptors for thromboxane A(2). J. Clin. Invest. 102: 1994–2001,1998.

[76436] 14766.Unoki, M.; Furuta, S.; Onouchi, Y.; Watanabe, O.; Doi, S.; Fujiwara, H.; Miyatake, A.; Fujita, K.; Tamari, M.; Nakamura, Y.: Associationstudies of 33 single nucleotide polymorphisms (SNPs) in 29 candidategenes for bronchial asthma: positive association a T924C polymorphism in the thromboxane A2 receptor gene. Hum. Genet. 106: 440–446, 2000.

[76437] 14767.Ushikubi, F.; Nakajima, M.; Hirata, M.; Okuma, M.; Fujiwara, M.; Narumiya, S.: Purification of the thromboxane A2/prostaglandin H2receptor from human blood platelets.. J. Biol. Chem. 264: 16496–16501,1989.

[76438] 14768.Ushikubi, F.; Okuma, M.; Kanaji, K.; Sugiyama, T.;

Ogorochi, T.; Narumiya, S.; Uchino, H.: Hemorrhagic thrombocytopathy with platelet thromboxane A<sub>2</sub> abnormality: defective signal transduction with normal binding activity. *Thromb. Haemost.* 57: 158–164, 1987.

- [76439] 14769. Ades, E. W.; Zwerner, R. K.; Acton, R. T.; Balch, C. M.: Isolation and partial characterisation of the human homologue of Thy-1. *J. Exp. Med.* 151: 400–406, 1980.
- [76440] 14770. Bonewald, L.; Ades, E. W.; Tung, E.; Marchalonis, J. J.; Wang, A. C.: Biochemical characterization of human Thy-1. *J. Immunogenet.* 11: 283–296, 1984.
- [76441] 14771. Gatti, R. A.; Lathrop, G. M.; Salser, W.; Silver, J.; Lalouel, J. M.; White, R.: Location of Thy-1 with respect to a primary linkage map of chromosome 11q. (Abstract) *Cytogenet. Cell Genet.* 46: 618 only, 1987.
- [76442] 14772. Gatti, R. A.; Shaked, R.; Wei, S.; Koyama, M.; Salser, W.; Silver, J.: DNA polymorphism in the human Thy-1 gene. *Hum. Immun.* 22: 145–150, 1988.
- [76443] 14773. Greenspan, R. J.; O'Brien, M. C.: Genetic evidence for the role of Thy-1 in neurite outgrowth in the mouse. *J. Neurogenet.* 5: 25–36, 1989.

- [76444] 14774.Grzeschik, K.-H.; Kazazian, H. H.: Report of the committee on the genetic constitution of chromosomes 10, 11, and 12. *Cytogenet.Cell Genet.* 40: 179–205, 1985.
- [76445] 14775.Letarte–Muirhead, M.; Barclay, A. N.; Williams, A. F.: Purification of the Thy–1 molecule, a major cell surface glycoprotein of rat thymocytes. *Biochem. J.* 151: 685–697, 1975.
- [76446] 14776.Mansour, M. H.; Negm, H. I.; Cooper, E. L.: Thy–1 evolution. *Dev.Comp. Immun.* 11: 3–15, 1987.
- [76447] 14777.McKenzie, J. L.; Fabre, J. W.: Human Thy–1: unusual localization and possible functional significance in lymphoid tissues. *J. Immun.* 126:843–850, 1981.
- [76448] 14778.Morris, R.: Thy–1 in developing nervous tissue. *Dev. Neurosci.* 7:133–160, 1985.
- [76449] 14779.Raff, M. C.: Surface antigenic markers for distinguishing T and B lymphocytes in mice. *Transplant. Rev.* 6: 52–80, 1971.
- [76450] 14780.Rettig, W. J.; Dracopoli, N. C.; Chesa, P. G.; Spengler, B. A.; Beresford, H. R.; Davies, P.; Biedler, J. L.; Old, L. J.: Role of human chromosome 11 in determining surface antigenic phenotype of normal and malignant cells: somatic cell genetic analysis of eight antigens, including putative human Thy–1. *J. Exp. Med.* 162: 1603–1619, 1985.

- [76451] 14781.Rettig, W. J.; Dracopoli, N. C.; Silver, J.; Old, L. J.: HumanTHY-1: regional mapping on chromosome 11 and comparison with otherchromosome 11-encoded cell surface glycoproteins. (Abstract) Cytogenet.Cell Genet. 40: 731 only, 1985.
- [76452] 14782.Rettig, W. J.; Dracopoli, N. C.; Spengler, B. A.; Biedler, J.L.; Old, L. J.: Somatic cell genetic analysis of human cell surfaceantigens, including putative human Thy-1: eight distinct antigenicsystems controlled by chromosome 11. (Abstract) Cytogenet. CellGenet. 40: 732 only, 1985.
- [76453] 14783.Seki, T.; Spurr, N.; Obata, F.; Goyert, S.; Goodfellow, P.; Silver,J.: The human Thy-1 gene: structure and chromosomal location. Proc.Nat. Acad. Sci. 82: 6657-6661, 1985.
- [76454] 14784.Tse, A. G. D.; Barclay, A. N.; Watts, A.; Williams, A. F.: Aglycophospholipid tail at the carboxyl terminus of the Thy-1 glycoproteinof neurons and thymocytes. Science 230: 1003-1008, 1985.
- [76455] 14785.Chen, Z.; Friedrich, G. A.; Soriano, P.: Transcriptional enhancerfactor 1 disruption by a retroviral gene trap leads to heart defectsand embryonic lethality in mice. Genes Dev. 8: 2293-2301, 1994.



- [76456] 14786.Aso, T.; Tsai, P.; Kawaguchi, T.; Menninger, J. C.; Kitajima, S.; Yasukochi, Y.; Ward, D. C.; Weissman, S. M.: Assignment of the humanGTF2F1 gene to chromosome 19p13.3. *Genomics* 16: 252–253, 1993.
- [76457] 14787.Aso, T.; Vasavada, H. A.; Kawaguchi, T.; Germino, F. J.; Ganguly, S.; Kitajima, S.; Weissman, S. M.; Yasukochi, Y.: Characterization of cDNA for the large subunit of the transcription initiation factorTFIIF. *Nature* 355: 461–464, 1992.
- [76458] 14788.Finkelstein, A.; Kostrub, C. F.; Li, J.; Chavez, D. P.; Wang, B.Q.; Fang, S. M.; Greenblatt, J.; Burton, Z. F.: A cDNA encoding RAP74, a general initiation factor for transcription by RNA polymerase II. *Nature* 355:464–467, 1992.
- [76459] 14789.Joliot, V.; Demma, M.; Prywes, R.: Interaction with RAP74 subunit of TFIIF is required for transcriptional activation by serum response factor. *Nature* 373: 632–635, 1995.
- [76460] 14790.Sopta, M.; Burton, Z. F.; Greenblatt, J.: Structure and associated DNA–helicase activity of a general transcription initiation factor that binds to RNA polymerase II. *Nature* 341: 410–414, 1989.
- [76461] 14791.Field, S. J.; Tsai, F.–Y.; Kuo, F.; Zubiaga, A. M.;

Kaelin, W.G., Jr.; Livingston, D. M.; Orkin, S. H.; Greenberg, M. E.: E2F-1 functions in mice to promote apoptosis and suppress proliferation. *Cell* 85:549–561, 1996.

[76462] 14792. Helin, K.; Lees, J. A.; Vidal, M.; Dyson, N.; Harlow, E.; Fattaey, A.: A cDNA encoding a pRB-binding protein with properties of the transcription factor E2F. *Cell* 70: 337–350, 1992.

[76463] 14793. Irwin, M.; Marin, M. C.; Phillips, A. C.; Seelan, R. S.; Smith, D. I.; Liu, W.; Flores, E. R.; Tsai, K. Y.; Jacks, T.; Vousden, K.H.; Kaelin, W. G., Jr.: Role for the p53 homologue p73 in E2F-1-induced apoptosis. *Nature* 407: 645–648, 2000.

[76464] 14794. Jacks, T.; Fazeli, A.; Schmitt, E. M.; Bronson, R. T.; Goodell, M. A.; Weinberg, R. A.: Effects of an Rb mutation in the mouse. *Nature* 359:295–300, 1992.

[76465] 14795. Lees, J. A.; Saito, M.; Valentine, M.; Look, T.; Harlow, E.; Dyson, N.; Helin, K.: The retinoblastoma protein binds to a family of E2F transcription factors. *Molec. Cell. Biol.* 13: 7813–7825, 1993.

[76466] 14796. Leone, G.; Sears, R.; Huang, E.; Rempel, R.; Nuckolls, F.; Park, C.-H.; Giangrande, P.; Wu, L.; Saavedra, H. I.; Field, S. J.; Thompson, M. A.; Yang, H.; Fujiwara, Y.; Greenberg, M. E.; Orkin, S.; Smith, C.; Nevins, J. R.: Myc requires

distinct E2F activities to induce S phase and apoptosis.

Molec. Cell 8: 105–113, 2001.

- [76467] 14797. Lissy, N. A.; Davis, P. K.; Irwin, M.; Kaelin, W. G.; Dowdy, S.F.: A common E2F–1 and p73 pathway mediates cell death induced by TCR activation. Nature 407: 642–645, 2000.
- [76468] 14798. Neuman, E.; Sellers, W. R.; McNeil, J. A.; Lawrence, J. B.; Kaelin, W. G., Jr.: Structure and partial genomic sequence of the human E2F1 gene. Gene 173: 163–169, 1996.
- [76469] 14799. Nevins, J. R.: The Rb/E2F pathway and cancer. Hum. Molec. Genet. 10:699–703, 2001.
- [76470] 14800. Nevins, J. R.: E2F: a link between the Rb tumor suppressor protein and viral oncoproteins. Science 258: 424–429, 1992.
- [76471] 14801. Ohtani, K.; DeGregori, J.; Nevins, J. R.: Regulation of the cyclin E gene by transcription factor E2F1. Proc. Nat. Acad. Sci. 92: 12146–12150, 1995.
- [76472] 14802. Arden, K. C.; Boutin, J.–M.; Djiane, J.; Kelly, P. A.; Cavenee, W. K.: The receptors for prolactin and growth hormone are localized in the same region of human chromosome 5. Cytogenet. Cell Genet. 53:161–165, 1990.
- [76473] 14803. Arden, K. C.; Cavenee, W. K.; Boutin, J.–M.; Kelly, P.

A.: The genes encoding the receptors for prolactin and growth hormone map to human chromosome 5. (Abstract) Am. J. Hum. Genet. 45 (suppl.):A129 only, 1989.

[76474] 14804. Boutin, J.-M.; Edery, M.; Shirota, M.; Jolicoeur, C.; Lesueur, L.; Ali, S.; Gould, D.; Djiane, J.; Kelly, P. A.: Identification of a cDNA encoding a long form of prolactin receptor in human hepatoma and breast cancer cells. Molec. Endocr. 3: 1455–1461, 1989.

[76475] 14805. Cunningham, B. C.; Bass, S.; Fuh, G.; Wells, J. A.: Zinc mediation of the binding of human growth hormone to the human prolactin receptor. Science 250:1709–1712, 1990.

[76476] 14806. Glasow, A.; Horn, L.-C.; Taymans, S. E.; Stratakis, C. A.; Kelly, P. A.; Kohler, U.; Gillespie, J.; Vonderhaar, B. K.; Bornstein, S. R.: Mutational analysis of the PRL receptor gene in human breast tumors with differential PRL receptor protein expression. J. Clin. Endocr. Metab. 86: 3826–3832, 2001.

[76477] 14807. Hu, Z.-Z.; Zhuang, L.; Meng, J.; Leondires, M.; Dufau, M. L.: The human prolactin receptor gene structure and alternative promoter utilization: the generic promoter hP<sub>III</sub> and a novel human promoter hP(N). J. Clin. Endocr. Metab. 84: 1153–1156, 1999.

- [76478] 14808.Ormandy, C. J.; Camus, A.; Barra, J.; Damotte, D.; Lucas, B.; Buteau,H.; Edery, M.; Brousse, N.; Babinet, C.; Binart, N.; Kelly, P. A.: Null mutation of the prolactin receptor gene produces multiple reproductivedefects in the mouse. *Genes Dev.* 11: 167–178, 1997.
- [76479] 14809.Perrot–Applanat, M.; Gualillo, O.; Pezet, A.; Vincent, V.; Edery,M.; Kelly, P. A.: Dominant negative and cooperative effects of mutantforms of prolactin receptor. *Molec. Endocr.* 11: 1020–1032, 1997.
- [76480] 14810.Tan, F.; Morris, P. W.; Skidgel, R. A.; Erdos, E. G.: Sequencingand cloning of human prolylcarboxypeptidase (angiotensinase C): similarityto both serine carboxypeptidase and prolylendopeptidase families. *J.Biol. Chem.* 268: 16631–16638, 1993.
- [76481] 14811.Watson, B., Jr.; Nowak, N. J.; Myracle, A. D.; Shows, T. B.; Warnock,D. G.: The human angiotensinase C gene (HUMPCP) maps to 11q14 within700 kb of D11S901: a candidate gene for essential hypertension. *Genomics* 44:365–367, 1997.
- [76482] 14812.Cheng, S. Y.; Gong, Q. H.; Parkinson, C.; Robinson, E. A.; Appella,E.; Merlino, G. T.; Pastan, I.: The nucleotide sequence of a humancellular thyroid hormone–binding protein present in endoplasmic reticulum. *J.Biol. Chem.*

262: 11221–11227, 1987.

- [76483] 14813.Koivu, J.; Myllyla, R.; Halaakoski, T.; Pihlajaniemi, T.; Tasanen,K.; Kivirikko, K. I.: A single polypeptide acts both as the betasubunit of prolyl 4–hydroxylase and as a protein disulfide–isomerase. *J.Biol. Chem.* 262: 6447–6449, 1987.
- [76484] 14814.Morris, J. I.; Varandani, P. T.: Characterization of a cDNA forhuman glutathione–insulin transhydrogenase (protein–disulfide isomerase/oxidoreductase). *Biochim.Biophys. Acta* 949: 169–180, 1988.
- [76485] 14815.Noiva, R.; Lennarz, W. J.: Protein disulfide isomerase: a multifunctionalprotein resident in the lumen of the endoplasmic reticulum. *J. Biol.Chem.* 267: 3553–3556, 1992.
- [76486] 14816.Pajunen, L.; Hoyhtya, M.; Tryggvason, K.; Kivirikko, K. I.; Myllyla,R.: Species–specific antibodies in the assignment of the gene forthe beta–subunit of human prolyl 4–hydroxylase. (Abstract) *Cytogenet.Cell Genet.* 40: 719 only, 1985.
- [76487] 14817.Pajunen, L.; Jones, T. A.; Goddard, A.; Sheer, D.; Solomon, E.;Pihlajaniemi, T.; Kivirikko, K. I.: Regional assignment of the humangene coding for a multifunctional polypeptide (P4HB) acting as thebeta–subunit of prolyl

4-hydroxylase and the enzyme protein disulfide isomerase to 17q25. *Cytogenet. Cell Genet.* 56: 165–168, 1991.

[76488] 14818. Pajunen, L.; Myllylä, R.; Helaakoski, T.; Pihlajaniemi, T.; Tasanen, K.; Hoyhtya, M.; Tryggvason, K.; Solomon, E.; Kivirikko, K. I.: Assignment of the gene coding for both the beta-subunit of prolyl 4-hydroxylase and protein disulphide isomerase to human chromosome region 17q23–25. (Abstract) *Cytogenet. Cell Genet.* 46: 674 only, 1987.

[76489] 14819. Pajunen, L.; Myllylä, R.; Helaakoski, T.; Pihlajaniemi, T.; Tasanen, K.; Hoyhtya, M.; Tryggvason, K.; Solomon, E.; Kivirikko, K. I.: Assignment of the gene coding for both the beta-subunit of prolyl 4-hydroxylase and the enzyme disulfide isomerase to human chromosome region 17p11–qter. *Cytogenet. Cell Genet.* 47: 37–41, 1988.

[76490] 14820. Pihlajaniemi, T.; Helaakoski, T.; Tasanen, K.; Myllylä, R.; Huhtala, M.-L.; Koivu, J.; Kivirikko, K. I.: Molecular cloning of the beta-subunit of human prolyl 4-hydroxylase: this subunit and protein disulphide isomerase are products of the same gene. *EMBO J.* 6: 643–649, 1987.

[76491] 14821. Popescu, N. C.; Cheng, S.; Pastan, I.: Chromosomal

localization of the gene for a human thyroid hormone-binding protein. *Am. J. Hum. Genet.* 42: 560–564, 1988.

- [76492] 14822. Tasanen, K.; Parkkonen, T.; Chow, L. T.; Kivirikko, K. I.; Pihlajaniemi, T.: Characterization of the human gene for a polypeptide that acts both as the beta-subunit of prolyl 4-hydroxylase and as protein disulfide isomerase. *J. Biol. Chem.* 263: 16218–16224, 1988.
- [76493] 14823. Kelly, A.; Powis, S. H.; Glynn, R.; Radley, E.; Beck, S.; Trowsdale, J.: Second proteasome-related gene in the human MHC class II region. *Nature* 353: 667–668, 1991.
- [76494] 14824. Martinez, C. K.; Monaco, J. J.: Homology of proteasome subunits to a major histocompatibility complex-linked LMP gene. *Nature* 353: 664–667, 1991.
- [76495] 14825. Petes, T. D.: Meiotic recombination hot spots and cold spots. *Nature Rev. Genet.* 2: 360–369, 2001.
- [76496] 14826. Van Kaer, L.; Ashton-Rickardt, P. G.; Eichelberger, M.; Gaczynska, M.; Nagashima, K.; Rock, K. L.; Goldberg, A. L.; Doherty, P. C.; Tonegawa, S.: Altered peptidase and viral-specific T cell response in LMP2 mutant mice. *Immunity* 1: 533–541, 1994.
- [76497] 14827. Zhou, P.; Zanelli, E.; Smart, M.; David, C.: Genomic organization and tissue expression of mouse proteasome gene *Lmp-2*. *Genomics* 16: 664–668, 1993.



- [76498] 14828.Bodmer, J. G.; Marsh, S. G. E.; Albert, E. D.; Bodmer, W. F.; Dupont, B.; Erlich, H. A.; Mach, B.; Mayr, W. R.; Parham, P.; Sasazuki, T.; Schreuder, G. M. T.; Strominger, J. L.; Svejgaard, A.; Terasaki, P. I.: Nomenclature for factors of the HLA system, 1991. *Tissue Antigens* 39:161–173, 1992.
- [76499] 14829.Deng, G. Y.; Muir, A.; Maclaren, N. K.; She, J.-X.: Association of LMP2 and LMP7 genes within the major histocompatibility complex with insulin-dependent diabetes mellitus: population and family studies. *Am. J. Hum. Genet.* 56: 528–534, 1995.
- [76500] 14830.Fehling, H. J.; Swat, W.; Laplace, C.; Kuhn, R.; Rajewsky, K.; Muller, U.; von Boehmer, H.: MHC class I expression in mice lacking the proteasome subunit LMP-7. *Science* 265: 1234–1237, 1994.
- [76501] 14831.Doolittle, R. F.; Hunkapiller, M. W.; Hood, L. E.; Devare, S. G.; Robbins, K. C.; Aaronson, S. A.; Antoniades, H. N.: Simian sarcomavirus onc gene, v-sis, is derived from the gene (or genes) encoding a platelet-derived growth factor. *Science* 221: 275–277, 1983.
- [76502] 14832.Frolova, L. Y.; Sudomoina, M. A.; Grigorieva, A. Y.; Zinovieva, O. L.; Kisselev, L. L.: Cloning and nucleotide sequence of the structural gene encoding for human trypto-

phanyl-tRNA synthetase. Gene 109:291-296, 1991.

- [76503] 14833.Vassart, G.: Personal Communication. Brussels, Belgium 1/15/1992.
- [76504] 14834.Wenger, G. D.: Personal Communication. Columbus, Ohio 8/3/1993.
- [76505] 14835.Gardner, T. L.; Elston, D. M.; Wotowic, P. J.: A familial dermatofibrosarcomaprotuberans. J. Am. Acad. Derm. 39: 504-505, 1998.
- [76506] 14836.Groffen, J.; Heisterkamp, N.; Stephenson, J. R.; Geurts van Kessel,A.; de Klein, A.; Grosveld, G.; Bootsma, D.: c-sis is translocatedfrom chromosome 22 to chromosome 9 in chronic myelocytic leukemia. J.Exp. Med. 158: 9-15, 1983.
- [76507] 14837.Hermansson, M.; Nister, M.; Betsholtz, C.; Heldin, C.-H.; Westermarck,B.; Funa, K.: Endothelial cell hyperplasia in human glioblastoma:coexpression of mRNA for platelet-derived growth factor (PDGF) B chainand PDGF receptor suggests autocrine growth stimulation. Proc. Nat.Acad. Sci. 85: 7748-7752, 1988.
- [76508] 14838.Josephs, S. F.; Dalla-Favera, R.; Gelmann, E. P.; Gallo, R. C.;Wong-Staal, F.: 5-prime viral and human cellular sequences correspondingto the transforming gene of simian sarcoma virus. Science 219: 503-505,1983.

- [76509] 14839. Josephs, S. F.; Guo, C.; Ratner, L.; Wong-Staal, F.: Human proto-oncogene nucleotide sequences corresponding to the transforming region of simian sarcoma virus. *Science* 223: 487–491, 1984.
- [76510] 14840. Josephs, S. F.; Ratner, L.; Clarke, M. F.; Westin, E. H.; Reitz, M. S.; Wong-Staal, F.: Transforming potential of human c-sis nucleotide sequences encoding platelet-derived growth factor. *Science* 225: 636–639, 1984.
- [76511] 14841. Kelly, J. D.; Raines, E. W.; Ross, R.; Murray, M. J.: The B chain of PDGF alone is sufficient for mitogenesis. *EMBO J.* 4: 3399–3405, 1985.
- [76512] 14842. Kiuru-Kuhlefelt, S.; El-Rifai, W.; Fanburg-Smith, J.; Kere, J.; Miettinen, M.; Knuutila, S.: Concomitant DNA copy number amplification at 17q and 22q in dermatofibrosarcoma protuberans. *Cytogenet. Cell Genet.* 92: 192–195, 2001.
- [76513] 14843. Kozak, C. A.; Sears, J. F.; Hoggan, M. D.: Genetic mapping of the mouse proto-oncogene c-sis to chromosome 15. *Science* 221: 867–869, 1983.
- [76514] 14844. Lindahl, P.; Johansson, B. R.; Leveen, P.; Betsholtz, C.: Pericyte loss and microaneurysm formation in PDGF-B-deficient mice. *Science* 277: 242–245, 1997.
- [76515] 14845. Owen, A. J.; Pantazis, P.; Antoniadou, H. N.: Simian

sarcomavirus-transformed cells secrete a mitogen identical to platelet-derived growth factor. *Science* 225: 54–56, 1984.

- [76516] 14846. Rao, C. D.; Igarashi, H.; Chiu, I.-M.; Robbins, K. C.; Aaronson, S. A.: Structure and sequence of the human c-sis/platelet-derived growth factor 2 (SIS/PDGF2) transcriptional unit. *Proc. Nat. Acad. Sci.* 83: 2392–2396, 1986.
- [76517] 14847. Robbins, K. C.; Antoniades, H. N.; Devare, S. G.; Hunkapiller, M. W.; Aaronson, S. A.: Structural and immunological similarities between simian sarcoma virus gene product(s) and human platelet-derived growth factor. *Nature* 305: 605–608, 1983.
- [76518] 14848. Robbins, K. C.; Devare, S. G.; Reddy, E. P.; Aaronson, S. A.: In vivo identification of the transforming gene product of simian sarcoma virus. *Science* 218: 1131–1133, 1982.
- [76519] 14849. Simon, M.-P.; Navarro, M.; Roux, D.; Pouyssegur, J.: Structural and functional analysis of a chimeric protein COL1A1-PDGFB generated by the translocation t(17;22)(q22;q13.1) in dermatofibrosarcoma protuberans (DP). *Oncogene* 20: 2965–2975, 2001.
- [76520] 14850. Smidt, M.; Kirsch, I.; Ratner, L.: Deletion of Alu sequences in the fifth c-sis intron in individuals with menin-

giomas. J. Clin. Invest. 86: 1151–1157, 1990.

- [76521] 14851. Turc-Carel, C.; Philip, I.; Berger, M. P.; Philip, T.; Lenoir, G. M.: Chromosomal translocations in Ewing's sarcoma. (Letter) New Eng. J. Med. 309: 497–498, 1983.
- [76522] 14852. Shimizu, A.; O'Brien, K. P.; Sjoblom, T.; Pietras, K.; Buchdunger, E.; Collins, V. P.; Heldin, C.-H.; Dumanski, J. P.; Ostman, A.: The dermatofibrosarcoma protuberans-associated collagen type I-alpha-1/platelet-derived growth factor (PDGF) beta-chain fusion gene generates a transforming protein that is processed to functional PDGF-BB. Cancer Res. 59:3719–3723, 1999.
- [76523] 14853. Simon, M.-P.; Pedeutour, F.; Sirvent, N.; Grosgeorge, J.; Minoletti, F.; Coindre, J.-M.; Terrier-Lacombe, M.-J.; Mandahl, N.; Craver, R.D.; Blin, N.; Sozzi, G.; Turc-Carel, C.; O'Brien, K. P.; Kedra, D.; Fransson, I.; Guilbaud, C.; Dumanski, J. P.: Deregulation of the platelet-derived growth factor B-chain gene via fusion with collagen gene COL1A1 in dermatofibrosarcoma protuberans and giant-cell fibroblastoma. Nature Genet. 15: 95–98, 1997.
- [76524] 14854. Waterfield, M. D.; Scrace, G. T.; Whittle, N.; Stroobant, P.; Johnsson, A.; Wasteson, A.; Westermark, B.; Heldin, C.-H.; Huang, J. S.; Deuel, T. F.: Platelet-derived growth factor is structurally related to the putative trans-

forming protein p28(sis) of simian sarcomavirus. Nature 304: 35–39, 1983.

[76525] 14855. Graphodatsky, A.; Frolova, L.; Biltueva, L.; Eremina, V.; Lushnikova, T.; Sudomoina, M.; Zinovieva, O.; Kisselev, L.: Localization of the tryptophanyl tRNA synthetase gene (WARS) on human and bovine chromosomes by in situ hybridization. Mammalian Genome 4: 183–184, 1993.

[76526] 14856. Jensen, L. L.; Nielsen, M. M.; Justesen, J.; Hansen, L. L.: Assignment of human NADH dehydrogenase (ubiquinone) 1 beta subcomplex 3 (NDUFB3) and of its four pseudogenes to human chromosomes 2q31.3, 1p13.3–p13.1, 9q32–q34.1, 14q22.3–q23.1 and 14q32.2 by radiation hybrid mapping. Cytogenet. Cell Genet. 93: 147–150, 2001.

[76527] 14857. Otani, A.; Slike, B. M.; Dorrell, M. I.; Hood, J.; Kinder, K.; Ewalt, K. L.; Cheresh, D.; Schimmel, P.; Friedlander, M.: A fragment of human TrpRS as a potent antagonist of ocular angiogenesis. Proc. Nat. Acad. Sci. 99: 178–183, 2002.

[76528] 14858. Shimizu, N.; Kucherlapati, R. S.; Ruddle, F. H.: Assignment of a human gene for tryptophanyl-tRNA synthetase to chromosome 14 using human-mouse somatic cell hybrids. Somat. Cell Genet. 2: 345–357, 1976.

- [76529] 14859.Tolstrup, A. B.; Bejder, A.; Fleckner, J.; Justesen, J.: Transcriptional regulation of the interferon-  
gamma-inducible tryptophanyl-tRNA synthetase includes  
alternative splicing. *J. Biol. Chem.* 270: 397-403, 1995.
- [76530] 14860.Turpaev, K. T.; Zakhariev, V. M.; Sokolova, I. V.;  
Narovlyansky, A. N.; Amchenkova, A. M.; Justesen, J.; Frol-  
ova, L. Y.: Alternative processing of the tryptophanyl-tRNA  
synthetase mRNA from interferon-treated human cells. *Eu-  
rop. J. Biochem.* 240: 732-737, 1996.
- [76531] 14861.Wakasugi, K.; Slike, B. M.; Hood, J.; Otani, A.; Ewalt,  
K. L.; Friedlander, M.; Cheresh, D. A.; Schimmel, P.: A hu-  
man aminoacyl-tRNA synthetase as a regulator of angio-  
genesis. *Proc. Nat. Acad. Sci.* 99:173-177, 2002.
- [76532] 14862.Miller, J. S.; Moxley, G.; Schwartz, L. B.: Cloning and  
characterization of a second complementary DNA for hu-  
man tryptase. *J. Clin. Invest.* 86:864-870, 1990.
- [76533] 14863.Miller, J. S.; Westin, E. H.; Schwartz, L. B.: Cloning  
and characterization of complementary DNA for human  
tryptase. *J. Clin. Invest.* 84: 1188-1195, 1989.
- [76534] 14864.Vanderslice, P.; Ballinger, S. M.; Tam, E. K.; Gold-  
stein, S. M.; Craik, C. S.; Caughey, G. H.: Human mast cell  
tryptase: multiple cDNAs and genes reveal a multigene  
serine protease family. *Proc. Nat. Acad. Sci.* 87:

3811–3815, 1990.

- [76535] 14865.Mach, B.; Steimle, V.; Reith, W.: MHC class II–deficient combinedimmunodeficiency: a disease of gene regulation. *Immun. Rev.* 138:207–221, 1994.
- [76536] 14866.Babcock, M.; de Silva, D.; Oaks, R.; Davis–Kaplan, S.; Jiralerspong,S.; Montermini, L.; Pandolfo, M.; Kaplan, J.: Regulation of mitochondrialiron accumulation by Yfh1p, a putative homolog of frataxin. *Science* 276:1709–1712, 1997.
- [76537] 14867.Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: The GAA triplet–repeatexpansion in Friedreich ataxia inter–feres with transcription and maybe associated with an unusual DNA structure. *Am. J. Hum. Genet.* 62:111–121, 1998.
- [76538] 14868.Bidichandani, S. I.; Ashizawa, T.; Patel, P. I.: Atypical Friedreichataxia caused by compound heterozygosity for a novel missense mutationand the GAA triplet–repeat expansion. (Letter) *Am. J. Hum. Genet.* 60:1251–1256, 1997.
- [76539] 14869.Cavadini, P.; Gellera, C.; Patel, P. I.; Isaya, G.: Human frataxinmaintains mitochondrial iron homeostasis in *Saccharomyces cerevisiae*. *Hum.Molec. Genet.* 9: 2523–2530, 2000.



- [76540] 14870.Sukhatme, V. P.; Vollmer, A. C.; Erikson, J.; Isobe, M.; Croce,C.; Parnes, J. R.: Gene for the human T cell differentiation antigenLeu-2/T8 is closely linked to the kappa light chain locus on chromosome2. J. Exp. Med. 161: 429-434, 1985.
- [76541] 14871.Traver, D.; Akashi, K.; Manz, M.; Merad, M.; Miyamoto, T.; Engleman,E. G.; Weissman, I. L.: Development of CD8-alpha-positive dendriticcells from a common myeloid progenitor. Science 290: 2152-2154,2000.
- [76542] 14872.Weichhold, G. M.; Huber, C.; Parnes, J. R.; Zachau, H. G.: TheCD8-alpha locus is located on the telomere side of the immunoglobulin-kappalocus at a distance of 2 Mb. Genomics 16: 512-514, 1993.
- [76543] 14873.Balk, S. P.; Burke, S.; Polischuk, J. E.; Frantz, M. E.; Yang,L.; Porcelli, S.; Colgan, S. P.; Blumberg, R. S.: Beta-2-microglobulin-independentMHC class Ib molecule expressed by human intestinal epithelium. Science 265:259-262, 1994.
- [76544] 14874.Benlagha, K.; Kyin, T.; Beavis, A.; Teyton, L.; Bendelac, A.:A thymic precursor to NK T cell lineage. Science 296: 553-555, 2002.
- [76545] 14875.Bradbury, A.; Milstein, C.; Kozak, C. A.: Chromosomal localizationof Cd1d genes in the mouse. Somat. Cell

Molec. Genet. 17: 93–96,1991.

- [76546] 14876.Honey, K.; Benlagha, K.; Beers, C.; Forbush, K.; Teyton, L.; Kleijmeer, M. J.; Rudensky, A. Y.; Bendelac, A.: Thymocyte expression of cathepsinL is essential for NKT cell development. Nature Immun. 7Oct, 2002.Note: Advance Electronic Publication.
- [76547] 14877.MacDonald, H. R.: T before NK. Science 296: 481–482, 2002.
- [76548] 14878.Pellicci, D. G.; Hammond, K. J. L.; Uldrich, A. P.; Baxter, A.G.; Smyth, M. J.; Godfrey, D. I.: A natural killer T (NKT) cell developmental pathway involving a thymus-dependent NK1.1(–)CD4(+) CD1d-dependent precursor stage. J. Exp. Med. 195: 835–844, 2002.
- [76549] 14879.Prigozy, T. I.; Naidenko, O.; Qasba, P.; Elewaut, D.; Brossay, L.; Khurana, A.; Natori, T.; Koezuka, Y.; Kulkarni, A.; Kronenberg, M.: Glycolipid antigen processing for presentation by CD1d molecules. Science 291:664–667, 2001.
- [76550] 14880.Shi, F.–D.; Flodstrom, M.; Balasa, B.; Kim, S. H.; Van Gunst, K.; Strominger, J. L.; Wilson, S. B.; Sarvetnick, N.: Germ line deletion of the CD1 locus exacerbates diabetes in the NOD mouse. Proc. Nat.Acad. Sci. 98: 6777–6782, 2001.

- [76551] 14881.Hashimoto, K.; Hirai, M.; Kurosawa, Y.: A gene outside the humanMHC related to classical HLA class I genes. Science 269: 693–695,1995.
- [76552] 14882.Shilo, B. Z.; Weinberg, R. A.: DNA sequences homologous to vertebrateoncogenes are conserved in *Drosophila melanogaster*. Proc. Nat. Acad.Sci. 78: 6789–6792, 1981.
- [76553] 14883.Takahashi, E.; Hori, T.; O'Connell, P.; Leppert, M.; White, R.: Mapping of the MYC gene to band 8q24.12–q24.13 by R–banding anddistal to fra(8)(q24.11), FRA8E, by fluorescence in situ hybridization. Cytogenet.Cell Genet. 57: 109–111, 1991.
- [76554] 14884.Taub, R.; Kirsch, I.; Morton, C.; Lenoir, G.; Swan, D.; Tronick,S.; Aaronson, S.; Leder, P.: Translocation of the c–myc gene intothe immunoglobulin heavy chain locus in human Burkitt lymphoma andmurine plasmacytoma cells. Proc. Nat. Acad. Sci. 79: 7837–7841,1982.
- [76555] 14885.Trumpp, A.; Refaeli, Y.; Oskarsson, T.; Gasser, S.; Murphy, M.;Martin, G. R.; Bishop, J. M.: c–Myc regulates mammalian body sizeby controlling cell number but not cell size. Nature 414: 768–773,2001.
- [76556] 14886.Vafa, O.; Wade, M.; Kern, S.; Beeche, M.; Pandita, T. K.; Hampton,G. M.; Wahl, G. M.: c–Myc can induce DNA

damage, increase reactiveoxygen species, and mitigate p53 function: a mechanism for oncogene–inducedgenetic instability. *Molec. Cell* 9: 1031–1044, 2002.

[76557] 14887.Watt, R.; Nishikura, K.; Sorrentino, J.; ar–Rushdi, A.; Croce,C. M.; Rovera, G.: The structure and nucleotide sequence of the 5–primeend of the human c–myc oncogene. *Proc. Nat. Acad. Sci.* 80: 6307–6311,1983.

[76558] 14888.Watt, R.; Stanton, L. W.; Marcu, K. B.; Gallo, R. C.; Croce, C.M.; Rovera, G.: Nucleotide sequence of cloned cDNA of human c–myconcogene. *Nature* 303: 725–728, 1983.

[76559] 14889.Azarnia, R.; Reddy, S.; Kmiecik, T. E.; Shalloway, D.; Loewenstein,W. R.: The cellular src gene product regulates junctional cell–to–cellcommunication. *Science* 239: 398–401, 1988.

[76560] 14890.Czernilofsky, A. P.; Levinson, A. D.; Varmus, H. E.; Bishop, J.M.; Tischer, E.; Goodman, H.: Correction to the nucleotide sequenceof the src gene of Rous sarcoma virus. *Nature* 301: 736–738, 1983.

[76561] 14891.Gibbs, C. P.; Tanaka, A.; Anderson, S. K.; Radul, J.; Baar, J.;Ridgway, A.; Kung, H.–J.; Fujita, D. J.: Isolation and structuralmapping of a human c–src gene homologous to the transforming gene(v–src) of Rous sarcoma virus. *J. Vi-*

rol. 53: 19–24, 1985.

[76562] 14892.Irby, R. B.; Mao, W.; Coppola, D.; Kang, J.; Loubeau, J. M.; Trudeau,W.; Karl, R.; Fujita, D. J.; Jove, R.; Yeatman, T. J.: ActivatingSRC mutation in a subset of advanced human colon cancers. *NatureGenet.* 21: 187–190, 1999.

[76563] 14893.Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J. D.:c-src is consistently conserved in the chromosomal deletion (20q)observed in myeloid disorders. *Proc. Nat. Acad. Sci.* 82: 6692–6696,1985.

[76564] 14894.Lowe, C.; Yoneda, T.; Boyce, B. F.; Chen, H.; Mundy, G. R.; Soriano,P.: Osteopetrosis in Src-deficient mice is due to an autonomous defectof osteoclasts. *Proc. Nat. Acad. Sci.* 90: 4485–4489, 1993.

[76565] 14895.Morris, C. M.; Honeybone, L. M.; Hollings, P. E.; Fitzgerald,P. H.: Localization of the SRC oncogene to chromosome band 20q11.2and loss of this gene with deletion (20q) in two leukemic patients. *Blood* 74:1768–1773, 1989.

[76566] 14896.Sakaguchi, A. Y.; Mohandas, T.; Naylor, S. L.: A human c-srcgene resides on the proximal long arm of chromosome 20 (cen–q13.1). *CancerGenet. Cytogenet.* 18: 123–129, 1985.

[76567] 14897.Sakaguchi, A. Y.; Naylor, S. L.; Weinberg, R. A.;

Shows, T. B.: Organization of human proto-oncogenes.  
(Abstract) Am. J. Hum. Genet. 34:175A, 1982.

[76568] 14898.Sakaguchi, A. Y.; Zabel, B. U.; Grzeschik, K. H.;  
Law, M. L.;Naylor, S. L.: Human proto-oncogene assign-  
ments. (Abstract) Cytogenet.Cell Genet. 37: 572–573,  
1984.

[76569] 14899.Wong, B. R.; Besser, D.; Kim, N.; Arron, J. R.; Volo-  
godskaia,M.; Hanafusa, H.; Choi, Y.: TRANCE, a TNF family  
member, activatesAkt/PKB through a signaling complex  
involving TRAF6 and c-Src. Molec.Cell 4: 1041–1049,  
1999.

[76570] 14900.Mangelsdorf, D. J.; Borgmeyer, U.; Heyman, R. A.;  
Zhou, J. Y.;Ong, E. S.; Oro, A. E.; Kakizuka, A.; Evans, R.  
M.: Characterizationof three RXR genes that mediate the  
action of 9-cis retinoic acid. GenesDev. 6: 329–344,  
1992.

[76571] 14901.Attard–Montalto, S.; Evans, N.; Sherwood, R. A.:  
Carotenaemiawith low vitamin A levels and low retinol-  
binding protein. J. Inherit.Metab. Dis. 15: 929–930, 1992.

[76572] 14902.Chainani, M.; Sampsell, B.; Elliott, R. W.: Localiza-  
tion of thegene for plasma retinol binding protein to the  
distal half of mousechromosome 19. Genomics 9:  
376–379, 1991.

- [76573] 14903.Matsuo, T.: Familial retinol-binding-protein deficiency? (Letter) *Lancet* II:910 only, 1987.
- [76574] 14904.Matsuo, T.; Matsuo, N.; Shiraga, F.; Koide, N.: Familial retinol-binding-protein deficiency. (Letter) *Lancet* II: 402-403, 1987.
- [76575] 14905.Morgan, F. F.; Canfield, R. E.; Goodman, D. S.: The partial structure of human plasma prealbumin and retinol-binding protein. *Biochim.Biophys. Acta* 236: 798-801, 1971.
- [76576] 14906.Rask, L.; Anundi, H.; Fohlman, J.; Peterson, P. A.: The complete amino acid sequence of human serum retinol-binding protein. *Upsala J. Med. Sci.* 92: 115-146, 1987.
- [76577] 14907.Rask, L.; Vahlquist, A.; Peterson, P. A.: Studies on two physiological forms of the human retinol-binding protein differing in vitamin A and arginine content. *J. Biol. Chem.* 246: 6638-6646, 1971.
- [76578] 14908.Rocchi, M.; Covone, A.; Romeo, G.; Faraonio, R.; Colantuoni, V.: Regional mapping of RBP4 to 10q23-q24 and RBP1 to 3q21-q22 in man. *Somat.Cell Molec. Genet.* 15: 185-190, 1989.
- [76579] 14909.Seeliger, M. W.; Biesalski, H. K.; Wissinger, B.; Gollnick, H.; Gielen, S.; Frank, J.; Beck, S.; Zrenner, E.: Pheno-

type in retinoldeficiency due to a hereditary defect in retinol binding protein synthesis. Invest.Ophthal. Vis. Sci. 40: 3–11, 1999.

[76580] 14910.Thuluvath, P. J.: Familial retinol–binding–protein deficiency?(Letter) Lancet II: 910 only, 1987.

[76581] 14911.Colantuoni, V.; Cortese, R.; Nilsson, M.; Lundvall, J.; Bavik,C. O.; Eriksson, U.; Peterson, P. A.; Sundelin, J.: Cloning and sequencingof a full length cDNA correspond–ing to human cellular retinol–bindingprotein. Biochem. Biophys. Res. Commun. 130: 431–439, 1985.

[76582] 14912.Colantuoni, V.; Rocchi, M.; Roncuzzi, L.; Romeo, G.: Mapping ofhuman cellular retinol–binding protein to chromosome 3. Cytogenet.Cell Genet. 43: 221–222, 1986.

[76583] 14913.De Baere, E.; Speleman, F.; Van Roy, N.; De Paepe, A.; Messiaen,L.: Assignment of the cellular retinol–binding protein 1 gene (RBP1)and of the coatomer beta–prime subunit gene (COPB2) to human chromosomeband 3q23 by in situ hybridization. Cytogenet. Cell Genet. 82: 226–227,1998.

[76584] 14914.Nilsson, M. H. L.; Spurr, N. K.; Lundvall, J.; Rask, L.; Peterson,P. A.: Human cellular retinol–binding protein gene organization andchromosomal location. Europ. J. Biochem. 173: 35–44, 1988.



- [76585] 14915. Carson, N. L.; Simpson, N. E.: A physical map of 13 markers on chromosome 10 from dosage studies on abnormal cell lines. (Abstract) *Cytogenet. Cell Genet.* 51: 974–975, 1989.
- [76586] 14916. Chin, K. S.; Mathew, C. G. P.; Fong, S. L.; Bridges, C. D. B.; Ponder, B. A. J.: Styl RFLP recognised by a human IRBP cDNA localised to chromosome 10. *Nucleic Acids Res.* 16: 1645 only, 1988.
- [76587] 14917. Danciger, M.; Kozak, C. A.; Nickerson, J.; Redmond, T. M.; Farber, D. B.: Localization of the gene for interphotoreceptor retinoid-binding protein to mouse chromosome 14 near Np-1. *Genomics* 8: 727–731, 1990.
- [76588] 14918. Farrer, L. A.; Castiglione, C. M.; Kidd, J. R.; Myers, S.; Carson, N.; Simpson, N. E.; Kidd, K. K.: A linkage group of five DNA markers on human chromosome 10. *Genomics* 3: 72–77, 1988.
- [76589] 14919. Fong, S.-L.; Liou, G. I.; Landers, R. A.; Alvarez, R. A.; Gonzalez-Fernandez, F.; Glazebrook, P. A.; Lam, D. M. K.; Bridges, C. D. B.: Characterization, localization, and biosynthesis of an interstitial retinol-binding glycoprotein in the human eye. *J. Neurochem.* 42: 1667–1676, 1984.
- [76590] 14920. Liou, G. I.; Fong, S.-L.; Beattie, W. G.; Cook, R. G.; Leone, J.; Landers, R. A.; Alvarez, R. A.; Wang, C.; Li, Y.;

Bridges, C. D. B.: Bovine interstitial retinol-binding protein (IRBP)--isolation and sequence analysis of cDNA clones, characterization and in vitro translation of mRNA. *Vision Res.* 26: 1645-1653, 1986.

[76591] 14921. Liou, G. I.; Fong, S.-L.; Gosden, J.; van Tuinen, P.; Ledbetter, D. H.; Christie, S.; Rout, D.; Bhattacharya, S.; Cook, R. G.; Li, Y.; Wang, C.; Bridges, C. D. B.: Human interstitial retinol-binding protein (IRBP): cloning, partial sequence, and chromosomal localization. *Somat. Cell Molec. Genet.* 13: 315-323, 1987.

[76592] 14922. Liou, G. I.; Li, Y.; Wang, C.; Fong, S.-L.; Bhattacharya, F. S.; Bridges, C. D. B.: Bgl II RFLP recognized by a human IRBP cDNA localized to chromosome 10. *Nucleic Acids Res.* 15: 3196 only, 1987.

[76593] 14923. McGuire, W.; Hill, A. V. S.; Allsopp, C. E. M.; Greenwood, B. M.; Kwiatkowski, D.: Variation in the TNF- $\alpha$  promoter region associated with susceptibility to cerebral malaria. *Nature* 371: 508-511, 1994.

[76594] 14924. Benbrook, D.; Pfahl, M.: A novel thyroid hormone receptor encoded by a cDNA clone from a human testis library. *Science* 238: 788-791, 1987.

[76595] 14925. Bernal, J.; Refetoff, S.; DeGroot, L. J.: Abnormalities of triiodothyronine binding to lymphocyte and fibroblast

nuclei from a patient with peripheral tissue resistance to thyroid hormone action. *J. Clin. Endocr. Metab.* 47:1266–1272, 1978.

- [76596] 14926. Dayton, A. I.; Selden, J. R.; Laws, G.; Dorney, D. J.; Finan, J.; Tripputi, P.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.; Croce, C.M.: A human c-erbA oncogene homologue is closely proximal to the chromosome 17 breakpoint in acute promyelocytic leukemia. *Proc. Nat. Acad. Sci.* 81: 4495–4499, 1984.
- [76597] 14927. Debuire, B.; Henry, C.; Benaissa, M.; Biserte, G.; Claverie, J.M.; Saule, S.; Martin, P.; Stehelin, D.: Sequencing the erbA gene of avian erythroblastosis virus reveals a new type of oncogene. *Science* 224:1456–1459, 1984.
- [76598] 14928. Ferro, M. T.; San Roman, C.: Constitutional t(15;17). *Cancer Genet. Cytogenet.* 4: 89–91, 1981.
- [76599] 14929. Gullberg, H.; Rudling, M.; Forrest, D.; Angelin, B.; Vennstrom, B.: Thyroid hormone receptor beta-deficient mice show complete loss of the normal cholesterol 7- $\alpha$ -hydroxylase (CYP7A) response to thyroid hormone but display enhanced resistance to dietary cholesterol. *Molec. Endocr.* 14: 1739–1749, 2000.
- [76600] 14930. Ichikawa, K.; Hughes, I. A.; Horwitz, A. L.; DeGroot, L. J.: Characterization of nuclear thyroid hormone recep-

tors of cultured skin fibroblasts from patients with resistance to thyroid hormone. *Metabolism* 36:392–399, 1987.

- [76601] 14931. Iskaros, J.; Pickard, M.; Evans, I.; Sinha, A.; Hardiman, P.; Ekins, R.: Thyroid hormone receptor gene expression in first trimester human fetal brain. *J. Clin. Endocr. Metab.* 85: 2620–2623, 2000.
- [76602] 14932. Jansson, M.; Philipson, L.; Vennstrom, B.: Isolation and characterization of multiple human genes homologous to the oncogenes of avian erythroblastosis virus. *EMBO J.* 2: 561–565, 1983.
- [76603] 14933. Jhanwar, S. C.; Chaganti, R. S. K.; Croce, C. M.: Germ-line chromosomal localization of human c-erb-A oncogene. *Somat. Cell Molec. Genet.* 11:99–102, 1985.
- [76604] 14934. Kaneshige, M.; Suzuki, H.; Kaneshige, K.; Cheng, J.; Wimbrow, H.; Barlow, C.; Willingham, M. C.; Cheng, S.: A targeted dominant negative mutation of the thyroid hormone alpha-1 receptor causes increased mortality, infertility, and dwarfism in mice. *Proc. Nat. Acad. Sci.* 98:15095–15100, 2001.
- [76605] 14935. Laudet, V.; Begue, A.; Henry-Duthoit, C.; Joubel, A.; Martin, P.; Stehelin, D.; Saule, S.: Genomic organization of the human thyroid hormone receptor alpha (c-erbA-1) gene. *Nucleic Acids Res.* 19: 1105–1112, 1991.

- [76606] 14936. Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Rowley, J. D.; Oren, M.: Translocation of the p53 gene in t(15;17) in acute promyelocytic leukaemia. *Nature* 316: 826–828, 1985.
- [76607] 14937. Mathieu–Mahul, D.; Xu, D. Q.; Saule, S.; Lidereau, R.; Galibert, F.; Berger, R.; Mauchauffe, M.; Larsen, C. J.: An EcoRI restriction fragment length polymorphism (RFLP) in the human c-erb A locus. *Hum. Genet.* 71: 41–44, 1985.
- [76608] 14938. McCabe, C. J.; Gittoes, N. J.; Sheppard, M. C.; Franklyn, J. A.: Thyroid receptor alpha–1 and alpha–2 mutations in nonfunctioning pituitary tumors. *J. Clin. Endocr. Metab.* 84: 649–653, 1999.
- [76609] 14939. Menezes–Ferreira, M. M.; Eil, C.; Wortsman, J.; Weintraub, B. D.: Decreased nuclear uptake of [125–I]triiodo–L–thyronine in fibroblasts from patients with peripheral thyroid hormone resistance. *J. Clin. Endocr. Metab.* 59: 1081–1087, 1984.
- [76610] 14940. Mitelman, F.; Manolov, G.; Manolova, Y.; Billstrom, R.; Heim, S.; Kristoffersson, U.; Mandahl, N.; Ferro, M. T.; San Roman, C.: High resolution chromosome analysis of constitutional and acquired t(15;17) maps c-erbA to sub-band 17q11.2. *Cancer Genet. Cytogenet.* 22:95–98, 1986.
- [76611] 14941. Miyajima, N.; Horiuchi, R.; Shibuya, Y.; Fukushige,

S.; Matsubara, K.; Toyoshima, K.; Yamamoto, T.: Two erbA homologs encoding proteins with different T(3) binding capacities are transcribed from opposite DNA strands of the same genetic locus. *Cell* 57: 31–39, 1989.

[76612] 14942. Nagaya, T.; Nomura, Y.; Fujieda, M.; Seo, H.: Heterodimerization preferences of thyroid hormone receptor alpha isoforms. *Biochem. Biophys. Res. Commun.* 226: 426–430, 1996.

[76613] 14943. Nakai, A.; Seino, S.; Sakurai, A.; Szilak, I.; Bell, G. I.; DeGroot, L. J.: Characterization of a thyroid hormone receptor expressed in human kidney and other tissues. *Proc. Nat. Acad. Sci.* 85: 2781–2785, 1988.

[76614] 14944. Ng, L.; Rusch, A.; Amma, L. L.; Nordstrom, K.; Erway, L. C.; Vennstrom, B.; Forrest, D.: Suppression of the deafness and thyroid dysfunction in *Thrb*-null mice by an independent mutation in the *Thra* thyroid hormone receptor gene. *Hum. Molec. Genet.* 10: 2701–2708, 2001.

[76615] 14945. Puzianowska-Kuznicka, M.; Krystyniak, A.; Madej, A.; Cheng, S.-Y.; Nauman, J.: Functionally impaired TR mutants are present in thyroid papillary cancer. *J. Clin. Endocr. Metab.* 87: 1120–1128, 2002.

[76616] 14946. Refetoff, S.; DeGroot, L. J.; Benard, B.; DeWind, L. T.: Studies of a sibship with apparent hereditary resistance

to the intracellular action of thyroid hormone. *Metabolism* 21: 723–756, 1972.

[76617] 14947. Rider, S. H.; Bailey, C. J.; Voss, R.; Sheer, D.; Hiorns, L. R.; Solomon, E.: RFLP for the human erb-A1 gene. *Nucleic Acids Res.* 15:863 only, 1987.

[76618] 14948. Sakurai, A.; Nakai, A.; DeGroot, L. J.: Expression of three forms of thyroid hormone receptor in human tissues. *Molec. Endocr.* 3:392–399, 1989.

[76619] 14949. Cerretti, D. P.; Lyman, S. D.; Kozlosky, C. J.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Valentine, V.; Kirstein, M. N.; Shapiro, D. N.; Morris, S. W.: The genes encoding the Eph-related receptor tyrosine kinase ligands LERK-1 (EPLG1, Epl1), LERK-3 (EPLG3, Epl3), and LERK-4 (EPLG4, Epl4) are clustered on human chromosome 1 and mouse chromosome 3. *Genomics* 33: 277–282, 1996.

[76620] 14950. Holzman, L. B.; Marks, R. M.; Dixit, V. M.: A novel immediate-early response gene of endothelium is induced by cytokines and encodes a secreted protein. *Molec. Cell. Biol.* 10: 5830–5838, 1990.

[76621] 14951. Pandey, A.; Lindberg, R. A.; Dixit, V. M.: Receptor orphans find a family. *Curr. Biol.* 5: 986–989, 1995.

[76622] 14952. Bunz, F.; Dutriaux, A.; Lengauer, C.; Waldman, T.; Zhou, S.; Brown, J. P.; Sedivy, J. M.; Kinzler, K. W.; Vogel-

stein, B.: Requirement for p53 and p21 to sustain G2 arrest after DNA damage. *Science* 282:1497–1501, 1998.

[76623] 14953. Colucci-Guyon, E.; Portier, M.-M.; Dunia, I.; Paulin, D.; Pournin, S.; Babinet, C.: Mice lacking vimentin develop and reproduce without an obvious phenotype. *Cell* 79: 679–694, 1994.

[76624] 14954. Ferrari, S.; Battini, R.; Kaczmarek, L.; Rittling, S.; Calabretta, B.; de Riel, J. K.; Philiponis, V.; Weil, J.-F.; Baserga, R.: Coding sequence and growth regulation of the human vimentin gene. *Molec. Cell. Biol.* 6: 3614–3620, 1986.

[76625] 14955. Gieser, L.; Swaroop, A.: Expressed sequence tags and chromosomal localization of cDNA clones from a subtracted retinal pigment epithelium library. *Genomics* 13: 873–876, 1992.

[76626] 14956. Lilienbaum, A.; Legagneux, V.; Portier, M.-M.; Delagi, K.; Paulin, D.: Vimentin gene: expression in human lymphocytes and in Burkitt's lymphoma cells. *EMBO J.* 5: 2809–2814, 1986.

[76627] 14957. Marcus, E. M.; Smith, B. A.; Telenius, H.; Landsvater, R. M.; Buys, C. H. C. M.; Ferrari, S.; Ponder, B. A. J.; Mathew, C. G. P.: BclIRFLP for the human vimentin gene. *Nucleic Acids Res.* 16: 9068 only, 1988.



- [76628] 14958.Mathew, C. G.; Wakeling, W.; Jones, E.; Easton, D.; Fisher, R.;Strong, C.; Smith, B.; Chin, K.; Little, P.; Nakamura, Y.; Shows,T. B.; Jones, C.; Goodfellow, P. J.; Povey, S.; Ponder, B. A. J.:Regional localization of polymorphic markers on chromosome 10 by physicaland genetic mapping. *Ann. Hum. Genet.* 54: 121–129, 1990.
- [76629] 14959.Perreau, J.; Lilienbaum, A.; Vasseur, M.; Paulin, D.: Nucleotidesequence of the human vimentin gene and regulation of its transcriptionin tissues and cultured cells. *Gene* 62: 7–16, 1988.
- [76630] 14960.Gimona, M.; Small, J. V.; Moeremans, M.; Van Damme, J.; Puype,M.; Vandekerckhove, J.: Porcine vinculin and metavinculin differby a 68–residue insert located close to the carboxy–terminal partof the molecule. *EMBO J.* 7: 2329–2334, 1988.
- [76631] 14961.Burglen, L.; Amiel. J.; Viollet, L.; Lefebvre, S.; Burlet, P.;Clermont, O.; Raclin, V.; Landrieu, P.; Verloes, A.; Munnich, A.;Melki, J.: Survival motor neuron gene deletion in the arthrogryposismultiplex congenita–spinal muscular atrophy association. *J. Clin.Invest.* 98: 1130–1132, 1996.
- [76632] 14962.Gabreels, B. A. T. F.; Swaab, D. F.; de Kleijn, D. P. V.; Seidah,N. G.; Van de Loo, J.–W.; Van de Ven, W. J. M.; Martens, G. J. M.;van Leeuwen, F. W.: Attenuation of the

polypeptide 7B2, prohormoneconvertase PC2, and vaso-pressin in the hypothalamus of some Prader-Willipatients: indications for a processing defect. J. Clin. Endocr. Metab. 83:591-599, 1998.

[76633] 14963.Martens, G. J. M.: Cloning and sequence analysis of human pituitarycDNA encoding the novel polypeptide 7B2. FEBS Lett. 234: 160-164,1988.

[76634] 14964.Mattei, M. G.; Mbikay, M.; Sylla, B. S.; Lenoir, G.; Mattei, J.F.; Seidah, N. G.; Chretien, M.: Assignment of the gene for neuroendocrineprotein 7B2 (SGNE1 locus) to mouse chromosome region 2[E3-F3] andto human chromosome region 15q11-q15. Genomics 6: 436-440, 1990.

[76635] 14965.Mbikay, M.; Seidah, N. G.; Chretien, M.: Neuroendocrine secretoryprotein 7B2: structure, expression and functions. Biochem. J. 357:329-342, 2001.

[76636] 14966.Roebroek, A. J. M.; Dehaen, M. R. M.; van Bokhoven, A.; Martens,G. J. M.; Marynen, P.; van den Berghe, H.; Van de Ven, W. J. M.:Regional mapping of the human gene encoding the novel pituitary polypeptide7B2 to chromosome 15q13-q14 by in situ hybridization. Cytogenet.Cell Genet. 50: 158-160, 1989.

[76637] 14967.Westphal, C. H.; Muller, L.; Zhou, A.; Zhu, X.; Bonner-Weir, S.;Schambelan, M.; Steiner, D. F.; Lindberg, I.;

Leder, P.: The neuroendocrine protein 7B2 is required for peptide hormone processing in vivo and provides a novel mechanism for pituitary Cushing's disease. *Cell* 96:689–700, 1999.

[76638] 14968. Bell, K.; Hopper, K. E.; McKenzie, H. A.; Murphy, W. H.; Shaw, D. C.: A comparison of bovine alpha-lactalbumin A and B of Droughtmaster. *Biochim. Biophys. Acta* 214: 437–444, 1970.

[76639] 14969. Bell, K.; McKenzie, H. A.; Murphy, W. H.; Shaw, D. C.: Beta-lactoglobulin (Droughtmaster): a unique protein variant. *Biochim. Biophys. Acta* 214: 427–436, 1970.

[76640] 14970. Julkunen, M.; Seppala, M.; Janne, O. A.: Complete amino acid sequence of human placental protein 14: a progesterone-regulated uterine protein homologous to beta-lactoglobulins. *Proc. Nat. Acad. Sci.* 85: 8845–8849, 1988.

[76641] 14971. Morris, H. R.; Dell, A.; Easton, R. L.; Panico, M.; Koistinen, H.; Koistinen, R.; Oehninger, S.; Patankar, M. S.; Seppala, M.; Clark, G. F.: Gender-specific glycosylation of human glycodelin affects its contraceptive activity. *J. Biol. Chem.* 271: 32159–32167, 1996.

[76642] 14972. Song, M.; Ramaswamy, S.; Ramachandran, S.; Flowers, L. C.; Horowitz, I. R.; Rock, J. A.; Parthasarathy, S.: An-

giogenic role for glycodeilin tumorigenesis. Proc. Nat. Acad. Sci. 98: 9265–9270, 2001.

- [76643] 14973. Van Cong, N.; Vaisse, C.; Gross, M.-S.; Slim, R.; Milgrom, E.; Bernheim, A.: The human placental protein 14 (PP14) gene is localized on chromosome 9q34. Hum. Genet. 86: 515–518, 1991.
- [76644] 14974. Lee, F. S.; Fox, E. A.; Zhou, H.-M.; Strydom, D. J.; Vallee, B. L.: Primary structure of human placental ribonuclease inhibitor. Biochemistry 27: 8545–8553, 1988.
- [76645] 14975. Weremowicz, S.; Fox, E. A.; Morton, C. C.; Vallee, B. L.: The placental ribonuclease inhibitor (RNH) gene is located on chromosome subband 11p15.5. Genomics 8: 717–721, 1990.
- [76646] 14976. Zneimer, S. M.; Crawford, D.; Schneider, N. R.; Beutler, B.: Mapping of the human ribonuclease inhibitor gene (RNH) to chromosome 11p15 by in situ hybridization. Genomics 8: 175–178, 1990.
- [76647] 14977. Coughlin, P.; Nicholl, J.; Sun, J.; Salem, H.; Bird, P.; Sutherland, G. R.: Chromosomal mapping of the human proteinase inhibitor 6 (PI6) gene to 6p25 by fluorescence in situ hybridization. Genomics 26: 431–433, 1995.
- [76648] 14978. Coughlin, P.; Sun, J.; Cerruti, L.; Salem, H. H.; Bird, P.: Cloning and molecular characterization of a human in-

tracellular serine proteinaseinhibitor. Proc. Nat. Acad. Sci. 90: 9417–9421, 1993.

[76649] 14979.Siu, G.; Wurster, A. L.; Duncan, D. D.; Soliman, T. M.; Hedrick,S. M.: A transcriptional silencer controls the developmental expressionof the CD4 gene. EMBO J. 13: 3570–3579, 1994.

[76650] 14980.Zhang, K.; Westberg, J. A.; Paetau, A.; von Boguslawsky, K.; Lindsberg,P.; Erlander, M.; Guo, H.; Su, J.; Olsen, H. S.; Andersson, L. C.: High expression of stannio-calcin in differentiated brain neurons. Am.J. Path. 153: 439–445, 1998.

[76651] 14981.Hamilton, S. E.; Hurley, J. B.: A phosphodiesterase inhibitorspecific to a subset of bovine retinal cones. J. Biol. Chem. 265:11259–11264, 1990.

[76652] 14982.Shimizu–Matsumoto, A.; Itoh, K.; Inazawa, J.; Nishida, K.; Matsumoto,Y.; Kinoshita, S.; Matsubara, K.; Okubo, K.: Isolation and chromosomallocalization of the human cone cGMP phosphodiesterase gamma cDNA (PDE6H). Genomics 32:121–124, 1996.

[76653] 14983.Chen, J.; Engle, S. J.; Seilhamer, J. J.; Tischfield, J. A.: Cloningand recombinant expression of a novel human low molecular weight Ca(2+)-dependentphospholipase A2. J. Biol. Chem. 269: 2365–2368, 1994.

- [76654] 14984.Tischfield, J. A.; Xia, Y.-R.; Shih, D. M.; Klisak, I.; Chen, J.;Engle, S. J.; Siakotos, A. N.; Winstead, M. V.; Seilhamer, J. J.;Allamand, V.; Gyapay, G.; Lusk, A. J.: Low molecular weight, calcium-dependentphospholipase A(2) genes are linked and map to homologous chromosomeregions in mouse and human. *Genomics* 32: 328–333, 1996.
- [76655] 14985.Alexopoulou, L.; Thomas, V.; Schnare, M.; Lobet, Y.; Anguita, J.;Schoen, R. T.; Medzhitov, R.; Fikrig, E.; Flavell, R. A.: Hyporesponsiveness to vaccination with *Borrelia burgdorferi* OspA in humans and in TLR1–and TLR2–deficient mice. *Nature Med.* 8: 878–884, 2002.
- [76656] 14986.Gay, N. J.; Keith, F. J.: *Drosophila* Toll and IL–1 receptor. (Letter) *Nature* 351:355–356, 1991.
- [76657] 14987.Muzio, M.; Bosisio, D.; Polentarutti, N.; D'amico, G.; Stoppacciaro, A.; Mancinelli, R.; van't Veer, C.; Penton–Rol, G.; Ruco, L. P.; Allavena, P.; Mantovani, A.: Differential expression and regulation of Toll–likereceptors (TLR) in human leukocytes: selective expression of TLR3 in dendritic cells. *J. Immun.* 164: 5998–6004, 2000.
- [76658] 14988.Rock, F. L.; Hardiman, G.; Timans, J. C.; Kastelein, R. A.; Bazan, J. F.: A family of human receptors structurally related to *Drosophila* Toll. *Proc. Nat. Acad. Sci.* 95:

588–593, 1998.

[76659] 14989.Taguchi, T.; Mitcham, J. L.; Dower, S. K.; Sims, J. E.; Testa, J. R.: Chromosomal localization of TIL, a gene encoding a protein related to the *Drosophila* transmembrane receptor Toll, to human chromosome 4p14. *Genomics* 32: 486–488, 1996.

[76660] 14990.Xu, Y.; Tao, X.; Shen, B.; Horng, T.; Medzhitov, R.; Manley, J.L.; Tong, L.: Structural basis for signal transduction by the Toll/interleukin-1 receptor domains. *Nature* 408: 111–115, 2000.

[76661] 14991.Perrimon, N.; Mahowald, A. P.: Multiple functions of segment polarity genes in *Drosophila*. *Dev. Biol.* 119: 587–600, 1987.

[76662] 14992.Pizzuti, A.; Novelli, G.; Mari, A.; Ratti, A.; Colosimo, A.; Amati, F.; Penso, D.; Sangiuolo, F.; Calabrese, G.; Palka, G.; Silani, V.; Gennarelli, M.; Mingarelli, R.; Scarlato, G.; Scambler, P.; Dallapiccola, B.: Human homologue sequences to the *Drosophila* dishevelled segment-polarity gene are deleted in the DiGeorge syndrome. *Am. J. Hum. Genet.* 58:722–729, 1996.

[76663] 14993.Brown, E. J.; Albers, M. W.; Shin, T. B.; Ichikawa, K.; Keith, C. T.; Lane, W. S.; Schreiber, S. L.: A mammalian protein targeted by G1-arresting rapamycin-receptor

complex. Nature 369: 756–758,1994.

- [76664] 14994.Dennis, P. B.; Jaeschke, A.; Saitoh, M.; Fowler, B.; Kozma, S.C.; Thomas, G.: Mammalian TOR: a homeostatic ATP sensor. Science 294:1102–1105, 2001.
- [76665] 14995.Fang, Y.; Vilella–Bach, M.; Bachmann, R.; Flanigan, A.; Chen, J.: Phosphatidic acid–mediated mitogenic activation of mTOR signaling. Science 294:1942–1945, 2001.
- [76666] 14996.Hara, K.; Maruki, Y.; Long, X.; Yoshino, K.; Oshiro, N.; Hidayat,S.; Tokunaga, C.; Avruch, J.; Yonezawa, K.: Raptor, a binding partnerof target of rapamycin, mediates TOR action. Cell 110: 177–189,2002.
- [76667] 14997.Kim, D.–H.; Sarbassov, D. D.; Ali, S. M.; King, J. E.; Latek, R.R.; Erdjument–Bromage, H.; Tempst, P.; Sabatini, D. M.: mTOR interactswith raptor to form a nutrient–sensitive complex that signals to thecell growth machinery. Cell 110: 163–175, 2002.
- [76668] 14998.Lench, N. J.; Macadam, R.; Markham, A. F.: The human gene encodingFKBP–rapamycin associated protein (FRAP) maps to chromosomal band1p36.2. Hum. Genet. 99: 547–549, 1997.
- [76669] 14999.Moore, P. A.; Rosen, C. A.; Carter, K. C.: Assignment of the humanFKBP12–rapamycin–associated protein (FRAP) gene to chromosome 1p36by fluorescence in situ



hybridization. *Genomics* 33: 331–332, 1996.

- [76670] 15000. Onyango, P.; Lubyova, B.; Gardellin, P.; Kurzbauer, R.; Weith, A.: Molecular cloning and expression analysis of five novel genes in chromosome 1p36. *Genomics* 50: 187–198, 1998.
- [76671] 15001. Sabatini, D. M.; Erdjument-Bromage, H.; Lui, M.; Tempst, P.; Snyder, S. H.: RAFT1: a mammalian protein that binds to FKBP12 in a rapamycin-dependent fashion and is homologous to yeast TORs. *Cell* 78: 35–43, 1994.
- [76672] 15002. Barbier, M.; Attoub, S.; Calvez, R.; Laffargue, M.; Jarry, A.; Mareel, M.; Altruda, F.; Gespach, C.; Wu, D.; Lu, B.; Hirsch, E.; Wymann, M. P.: Weakening link to colorectal cancer? *Nature* 413:796 only, 2001.
- [76673] 15003. Crackower, M. A.; Oudit, G. Y.; Kozieradzki, I.; Sarao, R.; Sun, H.; Sasaki, T.; Hirsch, E.; Suzuki, A.; Shioi, T.; Irie-Sasaki, J.; Sah, R.; Cheng, H.-Y. M.; and 13 others: Regulation of myocardial contractility and cell size by distinct PI3K–PTEN signaling pathways. *Cell* 110:737–749, 2002.
- [76674] 15004. Hirsch, E.; Katanaev, V. L.; Garlanda, C.; Azzolino, O.; Pirola, L.; Silengo, L.; Sozzani, S.; Mantovani, A.; Altruda, F.; Wymann, M. P.: Central role for G protein-coupled phosphoinositide 3-kinase  $\gamma$  in inflammation.

Science 287: 1049–1053, 2000.

- [76675] 15005.Jiang, K.; Zhong, B.; Gilvary, D. L.; Corliss, B. C. Hong–Geller,E.; Wei, S; Djeu, J. Y.: Pivotal role of phosphoinositide–3 kinasein regulation of cytotoxicity in natural killer cells. Nature Immun. 1:419–425, 2000.
- [76676] 15006.Kikuta, Y.; Kato, M.; Yamashita, Y.; Miyauchi, Y.; Tanaka, K.;Kamada, N.; Kusunose, M.: Human leukotriene B4 omega–hydroxylase(CYP4F3) gene: molecular cloning and chromosomal localization. DNACell Biol. 17: 221–230, 1998.
- [76677] 15007.Kikuta, Y.; Kusunose, E.; Endo, K.; Yamamoto, S.; Sogawa, K.; Fujii–Kuriyama,Y.; Kusunose, M.: A novel form of cytochrome P–450 family 4 in humanpolymorphonuclear leukocytes: cDNA cloning and expression of leukotrieneB4 omega–hydroxylase. J. Biol. Chem. 268: 9376–9380, 1993.
- [76678] 15008.Klingensmith, J.; Nusse, R.; Perrimon, N.: The Drosophila segmentpolarity gene dishevelled encodes a novel protein required for responseto wingless signal. Genes Dev. 8: 118–130, 1994.
- [76679] 15009.Holmes, S. E.; Riazzi, M. A.; Gong, W.; McDermid, H. E.; Sellinger,B. T.; Hua, A.; Chen, F.; Wang, Z.; Zhang, G.; Roe, B.; Gonzalez,I.; McDonald–McGinn, D. M.; Zackai, E.;

Emanuel, B. S.; Budarf, M.L.: Disruption of the clathrin heavy chain-like gene (CLTCL) associated with features of DGS/VCFS: a balanced (21;22)(p12;q11) translocation. Hum.Molec. Genet. 6: 357–367, 1997.

[76680] 15010.Kedra, D.; Peyrard, M.; Fransson, I.; Collins, J. E.; Dunham, I.; Roe, B. A.; Dumanski, J. P.: Characterization of a second human clathrin heavy chain polypeptide gene (CLH-22) from chromosome 22q11. Hum.Molec. Genet. 5: 625–631, 1996.

[76681] 15011.Long, K. R.; Trofatter, J. A.; Ramesh, V.; McCormick, M. K.; Buckler, A. J.: Cloning and characterization of a novel human clathrin heavy chain gene (CLTCL). Genomics 35: 466–472, 1996.

[76682] 15012.Lagenaur, C.; Kunemund, V.; Fischer, G.; Fushiki, S.; Schachner, M.: Monoclonal M6 antibody interferes with neurite extension of cultured neurons. J. Neurobiol. 23: 71–88, 1992.

[76683] 15013.Shimizu, F.; Watanabe, T. K.; Fujiwara, T.; Takahashi, E.; Nakamura, Y.; Maekawa, H.: Isolation and mapping of the human glycoprotein M6 gene (GPM6A) to 4q33–to–q34. Cytogenet. Cell Genet. 74: 138–139, 1996.

[76684] 15014.Bui, T. D.; Beier, D. R.; Jonssen, M.; Smith, K.; Dorrington, S.M.; Kaklamanis, L.; Kearney, L.; Regan, R.; Suss-

man, D. J.; Harris, A. L.: cDNA cloning of a human dishevelled DVL-3 gene, mapping to 3q27, and expression in human breast and colon carcinomas. *Biochem. Biophys. Res. Commun.* 239: 510–516, 1997.

[76685] 15015. Cheng, Y.-S. E.; Patterson, C. E.; Staeheli, P.: Interferon-induced guanylate-binding proteins lack an N(T)KXD consensus motif and bind GMP in addition to GDP and GTP. *Molec. Cell. Biol.* 11: 4717–4725, 1991.

[76686] 15016. Kumar, S.; Li, Q.; Dua, A.; Ying, Y.-K.; Bagchi, M. K.; Bagchi, I. C.: Messenger ribonucleic acid encoding interferon-inducible guanylate-binding protein 1 is induced in human endometrium within the putative window of implantation. *J. Clin. Endocr. Metab.* 86: 2420–2427, 2001.

[76687] 15017. Prochazka, M.; Staeheli, P.; Holmes, R. S.; Haller, O.: Interferon-induced guanylate-binding proteins: mapping of the murine Gbp-1 locus to chromosome 3. *Virology* 145: 273–279, 1985.

[76688] 15018. Strehlow, I.; Lohmann-Matthes, M. L.; Decker, T.: The interferon-inducible GBP1 gene: structure and mapping to human chromosome 1. *Gene* 144: 295–299, 1994.

[76689] 15019. Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek, F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.; Copeland, N. G.: Murine chromosomal location of four

hepatocyte-enriched transcription factors: HNF-3-alpha, HNF3-beta, HNF-3-gamma, and HNF-4. *Genomics* 13:264-268, 1992.

- [76690] 15020. Chartier, F. L.; Bossu, J.-P.; Laudet, V.; Fruchart, J.-C.: Cloning and sequencing of cDNAs encoding the human hepatocyte nuclear factor 4 indicate the presence of two isoforms in human liver. *Gene* 147:269-272, 1994.
- [76691] 15021. Eeckhoute, J.; Formstecher, P.; Laine, B.: Maturity-onset diabetes of the young type 1 (MODY1)-associated mutations R154X and E276Q in hepatocyte nuclear factor 4-alpha (HNF4-alpha) gene impair recruitment of p300, a key transcriptional coactivator. *Molec. Endocr.* 15: 1200-1210, 2001.
- [76692] 15022. Furuta, H.; Iwasaki, N.; Oda, N.; Hinokio, Y.; Horikawa, Y.; Yamagata, K.; Yano, N.; Sugahiro, J.; Ogata, M.; Ohgawara, H.; Omori, Y.; Iwamoto, Y.; Bell, G. I.: Organization and partial sequence of the hepatocyte nuclear factor-4-alpha/MODY1 gene and identification of a missense mutation, R127W, in a Japanese family with MODY. *Diabetes* 46: 1652-1657, 1997.
- [76693] 15023. Hani, E. H.; Suaud, L.; Boutin, P.; Chevre, J.-C.; Durand, E.; Philippi, A.; Demenais, F.; Vionnet, N.; Furuta, H.; Velho, G.; Bell, G. I.; Laine, B.; Froguel, P.: A missense mu-

tation in hepatocytenuclear factor-4-alpha, resulting in a reduced transactivation activity,in human late-onset non-insulin-dependent diabetes mellitus. J. Clin.Invest. 101: 521-526, 1998.

- [76694] 15024.Lausen, J.; Thomas, H.; Lemm, I.; Bulman, M.; Borgschulze, M.;Lingott, A.; Hattersley, A. T.; Ryffel, G. U.: Naturally occurringmutations in the human HNF4-alpha gene impair the function of thetranscription factor to a varying degree. Nucleic Acids Res. 28:430-437, 2000.
- [76695] 15025.Li, J.; Ning, G.; Duncan, S. A.: Mammalian hepatocyte differentiationrequires the transcription factor HNF-4-alpha. Genes Dev. 14: 464-474,2000.
- [76696] 15026.Stoffel, M.; Duncan, S. A.: The maturity-onset diabetes of theyoung (MODY1) transcription factor HNF4-alpha regulates expressionof genes required for glucose transport and metabolism. Proc. Nat.Acad. Sci. 94: 13209-13214, 1997.
- [76697] 15027.Yamagata, K.; Furuta, H.; Oda, N.; Kaisaki, P. J.; Menzel, S.;Cox, N. J.; Fajans, S. S.; Signorini, S.; Stoffel, M.; Bell, G. I.: Mutations in the hepatocyte nuclear factor-4-alpha gene in maturity-onsetdiabetes of the young (MODY1). Nature 384: 458-460, 1996.
- [76698] 15028.Zouali, H.; Hani, E. H.; Philippi, A.; Vionnet, N.;

Beckmann, J. S.; Demenais, F.; Froguel, P.: A susceptibility locus for early-onset non-insulin dependent (type 2) diabetes mellitus maps to chromosome 20q, proximal to the phosphoenolpyruvate carboxykinase gene. *Hum. Molec. Genet.* 6: 1401–1408, 1997.

[76699] 15029. Frolova, L.; Le Goff, X.; Rasmussen, H. H.; Cheperegina, S.; Drugeon, G.; Kress, M.; Arman, I.; Haenni, A.-L.; Celis, J. E.; Philippe, M.; Justesen, J.; Kisselev, L.: A highly conserved eukaryotic protein family possessing properties of polypeptide chain release factor. *Nature* 372:701–703, 1994.

[76700] 15030. Guenet, L.; Henry, C.; Toutain, B.; Dubourg, C.; Le Gall, J. Y.; David, V.; Le Treut, A.: Eukaryotic translation termination factor gene (ETF1/eRF1) maps at D5S500 in a commonly deleted region of chromosome 5q31 in malignant myeloid diseases. *Cytogenet. Cell Genet.* 88: 82–86, 2000.

[76701] 15031. Hansen, L. L.; Jakobsen, C. G.; Justesen, J.: Assignment of the human translation termination factor 1 (ETF1) to 5q31.1 and of the proximal marker D5S1995 by radiation hybrid mapping. *Cytogenet. Cell Genet.* 87: 256–257, 1999.

[76702] 15032. Le Goff, X.; Philippe, M.; Jean-Jean, O.: Overex-

pression of human release factor 1 alone has an antipressor effect in human cells. *Molec. Cell Biol.* 17: 3164–3172, 1997.

- [76703] 15033. Song, H.; Mugnier, P.; Das, A. K.; Webb, H. M.; Evans, D. R.; Tuite, M. F.; Hemmings, B. A.; Barford, D.: The crystal structure of human eukaryotic release factor eRF1—mechanism of stop codon recognition and peptidyl-tRNA hydrolysis. *Cell* 100: 311–321, 2000.
- [76704] 15034. DiDonato, C. J.; Chen, X.-N.; Noya, D.; Korenberg, J. R.; Nadeau, J. H.; Simard, L. R.: Cloning, characterization, and copy number of the murine survival motor neuron gene: homolog of the spinal muscular atrophy-determining gene. *Genome Res.* 7: 339–352, 1997.
- [76705] 15035. DiDonato, C. J.; Morgan, K.; Carpten, J. D.; Fuerst, P.; Ingraham, S. E.; Prescott, G.; McPherson, J. D.; Wirth, B.; Zerres, K.; Hurko, O.; Wasmuth, J. J.; Mendell, J. R.; Burghes, A. H. M.; Simard, L. R.: Association between Ag1-CA alleles and severity of autosomal recessive proximal spinal muscular atrophy. *Am. J. Hum. Genet.* 55: 1218–1229, 1994.
- [76706] 15036. Feldkotter, M.; Schwarzer, V.; Wirth, R.; Wienker, T. F.; Wirth, B.: Quantitative analyses of SMN1 and SMN2 based on real-time lightCyclerPCR: fast and highly reliable



carrier testing and prediction of severity of spinal muscular atrophy. *Am. J. Hum. Genet.* 70: 358–368, 2002.

- [76707] 15037. Friesen, W. J.; Massenet, S.; Paushkin, S.; Wyce, A.; Dreyfuss, G.: SMN, the product of the spinal muscular atrophy gene, binds preferentially to dimethylarginine-containing protein targets. *Molec. Cell* 7: 1111–1117, 2001.
- [76708] 15038. Frugier, T.; Tiziano, F. D.; Cifuentes-Diaz, C.; Miniou, P.; Roblot, N.; Dierich, A.; Le Meur, M.; Melki, J.: Nuclear targeting defect of SMN lacking the C-terminus in a mouse model of spinal muscular atrophy. *Hum. Molec. Genet.* 9: 849–858, 2000.
- [76709] 15039. Gambardella, A.; Mazzei, R.; Toscano, A.; Annesi, G.; Pasqua, A.; Annesi, F.; Quattrone, F.; Oliveri, R. L.; Valentino, P.; Bono, F.; Aguglia, U.; Zappia, M.; Vita, G.; Quattrone, A.: Spinal muscular atrophy due to an isolated deletion of exon 8 of the telomeric survival motor neuron gene. *Ann. Neurol.* 44: 836–839, 1998.
- [76710] 15040. Hahnen, E.; Schonling, J.; Rudnik-Schoneborn, S.; Raschke, H.; Zerres, K.; Wirth, B.: Missense mutations in exon 6 of the survival motor neuron gene in patients with spinal muscular atrophy (SMA). *Hum. Molec. Genet.* 6: 821–825, 1997.
- [76711] 15041. Hannus, S.; Buhler, D.; Romano, M.; Seraphin, B.;

Fischer, U.: The *Schizosaccharomyces pombe* protein Yab8p and a novel factor, Yip1p, share structural and functional similarity with the spinal muscular atrophy-associated proteins SMN and SIP1. *Hum. Molec. Genet.* 9:663–674, 2000.

[76712] 15042. Hsieh-Li, H. M.; Chang, J.-G.; Jong, Y.-J.; Wu, M.-H.; Wang, N.M.; Tsai, C. H.; Li, H.: A mouse model for spinal muscular atrophy. *Nature Genet.* 24: 66–70, 2000.

[76713] 15043. Jablonka, S.; Bandilla, M.; Wiese, S.; Buhler, D.; Wirth, B.; Sendtner, M.; Fischer, U.: Co-regulation of survival of motor neuron (SMN) protein and its interactor SIP1 during development and in spinal muscular atrophy. *Hum. Molec. Genet.* 10: 497–505, 2001.

[76714] 15044. Liu, Q.; Dreyfuss, G.: A novel nuclear structure containing the survival of motor neurons protein. *EMBO J.* 15: 3555–3565, 1996.

[76715] 15045. Liu, Q.; Fischer, U.; Wang, F.; Dreyfuss, G.: The spinal muscular atrophy disease gene product, SMN, and its associated protein SIP1 are in a complex with spliceosomal snRNP proteins. *Cell* 90: 1013–1021, 1997.

[76716] 15046. Lorson, C. L.; Androphy, E. J.: The domain encoded by exon 2 of the survival motor neuron protein mediates nucleic acid binding. *Hum. Molec. Genet.* 7: 1269–1275,

1998.

- [76717] 15047.Lorson, C. L.; Hahnen, E.; Androphy, E. J.; Wirth, B.: A singlenucleotide in the SMN gene regulates splicing and is responsible forspinal muscular atrophy. Proc. Nat. Acad. Sci. 96: 6307–6311, 1999.
- [76718] 15048.Lorson, C. L.; Strasswimmer, J.; Yao, J.–M.; Baleja, J. D.; Hahnen,E.; Wirth, B.; Le, T.; Burghes, A. H. M.; An–drophy, E. J.: SMN oligomerizationdefect correlates with spinal muscular atrophy severity. Nature Genet. 19:63–66, 1998.
- [76719] 15049.McAndrew, P. E.; Parsons, D. W.; Simard, L. R.; Ro–chette, C.;Ray, P. N.; Mendell, J. R.; Prior, T. W.; Burghes, A. H.: Identificationof proximal spinal muscular atrophy carriers and patients by analysisof SMNT and SMNC gene copy number. Am. J. Hum. Genet. 60: 1411–1422,1997.
- [76720] 15050.Meister, G.; Buhler, D.; Lagerbauer, B.; Zobawa, M.; Lottspeich,F.; Fischer, U.: Characterization of a nuclear 20S complex containingthe survival of motor neurons (SMN) protein and a specific subsetof spliceosomal Sm proteins. Hum. Molec. Genet. 9: 1977–1986, 2000.
- [76721] 15051.Mohaghegh, P.; Rodrigues, N. R.; Owen, N.; Ponting, C. P.; Le,T. T.; Burghes, A. H. M.; Davies, K. E.: Analysis of mutations inthe tudor domain of the survival

motor neuron protein SMN. *Europ.J. Hum. Genet.* 7: 519–525, 1999.

- [76722] 15052. Monani, U. R.; Lorson, C. L.; Parsons, D. W.; Prior, T. W.; Androphy, E. J.; Burghes, A. H. M.; McPherson, J. D.: A single nucleotide difference that alters splicing patterns distinguishes the SMA gene SMN1 from the copy gene SMN2. *Hum. Molec. Genet.* 8: 1177–1183, 1999.
- [76723] 15053. Mourelatos, Z.; Abel, L.; Yong, J.; Kataoka, N.; Dreyfuss, G.: SMN interacts with a novel family of hnRNP and spliceosomal proteins. *EMBO J.* 20: 5443–5452, 2001.
- [76724] 15054. Owen, N.; Doe, C. L.; Mellor, J.; Davies, K. E.: Characterization of the *Schizosaccharomyces pombe* orthologue of the human survival motor neuron (SMN) protein. *Hum. Molec. Genet.* 9: 675–684, 2000.
- [76725] 15055. Pagliardini, S.; Giavazzi, A.; Setola, V.; Lizier, C.; Di Luca, M.; DeBiasi, S.; Battaglia, G.: Subcellular localization and axonal transport of the survival motor neuron (SMN) protein in the developing rat spinal cord. *Hum. Molec. Genet.* 9: 47–56, 2000.
- [76726] 15056. Parsons, D. W.; McAndrew, P. E.; Iannaccone, S. T.; Mendell, J. R.; Burghes, A. H. M.; Prior, T. W.: Intragenic telSMN mutations: frequency, distribution, evidence of a founder effect, and modification of the spinal muscular at-

rophy phenotype by cenSMN copy number. *Am.J. Hum. Genet.* 63: 1712–1723, 1998.

- [76727] 15057.Parsons, D. W.; McAndrew, P. E.; Monani, U. R.; Mendell, J. R.;Burghes, A. H. M.; Prior, T. W.: An 11 base pair duplication in exon6 of the SMN gene produces a type I spinal muscular atrophy (SMA)phenotype: further evidence for SMN as the primary SMA–determininggene. *Hum. Molec. Genet.* 5: 1727–1732, 1996.
- [76728] 15058.Lewandoski, M.; Sun, X.; Martin, G. R.: Fgf8 signalling from theAER is essential for normal limb development. *Nature Genet.* 26:460–463, 2000.
- [76729] 15059.Lorenzi, M. V.; Long, J. E.; Miki, T.; Aaronson, S. A.: Expressioncloning, developmental expression and chromosomal localization offibroblast growth factor–8. *Oncogene* 10: 2051–2055, 1995.
- [76730] 15060.Meyers, E. N.; Martin, G. R.: Differences in left–right axispathways in mouse and chick: functions of FGF8 and SHH. *Science* 285:403–406, 1999.
- [76731] 15061.Moon, A. M.; Capecchi, M. R.: Fgf8 is required for outgrowthand patterning of the limbs. *Nature Genet.* 26: 455–459, 2000.
- [76732] 15062.Payson, R. A.; Wu, J.; Liu, Y.; Chiu, I.–M.: The human FGF–8gene localizes on chromosome 10q24 and is

subjected to induction by androgen in breast cancer cells. *Oncogene* 13: 47–53, 1996.

[76733] 15063. Streit, A.; Berliner, A. J.; Papanayotou, C.; Sirulnik, A.; Stern, C. D.: Initiation of neural induction by FGF signalling before gastrulation. *Nature* 406:74–78, 2000.

[76734] 15064. Sun, X.; Meyers, E. N.; Lewandoski, M.; Martin, G. R.: Targeted disruption of *Fgf8* causes failure of cell migration in the gastrulating mouse embryo. *Genes Dev.* 13: 1834–1846, 1999.

[76735] 15065. White, R. A.; Dowler, L. L.; Angeloni, S. V.; Pasztor, L. M.; MacArthur, C. A.: Assignment of FGF8 to human chromosome 10q25–q26: mutations in FGF8 may be responsible for some types of acrocephalosyndactyly linked to this region. *Genomics* 30: 109–111, 1995.

[76736] 15066. Yoshiura, K.; Leysens, N. J.; Chang, J.; Ward, D.; Murray, J. C.; Muenke, M.: Genomic structure, sequence, and mapping of human FGF8 with no evidence for its role in craniosynostosis/limb defects syndromes. *Am. J. Med. Genet.* 72: 354–362, 1997.

[76737] 15067. Zammit, C.; Coope, R.; Gomm, J. J.; Shousha, S.; Johnston, C. L.; Coombes, R. C.: Fibroblast growth factor 8 is expressed at higher levels in lactating human breast and in breast cancer. *Brit. J. Cancer* 86:1097–1103, 2002.

- [76738] 15068.Cao, H.; Mok, A.; Miskie, B.; Hegele, R. A.: Single-nucleotide polymorphisms of the proprotein convertase subtilisin/kexin type 5 (PCSK5) gene. *J. Hum. Genet.* 46: 730–732, 2001.
- [76739] 15069.van de Loo, J.-W. H. P.; Creemers, J. W. M.; Kas, K.; Roebroek, A. J. M.; Van de Ven, W. J. M.: Assignment of the human proprotein convertase gene PCSK5 to chromosome 9q21.3 *Cytogenet. Cell Genet.* 75:227–229, 1996.
- [76740] 15070.Aramburu, J.; Garcia-Cozar, F.; Raghavan, A.; Okamura, H.; Rao, A.; Hogan, P. G.: Selective inhibition of NFAT activation by a peptide spanning the calcineurin targeting site of NFAT. *Molec. Cell* 1:627–637, 1998.
- [76741] 15071.Castigli, E.; Pahwa, R.; Good, R. A.; Geha, R. S.; Chatila, T.A.: Molecular basis of a multiple lymphokine deficiency in a patient with severe combined immunodeficiency. *Proc. Nat. Acad. Sci.* 90:4728–4732, 1993.
- [76742] 15072.Chatila, T.; Castigli, E.; Pahwa, R.; Pahwa, S.; Chirmule, N.; Oyaizu, N.; Good, R. A.; Geha, R. S.: Primary combined immunodeficiency resulting from defective transcription of multiple T-cell lymphokine genes. *Proc. Nat. Acad. Sci.* 87: 10033–10037, 1990.
- [76743] 15073.Horsley, V.; Pavlath, G. K.: NFAT: ubiquitous regulator of cell differentiation and adaptation. *J. Cell Biol.* 156:

771–774, 2002.

- [76744] 15074.Jauliac, S.; Lopez–Rodriguez, C.; Shaw, L. M.; Brown, L. F.; Rao,A.; Toker, A.: The role of NFAT transcription factors in integrin–mediated carcinoma invasion. *Nature Cell Biol.* 4: 540–544, 2002.
- [76745] 15075.Li, X.; Ho, S. N.; Luna, J.; Giacalone, J.; Thomas, D. J.; Timmerman,L. A.; Crabtree, G. R.; Francke, U.: Cloning and chromosomal localization of the human and murine genes for the T–cell transcription factors NFATc and NFATp. *Cytogenet. Cell Genet.* 68: 185–191, 1995.
- [76746] 15076.Northrop, J. P.; Ho, S. N.; Chen, L.; Thomas, D. J.; Timmerman,L. A.; Nolan, G. P.; Admon, A.; Crabtree, G. R.: NF–AT components define a family of transcription factors targeted in T–cell activation. *Nature* 369:497–502, 1994.
- [76747] 15077.Okamura, H.; Aramburu, J.; Garcia–Rodriguez, C.; Viola, J. P. B.;Raghavan, A.; Tahiliani, M.; Zhang, X.; Qin, J.; Hogan, P. G.; Rao,A.: Concerted dephosphorylation of the transcription factor NFAT1 induces a conformational switch that regulates transcriptional activity. *Molec.Cell* 6: 539–550, 2000.
- [76748] 15078.Pahwa, R.; Chatila, T.; Pahwa, S.; Paradise, C.; Day, N. K.; Geha,R.; Schwartz, S. A.; Slade, H.; Oyaizu, N.; Good, R. A.: Recombinant interleukin 2 therapy in severe



combined immunodeficiency disease. *Proc.Nat. Acad. Sci.* 86: 5069–5073, 1989.

- [76749] 15079.Park, J.; Takeuchi, A.; Sharma, S.: Characterization of a new isoform of the NFAT (nuclear factor of activated T cells) gene family member NFATc. *J. Biol. Chem.* 271: 20914–20921, 1996.
- [76750] 15080.Peng, S. L.; Gerth, A. J.; Ranger, A. M.; Glimcher, L. H.: NFATc1 and NFATc2 together control both T and B cell activation and differentiation. *Immunity* 14:13–20, 2001.
- [76751] 15081.Aoki, K.; Suzuki, K.; Sugano, T.; Tasaka, T.; Nakahara, K.; Kuge, O.; Omori, A.; Kasai, M.: A novel gene, 'Translin,' encodes a recombination hotspot binding protein associated with chromosomal translocations. *Nature-Genet.* 10: 167–174, 1995.
- [76752] 15082.Badge, R. M.; Yardley, J.; Jeffreys, A. J.; Armour, J. A. L.: Crossover breakpoint mapping identifies a sub-telomeric hotspot for male meiotic recombination. *Hum. Molec. Genet.* 9: 1239–1244, 2000.
- [76753] 15083.Hosaka, T.; Kanoe, H.; Nakayama, T.; Murakami, H.; Yamamoto, H.; Nakamata, T.; Tsuboyama, T.; Oka, M.; Kasai, M.; Sasaki, M. S.; Nakamura, T.; Toguchida, J.: Translin binds to the sequences adjacent to the breakpoints of the TLS and CHOP genes in liposarcomas with

translocationt(12;16). *Oncogene* 19: 5821–5825, 2000.

[76754] 15084.Kasai, M.; Aoki, K.; Matsuo, Y.; Minowada, J.; Maziarz, R. T.;Strominger, J. L.: Recombination hotspot associated factors specificallyrecognize novel target sequences at the site of interchromosomal rearrangementsin T-ALL patients with t(8;14)(q24;q11)and (t(1;14)(p32;q11). *Int.Immun.* 6: 1017–1025, 1994.

[76755] 15085.Arceci, R. J.; King, A. A.; Simon, M. C.; Orkin, S. H.; Wilson,D. B.: Mouse GATA-4: a retinoic acid-inducible GATA-binding transcriptionfactor expressed in endodermally derived tissues and heart. *Molec.Cell. Biol.* 13: 2235–2246, 1993.

[76756] 15086.Cirillo, L. A.; Lin, F. R.; Cuesta, I.; Friedman, D.; Jarnik, M.;Zaret, K. S.: Opening of compacted chromatin by early developmentaltranscription factors HNF3(FoxA) and GATA-4. *Molec. Cell* 9: 279–289,2002.

[76757] 15087.Crispino, J. D.; Lodish, M. B.; Thurberg, B. L.; Litovsky, S. H.;Collins, T.; Molkentin, J. D.; Orkin, S. H.: Proper coronary vasculardevelopment and heart morphogenesis depend on interaction of GATA-4with FOG cofactors. *Genes Dev.* 15: 839–844, 2001.

[76758] 15088.Hasegawa, K.; Lee, S. J.; Jobe, S. M.; Markham, B. E.; Kitsis,R. N.: cis-acting sequences that mediate induction

of beta-myosinheavy chain gene expression during left ventricular hypertrophy dueto aortic constriction. Circulation 96: 3943–3953, 1997.

- [76759] 15089.Huang, W.-Y.; Cukerman, E.; Liew, C.-C.: Identification of a GATAmotif in the cardiac alpha-myosin heavy-chain-encoding gene and isolationof a human GATA-4 cDNA. Gene 155: 219–223, 1995.
- [76760] 15090.Huang, W.-Y.; Heng, H. H. Q.; Liew, C.-C.: Assignment of the humanGATA4 gene to 8p23.1–p22 using fluorescence in situ hybridizationanalysis. Cytogenet. Cell Genet. 72: 217–218, 1996.
- [76761] 15091.Kennedy, S. J.; Teebi, A. S.; Adatia, I.; Teshima, I.: Inheritedduplication, dup(8)(p23.1p23.1)Pat, in a father and daughter withcongenital heart defects. (Letter) Am. J. Med. Genet. 104: 79–80,2001.
- [76762] 15092.Ketola, I.; Pentikainen, V.; Vaskivuo, T.; Ilvesmaki, V.; Herva,R.; Dunkel, L.; Tapanainen, J. S.; Toppari, J.; Heikinheimo, M.:Expression of transcription factor GATA-4 during human testiculardevelopment and disease. J. Clin. Endocr. Metab. 85: 3925–3931,2000.
- [76763] 15093.Laitinen, M. P. E.; Anttonen, M.; Ketola, I.; Wilson, D. B.; Ritvos,O.; Butzow, R.; Heikinheimo, M.: Transcription factors GATA-4 andGATA-6 and a GATA family co-

factor, FOG-2, are expressed in human ovary and sex cord-derived ovarian tumors. *J. Clin. Endocr. Metab.* 85:3476–3483, 2000.

- [76764] 15094. Molkentin, J. D.; Kalvakolanu, D. V.; Markham, B. E.: Transcription factor GATA-4 regulates cardiac muscle-specific expression of the  $\alpha$ -myosin heavy-chain gene. *Molec. Cell. Biol.* 14: 4947–4957, 1994.
- [76765] 15095. Pehlivan, T.; Pober, B. R.; Brueckner, M.; Garrett, S.; Slauch, R.; Van Rheeden, R.; Wilson, D. B.; Watson, M. S.; Hing, A. V.: GATA4 haploinsufficiency in patients with interstitial deletion of chromosome region 8p23.1 and congenital heart disease. *Am. J. Med. Genet.* 83:201–206, 1999.
- [76766] 15096. White, R. A.; Dowler, L. L.; Pasztor, L. M.; Gatson, L. L.; Adkison, L. R.; Angeloni, S. V.; Wilson, D. B.: Assignment of the transcription factor GATA4 gene to human chromosome 8 and mouse chromosome 14: Gata4 is a candidate gene for Ds (disorganization). *Genomics* 27: 20–26, 1995.
- [76767] 15097. Brenneman, M. A.; Wagener, B. M.; Miller, C. A.; Allen, C.; Nickoloff, J. A.: XRCC3 controls the fidelity of homologous recombination: roles for XRCC3 in late stages of recombination. *Molec. Cell* 10: 387–395, 2002.

- [76768] 15098.Liu, N.; Lamerdin, J. E.; Tebbs, R. S.; Schild, D.; Tucker, J.D.; Shen, M. R.; Brookman, K. W.; Siciliano, M. J.; Walter, C. A.; Fan, W.; Narayana, L. S.; Zhou, Z.-Q.; Adamson, A. W.; Sorensen, K.J.; Chen, D. J.; Jones, N. J.; Thompson, L. H.: XRCC2 and XRCC3, new human Rad51-family members, promote chromosome stability and protect against DNA cross-links and other damages. *Molec. Cell* 1: 783–793, 1998.
- [76769] 15099.Tebbs, R. S.; Zhao, Y.; Tucker, J. D.; Scheerer, J. B.; Siciliano, M. J.; Hwang, M.; Liu, N.; Legerski, R. J.; Thompson, L. H.: Correction of chromosomal instability and sensitivity to diverse mutagens by a cloned cDNA of the XRCC3 DNA repair gene. *Proc. Nat. Acad. Sci.* 92:6354–6358, 1995.
- [76770] 15100.Winsey, S. L.; Haldar, N. A.; Marsh, H. P.; Bunce, M.; Marshall, S. E.; Harris, A. L.; Wojnarowska, F.; Welsh, K. I.: A variant within the DNA repair gene XRCC3 is associated with the development of melanoma skin cancer. *Cancer Res.* 60: 5612–5616, 2000.
- [76771] 15101.Hayashida, S.; Yamasaki, K.; Asada, Y.; Soeda, E.; Niikawa, N.; Kishino, T.: Construction of a physical and transcript map flanking the imprinted MEST/PEG1 region at 7q32. *Genomics* 66: 221–225, 2000.

- [76772] 15102.de Wind, N.; Dekker, M.; Claij, N.; Jansen, L.; van Klink, Y.;Radman, M.; Riggins, G.; van der Valk, M.; van't Wout, K.; te Riele,H.: HNPCC–like cancer predisposition in mice through simultaneousloss of Msh3 and Msh6 mismatch–repair protein functions. *Nature Genet.* 23:359–362, 1999.
- [76773] 15103.Braun–Dullaes, R. C.; Mann, M. J.; Ziegler, A.; von der Leyen,H. E.; Dzau, V. J.: A novel role for the cyclin–dependent kinaseinhibitor p27(Kip1) in angiotensin II–stimulated vascular smooth musclecell hypertrophy. *J. Clin. Invest.* 104: 815–823, 1999.
- [76774] 15104.Carrano, A. C.; Eytan, E.; Hershko, A.; Pagano, M.: SKP2 is requiredfor ubiquitin–mediated degradation of the CDK inhibitor p27. *NatureCell Biol.* 1: 193–199, 1999.
- [76775] 15105.Di Cristofano, A.; De Acetis, M.; Koff, A.; Cordon–Cardo, C.; Pandolfi,P. P.: Pten and p27(KIP1) cooperate in prostate cancer tumor suppressionin the mouse. *Nature Genet.* 27: 222–224, 2001.
- [76776] 15106.Fero, M. L.; Rivkin, M.; Tasch, M.; Porter, P.; Carow, C. E.; Firpo,E.; Polyak, K.; Tsai, L.–H.; Broudy, V.; Perlmutter, R. M.; Kaushansky,K.; Roberts, J. M.: A syndrome of multiorgan hyperplasia with featuresof gigantism, tumorigenesis, and female sterility in p27(Kip1)–deficientmice.

Cell 85: 733–744, 1996.

[76777] 15107. Albertella, M. R.; Campbell, R. D.: Characterization of a novel gene in the human major histocompatibility complex that encodes a potential new member of the I kappa B family of proteins. *Hum. Molec. Genet.* 3: 793–799, 1994.

[76778] 15108. Buyse, I. M.; Shao, G.; Huang, S.: The retinoblastoma protein binds to RIZ, a zinc-finger protein that shares an epitope with the adenovirus E1A protein. *Proc. Nat. Acad. Sci.* 92: 4467–4471, 1995.

[76779] 15109. Buyse, I. M.; Takahashi, E.; Huang, S.: Physical mapping of the retinoblastoma interacting zinc finger gene RIZ to D1S228 on chromosome 1p36. *Genomics* 34: 119–121, 1996.

[76780] 15110. Chadwick, R. B.; Jiang, G.-L.; Bennington, G. A.; Yuan, B.; Johnson, C. K.; Stevens, M. W.; Niemann, T. H.; Peltomaki, P.; Huang, S.; de la Chapelle, A.: Candidate tumor suppressor RIZ is frequently involved in colorectal carcinogenesis. *Proc. Nat. Acad. Sci.* 97: 2662–2667, 2000.

[76781] 15111. Mock, B. A.; Coleman, M. P.; Huang, S.: RIZ maps to distal chromosome 4 near genes involved in tumorigenesis and nerve degeneration. *Mammalian Genome* 7: 637 only, 1996.

- [76782] 15112. Poetsch, M.; Dittberner, T.; Woenckhaus, C.: Frameshift mutations of RIZ, but no point mutations in RIZ1 exons in malignant melanomas with deletions in 1p36. *Oncogene* 21: 3038–3042, 2002.
- [76783] 15113. Malek, N. P.; Sundberg, H.; McGrew, S.; Nakayama, K.; Kyriakidis, T. R.; Roberts, J. M.: A mouse knock-in model exposes sequential proteolytic pathways that regulate p27(Kip1) in G1 and S phase. *Nature* 413:323–327, 2001.
- [76784] 15114. Martin, E.; Cacheux, V.; Cave, H.; Lapierre, J. M.; Le Paslier, D.; Grandchamp, B.: Localization of the CDKN4/p27(Kip1) gene to human chromosome 12p12.3. *Hum. Genet.* 96: 668–670, 1995.
- [76785] 15115. Mitsuhashi, T.; Aoki, Y.; Eksioglu, Y. Z.; Takahashi, T.; Bhide, P. G.; Reeves, S. A.; Caviness, V. S., Jr.: Overexpression of p27(Kip1) lengthens the G1 phase in a mouse model that targets inducible gene expression to central nervous system progenitor cells. *Proc. Nat. Acad. Sci.* 98: 6435–6440, 2001.
- [76786] 15116. Polyak, K.; Lee, M.-H.; Erdjument-Bromage, H.; Koff, A.; Roberts, J. M.; Tempst, P.; Massague, J.: Cloning of p27(Kip1), a cyclin-dependent kinase inhibitor and a potential mediator of extracellular antimitogenic signals.



Cell 78: 59–66, 1994.

- [76787] 15117.Saito, T.; Seki, N.; Hattori, A.; Hayashi, A.; Abe, M.; Araki,R.; Fujimori, A.; Fukumura, R.; Kozuma, S.; Matsuda, Y.: Structure,expression profile, and chromosomal location of a mouse gene homologousto human DNA–PK(cs) interacting protein (KIP) gene. *Mammalian Genome* 10:315–317, 1999.
- [76788] 15118.Sherr, C. J.; Roberts, J. M.: Inhibitors of mammalian G1 cyclin–dependentkinases. *Genes Dev.* 9: 1149–1163, 1995.
- [76789] 15119.Toyoshima, H.; Hunter, T.: p27, a novel inhibitor of G1 cyclin–Cdkprotein kinase activity, is related to p21. *Cell* 78: 67–74, 1994.
- [76790] 15120.Hudson, A. W.; Birnbaum, M. J.: Identification of a nonneuronalisoform of synaptotagmin. *Proc. Nat. Acad. Sci.* 92: 5895–5899, 1995.
- [76791] 15121.Chen, C.–C.; Akopian, A. N.; Sivilotti, L.; Colquhoun, D.; Burnstock,G.; Wood, J. N.: A P2X purinoceptor expressed by a subset of sensoryneurons. *Nature* 377: 428–430, 1995.
- [76792] 15122.Cockayne, D. A.; Hamilton, S. G.; Zhu, Q.–M.; Dunn, P. M.; Zhong,Y.; Novakovic, S.; Malmberg, A. B.; Cain, G.; Berson, A.; Kassotakis,L.; Hedley, L.; Lachnit, W.

G.; Burnstock, G.; McMahon, S. B.; Ford, A. P. D. W.: Urinary bladder hyporeflexia and reduced pain-related behaviour in P2X<sub>3</sub>-deficient mice. *Nature* 407: 1011–1015, 2000.

[76793] 15123. Cook, S. P.; McCleskey, E. W.: ATP, pain and a full bladder. *Nature* 407: 951–952, 2000.

[76794] 15124. Garcia-Guzman, M.; Stuhmer, W.; Soto, F.: Molecular characterization and pharmacological properties of the human P2X<sub>3</sub> purinoceptor. *Molec. Brain Res.* 47: 59–66, 1997.

[76795] 15125. Kennedy, C.; Leff, P.: Painful connection for ATP. *Nature* 377: 385–386, 1995.

[76796] 15126. Lewis, C.; Neldhart, S.; Holy, C.; North, R. A.; Buell, G.; Surprenant, A.: Coexpression of P2X<sub>2</sub> and P2X<sub>3</sub> receptor subunits can account for ATP-gated currents in sensory neurons. *Nature* 377: 432–435, 1995.

[76797] 15127. Souslova, V.; Cesare, P.; Ding, Y.; Akopian, A. N.; Stanfa, L.; Suzuki, R.; Carpenter, K.; Dickenson, A.; Boyce, S.; Hill, R.; Nebenius-Oosthuizen, D.; Smith, A. J. H.; Kidd, E. J.; Wood, J. N.: Warm-coding deficits and aberrant inflammatory pain in mice lacking P2X<sub>3</sub> receptors. *Nature* 407: 1015–1017, 2000.

[76798] 15128. Souslova, V.; Ravenall, S.; Fox, M.; Wells, D.; Wood, J. N.; Akopian, A. N.: Structure and chromosomal mapping

of the mouse P2X(3) gene. *Gene* 195:101–111, 1997.

[76799] 15129.Khakh, B. S.; Zhou, X.; Sydes, J.; Galligan, J. J.; Lester, H.A.: State-dependent cross-inhibition between transmitter-gated cationchannels. *Nature* 406: 405–410, 2000.

[76800] 15130.Buell, G.; Lewis, C.; Collo, G.; North, R. A.; Surprenant, A.:An antagonist-insensitive P2X receptor expressed in epithelia andbrain. *EMBO J.* 15: 55–62, 1996.

[76801] 15131.Buell, G.; Lewis, C.; Collo, G.; North, R. A.; Surprenant, A.:Cloning and expression of a family of ATP-gated ion channels. (Abstract) *Soc.Neurosci.* 21: 1262, 1995.

[76802] 15132.Garcia-Guzman, M.; Soto, F.; Gomez-Hernandez, J. M.; Lund, P. E.;Stuhmer, W.: Characterization of the recombinant human P2X4 receptorreveals pharmacological differences to the rat homologue. *Molec.Pharm.* 51: 109–118, 1997.

[76803] 15133.Chen, J. D.; Evans, R. M.: A transcriptional co-repressor thatinteracts with nuclear hormone receptors. *Nature* 377: 454–457, 1995.

[76804] 15134.Fischle, W.; Dequiedt, F.; Hendzel, M. J.; Guenther, M. G.; Lazar,M. A.; Voelter, W.; Verdin, E.: Enzymatic activity associated withclass II HDACs is dependent on a multi-

protein complex containing HDAC3 and SMRT/N-CoR.

Molec. Cell 9: 45–57, 2002.

- [76805] 15135. Horlein, A. J.; Naar, A. M.; Heinzl, T.; Torchia, J.; Gloss, B.; Kurokawa, R.; Ryan, A.; Kamel, Y.; Soderstrom, M.; Glass, C. K.; Rosenfeld, M. G.: Ligand-independent repression by the thyroid hormone receptor mediated by a nuclear receptor co-repressor. Nature 377: 397–403, 1995.
- [76806] 15136. Hu, X.; Lazar, M. A.: The CoRNR motif controls the recruitment of corepressors by nuclear hormone receptors. Nature 402: 93–96, 1999.
- [76807] 15137. Jiang, Q.; Galiegue-Zouitina, S.; Roumier, C.; Hildebrand, M. P.; Thomas, S.; Coignet, L. J.: Genomic organization and refined mapping of the human nuclear corepressor 2 (NCOR2)/ silencing mediator of retinoid and thyroid hormone receptor (SMRT) gene on chromosome 12q24.3. Cytogenet. Cell Genet. 92: 217–220, 2001.
- [76808] 15138. Ordentlich, P.; Downes, M.; Xie, W.; Genin, A.; Spinner, N. B.; Evans, R. M.: Unique forms of human and mouse nuclear receptor corepressor SMRT. Proc. Nat. Acad. Sci. 96: 2639–2644, 1999.
- [76809] 15139. Colvin, J. S.; Green, R. P.; Schmahl, J.; Capel, B.; Ornitz, D. M.: Male-to-female sex reversal in mice lacking fi-

broblast growthfactor 9. Cell 104: 875–889, 2001.

[76810] 15140.Mattei, M.–G.; Penault–Llorca, F.; Coulier, F.; Birnbaum, D.:The human FGF9 gene maps to chromosomal region 13q11–q12. Genomics 29:811–812, 1995.

[76811] 15141.Miyamoto, M.; Naruo, K.–I.; Seko, C.; Matsumoto, S.; Kondo, T.;Kurokawa, T.: Molecular cloning of a novel cytokine cDNA encodingthe ninth member of the fibroblast growth factor family, which hasa unique secretion property. Molec. Cell. Biol. 13: 4251–4259, 1993.

[76812] 15142.Garcia, J. G.; Lazar, V.; Gilbert–McClain, L. I.; Gallagher, P.J.; Verin, A. D.: Myosin light chain kinase in endothelium: molecularcloning and regulation. Am. J. Resp. Cell Molec. Biol. 16: 489–494,1997.

[76813] 15143.Giorgi, D.; Brand–Arpon, V.; Rouquier, S.: The functional myosinlight chain kinase (MYLK) gene localizes with marker D3S3552 on humanchromosome 3q21 in a greater than 5–Mb yeast artificial chromosomeregion and is not linked to olfactory receptor genes. Cytogenet.Cell Genet. 92: 85–88, 2001.

[76814] 15144.Lazar, V.; Garcia, J. G. N.: A single human myosin light chainkinase gene (MLCK; MYLK) transcribes multiple nonmuscle isoforms. Genomics 57:256–267, 1999.

[76815] 15145.Potier, M.–C.; Chelot, E.; Pekarsky, Y.; Gardiner, K.;

Rossier,J.; Turnell, W. G.: The human myosin light chain kinase (MLCK) fromhippocampus: cloning, sequencing, expression, and localization to3cen-q21. Genomics 29: 562–570, 1995.

- [76816] 15146.Walker, L. A.; MacDonald, J. A.; Liu, X.; Nakamoto, R. K.; Haystead,T. A. J.; Somlyo, A. V.; Somlyo, A. P.: Site-specific phosphorylationand point mutations of telokin modulate its Ca(2+)-desensitizing effectin smooth muscle. J. Biol. Chem. 276: 24519–24524, 2001.
- [76817] 15147.Watterson, D. M.; Schavocky, J. P.; Guo, L.; Weiss, C.; Chlenski,A.; Shirinsky, V. P.; Van Eldik, L. J.; Haiech, J.: Analysis of thekinase-related protein gene found at human chromosome 3q21 in a multi-genecluster: organization, expression, alternative splicing, and polymorphic-marker. J. Cell. Biochem. 75: 481–491, 1999.
- [76818] 15148.Allcock, R. J. N.; Christiansen, F. T.; Price, P.: The centralMHC gene IKBL carries a structural polymorphism that is associatedwith HLA-A3,B7,DR15. Immunogenetics 49: 660–665, 1999.
- [76819] 15149.Druck, T.; Gu, Y.; Prabhala. G.; Cannizzaro, L. A.; Park, S.-H.;Huebner, K.; Keen, J. H.: Chromosome localization of human genesfor clathrin adaptor polypeptides AP2-beta and AP50 and the clathrin-bindingprotein, VCP.

Genomics 30: 94–97, 1995.

- [76820] 15150.Egerton, M.; Ashe, O. R.; Chen, D.; Druker, B. J.: Burgess, W.H.: Samelson, L. E.:VCP, the mammalian homolog of cdc48, is tyrosinephosphorylated in response to T cell antigen receptor activation. EMBOJ. 11: 3533–3540, 1992.
- [76821] 15151.Hoyle, J.; Tan, K. H.; Fisher, E. M. C.: Mapping the valosin-containingprotein (VCP) gene on human chromosome 9 and mouse chromosome 4, anda likely pseudo-gene on the mouse X chromosome. Mammalian Genome 8:778–780, 1997.
- [76822] 15152.Koller, K. J.; Brownstein, M. J.: Use of a cDNA clone to identifya supposed precursor protein containing valosin. Nature 325: 542–545,1987.
- [76823] 15153.Pleasure, I. T.; Black, M. M.; Keen, J. H.: Valosin-containingprotein, VCP, is a ubiquitous clathrin-binding protein. Nature 365:459–462, 1993.
- [76824] 15154.Ye, Y.; Meyer, H. H.; Rapoport, T. A.: The AAA ATPase Cdc48/p97and its partners transport proteins from the ER into the cytosol. Nature 414:652–656, 2001.
- [76825] 15155.Liu, Q.; Feng, Y.; Forgac, M.: Activity and in vitro reassemblyof the coated vesicle (H<sup>+</sup>)–ATPase requires the 50–kDa subunit of theclathrin assembly complex AP-2. J.

Biol. Chem. 269: 31592–31597,1994.

- [76826] 15156.Thurieu, C.; Brosius, J.; Burne, C.; Jolles, P.; Keen, J. H.;Mattaliano, R. J.; Chow, E. P.; Ramachandran, K. L.; Kirchhausen,T.: Molecular cloning and complete amino acid sequence of AP50, an assembly protein associated with clathrin-coated vesicles. DNA 7:663–669, 1988.
- [76827] 15157.Ponnambalam, S.; Robinson, M. S.; Jackson, A. P.; Peiperl, L.;Parham, P.: Conservation and diversity in families of coated vesicle adaptins. J. Biol. Chem. 265: 4814–4820, 1990.
- [76828] 15158.Robinson, M. S.: Cloning of cDNAs encoding two related 100-kD coated vesicle proteins (alpha-adaptins). J. Cell Biol. 108: 833–42,1989.
- [76829] 15159.Waelter, S.; Scherzinger, E.; Hasenbank, R.; Nordhoff, E.; Lurz,R.; Goehler, H.; Gauss, C.; Sathasivam, K.; Bates, G. P.; Lehrach,H.; Wanker, E. E.: The huntingtin interacting protein HIP1 is a clathrin and alpha-adaptin-binding protein involved in receptor-mediated endocytosis. Hum.Molec. Genet. 10: 1807–1817, 2001.
- [76830] 15160.Kaneko-Ishino, T.; Kuroiwa, Y.; Miyoshi, N.; Kohda, T.; Suzuki,R.; Yokoyama, M.; Viville, S.; Barton, S. C.; Ishino, F.; Surani,M. A.: Peg1/Mest imprinted gene on chromosome 6 identified by cDNA subtraction hybridiza-



tion. *Nature Genet.* 11: 52–59, 1995.

[76831] 15161.Kobayashi, S.; Kohda, T.; Miyoshi, N.; Kuroiwa, Y.; Aisaka, K.;Tsutsumi, O.; Kaneko–Ishino, T.; Ishino, F.: Human PEG1/MEST, an imprinted gene on chromosome 7. *Hum. Molec. Genet.* 6: 781–786, 1997.

[76832] 15162.Kosaki, K.; Kosaki, R.; Craigen, W. J.; Matsuo, N.: Isoform–specific imprinting of the human PEG1/MEST gene. (Letter) *Am. J. Hum. Genet.* 66:309–312, 2000.

[76833] 15163.Kotzot, D.; Schmitt, S.; Bernasconi, F.; Robinson, W. P.; Lurie, I. W.; Ilyina, H.; Mehes, K.; Hamel, B. C. J.; Otten, B. J.; Hergersberg, M.; Werder, E.; Shoenle, E.; Schinzel, A.: Uniparental disomy 7 in Silver–Russell syndrome and primordial growth retardation. *Hum. Molec. Genet.* 4: 583–587, 1995.

[76834] 15164.Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Keverne, E. B.; Surani, M. A.: Abnormal maternal behaviour and growth retardation associated with loss of the imprinted gene Mest. *Nature Genet.* 20:163–169, 1998.

[76835] 15165.Lefebvre, L.; Viville, S.; Barton, S. C.; Ishino, F.; Surani, M.A.: Genomic structure and parent–of–origin–specific methylation of Peg1. *Hum. Molec. Genet.* 6: 1907–1915, 1997.

[76836] 15166.Seki, T.; Tada, S.; Katada, T.; Enomoto, T.: Cloning

of a cDNA encoding a novel importin- $\alpha$  homologue, Qip1: discrimination of Qip1 and Rch1 from hSrp1 by their ability to interact with DNA helicase Q1/RecQL. *Biochem. Biophys. Res. Commun.* 234: 48–53, 1997.

- [76837] 15167. Ayala–Madrigal, M. L.; Doerr, S.; Ramirez–Duenas, M. L.; Hansmann, I.: Assignment of karyopherin  $\alpha$  1 (KPNA1) to human chromosome band 3q21 by in situ hybridization. *Cytogenet. Cell Genet.* 90: 58–59, 2000.
- [76838] 15168. Conti, E.; Uy, M.; Leighton, L.; Blobel, G.; Kuriyan, J.: Crystallographic analysis of the recognition of a nuclear localization signal by the nuclear import factor karyopherin  $\alpha$ . *Cell* 94: 193–204, 1998.
- [76839] 15169. Cortes, P.; Ye, Z.–S.; Baltimore, D.: RAG–1 interacts with the repeated amino acid motif of the human homologue of the yeast protein SRP1. *Proc. Nat. Acad. Sci.* 91: 7633–7637, 1994.
- [76840] 15170. Chen, H.; Antonarakis, S. E.: Localization of a human homolog of the mouse Tiam–1 gene to chromosome 21q22.1. *Genomics* 30: 123–127, 1995.
- [76841] 15171. Habets, G. G. M.; Scholtes, E. H. M.; Zuydgeest, D.; van der Kammen, R. A.; Stam, J. C.; Berns, A.; Collard, J. G.: Identification of an invasion–inducing gene, Tiam–1, that encodes a protein with homology to GDP–GTP exchangers

for Rho-like proteins. Cell 77: 537–549, 1994.

- [76842] 15172.Habets, G. G. M.; van der Kammen, R. A.; Jenkins, N. A.; Gilbert,D. J.; Copeland, N. G.; Hagemeijer, A.; Collard, J. G.: The invasion-inducingTIAM1 gene maps to human chromosome band 21q22 and mouse chromosome16. Cytogenet. Cell. Genet. 70: 48–51, 1995.
- [76843] 15173.Habets, G. G. M.; van der Kammen, R. A.; Stam, J. C.; Michiels,F.; Collard, J. G.: Sequence of the human invasion-inducing TIAM1gene, its conservation in evolution and its expression in tumor celllines of different tissue origin. Oncogene 10: 1371–1376, 1995.
- [76844] 15174.Malliri, A.; van der Kammen, R. A.; Clark, K.; van der Valk, M.;Michiels, F.; Collard, J. G.: Mice deficient in the Rac activatorTiam1 are resistant to Ras-induced skin tumours. Nature 417: 867–871,2002.
- [76845] 15175.Ghetti, A.; Pinol-Roma, S.; Michael, W. M.; Morandi, C.; Dreyfuss,G.: HNRNP I, the polypyrimidine tract-binding protein: distinct nuclearlocalization and association with hnRNAs. Nucleic Acids Res. 20:3671–3678, 1992.
- [76846] 15176.Raimondi, E.; Romanelli, M. G.; Moralli, D.; Gamberi, C.; Russo,M. P.; Morandi, C.: Assignment of the human gene encoding heterogeneousnuclear RNA ribonucleoprotein I (PTB) to chromosome 14q23–q24.1. Genomics

27:553–555, 1995.

- [76847] 15177. Romanelli, M. G.; Lorenzi, P.; Morandi, C.: Organization of the human gene encoding heterogeneous nuclear ribonucleoprotein type I (hnRNP I) and characterization of hnRNP I related pseudogene. *Gene* 255:267–272, 2000.
- [76848] 15178. Hoogenraad, C. C.; Koekkoek, B.; Akhmanova, A.; Krugers, H.; Dortland, B.; Miedema, M.; van Alphen, A.; Kistler, W. M.; Jaegle, M.; Koutsourakis, M.; Van Camp, N.; Verhoye, M.; van der Linden, A.; Kaverina, I.; Grosveld, F.; De Zeeuw, C. I.; Galjart, N.: Targeted mutation of *Cyln2* in the Williams syndrome critical region links CLIP–115 haploinsufficiency to neurodevelopmental abnormalities in mice. *Nature Genet.* 32: 116–127, 2002. Note: Erratum: *Nature Genet.* 32: 331 only, 2002.
- [76849] 15179. Gollob, M. H.; Green, M. S.; Roberts, R.: A gene responsible for familial Wolff–Parkinson–White syndrome. (Letter) *New Eng. J. Med.* 345:1063–1064, 2001.
- [76850] 15180. L'Allemand, D.; Tardy, V.; Gruters, A.; Schnabel, D.; Krude, H.; Morel, Y.: How a patient homozygous for a 30–kb deletion of the C4–CYP21 genomic region can have a nonclassic form of 21–hydroxylase deficiency. *J. Clin. Endocr. Metab.* 85: 4562–4567, 2000.
- [76851] 15181. Lako, M.; Ramsden, S.; Campbell, R. D.; Strachan,

T.: Mutationscreening in British 21-hydroxylase deficiency families and developmentof novel microsatellite based approaches to prenatal diagnosis. J.Med. Genet. 36: 119-124, 1999.

[76852] 15182.Lau, I. F.; Soardi, F. C.; Lemos-Marini, S. H. V.; Guerra, G.,Jr.; Baptista, M. T. M.; De Mello, M. P.: H28+C insertion in theCYP21 gene: a novel frameshift mutation in a Brazilian patient withthe classical form of 21-hydroxylase deficiency. J. Clin. Endocr.Metab. 86: 5877-5880, 2001.

[76853] 15183.Laue, L.; Merke, D. P.; Jones, J. V.; Barnes, K. M.; Hill, S.;Cutler, G. B.: A preliminary study of flutamide, testolactone, andreduced hydrocortisone dose in the treatment of congenital adrenalhyperplasia. J. Clin. Endocr. Metab. 81: 3535-3539, 1996.

[76854] 15184.Layrisse, A.; White, C.; Gunczler, P.; Gafaro Valera, L.; Arias,S.; Yunis, E. J.; Alper, C. A.; Awdeh, Z. L.: Sharing of MHC haplotypesamong apparently unrelated patients with congenital adrenal hyperplasiadue to 21-hydroxylase deficiency. Immunogenetics 25: 99-103, 1987.

[76855] 15185.Lee, H.-H.; Chao, H.-T.; Ng, H.-T.; Choo, K.-B.: Direct molecular diagnosis of CYP21 mutations in congenital adrenal hyperplasia. J.Med. Genet. 33: 371-375, 1996.

- [76856] 15186. Levine, L. S.; Dupont, B.; Lorenzen, F.; Pang, S.; Pollack, M.; Oberfield, S.; Kohn, B.; Lerner, A.; Cacciari, E.; Mantero, F.; Cassio, A.; Scaroni, C.; Chiumello, G.; Rondanini, G. F.; Gargantini, L.; Giovannelli, G.; Virdis, R.; Bartolotta, E.; Migliori, C.; Pintor, C.; Tato, L.; Barboni, F.; New, M. I.: Cryptic 21-hydroxylase deficiency in families of patients with classical congenital adrenal hyperplasia. *J. Clin. Endocr. Metab.* 51: 1316–1324, 1980.
- [76857] 15187. Levine, L. S.; Dupont, B.; Lorenzen, F.; Pang, S.; Pollack, M.; Oberfield, S. E.; Kohn, B.; Lerner, A.; Cacciari, E.; Mantero, F.; Cassio, A.; Scaroni, C.; Chiumello, G.; Rondanini, G. F.; Gargantini, L.; Giovannelli, G.; Virdis, R.; Bartolotta, E.; Migliori, C.; Pintor, C.; Tato, L.; Barboni, F.; New, M. I.: Genetic and hormonal characterization of cryptic 21-hydroxylase deficiency. *J. Clin. Endocr. Metab.* 53: 1193–1198, 1981.
- [76858] 15188. Levine, L. S.; Pang, S.; Dupont, B.; Pollack, M.; Lorenzen, F.; New, M. I.: Detection of heterozygote of 21-hydroxylase deficiency. (Letter) *Lancet* I: 603–604, 1980.
- [76859] 15189. Levine, L. S.; Zachmann, M.; New, M. I.; Prader, A.; Pollack, M. S.; O'Neill, G. J.; Yang, S. Y.; Oberfield, S. E.; Dupont, B.: Genetic mapping of the

21-hydroxylase-deficiency gene within the HLA linkage group. *New Eng. J. Med.* 299: 911–915, 1978.

[76860] 15190. Levo, A.; Partanen, J.: Mutation-haplotype analysis of steroid 21-hydroxylase (CYP21) deficiency in Finland. Implications for the population history of defective alleles. *Hum. Genet.* 99: 488–497, 1997.

[76861] 15191. Lewis, V. G.; Money, J.; Epstein, R.: Concordance of verbal and nonverbal ability in the adrenogenital syndrome. *Johns Hopkins Med. J.* 122: 192–195, 1968.

[76862] 15192. Libber, S. M.; Migeon, C. J.; Bias, W. B.: Ascertainment of 21-hydroxylase deficiency in individuals with HLA-B14 haplotype. *J. Clin. Endocr. Metab.* 60: 727–730, 1985.

[76863] 15193. Lo, J. C.; Schwitzgebel, V. M.; Tyrrell, J. B.; Fitzgerald, P. A.; Kaplan, S. L.; Conte, F. A.; Grumbach, M. M.: Normal female infants born of mothers with classic congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *J. Clin. Endocr. Metab.* 84: 930–936, 1999.

[76864] 15194. Lopez-Gutierrez, A. U.; Riba, L.; Ordonez-Sanchez, M. L.; Ramirez-Jimenez, S.; Cerrillo-Hinojosa, M.; Tusie-Luna, M. T.: Uniparental disomy for chromosome 6 results in steroid 21-hydroxylase deficiency: evidence of different genetic mechanisms involved in the production of the dis-

ease. J. Med. Genet. 35: 1014–1019, 1998.

- [76865] 15195. Lorenzen, F.; Pang, S.; New, M. I.; Dupont, B.; Pollack, M.; Chow, D. M.; Levine, L. S.: Hormonal phenotype and HLA-genotype in families of patients with congenital adrenal hyperplasia (21-hydroxylase deficiency). *Pediatr. Res.* 13: 1356–1360, 1979.
- [76866] 15196. Lorenzen, F.; Pang, S.; New, M. I.; Oberfield, S.; Dupont, B.; Chow, D.; Schneider, B.; Levine, L.: Studies of the C-21 and C-19 steroids and HLA genotyping in siblings and parents of patients with congenital adrenal hyperplasia due to 21-hydroxylase deficiency. *J. Clin. Endocr. Metab.* 50: 572–577, 1980.
- [76867] 15197. Lubani, M. M.; Issa, A.-R. A.; Bushnaq, R.; Al-Saleh, Q. A.; Dudin, K. I.; Reavey, P. C.; El-Khalifa, M. Y.; Manandhar, D. S.; Abdul Al, Y. K.; Ismail, E. A.; Teebi, A. S.: Prevalence of congenital adrenal hyperplasia in Kuwait. *Europ. J. Pediatr.* 149: 391–392, 1990.
- [76868] 15198. Matteson, K. J.; Phillips, J. A., III; Miller, W. L.; Chung, B.-C.; Orlando, P. J.; Frisch, H.; Ferrandez, A.; Burr, I. M.: P450XXI (steroid 21-hydroxylase) gene deletions are not found in family studies of congenital adrenal hyperplasia. *Proc. Nat. Acad. Sci.* 84: 5858–5862, 1987.
- [76869] 15199. McAlpine, P. J.: Personal Communication. Winnipeg,



Manitoba,Canada 2/26/1988.

- [76870] 15200.McGuire, L. S.; Omenn, G. S.: Congenital adrenal hyperplasia.I. Family studies of IQ. Behav. Genet. 5: 165–173, 1975.
- [76871] 15201.Merkatz, I. R.; New, M. I.; Peterson, R. E.; Seaman, M. P.: Prenataldiagnosis of adrenogenital syndrome by amniocentesis. J. Pediat. 75:977–982, 1969.
- [76872] 15202.Merke, D. P.; Chrousos, G. P.; Eisenhofer, G.; Weise, M.; Keil,M. F.; Rogol, A. D.; Van Wyk, J. J.; Bornstein, S. R.: Adrenomedullarydysplasia and hypofunction in patients with classic 21–hydroxylasedeficiency. New Eng. J. Med. 343: 1362–1368, 2000.
- [76873] 15203.Merke, D. P.; Keil, M. F.; Jones, J. V.; Fields, J.; Hill, S.;Cutler, G. B., Jr.: Flutamide, testolactone, and reduced hydrocortisonedose maintain normal growth velocity and bone maturation despite elevatedandrogen levels in children with congenital adrenal hyperplasia. J.Clin. Endocr. Metab. 85: 1114–1120, 2000.
- [76874] 15204.Meyer–Bahlburg, H. F. L.: What causes low rates of child–bearingin congenital adrenal hyperplasia? J. Clin. Endocr. Metab. 84: 1844–1847,1999.
- [76875] 15205.Miller, W. L.: Gene conversions, deletions and polymorphismsin congenital adrenal hyperplasia. Am. J. Hum.

Genet. 42: 4–7, 1988.

[76876] 15206. Miller, W. L.: Personal Communication. San Francisco, Calif. 1/3/1996.

[76877] 15207. Bannerman, R. M.; Edwards, J. A.; Kreimer-Birnbaum, M.; McFarland, E.; Russell, E. S.: Hereditary microcytic anaemia in the mouse; studies in iron distribution and metabolism. Brit. J. Haemat. 23: 235–245, 1972.

[76878] 15208. Vincent, J. B.; Herbrick, J.-A.; Gurling, H. M. D.; Bolton, P. F.; Roberts, W.; Scherer, S. W.: Identification of a novel gene on chromosome 7q31 that is interrupted by a translocation breakpoint in an autistic individual. Am. J. Hum. Genet. 67: 510–514, 2000.

[76879] 15209. Barichard, F.; Joulin, V.; Henry, I.; Garel, M.-C.; Valentin, C.; Rosa, R.; Cohen-Solal, M.; Junien, C.: Chromosomal assignment of the human 2,3-bisphosphoglycerate mutase gene (BPGM) to region 7q34–7q22. Hum. Genet. 77: 283–285, 1987.

[76880] 15210. Bowdler, A. J.; Prankerd, T. A. J.: Studies in congenital non-spherocytic haemolytic anaemias with specific enzyme defects. Acta Haemat. 31: 65–78, 1964.

[76881] 15211. Chen, S.-H.; Anderson, J. E.; Giblett, E. R.: 2,3-diphosphoglycerate mutase: its demonstration by electrophoresis and the detection of a genetic variant.

Biochem. Genet. 5: 481–486, 1971.

- [76882] 15212. Galacteros, F.; Rosa, R.; Prehu, M. O.; Najean, Y.; Calvin, M.C.: Deficit en diphosphoglycerate mutase: nouveaux cas associes a une polyglobulie. *Nouv. Rev. Franc. Hemat.* 26: 69–74, 1984.
- [76883] 15213. Joulin, V.; Barichard, F.; Henry, I.; Garel, M. C.; Valentin, C.; Rosa, R.; Cohen–Solal, M.; Junien, C.: Chromosomal assignment of the human 2,3–bisphosphoglycerate mutase gene (BPGM) to region 7q22–7q34. (Abstract) *Cytogenet. Cell Genet.* 46: 635 only, 1987.
- [76884] 15214. Joulin, V.; Garel, M.–C.; Le Boulch, P.; Valentin, C.; Rosa, R.; Rosa, J.; Cohen–Solal, M.: Isolation and characterization of the human 2,3–bisphosphoglycerate mutase gene. *J. Biol. Chem.* 263: 15785–15790, 1988.
- [76885] 15215. Joulin, V.; Peduzzi, J.; Romeo, P.–H.; Rosa, R.; Valentin, C.; Dubart, A.; Lapeyre, B.; Blouquit, Y.; Garel, M.–C.; Goossens, M.; Rosa, J.; Cohen–Solal, M.: Molecular cloning and sequencing of the human erythrocyte 2,3–bisphosphoglycerate mutase cDNA: revised amino acid sequence. *EMBO J.* 5: 2275–2283, 1986.
- [76886] 15216. Labie, D.; Leroux, J.–P.; Najman, A.; Reyrolle, C.: Familial diphosphoglycerate mutase deficiency: influence

on the oxygen affinity curves of hemoglobin. FEBS Lett. 9: 37–40, 1970.

- [76887] 15217. Lemarchandel, V.; Joulin, V.; Valentin, C.; Rosa, R.; Galacteros, F.; Rosa, J.; Cohen-Solal, M.: Compound heterozygosity in a complete erythrocyte bisphosphoglycerate mutase deficiency. Blood 80: 2643–2649, 1992.
- [76888] 15218. Rosa, R.; Audit, I.; Rosa, J.: Diphosphoglycerate mutase and 2,3-diphosphoglycerate phosphatase activities of red cells: comparative electrophoretic study. Biochem. Biophys. Res. Commun. 51: 536–542, 1973.
- [76889] 15219. Rosa, R.; Blouquit, Y.; Calvin, M.-C.; Prome, D.; Prome, J.-C.; Rosa, J.: Isolation, characterization, and structure of a mutant arg-to-cys bisphosphoglycerate mutase: implication of the active site in the mutation. J. Biol. Chem. 264: 7837–7843, 1989.
- [76890] 15220. Rosa, R.; Prehu, M.-O.; Beuzard, Y.; Rosa, J.: The first case of a complete deficiency of diphosphoglycerate mutase in human erythrocytes. J. Clin. Invest. 62: 907–915, 1978.
- [76891] 15221. Sasaki, R.; Ikura, K.; Sugimoto, E.; Chiba, H.: Purification of bisphosphoglyceromutase, 2,3-bisphosphoglycerate phosphatase and phosphoglyceromutase from human erythrocytes. Europ. J. Biochem.

50:581–593, 1975.

- [76892] 15222.Schroter, W.: Kongenitale nichtsphaerocytaere haemolytische Anaemie bei 2,3-Diphosphoglyceratmutasemangel der Erythrocyten im fruehenSaeuglingsalter. Klin. Wschr. 43: 1147–1153, 1965.
- [76893] 15223.Scott, E. M.; Wright, R. C.: An alternate method for demonstrationof bisphosphoglyceromutase (DPGM) on starch gels. Am. J. Hum. Genet. 34:1013–1015, 1982.
- [76894] 15224.Yanagawa, S.; Hitomi, K.; Sasaki, R.; Chiba, H.: Iso-lation andcharacterization of cDNA encoding rabbit reticulocyte 2,3-bisphosphoglyceratesynthase. Gene 44: 185–191, 1986.
- [76895] 15225.Cali, J. J.; Hsieh, C.-L.; Francke, U.; Russell, D. W.: Mutationsin the bile acid biosynthetic enzyme sterol 27-hydroxylase underliecerebrotendinous xanthomatosis. J. Biol. Chem. 266: 7779–7783, 1991.
- [76896] 15226.Cali, J. J.; Russell, D. W.: Characterization of human sterol27-hydroxylase: a mitochondrial cytochrome P-450 that catalyzes multipleoxidation reactions in bile acid biosynthesis. J. Biol. Chem. 266:7774–7778, 1991.
- [76897] 15227.Jaspers, M.; Zhang, Z.; Marynen, P.; Vekemans, S.; Aly, M. S.;Cuppens, H.; Hillicker, C.; Cassiman, J.-J.: Local-

ization of the genes encoding the alpha-2 and alpha-4 subunits of the human VLA-receptor to chromosome 5q23-31 and 2q31-32 respectively. (Abstract) Cytogenet. Cell Genet. 58: 1870 only, 1991.

[76898] 15228. Kaplan, C.; Morel-Kopp, M. C.; Kroll, H.; Kiefel, V.; Schlegel, N.; Chesnel, N.; Mueller-Eckhardt, C.: HPA-5b (Br-a) neonatal alloimmune thrombocytopenia: clinical and immunological analysis of 39 cases. Brit. J. Haemat. 78: 425-429, 1991.

[76899] 15229. Kiefel, V.; Santoso, S.; Katzmann, B.; Mueller-Eckhardt, C.: A new platelet-specific alloantigen Br(a): report of 4 cases with neonatal alloimmune thrombocytopenia. Vox Sang. 54: 101-106, 1988.

[76900] 15230. Kiefel, V.; Shechter, Y.; Atias, D.; Kroll, H.; Santoso, S.; Mueller-Eckhardt, C.: Neonatal alloimmune thrombocytopenia due to anti-Br(b) (HPA-5a): report of three cases in two families. Vox Sang. 60: 244-245, 1991.

[76901] 15231. Kritzik, M.; Savage, B.; Nugent, D. J.; Santoso, S.; Ruggeri, Z. M.; Kunicki, T. J.: Nucleotide polymorphisms in the alpha-2 gene define multiple alleles that are associated with differences in platelet alpha-2/beta-1 density. Blood 92: 2382-2388, 1998.

[76902] 15232. Kunicki, T. J.; Kritzik, M.; Annis, D. S.; Nugent, D. J.:

Hereditary variation in platelet integrin alpha-2-beta-1 density is associated with two silent polymorphisms in the alpha-2 gene coding sequence. *Blood* 89:1939-1943, 1997.

[76903] 15233. Mueller-Eckhardt, C.; Kiefel, V.; Grubert, A.; Kroll, H.; Weisheit, M.; Schmidt, S.; Mueller-Eckhardt, G.; Santos, S.: 348 cases of suspected neonatal alloimmune thrombocytopenia. *Lancet* I: 363-366, 1989.

[76904] 15234. Nieuwenhuis, H. K.; Akkerman, J. W. N.; Houdijk, W. P.; Sixma, J. J.: Human blood platelets showing no response to collagen fail to express surface glycoprotein Ia. *Nature* 318: 470-472, 1985.

[76905] 15235. Santos, S.; Amrhein, J.; Hofmann, H. A.; Sachs, U. J. H.; Walka, M. M.; Kroll, H.; Kiefel, V.: A point mutation thr799met on the alpha-2 integrin leads to the formation of new human platelet alloantigen Sit(a) and affects collagen-induced aggregation. *Blood* 94: 4103-4111, 1999.

[76906] 15236. Santos, S.; Kalb, R.; Walka, M.; Kiefel, V.; Mueller-Eckhardt, C.; Newman, P. J.: The human platelet alloantigens Br(a) and Br(b) are associated with a single amino acid polymorphism on glycoprotein Ia (integrin subunit alpha-2). *J. Clin. Invest.* 92: 2427-2432, 1993.

[76907] 15237. Santos, S.; Kunicki, T. J.; Kroll, H.; Haberbosch,

W.; Gardemann,A.: Association of the platelet glycoprotein Ia C807T gene polymorphism with nonfatal myocardial infarction in younger patients. Blood 93:2449–2453, 1999.

[76908] 15238.Takada, Y.; Hemler, M. E.: The primary structure of the VLA-2/collagen receptor alpha-2 subunit (platelet GPIa): homology to other integrins and the presence of a possible collagen-binding domain. J. Cell Biol. 109:397–407, 1989.

[76909] 15239.von Beckerath, N.; Koch, W.; Mehilli, J.; Bottiger, C.; Schomig,A.; Kastrati, A.: Glycoprotein Ia gene C807T polymorphism and risk for major adverse cardiac events within the first 30 days after coronary artery stenting. Blood 95: 3297–3301, 2000.

[76910] 15240.Fernandez-Ruiz, E.; Pardo-Manuel de Villena, F.; Rubio, M. A.;Corbi, A. L.; Rodriguez de Cordoba, S.; Sanchez-Madrid, F.: Mapping of the human VLA-alpha-4 gene to chromosome 2q31–q32. Europ. J. Immun. 22:587–590, 1992.

[76911] 15241.Roseblatt, M.; Vuillet-Gaugler, M. H.; Leroy, C.; Coulombel, L.: Coexpression of two fibronectin receptors, VLA-4 and VLA-5, by immature human erythroblastic precursor cells. J. Clin. Invest. 87: 6–11,1991.

[76912] 15242.Zhang, Z.; Vekemans, S.; Aly, M. S.; Jaspers, M.;



Marynen, P.;Cassiman, J.-J.: The gene for the alpha-4 subunit of the VLA-4 integrin maps to chromosome 2q31-32. *Blood* 78: 2396-2399, 1991.

- [76913] 15243.Hiesberger, T.; Trommsdorff, M.; Howell, B. W.; Goffinet, A.; Mumby, M. C.; Cooper, J. A.; Herz, J.: Direct binding of reelin to VLDL receptor and apoE receptor 2 induces tyrosine phosphorylation of disabled-1 and modulates tau phosphorylation. *Neuron* 24: 481-489, 1999.
- [76914] 15244.Oka, K.; Tzung, K.-W.; Sullivan, M.; Lindsay, E.; Baldini, A.; Chan, L.: Human very-low-density lipoprotein receptor complementary DNA and deduced amino acid sequence and localization of its gene (VLDLR) to chromosome band 9q24 by fluorescence in situ hybridization. *Genomics* 20:298-300, 1994.
- [76915] 15245.Okuizumi, K.; Onodera, O.; Namba, Y.; Ikeda, K.; Yamamoto, T.; Seki, K.; Ueki, A.; Nanko, S.; Tanaka, H.; Takahashi, H.; Oyanagi, K.; Mizusawa, H.; Kanazawa, I.; Tsuji, S.: Genetic association of the very low density lipoprotein (VLDL) receptor gene with sporadic Alzheimer's disease. *Nature Genet.* 11: 207-209, 1995.
- [76916] 15246.Gafvels, M. E.; Paavola, L. G.; Boyd, C. O.; Nolan, P. M.; Wittmaack, F.; Chawla, A.; Lazar, M. A.; Bucan, M.; Angelin, B.; Strauss, J.F., III: Cloning of a complementary de-

oxyribonucleic acid encoding the murine homolog of the very low density lipoprotein/apolipoprotein-E receptor: expression pattern and assignment of the gene to mouse chromosome 19. *Endocrinology* 135: 387–394, 1994.

[76917] 15247. Sakai, J.; Hoshino, A.; Takahashi, S.; Miura, Y.; Ishii, H.; Suzuki, H.; Kawarabayashi, Y.; Yamamoto, T.: Structure, chromosome location, and expression of the human very low density lipoprotein receptor gene. *J. Biol. Chem.* 269: 2173–2182, 1994.

[76918] 15248. Trommsdorff, M.; Gotthardt, M.; Hiesberger, T.; Shelton, J.; Stockinger, W.; Nimpf, J.; Hammer, R. E.; Richardson, J. A.; Herz, J.: Reeler/Disabled-like disruption of neuronal migration in knockout mice lacking the VLDL-receptor and ApoE receptor 2. *Cell* 97: 689–701, 1999.

[76919] 15249. Huebner, K.; Druck, T.; Croce, C. M.; Thiesen, H. J.: Twenty-seven nonoverlapping zinc finger cDNAs from human T cells map to nine different chromosomes with apparent clustering. *Am. J. Hum. Genet.* 48: 726–740, 1991.

[76920] 15250. Rousseau-Merck, M.-F.; Hillion, J.; Jonveaux, P.; Couillin, P.; Seite, P.; Thiesen, H.-J.; Berger, R.: Chromosomal localization of 9 KOX zinc finger genes: physical linkages suggest clustering of KOX genes on chromosomes 12, 16, and 19. *Hum. Genet.* 92: 583–587, 1993.

- [76921] 15251. Bray, P.; Lichter, P.; Thiesen, H.-J.; Ward, D. C.; Dawid, I. B.: Characterization and mapping of human genes encoding zinc fingerproteins. *Proc. Nat. Acad. Sci.* 88: 9563–9567, 1991.
- [76922] 15252. Lehmann, O. J.; El-ashry, M. F.; Ebenezer, N. D.; Ocaka, L.; Francis, P. J.; Wilkie, S. E.; Patel, R. J.; Ficker, L.; Jordan, T.; Khaw, P. T.; Bhattacharya, S. S.: A novel kerato- can mutation causing autosomalrecessive cornea plana. *Invest. Ophthal. Vis. Sci.* 42: 3118–3122, 2001.
- [76923] 15253. Bermingham, N.; Hernandez, D.; Balfour, A.; Gilmour, F.; Martin, J. E.; Fisher, E. M. C.: Mapping TNNC1, the gene that encodes cardiactroponin I in the human and mouse. *Genomics* 30: 620–622, 1995.
- [76924] 15254. Roher, A.; Lieska, N.; Spitz, W.: The amino acid se- quence of humancardiac troponin–C. *Muscle Nerve* 9: 73–77, 1986.
- [76925] 15255. Romero–Herrera, A. E.; Castillo, O.; Lehmann, H.: Human skeletal muscle proteins: the primary structure of troponin C. *J. Molec. Evol.* 8: 251–270, 1976.
- [76926] 15256. Schreier, T.; Kedes, L.; Gahlmann, R.: Cloning, structural analysis, and expression of the human slow twitch skeletal muscle/cardiac troponinC gene. *J. Biol. Chem.* 265: 21247–21253, 1990.

- [76927] 15257.Song, W.-J.; Van Keuren, M. L.; Drabkin, H. A.; Cypser, J. R.;Gemmill, R. M.; Kurnit, D. M.: Assignment of the human slow twitchskeletal muscle/cardiac troponin C gene (TNNC1) to human chromosome3p21.3–3p14.3 using somatic cell hybrids. *Cytogenet. Cell Genet.* 75:36–37, 1996.
- [76928] 15258.Song, W. J.: Personal Communication. Ann Arbor, Mich. 1/17/1997.
- [76929] 15259.Townsend, P. J.; Yacoub, M. H.; Barton, P. J. R.: Assignment ofthe human cardiac/slow skeletal muscle troponin C gene (TNNC1) betweenD3S3118 and GCT4B10 on the short arm of chromosome 3 by somatic cellhybrid analysis. *Ann. Hum. Genet.* 61: 375–377, 1997.
- [76930] 15260.Haldane, J. B. S.; Sprunt, A. D.; Haldane, N. M.: Reduplicationin mice. *J. Genet.* 5: 133–135, 1915.
- [76931] 15261.Lyon, M. F.; King, T. R.; Gondo, Y.; Gardner, J. M.; Nakatsu,Y.; Eicher, E. M.; Brilliant, M. H.: Genetic and molecular analysisof recessive alleles at the pink-eyed dilution (p) locus of the mouse. *Proc.Nat. Acad. Sci.* 89: 6968–6972, 1992.
- [76932] 15262.Stohl, W.; Kunkel, H. G.: Heterogeneity in expression of theT4 epitope in black individuals. *Scand. J. Immun.* 20: 273–278, 1984.

- [76933] 15263.Tishkoff, S. A.; Dietzsch, E.; Speed, W.; Pakstis, A. J.; Kidd, J. R.; Cheung, K.; Bonne-Tamir, B.; Santachiara-Benerecetti, A. S.; Moral, P.; Krings, M.; Paabo, S.; Watson, E.; Risch, N.; Jenkins, T.; Kidd, K. K.: Global patterns of linkage disequilibrium at the CD4 locus and modern human origins. *Science* 271: 1380–1387, 1996.
- [76934] 15264.van Dongen, J. J. M.; Wolvers-Tettero, I. L. M.; Versnel, M. A.; Westerveld, A.; Geurts van Kessel, A. H. M.: Assignment of the genes coding for the T-cell antigens CD7 (Tp41), CD5 (T1) and CD4 (T4) to human chromosome 17, 11, 12 respectively. (Abstract) *Cytogenet. Cell Genet.* 40: 767, 1985.
- [76935] 15265.Zou, Y.-R.; Sunshine, M.-J.; Taniuchi, I.; Hatam, F.; Killeen, N.; Littman, D. R.: Epigenetic silencing of CD4 in T cells committed to the cytotoxic lineage. *Nature Genet.* 332–336, 2001.
- [76936] 15266.DiLella, A. G.: Chromosomal assignment of the human immunophilin FKBP-12 gene. *Biochem. Biophys. Res. Commun.* 179: 1427–1433, 1991.
- [76937] 15267.DiLella, A. G.; Hawkins, A.; Craig, R. J.; Schreiber, S. L.; Griffin, C. A.: Chromosomal band assignments of the genes encoding human FKBP12 and FKBP13. *Biochem. Biophys. Res. Commun.* 189: 819–823, 1992.

- [76938] 15268. Goebel, M. G.: The peptidyl-prolyl isomerase, FK506-binding protein, is most likely the 12 kd endogenous inhibitor 2 of protein kinase C. (Letter) *Cell* 64: 1051-1052, 1991.
- [76939] 15269. Jin, Y.-J.; Albers, M. W.; Lane, W. S.; Bierer, B. E.; Schreiber, S. L.; Burakoff, S. J.: Molecular cloning of a membrane-associated human FK506- and rapamycin-binding protein, FKBP-13. *Proc. Nat. Acad. Sci.* 88: 6677-6681, 1991.
- [76940] 15270. Maki, N.; Sekiguchi, F.; Nishimaki, J.; Miwa, K.; Hayano, T.; Takahashi, N.; Suzuki, M.: Complementary DNA encoding the human T-cell FK506-binding protein, a peptidylprolyl cis-trans isomerase distinct from cyclophilin. *Proc. Nat. Acad. Sci.* 87: 5440-5443, 1990.
- [76941] 15271. Peattie, D. A.; Hsiao, K.; Benasutti, M.; Lippke, J. A.: Three distinct messenger RNAs can encode the human immunosuppressant-binding protein FKBP12. *Gene* 150: 251-257, 1994.
- [76942] 15272. Shou, W.; Aghdasi, B.; Armstrong, D. L.; Guo, Q.; Bao, S.; Charng, M.-J.; Mathews, L. M.; Schneider, M. D.; Hamilton, S. L.; Matzuk, M. M.: Cardiac defects and altered ryanodine receptor function in mice lacking FKBP12. *Nature* 391: 489-492, 1998.

- [76943] 15273.Standaert, R. F.; Galat, A.; Verdine, G. L.; Schreiber, S. L.:Molecular cloning and overexpression of the human FK506-binding proteinFKBP. *Nature* 346: 671–674, 1990.
- [76944] 15274.Wang, T.; Donahoe, P. K.; Zervos, A. S.: Specific interactionof type I receptors of the TGF-beta family with the immunophilin FKBP-12. *Science* 265:674–676, 1994.
- [76945] 15275.Jin, Y. J.; Albers, M. W.; Lane, W. S.; Bierer, B. E.; Schreiber,S. L.; Burakoff, S. J.: Molecular cloning of a membrane-associatedhuman FK506- and rapamycin-binding protein, FKBP-13. *Proc. Nat. Acad.Sci.* 88: 6677–6681, 1991.
- [76946] 15276.Howard, H. C.; Mount, D. B.; Rochefort, D.; Byun, N.; Dupre, N.;Lu, J.; Fan, X.; Song, L.; Riviere, J.-B.; Prevost, C.; Horst, J.;Simonati, A.; and 12 others: The K-Cl cotransporter KCC3 is mutantin a severe peripheral neuropathy associated with agenesis of thecorpus callosum. *Nature Genet.* 7Oct, 2002. Note: Advance ElectronicPublication.
- [76947] 15277.Haire, R. N.; Litman, G. W.: The murine form of TXK, a novel TECKinase expressed in thymus maps to chromosome 5. *Mammalian Genome* 6:476–480, 1995.
- [76948] 15278.Haire, R. N.; Ohta, Y.; Lewis, J. E.; Fu, S. M.; Kroisel, P.; Litman,G. W.: TXK, a novel human tyrosine kinase ex-

pressed in T cells shares sequence identity with Tec family kinases and maps to 4p12. *Hum.Molec. Genet.* 3: 897–901, 1994.

- [76949] 15279. Schneider, H.; Schwartzberg, P. L.; Rudd, C. E.: Resting lymphocyte kinase (Rlk/Txk) phosphorylates the YVKM motif and regulates PI 3-kinase binding to T-cell antigen CTLA-4. *Biochem. Biophys. Res. Commun.* 252:14–19, 1998.
- [76950] 15280. Takeba, Y.; Nagafuchi, H.; Takeno, M.; Kashiwakura, J.; Suzuki, N.: Txk, a member of nonreceptor tyrosine kinase of Tec family, acts as a Th1 cell-specific transcription factor and regulates IFN-gamma gene transcription. *J. Immun.* 168: 2365–2370, 2002.
- [76951] 15281. Al-Maghteh, M.; Vithana, E.; Tarttelin, E.; Jay, M.; Evans, K.; Moore, T.; Bhattacharya, S.; Inglehearn, C. F.: Evidence for a major retinitis pigmentosa locus on 19q13.4 (RP11), and association with a unique bimodal expressivity phenotype. *Am. J. Hum. Genet.* 59:864–871, 1996.
- [76952] 15282. Gasdaska, J. R.; Gasdaska, P. Y.; Gallegos, A.; Powis, G.: Human thioredoxin reductase gene localization to chromosomal position 12q23–q24.1 and mRNA distribution in human tissue. *Genomics* 37: 257–259, 1996.
- [76953] 15283. Gasdaska, P. Y.; Gasdaska, J. R.; Cochran, S.; Powis,



G.: Cloning and sequencing of a human thioredoxin reductase. FEBS Lett. 373:5–9, 1995.

[76954] 15284. Svaren, J.; Apel, E. D.; Simburger, K. S.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. A.; Milbrandt, J.: The Nab2 and Stat6 genes share a common transcription termination region. Genomics 41: 33–39, 1997.

[76955] 15285. Weichenrieder, O.; Wild, K.; Strub, K.; Cusack, S.: Structure and assembly of the Alu domain of the mammalian signal recognition particle. Nature 408: 167–173, 2000.

[76956] 15286. Nichols, R. C.; Blinder, J.; Pai, S. I.; Ge, Q.; Targoff, I. N.; Plotz, P. H.; Liu, P.: Assignment of two human autoantigen genes: isoleucyl-tRNA synthetase locates to 9q21 and lysyl-tRNA synthetase locates to 16q23–q24. Genomics 36: 210–213, 1996.

[76957] 15287. Nichols, R. C.; Raben, N.; Boerkoel, C. F.; Plotz, P. H.: Human isoleucyl-tRNA synthetase: sequence of the cDNA, alternative mRNA splicing, and the characteristics of an unusually long C-terminal extension. Gene 155: 299–304, 1995.

[76958] 15288. Dejgaard, K.; Leffers, H.; Rasmussen, H. H.; Madsen, P.; Kruse, T. A.; Gesser, B.; Nielsen, H.; Celis, J. E.: Identification, molecular cloning, expression and chromo-

some mapping of a family of transformation upregulated hnRNP-K proteins derived by alternative splicing. *J. Molec. Biol.* 236: 33–48, 1994.

[76959] 15289. Tommerup, N.; Leffers, H.: Assignment of human KH-box-containing genes by in situ hybridization: HN-RNPK maps to 9q21.32–q21.33, PCBP1 to 2p12–p13, and PCBP2 to 12q13.12–q13.13, distal to FRA12A. *Genomics* 32:297–298, 1996.

[76960] 15290. Agarwal, A. K.; Monder, C.; Eckstein, B.; White, P. C.: Cloning and expression of rat cDNA encoding corticosteroid 11- $\beta$ -dehydrogenase. *J. Biol. Chem.* 264: 18939–18943, 1989.

[76961] 15291. Agarwal, A. K.; Rogerson, F. M.; Mune, T.; White, P. C.: Gene structure and chromosomal localization of the human HSD11K gene encoding the kidney (type 2) isozyme of 11- $\beta$ -hydroxysteroid dehydrogenase. *Genomics* 29:195–199, 1995.

[76962] 15292. Lakshmi, V.; Monder, C.: Purification and characterization of the corticosteroid 11- $\beta$ -dehydrogenase component of the rat liver 11- $\beta$ -hydroxysteroid dehydrogenase complex. *Endocrinology* 123:2390–2398, 1988.

[76963] 15293. Kotelevtsev, Y.; Holmes, M. C.; Burchell, A.; Hous-

ton, P. M.; Schmoll, D.; Jamieson, P.; Best, R.; Brown, R.; Edwards, C. R. W.; Seckl, J. R.; Mullins, J. J.:

11-beta-hydroxysteroid dehydrogenase type 1 knockout-mice show attenuated glucocorticoid-inducible responses and resist hyperglycemia on obesity or stress. *Proc. Nat. Acad. Sci.* 94: 14924-14929, 1997.

[76964] 15294. Masuzaki, H.; Paterson, J.; Shinyama, H.; Morton, N. M.; Mullins, J. J.; Seckl, J. R.; Flier, J. S.: A transgenic model of visceral obesity and the metabolic syndrome. *Science* 294: 2166-2170, 2001.

[76965] 15295. Morton, N. M.; Holmes, M. C.; Fievet, C.; Staels, B.; Tailleux, A.; Mullins, J. J.; Seckl, J. R.: Improved lipid and lipoprotein profile, hepatic insulin sensitivity, and glucose tolerance in 11-beta-hydroxysteroid dehydrogenase type 1 null mice. *J. Biol. Chem.* 276: 41293-41300, 2001.

[76966] 15296. Ricketts, M. L.; Verhaeg, J. M.; Bujalska, I.; Howie, A. J.; Rainey, W. E.; Stewart, P. M.: Immunohistochemical localization of type 11-beta-hydroxysteroid dehydrogenase in human tissues. *J. Clin. Endocr. Metab.* 83: 1325-1335, 1998.

[76967] 15297. Alessi, D. R.; Smythe, C.; Keyse, S. M.: The human CL100 gene encodes a tyr/thr-protein phosphatase which potently and specifically inactivates MAP kinase and sup-

presses its activation by oncogenic ras in *Xenopus* oocyte extracts. *Oncogene* 8: 2015–2020, 1993.

- [76968] 15298.Brondello, J.-M.; Pouyssegur, J.; McKenzie, F. R.: Reduced MAPkinase phosphatase-1 degradation after p42/p44(MAPK)-dependent phosphorylation. *Science* 286:2514–2517, 1999.
- [76969] 15299.Emslie, E. A.; Jones, T. A.; Sheer, D.; Keyse, S. M.: The CL100 gene, which encodes a dual specificity (tyr/thr) MAP kinase phosphatase, is highly conserved and maps to human chromosome 5q34. *Hum. Genet.* 93:513–516, 1994.
- [76970] 15300.Keyse, S. M.; Emslie, E. A.: Oxidative stress and heat shock induce a human gene encoding a protein-tyrosine phosphatase. *Nature* 359:644–647, 1992.
- [76971] 15301.Martell, K. J.; Kwak, S.; Hakes, D. J.; Dixon, J. E.; Trent, J.M.: Chromosomal localization of four human VH1-like protein-tyrosine phosphatases. *Genomics* 22: 462–464, 1994.
- [76972] 15302.Christ, R.; Chen, H.; Morris, M. A.; Antonarakis, S. E.: Mapping of the human transcription factor GABPA (E4TF1-60) gene to chromosome 21. *Genomics* 28: 119–122, 1995.
- [76973] 15303.Gugneja, S.; Virbasius, J. V.; Scarpulla, R. C.: Four

structurally distinct, non-DNA-binding subunits of human nuclear respiratory factor 2 share a conserved transcriptional activation domain. *Molec. Cell Biol.* 15: 102–111, 1995.

[76974] 15304. Guo, A.; Nie, F.; Wong-Riley, M.: Human nuclear respiratory factor 2- $\alpha$  subunit cDNA: isolation, subcloning, sequencing, and in situ hybridization of transcripts in normal and monocularly deprived macaque visual system. *J. Comp. Neurol.* 417: 221–232, 2000.

[76975] 15305. Sawada, J.; Goto, M.; Watanabe, H.; Handa, H.; Yoshida, M. C.: Regional mapping of two subunits of transcription factor E4TF1 to human chromosome. *Jpn. J. Cancer Res.* 86: 10–12, 1995.

[76976] 15306. Virbasius, J. V.; Scarpulla, R. C.: Activation of the human mitochondrial transcription factor A gene by nuclear respiratory factors: a potential regulatory link between nuclear and mitochondrial gene expression in organelle biogenesis. *Proc. Nat. Acad. Sci.* 91: 1309–1313, 1994.

[76977] 15307. Watanabe, H.; Sawada, J.; Yano, K.-I.; Yamaguchi, K.; Goto, M.; Handa, H.: cDNA cloning of transcription factor E4TF1 subunits with Ets and notch motifs. *Molec. Cell Biol.* 13: 1385–1391, 1993.

- [76978] 15308.Becker, T.; Gerke, V.; Kube, E.; Weber, K.: S100P, a novel  $\text{Ca}(2+)$ -bindingprotein from human placenta: cDNA cloning, recombinant protein expressionand  $\text{Ca}(2+)$  binding properties. *Europ. J. Biochem.* 207: 541–547,1992.
- [76979] 15309.Dixon, J.; Loftus, S. K.; Gladwin, A. J.; Scambler, P. J.; Wasmuth,J. J.; Dixon, M. J.: Cloning of the human heparan sulfate–N–deacetylase/N–sulfotransferasegene from the Treacher Collins syndrome candidate region at 5q32–q33.1. *Genomics* 26:239–244, 1995.
- [76980] 15310.Gladwin, A. J.; Dixon, J.; Loftus, S. K.; Wasmuth, J. J.; Dixon,M. J.: Genomic organization of the human heparan sulfate–N–deacetylase/N–sulfotransferasegene: exclusion from a causative role in the pathogenesis of TreacherCollins syndrome. *Genomics* 32: 471–473, 1996.
- [76981] 15311.Hashimoto, Y.; Orellana, A.; Gil, G.; Hirschberg, C. B.: Molecularcloning and expression of rat liver N–heparan sulfate sulfotransferase. *J.Biol. Chem.* 267: 15744–15750, 1995.
- [76982] 15312.Chen, H.; Antonarakis, S. E.: Localisation of a human homologueof the *Drosophila* *mn*b and rat *Dyrk* genes to chromosome 21q22.2. *Hum.Genet.* 99: 262–265, 1997.
- [76983] 15313.Patil, N.; Cox, D. R.; Bhat, D.; Faham, M.; Myers, R. M.; Peterson,A. S.: A potassium channel mutation in

weaver mice implicates membrane excitability in granule cell differentiation. *Nature Genet.* 11:126–129, 1995.

[76984] 15314. Shindoh, N.; Kudoh, J.; Maeda, H.; Yamaki, A.; Minoshima, S.; Shimizu, Y.; Shimizu, N.: Cloning of a human homolog of the *Drosophila* minibrain/rat Dyrk gene from 'the Down syndrome critical region' of chromosome 21.

*Biochem. Biophys. Res. Commun.* 225: 92–99, 1996.

[76985] 15315. Smith, D. J.; Stevens, M. E.; Sudanagunta, S. P.; Bronson, R. T.; Makhinson, M.; Watabe, A. M.; O'Dell, T. J.; Fung, J.; Weier, H.-U. G.; Cheng, J.-F.; Rubin, E. M.: Functional screening of 2 Mb of human chromosome 21q22.2 in transgenic mice implicates minibrain in learning defects associated with Down syndrome. *Nature Genet.* 16: 28–36, 1997.

[76986] 15316. Song, W.-J.; Chung, S.-H.; Kurnit, D. M.: The murine Dyrk protein maps to chromosome 16, localizes to the nucleus, and can form multimers. *Biochem. Biophys. Res. Commun.* 231: 640–644, 1997.

[76987] 15317. Song, W.-J.; Sternberg, L. R.; Kasten-Sportes, C.; Van Keuren, M. L.; Chung, S.-H.; Slack, A. C.; Miller, D. E.; Glover, T. W.; Chiang, P.-W.; Lou, L.; Kurnit, D. M.: Isolation of human and murine homologues of the *Drosophila* minibrain gene: human homologue maps to 21q22.2 in the

Down syndrome 'critical region.' Genomics 38: 331–339, 1996.

[76988] 15318. Tejedor, F.; Zhu, X. R.; Kaltenbach, E.; Ackermann, A.; Baumann, A.; Canal, I.; Heisenberg, M.; Fischbach, K. F.; Pongs, O.: Minibrain: a new protein kinase family involved in postembryonic neurogenesis in *Drosophila*. Neuron 14: 287–301, 1995.

[76989] 15319. Niethammer, D.; Dieterle, U.; Kleihauer, E.; Wildfeuer, A.; Haferkamp, O.; Hitzig, W. H.: An inherited defect in granulocyte function: impaired chemotaxis, phagocytosis and intracellular killing of microorganisms. Helv. Paediat. Acta 30: 537–541, 1976.

[76990] 15320. Ostermann, G.; Weber, K. S. C.; Zerneck, A.; Schroder, A.; Weber, C.: JAM-1 is a ligand of the beta-2 integrin LFA-1 involved in transendothelial migration of leukocytes. Nature Immun. 3: 151–158, 2002.

[76991] 15321. Rivera-Matos, I. R.; Rakita, R. M.; Mariscalco, M. M.; Elder, F. F. B.; Dreyer, S. A.; Cleary, T. G.: Leukocyte adhesion deficiency mimicking Hirschsprung disease. J. Pediat. 127: 755–757, 1995.

[76992] 15322. Rosmarin, A. G.; Caprio, D.; Levy, R.; Simkevich, C.: CD18 (beta-2 leukocyte integrin) promoter requires PU.1 transcription factor for myeloid activity. Proc. Nat. Acad.



Sci. 92: 801–805, 1995.

[76993] 15323.Ross, G. D.: Clinical and laboratory features of patients with an inherited deficiency of neutrophil membrane complement receptor type 3 (CR3) and the related membrane antigens LFA-1 and p150,95. *J.Clin. Immun.* 6: 107–113, 1986.

[76994] 15324.Shuster, D. E.; Kehrli, M. E., Jr.; Ackermann, M. R.; Gilbert, R. O.: Identification and prevalence of a genetic defect that causes leukocyte adhesion deficiency in Holstein cattle. *Proc. Nat. Acad.Sci.* 89: 9225–9229, 1992.

[76995] 15325.Sligh, J. E., Jr.; Anderson, D. C.; Beaudet, A. L.: A mutation in the initiation codon of the CD18 gene in a patient with the moderate phenotype of leukocyte adhesion deficiency. (Abstract) *Am. J. Hum.Genet.* 45 (suppl.): A219, 1989.

[76996] 15326.Solomon, E.; Palmer, R. W.; Hing, S.; Law, S. K. A.: Regional localization of CD18, the beta-subunit of the cell surface adhesion molecule LFA-1, on human chromosome 21 by in situ hybridization. *Ann.Hum. Genet.* 52: 123–128, 1988.

[76997] 15327.Springer, T. A.; Miller, L. J.; Anderson, D. C.: p150,95, the third member of the Mac-1, LFA-1 human leukocyte adhesion glycoprotein family. *J. Immun.* 136:

240–245, 1986.

- [76998] 15328.Springer, T. A.; Thompson, W. S.; Miller, L. J.; Schmalstieg,F. C.; Anderson, D. C.: Inherited deficiency of the Mac-1, LFA-1,p150,95 glycoprotein family and its molecular basis. J. Exp. Med. 160:1901–1918, 1984.
- [76999] 15329.Ross, G. D.; Thompson, R. A.; Walport, M. J.; Springer, T. A.;Watson, J. V.; Ward, R. H. R.; Lida, J.; Newman, S. L.; Harrison,R. A.; Lachmann, P. J.: Characterization of patients with an increasedsusceptibility to bacterial infections and a genetic deficiency ofleukocyte membrane complement receptor type three (CR3) and the related-membrane antigen LFA-1. Blood 66: 882–890, 1985.
- [77000] 15330.Suomalainen, H. A.; Gahmberg, C. G.; Patarroyo, M.; Beatty, P.G.; Schroder, J.: Genetic assignment of GP90, leukocyte adhesionglycoprotein to human chromosome 21. Somat. Cell Molec. Genet. 12:297–302, 1986.
- [77001] 15331.Suomalainen, H. A.; Gahmberg, C. G.; Patarroyo, M.; Schroder,J.: GP90 (Leu-CAM antigen) is coded for by genes on chromosome 21.(Abstract) Cytogenet. Cell Genet. 40: 755, 1985.
- [77002] 15332.Taylor, G. M.; Williams, A.; D'Souza, S. W.; Ferguson, W. D.;Donnai, D.; Fennell, J.; Harris, R.: The expression of CD18 is increasedon trisomy 21 (Down syndrome)

lymphoblastoid cells. Clin. Exp. Immun. 71:324–328, 1988.

[77003] 15333.Todd, R. F., III; Freyer, D. R.: The CD11/CD18 leukocyte glycoprotein deficiency. Hemat. Oncol. Clin. North Am. 2: 13–31, 1988.

[77004] 15334.van der Meer, J. W. M.; van Zwet, T. L.; van Furth, R.: New familial defect in microbicidal function of polymorphonuclear leucocytes. Lancet II:630–632, 1975.

[77005] 15335.Vazquez–Torres, A.; Jones–Carson, J.; Baumler, A. J.; Falkow, S.; Valdivia, R.; Brown, W.; Le, M.; Berggren, R.; Parks, W. T.; Fang, F. C.: Extraintestinal dissemination of Salmonella by CD18–expressing phagocytes. Nature 401: 804–808, 1999.

[77006] 15336.Vedder, N. B.; Winn, R. K.; Rice, C. L.; Chi, E. Y.; Arfors, K.–E.; Harlan, J. M.: A monoclonal antibody to the adherence–promoting leukocyte glycoprotein, CD18, reduces organ injury and improves survival from hemorrhagic shock and resuscitation in rabbits. J. Clin. Invest. 81:939–944, 1988.

[77007] 15337.Wardlaw, A. J.; Hibbs, M. L.; Stacker, S. A.; Springer, T. A.: Distinct mutations in two patients with leukocyte adhesion deficiency and their functional correlates. J. Exp. Med. 172: 335–345, 1990.

- [77008] 15338. Weening, R. S.; Roos, D.; Weemaes, C. M. R.; Homan-Muller, J. W. T.; van Schaik, M. L. J.: Defective initiation of the metabolic stimulation in phagocytizing granulocytes: a new congenital defect. *J. Lab. Clin. Med.* 88: 757-768, 1976.
- [77009] 15339. Weitzman, J. B.; Wells, C. E.; Wright, A. H.; Clark, P. A.; Law, S. K. A.: The gene organisation of the human beta-2 integrin subunit (CD18). *FEBS Lett.* 294: 97-103, 1991.
- [77010] 15340. Wilson, J. M.; Ping, A. J.; Krauss, J. C.; Mayo-Bond, L.; Rogers, C. E.; Anderson, D. C.; Todd, R. F., III: Correction of CD18-deficient lymphocytes by retrovirus-mediated gene transfer. *Science* 248: 1413-1416, 1990.
- [77011] 15341. Wilson, R. W.; Ballantyne, C. M.; Smith, C. W.; Montgomery, C.; Bradley, A.; O'Brien, W. E.; Beaudet, A. L.: Gene targeting yields a CD18-mutant mouse for study of inflammation. *J. Immun.* 151: 1571-1578, 1993.
- [77012] 15342. Eichwald, E. J.; Silmsker, C. R.: *Skin. Transplant. Bull.* 2: 148-149, 1955.
- [77013] 15343. Gasser, D. L.; Silvers, W. K.: Genetics and immunology of sex-linked antigens. *Adv. Immun.* 15: 215-247, 1972.
- [77014] 15344. Kent-First, M. G.; Maffitt, M.; Muallem, A.; Brisco,

P.; Shultz, J.; Ekenberg, S.; Agulnik, A. I.; Agulnik, I.; Shramm, D.; Bavister, B.; Abdul-Mawgood, A.; VandeBerg, J.: Gene sequence and evolutionary conservation of human SMCY. (Letter) *Nature Genet.* 14: 128–129, 1996.

[77015] 15345. Scott, D. M.; Ehrmann, I. E.; Ellis, P. S.; Bishop, C. E.; Agulnik, A. I.; Simpson, E.; Mitchell, M. J.: Identification of a mouse male-specific transplantation antigen, H-Y. *Nature* 376: 695–698, 1995.

[77016] 15346. Wachtel, S. S.; Koo, G. C.; Breg, W. R.; Elias, S.; Boyse, E. A.; Miller, O. J.: Expression of H-Y antigen in human males with two Y chromosomes. *New Eng. J. Med.* 293: 1070–1072, 1975.

[77017] 15347. Wachtel, S. S.; Koo, G. C.; Zuckerman, E. E.; Hammerling, U.; Scheid, M. P.; Boyse, E. A.: Serological cross-reactivity between H-Y (male) antigens of mouse and man. *Proc. Nat. Acad. Sci.* 71: 1215–1218, 1974.

[77018] 15348. Wang, W.; Meadows, L. R.; den Haan, J. M. M.; Sherman, N. E.; Chen, Y.; Blokland, E.; Shabanowitz, J.; Agulnik, A. I.; Hendrickson, R. C.; Bishop, C. E.; Hunt, D. F.; Goulmy, E.; Engelhard, V. H.: Human H-Y: a male-specific histocompatibility antigen derived from the SMCY protein. *Science* 269: 1588–1590, 1995.

[77019] 15349. Redmond, T. M.; Yu, S.; Lee, E.; Bok, D.; Hamasaki,

D.; Chen,N.; Goletz, P.; Ma, J.-X.; Crouch, R. K.; Pfeifer, K.: Rpe65 is necessaryfor production of 11-cis-vitamin A in the retinal visual cycle. NatureGenet. 20: 344-351, 1998.

[77020] 15350.Seeliger, M. W.; Grimm, C.; Stahlberg, F.; Friedburg, C.; Jaissle,G.; Zrenner, E.; Guo, H.; Reme, C. E.; Humphries, P.; Hofmann, F.;Biel, M.; Fariss, R. N.; Redmond, T. M.; Wenzel, A.: New views onRPE65 deficiency: the rod system is the source of vision in a mousemodel of Leber congenital amaurosis. Nature Genet. 29: 70-74, 2001.

[77021] 15351.Thompson, D. A.; Gyurus, P.; Fleischer, L.; Bingham, E. L.; McHenry,C. L.; Apfelstedt-Sylla, E.; Zrenner, E.; Lorenz, B.; Richards, J.E.; Jacobson, S. G.; Sieving, P. A.; Gal, A.: Genetics and phenotypesof RPE65 mutations in inherited retinal degeneration. Invest. Ophthal.Vis. Sci. 41: 4293-4299, 2000.

[77022] 15352.Thompson, D. A.; McHenry, C. L.; Li, Y.; Richards, J. E.; Othman,M. I.; Schwinger, E.; Vollrath, D.; Jacobson, S. G.; Gal, A.: Retinaldystrophy due to paternal isodisomy for chromosome 1 or chromosome2, with homoallelism for mutations in RPE65 or MERTK, respectively. Am.J. Hum. Genet. 70: 224-229, 2002.

[77023] 15353.Wrigstad, A.; Narfstrom, K.; Nilsson, S. E.: Slowly

progressive changes of the retina and retinal pigment epithelium in Briard dogs with hereditary retinal dystrophy: a morphological study. *Doc. Ophthalmol.* 87:337–354, 1994.

- [77024] 15354. Bogenmann, E.; Lochrie, M. A.; Simon, M. I.: Cone cell-specific genes expressed in retinoblastoma. *Science* 240: 76–78, 1988.
- [77025] 15355. Bonaiti-Pellie, C.; Briard-Guillemot, M. L.: Segregation analysis in hereditary retinoblastoma. *Hum. Genet.* 57: 411–419, 1981.
- [77026] 15356. Harbitz, I.; Chowdhary, B.; Thomsen, P. D.; Davies, W.; Kaufmann, U.; Kran, S.; Gustavsson, I.; Christensen, K.; Hauge, J. G.: Assignment of the porcine calcium release channel gene, a candidate for the malignant hyperthermia locus, to the 6p11–q21 segment of chromosome 6. *Genomics* 8:243–248, 1990.
- [77027] 15357. Hunter, A. G. W.; Thompson, M. W.: Acromesomelic dwarfism: description of a patient and comparison with previously reported cases. *Hum. Genet.* 34: 107–113, 1976.
- [77028] 15358. Harbitz, I.; Kristensen, T.; Bosnes, M.; Kran, S.; Davies, W.: DNA sequence of the skeletal muscle calcium release channel cDNA and verification of the arg615-to-cys615 mutation, associated with porcine ma-

lignant hyperthermia, in Norwegian Landrace pigs. *AnimalGenet.* 23: 395–402, 1992.

- [77029] 15359. Keating, K. E.; Quane, K. A.; Manning, B. M.; Lehane, M.; Hartung, E.; Censier, K.; Urwyler, A.; Klausnitzer, M.; Muller, C. R.; Heffron, J. J. A.; McCarthy, T. V.: Detection of a novel RYR1 mutation in four malignant hyperthermia pedigrees. *Hum. Molec. Genet.* 3: 1855–1858, 1994.
- [77030] 15360. Lynch, P. J.; Tong, J.; Lehane, M.; Mallet, A.; Giblin, L.; Heffron, J. J. A.; Vaughan, P.; Zafra, G.; MacLennan, D. H.; McCarthy, T. V.: A mutation in the transmembrane/luminal domain of the ryanodine receptor is associated with abnormal  $\text{Ca}^{2+}$  release channel function and severe central core disease. *Proc. Nat. Acad. Sci.* 96: 4164–4169, 1999.
- [77031] 15361. MacKenzie, A. E.; Korneluk, R. G.; Duff, C.; Worton, R. G.; MacLennan, D. H.: Location of the human ryanodine receptor gene with respect to an established human chromosome 19 linkage group. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A149, 1989.
- [77032] 15362. MacKenzie, A. E.; Korneluk, R. G.; Zorzato, F.; Fujii, J.; Phillips, M.; Iles, D.; Wieringa, B.; Leblond, S.; Bailly, J.; Willard, H. F.; Duff, C.; Worton, R. G.; MacLennan, D. H.:



The human ryanodine receptorgene: its mapping to 19q13.1, placement in a chromosome 19 linkagegroup, and exclusion as the gene causing myotonic dystrophy. Am.J. Hum. Genet. 46: 1082–1089, 1990.

[77033] 15363.MacLennan, D. H.: Personal Communication. Toronto, Ontario,Canada 7/5/1991.

[77034] 15364.Manning, B. M.; Quane, K. A.; Lynch, P. J.; Urwyler, A.; Tegazzin,V.; Krivosic–Horber, R.; Censier, K.; Comi, G.; Adnet, P.; Wolz, W.;Lunardi, J.; Muller, C. R.; McCarthy, T. V.: Novel mutations at aCpG dinucleotide in the ryanodine receptor in malignant hyperthermia. Hum.Mutat. 11: 45–50, 1998.

[77035] 15365.Mattei, M. G.; Giannini, G.; Moscatelli, F.; Sorrentino, V.:Chromosomal localization of murine ryanodine receptor genes RYR1,RYR2, and RYR3 by in situ hybridization. Genomics 22: 202–204, 1994.

[77036] 15366.McCarthy, T. V.; Quane, K. A.; Lynch, P. J.: Ryanodine receptormutations in malignant hyperthermia and central core disease. Hum.Mutat. 15: 410–417, 2000.

[77037] 15367.Monnier, N.; Romero, N. B.; Lerale, J.; Landrieu, P.; Nivoche,Y.; Fardeau, M.; Lunardi, J.: Familial and sporadic forms of centralcore disease are associated with mutations in the C–terminal domainof the skeletal muscle

ryanodine receptor. Hum. Molec. Genet. 10:2581–2592, 2001.

- [77038] 15368. MacLennan, D. H.; Zorzato, F.; Fujii, J.; Otsu, K.; Phillips, M.; Lai, F. A.; Meissner, G.; Green, N. M.; Willard, H. F.; Britt, B. A.; Worton, R. G.; Korneluk, R. G.: Cloning and localization of the human calcium release channel (ryanodine receptor) gene to the proximal long arm (cen-q13.2) of human chromosome 19. (Abstract) Am. J. Hum. Genet. 45 (suppl.): A205, 1989.
- [77039] 15369. Manning, B. M.; Quane, K. A.; Ording, H.; Urwyler, A.; Tegazzin, V.; Lehane, M.; O'Halloran, J.; Hartung, E.; Giblin, L. M.; Lynch, P. J.; Vaughan, P.; Censier, K.; Bendixen, D.; Comi, G.; Heytens, L.; Monsieus, K.; Fagerlund, T.; Wolz, W.; Heffron, J. J. A.; Muller, C. R.; McCarthy, T. V.: Identification of novel mutations in the ryanodine-receptor gene (RYR1) in malignant hyperthermia: genotype-phenotype correlation. Am. J. Hum. Genet. 62: 599–609, 1998.
- [77040] 15370. Monnier, N.; Romero, N. B.; Lerale, J.; Nivoche, Y.; Qi, D.; MacLennan, D. H.; Fardeau, M.; Lunardi, J.: An autosomal dominant congenital myopathy with cores and rods is associated with a neomutation in the RYR1 gene encoding the skeletal muscle ryanodine receptor. Hum.

Molec.Genet. 9: 2599–2608, 2000.

- [77041] 15371.Monsieurs, K. G.; Van Broekhoven, C.; Martin, J.-J.; Van Hoof,V. O.; Heytens, L.: Gly341arg mutation indicating malignant hyperthermiasusceptibility: specific cause of chronically elevated serum creatinekinase activity. J. Neurol. Sci. 154: 62–65, 1998.
- [77042] 15372.O'Brien, R. O.; Taske, N. L.; Hansbro, P. M.; Matthaei, K. I.;Hogan, S. P.; Denborough, M. A.; Foster, P. S.: Exclusion of defects in the skeletal muscle specific regions of the DHPR alpha-1 subunit as frequent causes of malignant hyperthermia. J. Med. Genet. 32:913–914, 1995.
- [77043] 15373.Otsu, K.; Khanna, V. K.; Archibald, A. L.; MacLennan, D. H.:Cosegregation of porcine malignant hyperthermia and a probable causal mutation in the skeletal muscle ryanodine receptor gene in backcrossfamilies. Genomics 11: 744–750, 1991.
- [77044] 15374.Otsu, K.; Nishida, K.; Kimura, Y.; Kuzuya, T.; Hori, M.; Kamada,T.; Tada, M.: The point mutation arg615-to-cys in the Ca(2+) releasechannel of skeletal sarcoplasmic reticulum is responsible for hypersensitivityto caffeine and halothane in malignant hyperthermia. J. Biol. Chem. 269:9413–9415, 1994.

- [77045] 15375.Otsu, K.; Phillips, M. S.; Khanna, V. K.; de Leon, S.; MacLennan,D. H.: Refinement of diagnostic assays for a probable causal mutationfor porcine and human malignant hyperthermia. *Genomics* 13: 835–837,1992.
- [77046] 15376.Phillips, M. S.; Fujii, J.; Khanna, V. K.; DeLeon, S.; Yokobata,K.; De Jong, P. J.; MacLennan, D. H.: The structural organizationof the human skeletal muscle ryanodine receptor (RYR1) gene. *Genomics* 34:24–41, 1996.
- [77047] 15377.Quane, K. A.; Healy, J. M. S.; Keating, K. E.; Manning, B. M.;Couch, F. J.; Palmucci, L. M.; Doriguzzi, C.; Fagerlund, T. H.; Berg,K.; Ording, H.; Bendixen, D.; Mortier, W.; Linz, U.; Muller, C. R.;McCarthy, T. V.: Mutations in the ryanodine receptor gene in centralcore disease and malignant hyperthermia. *Nature Genet.* 5: 51–55,1993.
- [77048] 15378.Richter, M.; Schleithoff, L.; Deufel, T.; Lehmann–Horn, F.; Herrmann–Frank,A.: Functional characterization of a distinct ryanodine receptormutation in human malignant hyperthermia–susceptible muscle. *J. Biol.Chem.* 272: 5256–5260, 1997.
- [77049] 15379.Robinson, R. L.; Brooks, C.; Brown, S. L.; Ellis, F. R.; Halsall,P. J.; Quinnell, R. J.; Shaw, M.–A.; Hopkins, P. M.: RYR1 mutationscausing central core disease are associ–

ated with more severe malignant hyperthermia in vitro contracture test phenotypes. Hum. Mutat. 20:88–97, 2002.

- [77050] 15380. Sambuughin, N.; McWilliams, S.; de Bantel, A.; Sivakumar, K.; Nelson, T. E.: Single-amino-acid deletion in the RYR1 gene, associated with malignant hyperthermia susceptibility and unusual contraction phenotype. Am. J. Hum. Genet. 69: 204–208, 2001.
- [77051] 15381. Oesterling, J. E.; Chan, D. W.; Epstein, J. I.; Kimball, A. W., Jr.; Bruzek, D. J.; Rock, R. C.; Brendler, C. B.; Walsh, P. C.: Prostate specific antigen in the preoperative and postoperative evaluation of localized prostatic cancer treated with radical prostatectomy. J. Urol. 139: 766–772, 1988.
- [77052] 15382. Riegman, P. H. J.; Vlietstra, R. J.; van der Korput, J. A. G. M.; Brinkmann, A. O.; Trapman, J.: The promoter of the prostate-specific antigen gene contains a functional androgen responsive element. Molec. Endocr. 5: 1921–1930, 1991.
- [77053] 15383. van den Elsen, P.; Bruns, G.; Gerhard, D. S.; Pravtcheva, D.; Jones, C.; Housman, D.; Ruddle, F. A.; Orkin, S.; Terhorst, C.: Assignment of the gene coding for the T3–delta subunit of the T3–T-cell receptor complex

to the long arm of human chromosome 11 and to mouse chromosome 9. Proc. Nat. Acad. Sci. 82: 2920–2924, 1985.

[77054] 15384. van Rijs, J.; Giguere, V.; Hurst, J.; van Agthoven, T.; Geurts van Kessel, A.; Goyert, S.; Grosveld, F.: Chromosomal localization of the human Thy-1 gene. Proc. Nat. Acad. Sci. 82: 5832–5835, 1985.

[77055] 15385. Williams, A. F.; Gagnon, J.: Neuronal cell Thy-1 glycoprotein: homology with immunoglobulin. Science 216: 696–703, 1982.

[77056] 15386. Johansson, M.; Karlsson, A.: Cloning of the cDNA and chromosomal localization of the gene for human thymidine kinase 2. J. Biol. Chem. 272: 8454–8458, 1997.

[77057] 15387. Mandel, H.; Szargel, R.; Labay, V.; Elpeleg, O.; Saada, A.; Shalata, A.; Anbinder, Y.; Berkowitz, D.; Hartman, C.; Barak, M.; Eriksson, S.; Cohen, N.: The deoxyguanosine kinase gene is mutated in individuals with depleted hepatocerebral mitochondrial DNA. Nature Genet. 29: 337–341, 2001. Note: Erratum: Nature Genet. 29: 491 only, 2001.

[77058] 15388. Saada, A.; Shaag, A.; Mandel, H.; Nevo, Y.; Eriksson, S.; Elpeleg, O.: Mutant mitochondrial thymidine kinase in mitochondrial DNA depletion myopathy. Nature Genet.

29: 342–344, 2001.

- [77059] 15389.Wang, L.; Munch–Petersen, B.; Herrstrom Sjoberg, A.; Hellman, U.;Bergman, T.; Jornvall, H.; Eriksson, S.: Human thymidine kinase 2:molecular cloning and characterisation of the enzyme activity withantiviral and cytostatic nucleoside substrates. FEBS Lett. 443:170–174, 1999.
- [77060] 15390.Willecke, K.; Reuber, T.; Kucherlapati, R. S.; Ruddle, F. H.:Human mitochondrial thymidine kinase is coded for by a gene on chromosome16 of the nucleus. Somat. Cell Genet. 3: 237–245, 1977.
- [77061] 15391.Chang, J. G.; Scarpa, A.; Eddy, R. L.; Byers, M. G.; Harris, A.S.; Morrow, J. S.; Watkins, P.; Shows, T. B.; Forget, B. G.: Cloningof a portion of the chromosomal gene and cDNA for human beta–fodrin,the nonerythroid form of beta–spectrin. Genomics 17: 287–293, 1993.
- [77062] 15392.Watkins, P. C.; Eddy, R.; Forget, B. G.; Chang, J. G.; Rochelle,R.; Shows, T. B.: Assignment of a non–erythroid spectrin gene tohuman chromosome 2. (Abstract) Am. J. Hum. Genet. 43: A161, 1988.
- [77063] 15393.Miceli–Richard, C.; Lesage, S.; Rybojad, M.; Prieur, A.–M.; Manouvrier–Hanu,S.; Hafner, R.; Chamaillard, M.; Zouali, H.; Thomas, G.; Hugot, J.–P.: CARD15 mutations in Blau syndrome. Nature Genet. 29: 19–20, 2001.

- [77064] 15394.Cong, F.; Spencer, S.; Cote, J.-F.; Wu, Y.; Tremblay, M. L.; Lasky, L. A.; Goff, S. P.: Cytoskeletal protein PSTPIP1 directs the PEST-type protein tyrosine phosphatase to the c-Abl kinase to mediate Abl dephosphorylation. *Molec. Cell* 6: 1413–1423, 2000.
- [77065] 15395.Han, L. Wong, D.; Dhaka, A.; Afar, D.; White, M.; Xie, W.; Herschman, H.; Witte, O.; Colicelli, J.: Protein binding and signaling properties of RIN1 suggest a unique effector function. *Proc. Nat. Acad. Sci.* 94:4954–4959, 1997.
- [77066] 15396.Jhanwar, S. C.; Neel, B. G.; Hayward, W. S.; Chaganti, R. S. K.: Localization of the cellular oncogenes ABL, SIS, and FES on human germ-line chromosomes. *Cytogenet. Cell Genet.* 38: 73–75, 1984.
- [77067] 15397.Blank, V.; Andrews, N. C.: The Maf transcription factors: regulators of differentiation. *Trends Biochem. Sci.* 22: 437–441, 1997.
- [77068] 15398.Jamieson, R. V.; Perveen, R.; Kerr, B.; Carette, M.; Yardley, J.; Heon, E.; Wirth, M. G.; van Heyningen, V.; Donnai, D.; Munier, F.; Black, G. C. M.: Domain disruption and mutation of the bZIP transcription factor, MAF, associated with cataract, ocular anterior segment dysgenesis and coloboma. *Hum. Molec. Genet.* 11: 33–42, 2002.
- [77069] 15399.Kim, J. I.; Li, T.; Ho, I. C.; Grusby, M. J.; Glimcher, L.



H.:Requirement for the c-Maf transcription factor in crystallin generegulation and lens development. Proc. Nat. Acad. Sci. 96: 3781–3785,1999.

[77070] 15400.Nishizawa, M.; Kataoka, K.; Goto, N.; Fujiwara, K. T.; Kawai, S.: v-maf, a viral oncogene that encodes a 'leucine zipper' motif. Proc.Nat. Acad. Sci. 86: 7711–7715, 1989.

[77071] 15401.Yoshida, M. C.; Nishizawa, M.; Kataoka, K.; Goto, N.; Fujiwara,K. T.; Kawai, S.: Localization of the human MAF protooncogene onchromosome 16 to bands q22–q23. (Abstract) Cytogenet. Cell Genet. 58:2003 only, 1991.

[77072] 15402.Bao, S.; Tibbetts, R. S.; Brumbaugh, K. M.; Fang, Y.; Richardson,D. A.; Ali, A.; Chen, S. M.; Abraham, R. T.; Wang, X.–F.: ATR/ATM–mediatedphosphorylation of human Rad17 is required for genotoxic stress responses. Nature 411:969–974, 2001.

[77073] 15403.Barbosa, M. D. F. S.; Barrat, F. J.; Tchernev, V. T.; Nguyen, Q.A.; Mishra, V. S.; Colman, S. D.; Pastural, E.; Dufourcq–Lagelouse,R.; Fischer, A.; Holcombe, R. F.; Wallace, M. R.; Brandt, S. J.; deSaint Basile, G.; Kingsmore S. F.: Identification of mutations into two major mRNA isoforms of the Chediak–Higashi syndrome gene in human and

mouse. Hum. Molec. Genet. 6: 1091–1098, 1997.

- [77074] 15404. Barrat, F. J.; Auloge, L.; Pastural, E.; Dufourcq-Lagelouse, R.; Vilmer, E.; Cant, A. J.; Weissenbach, J.; Le Paslier, D.; Fischer, A.; de Saint Basile, G.: Genetic and physical mapping of the Chediak–Higashis syndrome on chromosome 1q42–43. Am. J. Hum. Genet. 59: 625–632, 1996.
- [77075] 15405. Dufourcq-Lagelouse, R.; Lambert, N.; Duval, M.; Viot, G.; Vilmer, E.; Fischer, A.; Prieur, M.; de Saint Basile, G.: Chediak–Higashis syndrome associated with maternal uniparental isodisomy of chromosome 1. Europ. J. Hum. Genet. 7: 633–637, 1999.
- [77076] 15406. Faigle, W.; Raposo, G.; Tenza, D.; Pinet, V.; Vogt, A. B.; Kropshofer, H.; Fischer, A.; de Saint-Basile, G.; Amigorena, S.: Deficient peptide loading and MHC class II endosomal sorting in a human genetic immunodeficiency-disease: the Chediak–Higashi syndrome. J. Cell Biol. 141: 1121–1134, 1998.
- [77077] 15407. Karim, M. A.; Suzuki, K.; Fukai, K.; Oh, J.; Nagle, D. L.; Moore, K. J.; Barbosa, E.; Falik-Borenstein, T.; Filipovich, A.; Ishida, Y.; Kivrikko, S.; Klein, C.; and 8 others: Apparent genotype–phenotype correlation in childhood, adolescent, and adult Chediak–Higashi syndrome. Am. J. Med. Genet.

108: 16–22, 2002.

- [77078] 15408.Kunieda, T.; Ide, H.; Nakagiri, M.; Yoneda, K.; Konfortov, B.;Ogawa, H.: Localization of the locus responsible for Chediak–Higashis syndrome in cattle to bovine chromosome 28. *Anim. Genet.* 31: 87–90,2000.
- [77079] 15409.Akama, T. O.; Nishida, K.; Nakayama, J.; Watanabe, H.; Ozaki, K.;Nakamura, T.; Dota, A.; Kawasaki, S.; Inoue, Y.; Maeda, N.; Yamamoto,S.; Fujiwara, T.; Thonar, E. J.–M. A.; Shimomura, Y.; Kinoshita, S.;Tanigami, A.; Fukuda, M. N.: Macular corneal dystrophy type I andtype II are caused by distinct mutations in a new sulphotransferasegene. *Nature Genet.* 26: 237–241, 2000.
- [77080] 15410.Enomoto, A.; Kimura, H.; Chairoungdua, A.; Shigeta, Y.; Jutabha,P.; Cha, S. H.; Hosoyamada, M.; Takeda, M.; Sekine, T.; Igarashi,T.; Matsuo, H.; Kikuchi, Y.; Oda, T.; Ichida, K.; Hosoya, T.; Shimokata,K.; Niwa, T.; Kanai, Y.; Endou, H.: Molecular identification of arenal urate–anion exchanger that regulates blood urate levels. *Nature* 417:447–452, 2002.
- [77081] 15411.Goedert, M.; Hasegawa, J.; Craxton, M.; Leversha, M. A.; Clegg,S.: Assignment of the human stress–activated protein kinase–3 gene(SAPK3) to chromosome 22q13.3 by fluorescence in situ hybridization. *Genomics* 41:501–502,

1997.

- [77082] 15412. Yang, H.; Jeffrey, P. D.; Miller, J.; Kinnucan, E.; Sun, Y.; Thoma, N. H.; Zheng, N.; Chen, P.-L.; Lee, W.-H.; Pavletich, N. P.: BRCA2 function in DNA binding and recombination from a BRCA2-DSS1-ssDNA structure. *Science* 297: 1837–1848, 2002.
- [77083] 15413. Koettnitz, K.; Kappel, B.; Baumruker, T.; Hauber, J.; Bevec, D.: The genomic structure encoding human initiation factor eIF-5A. *Gene* 144:249–252, 1994.
- [77084] 15414. Koettnitz, K.; Wohl, T.; Kappel, B.; Lottspeich, F.; Hauber, J.; Bevec, D.: Identification of a new member of the human eIF-5A gene family. *Gene* 159: 283–284, 1995.
- [77085] 15415. Steinkasserer, A.; Jones, T.; Sheer, D.; Koettnitz, K.; Hauber, J.; Bevec, D.: The eukaryotic cofactor for the human immunodeficiency virus type 1 (HIV-1) rev protein, eIF-5A, maps to chromosome 17p12–p13: three eIF-5A pseudogenes map to 10q23.3, 17q25, and 19q13.2. *Genomics* 25:749–752, 1995.
- [77086] 15416. Hartley, D. A.; Preiss, A.; Artavanis-Tsakonas, S.: A deduced gene product from the *Drosophila* neurogenic locus, enhancer of split, shows homology to mammalian G-protein beta subunit. *Cell* 55: 785–795, 1988.
- [77087] 15417. Hou, E. W.; Li, S.-L.: Genomic organization and

chromosome localization to band 19p13.3 of the human AES gene: gene product exhibits strong similarity to the N-terminal domain of Drosophila enhancer of split Groucho protein. DNA Cell Biol. 17: 911–913, 1998.

[77088] 15418. Miyasaka, H.; Choudhury, B. K.; Hou, E. W.; Li, S.

S.-L.: Molecular cloning and expression of mouse and human cDNA encoding AES and ESG proteins with strong similarity to Drosophila enhancer of split groucho protein. Europ. J. Biochem. 216: 343–352, 1993.

[77089] 15419. Imai, Y.; Kurokawa, M.; Tanaka, K.; Friedman, A. D.;

Ogawa, S.; Mitani, K.; Yazaki, Y.; Hirai, H.: TLE, the human homolog of groucho, interacts with AML1 and acts as a repressor of AML1-induced transactivation.

Biochem. Biophys. Res. Commun. 252: 582–589, 1998.

[77090] 15420. Liu, Y.; Dehni, G.; Purcell, K. J.; Sokolow, J.; Car-

cangiu, M. L.; Artavanis-Tsakonas, S.; Stifani, S.: Epithelial expression and chromosomal location of human TLE genes: implications for Notch signaling and neoplasia. Genomics 31: 58–64, 1996.

[77091] 15421. Stifani, S.; Blaumueller, C. M.; Redhead, N. J.; Hill,

R. E.; Artavanis-Tsakonas, S.: Human homologs of a Drosophila enhancer of split gene product define a novel family of nuclear proteins. Nature Genet. 2:

119–127,1992.

- [77092] 15422.de Bruijn, D. R. H.; Baats, E.; Zechner, U.; de Leeuw, B.; Balemans,M.; Olde Weghuis, D.; Hirning–Folz, U.; Geurts van Kessel, A.: Isolationand characterization of the mouse homolog of SYT, a gene implicatedin the development of human synovial sarcomas. *Oncogene* 13: 643–648,1996.
- [77093] 15423.de Bruijn, D. R. H.; Kater–Baats, E.; Eleveld, M.; Merkx, G.; vanKessel, A. G.: Mapping and characterization of the mouse and humanSS18 genes, two human SS18–like genes and a mouse Ss18 pseudogene. *Cytogenet.Cell Genet.* 92: 310–319, 2001.
- [77094] 15424.Thaete, C.; Brett, D.; Monaghan, P.; Whitehouse, S.; Rennie, G.;Rayner, E.; Cooper, C. S.; Goodwin, G.: Functional domains of theSYT and SYT–SSX synovial sarcoma translocation proteins and co–localizationwith the SNF protein BRM in the nucleus. *Hum. Molec. Genet.* 8: 585–591,1999.
- [77095] 15425.Ge, Q.; Trieu, E. P.; Targoff, I. N.: Primary structure and functionalexpression of human glycyl–tRNA synthetase, an autoantigen in myositis.*J. Biol. Chem.* 269: 28790–28797, 1994.
- [77096] 15426.Nichols, R. C.; Pai, S. I.; Ge, Q.; Targoff, I. N.; Plotz,

P. H.;Liu, P.: Localization of two human autoantigen genes by PCR screening and in situ hybridization--Glycyl-tRNA synthetase locates to 7p15 and alanyl-tRNA synthetase locates to 16q22. *Genomics* 30: 131-132, 1995.

[77097] 15427. Shiba, K.; Schimmel, P.; Motegi, H.; Noda, T.: Human glycyl-tRNA synthetase: wide divergence of primary structure from bacterial counterpart and species-specific aminoacylation. *J. Biol. Chem.* 269: 30049-30055, 1994.

[77098] 15428. Williams, J.; Osvath, S.; Khong, T. F.; Pearse, M.; Power, D.: Cloning, sequencing and bacterial expression of human glycine tRNA synthetase. *Nucleic Acids Res.* 23: 1307-1310, 1995.

[77099] 15429. Avraham, K. B.; Prezioso, V. R.; Chen, W. S.; Lai, E.; Sladek, F. M.; Zhong, W.; Darnell, J. E., Jr.; Jenkins, N. A.; Copeland, N. G.: Murine chromosomal location of four hepatocyte-enriched transcription factors: HNF-3-alpha, HNF-3-beta, HNF-3-gamma, and HNF-4. *Genomics* 13: 264-268, 1992.

[77100] 15430. Deleuze, J. F.; Dhorne, S.; Hazan, J.; Borghi, E.; Raynaud, N.; Pollet, N.; Meunier-Rotival, M.; Deschatrette, J.; Alagille, D.; Hadchouel, M.: Deleted chromosome 20 from a patient with Alagille syndrome isolated in a cell hybrid through leucine transport selection: study of three-

candidate genes. Mammalian Genome 5: 663–669, 1994.

[77101] 15431.Kaestner, K. H.; Hiemisch, H.; Luckow, B.; Schutz, G.: The HNF-3gene family of transcription factors in mice: gene structure, cDNAsequence, and mRNA distribution. Genomics 20: 377–385, 1994.

[77102] 15432.Mincheva, A.; Lichter, P.; Schutz, G.; Kaestner, K. H.: Assignmentof the human genes for hepatocyte nuclear factor 3-alpha, -beta, and-gamma (HNF3A, HNF3B, HNF3G) to 14q12-q13, 20p11, and 19q13.2-q13.4. Genomics 39:417–419, 1997.

[77103] 15433.Adams, R. H.; Porras, A.; Alonso, G.; Jones, M.; Vintersten, K.;Panelli, S.; Valladares, A.; Perez, L.; Klein, R.; Nebreda, A. R.: Essential role of p38-alpha MAP kinase in placental but not embryoniccardiovascular development. Molec. Cell 6: 109–116, 2000.

[77104] 15434.Ge, B.; Gram, H.; Di Padova, F.; Huang, B.; New, L.; Ulevitch,R. J.; Luo, Y.; Han, J.: MAPKK-independent activation of p38-alphamediated by TAB1-dependent autophosphorylation of p38-alpha. Science 295:1291–1294, 2002.

[77105] 15435.Han, J.; Lee, J.-D.; Bibbs, L.; Ulevitch, R. J.: A MAP kinasetargeted by endotoxin and hyperosmolarity in mammalian cells. Science 265:808–811, 1994.



- [77106] 15436.Kim, D. H.; Feinbaum, R.; Alloing, G.; Emerson, F. E.; Garsin,D. A.; Inoue, H.; Tanaka-Hino, M.; Hisamoto, N.; Matsumoto, K.; Tan,M.-W.; Ausubel, F. M.: A conserved p38 MAP kinase pathway in *Caenorhabditiselegans* innate immunity. *Science* 297: 623–626, 2002.
- [77107] 15437.Kumar, S.; McLaughlin, M. M.; McDonnell, P. C.; Lee, J. C.; Livi,G. P.; Young, P. R.: Human mitogen-activated protein kinase CSBP1,but not CSBP2, complements a *hog1* deletion in yeast. *J. Biol. Chem.* 270:29043–29046, 1995.
- [77108] 15438.Lee, J. C.; Laydon, J. T.; McDonnell, P. C.; Gallagher, T. F.;Kumar, S.; Green, D.; McNulty, D.; Blumenthal, M. J.; Heys, J. R.;Landvatter, S. W.; Stickler, J. E.; McLaughlin, M. M.; Siemens, I.R.; Fisher, S. M.; Livi, G. P.; White, J. R.; Adams, J. L.; Young,P. R.: A protein kinase involved in the regulation of inflammatorycytokine biosynthesis. *Nature* 372: 739–746, 1994.
- [77109] 15439.Liao, P.; Georgakopoulos, D.; Kovacs, A.; Zheng, M.; Lerner, D.;Pu, H.; Saffitz, J.; Chien, K.; Xiao, R.-P.; Kass, D. A.; Wang, Y.: The in vivo role of p38 MAP kinases in cardiac remodeling and restrictivecardiomyopathy. *Proc. Nat. Acad. Sci.* 98: 12283–12288, 2001.
- [77110] 15440.Maizels, E. T.; Mukherjee, A.; Sithanandam, G.; Pe-

ters, C. A.;Cottom, J.; Mayo, K. E.; Hunzicker–Dunn, M.: Developmental regulationof mitogen–activated protein kinase–activated kinases–2 and –3 (MAPKAPK–2/–3)in vivo during corpus luteum formation in the rat. *Molec. Endocr.* 15:716–733, 2001.

[77111] 15441.McDonnell, P. C.; DiLella, A. G.; Lee, J. C.; Young, P. R.: Localizationof the human stress responsive MAP kinase–like CSAIDs binding protein(CSBP) gene to chromosome 6p21.3/21.2. *Genomics* 29: 301–302, 1995.

[77112] 15442.New, L.; Jiang, Y.; Zhao, M.; Liu, K.; Zhu, W.; Flood, L. J.;Kato, Y.; Parry, G. C. N.; Han, J.: PRAK, a novel protein kinase regulated by the p38 MAP kinase. *EMBO J.* 17: 3372–3384, 1998.

[77113] 15443.Ni, H.; Wang, X. S.; Diener, K.; Yao, Z.: MAPKAPK5, a novel mitogen–activated protein kinase (MAPK)–activated protein kinase, is a substrate ofthe extracellular–regulated kinase (ERK) and p38 kinase. *Biochem.Biophys. Res. Commun.* 243: 492–496, 1998.

[77114] 15444.Takekawa, M.; Maeda, T.; Saito, H.: Protein phosphatase 2C–alphainhibits the human stress–responsive p38 and JNK MAPK pathways. *EMBOJ.* 17: 4744–4752, 1998.

[77115] 15445.Tamura, K.; Sudo, T.; Senftleben, U.; Dadak, A. M.;

Johnson, R.;Karin, M.: Requirement for p38-alpha in erythropoietin expression:a role for stress kinases in erythropoiesis. Cell 102: 221-231,2000.

[77116] 15446.Haber, N.; Stengel, D.; Defer, N.; Roeckel, N.; Mattei, M.-G.;Hanoune, J.: Chromosomal mapping of human adenylyl cyclase genestype III, type V and type VI. Hum. Genet. 94: 69-73, 1994.

[77117] 15447.Sinnarajah, S.; Dessauer, C. W.; Srikumar, D.; Chen, J.; Yuen,J.; Yilma, S.; Dennis, J. C.; Morrison, E. E.; Vodyanoy, V.; Kehrl,J. H.: RGS2 regulates signal transduction in olfactory neurons byattenuating activation of adenylyl cyclase III. Nature 409: 1051-1055,2001.

[77118] 15448.Pellizzoni, L.; Kataoka, N.; Charroux, B.; Dreyfuss, G.: A novelfunction for SMN, the spinal muscular atrophy gene product, in pre-mRNA splicing. Cell 95: 615-624, 1998.

[77119] 15449.Brown, K. A.; Leek, J. P.; Lench, N. J.; Moynihan, L. M.; Markham,A. F.; Mueller, R. F.: Human sequences homologous to the gene forthe cochlear protein Ocp-II do not map to currently known non-syndromichearing loss loci. Ann. Hum. Genet. 60: 385-389, 1996.

[77120] 15450.Chen, H.; Thalmann, I.; Adams, J. C.; Avraham, K. B.; Copeland,N. G.; Jenkins, N. A.; Beier, D. R.; Corey, D.

P.; Thalmann, R.; Duyk, G. M.: cDNA cloning, tissue distribution, and chromosomal localization of Ocp2, a gene encoding a putative transcription-associated factor predominantly expressed in the auditory organs. *Genomics* 27: 389–398, 1995.

[77121] 15451. Demetrick, D. J.; Zhang, H.; Beach, D. H.: Chromosomal mapping of the genes for the human CDK2/cyclin A-associated proteins p19 (SKP1A and SKP1B) and p45 (SKP2). *Cytogenet. Cell Genet.* 73: 104–107, 1996.

[77122] 15452. Liang, Y.; Chen, H.; Asher, J. H., Jr.; Chang, C.-C.; Friedman, T. B.: Human inner ear OCP2 cDNA maps to 5q22–5q35.2 with related sequences on chromosomes 4p16.2–4p14, 5p13–5q22, 7pter–q22, 10 and 12p13–12qter. *Gene* 184: 163–167, 1997.

[77123] 15453. Sowden, J.; Morrison, K.; Schofield, J.; Putt, W.; Edwards, Y.: A novel cDNA with homology to an RNA polymerase II elongation factor maps to human chromosome 5q31 (TCEB1L) and to mouse chromosome 11 (Tceb1l). *Genomics* 29: 145–151, 1995.

[77124] 15454. Thalmann, I.; Rosenthal, H. L.; Moore, B. W.; Thalmann, R.: Organ of Corti-specific polypeptides: OCP-I and OCP-II. *Arch. Oto-Rhino-Laryngol.* 226: 123–128, 1980.

[77125] 15455. Thalmann, I.; Takahashi, K.; Varghese, J.; Comegys,

T. H.; Thalmann,R.: Biochemical features of major organ of Corti proteins (OCP-I and OCP-II) including partial amino acid sequence. *Laryngoscope* 100:99–105, 1990.

- [77126] 15456.Zhang, H.; Kobayashi, R.; Galaktionov, K.; Beach, D.: p19Skp1 and p45Skp2 are essential elements of the cyclin A-CDK2 S phase kinase. *Cell* 82:915–925, 1995.
- [77127] 15457.Anand, A.; Chada, K.: In vivo modulation of Hmgic reduces obesity. *NatureGenet.* 24: 377–380, 2000.
- [77128] 15458.Arlotta, P.; Tai, A. K.-F.; Manfioletti, G.; Clifford, C.; Jay,G.; Ono, S. J.: Transgenic mice expressing a truncated form of the high mobility group I-C protein develop adiposity and an abnormally high prevalence of lipomas. *J. Biol. Chem.* 275: 14394–14400, 2000.
- [77129] 15459.Ashar, H. R.; Cherath, L.; Przybysz, K. M.; Chada, K.: Genomic characterization of human HMGI-C, a member of the accessory transcription factor family found at translocation breakpoints in lipomas. *Genomics* 31:207–214, 1996.
- [77130] 15460.Ashar, H. R.; Schoenberg Fejzo, M.; Tkachenko, A.; Zhou, X.; Fletcher,J. A.; Weremowicz, S.; Morton, C. C.; Chada, K.: Disruption of the architectural factor HMGI-C: DNA-binding AT hook motifs fused in lipoma to distinct transcriptional regulatory domains. *Cell* 82: 57–65,1995.

- [77131] 15461. Chau, K.-Y.; Patel, U. A.; Lee, K.-L. D.; Lam, H.-Y. P.; Crane-Robinson, C.: The gene for the human architectural transcription factor HMGI-C consists of five exons each coding for a distinct functional element. *Nucleic Acids Res.* 23: 4262-4266, 1995.
- [77132] 15462. Danforth, E., Jr.: Failure of adipocyte differentiation causes type II diabetes mellitus. *Nature Genet.* 26: 13 only, 2000.
- [77133] 15463. Friedmann, M.; Holth, L. T.; Zoghbi, H. Y.; Reeves, R.: Organization, inducible-expression and chromosome localization of the human HMG-I(Y) nonhistone protein gene. *Nucleic Acids Res.* 21: 4259-4267, 1993.
- [77134] 15464. Ishwad, C. S.; Shriver, M. D.; Lassig, D. M.; Ferrell, R. E.: The high mobility group I-C gene (HMGI-C): polymorphism and genetic localization. *Hum. Genet.* 99: 103-105, 1997.
- [77135] 15465. Kazmierczak, B.; Dal Cin, P.; Wanschura, S.; Bartschke, S.; Vanden Berghe, H.; Bullerdiek, J.: Cloning and molecular characterization of part of a new gene fused to HMGIC in mesenchymal tumors. *Am. J. Path.* 152: 431-435, 1998.
- [77136] 15466. Kazmierczak, B.; Pohnke, Y.; Bullerdiek, J.: Fusion transcripts between the HMGIC gene and RTVL-H-related

sequences in mesenchymal tumors without cytogenetic aberrations. *Genomics* 38: 223–226, 1996.

- [77137] 15467. Manfioletti, G.; Rustighi, A.; Mantovani, F.; Goodwin, G. H.; Giancotti, V.: Isolation and characterization of the gene coding for murine high-mobility-group protein HMGI-C. *Gene* 167: 249–253, 1995.
- [77138] 15468. Mine, N.; Kurose, K.; Nagai, H.; Doi, D.; Ota, Y.; Yoneyama, K.; Konishi, H.; Araki, T.; Emi, M.: Gene fusion involving HMGIC is a frequent aberration in uterine leiomyomas. *J. Hum. Genet.* 46: 408–412, 2001.
- [77139] 15469. Nucci, M. R.; Weremowicz, S.; Neskey, D. M.; Sornberger, K.; Tallini, G.; Morton, C. C.; Quade, B. J.: Chromosomal translocation t(8;12) induces aberrant HMGIC expression in aggressive angiomyxoma of the vulva. *Genes Chromosomes Cancer* 32: 172–176, 2001.
- [77140] 15470. Rodrigues, N. R.; Owen, N.; Talbot, K.; Ignatius, J.; Dubowitz, V.; Davies, K. E.: Deletions in the survival motor neuron gene on 5q13 in autosomal recessive spinal muscular atrophy. *Hum. Molec. Genet.* 4: 631–634, 1995.
- [77141] 15471. Rossoll, W.; Kroning, A.-K.; Ohndorf, U.-M.; Steegborn, C.; Jablonka, S.; Sendtner, M.: Specific interaction of Smn, the spinal muscular atrophy determining gene product, with hnRNP-R and grp-rbp/hnRNP-Q: a role for Smn

in RNA processing in motor axons? Hum. Molec. Genet. 11:93–105, 2002.

[77142] 15472.Simard, L. R.; Rochette, C.; Semionov, A.; Morgan, K.; Vanasse,M.: SMN(T) and NAIP mutations in Canadian families with spinal muscularatrophy (SMA): genotype/phenotype correlations with disease severity. Am.J. Med. Genet. 72: 51–58, 1997.

[77143] 15473.Sossi, V.; Giuli, A.; Vitali, T.; Tiziano, F.; Mirabella, M.;Antonelli, A.; Neri, G.; Brahe, C.: Premature termination mutationsin exon 3 of the SMN1 gene are associated with exon skipping and arelatively mild SMA phenotype. Europ. J. Hum. Genet. 9: 113–120,2001.

[77144] 15474.Stewart, H.; Wallace, A.; McGaughran, J.; Mountford, R.; Kingston,H.: Molecular diagnosis of spinal muscular atrophy. Arch. Dis. Child. 78:531–535, 1998.

[77145] 15475.Talbot, K.; Ponting, C. P.; Theodosiou, A. M.; Rodrigues, N. R.;Surtees, R.; Mountford, R.; Davies, K. E.: Missense mutation clusteringin the survival motor neuron gene: a role for a conserved tyrosineand glycine rich region of the protein in RNA metabolism? Hum. Molec.Genet. 6: 497–500, 1997.

[77146] 15476.van der Steege, G.; Grootschloten, P. M.; Cobben, J. M.; Zappata,S.; Scheffer, H.; den Dunnen, J. T.; van Om–



men, G.-J. B.; Brahe, C.; Buys, C. H. C. M.: Apparent gene conversions involving the SMN gene in the region of the spinal muscular atrophy locus on chromosome 5. *Am. J. Hum. Genet.* 59: 834–838, 1996.

[77147] 15477. Viollet, L.; Bertrand, S.; Bueno Brunialti, A. L.; Lefebvre, S.; Burlet, P.; Clermont, O.; Cruaud, C.; Guenet, J.-L.; Munnich, A.; Melki, J.: cDNA isolation, expression, and chromosomal localization of the mouse survival motor neuron gene (*Smn*). *Genomics* 40: 185–188, 1997.

[77148] 15478. Wang, C. H.; Xu, J.; Carter, T. A.; Ross, B. M.; Dominski, M. K.; Bellcross, C. A.; Penchaszadeh, G. K.; Munsat, T. L.; Gilliam, T. C.: Characterization of survival motor neuron (*SMNT*) gene deletions in asymptomatic carriers of spinal muscular atrophy. *Hum. Molec. Genet.* 5: 359–365, 1996.

[77149] 15479. Wang, J.; Dreyfuss, G.: A cell system with targeted disruption of the SMN gene: functional conservation of the SMN protein and dependence of Gemin2 on SMN. *J. Biol. Chem.* 276: 9599–9605, 2001.

[77150] 15480. Wirth, B.: An update of the mutation spectrum of the survival motor neuron gene (*SMN1*) in autosomal recessive spinal muscular atrophy (SMA). *Hum. Mutat.* 15: 228–237, 2000.

- [77151] 15481. Wirth, B.; Hahnen, E.; Morgan, K.; DiDonato, C. J.; Dadze, A.; Rudnik-Schoneborn, S.; Simard, L. R.; Zerres, K.; Burghes, A. H. M.: Allelic association and deletions in autosomal recessive proximal spinal muscular atrophy: association of marker genotype with disease severity and candidate cDNAs. *Hum. Molec. Genet.* 4: 1273–1284, 1995.
- [77152] 15482. Wirth, B.; Herz, M.; Wetter, A.; Moskau, S.; Hahnen, E.; Rudnik-Schoneborn, S.; Wienker, T.; Zerres, K.: Quantitative analysis of survival motor neuron copies: identification of subtle SMN1 mutations in patients with spinal muscular atrophy, genotype–phenotype correlation, and implications for genetic counseling. *Am. J. Hum. Genet.* 64: 1340–1356, 1999.
- [77153] 15483. Young, P. J.; Man, N.; Lorson, C. L.; Le, T. T.; Androphy, E. J.; Burghes, A. H. M.; Morris, G. E.: The exon 2b region of the spinal muscular atrophy protein, SMN, is involved in self-association and SIP1 binding. *Hum. Molec. Genet.* 9: 2869–2877, 2000.
- [77154] 15484. David, C.; Solimena, M.; De Camilli, P.: Autoimmunity in stiff-man syndrome with breast cancer is targeted to the C-terminal region of human amphiphysin, a protein similar to the yeast proteins, Rvs167 and Rvs161. *FEBS Lett.* 351: 73–79, 1994.

- [77155] 15485.De Camilli, P.; Thomas, A.; Cofield, R.; Folli, F.; Lichte, B.;Piccolo, G.; Meinck, H.-M.; Austoni, M.; Fassetta, G.; Bottazzo, G.;Bates, D.; Cartledge, N.; Solimena, M.; Kilimann, M. W.: The synapticvesicle-associated protein amphiphysin is the 128-kD autoantigen ofstiff-man syndrome with breast cancer. J. Exp. Med. 178: 2219-2223,1993.
- [77156] 15486.Jenkins, N. A.; Gilbert, D. J.; Yamamoto, R.; Kilimann, M. W.;Copeland, N. G.: Amphiphysin (Amph) maps to the proximal region ofmouse chromosome 13. Genomics 28: 363-365, 1995.
- [77157] 15487.Lichte, B.; Veh, R. W.; Meyer, H. E.; Kilimann, M. W.: Amphiphysin,a novel protein associated with synaptic vesicles. EMBO J. 11: 2521-2530,1992.
- [77158] 15488.Yamamoto, R.; Li, X.; Francke, U.; Kilimann, M. W.: Genetic mappingof the human amphiphysin gene (AMPH) at 7p14-p13 excludes its involvementin retinitis pigmentosa 9 or dominant cystoid macular dystrophy. (Letter) Am.J. Hum. Genet. 57: 970-972, 1995.
- [77159] 15489.Yamamoto, R.; Li, X.; Winter, S.; Francke, U.; Kilimann, M. W.: Primary structure of human amphiphysin, the dominant autoantigenof paraneoplastic Stiff-Man syndrome, and mapping of its gene (AMPH)to chromosome

7p13–p14. Hum. Molec. Genet. 4: 265–268, 1995.

- [77160] 15490.Kingsmore, S. F.; Suh, D.; Seldin, M. F.: Genetic mapping of the glycine receptor alpha-3 subunit on mouse chromosome 8. Mammalian Genome 5: 831–832, 1994.
- [77161] 15491.Kuhse, J.; Schmieden, V.; Betz, H.: Identification and functional expression of a novel ligand binding subunit of the inhibitory glycine receptor. J. Biol. Chem. 265: 22317–22320, 1990.
- [77162] 15492.Nikolic, Z.; Laube, B.; Weber, R. G.; Lichter, P.; Kioschis, P.; Poustka, A.; Mulhardt, C.; Becker, C.–M.: The human glycine receptor subunit alpha-3: GLRA3 gene structure, chromosomal localization, and functional characterization of alternative transcripts. J. Biol. Chem. 273: 19708–19714, 1998.
- [77163] 15493.Birkenmeier, E. H.; Rowe, L. B.; Crossman, M. W.; Gordon, J. I.: Ileal lipid-binding protein (Illbp) gene maps to mouse chromosome 11. Mammalian Genome 5: 805–806, 1994.
- [77164] 15494.Oelkers, P.; Dawson, P. A.: Cloning and chromosomal localization of the human ileal lipid-binding protein. Biochim. Biophys. Acta 1257:199–202, 1995.
- [77165] 15495.Johnson, D. W.; Berg, J. N.; Baldwin, M. A.; Gallione, C. J.; Marondel, I.; Yoon, S.–J.; Stenzel, T. T.; Speer, M.;

Pericak-Vance, M. A.;Diamond, A.; Guttmacher, A. E.; Jackson, C. E.; Attisano, L.; Kucherlapati,R.; Porteous, M. E. M.; Marchuk, D. A.: Mutations in the activinreceptor-like kinase 1 gene in hereditary haemorrhagic telangiectasi-atype 2. *Nature Genet.* 13: 189–195, 1996.

[77166] 15496.Seimiya, H.; Sawabe, T.; Inazawa, J.; Tsuruo, T.: Cloning, expressionand chromosomal localization of a novel gene for protein tyrosinephosphatase (PTP-U2) induced by various differentiation-inducing agents. *Onco-gene* 10:1731–1738, 1995.

[77167] 15497.Wiggins, R. C.; Wiggins, J. E.; Goyal, M.; Wharram, B. L.; Thomas,P. E.: Molecular cloning of cDNAs encoding human GLEPP1, a membraneprotein tyrosine phosphatase: characterization of the GLEPP1 proteindistribution in human kidney and assignment of the GLEPP1 gene tohuman chromosome 12p12–p13. *Genomics* 27: 174–181, 1995.

[77168] 15498.Borsani, G.; Rugarli, E. I.; Taglialatela, M.; Wong, C.; Ballabio,A.: Characterization of a human and murine gene (CLCN3) sharing similaritiesto voltage-gated chloride channels and to a yeast integral membraneprotein. *Genomics* 27: 131–141, 1995.

[77169] 15499.Mills, K. A.; Mathews, K. D.; Scherpbier-Heddema, T.; Buetow, K.H.; Baldini, A.; Ballabio, A.; Borsani, G.: Ge-

netic and physical mapping of a voltage-dependent chloride channel gene to human 4q32 and to mouse 8. Genomics 36: 374–376, 1996.

- [77170] 15500. Stobrawa, S. M.; Breiderhoff, T.; Takamori, S.; Engel, D.; Schweizer, M.; Zdebik, A. A.; Bosl, M. R.; Ruether, K.; Jahn, H.; Draguhn, A.; Jahn, R.; Jentsch, T. J.: Disruption of CLC-3, a chloride channel expressed on synaptic vesicles, leads to a loss of the hippocampus. Neuron 29:185–196, 2001.
- [77171] 15501. Taine, L.; Couprie, I.; Boisseau, P.; Saura, R.; Lacombe, D.; Arveiler, B.: Refined localisation of the voltage-gated chloride channel, CLCN3, to 4q33. Hum. Genet. 102: 178–181, 1998.
- [77172] 15502. Yoshikawa, M.; Uchida, S.; Ezaki, J.; Rai, T.; Hayama, A.; Kobayashi, K.; Kida, Y.; Noda, M.; Koike, M.; Uchiyama, Y.; Marumo, F.; Kominami, E.; Sasaki, S.: CLC-3 deficiency leads to phenotypes similar to human neuronal ceroid lipofuscinosis. Genes Cells 7: 597–605, 2002.
- [77173] 15503. Pagliuca, A.; Bartoli, P. C.; Saccone, S.; Valle, G. D.; Lania, L.: Molecular cloning of ID4, a novel dominant negative helix-loop-helix human gene on chromosome 6p21.3–p22. Genomics 27: 200–203, 1995.
- [77174] 15504. Rigolet, M.; Rich, T.; Gross-Morand, M.-S.; Molina-

Gomes, D.; Viegas-Pequignot, E.; Junien, C.: cDNA cloning, tissue distribution and chromosomal localization of the human ID4 gene. DNA Res. 5: 309–313, 1998.

[77175] 15505. Jia, S.; VanDusen, W. J.; Diehl, R. E.; Kohl, N. E.; Dixon, R. A. F.; Elliston, K. O.; Stern, A. M.; Friedman, P. A.: cDNA cloning and expression of bovine aspartyl (asparaginyl) beta-hydroxylase. J. Biol. Chem. 267: 14322–14327, 1992.

[77176] 15506. Koriath, F.; Gieffers, C.; Frey, J.: Cloning and characterization of the human gene encoding aspartyl beta-hydroxylase. Gene 150:395–399, 1994.

[77177] 15507. Lavaissiere, L.; Jia, S.; Nishiyama, M.; de la Monte, S.; Stern, A. M.; Wands, J. R.; Friedman, P. A.: Overexpression of human aspartyl (asparaginyl)-beta-hydroxylase in hepatocellular carcinoma and cholangiocarcinoma. J. Clin. Invest. 98:1313–1323, 1996.

[77178] 15508. Lim, K. Y.; Hong, C.-S.; Kim, D. H.: cDNA cloning and characterization of human cardiac junctin. Gene 255: 35–42, 2000.

[77179] 15509. Scott, A. F.: Personal Communication. Baltimore, Md. 2/19/2001.

[77180] 15510. Treves, S.; Feriotto, G.; Moccagatta, L.; Gambari, R.; Zorzato, F.: Molecular cloning, expression, functional

characterization, chromosomal localization, and gene structure of junctate, a novel integral calcium-binding protein of sarco(endo)plasmic reticulum membrane. *J. Biol. Chem.* 275: 39555–39568, 2000.

[77181] 15511. Wetzell, G. T.; Ding, S.; Chen, F.: Molecular cloning of junctin from human and developing rabbit heart. *Molec. Genet. Metab.* 69:252–258, 2000.

[77182] 15512. Zhou, P.; Sun, L. J.; Dotsch, V.; Wagner, G.; Verdine, G. L.: Solution structure of the core NFATC1/DNA complex. *Cell* 92: 687–696, 1998.

[77183] 15513. McCaffrey, P. G.; Luo, C.; Kerppola, T. K.; Jain, J.; Badalian, T. M.; Ho, A. M.; Burgeon, E.; Lane, W. S.; Lambert, J. N.; Curran, T.; Verdine, G. L.; Rao, A.; Hogan, P. G.: Isolation of the cyclosporin-sensitive T cell transcription factor NFATp. *Science* 262: 750–754, 1993.

[77184] 15514. Rengarajan, J.; Tang, B.; Glimcher, L. H.: NFATc2 and NFATc3 regulate TH2 differentiation and modulate TCR-responsiveness of naive TH cells. *Nature Immun.* 3: 48–54, 2002.

[77185] 15515. Chan, K.; Han, X.-D.; Kan, Y. W.: An important function of Nrf2 in combating oxidative stress: detoxification of acetaminophen. *Proc. Nat. Acad. Sci.* 98: 4611–4616, 2001.



- [77186] 15516. Baud, V.; Chissoe, S. L.; Viegas-Pequignot, E.; Diriong, S.; N'Guyen, V. C.; Roe, B. A.; Lipinski, M.: EMR1, an unusual member in the family of hormone receptors with seven transmembrane segments. *Genomics* 26:334–344, 1995.
- [77187] 15517. McKnight, A. J.; Macfarlane, A. J.; Seldin, M. F.; Gordon, S.: Chromosome mapping of the Emr1 gene. *Mammalian Genome* 8: 946–947, 1997.
- [77188] 15518. Atanasoski, S.; Toldo, S. S.; Malipiero, U.; Schreiber, E.; Fries, R.; Fontana, A.: Isolation of the human genomic brain-2/N-Oct 3 gene (POUF3) and assignment to chromosome 6q16. *Genomics* 26: 272–280, 1995.
- [77189] 15519. Schreiber, E.; Tobler, A.; Malipiero, U.; Schaffner, W.; Fontana, A.: cDNA cloning of human N-Oct 3, a nervous-system specific POU domain transcription factor binding to the octamer DNA motif. *Nucleic Acids Res.* 21: 253–258, 1993.
- [77190] 15520. Xia, Y.-R.; Andersen, B.; Mehrabian, M.; Diep, A. T.; Warden, C. H.; Mohandas, T.; McEvilly, R. J.; Rosenfeld, M. G.; Lusk, A. J.: Chromosomal organization of mammalian POU domain factors. *Genomics* 18:126–130, 1993.
- [77191] 15521. Brass, N.; Heckel, D.; Sahin, U.; Pfreundschuh, M.; Sybrecht, G. W.; Meese, E.: Translation initiation factor eIF-

4gamma is encoded by an amplified gene and induces an immune response in squamous cell carcinoma. Hum. Molec. Genet. 6: 33–39, 1997.

- [77192] 15522. Gradi, A.; Imataka, H.; Svitkin, Y. V.; Rom, E.; Raught, B.; Morino, S.; Sonenberg, N.: A novel functional human eukaryotic translation initiation factor 4G. Molec. Cell. Biol. 18: 334–342, 1998.
- [77193] 15523. Imataka, H.; Gradi, A.; Sonenberg, N.: A newly identified N-terminal amino acid sequence of human eIF4G binds poly(A)-binding protein and functions in poly(A)-dependent translation. EMBO J. 17: 7480–7489, 1998.
- [77194] 15524. Imataka, H.; Sonenberg, N.: Human eukaryotic translation initiation factor 4G (eIF4G) possesses two separate and independent binding sites for eIF4A. Molec. Cell. Biol. 17: 6940–6947, 1997.
- [77195] 15525. Yan, R.; Rhoads, R. E.: Human protein synthesis initiation factor eIF-4-gamma is encoded by a single gene (EIF4G) that maps to chromosome 3q27-qter. Genomics 26: 394–398, 1995.
- [77196] 15526. Yan, R.; Rychlik, W.; Etchison, D.; Rhoads, R. E.: Amino acid sequence of the human protein synthesis initiation factor eIF-4-gamma. J. Biol. Chem. 267:

23226–23231, 1992.

- [77197] 15527. Corrigan, A. V.; Hift, R. J.; Hancock, V.; Meissner, D.; Davids, L.; Kirsch, R. E.; Meissner, P. N.: Identification and characterisation of a deletion (537delAT) in the protoporphyrinogen oxidase gene in a South African variegate porphyria family. *Hum. Mutat.* 12: 403–407, 1998.
- [77198] 15528. Aso, T.; Haque, D.; Fukudome, K.; Brower, C. S.; Conaway, J. W.; Conaway, R. C.: A human cDNA encoding the 110-kDa A subunit of RNA polymerase II transcription factor elongin. *Gene* 168: 277–278, 1996.
- [77199] 15529. Aso, T.; Lane, W. S.; Conaway, J. W.; Conaway, R. C.: Elongin(SIII): a multisubunit regulator of elongation by RNA polymerase II. *Science* 269:1439–1443, 1995.
- [77200] 15530. Aso, T.; Mokady, N.; Haque, D.; Conaway, R. C.; Conaway, J. W.: Assignment of a human gene encoding the 110-kDa subunit of general transcription factor elongin (SIII) to chromosome 1p36.1. *Genomics* 30:393–394, 1995.
- [77201] 15531. Vikkula, M.; Boon, L. M.; Carraway, K. L., III; Calvert, J. T.; Diamonti, A. J.; Goumnerov, B.; Pasyk, K. A.; Marchuk, D. A.; Warman, M. L.; Cantley, L. C.; Mulliken, J. B.; Olsen, B. R.: Vascular dysmorphogenesis caused by an activating mutation in the receptor tyrosine kinase TIE2.

Cell 87:1181–1190, 1996.

- [77202] 15532.Iwata, T.; Kogame, K.; Toki, T.; Yokoyama, M.; Yamamoto, M.; Ito,E.: Structure and chromosome mapping of the human small maf genesMAFG and MAFK. Cytogenet. Cell Genet. 82: 88–90, 1998.
- [77203] 15533.Motohashi, H.; Katsuoka, F.; Shavit, J. A.; Engel, J. D.; Yamamoto,M.: Positive or negative MARE-dependent transcriptional regulationis determined by the abundance of small Maf proteins. Cell 103:865–875, 2000.
- [77204] 15534.Peters, L. L.; Eicher, E. M.: The ubiquitous subunit of the globin enhancer-binding protein NF-E2 (Nfe2u) maps to mouse chromosome 5. Genomics 22:490–491, 1994.
- [77205] 15535.Shavit, J. A.; Motohashi, H.; Onodera, K.; Akasaka, J.; Yamamoto,M.; Engel, J. D.: Impaired megakaryopoiesis and behavioral defects in mafG-null mutant mice. Genes Dev. 12: 2164–2174, 1998.
- [77206] 15536.Toki, T.; Itoh, J.; Kitazawa, J.; Arai, K.; Hatakeyama, K.; Akasaka,J.; Igarashi, K.; Nomura, N.; Yokoyama, M.; Yamamoto, M.; Ito, E.: Human small Maf proteins form heterodimers with CNC family transcription factors and recognize the NF-E2 motif. Oncogene 14: 1901–1910, 1997.

- [77207] 15537. Yang, B.; He, B.; Abdel-Halim, S. M.; Tibell, A.; Brendel, M. D.; Bretzel, R. G.; Efendic, S.; Hillert, J.: Molecular cloning of a full-length cDNA for human type 3 adenylyl cyclase and its expression in human islets. *Biochem. Biophys. Res. Commun.* 254: 548–551, 1999.
- [77208] 15538. Arden, K. C.; Viars, C. S.; Weiss, S.; Argentin, S.; Nemer, M.: Localization of the human B-type natriuretic peptide precursor (NPPB) gene to chromosome 1p36. *Genomics* 26: 385–389, 1995.
- [77209] 15539. Jordan, C. E.; White, R. I., Jr.; Fischer, K. C.; Neill, C.; Dorst, J. P.: The scoliosis of congenital heart disease. *Am. Heart J.* 84: 463–469, 1972.
- [77210] 15540. Ogawa, Y.; Itoh, H.; Yoshitake, Y.; Inoue, M.; Yoshimasa, T.; Serikawa, T.; Nakao, K.: Molecular cloning and chromosomal assignment of the mouse C-type natriuretic peptide (CNP) gene (Nppc): comparison with the human CNP gene (NPPC). *Genomics* 24: 383–387, 1994.
- [77211] 15541. Reckles, L. N.; Peterson, H. A.; Weidman, W. H.; Bianco, A. J., Jr. The association of scoliosis and congenital heart disease. *J. Bone Joint Surg.* 57A: 449–455, 1975.
- [77212] 15542. Steinhilber, M. E.: Structure, expression, and genomic mapping of the mouse natriuretic peptide type-B gene. *Circ. Res.* 72: 984–992, 1993.

- [77213] 15543. Tamura, N.; Ogawa, Y.; Chusho, H.; Nakamura, K.; Nakao, K.; Suda, M.; Kasahara, M.; Hashimoto, R.; Katsura, G.; Mukoyama, M.; Itoh, H.; Saito, Y.; Tanaka, I.; Otani, H.; Katsuki, M.; Nakao, K.: Cardiac fibrosis in mice lacking brain natriuretic peptide. *Proc. Nat. Acad. Sci.* 97: 4239–4244, 2000.
- [77214] 15544. Chawengsaksophak, K.; James, R.; Hammond, V. E.; Kontgen, F.; Beck, F.: Homeosis and intestinal tumours in *Cdx2* mutant mice. *Nature* 386:84–87, 1997.
- [77215] 15545. Drummond, F.; Putt, W.; Fox, M.; Edwards, Y. H.: Cloning and chromosome assignment of the human *CDX2* gene. *Ann. Hum. Genet.* 61: 393–400, 1997.
- [77216] 15546. German, M. S.; Wang, J.; Fernald, A. A.; Espinosa, R., III; LeBeau, M. M.; Bell, G. I.: Localization of the genes encoding two transcription factors, *LMX1* and *CDX3*, regulating insulin gene expression to human chromosomes 1 and 13. *Genomics* 24: 403–404, 1994.
- [77217] 15547. Woodford-Richens, K. L.; Halford, S.; Rowan, A.; Bevan, S.; Aaltonen, L. A.; Wasan, H.; Bicknell, D.; Bodmer, W. F.; Houlston, R. S.; Tomlinson, I. P. M.: *CDX2* mutations do not account for juvenile polyposis or Peutz–Jeghers syndrome and occur infrequently in sporadic colorectal cancers. *Brit. J. Cancer* 84: 1314–1316, 2001.

- [77218] 15548.Dubbink, H. J.; Verkaik, N. S.; Faber, P. W.; Trapman, J.; Schroder, F. H.; Romijn, J. C.: Tissue-specific and androgen-regulated expression of human prostate-specific transglutaminase. *Biochem. J.* 315: 901–908, 1996.
- [77219] 15549.Gentile, V.; Grant, F. J.; Porta, R.; Baldini, A.: Localization of the human prostate transglutaminase (type IV) gene (TGM4) to chromosome 3p21.33–p22 by fluorescence in situ hybridization. *Genomics* 27:219–220, 1995.
- [77220] 15550.Grant, F. J.; Taylor, D. A.; Sheppard, P. O.; Mathewes, S. L.; Lint, W.; Vanaja, E.; Bishop, P. D.; O'Hara, P. J.: Molecular cloning and characterization of a novel transglutaminase cDNA from a human prostate cDNA library. *Biochem. Biophys. Res. Commun.* 203: 1117–1123, 1994.
- [77221] 15551.Barker, H. M.; Brewis, N. D.; Street, A. J.; Spurr, N. K.; Cohen, P. T. W.: Three genes for protein phosphatase 1 map to different human chromosomes: sequence, expression and gene localisation of protein serine/threonine phosphatase 1 beta (PPP1CB). *Biochim. Biophys. Acta* 1220:212–218, 1994.
- [77222] 15552.Saadat, M.; Kakinoki, Y.; Mizuno, Y.; Kikuchi, K.; Yoshida, M.C.: Chromosomal localization of human, rat, and mouse protein phosphatase type 1 beta catalytic subunit genes (PPP1CB) by fluorescence in situ hybridization.

Jpn. J. Genet. 69: 697–700, 1994.

- [77223] 15553.King, L. S.; Nielsen, S.; Agre, P.: Aquaporins and the respiratory system: advice for a lung investigator. J. Clin. Invest. 105: 15–16, 2000.
- [77224] 15554.Lee, M. D.; Bhakta, K. Y.; Raina, S.; Yonescu, R.; Griffin, C.A.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Preston, G. M.; Agre, P.: The human aquaporin–5 gene: molecular characterization and chromosomal localization. J. Biol. Chem. 271: 8599–8604, 1996.
- [77225] 15555.Ma, T.; Fukuda, N.; Song, Y.; Matthay, M. A.; Verkman, A. S.: Lung fluid transport in aquaporin–5 knockout mice. J. Clin. Invest. 105: 93–100, 2000.
- [77226] 15556.Raina, S.; Preston, G. M.; Guggino, W. B.; Agre, P.: Molecular cloning and characterization of an aquaporin cDNA from salivary, lacrimal, and respiratory tissues. J. Biol. Chem. 270: 1908–1912, 1995.
- [77227] 15557.Padilla, C. A.; Bajalica, S.; Lagercrantz, J.; Holmgren, A.: The gene for human glutaredoxin (GLRX) is localized to human chromosome 5q14. Genomics 32: 455–457, 1996.
- [77228] 15558.Padilla, C. A.; Martinez–Galisteo, E.; Barcena, J. A.; Spyrou, G.; Holmgren, A.: Purification from placenta, amino acid sequence, structure comparisons and cDNA



cloning of human glutaredoxin. *Europ.J. Biochem.* 227: 27–34, 1995.

- [77229] 15559.Raghavachari, N.; Krysan, K.; Xing K.; Lou, M. F.: Regulation of thioltransferase expression in human lens epithelial cells. *Invest.Ophthal. Vis. Sci.* 42: 1002–1008, 2001.
- [77230] 15560.Corset, V.; Nguyen–Ba–Charvet, K. T.; Forcet, C.; Moyse, E.; Chedotal,A.; Mehlen, P.: Netrin–1–mediated axon outgrowth and cAMP production requires interaction with adenosine A2b receptor. *Nature* 407: 747–750,2000.
- [77231] 15561.Mino, R. P.; Spoerri, P. E.; Caballero, S.; Player, D.; Belardinelli,L.; Biaggioni, I.; Grant, M. B.: Adenosine receptor antagonists and retinal neovascularization in vivo. *Invest. Ophthal. Vis. Sci.* 42:3320–3324, 2001.
- [77232] 15562.Stein, E.; Zou, Y.; Poo, M.; Tessier–Lavigne, M.: Binding of DCC by netrin–1 to mediate axon guidance independent of adenosine A2B receptor activation. *Science* 291: 1976–1982, 2001.
- [77233] 15563.Townsend–Nicholson, A.; Baker, E.; Sutherland, G. R.; Schofield,P. R.: Localization of the adenosine A2b receptor subtype gene (ADORA2B) to chromosome 17p11.2–p12 by FISH and PCR screening of somatic cell–hybrids. *Genomics* 25: 605–607, 1995.

- [77234] 15564.von Gall, C.; Garabette, M. L.; Kell, C. A.; Frenzel, S.; Dehghani, F.; Schumm–Draeger, P.–M.; Weaver, D. R.; Korf, H.–W.; Hastings, M.H.; Stehle, J. H.: Rhythmic gene expression in pituitary depends on heterologous sensitization by the neurohormone melatonin. *Nature Neurosci.* 5: 234–238, 2002.
- [77235] 15565.Becq, F.; Hamon, Y.; Bajetto, A.; Gola, M.; Verrier, B.; Chimini, G.: ABC1, an ATP binding cassette transporter required for phagocytosis of apoptotic cells, generates a regulated anion flux after expression in *Xenopus laevis* oocytes. *J. Biol. Chem.* 272: 2695–2699, 1997.
- [77236] 15566.Decottignies, A.; Goffeau, A.: Complete inventory of the yeast ABC proteins. *Nature Genet.* 15: 137–145, 1997.
- [77237] 15567.Guo, Z.; Inazu, A.; Yu, W.; Suzumura, T.; Okamoto, M.; Nohara, A.; Higashikata, T.; Sano, R.; Wakasugi, K.; Hayakawa, T.; Yoshida, K.; Suehiro, T.; Schmitz, G.; Mabuchi, H.: Double deletions and missense mutations in the first nucleotide-binding fold of the ATP-binding cassette transporter A1 (ABCA1) gene in Japanese patients with Tangier disease. *J. Hum. Genet.* 47: 325–329, 2002.
- [77238] 15568.Hong, S. H.; Rhyne, J.; Zeller, K.; Miller, M.: Novel

ABCA1 compound variant associated with HDL cholesterol deficiency. *Biochim. Biophys. Acta* 1587: 60–64, 2002.

[77239] 15569. Huang, W.; Moriyama, K.; Koga, T.; Hua, H.; Ageta, M.; Kawabata, S.; Mawatari, K.; Imamura, T.; Eto, T.; Kawamura, M.; Teramoto, T.; Sasaki, J.: Novel mutations in ABCA1 gene in Japanese patients with Tangier disease and familial high density lipoprotein deficiency with coronary heart disease. *Biochim. Biophys. Acta* 1537: 71–78, 2001.

[77240] 15570. Ishii, J.; Nagano, M.; Kujiraoka, T.; Ishihara, M.; Egashira, T.; Takada, D.; Tsuji, M.; Hattori, H.; Emi, M.: Clinical variant of Tangier disease in Japan: mutation of the ABCA1 gene in hypoalphalipoproteinemia with corneal lipidoses. *J. Hum. Genet.* 47: 366–369, 2002.

[77241] 15571. Jennings, M. W.; Jones, R. W.; Wood, W. G.; Weatherall, D. J.: Analysis of an inversion within the human beta globin gene cluster. *Nucleic Acids Res.* 13: 2897–2906, 1985.

[77242] 15572. Kulozik, A. E.; Bellan-Koch, A.; Kohne, E.; Kleihauer, E.: A deletion/inversion rearrangement of the beta-globin gene cluster in a Turkish family with delta-beta(0)-thalassemia intermedia. *Blood* 79: 2455–2459, 1992.

[77243] 15573. Langmann, T.; Klucken, J.; Reil, M.; Liebisch, G.;

Luciani, M.-F.; Chimini, G.; Kaminski, W. E.; Schmitz, G.: Molecular cloning of the human ATP-binding cassette transporter 1 (hABC1): evidence for sterol-dependent regulation in macrophages. *Biochem. Biophys. Res. Commun.* 257: 29–33, 1999.

[77244] 15574. Lawn, R. M.; Wade, D. P.; Garvin, M. R.; Wang, X.; Schwartz, K.; Porter, J. G.; Seilhamer, J. J.; Vaughan, A. M.; Oram, J. F.: The Tangier disease gene product ABC1 controls the cellular apolipoprotein-mediated lipid removal pathway. *J. Clin. Invest.* 104: R25–R31, 1999.

[77245] 15575. Luciani, M. F.; Denizot, F.; Savary, S.; Mattei, M. G.; Chimini, G.: Cloning of two novel ABC transporters mapping on human chromosome 9. *Genomics* 21: 150–159, 1994.

[77246] 15576. Lapicka-Bodzioch, K.; Bodzioch, M.; Krull, M.; Kielar, D.; Probst, M.; Kiec, B.; Andrikovics, H.; Bottcher, A.; Hubacek, J.; Aslanidis, C.; Suttorp, N.; Schmitz, G.: Homogeneous assay based on 52 primer sets to scan for mutations of the ABCA1 gene and its application in genetic analysis of a new patient with familial high-density lipoprotein deficiency syndrome. *Biochim. Biophys. Acta* 1537: 42–48, 2001.

[77247] 15577. Marcil, M.; Boucher, B.; Krimbou, L.; Solymoss, B.

C.; Davignon, J.; Frohlich, J.; Genest, J., Jr.: Severe familial HDL deficiency in French-Canadian kindreds: clinical, biochemical, and molecular characterization. *Arterioscler. Thromb. Vasc. Biol.* 15: 1015-1024, 1995.

[77248] 15578. Marcil, M.; Yu, L.; Krimbou, L.; Boucher, B.; Oram, J. F.; Cohn, J. S.; Genest, J., Jr.: Cellular cholesterol transport and efflux in fibroblasts are abnormal in subjects with familial HDL deficiency. *Arterioscler. Thromb. Vasc. Biol.* 19: 159-169, 1999.

[77249] 15579. McNeish, J.; Aiello, R. J.; Guyot, D.; Turi, T.; Gabel, C.; Aldinger, C.; Hoppe, K. L.; Roach, M. L.; Royer, L. J.; de Wet, J.; Broccardo, C.; Chimini, G.; Francone, O. L.: High density lipoprotein deficiency and foam cell accumulation in mice with targeted disruption of ATP-binding cassette transporter-1. *Proc. Nat. Acad. Sci.* 97: 4245-4250, 2000.

[77250] 15580. Pullinger, C. R.; Hakamata, H.; Duchateau, P. N.; Eng, C.; Aouizerat, B. E.; Cho, M. H.; Fielding, C. J.; Kane, J. P.: Analysis of hABC1 gene 5-prime end: additional peptide sequence, promoter region, and four polymorphisms. *Biochem. Biophys. Res. Commun.* 271: 451-455, 2000.

[77251] 15581. Petit, M. M. R.; Schoenmakers, E. F. P. M.; Huysmans, C.; Geurts, J. M. W.; Mandahl, N.; Van de Ven, W. J. M.: LHFP, a novel translocation partner gene of HMGIC in a

lipoma, is a member of a new family of LHFP-like genes.

Genomics 57: 438–441, 1999.

[77252] 15582. Nishita, Y.; Yoshida, I.; Sado, T.; Takagi, N.: Genomic imprinting and chromosomal localization of the human MEST gene. Genomics 36:539–542, 1996.

[77253] 15583. Kohen, R.; Metcalf, M. A.; Khan, N.; Druck, T.; Huebner, K.; Lachowicz, J. E.; Meltzer, H. Y.; Sibley, D. R.; Roth, B. L.; Hamblin, M. W.: Cloning, characterization, and chromosomal localization of a human 5-HT-6 serotonin receptor. J. Neurochem. 66: 47–56, 1996.

[77254] 15584. Frank, J.; Aita, V. M.; Ahmad, W.; Lam, H.; Wolff, C.; Christiano, A. M.: Identification of a founder mutation in the protoporphyrinogen oxidase gene in variegate porphyria patients from Chile. Hum. Hered. 51:160–168, 2001.

[77255] 15585. Frank, J.; Jugert, F. K.; Breitkopf, C.; Goerz, G.; Merk, H. F.; Christiano, A. M.: Recurrent missense mutation in the protoporphyrinogen oxidase gene underlies variegate porphyria. Am. J. Med. Genet. 79:22–26, 1998.

[77256] 15586. Orso, E.; Broccardo, C.; Kaminski, W. E.; Bottcher, A.; Liebisch, G.; Drobnik, W.; Gotz, A.; Chambenoit, O.; Diederich, W.; Langmann, T.; Spruss, T.; Luciani, M.-F.; Rothe, G.; Lackner, K. J.; Chimini, G.; Schmitz, G.: Trans-

port of lipids from Golgi to plasma membrane is defective in Tangier disease patients and Abc1-deficient mice. *Nature Genet.* 24: 192–196, 2000.

[77257] 15587. Hift, R. J.; Meissner, P. N.; Corrigall, A. V.; Ziman, M. R.; Petersen, L. A.; Meissner, D. M.; Davidson, B. P.; Sutherland, J.; Dailey, H. A.; Kirsch, R. E.: Variegate porphyria in South Africa, 1688–1996: new developments in an old disease. *S. Afr. Med. J.* 87: 722–731, 1997.

[77258] 15588. Hift, R. J.; Meissner, P. N.; Todd, G.; Kirby, P.; Bilsland, D.; Collins, P.; Ferguson, J.; Moore, M. R.: Homozygous variegate porphyria: an evolving clinical syndrome. *Postgrad. Med. J.* 69: 781–786, 1993.

[77259] 15589. Lam, H.; Dragan, L.; Tsou, H. C.; Merk, H.; Peacocke, M.; Goerz, G.; Sassa, S.; Poh-Fitzpatrick, M.; Bickers, D. R.; Christiano, A. M.: Molecular basis of variegate porphyria: a de novo insertion mutation in the protoporphyrinogen oxidase gene. *Hum. Genet.* 99: 126–129, 1997.

[77260] 15590. Puy, H.; Robreau, A.-M.; Rosipal, R.; Nordmann, Y.; Deybach, J.-C.: Protoporphyrinogen oxidase: complete genomic sequence and polymorphisms in the human gene. *Biochem. Biophys. Res. Comm.* 226: 226–230, 1996.

[77261] 15591. Roberts, A. G.; Puy, H.; Dailey, T. A.; Morgan, R. R.;

Whatley,S. D.; Dailey, H. A.; Martasek, P.; Nordmann, Y.; Deybach, J.-C.;Elder, G. H.: Molecular characterization of homozygous variegateporphyria. Hum. Molec. Genet. 7: 1921–1925, 1998.

[77262] 15592.Honda, H.; Inazawa, J.; Nishida, J.; Yazaki, Y.; Hirai, H.: Molecularcloning, characterization, and chromosomal localization of a novelprotein–tyrosine phosphatase, HPTP eta. Blood 84: 4186–4194, 1994.

[77263] 15593.Ostman, A.; Yang, Q.; Tonks, N. K.: Expression of DEP–1, a receptor–likeprotein–tyrosine–phosphatase, is enhanced with increasing cell density. Proc.Nat. Acad. Sci. 91: 9680–9684, 1994.

[77264] 15594.Katoh, M.; Hirai, M.; Sugimura, T.; Terada, M.: Cloning and characterizationof MST, a novel (putative) serine/threonine kinase with SH3 domain. Oncogene 10:1447–1451, 1995.

[77265] 15595.Morgan, H. D.; Sutherland, H. G. E.; Martin, D. I. K.; Whitelaw,E.: Epigenetic inheritance at the agouti locus in the mouse. NatureGenet. 23: 314–318, 1999.

[77266] 15596.Jacoby, A. S.; Webb, G. C.; Liu, M. L.; Kofler, B.; Hort, Y. J.;Fathi, Z.; Bottema, C. D. K.; Shine, J.; Iismaa, T. P.: Structuralorganization of the mouse and human GALR1 galanin receptor genes (Galnrand GALNR) and chromoso–



mal localization of the mouse gene. Genomics  
45:496–508, 1997.

- [77267] 15597. Tomoda, T.; Kurashige, T.; Moriki, T.; Yamamoto, H.; Fujimoto, S.; Taniguchi, T.: Enhanced expression of poly(ADP-ribose) synthetase gene in malignant lymphoma. Am. J. Hemat. 37: 223–227, 1991.
- [77268] 15598. Vasquez, K. M.; Marburger, K.; Intody, Z.; Wilson, J. H.: Manipulating the mammalian genome by homologous recombination. Proc. Nat. Acad. Sci. 98: 8403–8410, 2001.
- [77269] 15599. Yu, S.-W.; Wang, H.; Poitras, M. F.; Coombs, C.; Bowers, W. J.; Federoff, H. J.; Poirier, G. G.; Dawson, T. M.; Dawson, V. L.: Mediation of poly(ADP-ribose) polymerase-1-dependent cell death by apoptosis-inducing factor. Science 297: 259–263, 2002.
- [77270] 15600. Zabel, B. U.; Herzog, H.; Schneider, R.; Auer, B.; Hirsch-Kauffmann, M.; Schweiger, M.: Chromosomal sublocalization of the gene for human poly (ADP-ribose) polymerase (NAD<sup>+</sup> ADP-ribosyltransferase) at 1q41–42. (Abstract) Cytogenet. Cell Genet. 51: 1115, 1989.
- [77271] 15601. Hood, L.; Kronenberg, M.; Hunkapiller, T.: T cell antigen receptors and the immunoglobulin supergene family. Cell 40: 225–229, 1985.

- [77272] 15602.de Villiers, J. N. P.; Hillermann, R.; Loubser, L.; Kotze, M. J.: Spectrum of mutations in the HFE gene implicated in haemochromatosis and porphyria. *Hum. Molec. Genet.* 8: 1517–1522, 1999.
- [77273] 15603.Deybach, J.-C.; Puy, H.; Robreau, A.-M.; Lamoril, J.; Da Silva, V.; Grandchamp, B.; Nordmann, Y.: Mutations in the protoporphyrinogen oxidase gene in patients with variegate porphyria. *Hum. Molec. Genet.* 5:407–410, 1996.
- [77274] 15604.Meissner, P. N.; Dailey, T. A.; Hift, R. J.; Ziman, M.; Corrigall, A. V.; Roberts, A. G.; Meissner, D. M.; Kirsch, R. E.; Dailey, H.A.: A R59W mutation in human protoporphyrinogen oxidase results in decreased enzyme activity and is prevalent in South Africans with variegate porphyria. *Nature Genet.* 13: 95–97, 1996.
- [77275] 15605.Kalman, K.; Nguyen, A.; Tseng-Crank, J.; Dukes, I. D.; Chandy, G.; Hustad, C. M.; Copeland, N. G.; Jenkins, N. A.; Mohrenweiser, H.; Brandriff, B.; Cahalan, M.; Gutman, G. A.; Chandy, K. G.: Genomic organization, chromosomal localization, tissue distribution, and biophysical characterization of a novel mammalian Shaker-related voltage-gated potassium channel, Kv1.7. *J. Biol. Chem.* 273: 5851–5857, 1998.
- [77276] 15606.Stewart, S. L.; Rosenberg, H.; Fletcher, J. E.: Failure

to identify the ryanodine receptor G1021A mutation in a large North American population with malignant hyperthermia. Clin. Genet. 54: 358–361, 1998.

[77277] 15607. Orkiszewski, K. G.; Tedesco, T. A.; Mellman, W. J.; Croce, C. M.: Linkage relationship between the genes for thymidine kinase and galactokinase in different primates. Somat. Cell Genet. 2: 21–26, 1976.

[77278] 15608. Dunham, I.; Collins, J.; Wadey, R.; Scambler, P.: Possible role for COMT in psychosis associated with velocardio-facial syndrome. (Letter) Lancet 340: 1361–1362, 1992.

[77279] 15609. Pulkka, A.; Ihalainen, R.; Suorsa, A.; Riviere, M.; Szpirer, J.; Pajunen, A.: Structures and chromosomal localizations of two rat genes encoding S-adenosylmethionine decarboxylase. Genomics 16: 342–349, 1993.

[77280] 15610. Radford, D. M.; Eddy, R.; Haley, L.; Henry, W. M.; Pegg, A. E.; Pajunen, A.; Shows, T. B.: Gene sequences coding for S-adenosylmethionine decarboxylase are present on human chromosome 6 and the X and are not amplified in colon neoplasia. Cytogenet. Cell Genet. 49: 285–288, 1989.

[77281] 15611. Grignani, F.; De Matteis, S.; Nervi, C.; Tomassoni, L.; Gelmetti, V.; Cioce, M.; Fanelli, M.; Ruthardt, M.; Fer-

rara, F. F.; Zamir, I.; Seiser, C.; Grignani, F.; Lazar, M. A.; Minucci, S.; Pelicci, P. G.: Fusion proteins of the retinoic acid receptor- $\alpha$  recruit histone deacetylase in promyelocytic leukaemia. *Nature* 391: 815–818, 1998.

[77282] 15612. Lin, R. J.; Nagy, L.; Inoue, S.; Shao, W.; Miller, W. H., Jr.; Evans, R. M.: Role of the histone deacetylase complex in acute promyelocytic leukaemia. *Nature* 391: 811–814, 1998.

[77283] 15613. Mendez, R.; Hake, L. E.; Andresson, T.; Littlepage, L. E.; Ruderman, J. V.; Richter, J. D.: Phosphorylation of CPE binding factor by Eg2 regulates translation of c-mos mRNA. *Nature* 404: 302–307, 2000.

[77284] 15614. Ahrendt, S. A.; Decker, P. A.; Alawi, E. A.; Zhu, Y.; Sanchez-Cespedes, M.; Yang, S. C.; Haasler, G. B.; Kadjacsy-Balla, A.; Demeure, M. J.; Sidransky, D.: Cigarette smoking is strongly associated with mutation of the K-ras gene in patients with primary adenocarcinoma of the lung. *Cancer* 92: 1525–1530, 2001.

[77285] 15615. Almoguera, C.; Shibata, D.; Forrester, K.; Martin, J.; Arnheim, N.; Perucho, M.: Most human carcinomas of the exocrine pancreas contain mutant c-K-ras genes. *Cell* 53: 549–554, 1988.

[77286] 15616. Andreyev, H. J. N.; Tilsed, J. V. T.; Cunningham, D.;

Sampson, S. A.; Norman, A. R.; Schneider, H. J.; Clarke, P. A.: K-ras mutations in patients with early colorectal cancers. *Gut* 41: 323–329, 1997.

[77287] 15617. Bollag, G.; Adler, F.; elMasry, N.; McCabe, P. C.; Connor, E., Jr.; Thompson, P.; McCormick, F.; Shannon, K.: Biochemical characterization of a novel KRAS insertion mutation from a human leukemia. *J. Biol. Chem.* 271: 32491–32494, 1996.

[77288] 15618. Burmer, G. C.; Loeb, L. A.: Mutations in the KRAS2 oncogene during progressive stages of human colon carcinoma. *Proc. Nat. Acad. Sci.* 86: 2403–2407, 1989.

[77289] 15619. Capon, D. J.; Seeburg, P. H.; McGrath, J. P.; Hayflick, J. S.; Edman, U.; Levinson, A. D.; Goeddel, D. V.: Activation of Ki-ras2 gene in human colon and lung carcinomas by two different point mutations. *Nature* 304: 507–513, 1983.

[77290] 15620. Cubilla, A. L.; Fitzgerald, P. J.: *Cancer Res.* 36: 2690–2698, 1976.

[77291] 15621. Der, C. J.; Cooper, G. M.: Altered gene products are associated with activation of cellular ras-k genes in human lung and colon carcinomas. *Cell* 32: 201–208, 1983.

[77292] 15622. Feig, L. A.; Bast, R. C., Jr.; Knapp, R. C.; Cooper, G. M.: Somatic activation of ras-K gene in a human ovarian

carcinoma. *Science* 223:698–701, 1984.

- [77293] 15623.Grimmond, S. M.; Raghavan, D.; Russell, P. J.: Detection of a rare point mutation in Ki-ras of a human bladder cancer xenograft by polymerase chain reaction and direct sequencing. *Urol. Res.* 20:121–126, 1992.
- [77294] 15624.Hayashi, N.; Sugai, S.; Ito, I.; Nakamori, S.; Ogawa, M.; Nakamura, Y.: Ethnic difference in the pattern of K-ras oncogene mutations in human colorectal cancers. *Hum. Mutat.* 8: 258–261, 1996.
- [77295] 15625.Tegazzin, V.; Accorsi, A.; Gritti, G.; Arcelli, L.; Di Giovanni, A.: MH fulminant reaction after eight anaesthetics: a case report. *Minerva Anesthesiol.* 60: 217–219, 1994.
- [77296] 15626.Tilgen, N.; Zorzato, F.; Halliger-Keller, B.; Muntoni, F.; Sewry, C.; Palumucci, L. M.; Schneider, C.; Hauser, E.; Lehmann-Horn, F.; Muller, C. R.; Treves, S.: Identification of four novel mutations in the C-terminal membrane spanning domain of the ryanodine receptor1: association with central core disease and alteration of calcium homeostasis. *Hum. Molec. Genet.* 10: 2879–2887, 2001.
- [77297] 15627.Tobin, J. R.; Jason, D. R.; Challa, V. R.; Nelson, T. E.; Sambuughin, N.: Malignant hyperthermia and apparent heat stroke. (Letter) *JAMA* 286:168–169, 2001.

- [77298] 15628.Wang, S.-Q.; Song, L.-S.; Lakatta, E. G.; Cheng, H.: Ca(2+) signalling between single L-type Ca(2+) channels and ryanodine receptors in heartcells. *Nature* 410: 592–596, 2001.
- [77299] 15629.Zhang, Y.; Chen, H. S.; Khanna, V. K.; De Leon, S.; Phillips, M. S.; Schappert, K.; Britt, B. A.; Brownell, A. K. W.; MacLennan, D. H.: A mutation in the human ryanodine receptor gene associated with central core disease. *Nature Genet.* 5: 46–50, 1993.
- [77300] 15630.Zorzato, F.; Fujii, J.; Otsu, K.; Phillips, M.; Green, N. M.; Lai, F. A.; Meissner, G.; MacLennan, D. H.: Molecular cloning of cDNA encoding human and rabbit forms of the Ca(2+) release channel (ryanodine receptor) of skeletal muscle sarcoplasmic reticulum. *J. Biol. Chem.* 265: 2244–2256, 1990.
- [77301] 15631.Marx, S. O.; Reiken, S.; Hisamatsu, Y.; Jayaraman, T.; Burkhoff, D.; Rosemlit, N.; Marks, A. R.: PKA phosphorylation dissociates FKBP12.6 from the calcium release channel (ryanodine receptor): defective regulation in failing hearts. *Cell* 101: 365–376, 2000.
- [77302] 15632.Mattei, M. G.; Giannini, G.; Moscatelli, F.; Sorrentino, V.: Chromosomal localization of murine ryanodine receptor genes RYR1, RYR2, and RYR3 by in situ hybridiza-

tion. *Genomics* 22: 202–204, 1994.

- [77303] 15633. Bertocchini, F.; Ovitt, C. E.; Conti, A.; Barone, V.; Scholer, H. R.; Bottinelli, R.; Reggiani, C.; Sorrentino, V.: Requirement for the ryanodine receptor type 3 for efficient contraction in neonatal skeletal muscles. *EMBO J.* 16: 6956–6963, 1997.
- [77304] 15634. Futatsugi, A.; Kato, K.; Ogura, H.; Li, S.-T.; Nagata, E.; Kuwajima, G.; Tanaka, K.; Itohara, S.; Mikoshiba, K.: Facilitation of NMDAR-independent LTP and spatial learning in mutant mice lacking ryanodine receptor type 3. *Neuron* 24: 701–713, 1999.
- [77305] 15635. Hakamata, Y.; Nakai, J.; Takeshima, H.; Imoto, K.: Primary structure and distribution of a novel ryanodine receptor/calcium release channel from rabbit brain. *FEBS Lett.* 312: 229–235, 1992.
- [77306] 15636. Leeb, T.; Brenig, B.: cDNA cloning and sequencing of the human ryanodine receptor type 3 (RYR3) reveals a novel alternative splice site in the RYR3 gene. *FEBS Lett.* 423: 367–370, 1998.
- [77307] 15637. Sorrentino, V.; Giannini, G.; Malzac, P.; Mattei, M. G.: Localization of a novel ryanodine receptor gene (RYR3) to human chromosome 15q14–q15 by in situ hybridization. *Genomics* 18: 163–165, 1993.



- [77308] 15638. Takeshima, H.; Ikemoto, T.; Nishi, M.; Nishiyama, N.; Shimuta, M.; Sugitani, Y.; Kuno, J.; Saito, I.; Saito, H.; Endo, M.; Iino, M.; Noda, T.: Generation and characterization of mutant mice lacking gamma-irradiation receptor type 3. *J. Biol. Chem.* 271: 19649–19652, 1996.
- [77309] 15639. Bigler, R. D.; Bushkin, Y.; Chiorazzi, N.: S152 (CD27): a modulating disulfide-linked T cell activation antigen. *J. Immun.* 141: 21–28, 1988.
- [77310] 15640. Hendriks, J.; Gravestien, L. A.; Tesselaar, K.; van Lier, R. A. W.; Schumacher, T. N. M.; Borst, J.: CD27 is required for generation and long-term maintenance of T cell immunity. *Nature Immun.* 1: 433–440, 2000.
- [77311] 15641. Prasad, K. V. S.; Ao, Z.; Yoon, Y.; Wu, M. X.; Rizk, M.; Jacquot, S.; Schlossman, S. F.: CD27, a member of the tumor necrosis factor receptor family, induces apoptosis and binds to Siva, a proapoptotic protein. *Proc. Nat. Acad. Sci.* 94: 6346–6351, 1997.
- [77312] 15642. Arnaiz-Villena, A.; Perez-Aciego, P.; Ballester, C.; Sotelo, T.; Perez-Seoane, C.; Martin-Villa, J. M.; Regueiro, J. R.: Biochemical basis of a novel T lymphocyte receptor immunodeficiency by immunohistochemistry: a possible CD3-gamma abnormality. *Lab. Invest.* 64: 675–681, 1991.

- [77313] 15643.Arnaiz-Villena, A.; Timon, M.; Corell, A.; Perez-Aciego, P.; Martin-Villa, J. M.; Regueiro, J. R.: Primary immunodeficiency caused by mutations in the gene encoding the CD3-gamma subunit of the T-lymphocyte receptor. *New Eng. J. Med.* 327: 529-533, 1992.
- [77314] 15644.Das, S.; Cotter, F. E.; Gibbons, B.; Dhut, S.; Young, B. D.: CD3G is within 200 kb of the leukemic t(4;11) translocation breakpoint. *Genes Chromosomes Cancer* 3: 44-47, 1991.
- [77315] 15645.Evans, G. A.; Lewis, K. A.; Lawless, G. M.: Molecular organization of the human CD3 gene family on chromosome 11q23. *Immunogenetics* 28:365-373, 1988.
- [77316] 15646.Krissansen, G. W.; Gorman, P. A.; Kozak, C. A.; Spurr, N. K.; Sheer, D.; Goodfellow, P. N.; Crumpton, M. J.: Chromosomal locations of the gene coding for the CD3 (T3) gamma-subunit of the human and mouse CD3/T-cell antigen receptor complexes. *Immunogenetics* 26: 258-266, 1987.
- [77317] 15647.Krissansen, G. W.; Owen, M. J.; Verbi, W.; Crumpton, M. J.: Primary structure of the T3 gamma subunit of the T3/T cell antigen receptor complex deduced from cDNA sequences: evolution of the T3 gamma and delta subunits. *EMBO J.* 5: 1799-1808, 1986.

- [77318] 15648.Saito, H.; Koyama, T.; Georgopoulos, K.; Clevers, H.; Haser, W.G.; LeBien, T.; Tonegawa, S.; Terhorst, C.: Close linkage of the mouse and human CD3 gamma- and delta-chain genes suggests that their transcription is controlled by common regulatory elements. *Proc.Nat. Acad. Sci.* 84: 9131–9134, 1987.
- [77319] 15649.Tunnacliffe, A.; Buluwela, L.; Rabbitts, T. H.: Physical linkage of three CD3 genes on human chromosome 11. *EMBO J.* 6: 2953–2957, 1987.
- [77320] 15650.Tunnacliffe, A.; McGuire, R. S.: A physical linkage group in human chromosome band 11q23 covering a region implicated in leukocyte neoplasia. *Genomics* 8:447–453, 1990.
- [77321] 15651.Tunnacliffe, A.; Olsson, C.; Buluwela, L.; Rabbitts, T. H.: Organization of the human CD3 locus on chromosome 11. *Europ. J. Immun.* 18: 1639–1642, 1988.
- [77322] 15652.Yunis, J. J.; Jones, C.; Madden, M. T.; Lu, D.; Mayer, M. G.: Gene order, amplification, and rearrangement of chromosome band 11q23 in hematologic malignancies. *Genomics* 5: 84–90, 1989.
- [77323] 15653.Park, S.-H.; Bendelac, A.: CD1-restricted T-cell responses and microbial infection. *Nature* 406: 788–792, 2000.

- [77324] 15654. Beckman, E. M.; Porcelli, S. A.; Morita, C. T.; Behar, S. M.; Furlong, S. T.; Brenner, M. B.: Recognition of a lipid antigen by CD1-restricted  $\alpha$ - $\beta$ (+) T cells. *Nature* 372: 691–694, 1994.
- [77325] 15655. Sieling, P. A.; Chatterjee, D.; Porcelli, S. A.; Prigozy, T. I.; Mazzaccaro, R. J.; Soriano, T.; Bloom, B. R.; Brenner, M. B.; Kronenberg, M.; Brennan, P. J.; Modlin, R. L.: CD1-restricted T cell recognition of microbial lipoglycan antigens. *Science* 269: 227–230, 1995.
- [77326] 15656. Sieling, P. A.; Jullien, D.; Dahlem, M.; Tedder, T. F.; Rea, T. H.; Modlin, R. L.; Porcelli, S. A.: CD1 expression by dendritic cells in human leprosy lesions: correlation with effective host immunity. *J. Immunol.* 162: 1851–1858, 1999.
- [77327] 15657. Nishimura, K.; Taketani, S.; Inokuchi, H.: Cloning of a human cDNA for protoporphyrinogen oxidase by complementation in vivo of a hemG mutant of *Escherichia coli*. *J. Biol. Chem.* 270: 8076–8080, 1995.
- [77328] 15658. Roberts, A. G.; Whatley, S. D.; Daniels, J.; Holmans, P.; Fenton, I.; Owen, M. J.; Thompson, P.; Long, C.; Elder, G. H.: Partial characterization and assignment of the gene for protoporphyrinogen oxidase and variegate porphyria to human chromosome 1q23. *Hum. Molec. Genet.* 4: 2387–2390, 1995.

- [77329] 15659.Taketani, S.; Inazawa, J.; Abe, T.; Furukawa, T.; Kohno, H.; Tokunaga,R.; Nishimura, K.; Inokuchi, H.: The human protoporphyrinogen oxidase gene (PPOX): organization and location to chromosome 1. *Genomics* 29:698–703, 1995.
- [77330] 15660.Warnich, L.; Kotze, M. J.; Groenewald, I. M.; Groenewald, J. Z.;van Brakel, M. G.; van Heerden, C. J.; de Villiers, J. N. P.; vande Ven, W. J. M.; Schoenmakers, E. F. P. M.; Taketani, S.; Retief,A. E.: Identification of three mutations and associated haplotypes in the protoporphyrinogen oxidase gene in South African families with variegate porphyria. *Hum. Molec. Genet.* 5: 981–984, 1996.
- [77331] 15661.Haas, M.; Ward, D. C.; Lee, J.; Roses, A. D.; Clarke, V.; D'Eustachio,P.; Lau, D.; Vega-Saenz de Miera, E.; Rudy, B.: Localization of Shaw-related K(+) channel genes on mouse and human chromosomes. *Mammalian Genome* 4:711–715, 1993.
- [77332] 15662.Albrecht, B.; Weber, K.; Pongs, O.: Characterization of a voltage-activated K-channel gene cluster on human chromosome 12p13. *Receptors Channels* 3:213–220, 1995.
- [77333] 15663.Grupe, A.; Schroter, K. H.; Ruppertsberg, J. P.; Stocker, M.; Drewes,T.; Beckh, S.; Pongs, O.: Cloning and

expression of a human voltage-gated potassium channel:  
a novel member of the RCK potassium channel family.  
EMBOJ. 9: 1749–1756, 1990.

- [77334] 15664.Klocke, R.; Roberds, S. L.; Tamkun, M. M.; Grone-  
meier, M.; Augustin, A.; Albrecht, B.; Pongs, O.; Jockusch,  
H.: Chromosomal mapping in the mouse of eight  
K(+)-channel genes representing the four Shaker-  
like subfamilies Shaker, Shab, Shaw, and Shal. Genomics  
18: 568–574, 1993.
- [77335] 15665.Grissmer, S.; Ghanshani, S.; Dethlefs, B.; McPherson,  
J. D.; Wasmuth, J. J.; Gutman, G. A.; Cahalan, M. D.;  
Chandy, K. G.: The Shaw-related potassium channel gene,  
Kv3.1, on human chromosome 11, encodes the type 1 K<sup>+</sup>  
channel in T cells. J. Biol. Chem. 267: 20971–20979,  
1992.
- [77336] 15666.Ried, T.; Rudy, B.; Vega-Saenz de Miera, E.; Lau, D.;  
Ward, D.C.; Sen, K.: Localization of a highly conserved hu-  
man potassium channel gene (NGK2-KV4; KCNC1) to chro-  
mosome 11p15. Genomics 15: 405–411, 1993.
- [77337] 15667.Stubbs, L.; Rinchik, E. M.; Goldberg, E.; Rudy, B.;  
Handel, M.A.; Johnson, D.: Clustering of six human 11p15  
gene homologs within a 500-kb interval of proximal  
mouse chromosome 7. Genomics 24: 324–332, 1994.

- [77338] 15668.Curran, M. E.; Landes, G. M.; Keating, M. T.: Molecular cloning, characterization, and genomic localization of a human potassium channel gene. *Genomics* 12: 729–737, 1992.
- [77339] 15669.Al-Chalabi, A.; Andersen, P. M.; Nilsson, P.; Chioza, B.; Andersson, J. L.; Russ, C.; Shaw, C. E.; Powell, J. F.; Leigh, P. N.: Deletion of the heavy neurofilament subunit tail in amyotrophic lateral sclerosis. *Hum. Molec. Genet.* 8: 157–164, 1999.
- [77340] 15670.Collard, J.-F.; Cote, F.; Julien, J.-P.: Defective axonal transport in a transgenic mouse model of amyotrophic lateral sclerosis. *Nature* 375:61–64, 1995.
- [77341] 15671.Figlewicz, D. A.; Krizus, A.; Martinoli, M. G.; Meininger, V.; Dib, M.; Rouleau, G. A.; Julien, J.-P.: Variants of the heavy neurofilament subunit are associated with the development of amyotrophic lateral sclerosis. *Hum. Molec. Genet.* 3: 1757–1761, 1994.
- [77342] 15672.Lees, J. F.; Shneiderman, P. S.; Skuntz, S. F.; Carden, M. J.; Lazzarini, R. A.: The structure and organization of the human heavy neurofilament subunit (NF-H) and the gene encoding it. *EMBO J.* 7: 1947–1955, 1988.
- [77343] 15673.Mattei, M.-G.; Dautigny, A.; Pham-Dinh, D.; Passage, E.; Mattei, J.-F.; Jolles, P.: The gene encoding the

large human neurofilament subunit (NF-H) maps to the q121-q131 region on human chromosome 22.

Hum.Genet. 80: 293-295, 1988.

[77344] 15674. Rooke, K.; Figlewicz, D. A.; Han, F.; Rouleau, G. A.: Analysis of the KSP repeat of the neurofilament heavy subunit in familial amyotrophic lateral sclerosis. Neurology 46: 789-790, 1996.

[77345] 15675. Rouleau, G. A.; Merel, P.; Lutchman, M.; Sanson, M.; Zucman, J.; Marineau, C.; Hoang-Xuan, K.; Demczuk, S.; Desmaze, C.; Plougastel, B.; Pulst, S. M.; Lenoir, G.; Bijlsma, E.; Fashold, R.; Dumanski, J.; de Jong, P.; Parry, D.; Eldridge, R.; Aurias, A.; Delattre, O.; Thomas, G.: Alteration in a new gene encoding a putative membrane-organizing protein causes neuro-fibromatosis type 2. Nature 363: 515-521, 1993.

[77346] 15676. Tomkins, J.; Usher, P.; Slade, J. Y.; Ince, P. G.; Curtis, A.; Bushby, K.; Shaw, P. J.: Novel insertion in the KSP region of the neurofilament heavy gene in amyotrophic lateral sclerosis. Neuroreport 9: 3967-3970, 1998.

[77347] 15677. Vechio, J. D.; Bruijn, L. I.; Xu, Z.; Brown, R. H., Jr.; Cleveland, D. W.: Sequence variants in human neurofilament proteins: absence of linkage to familial amyotrophic lateral sclerosis. Ann. Neurol. 40: 603-610, 1996.



- [77348] 15678. Watson, C. J.; Gaunt, L.; Evans, G.; Patel, K.; Harris, R.; Strachan, T.: A disease-associated germline deletion maps the type 2 neurofibromatosis (NF2) gene between the Ewing sarcoma region and the leukaemia inhibitory-factor locus. *Hum. Molec. Genet.* 2: 701–704, 1993.
- [77349] 15679. Levy, E.; Liem, R. K. H.; D'Eustachio, P.; Cowan, N. J.: Structure and evolutionary origin of the gene encoding NF-M, the middle-molecular-mass neurofilament protein. *Europ. J. Biochem.* 166: 71–77, 1987.
- [77350] 15680. Mersiyanova, I. V.; Perepelov, A. V.; Polyakov, A. V.; Sitnikov, V. F.; Dadali, E. L.; Oparin, R. B.; Petrin, A. N.; Evgrafov, O. V.: A new variant of Charcot-Marie-Tooth disease type 2 is probably the result of a mutation in the neurofilament-light gene. *Am. J. Hum. Genet.* 67: 37–46, 2000.
- [77351] 15681. Myers, M. W.; Lazzarini, R. A.; Lee, V. M.-Y.; Schlaepfer, W. W.; Nelson, D. L.: The human mid-size neurofilament subunit: a repeated protein sequence and the relationship of its gene to the intermediate filament gene family. *EMBO J.* 6: 1617–1626, 1987.
- [77352] 15682. Barker, P. E.; Besmer, P.; Ruddle, F. H.: Human c-kit oncogene on human chromosome 4. (Abstract) *Am. J. Hum. Genet.* 37: A143, 1985.

- [77353] 15683.Beghini, A.; Larizza, L.; Cairoli, R.; Morra, E.: c-kit activating mutations and mast cell proliferation in human leukemia. (Letter) *Blood* 92:701–702, 1998.
- [77354] 15684.Beghini, A.; Tibiletti, M.; Roversi, G.; Chiaravalli, A.; Serio, G.; Capella, C.; Larizza, L.: Germline mutation in the juxtamembrane domain of the KIT gene in a family with gastrointestinal stromal tumors and urticaria pigmentosa. *Cancer* 92: 657–662, 2001.
- [77355] 15685.Blume-Jensen, P.; Jiang, G.; Hyman, R.; Lee, K.-F.; O’Gorman, S.; Hunter, T.: Kit/stem cell factor receptor-induced activation of phosphatidylinositol 3-kinase is essential for male fertility. *Nature Genet.* 24:157–162, 2000.
- [77356] 15686.Bolognia, J. L.; Pawelek, J. M.: Biology of hypopigmentation. *J. Am. Acad. Derm.* 19: 217–255, 1988.
- [77357] 15687.Brannan, C. I.; Lyman, S. D.; Williams, D. E.; Eisenman, J.; Anderson, D. M.; Cosman, D.; Bedell, M. A.; Jenkins, N. A.; Copeland, N. G.: Steel-Dickie mutation encodes a c-kit ligand lacking transmembrane and cytoplasmic domains. *Proc. Nat. Acad. Sci.* 88: 4671–4674, 1991.
- [77358] 15688.Chabot, B.; Stephenson, D. A.; Chapman, V. M.; Besmer, P.; Bernstein, A.: The proto-oncogene c-kit encoding a transmembrane tyrosine kinase receptor maps to the

mouse W locus. *Nature* 335: 88–89, 1988.

[77359] 15689.d'Auriol, L.; Mattei, M.–G.; Andre, C.; Galibert, F.:

Localization of the human c-kit protooncogene on the q11–q12 region of chromosome 4. *Hum. Genet.* 78:

374–376, 1988.

[77360] 15690.De Miguel, M. P.; Cheng, L.; Holland, E. C.; Feder-

spiel, M. J.; Donovan, P. J.: Dissection of the c-Kit signaling pathway in mouse primordial germ cells by retroviral-mediated gene transfer. *Proc. Nat. Acad. Sci.* 99:

10458–10463, 2002.

[77361] 15691.Dubreuil, P.; Forrester, L.; Rottapel, R.; Reedijk, M.;

Fujita, J.; Bernstein, A.: The c-fms gene complements the mitogenic defect in mast cells derived from mutant W mice but not mi (microphthalmia) mice. *Proc. Nat. Acad. Sci.* 88:

2341–2345, 1991.

[77362] 15692.el-Omar, M.; Davies, J.; Gupta, S.; Ross, H.;

Thompson, R.: Leiomyosarcoma in leiomyomatosis of the small intestine. *Postgrad. Med. J.* 70:661–664, 1994.

[77363] 15693.Fleischman, R. A.: Human piebald trait resulting

from a dominant negative mutant allele of the c-kit membrane receptor gene. *J. Clin. Invest.* 89: 1713–1717, 1992.

[77364] 15694.Fleischman, R. A.; Saltman, D. L.; Stastny, V.;

Zneimer, S.: Deletion of the c-kit protooncogene in the

human developmental defect piebald trait. Proc. Nat. Acad. Sci. 88: 10885–10889, 1991.

[77365] 15695. Fritzsche-Polanz, R.; Jordan, J.-H.; Feix, A.; Sperr, W. R.; Sunder-Plassmann, G.; Valent, P.; Fodinger, M.: Mutation analysis of C-KIT in patients with myelodysplastic syndromes without mastocytosis and cases of systemic mastocytosis. Brit. J. Haemat. 113: 357–364, 2001.

[77366] 15696. Geissler, E. N.; Ryan, M. A.; Housman, D. E.: The dominant-whitespotting (W) locus of the mouse encodes the c-kit proto-oncogene. Cell 55: 185–192, 1988.

[77367] 15697. Giebel, L. B.; Spritz, R. A.: Mutation of the KIT (mast/stem cell growth factor receptor) protooncogene in human piebaldism. Proc. Nat. Acad. Sci. 88: 8696–8699, 1991.

[77368] 15698. Hirota, S.; Isozaki, K.; Moriyama, Y.; Hashimoto, K.; Nishida, T.; Ishiguro, S.; Kawano, K.; Hanada, M.; Kurata, A.; Takeda, M.; Tunio, G. M.; Matsuzawa, Y.; Kanakura, Y.; Shinomura, Y.; Kitamura, Y.: Gain-of-function mutations of c-kit in human gastrointestinal stromal tumors. Science 279: 577–580, 1998.

[77369] 15699. Huizinga, J. D.; Thuneberg, L.; Kluppel, M.; Malysz, J.; Mikkelsen, H. B.; Bernstein, A.: W/kit gene required for interstitial cells of Cajal and for intestinal pacemaker activ-

ity. *Nature* 373: 347–349,1995.

- [77370] 15700.Furitsu, T.; Tsujimura, T.; Tono, T.; Ikeda, H.; Kitayama, H.;Koshimizu, U.; Sugahara, H.; Butterfield, J. H.; Ashman, L. K.; Kanayama,Y.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Identification of mutations in the coding sequence of the proto-oncogene c-kit in a human mast cell leukemia cell line causing ligand-independent activation of c-kit product. *J. Clin. Invest.* 92: 1736–1744, 1993.
- [77371] 15701.Ingram, D. A.; Yang, F.-C.; Travers, J. B.; Wenning, M. J.; Hiatt,K.; New, S.; Hood, A.; Shannon, K.; Williams, D. A.; Clapp, D. W.: Genetic and biochemical evidence that haploinsufficiency of the Nf1 tumor suppressor gene modulates melanocyte and mast cell fates in vivo. *J. Exp. Med.* 191: 181–187, 2000.
- [77372] 15702.Isozaki, K.; Terris, B.; Belghiti, J.; Schiffmann, S.; Hirota,S.; Vanderwinden, J.-M.: Germline-activating mutation in the kinase domain of KIT gene in familial gastrointestinal stromal tumors. *Am.J. Path.* 157: 1581–1585, 2000.
- [77373] 15703.Joensuu, H.; Roberts, P. J.; Sarlomo-Rikala, M.; Andersson, L.C.; Tervahartiala, P.; Tuveson, D.; Silberman, S. L.; Capdeville,R.; Dimitrijevic, S.; Druker, B.; Demetri, G. D.: Effect of the tyrosine kinase inhibitor STI571 in a pa-

tient with a metastatic gastrointestinal stromal tumor. New Eng. J. Med. 344: 1052–1056, 2001.

- [77374] 15704. Johansson Moller, M.; Chaudhary, R.; Hellmen, E.; Hoyheim, B.; Chowdhary, B.; Andersson, L.: Pigs with the dominant white coat color phenotype carry a duplication of the KIT gene encoding the mast/stem cell growth factor receptor. Mammalian Genome 7: 822–830, 1996.
- [77375] 15705. Lasota, J.; Jasinski, M.; Sarlomo-Rikala, M.; Miettinen, M.: Mutations in exon 11 of c-kit occur preferentially in malignant versus benign gastrointestinal stromal tumors and do not occur in leiomyomas or leiomyosarcomas. Am. J. Path. 154: 53–60, 1999.
- [77376] 15706. Longley, B. J.; Tyrrell, L.; Lu, S.-Z.; Ma, Y.-S.; Langley, K.; Ding, T.; Duffy, T.; Jacobs, P.; Tang, L. H.; Modlin, I.: Somatic c-KIT activating mutation in urticaria pigmentosa and aggressive mastocytosis: establishment of clonality in a human mast cell neoplasm. Nature Genet. 12: 312–314, 1996.
- [77377] 15707. Longley, B. J., Jr.; Metcalfe, D. D.; Tharp, M.; Wang, X.; Tyrrell, L.; Lu, S.-Z.; Heitjan, D.; Ma, Y.: Activating and dominant inactivating c-KIT catalytic domain mutations in distinct clinical forms of human mastocytosis. Proc. Nat. Acad. Sci. 96: 1609–1614, 1999.

- [77378] 15708. Marklund, S.; Kijas, J.; Rodriguez-Martinez, H.; Ronnstrand, L.; Funa, K.; Moller, M.; Lange, D.; Edfors-Lilja, I.; Andersson, L.: Molecular basis for the dominant white phenotype in the domestic pig. *Genome Res.* 8: 826–833, 1998.
- [77379] 15709. Kiefer, M. C.; Tucker, J. E.; Joh, R.; Landsberg, K. E.; Saltman, D.; Barr, P. J.: Identification of a second human subtilisin-like protease gene in the fes/fps region of chromosome 15. *DNA Cell Biol.* 10:757–769, 1991.
- [77380] 15710. Amiel, J.; Audollent, S.; Joly, D.; Dureau, P.; Salomon, R.; Tellier, A.-L.; Auge, J.; Bouissou, F.; Antignac, C.; Gubler, M.-C.; Eccles, M. R.; Munnich, A.; Vekemans, M.; Lyonnet, S.; Attie-Bitach, T.: PAX2 mutations in renal-coloboma syndrome: mutational hotspot and germline mosaicism. *Europ. J. Hum. Genet.* 8: 820–826, 2000.
- [77381] 15711. Dehbi, M.; Ghahremani, M.; Lechner, M.; Dressler, G.; Pelletier, J.: The paired-box transcription factor, PAX2, positively modulates expression of the Wilms' tumor suppressor gene. *Oncogene* 13: 447–453, 1996.
- [77382] 15712. Devriendt, K.; Matthijs, G.; Van Damme, B.; Van Caesbroeck, D.; Eccles, M.; Vanrenterghem, Y.; Fryns, J.-P.; Leys, A.: Missense mutation and hexanucleotide duplication in the PAX2 gene in two unrelated families with renal-

coloboma syndrome (MIM 120330). Hum. Genet. 103: 149–153, 1998.

- [77383] 15713. Dressler, G. R.; Deutsch, U.; Chowdhury, K.; Nornes, H. O.; Gruss, P.: Pax2, a new murine paired-box-containing gene and its expression in the developing excretory system. Development 109: 787–795, 1990.
- [77384] 15714. Ford, B.; Rupps, R.; Lirenman, D.; Van Allen, M. I.; Farquharson, D.; Lyons, C.; Friedman, J. M.: Renal-coloboma syndrome: prenatal detection and clinical spectrum in a large family. Am. J. Med. Genet. 99: 137–141, 2001.
- [77385] 15715. Keller, S. A.; Jones, J. M.; Boyle, A.; Barrow, L. L.; Killen, P. D.; Green, D. G.; Kapousta, N. V.; Hitchcock, P. F.; Swank, R. T.; Meisler, M. H.: Kidney and retinal defects (Krd), a transgene-induced mutation with a deletion of mouse chromosome 19 that includes the Pax2 locus. Genomics 23: 309–320, 1994.
- [77386] 15716. Narahara, K.; Baker, E.; Ito, S.; Yokoyama, Y.; Yu, S.; Hewitt, D.; Sutherland, G. R.; Eccles, M. R.; Richards, R. I.: Localisation of a 10q breakpoint within the PAX2 gene in a patient with a de novo t(10;13) translocation and optic nerve coloboma-renal disease. J. Med. Genet. 34: 213–216, 1997.



- [77387] 15717. Pilz, A. J.; Povey, S.; Gruss, P.; Abbott, C. M.: Mapping of the human homologs of the murine paired-box-containing genes. *Mammalian Genome* 4: 78–82, 1993.
- [77388] 15718. Favor, J.; Sandulache, R.; Neuhauser-Klaus, A.; Pretsch, W.; Chatterjee, B.; Senft, E.; Wurst, W.; Blanquet, V.; Grimes, P.; Sporle, R.; Schughart, K.: The mouse Pax2(1Neu) mutation is identical to a human PAX2 mutation in a family with renal-coloboma syndrome and results in developmental defects of the brain, ear, eye, and kidney. *Proc. Nat. Acad. Sci.* 93:13870–13875, 1996.
- [77389] 15719. Liou, G. I.; Ma, D.-P.; Yang, Y.-W.; Geng, L.; Zhu, C.; Baehr, W.: Human interstitial retinoid-binding protein: gene structure and primary sequence. *J. Biol. Chem.* 264: 8200–8206, 1989.
- [77390] 15720. Nakamura, Y.; Lathrop, M.; Bragg, T.; Leppert, M.; O'Connell, P.; Jones, C.; Lalouel, J. M.; White, R.: An extended genetic linkage map of markers for human chromosome 10. *Genomics* 3: 389–392, 1988.
- [77391] 15721. Sap, J.; Munoz, A.; Damm, K.; Goldberg, Y.; Ghysdael, J.; Leutz, A.; Beug, H.; Vennstrom, B.: The c-erb-A protein is a high-affinity receptor for thyroid hormone. *Nature* 324: 635–640, 1986.

- [77392] 15722.Bayliss, W.; Starling, E. H.: The mechanism of pancreatic secretion. *J.Physiol. (London)* 28: 325–353, 1902.
- [77393] 15723.Chow, B. K.–C.: Molecular cloning and functional characterization of a human secretin receptor. *Biochem. Biophys. Res. Commun.* 212:204–211, 1995.
- [77394] 15724.Ishihara, T.; Nakamura, S.; Kaziro, Y.; Takahashi, T.; Takahashi,K.; Nagata, S.: Molecular cloning and expression of a cDNA encoding the secretin receptor. *EMBO J.* 10: 1635–1641, 1991.
- [77395] 15725.Mark, H. F. L.; Chow, B. K.–C.: Localization of the gene encoding the secretin receptor, SCTR, on human chromosome 2q14.1 by fluorescence in situ hybridization and chromosome morphometry. *Genomics* 29: 817–818,1995.
- [77396] 15726.Peel, A. L.; Rao, R. V.; Cottrell, B. A.; Hayden, M. R.; Ellerby,L. M.; Bredesen, D. E.: Double-stranded RNA-dependent protein kinase,PKR, binds preferentially to Huntington's disease (HD) transcripts and is activated in HD tissue. *Hum. Molec. Genet.* 10: 1531–1538,2001.
- [77397] 15727.Squire, J.; Meurs, E. F.; Chong, K. L.; McMillan, N. A. J.; Hovanessian,A. G.; Williams, B. R. G.: Localization of the human interferon-induced,ds-RNA activated p68 kinase gene (PRKR) to chromosome 2p21–p22. *Genomics*

16:768–770, 1993.

- [77398] 15728.Taylor, D. R.; Shi, S. T.; Romano, P. R.; Barber, G. N.; Lal, M.M. C.: Inhibition of the interferon–inducible protein kinase PKRby HCV E2 protein. *Science* 285: 107–110, 1999.
- [77399] 15729.Brott, B. K.; Alessandrini, A.; Largaespada, D. A.; Copeland, N.G.; Jenkins, N. A.; Crews, C. M.; Erikson, R. L.: MEK2 is a kinaserelated to MEK1 and is differentially expressed in murine tissues. *CellGrowth Differ.* 4: 921–929, 1993.
- [77400] 15730.Crews, C. M.; Alessandrini, A.; Erikson, R. L.: The primary structureof MEK, a protein kinase that phosphorylates the ERK gene product. *Science* 258:478–480, 1992.
- [77401] 15731.Favata, M. F.; Horiuchi, K. Y.; Manos, E. J.; Daulerio, A. J.;Stradley, D. A.; Feeser, W. S.; Van Dyk, D. E.; Pitts, W. J.; Earl,R. A.; Hobbs, F.; Copeland, R. A.; Magolda, R. L.; Scherle, P. A.;Trzaskos, J. M.: Identification of a novel inhibitor of mitogen–activatedprotein kinase kinase. *J. Biol. Chem.* 273: 18623–18632, 1998.
- [77402] 15732.Meloche, S.; Gopalbhai, K.; Beatty, B. G.; Scherer, S. W.; Pellerin,J.: Chromosome mapping of the human genes encoding the MAP kinasekinase MEK1 (MAP2K1) to 15q21 and MEK2 (MAP2K2) to 7q32. *Cytogenet.Cell Genet.* 88:

249–252, 2000.

- [77403] 15733.Orth, K.; Palmer, L. E.; Bao, Z. Q.; Stewart, S.; Rudolph, A. E.; Bliska, J. B.; Dixon, J. E.: Inhibition of the mitogen-activated protein kinase kinase superfamily by a Yersinia effector. *Science* 285:1920–1923, 1999.
- [77404] 15734.Perry, R. L. S.; Parker, M. H.; Rudnicki, M. A.: Activated MEK1 binds the nuclear MyoD transcriptional complex to repress transactivation. *Molec.Cell* 8: 291–301, 2001.
- [77405] 15735.Pleschka, S.; Wolff, T.; Ehrhardt, C.; Hobom, G.; Planz, O.; Rapp, U. R.; Ludwig, S.: Influenza virus propagation is impaired by inhibition of the Raf/MEK/ERK signalling cascade. *Nature Cell Biol.* 3: 301–305, 2001.
- [77406] 15736.Rampoldi, L.; Zimbello, R.; Bortoluzzi, S.; Tiso, N.; Valle, G.; Lanfranchi, G.; Danieli, G. A.: Chromosomal localization of four MAPK signaling cascade genes: MEK1, MEK3, MEK4 and MEK5. *Cytogenet.Cell Genet.* 78: 301–303, 1997.
- [77407] 15737.Ryan, K. M.; Ernst, M. K.; Rice, N. R.; Vousden, K. H.: Role of NF- $\kappa$ B in p53-mediated programmed cell death. *Nature* 404: 892–897, 2000.
- [77408] 15738.Sebolt-Leopold, J. S.; Dudley, D. T.; Herrera, R.; Van Becelaere, K.; Wiland, A.; Gowan, R. C.; Tecle, H.; Bar-

rett, S. D.; Bridges, A.; Przybranowski, S.; Leopold, W. R.; Saltiel, A. R.: Blockade of the MAP kinase pathway suppresses growth of colon tumors in vivo. *Nature Med.* 5: 810–816, 1999.

[77409] 15739. Seger, R.; Krebs, E. G.: The MAPK signaling cascade. *FASEB J.* 9:726–735, 1995.

[77410] 15740. Seger, R.; Seger, D.; Lozeman, F. J.; Ahn, N. G.; Graves, L. M.; Campbell, J. S.; Ericsson, L.; Harrylock, M.; Jensen, A. M.; Krebs, E. G.: Human T-cell mitogen-activated protein kinase kinases are related to yeast signal transduction kinases. *J. Biol. Chem.* 267:25628–25631, 1992.

[77411] 15741. Zheng, C. F.; Guan, K. L.: Cloning and characterization of two distinct human extracellular signal-regulated kinase activator kinases, MEK1 and MEK2. *J. Biol. Chem.* 268: 11435–11439, 1993.

[77412] 15742. Bunnell, B.; Heath, L. S.; Adams, D. E.; Lahti, J. M.; Kidd, V. J.: Elevated expression of a p58 protein kinase leads to changes in the CHO cell cycle. *Proc. Nat. Acad. Sci.* 87: 7467–7471, 1990.

[77413] 15743. Cornelis, S.; Bruynooghe, Y.; Denecker, G.; Van Huffel, S.; Tinton, S.; Beyaert, R.: Identification and characterization of a novel cell cycle-regulated internal ribosome

entry site. *Molec. Cell* 5: 597–605,2000.

- [77414] 15744.Eipers, P. G.; Barnoski, B. L.; Han, J.; Carroll, A. J.; Kidd,V. J.: Localization of the expressed human p58 protein kinase chromosomal gene to chromosome 1p36 and a highly related sequence to chromosome15. *Genomics* 11: 621–629, 1991.
- [77415] 15745.Eipers, P. G.; Lahti, J. M.; Kidd, V. J.: Structure and expression of the human p58(cdk-1) protein kinase chromosomal gene. *Genomics* 13:613–621, 1992.
- [77416] 15746.Gururajan, R.; Lahti, J. M.; Grenet, J.; Easton, J.; Gruber, I.; Ambros, P. F.; Kidd, V. J.: Duplication of a genomic region containing the Cdc2L1–2 and MMP21–22 genes on human chromosome 1p36.3 and their linkage to D1Z2. *Genome Res.* 8: 929–939, 1998.
- [77417] 15747.Lahti, J. M.; Valentine, M.; Xiang, J.; Jones, B.; Amann, J.; Grenet, J.; Richmond, G.; Look, A. T.; Kidd, V. J.: Alterations in the PITSLRE protein kinase gene complex on chromosome 1p36 in childhood neuroblastoma. *Nature-Genet.* 7: 370–375, 1994.
- [77418] 15748.White, P. S.; Maris, J. M.; Beltinger, C.; Sulman, E.; Marshall, H. N.; Fujimori, M.; Kaufman, B. A.; Biegel, J. A.; Allen, C.; Hilliard, C.; Valentine, M. B.; Look, A. T.; Enomoto, H.; Sakiyama, S.; Brodeur, G. M.: A region of

consistent deletion in neuroblastoma maps within human chromosome 1p36.2–36.3. *Proc. Nat. Acad. Sci.* 92: 5520–5524, 1995.

[77419] 15749. Xiang, J.; Lahti, J. M.; Grenet, J.; Easton, J.; Kidd, V. J.: Molecular cloning and expression of alternatively spliced PITSLRE protein kinase isoforms. *J. Biol. Chem.* 269: 15786–15794, 1994.

[77420] 15750. Collu, R.; Tang, J.; Castagne, J.; Lagace, G.; Masson, N.; Huot, C.; Deal, C.; Delvin, E.; Faccenda, E.; Eidne, K. A.; Van Vliet, G.: A novel mechanism for isolated central hypothyroidism: inactivating mutations in the thyrotropin-releasing hormone receptor gene. *J. Clin. Endocr. Metab.* 82: 1361–1365, 1997.

[77421] 15751. Amin, R. S.; Wert, S. E.; Baughman, R. P.; Tomashevski, J. F., Jr.; Nogee, L. M.; Brody, A. S.; Hull, W. M.; Whitsett, J. A.: Surfactant protein deficiency in familial interstitial lung disease. *J. Pediatr.* 139:85–92, 2001.

[77422] 15752. Glasser, S. W.; Burhans, M. S.; Korfhagen, T. R.; Na, C.-L.; Sly, P. D.; Ross, G. F.; Ikegami, M.; Whitsett, J. A.: Altered stability of pulmonary surfactant in SP-C-deficient mice. *Proc. Nat. Acad. Sci.* 98: 6366–6371, 2001.

[77423] 15753. Keller, D. M.; Zeng, X.; Wang, Y.; Zhang, Q. H.; Kapoor, M.; Shu, H.; Goodman, R.; Lozano, G.; Zhao, Y.; Lu,

H.: A DNA damage–induced p53 serine 392 kinase complex contains CK2, hSpt16, and SSRP1. *Molec. Cell* 283–292, 2001.

[77424] 15754. Maki, R. G.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Srivastava, P. K.: Localization of genes for the stress–inducible tumor rejection antigen gp96. (Abstract) *Cytogenet. Cell Genet.* 58: 1978 only, 1991.

[77425] 15755. Maki, R. G.; Old, L. J.; Srivastava, P. K.: Human homologue of murine tumor rejection antigen gp96: 5–prime–regulatory and coding regions and relationship to stress–induced proteins. *Proc. Nat. Acad. Sci.* 87: 5658–5662, 1990.

[77426] 15756. Schild, H.; Rammensee, H.–G.: gp96: the immune system's Swiss army knife. *Nature Immun.* 1: 100–101, 2000.

[77427] 15757. Lanske, B.; Karaplis, A. C.; Lee, K.; Luz, A.; Vortkamp, A.; Pirro, A.; Karperien, M.; Defize, L. H. K.; Ho, C.; Mulligan, R. C.; Abou–Samra, A.–B.; Juppner, H.; Segre, G. V.; Kronenberg, H. M.: PTH/PTHrP receptor in early development and Indian hedgehog–regulated bone growth. *Science* 273:663–666, 1996.

[77428] 15758. Albright, F.: Case records of the Massachusetts General Hospital (case 27461). *New Eng. J. Med.* 225:



789–791, 1941.

- [77429] 15759. Bakre, M. M.; Zhu, Y.; Yin, H.; Burton, D. W.; Terkeltaub, R.; Deftos, L. J.; Varner, J. A.: Parathyroid hormone-related peptide is a naturally occurring, protein kinase A-dependent angiogenesis inhibitor. *Nature Med.* 8: 995–1003, 2002.
- [77430] 15760. Broadus, A. E.; Mangin, M.; Ikeda, K.; Insogna, K. L.; Weir, E. C.; Burtis, W. J.; Stewart, A. F.: Humoral hypercalcemia of cancer: identification of a novel parathyroid hormone-like peptide. *New Eng. J. Med.* 319: 556–563, 1988.
- [77431] 15761. Hammonds, R. G., Jr.; McKay, P.; Winslow, G. A.; Diefenbach-Jagger, H.; Grill, V.; Glatz, J.; Rodda, C. P.; Moseley, J. M.; Wood, W. I.; Martin, T. J.: Purification and characterization of recombinant human parathyroid hormone-related protein. *J. Biol. Chem.* 264: 14806–14811, 1989.
- [77432] 15762. Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.; Yasuda, T.; Banville, D.; Goltzman, D.: Gene for parathyroid hormone-like peptide is on mouse chromosome 6. *Cytogenet. Cell Genet.* 53: 80–82, 1990.
- [77433] 15763. Hendy, G. N.; Sakaguchi, A. Y.; Lalley, P. A.; Martinez, L.; Yasuda, T.; Banville, D.; Goltzman, D.: Gene for

parathyroid hormone-like peptide is on mouse chromosome 6. (Abstract) Cytogenet. Cell Genet. 51:1003 only, 1989.

- [77434] 15764. Kamps, M. P.; Look, A. T.; Baltimore, D.: The human t(1:19) translocation in pre-B ALL produces multiple nuclear E2A-Pbx1 fusion proteins with differing transforming potentials. Genes Dev. 5: 358–368, 1991.
- [77435] 15765. Kim, S. K.; Selleri, L.; Lee, J. S.; Zhang, A. Y.; Gu, X.; Jacobs, Y.; Cleary, M. L.: Pbx1 inactivation disrupts pancreas development and in *lpl1*-deficient mice promotes diabetes mellitus. Nature Genet. 30:430–435, 2002.
- [77436] 15766. Mercader, N.; Leonardo, E.; Azpiazu, N.; Serrano, A.; Morata, G.; Martinez-A, C.; Torres, M.: Conserved regulation of proximal limb axis development by *Meis1/Hth*. Nature 402: 425–429, 1999.
- [77437] 15767. Monica, K.; Galili, N.; Nourse, J.; Saltman, D.; Cleary, M. L.: PBX2 and PBX3, new homeobox genes with extensive homology to the human proto-oncogene PBX1. Molec. Cell. Biol. 11: 6149–6157, 1991.
- [77438] 15768. Aguado, B.; Campbell, R. D.: The novel gene G17, located in the human major histocompatibility complex, encodes PBX2, a homeodomain-containing protein. Genomics 25: 650–659, 1995.

- [77439] 15769.Katsanis, N.; Fitzgibbon, J.; Fisher, E. M. C.: Paralog mapping: identification of a region in the human MHC triplicated onto human chromosomes 1 and 9 allows the prediction and isolation of novel PBX and NOTCH loci. *Genomics* 35: 101–108, 1996.
- [77440] 15770.Hourvitz, A.; Widger, A. E.; Filho, F. L. T.; Chang, R. J.; Adashi, E. Y.; Erickson, G. F.: Pregnancy-associated plasma protein-A gene expression in human ovaries is restricted to healthy follicles and corpora lutea. *J. Clin. Endocrin. Metab.* 85: 4916–4919, 2000.
- [77441] 15771.Lawrence, J. B.; Oxvig, C.; Overgaard, M. T.; Sottrup-Jensen, L.; Gleich, G. J.; Hays, L. G.; Yates, J. R., III; Conover, C. A.: The insulin-like growth factor (IGF)-dependent IGF binding protein-4 protease secreted by human fibroblasts is pregnancy-associated plasma protein-A. *Proc. Nat. Acad. Sci.* 96: 3149–3153, 1999.
- [77442] 15772.Silahtaroglu, A. N.; Tumer, Z.; Kristensen, T.; Sottrup-Jensen, L.; Tommerup, N.: Assignment of the human gene for pregnancy-associated plasma protein A (PAPPA) to 9q33.1 by fluorescence in situ hybridization to mitotic and meiotic chromosomes. *Cytogenet. Cell Genet.* 62: 214–216, 1993.
- [77443] 15773.Smith, G. C. S.; Stenhouse, E. J.; Crossley, J. A.;

Aitken, D.A.; Cameron, A. D.; Connor, J. M.: Early pregnancy levels of pregnancy-associated plasma protein A and the risk of intrauterine growth restriction, premature birth, preeclampsia, and stillbirth. *J. Clin. Endocr. Metab.* 87:1762–1767, 2002.

[77444] 15774. Smith, G. C. S.; Stenhouse, E. J.; Crossley, J. A.; Aitken, D.A.; Cameron, A. D.; Connor, J. M.: Early-pregnancy origins of low birth weight. *Nature* 417: 916 only, 2002.

[77445] 15775. Eisenhofer, G.; Walther, M. M.; Huynh, T.-T.; Li, S.-T.; Bornstein, S. R.; Vortmeyer, A.; Mannelli, M.; Goldstein, D. S.; Linehan, W.M.; Lenders, J. W. M.; Pacak, K.: Pheochromocytomas in von Hippel-Lindau syndrome and multiple endocrine neoplasia type 2 display distinct biochemical and clinical phenotypes. *J. Clin. Endocr. Metab.* 86:1999–2008, 2001.

[77446] 15776. Glasser, S. W.; Korfhagen, T. R.; Weaver, T. E.; Clark, J. C.; Pilot-Matias, T.; Meuth, J.; Fox, J. L.; Whitsett, J. A.: cDNA, deduced polypeptide structure and chromosomal assignment of human pulmonary surfactant protein lipid, SPL(pVal). *J. Biol. Chem.* 263: 9–12, 1988.

[77447] 15777. Zhang, S.-H.; Liu, J.; Kobayashi, R.; Tonks, N. K.: Identification of the cell cycle regulator VCP (p97/CDC48)

as a substrate of the band 4.1-related protein-tyrosine phosphatase PTPH1. *J. Biol. Chem.* 274:17806–17812, 1999.

[77448] 15778. Lorimer, D. D.; Benya, R. V.: Cloning and quantification of galanin-1 receptor expression by mucosal cells lining the human gastrointestinal tract. *Biochem. Biophys. Res. Commun.* 222: 379–385, 1996.

[77449] 15779. Nicholl, J.; Kofler, B.; Sutherland, G. R.; Shine, J.; Lismaa, T. P.: Assignment of the gene encoding human galanin receptor (GALNR) to 18q23 by in situ hybridization. *Genomics* 30: 629–630, 1995.

[77450] 15780. Parker, E. M.; Izzarelli, D. G.; Nowak, H. P.; Mahle, C. D.; Iben, L. G.; Wang, J.; Goldstein, M. E.: Cloning and characterization of the rat GALR1 galanin receptor from Rin14B insulinoma cells. *Molec. Brain Res.* 34: 179–189, 1995.

[77451] 15781. Simoneaux, D. K.; Leach, R. J.; O'Connell, P.: Galanin receptor1 gene (Galnr1) is tightly linked to the myelin basic protein gene on chromosome 18 in mouse. *Mammalian Genome* 8: 875 only, 1997.

[77452] 15782. Walli, R.; Schafer, H.; Morys-Wortmann, C.; Paetzold, G.; Nustede, R.; Schmidt, W. E.: Identification and biochemical characterization of the human brain galanin

receptor. *J. Molec. Endocr.* 13: 347–356, 1994.

- [77453] 15783. Janowski, B. A.; Grogan, M. J.; Jones, S. A.; Wisely, G. B.; Kliewer, S. A.; Corey, E. J.; Mangelsdorf, D. J.: Structural requirements of ligands for the oxysterol liver X receptors LXR- $\alpha$  and LXR- $\beta$ . *Proc. Nat. Acad. Sci.* 96: 266–271, 1999.
- [77454] 15784. Le Beau, M. M.; Song, C.; Davis, E. M.; Hiipakka, R. A.; Kokontis, J. M.; Liao, S.: Assignment of the human ubiquitous receptor gene (UNR) to 19q13.3 using fluorescence in situ hybridization. *Genomics* 26: 166–168, 1995.
- [77455] 15785. Repa, J. J.; Berge, K. E.; Pomajzl, C.; Richardson, J. A.; Hobbs, H.; Mangelsdorf, D. J.: Regulation of ATP-binding cassette sterol transporters ABCG5 and ABCG8 by the liver X receptors  $\alpha$  and  $\beta$ . *J. Biol. Chem.* 277: 18793–18800, 2002.
- [77456] 15786. Shinar, D. M.; Endo, N.; Rutledge, S. J.; Vogel, R.; Rodan, G. A.; Schmidt, A.: NER, a new member of the gene family encoding the human steroid hormone nuclear receptor. *Gene* 147: 273–276, 1994.
- [77457] 15787. Song, C.; Hiipakka, R. A.; Kokontis, J. M.; Liao, S.: Ubiquitous receptor: structures, immunocytochemical localization, and modulation of gene activation by receptors for retinoic acids and thyroid hormones. *Ann. N.Y. Acad.*

Sci. 761: 38–49, 1995.

- [77458] 15788.Tangirala, R. K.; Bischoff, E. D.; Joseph, S. B.; Wagner, B. L.;Walczak, R.; Laffitte, B. A.; Daige, C. L.; Thomas, D.; Heyman, R.A.; Mangelsdorf, D. J.; Wang, X.; Lusis, A. J.; Tontonoz, P.; Schulman,I. G.: Identification of macrophage liver X receptors as inhibitorsof atherosclerosis. Proc. Nat. Acad. Sci. 99: 11896–11901, 2002.
- [77459] 15789.Futterer, A.; Kruppa, G.; Kramer, B.; Lemke, H.; Kronke, M.: Molecularcloning and characterization of human kinectin. Molec. Biol. Cell 6:161–170, 1995.
- [77460] 15790.Plitz, T.; Pfeffer, K.: Intact lysosome transport and phagosomefunction despite kinectin deficiency. Molec. Cell. Biol. 21: 6044–6055,2001.
- [77461] 15791.Print, C. G.; Leung, E.; Harrison, J. E. B.; Watson, J. D.; Krissansen,G. W.: Cloning of a gene encoding a human leukocyte protein characterisedby extensive heptad repeats. Gene 144: 221–228, 1994.
- [77462] 15792.Print, C. G.; Morris, C. M.; Spurr, N. K.; Rooke, L.; Krissansen,G. W.: The CG–1 gene, a member of the kinectin and ES/130 family,maps to human chromosome band 14q22. Immunogenetics 43: 227–229,1996.

- [77463] 15793.Rao, P. N.; Yu, H.; Hodge, R.; Pettenati, M. J.; Sheetz, M. P.: Assignment of the human kinectin gene (KTN1), encoding a kinesin-binding protein, to chromosome 14 band q22.1 by in situ hybridization. *Cytogenet. Cell Genet.* 79: 196–197, 1997.
- [77464] 15794.Yu, H.; Nicchitta, C. V.; Kumar, J.; Becker, M.; Toyoshima, I.; Sheetz, M. P.: Characterization of kinectin, a kinesin-binding protein: primary sequence and N-terminal topogenic signal analysis. *Molec. Biol. Cell* 6: 171–183, 1995.
- [77465] 15795.Gordenin, D. A.; Kunkel, T. A.; Resnick, M. A.: Repeat expansion--all in a flap? *Nature Genet.* 16: 116–118, 1997.
- [77466] 15796.Greene, A. L.; Snipe, J. R.; Gordenin, D. A.; Resnick, M. A.: Functional analysis of human FEN1 in *Saccharomyces cerevisiae* and its role in genome stability. *Hum. Molec. Genet.* 8: 2263–2273, 1999.
- [77467] 15797.Augusseau, S.; Jouk, S.; Jalbert, P.; Prieur, M.: DiGeorge syndrome and 22q11 rearrangements. (Letter) *Hum. Genet.* 74: 206 only, 1986.
- [77468] 15798.Van Esch, H.; Groenen, P.; Daw, S.; Poffyn, A.; Holvoet, M.; Scambler, P.; Fryns, J.-P.; Van de Ven, W.; Devriendt, K.: Partial DiGeorge syndrome in two patients with



a 10p rearrangement. Clin. Genet. 55:269–276, 1999.

[77469] 15799.Wadey, R.; Daw, S.; Taylor, C.; Atif, U.; Kamath, S.; Halford,S.; O'Donnell, H.; Wilson, D.; Goodship, J.; Burn, J.; Scambler, P.: Isolation of a gene encoding an integral membrane protein from the vicinity of a balanced translocation breakpoint associated with DiGeorges syndrome.

Hum. Molec. Genet. 4: 1027–1033, 1995.

[77470] 15800.Contractur, A.; Rogers, C.; Maron, C.; Henkemeyer, M.; Swanson,G. T.; Heinemann, S. F.: Trans-synaptic Eph receptor–ephrin signaling in hippocampal mossy fiber LTP. Science 296: 1864–1869, 2002.

[77471] 15801.Tang, X. X.; Biegel, J. A.; Nycum, L. M.; Yoshioka, A.; Brodeur,G. M.; Pleasure, D. E.; Ikegaki, N.: cDNA cloning, molecular characterization, and chromosomal localization of NET (EPHT2), a human EPH-related receptor–protein–tyrosine kinase gene preferentially expressed in brain. :Genomics 29:426–437, 1995.

[77472] 15802.Kamp, C.; Hirschmann, P.; Voss, H.; Huellen, K.; Vogt, P. H.:Two long homologous retroviral sequence blocks in proximal Yq11 causeAZFa microdeletions as a result of intrachromosomal recombination events. Hum. Molec. Genet. 9: 2563–2572, 2000.

[77473] 15803.Krausz, C.; Meyts, E. R.–D.; Frydelund–Larson, L.;

Quintana-Murci, L.; McElreavey, K.; Skakkebaek, N. E.:  
Double-blind Y chromosomemicrodeletion analysis in  
men with known sperm parameters and reproductive hor-  
mone profiles: microdeletions are specific for spermato-  
genic failure. *J.Clin. Endocr. Metab.* 86: 2638-2642, 2001.

[77474] 15804. Simpson, E.; Chandler, P.; Goulmy, E.; Ma, K.; Har-  
greave, T. B.; Chandley, A. C.: Loss of the 'azoospermia  
factor' (AZF) on Yq in man is not associated with loss of  
HYA. *Hum. Molec. Genet.* 2: 469-471, 1993.

[77475] 15805. Sun, C.; Skaletsky, H.; Rozen, S.; Gromoll, J.; Ni-  
eschlag, E.; Oates, R.; Page, D. C.: Deletion of azoospermia  
factor a (AZFa) region of human Y chromosome caused by  
recombination between HERV15 proviruses. *Hum. Molec.*  
*Genet.* 9: 2291-2296, 2000.

[77476] 15806. Krausz, C.; Quintana-Murci, L.; Barboux, S.; Siffroi,  
J.-P.; Rouba, H.; Delafontaine, D.; Souleyreau-Therville, N.;  
Arvis, G.; Antoine, J. M.; Erdei, E.; Taar, J. P.; Tar, A.; Jean-  
didier, E.; Plessis, G.; Bourgeron, T.; Dadoune, J.-P.; Fel-  
lous, M.; McElreavey, K.: A high frequency of Y chromo-  
some deletions in males with nonidiopathic infertility.  
*J.Clin. Endocr. Metab.* 84: 3606-3612, 1999.

[77477] 15807. Ma, K.; Inglis, J. D.; Sharkey, A.; Bickmore, W. A.;  
Hill, R. E.; Prosser, E. J.; Speed, R. M.; Thomson, E. J.;

Jobling, M.; Taylor, K.; Wolfe, J.; Cooke, H. J.; Hargreave, T. B.; Chandley, A. C.: AY chromosome gene family with RNA-binding protein homology: candidates for the azoospermia factor AZF controlling human spermatogenesis. *Cell* 75:1287–1295, 1993.

[77478] 15808. Vergnaud, G.; Page, D. C.; Simmler, M. C.; Brown, L.; Rouyer, F.; Noel, B.; Botstein, D.; de la Chapelle, A.; Weissenbach, J.: A deletion map of the human Y chromosome based on DNA hybridization. *Am. J. Hum. Genet.* 38: 109–124, 1986.

[77479] 15809. Vogt, P.; Chandley, A. C.; Hargreave, T. B.; Keil, R.; Ma, K.; Sharkey, A.: Microdeletions in interval 6 of the Y chromosome of males with idiopathic sterility point to disruption of AZF, a human spermatogenesis gene. *Hum. Genet.* 89: 491–496, 1992.

[77480] 15810. Vogt, P. H.; Edelman, A.; Kirsch, S.; Henegariu, O.; Hirschmann, P.; Kieseewetter, F.; Kohn, F. M.; Schill, W. B.; Farah, S.; Ramos, C.; Hartmann, M.; Hartschuh, W.; Meschede, D.; Behre, H. M.; Castel, A.; Nieschlag, E.; Weidner, W.; Grone, H.-J.; Jung, A.; Engel, W.; Haidl, G.: Human Y chromosome azoospermia factors (AZF) mapped to different subregions in Yq11. *Hum. Molec. Genet.* 5: 933–943, 1996.

- [77481] 15811. Agulnik, A. I.; Bishop, C. E.; Lerner, J. L.; Agulnik, S. I.; Solovyev, V. V.: Analysis of mutation rates in the SMCY/SMCX genes shows that mammalian evolution is male driven. *Mammalian Genome* 8: 134–138, 1997.
- [77482] 15812. Hawkins, J. R.: Mutational analysis of SRY in XY females. *Hum. Mutat.* 2: 347–350, 1993.
- [77483] 15813. Hawkins, J. R.; Taylor, A.; Berta, P.; Levilliers, J.; Van der Auwera, B.; Goodfellow, P. N.: Mutational analysis of SRY: nonsense and missense mutations in XY sex reversal. *Hum. Genet.* 88: 471–474, 1992.
- [77484] 15814. Rohen, C.; Caselitz, J.; Stern, C.; Wanschura, S.; Schoenmakers, E. F.; Van de Ven, W. J.; Barnitzke, S.; Bullerdiek, J.: A hamartoma of the breast with an aberration of 12q mapped to the MAR region by fluorescence in situ hybridization. *Genes Chromosomes Cancer* 84:82–84, 1995.
- [77485] 15815. Schoenmakers, E. F. P. M.; Huysmans, C.; Van de Ven, W. J. M.: Allelic knockout of novel splice variants of human recombination repair gene RAD51B in t(12;14) uterine leiomyomas. *Cancer Res.* 59:19–23, 1999.
- [77486] 15816. Zhou, X.; Benson, K. F.; Ashar, H. R.; Chada, K.: Mutation responsible for the mouse pygmy phenotype in the developmentally regulated factor HMGI-C. *Nature* 377:

771–774, 1995.

- [77487] 15817. Agalioti, T.; Lomvardas, S.; Parekh, B.; Yie, J.; Maniatis, T.; Thanos, D.: Ordered recruitment of chromatin modifying and general transcription factors to the IFN- $\beta$  promoter. *Cell* 103: 667–678, 2000.
- [77488] 15818. Munshi, N.; Agalioti, T.; Lomvardas, S.; Merika, M.; Chen, G.; Thanos, D.: Coordination of a transcriptional switch by HMG(IY) acetylation. *Science* 293: 1133–1136, 2001.
- [77489] 15819. Thanos, D.; Maniatis, T.: Virus induction of human IFN- $\beta$  gene expression requires the assembly of an enhanceosome. *Cell* 83: 1091–1100, 1995.
- [77490] 15820. Wood, L. J.; Mukherjee, M.; Dolde, C. E.; Xu, Y.; Maher, J. F.; Bunton, T. E.; Williams, J. B.; Resar, L. M. S.: HMG-I/Y, a new c-Myb target gene and potential oncogene. *Molec. Cell Biol.* 20: 5490–5502, 2000.
- [77491] 15821. Burgess, D. L.; Kohrman, D. C.; Galt, J.; Plummer, N. W.; Jones, J. M.; Spear, B.; Meisler, M. H.: Mutation of a new sodium channel gene, *Scn8a*, in the mouse mutant 'motor endplate disease'. *Nature Genet.* 10: 461–465, 1995.
- [77492] 15822. DeRepentigny, Y.; Cote, P. D.; Pool, M.; Bernier, G.; Girard, S.; Vidal, S. M.; Kothary, R.: Pathological and ge-

netic analysis of the degenerating muscle (dmu) mouse: a new allele of Scn8a. Hum. Molec. Genet. 10: 1819–1827, 2001.

- [77493] 15823. Kohrman, D. C.; Harris, J. B.; Meisler, M. H.: Mutation detection in the med and med(J) alleles of the sodium channel Scn8a: unusual splicing due to a minor class AT-AC intron. J. Biol. Chem. 271:17576–17581, 1996.
- [77494] 15824. Kohrman, D. C.; Plummer, N. W.; Schuster, T.; Jones, J. M.; Jang, W.; Burgess, D. L.; Galt, J.; Spear, B. T.; Meisler, M. H.: Insertional mutation of the motor endplate disease (med) locus on mouse chromosome 15. Genomics 26: 171–177, 1995.
- [77495] 15825. Kohrman, D. C.; Smith, M. R.; Goldin, A. L.; Harris, J.; Meisler, M. H.: A missense mutation in the sodium channel Scn8a is responsible for cerebellar ataxia in the mouse mutant jolting. J. Neurosci. 16:5993–5999, 1996.
- [77496] 15826. Meisler, M. H.; Sprunger, L. K.; Plummer, N. W.; Escayg, A.; Jones, J. M.: Ion channel mutations in mouse models of inherited neurological disease. Ann. Med. 29: 569–574, 1997.
- [77497] 15827. Plummer, N. W.; Galt, J.; Jones, J. M.; Burgess, D. L.; Sprunger, L. K.; Kohrman, D. C.; Meisler, M. H.: Exon organization, coding sequence, physical mapping, and poly-

morphic intragenic markers for the human neuronal sodium channel gene SCN8A. Genomics 54: 287–296, 1998.

- [77498] 15828. Plummer, N. W.; McBurney, M. W.; Meisler, M. H.: Alternative splicing of the sodium channel SCN8A predicts a truncated two-domain protein in fetal brain and non-neuronal cells. J. Biol. Chem. 272: 24008–24015, 1997.
- [77499] 15829. Reid, E.; Escayg, A.; Dearlove, A. M.; Lee, D. D.; Meisler, M. H.; Rubinsztein, D. C.: The spastic paraplegia SPG10 locus: narrowing of critical region and exclusion of sodium channel gene SCN8A as a candidate. J. Med. Genet. 38: 65–67, 2001.
- [77500] 15830. Sprunger, L. K.; Escayg, A.; Tallaksen-Greene, S.; Albin, R. L.; Meisler, M. H.: Dystonia associated with mutation of the neuronal sodium channel Scn8a and identification of the modifier locus Scnm1 on mouse chromosome 3. Hum. Molec. Genet. 8: 471–479, 1999.
- [77501] 15831. Chang, D. Y.; Nelson, B.; Bilyeu, T.; Hsu, K.; Darlington, G. J.; Maraia, R. J.: A human Alu RNA-binding protein whose expression is associated with accumulation of small cytoplasmic Alu RNA. Molec. Cell. Biol. 14: 3949–3959, 1994.
- [77502] 15832. Chen, M.; She, H.; Davis, E. M.; Spicer, C. M.; Kim,

L.; Ren, R.; LeBeau, M. M.; Li, W.: Identification of Nck family genes, chromosomal localization, expression, and signaling specificity. *J. Biol. Chem.* 273:25171–25178, 1998.

[77503] 15833. Eden, S.; Rohatgi, R.; Podtelejnikov, A. V.; Mann, M.; Kirschner, M. W.: Mechanism of regulation of WAVE1-induced actin nucleation by Rac1 and Nck. *Nature* 418: 790–793, 2002.

[77504] 15834. Casey, J. L.; Di Jeso, B.; Rao, K.; Klausner, R. D.; Harford, J. B.: Two genetic loci participate in the regulation by iron of the gene for the human transferrin receptor. *Proc. Nat. Acad. Sci.* 85:1787–1791, 1988.

[77505] 15835. Enns, C. A.; Suomalainen, H. A.; Gebhardt, J. E.; Schroder, J.; Sussman, H. H.: Human transferrin receptor: expression of the receptor is assigned to chromosome 3. *Proc. Nat. Acad. Sci.* 79: 3241–3245, 1982.

[77506] 15836. Gareau, R.; Gagnon, M. G.; Thellend, C.; Chenard, C.; Audran, M.; Chanal, J.-L.; Ayotte, C.; Brisson, G. R.: Transferrin soluble receptor: a possible probe for detection of erythropoietin abuse by athletes. *Horm. Metab. Res.* 26: 311–312, 1994.

[77507] 15837. Goodfellow, P. N.; Banting, G.; Sutherland, R.; Greaves, M.; Solomon, E.; Povey, S.: Expression of human transferrin receptor is controlled by a gene on chromo-



some 3: assignment using species specificity of a monoclonal antibody. *Somat. Cell Genet.* 8: 197–206, 1982.

- [77508] 15838. Larrick, J. W.; Hyman, E. S.: Acquired iron-deficiency anemia caused by an antibody against the transferrin receptor. *New Eng. J. Med.* 311: 214–218, 1984.
- [77509] 15839. Levy, J. E.; Jin, O.; Fujiwara, Y.; Kuo, F.; Andrews, N. C.: Transferrin receptor is necessary for development of erythrocytes and the nervous system. *Nature Genet.* 21: 396–399, 1999.
- [77510] 15840. Miller, Y. E.; Jones, C.; Scoggin, C.; Morse, H.; Seligman, P.: Chromosome 3q (22–ter) encodes the human transferrin receptor. *Am. J. Hum. Genet.* 35: 573–583, 1983.
- [77511] 15841. Newman, R.; Schneider, C.; Sutherland, R.; Vodinich, L.; Greaves, M.: The transferrin receptor. *Trends Biochem. Sci.* 7: 397–400, 1982.
- [77512] 15842. Nikinmaa, B.; Schroder, J.: Two antigens, the transferrin receptor and p90 assigned to human chromosome 3, are probably the same protein. *Hereditas* 107:55–58, 1987.
- [77513] 15843. Omary, M. B.; Trowbridge, I. S.: Biosynthesis of the human transferrin receptor in cultured cells. *J. Biol. Chem.*

256: 12888–12892, 1981.

- [77514] 15844.Kashuba, V. I.; Gizatullin, R. Z.; Protopopov, A. I.; Allikmets,R.; Korolev, S.; Li, J.; Boldog, F.; Tory, K.; Zabarovska, V.; Marcsek,Z.; Sumegi, J.; Klein, G.; Zabarovsky, E. R.; Kisselev, L.: Notllinking/jumping clones of human chromosome 3: mapping of the TFRC,RAB7 and HAUSP genes to regions rearranged in leukemia and deletedin solid tumors. FEBS Lett. 419: 181–185, 1997.
- [77515] 15845.Rabin, M.; McClelland, A.; Kuhn, L.; Ruddle, F. H.: Regionallocalization of the human transferrin receptor gene to 3q26.2–qter. Am.J. Hum. Genet. 37: 1112–1116, 1985.
- [77516] 15846.Schneider, C.; Kurkinen, M.; Greaves, M.: Isolation of cDNA clonesfor the human transferrin receptor. EMBO J. 2: 2259–2263, 1983.
- [77517] 15847.Schneider, C.; Owen, M. J.; Banville, D.; Williams, J. G.: Primarystructure of human transferrin receptor deduced from the mRNA sequence. Nature 311:675–678, 1984.
- [77518] 15848.Valenzuela, C. Y.; Avendano, A.; Harb, Z.: Association betweenRh and plasma iron binding (transferrin). Hum. Genet. 87: 438–440,1991.
- [77519] 15849.Vodinelich, L.; Sutherland, R.; Schneider, C.; New–

man, R.; Greaves, M.: Receptor for transferrin may be a 'target' structure for natural killer cells. *Proc. Nat. Acad. Sci.* 80: 835–839, 1983.

- [77520] 15850. Carmeliet, P.; Ferreira, V.; Breier, G.; Pollefeyt, S.; Kleckens, L.; Gertsenstein, M.; Fahrig, M.; Vandenhoek, A.; Harpal, K.; Eberhardt, C.; Declercq, C.; Pawling, J.; Moons, L.; Collen, D.; Risau, W.; Nagy, A.: Abnormal blood vessel development and lethality in embryos lacking a single VEGF allele. *Nature* 380: 435–439, 1996.
- [77521] 15851. Crow, R. S.: Peripheral neuritis in myelomatosis. *Brit. Med. J.* 2: 802–804, 1956.
- [77522] 15852. Dantz, D.; Bewersdorf, J.; Fruehwald-Schultes, B.; Kern, W.; Jelkmann, W.; Born, J.; Fehm, H. L.; Peters, A.: Vascular endothelial growth factor: a novel endocrine defensive response to hypoglycemia. *J. Clin. Endocr. Metab.* 87: 835–840, 2002.
- [77523] 15853. Ferrara, N.; Carver-Moore, K.; Chen, H.; Dowd, M.; Lu, L.; O'Shea, K. S.; Powell-Braxton, L.; Hillan, K. J.; Moore, M. W.: Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene. *Nature* 380: 439–442, 1996.
- [77524] 15854. Folkman, J.: Angiogenesis in cancer, vascular, rheumatoid and other disease. *Nature Med.* 1: 27–31,

1995.

- [77525] 15855. Fukumura, D.; Xavier, R.; Sugiura, T.; Chen, Y.; Park, E.-C.; Lu, N.; Selig, M.; Nielsen, G.; Taksir, T.; Jain, R. K.; Seed, B.: Tumor induction of VEGF promoter activity in stromal cells. *Cell* 94: 715–725, 1998.
- [77526] 15856. Gerber, H.-P.; Malik, A. K.; Solar, G. P.; Sherman, D.; Liang, X. H.; Meng, G.; Hong, K.; Marsters, J. C.; Ferrara, N.: VEGF regulates haematopoietic stem cell survival by an internal autocrine loop mechanism. *Nature* 417: 954–958, 2002.
- [77527] 15857. Gerber, H.-P.; Vu, T. H.; Ryan, A. M.; Kowalski, J.; Werb, Z.; Ferrara, N.: VEGF couples hypertrophic cartilage remodeling, ossification and angiogenesis during endochondral bone formation. *Nature Med.* 5: 623–628, 1999.
- [77528] 15858. Giordano, F. J.; Gerber, H.-P.; Williams, S.-P.; Van-Bruggen, N.; Bunting, S.; Ruiz-Lozano, P.; Gu, Y.; Nath, A. K.; Huang, Y.; Hickey, R.; Dalton, N.; Peterson, K. L.; Ross, J., Jr.; Chien, K. R.; Ferrara, N.: A cardiac myocyte vascular endothelial growth factor paracrine pathway is required to maintain cardiac function. *Proc. Nat. Acad. Sci.* 98: 5780–5785, 2001.
- [77529] 15859. Helmlinger, G.; Endo, M.; Ferrara, N.; Hlatky, L.; Jain, R. K.: Formation of endothelial cell networks. *Nature*

405: 139–141, 2000.

- [77530] 15860.Hofman, P.; van Blijswijk, B. C.; Gaillard, P. J.; Vrensen, G.F. J. M.; Schlingemann, R. O.: Endothelial cell hypertrophy induced by vascular endothelial growth factor in the retina: new insights into the pathogenesis of capillary nonperfusion. *Arch. Ophthalmol.* 119:861–866, 2001.
- [77531] 15861.Holash, J.; Maisonpierre, P. C.; Compton, D.; Boland, P.; Alexander, C. R.; Zagzag, D.; Yancopoulos, G. D.; Wiegand, S. J.: Vessel cooption, regression, and growth in tumors mediated by angiopoietins and VEGF. *Science* 284:1994–1998, 1999.
- [77532] 15862.Jin, K.; Zhu, Y.; Sun, Y.; Mao, X. O.; Xie, L.; Greenberg, D.A.: Vascular endothelial growth factor (VEGF) stimulates neurogenesis in vitro and in vivo. *Proc. Nat. Acad. Sci.* 99: 11946–11950, 2002.
- [77533] 15863.Mattei, M.–G.; Borg, J.–P.; Rosnet, O.; Marme, D.; Birnbaum, D.: Assignment of vascular endothelial growth factor (VEGF) and placenta growth factor (PIGF) genes to human chromosome 6p12–p21 and 14q24–q31 regions, respectively. *Genomics* 32: 168–169, 1996.
- [77534] 15864.Miralles, G. D.; O'Fallon, J. R.; Talley, N. J.: Plasma-cell dyscrasia with polyneuropathy: the spectrum of POEMS syndrome. *New Eng. J. Med.* 327: 1919–1923, 1992.

- [77535] 15865. Mueller, M. D.; Vigne, J.-L.; Minchenko, A.; Lebovic, D. I.; Leitman, D. C.; Taylor, R. N.: Regulation of vascular endothelial growth factor (VEGF) gene transcription by estrogen receptors alpha and beta. *Proc. Nat. Acad. Sci.* 97: 10972–10977, 2000.
- [77536] 15866. Nakanishi, T.; Sobue, I.; Toyokura, Y.; Nishitani, H.; Kuroiwa, Y.; Satoyoshi, E.; Tsubaki, T.; Igata, A.; Ozaki, Y.: The Crow–Fukase syndrome: a study of 102 cases in Japan. *Neurology* 34: 712–720, 1984.
- [77537] 15867. Niimi, H.; Arimura, K.; Jonosono, M.; Hashiguchi, T.; Kawabata, M.; Osame, M.; Kitajima, I.: VEGF is causative for pulmonary hypertension in a patient with Crow–Fukase (POEMS) syndrome. *Intern. Med.* 39: 1101–1104, 2000.
- [77538] 15868. Oosthuyse, B.; Moons, L.; Storkebaum, E.; Beck, H.; Nuyens, D.; Brusselmans, K.; Van Dorpe, J.; Hellings, P.; Gorselink, M.; Heymans, S.; Theilmeier, G.; Dewerchin, M.; and 20 others: Deletion of the hypoxia–response element in the vascular endothelial growth factor promoter causes motor neuron degeneration. *Nature Genet.* 28: 131–138, 2001.
- [77539] 15869. Poltorak, Z.; Cohen, T.; Sivan, R.; Kandelis, Y.; Spira, G.; Vlodavsky, I.; Keshet, E.; Neufeld, G.: VEGF<sub>145</sub>, a secreted vascular endothelial growth factor isoform that

binds to extracellular matrix. J. Biol.Chem. 272:  
7151–7158, 1997.

[77540] 15870.Shimpo, S.: Solitary myeloma causing polyneuritis and endocrinedisorders. Nippon Rinsho 26: 2444–2456, 1968.

[77541] 15871.Soker, S.; Takashima, S.; Miao, H. Q.; Neufeld, G.; Klagsbrun,M.: Neuropilin–1 is expressed by endothelial and tumor cells as anisoform–specific receptor for vascular endothelial growth factor. Cell 92:735–745, 1998.

[77542] 15872.Sone, H.; Kawakami, Y.; Sakauchi, M.; Nakamura, Y.; Takahashi,A.; Shimano, H.; Okuda, Y.; Segawa, T.; Suzuki, H.; Yamada, N.: Neutralizationof vascular endothelial growth factor prevents collagen–induced arthritisand ameliorates established disease in mice. Biochem. Biophys. Res.Commun. 281: 562–568, 2001.

[77543] 15873.Springer, M. L.; Chen, A. S.; Kraft, P. E.; Bednarski, M.; Blau,H. M.: VEGF gene delivery to muscle: potential role for vasculogenesisin adults. Molec. Cell 2: 549–558, 1998.

[77544] 15874.Thurston, G.; Suri, C.; Smith, K.; McClain, J.; Sato, T. N.; Yancopoulos,G. D.; McDonald, D. M.: Leakage–resistant blood vessels in mice transgenicallyoverexpressing angiopoietin–1. Science 286: 2511–2514, 1999.

- [77545] 15875.Tischer, E.; Mitchell, R.; Hartman, T.; Silva, M.; Gospodarowicz,D.; Fiddes, J. C.; Abraham, J. A.: The human gene for vascular endothelialgrowth factor: multiple protein forms are encoded through alternativeexon splicing. *J. Biol. Chem.* 266: 11947–11954, 1991.
- [77546] 15876.Osborne, L. R.; Martindale, D.; Scherer, S. W.; Shi, X.–M.; Huizenga,J.; Heng, H. H. Q.; Costa, T.; Pober, B.; Lew, L.; Brinkman, J.; Rommens,J.; Koop, B.; Tsui, L.–C.: Identification of genes from a 500–kbregion at 7q11.23 that is commonly deleted in Williams syndrome patients. *Genomics* 36:328–336, 1996.
- [77547] 15877.Rampoldi, L.; Dobson–Stone, C.; Rubio, J. P.; Danek, A.; Chalmers,R. M.; Wood, N. W.; Verellen, C.; Ferrer, X.; Malandrini, A.; Fabrizi,G. M.; Brown, R.; Vance, J.; Pericak–Vance, M.; Rudolf, G.; Carre,S.; Alonso, E.; Manfredi, M.; Nemeth, A. H.; Monaco, A. P.: A conservedsort–ing–associated protein is mutant in chorea–acanthocytosis. *NatureGenet.* 28: 119–120, 2001.
- [77548] 15878.Weese–Mayer, D. E.; Bolk, S.; Silvestri, J. M.; Chakravarti, A.: Idiopathic congenital central hypoventilation syndrome: evaluationof brain–derived neurotrophic factor genomic DNA sequence variation. *Am.J. Med. Genet.* 107: 306–310, 2002.



- [77549] 15879.Belden, W. J.; Barlowe, C.: Role of Erv29p in collecting solublesecretory proteins into ER-derived transport vesicles. *Science* 294:1528–1531, 2001.
- [77550] 15880.Duhig, T.; Ruhrberg, C.; Mor, O.; Fried, M.: The human surfeitlocus. *Genomics* 52: 72–78, 1998.
- [77551] 15881.Reeves, J. E.; Fried, M.: The surf-4 gene encodes a novel 30 kDaintegral membrane protein. *Molec. Membr. Biol.* 12: 201–208, 1995.
- [77552] 15882.Archer, B. T., III; Ozcelik, T.; Jahn, R.; Francke, U.; Sudhof,T. C.: Structures and chromosomal localizations of two human genesencoding synaptobrevins 1 and 2. *J. Biol. Chem.* 265: 17267–17273,1990.
- [77553] 15883.Hu, K.; Carroll, J.; Fedorovich, S.; Rickman, C.; Sukhodub, A.;Davietov, B.: Vesicular restriction of synaptobrevin suggests a rolefor calcium in membrane fusion. *Nature* 415: 646–650, 2002.
- [77554] 15884.Franchi, F.; Biguzzi, E.; Cetin, I.; Facchetti, F.; Radaelli, T.;Bozzo, M.; Pardi, G.; Faioni, E. M.: Mutations in the thrombomodulinand endothelial protein C receptor genes in women with late fetalloss. *Brit. J. Haemat.* 114: 641–646, 2001.
- [77555] 15885.Perez, F.; Diamantopoulos, G. S.; Stalder, R.; Kreis, T. E.: CLIP-170highlights growing microtubule ends in

vivo. Cell 96: 517–527, 1999.

[77556] 15886. Pierre, P.; Pepperkok, R.; Kreis, T. E.: Molecular characterization of two functional domains of CLIP-170 in vivo. J. Cell Sci. 107:1909–1920, 1994.

[77557] 15887. Pierre, P.; Scheel, J.; Rickard, J. E.; Kreis, T. E.: CLIP-170 links endocytic vesicles to microtubules. Cell 70: 887–900, 1992.

[77558] 15888. Acland, G. M.; Aguirre, G. D.; Ray, J.; Zhang, Q.; Aleman, T. S.; Cideciyan, A. V.; Pearce-Kelling, S. E.; Anand, V.; Zeng, Y.; Maguire, A. M.; Jacobson, S. G.; Hauswirth, W. W.; Bennett, J.: Gene therapy restores vision in a canine model of childhood blindness. Nature 28:92–95, 2001.

[77559] 15889. Aguirre, G. D.; Baldwin, V.; Pearce-Kelling, S.; Narfstrom, K.; Ray, K.; Acland, G. M.: Congenital stationary night blindness in the dog: common mutation in the RPE65 gene indicates founder effect. Molec. Vision 4: 23, 1998.  
Note: Electronic Article.

[77560] 15890. Bavik, C.-O.; Busch, C.; Eriksson, U.: Characterization of a plasma retinol-binding protein membrane receptor expressed in the retinal pigment epithelium. J. Biol. Chem. 267: 23035–23042, 1992.

[77561] 15891. Feliuss, J.; Thompson, D. A.; Khan, N. W.; Bingham,

E. L.; Jamison, J. A.; Kemp, J. A.; Sieving P. A.: Clinical course and visual function in a family with mutations in the RPE65 gene. *Arch. Ophthalmol.* 120:55–61, 2002.

[77562] 15892. Grimm, C.; Wenzel, A.; Hafezi, F.; Yu, S.; Redmond, T. M.; Reme, C. E.: Protection of Rpe65-deficient mice identifies rhodopsin as a mediator of light-induced retinal degeneration. *Nature Genet.* 25:63–66, 2000.

[77563] 15893. Gu, S.; Thompson, D. A.; Srikumari, C. R. S.; Lorenz, B.; Finckh, U.; Nicoletti, A.; Murthy, K. R.; Rathmann, M.; Kumaramanickavel, G.; Denton, M. J.; Gal, A.: Mutations in RPE65 cause autosomal recessive childhood-onset severe retinal dystrophy. *Nature Genet.* 17: 194–197, 1997.

[77564] 15894. Hamel, C. P.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Redmond, T. M.: The gene for the retinal pigment epithelium-specific protein RPE65 is localized to human 1p31 and mouse 3. *Genomics* 20:509–512, 1994.

[77565] 15895. Hamel, C. P.; Tsilou, E.; Pfeiffer, B. A.; Hooks, J. J.; Detrick, B.; Redmond, T. M.: Molecular cloning and expression of RPE65, a novel retinal pigment epithelium-specific microsomal protein that is post-transcriptionally regulated in vitro. *J. Biol. Chem.* 268:15751–15757, 1993.

[77566] 15896. Marlhens, F.; Bareil, C.; Griffoin, J.-M.; Zrenner, E.;

Amalric,P.; Eliaou, C.; Liu, S.-Y.; Harris, E.; Redmond, T. M.; Arnaud, B.;Claustres, M.; Hamel, C. P.: Mutations in RPE65 cause Leber's congenitalamaurosis. (Letter) Nature Genet. 17: 139–141, 1997.

[77567] 15897.Morimura, H.; Fishman, G. A.; Grover, S. A.; Fulton, A. B.; Berson,E. L.; Dryja, T. P.: Mutations in the RPE65 gene in patients withautosomal recessive retinitis pigmentosa or Leber congenital amaurosis. Proc.Nat. Acad. Sci. 95: 3088–3093, 1998.

[77568] 15898.Narfstrom, K.; Wrigstad, A.; Nilsson, S. E.: The Briard dog:a new animal model of congenital stationary night blindness. Brit.J. Ophthal. 73: 750–756, 1989.

[77569] 15899.Nicoletti, A.; Wong, D. J.; Kawase, K.; Gibson, L. H.; Yang-Feng,T. L.; Richards, J. E.; Thompson, D. A.: Molecular characterizationof the human gene encoding an abundant 61 kDa protein specific tothe retinal pigment epithelium. Hum. Molec. Genet. 4: 641–649, 1995.

[77570] 15900.Adeokun, A. M.; West, S. P.; Ellis, F. R.; Halsall, P. J.; Hopkins,P. M.; Foroughmand, A. M.; Iles, D. E.; Robinson, R. L.; Stewart,A. D.; Curran, J. L.: The G1021A substitution in the RYR1 gene doesnot cosegregate with malignant hyperthermia susceptibility in a Britishpedigree. Am. J. Hum. Genet. 60: 833–341, 1997.

- [77571] 15901.Alestrom, A.; Fagerlund, T. H.; Berg, K.: A simple method to detect the RYR1 mutation G1021A, a cause of malignant hyperthermia susceptibility. *Clin.Genet.* 47: 274–275, 1995.
- [77572] 15902.Avila, G.; O'Brien, J. J.; Dirksen, R. T.: Excitation–contraction uncoupling by a human central core disease mutation in the ryanodine receptor. *Proc. Nat. Acad. Sci.* 98: 4215–4220, 2001.
- [77573] 15903.Barone, V.; Bertocchini, F.; Bottinelli, R.; Protasi, F.; Allen, P. D.; Armstrong, C. F.; Reggiani, C.; Sorrentino, V.: Contractile impairment and structural alterations of skeletal muscles from knockout mice lacking type 1 and type 3 ryanodine receptors. *FEBS Lett.* 422:160–164, 1998.
- [77574] 15904.Brandt, A.; Schleithoff, L.; Jurkat-Rott, K.; Klingler, W.; Baur, C.; Lehmann-Horn, F.: Screening of ryanodine receptor gene in 105 malignant hyperthermia families: novel mutations and concordance with the in vitro contracture test. *Hum. Molec. Genet.* 8: 2055–2062, 1999.
- [77575] 15905.Brown, R. L.; Pollock, A. N.; Couchman, K. G.; Hodges, M.; Hutchinson, D. O.; Waaka, R.; Lynch, P.; McCarthy, T. V.; Stowell, K. M.: A novel ryanodine receptor mutation and genotype–phenotype correlation in a large malignant hyperthermia New Zealand Maori pedigree.

Hum. Molec.Genet. 9: 1515–1524, 2000.

- [77576] 15906.Cavanna, J. S.; Greenfield, A. J.; Johnson, K. J.; Marks, A. R.;Nadal-Ginard, B.; Brown, S. D. M.: Establish-  
ment of the mouse chromosome7 region with homology  
to the myotonic dystrophy region of human chromo-  
some19q. Genomics 7: 12–18, 1990.
- [77577] 15907.Cheng, H.; Lederer, W. J.; Cannell, M. B.: Calcium  
sparks: elementaryevents underlying excitation–contrac-  
tion coupling in heart muscle. Science 262:740–744,  
1993.
- [77578] 15908.Deufel, T.; Sudbrak, R.; Feist, Y.; Rubsam, B.; Du  
Chesne, I.;Schafer, K.–L.; Roewer, N.; Grimm, T.;  
Lehmann–Horn, F.; Hartung,E. J.; Muller, C. R.: Discor-  
dance, in a malignant hyperthermia pedigree,between in  
vitro contracture–test phenotypes and haplotypes for  
theMHS1 region on chromosome 19q12–13.2, comprising  
the C1840T transitionin the RYR1 gene. Am. J. Hum.  
Genet. 56: 1334–1342, 1995.
- [77579] 15909.Eu, J. P.; Sun, J.; Xu, L.; Stamler, J. S.; Meissner, G.:  
Theskeletal muscle calcium release channel: coupled O<sub>2</sub>  
sensor and NOsignaling functions. Cell 102: 499–509,  
2000.
- [77580] 15910.Fagerlund, T.; Ording, H.; Bendixen, D.; Berg, K.:

Search for three known mutations in the RYR gene in 48 Danish families with malignant hyperthermia. Clin. Genet. 46: 401–404, 1994.

[77581] 15911.Fagerlund, T.; Ording, H.; Bendixen, D.; Islander, G.; Ranklev-Twetman, E.; Berg, K.: RYR1 mutation G1021A (gly341-to-arg) is not frequent in Danish and Swedish families with malignant hyperthermia susceptibility. Clin. Genet. 49: 186–188, 1996.

[77582] 15912.Fagerlund, T. H.; Islander, G.; Ranklev-Twetman, E.; Berg, K.: Recombination between the postulated CCD/MHE/MHS locus and RYR1 gene markers. Clin. Genet. 50: 455–458, 1996.

[77583] 15913.Fagerlund, T. H.; Islander, G.; Twetman, E. R.; Berg, K.: A search for three known RYR1 gene mutations in 41 Swedish families with predisposition to malignant hyperthermia. Clin. Genet. 48: 12–16, 1995.

[77584] 15914.Fagerlund, T. H.; Ording, H.; Bendixen, D.; Islander, G.; Ranklev-Twetman, E.; Berg, K.: Discordance between malignant hyperthermia susceptibility and RYR1 mutation C1840T in two Scandinavian MH families exhibiting this mutation. Clin. Genet. 52: 416–421, 1997.

[77585] 15915.Ferreiro, A.; Monnier, N.; Romero, N. B.; Leroy, J.-P.; Bonnemann, C.; Haenggeli, C.-A.; Straub, V.; Voss, W.

D.; Nivoche, Y.; Jungbluth, H.; Lemainque, A.; Voit, T.; Lunardi, J.; Fardeau, M.; Guicheney, P.: A recessive form of central core disease, transiently presenting as multiminicore disease, is associated with a homozygous mutation in the ryanodine receptor type 1 gene. *Ann. Neurol.* 51: 750–759, 2002.

[77586] 15916. Fujii, J.; Otsu, K.; Zorzato, F.; de Leon, S.; Khanna, V. K.; Weiler, J. E.; O'Brien, P. J.; MacLennan, D. H.: Identification of a mutation in porcine ryanodine receptor associated with malignant hyperthermia. *Science* 253: 448–451, 1991.

[77587] 15917. Gillard, E. F.; Otsu, K.; Fujii, J.; Duff, C.; de Leon, S.; Khanna, V. K.; Britt, B. A.; Worton, R. G.; MacLennan, D. H.: Polymorphisms and deduced amino acid substitutions in the coding sequence of the ryanodine receptor (RYR1) gene in individuals with malignant hyperthermia. *Genomics* 13: 1247–1254, 1992.

[77588] 15918. Gillard, E. F.; Otsu, K.; Fujii, J.; Khanna, V. K.; de Leon, S.; Derdemezi, J.; Britt, B. A.; Duff, C. L.; Worton, R. G.; MacLennan, D. H.: A substitution of cysteine for arginine 614 in the ryanodine receptor is potentially causative of human malignant hyperthermia. *Genomics* 11: 751–755, 1991.



- [77589] 15919.Hall–Curran, J. L.; Stewart, A. D.; Ball, S. P.; Halsall, J. P.;Hopkins, P. M.; Ellis, F. R.: No C1840 to T mutation in RYR1 in malignanthyperthermia.(Letter) Hum. Mutat. 2: 330, 1993.
- [77590] 15920.Quane, K. A.; Keating, K. E.; Manning, B. M.; Healy, J. M. S.;Monsieurs, K.; Heffron, J. J. A.; Lehane, M.; Heytens, L.; Krivosic–Horber,R.; Adnet, P.; Ellis, F. R.; Monnier, N.; Lunardi, J.; McCarthy, T.V.: Detection of a novel common mutation in the ryanodine receptorgene in malignant hyperthermia: implications for diagnosis and heterogeneitystudies. Hum. Molec. Genet. 3: 471–476, 1994.
- [77591] 15921.Vincenti, V.; Cassano, C.; Rocchi, M.; Persico, M. G.: Assignmentof the vascular endothelial growth factor gene to human chromosome6p21.3. Circulation 93: 1493–1495, 1996.
- [77592] 15922.Alonso, M.; Barton, D. E.; Francke, U.: MAL, a membrane proteinassociated with human T–cell differentiation, is encoded on humanchromosome 2, region cen–q13. (Abstract) Cytogenet. Cell Genet. 46:570 only, 1987.
- [77593] 15923.Alonso, M. A.; Barton, D. E.; Francke, U.: Assignment of the T–celldifferentiation gene MAL to human chromosome 2, region cen–q13. Immunogenetics

27:91–95, 1988.

- [77594] 15924.Alonso, M. A.; Weissman, S. M.: cDNA cloning and sequence of MAL,a hydrophobic protein associated with human T–cell differentiation. *Proc.Nat. Acad. Sci.* 84: 1997–2001, 1987.
- [77595] 15925.Millan, J.; Puertollano, R.; Fan, L.; Rancano, C.; Alonso, M. A.: The MAL proteolipid is a component of the detergent–insoluble membranesubdomains of human T–lymphocytes. *Biochem. J.* 321: 247–252, 1997.
- [77596] 15926.Rancano, C.; Rubio, T.; Alonso, M. A.: Alternative splicing ofhuman T–cell–specific MAL mRNA and its correlation with the exon/intronorganization of the gene. *Genomics* 21: 447–450, 1994.
- [77597] 15927.Wong–Staal, F.; Dalla–Favera, R.; Franchini, G.; Gelmann, E. P.;Gallo, R. C.: Three distinct genes in human DNA related to the transforminggenes of mammalian sarcoma retroviruses. *Science* 213: 226–228, 1981.
- [77598] 15928.Aurias, A.; Rimbaut, C.; Buffe, D.; Dubousset, J.; Mazabraud, A.: Chromosomal translocations in Ewing's sarcoma. (Letter) *New Eng.J. Med.* 309: 496–497, 1983.
- [77599] 15929.Bartram, C. R.; de Klein, A.; Hagemeijer, A.; Grosveld, G.; Heisterkamp,N.; Groffen, J.: Localization of the human c–sis oncogene in Ph–1–positiveand Ph–

1-negative chronic myelocytic leukemia by in situ hybridization. *Blood* 63:223-225, 1984.

[77600] 15930.Bechet, J.-M.; Bornkamm, G.; Freese, U.-K.; Lenoir, G. M.: The c-sis oncogene is not activated in Ewing's sarcoma. (Letter) *New Eng. J. Med.* 310: 393 only, 1984.

[77601] 15931.Bishop, J. M.: Enemies within: the genesis of retrovirus oncogenes. *Cell* 23:5-6, 1981.

[77602] 15932.Bolger, G. B.; Stamberg, J.; Kirsch, I. R.; Hollis, G. F.; Schwarz, D. F.; Thomas, G. H.: Chromosomal translocation t(14;22) and oncogene(c-sis) variant in a pedigree with familial meningioma. *New Eng.J. Med.* 312: 564-567, 1985.

[77603] 15933.Cohen, J. B.; Levinson, A. D.: A point mutation in the last intron responsible for increased expression and transforming activity of the c-Ha-ras oncogene. *Nature* 334: 119-124, 1988.

[77604] 15934.Collins, T.; Ginsburg, D.; Boss, J. M.; Orkin, S. H.; Pober, J.S.: Cultured human endothelial cells express platelet-derived growth factor B chain: cDNA cloning and structural analysis. *Nature* 316:748-750, 1985.

[77605] 15935.Dalla-Favera, R.; Gallo, R. C.; Giallongo, A.; Croce, C.: Chromosomal localization of the human homolog (c-sis) of the simian sarcoma virus onc gene. *Science* 218:

686–688, 1982.

- [77606] 15936.Dalla-Favera, R.; Gelmann, E. P.; Gallo, R. C.; Wong-Staal, F.: A human onc gene homologous to the transforming gene (v-sis) of simian sarcoma virus. *Nature* 292: 31–35, 1981.
- [77607] 15937.Deuel, T. F.; Huang, J. S.; Huang, S. S.; Stroobant, P.; Waterfield, M. D.: Expression of a platelet-derived growth factor-like protein in simian sarcoma virus transformed cells. *Science* 221: 1348–1350, 1983.
- [77608] 15938.Devare, S. G.; Reddy, E. P.; Law, J. D.; Robbins, K. C.; Aaronson, S. A.: Nucleotide sequence of the simian sarcoma virus genome: demonstration that its acquired cellular sequences encode the transforming gene product p28-sis. *Proc. Nat. Acad. Sci.* 80: 731–735, 1983.
- [77609] 15939.Charlet-B., N.; Logan, P.; Singh, G.; Cooper, T. A.: Dynamic antagonism between ETR-3 and PTB regulates cell type-specific alternative splicing. *Molec. Cell* 9: 649–658, 2002.
- [77610] 15940.Borglum, A. D.; Flint, T.; Tommerup, N.; Fleckner, J.; Justesen, J.; Kruse, T. A.: Assignment of the human tryptophanyl-tRNA synthetase gene (WARS) to chromosome 14q32.2–q32.32. *Cytogenet. Cell Genet.* 73:99–103, 1996.

- [77611] 15941.Denney, R. M.; Borgaonkar, D.; Ruddle, F. H.: Order of genes for NP and TRPRS on chromosome 14. *Cytogenet. Cell Genet.* 22: 493–497, 1978.
- [77612] 15942.Francke, U.; Denney, R. M.; Ruddle, F. H.: Intra-chromosomal genemapping in man: the gene for tryptophanyl-tRNA synthetase maps in region q21-qter of chromosome 14. *Somat. Cell Genet.* 3: 381–389, 1977.
- [77613] 15943.Benson, K. F.; Horwitz, M.; Wolff, J.; Friend, K.; Thompson, E.; White, S.; Richards, R. I.; Raskind, W. H.; Bird, T. D.: CAG repeat expansion in autosomal dominant familial spastic paraparesis: novel expansion in a subset of patients. *Hum. Molec. Genet.* 7: 1779–1786, 1998.
- [77614] 15944.Fonknechten, N.; Mavel, D.; Byrne, P.; Davoine, C.-S.; Cruaud, C.; Boentsch, D.; Samson, D.; Coutinho, P.; Hutchinson, M.; McMonagle, P.; Burgunder, J.-M.; Tartaglione, A.; and 10 others: Spectrum of SPG4 mutations in autosomal dominant spastic paraplegia. *Hum. Molec. Genet.* 9: 637–644, 2000.
- [77615] 15945.Nielsen, J. E.; Koefoed, P.; Abell, K.; Hasholt, L.; Eiberg, H.; Fenger, K.; Niebuhr, E.; Sorensen, S. A.: CAG repeat expansion in autosomal dominant pure spastic paraplegia linked to chromosome 2p21-p24. *Hum. Molec. Genet.* 6: 1811–1816, 1997.

- [77616] 15946.Svenson, I. K.; Ashley-Koch, A. E.; Gaskell, P. C.; Riney, T.J.; Cumming, W. J. K.; Kingston, H. M.; Hogan, E. L.; Boustany, R.-M.N.; Vance, J. M.; Nance, M. A.; Pericak-Vance, M. A.; Marchuk, D.A.: Identification and expression analysis of spastin gene mutations in hereditary spastic paraplegia. *Am. J. Hum. Genet.* 68: 1077–1085, 2001.
- [77617] 15947.Ala-Kapee, M.; Nevanlinna, H.; Mali, M.; Jalkanen, M.; Schroder, J.: Localization of gene for human syndecan, an integral membrane proteoglycan and a matrix receptor, to chromosome 2. *Somat. Cell Molec. Genet.* 16: 501–505, 1990.
- [77618] 15948.Alexander, C. M.; Reichsman, F.; Hinkes, M. T.; Lincecum, J.; Becker, K. A.; Cumberledge, S.; Bernfield, M.: Syndecan-1 is required for Wnt-1-induced mammary tumorigenesis in mice. *Nature Genet.* 25: 329–332, 2000.
- [77619] 15949.Mali, M.; Jaakkola, P.; Arvilommi, A.-M.; Jalkanen, M.: Sequence of human syndecan indicates a novel gene family of integral membrane proteoglycans. *J. Biol. Chem.* 265: 6884–6889, 1990.
- [77620] 15950.Oettinger, H. F.; Streeter, H.; Lose, E.; Copeland, N. G.; Gilbert, D. J.; Justice, M. J.; Jenkins, N. A.; Mohandas, T.; Bernfield, M.: Chromosome mapping of the murine syndecan gene. *Genomics* 11: 334–338, 1991.

- [77621] 15951.Reizes, O.; Lincecum, J.; Wang, Z.; Goldberger, O.; Huang, L.;Kaksonen, M.; Ahima, R.; Hinkes, M. T.; Barsh, G. S.; Rauvala, H.;Bernfield, M.: Transgenic expression of syndecan-1 uncovers a physiological control of feeding behavior by syndecan-3. *Cell* 106: 105-116, 2001.
- [77622] 15952.Sanderson, R. D.; Lalor, P.; Bernfield, M.: B lymphocytes express and lose syndecan at specific stages of differentiation. *Cell Regulation* 1:27-35, 1989.
- [77623] 15953.Westman, P.; Hsieh, C.-L.; Mali, M.; Jalkanen, M.; Francke, U.;Schroder, J.: Assignment of the human syndecan (SDC) gene to short arm of chromosome 2. (Abstract) *Cytogenet. Cell Genet.* 58: 1873-1874, 1991.
- [77624] 15954.Asch, A. S.; Barnwell, J.; Silverstein, R. L.; Nachman, R. L.: Isolation of the thrombospondin membrane receptor. *J. Clin. Invest.* 79:1054-1061, 1987.
- [77625] 15955.de Fraipont, F.; El Atifi, M.; Gicquel, C.; Bertagna, X.; Chambaz, E. M.; Feige, J. J.: Expression of the angiogenesis markers vascular endothelial growth factor-A, thrombospondin-1, and platelet-derived endothelial cell growth factor in human sporadic adrenocortical tumors: correlation with genotypic alterations. *J. Clin. Endocr. Metab.* 85:4734-4741, 2000.
- [77626] 15956.Dixit, V. M.; Hennessy, S. W.; Grant, G. A.; Rotwein,

P.; Frazier, W. A.: Characterization of a cDNA encoding the heparin and collagenbinding domains of human thrombospondin. *Proc. Nat. Acad. Sci.* 83:5449–5453, 1986.

[77627] 15957. Frazier, W. A.: Thrombospondin: a modular adhesive glycoprotein of platelets and nucleated cells. *J. Cell Biol.* 105: 625–632, 1987.

[77628] 15958. Jaffe, E.; Bornstein, P.; Disteché, C. M.: Mapping of the thrombospondin gene to human chromosome 15 and mouse chromosome 2 by in situ hybridization. *Genomics* 7:123–126, 1990.

[77629] 15959. Lawler, J.; Sunday, M.; Thibert, V.; Duquette, M.; George, E. L.; Rayburn, H.; Hynes, R. O.: Thrombospondin-1 is required for normal murine pulmonary homeostasis and its absence causes pneumonia. *J. Clin. Invest.* 101: 982–992, 1998.

[77630] 15960. Rodríguez-Manzanique, J. C.; Lane, T. F.; Ortega, M. A.; Hynes, R. O.; Lawler, J.; Iruela-Arispe, M. L.: Thrombospondin-1 suppresses spontaneous tumor growth and inhibits activation of matrix metalloproteinase-9 and mobilization of vascular endothelial growth factor. *Proc. Nat. Acad. Sci.* 98: 12485–12490, 2001.

[77631] 15961. Wolf, F. W.; Eddy, R. L.; Shows, T. B.; Dixit, V. M.: Structure and chromosomal localization of the human



thrombospondin gene. *Genomics* 6:685–691, 1990.

[77632] 15962.Imai, K.; Sarker, A. H.; Akiyama, K.; Ikeda, S.; Yao, M.; Tsutsui,K.; Shohmori, T.; Seki, S.: Genomic structure and sequence of a humanhomologue (NTHL1/NTH1) of *Escherichia coli* endonuclease III with thoseof the adjacent parts of TSC2 and SLC9A3R2 genes. *Gene* 222:

287–295,1998.

[77633] 15963.Mira, J.–P.; Cariou, A.; Grall, F.; Delclaux, C.; Losser, M.–R.;Heshmati, F.; Cheval, C.; Monchi, M.; Teboul, J.–L.; Riche, F.; Leleu,G.; Arbibe, L.; Mignon, A.; Delpech, M.; Dhainaut, J.–F.: Associationof TNF2, a TNF–alpha promoter polymorphism, with septic shock susceptibilityand mortality: a multicenter study. *J.A.M.A.* 282: 561–568, 1999.

[77634] 15964.Moffatt, M. F.; Cookson, W. O. C. M.: Tumour necrosis factorhaplotypes and asthma. *Hum. Molec. Genet.* 6: 551–554, 1997.

[77635] 15965.Moraes, M. O.; Duppre, N. C.; Suffys, P. N.; Santos, A. R.; Almeida,A. S.; Nery, J. A. C.; Sampaio, E. P.; Sarno, E. N.: Tumor necrosisfactor–alpha promoter polymorphism TNF2 is associated with a strongerdelayed–type hypersensitivity reaction in the skin of borderline tuberculoi–dleprosy patients. *Immunogenetics* 53: 45–47, 2001.

- [77636] 15966. Mulcahy, B.; Waldron-Lynch, F.; McDermott, M. F.; Adams, C.; Amos, C. I.; Zhu, D. K.; Ward, R. H.; Clegg, D. O.; Shanahan, F.; Molloy, M. G.; O'Gara, F.: Genetic variability in the tumor necrosis factor-lymphotoxin region influences susceptibility to rheumatoid arthritis. *Am. J. Hum. Genet.* 59: 676-683, 1996.
- [77637] 15967. Muller, U.; Jongeneel, C. V.; Nedospasov, S. A.; Lindahl, K. F.; Steinmetz, M.: Tumour necrosis factor and lymphotoxin genes map close to H-2D in the mouse major histocompatibility complex. *Nature* 325:265-267, 1987.
- [77638] 15968. Nadel, S.; Newport, M. J.; Booy, R.; Levin, M.: Variation in the tumor necrosis factor-alpha gene promoter region may be associated with death from meningococcal disease. *J. Infect. Dis.* 174: 878-880, 1996.
- [77639] 15969. Nedospasov, S. A.; Hirt, B.; Shakhov, A. N.; Dobrynin, V. N.; Kawashima, E.; Accolla, R. S.; Jongeneel, C. V.: The genes for tumor necrosis factor (TNF-alpha) and lymphotoxin (TNF-beta) are tandemly arranged on chromosome 17 of the mouse. *Nucleic Acids Res.* 14: 7713-7725, 1986.
- [77640] 15970. Nedwin, G. E.; Naylor, S. L.; Sakaguchi, A. Y.; Smith, D.; Jarrett-Nedwin, J.; Pennica, D.; Goeddel, D. V.; Gray, P. W.: Human lymphotoxin and tumor necrosis factor genes:

structure, homology and chromosomal localization. *Nucleic Acids Res.* 13: 6361–6373, 1985.

[77641] 15971. Norman, R. A.; Bogardus, C.; Ravussin, E.: Linkage between obesity and a marker near the tumor necrosis factor- $\alpha$  locus in Pima Indians. *J. Clin. Invest.* 96: 158–162, 1995.

[77642] 15972. Obayashi, H.; Hasegawa, G.; Fukui, M.; Kamiuchi, K.; Kitamura, A.; Ogata, M.; Kanatsuka, T.; Shigeta, H.; Kitagawa, Y.; Nakano, K.; Nishimura, M.; Ohta, M.; Nakamura, N.: Tumor necrosis factor microsatellite polymorphism influences the development of insulin dependency in adult-onset diabetes patients with the DRB1\*1502–DQB1\*0601 allele and anti-glutamic acid decarboxylase antibodies. *J. Clin. Endocr. Metab.* 85: 3348–3351, 2000.

[77643] 15973. Old, L. J.: Tumor necrosis factor (TNF). *Science* 230: 630–632, 1985.

[77644] 15974. Pennica, D.; Nedwin, G. E.; Hayflick, J. S.; Seeburg, P. H.; Derynck, R.; Palladino, M. A.; Kohr, W. J.; Aggarwal, B. B.; Goeddel, D. V.: Human tumour necrosis factor: precursor structure, expression and homology to lymphotoxin. *Nature* 312: 724–729, 1984.

[77645] 15975. Ragoussis, J.; Bloemer, K.; Weiss, E. H.; Ziegler, A.:

Localization of the genes for tumor necrosis factor and lymphotoxin between the HLA class I and III regions by field inversion gel electrophoresis. *Immunogenetics* 27:66–69, 1988.

- [77646] 15976. Rasmussen, S. K.; Urhammer, S. A.; Jensen, J. N.; Hansen, T.; Borch-Johnsen, K.; Pedersen, O.: The –238 and –308 G6A polymorphisms of the tumor necrosis factor alpha gene promoter are not associated with features of the insulin resistance syndrome or altered birthweight in Danish Caucasians. *J. Clin. Endocr. Metab.* 85: 1731–1734, 2000.
- [77647] 15977. Rosmond, R.; Chagnon, M.; Bouchard, C.; Bjorntorp, P.: G–308 A polymorphism of the tumor necrosis factor alpha gene promoter and salivary cortisol secretion. *J. Clin. Endocr. Metab.* 86: 2178–2180, 2001.
- [77648] 15978. Roy, S.; McGuire, W.; Mascie-Taylor, C. G.; Saha, B.; Hazra, S. K.; Hill, A. V.; Kwiatkowski, D.: Tumor necrosis factor promoter polymorphism and susceptibility to lepromatous leprosy. *J. Infect. Dis.* 176: 530–532, 1997.
- [77649] 15979. Ruuls, S. R.; Sedgwick, J. D.: Unlinking tumor necrosis factor biology from the major histocompatibility complex: lessons from human genetics and animal models. *Am. J. Hum. Genet.* 65: 294–301, 1999.

- [77650] 15980.Spies, T.; Morton, C. C.; Nedospasov, S. A.; Fiers, W.; Pious,D.; Strominger, J. L.: Genes for the tumor necrosis factors alphaand beta are linked to the human major histocompatibility complex. Proc.Nat. Acad. Sci. 83: 8699–8702, 1986.
- [77651] 15981.van Hensbroek, M. B.; Palmer, A.; Onyiorah, E.; Schneider, G.;Jaffar, S.; Dolan, G.; Memming, H.; Frenkel, J.; Enwere, G.; Bennett,S.; Kwiatkowski, D.; Greenwood, B.: The effect of a monoclonal antibodyto tumor necrosis factor on survival from childhood cerebral malaria. J.Infect. Dis. 174: 1091–1097, 1996.
- [77652] 15982.Sashio, H.; Tamura, K.; Ito, R.; Yamamoto, Y.; Bamba, H.; Kosaka,T.; Fukui, S.; Sawada, K.; Fukuda, Y.; Tamura, K.; Satomi, M.; Shimoyama,T.; Furuyama, J.: Polymorphisms of the TNF gene and the TNF receptorsuperfamily member 1B gene are associated with susceptibility to ulcerativecolitis and Crohn's disease, respectively. Immunogenetics 53: 1020–1027,2002.
- [77653] 15983.Van Ostade, X.; Vandenabeele, P.; Everaerd, B.; Loetscher, H.;Gentz, R.; Brockhaus, M.; Lesslauer, W.; Tavernier, J.; Brouckaert,P.; Fiers, W.: Human TNF mutants with selective activity on the p55receptor. Nature 361: 266–269, 1993.

- [77654] 15984.Wang, A. M.; Creasey, A. A.; Ladner, M. B.; Lin, L. S.; Strickler,J.; Van Arsdell, J. N.; Yamamoto, R.; Mark, D. F.: Molecular cloningof the complementary DNA for human tumor necrosis factor. *Science* 228:149–154, 1985.
- [77655] 15985.Wilson, A. G.; Symons, J. A.; McDowell, T. L.; et al: Effectsof a polymorphism in the human tumor necrosis factor alpha promoter on transcriptional activation. *Proc. Nat. Acad. Sci.* 94: 3195–3199,1997.
- [77656] 15986.Winchester, E. C.; Millwood, I. Y.; Rand, L.; Penny, M. A.; Kessling,A. M.: Association of the TNF–alpha–308 (G–A) polymorphism with self–reportedhistory of childhood asthma. *Hum. Genet.* 107: 591–596, 2000.
- [77657] 15987.Witte, J. S.; Palmer, L. J.; O'Connor, R. D.; Hopkins, P. J.;Hall, J. M.: Relation between tumour necrosis factor polymorphismTNF–alpha–308 and risk of asthma. *Europ. J. Hum. Genet.* 10: 82–85,2002.
- [77658] 15988.Peter, D.; Finn, J. P.; Klisak, I.; Liu, Y.; Kojis, T.; Heinzmann,C.; Roghani, A.; Sparkes, R. S.; Edwards, R. H.: Chromosomal localizationof the human vesicular amine transporter genes. *Genomics* 18: 720–723,1993.
- [77659] 15989.Roghani, A.; Welch, C.; Xia, Y.–R.; Liu, Y.; Peter, D.; Finn, J.P.; Edwards, R. H.; Lusi, A. J.: Assignment of the mouse vesicularmonoamine transporter genes, *Slc18a1*

and Slc18a2, to chromosomes 8 and 19 by linkage analysis. *Mammalian Genome* 7: 393–394, 1996.

- [77660] 1990. Lania, L.; Donti, E.; Pannuti, A.; Pascucci, A.; Pengue, G.; Feliciello, I.; La Mantia, G.; Lanfranccone, L.; Pelicci, P.-G.: cDNA isolation, expression analysis, and chromosomal localization of two human zinc finger genes. *Genomics* 6: 333–340, 1990.
- [77661] 1991. Thiesen, H.-J.: Multiple genes encoding zinc finger domains are expressed in human T cells. *New Biologist* 2: 363–374, 1990.
- [77662] 1992. Seite, P.; Huebner, K.; Rousseau-Merck, M. F.; Berger, R.; Thiesen, H. J.: Two human genes encoding zinc finger proteins, ZNF12 (KOX3) and ZNF26 (KOX20), map to chromosomes 7p22–p21 and 12q24.33, respectively. *Hum. Genet.* 86: 585–590, 1991.
- [77663] 1993. Gaynor, R. B.; Muchardt, C.; Diep, A.; Mohandas, T. K.; Sparkes, R. S.; Lusk, A. J.: Localization of the zinc finger DNA-binding protein HIV-EP1/MBP-1/PRDII-BF1 to human chromosome 6p22.3–p24. *Genomics* 9: 758–761, 1991.
- [77664] 1994. Tripathi, R. K.; Droetto, S.; Spritz, R. A.: Many patients with 'tyrosinase-positive' oculocutaneous albinism have tyrosinase gene mutations. (Abstract) *Am. J. Hum.*

Genet. 51 (suppl.): A179, 1992.

- [77665] 15995.Brilliant, M. H.: The mouse pink-eyed dilution locus: a model for aspects of Prader-Willi syndrome, Angelman syndrome, and a form of hypomelanosis of Ito. *Mammalian Genome* 3: 187-191, 1992.
- [77666] 15996.Castle, D.; Kromberg, J.; Kowalsky, R.; Moosa, R.; Gillman, N.; Zwane, E.; Fritz, V.: Visual evoked potentials in Negro carriers of the gene for tyrosinase positive oculocutaneous albinism. *J. Med. Genet.* 25: 835-837, 1988.
- [77667] 15997.Colman, M. A.; Stevens, G.; Ramsay, M.; Kwon, B.; Jenkins, T.: Exclusion of two candidate pigment loci, c and b, part of chromosome 11p, and 33 random polymorphic markers as the locus for tyrosinase-positive oculocutaneous albinism. *Hum. Genet.* 90: 556-560, 1993.
- [77668] 15998.Donlon, T. A.; Lalande, M.; Wyman, A.; Bruns, G.; Latt, S. A.: Isolation of molecular probes associated with the chromosome 15 instability in the Prader-Willi syndrome. *Proc. Nat. Acad. Sci.* 83: 4408-4412, 1986.
- [77669] 15999.Durham-Pierre, D.; Gardner, J. M.; Nakatsu, Y.; King, R. A.; Francke, U.; Ching, A.; Aquaron, R.; del Mar-mol, V.; Brilliant, M. H.: African origin of an intragenic deletion of the human P gene in tyrosinase positive oculo-



cutaneous albinism. *Nature Genet.* 7: 176–179, 1994.

[77670] 16000.Durham–Pierre, D.; King, R. A.; Naber, J. M.; Laken, S.; Brilliant, M. H.: Estimation of carrier frequency of a 2.7 kb deletion allele of the P gene associated with OCA2 in African–Americans. *Hum. Mutat.* 7:370–373, 1996.

[77671] 16001.Gardner, J. M.; Nakatsu, Y.; Gondo, Y.; Lee, S.; Lyon, M. F.; King, R. A.; Brilliant, M. H.: The mouse pink-eyed dilution gene: association with human Prader–Willi and Angelman syndromes. *Science* 257: 1121–1124, 1992.

[77672] 16002.Gondo, Y.; Gardner, J. M.; Nakatsu, Y.; Durham–Pierre, D.; Deveau, S. A.; Kuper, C.; Brilliant, M. H.: High-frequency genetic reversion mediated by a DNA duplication: the mouse pink-eyed unstable mutation. *Proc. Nat. Acad. Sci.* 90: 297–301, 1993.

[77673] 16003.Hagiwara, N.; Klewer, S. E.; Samson, R. A.; Erickson, D. T.; Lyon, M. F.; Brilliant, M. H.: Sox6 is a candidate gene for p100H myopathy, heart block, and sudden neonatal death. *Proc. Nat. Acad. Sci.* 97:4180–4185, 2000.

[77674] 16004.Hamabe, J.; Fukushima, Y.; Harada, N.; Abe, K.; Matsuo, N.; Nagai, T.; Yoshioka, A.; Tonoki, H.; Tsukino, R.; Niikawa, N.: Molecular study of the Prader–Willi syndrome: deletion, RFLP, and phenotype analyses of 50 patients. *Am. J. Med. Genet.* 41: 54–63, 1991.

- [77675] 16005.Heim, R. A.; Dunn, D. S.; Candy, S. E.; Zwane, E.; Kromberg, J.G. R.; Jenkins, T.: The tyrosine-positive oculocutaneous albinism locus is not linked to the beta-globin locus in man. *Hum. Genet.* 79:89, 1988.
- [77676] 16006.Kagore, F.; Lund, P. M.: Oculocutaneous albinism among schoolchildren in Harare, Zimbabwe. *J. Med. Genet.* 32: 859–861, 1995.
- [77677] 16007.Kedda, M. A.; Stevens, G.; Manga, P.; Viljoen, C.; Jenkins, T.; Ramsay, M.: The tyrosinase-positive oculocutaneous albinism gene shows locus homogeneity on chromosome 15q11–q13 and evidence of multiple mutations in Southern African negroids. *Am. J. Hum. Genet.* 54: 1078–1084, 1994.
- [77678] 16008.Kerr, R.; Stevens, G.; Manga, P.; Salm, S.; John, P.; Haw, T.; Ramsay, M.: Identification of P gene mutations in individuals with oculocutaneous albinism in sub-Saharan Africa. *Hum. Mutat.* 15: 166–172, 2000.
- [77679] 16009.King, R. A.: Personal Communication. Minneapolis, Minn. 12/31/1992.
- [77680] 16010.Ansari-Lari, M. A.; Muzny, D. M.; Lu, J.; Lu, F.; Lilley, C. E.; Spanos, S.; Malley, T.; Gibbs, R. A.: A gene-rich cluster between the CD4 and triosephosphate isomerase genes at human chromosome 12p13. *Genome Res.* 6:

314–326, 1996.

- [77681] 16011. Bach, M. A.; Phan–Dinh–Tuy, F.; Bach, J. F.; Wallach, D.; Biddison, W. E.; Sharrow, S. O.; Goldstein, G.; Kung, P. C.: Unusual phenotypes of human inducer T cells as measured by OKT4 and related monoclonal antibodies. *J. Immunol.* 127: 980–982, 1981.
- [77682] 16012. Browning, J.; Horner, J. W.; Pettoello–Mantovani, M.; Raker, C.; Yurasov, S.; DePinho, R. A.; Goldstein, H.: Mice transgenic for human CD4 and CCR5 are susceptible to HIV infection. *Proc. Nat. Acad. Sci.* 94:14637–14641, 1997.
- [77683] 16013. Buttini, M.; Westland, C. E.; Masliah, E.; Yafeh, A. M.; Wyss–Coray, T.; Mucke, L.: Novel role of human CD4 molecule identified in neurodegeneration. *Nature Med.* 4: 441–446, 1998.
- [77684] 16014. Committee on Human Leukocyte Differentiation Antigens, IUIS WHO Nomenclature Subcommittee: Proposed nomenclature for human leukocyte differentiation antigens. *Bull. WHO* 5: 809–811, 1984. Note: *Alternate: Immunology Today* 5: 280 only, 1984.
- [77685] 16015. Fukuda, T.; Matsunaga, M.; Kurata, A.; Mine, M.; Ikari, N.; Katamine, S.; Kanazawa, H.; Eguchi, K.; Nagataki, S.: Hereditary deficiency of OKT4–positive cells: studies for

mode of inheritance and lymphocyte functions. Immunology 53: 643–649, 1984.

[77686] 16016.Fuller, T. C.; Trevithick, J. E.; Fuller, A. A.; Colvin, R. B.;Cosimi, A. B.; Kung, P. C.: Antigenic polymorphism of the T4 differentiation antigen expressed on human T helper/inducer lymphocytes. Hum. Immun. 9:89–102, 1984.

[77687] 16017.Gill, J. C.; Maples, J.; Nikaein, A.; Kirchner, P.; Lockhart, D.;Snyder, A. J.; Montgomery, R. R.; Casper, J. T.: Inherited absence of OKT4 lymphocyte antigen in a chronically transfused patient with homozygous sickle cell disease. J. Pediat. 107: 251–253, 1985.

[77688] 16018.Irvine, D. J.; Purbhoo, M. A.; Krogsgaard, M.; Davis, M. M.:Direct observation of ligand recognition by T cells. Nature 419:845–849, 2002.

[77689] 16019.Isobe, M.; Huebner, K.; Maddon, P. J.; Littman, D. R.; Axel, R.;Croce, C. M.: The gene encoding the T-cell surface protein T4 is located on human chromosome 12. Proc. Nat. Acad. Sci. 83: 4399–4402, 1986.

[77690] 16020.Karol, R. A.; Eng, J.; Dennison, D. K.; Faris, E.; Marcus, D.M.: Hereditary abnormalities of the OKT4 human lymphocyte epitope in two families. J. Clin. Immun. 4: 71–74, 1984.

- [77691] 16021.Kozbor, D.; Finan, J.; Nowell, P. C.; Croce, C. M.: The gene encoding the T4 antigen maps to human chromosome 12. *J. Immun.* 136:1141–1143, 1986.
- [77692] 16022.Maddon, P. J.; Molineaux, S. M.; Maddon, D. E.; Zimmerman, K.A.; Godfrey, M.; Alt, F. W.; Chess, L.; Axel, R.: Structure and expression of the human and mouse T4 genes. *Proc. Nat. Acad. Sci.* 84: 9155–9159, 1987.
- [77693] 16023.Piguet, V.; Gu, F.; Foti, M.; Demarex, N.; Gruenberg, J.; Carpentier, J.-L.; Trono, D.: Nef-induced CD4 degradation: a diacidic-based motif in Nef functions as a lysosomal targeting signal through the binding of beta-COP in endosomes. *Cell* 97: 63–73, 1999.
- [77694] 16024.Sato, M.; Hayashi, Y.; Yoshida, H.; Yanagawa, T.; Yura, Y.: A family with hereditary lack of T4+ inducer/helper T cell subsets in peripheral blood lymphocytes. *J. Immun.* 132: 1071–1073, 1984.
- [77695] 16025.Sawada, S.; Scarborough, J. D.; Killeen, N.; Littman, D. R.: A lineage-specific transcriptional silencer regulates CD4 gene expression during T lymphocyte development. *Cell* 77: 917–929, 1994.
- [77696] 16026.Zinman, B.; Hanley, A. J. G.; Harris, S. B.; Kwan, J.; Fantus, I. G.: Circulating tumor necrosis factor-alpha concentrations in a Native Canadian population with high rates

of type 2 diabetes mellitus. J.Clin. Endocr. Metab. 84: 272–278, 1999.

[77697] 16027.Spurr, N. K.; Goodfellow, P. N.; Sheer, D.; Bodmer, W. F.; Vennstrom,B.: Mapping of cellular oncogenes: ERBA1 is on chromosome 17. (Abstract) Cytogenet.Cell Genet. 37: 591 only, 1984.

[77698] 16028.Thompson, C. C.; Weinberger, C.; Lebo, R.; Evans, R. M.: Identificationof a novel thyroid hormone receptor expressed in the mammalian centralnervous system. Science 237: 1610–1614, 1987.

[77699] 16029.Weinberger, C.; Thompson, C. C.; Ong, E. S.; Lebo, R.; Gruol,D. J.; Evans, R. M.: The c-erb-A gene encodes a thyroid hormone receptor. Nature 324:641–646, 1986.

[77700] 16030.Zabel, B. U.; Fournier, R. E. K.; Lalley, P. A.; Naylor, S. L.;Sakaguchi, A. Y.: Cellular homologs of the avian erythroblastosisvirus erb-A and erb-B genes are syntenic in mouse but asyntenic inman. Proc. Nat. Acad. Sci. 81: 4874–4878, 1984.

[77701] 16031.Carraway, K. L., III; Sliwkowski, M. X.; Akita, R.; Platko, J.V.; Guy, P. M.; Nuijens, A.; Diamonti, A. J.; Vandlen, R. L.; Cantley,L. C.; Cerione, R. A.: The erbB3 gene product is a receptor for heregulin. J.Biol. Chem. 269: 14303–14306, 1994.

- [77702] 16032.Cho, H.-S.; Leahy, D. J.: Structure of the extracellular region of HER3 reveals an interdomain tether. *Science* 297: 1330–1333, 2002.
- [77703] 16033.Kraus, M. H.; Issing, W.; Miki, T.; Popescu, N. C.; Aaronson, S.A.: Isolation and characterization of ERBB3, a third member of the ERBB/epidermal growth factor receptor family: evidence for overexpression in a subset of human mammary tumors. *Proc. Nat. Acad. Sci.* 86: 9193–9197, 1989.
- [77704] 16034.Plowman, G. D.; Whitney, G. S.; Neubauer, M. G.; Green, J. M.; McDonald, V. L.; Todaro, G. J.; Shoyab, M.: Molecular cloning and expression of an additional epidermal growth factor receptor-related gene. *Proc. Nat. Acad. Sci.* 87: 4905–4909, 1990.
- [77705] 16035.Elchebly, M.; Payette, P.; Michaliszyn, E.; Cromlish, W.; Collins, S.; Loy, A. L.; Normandin, D.; Cheng, A.; Himms-Hagen, J.; Chan, C.-C.; Ramachandran, C.; Gresser, M. J.; Tremblay, M. L.; Kennedy, B. P.: Increased insulin sensitivity and obesity resistance in mice lacking the protein tyrosine phosphatase-1B gene. *Science* 283: 1544–1548, 1999.
- [77706] 16036.Forsell, P. K. A. L.; Boie, Y.; Montalibet, J.; Collins, S.; Kennedy, B. P.: Genomic characterization of the human

and mouse protein tyrosinephosphatase-1B genes. *Gene* 260: 145–153, 2000.

[77707] 16037.Gu, H. F.; Almgren, P.; Lindholm, E.; Frittitta, L.; Pizzuti, A.;Trischitta, V.; Groop, L. C.: Association between the human glycoproteinPC-1 gene and elevated glucose and insulin levels in paired-siblinganalysis. *Diabetes* 49: 1601–1603, 2000.

[77708] 16038.Haj, F. G.; Verveer, P. J.; Squire, A.; Neel, B. G.; Bastiaens,P. I. H.: Imaging sites of receptor dephosphorylation by PTP1B onthe surface of the endoplasmic reticulum. *Science* 295: 1708–1711,2002.

[77709] 16039.Jia, Z.; Barford, D.; Flint, A. J.; Tonks, N. K.: Structuralbasis for phosphotyrosine peptide recognition by protein tyrosinephosphatase 1B. *Science* 268: 1754–1758, 1995.

[77710] 16040.Kennedy, B. P.; Ramachandran, C.: Protein tyrosine phosphatase-1Bin diabetes. *Biochem. Pharm.* 60: 877–883, 2000.

[77711] 16041.Mok, A.; Cao, H.; Zinman, B.; Hanley, A. J. G.; Harris, S. B.;Kennedy, B. P.; Hegele, R. A.: A single nucleotide polymorphism inprotein tyrosine phosphatase PTP-1B is associated with protectionfrom diabetes or impaired glucose tolerance in Oji-Cree. *J. Clin.Endocr. Metab.* 87:



724–727, 2002.

- [77712] 16042. Tonks, N. K.; Diltz, C. D.; Fischer, E. H.: Purification of the major protein-tyrosine-phosphatases of human placenta. *J. Biol. Chem.* 263:6722–6730, 1988.
- [77713] 16043. Barnea, G.; Silvennoinen, O.; Shaanan, B.; Honegger, A. M.; Canoll, P. D.; D'Eustachio, P.; Morse, B.; Levy, J. B.; Laforgia, S.; Huebner, K.; Musacchio, J. M.; Sap, J.; Schlessinger, J.: Identification of a carbonic anhydrase-like domain in the extracellular region of RPTP-gamma defines a new subfamily of receptor tyrosine phosphatases. *Molec. Cell. Biol.* 13: 1497–1506, 1993.
- [77714] 16044. Kastury, K.; Ohta, M.; Lasota, J.; Moir, D.; Dorman, T.; LaForgia, S.; Druck, T.; Huebner, K.: Structure of the human receptor tyrosine phosphatase gamma gene (PTPRG) and relation to the familial RCC t(3;8) chromosome translocation. *Genomics* 32: 225–235, 1996.
- [77715] 16045. LaForgia, S.; Morse, B.; Levy, J.; Barnea, G.; Cannizzaro, L. A.; Li, F.; Nowell, P. C.; Boghosian-Sell, L.; Glick, J.; Weston, A.; Harris, C. C.; Drabkin, H.; Patterson, D.; Croce, C. M.; Schlessinger, J.; Huebner, K.: Receptor protein-tyrosine phosphatase gamma is a candidate tumor suppressor gene at human chromosome region 3p21. *Proc. Nat. Acad. Sci.* 88: 5036–5040, 1991.

- [77716] 16046.Latif, F.; Tory, K.; Modi, W.; Geil, L.; LaForgia, S.; Huebner,K.; Zbar, B.; Lerman, M. I.: A MspI polymorphism and linkage mapping of the human protein-tyrosine phosphatase G (PTPRG) gene. Hum. Molec.Genet. 2: 91, 1993.
- [77717] 16047.Ariyama, T.; Hasegawa, K.; Inazawa, J.; Mizuno, K.; Ogimoto, M.;Katagiri, T.; Yakura, H.: Assignment of the human protein tyrosinephosphatase, receptor-type, zeta (PTPRZ) gene to chromosome band 7q31.3. Cytogenet.Cell Genet. 70: 52-54, 1995.
- [77718] 16048.Schols, L.; Szymanski, S.; Peters, S.; Przuntek, H.; Epplen, J.T.; Hardt, C.; Riess, O.: Genetic background of apparently idiopathicsporadic cerebellar ataxia. Hum. Genet. 107: 132-137, 2000.
- [77719] 16049.Chan, J. Y.; Cheung, M.-C.; Moi, P.; Chan, K.; Kan, Y. W.: Chromosomallocalization of the human NF-E2 family of bZIP transcription factorsby fluorescence in situ hybridization. Hum. Genet. 95: 265-269,1995.
- [77720] 16050.Chan, J. Y.; Han, X.-L.; Kan, Y. W.: Cloning of Nrf1, an NF-E2-relatedtranscription factor, by genetic selection in yeast. Proc. Nat. Acad.Sci. 90: 11371-11375, 1993.
- [77721] 16051.Chan, J. Y.; Kwong, M.; Lu, R.; Chang, J.; Wang, B.; Yen, T. S.B.; Kan, Y. W.: Targeted disruption of the ubiquitous CNC-bZIP transcriptionfactor, Nrf-1, results in ane-

mia and embryonic lethality in mice. EMBOJ. 17:  
1779–1787, 1998.

[77722] 16052.Luna, L.; Johnsen, O.; Skartlien, A.; Pedeutour, F.;  
Turc-Carel,C.; Prydz, H.; Kolsto, A.–B.: Molecular cloning  
of a putative novelhuman bZIP transcription factor on  
chromosome 17q22. Genomics 22:553–562, 1994.

[77723] 16053.Luna, L.; Johnsen, O.; Skartlien, A. H.; Pedeutour,  
F.; Turc-Carel,C.; Prydz, H.; Kolsto, A.–B.: Molecular  
cloning of a putative novelhuman bZIP transcription factor  
on chromosome 17q22. Genomics 22:553–562, 1994.

[77724] 16054.McKie, J.; Johnstone, K.; Mattei, M.–G.; Scambler,  
P.: Cloningand mapping of murine Nfe2l1. Genomics 25:  
716–719, 1995.

[77725] 16055.McKie, J.; Scambler, P. J.: The Nfe2l1 gene maps to  
distal mousechromosome 11. Mammalian Genome 7:  
89–90, 1996.

[77726] 16056.Kas, K.; Wlodarska, I.; Meyen, E.; Van den Berghe,  
H.; Van de Ven,W. J. M.: Assignment of the gene encoding  
human Kruppel–related zincfinger protein 4 (GLI4) to  
8q24.3 by fluorescent in situ hybridization. Cytogenet.Cell  
Genet. 72: 297–298, 1996.

[77727] 16057.Goldblum, S. E.; Ding, X.; Funk, S. E.; Sage, E. H.:  
SPARC (secretedprotein acidic and rich in cysteine) regu–

lates endothelial cell shape and barrier function. *Proc. Nat. Acad. Sci.* 91: 3448–3452, 1994.

- [77728] 16058. Lossie, A. C.; Gordon, D. F.; Camper, S. A.: Localization of thyrotropin-releasing hormone receptor and thyrotroph embryonic factor on mouse chromosome 15. *Mammalian Genome* 4: 621–623, 1993.
- [77729] 16059. Matre, V.; Karlsen, H. E.; Wright, M. S.; Lundell, I.; Fjeldheim, A. K.; Gabrielsen, O. S.; Larhammar, D.; Gautvik, K. M.: Molecular cloning of a functional human thyrotropin-releasing hormone receptor. *Biochem. Biophys. Res. Commun.* 195: 179–185, 1993.
- [77730] 16060. Morrison, N.; Duthie, S. M.; Boyd, E.; Eidne, K. A.; Connor, J. M.: Assignment of the gene encoding the human thyrotropin-releasing hormone receptor to 8q23 by fluorescence in situ hybridization. *Hum. Genet.* 93: 716–718, 1994.
- [77731] 16061. Straub, R. E.; Frech, G. C.; Joho, R. H.; Gershengorn, M. C.: Expression cloning of a cDNA encoding the mouse pituitary thyrotropin-releasing hormone receptor. *Proc. Nat. Acad. Sci.* 87: 9514–9518, 1990.
- [77732] 16062. Yamada, M.; Monden, T.; Konaka, S.; Mori, M.: Assignment of human thyrotropin-releasing hormone (TRH) receptor gene to chromosome 8. *Somat. Cell Molec. Genet.*

19: 577–580, 1993.

[77733] 16063.Zhao, D.; Yang, J.; Jones, K. E.; Gerald, C.; Suzuki, Y.; Hogan, P. G.; Chin, W. W.; Tashjian, A. H., Jr.: Molecular cloning of a complementary deoxyribonucleic acid encoding the thyrotropin-releasing hormone receptor and regulation of its messenger ribonucleic acid in rat GH cells. *Endocrinology* 130: 3529–3536, 1992.

[77734] 16064.Bongarzone, I.; Butti, M. G.; Coronelli, S.; Borrello, M. G.; Santoro, M.; Mondellini, P.; Pilotti, S.; Fusco, A.; Della Porta, G.; Pierotti, M. A.: Frequent activation of ret protooncogene by fusion with a new activating gene in papillary thyroid carcinomas. *Cancer Res.* 54:2979–2985, 1994.

[77735] 16065.Bongarzone, I.; Butti, M. G.; Fugazzola, L.; Pacini, F.; Pinchera, A.; Vorontsova, T. V.; Demidchik, E. P.; Pierotti, M. A.: Comparison of the breakpoint regions of ELE1 and RET genes involved in the generation of RET/PTC3 oncogene in sporadic and in radiation-associated papillary thyroid carcinomas. *Genomics* 42: 252–259, 1997.

[77736] 16066.Klein, M.; Vignaud, J.-M.; Hennequin, V.; Toussaint, B.; Bresler, L.; Plenat, F.; Leclere, J.; Duprez, A.; Weryha, G.: Increased expression of the vascular endothelial growth

factor is a pejorative prognosismarker in papillary thyroid carcinoma. *J. Clin. Endocr. Metab.* 86:656–658, 2001.

[77737] 16067.Katyal, S. L.; Singh, G.; Locker, J.: Characterization of a secondhuman pulmonary surfactant-associated protein SP-A gene. *Am. J. Resp.Cell Molec. Biol.* 6: 446–452, 1992.

[77738] 16068.Ramet, M.; Lofgren, J.; Albo, O.–P.; Hallman, M.: Surfactant protein–A gene locus associated with recurrent otitis media. *J. Pediat.* 138:266–268, 2001.

[77739] 16069.Robinson, P. R.; Cohen, G. B.; Zhukovsky, E. A.; Oprian, D. D.: Constitutively active mutants of rhodopsin. *Neuron* 9: 719–725,1992.

[77740] 16070.Mason, I. J.; Murphy, D.; Munke, M.; Francke, U.; Elliott, R. W.;Hogan, B. L. M.: Developmental and transformation-sensitive expressionof the SPARC gene on mouse chromosome 11. *EMBO J.* 5: 1831–1837,1986.

[77741] 16071.Mason, I. J.; Taylor, A.; Williams, J. G.; Sage, H.; Hogan, B.L. M.: Evidence from molecular cloning that SPARC, a major productof mouse embryo parietal endoderm, is related to an endothelial cell'culture shock' glycoprotein of Mr 43,000. *EMBO J.* 5: 1465–1472,1986.

[77742] 16072.Naylor, S. L.; Helen–Davis, D.; Villarreal, X. C.; Long, G. L.: The human osteonectin gene on chromosome

5 is polymorphic. (Abstract) Cytogenet.Cell Genet. 51: 1051 only, 1989.

[77743] 16073.Schwartz, R. C.; Young, M. F.; Tsipouras, P.: Two RFLPs in the 5-prime end of the human osteonectin (ON) gene. Nucleic Acids Res. 16:9076 only, 1988.

[77744] 16074.Stenner, D. D.; Tracy, R. P.; Riggs, B. L.; Mann, K. G.: Human platelets contain and secrete osteonectin, a major protein of mineralized bone. Proc. Nat. Acad. Sci. 83: 6892-6896, 1986.

[77745] 16075.Swaroop, A.; Francke, U.: Molecular cloning, cDNA sequence, and expression of human SPARC (osteonectin). (Abstract) Am. J. Hum. Genet. 41:A240 only, 1987.

[77746] 16076.Swaroop, A.; Hogan, B. L. M.; Francke, U.: Molecular analysis of the cDNA for human SPARC/osteonectin/BM-40: sequence, expression, and localization of the gene to chromosome 5q31-q33. Genomics 2:37-47, 1988.

[77747] 16077.Termin, J. D.; Kleinman, H. K.; Whitson, S. W.; Conn, K. M.; McGarvey, M. L.; Martin, G. R.: Osteonectin, a bone-specific protein linking mineral to collagen. Cell 26: 99-105, 1981.

[77748] 16078.Blau, N.; Thony, B.; Renneberg, A.; Arnold, L. A.; Hyland, K.: Dihydropteridine reductase deficiency localized

to the central nervous system. J. Inherit. Metab. Dis. 21: 433–434, 1998.

[77749] 16079. Blau, N.; Thony, B.; Renneberg, A.; Penzien, J. M.; Hyland, K.; Hoffmann, G.: Variant of dihydropteridine reductase deficiency without hyperphenylalaninemia: effect of oral phenylalanine loading. J. Inherit. Metab. Dis. 22: 216–220, 1999.

[77750] 16080. Bonafe, L.; Thony, B.; Penzien, J. M.; Czarnecki, B.; Blau, N.: Mutations in the sepiapterin reductase gene cause a novel tetrahydrobiopterin-dependent-monoamine-neurotransmitter deficiency without hyperphenylalaninemia. Am. J. Hum. Genet. 69: 269–277, 2001.

[77751] 16081. Ichinose, H.; Katoh, S.; Sueoka, T.; Titani, K.; Fujita, K.; Nagatsu, T.: Cloning and sequencing of cDNA encoding human sepiapterin reductase: an enzyme involved in tetrahydrobiopterin biosynthesis. Biochem. Biophys. Res. Commun. 179: 183–189, 1991.

[77752] 16082. Murdoch, J. N.; Eddleston, J.; Stanier, P.; Copp, A. J.: Localization of the mouse gene encoding tyrosine kinase receptor type 10 on distal chromosome 1. Mammalian Genome 8: 941–952, 1997.

[77753] 16083. Thony, B.; Heizmann, C. W.; Mattei, M.-G.: Human GTP-cyclohydrolase I gene and sepiapterin reductase gene



map to region 14q21–q22 and 2p14–p12, respectively, by in situ hybridization. *Genomics* 26: 168–170, 1995.

- [77754] 16084. Jin, H.; Oksenberg, D.; Ashkenazi, A.; Peroutka, S. J.; Duncan, A. M. V.; Rozmahel, R.; Yang, Y.; Mengod, G.; Palacios, J. M.; O'Dowd, B. F.: Characterization of the human 5-hydroxytryptamine(1B) receptor. *J. Biol. Chem.* 267: 5735–5738, 1992.
- [77755] 16085. Levy, F. O.; Gudermann, T.; Birnbaumer, M.; Kaumann, A. J.; Birnbaumer, L.: Molecular cloning of a human gene (S31) encoding a novel serotonin receptor mediating inhibition of adenylyl cyclase. *FEBS Lett.* 296: 201–206, 1992.
- [77756] 16086. Levy, F. O.; Holtgreve-Grez, H.; Tasken, K.; Solberg, R.; Ried, T.; Gudermann, T.: Assignment of the gene encoding the 5-HT-1E serotonin receptor (S31) (locus HTR1E) to human chromosome 6q14–q15. *Genomics* 22: 637–640, 1994.
- [77757] 16087. McAllister, G.; Charlesworth, A.; Snodin, C.; Beer, M. S.; Noble, A. J.; Middlemiss, D. N.; Iversen, L. L.; Whiting, P.: Molecular cloning of a serotonin receptor from human brain (5HT1E): a fifth 5HT1-like subtype. *Proc. Nat. Acad. Sci.* 89: 5517–5521, 1992.
- [77758] 16088. Weinshank, R. L.; Zgombick, J. M.; Macchi, M. J.;

Branchek, T.A.; Hartig, P. R.: Human serotonin 1D receptor is encoded by a subfamily of two distinct genes: 5-HT(1D-alpha) and 5-HT(1D-beta). Proc. Nat. Acad. Sci. 89: 3630-3634, 1992.

[77759] 16089. Evans, T.; Reitman, M.; Felsenfeld, G.: An erythrocyte-specific DNA-binding factor recognizes a regulatory sequence common to all chicken globin genes. Proc. Nat. Acad. Sci. 85: 5976-5980, 1988.

[77760] 16090. Chen, Z.-Y.; Battinelli, E. M.; Fielder, A.; Bunday, S.; Sims, K.; Breakefield, X. O.; Craig, I. W.: A mutation in the Norrie disease gene (NDP) associated with X-linked familial exudative vitreoretinopathy. Nature Genet. 5: 180-183, 1993.

[77761] 16091. Dudgeon, J.: Familial exudative vitreo-retinopathy. Trans. Ophthal. Soc. U.K. 99: 45-49, 1979.

[77762] 16092. Yang-Feng, T. L.; DeGennaro, L. J.; Francke, U.: Genes for synapsin I, a neuronal phosphoprotein, map to conserved regions of human and murine X chromosomes. Proc. Nat. Acad. Sci. 83: 8679-8683, 1986.

[77763] 16093. Sinke, R. J.; de Leeuw, B.; Janssen, H. A. P.; Olde Weghuis, D.; Suijkerbuijk, R. F.; Meloni, A. M.; Gilgenkrantz, S.; Berger, W.; Ropers, H. H.; Sandberg, A. A.; Geurts van Kessel, A.: Localization of X chromosome short

arm markers relative to synovial sarcoma– adrenal adenocarcinoma–associated translocation breakpoints. Hum. Genet. 92:305–308, 1993.

[77764] 16094.Brennan, T. J.; Seeley, W. W.; Kilgard, M.; Schreiner, C. E.; Tecott,L. H.: Sound–induced seizures in serotonin 5–HT–2C receptor mutantmice. Nature Genet. 16: 387–390, 1997.

[77765] 16095.Gurevich, I.; Tamir, H.; Arango, V.; Dwork, A. J.; Mann, J. J.;Schmauss, C.: Altered editing of serotonin 2C receptor pre–mRNA inthe prefrontal cortex of depressed suicide victims. Neuron 34: 349–356,2002.

[77766] 16096.Hall, C. S.: Genetic differences in fatal audiogenic seizures:between two inbred strains of house mice. J. Hered. 38: 3–6, 1947.

[77767] 16097.Lappalainen, J.; Zhang, L.; Dean, M.; Oz, M.; Ozaki, N.; Yu, D.;Virkkunen, M.; Weight, F.; Linnoila, M.; Gold–man, D.: Identification,expression, and pharmacology of a cys(23)–ser(23) substitution inthe human 5–HT(2C) receptor gene (HTR2C). Genomics 27: 274–279,1995.

[77768] 16098.Milatovich, A.; Hsieh, C.–L.; Bonaminio, G.; Tecott, L.; Julius,D.; Francke, U.: Serotonin receptor 1c gene assigned to X chromosomein human (band q24) and mouse (bands D–F4). Hum. Molec. Genet. 1:681–684, 1992.

- [77769] 16099.Tecott, L. H.; Sun, L. M.; Akana, S. F.; Strack, A. M.; Lowenstein,D. H.; Dallman, M. F.; Julius, D.: Eating disorder and epilepsy in mice lacking 5-HT<sub>2C</sub> serotonin receptors. *Nature* 374: 542–546, 1995.
- [77770] 16100.Blaschke, R. J.; Monaghan, A. P.; Schiller, S.; Schechinger, B.;Rao, E.; Padilla–Nash, H.; Ried, T.; Rappold, G. A.: SHOT, a SHOX–related homeobox gene, is implicated in craniofacial, brain, heart, and limb development. *Proc. Nat. Acad. Sci.* 95: 2406–2411, 1998.
- [77771] 16101.Clement–Jones, M.; Schiller, S.; Rao, E.; Blaschke, R. J.; Zuniga,A.; Zeller, R.; Robson, S. C.; Binder, G.; Glass, I.; Strachan, T.;Lindsay, S.; Rappold, G. A.: The short stature homeobox gene SHOX is involved in skeletal abnormalities in Turner syndrome. *Hum. Molec.Genet.* 9: 695–702, 2000.
- [77772] 16102.Affara, N. A.; Chalmers, I. J.; Ferguson–Smith, M. A.: Analysis of the SRY gene in 22 sex–reversed XY females identifies four new point mutations in the conserved DNA binding domain. *Hum. Molec.Genet.* 2: 785–789, 1993.
- [77773] 16103.Behlke, M. A.; Bogan, J. S.; Beer–Romero, P.; Page, D. C.: Evidence that the SRY protein is encoded by a single exon on the human Y chromosome. *Genomics* 17:736–739, 1993.

- [77774] 16104. Berkovitz, G. D.; Fechner, P. Y.; Zacur, H. W.; Rock, J. A.; Snyder, H. M., III; Migeon, C. J.; Perlman, E. J.: Clinical and pathologic spectrum of 46,XY gonadal dysgenesis: its relevance to the understanding of sex differentiation. *Medicine* 70: 375–383, 1991.
- [77775] 16105. Bowles, J.; Cooper, L.; Berkman, J.; Koopman, P.: Sry requires a CAG repeat domain for male sex determination in *Mus musculus*. *Nature Genet.* 22: 405–408, 1999.
- [77776] 16106. Braun, A.; Kammerer, S.; Cleve, H.; Lohrs, U.; Schwarz, H.-P.; Kuhnle, U.: True hermaphroditism in a 46,XY individual, caused by a postzygotic somatic point mutation in the male gonadal sex-determining locus (SRY): molecular genetics and histological findings in a sporadic case. *Am. J. Hum. Genet.* 52: 578–585, 1993.
- [77777] 16107. Brown, S.; Yu, C. C.; Lanzano, P.; Heller, D.; Thomas, L.; Warburton, D.; Kitajewski, J.; Stadtmauer, L.: A de novo mutation (Gln2stop) at the 5-prime end of the SRY gene leads to sex reversal with partial ovarian function. (Letter) *Am. J. Hum. Genet.* 62: 189–192, 1998.
- [77778] 16108. Canto, P.; de la Chesnaye, E.; Lopez, M.; Cervantes, A.; Chavez, B.; Vilchis, F.; Reyes, E.; Ulloa-Aguirre, A.; Kofman-Alfaro, S.; Mendez, J. P.: A mutation in the 5-prime non-high mobility group box region of the SRY gene in

patients with Turner syndrome and Ymosaicism. *J. Clin. Endocr. Metab.* 85: 1908–1911, 2000.

[77779] 16109.Cherfas, J.: Sex and the single gene. *Science* 252: 782, 1991.

[77780] 16110.Clepet, C.; Schafer, A. J.; Sinclair, A. H.; Palmer, M. S.; Lovell–Badge,R.; Goodfellow, P. N.: The human SRY transcript. *Hum. Molec. Genet.* 2:2007–2012, 1993.

[77781] 16111.Davis, R. M.: Localisation of male determining factors in man:a thorough review of structural anomalies of the Y chromosome. *J.Med. Genet.* 18: 161–195, 1981.

[77782] 16112.Dork, T.; Stuhrmann, M.; Miller, K.; Schmidtke, J.: Independentobservation of SRY mutation I90M in a patient with complete gonadaldysgenesis. (Letter) *Hum. Mutat.* 11: 90–91, 1998.

[77783] 16113.Eicher, E. M.; Washburn, L. L.: Genetic control of primary sexdetermination in mice. *Annu. Rev. Genet.* 20: 327–360, 1986.

[77784] 16114.Foster, J. W.; Brennan, F. E.; Hampikian, G. K.; Goodfellow, P.N.; Sinclair, A. H.; Lovell–Badge, R.; Selwood, L.; Renfree, M. B.;Cooper, D. W.; Graves, J. A. M.: Evolution of sex determination andthe Y chromosome: SRY–related sequences in marsupials. *Nature* 359:531–533, 1992.

- [77785] 16115.Fuqua, J. S.; McLaughlin, J.; Perlman, E. J.; Berkowitz, G. D.: Analysis of the SRY gene in gonadal tissue of subjects with 46,XYgonadal dysgenesis (Letter) J. Clin. Endocr. Metab. 82: 701–702,1997.
- [77786] 16116.Domenice, S.; Nishi, M. Y.; Billerbeck, A. E. C.; Latronico, A.C.; Medeiros, M. A.; Russell, A. J.; Vass, K.; Carvalho, F. M.; Frade,E. M. C.; Arnhold, I. J. P.; Mendonca, B. B.: A novel missense mutation(S18N) in the 5–prime non–HMG box region of the SRY gene in a patientwith partial gonadal dysgenesis and his normal male relatives. Hum.Genet. 102: 213–215, 1998.
- [77787] 16117.Giese, K.; Pagel, J.; Grosschedl, R.: Distinct DNA–binding propertiesof the high mobility group domain of murine and human SRY sex–determiningfactors. Proc. Nat. Acad. Sci. 91: 3368–3372, 1994.
- [77788] 16118.Goodfellow, P. N.; Lovell–Badge, R.: SRY and sex determinationin mammals. Annu. Rev. Genet. 27: 71–92, 1993.
- [77789] 16119.Gubbay, J.; Collignon, J.; Koopman, P.; Capel, B.; Economou, A.;Munsterberg, A.; Vivian, N.; Goodfellow, P. N.; Lovell–Badge, R.:A gene mapping to the sex–determining region of the mouse Y chromosomeis a member of a novel family of embryonically expressed

genes. *Nature* 346:245–250, 1990.

- [77790] 16120. Haqq, C. M.; King, C.-Y.; Ukiyama, E.; Falsafi, S.; Haqq, T. N.; Donahoe, P. K.; Weiss, M. A.: Molecular basis of mammalian sexual determination: activation of Mullerian inhibiting substance gene expression by SRY. *Science* 266: 1494–1500, 1994.
- [77791] 16121. Clark, J.; Rocques, P. J.; Crew, A. J.; Gill, S.; Shipley, J.; Chan, A. M.-L.; Gusterson, B. A.; Cooper, C. S.: Identification of novel genes, SYT and SSX, involved in the t(X;18)(p11.2;q11.2) translocation found in human synovial sarcoma. *Nature Genet.* 7: 502–508, 1994.
- [77792] 16122. Yorifuji, T.; Wilson, R. W.; Beaudet, A. L.: Retroviral mediated expression of CD18 in normal and deficient human bone marrow progenitor cells. *Hum. Molec. Genet.* 2: 1443–1448, 1993.
- [77793] 16123. Ramulu, P.; Kennedy, M.; Xiong, W.-H.; Williams, J.; Cowan, M.; Blesh, D.; Yau, K.-W.; Hurley, J. B.; Nathans, J.: Normal light response, photoreceptor integrity, and rhodopsin dephosphorylation in mice lacking both protein phosphatases with EF hands (PPEF-1 and PPEF-2). *Molec. Cell. Biol.* 21: 8605–8614, 2001.
- [77794] 16124. van de Vosse, E.; Franco, B.; van der Bent, P.; Montini, E.; Orth, U.; Hanauer, A.; Tijmes, N.; van Ommen, G.-J.



B.; Ballabio, A.; denDunnen, J. T.; Bergen, A. A. B.: Exclusion of PPEF as the gene causing X-linked juvenile retinoschisis. *Hum. Genet.* 101: 235–237, 1997.

[77795] 16125. Fisher, S. E.; Ciccodicola, A.; Tanaka, K.; Curci, A.; Desicato, S.; D'Urso, M.; Craig, I. W.: Sequence-based exon prediction around the synaptophysin locus reveals a gene-rich area containing novel genes in human proximal Xp. *Genomics* 45: 340–347, 1997.

[77796] 16126. Oliva, M. M.; Wu, T. C.; Yang, V. W.: Isolation and characterization of a differentiation-dependent gene in the human colonic cell line HT29–18. *Arch. Biochem. Biophys.* 302: 183–192, 1993.

[77797] 16127. Prakash, S. K.; Paylor, R.; Jenna, S.; Lamarche-Vane, N.; Armstrong, D. L.; Xu, B.; Mancini, M. A.; Zoghbi, H. Y.: Functional analysis of ARHGAP6, a novel GTPase-activating protein for RhoA. *Hum. Molec. Genet.* 9: 477–488, 2000.

[77798] 16128. Schaefer, L.; Prakash, S.; Zoghbi, H. Y.: Cloning and characterization of a novel rho-type GTPase-activating protein gene (ARHGAP6) from the critical region for microphthalmia with linear skin defects. *Genomics* 46: 268–277, 1997.

[77799] 16129. Aman, M. J.; Tayebi, N.; Obiri, N. I.; Puri, R. K.;

Modi, W. S.;Leonard, W. J.: cDNA cloning and characterization of the human interleukin13 receptor alpha chain. J. Biol. Chem. 271: 29265–29270, 1996.

- [77800] 16130.Guo, J.; Apiou , F.; Mellerin, M.–P.; Lebeau, B.; Jacques, Y.;Minvielle, S.: Chromosome mapping and expression of the human interleukin–13receptor. Genomics 42: 141–5, 1997.
- [77801] 16131.Hilton, D. J.; Zhang, J.–G.; Metcalf, D.; Alexander, W. S.; Nicola,N. A.; Willson, T. A.: Cloning and characterization of a bindingsubunit of the interleukin 13 receptor that is also a component ofthe interleukin 4 receptor. Proc. Nat. Acad. Sci. 93: 497–501, 1996.
- [77802] 16132.Kaye, F. J.; Modi, S.; Ivanovska, I.; Koonin, E. V.; Thress, K.;Kubo, A.; Kornbluth, S.; Rose, M. D.: A family of ubiquitin–likeproteins binds the ATPase domain of Hsp70–like Stch. FEBS Lett. 467:348–352, 2000.
- [77803] 16133.Kaye, F. J.; Shows, T. B.: Assignment of ubiquilin 2 (UBQLN2)to human chromosome xp11.23–p11.1 by Gene–Bridge radiation hybrids. Cytogenet.Cell Genet. 89: 116–117, 2000.
- [77804] 16134.Kleijnen, M. F.; Shih, A. H.; Zhou, P.; Kumar, S.; Soccio, R. E.;Kedersha, N. L.; Gill, G.; Howley, P. M.: The hPLIC proteins mayprovide a link between the ubiquitina–

tion machinery and the proteasome. *Molec.Cell* 6:  
409–419, 2000.

[77805] 16135.Marchand, J.-B.; Kaiser, D. A.; Pollard, T. D.; Higgs, H. N.:Interaction of WASP/Scar proteins with actin and vertebrate Arp2/3complex. *Nature Cell Biol.* 3: 76–82, 2001.

[77806] 16136.Lee, D. K.; Nguyen, T.; Lynch, K. R.; Cheng, R.; Vanti, W. B.;Arkhitko, O.; Lewis, T.; Evans, J. F.; George, S. R.; O'Dowd, B. F.: Discovery and mapping of ten novel G protein–coupled receptor genes. *Gene* 275:83–91, 2001.

[77807] 16137.Furneaux, H. M.; Rosenblum, M. K.; Dalmau, J.; Wong, E.; Woodruff,P.; Graus, F.; Posner, J. B.: Selective expression of Purkinje–cellantigens in tumor tissue from patients with paraneoplastic cerebellardegeneration. *New Eng. J. Med.* 322: 1844–1851, 1990.

[77808] 16138.Furneaux, H. M.; Wong, E.; Posner, J. B.: Isolation of cDNA clonesencoding the major Yo paraneoplastic anti–gen. (Abstract) *Neurology* 40(suppl. 1): 166 only, 1990.

[77809] 16139.Knight, J. C.; Renwick, P. J.; Downing, J. R.; Okuda, T.: Physicallinkage of the cdc2–related gene (PCTK1) and the ubiquitin–activatingenzyme E1 gene (UBE1) on human Xp11.3. *Cytogenet. Cell Genet.* 71:155–157, 1995.

[77810] 16140.Aarskog, D.: A familial syndrome of short stature

associated with facial dysplasia and genital anomalies. J. Pediat. 77: 856–861, 1970.

[77811] 16141. Baldellou, A.; Galve, L.; Bassecourt, M.: Risk of medullary damage in Aarskog–Scott syndrome. (Abstract) Clin. Genet. 23: 225 only, 1983.

[77812] 16142. Bawle, E.; Tyrkus, M.; Lipman, S.; Bozimowski, D.: Aarskog syndrome: full male and female expression associated with an X–autosome translocation. Am. J. Med. Genet. 17: 595–602, 1984.

[77813] 16143. Berman, P. A.; Desjardins, C.; Fraser, F. C.: Inheritance of the Aarskog syndrome. Birth Defects Orig. Art. Ser. X(7): 151–159, 1974.

[77814] 16144. Buechner, M.; Hall, D. H.; Bhatt, H.; Hedgecock, E. M.: Cystic canal mutants in *Caenorhabditis elegans* are defective in the apical membrane domain of the renal (excretory) cell. Dev. Biol. 214: 227–241, 1999.

[77815] 16145. Casteels, M.; Samain, H.; Penninckx, F.; Coremans, G.; Beirinckx, J.; Fryns, J. P.: Megadolichosigmoid in a young male with Aarskog syndrome. Genet. Counsel. 5: 81–83, 1994.

[77816] 16146. Estrada, L.; Caron, E.; Gorski, J. L.: Fgd1, the Cdc42 guanine nucleotide exchange factor responsible for facio-genital dysplasia, is localized to the subcortical actin cy-

toskeleton and Golgi membrane. Hum.Molec. Genet. 10: 485–495, 2001.

- [77817] 16147.Fernandez, I.; Tsukahara, M.; Mito, H.; Yoshii, H.; Uchida, M.; Matsuo, K.; Kajii, T.: Congenital heart defects in Aarskog syndrome. Am.J. Med. Genet. 50: 318–322, 1994.
- [77818] 16148.Fryns, J.–P.: Dolichomegasigmoid in Aarskog syndrome. (Letter) Am.J. Med. Genet. 45: 122 only, 1993.
- [77819] 16149.Fryns, J. P.: Aarskog syndrome: the changing phenotype with age. Am.J. Med. Genet. 43: 420–427, 1992.
- [77820] 16150.Fryns, J. P.; Macken, J.; Vinken, L.; Igodt–Ameye, L.; van denBerghe, H.: The Aarskog syndrome. Hum. Genet. 42: 129–135, 1978.
- [77821] 16151.Funderburk, S. J.; Crandall, B. F.: The Aarskog syndrome in threebrothers. Clin. Genet. 6: 119–124, 1974.
- [77822] 16152.Furukawa, C. T.; Hall, B. D.; Smith, D. W.: The Aarskog syndrome. J.Pediat. 81: 1117–1122, 1972.
- [77823] 16153.Gao, J.; Estrada, L.; Cho, S.; Ellis, R. E.; Gorski, J. L.: TheCaenorhabditis elegans homolog of FGD1, the human Cdc42 GEF gene responsiblefor faciogenital dysplasia, is critical for excretory cell morphogenesis. Hum.Molec. Genet. 10: 3049–3062, 2001.
- [77824] 16154.Glover, T. W.; Verga, V.; Rafael, J.; Barcroft, C.;

Gorski, J.L.; Bawle, E. V.; Higgins, J. V.: Translocation breakpoint in Aarskogsyndrome maps to Xp11.21 between ALAS2 and DXS323. Hum. Molec. Genet. 2:1717–1718, 1993.

[77825] 16155.Grier, R. E.; Farrington, F. H.; Kendig, R.; Mamunes, P.: Autosomaldominant inheritance of the Aarskog phenotype. (Abstract) Am. J.Hum. Genet. 33: 64A only, 1981.

[77826] 16156.Hoo, J. J.: The Aarskog (facio-digito-genital) syndrome. Clin.Genet. 16: 269–276, 1979.

[77827] 16157.Kodama, M.; Fujimoto, S.; Namikawa, T.; Matsuda, I.: Aarskogsyndrome with isolated growth hormone deficiency. Europ. J. Pediat. 135:273–276, 1981.

[77828] 16158.Lebel, R. R.; May, M.; Pouls, S.; Lubs, H. A.; Stevenson, R. E.;Schwartz, C. E.: Non-syndromic X-linked mental retardation associatedwith a missense mutation (P312L) in the FGD1 gene. Clin. Genet. 61:139–145, 2002.

[77829] 16159.Logie, L. J.; Porteous, M. E. M.: Intelligence and developmentin Aarskog syndrome. Arch. Dis. Child. 79: 359–360, 1998.

[77830] 16160.Mikelsaar, R. V.–A.; Lurie, I. W.: Atypical case of Aarskog syndrome. J.Med. Genet. 29: 349–350, 1992.

[77831] 16161.Nielsen, K. B.: Aarskog syndrome in a Danish family: an illustrationof the need for dysmorphology in paedi-

atrics. Clin. Genet. 33: 315–317,1988.

[77832] 16162.Oberiter, V.; Lovrencic, M. K.; Schmutzer, L.; Kraus, O.: TheAarskog syndrome. Acta Paediat. Scand. 69: 567–570, 1980.

[77833] 16163.Orrico, A.; Galli, L.; Falciani, M.; Bracci, M.; Cavaliere, M.L.; Rinaldi, M. M.; Musacchio, A.; Sorrentino, V.: A mutation inthe pleckstrin homology (PH) domain of the FGD1 gene in an Italianfamily with faciogenital dysplasia (Aarskog–Scott syndrome). FEBSLett. 478: 216–220, 2000.

[77834] 16164.Pasteris, N. G.; Cadle, A.; Logie, L. J.; Porteous, M. E. M.;Schwartz, C. E.; Stevenson, R. E.; Glover, T. W.; Wilroy, R. S.; Gorski,J. L.: Isolation and analysis of the faciogenital dysplasia (Aarskog–Scottsyndrome) gene: a putative, rho/rac guanine nucleotide exchange factor. Cell 79:669–678, 1994.

[77835] 16165.Pasteris, N. G.; de Gouyon, B.; Cadle A. B.; Campbell, K.; Herman,G. E.; Gorski, J. L.: Cloning and regional localization of the mousefaciogenital dysplasia (Fgd1) gene. Mammalian Genome 6: 658–661,1995.

[77836] 16166.Pedersen, J. C.; Fryns, J. P.; Bracke, P.; Geeraert, M.; Van DenBerghe, H.: The Aarskog syndrome. Ann. Genet. 23: 108–110, 1980.

[77837] 16167.Porteous, M. E. M.; Goudie, D. R.: Aarskog syn–

drome. J. Med.Genet. 28: 44–47, 1991.

[77838] 16168.Rafael, J.; Verga, V.; Hall, B.; Burrigh, E.; Gorski, J.; Bawle,J.; Higgins, J. V.; Glover, T. W.: Assignment of the translocationbreakpoint in a patient with Aarskog syndrome to Xp11.21. (Abstract) Am.J. Hum. Genet. 51 (suppl.): A116 only, 1992.

[77839] 16169.Schwartz, C. E.; Gillessen–Kaesbach, G.; May, M.; Cappa, M.; Gorski,J.; Steindl, K.; Neri, G.: Two novel mutations confirm FGD1 is responsiblefor the Aarskog syndrome. Europ. J. Hum. Genet. 8: 869–874, 2000.

[77840] 16170.Scott, C. I., Jr.: Unusual facies, joint hypermobility, genitalanomaly and short stature: a new dysmorphic syndrome. Birth DefectsOrig. Art. Ser. VII(6): 240–246, 1971.

[77841] 16171.Demarest, S. J.; Martinez–Yamout, M.; Chung, J.; Chen, H.; Xu,W.; Dyson, H. J.; Evans, R. M.; Wright, P. E.: Mutual synergisticfolding in recruitment of CBP/p300 by p160 nuclear receptor coactivators. Nature 415:549–553, 2002.

[77842] 16172.Eckner, R.; Ewen, M. E.; Newsome, D.; Gerdes, M.; DeCaprio, J.A.; Lawrence, J. B.; Livingston, D. M.: Molecular cloning and functionalanalysis of the adenovirus E1A-associated 300–kD protein (p300) reveals a protein with properties of a transcriptional adaptor. Genes Dev.



15:869–884, 1994.

- [77843] 16173.Lin, C. H.; Hare, B. J.; Wagner, G.; Harrison, S. C.; Maniatis,T.; Fraenkel, E.: A small domain of CBP/p300 binds diverse proteins:solution structure and functional studies. *Molec. Cell* 8: 581–590,2001.
- [77844] 16174.Cohen–Haguenauer, O.; Picard, J. Y.; Mattei, M.–G.; Serero, S.;Van Cong, N.; de Tand, M.–F.; Guerrier, D.; Hors–Cayla, M.–C.; Josso,N.; Frezal, J.: Mapping of the gene for anti–mullerian hormone tothe short arm of human chromosome 19. *Cytogenet. Cell Genet.* 44:2–6, 1987.
- [77845] 16175.Forest, M. G.: Serum Mullerian inhibiting substance assay -- anew diagnostic test for disorders of gonadal development. (Editorial) *NewEng. J. Med.* 336: 1519–1521, 1997.
- [77846] 16176.Cate, R. L.; Mattaliano, R. J.; Hession, C.; Tizard, R.; Farber,N. M.; Cheung, A.; Ninfa, E. G.; Frey, A. Z.; Gash, D. J.; Chow, E.P.; Fisher, R. A.; Bertoni, J. M.; Torres, G.; Wallner, B. P.; Ramachandran,K. L.; Ragin, R. C.; Mangano, T. F.; MacLaughlin, D. T.; Donahoe,P. K.: Isolation of the bovine and human genes for mullerian inhibiting substance and expression of the human gene in animal cells. *Cell* 45:685–698, 1986.

- [77847] 16177.Guerrier, D.; Tran, D.; Vanderwinden, J. M.; Hideux, S.; Van Outryve,L.; Legeai, L.; Bouchard, M.; Van Vliet, G.; De Laet, M. H.; Picard,J. Y.; Kahn, A.; Josso, N.: The persistent mullerian duct syndrome:a molecular approach. J. Clin. Endocr. Metab. 68: 46–52, 1989.
- [77848] 16178.Harbison, M. D.; Magid, M. L. S.; Josso, N.; Mininberg, D. T.;New, M. I.: Anti–Mullerian hormone in three intersex conditions. Ann.Genet. 34: 226–232, 1991.
- [77849] 16179.King, T. R.; Lee, B. K.; Behringer, R. R.; Eicher, E. M.: Mappinganti–Mullerian hormone (Amh) and related sequences in the mouse: identificationof a new region of homology between MMU10 and HSA19p. Genomics 11:273–283, 1991.
- [77850] 16180.Knebelmann, B.; Boussin, L.; Guerrier, D.; Legeai, L.; Kahn, A.;Josso, N.; Picard, J.–Y.: Anti–Mullerian hormone Bruxelles: a nonsensemutation associated with the persistent Mullerian duct syndrome. Proc.Nat. Acad. Sci. 88: 3767–3771, 1991.
- [77851] 16181.Lee, M. M.; Donahoe, P. K.; Silverman, B. L.; Hasegawa, T.; Hasegawa,Y.; Gustafson, M. L.; Chang, Y.; MacLaughlin, D. T.: Measurementsof serum Mullerian inhibiting substance in the evaluation of childrenwith non-palpable gonads. New Eng. J. Med. 336: 1480–1486,

1997.

- [77852] 16182.Rey, R. A.; Belville, C.; Nihoul-Fekete, C.; Michel-Calemard,L.; Forest, M. G.; Lahlou, N.; Jaubert, F.; Mowszowicz, I.; David,M.; Saka, N.; Bouvattier, C.; Bertrand, A.-M.; and 16 others: Evaluationof gonadal function in 107 intersex patients by means of serum antimullerian-hormone measurement. *J. Clin. Endocr. Metab.* 84: 627-631, 1999.
- [77853] 16183.Rogers, D. S.; Gallagher, D. S.; Womack, J. E.: Somatic cellmapping of the genes for anti-mullerian hormone and osteonectin incattle: identification of a new bovine syntenic group. *Genomics* 9:298-300, 1991.
- [77854] 16184.Riesewijk, A. M.; Blagitko, N.; Schinzel, A. A.; Hu, L.; Schulz,U.; Hamel, B. C. J.; Ropers, H.-H.; Kalscheuer, V. M.: Evidence againsta major role of PEG1/MEST Silver-Russell syndrome. *Europ. J. Hum.Genet.* 6: 114-120, 1998.
- [77855] 16185.Riesewijk, A. M.; Hu, L.; Schulz, U.; Tariverdian, G.; Hoglund,P.; Kere, J.; Ropers, H.-H.; Kalscheuer, V. M.: Monoallelic expressionof human PEG1/MEST is paralleled by parent-specific methylation infetuses. *Genomics* 42: 236-244, 1997.
- [77856] 16186.Sado, T.; Nakajima, N.; Tada, M.; Takagi, N.: A

novel mesoderm-specific cDNA isolated from a mouse embryonal carcinoma cell line. *Dev. Growth Differ.* 35: 551–560, 1993.

- [77857] 16187. Jayakumar, A.; Chirala, S. S.; Chinault, A. C.; Baldini, A.; Abu-Elheiga, L.; Wakil, S. J.: Isolation and chromosomal mapping of genomic clones encoding the human fatty acid synthase gene. *Genomics* 23: 420–424, 1994.
- [77858] 16188. Jayakumar, A.; Tai, M.-H.; Huang, W.-Y.; Al-Feel, W.; Hsu, M.; Abu-Elheiga, L.; Chirala, S. S.; Wakil, S. J.: Human fatty acid synthase: properties and molecular cloning. *Proc. Nat. Acad. Sci.* 92: 8695–8699, 1995.
- [77859] 16189. Loftus, T. M.; Jaworsky, D. E.; Frehywot, G. L.; Townsend, C. A.; Ronnett, G. V.; Lane, M. D.; Kuhajda, F. P.: Reduced food intake and body weight in mice treated with fatty acid synthase inhibitors. *Science* 288: 2379–2381, 2000.
- [77860] 16190. Wakil, S. J.: Fatty acid synthase, a proficient multifunctional enzyme. *Biochemistry* 28: 4523–4530, 1989.
- [77861] 16191. Ye, Q.; Chung, L. W. K.; Li, S.; Zhau, H. E.: Identification of a novel FAS/ER- $\alpha$  fusion transcript expressed in human cancer cells. *Biochim. Biophys. Acta* 1493: 373–377, 2000.
- [77862] 16192. Hofmann, M. A.; Drury, S.; Fu, C.; Qu, W.; Taguchi,

A.; Lu, Y.; Avila, C.; Kambham, N.; Bierhaus, A.; Nawroth, P.; Neurath, M. F.; Slattery, T.; Beach, D.; McClary, J.; Nagashima, M.; Morser, J.; Stern, D.; Schmidt, A. M.: RAGE mediates a novel proinflammatory axis: a central cell surface receptor for S100/calgranulin polypeptides. *Cell* 97:889–901, 1999.

[77863] 16193. Yan, S. D.; Chen, X.; Fu, J.; Chen, M.; Zhu, H.; Roher, A.; Slattery, T.; Zhao, L.; Nagashima, M.; Morser, J.; Migheli, A.; Nawroth, P.; Stern, D.; Schmidt, A. M.: RAGE and amyloid-beta peptide neurotoxicity in Alzheimer's disease. *Nature* 382: 685–691, 1996.

[77864] 16194. Argeson, A. C.; Druck, T.; Veronese, M. L.; Knopf, J. L.; Buchberg, A. M.; Huebner, K.; Siracusa, L. D.: Phospholipase C gamma-2 (Plcg2) and phospholipase C gamma-1 (Plcg1) map to distinct regions in the human and mouse genomes. *Genomics* 25: 29–35, 1995.

[77865] 16195. Hernandez, D.; Egan, S. E.; Yulug, I. G.; Fisher, E. M. C.: Mapping the gene that encodes phosphatidylinositol-specific phospholipase C-gamma-2 in the human and the mouse. *Genomics* 23: 504–507, 1994.

[77866] 16196. Kang, J. S.; Kohlhuber, F.; Hug, H.; Marme, D.; Eick, D.; Ueffing, M.: Cloning and functional analysis of the hematopoietic cell-specific phospholipase C-gamma-2

promoter. FEBS Lett. 399: 14–20, 1996.

- [77867] 16197. Dumont, D. J.; Anderson, L.; Breitman, M. L.; Duncan, A. M. V.: Assignment of the endothelial-specific protein receptor tyrosine kinase gene (TEK) to human chromosome 9p21. Genomics 23: 512–513, 1994.
- [77868] 16198. Delot, E.; King, L. M.; Briggs, M. D.; Wilcox, W. R.; Cohn, D. H.: Trinucleotide expansion mutations in the cartilage oligomeric matrix protein (COMP) gene. Hum. Molec. Genet. 8: 123–128, 1999.
- [77869] 16199. Back, A. L.; Kerkerling, M.; Baker, D.; Bauer, T. R.; Embree, L. J.; Hickstein, D. D.: A point mutation associated with leukocyte adhesion deficiency type 1 of moderate severity. Biochem. Biophys. Res. Commun. 193: 912–918, 1993.
- [77870] 16200. Back, A. L.; Kwok, W. W.; Hickstein, D. D.: Identification of two molecular defects in a child with leukocyte adherence deficiency. J. Biol. Chem. 267: 5482–5487, 1992.
- [77871] 16201. Barclay, A. N.; Birkeland, M. L.; Brown, M. H.; Beyers, A. D.; Davis, S. J.; Somoza, C.; Williams, A. F.: The Leukocyte Antigen Facts Book. New York: Academic Press (pub.) 1993. Pp. 124–127 and 140–141.
- [77872] 16202. Beatty, P. G.; Ochs, H. D.; Harlan, J. M.; Price, T. H.;

Rosen,H.; Taylor, R. F.; Hansen, J. A.; Klebanoff, S. J.: Absence of monoclonal-antibody-defined protein complex in a boy with abnormal leucocyte function. *Lancet* I:535-537, 1984.

[77873] 16203.Bianchi, E.; Denti, S.; Granata, A.; Bossi, G.; Geginat, J.; Villa,A.; Rogge, L.; Pardi, R.: Integrin LFA-1 interacts with the transcriptional co-activator JAB1 to modulate AP-1 activity. *Nature* 404: 617-621,2000.

[77874] 16204.Bissenden, J. G.; Haeney, M. R.; Tarlow, M. J.; Thompson, R. A.: Delayed separation of the umbilical cord, severe widespread infections, and immunodeficiency. *Arch. Dis. Child.* 56: 397-399, 1981.

[77875] 16205.Boucheix, C.: Personal Communication. Villejuif, France 1/31/1987.

[77876] 16206.Bowen, T. J.; Ochs, H. D.; Altman, L. C.; Price, T. H.; Van Epps,D. E.; Brautigan, D. L.; Rosin, R. E.; Perkins, W. D.; Babior, B.M.; Klebanoff, S. J.; Wedgwood, R. J.: Severe recurrent bacterial infections associated with defective adherence and chemotaxis in two patients with neutrophils deficient in a cell-associated glycoprotein. *J.Pediat.* 101: 932-940, 1982.

[77877] 16207.Bullard, D. C.; Scharffetter-Kochanek, K.; McArthur, M. J.; Chosay,J. G.; McBride, M. E.; Montgomery, C. A.;

Beaudet, A. L.: A polygenic mouse model of psoriasiform skin disease in CD18-deficient mice. *Proc. Nat. Acad. Sci.* 93: 2116–2121, 1996.

[77878] 16208. Crowley, C. A.; Curnutte, J. T.; Rosin, R. E.; Andre-Schwartz, J.; Gallin, J. I.; Klempner, M.; Snyderman, R.; Southwick, F. S.; Stossel, T. P.; Babior, B. M.: An inherited abnormality of neutrophil adhesion: its genetic transmission and its association with a missing protein. *New Eng. J. Med.* 302: 1163–1168, 1980.

[77879] 16209. Dana, N.; Clayton, L. K.; Tennen, D. G.; Pierce, M. W.; Lachmann, P. J.; Law, S. A.; Arnaout, M. A.: Leukocytes from four patients with complete or partial Leu-CAM deficiency contain the common beta-subunit precursor and beta-subunit messenger RNA. *J. Clin. Invest.* 79: 1010–1015, 1987.

[77880] 16210. Dana, N.; Todd, R. F., III; Pitt, J.; Springer, T. A.; Arnaout, M. A.: Deficiency of a surface membrane glycoprotein (Mo1) in man. *J. Clin. Invest.* 73: 153–159, 1984.

[77881] 16211. Fischer, A.; Griscelli, C.; Friedrich, W.; Kubanek, B.; Levinsky, R.; Morgan, G.; Vossen, J.; Wagemaker, G.; Landais, P.: Bone-marrow transplantation for immunodeficiencies and osteopetrosis: European survey, 1968–1985. *Lancet* II: 1080–1084, 1986.



- [77882] 16212.Fujita, K.; Kobayashi, K.; Kajii, T.: Impaired neutrophil adhesion:a new patient in a previously reported family. *Acta Paediat. Jpn.* 27:527–534, 1985.
- [77883] 16213.Fujita, K.; Kobayashi, K.; Okino, F.: Juvenile rheumatoid arthritis in two siblings with congenital leukocyte adhesion deficiency. *Europ.J. Pediat.* 148: 118–119, 1988.
- [77884] 16214.Harvath, L.; Andersen, B. R.: Defective initiation of oxidativemetabolism in polymorphonuclear leukocytes. *New Eng. J. Med.* 300:1130–1135, 1979.
- [77885] 16215.Hayward, A. R.; Leonard, J.; Harvey, B. A. M.; Greenwood, M. C.;Wood, C. B. S.; Soothill, J. F.: Delayed separation of the umbilicalcord, widespread infections, and defective neutrophil mobility. *Lancet I*:1099–1101, 1979.
- [77886] 16216.Hibbs, M. L.; Wardlaw, A. J.; Stacker, S. A.; Anderson, D. C.;Lee, A.; Roberts, T. M.; Springer, T. A.: Transfection of cells frompatients with leukocyte adhesion deficiency with an integrin betasubunit (CD18) restores lymphocyte function–associated antigen–1 expressionand function. *J. Clin. Invest.* 85: 674–681, 1990.
- [77887] 16217.Hogg, N.; Stewart, M. P.; Scarth, S. L.; Newton, R.; Shaw, J.M.; Law, S. K. A.; Klein, N.: A novel leukocyte ad–

hesion deficiency caused by expressed but nonfunctional beta-2 integrins Mac-1 and LFA-1. *J. Clin. Invest.* 103: 97-106, 1999.

[77888] 16218. Hynes, R. O.: Integrins: versatility, modulation and signaling in cell adhesion. *Cell* 69: 11-25, 1992.

[77889] 16219. Kehrli, M. E., Jr.; Ackermann, M. R.; Shuster, D. E.; van der Maaten, M. J.; Schmalstieg, F. C.; Anderson, D. C.; Hughes, B. J.: Bovine leukocyte adhesion deficiency: beta(2) integrin deficiency in young Holstein cattle. *Am. J. Path.* 140: 1489-1492, 1992.

[77890] 16220. Kishimoto, T. K.; Hollander, N.; Roberts, T. M.; Anderson, D. C.; Springer, T. A.: Heterogeneous mutations in the beta subunit common to the LFA-1, Mac-1, and p150,95 glycoproteins cause leukocyte adhesion deficiency. *Cell* 50: 193-202, 1987.

[77891] 16221. Kishimoto, T. K.; O'Connor, K.; Lee, A.; Roberts, T. M.; Springer, T. A.: Cloning of the beta subunit of the leukocyte adhesion proteins: homology to an extracellular matrix receptor defines a novel supergene family. *Cell* 48: 681-690, 1987.

[77892] 16222. Kobayashi, K.; Fujita, K.; Okino, F.; Kajii, T.: An abnormality of neutrophil adhesion: autosomal recessive inheritance associated with missing neutrophil glycopro-

teins. *Pediatrics* 73: 606–610, 1984.

- [77893] 16223.Krauss, J. C.; Mayo–Bond, L. A.; Rogers, C. E.; Weber, K. L.; Todd, R. F., III; Wilson, J. M.: An in vivo animal model of genetherapy for leukocyte adhesion deficiency. *J. Clin. Invest.* 88:1412–1417, 1991.
- [77894] 16224.Matsuura, S.; Kishi, F.; Tsukahara, M.; Nunoi, H.; Matsuda, I.; Kobayashi, K.; Kajii, T.: Leukocyte adhesion deficiency: identification of novel mutations in two Japanese patients with a severe form. *Biochem. Biophys. Res. Commun.* 184: 1460–1467, 1992.
- [77895] 16225.Nelson, C.; Rabb, H.; Arnaout, M. A.: Genetic cause of leukocyte adhesion molecule deficiency: abnormal splicing and a missense mutation in a conserved region of CD18 impair cell surface expression of beta-2 integrins. *J. Biol. Chem.* 267: 3351–3357, 1992.
- [77896] 16226.Tini, M.; Benecke, A.; Um, S.-J.; Torchia, J.; Evans, R. M.; Chambon, P.: Association of CBP/p300 acetylase and thymine DNA glycosylase links DNA repair and transcription. *Molec. Cell* 9: 265–277, 2002.
- [77897] 16227.Weaver, B. K.; Kumar, K. P.; Reich, N. C.: Interferon regulatory factor 3 and CREB-binding protein/p300 are subunits of double-stranded RNA-activated transcription factor DRAF1. *Molec. Cell. Biol.* 18:1359–1368, 1998.

- [77898] 16228.Hansen, J. J.; Durr, A.; Cournu-Rebeix, I.; Georgopoulos, C.; Ang, D.; Nielsen, M. N.; Davoine, C.-S.; Brice, A.; Fontaine, B.; Gregersen, N.; Bross, P.: Hereditary spastic paraplegia SPG13 is associated with a mutation in the gene encoding the mitochondrial chaperonin Hsp60. *Am. J. Hum. Genet.* 70: 1328–1332, 2002.
- [77899] 16229.Todd, M. J.; Viitanen, P. V.; Lorimer, G. H.: Dynamics of the chaperonin ATPase cycle: implications for facilitated protein folding. *Science* 265:659–666, 1994.
- [77900] 16230.Hellevuo, K.; Berry, R.; Sikela, J. M.; Tabakoff, B.: Localization of the gene for a novel human adenylyl cyclase (ADCY7) to chromosome 16. *Hum. Genet.* 95: 197–200, 1995.
- [77901] 16231.Hellevuo, K.; Yoshimura, M.; Kao, M.; Hoffman, P. L.; Cooper, D.M. F.; Tabakoff, B.: A novel adenylyl cyclase sequence cloned from the human erythroleukemia cell line. *Biochem. Biophys. Res. Commun.* 192:311–318, 1993.
- [77902] 16232.Iavarone, A.; Garg, P.; Lasorella, A.; Hsu, J.; Israel, M. A.: The helix-loop-helix protein Id-2 enhances cell proliferation and binds to the retinoblastoma protein. *Genes Dev.* 8: 1270–1284, 1994.
- [77903] 16233.Lasorella, A.; Iavarone, A.; Israel, M. A.: Id2 specifically alters regulation of the cell cycle by tumor suppressor

proteins. *Molec.Cell. Biol.* 16: 2570–2578, 1996.

- [77904] 16234.Lasorella, A.; Nosedà, M.; Beyna, M.; Iavarone, A.: Id2 is a retinoblastoma protein target and mediates signalling by Myc oncoproteins. *Nature* 407:592–598, 2000.
- [77905] 16235.Moerman, P.; Fryns, J.-P.; Sastrowijoto, S. H.; Vandenberghe, K.; Lauweryns, J. M.: Hereditary renal adysplasia: new observations and hypotheses. *Pediat. Path.* 14: 405–410, 1994.
- [77906] 16236.Stapleton, D.; Woollatt, E.; Mitchell, K. I.; Nicholl, J. K.; Fernandez, C. S.; Michell, B. J.; Witters, L. A.; Power, D. A.; Sutherland, G. R.; Kemp, B. E.: AMP-activated protein kinase isoenzyme family: subunit structure and chromosomal location. *FEBS Lett.* 409: 452–456, 1997.
- [77907] 16237.Budarf, M. L.; Perier, F.; Barnoski, B. L.; Bell, C. J.; Vandenberg, C. A.: Assignment of the human hippocampal inward rectifier potassium channel (HIR) gene to 22q13.1. *Genomics* 26: 625–629, 1995.
- [77908] 16238.Makhina, E. N.; Kelly, A. J.; Lopatin, A. N.; Mercer, R. W.; Nichols, C. G.: Cloning and expression of a novel human brain inward rectifier potassium channel. *J. Biol. Chem.* 269: 20468–20474, 1994.
- [77909] 16239.Perier, F.; Radeke, C. M.; Vandenberg, C. A.: Primary structure and characterization of a small-

conductance inwardly rectifying potassium channel from human hippocampus. Proc. Nat. Acad. Sci. 91: 6240–6244, 1994.

- [77910] 16240. Derry, J. M.; Barnard, P. J.: The gene for tissue inhibitor of metalloproteinases (TIMP) is located within an intron of the synapsin I gene on the X chromosome. (Abstract) Cytogenet. Cell Genet. 58:2061–2062, 1991.
- [77911] 16241. Gal, A.; Wieringa, B.; Smeets, D. F. C. M.; Bleeker-Wagemakers, L.; Ropers, H. H.: Submicroscopic interstitial deletion of the X chromosome explains a complex genetic syndrome dominated by Norrie disease. Cytogenet. Cell Genet. 42: 219–224, 1986.
- [77912] 16242. Simpson, E.: The H-Y antigen and sex reversal. Cell 44: 813–814, 1986.
- [77913] 16243. Di Paola, R.; Frittitta, L.; Miscio, G.; Bozzali, M.; Baratta, R.; Centra, M.; Spampinato, D.; Santagati, M. G.; Ercolino, T.; Cisternino, C.; Soccio, T.; Mastroianno, S.; Tassi, V.; Almgren, P.; Pizzuti, A.; Vigneri, R.; Trischitta, V.: A variation in 3-prime UTR of hPTP1B increases specific gene expression and associates with insulin resistance. Am. J. Hum. Genet. 70: 806–812, 2002.
- [77914] 16244. Eidne, K.; Taylor, P.; Connor, M.; Duthie, S.: Isolation, characterization and chromosomal localization of the

human thyrotropin releasing hormone receptor. (Abstract)  
75th Annual Meeting of the Endocr. Soc. 437only, 1993.

- [77915] 16245.Castedo, M.; Ferri, K. F.; Blanco, J.; Roumier, T.; Larochette, N.; Barretina, J.; Amendola, A.; Nardacci, R.; Metivier, D.; Este, J. A.; Piacentini, M.; Kroemer, G.: Human immunodeficiency virus 1 envelope glycoprotein complex-induced apoptosis involves mammalian target of rapamycin/FKBP12-rapamycin-associated protein-mediated p53 phosphorylation. *J. Exp. Med.* 194: 1097-1110, 2001.
- [77916] 16246.El-Deiry, W. S.; Tokino, T.; Velculescu, V. E.; Levy, D. B.; Parsons, R.; Trent, J. M.; Lin, D.; Mercer, E.; Kinzler, K. W.; Vogelstein, B.: WAF1, a potential mediator of p53 tumor suppression. *Cell* 75:817-825, 1993.
- [77917] 16247.Moore, K. J.; D'Amore-Bruno, M. A.; Korfhagen, T. R.; Glasser, S. W.; Whitsett, J. A.; Jenkins, N. A.; Copeland, N. G.: Chromosomal localization of three pulmonary surfactant protein genes in the mouse. *Genomics* 12:388-393, 1992.
- [77918] 16248.Nogee, L. M.; Dunbar, A. E., III; Wert, S. E.; Askin, F.; Hamvas, A.; Whitsett, J. A.: A mutation in the surfactant protein C gene associated with familial interstitial lung disease. *New Eng. J. Med.* 344:573-579, 2001.
- [77919] 16249.Warr, R. G.; Hawgood, S.; Buckley, D. I.; Crisp, T.

M.; Schilling, J.; Benson, B. J.; Ballard, P. L.; Clements, J. A.; White, R. T.: Low molecular weight human pulmonary surfactant protein (SP5): isolation, characterization, and cDNA and amino acid sequences. *Proc. Nat. Acad. Sci.* 84: 7915–7919, 1987.

[77920] 16250. Wood, S.; Yaremko, M. L.; Schertzer, M.; Kelemen, P. R.; Minna, J.; Westbrook, C. A.: Mapping of the pulmonary surfactant SP5 (SFTP2) locus to 8p21 and characterization of a microsatellite repeat marker that shows frequent loss of heterozygosity in human carcinomas. *Genomics* 24:597–600, 1994.

[77921] 16251. Kolble, K.; Lu, J.; Mole, S. E.; Kaluz, S.; Reid, K. B. M.: Assignment of the human pulmonary surfactant protein D gene (SFTP4) to 10q22–q23 close to the surfactant protein A gene cluster. *Genomics* 17: 294–298, 1993.

[77922] 16252. Botas, C.; Poulain, F.; Akiyama, J.; Brown, C.; Allen, L.; Goerke, J.; Clements, J.; Carlson, E.; Gillespie, A. M.; Epstein, C.; Hawgood, S.: Altered surfactant homeostasis and alveolar type II cell morphology in mice lacking surfactant protein D. *Proc. Nat. Acad. Sci.* 95:11869–11874, 1998.

[77923] 16253. Crouch, E.; Rust, K.; Veile, R.; Donis-Keller, H.; Grosso, L.: Genomic organization of human surfactant protein D (SP-D): SP-D is encoded on chromosome



10q22.2–23.1. J. Biol. Chem. 268: 2976–2983,1993.

- [77924] 16254.Holmskov, U.; Lawson, P.; Teisner, B.; Tornoe, I.; Willis, A. C.;Morgan, C.; Koch, C.; Reid, K. B. M.: Isolation and characterizationof a new member of the scavenger receptor superfamily, glycoprotein–340(gp–340), as a lung surfactant protein–D binding molecule. J. Biol.Chem. 272: 13743–13749, 1997.
- [77925] 16255.Holmskov, U.; Mollenhauer, J.; Madsen, J.; Vitved, L.; Gronlund,J.; Tornoe, I.; Kliem, A.; Reid, K. B. M.; Poustka, A.; Skjodt, K.: Cloning of gp–340, a putative opsonin receptor for lung surfactantprotein D. Proc. Nat. Acad. Sci. 96: 10794–10799, 1999.
- [77926] 16256.LeVine, A. M.; Whitsett, J. A.; Hartshorn, K. L.; Crouch, E. C.;Korfhagen, T. R.: Surfactant protein D enhances clearance of influenzaA virus from the lung in vivo. J. Immun. 167: 5868–5873, 2001.
- [77927] 16257.Lu, J.; Willis, A. C.; Reid, K. B. M.: Purification, characterizationand cDNA cloning of human lung surfactant protein D. Biochem. J. 284:795–802, 1992.
- [77928] 16258.Rust, K.; Grosso, L.; Zhang, V.; Chang, D.; Persson, A.; Longmore,W.; Cai, G.–Z.; Crouch, E.: Human surfactant protein D: SP–D containsa C–type lectin carbohydrate recognition domain. Arch. Biochem. Biophys.

290:116–126, 1991.

- [77929] 16259. Wert, S. E.; Yoshida, M.; LeVine, A. M.; Ikegami, M.; Jones, T.; Ross, G. F.; Fisher, J. H.; Korfhagen, T. R.; Whitsett, J. A.: Increased metalloproteinase activity, oxidant production, and emphysema in surfactant protein D gene-inactivated mice. *Proc. Nat. Acad. Sci.* 97: 5972–5977, 2000.
- [77930] 16260. Gedde-Dahl, T., Jr.; Olaisen, B.; Teisberg, P.; Wilhelm, M. C.; Mevag, B.; Helland, R.: The locus for apolipoprotein E (apoE) is close to the Lutheran (Lu) blood group locus on chromosome 19. *Hum. Genet.* 67: 178–182, 1984.
- [77931] 16261. Grubb, R.: Zur Genetik des Lewis-Systems. *Naturwissenschaften* 21:560–561, 1953.
- [77932] 16262. Kolanus, W.; Nagel, W.; Schiller, B.; Zeitlmann, L.; Godar, S.; Stockinger, H.; Seed, B.: Alpha-L-beta-2 integrin/LFA-1 binding to ICAM-1 induced by cytohesin-1, a cytoplasmic regulatory molecule. *Cell* 86:233–242, 1996.
- [77933] 16263. Ogasawara, M.; Kim, S.-C.; Adamik, R.; Togawa, A.; Ferrans, V.J.; Takeda, K.; Kirby, M.; Moss, J.; Vaughan, M.: Similarities in function and gene structure of cytohesin-4 and cytohesin-1, guanine nucleotide-exchange proteins for ADP-ribosylation factors. *J. Biol. Chem.* 275:

3221–3230, 2000.

- [77934] 16264. Findlay, D. M.; Fisher, L. W.; McQuillan, C. I.; Termine, J. D.; Young, M. F.: Isolation of the osteonectin gene: evidence that a variable region of the osteonectin molecule is encoded within one exon. *Biochemistry* 27: 1483–1489, 1988.
- [77935] 16265. Gilmour, D. T.; Lyon, G. J.; Carlton, M. B. L.; Sanes, J. R.; Cunningham, J. M.; Anderson, J. R.; Hogan, B. L. M.; Evans, M. J.; Colledge, W. H.: Mice deficient for the secreted glycoprotein SPARC/osteonectin/BM40 develop normally but show severe age-onset cataract formation and disruption of the lens. *EMBO J.* 17: 1860–1870, 1998.
- [77936] 16266. Sanyanusin, P.; McNoe, L. A.; Sullivan, M. J.; Weaver, R. G.; Eccles, M. R.: Mutation of PAX2 in two siblings with renal-coloboma syndrome. *Hum. Molec. Genet.* 4: 2183–2184, 1995.
- [77937] 16267. Sanyanusin, P.; Norrish, J. H.; Ward, T. A.; Nebel, A.; McNoe, L. A.; Eccles, M. R.: Genomic structure of the human PAX2 gene. *Genomics* 35: 258–261, 1996.
- [77938] 16268. Sanyanusin, P.; Schimmenti, L. A.; McNoe, L. A.; Ward, T. A.; Pierpont, M. E. M.; Sullivan, M. J.; Dobyns, W. B.; Eccles, M. R.: Mutation of the PAX2 gene in a family with optic nerve colobomas, renal anomalies and vesi-

coureteral reflux. *Nature Genet.* 9: 358–364,1995.

- [77939] 16269.Schimmenti, L. A.; Cunliffe, H. E.; McNoe, L. A.; Ward, T. A.;French, M. C.; Shim, H. H.; Zhang, Y.–H.; Proesmans, W.; Leys, A.;Byerly, K. A.; Braddock, S. R.; Masuno, M.; Imaizumi, K.; Devriendt,K.; Eccles, M. R.: Further delineation of renal–coloboma syndromein patients with extreme variability of phenotype and identical PAX2mutations. *Am. J. Hum. Genet.* 60: 869–878, 1997.
- [77940] 16270.Schimmenti, L. A.; Pierpont, M. E.; Carpenter, B. L. M.; Kashtan,C. E.; Johnson, M. R.; Dobyns, W. B.: Autosomal dominant optic nervecolobomas, vesicoureteral reflux, and renal anomalies. *Am. J. Med.Genet.* 59: 204–208, 1995.
- [77941] 16271.Schimmenti, L. A.; Shim, H. H.; Wirtschafter, J. D.; Panzarino,V. A.; Kashtan, C. E.; Kirkpatrick, S. J.; Wargowski, D. S.; France,T. D.; Michel, E.; Dobyns, W. B.: Homonucleotide expansion and contractionmutations of PAX2 and inclusion of Chiari 1 malformation as part ofrenal–coloboma syndrome. *Hum. Mutat.* 14: 369–376, 1999.
- [77942] 16272.Stapleton, P.; Weith, A.; Urbanek, P.; Kozmik, Z.; Busslinger,M.: Chromosomal localization of seven PAX genes and cloning of a novel family member, PAX–9. *Na–*

ture Genet. 3: 292–298, 1993.

[77943] 16273.Tellier, A.–L.; Amiel, J.; Delezoide, A.–L.; Audollent, S.; Auge,J.; Esnault, D.; Encha–Razavi, F.; Munnich, A.; Lyonnet, S.; Vekemans,M.; Attie–Bitach, T.: Expression of the PAX2 gene in human embryosand exclusion in the CHARGE syndrome. Am. J. Med. Genet. 93: 85–88,2000.

[77944] 16274.Tellier, A.–L.; Amiel, J.; Salomon, R.; Jolly, D.; Delezoide,A.–L.; Auge, J.; Gubler, M.–C.; Munnich, A.; Lyonnet, S.; Antignac,C.; Vekemans, M.; Broyer, M.; Attie–Bitach, T.: PAX2 expression duringearly human development and its mutations in renal hypoplasia withor without coloboma. (Abstract) Am. J. Hum. Genet. 63 (supl.):A7 only, 1998.

[77945] 16275.Ward, T. A.; Nebel, A.; Reeve, A. E.; Eccles, M. R.: Alternativemessenger RNA forms and open reading frames within an additional conservedregion of the human PAX–2 gene. Cell Growth Differ. 5: 1015–1021,1994.

[77946] 16276.Weaver, R. G.; Cashwell, L. F.; Lorentz, W.; White–man, D.; Geisinger,K. R.; Ball, M.: Optic nerve coloboma associated with renal disease. Am.J. Med. Genet. 29: 597–605, 1988.

[77947] 16277.Barr, F. G.; Nauta, L. E.; Davis, R. J.; Schafer, B. W.; Nycum,L. M.; Biegel, J. A.: In vivo amplification of the PAX3–FKHR andPAX7–FKHR fusion genes in alveolar rhab–

domyosarcoma. Hum. Molec.Genet. 5: 15–21, 1996.

[77948] 16278.Burri, M.; Tromvoukis, Y.; Bopp, D.; Frigerio, G.; Noll, M.: Conservation of the paired domain in metazoans and its structure in three isolated human genes. EMBO J. 8: 1183–1190, 1989.

[77949] 16279.Gruss, P.; Walther, C.: Pax in development. Cell 69: 719–722, 1992.

[77950] 16280.Schafer, B. W.; Mattei, M. G.: The human paired domain gene PAX7(Hup1) maps to chromosome 1p35–1p36.2. Genomics 17: 249–251, 1993.

[77951] 16281.Seale, P.; Sabourin, L. A.; Girgis–Gabardo, A.; Mansouri, A.; Gruss, P.; Rudnicki, M. A.: Pax7 is required for the specification of myogenic satellite cells. Cell 102: 777–786, 2000.

[77952] 16282.Shapiro, D. N.; Sublett, J. E.; Li, B.; Valentine, M. B.; Morris, S. W.; Noll, M.: The gene for PAX7, a member of the paired–box–containing genes, is localized on human chromosome arm 1p36. Genomics 17: 767–769, 1993.

[77953] 16283.Helwig, U.; Imai, K.; Schmahl, W.; Thomas, B. E.; Varnum, D. S.; Nadeau, J. H.; Balling, R.: Interaction between undulated and Patch leads to an extreme form of spina bifida in double–mutant mice. Nature Genet. 11: 60–63, 1995.

- [77954] 16284.Sun, J.; Rose, J. B.; Bird, P.: Gene structure, chromosomal localization, and expression of the murine homologue of human proteinase inhibitor6 (PI-6) suggests divergence of PI-6 from the ovalbumin serpins. *J.Biol. Chem.* 270: 16089–16096, 1995.
- [77955] 16285.Aberle, H.; Bierkamp, C.; Torchard, D.; Serova, O.; Wagner, T.; Natt, E.; Wirsching, J.; Heidkamper, C.; Montagna, M.; Lynch, H. T.; Lenoir, G. M.; Scherer, G.; Feunteun, J.; Kemler, R.: The human plakoglobingene localizes on chromosome 17q21 and is subjected to loss of heterozygosity in breast and ovarian cancers. *Proc. Nat. Acad. Sci.* 92: 6384–6388, 1995.
- [77956] 16286.Bierkamp, C.; Mclaughlin, K. J.; Schwarz, H.; Huber, O.; Kemler, R.: Embryonic heart and skin defects in mice lacking plakoglobin. *Dev.Biol.* 180: 780–785, 1996.
- [77957] 16287.Cowley, C. M. E.; Simrak, D.; Spurr, N. K.; Arneemann, J.; Buxton, R. S.: The plakophilin 1 (PKP1) and plakoglobin (JUP) genes map to human chromosomes 1q and 17, respectively. *Hum. Genet.* 100: 486–488, 1997.
- [77958] 16288.Franke, W. W.; Goldschmidt, M. D.; Zimbelmann, R.; Mueller, H.M.; Schiller, D. L.; Cowin, P.: Molecular cloning and amino acid sequence of human plakoglobin, the common junctional plaque protein. *Proc.Nat. Acad. Sci.* 86:

4027–4031, 1989.

- [77959] 16289.Guenet, J.-L.; Simon-Chazottes, D.; Ringwald, M.; Kemler, R.:The genes coding for alpha and beta catenin (Catna1 and Catnb) and plakoglobin (Jup) map to mouse chromosomes 18, 9, and 11, respectively. *Mammalian Genome* 6: 363–366, 1995.
- [77960] 16290.Knudsen, K. A.; Wheelock, M. J.: Plakoglobin, or an 83-kD homolog distinct from beta-catenin, interacts with E-cadherin and N-cadherin. *J.Cell Biol.* 118: 671–679, 1992.
- [77961] 16291.Kolligs, F. T.; Kolligs, B.; Hajra, K. M.; Hu, G.; Tani, M.; Cho, K. R.; Fearon, E. R.: Gamma-catenin is regulated by the APC tumor suppressor and its oncogenic activity is distinct from that of beta-catenin. *Genes Dev.* 14: 1319–1331, 2000.
- [77962] 16292.Mathur, M.; Goodwin, L.; Cowin, P.: Interactions of the cytoplasmic domain of the desmosomal cadherin Dsg1 with plakoglobin. *J. Biol.Chem.* 269: 14075–14080, 1994.
- [77963] 16293.McKoy, G.; Protonotarios, N.; Crosby, A.; Tsatsopoulou, A.; Anastasakis, A.; Coonar, A.; Norman, M.; Baboonian, C.; Jeffery, S.; McKenna, W.J.: Identification of a deletion in plakoglobin in arrhythmogenic right ventricular cardiomyopathy with palmoplantar keratoderma and–



woolly hair (Naxos disease). *Lancet* 355: 2119–2124, 2000.

[77964] 16294. Ruiz, P.; Brinkmann, V.; Ledermann, B.; Behrend, M.; Grund, C.; Thalhammer, C.; Vogel, F.; Birchmeier, C.; Gunthert, U.; Franke, W.W.; Birchmeier, W.: Targeted mutation of plakoglobin in mice reveals essential functions of desmosomes in the embryonic heart. *J. Cell Biol.* 135: 215–225, 1996.

[77965] 16295. Buckley, M. F.; Goding, J. W.: Plasma cell membrane glycoprotein gene Pca-1 (alkaline phosphodiesterase I) is linked to the proto-oncogene Myb on mouse chromosome 10. *Immunogenetics* 36: 199–201, 1992.

[77966] 16296. Buckley, M. F.; Loveland, K. A.; McKinstry, W. J.; Garson, O. M.; Goding, J. W.: Plasma cell membrane glycoprotein PC-1: cDNA cloning of the human molecule, amino acid sequence, and chromosomal location. *J. Biol. Chem.* 265: 17506–17511, 1990.

[77967] 16297. Frittitta, L.; Baratta, R.; Spampinato, D.; Di Paola, R.; Pizzuti, A.; Vigneri, R.; Trischitta, V.: The Q121 PC-1 variant and obesity have additive and independent effects in causing insulin resistance. *J. Clin. Endocr. Metab.* 86: 5888–5891, 2001.

[77968] 16298. Harahap, A. R.; Goding, J. W.: Distribution of the

murine plasmacell antigen PC-1 in non-lymphoid tissues. J. Immun. 141: 2317-2320,1988.

[77969] 16299.Huang, R.; Rosenbach, M.; Vaughn, R.; Provvedini, D.; Rebbe, N.;Hickman, S.; Goding, J.; Terkeltaub, R.: Expression of the murineplasma cell nucleotide pyrophosphohydrolase PC-1 is shared by humanliver, bone, and cartilage cells: regulation of PC-1 expression inosteosarcoma cells by transforming growth factor-beta. J. Clin. Invest. 94:560-567, 1994.

[77970] 16300.Kahn, C. R.: Causes of insulin resistance. Nature 373: 384-385,1995.

[77971] 16301.Maddux, B. A.; Goldfine, I. D.: Membrane glycoprotein PC-1 inhibitionof insulin receptor function occurs via direct interaction with thereceptor alpha-subunit. Diabetes 49: 13-19, 2000.

[77972] 16302.Maddux, B. A.; Sbraccia, P.; Kumakura, S.; Sasson, S.; Youngren,J.; Fisher, A.; Spencer, S.; Grupe, A.; Henzel, W.; Stewart, T. A.;Reaven, G. M.; Goldfine, I. D.: Membrane glycoprotein PC-1 and insulinresistance in non-insulin-dependent diabetes mellitus. Nature 373:448-451, 1995.

[77973] 16303.Nakamura, I.; Ikegawa, S.; Okawa, A.; Okuda, S.; Koshizuka, Y.;Kawaguchi, H.; Nakamura, K.; Koyama, T.;

Goto, S.; Toguchida, J.; Matsushita, M.; Ochi, T.; Takaoka, K.; Nakamura, Y.: Association of the human NPPS gene with ossification of the posterior longitudinal ligament of the spine (OPLL). *Hum. Genet.* 104: 492–497, 1999.

[77974] 16304. Okawa, A.; Nakamura, I.; Goto, S.; Moriya, H.; Nakamura, Y.; Ikegawa, S.: Mutation in *Npps* in a mouse model of ossification of the posterior longitudinal ligament of the spine. *Nature Genet.* 19: 271–273, 1998.

[77975] 16305. Pizzuti, A.; Frittitta, L.; Argiolas, A.; Baratta, R.; Goldfine, I. D.; Bozzali, M.; Ercolino, T.; Scarlato, G.; Iacoviello, L.; Vigneri, R.; Tassi, V.; Trischitta, V.: A polymorphism (K121Q) of the human glycoprotein PC-1 gene coding region is strongly associated with insulin resistance. *Diabetes* 48: 1881–1884, 1999.

[77976] 16306. Rebbe, N. F.; Tong, B. D.; Finley, E. M.; Hickman, S.: Identification of nucleotide pyrophosphatase/alkaline phosphodiesterase I activity associated with the mouse plasma cell differentiation antigen PC-1. *Proc. Nat. Acad. Sci.* 88: 5192–5196, 1991.

[77977] 16307. Sakaguchi, A. Y.; Lalley, P. A.; Zabel, B. U.; Ellis, R. W.; Scolnick, E. M.; Naylor, S. L.: Chromosome assignments of four mouse cellular homologs of sarcoma and leukemia virus oncogenes. *Proc. Nat. Acad. Sci.* 81: 525–529, 1984.

- [77978] 16308.Takahashi, T.; Old, L. J.; Boyse, E. A.: Surface alloantigen of plasma cells. *J. Exp. Med.* 131: 1325–1341, 1970.
- [77979] 16309.van Driel, I. R.; Goding, J. W.: Plasma cell membrane glycoprotein PC-1: primary structure deduced from cDNA clones. *J. Biol. Chem.* 262:4882–4887, 1987.
- [77980] 16310.Frank, S. L.; Klisak, I.; Sparkes, R. S.; Lusk, A. J.: A gene homologous to plasminogen located on human chromosome 2q11–p11. *Genomics* 4:449–451, 1989.
- [77981] 16311.Westergaard, J. G.; Chemnitz, J.; Teisner, B.; Poulsen, H. K.; Ipsen, L.; Beck, B.; Grudzinski, J. G.: Pregnancy-associated plasma protein A: a possible marker in the classification and prenatal diagnosis of Cornelia de Lange syndrome. *Prenatal Diag.* 3: 225–232, 1983.
- [77982] 16312.Barnett, T.; Pickle, W., II; Rae, P. M. M.; Hart, J.; Kamarck, M.; Elting, J.: Pregnancy-specific beta-1-glycoproteins are related to carcinoembryonic antigens and map to chromosome 19. (Abstract) *Cytogenet. Cell Genet.* 51: 958, 1989.
- [77983] 16313.Barnett, T. R.; Pickle, W., II; Rae, P. M. M.; Hart, J.; Kamarck, M.; Elting, J.: Human pregnancy-specific beta(1)-glycoproteins are coded within chromosome 19. *Am. J. Hum. Genet.* 44: 890–893, 1989.

- [77984] 16314. Bartels, I.; Lindemann, A.: Maternal levels of pregnancy-specific beta-1-glycoprotein (SP-1) are elevated in pregnancies affected by Down's syndrome. *Hum. Genet.* 80: 46-48, 1988.
- [77985] 16315. Brandriff, B. F.; Gordon, L. A.; Tynan, K. T.; Olsen, A. S.; Mohrenweiser, H. W.; Fertitta, A.; Carrano, A. V.; Trask, B. J.: Order and genomic distances among members of the carcinoembryonic antigen (CEA) gene family determined by fluorescence in situ hybridization. *Genomics* 12: 773-779, 1992.
- [77986] 16316. Chan, W.-Y.; Qiu, W.-R.: Human pregnancy-specific beta-1 glycoprotein is encoded by multiple genes localized on two chromosomes. *Am. J. Hum. Genet.* 43: 152-159, 1988.
- [77987] 16317. Khan, W. N.; Teglund, S.; Bremer, K.; Hammarstrom, S.: The pregnancy-specific glycoprotein family of the immunoglobulin superfamily: identification of new members and estimation of family size. *Genomics* 12: 780-787, 1992.
- [77988] 16318. Niemann, S. C.; Flake, A.; Bohn, H.; Bartels, I.: Pregnancy-specific beta-1-glycoprotein: cDNA cloning, tissue expression, and species specificity of one member of the PSBG family. *Hum. Genet.* 82: 239-243, 1989.

- [77989] 16319.Niemann, S. C.; Schonk, D.; van Dijk, P.; Wieringa, B.; Grzeschik,K.-H.; Bartels, I.: Regional localization of the gene encoding pregnancyspecific beta-1-glycoprotein 1 (PSBG1) to human chromosome 19q13.1. Cytogenet.Cell Genet. 52: 95-97, 1989.
- [77990] 16320.Niemann, S. C.; Schonk, D.; van Dijk, P. E.; Grzeschik, K.-H.;Bartels, I.: Chromosomal assignment of a cDNA clone encoding pregnancy-specific-beta-1-glycoprotein to chromosome 19. (Abstract) Cytogenet. CellGenet. 51: 1053, 1989.
- [77991] 16321.Olsen, A.; Teglund, S.; Nelson, D.; Gordon, L.; Copeland, A.;Georgescu, A.; Carrano, A.; Hammarstrom, S.: Gene organization ofthe pregnancy-specific glycoprotein region on human chromosome 19:assembly and analysis of a 700-kb cosmid contig spanning the region. Genomics 23:659-668, 1994.
- [77992] 16322.Streydio, C.; Swillens, S.; Georges, M.; Szpirer, C.; Vassart,G.: Structure, evolution and chromosomal localization of the humanpregnancy-specific beta-1 glycoprotein gene family. Genomics 6:579-592, 1990. Note: Erratum: Genomics 7: 661-662, 1990.
- [77993] 16323.Teglund, S.; Olsen, A.; Khan, W. N.; Frangsmyr, L.; Hammarstrom,S.: The pregnancy-specific glycoprotein

(PSG) gene cluster on human chromosome 19: fine structure of the 11 PSG genes and identification of 6 new genes forming a third subgroup within the carcinoembryonic antigen (CEA) family. *Genomics* 23: 669–684, 1994.

[77994] 16324. Thompson, J.; Koumari, R.; Wagner, K.; Barnert, S.; Schleussner, C.; Schrewe, H.; Zimmermann, W.; Muller, G.; Schempp, W.; Zaninetta, D.; Ammaturo, D.; Hardman, N.: The human pregnancy-specific glycoprotein genes are tightly linked on the long arm of chromosome 19 and are coordinately expressed. *Biochem. Biophys. Res. Commun.* 167: 848–859, 1990.

[77995] 16325. Watanabe, S.; Chou, J. Y.: Isolation and characterization of complementary DNAs encoding human pregnancy-specific beta-1-glycoprotein. *J. Biol. Chem.* 263: 2049–2054, 1988.

[77996] 16326. Leslie, K. K.; Watanabe, S.; Lei, K.-J.; Chou, D. Y.; Plouzek, C. A.; Deng, H.-C.; Torres, J.; Chou, J. Y.: Linkage of two human pregnancy-specific beta-1-glycoprotein genes: one is associated with hydatidiform mole. *Proc. Nat. Acad. Sci.* 87: 5822–5826, 1990.

[77997] 16327. Olsen, A.; Teglund, S.; Nelson, D.; Gordon, L.; Copeland, A.; Georgescu, A.; Carrano, A.; Hammarstrom, S.: Gene organization of the pregnancy-speci-

ficglycoprotein region on human chromosome 19: assembly and analysis of a 700-kb cosmid contig spanning the region. *Genomics* 23: 659–668, 1994.

[77998] 16328. Thompson, J.; Zimmermann, W.; Osthus-Bugat, P.; Schleussner, C.; Eades-Perner, A.-M.; Barnert, S.; von Kleist, S.; Willcocks, T.; Craig, I.; Tynan, K.; Olsen, A.; Mohrenweiser, H.: Long-range chromosomal mapping of the carcinoembryonic antigen (CEA) gene family cluster. *Genomics* 12: 761–772, 1992.

[77999] 16329. Tynan, K.; Olsen, A.; Trask, B.; de Jong, P.; Thompson, J.; Zimmermann, W.; Carrano, A.; Mohrenweiser, H.: Assembly and analysis of cosmid contigs in the CEA-gene family region of human chromosome 19. *Nucleic Acids Res.* 20: 1629–1636, 1992.

[78000] 16330. Sheer, D.; Sheppard, D. M.; Le Beau, M.; Rowley, J. D.; San Roman, C.; Solomon, E.: Localization of the oncogene c-erbA1 immediately proximal to the acute promyelocytic leukaemia breakpoint on chromosome 17. *Ann. Hum. Genet.* 49: 167–171, 1985.

[78001] 16331. Yoshitake, H.; Rittling, S. R.; Denhardt, D. T.; Noda, M.: Osteopontin-deficient mice are resistant to ovariectomy-induced bone resorption. *Proc. Nat. Acad. Sci.* 96: 8156–8160, 1999.



- [78002] 16332. Edelman, W.; Zervas, M.; Costello, P.; Roback, L.; Fischer, I.; Hammarback, J. A.; Cowan, N.; Davies, P.; Wainer, B.; Kucherlapati, R.: Neuronal abnormalities in microtubule-associated protein 1B mutant mice. *Proc. Nat. Acad. Sci.* 93: 1270–1275, 1996.
- [78003] 16333. Hammarback, J. A.; Obar, R. A.; Hughes, S. M.; Vallee, R. B.: MAP1B is encoded as a polyprotein that is processed to form a complex N-terminal microtubule-binding domain. *Neuron* 7: 129–139, 1991.
- [78004] 16334. Lien, L. L.; Boyce, F. M.; Kleyn, P.; Brzustowicz, L. M.; Menninger, J.; Ward, D. C.; Gilliam, T. C.; Kunkel, L. M.: Mapping of human microtubule-associated protein 1B in proximity to the spinal muscular atrophy locus at 5q13. *Proc. Nat. Acad. Sci.* 88: 7873–7876, 1991.
- [78005] 16335. Lien, L. L.; Feener, C. A.; Fischbach, N.; Kunkel, L. M.: Cloning of human microtubule-associated protein 1B and the identification of a related gene on chromosome 15. *Genomics* 22: 273–280, 1994.
- [78006] 16336. Wirth, B.; Voosen, B.; Rohrig, D.; Knapp, M.; Piechaczek, B.; Rudnik-Schoneborn, S.; Zerres, K.: Fine mapping and narrowing of the genetic interval of the spinal muscular atrophy region by linkage studies. *Genomics* 15: 113–118, 1993.

- [78007] 16337.Zhang, Y. Q.; Bailey, A. M.; Matthies, H. J. G.; Renden, R. B.;Smith, M. A.; Speese, S. D.; Rubin, G. M.; Broadie, K.: *Drosophila* fragile X-related gene regulates the MAP1B homolog Futsch to control synaptic structure and function. *Cell* 107: 591–603, 2001.
- [78008] 16338.Garner, C. C.; Tucker, R. P.; Matus, A.: Selective localization of messenger RNA for cytoskeletal protein MAP2 in dendrites. *Nature* 336:674–677, 1988.
- [78009] 16339.Kalcheva, N.; Albala, J.; O'Guin, K.; Rubino, H.; Garner, C.; Shafit-Zagardo, B.: Genomic structure of human microtubule-associated protein 2 (MAP-2) and characterization of additional MAP-2 isoforms. *Proc. Nat. Acad. Sci.* 92: 10894–10898, 1995.
- [78010] 16340.Kindler, S.; Garner, C. C.: Four repeat MAP2 isoforms in human and rat brain. *Molec. Brain Res.* 26: 218–224, 1994.
- [78011] 16341.Marsden, K. M.; Doll, T.; Ferralli, J.; Botteri, F.; Matus, A.: Transgenic expression of embryonic MAP2 in adult mouse brain: implications for neuronal polarization. *J. Neurosci.* 16: 3265–3273, 1996.
- [78012] 16342.Neve, R. L.; Harris, P.; Kosik, K. S.; Kurnit, D. M.; Donlon, T.A.: Identification of cDNA clones for the human microtubule-associated protein tau and chromosomal lo-

calization of the genes for tau and microtubule-associated protein 2. *Molec. Brain Res.* 1: 271–280, 1986.

[78013] 16343. Chapin, S. J.; Bulinski, J. C.: Non-neuronal 210 x 10(3) M(r) microtubule-associated protein (MAP4) contains a domain homologous to the microtubule-binding domains of neuronal MAP2 and tau. *J. Cell Sci.* 98: 27–36, 1991.

[78014] 16344. Adriaansen, H. J.; Geurts Van Kessel, A. H. M.; Wijdenes–De Bresser, J. H. F. M.; Van Drunen–Schoenmaker, E.; Van Dongen, J. J. M.: Expression of the myeloid differentiation antigen CD33 depends on the presence of human chromosome 19 in human–mouse hybrids. *Ann. Hum. Genet.* 54: 115–119, 1990.

[78015] 16345. Bonthron, D. T.; Dunlop, N.; Barr, D. G. D.; El Sanousi, A. A.; Al–Gazali, L. I.: Organisation of the human PAX4 gene and its exclusion as a candidate for the Wolcott–Rallison syndrome. *J. Med. Genet.* 35: 288–292, 1998.

[78016] 16346. Inoue, H.; Nomiyama, J.; Nakai, K.; Matsutani, A.; Tanizawa, Y.; Oka, Y.: Isolation of full-length cDNA of mouse PAX4 gene and identification of its human homologue. *Biochem. Biophys. Res. Commun.* 243: 628–633, 1998.

[78017] 16347. Mansouri, A.; St–Onge, L.; Gruss, P.: Role of Pax

genes in endoderm-derived organs. Trends Endocr. Metab. 10: 164–167, 1999.

- [78018] 16348. Matsushita, T.; Yamaoka, T.; Otsuka, S.; Moritani, M.; Matsumoto, T.; Itakura, M.: Molecular cloning of mouse paired-box-containing gene (Pax)-4 from an islet beta cell line and deduced sequence of human Pax-4. Biochem. Biophys. Res. Commun. 242: 176–180, 1998.
- [78019] 16349. Sosa-Pineda, B.; Chowdhury, K.; Torres, M.; Oliver, G.; Gruss, P.: The Pax4 gene is essential for differentiation of insulin-producing beta cells in the mammalian pancreas. Nature 386: 399–402, 1997.
- [78020] 16350. St-Onge, L.; Sosa-Pineda, B.; Chowdhury, K.; Mansouri, A.; Gruss, P.: Pax6 is required for differentiation of glucagon-producing alpha-cells in mouse pancreas. Nature 387: 406–409, 1997.
- [78021] 16351. Tamura, T.; Izumikawa, Y.; Kishino, T.; Soejima, H.; Jinno, Y.; Niikawa, N.: Assignment of the human PAX4 gene to chromosome band 7q32 by fluorescence in situ hybridization. Cytogenet. Cell Genet. 66:132–134, 1994.
- [78022] 16352. Adams, B.; Dorfler, P.; Aguzzi, A.; Kozmik, Z.; Urbanek, P.; Maurer-Fogy, I.; Busslinger, M.: Pax-5 encodes the transcription factor BSAP and is expressed in B lymphocytes, the developing CNS, and adult testis. Genes Dev.

6: 1589–1607, 1992.

- [78023] 16353. Busslinger, M.; Klix, N.; Pfeffer, P.; Graninger, P. G.; Kozmik, Z.: Deregulation of PAX-5 by translocation of the E- $\mu$  enhancer of the IgH locus adjacent to two alternative PAX-5 promoters in a diffuse large-cell lymphoma. *Proc. Nat. Acad. Sci.* 93: 6129–6134, 1996.
- [78024] 16354. Mikkola, I.; Heavey, B.; Horcher, M.; Busslinger, M.: Reversion of B cell commitment upon loss of Pax5 expression. *Science* 297:110–113, 2002.
- [78025] 16355. Nutt, S. L.; Heavey, B.; Rolink, A. G.; Busslinger, M.: Commitment to the B-lymphoid lineage depends on the transcription factor Pax5. *Nature* 401:556–562, 1999.
- [78026] 16356. Nutt, S. L.; Vambrie, S.; Steinlein, P.; Kozmik, Z.; Rolink, A.; Weith, A.; Busslinger, M.: Independent regulation of the two Pax5 alleles during B-cell development. *Nature Genet.* 21: 390–395, 1999.
- [78027] 16357. Ohno, H.; Furukawa, T.; Fukuhara, S.; Zong, S. Q.; Kamesaki, H.; Shows, T. B.; Le Beau, M. M.; McKeithan, T. W.; Kawakami, T.; Honjo, T.: Molecular analysis of a chromosomal translocation, t(9;14)(p13;q32), in a diffuse large-cell lymphoma cell line expressing the Ki-1 antigen. *Proc. Nat. Acad. Sci.* 87: 628–632, 1990.
- [78028] 16358. Rolink, A. G.; Nutt, S. L.; Melchers, F.; Busslinger,

M.: Long-term *in vivo* reconstitution of T-cell development by Pax5-deficient B-cell progenitors. *Nature* 401: 603–606, 1999.

- [78029] 16359. Vorechovsky, I.; Koskinen, S.; Paganelli, R.; Smith, C. I. E.; Busslinger, M.; Hammarstrom, L.: The PAX5 gene: a linkage and mutation analysis in candidate human primary immunodeficiencies. *Immunogenetics* 42:149–152, 1995.
- [78030] 16360. Walther, C.; Guenet, J.-L.; Simon, D.; Deutsch, U.; Jostes, B.; Goulding, M. D.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a murine multigene family of paired box containing genes. *Genomics* 11: 424–434, 1991.
- [78031] 16361. Koseki, H.; Zachgo, J.; Mizutani, Y.; Simon-Chazottes, D.; Guenet, J.-L.; Balling, R.; Gossler, A.: Fine genetic mapping of the proximal part of mouse chromosome 2 excludes Pax-8 as a candidate gene for Danforth's short tail (Sd). *Mammalian Genome* 4: 324–327, 1993.
- [78032] 16362. Kroll, T. G.; Sarraf, P.; Pecciarini, L.; Chen, C.-J.; Mueller, E.; Splegeman, B. M.; Fletcher, J. A.: PAX8-PPAR-gamma-1 fusion in oncogene human thyroid carcinoma. *Science* 289: 1357–1360, 2000.
- [78033] 16363. Macchia, P. E.; Lapi, P.; Krude, H.; Pirro, M. T.; Missero, C.; Chiovato, L.; Souabni, A.; Baserga, M.; Tassi, V.; Pinchera, A.; Fenzi, G.; Gruters, A.; Busslinger, M.; Di

Lauro, R.: PAX8 mutations associated with congenital hypothyroidism caused by thyroid dysgenesis. *Nature Genet.* 19: 83–86, 1998.

[78034] 16364. Mansouri, A.; Chowdhury, K.; Gruss, P.: Follicular cells of the thyroid gland require Pax8 gene function. *Nature Genet.* 19: 87–90, 1998.

[78035] 16365. Pasca di Magliano, M.; Di Lauro, R.; Zannini, M.: Pax8 has a key role in thyroid cell differentiation. *Proc. Nat. Acad. Sci.* 97: 13144–13149, 2000.

[78036] 16366. Plachov, D.; Chowdhury, K.; Walther, C.; Simon, D.; Guenet, J.-L.; Gruss, P.: Pax-8, a murine paired box gene expressed in the developing excretory system and thyroid gland. *Development* 110: 643–651, 1990.

[78037] 16367. Tell, G.; Pellizzari, L.; Esposito, G.; Pucillo, C.; Macchia, P. E.; Di Lauro, R.; Damante, G.: Structural defects of a Pax8 mutant that give rise to congenital hypothyroidism. *Biochem. J.* 341: 89–93, 1999.

[78038] 16368. Bongarzone, I.; Vigano, E.; Alberti, L.; Borrello, M. G.; Pasini, B.; Greco, A.; Mondellini, P.; Smith, D. P.; Ponder, B. A. J.; Romeo, G.; Pierotti, M. A.: Full activation of MEN2B mutant RET by an additional MEN2A mutation or by ligand GDNF stimulation. *Oncogene* 16: 2295–2301, 1998.

[78039] 16369. Carlson, K. M.; Bracamontes, J.; Jackson, C. E.;

Clark, R.; Lacroix, A.; Wells, S. A., Jr.; Goodfellow, P. J.: Parent-of-origin effects in multiple endocrine neoplasia type 2B. *Am. J. Hum. Genet.* 55:1076–1082, 1994.

[78040] 16370. Carlson, K. M.; Dou, S.; Chi, D.; Scavarda, N.; Toshima, K.; Jackson, C. E.; Wells, S. A., Jr.; Goodfellow, P. J.; Donis-Keller, H.: Single missense mutation in the tyrosine kinase catalytic domain of the RET protooncogene is associated with multiple endocrine neoplasia type 2B. *Proc. Nat. Acad. Sci.* 91: 1579–1583, 1994.

[78041] 16371. Mulligan, L. M.; Eng, C.; Healey, C. S.; Clayton, D.; Kwok, J. B. J.; Gardner, E.; Ponder, M. A.; Frilling, A.; Jackson, C. E.; Lehnert, H.; Neumann, H. P. H.; Thibodeau, S. N.; Ponder, B. A. J.: Specific mutations of the RET protooncogene are related to disease phenotype in MEN 2A and FMTC. *Nature Genet.* 6: 70–74, 1994.

[78042] 16372. Bruss, M.; Kunz, J.; Linggen, B.; Bonisch, H.: Chromosomal mapping of the human gene for the tricyclic antidepressant-sensitive noradrenaline transporter. *Hum. Genet.* 91: 278–280, 1993.

[78043] 16373. Esler, M.; Jennings, G.; Lambert, G.; Meredith, I.; Horne, M.; Eisenhofer, G.: Overflow of catecholamine neurotransmitters to the circulation: source, fate, and functions. *Physiol. Rev.* 70: 963–985, 1990.



- [78044] 16374.Fritz, J. D.; Jayanthi, L. D.; Thoreson, M. A.; Blakely, R. D.: Cloning and chromosomal mapping of the murine norepinephrine transporter. *J.Neurochem.* 70: 2241–2251, 1998.
- [78045] 16375.Gelernter, J.; Kruger, S.; Kidd, K. K.; Amara, S.: TaqI RFLP at norepinephrine transporter protein (NET) locus. *Hum. Molec. Genet.* 2:820 only, 1993.
- [78046] 16376.Gelernter, J.; Kruger, S.; Pakstis, A. J.; Pacholczyk, T.; Sparkes, R. S.; Kidd, K. K.; Amara, S.: Assignment of the norepinephrine transporter protein (NET1) locus to chromosome 16. *Genomics* 18: 690–692, 1993.
- [78047] 16377.Kaye, W. H.; Jimerson, D. C.; Lake, C. R.; Ebert, M. H.: Altered norepinephrine metabolism following long-term weight recovery in patients with anorexia nervosa. *Psychiat. Res.* 14: 333–342, 1985.
- [78048] 16378.Pacholczyk, T.; Blakely, R. D.; Amara, S. G.: Expression cloning of a cocaine- and antidepressant-sensitive human noradrenaline transporter. *Nature* 350:350–354, 1991.
- [78049] 16379.Paczkowski, F. A.; Bonisch, H.; Bryan-Lluka, L. J.: Pharmacological properties of the naturally occurring ala457pro variant of the human norepinephrine transporter. *Pharmacogenetics* 12: 165–173, 2002.

- [78050] 16380.Pirke, K. M.; Kellner, M.; Philipp, E.; Laessle, R.; Krieg, J.C.; Fichter, M. M.: Plasma norepinephrine after a standardized testmeal in acute and remitted patients with anorexia nervosa and in healthycontrols. Biol. Psychiat. 31: 1074–1077, 1992.
- [78051] 16381.Porzgen, P.; Bonisch, H.; Bruss, M.: Molecular cloning and organizationof the coding region of the human norepinephrine transporter gene. Biochem.Biophys. Res. Commun. 215: 1145–1150, 1995. Notes: Erratum: Biochem.Biophys. Res. Commun. 227: 642–643, 1996.
- [78052] 16382.Shannon, J. R.; Flattem, N. L.; Jordan, J.; Jacob, G.; Black,B. K.; Biaggioni, I.; Blakely, R. D.; Robertson, D.: Orthostaticintolerance and tachycardia associated with norepinephrine–transporterdeficiency. New Eng. J. Med. 342: 541–549, 2000.
- [78053] 16383.Urwin, R. E.; Bennetts, B.; Wilcken, B.; Lampropoulos, B.; Beumont,P.; Clarke, S.; Russell, J.; Tanner, S.; Nunn, K. P.: Anorexia nervosa(restrictive subtype) is associated with a polymorphism in the novelnorepinephrine transporter gene promoter polymorphic region. Molec.Psychiat. 7: 652–657, 2002.
- [78054] 16384.Tomilin, A.; Remenyi, A.; Lins, K.; Bak, H.; Leidel, S.; Vriend,G.; Wilmanns, M.; Scholer, H. R.: Synergism with

the coactivator OBF-1 (OCA-B, BOB-1) is mediated by a specific POU dimer configuration. *Cell* 103:853–864, 2000.

- [78055] 16385. Ko, H.-S.; Fast, P.; McBride, W.; Staudt, L. M.: A human protein specific for the immunoglobulin octamer DNA motif contains a functional homeobox domain. *Cell* 55: 135–144, 1988.
- [78056] 16386. Schubart, K.; Massa, S.; Schubart, D.; Corcoran, L. M.; Rolink, A. G.; Matthias, P.: B cell development and immunoglobulin gene transcription in the absence of Oct-2 and OBF-1. *Nature Immun.* 2: 69–74, 2001.
- [78057] 16387. Staudt, L. M.; Clerc, R. G.; Singh, H.; LeBowitz, J. H.; Sharp, P. A.; Baltimore, D.: Cloning of a lymphoid-specific cDNA encoding a protein binding the regulatory octamer DNA motif. *Science* 241:577–580, 1988.
- [78058] 16388. Crouau-Roy, B.; Amadou, C.; Bouissou, C.; Clayton, J.; Vernet, C.; Ribouchon, M.-T.; Pontarotti, P.: Localization of the OTF3 gene within the human MHC class I region by physical and meiotic mapping. *Genomics* 21:241–243, 1994.
- [78059] 16389. Guillaudeux, T.; Mattei, M. G.; Depetris, D.; Le Bouteiller, P.; Pontarotti, P.: In situ hybridization localizes the human OTF3 to chromosome 6p21.3–p22 and OTF3L

to 12p13. Cytogenet. Cell Genet. 63:212–214, 1993.

[78060] 16390.Nichols, J.; Zevnik, B.; Anastassiadis, K.; Niwa, H.; Klewe–Nebenius,D.; Chambers, I.; Scholer, H.; Smith, A.: Formation of pluripotentstem cells in the mammalian embryo depends on the POU transcriptionfactor Oct4. Cell 95: 379–391, 1998.

[78061] 16391.Niwa, H.; Miyazaki, J.; Smith, A. G.: Quantitative expressionof Oct–3/4 defines differentiation, dedifferentiation or self–renewalof ES cells. Nature Genet. 24: 372–376, 2000.

[78062] 16392.Rosner, M. H.; De Santo, R. J.; Arnheiter, H.; Staudt, L. M.:Oct–3 is a maternal factor required for the first mouse embryonicdivision. Cell 64: 1103–1110, 1991.  
Note: Retraction: Cell 69: 724only, 1992.

[78063] 16393.Takeda, J.; Seino, S.; Bell, G. I.: Human Oct3 gene family: cDNAsequences, alternative splicing, gene organization, chromosomal location,and expression at low levels in adult tissues. Nucleic Acids Res. 20:4613–4620, 1992.

[78064] 16394.Parker, R. C.; Mardon, G.; Lebo, R. V.; Varmus, H. E.; Bishop,J. M.: Isolation of duplicated human c–src genes located on chromosomes1 and 20. Molec. Cell. Biol. 5: 831–838, 1985.

[78065] 16395.Tronick, S. R.; Popescu, N. C.; Cheah, M. S. C.;

Swan, D. C.; Amsbaugh, S. C.; Lengel, C. R.; DiPaolo, J. A.; Robbins, K. C.: Isolation and chromosomal localization of the human *fgr* protooncogene, a distinct member of the tyrosine kinase gene family. *Proc. Nat. Acad. Sci.* 82:6595–6599, 1985.

[78066] 16396. Krebs, L. T.; Xue, Y.; Norton, C. R.; Shutter, J. R.; Maguire, M.; Sundberg, J. P.; Gallahan, D.; Closson, V.; Kitajewski, J.; Callahan, R.; Smith, G. H.; Stark, K. L.; Gridley, T.: Notch signaling is essential for vascular morphogenesis in mice. *Genes Dev.* 14: 1343–1352, 2000.

[78067] 16397. Li, L.; Huang, G. M.; Banta, A. B.; Deng, Y.; Smith, T.; Dong, P.; Friedman, C.; Chen, L.; Trask, B. J.; Spies, T.; Rowen, L.; Hood, L.: Cloning, characterization, and the complete 56.8-kilobase DNA sequence of the human NOTCH4 gene. *Genomics* 51: 45–58, 1998.

[78068] 16398. McGinnis, R. E.; Fox, H.; Yates, P.; Cameron, L.-A.; Barnes, M. R.; Gray, I. C.; Spurr, N. K.; Hurko, O.; St. Clair, D.: Failure to confirm NOTCH4 association with schizophrenia in a large population-based sample from Scotland. *Nature Genet.* 28: 128–129, 2001.

[78069] 16399. Sklar, P.; Schwab, S. G.; Williams, N. M.; Daly, M.; Schaffner, S.; Maier, W.; Albus, M.; Trixler, M.; Eichhammer, P.; Lerer, B.; Hallmayer, J.; Norton, N.; and 11 others:

Association analysis of NOTCH4 loci in schizophrenia using family and population-based controls. *Nature Genet.* 28: 126–128, 2001.

- [78070] 16400. Sugaya, K.; Sasanuma, S.; Nohata, J.; Kimura, T.; Fukagawa, T.; Nakamura, Y.; Ando, A.; Inoko, H.; Ikemura, T.; Mita, K.: Gene organization of human NOTCH4 and (CTG)<sub>n</sub> polymorphism in this human counterpart gene of mouse proto-oncogene *Int3*. *Gene* 189: 235–244, 1997.
- [78071] 16401. Uyttendaele, H.; Marazzi, G.; Wu, G.; Yan, Q.; Sassoon, D.; Kitajewski, J.: *Notch4/int-3*, a mammary proto-oncogene, is an endothelial cell-specific mammalian Notch gene. *Development* 122: 2251–2259, 1996.
- [78072] 16402. Wei, J.; Hemmings, G. P.: The NOTCH4 locus is associated with susceptibility to schizophrenia. (Letter) *Nature Genet.* 25: 376–377, 2000.
- [78073] 16403. Bork, P.: The modular architecture of a new family of growth regulators related to connective tissue growth factor. *FEBS Lett.* 327: 125–130, 1993.
- [78074] 16404. Sugaya, K.; Fukagawa, T.; Matsumoto, K.; Mita, K.; Takahashi, E.; Ando, A.; Inoko, H.; Ikemura, T.: Three genes in the human MHC class III region near the junction with the class II: gene for receptor of advanced glycosylation end products, PBX2 homeobox gene and a Notch ho-

molog, human counterpart of mouse mammary tumor gene int-3. *Genomics* 23:408–419, 1994.

[78075] 16405.Joliot, V.; Martinerie, C.; Dambrine, G.; Plassiart, G.; Brisac,M.; Crochet, J.; Perbal, B.: Proviral rearrangements and overexpressionof a new cellular gene (nov) in myeloblastosis–associated virus type1–induced nephroblastomas. *Molec. Cell. Biol.* 12: 10–21, 1992.

[78076] 16406.Kim, H.–S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T.,Jr.; Rosenfeld, R. G.: Identification of a family of low–affinityinsulin–like growth factor binding proteins (IFGBPs): characterizationof connective tissue growth factor as a member of the IGFBP superfamily. *Proc.Nat. Acad. Sci.* 94: 12981–12986, 1997.

[78077] 16407.Martinerie, C.; Perbal, B.: Expression of a gene encoding a novelpotential IGF binding protein in human tissues. *C. R. Acad. Sci.(Paris)* 313 (ser. 3): 345–351, 1991.

[78078] 16408.Snaith, M. R.; Natarajan, D.; Taylor, L. B.; Choi, C.–P.; Martinerie,C.; Perbal, B.; Schofield, P. N.; Boulter, C. A.: Genomic structureand chromosomal mapping of the mouse nov gene. *Genomics* 38: 425–428,1996.

[78079] 16409.Soret, J.; Dambrine, G.; Perbal, B.: Induction of nephroblastomaby myeloblastosis–associated virus type 1: state of proviral DNAsin tumor cells. *J. Virol.* 63:

1803–1807, 1989.

- [78080] 16410.Amson, R.; Sigaux, F.; Przedborski, S.; Flandrin, G.; Givol, D.;Telerman, A.: The human protooncogene product p33pim is expressedduring fetal hematopoiesis and in diverse leukemias. *Proc. Nat. Acad.Sci.* 86: 8857–8861, 1989.
- [78081] 16411.Ark, B.; Gummere, G.; Bennett, D.; Artzt, K.: Mapping of the Pim-1oncogene in mouse t-haplotypes and its use to define the relativemap positions of the tcl loci t-0(t-6) and t-w12 and the marker tf(tufted). *Genomics* 10: 385–389, 1991.
- [78082] 16412.Cuypers, H. T.; Selten, G.; Berns, A.; Geurts van Kessel, A. H.M.: Assignment of the human homologue of Pim-1, a mouse gene implicatedin leukemogenesis, to the pter-q12 region of chromosome 6. *Hum. Genet.* 72:262–265, 1986.
- [78083] 16413.Domen, J.; von Lindern, M.; Hermans, A.; Breuer, M.; Grosveld,G.; Berns, A.: Comparison of the human and mouse PIM-1 cDNAs: nucleotidesequence and immunological identification of the in vitro synthesizedPIM-1 protein. *Oncogene Res.* 1: 103–112, 1987.
- [78084] 16414.Boitard, C.; Villa, M. C.; Becourt, C.; Pham Gia, H.; Huc, C.;Sempe, P.; Portier, M. M.; Bach, J. F.: Peripherin: an



islet antigen that is cross-reactive with nonobese diabetic mouse class II gene products. *Proc. Nat. Acad. Sci.* 89: 172–176, 1992.

[78085] 16415. Leonard, D. G.; Gorham, J. D.; Cole, P.; Greene, L. A.; Ziff, E. B.: A nerve growth factor-regulated messenger RNA encodes a new intermediate filament protein. *J. Cell Biol.* 106: 181–193, 1988.

[78086] 16416. Moncla, A.; Landon, F.; Mattei, M.-G.; Portier, M.-M.: Chromosomal localisation of the mouse and human peripherin genes. *Genet. Res.* 59: 125–129, 1992.

[78087] 16417. Pendleton, J. W.; Violette, S. M.; Hunihan, L. W.; Greene, L. A.; Ruddle, F. H.: The peripherin gene maps to mouse chromosome 15. *Genomics* 9: 369–372, 1991.

[78088] 16418. Portier, M.-M.; de Nechaud, B.; Gros, F.: Peripherin, a new member of the intermediate filament protein family. *Dev. Neurosci.* 6: 335–344, 1984.

[78089] 16419. Thompson, M. A.; Ziff, E. B.: Structure of the gene encoding peripherin, an NGF-regulated neuronal-specific type III intermediate filament protein. *Neuron* 2: 1043–1053, 1989.

[78090] 16420. Ben Othmane, K.; Loeb, D.; Hayworth-Hodgite, R.; Hentati, F.; Rao, N.; Roses, A. D.; Ben Hamida, M.; Pericak-Vance, M. A.; Vance, J. M.: Physical and genetic mapping of

the CMT4A locus and exclusion of PMP-2 as the defect in CMT4A. *Genomics* 28: 286–290, 1995.

- [78091] 16421. Hayasaka, K.; Himoro, M.; Takada, G.; Takahashi, E.; Minoshima, S.; Shimizu, N.: Structure and localization of the gene encoding human peripheral myelin protein 2 (PMP2). *Genomics* 18: 244–248, 1993.
- [78092] 16422. Hayasaka, K.; Nanao, K.; Tahara, M.; Sato, W.; Takada, G.; Miura, M.; Uyemura, K.: Isolation and sequence determination of cDNA encoding P2 protein of human peripheral myelin. *Biochem. Biophys. Res. Commun.* 181: 204–207, 1991.
- [78093] 16423. Narayanan, V.; Ripepi, B.; Jabs, E. W.; Hawkins, A.; Griffin, C.; Tennekoon, G.: Partial structure and mapping of the human myelin P2 protein gene. *J. Neurochem.* 63: 2010–2013, 1994.
- [78094] 16424. Manser, E.; Loo, T.-H.; Koh, C.-G.; Zhao, Z.-S.; Chen, X.-Q.; Tan, L.; Tan, I.; Leung, T.; Lim, L.: PAK kinases are directly coupled to the PIX family of nucleotide exchange factors. *Molec. Cell* 1: 183–192, 1998.
- [78095] 16425. Nomura, N.; Miyajima, N.; Sazuka, T.; Tanaka, A.; Kawarabayashi, Y.; Sato, S.; Nagase, T.; Seki, N.; Ishikawa, K.; Tabata, S.: Prediction of the coding sequences of unidentified human genes. I. The coding sequences of 40

new genes (KIAA0001–KIAA0040) deduced by analysis of randomly sampled cDNA clones from human immature myeloid cell line, KG–1. DNA Res. 1: 27–35, 1994.

- [78096] 16426. Buggy, J. J.; Sideris, M. L.; Mak, P.; Lorimer, D. D.; McIntosh, B.; Clark, J. M.: Cloning and characterization of a novel human histone deacetylase, HDAC8. Biochem. J. 350: 199–205, 2000.
- [78097] 16427. Hu, E.; Chen, Z.; Fredrickson, T.; Zhu, Y.; Kirkpatrick, R.; Zhang, G.–F.; Johanson, K.; Sung, C.–M.; Liu, R.; Winkler, J.: Cloning and characterization of a novel human class I histone deacetylase that functions as a transcription repressor. J. Biol. Chem. 275: 15254–15264, 2000.
- [78098] 16428. Van den Wyngaert, I.; de Vries, W.; Kremer, A.; Neefs, J.–M.; Verhasselt, P.; Luyten, W. H. M. L.; Kass, S. U.: Cloning and characterization of human histone deacetylase 8. FEBS Lett. 478: 77–83, 2000.
- [78099] 16429. Grozinger, C. M.; Hassig, C. A.; Schreiber, S. L.: Three proteins define a class of human histone deacetylases related to yeast Hda1p. Proc. Nat. Acad. Sci. 96: 4868–4873, 1999.
- [78100] 16430. Hubbert, C.; Guardiola, A.; Shao, R.; Kawaguchi, Y.; Ito, A.; Nixon, A.; Yoshida, M.; Wang, X.–F.; Yao, T.–P.: HDAC6 is a microtubule-associated deacetylase. Nature

417: 455–458, 2002.

- [78101] 16431.Mahlknecht, U.; Schnittger, S.; Landgraf, F.; Schoch, C.; Ottmann,O. G.; Hiddemann, W.; Hoelzer, D.: Assignment of the human histonedeacetylase 6 gene (HDAC6) to X chromosome p11.23 by in situ hybridization. *Cytogenet.Cell Genet.* 93: 135–136, 2001.
- [78102] 16432.Nagase, T.; Isikawa, K.; Suyama, M.; Kikuno, R.; Hirose, M.;Miyajima, N.; Tanaka, A.; Kotani, H.; Nomura, N.; Ohara, O.: Predictionof the coding sequences of unidentified human genes. XII. The complete sequences of 100 new cDNA clones from brain which code for large proteinsin vitro. *DNA Res.* 5: 355–364, 1998.
- [78103] 16433.Disteche, C. M.; Brannan, C. I.; Larsen, A.; Adler, D. A.; Schorderet,D. F.; Gearing, D.; Copeland, N. G.; Jenkins, N. A.; Park, L. S.:The human pseudoautosomal GM-CSF receptor alpha subunit gene is autosomal in mouse. *Nature Genet.* 1: 333–336, 1992.
- [78104] 16434.Sugarman, G. I.; Rimoin, D. L.; Lachman, R. S.: The facial–digital–genital(Aarskog) syndrome. *Am. J. Dis. Child.* 126: 248–252, 1973.
- [78105] 16435.Tsukahara, M.; Fernandez, G. I.: Umbilical findings in Aarskogsyndrome. *Clin. Genet.* 45: 260–265, 1994.
- [78106] 16436.Tyrkus, M.; Bawle, E.; Lipman, S.; Bozimowski, D.;

Woolley, P.V., Jr.: Aarskog–Scott syndrome inherited as an X-linked dominant with full male–female expression.

(Abstract) Am. J. Hum. Genet. 32:134A only, 1980.

- [78107] 16437. van den Bergh, P.; Fryns, J. P.; Wilms, G.; Piot, R.; Dralands, G.; van den Bergh, R.: Anomalous cerebral venous drainage in Aarskog syndrome. Clin. Genet. 25: 288–294, 1984.
- [78108] 16438. Zheng, Y.; Fischer, D. J.; Santos, M. F.; Tigyi, G.; Pasteris, N. G.; Gorski, J. L.; Xu, Y.: The faciogenital dysplasia gene product FGD1 functions as a Cdc42Hs–specific guanine–nucleotide exchange factor. J. Biol. Chem. 271: 33169–33172, 1996.
- [78109] 16439. Inaba, H.; Fujimaki, M.; Kazazian, H. H., Jr.; Antonarakis, S. E.: MspI polymorphic site in intron 22 of the factor VIII gene in the Japanese population. Hum. Genet. 84: 214–215, 1990.
- [78110] 16440. Levinson, B.; Bermingham, J. R., Jr.; Metzenberg, A.; Kenwrick, S.; Chapman, V.; Gitschier, J.: Sequence of the human factor VIII–associated gene is conserved in mouse. Genomics 13: 862–865, 1992.
- [78111] 16441. Levinson, B.; Kenwrick, S.; Lakich, D.; Hammonds, G.; Gitschier, J.: A transcribed gene in an intron of the human factor VIII gene. Genomics 7: 1–11, 1990.

- [78112] 16442.Peters, M. F.; Ross, C. A.: Isolation of a 40-kDa huntingtin-associated protein. *J. Biol. Chem.* 276: 3188–3194, 2001.
- [78113] 16443.Berra, B.; Gornati, R.; Rapelli, S.; Gatti, R.; Mancini, G. M.S.; Ciana, G.; Bembi, B.: Infantile sialic acid storage disease:biochemical studies. *Am. J. Med. Genet.* 58: 24–31, 1995.
- [78114] 16444.Cameron, P. D.; Dubowitz, V.; Besley, G. T. N.; Fensom, A. H.:Sialic acid storage disease. *Arch. Dis. Child.* 65: 314–315, 1990.
- [78115] 16445.Haataja, L.; Schleutker, J.; Laine, A.-P.; Renlund, M.; Savontaus,M.-L.; Dib, C.; Weissenbach, J.; Peltonen, L.; Aula, P.: The genetic locus for free sialic acid storage disease maps to the long arm of chromosome 6. *Am. J. Hum. Genet.* 54: 1042–1049, 1994.
- [78116] 16446.Fojo, A.; Lebo, R.; Shimizu, N.; Chin, J. E.; Roninson, I. B.;Merlino, G. T.; Gottesman, M. M.; Pastan, I.: Localization of multidrug resistance-associated DNA sequences to human chromosome 7. *Somat.Cell Molec. Genet.* 12: 415–420, 1986.
- [78117] 16447.Fojo, A. T.; Ueda, K.; Slamon, D. J.; Poplack, D. G.; Gottesman,M. M.; Pastan, I.: Expression of a multidrug-resistance gene in human tumors and tissues. *Proc. Nat.*

Acad. Sci. 84: 265–269, 1987.

- [78118] 16448.Fujii, J.; Zarain-Herzberg, A.; Willard, H. F.; Tada, M.; MacLennan, D. H.: Structure of the rabbit phospholamban gene, cloning of the human cDNA, and assignment of the gene to human chromosome 6. J. Biol. Chem. 266: 11669–11675, 1991.
- [78119] 16449.McTiernan, C. F.; Frye, C. S.; Lemster, B. H.; Kinder, E. A.; Ogletree-Hughes, M. L.; Moravec, C. S.; Feldman, A. M.: The human phospholamban gene: structure and expression. J. Molec. Cell Cardiol. 31: 679–692, 1999.
- [78120] 16450.Otsu, K.; Fujii, J.; Periasamy, M.; Difilippantonio, M.; Uppender, M.; Ward, D. C.; MacLennan, D. H.: Chromosome mapping of five human cardiac and skeletal muscle sarcoplasmic reticulum protein genes. Genomics 17:507–509, 1993.
- [78121] 16451.Barhanin, J.; Lesage, F.; Guillemare, E.; Fink, M.; Lazdunski, M.; Romey, G.: K(v)LQT1 and IsK (minK) proteins associate to form the I(Ks) cardiac potassium current. Nature 384: 78–80, 1996.
- [78122] 16452.Marx, S. O.; Kurokawa, J.; Reiken, S.; Motoike, H.; D'Armiento, J.; Marks, A. R.; Kass, R. S.: Requirement of a macromolecular signaling complex for beta adrenergic receptor modulation of the KCNQ1–KCNE1 potassium chan-

nel. Science 295: 496–499, 2002.

- [78123] 16453.Sanguinetti, M. C.; Curran, M. E.; Zou, A.; Shen, J.; Spector, P. S.; Atkinson, D. L.; Keating, M. T.: Coassembly of K(v)LQT1 and minK (IsK) proteins to form cardiac I(Ks) potassium channel. Nature 384:80–83, 1996.
- [78124] 16454.Coleman, R. A.; Smith, W. L.; Narumiya, S.: VIII. International union of pharmacology classification of prostanoid receptors: properties, distribution, and structure of the receptors and their subtypes. Pharm.Rev. 46: 205–229, 1994.
- [78125] 16455.Duncan, A. M. V.; Anderson, L. L.; Funk, C. D.; Abramovitz, M.; Adam, M.: Chromosomal localization of the human prostanoid receptor gene family. Genomics 25: 740–742, 1995.
- [78126] 16456.Nagata, A.; Suzuki, Y.; Igarashi, M.; Eguchi, N.; Toh, H.; Urade, Y.; Hayaishi, O.: Human brain prostaglandin D synthase has been evolutionarily differentiated from lipophilic–ligand carrier proteins. Proc. Nat.Acad. Sci. 88: 4020–4024, 1991.
- [78127] 16457.Onoe, H.; Ueno, R.; Fujita, I.; Nishino, H.; Oomura, Y.; Hayashi, O.: Proc. Nat. Acad. Sci. 85: 4082–4086, 1988.
- [78128] 16458.Pinzar, E.; Kanaoka, Y.; Inui, T.; Eguchi, N.; Urade, Y.; Hayaishi, O.: Prostaglandin D synthase gene is involved



in the regulation of non-rapid eye movement sleep. Proc. Nat. Acad. Sci. 97: 4903–4907, 2000.

- [78129] 16459. Tanaka, T.; Urade, Y.; Kimura, H.; Eguchi, N.; Nishikawa, A.; Hayaishi, O.: Lipocalin-type prostaglandin D synthase (beta-trace) is a newly recognized type of retinoid transporter. J. Biol. Chem. 272: 15789–15795, 1997.
- [78130] 16460. White, D. M.; Mikol, D. D.; Espinosa, R.; Weimer, B.; Le Beau, M. M.; Stefansson, K.: Structure and chromosomal localization of the human gene for a brain form of prostaglandin D-2 synthase. J. Biol. Chem. 267: 23202–23208, 1992.
- [78131] 16461. Sales, K. J.; Katz, A. A.; Davis, M.; Hinz, S.; Soeters, R. P.; Hofmeyr, M. D.; Millar, R. P.; Jabbour, H. N.: Cyclooxygenase-2 expression and prostaglandin E2 synthesis are up-regulated in carcinomas of the cervix: a possible autocrine/paracrine regulation of neoplastic cell function via EP2/EP4 receptors. J. Clin. Endocr. Metab. 86: 2243–2249, 2001.
- [78132] 16462. Smock, S. L.; Pan, L. C.; Castleberry, T. A.; Lu, B.; Mather, R. J.; Owen, T. A.: Cloning, structural characterization, and chromosomal localization of the gene encoding the human prostaglandin E2 receptor EP2 subtype. Gene

237: 393–402, 1999.

[78133] 16463.Taketo, M.; Rochelle, J. M.; Sugimoto, Y.; Namba, T.; Honda, A.;Negishi, M.; Ichikawa, A.; Narumiya, S.; Seldin, M. F.: Mapping ofthe genes encoding mouse thromboxane A<sub>2</sub> receptor and prostaglandinE receptor subtypes EP<sub>2</sub> and EP<sub>3</sub>. Genomics 19: 585–588, 1994.

[78134] 16464.Tilley, S. L.; Audoly, L. P.; Hicks, E. H.; Kim, H.–S.; Flannery,P. J.; Coffman, T. M.; Koller, B. H.: Reproductive failure and reducedblood pressure in mice lacking the EP<sub>2</sub> prostaglandin E–2 receptor. J.Clin. Invest. 103: 1539–1545, 1999.

[78135] 16465.Catella–Lawson, F.; Reilly, M. P.; Kapoor, S. C.; Cucchiara, A.J.; DeMarco, S.; Tournier, B.; Vyas, S. N.; FitzGerald, G. A.: Cyclooxygenaseinhibitors and the antiplatelet effects of aspirin. New Eng. J. Med. 345:1809–1817, 2001.

[78136] 16466.Dube, J.–N.; Drouin, J.; Aminian, M.; Plant, M. H.; Laneuville,O.: Characterization of a partial prostaglandin endoperoxide H synthase–1deficiency in a patient with a bleeding disorder. Brit. J. Haemat. 113:878–885, 2001.

[78137] 16467.Funk, C. D.; Funk, L. B.; Kennedy, M. E.; Pong, A. S.; Fitzgerald,G. A.: Human platelet/erythro leukemia cell prostaglandin G/H synthase:cDNA cloning, expression,

and gene chromosomal assignment. FASEBJ. 5:  
2304–2312, 1991.

- [78138] 16468. Gavett, S. H.; Madison, S. L.; Chulada, P. C.; Scarborough, P. E.; Qu, W.; Boyle, J. E.; Tiano, H. F.; Lee, C. A.; Langenbach, R.; Roggli, V. L.; Zeldin, D. C.: Allergic lung responses are increased in prostaglandin H synthase-deficient mice. J. Clin. Invest. 104:721–732, 1999.
- [78139] 16469. Kirschenbaum, A.; Liotta, D. R.; Yao, S.; Liu, X.-H.; Klausner, A. P.; Unger, P.; Shapiro, E.; Leav, I.; Levine, A. C.: Immunohistochemical localization of cyclooxygenase-1 and cyclooxygenase-2 in the human fetal and adult male reproductive tracts. J. Clin. Endocr. Metab. 85:3436–3441, 2000.
- [78140] 16470. Langenbach, R.; Morham, S. G.; Tiano, H. F.; Loftin, C. D.; Ghanayem, B. I.; Chulada, P. C.; Mahler, J. F.; Lee, C. A.; Goulding, E. H.; Kluckman, K. D.; Kim, H. S.; Smithies, O.: Prostaglandin synthase 1 gene disruption in mice reduces arachidonic acid-induced inflammation and indomethacin-induced gastric ulceration. Cell 83: 483–492, 1995.
- [78141] 16471. Malkowski, M. G.; Ginell, S. L.; Smith, W. L.; Garavito, R. M.: The productive conformation of arachidonic acid bound to prostaglandin synthase. Science 289:

1933–1937, 2000.

- [78142] 16472.Pareti, F. I.; Manucci, P. M.; D'Angelo, A.; Smith, J. B.; Sautebin,L.; Galli, G.: Congenital deficiency of thromboxane and prostacyclin. *Lancet* I:898–900, 1980.
- [78143] 16473.Picot, D.; Loll, P. J.; Garavito, R. M.: The x-ray crystal structureof the membrane protein prostaglandin H-2 synthase-1. *Nature* 367:243–249, 1994.
- [78144] 16474.Allen, J. M.; Bloom, S. R.: Neuropeptide Y: a putative neurotransmitter. *Neurochem.Int.* 8: 1–8, 1986.
- [78145] 16475.Niswender, K. D.; Morton, G. J.; Stearns, W. H.; Rhodes, C. J.;Myers, M. G., Jr.; Schwartz, M. W.: Key enzyme in leptin-inducedanorexia. *Nature* 413: 794–795, 2001.
- [78146] 16476.Shimomura, I.; Matsuda, M.; Hammer, R. E.; Bashmakov, Y.; Brown,M. S.; Goldstein, J. L.: Decreased IRS-2 and increased SREBP-1c leadto mixed insulin resistance and sensitivity in livers of lipodystrophicand ob/ob mice. *Molec. Cell* 6: 77–86, 2000.
- [78147] 16477.Siracusa, L. D.; Rosner, M. H.; Vigano, M. A.; Gilbert, D. J.;Staudt, L. M.; Copeland, N. G.; Jenkins, N. A.: Chromosomal locationof the octamer transcription factors, Otf-1, Otf-2, and Otf-3, definesmultiple Otf-3-related sequences dispersed in the mouse genome. *Ge-*

nomics 10:313–326, 1991.

[78148] 16478.Sturm, R. A.; Eyre, H. J.; Baker, E.; Sutherland, G. R.: The humanOTF1 locus which overlaps the CD3Z gene is located at 1q22–q23. Cytogenet.Cell Genet. 68: 231–232, 1995.

[78149] 16479.Marshall, J. B.; Diaz–Arias, A. A.; Bochna, G. S.; Vo–gele, K.A.: Achalasia due to diffuse esophageal leiomy–omatosis and inheritedas an autosomal dominant disorder. Gastroenterology 98: 1358–1365,1990.

[78150] 16480.Mattei, M. G.; d'Auriol, L.; Andre, C.; Passage, E.; Mattei, J.F.; Galibert, F.: Assignment of the human c–kit proto–oncogene tothe q11–q12 region of chromosome 4, using in situ hybridization. (Abstract) Cytogenet.Cell Genet. 46: 657, 1987.

[78151] 16481.Nagata, H.; Worobec, A. S.; Oh, C. K.; Chowdhury, B. A.; Tannenbaum,S.; Suzuki, Y.; Metcalfe, D. D.: Identifi–cation of a point mutationin the catalytic domain of the protooncogene c–kit in peripheral bloodmononuclear cells of patients who have mastocytosis with an associatedhe–matologic disorder. Proc. Nat. Acad. Sci. 92: 10560–10564, 1995.

[78152] 16482.Nishida, T.; Hirota, S.; Taniguchi, M.; Hashimoto, K.; Isozaki,K.; Nakamura, H.; Kanakura, Y.; Tanaka, T.;

Takabayashi, A.; Matsuda, H.; Kitamura, Y.: Familial gastrointestinal stromal tumours with germline mutation of the KIT gene. (Letter) *Nature Genet.* 19: 323–324, 1998.

[78153] 16483. Nocka, K.; Tan, J. C.; Chiu, E.; Chu, T. Y.; Ray, P.; Traktman, P.; Besmer, P.: Molecular bases of dominant negative and loss of function mutations at the murine *c-kit*/white spotting locus: W-37, W-v, W-41 and W. *EMBO J.* 9: 1805–1813, 1990.

[78154] 16484. Nomura, K.; Hatayama, I.; Narita, T.; Kaneko, T.; Shiraishi, M.: A novel KIT gene missense mutation in a Japanese family with piebaldism. (Letter) *J. Invest. Derm.* 111: 337–338, 1998.

[78155] 16485. Pignon, J.-M.; Giraudier, S.; Duquesnoy, P.; Jouault, H.; Imbert, M.; Vainchenker, W.; Vernant, J.-P.; Tulliez, M.: A new *c-kit* mutation in a case of aggressive mast cell disease. *Brit. J. Haemat.* 96:374–376, 1997.

[78156] 16486. Reinsch, N.; Thomsen, H.; Xu, N.; Brink, M.; Looft, C.; Kalm, E.; Brockmann, G. A.; Grupe, S.; Kuhn, C.; Schwerin, M.; Leyhe, B.; Hiendleder, S.; Erhardt, G.; Medjugorac, I.; Russ, I.; Forster, M.; Reents, R.; Averdunk, G.: A QTL for the degree of spotting in cattle shows synteny with the KIT locus on chromosome 6. *J. Hered.* 90:629–634, 1999.

[78157] 16487. Selmanowitz, V. J.; Rabinowitz, A. D.; Orentreich,

N.; Wenk, E.: Pigmentary correction of piebaldism by autografts. I. Procedures and clinical findings. J. Derm. Surg. Oncol. 3: 615–622, 1977.

[78158] 16488. Spritz, R. A.; Beighton, P.: Piebaldism with deafness: molecular evidence for an expanded syndrome. Am. J. Med. Genet. 75: 101–103, 1998.

[78159] 16489. Spritz, R. A.; Droetto, S.; Fukushima, Y.: Deletion of the KIT and PDGFRA genes in a patient with piebaldism. Am. J. Med. Genet. 44: 492–495, 1992.

[78160] 16490. Spritz, R. A.; Giebel, L. B.: Mutation of the c-kit (mast/stem cell growth factor receptor) proto-oncogene in human piebaldism. (Abstract) Am. J. Hum. Genet. 49 (suppl.): 38, 1991.

[78161] 16491. Spritz, R. A.; Giebel, L. B.; Holmes, S. A.: Dominant negative and loss of function mutations of the c-kit (mast/stem cell growth factor receptor) proto-oncogene in human piebaldism. Am. J. Hum. Genet. 50: 261–269, 1992.

[78162] 16492. Spritz, R. A.; Holmes, S. A.; Ramesar, R.; Greenberg, J.; Curtis, D.; Beighton, P.: Mutations of the KIT (mast/stem cell growth factor receptor) proto-oncogene account for a continuous range of phenotypes in human piebaldism. Am. J. Hum. Genet. 51: 1058–1065, 1992.

- [78163] 16493.Syrris, P.; Malik, N. M.; Murday, V. A.; Patton, M. A.; Carter,N. D.; Hughes, H. E.; Metcalfe, K.: Three novel mutations of theproto-oncogene KIT cause human piebaldism. (Letter) Am. J. Med. Genet. 95:79–81, 2000.
- [78164] 16494.Tan, J. C.; Nocka, K.; Ray, P.; Traktman, P.; Besmer, P.: Thedominant W42 spotting phenotype results from a missense mutation inthe c-kit receptor kinase. Science 247: 209–212, 1990.
- [78165] 16495.Taylor, M. L.; Dastych, J.; Sehgal, D.; Sundstrom, M.; Nilsson,G.; Akin, C.; Mage, R. G.; Metcalfe, D. D.: The Kit-activating mutationD816V enhances stem cell factor-dependent chemotaxis. Blood 98:1195–1199, 2001.
- [78166] 16496.Thomsen, L.; Robinson, T. L.; Lee, J. C. F.; Farraway, L. A.;Hughes, M. J. G.; Andrews, D. W.; Huizinga, J. D.: Interstitial cellsof Cajal generate a rhythmic pacemaker current. Nature Med. 4: 848–851,1998.
- [78167] 16497.Tian, Q.; Frierson, H. F., Jr.; Krystal, G. W.; Moskaluk, C. A.: Activating c-kit gene mutations in human germ cell tumors. Am.J. Path. 154: 1643–1647, 1999.
- [78168] 16498.Tsujimura, T.; Furitsu, T.; Morimoto, M.; Isozaki, K.; Nomura,S.; Matsuzawa, Y.; Kitamura, Y.; Kanakura, Y.: Ligand-independentactivation of c-kit receptor tyrosine kinase in a murine mastocytomacell line P-815 generated



by a point mutation. *Blood* 83: 2619–2626, 1994.

- [78169] 16499. Vandenbark, G. R.; deCastro, C. M.; Taylor, H.; Dew-Knight, S.; Kaufman, R. E.: Cloning and structural analysis of the human c-kit gene. *Oncogene* 7: 1259–1266, 1992.
- [78170] 16500. Worobec, A. S.; Semere, T.; Nagata, H.; Metcalfe, D. D.: Clinical correlates of the presence of the asp816val c-kit mutation in the peripheral blood mononuclear cells of patients with mastocytosis. *Cancer* 83: 2120–2129, 1998.
- [78171] 16501. Yarden, Y.; Kuang, W.-J.; Yang-Feng, T.; Coussens, L.; Munemitsu, S.; Dull, T. J.; Chen, E.; Schlessinger, J.; Francke, U.; Ullrich, A.: Human proto-oncogene c-kit: a new cell surface receptor tyrosine kinase for an unidentified ligand. *EMBO J.* 6: 3341–3351, 1987.
- [78172] 16502. Cheah, M. S. C.; Ley, T. J.; Tronick, S. R.; Robbins, K. C.: Fgr proto-oncogene mRNA induced in B lymphocytes by Epstein-Barr virus infection. *Nature* 319: 238–240, 1986.
- [78173] 16503. Dracopoli, N. C.; Stanger, B. Z.; Lager, M.; Housman, D. E.: Localization of the FGR protooncogene on the genetic linkage map of human chromosome 1p. *Genomics* 3: 124–128, 1988.
- [78174] 16504. Le Beau, M. M.; Westbrook, C. A.; Diaz, M. O.; Row-

ley, J. D.:Evidence for two distinct c-src loci on human chromosomes 1 and 20. *Nature* 312:70–71, 1984.

[78175] 16505.Lebo, R. V.; Cheung, M.-C.; Bruce, B. D.: Rapid gene mapping by dual laser chromosome sorting and spot blot DNA analysis. (Abstract) *Am.J. Hum. Genet.* 36: 101S, 1984.

[78176] 16506.Nishizawa, M.; Semba, K.; Yoshida, M. C.; Yamamoto, T.; Sasaki, M.; Toyoshima, K.: Structure, expression, and chromosomal location of the human c-fgr gene. *Molec. Cell. Biol.* 6: 511–517, 1986.

[78177] 16507.Muyan, M.; Furuhashi, M.; Sugahara, T.; Boime, I.: The carboxy-terminal region of the beta-subunits of luteinizing hormone and chorionic gonadotropin differentially influence secretion and assembly of the heterodimers. *Molec.Endocr.* 10: 1678–1687, 1996.

[78178] 16508.Ye, K.; Aghdasi, B.; Luo, H. R.; Moriarity, J. L.; Wu, F. Y.; Hong, J. J.; Hurt, K. J.; Bae, S. S.; Suh, P.-G.; Snyder, S. H.: Phospholipase C-gamma-1 is a physiological guanine nucleotide exchange factor for the nuclear GTPase PIKE. *Nature* 415: 541–544, 2002.

[78179] 16509.Tsujii, M.; Kawano, S.; Tsuji, S.; Sawaoka, H.; Hori, M.; DuBois, R. N.: Cyclooxygenase regulates angiogenesis induced by colon cancer cells. *Cell* 93: 705–716, 1998.

- [78180] 16510.Vane, J. R.; Mitchell, J. A.; Appleton, I.; Tomlinson, A.; Bishop-Bailey, D.; Croxtall, J.; Willoughby, D. A.: Inducible isoforms of cyclooxygenase and nitric-oxide synthase in inflammation. *Proc. Nat. Acad. Sci.* 91:2046–2050, 1994.
- [78181] 16511.Yokoyama, C.; Tanabe, T.: Cloning of human gene encoding prostaglandin endoperoxide synthase and primary structure of the enzyme. *Biochem. Biophys. Res. Commun.* 165: 888–894, 1989.
- [78182] 16512.Adam, M.; Boie, Y.; Rushmore, T. H.; Muller, G.; Bastien, L.; McKee, K. T.; Metters, K. M.; Abramovitz, M.: Cloning and expression of three isoforms of the human EP(3) prostanoid receptor. *FEBS Lett.* 338:170–174, 1994.
- [78183] 16513.Kotani, M.; Tanaka, I.; Ogawa, Y.; Suganami, T.; Matsumoto, T.; Muro, S.; Yamamoto, Y.; Sugawara, A.; Yoshimasa, Y.; Sagawa, N.; Narumiya, S.; Nakao, K.: Multiple signal transduction pathways through two prostaglandin E receptor EP3 subtype isoforms expressed in human uterus. *J. Clin. Endocr. Metab.* 85: 4315–4322, 2000.
- [78184] 16514.Kotani, M.; Tanaka, I.; Ogawa, Y.; Usui, T.; Mori, K.; Ichikawa, A.; Narumiya, S.; Yoshimi, T.; Nakao, K.: Molecular cloning and expression of multiple isoforms of human

prostaglandin E receptor EP(3) subtype generated by alternative messenger RNA splicing: multiple second messenger systems and tissue-specific distributions. *Molec. Pharm.* 48: 869–879, 1995.

[78185] 16515. Kotani, M.; Tanaka, I.; Ogawa, Y.; Usui, T.; Tamura, N.; Mori, K.; Narumiya, S.; Yoshimi, T.; Nakao, K.: Structural organization of the human prostaglandin EP(3) receptor subtype gene (PTGER3). *Genomics* 40: 425–434, 1997.

[78186] 16516. Schmid, A.; Thierauch, K.-H.; Schleuning, W.-D.; Dinter, H.: Splice variants of the human EP(3) receptor for prostaglandin E(2). *Europ. J. Biochem.* 228: 23–30, 1995.

[78187] 16517. Ushikubi, F.; Segi, E.; Sugimoto, Y.; Murata, T.; Matsuoka, T.; Kobayashi, T.; Hizaki, H.; Tuboi, K.; Katsuyama, M.; Ichikawa, A.; Tanaka, T.; Yoshida, N.; Narumiya, S.: Impaired febrile response in mice lacking the prostaglandin E receptor subtype EP(3). *Nature* 395: 281–284, 1998.

[78188] 16518. Wurz, H.; Geiger, W.; Kunzig, H. J.; Jabs-Lehmann, A.; Bohn, H.; Luben, G.: Radioimmunoassay of SP1 (pregnancy-specific beta-1-glycoprotein) in maternal blood and in amniotic fluid in normal and pathologic pregnancies. *J. Perinat. Med.* 2: 67–78, 1981.

[78189] 16519. Kaplan, R.; Morse, B.; Huebner, K.; Croce, C.; Howk,

R.; Ravera, M.; Ricca, G.; Jaye, M.; Schlessinger, J.: Cloning of three human tyrosine phosphatases reveals a multigene family of receptor-linked protein-tyrosine-phosphatases expressed in brain. *Proc. Nat. Acad. Sci.* 87: 7000–7004, 1990.

[78190] 16520. Banville, D.; Stocco, R.; Shen, S.-H.: Human protein tyrosine phosphatase 1C (PTPN6) gene structure: alternate promoter usage and exon skipping generate multiple transcripts. *Genomics* 27: 165–173, 1995.

[78191] 16521. Beghini, A.; Ripamonti, C. B.; Peterlongo, P.; Roversi, G.; Cairoli, R.; Morra, E.; Larizza, L.: RNA hyperediting and alternative splicing of hematopoietic cell phosphatase (PTPN6) gene in acute myeloid leukemia. *Hum. Molec. Genet.* 9: 2297–2304, 2000.

[78192] 16522. Matthews, R. J.; Bowne, D. B.; Flores, E.; Thomas, M. L.: Characterization of hematopoietic intracellular protein tyrosine phosphatases: description of a phosphatase containing an SH2 domain and another enriched in proline-, glutamic acid-, serine-, and threonine-rich sequences. *Molec. Cell. Biol.* 12: 2396–2405, 1992.

[78193] 16523. Plutzky, J.; Neel, B. G.; Rosenberg, R. D.; Eddy, R. L.; Byers, M. G.; Jani-Sait, S.; Shows, T. B.: Chromosomal localization of an SH2-containing tyrosine phosphatase

(PTPN6). Genomics 13: 869–872,1992.

- [78194] 16524.Shen, S.–H.; Bastien, L.; Posner, B. I.; Chretien, P.: A protein–tyrosinephosphatase with sequence similarity to the SH2 domain of the protein–tyrosinekinases. Nature 352: 736–739, 1991.
- [78195] 16525.Shultz, L. D.; Schweitzer, P. A.; Rajan, T. V.; Yi, T.; Ihle, J.N.; Matthews, R. J.; Thomas, M. L.; Beier, D. R.: Mutations at the murine motheaten locus are within the hematopoietic cell protein–tyrosinephosphatase (Hcph) gene. Cell 73: 1445–1454, 1993.
- [78196] 16526.Tsui, H. W.; Siminovitch, K. A.; de Souza, L.; Tsui, F. W. L.: Motheaten and viable motheaten mice have mutations in the haematopoietic cell phosphatase gene. Nature Genet. 4: 124–129, 1993.
- [78197] 16527.Yi, T.; Cleveland, J. L.; Ihle, J. N.: Identification of novel protein tyrosine phosphatases of hematopoietic cells by PCR amplification. Blood 78:2222–2228, 1991.
- [78198] 16528.Yi, T.; Cleveland, J. L.; Ihle, J. N.: Protein tyrosine phosphatase containing SH2 domains: characterization, preferential expression in hematopoietic cells, and localization to human chromosome 12p12–p13. Molec.Cell. Biol. 12: 836–846, 1992.
- [78199] 16529.Yi, T.; Gilbert, D. J.; Jenkins, N. A.; Copeland, N. G.;

Ihle, J. N.: Assignment of a novel protein tyrosine phosphatase gene (Hcph) to mouse chromosome 6. *Genomics* 14: 793–795, 1992.

[78200] 16530. Jirik, F. R.; Anderson, L. L.; Duncan, A. M. V.: The human protein-tyrosine phosphatase PTP-alpha/LRP gene (PTPA) is assigned to chromosome 20p13. *Cytogenet. Cell Genet.* 60: 117–118, 1992.

[78201] 16531. Jirik, F. R.; Janzen, N. M.; Melhado, I. G.; Harder, K. W.: Cloning and chromosomal assignment of a widely expressed human receptor-like protein-tyrosine phosphatase. *FEBS Lett.* 273: 239–242, 1990.

[78202] 16532. Matthews, R. J.; Cahir, E. D.; Thomas, M. L.: Identification of an additional member of the protein-tyrosine-phosphatase family: evidence for alternative splicing in the tyrosine phosphatase domain. *Proc. Nat. Acad. Sci.* 87: 4444–4448, 1990.

[78203] 16533. Rao, V. V. N. G.; Löffler, C.; Sap, J.; Schlessinger, J.; Hansmann, I.: The gene for receptor-linked protein-tyrosine-phosphatase (PTPA) is assigned to human chromosome 20p12-pter by in situ hybridization (ISH and FISH). *Genomics* 13: 906–907, 1992.

[78204] 16534. Schnittger, S.; Rao, V. V. N. G.; Deutsch, U.; Gruss, P.; Balling, R.; Hansmann, I.: PAX1, a member of the paired

box-containing class of developmental control genes, is mapped to human chromosome 20p11.2 by *in situ* hybridization (ISH and FISH). *Genomics* 14: 740–744, 1992.

- [78205] 16535. Ahmad, F.; Azevedo, J. L., Jr.; Cortright, R.; Dohm, G. L.; Goldstein, B. J.: Alternations in skeletal muscle protein-tyrosine phosphatase activity and expression in insulin-resistant human obesity and diabetes. *J. Clin. Invest.* 100: 449–458, 1997.
- [78206] 16536. Brown-Shimer, S.; Johnson, K. A.; Lawrence, J. B.; Johnson, C.; Bruskin, A.; Green, N. R.; Hill, D. E.: Molecular cloning and chromosomal mapping of the human gene encoding protein phosphotyrosyl phosphatase 1B. *Proc. Nat. Acad. Sci.* 87: 5148–5152, 1990.
- [78207] 16537. Charbonneau, H.; Tonks, N. K.; Kumar, S.; Diltz, C. D.; Harrylock, M.; Cool, D. E.; Krebs, E. G.; Fischer, E. H.; Walsh, K. A.: Human placenta protein-tyrosine-phosphatase: amino acid sequence and relationship to a family of receptor-like proteins. *Proc. Nat. Acad. Sci.* 86: 5252–5256, 1989.
- [78208] 16538. Chernoff, J.; Schievella, A. R.; Jost, C. A.; Erikson, R. L.; Neel, B. G.: Cloning of a cDNA for a major human protein-tyrosine-phosphatase. *Proc. Nat. Acad. Sci.* 87: 2735–2739, 1990.



- [78209] 16539. Matilla, A.; Gorbea, C.; Einum, D. D.; Townsend, J.; Michalik, A.; van Broeckhoven, C.; Jensen, C. C.; Murphy, K. J.; Ptacek, L. J.; Fu, Y.-H.: Association of ataxin-7 with the proteasome subunit S4 of the 19S regulatory complex. *Hum. Molec. Genet.* 10: 2821-2831, 2001.
- [78210] 16540. Cazzaniga, G.; Tosi, S.; Aloisi, A.; Giudici, G.; Dan-  
iotti, M.; Pioltelli, P.; Kearney, L.; Biondi, A.: The tyrosine  
kinase Abl-related gene ARG is fused to ETV6 in an AML-  
M4Eo patient with a t(1;12)(q25;p13): molecular cloning of  
both reciprocal transcripts. *Blood* 94: 4370-4373, 1999.
- [78211] 16541. Iijima, Y.; Ito, T.; Oikawa, T.; Eguchi, M.; Eguchi-  
Ishimae, M.; Kamada, N.; Kishi, K.; Asano, S.; Sakaki, Y.;  
Sato, Y.: A new ETV6/TEL partner gene, ARG (ABL-related  
gene or ABL2), identified in an AML-M3 cell line with a  
t(1;12)(q25;p13) translocation. *Blood* 95:  
2126-2131, 2000.
- [78212] 16542. Joazeiro, C. A. P.; Wing, S. S.; Huang, H.; Levenson,  
J. D.; Hunter, T.; Liu, Y.-C.: The tyrosine kinase negative  
regulator c-Cbl as a RING-type, E2-dependent ubiquitin-  
protein ligase. *Science* 286: 309-312, 1999.
- [78213] 16543. Langdon, W. Y.; Hartley, J. W.; Klinken, S. P.;  
Ruscetti, S. K.; Morse, H. C., III: v-cbl, an oncogene from a  
dual-recombinant murine retrovirus that induces early B-

lineage lymphomas. Proc. Nat. Acad.Sci. 86: 1168–1172, 1989.

[78214] 16544.Mastick, C. C.; Brady, M. J.; Saltiel, A. R.: Insulin stimulates the tyrosine phosphorylation of caveolin. J. Cell Biol. 129: 1523–1531, 1995.

[78215] 16545.Petrelli, A.; Gilestro, G. F.; Lanzardo, S.; Comoglio, P. M.; Migone, N.; Giordano, S.: The endophilin–CIN85–Cbl complex mediates ligand–dependent downregulation of c–Met. Nature 416: 187–190, 2002.

[78216] 16546.Savage, P. D.; Shapiro, M.; Langdon, W. Y.; Geurts van Kessel, A. D.; Seunavaz, H. N.; Akao, Y.; Croce, C.; Morse, H. C., III; Kersey, J. H.: Relationship of the human protooncogene CBL2 on 11q23 to the t(4;11), t(11;22), and t(11;14) breakpoints. Cytogenet. Cell Genet. 56:112–115, 1991.

[78217] 16547.Soubeyran, P.; Kowanetz, K.; Szymkiewicz, I.; Langdon, W. Y.; Dikic, I.: Cbl–CIN85–endophilin complex mediates ligand–induced downregulation of EGF receptors. Nature 416: 183–187, 2002.

[78218] 16548.Thien, C. B. F.; Walker, F.; Langdon, W. Y.: RING finger mutations that abolish c–Cbl–directed polyubiquitination and downregulation of the EGF receptor are insufficient for cell transformation. Molec.Cell 7: 355–365,

2001.

- [78219] 16549.Wei, S.; Rocchi, M.; Archidiacono, N.; Sacchi, N.; Romeo, G.;Gatti, R. A.: Physical mapping of the human chromosome 11q23 regioncontaining the ataxia-telangiectasia locus. *Cancer Genet. Cytogenet.* 46:1-8, 1990.
- [78220] 16550.Yoon, C. H.; Lee, J.; Jongeward, G. D.; Sternberg, P. W.: Similarityof sli-1, a regulator of vulval development in *C. elegans*, to themammalian proto-oncogene c-cbl. *Science* 269: 1102-1105, 1995.
- [78221] 16551.Cannizzaro, L. A.; Madaule, P.; Hecht, F.; Axel, R.; Croce, C.M.; Huebner, K.: Chromosome localization of human ARH genes, a ras-relatedgene family. *Genomics* 6: 197-203, 1990.
- [78222] 16552.Chardin, P.; Madaule, P.; Tavitian, A.: Coding sequence of humanrho cDNAs clone 6 and clone 9. *Nucleic Acid Res.* 16: 2717 only,1988.
- [78223] 16553.Madaule, P.; Axel, R.: A novel ras-related gene family. *Cell* 41:31-40, 1985.
- [78224] 16554.Maekawa, M.; Ishizaki, T.; Boku, S.; Watanabe, N.; Fujita, A.;Iwamatsu, A.; Obinata, T.; Ohashi, K.; Mizuno, K.; Narumiya, S.:Signaling from Rho to the actin cytoskeleton through protein kinasesROCK and LIM-kinase.

Science 285: 895–898, 1999.

- [78225] 16555.Ridley, A. J.; Hall, A.: The small GTP-binding protein rho regulates the assembly of focal adhesions and actin stress fibers in response to growth factors. Cell 70: 389–399, 1992.
- [78226] 16556.Fagan, K. P.; Oliveira, L.; Pittler, S. J.: Sequence of rho small GTP-binding protein cDNAs from human retina and identification of novel 5-prime end cloning artifacts. Exp. Eye Res. 59: 235–237, 1994.
- [78227] 16557.Morris, S. W.; Valentine, M. B.; Kirstein, M. N.; Huebner, K.: Reassignment of the human ARH9 RAS-related gene to chromosome 1p13–p21. Genomics 15: 677–679, 1993.
- [78228] 16558.Maesaki, R.; Ihara, K.; Shimizu, T.; Kuroda, S.; Kaibuchi, K.; Hakoshima, T.: The structural basis of Rho effector recognition revealed by the crystal structure of human RhoA complexed with the effector domain of PKN/PRK1. Molec. Cell 4: 793–803, 1999.
- [78229] 16559.Nakamura, M.; Nagano, T.; Chikama, T.; Nishida, T.: Role of the small GTP-binding protein Rho in epithelial cell migration in the rabbit cornea. Invest. Ophthalmol. Vis. Sci. 42: 941–947, 2001.
- [78230] 16560.Rao, P. V.; Deng, P.-F.; Kumar, J.; Epstein, D. L.:

Modulation of aqueous humor outflow facility by the Rho kinase-specific inhibitor Y-27632. *Invest. Ophthalmol. Vis. Sci.* 42: 1029–1037, 2001.

[78231] 16561. Akiyama, T.; Takasawa, S.; Nata, K.; Kobayashi, S.; Abe, M.; Shervani, N. J.; Ikeda, T.; Nakagawa, K.; Unno, M.; Matsuno, S.; Okamoto, H.: Activation of Reg gene, a gene for insulin-producing beta-cell regeneration: poly(ADP-ribose) polymerase binds Reg promoter and regulates the transcription by autopoly(ADP-ribosylation). *Proc. Nat. Acad. Sci.* 98: 48–53, 2001.

[78232] 16562. Bartoli, C.; Gharib, B.; Giorgi, D.; Sansonetti, A.; Dagorn, J.-C.; Berge-Lefranc, J.-L.: A gene homologous to the reg gene is expressed in human pancreas. *FEBS Lett.* 327: 289–293, 1993.

[78233] 16563. Gharib, B.; Fox, M. F.; Bartoli, C.; Giorgi, D.; Sansonetti, A.; Swallow, D. M.; Dagorn, J. C.; Berge-Lefranc, J. L.: Human regeneration protein/lithostathine genes map to chromosome 2p12. *Ann. Hum. Genet.* 57: 9–16, 1993.

[78234] 16564. Giorgi, D.; Bernard, J.-P.; Rouquier, S.; Iovanna, J.; Sarles, H.; Dagorn, J.-C.: Secretory pancreatic stone protein messenger RNA: nucleotide sequence and expression in chronic calcifying pancreatitis. *J. Clin. Invest.* 84: 100–106, 1989.

- [78235] 16565.Miyashita, H.; Nakagawara, K.; Mori, M.; Narushima, Y.; Noguchi,N.; Moriizumi, S.; Takasawa, S.; Yonekura, H.; Takeuchi, T.; Okamoto,H.: Human REG family genes are tandemly ordered in a 95-kilobaseregion of chromosome 2p12. FEBS Lett. 377: 429–433, 1995.
- [78236] 16566.Perfetti, R.; Hawkins, A. L.; Griffin, C. A.; Egan, J. M.; Zenilman,M. E.; Shuldiner, A. R.: Assignment of the human pancreatic regenerating(REG) gene to chromosome 2p12. Genomics 20: 305–307, 1994.
- [78237] 16567.Sarles, H.; Dagorn, J. C.; Giorgi, D.; Bernard, J. P.: Renamingpancreatic stone protein as 'lithostathine'. (Letter) Gastroenterology 99:900–901, 1990.
- [78238] 16568.Stewart, T. A.: The human REG gene encodes pancreatic stone protein.(Letter) Biochem. J. 260: 622–623, 1989.
- [78239] 16569.Terazono, K.; Yamamoto, H.; Takasawa, S.; Shiga, K.; Yonemura,Y.; Tochino, Y.; Okamoto, H.: A novel gene activated in regeneratingislets. J. Biol. Chem. 263: 2111–2114, 1988.
- [78240] 16570.Unno, M.; Yonekura, H.; Nakagawara, K.; Watanabe, T.; Miyashita,H.; Moriizumi, S.; Okamoto, H.: Structure, chromosomal localization,and expression of mouse reg genes, reg I and reg II: a novel typeof reg gene, reg II, ex–

ists in the mouse genome. J. Biol. Chem.

268:15974–15982, 1993.

[78241] 16571. Verdier, J. M.; Dussol, B.; Casanova, P.; Daudon, M.; Dupuy, P.; Berthezene, P.; Boistelle, R.; Berland, Y.; Dagorn, J. C.: Evidence that human kidney produces a protein similar to lithostathine, the pancreatic inhibitor of  $\text{CaCO}_3$  crystal growth. *Europ. J. Clin. Invest.* 22:469–474, 1992.

[78242] 16572. Moriizumi, S.; Watanabe, T.; Unno, M.; Nakagawara, K.; Suzuki, Y.; Miyashita, H.; Yonekura, H.; Okamoto, H.: Isolation, structural determination and expression of a novel reg gene, human regI-beta. *Biochim. Biophys. Acta* 1217: 199–202, 1994.

[78243] 16573. Boel, E.; Schwartz, T. W.; Norris, K. E.; Fiil, N. P.: A cDNA encoding a small common precursor for human pancreatic polypeptide and pancreatic icosapeptide. *EMBO J.* 3: 909–912, 1984.

[78244] 16574. Chandrasekharappa, S. C.; Friedman, L.; King, S. E.; Lee, Y.-H.; Welsch, P.; Bowcock, A. M.; Weber, B. L.; King, M.-C.; Collins, F. S.: The gene for pancreatic polypeptide (PPY) and the anonymous marker D17S78 are within 45 kb of each other on chromosome 17q21. *Genomics* 21:458–460, 1994.

[78245] 16575. Hort, Y.; Baker, E.; Sutherland, G. R.; Shine, J.; Her-

zog, H.:Gene duplication of the human peptide YY gene (PYY) generated thepancreatic polypeptide gene (PPY) on chromosome 17q21.1. *Genomics* 26:77–83, 1995.

[78246] 16576.Kimmel, J. R.; Hayden, L. J.; Pollock, H. G.: Isolation and characterizationof a new pancreatic polypeptide hormone. *J. Biol. Chem.* 250: 9369–9376,1975.

[78247] 16577.Larsson, L.–I.; Sundler, F.; Hakanson, R.: Pancreatic polypeptide--apostulated new hormone: identification of its cellular storage siteby light and electron microscopic immunocytochemistry. *Diabetologia* 12:211–216, 1976.

[78248] 16578.Schwartz, T. W.: Pancreatic polypeptide: a hormone under vagalcontrol. *Gastroenterology* 85: 1411–1425, 1983.

[78249] 16579.Schwartz, T. W.; Tager, H. S.: Isolation and biogenesis of a newpeptide from pancreatic islets. *Nature* 294: 589–591, 1981.

[78250] 16580.Takeuchi, T.; Gumucio, D.; Eddy, R.; Meisler, M.; Minth, C.; Dixon,J.; Yamada, T.; Shows, T.: Assignment of the related pancreatic polypeptide(PNP or PPY) and neuropeptide Y (NPY) genes to regions on human chromosomes17 and 7. (Abstract) *Cytogenet. Cell Genet.* 40: 759 only, 1985.

[78251] 16581.Zipf, W. B.; O'Dorisio, T. M.; Cataland, S.; Sotos, J.:



Blunted pancreatic polypeptide responses in children with obesity of Prader–Willi syndrome. *J. Clin. Endocr. Metab.* 52: 1264–1265, 1981.

[78252] 16582. Chen, K.; Yang, W.; Grimsby, J.; Shih, J. C.: The human 5-HT<sub>2</sub> receptor is encoded by a multiple intron–exon gene. *Brain Res. Molec. Brain Res.* 14: 20–26, 1992.

[78253] 16583. Holmes, C.; Arranz, M. J.; Powell, J. F.; Collier, D. A.; Lovestone, S.: 5-HT<sub>2A</sub> and 5-HT<sub>2C</sub> receptor polymorphisms and psychopathology in late onset Alzheimer's disease. *Hum. Molec. Genet.* 7: 1507–1509, 1998.

[78254] 16584. Hsieh, C.–L.; Bowcock, A. M.; Farrer, L. A.; Hebert, J. M.; Huang, K. N.; Cavalli–Sforza, L. L.; Julius, D.; Francke, U.: The serotonin receptor subtype 2 locus HTR2 is on human chromosome 13 near genes for esterase D and retinoblastoma–1 and on mouse chromosome 14. *Somat. Cell Molec. Genet.* 16: 567–574, 1990.

[78255] 16585. Kato, M. V.; Shimizu, T.; Nagayoshi, M.; Kaneko, A.; Sasaki, M. S.; Ikawa, Y.: Genomic imprinting of the human serotonin–receptor (HTR2) gene involved in development of retinoblastoma. *Am. J. Hum. Genet.* 59: 1084–1090, 1996.

[78256] 16586. Liu, J.; Chen, Y.; Kozak, C. A.; Yu, L.: The 5-HT(2) serotonin receptor gene Htr–2 is tightly linked to Es–10 on

mouse chromosome14. Genomics 11: 231–234, 1991.

[78257] 16587.Peroutka, S. J.: 5-Hydroxytryptamine receptor subtypes. Ann.Rev. Neurosci. 11: 45–60, 1988.

[78258] 16588.Sparkes, R. S.; Lan, N.; Klisak, I.; Mohandas, T.; Diep, A.; Kojis,T.; Heinzmann, C.; Shih, J. C.: Assignment of a serotonin 5HT–2 receptorgene (HTR2) to human chromosome 13q14–q21 and mouse chromosome 14. Genomics 9:461–465, 1991.

[78259] 16589.Williams, J.; McGuffin, P.; Nothen, M.; Owen, M. J.; EMASS CollaborativeGroup: Meta-analysis of association between the 5-HT (2A) receptorT102C polymorphism and schizophrenia. (Letter) Lancet 349: 1221only, 1997.

[78260] 16590.Williams, J.; Spurlock, G.; McGuffin, P.; Mallet, J.; Nothen, M.M.; Gill, M.; Aschauer, H.; Nylander, P. O.; Macciardi, F.; Owen,M. J.: Association between schizophrenia and T102C polymorphism ofthe 5-hydroxytryptophan type 2a–receptor gene. European MulticentreAssociation Study of Schizophrenia (EMASS) Group. Lancet 347: 1294–1296,1996.

[78261] 16591.Philipson, L. H.; Hice, R. E.; Schaefer, K.; LaMendola, J.; Bell,G. I.; Neldon, D. J.; Steiner, D. F.: Sequence and functional expressionin Xenopus oocytes of a human insulinoma and islet potassium channel. Proc.Nat. Acad.

Sci. 88: 53–57, 1991.

- [78262] 16592. Ghanshani, S.; Pak, M.; McPherson, J. D.; Strong, M.; Dethlefs, B.; Wasmuth, J. J.; Salkoff, L.; Gutman, G. A.; Chandy, K. G.: Genomic organization, nucleotide sequence, and cellular distribution of a Shaw-related potassium channel gene, Kv3.3, and mapping of Kv3.3 and Kv3.4 to human chromosomes 19 and 1. *Genomics* 12: 190–196, 1992.
- [78263] 16593. McPherson, J. D.; Wasmuth, J. J.; Chandy, K. G.; Swanson, R.; Dethlefs, B.; Chandy, G.; Wymore, R.; Ghanshani, S.: Chromosomal localization of 7 potassium channel genes. (Abstract) *Cytogenet. Cell Genet.* 58:1979 only, 1991.
- [78264] 16594. Gessler, M.; Grupe, A.; Grzeschik, K.-H.; Pongs, O.: The potassium channel gene HK1 maps to human chromosome 11p14.1, close to the FSHB gene. *Hum. Genet.* 90: 319–321, 1992.
- [78265] 16595. Grandy, D.; Mathew, M. K.; Ramaswami, M.; Tanouye, M.; Sheffield, V.; Jones, C. A.; Al-Dhalimi, M.; Zhang, Y.; Saez, C.; Litt, M.: A human voltage-gated potassium channel gene, HuKII, maps to chromosome 11p14–p13. (Abstract) *Am. J. Hum. Genet.* 51 (suppl.): A396 only, 1992.

- [78266] 16596.Philipson, L. H.; Eddy, R. L.; Shows, T. B.; Bell, G. I.: Assignment of human potassium channel gene KCNA4 (Kv1.4, PCN2) to chromosome 11q13.4–q14.1. *Genomics* 15: 463–464, 1993.
- [78267] 16597.Philipson, L. H.; Schaefer, K.; LaMendola, J.; Bell, G. I.; Steiner, D. F.: Sequence of a human fetal skeletal muscle potassium channel cDNA related to RCK4. *Nucleic Acids Res.* 18: 7160 only, 1990.
- [78268] 16598.Stuehmer, W.; Ruppersberg, J. P.; Schroter, K. H.; Sakmann, B.; Stocker, M.; Giese, K. P.; Perschke, A.; Bauermann, A.; Pongs, O.: Molecular basis of functional diversity of voltage-gated potassium channels in mammalian brain. *EMBO J.* 8: 3235–3244, 1989.
- [78269] 16599.Tamkun, M. M.; Knoth, K. M.; Walbridge, J. A.; Kroemer, H.; Roden, D. M.; Glover, D. M.: Molecular cloning and characterization of two voltage-gated K<sup>+</sup> channel cDNAs from human ventricle. *FASEB J.* 5:331–337, 1991.
- [78270] 16600.Phromchotikul, T.; Browne, D. L.; Curran, M. E.; Keating, M. T.; Litt, M.: Dinucleotide repeat polymorphism at the KCNA5 locus. *Hum.Molec. Genet.* 2: 1512 only, 1993.
- [78271] 16601.Bardien-Kruger, S.; Wulff, H.; Arieff, Z.; Brink, P.; Chandy, K.G.; Corfield, V.: Characterisation of the human

voltage-gated potassium channel gene, KCNA7, a candidate gene for inherited cardiac disorders, and its exclusion as cause of progressive familial heart block I (PFHBI). *Eur. J. Hum. Genet.* 10: 36–43, 2002.

[78272] 16602. Day, J. R.; Albers, J. J.; Lofton-Day, C. E.; Gilbert, T. L.; Ching, A. F. T.; Grant, F. J.; O'Hara, P. J.; Marcovina, S. M.; Adolphson, J. L.: Complete cDNA encoding human phospholipid transfer protein from human endothelial cells. *J. Biol. Chem.* 269: 9388–9391, 1994.

[78273] 16603. Jiang, X.; Bruce, C.; Mar, J.; Lin, M.; Ji, Y.; Francone, O. L.; Tall, A. R.: Targeted mutation of plasma phospholipid transfer protein gene markedly reduces high-density lipoprotein levels. *J. Clin. Invest.* 103: 907–914, 1999.

[78274] 16604. LeBoeuf, R. C.; Caldwell, M.; Tu, A.-Y.; Albers, J. J.: Phospholipid transfer protein maps to distal mouse chromosome 2. *Genomics* 34: 259–260, 1996.

[78275] 16605. Tu, A.-Y.; Wolfbauer, G.; Albers, J. J.: Functional characterization of the promoter region of the human phospholipid transfer protein gene. *Biochem. Biophys. Res. Commun.* 217: 705–711, 1995.

[78276] 16606. Whitmore, T. E.; Day, J. R.; Albers, J. J.: Localization of the human phospholipid transfer protein gene to chromosome 20q12–q13.1. *Genomics* 28: 599–600, 1995.

- [78277] 16607.Comings, D. E.: Evidence for ancient tetraploidy and conservation of linkage groups in mammalian chromosomes. *Nature* 238: 455–457, 1972.
- [78278] 16608.D'Ancona, G. G.; Chern, C. J.; Benn, P.; Croce, C. M.: Assignment of the human gene for enolase 1 to region pter–p36 of chromosome 1. *Cytogenet. Cell Genet.* 18: 327–332, 1977.
- [78279] 16609.D'Ancona, G. G.; Croce, C. M.: Assignment of the gene for enolase to mouse chromosome 4 using somatic cell hybrids. *Cytogenet. Cell Genet.* 19: 1–6, 1977.
- [78280] 16610.Feo, S.; Oliva, D.; Arico, B.; Barba, G.; Cali, L.; Giallongo, A.: The human genome contains a single processed pseudogene for alpha enolase located on chromosome 1. *DNA Sequence* 1: 79–83, 1990.
- [78281] 16611.Giallongo, A.; Feo, S.; Moore, R.; Croce, C. M.; Showe, L. C.: Molecular cloning and nucleotide sequence of a full-length cDNA for human alpha enolase. *Proc. Nat. Acad. Sci.* 83: 6741–6745, 1986.
- [78282] 16612.Giblett, E. R.; Chen, S.–H.; Anderson, J. E.; Lewis, M.: A family study suggesting genetic linkage of phosphopyruvate hydratase (enolase) to the Rh blood group system. *Cytogenet. Cell Genet.* 13: 91–92, 1974.
- [78283] 16613.Lachant, N. A.; Jennings, M. A.; Tanaka, K. R.: Par–

tial erythrocyte enolase deficiency: a hereditary disorder with variable clinical expression. (Abstract) Blood 68: 55a only, 1986.

[78284] 16614. Lachant, N. A.; Tanaka, K. R.: Enolase kinetic properties in partial erythrocyte enolase deficiency. (Abstract) Clin. Res. 35:426A only, 1987.

[78285] 16615. Lalley, P. A.; Francke, U.; Minna, J. D.: Homologous genes for enolase, phosphogluconate dehydrogenase, phosphoglucomutase, and adenylate kinase are syntenic on mouse chromosome 4 and human chromosome 1p. Proc. Nat. Acad. Sci. 75: 2382–2386, 1978.

[78286] 16616. Ribaud, M. R.; Di Leonardo, A.; Rubino, P.; Giallongo, A.; Feo, S.: Assignment of enolase processed pseudogene (ENO1P) to human chromosome 1 bands 1q41–q42. Cytogenet. Cell Genet. 74: 201–202, 1996.

[78287] 16617. Van Cong, N.; Weil, D.; Rebourcet, R.; Frezal, J.: Localisation des enolases 1 et 2 respectivement sur les chromosomes 1 et 12 par l'analyse des hybrides homme-souris. Ann. Genet. 20: 153–157, 1977.

[78288] 16618. Wistow, G. J.; Lietman, T.; Williams, L. A.; Stapel, S. O.; de Jong, W. W.; Horwitz, J.; Piatigorsky, J.: Tau-crystallin/alpha-enolase: one gene encodes both an enzyme and a lens structural protein. J. Cell Biol. 107:

2729–2736, 1988.

- [78289] 16619. Barton, J. W.; Hart, I. M.; Patterson, D.: Mapping of a locus correcting lack of phosphoribosylaminoimidazole carboxylase activity in Chinese hamster ovary cell Ade(–)D mutants to human chromosome 4. *Genomics* 9: 314–321, 1991.
- [78290] 16620. Brayton, K. A.; Chen, Z.; Zhou, G.; Nagy, P. L.; Gavalas, A.; Trent, J. M.; Deaven, L. L.; Dixon, J. E.; Zalkin, H.: Two genes for de novo purine nucleotide synthesis on human chromosome 4 are closely linked and divergently transcribed. *J. Biol. Chem.* 269: 5313–5321, 1994.
- [78291] 16621. Torban, E.; Pelletier, J.; Goodyer, P.: F329L polymorphism in the human PAX8 gene. *Am. J. Med. Genet.* 72: 186–187, 1997.
- [78292] 16622. Gartner, J.; Kearns, W.; Pearson, P.; Valle, D.: Characterization and localization of the human 70-kD peroxisomal membrane protein (PMP70) gene. (Abstract) *Am. J. Hum. Genet.* 51 (suppl.): A168, 1992.
- [78293] 16623. Gartner, J.; Kearns, W.; Rosenberg, C.; Pearson, P.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Valle, D.: Localization of the 70-kDa peroxisomal membrane protein to human 1p21–p22 and mouse
- [78294] 16624. *Genomics* 15: 412–414, 1993.3. Gartner, J.; Moser,



H.; Valle, D.: Mutations in the 70K peroxisomal membrane protein gene in Zellweger syndrome. *Nature Genet.* 1: 16–23, 1992.

[78295] 16625. Vilain, C.; Rydlewski, C.; Duprez, L.; Heinrichs, C.; Abramowicz, M.; Malvaux, P.; Renneboog, B.; Parma, J.; Costagliola, S.; Vassart, G.: Autosomal dominant transmission of congenital thyroid hypoplasia due to loss-of-function mutation of PAX8. *J. Clin. Endocr. Metab.* 86:234–238, 2001.

[78296] 16626. Walther, C.; Guenet, J.-L.; Simon, D.; Deutsch, U.; Jostes, B.; Goulding, M.; Plachov, D.; Balling, R.; Gruss, P.: Pax: a murine multigene family of paired box-containing genes. *Genomics* 11: 424–434, 1991.

[78297] 16627. Das, P.; Stockton, D. W.; Bauer, C.; Shaffer, L. G.; D'Souza, R. N.; Wright, J. T.; Patel, P. I.: Haploinsufficiency of PAX9 is associated with autosomal dominant hypodontia. *Hum. Genet.* 110: 371–376, 2002.

[78298] 16628. Nieminen, P.; Arte, S.; Tanner, D.; Paulin, L.; Alaluusua, S.; Thesleff, I.; Pirinen, S.: Identification of a nonsense mutation in the PAX9 gene in molar oligodontia. *Europ. J. Hum. Genet.* 9:743–746, 2001.

[78299] 16629. Peters, H.; Neubuser, A.; Kratochwil, K.; Balling, R.: Pax9-deficient mice lack pharyngeal pouch derivatives and

teeth and exhibit craniofacial and limb abnormalities.

Genes Dev. 12: 2735–2747, 1998.

- [78300] 16630. Santagati, F.; Gerber, J.-K.; Blusch, J. H.; Kokubu, C.; Peters, H.; Adamski, J.; Werner, T.; Balling, R.; Imai, K.: Comparative analysis of the genomic organization of Pax9 and its conserved physical association with Nkx2-9 in the human, mouse, and pufferfish genomes. Mammalian Genome 12: 232–237, 2001.
- [78301] 16631. Stockton, D. W.; Das, P.; Goldenberg, M.; D'Souza, R. N.; Patel, P. I.: Mutation of PAX9 is associated with oligodontia. (Letter) Nature Genet. 24: 18–19, 2000.
- [78302] 16632. Wallin, J.; Mizutani, Y.; Imai, K.; Miyashita, N.; Moriwaki, K.; Taniguchi, M.; Koseki, H.; Balling, R.: A new Pax gene, Pax-9, maps to mouse chromosome 12. Mammalian Genome 4: 354–358, 1993.
- [78303] 16633. Grueneberg, D. A.; Natesan, S.; Alexandre, C.; Gilman, M. Z.: Human and Drosophila homeodomain proteins that enhance the DNA-binding activity of serum response factor. Science 257: 1089–1095, 1992.
- [78304] 16634. Kern, M. J.; Argao, E. A.; Birkenmeier, E. H.; Rowe, L. B.; Potter, S. S.: Genomic organization and chromosome localization of the murine homeobox gene Pmx. Genomics 19: 334–340, 1994.

- [78305] 16635.Nakamura, T.; Yamazaki, Y.; Hatano, Y.; Miura, I.: NUP98 is fused to PMX1 homeobox gene in human acute myelogenous leukemia with chromosomal translocation t(1;11)(q23;p15). *Blood* 94: 741–747, 1999.
- [78306] 16636.Norris, R. A.; Scott, K. K.; Moore, C. S.; Stetten, G.; Brown, C. R.; Jabs, E. W.; Wulfsberg, E. A.; Yu, J.; Kern, M. J.: Human PRRX1 and PRRX2 genes: cloning, expression, genomic localization, and exclusion as disease genes for Nager syndrome. *Mammalian Genome* 11:1000–1005, 2000.
- [78307] 16637.Riazuddin, S.; Castelein, C. M.; Ahmed, Z. M.; Lalwani, A. K.; Mastroianni, M. A.; Naz, S.; Smith, T. N.; Liburd, N. A.; Friedman, T. B.; Griffith, A. J.; Riazuddin, S.; Wilcox, E. R.: Dominant modifier DFNM1 suppresses recessive deafness DFNB26. *Nature Genet.* 26: 431–434, 2000.
- [78308] 16638.Hebrok, M.; Kim, S. K.; St Jacques, B.; McMahon, A. P.; Melton, D. A.: Regulation of pancreas development by hedgehog signaling. *Development* 127:4905–4913, 2000.
- [78309] 16639.Reidenberg, M. M.; Levy, M.; Drayer, D. E.; Zylber-Katz, E.; Robbins, W. C.: Acetylator phenotype in idiopathic systemic lupus erythematosus. *Arthritis Rheum.* 23: 569–573, 1980.
- [78310] 16640.Stohl, W.; Crow, M. K.; Kunkel, H. G.: Systemic lu-

pus erythematosus with deficiency of the T4 epitope on T helper/inducer cells. *New Eng. J. Med.* 312: 1671–1678, 1985.

[78311] 16641. Theofilopoulos, A. N.; Dixon, F. J.: Murine models of systemic lupus erythematosus. *Adv. Immun.* 37: 269–390, 1985.

[78312] 16642. Friedl, W.; Uhlhaas, S.; Schulmann, K.; Stolte, M.; Loff, S.; Back, W.; Mangold, E.; Stern, M.; Knaebel, H. P.; Sutter, C.; Weber, R. G.; Pistorius, S.; Burger, B.; Propping, P.: Juvenile polyposis: massive gastric polyposis is more common in MADH4 mutation carriers than in BMPR1A mutation carriers. *Hum. Genet.* 111: 108–111, 2002.

[78313] 16643. Houlston, R.; Bevan, S.; Williams, A.; Young, J.; Dunlop, M.; Rozen, P.; Eng, C.; Markie, D.; Woodford-Richens, K.; Rodriguez-Bigas, M. A.; Leggett, B.; Neale, K.; Phillips, R.; Sheridan, E.; Hodgson, S.; Iwama, T.; Eccles, D.; Bodmer, W.; Tomlinson, I.: Mutations in DPC4 (SMAD4) cause juvenile polyposis syndrome, but only account for a minority of cases. *Hum. Molec. Genet.* 7: 1907–1912, 1998.

[78314] 16644. Howe, J. R.; Ringold, J. C.; Summers, R. W.; Mitros, F. A.; Nishimura, D. Y.; Stone, E. M.: A gene for familial juvenile polyposis maps to chromosome 18q21.1. *Am. J.*

Hum. Genet. 62: 1129–1136, 1998.

[78315] 16645. Howe, J. R.; Roth, S.; Ringold, J. C.; Summers, R. W.; Jarvinen, H. J.; Sistonen, P.; Tomlinson, I. P. M.; Houlston, R. S.; Bevan, S.; Mitros, F. A.; Stone, E. M.; Aaltonen, L. A.: Mutations in the SMAD4/DPC4 gene in juvenile polyposis. Science 280: 1086–1088, 1998.

[78316] 16646. Leslie, K. K.; Watanabe, S.; Lei, K.-J.; Chou, D. Y.; Plouzek, C. A.; Deng, H.-C.; Torres, J.; Chou, J. Y.: Linkage of two human pregnancy-specific beta 1-glycoprotein genes: one is associated with hydatidiform mole. Proc. Nat. Acad. Sci. 87: 5822–5826, 1990.

[78317] 16647. Tatarinov, Y. S.; Falaleeva, D. M.; Kalashnikov, V. V.: Human pregnancy-specific beta-1-globulin and its relation to chorioepithelioma. Int. J. Cancer 17: 626–632, 1976.

[78318] 16648. Berry, C.; Cree, J.; Mann, T.: Aarskog's syndrome. Arch. Dis. Child. 55: 706–710, 1980.

[78319] 16649. Escobar, V.; Weaver, D. D.: Aarskog syndrome: new findings and genetic analysis. J. A. M. A. 240: 2638–2641, 1978.

[78320] 16650. Hanley, W. B.; McKusick, V. A.; Barranco, F. T.: Osteochondritis dissecans and associated malformations in brothers: a review of familial aspects. J. Bone Joint Surg.

49A: 925–937, 1967.

- [78321] 16651.Nagase, T.; Seki, N.; Tanaka, A.; Ishikawa, K.; No-mura, N.: Prediction of the coding sequences of unidentified human genes. IV. The coding sequences of 40 new genes (KIAA0121–KIAA0160) deduced by analysis of cDNA clones from human cell line KG–1. *DNA Res.* 2: 167–174, 1995.
- [78322] 16652.Parrish, J. E.; Ciccodicola, A.; Wehnert, M.; Cox, G. F.; Chen, E.; Nelson, D. L.: A muscle-specific DNase I-like gene in human Xq28. *Hum.Molec. Genet.* 4: 1557–1564, 1995.
- [78323] 16653.Pergolizzi, R.; Appierto, V.; Bosetti, A.; DeBellis, G. L.; Rovidia, E.; Biunno, I.: Cloning of a gene encoding a DNase I-like endonuclease in the human Xq28 region. *Gene* 168: 267–270, 1996.
- [78324] 16654.Klink, A.; Schiebel, K.; Winkelmann, M.; Rao, E.; Horsthemke, B.; Ludecke, H.–J.; Claussen, U.; Scherer, G.; Rappold, G.: The human protein kinase gene PKX1 on Xp22.3 displays Xp/Yp homology and is a site of chromosomal instability. *Hum. Molec. Genet.* 4: 869–878, 1995.
- [78325] 16655.Schiebel, K.; Mertz, A.; Winkelmann, B.; Glaser, B.; Schempp, W.; Rappold, G.: FISH localization of the human Y-homolog of protein kinase PRKX (PRKY) to Yp11.2 and

two pseudogenes to 15q26 and Xq12–q13. *Cytogenet. Cell Genet.* 76: 49–52, 1997.

[78326] 16656. Brown, C. J.; Carrel, L.; Willard, H. F.: Expression of genes from the human active and inactive X chromosomes. *Am. J. Hum. Genet.* 60:1333–1343, 1997.

[78327] 16657. Dong, B.; Horowitz, D. S.; Kobayashi, R.; Krainer, A. R.: Purification and cDNA cloning of HeLa cell p54(nrb), a nuclear protein with two RNA recognition motifs and extensive homology to human splicing factor PSF and *Drosophila* NONA/BJ6. *Nucleic Acids Res.* 21: 4085–4092, 1993.

[78328] 16658. Nelson, D. L.; Ballabio, A.; Cremers, F.; Monaco, A. P.; Schlessinger, D.: Report of the sixth international workshop on X chromosome mapping 1995. *Cytogenet. Cell Genet.* 71: 308–342, 1995.

[78329] 16659. Janssens, R.; Boeynaems, J.–M.; Godart, M.; Comuni, D.: Cloning of a human heptahelical receptor closely related to the P2Y(5) receptor. *Biochem. Biophys. Res. Commun.* 236: 106–112, 1997.

[78330] 16660. O'Dowd, B. F.; Nguyen, T.; Jung, B. P.; Marchese, A.; Cheng, R.; Heng, H. H. Q.; Kolakowski, L. F., Jr.; Lynch, K. R.; George, S. R.: Cloning and chromosomal mapping of four putative novel human G–protein–coupled receptor

genes. *Gene* 187: 75–81, 1997.

- [78331] 16661.Knight, S. W.; Heiss, N. S.; Vulliamy, T. J.; Aalfs, C. M.; McMahon,C.; Richmond, P.; Jones, A.; Hennekam, R. C. M.; Poustka, A.; Mason,P. J.; Dokal, I.: Unexplained aplastic anaemia, immunodeficiency,and cerebellar hypoplasia (Hoyeraal–Hreidarsson syndrome) due to mutationsin the dyskeratosis congenita gene, DKC1. *Brit. J. Haemat.* 107:335–339, 1999.
- [78332] 16662.Knight, S. W.; Heiss, N. S.; Vulliamy, T. J.; Greschner, S.; Stavrides,G.; Pai, G. S.; Lestringant, G.; Varma, N.; Mason, P. J.; Dokal, I.;Poustka, A.: X–linked dyskeratosis congenita is predominantly causedby mis–sense mutations in the DKC1 gene. *Am. J. Hum. Genet.* 65: 50–58,1999.
- [78333] 16663.Knight, S. W.; Vulliamy, T. J.; Morgan, B.; Devriendt, K.; Mason,P. J.; Dokal, I.: Identification of novel DKC1 mutations in patientswith dyskeratosis congenita: implications for pathophysiology anddiagnosis. *Hum. Genet.* 108: 299–303, 2001.
- [78334] 16664.Luzzatto, L.; Karadimitris, A.: Dyskeratosis and ribosomal rebellion. *NatureGenet.* 19: 6–7, 1998.
- [78335] 16665.McGrath, J. A.: Dyskeratosis congenita: new clinical and molecularinsights into ribosome function. *Lancet* 353:



1204–1205, 1999.

- [78336] 16666.Mitchell, J. R.; Wood, E.; Collins, K.: A telomerase component is defective in the human disease dyskeratosis congenita. *Nature* 402:551–555, 1999.
- [78337] 16667.Salowsky, R.; Heiss, N. S.; Benner, A.; Wittig, R.; Poustka, A.: Basal transcription activity of the dyskeratosis congenita genes is mediated by Sp1 and Sp3 and a patient mutation in a Sp1 binding site is associated with decreased promoter activity. *Gene* 293: 9–19, 2002.
- [78338] 16668.Vuliamy, T. J.; Knight, S. W.; Heiss, N. S.; Smith, O. P.; Poustka, A.; Dokal, I.; Mason, P. J.: Dyskeratosis congenita caused by a 3–prime deletion: germline and somatic mosaicism in a female carrier. *Blood* 94:1254–1260, 1999.
- [78339] 16669.Yaghmai, R.; Kimyai-Asadi, A.; Rostamiani, K.; Heiss, N. S.; Poustka, A.; Eyaid, W.; Bodurtha, J.; Nousari, H. C.; Hamosh, A.; Metzenberg, A.: Overlap of dyskeratosis congenita with the Hoyeraal–Hreidarsson syndrome. *J. Pediatr.* 136: 390–393, 2000.
- [78340] 16670.Greenfield, A.; Carrel, L.; Pennisi, D.; Philippe, C.; Quaderi, N.; Siggers, P.; Steiner, K.; Tam, P. P. L.; Monaco, A. P.; Willard, H. F.; Koopman, P.: The UTX gene escapes X inactivation in mice and humans. *Hum. Molec. Genet.* 7:

737–742, 1998.

- [78341] 16671.Greenfield, A. J.; Scott, D.; Pennisi, D.; Ehrmann, I.; Ellis,P.; Cooper, L.; Simpson, E.; Koopman, P.: An H–YDb epitope is encodedby a novel mouse Y chromosome gene. Nature Genet. 14: 474–478, 1996.
- [78342] 16672.Caput, D.; Laurent, P.; Kaghad, M.; Lelias, J.–M.; Lefort, S.;Vita, N.; Ferrara, P.: Cloning and characterization of a specificinterleukin (IL)–13 binding protein structurally related to the IL–5receptor alpha chain. J. Biol. Chem. 271: 16921–16926, 1996.
- [78343] 16673.Guo, J.; Apiou, F.; Mellerin, M.–P.; Lebeau, B.; Jacques, Y.; Minvielle,S.: Chromosome mapping and ex–pression of the human interleukin–13receptor. Genomics 42: 141–145, 1997.
- [78344] 16674.Lin, C.–S.; Aebersold, R. H.; Leavitt, J.: Correction of the N–terminalsequences of the human plastin iso–forms by using anchored polymerasechain reaction: iden–tification of a potential calcium–binding domain. Molec.Cell. Biol. 10: 1818–1821, 1990.
- [78345] 16675.Fukuda, M. N.; Sato, T.; Nakayama, J.; Klier, G.; Mikami, M.; Aoki,D.; Nozawa, S.: Trophinin and tastin, a novel cell adhesion moleculecomplex with potential in–volvement in embryo implantation. GenesDev. 9:

1199–1210, 1995.

- [78346] 16676.Pack, S. D.; Tanigami, A.; Ledbetter, D. H.; Sato, T.; Fukuda,M. N.: Assignment of trophoblast/endometrial epithelium cell adhesionmolecule trophinin gene TRO to human chromosome bands Xp11.22–p11.21by in situ hybridization. Cytogenet. Cell Genet. 79: 123–124, 1997.
- [78347] 16677.Suzuki, N.; Nakayama, J.; Shih, I. M.; Aoki, D.; Nozawa, S.; Fukuda,M. N.: Expression of trophinin, tastin, and bystin by trophoblastand endometrial cells in human placenta. Biol. Reprod. 60: 621–627,1999.
- [78348] 16678.Sana, T. R.; Debets, R.; Timans, J. C.; Bazan, J. F.; Kastelein,R. A.: Computational identification, cloning, and characterizationof IL1R9, a novel interleukin–1 receptor–like gene encoded over anunusually large interval of human chromosome Xq22.2–q22.3. Genomics 69:252–262, 2000.
- [78349] 16679.Bech–Hansen, N. T.; Naylor, M. J.; Maybaum, T. A.; Sparkes, R.L.; Koop, B.; Birch, D. G.; Bergen, A. A. B.; Prinsen, C. F. M.; Polomeno,R. C.; Gal, A.; Drack, A. V.; Musarella, M. A.; Jacobson, S. G.; Young,R. S. L.; Weleber, R. G.: Mutations in NYX, encoding the leucine–richproteoglycan nyctalopin, cause X–linked complete congenital stationarynight blindness. Nature Genet. 26:

319–323, 2000.

- [78350] 16680. Pusch, C. M.; Zeitz, C.; Brandau, O.; Pesch, K.; Achatz, H.; Feil, S.; Scharfe, C.; Maurer, J.; Jacobi, F. K.; Pinckers, A.; Andreasson, S.; Hardcastle, A.; Wissinger, B.; Berger, W.; Meindl, A.: The complete form of X-linked congenital stationary night blindness is caused by mutations in a gene encoding a leucine-rich repeat protein. *Nature-Genet.* 26: 324–327, 2000.
- [78351] 16681. Isbrandt, D.; Leicher, T.; Waldschutz, R.; Zhu, X.; Luhmann, U.; Michel, U.; Sauter, K.; Pongs, O.: Gene structures and expression profiles of three human KCND (Kv4) potassium channels mediating A-type currents  $I_{(to)}$  and  $I_{(sa)}$ . *Genomics* 64: 144–154, 2000.
- [78352] 16682. Davies, J. P.; Cotter, P. D.; Ioannou, Y. A.: Cloning and mapping of human Rab7 and Rab9 cDNA sequences and identification of a Rab9 pseudogene. *Genomics* 41: 131–134, 1997.
- [78353] 16683. Brinkmann, U.; Vasmatazis, G.; Lee, B.; Pastan, I.: Novel genes in the PAGE and GAGE family of tumor antigens found by homology walking in the dbEST database. *Cancer Res.* 59: 1445–1448, 1999.
- [78354] 16684. Liu, X. F.; Helman, L. J.; Yeung, C.; Bera, T. K.; Lee, B.; Pastan, I.: XAGE-1, a new gene that is frequently ex-

pressed in Ewing's sarcoma. *CancerRes.* 60: 4752–4755, 2000.

[78355] 16685.Lahn, B. T.; Page, D. C.: Functional coherence of the human Ychromosome. *Science* 278: 675–680, 1997.

[78356] 16686.Chung, J.; Lee, S.–G.; Song, K.: Identification of a human homolog of a putative RNA helicase gene (mDEAD3) expressed in mouse erythroid cells. *Korean J. Biochem.* 27: 193–197, 1995.

[78357] 16687.Park, S. H.; Lee, S.–G.; Kim, Y.; Song, K.: Assignment of a human putative RNA helicase gene, DDX3, to human X chromosome bands p11.3–p11.23. *Cytogenet. Cell Genet.* 81: 178–179, 1998.

[78358] 16688.Gasper, N. J.; Kinzy, T. G.; Scherer, B. J.; Humbelin, M.; Hershey, J. W. B.; Merrick, W. C.: Translation initiation factor eIF–2: cloning and expression of the human cDNA encoding the gamma–subunit. *J. Biol. Chem.* 269: 3415–3422, 1994.

[78359] 16689.Ehrmann, I. E.; Ellis, P. S.; Mazeyrat, S.; Duthie, S.; Brockdorff, N.; Mattei, M. G.; Gavin, M. A.; Affara, N. A.; Brown, G. M.; Simpson, E.; Mitchell, M. J.; Scott, D. M.: Characterization of genes encoding translation initiation factor eIF–2–gamma in mouse and human: sex chromosome localization, escape from X–inactivation and evolu–

tion. Hum.Molec. Genet. 7: 1725–1737, 1998.

- [78360] 16690.Lee, S. M. Y.; Tsui, S. K. W.; Chan, K. K.; Garcia-Barcelo, M.;Waye, M. M. Y.; Fung, K. P.; Liew, C. C.; Lee, C. Y.: Chromosomalmapping, tissue distribution and cDNA sequence of four-and-a-halfLIM domain protein 1 (FHL1). Gene 216: 163–170, 1998.
- [78361] 16691.Morgan, M. J.; Madgwick, A. J.; Charleston, B.; Pell, J. M.; Loughna,P. T.: The developmental regulation of a novel muscle LIM–protein. Biochem.Biophys. Res. Commun. 212: 840–846, 1995.
- [78362] 16692.Morgan, M. J.; Madgwick, A. J. A.: Slim defines a novel familyof LIM–proteins expressed in skeletal muscle. Biochem. Biophys. Res.Communic. 225: 632–638, 1996.
- [78363] 16693.Bard, J. A.; Zgombick, J.; Adham, N.; Vaysse, P.; Branchek, T.A.; Weinshank, R. L.: Cloning of a novel human serotonin receptor(5–HT–7) positively linked to adenylate cyclase. J. Biol. Chem. 268:23422–23426, 1993.
- [78364] 16694.Gelernter, J.; Rao, P. A.; Pauls, D. L.; Hamblin, M. W.; Sibley,D. R.; Kidd, K. K.: Assignment of the 5HT7 receptor gene (HTR7) to chromosome 10q and exclusion of genetic linkage with Tourette syndrome. Genomics 26:207–209, 1995.
- [78365] 16695.Kenakin, T. P.; Bond, R. A.; Bonner, T. I.: II. Defini–

tion of pharmacological receptors. *Pharm. Rev.* 44: 351–362, 1992.

- [78366] 16696. Lassig, J. P.; Vachirasomtoon, K.; Hartzell, K.; Leventhal, M.; Courchesne, E.; Courchesne, R.; Lord, C.; Leventhal, B. L.; Cook, E. H., Jr.: Physical mapping of the serotonin 5-HT(7) receptor gene (HTR7) to chromosome 10 and pseudogene (HTR7P) to chromosome 12, and testing of linkage disequilibrium between HTR7 and autistic disorder. *Am. J. Med. Genet.* 88: 472–475, 1999.
- [78367] 16697. Ruat, M.; Traiffort, E.; Leurs, R.; Tardivel-Lacombe, J.; Diaz, J.; Arrang, J.-M.; Schwartz, J.-C.: Molecular cloning, characterization, and localization of a high-affinity serotonin receptor (5-HT-7) activating cAMP formation. *Proc. Nat. Acad. Sci.* 90: 8547–8551, 1993.
- [78368] 16698. Tsou, A.; Kosaka, A.; Bach, C.; Zuppan, P.; Yee, C.; Tom, L.; Alvarez, R.; Ramsey, S.; Bonhaus, D. W.; Stefanich, E.; Jakeman, L.; Eglen, R. M.; Chan, H. W.: Cloning and expression of a 5-hydroxytryptamine-7 receptor positively coupled to adenylyl cyclase. *J. Neurochem.* 63: 456–464, 1994.
- [78369] 16699. Gelernter, J.; Kruger, S.; Pakstis, A. J.; Amara, S.; Kidd, K. K.: Linkage mapping of norepinephrine transporter protein and serotonin transporter protein genes.

(Abstract) Am. J. Hum. Genet. 53 (suppl.):A1003, 1993.

- [78370] 16700.Gelernter, J.; Pakstis, A. J.; Kidd, K. K.: Linkage mapping of serotonin transporter protein gene SLC6A4 on chromosome 17. Hum.Genet. 95: 677–680, 1995.
- [78371] 16701.Gregor, P.; Patel, A.; Shimada, S.; Lin, C.–L.; Rochelle, J. M.; Kitayama, S.; Seldin, M. F.; Uhl, G. R.: Murine serotonin transporter: sequence and localization to chromosome 11. Mammalian Genome 4:283–284, 1993.
- [78372] 16702.Hariri, A. R.; Mattay, V. S.; Tessitore, A.; Kolachana, B.; Fera, F.; Goldman, D.; Egan, M. F.; Weinberger, D. R.: Serotonin transporter genetic variation and the response of the human amygdala. Science 297:400–403, 2002.
- [78373] 16703.Heils, A.; Teufel, A.; Petri, S.; Seemann, M.; Bengel, D.; Balling, U.; Riederer, P.; Lesch, K. P.: Functional promoter and polyadenylation site mapping of the human serotonin (5-HT) transporter gene. J. Neural Transm. 102: 247–254, 1995.
- [78374] 16704.Heils, A.; Teufel, A.; Petri, S.; Stober, G.; Riederer, P.; Bengel, D.; Lesch, K. P.: Allelic variation of human serotonin transporter gene expression. J. Neurochem. 66: 2621–2624, 1996.
- [78375] 16705.Kim, S.–J.; Cox, N.; Courchesne, R.; Lord, C.; Corsello, C.; Akshoomoff, N.; Guter, S.; Leventhal, B. L.;



Courchesne, E.; Cook, E. H., Jr.: Transmission disequilibrium mapping at the serotonin transporter gene (SLC6A4) region in autistic disorder. *Molec. Psychiat.* 7: 278–288, 2002.

[78376] 16706. Lesch, K. P.; Balling, U.; Gross, J.; Strauss, K.; Wolozin, B. L.; Murphy, D. L.: Organisation of the human serotonin transporter gene. *J. Neural Trans.* 95: 157–162, 1994.

[78377] 16707. Chern, C. J.; Beutler, E.: Biochemical and electrophoretic studies of erythrocyte pyridoxine kinase in white and black Americans. *Am. J. Hum. Genet.* 28: 9–17, 1976.

[78378] 16708. Hanna, M. C.; Turner, A. J.; Kirkness, E. F.: Human pyridoxalkinase: cDNA cloning, expression, and modulation by ligands of the benzodiazepine receptor. *J. Biol. Chem.* 272: 10756–10760, 1997.

[78379] 16709. Dougherty, K. M.; Brandriss, M. C.; Valle, D.: Cloning human pyrroline-5-carboxylate reductase cDNA by complementation in *Saccharomyces cerevisiae*. *J. Biol. Chem.* 267: 871–875, 1992.

[78380] 16710. Merrill, M. J.; Yeh, G. C.; Phang, J. M.: Purified human erythrocyte pyrroline-5-carboxylate reductase: preferential oxidation of NADPH. *J. Biol. Chem.* 264:

9352–9358, 1989.

- [78381] 16711.Brissenden, J. E.; Caras, I.; Thelander, L.; Francke, U.: The structural gene for the M1 subunit of human ribonucleotide reductase maps to chromosome 11, band p15, in human and to chromosome 7 in mouse. *Exp.Cell Res.* 174: 302–308, 1988.
- [78382] 16712.Byrne, J.; Smith, P.: Human polymorphic probe pE1.8 detects SacI polymorphism in the ribonucleotide reductase M1 subunit gene. *Hum.Genet.* 87: 376 only, 1991.
- [78383] 16713.Barletta, C.; Druck, T.; LaForgia, S.; Calabretta, B.; Drabkin, H.; Patterson, D.; Croce, C. M.; Huebner, K.: Chromosome locations of the MYB related genes, AMYB and BMYB. *Cancer Res.* 51: 3821–3824, 1991.
- [78384] 16714.Nomura, N.; Takahashi, M.; Matsui, M.; Ishii, S.; Date, T.; Sasamoto, S.; Ishizaki, R.: Isolation of human cDNA clones of MYB-related genes, A-MYB and B-MYB. *Nucleic Acids Res.* 16: 11075–11089, 1988.
- [78385] 16715.Takahashi, T.; Nakagoshi, H.; Sarai, A.; Nomura, N.; Yamamoto, T.; Ishii, S.: Human A-myb gene encodes a transcriptional activator containing the negative regulatory domains. *FEBS Lett.* 358: 89–96, 1995.
- [78386] 16716.Boylan, K. B.; Takahashi, N.; Diamond, M.; Hood, L.

E.; Prusiner, S. B.: DNA length polymorphism located 5-prime to the human myelin basic protein gene. *Am. J. Hum. Genet.* 40: 387–400, 1987.

[78387] 16717. Dayhoff, M. O.: *Atlas of Protein Sequence and Structure*. Myelin membrane encephalitogenic protein. Washington: National Biomedical Research Foundation (pub.) 5: 1972. Pp. D324 only.

[78388] 16718. Eylar, E. H.; Brostoff, S.; Hashim, G.; Westall, F. C.: Basic A1 protein of the myelin membrane: the complete amino acid sequence. *J. Biol. Chem.* 246: 5770–5784, 1971.

[78389] 16719. Gomez, C. M.; Muggleton-Harris, A. L.; Whittingham, D. G.; Hood, L. E.; Readhead, C.: Rapid preimplantation detection of mutant (shiverer) and normal alleles of the mouse myelin basic protein gene allowing selective implantation and birth of live young. *Proc. Nat. Acad. Sci.* 87: 4481–4484, 1990.

[78390] 16720. Kamholz, J.; de Ferra, F.; Puckett, C.; Lazzarini, R.: Identification of three forms of human myelin basic protein by cDNA cloning. *Proc. Nat. Acad. Sci.* 83: 4962–4966, 1986.

[78391] 16721. Kamholz, J.; Spielman, R.; Gogolin, K.; Modi, W.; O'Brien, S.; Lazzarini, R.: The human myelin–basic–protein

gene: chromosomal localization and RFLP analysis. *Am. J. Hum. Genet.* 40: 365–373, 1987.

- [78392] 16722. Lalley, P. A.; McKusick, V. A.: Report of the committee on comparative mapping (HGM8). *Cytogenet. Cell Genet.* 40: 536–566, 1985.
- [78393] 16723. Martensen, R. E.: Myelin basic protein speciation. *Prog. Clin. Biol. Res.* 146: 511–521, 1984.
- [78394] 16724. Marty, M. C.; Alliot, F.; Rutin, J.; Fritz, R.; Trisler, D.; Pessac, B.: The myelin basic protein gene is expressed in differentiated blood cell lineages and in hemopoietic progenitors. *Proc. Nat. Acad. Sci.* 99: 8856–8861, 2002.
- [78395] 16725. Molineaux, S. M.; Engh, H.; de Ferra, F.; Hudson, L.; Lazzarini, R. A.: Recombination within the myelin basic protein gene created the dysmyelinating shiverer mouse mutation. *Proc. Nat. Acad. Sci.* 83: 7542–7546, 1986.
- [78396] 16726. Popko, B.; Puckett, C.; Lai, E.; Shine, H. D.; Readhead, C.; Takahashi, N.; Hunt, S. W., III; Sidman, R. L.; Hood, L.: Myelin deficient mice: expression of myelin basic protein and generation of mice with varying levels of myelin. *Cell* 48: 713–721, 1987.
- [78397] 16727. Readhead, C.; Popko, B.; Takahashi, N.; Shine, H. D.; Saavedra, R. A.; Sidman, R. L.; Hood, L.: Expression of a myelin basic protein gene in transgenic shiverer mice: cor-

rection of the dysmyelinating phenotype. Cell 48: 703–712, 1987.

- [78398] 16728.Roach, A.; Boylan, K.; Horvath, S.; Prusiner, S. B.; Hood, L.E.: Characterization of cloned cDNA representing rat myelin basic protein: absence of expression in shiverer mutant mice. Cell 34:799–806, 1983.
- [78399] 16729.Roach, A.; Takahashi, N.; Pravtcheva, D.; Ruddle, F.; Hood, L.: Chromosomal mapping of mouse myelin basic protein gene and structure and transcription of the partially deleted gene in shiverer mutant mice. Cell 42: 149–155, 1985.
- [78400] 16730.Saxe, D. F.; Takahashi, N.; Hood, L.; Simon, M. I.: Localization of the human myelin basic protein gene (MBP) to region 18q22–qter by in situ hybridization. Cytogenet. Cell Genet. 39: 246–249, 1985.
- [78401] 16731.Sheremata, W.; Cosgrove, J. B. R.; Hylar, E. H.: Cellular hypersensitivity to basic myelin (A1) protein and clinical multiple sclerosis. New Eng. J. Med. 291: 14–17, 1974.
- [78402] 16732.Sidman, R.: Personal Communication. Boston, Mass. 1983.
- [78403] 16733.Sidman, R. L.; Conover, C. S.; Carson, J. H.: Shiverer gene maps near the distal end of chromosome 18 in the house mouse. Cytogenet. Cell Genet. 39: 241–245, 1985.

- [78404] 16734. Sparkes, R. S.; Mohandas, T.; Heinzmann, C.; Roth, H. J.; Klisak, I.; Campagnoni, A. T.: Assignment of the myelin basic protein gene to human chromosome 18q22-qter. *Hum. Genet.* 75: 147–150, 1987.
- [78405] 16735. Takahashi, N.; Roach, A.; Teplow, D. B.; Prusiner, S. B.; Hood, L.: Cloning and characterization of the myelin basic protein gene from mouse: one gene can encode both 14 kd and 18.5 kd MBPs by alternate use of exons. *Cell* 42: 139–148, 1985.
- [78406] 16736. Barouch, L. A.; Harrison, R. W.; Skaf, M. W.; Rosas, G. O.; Cappola, T. P.; Kobeissi, Z. A.; Hobai, I. A.; Lemmon, C. A.; Burnett, A. L.; O'Rourke, B.; Rodriguez, E. R.; Huang, P. L.; Lima, J. A. C.; Berkowitz, D. E.; Hare, J. M.: Nitric oxide regulates the heart by spatial confinement of nitric oxide synthase isoforms. *Nature* 416: 337–340, 2002.
- [78407] 16737. Dimmeler, S.; Fleming, I.; Fisslthaler, B.; Hermann, C.; Busse, R.; Zeiher, A. M.: Activation of nitric oxide synthase in endothelial cells by Akt-dependent phosphorylation. *Nature* 399: 601–605, 1999.
- [78408] 16738. Fulton, D.; Gratton, J.-P.; McCabe, T. J.; Fontana, J.; Fujio, Y.; Walsh, K.; Franke, T. F.; Papapetropoulos, A.; Sessa, W. C.: Regulation of endothelium-derived nitric oxide production by the protein kinase Akt. *Nature* 399:

597–601, 1999.

- [78409] 16739.Kruh, G. D.; King, C. R.; Kraus, M. H.; Popescu, N. C.; Amsbaugh, S. C.; McBride, W. O.; Aaronson, S. A.: A novel human gene closely related to the abl proto-oncogene. *Science* 234: 1545–1548, 1986.
- [78410] 16740.Abe, T.; Kikuchi, T.; Shinohara, T.: The sequence of the human phosducin gene (PDC) and its 5-prime-flanking region. *Genomics* 19:369–372, 1994.
- [78411] 16741.Craft, C. M.; Lolley, R. N.; Seldin, M. F.; Lee, R. H.: Rat pineal gland phosducin: cDNA isolation, nucleotide sequence, and chromosomal assignment in the mouse. *Genomics* 10: 400–409, 1991.
- [78412] 16742.Ding, C.; Li, X.; Griffin, C. A.; Jabs, E. W.; Hawkins, A. L.; Levine, M. A.: The gene for human phosducin (PDC), a soluble protein that binds G-protein beta gamma dimers, maps to 1q25–q31.1. *Genomics* 18:457–459, 1993.
- [78413] 16743.Lee, R. H.; Fowler, A.; McGinnis, J. F.; Lolley, R. N.; Craft, C. M.: Amino acid and cDNA sequence of bovine phosducin, a soluble phosphoprotein from photoreceptor cells. *J. Biol. Chem.* 265: 15867–15873, 1990.
- [78414] 16744.Reiter, R. J.: The mammalian pineal gland: structure and function. *Am.J. Anat.* 162: 287–313, 1981.
- [78415] 16745.Sparkes, R. S.; Lee, R. H.; Shinohara, T.; Craft, C.

M.; Kojis,T.; Klisak, I.; Heinzmann, C.; Bateman, J. B.: Assignment of the phosphatidylethanolamine phosphatase (PEP) gene to human chromosome 1q25–1q32.1 by somatic cell hybridization and in situ hybridization. *Genomics* 18: 426–428, 1993.

[78416] 16746.Arnaud, J.; Vavrusa, B.; Sevin, J.; Constans, J.: Human red-cell acid phosphatase (ACP1): a new mutant (ACP1\*KUK) detected by isoelectric focusing, kinetics of thermostability and substrate activity. *Hum.Hered.* 39: 288–293, 1989.

[78417] 16747.Arnaud, J.; Vavrusa, B.; Wiederanders, G.; Constans, J.: Human red-cell acid phosphatase (ACP1): kinetic and thermodynamic characterization of the KUK variant. *Hum. Hered.* 42: 140–142, 1992.

[78418] 16748.Beemer, F. A.; van der Heiden, C.; Van Hemel, J. O.; Jansen, M.: Letter to the editors. (Letter) *Clin. Genet.* 24: 151, 1983.

[78419] 16749.Berg, K.: Close linkage between APOB and ACP1 excluded. (Abstract) *Cytogenet.Cell Genet.* 46: 580, 1987.

[78420] 16750.Bottini, E.; Carapella, E.; Orzalesi, M.; Lucarelli, P.; Pascone,R.; Gloria-Bottini, F.; Coccia, M.: Is there a role of erythrocyte acid phosphatase polymorphism in intrauterine development? (Letter) *Am.J. Hum. Genet.* 32: 764–767, 1980.



- [78421] 16751.Seldin, M. F.; Kruh, G. D.: Mapping of Abll within a conservedlinkage group on distal mouse chromosome 1 syntenic with human chromosome1 using an interspecific cross. *Genomics* 4: 221–223, 1989.
- [78422] 16752.Bello, M. J.; Salagnon, N.; Rey, J. A.; Guichaoua, M. R.; Berge–Lefranc,J. L.; Jordan, B. R.; Luciani, J. M.: Precise in situ localizationof NCAM, ETS1, and D11S29 on human meiotic chromosomes. *Cytogenet.Cell Genet.* 52: 7–10, 1989.
- [78423] 16753.Gold, D. P.; van Dongen, J. J. M.; Morton, C. C.; Bruns, G. A.P.; van den Elsen, P.; Geurts van Kessel, A. H. M.; Terhorst, C.:The gene encoding the epsilon subunit of the T3/T–cell receptor complexmaps to chromosome 11 in humans and to chromosome 9 in mice. *Proc.Nat. Acad. Sci.* 84: 1664–1668, 1987.
- [78424] 16754.Ohtani, N.; Zebedee, Z.; Huot, T. J. G.; Stinson, J. A.; Sugimoto,M.; Ohashi, Y.; Sharrocks, A. D.; Peters, G.; Hara, E.: Opposingeffects of Ets and Id proteins on p16(INK4A) expression during cellularsenescence. *Nature* 409: 1067–1070, 2001.
- [78425] 16755.Sacchi, N.; Watson, D. K.; Geurts van Kessel, A. H. M.; Hagemeijer,A.; Kersey, J.; Drabkin, H. D.; Patterson, D.; Papas, T. S.: Hu–ets–1and Hu–ets–2 genes are transposed

in acute leukemias with (4;11) and(8;21) translocations.

Science 231: 379–382, 1986.

- [78426] 16756.Watson, D. K.; McWilliams–Smith, M. J.; Kozak, C.; Reeves, R.;Gearhart, J.; Nunn, M. F.; Nash, W.; Fowle, J. R., III; Duesberg,P.; Papas, T. S.; O'Brien, S. J.: Conserved chromosomal positionsof dual domains of the ets protooncogene in cats, mice, and humans. Proc.Nat. Acad. Sci. 83: 1792–1796, 1986.
- [78427] 16757.Watson, D. K.; McWilliams–Smith, M. J.; Nunn, M. F.; Duesberg,P. H.; O'Brien, S. J.; Papas, T. S.: The ets sequence from the transforminggene of avian erythroblastosis virus, E26, has unique domains on humanchromosomes 11 and 21: both loci are transcriptionally active. Proc.Nat. Acad. Sci. 82: 7294–7298, 1985.
- [78428] 16758.Bellacosa, A.; Franke, T. F.; Gonzalez–Portal, M. E.; Datta, K.;Taguchi, T.; Gardner, J.; Cheng, J. Q.; Testa, J. R.; Tsichlis, P.N.: Structure, expression and chromosomal mapping of c–akt: relationship to v–akt and its implications. Oncogene 8: 745–754, 1993.
- [78429] 16759.Brunet, A.; Bonni, A.; Zigmond, M. J.; Lin, M. Z.; Juo, P.; Hu,L. S.; Anderson, M. J.; Arden, K. C.; Blenis, J.; Greenberg, M. E.: Akt promotes cell survival by phosphorylating and inhibiting a Forkheadtranscription factor. Cell

96: 857–868, 1999.

- [78430] 16760.Chen, W. S.; Xu, P.–Z.; Gottlob, K.; Chen, M.–L.; Sokol, K.; Shiyanova,T.; Roninson, I.; Weng, W.; Suzuki, R.; Tobe, K.; Kadowaki, T.; Hay,N.: Growth retardation and increased apoptosis in mice with homozygousdisruption of the akt1 gene. *Genes Dev.* 15: 2203–2208, 2001.
- [78431] 16761.Condorelli, G.; Drusco, A.; Stassi, G.; Bellacosa, A.; Roncarati,R.; Iaccarino, G.; Russo, M. A.; Gu, Y.; Dalton, N.; Chung, C.; Latronico,M. V. G.; Napoli, C.; Sadoshima, J.; Croce, C. M.; Ross, J., Jr.:Akt induces enhanced myocardial contractility and cell size in vivo in transgenic mice. *Proc. Nat. Acad. Sci.* 99: 12333–12338, 2002.
- [78432] 16762.Ellis, N. A.: Ecce Ohno. *Nature Genet.* 10: 373–375, 1995.
- [78433] 16763.Haldane, J. B. S.: Sex ratio and unisexual sterility in hybrid animals. *J. Genet.* 12: 101–109, 1922.
- [78434] 16764.Milatovich, A.; Kitamura, T.; Miyajima, A.; Francke, U.: Gene for the alpha–subunit of the human interleukin–3 receptor (IL3RA) localized to the X–Y pseudoautosomal region. *Am. J. Hum. Genet.* 53:1146–1153, 1993.
- [78435] 16765.Ohno, S.: Sex Chromosomes and Sex–linked Genes. Berlin and New York: Springer (pub.) 1967.
- [78436] 16766.Palmer, S.; Perry, J.; Ashworth, A.: A contravention

of Ohno's law in mice. *Nature Genet.* 10: 472–476, 1995.

[78437] 16767. Rugarli, E. I.; Adler, D. A.; Borsani, G.; Tsuchiya, K.; Franco, B.; Hauge, X.; Distèche, C.; Chapman, V.; Ballabio, A.: Different chromosomal localization of the *Clcn4* gene in *Mus spretus* and C57BL/6J mice. *Nature Genet.* 10: 466–471, 1995.

[78438] 16768. Schnur, R. E.; Wick, P. A.: Intragenic *TaqI* restriction fragment length polymorphism (RFLP) in *CLCN4*, between the loci for X-linked ocular albinism (OA1) and microphthalmia with linear skin defect syndrome (MLS). *Hum. Genet.* 95: 594–595, 1995.

[78439] 16769. van Slegtenhorst, M. A.; Bassi, M. T.; Borsani, G.; Wapenaar, M. C.; Ferrero, G. B.; de Conciliis, L.; Rugarli, E. I.; Grillo, A.; Franco, B.; Zoghbi, H. Y.; Ballabio, A.: A gene from the Xp22.3 region shares homology with voltage-gated chloride channels. *Hum. Molec. Genet.* 3: 547–552, 1994.

[78440] 16770. Curry, C. J. R.; Magenis, R. E.; Brown, M.; Lanman, J. T., Jr.; Tsai, J.; O'Lague, P.; Goodfellow, P.; Mohandas, T.; Bergner, E. A.; Shapiro, L. J.: Inherited chondrodysplasia punctata due to a deletion of the terminal short arm of an X chromosome. *New Eng. J. Med.* 311: 1010–1015, 1984.

[78441] 16771. Maslen, G. L.; Boyd, Y.: Comparative mapping of

the Grpr locus on the X chromosomes of man and mouse. Genomics 17: 106–109, 1993.

[78442] 16772.Schantz, L. J.; Naylor, S. L.; Giladi, E.; Spindel, E. R.: Assignment of the GRP receptor gene to the human X chromosome. (Abstract) Cytogenet. Cell Genet. 58: 2085–2086, 1991.

[78443] 16773.Shiraishi, M.; Alitalo, T.; Sekiya, T.: The chromosomal organization of the human endogenous retrovirus-like sequence HERV-H: clustering of the HERV-H sequences in a 300-kb region close to the GRPR locus on the X chromosome. DNA Res. 3: 425–429, 1996.

[78444] 16774.Shriver, S. P.; Bourdeau, H. A.; Gubish, C. T.; Tirpak, D. L.; Davis, A. L. G.; Luketich, J. D.; Siegfried, J. M.: Sex-specific expression of gastrin-releasing peptide receptor: relationship to smoking history and risk of lung cancer. J. Nat. Cancer Inst. 92:24–33, 2000.

[78445] 16775.Spindel, E. R.; Giladi, E.; Brehm, P.; Goodman, R. H.; Segerson, T. P.: Cloning and functional characterization of a complementary DNA encoding the murine fibroblast bombesin/gastrin-releasing peptide receptor. Molec. Endocr. 4: 1956–1963, 1990.

[78446] 16776.Ma, K.; Sharkey, A.; Kirsch, S.; Vogt, P.; Keil, R.; Hargreave, T. B.; McBeath, S.; Chandley, A. C.: Towards the

molecular localisation of the AZF locus: mapping of microdeletions in azoospermic men within 14 subintervals of interval 6 of the human Y chromosome. Hum.

Molec.Genet. 1: 29–33, 1992.

[78447] 16777. Adam, A.; Tippett, P.; Gavin, J.; Noades, J.; Sanger, R.; Race, R. R.: The linkage relation of Xg to G6PD in Israelis: the evidence of a second series of families. Am. J. Hum. Genet. 30: 211–218, 1966.

[78448] 16778. Ahluwalia, A.; Corcoran, C. M.; Vulliamy, T. J.; Ishwad, C. S.; Naidu, J. M.; Argusti, A.; Stevens, D. J.; Mason, P. J.; Luzzatto, L.: G6PD Kalyan and G6PD Kerala; two deficient variants in India caused by the same 317 glu-to-lys mutation. Hum. Molec. Genet. 3: 209–210, 1992.

[78449] 16779. Aksoy, K.; Yuregir, G. T.; Dikmen, N.; Unlukurt, I.: Three new G6PD variants, G6PD Adana, G6PD Samandag, and G6PD Balcali in Cukurova, Turkey. Hum. Genet. 76: 199–201, 1987.

[78450] 16780. Alfinito, F.; Cimmino, A.; Ferraro, F.; Cubellis, M. V.; Vitagliano, L.; Francese, M.; Zagari, A.; Rotoli, B.; Filosa, S.; Martini, G.: Molecular characterization of G6PD deficiency in Southern Italy: heterogeneity, correlation genotype–phenotype and description of a new variant (G6PD Neapolis). Brit. J. Haemat. 98: 41–46, 1997.

- [78451] 16781.Alperin, J. B.; Mills, G. C.: New variants of glucose-6-phosphatedehydrogenase (G6PD). Clin. Res. 20: 76, 1972.
- [78452] 16782.Azevedo, E.; Kirkman, H. N.; Morrow, A. C.; Motulsky, A. G.: Variantsof red cell glucose-6-phosphate dehydrogenase among Asiatic Indians. Ann.Hum. Genet. 31: 373-379, 1968.
- [78453] 16783.Azevedo, E. S.; Yoshida, A.: Brazilian variant of glucose-6-phosphatedehydrogenase (GD Minas Gerais). Nature 222: 380-382, 1969.
- [78454] 16784.Babalola, A. O. G.; Beetlestone, J. G.; Luzzatto, L.: Geneticvariants of human erythrocyte glucose-6-phosphate dehydrogenase: kineticand thermodynamic parameters of variants A, B, and A- in relationto quaternary structure. J. Biol. Chem. 251: 2993-3002, 1976.
- [78455] 16785.Balinsky, D.; Cayanis, E.; Carter, G.; Jenkins, T.; Bersohn, I.: A new variant of human erythrocyte glucose-6-phosphate dehydrogenase:G6PD Port Elizabeth. Int. J. Biochem. 4: 235-244, 1973.
- [78456] 16786.Balinsky, D.; Gomperts, E.; Cayanis, E.; Jenkins, T.; Bryer, D.;Bersohn, I.; Metz, J.: Glucose 6-phosphate dehydrogenase Johannesburg:a new variant with reduced activity in a patient with congenital non-

spherocythaemolytic anaemia. Brit. J. Haemat. 25: 385–392, 1973.

- [78457] 16787. Balinsky, D.; Rootman, A. J.; Nurse, G. T.; Cayanis, E.; Lane, A.; Jenkins, T.; Bersohn, I.: G6PD Kuanyama: a new variant of human erythrocyte glucose 6-phosphate dehydrogenase showing slower than normal electrophoretic mobility. S. Afr. J. Med. Sci. 39: 5–13, 1974.
- [78458] 16788. Barretto, O. C.; Nonoyama, K.: Gd(+) Cuiaba, a new rare glucose-6-phosphate dehydrogenase variant presenting normal activity. Hum. Genet. 77: 201–202, 1987.
- [78459] 16789. Kornak, U.; Kasper, D.; Bosl, M. R.; Kaiser, E.; Schweizer, M.; Schulz, A.; Friedrich, W.; Delling, G.; Jentsch, T. J.: Loss of the ClC-7 chloride channel leads to osteopetrosis in mice and man. Cell 104: 205–215, 2001.
- [78460] 16790. Kornak, U.; Schulz, A.; Friedrich, W.; Uhlhaas, S.; Kremens, B.; Voit, T.; Hasan, C.; Bode, U.; Jentsch, T. J.; Kubisch, C.: Mutations in the  $\alpha 3$  subunit of the vacuolar H(+)–ATPase cause infantile malignant osteopetrosis. Hum. Molec. Genet. 9: 2059–2063, 2000.
- [78461] 16791. Rosenberg, C. L.; Wong, E.; Petty, E. M.; Bale, A. E.; Tsujimoto, Y.; Harris, N. L.; Arnold, A.: PRAD1, a candidate BCL1 oncogene: mapping and expression in centrocytic lymphoma. Proc. Nat. Acad. Sci. 88: 9638–9642, 1991.



- [78462] 16792. Tsujimoto, Y.; Jaffe, E.; Cossman, J.; Gorham, J.; Nowell, P.C.; Croce, C. M.: Clustering of breakpoints on chromosome 11 in human B-cell neoplasms with the t(11;14) chromosome translocation. *Nature* 315:340–343, 1985.
- [78463] 16793. Tsujimoto, Y.; Yunis, J.; Onorato-Showe, L.; Erikson, J.; Nowell, P. C.; Croce, C. M.: Molecular cloning of the chromosomal breakpoint of B-cell lymphomas and leukemias with the t(11;14) chromosome translocation. *Science* 224:1403–1406, 1984.
- [78464] 16794. Barbany, G.; Hoglund, M.; Simonsson, B.: Complete molecular remission in chronic myelogenous leukemia after imatinib therapy. (Letter) *New Eng. J. Med.* 347: 539–540, 2002.
- [78465] 16795. Budarf, M.; Canaani, E.; Emanuel, B. S.: Linear order of the four BCR-related loci in 22q11. *Genomics* 3: 168–171, 1988.
- [78466] 16796. Castellanos, A.; Pintado, B.; Weruaga, E.; Arevalo, R.; Lopez, A.; Orfao, A.; Sanchez-Garcia, I.: A BCR-ABL(p190) fusion gene made by homologous recombination causes B-cell acute lymphoblastic leukemia in chimeric mice with independence of the endogenous bcr product. *Blood* 90:2168–2174, 1997. 1. Brunning, R. D.:

Philadelphia chromosome positive leukemia. *Hum. Path.* 11: 307–309, 1980.

[78467] 16797. Chisoe, S. L.; Bodenteich, A.; Wang, Y.-F.; Wang, Y.-P.; Burian, D.; Clifton, S. W.; Crabtree, J.; Freeman, A.; Iyer, K.; Jian, L.; Ma, Y.; McLaury, H.-J.; Pan, H.-Q.; Sarhan, O. H.; Toth, S.; Wang, Z.; Zhang, G.; Heisterkamp, N.; Groffen, J.; Roe, B. A.: Sequence and analysis of the human ABL gene, the BCR gene, and regions involved in the Philadelphia chromosomal translocation. *Genomics* 27: 67–82, 1995.

[78468] 16798. Court Brown, W. M.; Doll, R.: Mortality from cancer and other causes after radiotherapy for ankylosing spondylitis. *Brit. Med. J.* 2: 1327–1332, 1965.

[78469] 16799. Croce, C. M.; Huebner, K.; Isobe, M.; Fainstein, E.; Lifshitz, B.; Shtivelman, E.; Canaani, E.: Mapping of four distinct BCR-related loci to chromosome region 22q11: order of BCR loci relative to chronic myelogenous leukemia and acute lymphoblastic leukemia breakpoints. *Proc. Nat. Acad. Sci.* 84: 7174–7178, 1987.

[78470] 16800. Daley, G. Q.; Van Etten, R. A.; Baltimore, D.: Induction of chronic myelogenous leukemia in mice by the P210(bcr/abl) gene of the Philadelphia chromosome. *Science* 247: 824–830, 1990.

- [78471] 16801.de Klein, A.; Geurts van Kessel, A.; Grosveld, G.; Bartram, C.R.; Hagemeijer, A.; Bootsma, D.; Spurr, N. K.; Heisterkamp, N.; Groffen, J.; Stephenson, J. R.: A cellular oncogene is translocated to the Philadelphia chromosome in chronic myelocytic leukaemia. *Nature* 300:765–767, 1982.
- [78472] 16802.Diekmann, D.; Brill, S.; Garrett, M. D.; Totty, N.; Hsuan, J.; Monfries, C.; Hall, C.; Lim, L.; Hall, A.: Bcr encodes a GTPase-activating protein for p21(rac). *Nature* 351: 400–402, 1991.
- [78473] 16803.Druker, B. J.; Sawyers, C. L.; Kantarjian, H.; Resta, D. J.; Reese, S. F.; Ford, J. M.; Capdeville, R.; Talpaz, M.: Activity of a specific inhibitor of the BCR–ABL tyrosine kinase in the blast crisis of chronic myeloid leukemia and acute lymphoblastic leukemia with the Philadelphia chromosome. *New Eng. J. Med.* 344: 1038–1042, 2001.
- [78474] 16804.Druker, B. J.; Talpaz, M.; Resta, D. J.; Peng, B.; Buchdunger, E.; Ford, J. M.; Lydon, N. B.; Kantarjian, H.; Capdeville, R.; Ohno–Jones, S.; Sawyers, C. L.: Efficacy and safety of a specific inhibitor of the BCR–ABL tyrosine kinase in chronic myeloid leukemia. *New Eng. J. Med.* 344: 1031–1037, 2001.
- [78475] 16805.Emanuel, B. S.; Selden, J. R.; Wang, E.; Nowell, P. C.;

Croce, C. M.: In situ hybridization and translocation breakpoint mapping. I. Non-identical 22q11 breakpoints for the t(9;22) of CML and t(8;22) of Burkitt lymphoma. *Cytogenet. Cell Genet.* 38: 127–131, 1984.

[78476] 16806. Grandori, C.; Mac, J.; Siebelt, F.; Ayer, D. E.; Eisenman, R. N.: Myc–Max heterodimers activate a DEAD box gene and interact with multiple E box–related sites in vivo. *EMBO J.* 15: 4344–4357, 1996.

[78477] 16807. Gallagher, A. R.; Cedzich, A.; Gretz, N.; Somlo, S.; Witzgall, R.: The polycystic kidney disease protein PKD2 interacts with Hax–1, a protein associated with the actin cytoskeleton. *Proc. Nat. Acad. Sci.* 97: 4017–4022, 2000.

[78478] 16808. Cleutjens, K. B. J. M.; van Eekelen, C. C. E. M.; van der Korput, H. A. G. M.; Brinkmann, A. O.; Trapman, J.: Two androgen response regions cooperate in steroid hormone regulated activity of the prostate–specific antigen promoter. *J. Biol. Chem.* 271: 6379–6388, 1996.

[78479] 16809. Evans, B. A.; Drinkwater, C. C.; Richards, R. I.: Mouse glandular kallikrein genes: structure and partial sequence analysis of the kallikrein gene locus. *J. Biol. Chem.* 262: 8027–8034, 1987.

[78480] 16810. Lundwall, A.; Lilja, H.: Molecular cloning of human prostate specific antigen cDNA. *FEBS Lett.* 214: 317–322,

1987.

[78481] 16811.Melegos, D. N.; Yu, H.; Ashok, M.; Wang, C.;  
Stanczyk, F.; Diamandis,E. P.: Prostate-specific antigen in  
female serum, a potential newmarker of androgen excess.  
J. Clin. Endocr. Metab. 82: 777–780,1997.

- [78482] 16812.Prendergast, G. C.; Lawe, D.; Ziff, E. B.: Association of Myn,the murine homolog of Max, with c-Myc stimulates methylation-sensitiveDNA binding and Ras cotransformation. *Cell* 65: 395–408, 1991.
- [78483] 16813.Wagner, A. J.; Le Beau, M. M.; Diaz, M. O.; Hay, N.: Expression,regulation, and chromosomal localization of the Max gene. *Proc. Nat.Acad. Sci.* 89: 3111–3115, 1992.
- [78484] 16814.Zervos, A. S.; Faccio, L.; Gatto, J. P.; Kyriakis, J. M.; Brent,R.: Mxi2, a mitogen-activated protein kinase that recognizes andphosphorylates Max protein. *Proc. Nat. Acad. Sci.* 92: 10531–10534,1995.
- [78485] 16815.Emi, M.; Katagiri, T.; Harada, Y.; Saito, H.; Inazawa, J.; Ito,I.; Kasumi, F.; Nakamura, Y.: A novel metalloprotease/disintegrin-likegene at 17q21.3 is somatically rearranged in two primary breast cancers. *NatureGenet.* 5: 151–157, 1993.
- [78486] 16816.Katagiri, T.; Harada, Y.; Emi, M.; Nakamura, Y.: Human metalloprotease/disintegrin-like(MDC) gene: exon-intron organization and alternative splicing. *Cytogenet.Cell Genet.* 68: 39–44, 1995.
- [78487] 16817.Begley, C. G.; Lipkowitz, S.; Gobel, V.; Mahon, K. A.; Bertness,V.; Green, A. R.; Gough, N. M.; Kirsch, I. R.: Molecular characterizationof NSCL, a gene encoding a he-

lix-loop-helix protein expressed in the developing nervous system. *Proc. Nat. Acad. Sci.* 89: 38–42, 1992.

[78488] 16818. Brown, L.; Espinosa, R., III; Le Beau, M. M.; Siciliano, M. J.; Baer, R.: HEN1 and HEN2: a subgroup of basic helix-loop-helix genes that are coexpressed in a human neuroblastoma. *Proc. Nat. Acad. Sci.* 89: 8492–8496, 1992.

[78489] 16819. Cogliati, T.; Good, D. J.; Haigney, M.; Delgado-Romero, P.; Eckhaus, M. A.; Koch, W. J.; Kirsch, I. R.: Pre-disposition to arrhythmia and autonomic dysfunction in *Nhlh1*-deficient mice. *Molec. Cell. Biol.* 22: 4977–4983, 2002.

[78490] 16820. Lipkowitz, S.; Gobel, V.; Varterasian, M. L.; Nakahara, K.; Tchorz, K.; Kirsch, I. R.: A comparative structural characterization of the human NSCL-1 and NSCL-2 genes: two basic helix-loop-helix genes expressed in the developing nervous system. *J. Biol. Chem.* 267: 21065–21071, 1992.

[78491] 16821. Mullick, A.; Groulx, N.; Trasler, D.; Gros, P.: *Nhlh1*, a basic helix-loop-helix transcription factor, is very tightly linked to the mouse looptail (Lp) mutation. *Mammalian Genome* 6: 700–704, 1995.

[78492] 16822. Meeker, T. C.; Nagarajan, L.; ar-Rushdi, A.; Rovera, G.; Huebner, K.; Croce, C. M.: Characterization of the hu-

man PIM-1 gene: a putative proto-oncogene coding for a tissue specific member of the protein kinase family. *Oncogene Res.* 1: 87-101, 1987.

- [78493] 16823. Pasqualucci, L.; Neumeister, P.; Goossens, T.; Nangud, G.; Chaganti, R. S. K.; Kuppers, R.; Dalla-Favera, R.: Hypermutation of multiple proto-oncogenes in B-cell diffuse large-cell lymphomas. *Nature* 412:341-346, 2001.
- [78494] 16824. Ragoussis, J.; Senger, G.; Mockridge, I.; Sanseau, P.; Ruddy, S.; Dudley, K.; Sheer, D.; Trowsdale, J.: A testis-expressed Zn finger gene (ZNF76) in human 6p21.3 centromeric to the MHC is closely linked to the human homolog of the t-complex gene *tcp-11*. *Genomics* 14:673-679, 1992.
- [78495] 16825. Saris, C. J. M.; Domen, J.; Berns, A.: The *pim-1* oncogene encodes two related protein-serine/threonine kinases by alternative initiation at AUG and CUG. *EMBO J.* 10: 655-664, 1991.
- [78496] 16826. Selten, G.; Cuypers, H. T.; Boelens, W.; Robanus-Maandag, E.; Verbeek, J.; Domen, J.; van Beveren, C.; Berns, A.: The primary structure of the putative oncogene *pim-1* shows extensive homology with protein kinases. *Cell* 46: 603-611, 1986.
- [78497] 16827. Zoghbi, H. Y.; Ballantyne, C. M.; O'Brien, W. E.; Mc-



Call, A. E.; Kwiatkowski, T. J., Jr.; Ledbetter, S. A.; Beaudet, A. L.: Deletion and linkage mapping of eight markers from the proximal short arm of chromosome 6. *Genomics* 6: 352–357, 1990.

- [78498] 16828. Chapin, S. J.; Lue, C.-M.; Yu, M. T.; Bulinski, J. C.: Differential expression of alternatively spliced forms of MAP4: a repertoire of structurally different microtubule-binding domains. *Biochemistry* 34:2289–2301, 1995.
- [78499] 16829. Laird, P. W.; van der Lugt, N. M. T.; Clarke, A.; Domen, J.; Linders, K.; McWhir, J.; Berns, A.; Hooper, M.: In vivo analysis of Pim-1 deficiency. *Nucleic Acids Res.* 21: 4750–4755, 1993.
- [78500] 16830. Gartner, J.; Obie, C.; Watkins, P.; Valle, D.: Restoration of peroxisome biogenesis in a peroxisome-deficient mammalian cell line by expression of either the 35 kDa or the 70 kDa peroxisomal membrane proteins. *J. Inher. Metab. Dis.* 17: 327–329, 1994.
- [78501] 16831. Maric, S. C.; Crozat, A.; Janne, O. A.: Structure and organization of the human S-adenosylmethionine decarboxylase gene. *J. Biol. Chem.* 267:18915–18923, 1992.
- [78502] 16832. Maric, S. C.; Crozat, A.; Louhimo, J.; Knuutila, S.; Janne, O. A.: The human S-adenosylmethionine decarboxylase gene: nucleotide sequence of a pseudogene and

chromosomal localization of the activegene (AMD1) and the pseudogene (AMD2). Cytogenet. Cell Genet. 70:195–199, 1995.

- [78503] 16833.Mangan, M. E.; Olmsted, J. B.: The gene for micro-tubule-associatedprotein 4 (Mtap4) maps to the distal re-gion of mouse chromosome 9. MammalianGenome 7: 918–925, 1996.
- [78504] 16834.West, R. R.; Tenbarger, K. M.; Olmsted, J. B.: A model for microtubule-associatedprotein 4 structure: do-mains defined by comparisons of human, mouse,and bovine sequences. J. Biol. Chem. 266: 21886–21896, 1991.
- [78505] 16835.Abel, K. J.; Boehnke, M.; Prahalad, M.; Ho, P.; Fle-jter, W. L.;Watkins, M.; VanderStoep, J.; Chan-drasekharappa, S. C.; Collins, F.S.; Glover, T. W.; Weber, B. L.: A radiation hybrid map of the BRCA1region of chromo-some 17q12–q21. Genomics 17: 632–641, 1993.
- [78506] 16836.Alonso, A. D. C.; Grundke-Iqbal, I.; Iqbal, K.: Alzheimer's diseas hyperphosphorylated tau sequesters normal tau into tangles of filamentsand disassembles mi-crotubules. Nature Med. 2: 783–787, 1996.
- [78507] 16837.Arima, K.; Kowalska, A.; Hasegawa, M.; Mukoyama, M.; Watanabe,R.; Kawai, M.; Takahashi, K.; Iwatsubo, T.;

Tabira, T.; Sunohara, N.: Two brothers with frontotemporal dementia and parkinsonism with an N279K mutation of the tau gene. *Neurology* 54: 1787–1795, 2000.

[78508] 16838. Baker, M.; Litvan, I.; Houlden, H.; Adamson, J.; Dickson, D.; Perez-Tur, J.; Hardy, J.; Lynch, T.; Bigio, E.; Hutton, M.: Association of an extended haplotype in the tau gene with progressive supranuclear palsy. *Hum. Molec. Genet.* 8: 711–715, 1999.

[78509] 16839. Brown, J.; Lantos, P. L.; Roques, P.; Fidani, L.; Rossor, M. N.: Familial dementia with swollen achromatic neurons and corticobasal inclusion bodies: a clinical and pathological study. *J. Neurol. Sci.* 135:21–30, 1996.

[78510] 16840. Clark, L. N.; Poorkaj, P.; Wszolek, Z.; Geschwind, D. H.; Nasreddine, Z. S.; Miller, B.; Li, D.; Payami, H.; Awert, F.; Markopoulou, K.; Andreadis, A.; D'Souza, I.; Lee, V. M.-Y.; Reed, L.; Trojanowski, J. Q.; Zhukareva, V.; Bird, T.; Schellenberg, G.; Wilhelmsen, K. C.: Pathogenic implications of mutations in the tau gene in pallido-ponto-nigral degeneration and related neurodegenerative disorders linked to chromosome 17. *Proc. Nat. Acad. Sci.* 95: 13103–13107, 1998.

[78511] 16841. Connell, J. W.; Gibb, G. M.; Betts, J. C.; Blackstock, W. P.; Gallo, J.-M.; Lovestone, S.; Hutton, M.; Anderton, B.

H.: Effects of FTDP-17 mutations on the in vitro phosphorylation of tau by glycogen synthase kinase 3-beta identified by mass spectrometry demonstrate certain mutations exert long-range conformational changes. FEBS Lett. 493:40-44, 2001.

[78512] 16842. Conrad, C.; Andreadis, A.; Trojanowski, J. Q.; Dickson, D. W.; Kang, D.; Chen, X.; Weiderholt, W.; Hansen, L.; Masliah, E.; Thal, L. J.; Katzman, R.; Xia, Y.; Saitoh, T.: Genetic evidence for the involvement of tau in progressive supranuclear palsy. Ann. Neurol. 41: 277-281, 1997.

[78513] 16843. Conrad, C.; Vianna, C.; Freeman, M.; Davies, P.: A polymorphic gene nested within an intron of the tau gene: implications for Alzheimer's disease. Proc. Nat. Acad. Sci. 99: 7751-7756, 2002.

[78514] 16844. Delisle, M. B.; Murrell, J. R.; Richardson, R.; Trofatter, J. A.; Rascol, O.; Soulages, X.; Mohr, M.; Calvas, P.; Ghetti, B.: A mutation at codon 279 (N279K) in exon 10 of the tau gene causes a tauopathy with dementia and supranuclear palsy. Acta Neuropath. 98:62-77, 1999.

[78515] 16845. Donlon, T. A.; Harris, P.; Neve, R. L.: Localization of microtubule-associated protein tau (MTBT1) to chromosome 17q21. (Abstract) Cytogenet. Cell Genet. 46: 607, 1987.

- [78516] 16846.Goedert, M.; Crowther, R. A.; Spillantini, M. G.: Tau mutations cause frontotemporal dementias. *Neuron* 21: 955–958, 1998.
- [78517] 16847.Goedert, M.; Spillantini, M. G.; Potier, M. C.; Ulrich, J.; Crowther, R. A.: Cloning and sequencing of the cDNA encoding an isoform of microtubule-associated protein tau containing four tandem repeats: differential expression of tau protein mRNAs in human brain. *EMBO J.* 8: 393–399, 1989.
- [78518] 16848.Goedert, M.; Wischik, C. M.; Crowther, R. A.; Walker, J. E.; Klug, A.: Cloning and sequencing of the cDNA encoding a core protein of the paired helical filament of Alzheimer disease: identification as the microtubule-associated protein tau. *Proc. Nat. Acad. Sci.* 85:4051–4055, 1988.
- [78519] 16849.Goode, B. L.; Chau, M.; Denis, P. E.; Feinstein, S. C.: Structural and functional differences between 3-repeat and 4-repeat tau isoforms: implications for normal tau function and the onset of neurodegenerative disease. *J. Biol. Chem.* 275: 38182–38189, 2000.
- [78520] 16850.Heutink, P.: Untangling tau-related dementia. *Hum. Molec. Genet.* 9:979–986, 2000.
- [78521] 16851.Heutink, P.; Stevens, M.; Rizzu, P.; Bakker, E.; Kros,

J. M.;Tibben, A.; Niermeijer, M. F.; van Duijn, C. M.; Oos-  
tra, B. A.; vanSwieten, J. C.: Hereditary frontotemporal de-  
mentia is linked to chromosome17q21-q22: a genetic and  
clinicopathological study of three Dutchfamilies. Ann.  
Neurol. 41: 150-159, 1997.

- [78522] 16852.Hiesberger, T.; Trommsdorff, M.; Howell, B. W.;  
Goffinet, A.;Mumby, M. C.; Cooper, J. A.; Herz, J.: Direct  
binding of reelin toVLDL receptor and apoE receptor 2 in-  
duces tyrosine phosphorylationof disabled-1 and modu-  
lates tau phosphorylation. Neuron 24: 481-489,1999.
- [78523] 16853.Hong, M.; Zhukareva, V.; Vogelsberg-Ragaglia, V.;  
Wszolek, Z.;Reed, L.; Miller, B. I.; Geschwind, D. H.; Bird, T.  
D.; McKeel, D.;Goate, A.; Morris, J. C.; Wilhelmsen, K. C.;  
Schellenberg, G. D.;Trojanowski, J. Q.; Lee, V. M.-Y.: Mu-  
tation-specific functional impairmentsin distinct tau iso-  
forms of hereditary FTDP-17. Science 282:  
1914-1917,1998.
- [78524] 16854.Hutton, M.: Missense and splice site mutations in  
tau associatedwith FTDP-17: multiple pathogenic mecha-  
nisms. Neurology 56 (suppl.4): S21-S25, 2001.
- [78525] 16855.Hutton, M.; Lendon, C. L.; Rizzu, P.; Baker, M.;  
Froelich, S.;Houlden, H.; Pickering-Brown, S.; Chakraverty,  
S.; Isaacs, A.; Grover,A.; Hackett, J.; Adamson, J.; and 39

others: Association of missense and 5-prime-splice-site mutations in tau with the inherited dementia FTDP-17. *Nature* 393: 702–705, 1998.

[78526] 16856. Iijima, M.; Tabira, T.; Poorkaj, P.; Schellenberg, G. D.; Trojanowski, J. Q.; Lee, V. M.; Schmidt, M. L.; Takahashi, K.; Nabika, T.; Matsumoto, T.; Yamashita, Y.; Yoshioka, S.; Ishino, H.: A distinct familial presenile dementia with a novel missense mutation in the tau gene. *Neuroreport* 10:497–501, 1999.

[78527] 16857. Ishihara, T.; Hong, M.; Zhang, B.; Nakagawa, Y.; Lee, M. K.; Trojanowski, J. Q.; Lee, V. M.-Y.: Age-dependent emergence and progression of a tauopathy in transgenic mice overexpressing the shortest human tau isoform. *Neuron* 24: 751–762, 1999.

[78528] 16858. Janssen, J. C.; Warrington, E. K.; Morris, H. R.; Lantos, P.; Brown, J.; Revesz, T.; Wood, N.; Khan, M. N.; Cipolotti, L.; Fox, N. C.; Rossor, M. N.: Clinical features of frontotemporal dementia due to the intronic tau 10 + 16 mutation. *Neurology* 58: 1161–1168, 2002.

[78529] 16859. Bergstrom, D. A.; Penn, B. H.; Strand, A.; Perry, R. L. S.; Rudnicki, M. A.; Tapscott, S. J.: Promoter-specific regulation of MyoD binding and signal transduction cooperate to pattern gene expression. *Mol. Cell* 9: 587–600,

2002.

- [78530] 16860. Braun, T.; Grzeschik, K.-H.; Bober, E.; Arnold, H.-H.: The MYF genes, a group of human muscle determining factors, are localized on different human chromosomes. (Abstract) Cytogenet. Cell Genet. 51:969 only, 1989.
- [78531] 16861. Davis, R. L.; Weintraub, H.; Lassar, A. B.: Expression of a single transfected cDNA converts fibroblasts to myoblasts. Cell 51: 987–1000, 1987.
- [78532] 16862. de la Serna, I. L.; Carlson, K. A.; Imbalzano, A. N.: Mammalian SWI/SNF complexes promote MyoD-mediated muscle differentiation. Nature Genet. 27: 187–190, 2001.
- [78533] 16863. Gessler, M.; Hameister, H.; Henry, I.; Junien, C.; Braun, T.; Arnold, H. H.: The human MyoD1 (MYF3) gene maps on the short arm of chromosome 11 but is not associated with the WAGR locus or the region for the Beckwith-Wiedemann syndrome. Hum. Genet. 86: 135–138, 1990.
- [78534] 16864. Guttridge, D. C.; Mayo, M. W.; Madrid, L. V.; Wang, C.-Y.; Baldwin, A. S, Jr.: NF- $\kappa$ B-induced loss of MyoD messenger RNA: possible role in muscle decay and cachexia. Science 289: 2363–2366, 2000.
- [78535] 16865. Henry, I.; Puech, A.; Antignac, C.; Couillin, P.; Jeanpierre, M.; Ahnine, L.; Barichard, F.; Boehm, T.; Augereau,



P.; Scrable, H.; Rabbitts, T. H.; Rochefort, H.; Cavenee, W.; Junien, C.: Subregional mapping of BWS, CTSD, MYOD1, and a T-ALL breakpoint in 11p15. (Abstract) Cytogenet. Cell Genet. 51: 1013 only, 1989.

[78536] 16866. Kim, Y.-J.; Noguchi, S.; Hayashi, Y. K.; Tsukahara, T.; Shimizu, T.; Arahata, K.: The product of an oculopharyngeal muscular dystrophy gene, poly(A)-binding protein 2, interacts with SKIP and stimulates muscle-specific gene expression. Hum. Molec. Genet. 10: 1129-1139, 2001.

[78537] 16867. Olson, E. N.: MyoD family: a paradigm for development. Genes Dev. 4: 1454-1461, 1990.

[78538] 16868. Sartorelli, V.; Puri, P. L.; Hamamori, Y.; Ogryzko, V.; Chung, G.; Nakatani, Y.; Wang, J. Y. J.; Kedes, L.: Acetylation of MyoD directed by PCAF is necessary for the execution of the muscle program. Molec. Cell 4: 725-734, 1999.

[78539] 16869. Tapscott, S. J.; Davis, R. L.; Thayer, M. J.; Cheng, P. F.; Weintraub, H.; Lassar, A. B.: MyoD1: a nuclear phosphoprotein requiring a Myc homology region to convert fibroblasts to myoblasts. Science 242: 405-411, 1988.

[78540] 16870. Bonner, T. I.; Young, A. C.; de Miguel, C.; Detera-Wadleigh, S.; Modi, W. S.; O'Brien, S. J.: The chromosomal location of the two human tachykinin (sic) genes: human substance P and neurokinin K. (Abstract) Cytogenet. Cell

Genet. 46: 584 only, 1987.

- [78541] 16871.Cao, Y. Q.; Mantyh, P. W.; Carlson, E. J.; Gillespie, A.-M.; Epstein,C. J.; Basbaum, A. I.: Primary afferent tachykinins are required to experience moderate to intense pain. Nature 392: 390–393, 1998.
- [78542] 16872.Krause, J. E.; Chirgwin, J. M.; Carter, M. S.; Xu, Z. S.; Hershey,A. D.: Three rat preprotachykinin mRNAs encode the neuropeptidessubstance P and neurokinin A. Proc. Nat. Acad. Sci. 84: 881–885,1987.
- [78543] 16873.Liu, H.; Cao, Y.; Basbaum, A. I.; Mazarati, A. M; Sankar, R.; Wasterlain,C. G.: Resistance to excitotoxin–induced seizures and neuronal deathin mice lacking the preprotachykinin A gene. Proc. Nat. Acad. Sci. 96:12096–12101, 1999.
- [78544] 16874.Troger, J.; Neyer, S.; Heufler, C.; Huemer, H.; Schmid, E.; Griesser,U.; Kralinger, M.; Kremser, B.; Baldissera, I.; Kieselbach, G.: SubstanceP and vasoactive intestinal polypeptide in the streptozotocin–induceddiabetic rat retina. Invest. Ophthal. Vis. Sci. 42: 1045–1050, 2001.
- [78545] 16875.Zimmer, A.; Zimmer, A. M.; Baffi, J.; Usdin, T.; Reynolds, K.;Konig, M.; Palkovits, M.; Mezey, E.: Hypoalgesia in mice with a targeteddeletion of the tachykinin 1 gene. Proc. Nat. Acad. Sci. 95: 2630–2635,1998.

- [78546] 16876.Gregory, C. A.; Schwartz, J. S.: The cDNA of the human neuromedinB gene (NMB) mapped to 15q11-qter recognizes an XbaI RFLP. *NucleicAcids Res.* 19: 1167 only, 1991.
- [78547] 16877.Krane, I. M.; Naylor, S. L.; Helin-Davis, D.; Chin, W. W.; Spindel,E. R.: Molecular cloning of cDNAs encoding the human bombesin-likepeptide neuromedin B: chromosomal localization and comparison to cDNAsencoding its amphibian homolog ranatensin. *J. Biol. Chem.* 263:13317-13323, 1988. Note: Erratum: *J. Biol. Chem.* 265: 7091 only, 1990.
- [78548] 16878.Mattei, M.-G.; Riviere, M.; Krust, A.; Ingvarsson, S.; Vennstrom,B.; Islam, M. Q.; Levan, G.; Kautner, P.; Zelent, A.; Chambon, P.;Szpirer, J.; Szpirer, C.: Chromosomal assignment of retinoic acidreceptor (RAR) genes in the human, mouse, and rat genomes. *Genomics* 10:1061-1069, 1991.
- [78549] 16879.Abramson, D. H.; Ellsworth, R. M.; Zimmerman, L. E.: Monocularcancer in retinoblastoma survivors. *Trans. Am. Acad. Ophthal. Otolaryng.* 81:454-457, 1976.
- [78550] 16880.Aherne, G. E. S.; Roberts, D. F.: Retinoblastoma--a clinical surveyand its genetic implications. *Clin. Genet.* 8: 275-290, 1975.

- [78551] 16881.Alonso, J.; Garcia-Miguel, P.; Abelairas, J.; Mendiola, M.; Sarret,E.; Vendrell, M. T.; Navajas, A.; Pestana, A.: Spectrum of germlineRB1 gene mutations in Spanish retinoblastoma patients: phenotypicand molecular epidemiological implications. Hum. Mutat. 17: 412–422,2001.
- [78552] 16882.Amoaku, W. M. K.; Willshaw, H. E.; Parkes, S. E.; Shah, K. J.;Mann, J. R.: Trilateral retinoblastoma: a report of five patients. Cancer 78:858–863, 1996.
- [78553] 16883.Bader, J. L.; Meadows, A. T.; Zimmerman, L. E.; Rorke, L. B.; Voute,P. A.; Champion, L. A. A.; Miller, R. W.: Bilateral retinoblastomawith ectopic intracranial retinoblastoma: trilateral retinoblastoma. CancerGenet. Cytogenet. 5: 203–213, 1982.
- [78554] 16884.Balaban–Malenbaum, G.; Gilbert, F.; Nichols, W. W.; Hill, R.; Shields,J.; Meadows, A. T.: A deleted chromosome no. 13 in human retinoblastomacells: relevance to tumorigenesis. Cancer Genet. Cytogenet. 3: 243–250,1981.
- [78555] 16885.Bandara, L. R.; Adamczewski, J. P.; Hunt, T.; La Thangue, N. B.: Cyclin A and the retinoblastoma gene product complex with a commontranscription factor. Nature 352: 249–251, 1991.
- [78556] 16886.Benedict, W. F.; Murphree, A. L.; Banerjee, A.;

Spina, C. A.; Sparkes, M. C.; Sparkes, R. S.: Patient with 13 chromosome deletion: evidence that the retinoblastoma gene is a recessive cancer gene. *Science* 219:973–975, 1983.

[78557] 16887. Benedict, W. F.; Xu, H.-J.; Hu, S.-X.; Takahashi, R.: Role of the retinoblastoma gene in the initiation and progression of human cancer. *J. Clin. Invest.* 85: 988–993, 1990.

[78558] 16888. Bia, B.; Cowell, J. K.: Independent constitutional germline mutations occurring in the RB1 gene in cousins with bilateral retinoblastoma. *Oncogene* 11:977–979, 1995.

[78559] 16889. Blanquet, V.; Turleau, C.; de Grouchy, J.; Creau-Goldberg, N.: Physical map around the retinoblastoma gene: possible genomic imprinting suggested by Nrul digestion. *Genomics* 10: 350–355, 1991.

[78560] 16890. Blanquet, V.; Turleau, C.; Gross-Morand, M. S.; Senamaud-Beaufort, C.; Doz, F.; Besmond, C.: Spectrum of germline mutations in the RB1 gene: a study of 232 patients with hereditary and non hereditary retinoblastoma. *Hum. Molec. Genet.* 4: 383–388, 1995.

[78561] 16891. Jiang, Z.; Cote, J.; Kwon, J. M.; Goate, A. M.; Wu, J. Y.: Aberrant splicing of tau pre-mRNA caused by intronic

mutations associated with the inherited dementia from  
to temporal dementia with Parkinsonism linked to chromo-  
some 17. *Molec. Cell. Biol.* 20: 4036–4048, 2000.

[78562] 16892. Tapscott, S. J.; Weintraub, H.: MyoD and the regu-  
lation of myogenesis by helix–loop–helix proteins. *J. Clin.*  
*Invest.* 87: 1133–1138, 1991.

[78563] 16893. Futterweit, W.; Ritch, R.; Teekhasaene, C.; Nelson,  
E. S.: Coexistence of Prader–Willi syndrome, congenital ec-  
tropion uveae with glaucoma, and factor XI deficiency.  
*J.A.M.A.* 255: 3280–3282, 1986.

[78564] 16894. Trapani, J. A.; Browne, K. A.; Dawson, M. J.; Ram-  
say, R. G.; Eddy, R. L.; Shows, T. B.; White, P. C.; Dupont,  
B.: A novel gene constitutively expressed in human lym-  
phoid cells is inducible with interferon–gamma in myeloid  
cells. *Immunogenetics* 36: 369–376, 1992.

[78565] 16895. Bowcock, A. M.; Kidd, J. R.; Lathrop, G. M.; Danesh-  
var, L.; May, L. T.; Ray, A.; Sehgal, P. B.; Kidd, K. K.; Cav-  
alli-Sforza, L. L.: The human 'interferon-  
beta-2/hepatocyte stimulating factor/interleukin-6' gene:  
DNA polymorphism studies and localization to chromo-  
some 7p21. *Genomics* 3:8–16, 1988.

[78566] 16896. Chen, Y.; Ferguson-Smith, A. C.; Newman, M. S.;  
May, L. T.; Sehgal, P. B.; Ruddle, F. H.: Regional localization

of the human beta 2–interferongene. (Abstract) Am. J. Hum. Genet. 41: A161, 1987.

[78567] 16897. Chow, D.; He, X.; Snow, A. L.; Rose–John, S.; Garcia, K. C.: Structure of an extracellular gp130 cytokine receptor signaling complex. Science 291:2150–2155, 2001.

[78568] 16898. Chung, U.; Tanaka, Y.; Fujita, T.: Association of interleukin–6 and hypoaldosteronism in patients with cancer. (Letter) New Eng. J. Med. 334: 473, 1996.

[78569] 16899. Cressman, D. E.; Greenbaum, L. E.; DeAngelis, R. A.; Ciliberto, G.; Furth, E. E.; Poli, V.; Taub, R.: Liver failure and defective hepatocyte regeneration in interleukin–6–deficient mice. Science 274:1379–1382, 1996.

[78570] 16900. De Benedetti, F.; Alonzi, T.; Moretta, A.; Lazzaro, D.; Costa, P.; Poli, V.; Martini, A.; Ciliberto, G.; Fattori, E.: Interleukin6 causes growth impairment in transgenic mice through a decrease in insulin–like growth factor–I. J. Clin. Invest. 99: 643–650, 1997.

[78571] 16901. Ferguson–Smith, A. C.; Chen, Y.–F.; Newman, M. S.; May, L. T.; Sehgal, P. B.; Ruddle, F. H.: Regional localization of the interferon beta–2/B–cell stimulatory factor 2/hepatocyte stimulating factor gene to human chromosome 7p15–p21. Genomics 2: 203–208, 1988.

[78572] 16902. Fernandez–Real, J.–M.; Broch, M.; Vendrell, J.;

Richart, C.; Ricart,W.: Interleukin-6 gene polymorphism and lipid abnormalities in healthysubjects. J. Clin. Endocr. Metab. 85: 1334-1339, 2000.

[78573] 16903.Fishman, D.; Faulds, G.; Jeffery, R.; Mohamed-Ali, V.; Yudkin,J. S.; Humphries, S.; Woo, P.: The effect of novel polymorphismsin the interleukin-6 (IL-6) gene on IL-6 transcription and plasmall-6 levels, and an association with systemic-onset juvenile chronicarthritis. J. Clin. Invest. 102: 1369-1376, 1998.

[78574] 16904.Foster, C. B.; Lehrnbecher, T.; Samuels, S.; Stein, S.; Mol, F.;Metcalf, J. A.; Wyvill, K.; Steinberg, S. M.; Kovacs, J.; Blauvelt,A.; Yarchoan, R.; Chanock, S. J.: An IL6 promoter polymorphism isassociated with a lifetime risk of development of Kaposi sarcoma inmen infected with human immunodeficiency virus. Blood 96: 2562-2567,2000.

[78575] 16905.Funatsu, H.; Yamashita, H.; Noma, H.; Mimura, T.; Yamashita, T.;Hori, S.: Increased levels of vascular endothelial growth factorand interleukin-6 in the aqueous humor of diabetics with macular edema. Am.J. Ophthal. 133: 70-77, 2002.

[78576] 16906.Hirano, T.; Yasukawa, K.; Harada, H.; Taga, T.; Watanabe, Y.;Matsuda, T.; Kashiwamura, S.; Nakajima, K.; Koyama, K.; Iwamatsu,A.; Tsunasawa, S.; Sakiyama, F.;



Matsui, H.; Takahara, Y.; Taniguchi, T.; Kishimoto, T.:  
Complementary DNA for a novel human interleukin (BSF-2)  
that induces B lymphocytes to produce immunoglobulin.  
Nature 324:73–76, 1986.

- [78577] 16907. Kawano, M.; Hirano, T.; Matsuda, T.; Taga, T.;  
Horii, Y.; Iwato, K.; Asaoku, H.; Tang, B.; Tanabe, O.;  
Tanaka, H.; Kuramoto, A.; Kishimoto, T.: Autocrine gener-  
ation and requirement of BSF-2/IL-6 for human multiple  
myelomas. Nature 332: 83–85, 1988.
- [78578] 16908. Kovalchuk, A. L.; Kim, J. S.; Park, S. S.; Coleman, A.  
E.; Ward, J. M.; Morse, H. C, III; Kishimoto, T.; Potter, M.;  
Janz, S.: IL-6 transgenic mouse model for extraosseous  
plasmacytoma. Proc. Nat. Acad. Sci. 99: 1509–1514, 2002.
- [78579] 16909. May, L. T.; Ghrayeb, J.; Santhanam, U.; Tatter, S. B.;  
Sthoeger, Z.; Helfgott, D. C.; Chiorazzi, N.; Grieninger, G.;  
Sehgal, P. B.: Synthesis and secretion of multiple forms of  
beta-2-interferon/B-cell differentiation factor  
2/hepatocyte-stimulating factor by human fibroblasts and  
monocytes. J. Biol. Chem. 263: 7760–7766, 1988.
- [78580] 16910. Ota, N.; Hunt, S. C.; Nakajima, T.; Suzuki, T.; Hosoi,  
T.; Orimo, H.; Shirai, Y.; Emi, M.: Linkage of interleukin 6  
locus to human osteopenia by sibling pair analysis. Hum.  
Genet. 105: 253–257, 1999.

- [78581] 16911.Ota, N.; Nakajima, T.; Nakazawa, I.; Suzuki, T.; Hosoi, T.; Orimo,H.; Inoue, S.; Shirai, Y.; Emi, M.: A nucleotide variant in the promoterregion of the interleukin-6 gene associated with decreased bone mineraldensity. J. Hum. Genet. 46: 267-272, 2001.
- [78582] 16912.Redwine, L.; Hauger, R. L.; Gillin, J. C.; Irwin, M.: Effectsof sleep and sleep deprivation on interleukin-6, growth hormone, cortisol,and melatonin levels in humans. J. Clin. Endocr. Metab. 85: 3597-3603,2000.
- [78583] 16913.Roodman, G. D.; Kurihara, N.; Ohsaki, Y.; Kukita, A.; Hosking,D.; Demulder, A.; Smith, J. F.; Singer, F. R.: Interleukin 6: a potentialautocrine/paracrine factor in Paget's disease of bone. J. Clin. Invest. 89:46-52, 1992.
- [78584] 16914.Rooney, M.; David, J.; Symons, J.; Di Giovine, F.; Varsani, H.;Woo, P.: Inflammatory cytokine responses in juvenile chronic arthritis. Brit.J. Rheum. 34: 454-460, 1995.
- [78585] 16915.Santhanam, U.; Ray, A.; Sehgal, P. B.: Repression of the interleukin6 gene promoter by p53 and the retinoblastoma susceptibility geneproduct. Proc. Nat. Acad. Sci. 88: 7605-7609, 1991.
- [78586] 16916.Scheidt-Nave, C.; Bismar, H.; Leidig-Bruckner, G.; Woitge, H.;Seibel, M. J.; Ziegler, R.; Pfeilschifter, J.: Serum

interleukin6 is a major predictor of bone loss in women specific to the firstdecade past menopause. J. Clin. Endocr. Metab. 86: 2032–2042, 2001.

[78587] 16917.Sehgal, P. B.; May, L. T.; Tamm, I.; Vilcek, J.: Human beta-2interferon and B-cell differentiation factor BSF-2 are identical. Science 235:731–732, 1987.

[78588] 16918.Sehgal, P. B.; Walther, Z.; Tamm, I.: Rapid enhancement of beta(2)-interferon/B-celldifferentiation factor BSF-2 gene expression in human fibroblastsby diacylglycerols and the calcium ionophore A23187. Proc. Nat. Acad.Sci. 84: 3663–3667, 1987.

[78589] 16919.Sehgal, P. B.; Zilberstein, A.; Ruggieri, R.–M.; May, L. T.; Ferguson–Smith,A.; Slate, D. L.; Revel, M.; Ruddle, F. H.: Human chromosome 7 carriesthe beta-2 interferon gene. Proc. Nat. Acad. Sci. 83: 5219–5222,1986.

[78590] 16920.Lewis, J.; McGowan, E.; Rockwood, J.; Melrose, H.; Nacharaju,P.; Van Slegtenhorst, M.; Gwinn–Hardy, K.; Murphy, M. P.; Baker, M.;Yu, X.; Duff, K.; Hardy, J.; Corral, A.; Lin, W.–L.; Yen, S.–H.; Dickson,D. W.; Davies, P.; Hutton, M.: Neurofibrillary tangles, amyotrophyand progressive motor disturbance in mice expressing mutant (P301L)tau protein. Nature Genet. 25: 402–405, 2000. Note: Erratum: NatureGenet. 26: 127 only, 2000.

- [78591] 16921.Lippa, C. F.; Zhukareva, V.; Kawarai, T.; Uryu, K.; Shafiq, M.;Nee, L. E.; Grafman, J.; Liang, Y.; St George-Hyslop, P. H.; Trojanowski,J. Q.; Lee, V. M.-Y.: Frontotemporal dementia with novel tau pathologyand a glu342val tau mutation. *Ann. Neurol.* 48: 850–858, 2000.
- [78592] 16922.Litvan, I.; Baker, M.; Hutton, M.: Tau genotype: no effect ononset, symptom severity, or survival in progressive supranuclear palsy. *Neurology* 57:138–140, 2001.
- [78593] 16923.Martin, E. R.; Scott, W. K.; Nance, M. A.; Watts, R. L.; Hubble,J. P.; Koller, W. C.; Lyons, K.; Pahwa, R.; Stern, M. B.; Colcher,A.; Hiner, B. C.; Jankovic, J.; and 20 others: Association of single-nucleotidepolymorphisms of the Tau gene with late-onset Parkinson disease. *J.A.M.A.* 286:2245–2250, 2001.
- [78594] 16924.Moser, A. B.; Rasmussen, M.; Naidu, S.; Watkins, P. A.; McGuinness,M.; Hajra, A. K.; Chen, G.; Raymond, G.; Liu, A.; Gordon, D.; Garnaas,K.; Walton, D. S.; Skjedal, O. H.; Guggenheim, M. A.; Jackson, L.G.; Elias, E. R.; Moser, H. W.: Phenotype of patients with peroxisomaldisorders subdivided into sixteen complementation groups. *J. Pe-diat.* 127:13–22, 1995.
- [78595] 16925.Paton, B. C.; Heron, S. E.; Nelson, P. V.; Morris, C. P.; Poulos,A.: Absence of mutations raises doubts about

the role of the 70-kD peroxisomal membrane protein in Zellweger syndrome. (Letter) *Am. J. Hum. Genet.* 60: 1535–1539, 1997.

- [78596] 16926. Aljada, A.; Ghanim, H.; Friedman, J.; Garg, R.; Mohanty, P.; Dandona, P.: Troglitazone reduces the expression of PPAR- $\gamma$  while stimulating that of PPAR- $\alpha$  in mononuclear cells in obese subjects. *J. Clin. Endocr. Metab.* 86: 3130–3133, 2001.
- [78597] 16927. Costet, P.; Legendre, C.; More, J.; Edgar, A.; Galtier, P.; Pineau, T.: Peroxisome proliferator-activated receptor  $\alpha$ -isoform deficiency leads to progressive dyslipidemia with sexually dimorphic obesity and steatosis. *J. Biol. Chem.* 273: 29577–29585, 1998.
- [78598] 16928. Djouadi, F.; Weinheimer, C. J.; Saffitz, J. E.; Pitchford, C.; Bastin, J.; Gonzalez, F. J.; Kelly, D. P.: A gender-related defect in lipid metabolism and glucose homeostasis in peroxisome proliferator-activated receptor  $\alpha$ -deficient mice. *J. Clin. Invest.* 102: 1083–1091, 1998.
- [78599] 16929. Kersten, S.; Desvergne, B.; Wahli, W.: Roles of PPARs in health and disease. *Nature* 405: 421–424, 2000.
- [78600] 16930. Kersten, S.; Seydoux, J.; Peters, J. M.; Gonzalez, F. J.; Desvergne, B.; Wahli, W.: Peroxisome proliferator-activated receptor  $\alpha$  mediates the adaptive response to

fasting. *J. Clin. Invest.* 103: 1489–1498,1999.

[78601] 16931.Lee, S. S.–T.; Pineau, T.; Drago, J.; Lee, E. J.; Owens, J. W.;Kroetz, D. L.; Fernandez–Salguero, P. M.; Westphal, H.; Gonzalez,F. J.: Targeted disruption of the alpha iso–form of the peroxisomeproliferator–activated receptor gene in mice results in abolishmentof the pleiotropic effects of peroxisome proliferators. *Molec. Cell.Biol.* 15: 3012–3022, 1995.

[78602] 16932.Lee, Y.; Yu, X.; Gonzales, F.; Mangelsdorf, D. J.; Wang, M.–Y.;Richardson, C.; Witters, L. A.; Unger, R. H.: PPAR–alpha is necessaryfor the lipopenic action of hyper–leptinemia on white adipose and livertissue. *Proc. Nat. Acad. Sci.* 99: 11848–11853, 2002.

[78603] 16933.Leone, T. C.; Weinheimer, C. J.; Kelly, D. P.: A critical rolefor the peroxisome proliferator–activated receptor alpha (PPAR–alpha)in the cellular fasting response: the PPAR–alpha–null mouse as a modelof fatty acid oxidation disorders. *Proc. Nat. Acad. Sci.* 96: 7473–7478,1999.

[78604] 16934.Michalik, L.; Desvergne, B.; Tan, N. S.; Basu–Modak, S.; Escher,P.; Rieusset, J.; Peters, J. M.; Kaya, G.; Gonzalez, F. J.; Zakany,J.; Metzger, D.; Chambon, P.; Duboule, D.; Wahli, W.: Impaired skinwound healing in peroxisome proliferator–activated receptor (PPAR)–alphaand PPAR–

beta mutant mice. *J. Cell Biol.* 154: 799–814, 2001.

- [78605] 16935.Sher, T.; Yi, H.–F.; McBride, O. W.; Gonzalez, F. J.: cDNA cloning,chromosomal mapping, and functional characterization of the humanperoxisome proliferator activated receptor. *Biochemistry* 32: 5598–5604,1993.
- [78606] 16936.Xu, H. E.; Stanley, T. B.; Montana, V. G.; Lambert, M. H.; Shearer,B. G.; Cobb, J. E.; McKee, D. D.; Galardi, C. M.; Plunket, K. D.;Nolte, R. T.; Parks, D. J.; Moore, J. T.; Kliewer, S. A.; Willson,T. M.; Stimmel, J. B.: Structural basis for antagonist–mediated recruitmentof nuclear co–repressors by PPAR–alpha. *Nature* 415: 813–817, 2002.
- [78607] 16937.Ames, G. F.–L.: The basis of multidrug resistance in mammaliancells: homology with bacterial transport. *Cell* 47: 323–324, 1986.
- [78608] 16938.Baldini, N.; Scotlandi, K.; Barbanti–Brodano, G.; Manara, M. C.;Maurici, D.; Bacci, G.; Bertoni, F.; Picci, P.; Sottili, S.; Campanacci,M.; Serra, M.: Expression of P–glycoprotein in high–grade osteosarcomasin relation to clinical outcome. *New Eng. J. Med.* 333: 1380–1385,1995.
- [78609] 16939.Bell, D. R.; Trent, J. M.; Willard, H. F.; Riordan, J. R.; Ling,V.: Chromosomal location of human P–glycoprotein gene sequences. *CancerGenet. Cytogenet.* 25: 141–148, 1987.

- [78610] 16940.Callen, D. F.; Baker, E.; Simmers, R. N.; Seshadri, R.; Roninson, I. B.: Localization of the human multiple drug resistance gene, MDR1, to 7q21.1. *Hum. Genet.* 77: 142–144, 1987.
- [78611] 16941.Chan, H. S. L.; Haddad, G.; Thorner, P. S.; DeBoer, G.; Lin, Y.P.; Ondrusek, N.; Yeger, H.; Ling, V.: P-glycoprotein expression as a predictor of the outcome of therapy for neuroblastoma. *New Eng.J. Med.* 325: 1608–1614, 1991.
- [78612] 16942.Chang, G.; Roth, C. B.: Structure of MsbA from *E. coli*: a homolog of the multidrug resistance ATP binding cassette (ABC) transporters. *Science* 293:1793–1800, 2001.
- [78613] 16943.Chen, C.; Chin, J. E.; Ueda, K.; Clark, D. P.; Pastan, I.; Gottesman, M. M.; Roninson, I. B.: Internal duplication and homology with bacterial transport proteins in the *mdr1* (P-glycoprotein) gene from multidrug-resistant human cells. *Cell* 47: 381–389, 1986.
- [78614] 16944.Choi, K. H.; Chen, C.-J.; Kriegler, M.; Roninson, I. B.: An altered pattern of cross-resistance in multidrug-resistant human cells results from spontaneous mutations in the *mdr1* (P-glycoprotein) gene. *Cell* 53:519–529, 1988.
- [78615] 16945.Croop, J. M.; Gros, P.; Housman, D. E.: Genetics of



multidrugresistance. J. Clin. Invest. 81: 1303–1309, 1988.

[78616] 16946.de Lannoy, I. A. M.; Silverman, M.: The MDR1 gene product, P-glycoprotein,mediates the transport of the cardiac glycoside, digoxin. Biochem.Biophys. Res. Commun. 189: 551–557, 1992.

[78617] 16947.Cameron, P.; Limjoco, G.; Rodkey, J.; Bennett, C.; Schmidt, J.A.: Amino acid sequence analysis of human interleukin 1 (IL-1): evidencefor biochemically distinct forms of IL-1. J. Exp. Med. 162: 790–801,1985.

[78618] 16948.D'Eustachio, P.; Jadidi, S.; Fuhlbrigge, R. C.; Gray, P. W.; Chaplin,D. D.: Interleukin-1 alpha and beta genes: linkage on chromosome2 in the mouse. Immunogenetics 26: 339–343, 1987.

[78619] 16949.Dinarello, C. A.: An update on human interleukin-1: from molecularbiology to clinical relevance. J. Clin. Immun. 5: 287–297, 1985.

[78620] 16950.El-Omar, E. M.; Carrington, M.; Chow, W.-H.; McColl, K. E. L.;Bream, J. H.; Young, H. A.; Herrera, J.; Lisowska, J.; Yuan, C.-C.;Rothman, N.; Lanyon, G.; Martin, M.; Fraumeni, J. F., Jr.; Rabkin,C. S.: Interleukin-1 polymorphisms associated with increased riskof gastric cancer. Nature 404: 398–402, 2000. Note: Erratum: Nature412: 99 only, 2001.

- [78621] 16951.Furutani, Y.; Notake, M.; Yamayoshi, M.; Yamagishi, J.; Nomura,H.; Ohue, M.; Furuta, R.; Fukui, T.; Yamada, M.; Nakamura, S.: Cloningand characterization of the cDNAs for human and rabbit interleukin-1precursor. *Nucleic Acids Res.* 13: 5869–5882, 1985.
- [78622] 16952.Hamajima, N.; Matsuo, K.; Saito, T.; Tajima, K.; Okuma, K.; Yamao,K.; Tominaga, S.: Interleukin 1 polymorphisms, lifestyle factors,and helicobacter pylori infection. *Jpn. J. Cancer Res.* 92: 383–389,2001.
- [78623] 16953.Le Beau, M. M.; Rowley, J. D.: Personal Communi-  
cation. Chicago,Ill. 6/18/1986.
- [78624] 16954.March, C. J.; Mosley, B.; Larsen, A.; Cerretti, D. P.; Braedt,G.; Price, V.; Gillis, S.; Henney, C. S.; Kronheim, S. R.; Grabstein,K.; Conlon, P. J.; Hopp, T. P.; Cosman, D.: Cloning, sequence andexpression of two distinct human interleukin-1 complementary DNAs. *Nature* 315:641–647, 1985.
- [78625] 16955.Nicklin, M. J. H.; Weith, A.; Duff, G. W.: A physical map ofthe region encompassing the human interleukin-1-alpha, interleukin-1-beta,and interleukin-1 receptor antagonist genes. *Genomics* 19: 382–384,1994.
- [78626] 16956.Samad, T. A.; Moore, K. A.; Sapirstein, A.; Billet, S.; Allchorne,A.; Poole, S.; Bonventre, J. V.; Woolf, C. J.: Inter-

leukin-1-beta-mediated induction of Cox-2 in the CNS contributes to inflammatory pain hypersensitivity. *Nature* 410:471-475, 2001.

- [78627] 16957. Vidal-Vanaclocha, F.; Fantuzzi, G.; Mendoza, L.; Fuentes, A. M.; Anasagasti, M. J.; Martin, J.; Carrascal, T.; Walsh, P.; Reznikov, L. L.; Kim, S.-H.; Novick, D.; Rubinstein, M.; Dinarello, C. A.: IL-18 regulates IL-1-beta-dependent hepatic melanoma metastasis via vascular cell adhesion molecule-1. *Proc. Nat. Acad. Sci.* 97: 734-739, 2000.
- [78628] 16958. Webb, A.; Collins, K.; Auron, P.; Eddy, R.; Nakai, H.; Byers, M.; Shows, T. B.: Genetics of acute phase response: a gene for interleukin-1 is on chromosome 2. (Abstract) *Am. J. Hum. Genet.* 37: A142 only, 1985.
- [78629] 16959. Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai, H.; Byers, M.; Shows, T. B.: The gene for interleukin-1 (IL1) is on human chromosome 2. (Abstract) *Cytogenet. Cell Genet.* 40: 774 only, 1985.
- [78630] 16960. Webb, A. C.; Collins, K. L.; Auron, P. E.; Eddy, R. L.; Nakai, H.; Byers, M. G.; Haley, L. L.; Henry, W. M.; Shows, T. B.: Interleukin-1 gene (IL1) assigned to long arm of human chromosome 2. *Lymphokine Res.* 5: 77-85, 1986.
- [78631] 16961. Ansano, M.; Toda, M.; Sakaguchi, N.; Sakaguchi, S.:

Autoimmunedisease as a consequence of developmental abnormality of a T cellsubpopulation. J. Exp. Med. 184: 387–396, 1996.

[78632] 16962.Cosman, D.; Cerretti, D. P.; Larsen, A.; Park, L.; March, C.; Dower,S.; Gillis, S.; Urdal, D.: Cloning, sequence and expression of humaninterleukin–2 receptor. Nature 312: 768–771, 1984.

[78633] 16963.Ferrari, S.; Cannizzaro, L. A.; Battini, R.; Huebner, K.; Baserga,R.: The gene encoding human vimentin is located on the short armof chromosome 10. Am. J. Hum. Genet. 41: 616–626, 1987.

[78634] 16964.Greene, W. C.; Leonard, W. J.; Depper, J. M.; Nelson, D. L.; Waldmann,T. A.: The human interleukin–2 receptor: normal and abnormal expressionin T cells and in leukemias induced by the human T–lymphotropic retro–viruses. Ann.Intern. Med. 105: 560–572, 1986.

[78635] 16965.Hatakeyama, M.; Minamoto, S.; Taniguchi, T.: Intracytoplasmicphosphorylation sites of Tac antigen (p55) are not essential for theconformation, function, and regulation of the human interleukin 2receptor. Proc. Nat. Acad. Sci. 83: 9650–9654, 1986.

[78636] 16966.Hatakeyama, M.; Minamoto, S.; Uchiyama, T.; Hardy, R. R.; Yamada,G.; Taniguchi, T.: Reconstitution of

functional receptor for human interleukin-2 in mouse cells. *Nature* 318: 467-470, 1985.

[78637] 16967. Ihle, J. N.; Kerr, I. M.: Jaks and Stats in signaling by the cytokine receptor superfamily. *Trends Genet.* 11: 69-74, 1995.

[78638] 16968. Ishida, N.; Kanamori, H.; Noma, T.; Nikaido, T.; Sabe, H.; Suzuki, N.; Shimizu, A.; Honjo, T.: Molecular cloning and structure of the human interleukin 2 receptor gene. *Nucleic Acids Res.* 13: 7579-7589, 1985.

[78639] 16969. Kondo, S.; Shimizu, A.; Maeda, M.; Tagaya, Y.; Yodoi, J.; Honjo, T.: Expression of functional human interleukin-2 receptor in mouse T cells by cDNA transfection. *Nature* 320: 75-77, 1986.

[78640] 16970. Leonard, W. J.; Depper, J. M.; Crabtree, G. R.; Rudikoff, S.; Pumphrey, J.; Robb, R. J.; Kronke, M.; Svetlik, P. B.; Pfeffer, N. J.; Waldmann, T. A.; Greene, W. C.: Molecular cloning and expression of cDNAs for the human interleukin-2 receptor. *Nature* 311: 626-631, 1984.

[78641] 16971. Dahia, P. L. M.; Aguiar, R. C. T.; Alberta, J.; Kum, J. B.; Caron, S.; Sill, H.; Marsh, D. J.; Ritz, J.; Freedman, A.; Stiles, C.; Eng, C.: PTEN is inversely correlated with the cell survival factor Akt/PKB and is inactivated via multiple mechanisms in haematological malignancies. *Hum. Molec.*

Genet. 8: 185–193, 1999.

[78642] 16972.Bryson, G. L. M.; Massa, H.; Trask, B. J.; Van Etten, R. L.: Genestructure, sequence, and chromosomal localization of the human redcell-type low molecular weight acid phosphotyrosyl phosphatase gene,ACP1. Genomics 30: 133–140, 1995.

[78643] 16973.Scrable, H. J.; Johnson, D. K.; Rinchik, E. M.; Cavenee, W. K.: Rhabdomyosarcoma-associated locus and MYOD1 are syntenic but separateloci on the short arm of human chromosome 11. Proc. Nat. Acad. Sci. 87:2182–2186, 1990.

[78644] 16974.Chen, T.–R.; McMorris, F. A.; Creagan, R.; Ricciuti, F. C.; Tischfield,J.; Ruddle, F. H.: Assignment of the genes for malate oxidoreductasedecarboxylating to chromosome 6 and peptidase B and lactate dehydrogenaseB to chromosome 12 in man. Am. J. Hum. Genet. 25: 200–207, 1973.

[78645] 16975.Hamerton, J. L.; Mohandas, T.; McAlpine, P. J.; Douglas, G. R.: Localization of human gene loci using spontaneous chromosome rearrangementsin human–Chinese hamster somatic cell hybrids. Am. J. Hum. Genet. 27:595–608, 1975.

[78646] 16976.Herbschleb–Voogt, E.; Meera Khan, P.: Defining the locus of originof a genetically determined electrophoretic

variant of a multilocus enzyme system; the Calcutta-1 of human LDH system is a B-locus variant. Hum.Genet. 57: 290-295, 1981.

- [78647] 16977.Houki, N.; Matsushima, Y.; Kitamura, M.; Tukada, T.; Nishina, T.; Nakayama, T.: A case of deficiency of lactate dehydrogenase H-subunit. Jpn.J. Clin. Chem. 15: 85-90, 1986.
- [78648] 16978.Kitamura, M.; Iijima, N.; Hashimoto, F.; Hiratsuka, A.: Hereditary deficiency of subunit H of lactate dehydrogenase. Clin. Chim. Acta 34:419-423, 1971.
- [78649] 16979.Maekawa, M.; Sudo, K.; Kitajima, M.; Matsuura, Y.; Li, S. S.-L.; Kanno, T.: Analysis of a genetic mutation in an electrophoretic variant of the human lactate dehydrogenase-B(H) subunit. Hum. Genet. 91:423-426, 1993.
- [78650] 16980.Maekawa, M.; Sudo, K.; Nagura, K.; Li, S. S.-L.; Kanno, T.: Population screening of lactate dehydrogenase deficiencies in Fukuoka Prefecture in Japan and molecular characterization of three independent mutations in the lactate dehydrogenase-B(H) gene. Hum. Genet. 93: 74-76, 1994.
- [78651] 16981.Malpuech, G.; Kaplan, J. C.; Rethore, M. O.; Junien, C.; Geneix, A.: Une observation de deletion partielle du bras court du chromosome 12: localisation du gene de la

lacticodehydrogenase B. *Lyon Med.* 233:275–279, 1975.

[78652] 16982.Mayeda, K.; Weiss, L.; Lindahl, R.; Dully, M.: Localization of the human lactate dehydrogenase B gene on the short arm of chromosome 12. *Am. J. Hum. Genet.* 26: 59–64, 1974.

[78653] 16983.Miwa, S.; Nishima, T.; Kanehashi, Y.; Kitamura, M.; Hiratsuka, A.; Shizume, K.: Studies on erythrocyte metabolism in a case with hereditary deficiency of H-subunit of lactate dehydrogenase. *Acta Haemat. Jpn.* 34: 228–232, 1971.

[78654] 16984.Bell, D. W.; Varley, J. M.; Szydlo, T. E.; Kang, D. H.; Wahrer, D. C. R.; Shannon, K. E.; Lubratovich, M.; Verselis, S. J.; Isselbacher, K. J.; Fraumeni, J. F.; Birch, J. M.; Li, F. P.; Garber, J. E.; Haber, D. A.: Heterozygous germ line hCHK2 mutations in Li–Fraumeni syndrome. *Science* 286:2528–2531, 1999.

[78655] 16985.Goy, A.; Gilles, F.; Remache, Y.; Zelenetz, A. D.: Physical linkage of the lysyl oxidase-like (LOXL1) gene to the PML gene on human chromosome 15q22. *Cytogenet. Cell Genet.* 88: 22–24, 2000.

[78656] 16986.Kenyon, K.; Modi, W. S.; Contente, S.; Friedman, R. M.: A novel human cDNA with a predicted protein similar to lysyl oxidase maps to chromosome 15q24–q25. *J. Biol.*



Chem. 268: 18435–18437, 1993.

[78657] 16987.Kim, Y.; Boyd, C. D.; Csiszar, K.: A new gene with sequence and structural similarity to the gene encoding human lysyl oxidase. J.Biol. Chem. 270: 7176–7182, 1995.

[78658] 16988.Wydner, K. S.; Kim, Y.; Csiszar, K.; Boyd, C. D.; Passmore, H.C.: An intron capture strategy used to identify and map a lysyl oxidase–like gene on chromosome 9 in the mouse. Genomics 40: 342–345, 1997.

[78659] 16989.Dudek, H.; Datta, S. R.; Franke, T. F.; Birnbaum, M. J.; Yao, R.; Cooper, G. M.; Segal, R. A.; Kaplan, D. R.; Greenberg, M. E.: Regulation of neuronal survival by the serine–threonine protein kinase Akt. Science 275:661–663, 1997.

[78660] 16990.Franke, T. F.; Kaplan, D. R.; Cantley, L. C.; Toker, A.: Direct regulation of the Akt proto–oncogene product by phosphatidylinositol–3,4–bisphosphate. Science 275:665–667, 1997.

[78661] 16991.Franke, T. F.; Yang, S.–I.; Chan, T. O.; Datta, K.; Kaziauskas, A.; Morrison, D. K.; Kaplan, D. R.; Tsichlis, P. N.: The protein kinase encoded by the Akt proto–oncogene is a target of the PDGF–activated phosphatidylinositol 3–kinase. Cell 81: 727–736, 1995.

- [78662] 16992.Furnari, F. B.; Huang, H. J.; Cavenee, W. K.: The phosphoinositolphosphatase activity of PTEN mediates a serum-sensitive G1 growtharrest in glioma cells. *Cancer Res.* 58: 5002–5008, 1998.
- [78663] 16993.Hajduch, E.; Litherland, G. J.; Hundal, H. S.: Protein kinaseB (PKB/Akt)--a key regulator of glucose transport? *FEBS Lett.* 492:199–203, 2001.
- [78664] 16994.Hemmings, B. A.: Akt signaling: linking membrane events to lifeand death decisions. *Science* 275: 628–630, 1997.
- [78665] 16995.Jones, P. F.; Jakubowicz, T.; Pitossi, F. J.; Maurer, F.; Hemmings,B. A.: Molecular cloning and identification of a serine/threonineprotein kinase of the second-messenger subfamily. *Proc. Nat. Acad.Sci.* 88: 4171–4175, 1991.
- [78666] 16996.Lee, M.-J.; Thangada, S.; Paik, J.-H.; Sapkota, G. P.; Ancellin,N.; Chae, S.-S.; Wu, M.; Morales-Ruiz, M.; Sessa, W. C.; Alessi, D.R.; Hla, T.: Akt-mediated phosphorylation of the G protein-coupledreceptor EDG-1 is required for endothelial cell chemotaxis. *Molec.Cell* 8: 693–704, 2001.
- [78667] 16997.Maira, S.-M.; Galetic, I.; Brazil, D. P.; Kaech, S.; Ingley, E.;Thelen, M.; Hemmings, B. A.: Carboxyl-terminal modulator protein(CTMP), a negative regulator of PKB/Akt and v-Akt at the plasma membrane. *Science* 294:

374–380, 2001.

- [78668] 16998.Powell, D. W.; Rane, M. J.; Chen, Q.; Singh, S.; McLeish, K. R.: Identification of 14–3–3–zeta as a protein kinase B/Akt substrate. *J.Biol. Chem.* 277: 21639–21642, 2002.
- [78669] 16999.Staal, S. P.: Molecular cloning of the akt oncogene and its humanhomologues AKT1 and AKT2: amplification of AKT1 in a primary humangastric adenocarcinoma. *Proc. Nat. Acad. Sci.* 84: 5034–5037, 1987.
- [78670] 17000.Staal, S. P.; Huebner, K.; Croce, C. M.; Parsa, N. Z.; Testa,J. R.: The AKT1 proto–oncogene maps to human chromosome 14, bandq32. *Genomics* 2: 96–98, 1988.
- [78671] 17001.Vanhaesebroeck, B.; Alessi, D. R.: The PI3K–PDK1 connection:more than just a road to PKB. *Biochem. J.* 346: 561–576, 2000.
- [78672] 17002.Weng, L.–P.; Brown, J. L.; Eng, C.: PTEN induces apoptosis andcell cycle arrest through phosphoinositol–3–kinase/Akt–dependent and–independent pathways. *Hum. Molec. Genet.* 10: 237–242, 2001.
- [78673] 17003.Yang, J.; Cron, P.; Thompson, V.; Good, V. M.; Hess, D.; Hemmings,B. A.; Barford, D.: Molecular mechanism for the regulation of proteinkinase B/Akt by hydrophobic motif phosphorylation. *Molec. Cell*

9:1227–1240, 2002.

- [78674] 17004.Le Beau, M. M.; Rowley, J. D.; Sacchi, N.; Watson, D. K.; Papas, T. S.; Diaz, M. O.: Hu-ets-2 is translocated to chromosome 8 in the (8;21) in acute myelogenous leukemia. *Cancer Genet. Cytogenet.* 23:269–274, 1986.
- [78675] 17005.Mavrothalassitis, G. J.; Watson, D. K.; Papas, T. S.: Molecular and functional characterization of the promoter of ETS2, the human c-ets-2 gene. *Proc. Nat. Acad. Sci.* 87: 1047–1051, 1990.
- [78676] 17006.Sacchi, N.; Cheng, S. V.; Tanzi, R. E.; Gusella, J. F.; Drabkin, H. A.; Patterson, D.; Haines, J. H.; Papas, T. S.: The ETS genes on chromosome 21 are distal to the breakpoint of the acute myelogenous leukemia translocation (8;21). *Genomics* 3: 110–116, 1988.
- [78677] 17007.Sumarsono, S. H.; Wilson, T. J.; Tymms, M. J.; Venter, D. J.; Corrick, C. M.; Kola, R.; Lahoud, M. H.; Papas, T. S.; Seth, A.; Kola, I.: Down's syndrome-like skeletal abnormalities in Ets2 transgenic mice. *Nature* 379:534–540, 1996.
- [78678] 17008.Chu, E. H. Y.; Chang, C. C.; Sun, N. C.: Synteny of the human genes for gal-1-PT, ACP-1, MDH-1, and gal-plus-activator and assignment to chromosome 2. *Birth Defects Orig. Art. Ser.* XI(3): 103–106, 1975. Note: Alternate:

Cytogenet. Cell Genet. 14: 273–276, 1975...

- [78679] 17009.Rasmussen, S. K.; Lautier, C.; Hansen, L.; Echwald, S. M.; Hansen, T.; Ekstrom, C. T.; Urhammer, S. A.; Borch-Johnsen, K.; Grigorescu, F.; Smith, R. J.; Pedersen, O.: Studies of the variability of the genes encoding the insulin-like growth factor I receptor and its ligand in relation to type 2 diabetes mellitus. J. Clin. Endocr. Metab. 85:1606–1610, 2000.
- [78680] 17010.Dissing, J.; Johnsen, A. H.: Human red cell acid phosphatase (ACP1): the primary structure of the two pairs of isozymes encoded by the ACP1\*A and ACP1\*C alleles. Biochim. Biophys. Acta 1121: 261–268, 1992.
- [78681] 17011.Dissing, J.; Johnsen, A. H.; Sensabaugh, G. F.: Human red cell acid phosphatase (ACP1): the amino acid sequence of the two isozymes Bf and Bs encoded by the ACP1\*B allele. J. Biol. Chem. 266: 20619–20625, 1991.
- [78682] 17012.Emanuel, B. S.; Zackai, E. H.; Van Dyke, D. C.; Swallow, D. M.; Allen, F. H.; Mellman, W. J.: Deletion mapping: further evidence for the location of acid phosphatase (ACP-1) within 2p23. Am. J. Med. Genet. 4: 167–172, 1979.
- [78683] 17013.Ferguson-Smith, M. A.; Newman, B. F.; Ellis, P. M.; Thomson, D. M. G.; Riley, I. D.: Assignment by deletion of

human red cell acidphosphatase gene locus to the short arm of chromosome 2. *Nature* 243:271–273, 1973.

[78684] 17014.Fisher, R. A.; Harris, H.: Studies on the separate isoenzymes of red cell acid phosphatase phenotypes A and B: chromatographic separation of the isoenzymes. *Ann. Hum. Genet.* 34: 431–438, 1971.

[78685] 17015.Fuhrmann, W.; Lichte, K. H.: Human red cell acid phosphatase polymorphism: a study on gene frequency and forensic use of the system in cases of disputed paternity. *Humangenetik* 3: 121–126, 1966.

[78686] 17016.Giblett, E. R.; Scott, N. M.: Red cell acid phosphatase: racial distribution and report of a new phenotype. *Am. J. Hum. Genet.* 17:425–432, 1965.

[78687] 17017.Herbich, J.; Fisher, R. A.; Hopkinson, D. A.: Atypical segregation of human red cell acid phosphatase phenotypes: evidence for a rare 'silent' allele P(O). *Ann. Hum. Genet.* 34: 145–152, 1970.

[78688] 17018.Herbich, J.; Meinhardt, K.: The rare 'silent' allele P(O) or P(V)(P Vienna) of human red cell acid phosphatase, typed in a second family. *Humangenetik* 15:345–348, 1972.

[78689] 17019.Hopkinson, D. A.; Spencer, N.; Harris, H.: Red cell acid phosphatase variants: a new human polymorphism.

Nature 199: 969–971, 1963.

- [78690] 17020.Hulten, M.; Lindsten, J.; Pen-Ming, L. M.; Fraccaro, M.; Mannini,A.; Tiepolo, L.; Robson, E. B.; Heiken, A.; Tillinger, K. G.: Possible localization of the genes for the Kidd blood group on an autosome involved in a reciprocal translocation. Nature 211: 1067–1068, 1966.
- [78691] 17021.Junien, C.; Kaplan, J.-C.; Bernheim, A.; Berger, R.: Regional assignment of red cell acid phosphatase locus to band 2p25. Hum.Genet. 48: 17–21, 1979.
- [78692] 17022.Karp, G. W., Jr.; Sutton, H. E.: Some new phenotypes of human red cell acid phosphatase. Am. J. Hum. Genet. 19: 54–62, 1967.
- [78693] 17023.Lothe, R. A.; Gedde-Dahl, T.; Olaisen, B.; Bakker, E.; Pearson,P.: Very close linkage between D2S1 and ACP1 on chromosome 2p. Ann.Hum. Genet. 50: 361–367, 1986.
- [78694] 17024.Mace, M. A.; Cook, P. J. L.; Robson, E. B.: Linkage data on red cell acid phosphatase from family studies. Ann. Hum. Genet. 38:471–477, 1975.
- [78695] 17025.Mace, M. A.; Robson, E. B.: Linkage data on ACP-1 and MNSs. Cytogenet.Cell Genet. 13: 123–125, 1974.
- [78696] 17026.Magenis, R. E.; Koler, R. D.; Lovrien, E. W.; Bigley, R. H.; Duval,M. C.; Overton, K. M.: Gene dosage: evidence for assignment of erythrocyte acid phosphatase locus to

chromosome 2. Proc. Nat. Acad. Sci. 72:4526–4530, 1975.

[78697] 17027.Mayr, W. R.: No close linkage between MNSs and red cell acidphosphatase. Hum. Hered. 26: 1–3, 1976.

[78698] 17028.Miller, S. A.; Nelson, M. S.; Dykes, D. D.; Polesky, H. F.: Comparison of acid phosphatase ACP1 variants by isoelectric focusing and conventional electrophoresis: identification of three new alleles, ACP1\*N, ACP1\*P and ACP1\*S. Hum. Hered. 37: 371–375, 1987.

[78699] 17029.Mohrenweiser, H. W.; Novotny, J. E.: ACP–1–GUA–1: a low–activity variant of human erythrocyte acid phosphatase: association with increased glutathione reductase activity. Am. J. Hum. Genet. 34: 425–433, 1982.

[78700] 17030.Nezbeda, P.: Occurrence of the ACP–1 null allele in Czechoslovakia. Hum. Genet. 46: 227–229, 1979.

[78701] 17031.Palmarino, R.; Agostino, R.; Gloria, F.; Lucarelli, P.; Businco, L.; Antognoni, G.; Maggioni, G.; Workman, P. L.; Bottini, E.: Red cell acid phosphatase: another polymorphism correlated with malaria? Am. J. Phys. Anthropol. 43: 177–186, 1975.

[78702] 17032.Povey, S.; Swallow, D. M.; Bobrow, M.; Craig, I.; Van Heyningen, V.: Probable assignment of the locus determining human red cell acid phosphatase ACP(1) to chromo–



some 2 using somatic cell hybrids. *Ann.Hum. Genet.* 38: 1–5, 1974.

[78703] 17033.Renwick, J. H.: Assignment and map-positioning of human loci using chromosomal variation. *Ann. Hum. Genet.* 35: 79–97, 1971.

[78704] 17034.Sensabaugh, G. F.; Golden, V. L.: Phenotype dependence in the inhibition of red cell acid phosphatase (ACP) by folates. *Am. J.Hum. Genet.* 30: 553–560, 1978.

[78705] 17035.Swallow, D. M.; Povey, S.; Harris, H.: Activity of the 'red cell' acid phosphatase locus in other tissues. *Ann. Hum. Genet.* 37: 31–38, 1973.

[78706] 17036.Wu, R. M.; Cheng, C. W.; Chen, K. H.; Lu, S. L.; Shan, D. E.; Ho, Y. F.; Chern, H. D.: The COMT L allele modifies the association between MAOB polymorphism and PD in Taiwanese. *Neurology* 56: 375–382, 2001.

[78707] 17037.Levy, J. B.; Canoll, P. D.; Silvennoinen, O.; Barnea, G.; Morse, B.; Honegger, A. M.; Huang, J.–T.; Cannizzaro, L. A.; Park, S.–H.; Druck, T.; Huebner, K.; Sap, J.; Ehrlich, M.; Musacchio, J. M.; Schlessinger, J.: The cloning of a receptor-type protein tyrosine phosphatase expressed in the central nervous system. *J. Biol. Chem.* 268: 10573–10581, 1993.

[78708] 17038.Berube, D.; Simard, J.; Sandberg, M.; Grzeschik,

K.-H.; Gagne,R.; Hansson, V.; Jahnsen, T.: Assignment of the gene encoding the catalytic subunit C(beta) of cAMP-dependent protein kinase to the p36 band on chromosome 1. (Abstract) Cytogenet. Cell Genet. 58:1850 only, 1991.

[78709] 17039. Leonard, W. J.; Depper, J. M.; Kanehisa, M.; Kronke, M.; Pfeffer, N. J.; Svetlik, P. B.; Sullivan, M.; Greene, W. C.: Structure of the human interleukin-2 receptor gene. Science 230: 633-639, 1985.

[78710] 17040. Leonard, W. J.; Depper, J. M.; Robb, R. J.; Waldmann, T. A.; Greene, W. C.: Characterization of the human receptor for T-cell growth factor. Proc. Nat. Acad. Sci. 80: 6957-6961, 1983.

[78711] 17041. Marx, J. L.: The interleukin-2 receptor gene is cloned. Science 226:1064-1065, 1985.

[78712] 17042. Nikaido, T.; Shimizu, A.; Ishida, N.; Sabe, H.; Teshigawara, K.; Maeda, M.; Uchiyama, T.; Yodoi, J.; Honjo, T.: Molecular cloning of a cDNA encoding human interleukin-2 receptor. Nature 311: 631-635, 1984.

[78713] 17043. Robb, R. J.; Rusk, C. M.; Neeper, M. P.: Structure-function relationships for the interleukin 2 receptor: location of ligand and antibody binding sites on the Tac receptor chain by mutational analysis. Proc. Nat. Acad. Sci. 85: 5654-5658, 1988.

- [78714] 17044.Sharfe, N.; Dadi, H. K.; Shahar, M.; Roifman, C. M.: Human immunodeficiency disorder arising from mutation of the alpha chain of the interleukin-2 receptor. Proc. Nat. Acad. Sci. 94: 3168–3171, 1997.
- [78715] 17045.Shevach, E. M.: Certified professionals: CD4(+)CD25(+) suppressor T cells. J. Exp. Med. 193: F41–F45, 2001.
- [78716] 17046.Tsuda, M.; Kozak, R. W.; Goldman, C. K.; Waldmann, T. A.: Demonstration of a non-Tac peptide that binds interleukin 2: a potential participant in a multichain interleukin 2 receptor complex. Proc. Nat. Acad. Sci. 83: 9694–9698, 1986.
- [78717] 17047.Urdal, D. L.; March, C. J.; Gillis, S.; Larsen, A.; Dower, S.K.: Purification and chemical characterization of the receptor for interleukin 2 from activated human T lymphocytes and from a human T-cell lymphoma cell line. Proc. Nat. Acad. Sci. 81: 6481–6485, 1984.
- [78718] 17048.Celestin, J.; Rotschke, O.; Falk, K.; Ramesh, N.; Jabara, H.; Strominger, J.; Geha, R. S.: IL-3 induces B7.2 (CD86) expression and costimulatory activity in human eosinophils. J. Immun. 167: 6097–6104, 2001.
- [78719] 17049.Mohrenweiser, H. W.; Neel, J. V.: Frequency of thermostability variants: estimation of total 'rare' variant fre-

quency in human populations. *Proc.Nat. Acad. Sci.* 78: 5729–5733, 1981.

[78720] 17050.Gervais, F. G.; Singaraja, R.; Xanthoudakis, S.; Gutekunst, C.-A.;Leavitt, B. R.; Metzler, M.; Hackam, A. S.; Tam, J.; Vaillancourt,J. P.; Houtzager, V.; Rasper, D. M.; Roy, S.; Hayden, M. R.; Nicholson,D. W.: Recruitment and activation of caspase-8 by the huntingtin-inter-actingprotein Hip-1 and a novel partner Hippi. *Nature Cell Biol.* 4: 95–105,2002.

[78721] 17051.Abbott, A. M.; Bueno, R.; Pedrini, M. T.; Murray, J. M.; Smith,R. J.: Insulin-like growth factor I receptor gene structure. *J.Biol. Chem.* 267: 10759–10763, 1992.

[78722] 17052.All-Ericsson, C.; Girnita, L.; Seregard, S.; Bartolazzi, A.; Jager,M. J.; Larsson, O.: Insulin-like growth factor-1 receptor in uvealmelanoma: a predictor for metastatic disease and a potential therapeutictarget. *Invest. Ophthal. Vis. Sci.* 43: 1–8, 2002.

[78723] 17053.Cooke, D. W.; Bankert, L. A.; Roberts, C. T., Jr.; LeRoith, D.;Casella, S. J.: Analysis of the human type I insulin-like growthfactor receptor promoter region. *Biochem. Biophys. Res. Commun.* 177:1113–1120, 1991.

[78724] 17054.Dey, B. R.; Furlanetto, R. W.; Nissley, S. P.: Cloning of humanp55-gamma, a regulatory subunit of phos-

phatidylinositol 3-kinase, by a yeast two-hybrid library screen with the insulin-like growth factor-I receptor. *Gene* 209: 175-183, 1998.

[78725] 17055. Fernandez, A. M.; Kim, J. K.; Yakar, S.; Dupont, J.; Hernandez-Sanchez, C.; Castle, A. L.; Filmore, J.; Shulman, G. I.; Le Roith, D.: Functional inactivation of the IGF-I and insulin receptors in skeletal muscle causes type 2 diabetes. *Genes Dev.* 15: 1926-1934, 2001.

[78726] 17056. Flier, J. S.; Usher, P.; Moses, A. C.: Monoclonal antibody to the type I insulin-like growth factor (IGF-I) receptor blocks IGF-I receptor-mediated DNA synthesis: clarification of the mitogenic mechanisms of IGF-I and insulin in human skin fibroblasts. *Proc. Nat. Acad. Sci.* 83: 664-668, 1986.

[78727] 17057. Francke, U.; Yang-Feng, T. L.; Brissenden, J. E.; Ullrich, A.: Chromosomal mapping of genes involved in growth control. *Cold Spring Harbor Symp. Quant. Biol.* 51: 855-866, 1986.

[78728] 17058. Grant, E. S.; Ross, M. B.; Ballard, S.; Naylor, A.; Habib, F. K.: The insulin-like growth factor type I receptor stimulates growth and suppresses apoptosis in prostatic stromal cells. *J. Clin. Endocr. Metab.* 83: 3252-3257, 1998.

[78729] 17059. Herskowitz, I.: Functional inactivation of genes by

dominant negative mutations. *Nature* 329: 219–222, 1987.

- [78730] 17060. Howard, T. K.; Algar, E. M.; Glatz, J. A.; Reeve, A. E.; Smith, P. J.: The insulin-like growth factor 1 receptor gene is normally biallelically expressed in human juvenile tissue and tumours. *Hum. Molec. Genet.* 2: 2089–2092, 1993.
- [78731] 17061. Kulkarni, R. N.; Holzenberger, M.; Shih, D. Q.; Ozcan, U.; Stoffel, M.; Magnuson, M. A.; Kahn, C. R.: Beta-cell-specific deletion of the Igf1 receptor leads to hyperinsulinemia and glucose intolerance but does not alter beta-cell mass. *Nature Genet.* 31: 111–115, 2002.
- [78732] 17062. Maor, S. B.; Abramovitch, S.; Erdos, M. R.; Brody, L. C.; Werner, H.: BRCA1 suppresses insulin-like growth factor-I receptor promoter activity: potential interaction between BRCA1 and Sp1. *Molec. Genet. Metab.* 69: 130–136, 2000.
- [78733] 17063. Poduslo, S. E.; Dean, M.; Kolch, U.; O'Brien, S. J.: Detecting high-resolution polymorphisms in human coding loci by combining PCR and single-strand conformation polymorphism (SSCP) analysis. *Am. J. Hum. Genet.* 49: 106–111, 1991.
- [78734] 17064. Prager, D.; Yamasaki, H.; Weber, M. M.; Gebremedhin, S.; Melmed, S.: Human insulin-like growth factor I re-

ceptor function in pituitary cells is suppressed by a dominant negative mutant. *J. Clin. Invest.* 90:2117–2122, 1992.

- [78735] 17065. Roback, E. W.; Barakat, A. J.; Dev, V. G.; Mbikay, M.; Chretien, M.; Butler, M. G.: An infant with deletion of the distal long arm of chromosome 15 (q26.1–qter) and loss of insulin-like growth factor 1 receptor gene. *Am. J. Med. Genet.* 38: 74–79, 1991.
- [78736] 17066. Okumura, N.; Terasawa, F.; Ueno, I.; Oki, K.; Yamauchi, K.; Hidaka, H.; Tozuka, M.; Okura, M.; Katsuyama, T.: Genetic analyses in homozygous and heterozygous variants of lactate dehydrogenase-B (H) subunit: LD-B Matsumoto I and II (LD-B W323R). *Clin. Chim. Acta* 287: 163–171, 1999.
- [78737] 17067. Rethore, M.-O.; Junien, C.; Malpuech, G.; Baccichetti, C.; Tenconi, R.; Kaplan, J. C.; de Romeuf, J.; Lejeune, J.: Localisation du gène de la glyceraldehyde 3-phosphate dehydrogenase (G3PD) sur le segment distal du bras court du chromosome 12. *Ann. Genet.* 19: 140–142, 1976.
- [78738] 17068. Rethore, M.-O.; Kaplan, J.-C.; Junien, C.; Cruveiller, J.; Dutrillaux, B.; Aurias, A.; Carpentier, S.; Lafourcade, J.; Lejeune, J.: Augmentation de l'activité de la LDH-B chez un garçon trisomique 12p par mauvaise ségrégation d'une translo-

cation maternelle t(12;14)(q12;p11). Ann. Genet.  
18:81–87, 1975.

- [78739] 17069.Sakai, I.; Sharief, F. S.; Pan, Y.–C. E.; Li, S. S.–L.: The cDNA and protein sequences of human lactate dehydrogenase B. Biochem.J. 248: 933–936, 1987.
- [78740] 17070.Santachiara, A. S.; Nabholz, M.; Miggiano, V.; Darlington, A.J.; Bodmer, W. F.: Linkage between human lactate dehydrogenase B and peptidase B genes. Nature 227: 248–251, 1970.
- [78741] 17071.Steinbach, P.; Rehder, H.: Tetrasomy for the short arm of chromosome 12 with accessory isochromosome (+i[12p]) and a marked LDH–B gene dosage effect. Clin. Genet. 32: 1–4, 1987.
- [78742] 17072.Sudo, K.: Personal Communication. Komae City, Japan 7/12/1993.
- [78743] 17073.Sudo, K.; Maekawa, M.; Ikawa, S.; Machida, K.; Kitamura, M.; Li, S. S.–L.: A missense mutation found in human lactate dehydrogenase–B(H) variant gene. Biochem. Biophys. Res. Commun. 168: 672–676, 1990.
- [78744] 17074.Sudo, K.; Maekawa, M.; Luedemann, M. M.; Deaven, L. L.; Li, S.S.–L.: Human lactate dehydrogenase–B processed pseudogene: nucleotide sequence analysis and assignment to the X–chromosome. Biochem. Biophys. Res.



Commun. 171: 67–74, 1990.

- [78745] 17075.Sudo, K.; Maekawa, M.; Tomonaga, A.; Tsukada, T.; Nakayama, T.; Kitamura, M.; Li, S. S.–L.; Kanno, T.; Toriumi, J.: Molecular characterization of genetic mutations in human lactate dehydrogenase (LDH)B (H) variant. Hum.Genet. 89: 158–162, 1992.
- [78746] 17076.Van Someren, H.; Meera Khan, P.; Westerveld, A.; Bootsma, D.: Human genetics--two new linkage groups carrying different loci for LDH and glutamic-pyruvic transaminase found. Nature 240: 221–222, 1972.
- [78747] 17077.Weiss, L.; Mayeda, K.; Lindahl, R.; Dully, M.: Localization of human LDH–B gene of the short arm of chromosome 12. (Abstract) Am.J. Hum. Genet. 25: 85A only, 1973.
- [78748] 17078.Davies, H.; Bignell, G. R.; Cox, C.; Stephens, P.; Edkins, S.; Clegg, S.; Teague, J.; Woffendin, H.; Garnett, M. J.; Bottomley, W.; Davis, N.; Dicks, E.; and 40 others: Mutations of the BRAF gene in human cancer. Nature 417: 949–954, 2002.
- [78749] 17079.Eychene, A.; Barnier, J. V.; Apiou, F.; Dutrillaux, B.; Calothy, G.: Chromosomal assignment of two human B–raf(Rmil) proto-oncogene loci: B–raf–1 encoding the p94(Braf/Rmil) and B–raf–2, a processed pseudogene.

Oncogene 7: 1657–1660, 1992.

- [78750] 17080.Rajagopalan, H.; Bardelli, A.; Lengauer, C.; Kinzler, K. W.; Vogelstein, B.; Velculescu, V. E.: RAF/RAS oncogenes and mismatch–repair status.(Letter) Nature 418: 934 only, 2002.
- [78751] 17081.Sithanandam, G.; Druck, T.; Cannizzaro, L. A.; Leuzzi, G.; Huebner, K.; Rapp, U. R.: B–raf and a B–raf pseudogene are located on 7q in man. Oncogene 7: 795–799, 1992.
- [78752] 17082.Wojnowski, L.; Zimmer, A. M.; Beck, T. W.; Hahn, H.; Bernal, R.; Rapp, U. R.; Zimmer, A.: Endothelial apoptosis in Braf–deficient mice. Nature Genet. 16: 293–297, 1997.
- [78753] 17083.Zhong, S.; Delva, L.; Rachez, C.; Cenciarelli, C.; Gandini, D.; Zhang, H.; Kalantry, S.; Freedman, L. P.; Pandolfi, P. P.: A RA–dependent, tumour–growth suppressive transcription complex is the target of the PML–RAR–alpha and T18 oncoproteins. Nature Genet. 23: 287–295, 1999.
- [78754] 17084.Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura, N.; Shimizu, N.: Assignment of hepatocyte growth factor (HGF) gene to chromosome 7q21.1. (Abstract) Cytogenet. Cell Genet. 58: 1921 only, 1991.
- [78755] 17085.Fukuyama, R.; Ichijoh, Y.; Minoshima, S.; Kitamura,

N.; Shimizu, N.: Regional localization of the hepatocyte growth factor (HGF) gene to human chromosome 7 band q21.1. *Genomics* 11: 410–415, 1991.

[78756] 17086. Gherardi, E.; Stoker, M.: Hepatocytes and scatter factor. *Nature* 346:228 only, 1990.

[78757] 17087. Gohda, E.; Tsubouchi, H.; Nakayama, H.; Hirono, S.; Sakiyama, O.; Takahashi, K.; Miyazaki, H.; Hashimoto, S.; Daikuhara, Y.: Purification and partial characterization of hepatocyte growth factor from plasma of a patient with fulminant hepatic failure. *J. Clin. Invest.* 81:414–419, 1988.

[78758] 17088. Kilby, M. D.; Afford, S.; Li, X. F.; Strain, A. J.; Ahmed, A.; Whittle, M. J.: Localisation of hepatocyte growth factor and its receptor (c-met) protein and mRNA in human term placenta. *Growth Factors* 13: 133–139, 1996.

[78759] 17089. Lai, L.; Goldschneider, I.: Cutting edge: identification of a hybrid cytokine consisting of IL-7 and the beta-chain of the hepatocyte growth factor/scatter factor. *J. Immun.* 167: 3550–3554, 2001.

[78760] 17090. Maina, F.; Casagrande, F.; Audero, E.; Simeone, A.; Comoglio, P. M.; Klein, R.; Ponzetto, C.: Uncoupling of Grb2 from the Met receptor in vivo reveals complex roles in muscle development. *Cell* 87: 531–542, 1996.

- [78761] 17091.Miyazawa, K.; Tsubouchi, H.; Naka, D.; Takahashi, K.; Okigaki,M.; Arakaki, N.; Nakayama, H.; Hirono, S.; Sakiyama, O.; Takahashi,K.; Gohda, E.; Daikuhara, Y.; Kitamura, N.: Molecular cloning andsequence analysis of cDNA for human hepatocyte growth factor. Biochem.Biophys. Res. Commun. 163: 967–973, 1989.
- [78762] 17092.Nakamura, T.; Nishizawa, T.; Hagiya, M.; Seki, T.; Shimonishi,M.; Sugimura, A.; Tashiro, K.; Shimizu, S.: Molecular cloning andexpression of human hepatocyte growth factor. Nature 342: 440–443,1989.
- [78763] 17093.Noonan, F. P.; Recio, J. A.; Takayama, H.; Duray, P.; Anver, M.R.; Rush, W. L.; De Fabo, E. C.; Merlino, G.: Neonatal sunburn andmelanoma in mice: severe sunburn in newborn, but not adult, mice islinked with melanoma in later life. Nature 413: 271–272, 2001.
- [78764] 17094.Powell, E. M.; Mars, W. M.; Levitt, P.: Hepatocyte growth factor/scatterfactor is a motogen for interneurons migrating from the ventral todorsal telencephalon. Neuron 30: 79–89, 2001.
- [78765] 17095.Rubin, J. S.; Chan, A. M.–L.; Bottaro, D. P.; Burgess, W. H.;Taylor, W. G.; Cech, A. C.; Hirschfield, D. W.; Wong, J.; Miki, T.;Finch, P. W.; Aaronson, S. A.: A broad–spectrum human lung fibroblast–derivedmitogen is a variant of

hepatocyte growth factor. Proc. Nat. Acad.Sci. 88:  
415–419, 1991.

[78766] 17096.Saccone, S.; Narsimhan, R. P.; Gaudino, G.; Dalpra, L.; Comoglio,P. M.; Della Valle, G.: Regional mapping of the human hepatocytegrowth factor (HGF)–scatter factor gene to chromosome 7q21.1. Genomics 13:912–914, 1992.

[78767] 17097.Schmidt, C.; Bladt, F.; Goedecke, S.; Brinkmann, V.; Zschiesche,W.; Sharpe, M.; Gherardi, E.; Birchmeier, C.: Scatter factor/hepatocytegrowth factor is essential for liver development. Nature 373: 699–702,1995.

[78768] 17098.Uehara, Y.; Minowa, O.; Mori, C.; Shiota, K.; Kuno, J.; Noda,T.; Kitamura, N.: Placental defect and embryonic lethality in micelacking hepatocyte growth factor/scatter factor. Nature 373: 702–705,1995.

[78769] 17099.Weidner, K. M.; Arakaki, N.; Hartmann, G.; Vandekerckhove, J.;Weingart, S.; Rieder, H.; Fonatsch, C.; Tsubouchi, H.; Hishida, T.;Daikuhara, Y.; Birchmeier, W.: Evidence for the identity of humanscatter factor and human hepatocyte growth factor. Proc. Nat. Acad.Sci. 88: 7001–7005, 1991.

[78770] 17100.Zarnegar, R.; Petersen, B.; DeFrances, M. C.; Michalopoulos, G.: Localization of hepatocyte growth fac–

tor (HGF) gene on human chromosome7. Genomics 12: 147–150, 1992.

[78771] 17101.Calandro, L. M.; Baer, D. M.; Sensabaugh, G. F.: Characterization of a recombinant that locates the hereditary hemochromatosis gene telomeric to HLA-F. Hum. Genet. 96: 339–342, 1995.

[78772] 17102.Geraghty, D. E.; Wei, X. H.; Orr, H. T.; Koller, B. H.: Human leukocyte antigen F (HLA-F): an expressed HLA gene composed of a class I coding sequence linked to a novel transcribed repetitive element. J.Exp. Med. 171: 1–18, 1990.

[78773] 17103.Shiina, T.; Tamiya, G.; Oka, A.; Takishima, N.; Yamagata, T.; Kikkawa, E.; Iwata, K.; Tomizawa, M.; Okuaki, N.; Kuwano, Y.; Watanabe, K.; Fukuzumi, Y.; and 11 others: Molecular dynamics of MHC genes unraveled by sequence analysis of the 1,796,938-bp HLA class I region. Proc.Nat. Acad. Sci. 96: 13282–13287, 1999.

[78774] 17104.Gasparini, P.; Borgato, L.; Piperno, A.; Girelli, D.; Olivieri, O.; Gottardi, E.; Roetto, A.; Dianzani, I.; Fargion, S.; Schinaia, G.; Cappellini, M. D.; Gandini, G.; Pignatti, P. F.; Fiorelli, G.; De Sandre, G.; Camaschella, C.: Linkage analysis of 6p21 polymorphic markers and the hereditary hemochromatosis: localization of the gene centromeric to

HLA-F. Hum. Molec. Genet. 2: 571-576, 1993.

- [78775] 17105.Engel, W.; Klemme, B.; Ebrecht, A.: Serological evidence for H-Yantigen in XO-female mice. Hum. Genet. 57: 68-70, 1981.
- [78776] 17106.Lau, Y.-F.: Personal Communication. San Francisco, Calif. 5/4/1987.
- [78777] 17107.Lau, Y.-F.; Chan, K.; Kan, Y. W.; Goldberg, E.: Isolation of amale-specific and conserved gene using an anti-H-Y antibody. (Abstract) Am.J. Hum. Genet. 39: A142 only, 1986.
- [78778] 17108.Lau, Y.-F.; Chan, K.; Kan, Y. W.; Goldberg, E.: Structure and expression of a gene isolated with an anti-H-Y antibody. (Abstract) Clin.Res. 35: 647A only, 1987.
- [78779] 17109.Muller, U.: Identification and function of serologically detectable H-Y antigen. Hum. Genet. 61: 91-94, 1982.
- [78780] 17110.Shapiro, M.; Erickson, R. P.: Evidence that the serological determinant of H-Y antigen is carbohydrate. Nature 290: 503-505, 1981.
- [78781] 17111.Wiberg, U.; Mayerova, A.; Muller, U.; Fredga, K.; Wolf, U.: X-linked genes of the H-Y antigen system in the wood lemming (*Myopus schisticolor*). Hum.Genet. 60: 163-166, 1982.

- [78782] 17112.Wolf, U.: Zum Mechanismus der Gonadendifferenzierung. Bull. Schweiz.Akad. Med. Wiss. 34: 357–368, 1978.
- [78783] 17113.van den Boogaard, M.–J. H.; Dorland, M.; Beemer, F. A.; Ploosvan Amstel, H. K.: MSX1 mutation is associated with orofacial clefting and tooth agenesis in humans. (Letter) Nature Genet. 24: 342–343, 2000. Note: Erratum: Nature Genet. 25: 125 only, 2000.
- [78784] 17114.Del Campo, M.; Jones, M. C.; Veraksa, A. N.; Curry, C. J.; Jones, K. L.; Mascarello, J. T.; Ali-Kahn-Catts, Z.; Drumheller, T.; McGinnis, W.: Monodactylous limbs and abnormal genitalia are associated with hemizyosity for the human 2q31 region that includes the HOXD cluster. Am.J. Hum. Genet. 65: 104–110, 1999.
- [78785] 17115.Zakany, J.; Kmita, M.; Alarcon, P.; de la Pompa, J.–L.; Duboule, D.: Localized and transient transcription of Hox genes suggests a link between patterning and the segmentation clock. Cell 106: 207–217, 2001.
- [78786] 17116.Perez, O. D.; Kinoshita, S.; Hitoshi, Y.; Payan, D. G.; Kitamura, T.; Nolan, G. P.; Lorens, J. B.: Activation of the PKB/AKT pathway by ICAM–2. Immunity 16: 51–65, 2002.
- [78787] 17117.Sansom, D.; Borrow, J.; Solomon, E.; Trowsdale, J.: The human ICAM2 gene maps to 17q23–25. Genomics 11:



462–464, 1991.

- [78788] 17118.Staunton, D. E.; Dustin, M. L.; Springer, T. A.: Functional cloning of ICAM–2, a cell adhesion ligand for LFA–1 homologous to ICAM–1. *Nature* 339:61–64, 1989.
- [78789] 17119.Rotem–Yehudar, R.; Galperin, E.; Horowitz, M.: Association of insulin–like growth factor 1 receptor with EHD1 and SNAP29. *J. Biol.Chem.* 276: 33054–33060, 2001.
- [78790] 17120.Scott, B. A.; Avidan, M. S.; Crowder, C. M.: Regulation of hypoxic death in *C. elegans* by the insulin/IGF receptor homolog DAF–2. *Science* 296:2388–2391, 2002.
- [78791] 17121.Tatar, M.; Kopelman, A.; Epstein, D.; Tu, M.–P.; Yin, C.–M.; Garofalo, R. S.: A mutant *Drosophila* insulin receptor homolog that extends life–span and impairs neuroendocrine function. *Science* 292: 107–110, 2001.
- [78792] 17122.Ullrich, A.; Gray, A.; Tam, A. W.; Yang–Feng, T.; Tsubokawa, M.; Collins, C.; Henzel, W.; Le Bon, T.; Kathuria, S.; Chen, E.; Jacobs, S.; Francke, U.; Ramachandran, J.; Fujita–Yamaguchi, Y.: Insulin–like growth factor I receptor primary structure: comparison with insulin receptor suggests structural determinants that define functional specificity. *EMBOJ.* 5: 2503–2512, 1986.
- [78793] 17123.Werner, H.; Karnieli, E.; Rauscher, F. J., III; LeRoth,

D.: Wild-type and mutant p53 differentially regulate transcription of the insulin-like growth factor I receptor gene. Proc. Nat. Acad. Sci. 93: 8318–8323, 1996.

- [78794] 17124. Barton, D. E.; Yang-Feng, T. L.; Mason, A. J.; Seeburg, P. H.; Francke, U.: INHA, INHBA and INHBB, the loci for the three subunits of inhibin, mapped in mouse and man. (Abstract) Cytogenet. Cell Genet. 46:578 only, 1987.
- [78795] 17125. Barton, D. E.; Yang-Feng, T. L.; Mason, A. J.; Seeburg, P. H.; Francke, U.: Mapping of genes for inhibin subunits alpha, beta(A) and beta(B) on human and mouse chromosomes and studies of jsd mice. Genomics 5:91–99, 1989.
- [78796] 17126. Bremner, W. J.: Inhibin: from hypothesis to clinical application. (Editorial) New Eng. J. Med. 321: 826–827, 1989.
- [78797] 17127. Lappohn, R. E.; Burger, H. G.; Bouma, J.; Bangah, M.; Krans, M.; de Bruijn, H. W. A.: Inhibin as a marker for granulosa-cell tumors. New Eng. J. Med. 321: 790–793, 1989.
- [78798] 17128. Mason, A. J.; Hayflick, J. S.; Ling, N.; Esch, F.; Ueno, N.; Ying, S.-Y.; Guillemin, R.; Niall, H.; Seeburg, P. H.: Complementary DNA sequences of ovarian follicular fluid inhibin show precursor structure and homology with

transforming growth factor-beta. *Nature* 318:  
659-663, 1985.

[78799] 17129. Mason, A. J.; Niall, H. D.; Seeburg, P. H.: Structure of two human ovarian inhibins. *Biochem. Biophys. Res. Commun.* 135: 957-964, 1986.

[78800] 17130. Matzuk, M. M.; Finegold, M. J.; Mather, J. P.; Krummen, L.; Lu, H.; Bradley, A.: Development of cancer cachexia-like syndrome and adrenal tumors in inhibin-deficient mice. *Proc. Nat. Acad. Sci.* 91: 8817-8821, 1994.

[78801] 17131. Matzuk, M. M.; Finegold, M. J.; Su, J.-G. J.; Hsueh, A. J. W.; Bradley, A.: Alpha-inhibin is a tumour-suppressor gene with gonadal specificity in mice. *Nature* 360: 313-319, 1992.

[78802] 17132. Mellor, S. L.; Richards, M. G.; Pedersen, J. S.; Robertson, D. M.; Risbridger, G. P.: Loss of the expression and localization of inhibin alpha-subunit in high grade prostate cancer. *J. Clin. Endocr. Metab.* 83: 969-975, 1998.

[78803] 17133. Meunier, H.; Rivier, C.; Evans, R. M.; Vale, W.: Gonadal and extragonadal expression of inhibin alpha, beta-A and beta-B subunits in various tissues predicts diverse functions. *Proc. Nat. Acad. Sci.* 85: 247-251, 1988.

[78804] 17134. Ramasharma, K.; Li, C. H.: Characteristics of bind-

ing of human seminal alpha-inhibin-92 to human pituitary membranes. *Proc. Nat. Acad. Sci.* 84: 3595–3598, 1987.

[78805] 17135. Schmitt, J. F.; Millar, D. S.; Pedersen, J. S.; Clark, S. L.; Venter, D. J.; Frydenberg, M.; Molloy, P. L.; Risbridger, G. P.: Hypermethylation of the inhibin alpha-subunit gene in prostate carcinoma. *Molec. Endocr.* 16: 213–220, 2002.

[78806] 17136. Thirunavukarasu, P.; Stephenson, T.; Forray, J.; Stanton, P. G.; Groome, N.; Wallace, E.; Robertson, D. M.: Changes in molecular weight forms of inhibin A and pro-alpha-C in maternal serum during human pregnancy. *J. Clin. Endocr. Metab.* 86: 5794–5804, 2001.

[78807] 17137. Lafuse, W. P.; Zwilling, B. S.: Localization of the inhibin beta-B gene on mouse chromosome 1. *Mammalian Genome* 4: 399–400, 1993.

[78808] 17138. Salmenkivi, K.; Arola, J.; Voutilainen, R.; Ilvesmaki, V.; Haglund, C.; Kahri, A. I.; Heikkila, P.; Liu, J.: Inhibin/activin beta-B-subunit expression in pheochromocytomas favors benign diagnosis. *J. Clin. Endocr. Metab.* 86: 2231–2235, 2001.

[78809] 17139. Schrewe, H.; Gendron-Maguire, M.; Harbison, M. L.; Gridley, T.: Mice homozygous for a null mutation of activin beta B are viable and fertile. *Mech. Dev.* 47: 43–51, 1994.

- [78810] 17140.Nicoll, J. A. R.; Mrak, R. E.; Graham, D. I.; Stewart, J.; Wilcock,G.; MacGowan, S.; Esiri, M. M.; Murray, L. S.; Dewar, D.; Love, S.;Moss, T.; Griffin, W. S. T.: Association of interleukin-1 gene polymorphisms with Alzheimer's disease. *Ann. Neurol.* 47: 365-368, 2000.
- [78811] 17141.Rogers, J.: An IL-1-alpha susceptibility polymorphism in Alzheimer's disease: new fuel for the inflammation hypothesis. (Editorial) *Neurology* 55:464-465, 2000.
- [78812] 17142.Sabatino, M.; Boyce, B.; Aufdemorte, T.; Bonewald, L.; Mundy,G. R.: Infusions of recombinant human interleukins 1 alpha and 1beta cause hypercalcemia in normal mice. *Proc. Nat. Acad. Sci.* 85:5235-5239, 1988.
- [78813] 17143.Silver, A. R. J.; Masson, W. K.; George, A. M.; Adam, J.; Cox,R.: The IL-1 alpha and beta genes are closely linked (less than 70kb) on mouse chromosome 2. *Somat. Cell Molec. Genet.* 16: 549-556,1990.
- [78814] 17144.Dickensheets, H. L.; Venkataraman, C.; Schindler, U.; Donnelly,R. P.: Interferons inhibit activation of STAT6 by interleukin 4 in human monocytes by inducing SOCS-1 gene expression. *Proc. Nat. Acad.Sci.* 96: 10800-10805, 1999.
- [78815] 17145.Kotanides, H.; Reich, N. C.: Interleukin-4-induced STAT6 recognizes and activates a target site in the pro-

moter of the interleukin-4 receptorgene. J. Biol. Chem. 271: 25555-25561, 1996.

[78816] 17146.Takahashi, M.; Yoshida, M. C.; Satoh, H.; Hilgers, J.; Yaoita,Y.; Honjo, T.: Chromosomal mapping of the mouse IL-4 and human IL-5genes. Genomics 4: 47-52, 1989.

[78817] 17147.Caggana, M.; Walker, K.; Reilly, A. A.; Conroy, J. M.; Duva, S.;Walsh, A. C.: Population-based studies reveal differences in theallelic frequencies of two functionally significant human interleukin-4receptor polymorphisms in several ethnic groups. Genet. Med. 1:267-271, 1999.

[78818] 17148.Deichmann, K. A.; Heinzmann, A.; Forster, J.; Dischinger, S.; Mehl,C.; Brueggenolte, E.; Hildebrandt, F.; Moseler, M.; Kuehr, J.: Linkageand allelic association of atopy and markers flanking the IL4-receptorgene. Clin. Exp. Allergy 28: 151-155, 1998.

[78819] 17149.MacLeod, J. N.; Lee, A. K.; Liebhaber, S. A.; Cooke, N. E.: Developmentalcontrol and alternative splicing of the placentally expressed transcriptsfrom the human growth hormone gene cluster. J. Biol. Chem. 267:14219-14226, 1992.

[78820] 17150.Wurzel, J. M.; Parks, J. S.; Herd, J. E.; Nielsen, P. V.: A genedeletion is responsible for absence of human chorionic somatomammotropin. DNA 1:251-257, 1982.

- [78821] 17151.Gallin, J. I.: Personal Communication. Bethesda, Md. 10/12/1990.
- [78822] 17152.McCombs, J. L.; Teng, C. T.; Pentecost, B. T.; Magnuson, V. L.; Moore, C. M.; McGill, J. R.: Chromosomal localization of human lactotransferringene (LTF) by in situ hybridization. *Cytogenet. Cell Genet.* 47:16–17, 1988.
- [78823] 17153.Moriuchi, M.; Moriuchi, H.: A milk protein lactoferrin enhances human T cell leukemia virus type I and suppresses HIV–1 infection. *J.Immun.* 166: 4231–4236, 2001.
- [78824] 17154.Naylor, S. L.; Marshall, A.; Solomon, A.; McGill, J. R.; McCombs, J.; Magnuson, V. L.; Moore, C. M.; Lalley, P. A.; Pentecost, B. T.; Teng, C.: Lactoferrin maps to human chromosome 3(q21–q23) and mouse chromosome 9. (Abstract) *Cytogenet. Cell Genet.* 46: 669 only, 1987.
- [78825] 17155.Powell, M. J.; Ogden, J. E.: Nucleotide sequence of human lactoferrin cDNA. *Nucleic Acids Res.* 18: 4013 only, 1990.
- [78826] 17156.Qiu, J.; Hendrixson, D. R.; Baker, E. N.; Murphy, T. F.; St. Geme, J. W., III; Plaut, A. G.: Human milk lactoferrin inactivates two putative colonization factors expressed by *Haemophilus influenzae*. *Proc.Nat. Acad. Sci.* 95: 12641–12646, 1998.
- [78827] 17157.Rado, T. A.; Wei, X.; Benz, E. J., Jr.: Isolation of

lactoferrin cDNA from a human myeloid library and expression of mRNA during normal and leukemic myelopoiesis. *Blood* 70: 989–993, 1987.

- [78828] 17158. Singh, P. K.; Parsek, M. R.; Greenberg, E. P.; Welsh, M. J.: A component of innate immunity prevents bacterial biofilm development. *Nature* 417:552–555, 2002.
- [78829] 17159. Teng, C. T.; Pentecost, B. T.; Marshall, A.; Solomon, A.; Bowman, B. H.; Lalley, P. A.; Naylor, S. L.: Assignment of the lactotransferrin gene to human chromosome 3 and to mouse chromosome 9. *Somat. Cell Molec. Genet.* 13: 689–693, 1987.
- [78830] 17160. Yang, F.; Lum, J.; Baldwin, W. D.; Brune, J. L.; van Bragt, P.; Bowman, B. H.: Genetic analysis of human iron binding glycoproteins. (Abstract) *Am. J. Hum. Genet.* 35: 184A only, 1983.
- [78831] 17161. Tanaka, A. R.; Ikeda, Y.; Abe-Dohmae, S.; Arakawa, R.; Sadanami, K.; Kidera, A.; Nakagawa, S.; Nagase, T.; Aoki, R.; Kioka, N.; Amachi, T.; Yokoyama, S.; Ueda, K.: Human ABCA1 contains a large amino terminal extracellular domain homologous to an epitope of Sjogren's syndrome. *Biochem. Biophys. Res. Commun.* 283: 1019–1025, 2001.
- [78832] 17162. Xu, W.; Gelber, S.; Orr-Urtreger, A.; Armstrong, D.;



Lewis, R.A.; Ou, C.-N.; Patrick, J.; Role, L.; De Biasi, M.; Beaudet, A. L.: Megacystis, mydriasis, and ion channel defect in mice lacking the  $\alpha$ -3 neuronal nicotinic acetylcholine receptor. *Proc. Nat. Acad.Sci.* 96: 5746–5751, 1999.

[78833] 17163. Gallardo, M. E.; Lopez-Rios, J.; Fernaud-Espinosa, I.; Granadino, B.; Sanz, R.; Ramos, C.; Ayuso, C.; Seller, M. J.; Brunner, H. G.; Bovolenta, P.; Rodriguez de Cordoba, S.: Genomic cloning and characterization of the human homeobox gene SIX6 reveals a cluster of SIX genes in chromosome 14 and associates SIX6 hemizygosity with bilateral anophthalmia and pituitary anomalies. *Genomics* 61: 82–91, 1999.

[78834] 17164. Sun, X.-H.: Constitutive expression of the *Id1* gene impairs mouse B cell development. *Cell* 79: 893–900, 1994.

[78835] 17165. Nowell, P. C.; Hungerford, D. A.: Chromosome studies on normal and leukemic human leukocytes. *J. Nat. Cancer Inst.* 25: 85–109, 1960.

[78836] 17166. Dowbenko, D. J.; Diep, A.; Taylor, B. A.; Lysis, A. J.; Lasky, L. A.: Characterization of the murine homing receptor gene reveals correspondence between protein domains and coding exons. *Genomics* 9: 270–277, 1991.

- [78837] 17167.Couturier, J.; Sastre-Garau, X.; Schneider-Maunoury, S.; Labib,A.; Orth, G.: Integration of papillomavirus DNA near myc genes ingenital carcinomas and its consequences for proto-oncogene expression. J.Virol. 65: 4534-4538, 1991.
- [78838] 17168.Habraken, Y.; Sung, P.; Prakash, S.; Prakash, L.: Transcriptionfactor TFIIH and DNA endonuclease Rad2 constitute yeast nucleotideexcision repair factor 3: implications for nucleotide excision repairand Cockayne syndrome. Proc. Nat. Acad. Sci. 93: 10718-10722, 1996.
- [78839] 17169.Abruzzo, L. V.; Jaffe, E. S.; Cotelingam, J. D.; Whang-Peng, J.;Del Duca, V., Jr.; Medeiros, L. J.: T-cell lymphoblastic lymphomawith eosinophilia associated with subsequent myeloid malignancy. Am.J. Surg. Path. 16: 236-245, 1992.
- [78840] 17170.Chesi, M.; Nardini, E.; Brents, L. A.; Schrock, E.; Ried, T.; Kuehl,W. M.; Bergsagel, P. L.: Frequent translocation t(4;14)(p16.3;q32.3)in multiple myeloma is associated with increased expression and activatingmutations of fibroblast growth factor receptor 3. Nature Genet. 16:260-264, 1997.
- [78841] 17171.Demiroglu, A.; Steer, E. J.; Heath, C.; Taylor, K.; Bentley, M.;Allen, S. L.; Koduru, P.; Brody, J. P.; Hawson,

G.; Rodwell, R.; Doody, M.-L.; Carnicero, F.; Reiter, A.; Goldman, J. M.; Melo, J. V.; Cross, N. C. P.: The t(8;22) in chronic myeloid leukemia fuses BCR to FGFR1: transforming activity and specific inhibition of FGFR1 fusion proteins. *Blood* 98:3778–3783, 2001.

[78842] 17172. Guasch, G.; Mack, G. J.; Popovici, C.; Dastugue, N.; Birnbaum, D.; Rattner, J. B.; Pebusque, M.-J.: FGFR1 is fused to the centrosome-associated protein CEP110 in the 8p12 stem cell myeloproliferative disorder with t(8;9)(p12;q33). *Blood* 95: 1788–1796, 2000.

[78843] 17173. Baroni, M. G.; Oelbaum, R. S.; Pozzilli, P.; Stocks, J.; Li, S.-R.; Fiore, V.; Galton, D. J.: Polymorphisms at the GLUT1 (HepG2) and GLUT4 (muscle/adipocyte) glucose transporter genes and non-insulin-dependent diabetes mellitus (NIDDM). *Hum. Genet.* 88: 557–561, 1992.

[78844] 17174. Kelley, M. J.; Pech, M.; Seuanez, H. N.; Rubin, J. S.; O'Brien, S. J.; Aaronson, S. A.: Emergence of the keratinocyte growth factor multigene family during the great ape radiation. *Proc. Nat. Acad. Sci.* 89: 9287–9291, 1992.

[78845] 17175. Mattei, M.-G.; deLapeyriere, O.; Bresnick, J.; Dickson, C.; Birnbaum, D.; Mason, I.: Mouse Fgf7 (fibroblast growth factor 7) and Fgf8 (fibroblast growth factor 8) genes map to chromosomes 2 and 19 respectively. *Mam-*

malianGenome 6: 196–197, 1995.

- [78846] 17176. Rubin, J. S.; Osada, H.; Finch, P. W.; Taylor, W. G.; Rudikoff, S.; Aaronson, S. A.: Purification and characterization of a newly identified growth factor specific for epithelial cells. *Proc. Nat. Acad. Sci.* 86: 802–806, 1989.
- [78847] 17177. Werner, S.; Smola, H.; Liao, X.; Longaker, M. T.; Krieg, T.; Hofschneider, P. H.; Williams, L. T.: The function of KGF in morphogenesis of epithelium and reepithelialization of wounds. *Science* 266: 819–822, 1994.
- [78848] 17178. Zimonjic, D. B.; Kelley, M. J.; Rubin, J. S.; Aaronson, S. A.; Popescu, N. C.: Fluorescence in situ hybridization analysis of keratinocyte growth factor gene amplification and dispersion in evolution of great apes and humans. *Proc. Nat. Acad. Sci.* 94: 11461–11465, 1997.
- [78849] 17179. Edwards, Y. H.; Parkar, M.; Povey, S.; West, L. F.; Parrington, J. M.; Solomon, E.: Human myosin heavy chain genes assigned to chromosome 17 using a human cDNA clone as probe. *Ann. Hum. Genet.* 49: 101–109, 1985.
- [78850] 17180. Rappold, G. A.; Vosberg, H.-P.: Chromosomal localization of a human myosin heavy-chain gene by in situ hybridization. *Hum. Genet.* 65: 195–197, 1983.
- [78851] 17181. Sparkes, R. S.; Zollman, S.; Klisak, I.; Kirchgessner, T. G.; Komaromy, M. C.; Mohandas, T.; Schotz, M. C.; Lu-

sis, A. J.: Humangenom involved in lipolysis of plasma lipoproteins: mapping of loci for lipoprotein lipase to 8p22 and hepatic lipase to 15q21. *Genomics* 1:138–144, 1987.

[78852] 17182. Baumgartner, M.; Holzfeind, P.; Redl, B.: Assignment of the gene for human tear prealbumin (LCN1), a member of the lipocalin superfamily, to chromosome 8q24. *Cytogenet. Cell Genet.* 65: 101–103, 1994.

[78853] 17183. Blaker, M.; Kock, K.; Ahlers, C.; Buck, F.; Schmale, H.: Molecular cloning of human von Ebner's gland protein, a member of the lipocalin superfamily highly expressed in lingual salivary glands. *Biochim. Biophys. Acta* 1172: 131–137, 1993.

[78854] 17184. Liu, Y.; Wu, Y.-P.; Wada, R.; Neufeld, E. B.; Mullin, K. A.; Howard, A. C.; Pentchev, P. G.; Vanier, M. T.; Suzuki, K.; Proia, R. L.: Alleviation of neuronal ganglioside storage does not improve the clinical course of the Niemann–Pick C disease mouse. *Hum. Molec. Genet.* 9:1087–1092, 2000.

[78855] 17185. Ito, M.; Yuan, C.-X.; Malik, S.; Gu, W.; Fondell, J. D.; Yamamura, S.; Fu, Z.-Y.; Zhang, X.; Qin, J.; Roeder, R. G.: Identity between TRAP and SMCC complexes indicates novel pathways for the function of nuclear receptors and diverse mammalian activators. *Molec. Cell* 3:361–370,

1999.

- [78856] 17186. Nagase, T.; Seki, N.; Ishikawa, K.; Tanaka, A.; Nomura, N.: Prediction of the coding sequences of unidentified human genes. V. The coding sequences of 40 new genes (KIAA0161–KIAA0200) deduced by analysis of cDNA clones from human cell line KG-1. *DNA Res.* 3: 17–24, 1996.
- [78857] 17187. Makino, K.; Kuwahara, H.; Masuko, N.; Nishiyama, Y.; Morisaki, T.; Sasaki, J.; Nakao, M.; Kuwano, A.; Nakata, M.; Ushio, Y.; Saya, H.: Cloning and characterization of NE-dlg: a novel human homolog of the *Drosophila* discs large (dlg) tumor suppressor protein interacts with the APC protein. *Oncogene* 14: 2425–2433, 1997.
- [78858] 17188. Stathakis, D. G.; Lee, D.; Bryant, P. J.: DLG3, the gene encoding human neuroendocrine Dlg (NE-Dlg), is located within the 1.8-Mb dystonia-parkinsonism region at Xq13.1. *Genomics* 49: 310–313, 1998.
- [78859] 17189. Maeda, K.; Matsushashi, S.; Hori, K.; Xin, Z.; Mukai, T.; Tabuchi, K.; Egashira, M.; Niikawa, N.: Cloning and characterization of a novel human gene, TM4SF6, encoding a protein belonging to the transmembrane 4 superfamily, and mapped to Xq22. *Genomics* 52: 240–242, 1998.
- [78860] 17190. Todd, S. C.; Doctor, V. S.; Levy, S.: Sequences and

expression of six new members of the tetraspanin/TM4SF family. *Biochim. Biophys. Acta* 1399: 101–104, 1998.

- [78861] 17191. Crew, A. J.; Clark, J.; Fisher, C.; Gill, S.; Grimer, R.; Chand, A.; Shipley, J.; Gusterson, B. A.; Cooper, C. S.: Fusion of SYT to two genes, SSX1 and SSX2, encoding proteins with homology to the Kruppel-associated box in human synovial sarcoma. *EMBO J.* 14: 2333–2340, 1995.
- [78862] 17192. Vaccari, T.; Beltrame, M.; Ferrari, S.; Bianchi, M. E.: Hmg4, a new member of the Hmg1/2 gene family. *Genomics* 49: 247–252, 1998.
- [78863] 17193. Wilke, K.; Wiemann, S.; Gaul, R.; Gong, W.; Poustka, A.: Isolation of human and mouse HMG2a cDNAs: evidence for an HMG2a-specific 3'-prime untranslated region. *Gene* 198: 269–274, 1997.
- [78864] 17194. Anson, D. S.; Blake, D. J.; Winship, P. R.; Birnbaum, D.; Brownlee, G. G.: Nullisomic deletion of the mcf.2 transforming gene in two haemophilia B patients. *EMBO J.* 7: 2795–2799, 1988.
- [78865] 17195. Imbeaud, S.; Belville, C.; Messika-Zeitoun, L.; Rey, R.; di Clemente, N.; Josso, N.; Picard, J.-Y.: A 27 base-pair deletion of the anti-Müllerian type II receptor gene is the most common cause of the persistent Müllerian duct syndrome. *Hum. Molec. Genet.* 5: 1269–1277, 1996.

- [78866] 17196.Imbeaud, S.; Carre-Eusebe, D.; Rey, R.; Belville, C.; Josso, N.;Picard, J.-Y.: Molecular genetics of the persistent Mullerian ductsyndrome: a study of 19 families. Hum. Molec. Genet. 3: 125-131,1994.
- [78867] 17197.Imbeaud, S.; Faure, E.; Lamarre, I.; Mattei, M.-G.; di Clemente,N.; Tizard, R.; Carre-Eusebe, D.; Belville, C.; Tragethon, L.; Tonkin,C.; Nelson, J.; McAuliffe, M.; Bidart, J.-M.; Lababidi, A.; Josso,N.; Cate, R. L.; Picard, J.-V.: In-sensitivity to anti-Mullerian hormonedue to a mutation in the human anti-Mullerian hormone receptor. NatureGenet. 11: 382-388, 1995.
- [78868] 17198.Lang-Muritano, M.; Biason-Lauber, A.; Gitzelmann, C.; Belville,C.; Picard, Y.; Schoenle, E. J.: A novel mutation in the anti-mullerianhormone gene as cause of persistent mullerian duct syndrome. Europ.J. Pediat. 160: 652-654, 2001.
- [78869] 17199.Picard, J.-Y.; Benarous, R.; Guerrier, D.; Josso, N.; Kahn, A.: Cloning and expression of cDNA for anti-mullerian hormone. Proc.Nat. Acad. Sci. 83: 5464-5468, 1986.
- [78870] 17200.Tezcan, I.; Xu, W.; Gurgey, A.; Tuncer, M.; Cetin, M.; Oner, C.;Yetgin, S.; Ersoy, F.; Aizencang, G.; Astrin, K. H.; Desnick, R. J.: Congenital erythropoietic porphyria suc-



cessfully treated by allogeneic bone marrow transplantation. *Blood* 92: 4053–4058, 1998.

- [78871] 17201. Xu, W.; Astrin, K. H.; Desnick, R. J.: Molecular basis of congenital erythropoietic porphyria: mutations in the human uroporphyrinogen III synthase gene. *Hum. Mutat.* 7: 187–192, 1996.
- [78872] 17202. Pennarun, G.; Escudier, E.; Chapelin, C.; Bridoux, A.-M.; Cacheux, V.; Roger, G.; Clement, A.; Goossens, M.; Amselem, S.; Duriez, B.: Loss-of-function mutations in a human gene related to *Chlamydomonas reinhardtii* dynein IC78 result in primary ciliary dyskinesia. *Am. J. Hum. Genet.* 65: 1508–1519, 1999.
- [78873] 17203. Weiss, P.; Tietze, F.; Gahl, W. A.; Seppala, R.; Ashwell, G.: Identification of the metabolic defect in sialuria. *J. Biol. Chem.* 264: 17635–17636, 1989.
- [78874] 17204. Bassi, M. T.; Ramesar, R. S.; Caciotti, B.; Winship, I. M.; DeGrandi, A.; Riboni, M.; Townes, P. L.; Beighton, P.; Ballabio, A.; Borsani, G.: X-linked late-onset sensorineural deafness caused by a deletion involving OA1 and a novel gene containing WD-40 repeats. *Am. J. Hum. Genet.* 64: 1604–1616, 1999.
- [78875] 17205. Converse, P. J.: Personal Communication. Baltimore, Md. 8/24/2001.

- [78876] 17206.Disteche, C. M.; Dinulos, M. B.; Bassi, M. T.; Elliott, R. W.;Rugarli, E. I.: Mapping of the murine Tbl1 gene reveals a new rearrangementbetween mouse and human X chromosomes. *Mammalian Genome* 9: 1062–1064,1998.
- [78877] 17207.Dong, X.; Tsuda, L.; Zavitz, K. H.; Lin, M.; Li, S.; Carthew, R.W.; Zipursky, S. L.: ebi regulates epidermal growth factor receptorsignaling pathways in *Drosophila*. *Genes Dev.* 13: 954–965, 1999.
- [78878] 17208.Matsuzawa, S.; Reed, J. C.: Siah–1, SIP, and Ebi collaborate ina novel pathway for beta–catenin degradation linked to p53 responses. *Molec.Cell* 7: 915–926, 2001.
- [78879] 17209.Zhang, J.; Kalkum, M.; Chait, B. T.; Roeder, R. G.: The N–CoR–HDAC3nuclear receptor corepressor complex inhibits the JNK pathway throughthe integral subunit GPS2. *Molec. Cell* 9: 611–623, 2002.
- [78880] 17210.Mu, J.; Roach, P. J.: Characterization of human glycogenin–2,a self–glucosylating initiator of liver glycogen metabolism. *J. Biol.Chem.* 273: 34850–34856, 1998.
- [78881] 17211.Mu, J.; Skurat, A. V.; Roach, P. J.: Glycogenin–2, a novel self–glucosylatingprotein involved in liver glycogen biosynthesis. *J. Biol. Chem.* 272:27589–27597, 1997.
- [78882] 17212.Delbridge, M. L.; Lingenfelter, P. A.; Disteche, C. M.; MarshallGraves, J. A.: The candidate spermatogenesis

gene RBMY has a homologue on the human X chromosome. (Letter) *Nature Genet.* 22: 223–224, 1999.

[78883] 17213. Le Coniat, M.; Soulard, M.; Della Valle, V.; Larsen, C.-J.; Berger, R.: Localization of the human gene encoding heterogeneous nuclear RNA ribonucleoprotein G (hnRNP-G) to chromosome 6p12. *Hum. Genet.*

88:593–595, 1992.

[78884] 17214. Mazeyrat, S.; Saut, N.; Mattei, M.-G.; Mitchell, M. J.: RBMY evolved on the Y chromosome from a ubiquitously transcribed X-Y identical gene. (Letter) *Nature Genet.* 22: 224–226, 1999.

[78885] 17215. Venables, J. P.; Elliott, D. J.; Makarova, O. V.; Makarov, E. M.; Cooke, H. J.; Eperon, I. C.: RBMY, a probable human spermatogenesis factor, and other hnRNP G proteins interact with Tra2-beta and affect splicing. *Hum. Molec. Genet.* 9: 685–694, 2000.

[78886] 17216. Vogt, P. H.; Affara, N.; Davey, P.; Hammer, M.; Jobling, M. A.; Lau, Y. F.; Mitchell, M.; Schempp, W.; Tyler-Smith, C.; Williams, G.; Yen, P.; Rappold, G. A: Report of the Third International Workshop on Y Chromosome Mapping 1997. Heidelberg, Germany, April 13–16, 1997. *Cytogenet. Cell Genet.* 79: 1–20, 1997.

[78887] 17217. Wilcken, B.; Don, N.; Greenaway, R.; Hammond, J.;

Sosula, L.:Sialuria: a second case. J. Inherit. Metab. Dis. 10: 97–102, 1987.

[78888] 17218.Sutherland, G. R.; Baker, E.; Callen, D. F.; Hyland, V. J.; Wong,G.; Clark, S.; Jones, S. S.; Eglinton, L. K.; Shannon, M. F.; Lopez,A. F.; Vadas, M. A.: Interleukin 4 is at 5q31 and interleukin 6 isat 7p15. Hum. Genet. 79: 335–337, 1988.

[78889] 17219.Symmons, D. P.; Jones, M.; Osborne, J.; Sills, J.; Southwood,T. R.; Woo, P.: Pediatric rheumatology in the United Kingdom: datafrom the British Pediatric Rheumatology Group National DiagnosticRegister. J. Rheum. 23: 1975–1980, 1996.

[78890] 17220.Tosato, G.; Seamon, K. B.; Goldman, N. D.; Sehgal, P. B.; May,L. T.; Washington, G. C.; Jones, K. D.; Pike, S. E.: Monocyte–derivedhuman B–cell growth factor identified as interferon–beta–2 (BSF–2,IL–6). Science 239: 502–504, 1988.

[78891] 17221.Villuendas, G.; San Millan, J. L.; Sancho, J. and Escobar–Morreale,H. F.: The –597 G–A and –174 G–C polymorphisms in the promoter ofthe IL–6 gene are associated with hyperandrogenism. J. Clin. Endocr.Metab. 87: 1134–1141, 2002.

[78892] 17222.Zilberstein, A.; Ruggieri, R.; Korn, J. H.; Revel, M.:

Structure and expression of cDNA and genes for human interferon- $\beta$ -2, a distinct species inducible by growth-stimulatory cytokines. *EMBO J.* 5: 2529–2537, 1986.

[78893] 17223. Gaedigk, R.; Duncan, A. M. V.; Miyazaki, I.; Robinson, B. H.; Dosch, H.-M.: ICA1 encoding p69, a protein linked to the development of type 1 diabetes, maps to human chromosome 7p22. *Cytogenet. Cell Genet.* 66:274–276, 1994.

[78894] 17224. Gaedigk, R.; Karges, W.; Hui, M. F.; Scherer, S. W.; Dosch, H.-M.: Genomic organization and transcript analysis of ICAp69, a target antigen in diabetic autoimmunity. *Genomics* 38: 382–391, 1996.

[78895] 17225. Pietropaolo, M.; Castano, L.; Babu, S.; Buelow, R.; Kuo, Y.-L.S.; Martin, S.; Martin, A.; Powers, A. C.; Prochazka, M.; Naggert, J.; Leiter, E. H.; Eisenbarth, G. S.: Islet cell autoantigen 69 kD (ICA69): molecular cloning and characterization of a novel diabetes-associated autoantigen. *J. Clin. Invest.* 92: 359–371, 1993.

[78896] 17226. Cavalieri, R. L.; Havell, E. A.; Vilcek, J.; Pestka, S.: Synthesis of human interferon by *Xenopus laevis* oocytes: two structural genes for interferons in human cells. *Proc. Nat. Acad. Sci.* 74: 3287–3291, 1977.

[78897] 17227. Chany, C.; Finaz, C.; Weil, D.; Vignal, M.; Van Cong,

N.; de Grouchy,J.: Investigations on the chromosomal localizations of the human and chimpanzee interferon genes: possible role of chromosomes 9 and 13. *Ann. Genet.* 23: 201–207, 1980.

[78898] 17228.Derynck, R.; Content, J.; DeClercq, E.; Volckaert, G.; Tavernier,J.; Devos, R.; Fiers, W.: Isolation and structure of a human fibroblast interferon gene. *Nature* 285: 542–547, 1980.

[78899] 17229.Diaz, M. O.; Le Beau, M. M.; Pitha, P.; Rowley, J. D.: Interferon and c-ets-1 genes in the translocation (9;11)(p22;q23) in human acute monocytic leukemia. *Science* 231: 265–267, 1986.

[78900] 17230.Diaz, M. O.; Ziemin, S.; Le Beau, M. M.; Pitha, P.; Smith, S. D.; Chilcote, R. R.; Rowley, J. D.: Homozygous deletion of the alpha- and beta(1)-interferon genes in human leukemia and derived cell lines. *Proc.Nat. Acad. Sci.* 85: 5259–5263, 1988.

[78901] 17231.Erickson, B. W.; May, L. T.; Sehgal, P. B.: Internal duplication in human alpha-1 and beta-1 interferons. *Proc. Nat. Acad. Sci.* 81:7171–7175, 1984.

[78902] 17232.Henry, L.; Sizun, J.; Turleau, C.; Boue, J.; Azoulay, M.; Junien,C.: The gene for human fibroblast interferon (IFB) maps to 9p21. *Hum.Genet.* 68: 67–69, 1984.

- [78903] 17233.Houghton, M.; Eaton, M. A. W.; Stewart, A. G.; Smith, J. C.; Doel, S. M.; Catlin, G. H.; Lewis, H. M.; Patel, T. P.; Emtage, J. S.; Carey, N. H.; Porter, A. G.: The complete amino acid sequence of human fibroblast interferon as deduced using synthetic oligodeoxyribonucleotide primers of reverse transcriptase. *Nucleic Acids Res.* 8: 2885–2894, 1980.
- [78904] 17234.Houghton, M.; Jackson, I. J.; Porter, A. G.; Doel, S. M.; Catlin, G. H.; Barber, C.; Carey, N. H.: The absence of introns within a human fibroblast interferon gene. *Nucleic Acids Res.* 9: 247–266, 1981.
- [78905] 17235.Knight, E., Jr.: Human fibroblast interferon: amino acid analysis and amino terminal amino acid sequence. *Science* 207: 525–527, 1980.
- [78906] 17236.May, L. T.; Landsberger, F. R.; Inouye, M.; Sehgal, P. B.: Significance of similarities in patterns: an application to beta interferon-related DNA on human chromosome 2. *Proc. Nat. Acad. Sci.* 82: 4090–4094, 1985.
- [78907] 17237.Meager, A.; Graves, H.; Burke, D. C.; Swallow, D. M.: Involvement of a gene on chromosome 9 in human fibroblast interferon production. *Nature* 280:493–495, 1979.
- [78908] 17238.Meager, A.; Graves, H. E.; Walker, J. R.; Burke, D.

C.; Swallow,D. M.; Westerveld, A.: Tentative assignment of the gene for humanfibroblast interferon to chromosome 9. (Abstract) Cytogenet. CellGenet. 25: 183–184, 1979.

- [78909] 17239.Ohlsson, M.; Feder, J.; Cavalli-Sforza, L. L.; von Gabain, A.: Close linkage of alpha and beta interferons and infrequent duplicationof beta interferon in humans. Proc. Nat. Acad. Sci. 82: 4473–4476,1985.
- [78910] 17240.Ohno, S.; Taniguchi, T.: Structure of a chromosomal gene forhuman interferon beta. Proc. Nat. Acad. Sci. 78: 5305–5309, 1981.
- [78911] 17241.Owerbach, D.; Rutter, W. J.; Shows, T. B.; Gray, P.; Goeddel,D. V.; Lawn, R. M.: Leukocyte and fibroblast interferon genes arelocated on human chromosome 9. Proc. Nat. Acad. Sci. 78: 3123–3127,1981.
- [78912] 17242.Pitha, P. M.; Slate, D. L.; Raj, N. B. K.; Ruddle, F. H.: Humanbeta interferon gene localization and expression in somatic cell hybrids. Molec.Cell. Biol. 2: 564–570, 1982.
- [78913] 17243.Sagar, A. D.; Sehgal, P. B.; May, L. T.; Inouye, M.; Slate, D.L.; Shulman, L.; Ruddle, F. H.: Interferon–beta–related DNA is dispersedin the human genome. Science 223: 1312–1315, 1984.
- [78914] 17244.Sehgal, P. B.; May, L. T.; Sagar, A. D.; LaForge, K. S.; Inouye,M.: Isolation of novel human genomic DNA



clones related to humaninterferon-beta(1) cDNA. Proc. Nat. Acad. Sci. 80: 3632-3636, 1983.

- [78915] 17245. Shepard, H. M.; Leung, D.; Stebbing, N.; Goeddel, D. V.: A single amino acid change in IFN-beta(1) abolishes its antiviral activity. Nature 294:563-567, 1981.
- [78916] 17246. Siegal, F. P.; Kadowaki, N.; Shodell, M.; Fitzgerald-Bocarsly, P. A.; Shah, K.; Ho, S.; Antonenko, S.; Liu, Y.-J.: The nature of the principal type 1 interferon-producing cells in human blood. Science 284:1835-1837, 1999.
- [78917] 17247. Erikson, J.; Griffin, C.; ar-Rushdi, A.; Valtieri, M.; Hoxie, J.; Finan, J.; Emanuel, B. S.; Rovera, G.; Nowell, P. C.; Croce, C. M.: Heterogeneity of chromosome 22 breakpoint in Philadelphia-positive (Ph+) acute lymphocytic leukemia. Proc. Nat. Acad. Sci. 83: 1807-1811, 1986.
- [78918] 17248. Fioretos, T.; Heisterkamp, N.; Groffen, J.: No evidence for genomic imprinting of the human BCR gene. Blood 83: 3441-3444, 1994.
- [78919] 17249. Fitzgerald, P. H.: Evidence that chromosome band 22q12 is concerned with cell proliferation in chronic myeloid leukemia. Hum. Genet. 33:269-274, 1976.
- [78920] 17250. Ganesan, T. S.; Rassool, F.; Guo, A.-P.; Th'ng, K. H.; Dowding, C.; Hibbin, J. A.; Young, B. D.; White, H.; Kumaran, T. O.; Galton, D. A. G.; Goldman, J. M.: Rearrange-

ment of the bcr gene in Philadelphia chromosome-negative chronic myeloid leukemia. Blood 68: 957-960,1986.

[78921] 17251. Goldman, J. M.; Melo, J. V.: Targeting the BCR-ABL tyrosine kinase in chronic myeloid leukemia. (Editorial) New Eng. J. Med. 344: 1084-1086,2001.

[78922] 17252. Gorre, M. E.; Mohammed, M.; Ellwood, K.; Hsu, N.; Paquette, R.; Rao, P. N.; Sawyers, C. L.: Clinical resistance to STI-571 cancer therapy caused by BCR-ABL gene mutation or amplification. Science 293:876-883, 2001.

[78923] 17253. Groffen, J.; Stephenson, J. R.; Heisterkamp, N.; de Klein, A.; Bartram, C. R.; Grosveld, G.: Philadelphia chromosomal breakpoints are clustered within a limited region, bcr, on chromosome 22. Cell 36:93-99, 1984.

[78924] 17254. Grossman, A.; Silver, R. T.; Arlin, Z.; Coleman, M.; Camposano, E.; Gascon, P.; Benn, P. A.: Fine mapping of chromosome 22 breakpoints within the breakpoint cluster region (bcr) implies a role for bcr exon 3 in determining disease duration in chronic myeloid leukemia. Am. J. Hum. Genet. 45: 729-738, 1989.

[78925] 17255. Grosveld, G.; Verwoerd, T.; van Agthoven, T.; de Klein, A.; Ramachandran, K. L.; Heisterkamp, N.; Stam, K.; Groffen, J.: The chronic myelocytic cell line K562 contains a breakpoint in bcr and produces a chimeric bcr/c-abl

transcript. *Molec. Cell. Biol.* 6: 607–616, 1986.

[78926] 17256.Haas, O. A.: Are ABL and BCR imprinted? No definitive answers, but more questions. *Leukemia* 9: 740–745, 1995.

[78927] 17257.Haas, O. A.; Argyriou-Tirita, A.; Lion, T.: Parental origin of chromosomes involved in the translocation t(9;22). *Nature* 359: 414–416, 1992.

[78928] 17258.Hariharan, I. K.; Adams, J. M.: cDNA sequence for human bcr, the gene that translocates to the abl oncogene in chronic myeloid leukaemia. *EMBO J.* 6: 115–119, 1987.

[78929] 17259.Heisterkamp, N.; Jenster, G.; ten Hoeve, J.; Zovich, D.; Pattengale, P. K.; Groffen, J.: Acute leukaemia in bcr/abl transgenic mice. *Nature* 344:251–253, 1990.

[78930] 17260.Heisterkamp, N.; Stam, K.; Groffen, J.; de Klein, A.; Grosveld, G.: Structural organization of the bcr gene and its role in the Ph-1 translocation. *Nature* 315: 758–761, 1985.

[78931] 17261.Hermans, A.; Heisterkamp, N.; von Lindern, M.; van Baal, S.; Meijer, D.; van der Plas, D.; Wiedemann, L. M.; Groffen, J.; Bootsma, D.; Grosveld, G.: Unique fusion of bcr and c-abl genes in Philadelphia chromosome positive acute lymphoblastic leukemia. *Cell* 51: 33–40, 1987.

[78932] 17262.Huettner, C. S.; Zhang, P.; Van Etten, R. A.; Tenen,

D. G.: Reversibility of acute B-cell leukaemia induced by BCR-ABL1. *Nature Genet.* 24:57-60, 2000.

[78933] 17263. Jacobs, A.: Benzene and leukemia. *Brit. J. Haemat.* 72: 119-121, 1989.

[78934] 17264. Klein, G.: Specific chromosomal translocations and the genesis of the B-cell-derived tumors in mice and men. *Cell* 32: 311-315, 1983.

[78935] 17265. Koeffler, H. P.; Golde, D. W.: Chronic myelogenous leukemia--new concepts. *New Eng. J. Med.* 304: 1201-1209 and 1269-1274, 1981.

[78936] 17266. Kohno, S.-I.; Sandberg, A. A.: Chromosomes and causation of human cancer and leukemia. XXXIX. Usual and unusual findings in Ph(1)-positive CML. *Cancer* 46: 2227-2237, 1980.

[78937] 17267. Laurent, E.; Talpaz, M.; Kantarjian, H.; Kurzrock, R.: The BCR gene and Philadelphia chromosome-positive leukemogenesis. *Cancer Res.* 61: 2343-2355, 2001.

[78938] 17268. Lillicrap, D. A.; Sterndale, H.: Familial chronic myeloid leukaemia. (Letter) *Lancet* II: 699, 1984.

[78939] 17269. Lim, Y.-M.; Wong, S.; Lau, G.; Witte, O. N.; Colicelli, J.: BCR/ABL inhibition by an escort/phosphatase fusion protein. *Proc. Nat. Acad. Sci.* 97: 12233-12238, 2000.

[78940] 17270. Litz, C. E.; Copenhaver, C. M.: Paternal origin of the

rearranged major breakpoint cluster region in chronic myeloid leukemia. *Blood* 83:3445–3448, 1994.

[78941] 17271. Maru, Y.; Witte, O. N.: The BCR gene encodes a novel serine/threonine kinase activity within a single exon. *Cell* 67: 459–468, 1991.

[78942] 17272. Melo, J. V.; Yan, X.-H.; Diamond, J.; Goldman, J. M.: Lack of imprinting of the ABL gene. (Letter) *Nature Genet.* 8: 318–319, 1994.

[78943] 17273. Melo, J. V.; Yan, X.-H.; Diamond, J.; Goldman, J. M.: Balanced parental contribution to the ABL component of the BCR–ABL gene in chronic myeloid leukemia. *Leukemia* 9: 734–745, 1995.

[78944] 17274. Mes-Masson, A.-M.; McLaughlin, J.; Daley, G. Q.; Paskind, M.; Witte, O. N.: Overlapping cDNA clones define the complete coding region for the P210(c-abl) gene product associated with chronic myelogenous leukemia cells containing the Philadelphia chromosome. *Proc. Nat. Acad. Sci.* 83: 9768–9772, 1986.

[78945] 17275. Mills, K. I.; MacKenzie, E. D.; Birnie, G. D.: The site of the breakpoint within the bcr is a prognostic factor in Philadelphia–positive CML patients. *Blood* 72: 1237–1241, 1988.

[78946] 17276. Mittelman, F.; Levan, G.: Clustering of aberrations

to specific chromosomes in human neoplasms. III. Incidence and geographic distribution of chromosome aberrations in 856 cases. *Hereditas* 89: 207–232, 1978.

[78947] 17277. Nowell, P. C.; Hungerford, D. A.: A minute chromosome in human chronic granulocytic leukemia. *Science* 132: 1497, 1960.

[78948] 17278. Kusari, J.; Takata, Y.; Hatada, E.; Freidenberg, G.; Kolterman, O.; Olefsky, J. M.: Insulin resistance and diabetes due to different mutations in the tyrosine kinase domain of both insulin receptor gene alleles. *J. Biol. Chem.* 266: 5260–5267, 1991.

[78949] 17279. Lander, E. S.; Botstein, D.: Homozygosity mapping: a way to map human recessive traits with the DNA of inbred children. *Science* 236: 1567–1570, 1987.

[78950] 17280. Kramer, F.; White, K.; Pauleikhoff, D.; Gehrig, A.; Passmore, L.; Rivera, A.; Rudolph, G.; Kellner, U.; Andrassi, M.; Lorenz, B.; Rohrschneider, K.; Blankenagel, A.; Jurklies, B.; Schilling, H.; Schutt, F.; Holz, F. G.; Weber, B. H. F.: Mutations in the VMD2 gene are associated with juvenile-onset vitelliform macular dystrophy (Best disease) and adult vitelliform macular dystrophy but not age-related macular degeneration. *Europ. J. Hum. Genet.* 8: 286–292, 2000.

[78951] 17281. Sagane, K.; Ohya, Y.; Hasegawa, Y.; Tanaka, I.:

Metalloproteinase-like, disintegrin-like, cysteine-rich proteins MDC2 and MDC3: novel human cellular disintegrins highly expressed in the brain. *Biochem. J.* 334:93–98, 1998.

[78952] 17282. Alsobrook, J. P., II; Zohar, A. H.; Leboyer, M.; Chabane, N.; Ebstein, R. P.; Pauls, D. L.: Association between the COMT locus and obsessive-compulsive disorder in females but not males. *Am. J. Med. Genet.* (Neuropsychiat. Genet.) 114: 116–120, 2002.

[78953] 17283. Karayiorgou, M.; Sobin, C.; Blundell, M. L.; Galke, B. L.; Malinova, L.; Goldberg, P.; Ott, J.; Gogos, J. A.: Family-based association studies support a sexually dimorphic effect of COMT and MAOA on genetic susceptibility to obsessive-compulsive disorder. *Biol. Psychiat.* 45:1178–1189, 1999.

[78954] 17284. Taira, M.; Yoshida, T.; Miyagawa, K.; Sakamoto, H.; Terada, M.; Sugimura, T.: cDNA sequence of human transforming gene hst and identification of the coding sequence required for transforming activity. *Proc. Nat. Acad. Sci.* 84: 2980–2984, 1987.

[78955] 17285. Krill, A. E.; Morse, P. A.; Potts, A. M.; Klien, B. A.: Hereditary vitelliruptive macular degeneration. *Am. J. Ophthalm.* 61: 1405–1415, 1966.

- [78956] 17286.Maloney, W. F.; Robertson, D. M.; Miller, S. A.: Hereditary vitelliformmacular degeneration--variable fundus findings within a single pedigree. Arch.Ophthal. 95: 979-983, 1977.
- [78957] 17287.Mansergh, F. C.; Kenna, P. F.; Rudolph, G.; Meitinger, T.; Farrar,G. J.; Kumar-Singh, R.; Scorer, J.; Hally, A. M.; Mynett-Johnson,L.; Humphries, M. M.; Kiang, S.; Humphries, P.: Evidence for geneticheterogeneity in Best's vitelliform macular dystrophy. J. Med. Genet. 32:855-858, 1995.
- [78958] 17288.Marmorstein, A. D.; Marmorstein, L. Y.; Rayborn, M.; Wang, X.;Hollyfield, J. G.; Petrukhin, K.: Bestrophin, the product of theBest vitelliform macular dystrophy gene (VMD2), localizes to the basolateralplasma membrane of the retinal pigment epithelium. Proc. Nat. Acad.Sci. 97: 12758-12763, 2000.
- [78959] 17289.Marquardt, A.; Stohr, H.; Passmore, L. A.; Kramer, F.; Rivera,A.; Weber, B. H. F.: Mutations in a novel gene, VMD2, encoding aprotein of unknown properties cause juvenile-onset vitelliform maculardystrophy (Best's disease). Hum. Molec. Genet. 7: 1517-1525, 1998.
- [78960] 17290.Nichols, B. E.; Bascom, R.; Litt, M.; McInnes, R.; Sheffield,V. C.; Stone, E. M.: Refining the locus for Best



vitelliform maculardystrophy and mutation analysis of the candidate gene ROM1. *Am. J. Hum. Genet.* 54: 95–103, 1994.

- [78961] 17291. Nordstrom, S.: Epidemiological studies of hereditary macular degeneration (Best's disease) in Swedish and Swedish–American populations. In: Eriksson, A. W.; Forsius, H. R.; Nevanlinna, H. R.; Workman, P. L.; Norio, R. K.: *Population Structure and Genetic Disorders*. New York: Academic Press (pub.) 1980. Pp. 431–443.
- [78962] 17292. Nordstrom, S.: Personal Communication. Umea, Sweden 1978.
- [78963] 17293. Nordstrom, S.; Thorburn, W.: Dominantly inherited macular degeneration (Best's disease) in a homozygous father with 11 children. *Clin. Genet.* 18: 211–216, 1980.
- [78964] 17294. O'Gorman, S.; Flaherty, W. A.; Fishman, G. A.; Berson, E. L.: Histopathologic findings in Best's vitelliform macular dystrophy. *Arch. Ophthalmol.* 106: 1261–1268, 1988.
- [78965] 17295. Petrukhin, K.; Koisti, M. J.; Bakall, B.; Li, W.; Xie, G.; Marknell, T.; Sandgren, O.; Forsman, K.; Holmgren, G.; Andreasson, S.; Vujic, M.; Bergen, A. A. B.; McGarty–Dugan, V.; Figueroa, D.; Austin, C. P.; Metzker, M. L.; Caskey, C. T.; Wadelius, C.: Identification of the gene responsible for Best macular dystrophy. *Nature Genet.* 19: 241–247, 1998.

- [78966] 17296.Rivas, F.; Ruiz, C.; Rivera, H.; Moller, M.; Serrano-Lucas, J.I.; Cantu, J. M.: De novo del(6)(q25) associated with macular degeneration. *Ann.Genet.* 29: 42–44, 1986.
- [78967] 17297.Rosas, F. E.: Maculopatía hereditaria viteliforme de Best. *Ann.Soc. Mex. Oft.* 50: 157–171, 1976.
- [78968] 17298.Sorsby, A.; Savory, M.; Davey, J. B.; Fraser, R. J. L.: Macular cysts: a dominantly inherited affection with a progressive course. *Brit.J. Ophthal.* 40: 144–158, 1956.
- [78969] 17299.Stohr, H.; Marquardt, A.; Rivera, A.; Cooper, P. R.; Nowak, N.J.; Shows, T. B.; Gerhard, D. S.; Weber, B. H. F.: A gene map of the Best's vitelliform macular dystrophy region in chromosome 11q12–q13.1. *GenomeRes.* 8: 48–56, 1998.
- [78970] 17300.Stone, E. M.; Nichols, B. E.; Streb, L. M.; Kimura, A. E.; Sheffield, V. C.: Genetic linkage of vitelliform macular degeneration (Best's disease) to chromosome 11q13. *Nature Genet.* 1: 246–250, 1992.
- [78971] 17301.Sun, H.; Tsunenari, T.; Yau, K.-W.; Nathans, J.: The vitelliform macular dystrophy protein defines a new family of chloride channels. *Proc.Nat. Acad. Sci.* 99: 4008–4013, 2002.
- [78972] 17302.Vail, D.; Shoch, D.: Hereditary degeneration of the macula. II. Follow-up report and histopathologic study.

Trans. Am. Ophthal. Soc. 63:51–63, 1965.

- [78973] 17303.Vossius, A.: Ueber die Bestsche familiaere Maculadegeneration. Arch.Ophthal. 105: 1050–1059, 1921.
- [78974] 17304.Weber, B. H. F.; Walker, D.; Muller, B.: Molecular evidence for non-penetrance in Best's disease. J. Med. Genet. 31: 388–392, 1994.
- [78975] 17305.Weber, B. H. F.; Walker, D.; Muller, B.; Mar, L.: Best's vitelliform dystrophy (VMD2) maps between D11S903 and PYGM: no evidence for locus heterogeneity. Genomics 20: 267–274, 1994.
- [78976] 17306.White, K.; Marquardt, A.; Weber, B. H. F.: VMD2 mutations in vitelliform macular dystrophy (Best disease) and other maculopathies. Hum.Mutat. 15: 301–308, 2000.
- [78977] 17307.Yoder, F. E.; Cross, H. E.; Chase, G. A.; Fine, S. L.; Freidhoff, L.; Machan, C. H.; Bias, W. B.: Linkage studies of Best's macular dystrophy. Clin. Genet. 34: 26–30, 1988.
- [78978] 17308.Farndon, J. R.; Leight, G. S.; Dilley, W. G.; Baylin, S. B.; Smallridge, R. C.; Harrison, T. S.; Wells, S. A., Jr.: Familial medullary thyroid carcinoma without associated endocrinopathies: a distinct clinical entity. Brit. J. Surg. 73: 278–281, 1986.
- [78979] 17309.Wallis, D. E.; Roessler, E.; Hehr, U.; Nanni, L.; Wiltshire, T.; Richieri-Costa, A.; Gillesen-Kaesbach, G.; Za-

ckai, E. H.; Rommens, J.; Muenke, M.: Mutations in the homeodomain of the human SIX3 gene cause holoprosencephaly. *Nature Genet.* 22: 196–198, 1999.

[78980] 17310. McMichael, A.; Makgoba, W.: Complexity in human histocompatibility loci. *Nature* 293: 701–702, 1981.

[78981] 17311. Moen, T.; Albrechtsen, D.; Flatmark, A.; Jakobsen, A.; Jervell, J.; Halvorsen, S.; Solheim, B. G.; Thorsby, E.: Importance of HLA-DR matching in cadaveric renal transplantation: a prospective one-center study of 170 transplants. *New Eng. J. Med.* 303: 850–854, 1980.

[78982] 17312. Olerup, O.; Troye-Blomberg, M.; Schreuder, G. M. T.; Riley, E. M.: HLA-DR and -DQ gene polymorphism in West Africans is twice as extensive as in North European Caucasians: evolutionary implications. *Proc. Nat. Acad. Sci.* 88: 8480–8484, 1991.

[78983] 17313. Park, M. S.; Terasaki, P. I.; Bernoco, D.: Relationship between MT and DR antigens. In: Terasaki, P. I.: *Histocompatibility Testing 1980*. Los Angeles: UCLA Press (pub.) 1980.

[78984] 17314. Park, M. S.; Terasaki, P. I.; Bernoco, D.; Iwaki, Y.: Evidence for a second B-cell locus separate from the DR locus. *Transplant. Proc.* 10: 823–828, 1978.

[78985] 17315. Pisella, P.-J.; Brignole, F.; Debbasch, C.; Lozato,

P.-A.; Creuzot-Garcher, C.; Bara, J.; Saiag, P.; Warnet, J.-M.; Baudouin, C.: Flow cytometric analysis of conjunctival epithelium in ocular rosacea and keratoconjunctivitis sicca. *Ophthalmology* 107: 1841–1849, 2000.

[78986] 17316. Rollini, P.; Mach, B.; Gorski, J.: Linkage map of three HLA-DR beta-chain genes: evidence for a recent duplication event. *Proc. Nat. Acad. Sci.* 82: 7197–7201, 1985.

[78987] 17317. Sachs, J. A.; Jaraquemada, D.; Festenstein, H.: Intra HLA-D region recombinant maps HLA-DR between HLA-B and HLA-D. *Tissue Antigens* 17:43–56, 1981.

[78988] 17318. Schwartz, B. D.: Personal Communication. St. Louis, Mo. 2/22/1983.

[78989] 17319. Shackelford, D. A.; Mann, D. L.; van Rood, J. J.; Ferrara, G. B.; Strominger, J. L.: Human B-cell alloantigens DC1, MT1, and LB12 are identical to each other but distinct from the HLA-DR antigen. *Proc. Nat. Acad. Sci.* 78: 4566–4570, 1981.

[78990] 17320. Strominger, J. L.: Biology of the human histocompatibility leukocyte antigen (HLA) system and a hypothesis regarding the generation of autoimmune diseases. *J. Clin. Invest.* 77: 1411–1415, 1986.

[78991] 17321. Suciú-Foca, N.; Weiner, J.; Rohowsky, C.; McKiernan, P.; Susinno, E.; Rubinstein, P.: Indications that Dw de-

terminants are controlled by distinct (but closely linked) genes. Transplant. Proc. 10: 799–804, 1978.

[78992] 17322. Thursz, M. R.; Thomas, H. C.; Greenwood, B. M.; Hill, A. V. S.: Heterozygote advantage for HLA class-II type in hepatitis B virus infection. (Letter) Nature Genet. 17: 11–12, 1997.

[78993] 17323. Tiercy, J.-M.; Gorski, J.; Jeannet, M.; Mach, B.: Identification and distribution of three serologically undetected alleles of HLA-DR by oligonucleotide-DNA typing analysis. Proc. Nat. Acad. Sci. 85: 198–202, 1988.

[78994] 17324. Tiwari, J. L.; Terasaki, P. I.: HLA and Disease Associations. New York: Springer-Verlag (pub.) 1985.

[78995] 17325. Trowsdale, J.: Genetics and polymorphism: class II antigens. Brit. Med. Bull. 43: 15–36, 1987.

[78996] 17326. Tsubota, K.; Fukagawa, K.; Fujihara, T.; Shimmura, S.; Saito, I.; Saito, K.; Takeuchi, T.: Regulation of human leukocyte antigen expression in human conjunctival epithelium. Invest. Ophthalm. Vis. Sci. 40: 28–34, 1999.

[78997] 17327. Walker, L. E.; Hewick, R.; Hunkapiller, M. W.; Hood, L. E.; Dreyer, W. J.; Reisfeld, R. A.: N-terminal amino acid sequences of the alpha and beta chains of HLA-DR1 and HLA-DR2 antigens. Biochemistry 23: 185–188, 1983.

[78998] 17328. Geraghty, D. E.; Pei, J.; Lipsky, B.; Hansen, J. A.;

Taillon-Miller, P.; Bronson, S. K.; Chaplin, D. D.: Cloning and physical mapping of the HLA class I region spanning the HLA-E-to-HLA-F interval by using yeast artificial chromosomes. *Proc. Nat. Acad. Sci.* 89: 2669–2673, 1992.

[78999] 17329. McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba, Canada 6/22/1988.

[79000] 17330. Stewart, W. E., II: *The Interferon System*. Berlin: Springer(pub.) 1979.

[79001] 17331. Farfel, Z.; Iiri, T.; Shapira, H.; Roitman, A.; Mouallem, M.; Bourne, H. R.: Pseudohypoparathyroidism, a novel mutation in the beta/gamma-contact region of Gs-alpha impairs receptor stimulation. *J. Biol. Chem.* 271: 19653–19655, 1996.

[79002] 17332. Heisler, L. K.; Cowley, M. A.; Tecott, L. H.; Fan, W.; Low, M. J.; Smart, J. L.; Rubinstein, M.; Tatro, J. B.; Marcus, J. N.; Holstege, H.; Lee, C. E.; Cone, R. D.; Elmquist, J. K.: Activation of central melanocortin pathways by fenfluramine. *Science* 297: 609–611, 2002.

[79003] 17333. Folster-Holst, R.; Moises, H. W.; Yang, L.; Fritsch, W.; Weissenbach, J.; Christophers, E.: Linkage between atopy and the IgE high-affinity receptor gene at 11q13 in atopic dermatitis families. *Hum. Genet.* 102:236–239, 1998.

- [79004] 17334.Hill, M. R.; Cookson, W. O. C. M.: A new variant of the beta subunit of the high-affinity receptor for immunoglobulin E (Fc-epsilon-RI-betaE237G): associations with measures of atopy and bronchial hyper-responsiveness. *Hum.Molec. Genet.* 5: 959-962, 1996.
- [79005] 17335.Hupp, K.; Siwarski, D.; Mock, B. A.; Kinet, J.-P.: Gene mapping of the three subunits of the high affinity FcR for IgE to mouse chromosomes 1 and 19. *J. Immun.* 143: 3787-3791, 1989.
- [79006] 17336.Kuster, H.; Zhang, L.; Brini, A. T.; MacGlashan, D. W. J.; Kinet, J.-P.: The gene and cDNA for the human high affinity immunoglobulin E receptor beta chain and expression of the complete human receptor. *J.Biol. Chem.* 267: 12782-12787, 1992.
- [79007] 17337.Nagata, H.; Mutoh, H.; Kumahara, K.; Arimoto, Y.; Tomemori, T.; Sakurai, D.; Arase, K.; Ohno, K.; Yamakoshi, T.; Nakano, K.; Okawa, T.; Numata, T.; Konno, A.: Association between nasal allergy and a coding variant of the Fc-epsilon-RI-beta gene Glu237Gly in a Japanese population. *Hum. Genet.* 109: 262-266, 2001.
- [79008] 17338.Shirakawa, T.; Li, A.; Dubowitz, M.; Dekker, J. W.; Shaw, A. E.; Faux, J. A.; Ra, C.; Cookson, W. O. C. M.; Hopkin, J. M.: Association between atopy and variants of the



beta subunit of the high-affinity immunoglobulin E receptor. *Nature Genet.* 7: 125–130, 1994.

- [79009] 17339. Shirakawa, T.; Mao, X.-Q.; Sasaki, S.; Enomoto, T.; Kawai, M.; Morimoto, K.; Hopkin, J.: Association between atopic asthma and a coding variant of Fc-epsilon-RI-beta in a Japanese population. *Hum. Molec. Genet.* 5: 1129–1130, 1996.
- [79010] 17340. Szepietowski, P.; Gaudray, P.: FCER1B, a candidate gene for atopy, is located in 11q13 between CD20 and TCN1. *Genomics* 19: 399–400, 1994.
- [79011] 17341. Kuster, H.; Thompson, H.; Kinet, J.-P.: Characterization and expression of the gene for the human Fc receptor gamma subunit: definition of a new gene family. *J. Biol. Chem.* 265: 6448–6452, 1990.
- [79012] 17342. Takayanagi, H.; Kim, S.; Matsuo, K.; Suzuki, H.; Suzuki, T.; Sato, K.; Yokochi, T.; Oda, H.; Nakamura, K.; Ida, N.; Wagner, E. F.; Taniguchi, T.: RANKL maintains bone homeostasis through c-Fos-dependent induction of interferon- $\gamma$ . *Nature* 416: 744–749, 2002.
- [79013] 17343. Taniguchi, T.; Fujii-Kuriyama, Y.; Muramatsu, M.: Molecular cloning of human interferon cDNA. *Proc. Nat. Acad. Sci.* 77: 4003–4006, 1980.
- [79014] 17344. Taniguchi, T.; Mantei, N.; Schwarzenstein, M.; Nagata,

S.; Muramatsu, M.; Weissmann, C.: Human leukocyte and fibroblast interferons are structurally related. *Nature* 285: 547–549, 1980.

[79015] 17345. Taniguchi, T.; Ohno, S.; Fujii-Kuriyama, Y.; Muramatsu, M.: The nucleotide sequence of human fibroblast interferon cDNA. *Gene* 10:11–15, 1980.

[79016] 17346. Tavernier, J.; Derynck, R.; Fiers, W.: Evidence for a unique human fibroblast interferon (IFN- $\beta$ 1) chromosomal gene devoid of intervening sequences. *Nucleic Acids Res.* 9: 461–471, 1981.

[79017] 17347. Tavernier, J.; Gheysen, D.; Duerinck, F.; Van der Heyden, J.; Fiers, W.: Deletion mapping of the inducible promoter of human IFN- $\beta$  gene. *Nature* 301: 634–636, 1983.

[79018] 17348. Trent, J. M.; Olson, S.; Lawn, R.: Chromosomal localization of human leukocyte, fibroblast, and immune interferon genes by means of in situ hybridization. *Proc. Nat. Acad. Sci.* 79: 7809–7813, 1982.

[79019] 17349. Weissenbach, J.; Chernajovsky, Y.; Zeevi, M.; Shulman, L.; Soreq, H.; Nir, U.; Wallach, D.; Perricaudet, M.; Tiollais, P.; Revel, M.: Two interferon mRNAs in human fibroblasts: in vitro translation and *Escherichia coli* cloning studies. *Proc. Nat. Acad. Sci.* 77:7152–7156, 1980.

- [79020] 17350.Zinn, K.; DiMaio, D.; Maniatis, T.: Identification of two distinct regulatory regions adjacent to the human beta-interferon gene. *Cell* 34:865–879, 1983.
- [79021] 17351.Bruns, G. A. P.; Eisenman, R. E.; Gerald, P. S.: Human mitochondrial NADP-dependent isocitrate dehydrogenase in man-mouse somatic cell hybrids. *Cytogenet. Cell Genet.* 17: 200–211, 1976.
- [79022] 17352.Champion, M. J.; Brown, J. A.; Shows, T. B.: Assignment of cytoplasmic alpha-mannosidase (MAN-A) and confirmation of the mitochondrial isocitrate dehydrogenase (IDH-M) genes to the q11-qter region of chromosome 15 in man. *Cytogenet. Cell Genet.* 22: 498–502, 1978.
- [79023] 17353.Grzeschik, K.-H.: Assignment of a gene for human mitochondrial isocitrate dehydrogenase (ICD-M, EC 1.1.1.41.) to chromosome 15. *Hum.Genet.* 34: 23–28, 1976.
- [79024] 17354.Huh, T.-L.; Kim, Y.-O.; Oh, I.-U.; Song, B. J.; Inazawa, J.: Assignment of the human mitochondrial NAD(+)-specific isocitrate dehydrogenase alpha subunit (IDH3A) gene to 15q25.1-q25.2 by in situ hybridization. *Genomics* 32:295–296, 1996.
- [79025] 17355.Oh, I.-U.; Inazawa, J.; Kim, Y.-O.; Song, B. J.; Huh,

T.-L.: Assignment of the human mitochondrial NADP(+)-specific isocitrate dehydrogenase (IDH2) gene to 15q26.1 by in situ hybridization. *Genomics* 38: 104–106, 1996.

[79026] 17356. Shimizu, N.; Giles, R. E.; Kucherlapati, R. S.; Shimizu, Y.; Ruddle, F. H.: Somatic cell genetic assignment of the human gene for mitochondrial NADP-linked isocitrate dehydrogenase to the long arm of chromosome 15. *Somat. Cell Genet.* 3: 47–60, 1977.

[79027] 17357. Appel, S.; Filter, M.; Reis, A.; Hennies, H. C.; Bergheim, A.; Ogilvie, E.; Arndt, S.; Simmons, A.; Lovett, M.; Hide, W.; Ramsay, M.; Reichwald, K.; Zimmermann, W.; Rosenthal, A.: Physical and transcriptional map of the critical region for keratolytic winter erythema (KWE) on chromosome 8p22–p23 between D8S550 and D8S1759. *Europ. J. Hum. Genet.* 10:17–25, 2002.

[79028] 17358. Olavarria, E.; Craddock, C.; Dazzi, F.; Marin, D.; Markt, S.; Apperley, J. F.; Goldman, J. M.: Imatinib mesylate (STI571) in the treatment of relapse of chronic myeloid leukemia after allogeneic stem cell transplantation. *Blood* 99: 3861–3862, 2002.

[79029] 17359. Pegoraro, L.; Matera, L.; Ritz, J.; Levis, A.; Palumbo, A.; Biagini, G.: Establishment of a Ph(1)-positive human

cell line (BV173). J.Nat. Cancer Inst. 70: 447–451, 1983.

[79030] 17360.Perrotti, D.; Cesi, V.; Trotta, R.; Guerzoni, C.; Santilli, G.;Campbell, K.; Iervolino, A.; Condorelli, F.; Gambacorti-Passerini,C.; Caligiuri, M. A.; Calabretta, B.: BCR–ABL suppresses C/EBP–alphaexpression through inhibitory action of hnRNP E2. Nature Genet. 30:48–58, 2002.

[79031] 17361.Prakash, O.; Yunis, J. J.: High resolution chromosomes of thet(9;22) positive leukemias. Cancer Genet. Cytogenet. 11: 361–367,1984.

[79032] 17362.Priest, J. R.; Robison, L. L.; McKenna, R. W.; Lindquist, L. L.;Warkentin, P. I.; LeBien, T. W.; Woods, W. G.; Kersey, J. H.; Coccia,P. F.; Nesbit, M. E., Jr.: Philadelphia chromosome positive childhoodacute lymphoblastic leukemia. Blood 56: 15–22, 1980.

[79033] 17363.Rowley, J. D.: A new consistent chromosomal abnormality in chronicmyelogenous leukemia identified by quinacrine fluorescence and Giemsa staining. Nature 243: 290–293, 1973.

[79034] 17364.Rubin, C. M.; Carrino, J. J.; Dickler, M. N.; Leibowitz, D.; Smith,S. D.; Westbrook, C. A.: Heterogeneity of genomic fusion of BCR andABL in Philadelphia chromosome–positive acute lymphoblastic leukemia. Proc.Nat. Acad. Sci. 85: 2795–2799, 1988.

- [79035] 17365.Saglio, G.; Storlazzi, C. T.; Giugliano, E.; Surace, C.; Anelli,L.; Rege-Cambrin, G.; Zagaria, A.; Velasco, A. J.; Heiniger, A.; Scaravaglio,P.; Gomez, A. T.; Gomez, J. R.; Archidiacono, N.; Banfi, S.; Rocchi,M.: A 76-kb duplicon maps close to the BCR gene on chromosome 22and the ABL gene on chromosome 9: possible involvement in the genesisof the Philadelphia chromosome translocation. Proc. Nat. Acad. Sci. 99:9882–9887, 2002.
- [79036] 17366.Savage, D. G.; Antman, K. H.: Imatinib mesylate--a new oral targetedtherapy. New Eng. J. Med. 346: 683–693, 2002.
- [79037] 17367.Sawyers, C. L.: Chronic myeloid leukemia. New Eng. J. Med. 340:1330–1340, 1999.
- [79038] 17368.Schaefer-Rego, K.; Dudek, H.; Popenoe, D.; Arlin, Z.; Mears, J.G.; Bank, A.; Leibowitz, D.: CML patients in blast crisis have breakpointslocalized to a specific region of the BCR. Blood 70: 448–455, 1987.
- [79039] 17369.Shtivelman, E.; Gale, R. P.; Drezzen, O.; Berrebi, A.; Zaizov,R.; Kubonishi, I.; Miyoshi, I.; Canaani, E.: bcr-abl RNA in patientswith chronic myelogenous leukemia. Blood 69: 971–973, 1987.
- [79040] 17370.Shtivelman, E.; Lifshitz, B.; Gale, R. P.; Canaani, E.: Fusedtranscript of abl and bcr genes in chronic myeloge-

nous leukaemia. Nature 315:550–554, 1985.

[79041] 17371. Skorski, T.; Nieborowska-Skorska, M.; Nicolaides, N. C.; Szczalik, C.; Iversen, P.; Iozzo, R. V.; Zon, G.; Calabretta, B.: Suppression of Philadelphia-1 leukemia cell growth in mice by BCR-ABL antisense oligodeoxynucleotide. Proc. Nat. Acad. Sci. 91: 4504–4508, 1994.

[79042] 17372. Stam, K.; Heisterkamp, N.; Grosveld, G.; de Klein, A.; Verma, R. S.; Coleman, M.; Dosik, H.; Groffen, J.: Evidence of a new chimeric bcr/c-abl mRNA in patients with chronic myelocytic leukemia and the Philadelphia chromosome. New Eng. J. Med. 313: 1429–1433, 1985.

[79043] 17373. Stam, K.; Heisterkamp, N.; Reynolds, F. H., Jr.; Groffen, J.: Evidence that the phl gene encodes a 160,000-dalton phosphoprotein with associated kinase activity. Molec. Cell. Biol. 7: 1955–1960, 1987.

[79044] 17374. Swan, D. C.; McBride, O. W.; Robbins, K. C.; Keithley, D. A.; Reddy, E. P.; Aaronson, S. A.: Chromosomal mapping of the simian sarcoma virus onc gene analogue in human cells. Proc. Nat. Acad. Sci. 79: 4691–4695, 1982.

[79045] 17375. Tanabe, T.; Kuwabara, T.; Warashina, M.; Tani, K.; Taira, K.; Asano, S.: Oncogene inactivation in a mouse model: tissue invasion by leukaemic cells is stalled by loading them with a designer ribozyme. Nature

406:473–474, 2000.

[79046] 17376.Teyssier, J. R.; Bartram, C. R.; Deville, J.; Potron, G.; Pigeon, F.: C-abl oncogene and chromosome 22 'bcr' juxtaposition in chronic myelogenous leukemia. *New Eng. J. Med.* 312: 1393–1394, 1985.

[79047] 17377.Tkachuk, D. C.; Westbrook, C. A.; Andreeff, M.; Donlon, T. A.; Cleary, M. L.; Suryanarayan, K.; Homge, M.; Redner, A.; Gray, J.; Pinkel, D.: Detection of bcr–abl fusion in chronic myelogeneous leukemia by in situ hybridization. *Science* 250: 559–562, 1990.

[79048] 17378.Verhest, A.; Monsieur, R.: Philadelphia chromosome–positive thrombocythemia with leukemic transformation. (Letter) *New Eng. J. Med.* 308: 1603, 1983.

[79049] 17379.Verma, R. S.; Dosik, H.: Heteromorphisms of the Philadelphia(Ph–1) chromosome in patients with chronic myelogenous leukaemia (CML).I. Classification and clinical significance. *Brit. J. Haemat.* 45:215–222, 1980.

[79050] 17380.Cockerham, G. C.; Hidayat, A. A.; Bijwaard, K. E.; Sheng, Z.–M.: Re–evaluation of 'reactive lymphoid hyperplasia of the uvea': an immunohistochemical and molecular analysis of 10 cases. *Ophthalmology* 107:151–158, 2000.

[79051] 17381.Guru, S. C.; Agarwal, S. K.; Manickam, P.; Olufemi, S.–E.; Crabtree, J. S.; Weisemann, J. M.; Kester, M. B.; Kim,



Y. S.; Wang, Y.; Emmert-Buck, M. R.; Liotta, L. A.; Spiegel, A. M.; Boguski, M. S.; Roe, B. A.; Collins, F. S.; Marx, S. J.; Burns, L.; Chandrasekharappa, S. C.: A transcript map for the 2.8-Mb region containing the multiple endocrine neoplasia type 1 locus. *Genome Res.* 7: 725–735, 1997.

[79052] 17382. Frossard, P. M.; Lestringant, G. G.: Association between a dimorphic site on chromosome 12 and clinical diagnosis of hypertension in three independent populations. *Clin. Genet.* 48: 284–287, 1995.

[79053] 17383. Glenn, C. L.; Wang, W. Y. S.; Benjafield, A. V.; Morris, B. J.: Linkage and association of tumor necrosis factor receptor 2 locus with hypertension, hypercholesterolemia and plasma shed receptor. *Hum. Molec. Genet.* 9: 1943–1949, 2000.

[79054] 17384. Lasky, L. A.; Singer, M. S.; Dowbenko, D.; Imai, Y.; Henzel, W. J.; Grimley, C.; Fennie, C.; Gillett, N.; Watson, S. R.; Rosen, S. D.: An endothelial ligand for L-selectin is a novel mucin-like molecule. *Cell* 69: 927–938, 1992.

[79055] 17385. Lasky, L. A.; Singer, M. S.; Yednock, T. A.; Dowbenko, D.; Fennie, C.; Rodriguez, H.; Nguyen, T.; Stachel, S.; Rosen, S. D.: Cloning of a lymphocyte homing receptor reveals a lectin domain. *Cell* 56: 1045–1055, 1989.

[79056] 17386. Ord, D. C.; Ernst, T. J.; Zhou, L.-J.; Rambaldi, A.;

Spertini, O.; Griffin, J.; Tedder, T. F.: Structure of the gene encoding the human leukocyte adhesion molecule-1 (TQ1, Leu-8) of lymphocytes and neutrophils. *J. Biol. Chem.* 265: 7760-7767, 1990.

[79057] 17387. Siegelman, M. H.; Weissman, I. L.: Human homologue of mouse lymphnode homing receptor: evolutionary conservation at tandem cell interaction domains. *Proc. Nat. Acad. Sci.* 86: 5562-5566, 1989.

[79058] 17388. Sitrin, R. G.; Pan, P. M.; Blackwood, R. A.; Huang, J.; Petty, H. R.: Cutting edge: evidence for a signaling partnership between urokinase receptors (CD87) and L-selectin (CD62L) in human polymorphonuclear neutrophils. *J. Immun.* 166: 4822-4825, 2001.

[79059] 17389. Tedder, T. F.; Isaacs, C. M.; Ernst, T. J.; Demetri, G. D.; Adler, D. A.; Disteche, C. M.: Isolation and chromosomal localization of cDNAs encoding a novel human lymphocyte cell surface molecule, LAM-1: homology with the mouse lymphocyte homing receptor and other human adhesion proteins. *J. Exp. Med.* 170: 123-133, 1989.

[79060] 17390. Aizawa, S.; Nakano, H.; Ishida, T.; Horie, R.; Nagai, M.; Ito, K.; Yagita, H.; Okumura, K.; Inoue, J.; Watanabe, T.: Tumor necrosis factor receptor-associated factor (TRAF) 5 and TRAF2 are involved in CD30-mediated NF-kappa-B

activation. J. Biol. Chem. 272: 2042–2045,1997.

- [79061] 17391.Durkop, H.; Latza, U.; Hummel, M.; Eitelbach, F.; Seed, B.; Stein,H.: Molecular cloning and expression of a new member of the nervegrowth factor receptor family that is characteristic for Hodgkin'sdisease. Cell 68: 421–427, 1992.
- [79062] 17392.Kurts, C.; Carbone, F. R.; Krummel, M. F.; Koch, K. M.; Miller,J. F. A. P.; Heath, W. R.: Signalling through CD30 protects againstautoimmune diabetes mediated by CD8 T cells. Nature 398: 341–344,1999.
- [79063] 17393.McClive, P. J.; Morahan, G.: Assignment of the mouse homologuesof 6 loci from HSA1p to chromosomes 3 and 4. Genomics 23: 243–246,1994.
- [79064] 17394.Stein, H.; Gerdes, J.; Schwab, U.; Lemke, H.; Mason, D. Y.; Ziegler,A.; Schienle, W.; Diehl, V.: Identification of Hodgkin and Sternberg–Reedcells as a unique cell type derived from a newly detected small cellpopulation. Int. J. Cancer 30: 445–449, 1982.
- [79065] 17395.de Lau, W.; Clevers, H.: LEF1 turns over a new leaf. Nature Genet. 28:3–5, 2001.
- [79066] 17396.Hovanes, K.; Li, T. W. H.; Munguia, J. E.; Truong, T.; Milovanovic,T.; Marsh, J. L.; Holcombe, R. F.; Waterman, M. L.: Beta–catenin–sensitiveisoforms of lymphoid enhancer

factor-1 are selectively expressed in colon cancer. *Nature Genet.* 28: 53-57, 2001.

[79067] 17397. Love, J. J.; Li, X.; Case, D. A.; Giese, K.; Grosschedl, R.; Wright, P. E.: Structural basis for DNA bending by the architectural transcription factor LEF-1. *Nature* 376: 791-795, 1995.

[79068] 17398. Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.: LEF1, the gene for lymphoid enhancer-binding factor 1, mapped to human chromosome 4 (q23-q25) and distal mouse chromosome 3 by Southern blot analysis and fluorescence in situ hybridization. (Abstract) *Cytogenet. Cell-Genet.* 58: 1888 only, 1991.

[79069] 17399. Milatovich, A.; Travis, A.; Grosschedl, R.; Francke, U.: Gene for lymphoid enhancer-binding factor 1 (LEF1) mapped to human chromosome 4 (q23-q25) and mouse chromosome 3 near *Egf*. *Genomics* 11: 1040-1048, 1991.

[79070] 17400. van Genderen, C.; Okamura, R. M.; Farinas, I.; Quo, R. G.; Parslow, T. G.; Bruhn, L.; Grosschedl, R.: Development of several organs that require inductive epithelial-mesenchymal interactions is impaired in LEF-1-deficient mice. *Genes Dev.* 8: 2691-2703, 1994.

[79071] 17401. Waterman, M. L.; Fischer, W. H.; Jones, K. A.: A thymus-specific member of the HMG protein family regulates

the human T cell receptorC alpha enhancer. *Genes Dev.* 5: 656–669, 1991.

[79072] 17402.Zhou, P.; Byrne, C.; Jacobs, J.; Fuchs, E.: Lymphoid enhancerfactor 1 directs hair follicle patterning and epithelial cell fate. *GenesDev.* 9: 700–713, 1995.

[79073] 17403.White, P. S.; Jensen, S. J.; Rajalingam, V.; Stairs, D.; Sulman,E. P.; Maris, J. M.; Biegel, J. A.; Wooster, R.; Brodeur, G. M.:Physical mapping of the CA6, ENO1, and SLC2A5 (GLUT5) genes and reassignmentof SLC2A5 to 1p36.2. *Cytogenet. Cell Genet.* 81: 60–64, 1998.

[79074] 17404.Contractor, A.; Swanson, G.; Heinemann, S. F.: Kainate receptorsare involved in short- and long-term plasticity at mossy fiber synapsesin the hippocampus. *Neuron* 29: 209–216, 2001.

[79075] 17405.Paschen, W.; Hedreen, J. C.; Ross, C. A.: RNA editing of the glutamatereceptor subunits GluR2 and GluR6 in human brain tissue. *J. Neurochem.* 63:1596–1602, 1994.

[79076] 17406.Barbon, A.; Barlati, S.: Genomic organization, proposed alternativesplicing mechanisms, and RNA editing structure of GRIK1. *Cytogenet.Cell Genet.* 88: 236–239, 2000.

[79077] 17407.Eubanks, J. H.; Puranam, R. S.; Kleckner, N. W.; Bettler, B.; Heinemann,S. F.; McNamara, J. O.: The gene en-

coding the glutamate receptorsubunit GluR5 is located on human chromosome 21q21.1–22.1 in the vicinityof the gene for familial amyotrophic lateral sclerosis. Proc. Nat.Acad. Sci. 90: 178–182, 1993.

- [79078] 17408.Gregor, P.; Gaston, S. M.; Yang, X.; O'Regan, J. P.; Rosen, D.R.; Tanzi, R. E.; Patterson, D.; Haines, J. L.; Horvitz, H. R.; Uhl,G. R.; Brown, R. H., Jr.: Genetic and physical mapping of the GLUR5glutamate receptor gene on human chromosome 21. Hum. Genet. 94:565–570, 1994.
- [79079] 17409.Gregor, P.; Reeves, R. H.; Jabs, E. W.; Yang, X.; Dackowski, W.;Rochelle, J. M.; Brown, R. H., Jr.; Haines, J. L.; O'Hara, B. F.;Uhl, G. R.; Seldin, M. F.: Chromosomal localization of glutamatereceptor genes: relationship to familial amyotrophic lateral sclerosisand other neurological disorders of mice and humans. Proc. Nat. Acad.Sci. 90: 3053–3057, 1993.
- [79080] 17410.Gregor, P.; Seldin, M. F.; Reeve, R.; Jabs, E.; Yang, X.; Rochelle,J. M.; O'Hara, B. F.; Uhl, G. R.: Chromosomal localization of glutamatereceptor genes: candidates for familial amyotrophic lateral sclerosisand other neurological disorders of mice and man. (Abstract) Neurosci.Abstracts 18: 395 only, 1992.

- [79081] 17411.Paschen, W.; Djuricic, B.: Extent of RNA editing of glutamate receptor subunit GluR5 in different brain regions of the rat. *Cell.Molec. Neurobiol.* 14: 259–270, 1994.
- [79082] 17412.Potier, M.–C.; Dutriaux, A.; Lambolez, B.; Bochet, P.; Rossier,J.: Assignment of the human glutamate receptor gene GLUR5 to 21q22by screening a chromosome 21 YAC library. *Genomics* 15: 696–697,1993.
- [79083] 17413.Sander, T.; Hildmann, T.; Kretz, R.; Furst, R.; Sailer, U.; Bauer,G.; Schmitz, B.; Beck–Mannagetta, G.; Wienker, T. F.; Janz, D.: Allelicassociation of juvenile absence epilepsy with a gluR5 kainate receptorgene (GRIK1) polymorphism. *Am. J. Med. Genet.* 74: 416–421, 1997.
- [79084] 17414.Smolders, I.; Bortolotto, Z. A.; Clarke, V. R. J.; Warre, R.;Khan, G. M.; O'Neill, M. J.; Ornstein, P. L.; Bleakman, D.; Ogden,A.; Weiss, B.; Stables, J. P.; Ho, K. H.; Ebinger, G.; Collingridge,G. L.; Lodge, D.; Michotte, Y.: Antagonists of GLU–K5–containingkainate receptors prevent pilocarpine–induced limbic seizures. *NatureNeurosci.* 5: 796–804, 2002.
- [79085] 17415.McNamara, J. O.; Eubanks, J. H.; McPherson, J. D.; Wasmuth, J.J.; Evans, G. A.; Heinemann, S. F.: Chromosomal localization of humanglutamate receptor genes. *J.*

Neurosci. 12: 2555–2562, 1992.

[79086] 17416. Armstrong, N.; Gouaux, E.: Mechanisms for activation and antagonism of an AMPA-sensitive glutamate receptor: crystal structures of the GluR2 ligand binding core. Neuron 28: 165–181, 2000.

[79087] 17417. Brusa, R.; Zimmermann, F.; Koh, D.-S.; Feldmeyer, D.; Gass, P.; Seeburg, P. H.; Sprengel, R.: Early-onset epilepsy and postnatal lethality associated with an editing-deficient GluR-B allele in mice. Science 270: 1677–1680, 1995.

[79088] 17418. Hollmann, M.; Hartley, M.; Heinemann, S.:  $\text{Ca}^{2+}$  permeability of KA-AMPA-gated glutamate receptor channels depends on subunit composition. Science 252: 851–853, 1991.

[79089] 17419. Kohler, M.; Kornau, H. C.; Seeburg, P. H.: The organization of the gene for the functionally dominant  $\alpha$ -amino-3-hydroxy-5-methylisoxazole-4-propionic acid receptor subunit Glu4-B. J. Biol. Chem. 269: 17367–17370, 1994.

[79090] 17420. Sun, W.; Ferrer-Montiel, A. V.; Schinder, A. F.; McPherson, J. P.; Evans, G. A.; Montal, M.: Molecular cloning, chromosomal mapping, and functional expression of human brain glutamate receptors. Proc. Nat. Acad. Sci.



89: 1443–1447, 1992.

- [79091] 17421.Sun, Y.; Olson, R.; Horning, M.; Armstrong, N.; Mayer, M.; Gouaux,E.: Mechanism of glutamate receptor desensitization. *Nature* 417:245–253, 2002.
- [79092] 17422.Chui, D.; Oh–Eda, M.; Liao, Y.–F.; Panneerselvam, K.; Lai, A.;Marek, K. W.; Freeze, H. H.; Moremen, K. W.; Fukuda, M. N.; Marth,J. D.: Alpha–mannosidase–II deficiency results in dyserythropoiesisand unveils an alternate pathway in oligosaccharide biosynthesis. *Cell* 90:157–167, 1997.
- [79093] 17423.Chui, D.; Sellakumar, G.; Green, R. S.; Sutton–Smith, M.; McQuistan,T.; Marek, K. W.; Morris, H. R.; Dell, A.; Marth, J. D.: Geneticremodeling of protein glycosylation in vivo induces autoimmune disease. *Proc.Nat. Acad. Sci.* 98: 1142–1147, 2001.
- [79094] 17424.Gasparini, P.; del Giudice, E. M.; Delaunay, J.; Totaro, A.; Granatiero,M.; Melchionda, S.; Zelante, L.; Iolascon, A.: Localization of thecongenital dyserythropoietic anemia II locus to chromosome 20q11.2by genomewide search. *Am. J. Hum. Genet.* 61: 1112–1116, 1997.
- [79095] 17425.Misago, M.; Liao, Y.–F.; Kudo, S.; Eto, S.; Mattei, M.–G.; Moremen,K. W.; Fukuda, M. N.: Molecular cloning and expression of cDNAs encodinghuman alpha–

mannosidase II and a previously unrecognized alpha-mannosidasell(X) isozyme. Proc. Nat. Acad. Sci. 92: 11766–11770, 1995.

[79096] 17426. Moremen, K. W.; Robbins, P. W.: Isolation, characterization, and expression of cDNAs encoding murine alpha-mannosidase II, a Golgi enzyme that controls conversion of high mannose to complex N-glycans. J. Cell Biol. 115: 1521–1534, 1991.

[79097] 17427. Wada, A.; Sakamoto, H.; Katoh, O.; Yoshida, T.; Yokota, J.; Little, P. F. R.; Sugimura, T.; Terada, M.: Two homologous oncogenes, HST1 and INT2, are closely located in human genome. Biochem. Biophys. Res. Commun. 157: 828–835, 1988.

[79098] 17428. Yoshida, T.; Tsutsumi, M.; Sakamoto, H.; Miyagawa, K.; Teshima, S.; Sugimura, T.; Terada, M.: Expression of the HST1 oncogene in human germ cell tumors. Biochem. Biophys. Res. Commun. 155: 1324–1329, 1988.

[79099] 17429. Nimmo, E.; Padua, R.-A.; Hughes, D.; Brook, J. D.; Williamson, R.; Johnson, K. J.: Confirmation and refinement of the localisation of the c-MEL locus on chromosome 19 by physical and genetic mapping. Hum. Genet. 81: 382–384, 1989.

[79100] 17430. Nimmo, E.; Williamson, R.; Johnson, K.: Localization

of the c-MELgene to 19(cen-p13.2). (Abstract) Cytogenet. Cell Genet. 51: 1053only, 1989.

[79101] 17431.Nimmo, E. R.; Sanders, P. G.; Padua, R. A.; Hughes, D.; Williamson,R.; Johnson, K. J.: The MEL gene: a new member of the RAB/YPT classof RAS-related genes. Onco-gene 6: 1347-1351, 1991.

[79102] 17432.Padua, R. A.; Barrass, N.; Currie, G. A.: A novel transforminggene in a human malignant melanoma cell line. Nature 311: 671-673,1984.

[79103] 17433.Spurr, N. K.; Hughes, D.; Goodfellow, P. N.; Brook, J. D.; Padua,R. A.: Chromosomal assignment of c-MEL, a human transforming oncogene,to chromosome 19(p13.2-q13.2). Somat. Cell Molec. Genet. 12: 637-640,1986.

[79104] 17434.Dabiri, G. A.; Young, C. L.; Rosenbloom, J.; South-wick, F. S.:Molecular cloning of human macrophage cap-ping protein cDNA: a uniquemember of the gelsolin/villin family expressed primarily in macrophages. J.Biol. Chem. 267: 16545-16552, 1992.

[79105] 17435.Mishra, V. S.; Henske, E. P.; Kwiatkowski, D. J.; Southwick, F.S.: The human actin-regulatory protein cap G: gene structure andchromosome location. Genomics 23: 560-565, 1994.

- [79106] 17436.Southwick, F. S.; DiNubile, M. J.: Rabbit alveolar macrophages contain a  $\text{Ca}^{2+}$ -sensitive, 41,000-Dalton protein which reversibly blocks the 'barbed' ends of actin filaments but does not sever them. *J.Biol. Chem.* 261: 14191-14195, 1986.
- [79107] 17437.Eichbaum, Q.; Clerc, P.; Bruns, G.; McKeon, F.; Ezekowitz, R. A.B.: Assignment of the human macrophage mannose receptor gene (MRC1) to 10p13 by in situ hybridization and PCR-based somatic cell hybrid mapping. *Genomics* 22: 656-658, 1994.
- [79108] 17438.Harris, N.; Peters, L. L.; Eicher, E. M.; Rits, M.; Raspberry, D.; Eichbaum, Q. G.; Super, M.; Ezekowitz, R. A. B.: The exon-intron structure and chromosomal localization of the mouse macrophage mannose receptor gene, *Mcr1*: identification of a ricin-like domain at the N-terminus of the receptor. *Biochem. Biophys. Res. Commun.* 198:682-692, 1994.
- [79109] 17439.Kim, S. J.; Ruiz, N.; Bezouska, K.; Drickamer, K.: Organization of the gene encoding the human macrophage mannose receptor (MRC1). *Genomics* 14:721-727, 1992.
- [79110] 17440.Lee, S. J.; Evers, S.; Roeder, D.; Parlow, A. F.; Risteli, J.; Risteli, L.; Lee, Y. C.; Feizi, T.; Langen, H.; Nussenzweig, M. C.: Mannose receptor-mediated regulation of serum

glycoprotein homeostasis. *Science* 295:1898–1901, 2002.

- [79111] 17441. Bernhagen, J.; Calandra, T.; Mitchell, R. A.; Martin, S. B.; Tracey, K. J.; Voelter, W.; Manogue, K. R.; Cerami, A.; Bucala, R.: MIF is a pituitary-derived cytokine that potentiates lethal endotoxaemia. *Nature* 365:756–759, 1993.
- [79112] 17442. Bloom, B. R.; Bennett, B.: Mechanism of a reaction in vitro associated with delayed-type hypersensitivity. *Science* 153: 80–82, 1966.
- [79113] 17443. Bozza, M.; Kolakowski, L. F., Jr.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; David, J. R.; Gerard, C.: Structural characterization and chromosomal location of the mouse macrophage migration inhibitory factor gene and pseudogenes. *Genomics* 27: 412–419, 1995.
- [79114] 17444. Bucala, R.: MIF rediscovered: cytokine, pituitary hormone, and glucocorticoid-induced regulator of the immune response. *FASEB J.* 10:1607–1613, 1996.
- [79115] 17445. Budarf, M.; McDonald, T.; Sellinger, B.; Kozak, C.; Graham, C.; Wistow, G.: Localization of the human gene for macrophage migration inhibitory factor (MIF) to chromosome 22q11.2. *Genomics* 39: 235–236, 1997.
- [79116] 17446. Cato, A. C. B.; Sillar, G. M.; Kioussis, J.; Burdon, R. H.: Molecular cloning of cDNA sequences coding for the major (beta-, gamma-, delta-, and epsilon) heat-shock

polypeptides of HeLa cells. *Gene* 16: 27–34, 1981.

- [79117] 17447. Cummings, C. J.; Sun, Y.; Opal, P.; Antalffy, B.; Mestril, R.; Orr, H. T.; Dillmann, W. H.; Zoghbi, H. Y.: Over-expression of inducible HSP70 chaperone suppresses neuropathology and improves motor function in SCA1 mice. *Hum. Molec. Genet.* 10: 1511–1518, 2001.
- [79118] 17448. Gaskins, H. R.; Prochazka, M.; Nadeau, J. H.; Henson, V. W.; Leiter, E. H.: Localization of a mouse heat shock Hsp70 gene within the H-2 complex. *Immunogenetics* 32: 286–289, 1990.
- [79119] 17449. Goate, A. M.; Cooper, D. N.; Hall, C.; Leung, T. K. C.; Solomon, E.; Lim, L.: Localization of a human heat-shock HSP70 gene sequence to chromosome 6 and detection of two other loci by somatic-cell hybrid and restriction fragment length polymorphism analysis. *Hum. Genet.* 75: 123–128, 1987.
- [79120] 17450. Fukui, K.; Oh, J.; Frenk, E.; Almodovar, C.; Spritz, R. A.: Linkage disequilibrium mapping of the gene for Hermansky-Pudlak syndrome to chromosome 10q23.1–q23.3. *Hum. Molec. Genet.* 4: 1665–1669, 1995.
- [79121] 17451. Gwynn, B.; Eicher, E. M.; Peters, L. L.: Genetic localization of Cd63, a member of the transmembrane 4 superfamily, reveals two distinct loci in the mouse genome. *Ge-*

nomics 35: 389–391, 1996.

- [79122] 17452.Hotta, H.; Ross, A. H.; Huebner, K.; Isobe, M.; Wendeborn, S.;Chao, M. V.; Ricciardi, R. P.; Tsujimoto, Y.; Croce, C. M.; Koprowski,H.: Molecular cloning and characterization of an antigen associatedwith early stages of melanoma tumor progression. Cancer Res. 48:2955–2962, 1988.
- [79123] 17453.Metzelaar, M. J.; Wijngaard, P. L. J.; Peters, P. J.; Sixma, J.J.; Nieuwenhuis, H. K.; Clevers, H. C.: CD63 antigen: a novel lysosomalmembrane glycoprotein, cloned by a screening procedure for intracellularantigens in eukaryotic cells. J. Biol. Chem. 266: 3239–3245, 1991.
- [79124] 17454.Nishibori, M.; Cham, B.; McNicol, A.; Shalev, A.; Jain, N.; Gerrard,J. M.: The protein CD63 is in platelet dense granules, is deficientin a patient with Hermansky–Pudlak syndrome, and appears identicalto granulophysin. J. Clin. Invest. 91: 1775–1782, 1993.
- [79125] 17455.Azizi, E.; Friedman, J.; Pavlovsky, F.; Iscovich, J.; Bornstein,A.; Shafir, R.; Trau, H.; Brenner, H.; Nass, D.: Familial cutaneousmalignant melanoma and tumors of the nervous system. Cancer 76:1571–1578, 1995.
- [79126] 17456.Doege, K.; Rhodes, C.; Sasaki, M.; Hassell, J. R.; Yamada, Y.:Molecular biology of cartilage proteoglycan

(aggrecan) and link protein. In: Sandel, L. J.; Boyd, C. D.: Extracellular Matrix Genes. New York: Academic Press (pub.) 1990. Pp. 137–152.

[79127] 17457. Finkelstein, J. E.; Doege, K.; Yamada, Y.; Pyeritz, R. E.; Graham, J. M., Jr.; Moeschler, J. B.; Pauli, R. M.; Hecht, J. T.; Francomano, C. A.: Analysis of the chondroitin sulfate proteoglycan core protein (CSPGCP) gene in achondroplasia and pseudoachondroplasia. *Am. J. Hum. Genet.* 48: 97–102, 1991.

[79128] 17458. Just, W.; Klett, C.; Vetter, U.; Vogel, W.: Assignment of the human aggrecan gene AGC1 to 15q25–q26.2 by in situ hybridization. *Hum. Genet.* 92: 516–518, 1993.

[79129] 17459. Kimata, K.; Barrach, H. J.; Brown, K. S.; Penny-packer, J. P.: Absence of proteoglycan core protein in cartilage from the cmd/cmd (cartilage deficiency) mouse. *J. Biol. Chem.* 256: 6961–6968, 1981.

[79130] 17460. Korenberg, J. R.; Chen, X. N.; Doege, K.; Grover, J.; Roughley, P. J.: Assignment of the human aggrecan gene (AGC1) to 15q26 using fluorescence in situ hybridization analysis. *Genomics* 16: 546–548, 1993.

[79131] 17461. Lane, P. W.: New mutants and linkages: small with kinky tail. *Mouse Newsletter* 80: 165, 1988.

[79132] 17462. Watanabe, H.; Kimata, K.; Line, S.; Strong, D.; Gao,



L.; Kozak, C. A.; Yamada, Y.: Mouse cartilage matrix deficiency (cmd) caused by a 7 bp deletion in the aggrecan gene. *Nature Genet.* 7: 154–157, 1994.

- [79133] 17463. Watanabe, H.; Nakata, K.; Kimata, K.; Nakanishi, I.; Yamada, Y.: Dwarfism and age-associated spinal degeneration of heterozygote cmd mice defective in aggrecan. *Proc. Nat. Acad. Sci.* 94: 6943–6947, 1997.
- [79134] 17464. Fong, L. Y. Y.; Fidanza, V.; Zanesi, N.; Lock, L. F.; Siracusa, L. D.; Mancini, R.; Siprashvili, Z.; Ottey, M.; Martin, S. E.; Druck, T.; McCue, P. A.; Croce, C. M.; Huebner, K.: Muir–Torre–like syndrome in Fhit-deficient mice. *Proc. Nat. Acad. Sci.* 97: 4742–4747, 2000.
- [79135] 17465. Azen, E. A.; Carlson, D. M.; Clements, S.; Lalley, P. A.; Vanin, E.: Salivary proline-rich protein genes on chromosome 8 of mouse. *Science* 226:967–969, 1984.
- [79136] 17466. Azen, E. A.; Davisson, M. T.; Cherry, M.; Taylor, B. A.: Prp (proline-rich protein) genes linked to markers Es-12 (esterase-12), Ea-10 (erythrocyte alloantigen), and loci on distal mouse chromosome 6. *Genomics* 5:415–422, 1989.
- [79137] 17467. Azen, E. A.; Denniston, C. L.: Genetic polymorphism of human salivary proline-rich proteins, further genetic analysis. *Biochem. Genet.* 12:109–120, 1974.

- [79138] 17468.Azen, E. A.; Oppenheim, F. G.: Genetic polymorphism of proline-rich human salivary proteins. *Science* 180: 1067–1069, 1973.
- [79139] 17469.Baird, M.; Neuweiler, J.; Balazs, I.: Linkage studies between proline rich-protein (SPC) genes and RFLPs from chromosome 12. (Abstract) *Cytogenet.Cell Genet.* 40: 573–574, 1985.
- [79140] 17470.Degand, P.; Aubert, J. P.; Boersma, A.; Richet, C.; Loucheux-Lefebvre, M. H.; Biserte, G.: Parotid alpha-amylase activity: a possible role for proline-rich proteins. *FEBS Lett.* 63: 137–140, 1976.
- [79141] 17471.Goodman, P. A.; Yu, P.-L.; Azen, E. A.; Karn, R. C.: The human salivary protein complex (SPC): a large block of related genes. *Am.J. Hum. Genet.* 37: 785–797, 1985.
- [79142] 17472.Oppenheim, F. G.; Hay, D. I.; Fraublau, C.: Proline-rich proteins from human parotid saliva. Isolation and partial characterization. *Biochemistry* 10:4233–4238, 1971.
- [79143] 17473.Adkins, S.; Gan, K. N.; Mody, M.; La Du, B. N.: Molecular basis for the polymorphic forms of human serum paraoxonase/arylesterase: glutamine or arginine at position 191, for the respective A and B allozymes. *Am. J. Hum. Genet.* 52: 598–608, 1993.
- [79144] 17474.Antikainen, M.; Murtomaki, S.; Syvanne, M.;

Pahlman, R.; Tahvanainen, E.; Jauhiainen, M.; Frick, M. H.; Ehnholm, C.: The gln-arg191 polymorphism of the human paraoxonase gene (HUMPONA) is not associated with the risk of coronary artery disease in Finns. *J. Clin. Invest.* 98: 883–885, 1996.

[79145] 17475. Augustinsson, K.-B.; Henricson, B.: A genetically controlled esterase in rat plasma. *Biochim. Biophys. Acta* 124: 323–331, 1966.

[79146] 17476. Barbieri, M.; Bonafe, M.; Marfella, R.; Ragno, E.; Giugliano, D.; Franceschi, C.; Paolisso, G.: LL-paraoxonase genotype is associated with a more severe degree of homeostasis model assessment IR in healthy subjects. *J. Clin. Endocr. Metab.* 87: 222–225, 2002.

[79147] 17477. Brophy, V. H.; Jampsa, R. L.; Clendenning, J. B.; McKinstry, L. A.; Jarvik, G. P.; Furlong, C. E.: Effects of 5-prime regulatory-region polymorphisms on paraoxonase-gene (PON1) expression. *Am. J. Hum. Genet.* 68: 1428–1436, 2001.

[79148] 17478. Cherry, N.; Mackness, M.; Durrington, P.; Povey, A.; Dippnall, M.; Smith, T.; Mackness, B.: Paraoxonase (PON1) polymorphisms in farmers attributing ill health to sheep dip. *Lancet* 359: 763–764, 2002.

[79149] 17479. Clendenning, J. B.; Humbert, R.; Green, E. D.;

Wood, C.; Traver,D.; Furlong, C. E.: Structural organization of the human PON1 gene. *Genomics* 35:586–589, 1996.

[79150] 17480.Coates, P. M.; Mestriner, M. A.; Hopkinson, D. A.: A preliminaryinterpretation of the esterase isozymes of human tissues. *Ann. Hum.Genet.* 39: 1–20, 1975.

[79151] 17481.Davies, H. G.; Richter, R. J.; Keifer, M.; Broomfield, C. A.; Sowalla,J.; Furlong, C. E.: The effect of the human serum paraoxonase polymorphismis reversed with diazoxon, soman and sarin. *Nature Genet.* 14: 334–336,1996.

[79152] 17482.Deakin, S.; Leviev, I.; Nicaud, V.; Meynet, M.–C. B.; Tiret, L.;James, R. W.: Paraoxonase–1 L55M polymorphism is associated withan abnormal oral glucose tolerance test and differentiates high riskcoronary disease families. *J. Clin. Endocr. Metab.* 87: 1268–1273,2002.

[79153] 17483.Eckerson, H. W.; Romson, J.; Wyte, C.; La Du, B. N.: The humanserum paraoxonase polymorphism: identification of phenotypes by theirresponse to salts. *Am. J. Hum. Genet.* 35: 214–227, 1983.

[79154] 17484.Eckerson, H. W.; Wyte, C. M.; La Du, B. N.: The human serum paraoxonase/arylesterasepolymorphism. *Am. J. Hum. Genet.* 35: 1126–1136, 1983.

[79155] 17485.Eiberg, H.; Mohr, J.: Linkage relations of the

paraoxonase polymorphism with 43 marker systems.

(Abstract) Cytogenet. Cell Genet. 25: 150, 1979.

[79156] 17486. Rey-Campos, J.; Chouard, T.; Yaniv, M.; Cereghini, S.: vHNF1 is a homeoprotein that activates transcription and forms heterodimers with HNF1. EMBO J. 10: 1445–1457, 1991.

[79157] 17487. Shih, D. Q.; Bussen, M.; Sehayek, E.; Ananthanarayanan, M.; Shneider, B. L.; Suchy, F. J.; Shefer, S.; Bollileni, J. S.; Gonzalez, F. J.; Breslow, J. L.; Stoffel, M.: Hepatocyte nuclear factor-1- $\alpha$  is an essential regulator of bile acid and plasma cholesterol metabolism. Nature-Genet. 27: 375–382, 2001.

[79158] 17488. Urhammer, S. A.; Hansen, T.; Ekstrom, C. T.; Eiberg, H.; Pederson, O.: The Ala/Val98 polymorphism of the hepatocyte nuclear factor-1- $\alpha$  gene contributes to the interindividual variation in serum C-peptide response during an oral glucose tolerance test: evidence from studies of 231 glucose-tolerant first degree relatives of type 2 diabetic probands. J. Clin. Endocr. Metab. 83: 4506–4509, 1998.

[79159] 17489. Urhammer, S. A.; Moller, A. M.; Nyholm, B.; Ekstrom, C. T.; Eiberg, H.; Clausen, J. O.; Hansen, T.; Pedersen, O.; Schmitz, O.: The effect of two frequent amino acid

variants of the hepatocyte nuclear factor-1- $\alpha$  gene on estimates of the pancreatic beta-cell function in Caucasian glucose-tolerant first-degree relatives of type 2 diabetic patients. *J. Clin. Endocr. Metab.* 83: 3992–3995, 1998.

[79160] 17490. Urhammer, S. A.; Rasmussen, S. K.; Kaisaki, P. J.; Oda, N.; Yamagata, K.; Moller, A. M.; Fridberg, M.; Hansen, L.; Hansen, T.; Bell, G. I.; Pedersen, O.: Genetic variation in the hepatocyte nuclear factor-1- $\alpha$  gene in Danish Caucasians with late-onset NIDDM. *Diabetologia* 40:473–475, 1997.

[79161] 17491. Vaxillaire, M.; Rouard, M.; Yamagata, K.; Oda, N.; Kaisaki, P. J.; Boriraj, V. V.; Chevre, J.-C.; Boccio, V.; Cox, R. D.; Lathrop, G. M.; Dussoix, P.; Philippe, J.; Timsit, J.; Charpentier, G.; Velho, G.; Bell, G. I.; Froguel, P.: Identification of nine novel mutations in the hepatocyte nuclear factor 1  $\alpha$  gene associated with maturity-onset diabetes of the young (MODY3). *Hum. Molec. Genet.* 6: 583–586, 1997.

[79162] 17492. Yamada, S.; Nishigori, H.; Onda, H.; Utsugi, T.; Yanagawa, T.; Maruyama, T.; Onigata, K.; Nagashima, K.; Nagai, R.; Morikawa, A.; Takeuchi, T.; Takeda, J.: Identification of mutations in the hepatocyte nuclear factor

(HNF)-1-alpha gene in Japanese subjects with IDDM. Diabetes 46:1643-1647, 1997.

- [79163] 17493. Yamagata, K.; Oda, N.; Kalsaki, P. J.; Menzel, S.; Furuta, H.; Vaxillaire, M.; Southam, L.; Cox, R. D.; Lathrop, G. M.; Borhaj, V.V.; Chen, X.; Cox, N. J.; Oda, Y.; Yano, H.; Le Beau, M. M.; Yamada, S.; Nishigori, H.; Takeda, J.; Fajans, S. S.; Hattersley, A. T.; Iwasaki, N.; Hansen, T.; Pedersen, O.; Polonsky, K. S.; Turner, R. C.; Velho, G.; Chevre, J.-C.; Froguel, P.; Bell, G. I.: Mutations in the hepatocytenuclear factor-1-alpha gene in maturity-onset diabetes of the young (MODY3). Nature 384: 455-457, 1996.
- [79164] 17494. Yoshiuchi, I.; Yamagata, K.; Yoshimoto, M.; Zhu, Q.; Yang, Q.; Nammo, T.; Uenaka, R.; Kinoshita, E.; Hanafusa, T.; Miyagawa, J.; Matsuzawa, Y.: Analysis of a non-functional HNF-1-alpha (TCF1) mutation in Japanese subjects with familial type 1 diabetes. Hum. Mutat. 18:345-351, 2001.
- [79165] 17495. Dhanasekaran, S. M.; Barrette, T. R.; Ghosh, D.; Shah, R.; Varambally, S.; Kurachi, K.; Pienta, K. J.; Rubin, M. A.; Chinnaiyan, A. M.: Delineation of prognostic biomarkers in prostate cancer. Nature 412:822-826, 2001.
- [79166] 17496. Leytus, S. P.; Loeb, K. R.; Hagen, F. S.; Kurachi, K.; Davie, E.W.: A novel trypsin-like serine protease (hepsin)

with a putative transmembrane domain expressed by human liver and hepatoma cells. *Biochemistry* 27:1067–1074, 1988.

- [79167] 17497. Tsuji, A.; Torres–Rosado, A.; Arai, T.; Le Beau, M. M.; Lemons, R. S.; Chou, S.–H.; Kurachi, K.: Hepsin, a cell membrane–associated protease: characterization, tissue distribution, and gene localization. *J. Biol. Chem.* 266: 16948–16953, 1991.
- [79168] 17498. Wu, Q.; Yu, D.; Post, J.; Halks–Miller, M.; Sadler, J. E.; Morser, J.: Generation and characterization of mice deficient in hepsin, a hepatic transmembrane serine protease. *J. Clin. Invest.* 101: 321–326, 1998.
- [79169] 17499. Cannella, B.; Hoban, C. J.; Gao, Y.–L.; Garcia–Arenas, R.; Lawson, D.; Marchionni, M.; Gwynne, D.; Raine, C. S.: The neuregulin, glial growth factor 2, diminishes autoimmune demyelination and enhances remyelination in a chronic relapsing model for multiple sclerosis. *Proc. Nat. Acad. Sci.* 95: 10100–10105, 1998.
- [79170] 17500. Falls, D. L.; Rosen, K. M.; Corfas, G.; Lane, W. S.; Fischbach, G. D.: ARIA, a protein that stimulates acetylcholine receptor synthesis, is a member of the Neu ligand family. *Cell* 72: 801–815, 1993.
- [79171] 17501. Fernandez, P.–A.; Tang, D. G.; Cheng, L.; Prochi–



antz, A.; Mudge, A. W.; Raff, M. C.: Evidence that axon-derived neuregulin promotes oligodendrocyte survival in the developing rat optic nerve. *Neuron* 28:81–90, 2000.

[79172] 17502. Holmes, W. E.; Sliwkowski, M. X.; Akita, R. W.; Henzel, W. J.; Lee, J.; Park, J. W.; Yansura, D.; Abadi, N.; Raab, H.; Lewis, G. D.; Shepard, H. M.; Kuang, W.-J.; Wood, W. I.; Goeddel, D. V.; Vandlen, R. L.: Identification of heregulin, a specific activator of p185(erbB2). *Science* 256:1205–1210, 1992.

[79173] 17503. Huang, Y. Z.; Won, S.; Ali, D. W.; Wang, Q.; Tanowitz, M.; Du, Q. S.; Pelkey, K. A.; Yang, D. J.; Xiong, W. C.; Salter, M. W.; Mei, L.: Regulation of neuregulin signaling by PSD-95 interacting with ErbB4 at CNS synapses. *Neuron* 26: 443–455, 2000.

[79174] 17504. Lee, J.; Wood, W. I.: Assignment of heregulin (HGL) to human chromosome 8p22–p11 by PCR analysis of somatic cell hybrid DNA. *Genomics* 16:790–791, 1993.

[79175] 17505. Martinou, J.-C.; Falls, D. L.; Fischbach, G. D.; Merlie, J. P.: Acetylcholine receptor-inducing activity stimulates expression of the epsilon-subunit gene of the muscle acetylcholine receptor. *Proc. Nat. Acad. Sci.* 88: 7669–7673, 1991.

[79176] 17506. Ni, C.-Y.; Murphy, M. P.; Golde, T. E.; Carpenter,

G.: Gamma-secretase cleavage and nuclear localization of ErbB-4 receptor tyrosine kinase. *Science* 294:2179–2181, 2001.

[79177] 17507. Orr-Urtreger, A.; Trakhtenbrot, L.; Ben-Levy, R.; Wen, D.; Rechavi, G.; Lonai, P.; Yarden, Y.: Neural expression and chromosomal mapping of NEU differentiation factor to 8p12–p21. *Proc. Nat. Acad. Sci.* 90:1867–1871, 1993.

[79178] 17508. Boyd, D.; Jain, S. K.; Crampton, J.; Barrett, K. J.; Drysdale, J.: Isolation and characterization of a cDNA clone for human ferritin heavy chain. *Proc. Nat. Acad. Sci.* 81: 4751–4755, 1984.

[79179] 17509. Caskey, J. H.; Jones, C.; Miller, Y. E.; Seligman, P. A.: Human ferritin gene is assigned to chromosome 19. *Proc. Nat. Acad. Sci.* 80:482–486, 1983.

[79180] 17510. Costanzo, F.; Colombo, M.; Staempfli, S.; Santoro, C.; Marone, M.; Frank, R.; Delius, H.; Cortese, R.: Structure of gene and pseudogenes of human apoferritin H. *Nucleic Acids Res.* 14: 721–736, 1986.

[79181] 17511. Cragg, S. J.; Drysdale, J.; Worwood, M.: Genes for the 'H' subunit of human ferritin are present on a number of human chromosomes. *Hum. Genet.* 71: 108–112, 1985.

[79182] 17512. Eisenstein, R. S.: Iron regulatory proteins and the

molecular control of mammalian iron metabolism. *Annu. Rev. Nutr.* 20: 627–662, 2000.

- [79183] 17513. Ferreira, C.; Bucchini, D.; Martin, M.–E.; Levi, S.; Arosio, P.; Grandchamp, B.; Beaumont, C.: Early embryonic lethality of H ferritin gene deletion in mice. *J. Biol. Chem.* 275: 3021–3024, 2000.
- [79184] 17514. Ferreira, C.; Santambrogio, P.; Martin, M.–E.; Andrieu, V.; Feldmann, G.; Henin, D.; Beaumont, C.: H ferritin knockout mice: a model of hyperferritinemia in the absence of iron overload. *Blood* 98: 525–532, 2001.
- [79185] 17515. Gailani, M. R.; Petty, E. M.; Horsthemke, B.; Arnold, A.; Marx, S. J.; Bale, A. E.: Physical mapping of chromosome 11q12–13 by pulsed field gel electrophoresis (PFGE). (Abstract) *Cytogenet. Cell Genet.* 58:1959 only, 1991.
- [79186] 17516. Gatti, R. A.; Shaked, R.; Mohandas, T. K.; Salser, W.: Human ferritin genes: chromosomal assignments and polymorphisms. *Am. J. Hum. Genet.* 41: 654–667, 1987.
- [79187] 17517. Harrison, P. M.; Arosio, P.: The ferritins: molecular properties, iron storage function and cellular regulation. *Biochim. Biophys. Acta* 1275: 161–203, 1996.
- [79188] 17518. Hentze, M. W.; Keim, S.; Papadopoulos, P.; O'Brien, S.; Modi, W.; Drysdale, J.; Leonard, W. J.; Harford, J. B.; Klausner, R. D.: Cloning, characterization, expression, and

chromosomal localization of a human ferritin heavy-chain gene. *Proc. Nat. Acad. Sci.* 83:7226–7230, 1986.

- [79189] 17519. Kato, J.; Fujikawa, K.; Kanda, M.; Fukuda, N.; Sasaki, K.; Takayama, T.; Kobune, M.; Takada, K.; Takimoto, R.; Hamada, H.; Ikeda, T.; Niitsu, Y.: A mutation, in the iron-responsive element of H ferritin mRNA, causing autosomal dominant iron overload. *Am. J. Hum. Genet.* 69:191–197, 2001.
- [79190] 17520. Leibold, E. A.; Munro, H. N.: Cytoplasmic protein binds in vitro to a highly conserved sequence in the 5-prime untranslated region of ferritin heavy- and light-subunit mRNAs. *Proc. Nat. Acad. Sci.* 85:2171–2175, 1988.
- [79191] 17521. Anderson, L. A.; Friedman, L.; Osborne-Lawrence, S.; Lynch, E.; Weissenbach, J.; Bowcock, A.; King, M.-C.: High-density genetic map of the BRCA1 region of chromosome 17q12–q21. *Genomics* 17: 618–623, 1993.
- [79192] 17522. Fryns, J. P.; Kleczkowska, A.; Moerman, P.; Vandenberghe, K.: Hereditary hydronephrosis and the short arm of chromosome 6. (Letter) *Hum. Genet.* 91: 514–515, 1993.
- [79193] 17523. Groenen, P. M. A.; Garcia, E.; Debeer, P.; Devriendt, K.; Fryns, J. P.; Van de Ven, W. J. M.: Structure, sequence,

and chromosome19 localization of human USF2 and its rearrangement in a patient withmulticystic renal dysplasia. Genomics 38: 141–148, 1996.

- [79194] 17524.Groenen, P. M. A.; Vanderlinden, G.; Devriendt, K.; Fryns, J.-P.;Van de Ven, W. J. M.: Rearrangement of the human CDC5L gene by at(6;19)(p21;q13.1) in a patient with multicystic renal dysplasia. Genomics 49:218–229, 1998.
- [79195] 17525.Galizzi, J.-P.; Zuber, C. E.; Harada, N.; Gorman, D. M.; Kastelein,R.; Banchereau, J.; Howard, M.; Miyajima, A.: Molecular cloning of a cDNA encoding the human interleukin 4 receptor. Int. Immun. 2:669–675, 1990.
- [79196] 17526.Grimbacher, B.; Holland, S. M.; Puck, J. M.: The interleukin–4receptor variant Q576R in hyper-IgE syndrome. (Letter) New Eng. J.Med. 338: 1073–1074, 1998.
- [79197] 17527.Idzerda, R. L.; March, C. J.; Mosley, B.; Lyman, S. D.; VandenBos,T.; Gimpel, S. D.; Din, W. S.; Grabstein, K. H.; Widmer, M. B.; Park,L. S.; Cosman, D.; Beckmann, M. P.: Human interleukin–4 receptorconfers biological responsiveness and defines a novel receptor superfamily. J.Exp. Med. 171: 861–873, 1990.
- [79198] 17528.Khurana Hershey, G. K.; Friedrich, M. F.; Esswein, L. A.; Thomas,M. L.; Chatila, T. A.: The association of atopy with a gain-of-functionmutation in the alpha subunit of

the interleukin-4 receptor. *NewEng. J. Med.* 337:  
1720-1725, 1997.

- [79199] 17529. Kruse, S.; Japha, T.; Tedner, M.; Sparholt, S. H.; Forster, J.; Kuehr, J.; Deichmann, K. A.; Abello, A.: The polymorphisms S503P and Q576R in the interleukin-4 receptor alpha gene are associated with atopy and influence the signal transduction. *Immunology* 96:365-371, 1999.
- [79200] 17530. Mitsuyasu, H.; Izuhara, K.; Mao, X.-Q.; Gao, P.-S.; Arinobu, Y.; Enomoto, T.; Kawai, M.; Sasaki, S.; Dake, Y.; Hamasaki, N.; Shirakawa, T.; Hopkin, J. M.: Ile50val variant of IL4R-alpha upregulates IgE synthesis and associates with atopic asthma. (Letter) *Nature Genet.* 19:119-120, 1998.
- [79201] 17531. Olson, M.; Hood, L.; Cantor, C.; Botstein, D.: A common language for physical mapping of the human genome. *Science* 245: 1434-1435, 1989.
- [79202] 17532. Sutherland, G. R.; Baker, E.; Fernandez, K. E. W.; Callen, D. F.; Goodwin, R. G.; Lupton, S.; Namen, A. E.; Shannon, M. F.; Vadas, M. A.: The gene for human interleukin 7 (IL7) is at 8q12-13. *Hum. Genet.* 82:371-372, 1989.
- [79203] 17533. Watanabe, M.; Ueno, Y.; Yajima, T.; Iwao, Y.; Tsuchiya, M.; Ishikawa, H.; Aiso, S.; Hibi, T.; Ishii, H.: Inter-

leukin 7 is produced by humanintestinal epithelial cells and regulates the proliferation of intestinalmucosal lymphocytes. J. Clin. Invest. 95: 2945–2953, 1995.

- [79204] 17534. Corcoran, A. E.; Riddell, A.; Krooshoop, D.; Venkataraman, A. R.: Impaired immunoglobulin gene rearrangement in mice lacking the IL–7receptor. Nature 391: 904–907, 1998.
- [79205] 17535. Ober, C.; Leavitt, S. A.; Tsalenko, A.; Howard, T. D.; Hoki, D.M.; Daniel, R.; Newman, D. L.; Wu, X.; Parry, R.; Lester, L. A.; Solway, J.; Blumenthal, M.; King, R. A.; Xu, J.; Meyers, D. A.; Bleecker, E. R.; Cox, N. J.: Variation in the interleukin 4–receptor alphas gene confers susceptibility to asthma and atopy in ethnically diverse populations. Am. J. Hum. Genet. 66: 517–526, 2000.
- [79206] 17536. Ober, C.; Tsalenko, A.; Willadsen, S.; Newman, D.; Daniel, R.; Wu, X.; Andal, J.; Hoki, D.; Schneider, D.; True, K.; Schou, C.; Parry, R.; Cox, N.: Genome–wide screen for atopy susceptibility alleles in the Hutterites. Clin. Exp. Allergy 29 (suppl. 4): 11–15, 1999.
- [79207] 17537. Patuzzo, C.; Trabetti, E.; Malerba, G.; Martinati, L. C.; Boner, A. L.; Pesciollderungg, L.; Zanoni, G.; Pignatti, P. F.: No linkage or association of the IL–4R–alpha gene Q576R mutation with atopic asthma in Italian families.

(Letter) J. Med. Genet. 37: 382–384,2000.

- [79208] 17538.Pritchard, M. A.; Baker, E.; Whitmore, S. A.; Sutherland, G. R.; Idzerda, R. L.; Park, L. S.; Cosman, D.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Beckmann, M. P.: The interleukin-4 receptor gene (IL4R) maps to 16p11.2–16p12.1 in human and to the distal region of mouse chromosome 7. Genomics 10: 801–806, 1991.
- [79209] 17539.Suzuki, H.; Chung, F.; Palmer, E.; Sasaki, T.; Ohara, N.; Taylor, B. A.; Ohara, J.-I.: Gene mapping of mouse IL-4 receptor: the locus of interleukin 4 (IL-4) receptor gene and lymphocyte function associated antigen 1 (LFA-1) gene are closely linked on chromosome 7. Immunogenetics 34:252–256, 1991.
- [79210] 17540.Zurawski, S. M.; Vega, F., Jr.; Huyghe, B.; Zurawski, G.: Receptors for interleukin-13 and interleukin-4 are complex and share a novel component that functions in signal transduction. EMBO J. 12: 2663–2670,1993.
- [79211] 17541.Koshland, M. E.: The coming of age of the immunoglobulin J chain. Annu.Rev. Immun. 3: 425–453, 1985.
- [79212] 17542.Max, E. E.; McBride, O. W.; Morton, C. C.; Robinson, M. A.: Human J chain gene: chromosomal localization and associated restriction fragment length polymorphisms.



Proc. Nat. Acad. Sci. 83: 5592–5596,1986.

- [79213] 17543.Jones, C.; Penny, L.; Mattina, T.; Yu, S.; Baker, E.; Voullaire,L.; Langdon, W. Y.; Sutherland, G. R.; Richards, R. I.; Tunnacliffe,A.: Association of a chromosome deletion syndrome with a fragilesite within the proto-oncogene CBL2. Nature 376: 145–149, 1995.
- [79214] 17544.Brunton, L. L.; Lupton, S. D.: An STS in the human IL7 gene locatedat 18q12–13. Nucleic Acids Res. 18: 1315 only, 1990.
- [79215] 17545.Carvalho, T. L.; Mota–Santos, T.; Cumano, A.; Demengeot, J.; Vieira,P.: Arrested B lymphopoiesis and persistence of activated B cellsin adult interleukin 7–/– mice. J. Exp. Med. 194: 1141–1150, 2001.
- [79216] 17546.Goodwin, R. G.; Lupton, S.; Schmierer, A.; Hjerrild, K. J.; Jerzy,R.; Clevenger, W.; Gillis, S.; Cosman, D.; Namen, A. E.: Human interleukin7: molecular cloning and growth factor activity on human and murineB–lineage cells. Proc. Nat. Acad. Sci. 86: 302–306, 1989.
- [79217] 17547.Namen, A. E.; Lupton, S.; Hjerrild, K.; Wignall, J.; Mochizuki,D. Y.; Schmierer, A.; Mosley, B.; March, C. J.; Urdal, D.; Gillis,S.; Cosman, D.; Goodwin, R. G.: Stimulation of B–cell progenitorsby cloned murine interleukin–7. Nature 333: 571–573, 1988.

- [79218] 17548.Namen, A. E.; Schmierer, A. E.; March, C. J.; Overell, R. W.; Park,L. S.; Urdal, D. L.; Mochizuki, D. Y.: B-cell precursor growth-promoting activity: purification and characterization of a growth factor active on lymphocyte precursors. *J. Exp. Med.* 167: 988–1002, 1988.
- [79219] 17549.Goodwin, R. G.; Friend, D.; Ziegler, S. F.; Jerzy, R.; Falk, B.A.; Gimpel, S.; Cosman, D.; Dower, S. K.; March, C. J.; Namen, A.E.; Park, L. S.: Cloning of the human and murine interleukin–7 receptors:demonstration of a soluble form and homology to a new receptor superfamily. *Cell* 60:941–951, 1990.
- [79220] 17550.Kondo, M.; Takeshita, T.; Higuchi, M.; Nakamura, M.; Sudo, T.;Nishikawa, S.–I.; Sugamura, K.: Functional participation of the IL–2receptor gamma chain in IL–7 receptor complexes. *Science* 263: 1453–1454,1994.
- [79221] 17551.Lai, S. Y.; Molden, J.; Goldsmith, M. A.: Shared gamma(c) subunit within the human interleukin–7 receptor complex: a molecular basis for the pathogenesis of X-linked severe combined immunodeficiency. *J.Clin. Invest.* 99: 169–177, 1997.
- [79222] 17552.Lynch, M.; Baker, E.; Park, L. S.; Sutherland, G. R.; Goodwin,R. G.: The interleukin–7 receptor gene is at 5p13. *Hum. Genet.* 89:566–568, 1992.

- [79223] 17553.Maki, K.; Sunaga, S.; Komagata, Y.; Kodaira, Y.; Mabuchi, A.; Karasuyama,H.; Yokomuro, K.; Miyazaki, J.; Ikuta, K.: Interleukin 7 receptor-deficientmice lack gamma/delta T cells. Proc. Nat. Acad. Sci. 93: 7172-7177,1996.
- [79224] 17554.Noguchi, M.; Nakamura, Y.; Russell, S. M.; Ziegler, S. F.; Tsang,M.; Cao, X.; Leonard, W. J.: Interleukin-2 receptor gamma chain:a functional component of the interleukin-7 receptor. Science 262:1877-1880, 1993.
- [79225] 17555.Peschon, J. J.; Morrissey, P. J.; Grabstein, K. H. et al.: Earlylymphocyte expansion is severely impaired in interleukin 7 receptor-deficientmice. J. Exp. Med. 180: 1955-1960, 1994.
- [79226] 17556.Puel, A.; Ziegler, S. F.; Buckley, R. H.; Leonard, W. J.: DefectiveIL7R expression in T-B+NK+ severe combined immunodeficiency. NatureGenet. 20: 394-397, 1998.
- [79227] 17557.Mezei, M. M.; Mankodi, A.; Brais, B.; Marineau, C.; Thornton, C.A.; Rouleau, G. A.; Karpati, G.: Minimal expansion of the GCG repeatin the PABP2 gene does not predispose to sporadic inclusion body myositis. Neurology 52:669-670, 1999.
- [79228] 17558.Aleman, A.; Verhaar, H. J. J.; De Haan, E. H. F.; De Vries, W.R.; Samson, M. M.; Drent, M. L.; Van Der Veen, E.

A.; Koppeschaar, H. P. F.: Insulin-like growth factor-I and cognitive function in healthy older men. *J. Clin. Endocr. Metab.* 84: 471–475, 1999.

[79229] 17559. Baker, J.; Liu, J.-P.; Robertson, E. J.; Efstratiadis, A.: Role of insulin-like growth factors in embryonic and post-natal growth. *Cell* 75:73–82, 1993.

[79230] 17560. Bowcock, A.; Sartorelli, V.: Polymorphism and mapping of the IGF1 gene, and absence of association with stature among African Pygmies. *Hum. Genet.* 85: 349–354, 1990.

[79231] 17561. Camacho-Hubner, C.; Woods, K. A.; Miraki-Moud, F.; Hindmarsh, P. C.; Clark, A. J.; Hansson, Y.; Johnston, A.; Baxter, R. C.; Savage, M. O.: Effects of recombinant human insulin-like growth factor I (IGF-I) therapy on the growth hormone-IGF system of a patient with a partial IGF-I gene deletion. *J. Clin. Endocr. Metab.* 84: 1611–1616, 1999.

[79232] 17562. Goddard, A. D.; Covello, R.; Luoh, S.-M.; Clackson, T.; Attie, K. M.; Gesundheit, N.; Rundle, A. C.; Wells, J. A.; Carlsson, L. M. S.: Mutations of the growth hormone receptor in children with idiopathic short stature. *New Eng. J. Med.* 333: 1093–1098, 1995.

[79233] 17563. Guler, H.-P.; Binz, K.; Eigenmann, E.; Jaggi, S.; Zimmermann, D.; Zapf, J.; Froesch, E. R.: Small stature and in-

sulin-like growth factors:prolonged treatment of mini-poodles with recombinant human insulin-likegrowth factor I. *Acta Endocr.* 121: 456-464, 1989.

[79234] 17564.Fajans, S. S.: Maturity-onset diabetes of the young (MODY). *DiabetesMetab. Rev.* 5: 579-606, 1989.

[79235] 17565.Grosz, M. D.; Womack, J. E.; Skow, L. C.: Syntenic conservationof HSP70 genes in cattle and humans. *Genomics* 14: 863-868, 1992.

[79236] 17566.Harrison, G. S.; Drabkin, H. A.; Kao, F.-T.; Hartz, J.; Hart, I.M.; Chu, E. H. Y.; Wu, B. J.; Morimoto, R. I.: Chromosomal locationof human genes encoding major heat-shock protein HSP70. *Somat. CellMolec. Genet.* 13: 119-130, 1987.

[79237] 17567.Harrison, G. S.; Morimoto, R.; Kao, F.-T.; Chu, E. H. Y.; Wu, B.J.; Drabkin, H.: Chromosomal location of human genes encoding themajor heat shock protein HSP70. (Abstract) *Am. J. Hum. Genet.* 39:A157 only, 1986.

[79238] 17568.Hickey, E.; Brandon, S. E.; Sadis, S.; Smale, G.; Weber, L. A.: Molecular cloning of sequences encoding the human heat-shock proteinsand their expression during hyperthermia. *Gene* 43: 147-154, 1986.

[79239] 17569.Imai, Y.; Soda, M.; Hatakeyama, S.; Akagi, T.; Hashikawa, T.; Nakayama,K.; Takahashi, R.: CHIP is asso-

ciated with Parkin, a gene responsible for familial Parkinson's disease, and enhances its ubiquitin ligase activity. *Molec. Cell* 10: 55–67, 2002.

- [79240] 17570. Laroia, G.; Cuesta, R.; Brewer, G.; Schneider, R. J.: Control of mRNA decay by heat shock–ubiquitin–proteasome pathway. *Science* 284:499–502, 1999.
- [79241] 17571. Leung, T. K. C.; Hall, C.; Rajendran, M.; Spurr, N. K.; Lim, L.: The human heat–shock genes HSPA6 and HSPA7 are both expressed and localize to chromosome 1. *Genomics* 12: 74–79, 1992.
- [79242] 17572. Milner, C. M.; Campbell, R. D.: Polymorphic analysis of the three MHC–linked HSP70 genes. *Immunogenetics* 36: 357–362, 1992.
- [79243] 17573. Milner, C. M.; Campbell, R. D.: Structure and expression of the three MHC–linked HSP70 genes. *Immunogenetics* 32: 242–251, 1990.
- [79244] 17574. Pelham, H. R. B.: Speculations on the functions of the major heat shock and glucose–regulated proteins. *Cell* 46: 959–961, 1986.
- [79245] 17575. Sargent, C. A.; Dunham, I.; Trowsdale, J.; Campbell, R. D.: Human major histocompatibility complex contains genes for the major heat shock protein HSP70. *Proc. Nat. Acad. Sci.* 86: 1968–1972, 1989.

- [79246] 17576.Slater, A.; Cato, A. C. B.; Sillar, G. M.; Kioussis, J.; Burdon,R. H.: The pattern of protein synthesis induced by heat shock ofHeLa cells. *Europ. J. Biochem.* 117: 341–346, 1981.
- [79247] 17577.Leung, T. K. C.; Rajendran, M. Y.; Monfries, C.; Hall, C.; Lim,L.: The human heat–shock protein family: expres–sion of a novel heat–inducibleHSP70 (HSP70B–prime) and isolation of its cDNA and genomic DNA. *Biochem.J.* 267: 125–132, 1990.
- [79248] 17578.Schiller, P.; Amin, J.; Ananthan, J.; Brown, M. E.; Scott, W. A.;Voellmy, R.: Cis–acting elements involved in the regulated expressionof a human HSP70 gene. *J. Molec. Biol.* 203: 97–105, 1988.
- [79249] 17579.Voellmy, R.; Ahmed, A.; Schiller, P.; Bromley, P.; Rungger, D.: Isolation and functional analysis of a human 70,000–dalton heatshock protein gene segment. *Proc. Nat. Acad. Sci.* 82: 4949–4953,1985.
- [79250] 17580.Rebbe, N. F.; Hickman, W. S.; Ley, T. J.; Stafford, D. W.; Hickman,S.: Nucleotide sequence and regulation of a human 90–kDa heat shockprotein gene. *J. Biol. Chem.* 264: 15006–15011, 1989.
- [79251] 17581.Wolpowitz, D.; Mason, T. B. A.; Dietrich, P.; Mendelsohn, M.;Talmage, D. A.; Role, L. W.: Cysteine–rich

domain isoforms of the neuregulin-1 gene are required for maintenance of peripheral synapses. *Neuron* 25:79–91, 2000.

[79252] 17582. Bolton, P.; Powell, J.; Rutter, M.; Buckle, V.; Yates, J. R. W.; Ishikawa-Brush, Y.; Monaco, A. P.: Autism, mental retardation, multiple exostoses and short stature in a female with 46,X,t(X;8)(p22.13;q22.1). *Psychiat. Genet.* 5: 51–55, 1995.

[79253] 17583. David, G.; van der Schueren, B.; Marynen, P.; Cassiman, J.-J.; van den Berghe, H.: Molecular cloning of amphiglycan, a novel integral membrane heparan sulfate proteoglycan expressed by epithelial and fibroblastic cells. *J. Cell Biol.* 118: 961–969, 1992.

[79254] 17584. Ishikawa-Brush, Y.; Powell, J. F.; Bolton, P.; Miller, A. P.; Francis, F.; Willard, H. F.; Lehrach, H.; Monaco, A. P.: Autism and multiple exostoses associated with an X;8 translocation occurring within the GRPR gene and 3-prime to the SDC2 gene. *Hum. Molec. Genet.* 6: 1241–1250, 1997.

[79255] 17585. Spring, J.; Goldberger, O. A.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Bernfield, M.: Mapping of the syndecan genes in the mouse: linkage with members of the Myc gene family. *Genomics* 21:597–601, 1994.



- [79256] 17586.Grenningloh, G.; Pribilla, I.; Prior, P.; Multhaup, G.; Beyreuther,K.; Taleb, O.; Betz, H.: Cloning and expression of the 58 kd betasubunit of the inhibitory glycine receptor. *Neuron* 4: 963–970, 1990.
- [79257] 17587.Handford, C. A.; Lynch, J. W.; Baker, E.; Webb, G. C.; Ford, J.H.; Sutherland, G. R.; Schofield, P. R.: The human glycine receptorbeta subunit: primary structure, functional characterisation and chromosomallocalisation of the human and murine genes. *Molec. Brain Res.* 35:211–219, 1996.
- [79258] 17588.Milani, N.; Mulhardt, C.; Weber, R. G.; Lichter, P.; Kioschis,P.; Poustka, A.; Becker, C.–M.: The human glycine receptor beta subunitgene (GLRB): structure, refined chromosomal localization, and populationpolymorphism. *Genomics* 50: 341–345, 1998.
- [79259] 17589.Mulhardt, C.; Fischer, M.; Gass, P.; Simon–Chazottes, D.; Guenet,J.–L.; Kuhse, J.; Betz, H.; Becker, C.–M.: The spastic mouse: aberrantsplicing of glycine receptor beta subunit mRNA caused by intronicinsertion of L1 element. *Neuron* 13: 1003–1015, 1994.
- [79260] 17590.Browner, M. F.; Nakano, K.; Bang, A. G.; Fletterick, R. J.: Humanmuscle glycogen synthase cDNA sequence: a negatively charged proteinwith an asymmetric charge dis–

tribution. Proc. Nat. Acad. Sci. 86:1443–1447, 1989.

- [79261] 17591.Groop, L. C.; Kankuri, M.; Schalin-Jantti, C.; Ekstrand, A.; Nikula-Ijas, P.; Widen, E.; Kuusmanen, E.; Eriksson, J.; Franssila-Kallunki, A.; Saloranta, C.; Koskimies, S.: Association between polymorphism of the glycogen synthase gene and non-insulin-dependent diabetes mellitus. NewEng. J. Med. 328: 10–14, 1993.
- [79262] 17592.Huang, X.; Vaag, A.; Hansson, M.; Weng, J.; Laurila, E.; Groop, L.: Impaired insulin-stimulated expression of the glycogen synthase gene in skeletal muscle of type 2 diabetic patients is acquired rather than inherited. J. Clin. Endocr. Metab. 85: 1584–1590, 2000.
- [79263] 17593.Kadowaki, T.; Kadowaki, H.; Yazaki, Y.: Polymorphism of the glycogen synthase gene and non-insulin-dependent diabetes mellitus. (Letter) NewEng. J. Med. 328: 1569, 1993.
- [79264] 17594.Lehto, M.; Stoffel, M.; Groop, L.; Espinosa, R., III; Le Beau, M. M.; Bell, G. I.: Assignment of the gene encoding glycogen synthase (GYS) to human chromosome 19, band q13.3. Genomics 15: 460–461, 1993.
- [79265] 17595.Seldin, M. F.; Mott, D.; Bhat, D.; Petro, A.; Kuhn, C. M.; Kingsmore, S. F.; Bogardus, C.; Opara, E.; Feinglos, M. N.; Surwit, R. S.: Glycogen synthase: a putative locus for

diet-induced hyperglycemia. *J. Clin. Invest.* 94: 269–276, 1994.

[79266] 17596. Zouali, H.; Velho, G.; Froguel, P.: Polymorphism of the glycogen synthase gene and non-insulin-dependent diabetes mellitus. (Letter) *New Eng. J. Med.* 328: 1568, 1993.

[79267] 17597. David, J. R.: Delayed hypersensitivity in vitro: its mediation by cell-free substances formed by lymphoid cell-antigen interaction. *Proc. Nat. Acad. Sci.* 56: 72–77, 1966.

[79268] 17598. Reche, P. A.; Soumelis, V.; Gorman, D. M.; Clifford, T.; Liu, M.; Travis, M.; Zurawski, S. M.; Johnston, J.; Liu, Y.-J.; Spits, H.; de Waal Malefyt, R.; Kastelein, R. A.; Bazan, J. F.: Human thymic stromal lymphopoietin preferentially stimulates myeloid cells. *J. Immun.* 167: 336–343, 2001.

[79269] 17599. Kalmar, G. B.; Kay, R. J.; LaChance, A. C.; Cornell, R. B.: Primary structure and expression of a human CTP:phosphocholine cytidylyltransferase. *Biochim. Biophys. Acta* 1219: 328–334, 1994.

[79270] 17600. Rutherford, M. S.; Rock, C. O.; Jenkins, N. A.; Gilbert, D. J.; Tessner, T. G.; Copeland, N. G.; Jackowski, S.: The gene for murine CTP:phosphocholine cytidylyltransferase (Ctpct) is located on mouse chromosome 16. *Ge-*

nomics 18: 698–701, 1993.

[79271] 17601.Tabas, J. A.; Zasloff, M.; Wasmuth, J. J.; Emanuel, B. S.; Altherr, M. R.; McPherson, J. D.; Wozney, J. M.; Kaplan, F. S.: Bone morphogenetic protein: chromosomal localization of human genes for BMP1, BMP2A, and BMP3. *Genomics* 9: 283–289, 1991.

[79272] 17602.Gopal Rao, V. V. N.; Loffler, C.; Wozney, J. M.; Hansmann, I.: The gene for bone morphogenetic protein 2A (BMP2A) is localized to human chromosome 20p12 by radioactive and nonradioactive in situ hybridization. *Hum. Genet.* 90: 299–302, 1992.

[79273] 17603.Wang, E. A.; Rosen, V.; D'Alessandro, J. S.; Bauduy, M.; Cordes, P.; Harada, T.; Israel, D. I.; Hewick, R. M.; Kerns, K. M.; LaPan, P.; Luxenberg, D. P.; McQuaid, D.; Moutsatsos, I. K.; Nove, J.; Wozney, J. M.: Recombinant human bone morphogenetic protein induces bone formation. *Proc. Nat. Acad. Sci.* 87: 2220–2224, 1990.

[79274] 17604.White, P. M.; Morrison, S. J.; Orimoto, K.; Kubu, C. J.; Verdi, J. M.; Anderson, D. J.: Neural crest stem cells undergo cell-intrinsic developmental changes in sensitivity to instructive differentiation signals. *Neuron* 29: 57–71, 2001.

[79275] 17605.Wozney, J. M.; Rosen, V.; Celeste, A. J.; Mitsock, L.

M.; Whitters, M. J.; Kriz, R. W.; Hewick, R. M.; Wang, E. A.: Novel regulators of bone formation: molecular clones and activities. *Science* 242:1528–1534, 1988.

[79276] 17606. Connor, J. M.: Fibrodysplasia ossificans progressiva: lessons from rare maladies. (Editorial) *New Eng. J. Med.* 335: 591–593, 1996.

[79277] 17607. Dooley, C. A.; Attia, G. R.; Rainey, W. E.; Moore, D. R.; Carr, B. R.: Bone morphogenetic protein inhibits ovarian androgen production. *J. Clin. Endocr. Metab.* 85: 3331–3337, 2000.

[79278] 17608. McAlpine, P. J.: Personal Communication. Winnipeg, Manitoba, Canada 7/15/1992.

[79279] 17609. Monsoro-Burq, A.-H.; le Douarin, N. M.: BMP4 plays a key role in left–right patterning in chick embryos by maintaining Sonic Hedgehog asymmetry. *Molec. Cell* 7: 789–799, 2001.

[79280] 17610. Shafritz, A. B.; Shore, E. M.; Gannon, F. H.; Zasloff, M. A.; Taub, R.; Muenke, M.; Kaplan, F. S.: Overexpression of an osteogenic morphogen in fibrodysplasia ossificans progressiva. *New Eng. J. Med.* 335:555–561, 1996.

[79281] 17611. Tabas, J. A.; Hahn, G. V.; Cohen, R. B.; Seaunez, H. N.; Modi, W. S.; Wozney, J. M.; Zasloff, M.; Kaplan, F. S.: Chromosomal assignment of the human gene for bone

morphogenetic protein 4. Clin. Orthop.Rel. Res. 293: 310–316, 1993.

[79282] 17612.Tucker, A. S.; Matthews, K. L.; Sharpe, P. T.: Transformation of tooth type induced by inhibition of BMP signaling. Science 282:1136–1138, 1998.

[79283] 17613.van den Wijngaard, A.; Olde Weghuis, D.; Boersma, C. J. C.; van Zoelen, E. J. J.; Geurts van Kessel, A.; Olijve, W.: Fine mapping of the human bone morphogenetic protein-4 gene (BMP4) to chromosome 14q22–q23 by in situ hybridization. Genomics 27: 559–560, 1995.

[79284] 17614.van den Wijngaard, A.; van Kraay, M.; van Zoelen, E. J. J.; Olijve, W.; Boersma, C. J. C.: Genomic organization of the human bone morphogenetic protein-4 gene: molecular basis for multiple transcripts. Biochem.Biophys. Res. Commun. 219: 789–794, 1996.

[79285] 17615.Amano, S.; Scott, I. C.; Takahara, K.; Koch, M.; Champlaud, M.-F.; Gerecke, D. R.; Keene, D. R.; Hudson, D. L.; Nishiyama, T.; Lee, S.; Greenspan, D. S.; Burgeson, R. E.: Bone morphogenetic protein 1 is an extracellular processing enzyme of the laminin 5 gamma-2 chain. J.Biol. Chem. 275: 22728–22735, 2000.

[79286] 17616.Bond, J. S.; Beynon, R. J.: The astacin family of metalloendopeptidases. Protein Sci. 4: 1247–1261, 1995.

- [79287] 17617.Ceci, J. D.; Kingsley, D. M.; Silan, C. M.; Copeland, N. G.; Jenkins, N. A.: An interspecific backcross linkage map of the proximal half of mouse chromosome 14. *Genomics* 6: 673–678, 1990.
- [79288] 17618.Kessler, E.; Takahara, K.; Biniaminov, L.; Brusel, M.; Greenspan, D.: Bone morphogenic protein–1: the type I procollagen C–proteinase. *Science* 271:360–362, 1996.
- [79289] 17619.Donalies, M.; Cramer, M.; Ringwald, M.; Starzinski–Powitz, A.: Expression of M–cadherin, a member of the cadherin multigene family, correlates with differentiation of skeletal muscle cells. *Proc. Nat. Acad. Sci.* 88: 8024–8028, 1991.
- [79290] 17620.Kaupmann, K.; Becker–Follmann, J.; Scherer, G.; Jockusch, H.; Starzinski–Powitz, A.: The gene for the cell adhesion molecule M–cadherin maps to mouse chromosome 8 and human chromosome 16q24.1–qter and is near the E–cadherin (uvomorulin) locus in both species. *Genomics* 14: 488–490, 1992.
- [79291] 17621.Kremmidiotis, G.; Baker, E.; Crawford, J.; Eyre, H. J.; Nahmias, J.; Callen, D. F.: Localization of human cadherin genes to chromosomal regions exhibiting cancer–related loss of heterozygosity. *Genomics* 49:467–471, 1998.
- [79292] 17622.Garcia–Castro, M. I.; Vielmetter, E.; Bronner–Fraser,

M.: N-cadherin, a cell adhesion molecule involved in establishment of embryonic left-right asymmetry. *Science* 288: 1047–1051, 2000.

[79293] 17623. Hermiston, M. L.; Gordon, J. I.: Inflammatory bowel disease and adenomas in mice expressing a dominant negative N-cadherin. *Science* 270:1203–1206, 1995.

[79294] 17624. Miyatani, S.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Takeichi, M.: Genomic structure and chromosomal mapping of the mouse N-cadherin gene. *Proc. Nat. Acad. Sci.* 89: 8443–8447, 1992.

[79295] 17625. Takeichi, M.: Cadherins: a molecular family essential for selective cell–cell adhesion and animal morphogenesis. *Trends Genet.* 3: 213–217, 1987.

[79296] 17626. Tanaka, H.; Shan, W.; Phillips, G. R.; Arndt, K.; Bozdagi, O.; Shapiro, L.; Huntley, G. W.; Benson, D. L.; Coleman, D. R.: Molecular modification of N-cadherin in response to synaptic activity. *Neuron* 25:93–107, 2000.

[79297] 17627. Wallis, J.; Fox, M. F.; Walsh, F. S.: Structure of the human N-cadherin gene: YAC analysis and fine chromosomal mapping to 18q11.2. *Genomics* 22:172–179, 1994.

[79298] 17628. Walsh, F. S.; Barton, C. H.; Putt, W.; Moore, S. E.; Kelsell, D.; Spurr, N.; Goodfellow, P. N.: N-cadherin gene maps to human chromosome 18 and is not linked to the



E-cadherin gene. J. Neurochem. 55: 805–812,1990.

- [79299] 17629.Claverie, J.–M.; Hardelin, J.–P.; Legouis, R.; Levil-  
liers, J.;Bougueleret, L.; Mattei, M.–G.; Petit, C.: Character-  
ization and chromosomal assignment of a human cDNA  
encoding a protein related to the murine 102–kDa cad-  
herin-associated protein (alpha-catenin). Genomics  
15:13–20, 1993.
- [79300] 17630.Cook, S. A.; Bronson, R. T.; Donahue, L. R.; Ben-  
Arie, N.; Davisson, M. T.: Cerebellar deficient folia (cdf): a  
new mutation on mouse chromosome 6. Mammalian  
Genome 8: 108–112, 1997.
- [79301] 17631.Herrenknecht, K.; Ozawa, M.; Eckerskorn, C.;  
Lottspeich, F.; Lenter, M.; Kemler, R.: The uvomorulin-an-  
chorage protein alpha-catenin is a vinculin homologue.  
Proc. Nat. Acad. Sci. 88: 9156–9160, 1991.
- [79302] 17632.Park, C.; Falls, W.; Finger, J. H.; Longo-Guess, C.  
M.; Ackerman, S. L.: Deletion in Catna2, encoding alpha-  
N-catenin, causes cerebellar and hippocampal lamination  
defects and impaired startle modulation. Nature Genet. 31:  
279–284, 2002.
- [79303] 17633.Magnuson, V. L.; McCombs, J. L.; Lee, C.–C.; Yang,  
F.; Bowman, B. H.; McGill, J. R.: Human alpha-  
2-HS-glycoprotein localized to 3q27–q29 by in situ hy-

bridization. Cytogenet. Cell Genet. 47: 72–74,1988.

- [79304] 17634.Seroussi, E.; Kedra, D.; Kost-Alimova, M.; Sandberg-Nordqvist,A.-C.; Fransson, I.; Jacobs, J. F. M.; Fu, Y.; Pan, H.-Q.; Roe, B.A.; Imreh, S.; Dumanski, J. P.: TOM1 genes map to human chromosome22q13.1 and mouse chromosome 8C1 and encode proteins similar to theendosomal proteins HGS and STAM. Genomics 57: 380–388, 1999.
- [79305] 17635.Hanson, I.; Churchill, A.; Love, J.; Axton, R.; Moore, T.; Clarke,M.; Meire, F.; van Heyningen, V.: Missense mutations in the mostancient residues of the PAX6 paired domain underlie a spectrum ofhuman congenital eye malformations. Hum. Molec. Genet. 8: 165–172,1999.
- [79306] 17636.O'Farrell, P. H.: High resolution two-dimensional electrophoresisof proteins. J. Biol. Chem. 250: 4007–4021, 1975.
- [79307] 17637.Osawa, M.; Umetsu, K.; Ohki, T.; Nagasawa, T.; Suzuki, T.; Takeichi,S.: Molecular evidence for human alpha2-HS glycoprotein (AHSG) polymorphism. Hum.Genet. 99: 18–21, 1997.
- [79308] 17638.Pedersen, K. O.: Fetuin, a new globulin isolated from serum.(Letter) Nature 154: 575, 1944.
- [79309] 17639.Rizzu, P.; Baldini, A.: Three members of the human

cystatin genesuperfamily, AHSG, HRG, and KNG, map within one megabase of genomicDNA at 3q27. Cytogenet. Cell Genet. 70: 26–28, 1995.

- [79310] 17640.Terkelsen, O. B. F.; Jahnen–Dechent, W.; Nielsen, H.; Moos, T.;Fink, E.; Nawratil, P.; Muller–Esterl, W.; Mollgard, K.: Rat fetuin:distribution of protein and mRNA in embryonic and neonatal rat tissues. Anat.Embryol. 197: 125–133, 1998.
- [79311] 17641.Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha–2–HS–glycoproteintyping by isoelectric focusing. Hum. Genet. 67: 70–71, 1984.
- [79312] 17642.Umetsu, K.; Kashimura, S.; Ikeda, N.; Suzuki, T.: A new alpha–2–HS–glycoproteinallele (AHS\*5\*) in two Japanese families. Hum. Genet. 68: 264–265,1984.
- [79313] 17643.Umetsu, K.; Yuasa, I.; Ikuta, M.; Suzuki, T.: Alpha–2–HS–glycoprotein(A2HS) polymorphism in a Japanese population: existence of two newvariants. Jpn. J. Hum. Genet. 32: 263–266, 1987.
- [79314] 17644.Umetsu, K.; Yuasa, I.; Suzuki, T.: The polymorphism of desialyzedalpha–2–HS–glycoprotein (AHS): isoelectric focusing in 2.5M urea asa method for identification of genetic variants. Hum. Genet. 73:372–373, 1986.
- [79315] 17645.Yoshioka, Y.; Gejyo, F.; Marti, T.; Rickli, E. E.; Burgi,

W.;Offner, G. D.; Troxler, R. F.; Schmid, K.: The complete amino acidsequence of the A-chain of human plasma al-  
pha-2HS-glycoprotein. J.Biol. Chem. 261: 1665-1676,  
1986.

[79316] 17646.Zelinski, T.; Kaita, H.; Lewis, M.; Coghlan, G.; Craig,  
D.: Thesequence of chromosome 3 loci AHSG:TF:CHE1.  
Hum. Hered. 37: 1-6,1987.

[79317] 17647.Elpeleg, O. N.; Shaag, A.; Glustein, J. Z.; Anikster,  
Y.; Joseph,A.; Saada, A.: Lipoamide dehydrogenase defi-  
ciency in Ashkenazi Jews:an insertion mutation in the mi-  
tochondrial leader sequence. Hum.Mutat. 10: 256-257,  
1997.

[79318] 17648.Feigenbaum, A. S.; Robinson, B. H.: The structure  
of the humandihydrolipoamide dehydrogenase gene (DLD)  
and its upstream elements. Genomics 17:376-381, 1993.

[79319] 17649.Hong, Y. S.; Kerr, D. S.; Craigen, W. J.; Tan, J.; Pan,  
Y.; Lusk,M.; Patel, M. S.: Identification of two mutations in  
a compound heterozygouschild with dihydrolipoamide  
dehydrogenase deficiency. Hum. Molec.Genet. 5:  
1925-1930, 1996.

[79320] 17650.Johnson, M.; Yang, H.-S.; Johanning, G. L.; Patel, M.  
S.: Characterizationof the mouse dihydrolipoamide dehy-  
drogenase (Dld) gene: genomic structure,promoter se-

quence, and chromosomal localization. *Genomics* 41: 320–326, 1997.

- [79321] 17651. Liu, T.-C.; Kim, H.; Arizmendi, C.; Kitano, A.; Patel, M. S.: Identification of two missense mutations in a dihydrolipoamide dehydrogenase-deficient patient. *Proc. Nat. Acad. Sci.* 90: 5186–5190, 1993.
- [79322] 17652. Matalon, R.; Michals, K.; Stumpf, D.; Goodman, S.; Parks, J.: Lactic acidosis due to lipoamide dehydrogenase (LAD) deficiency: improvement after oral lipoic acid. (Abstract) *Am. J. Hum. Genet.* 33: 48A, 1981.
- [79323] 17653. Munnich, A.; Saudubray, J. M.; Taylor, J.; Charpentier, C.; Marsac, C.; Rocchiccioli, F.; Amedee-Manesme, O.; Coude, F. X.; Frezal, J.; Robinson, B. H.: Congenital lactic acidosis, alpha-ketoglutaric aciduria and variant form of maple syrup urine disease due to a single enzyme defect: dihydrolipoyl dehydrogenase deficiency. *Acta Paediat. Scand.* 71: 167–171, 1982.
- [79324] 17654. Olson, S.; Song, B. J.; Huh, T.-L.; Chi, Y.-T.; Veech, R. L.; McBride, O. W.: Three genes for enzymes of the pyruvate dehydrogenase complex map to human chromosomes 3, 7, and X. *Am. J. Hum. Genet.* 46: 340–349, 1990.
- [79325] 17655. Otulakowski, G.; Robinson, B. H.: Isolation and sequence determination of cDNA clones for porcine and hu-

man lipoamide dehydrogenase. J.Biol. Chem. 262: 17313–17318, 1987.

- [79326] 17656.Otulakowski, G.; Robinson, B. H.; Willard, H. F.: Gene for lipoamidedehydrogenase maps to human chromosome 7. Somat. Cell Molec. Genet. 14:411–414, 1988.
- [79327] 17657.Pons, G.; Raefsky–Estrin, C.; Carothers, D. J.; Pepin, R. A.;Javed, A. A.; Jesse, B. W.; Ganapathi, M. K.; Samols, D.; Patel, M.S.: Cloning and cDNA sequence of the dihydrolipoamide dehydrogenasecomponent of human alpha-ketoacid dehydrogenase complexes. Proc.Nat. Acad. Sci. 85: 1422–1426, 1988.
- [79328] 17658.Robinson, B. H.; Taylor, J.; Kahler, S. G.; Kirkman, H. N.: Lacticacidemia, neurologic deterioration and carbohydrate dependence ina girl with dihydrolipoyl dehydrogenase deficiency. Europ. J. Pediat. 136:35–39, 1981.
- [79329] 17659.Robinson, B. H.; Taylor, J.; Sherwood, W. G.: Deficiency of dihydrolipoyldehydrogenase (a component of the pyruvate and alpha-ketoglutaratedehydrogenase complexes): a cause of congenital lactic acidosis ininfancy. Pediat. Res. 11: 1198–1202, 1977.
- [79330] 17660.Sakaguchi, Y.; Yoshino, M.; Aramaki, S.; Yoshida, I.; Yamashita,F.; Kuhara, T.; Matsumoto, I.; Hayashi, T.: Dihydrolipoyl dehydrogenasedeficiency: a therapeutic trial with

branched-chain amino acid restriction. *Europ.J. Pediat.* 145: 271–274, 1986.

- [79331] 17661.Scherer, S. W.; Otulakowski, G.; Robinson, B. H.; Tsui, L.–C.: Localization of the human dihydrolipoamide dehydrogenase gene (DLD)to 7q31–q32. *Cytogenet. Cell Genet.* 56: 176–177, 1991.
- [79332] 17662.Shaag, A.; Saada, A.; Berger, I.; Mandel, H.; Joseph, A.; Feigenbaum,A.; Elpeleg, O. N.: Molecular basis of lipoamide dehydrogenase deficiencyin Ashkenazi Jews. *Am. J. Med. Genet.* 82: 177–182, 1999.
- [79333] 17663.Martin–Subero, J. I.; Gesk, S.; Harder, L.; Sonoki, T.; Tucker,P. W.; Schlegelberger, B.; Grote, W.; Novo, F. J.; Calasanz, M. J.;Hansmann, M. L.; Dyer, M. J. S.; Siebert, R.: Recurrent involvementof the REL and BCL11A loci in classical Hodgkin lymphoma. *Blood* 99:1474–1477, 2002.
- [79334] 17664.Hankinson, S. E.; Willett, W. C.; Colditz, G. A.; Hunter, D. J.;Michaud, D. S.; Deroo, B.; Rosner, B.; Speizer, F. E.; Pollak, M.: Circulating concentrations of insulin–like growth factor–I and riskof breast cancer. *Lancet* 351: 1393–1396, 1998.
- [79335] 17665.Jones, C.; Slijepcevic, P.; Marsh, S.; Baker, E.; Langdon, W. Y.;Richards, R. I.; Tunnacliffe, A.: Physical linkage of the fragilesite FRA11B and a Jacobsen syndrome chro–

mosome deletion breakpoint in 11q23.3. Hum. Molec. Genet. 3: 2123–2130, 1994.

[79336] 17666. Esumi, N.; Budarf, M.; Ciccarelli, L.; Sellinger, B.; Kozak, C.A.; Wistow, G.: Conserved gene structure and genomic linkage for D–dopachrome tautomerase (DDT) and MIF. Mammalian Genome 9: 753–757, 1998.

[79337] 17667. Kleemann, R.; Hausser, A.; Geiger, G.; Mischke, R.; Burger–Kentischer, A.; Flieger, O.; Johannes, F.–J.; Roger, T.; Calandra, T.; Kapurniotu, A.; Grell, M.; Finkelmeier, D.; Brunner, H.; Bernhagen, J.: Intracellular action of the cytokine MIF to modulate AP–1 activity and the cell cycle through Jab1. Nature 408: 211–216, 2000.

[79338] 17668. Kozak, C. A.; Adamson, M. C.; Buckler, C. E.; Segovia, L.; Paralkar, V.; Wistow, G.: Genomic cloning of mouse MIF (macrophage inhibitory factor) and genetic mapping of the human and mouse expressed gene and nine mouse pseudogenes. Genomics 27: 405–411, 1995.

[79339] 17669. Paralkar, V.; Wistow, G.: Cloning the human gene for macrophage migration inhibitory factor (MIF). Genomics 19: 48–51, 1994.

[79340] 17670. Pastrana, D. V.; Raghavan, N.; Fitzgerald, P.; Eisinger, S. W.; Metz, C.; Bucala, R.; Schleimer, R. P.; Bickel, C.; Scott, A. L.: Filarial nematode parasites secrete a ho–



mologue of the human cytokinemacrophage migration inhibitory factor. *Infect. Immun.* 66: 5955–5963,1998.

- [79341] 17671.Roger, T.; David, J.; Glauser, M. P.; Calandra, T.: MIF regulates innate immune response through modulation of Toll-like receptor 4. *Nature* 414:920–924, 2001.
- [79342] 17672.Weiser, W. Y.; Temple, P. A.; Witek-Giannotti, J. S.; Remold, H. G.; Clark, S. C.; David, J. R.: Molecular cloning of a cDNA encoding a human macrophage migration inhibitory factor. *Proc. Nat. Acad.Sci.* 86: 7522–7526, 1989.
- [79343] 17673.Emi, M.; Asaoka, H.; Matsumoto, A.; Itakura, H.; Kurihara, Y.; Wada, Y.; Kanamori, H.; Yazaki, Y.; Takahashi, E.; Lepert, M.; Lalouel, J.-M.; Kodama, T.; Mukai, T.: Structure, organization, and chromosomal mapping of the human macrophage scavenger receptor gene. *J. Biol.Chem.* 268: 2120–2125, 1993.
- [79344] 17674.Latil, A.; Lidereau, R.: Genetic aspects of prostate cancer. *VirchowsArch.* 432: 389–406, 1998.
- [79345] 17675.Matsumoto, A.; Naito, M.; Itakura, H.; Ikemoto, S.; Asaoka, H.; Hayakawa, I.; Kanamori, H.; Aburatani, H.; Takaku, F.; Suzuki, H.; Kobari, Y.; Miyai, T.; Takahashi, K.; Cohen, E. H.; Wydro, R.; Housman, D. E.; Kodama, T.: Human macrophage scavenger receptors: primary structure, expression, and localization in atherosclerotic lesions.

Proc.Nat. Acad. Sci. 87: 9133–9137, 1990.

- [79346] 17676.Xu, J.; Zheng, S. L.; Hawkins, G. A.; Faith, D. A.; Kelly, B.; Isaacs, S. D.; Wiley, K. E.; Chang, B.; Ewing, C. M.; Bujnovszky, P.; Carpten, J. D.; Bleecker, E. R.; Walsh, P. C.; Trent, J. M.; Meyers, D. A.; Isaacs, W. B.: Linkage and association studies of prostate cancer susceptibility: evidence for linkage at 8p22–23. *Am. J. Hum. Genet.* 69: 341–350, 2001.
- [79347] 17677.Xu, J.; Zheng, S. L.; Komiya, A.; Mychaleckyj, J. C.; Isaacs, S. D.; Hu, J. J.; Sterling, D.; Lange, E. M.; Hawkins, G. A.; Turner, A.; Ewing, C. M.; Faith, D. A.; and 19 others: Germline mutations and sequence variants of the macrophage scavenger receptor 1 gene are associated with prostate cancer risk. *Nature Genet.* 32: 321–325, 2002.
- [79348] 17678.Holness, C. L.; Simmons, D. L.: Molecular cloning of CD68, a human macrophage marker related to lysosomal glycoproteins. *Blood* 81:1607–1613, 1993.
- [79349] 17679.Jones, E.; Quinn, C. M.; See, C. G.; Montgomery, D. S.; Ford, M. J.; Kolbe, K.; Gordon, S.; Greaves, D. R.: The linked human elongation initiation factor 4A1 (EIF4A1) and CD68 genes map to chromosome 17p13. *Genomics* 53:248–250, 1998.
- [79350] 17680.Chavrier, P.; Janssen–Timmen, U.; Mattei, M.–G.;

Zerial, M.; Bravo,R.; Charnay, P.: Structure, chromosome location, and expression of the mouse zinc finger gene Krox-20: multiple gene products and coregulation with the proto-oncogene c-fos. *Molec. Cell. Biol.* 9: 787-797, 1989.

[79351] 17681. Le Beau, M. M.; Espinosa, R., III; Neuman, W. L.; Stock, W.; Roulston, D.; Larson, R. A.; Keinanen, M.; Westbrook, C. A.: Cytogenetic and molecular delineation of the smallest commonly deleted region of chromosome 5 in malignant myeloid diseases. *Proc. Nat. Acad. Sci.* 90: 5484-5488, 1993.

[79352] 17682. Barrow, L. L.; Simin, K.; Jones, J. M.; Lee, D. C.; Meisler, M. H.: Conserved linkage of early growth response 4, annexin 4, and transforming growth factor alpha on mouse chromosome 6. *Genomics* 19:388-390, 1994.

[79353] 17683. Crosby, S. D.; Veile, R. A.; Donis-Keller, H.; Baraban, J. M.; Bhat, R. V.; Simburger, K. S.; Milbrandt, J.: Neural-specific expression, genomic structure, and chromosomal localization of the gene encoding the zinc-finger transcription factor NGFI-C. *Proc. Nat. Acad. Sci.* 89:4739-4743, 1992.

[79354] 17684. Boerkoel, C. F.; Takashima, H.; Bacino, C. A.; Daentl, D.; Lupski, J. R.: EGR2 mutation R359W causes a

spectrum of Dejerine–Sottas neuropathy. *Neurogenetics* 3:153–157, 2001.

- [79355] 17685. Willcutt, E. G.; Pennington, B. F.; Smith, S. D.; Cardon, L. R.; Gayán, J.; Knopik, V. S.; Olson, R. K.; DeFries, J. C.: Quantitative trait locus for reading disability on chromosome 6p is pleiotropic for attention-deficit/hyperactivity disorder. *Am. J. Med. Genet.* 114:260–268, 2002.
- [79356] 17686. Inazu, A.; Brown, M. L.; Hesler, C. B.; Agellon, L. B.; Koizumi, J.; Takata, K.; Maruhama, Y.; Mabuchi, H.; Tall, A. R.: Increased high-density lipoprotein levels caused by a common cholesteryl-ester transfer protein gene mutation. *New Eng. J. Med.* 323: 1234–1238, 1990.
- [79357] 17687. Saito, F.: A pedigree of homozygous familial hyperalphalipoproteinemia. *Metabolism* 33:629–633, 1984.
- [79358] 17688. Andres, D. A.; Milatovich, A.; Ozcelik, T.; Wenzlau, J. M.; Brown, M. S.; Goldstein, J. L.; Francke, U.: cDNA cloning of the two subunits of human CAAX farnesyl transferase and chromosomal mapping of FNTA and FNTB loci and related sequences. *Genomics* 18: 105–112, 1993.
- [79359] 17689. Epstein, W. W.; Lever, D.; Leining, L. M.; Bruenger, E.; Rilling, H. C.: Quantitation of prenylcysteines by a selective cleavage reaction. *Proc. Nat. Acad. Sci.* 88:

9668–9670, 1991.

- [79360] 17690.Long, S. B.; Casey, P. J.; Beese, L. S.: Reaction path of proteinfarnesyltransferase at atomic resolution. *Nature* 419: 645–650, 2002.
- [79361] 17691.Porter, J. C.; Messer, A.: Genetic mapping of farne-syltransferasealpha(Fnta) to mouse chromosome 8. *Mam-malian Genome* 7: 622–623,1996.
- [79362] 17692.Schafer, W. R.; Rine, J.: Protein prenylation: genes, enzymes,targets, and functions. *Annu. Rev. Genet.* 26: 209–237, 1992.
- [79363] 17693.Wang, T.; Danielson, P. D.; Li, B.; Shah, P. C.; Kim, S. D.; Donahoe,P. K.: The p21ras farnesyltransferase alpha subunit in TGF-beta andactivin signaling. *Science* 271: 1120–1122, 1996.
- [79364] 17694.Zhang, F. L.; Diehl, R. E.; Kohl, N. E.; Gibbs, J. B.; Giros, B.;Casey, P. J.; Omer, C. A.: cDNA cloning and ex-pression of rat andhuman protein geranylgeranyltrans-ferase type-I. *J. Biol. Chem.* 269:3175–3180, 1994.
- [79365] 17695.Roifman, C. M.; Zhang, J.; Chitayat, D.; Sharfe, N.: A partialdeficiency of interleukin-7R-alpha is sufficient to abrogate T-celldevelopment and cause severe combined immunodeficiency. *Blood* 96:2803–2807, 2000.
- [79366] 17696.Abraham, R.; Myers, A.; Wavrant-DeVrieze, F.;

Hamshire, M. L.; Thomas, H. V.; Marshall, H.; Compton, D.; Spurlock, G.; Turic, D.; Hoogendoorn, B.; Kwon, J. M.; Petersen, R. C.; and 12 others: Substantial linkage disequilibrium across the insulin-degrading enzyme locus but no association with late-onset Alzheimer's disease. *Hum. Genet.* 109:646–652, 2001.

[79367] 17697. Affholter, J. A.; Fried, V. A.; Roth, R. A.: Human insulin-degrading enzyme shares structural and functional homologies with *E. coli* protease III. *Science* 242: 1415–1418, 1988.

[79368] 17698. Affholter, J. A.; Hsieh, C.-L.; Francke, U.; Roth, R. A.: Insulin-degrading enzyme: stable expression of the human complementary DNA, characterization of its protein product, and chromosomal mapping of the human and mouse genes. *Molec. Endocr.* 4: 1125–1135, 1990.

[79369] 17699. Edbauer, D.; Willem, M.; Lammich, S.; Steiner, H.; Haass, C.: Insulin-degrading enzyme rapidly removes the beta-amyloid precursor protein intracellular domain (AICD). *J. Biol. Chem.* 277: 13389–13393, 2002.

[79370] 17700. Espinosa, R., III; Lemons, R. S.; Perlman, R. K.; Kuo, W.-L.; Rosner, M. R.; Le Beau, M. M.: Localization of the gene encoding insulin-degrading enzyme to human chromosome 10, bands q23–q25. *Cytogenet. Cell Genet.*

57:184–186, 1991.

- [79371] 17701.Qiu, W. Q.; Walsh, D. M.; Ye, Z.; Vekrellis, K.; Zhang, J.; Podlisny, M. B.; Rosner, M. R.; Safavi, A.; Hersh, L. B.; Selkoe, D. J.: Insulin-degrading enzyme regulates extracellular levels of amyloid beta-protein by degradation. *J. Biol. Chem.* 273: 32730–32738, 1998.
- [79372] 17702.Qiu, W. Q.; Ye, Z.; Kholodenko, D.; Seubert, P.; Selkoe, D. J.: Degradation of amyloid beta-protein by a metalloprotease secreted by microglia and other neural and non-neural cells. *J. Biol. Chem.* 272:6641–6646, 1997.
- [79373] 17703.Glasgow, B. J.; Heinzmann, C.; Kojis, T.; Sparkes, R. S.; Mohandas, T.; Bateman, J. B.: Assignment of tear lipocalin gene to human chromosome 9q34–9qter. *Curr. Eye Res.* 12: 1019–1023, 1993.
- [79374] 17704.Herzog, H.; Baumgartner, M.; Holzfeind, P.; Redl, B.: Confirmation of 9q34 as the chromosomal site of the human lipocalin LCN1 gene. *Cytogenet. Cell Genet.* 69: 108–109, 1995.
- [79375] 17705.Holzfeind, P.; Merschak, P.; Rogatsch, H.; Culig, Z.; Feichtinger, H.; Klocker, H.; Redl, B.: Expression of the gene for tear lipocalin/von Ebner's gland protein in human prostate. *FEBS Lett.* 395: 95–98, 1996.

- [79376] 17706.Lacazette, E.; Pitiot, G.; Jobert, S.; Mallet, J.; Gachon, A. M.F.: Fine genetic mapping of LCN1/D9S1826 within 9q34. *Ann. Hum.Genet.* 61: 449–455, 1997.
- [79377] 17707.Lassagne, H.; Gachon, A. M. F.: Cloning of a human lacrimal lipocalinsecreted in tears. *Exp. Eye Res.* 56: 605–609, 1993.
- [79378] 17708.Lassagne, H.; Nguyen, V. C.; Mattei, M. G.; Gachon, A. M. F.:Assignment of LCN1 to human chromosome 9 is confirmed. *Cytogent.Cell Genet.* 71: 104, 1995.
- [79379] 17709.Lassagne, H.; Ressot, C.; Mattei, M. G.; Gachon, A. M. F.: Assignmentof the human tear lipocalin gene (LCN1) to 9q34 by in situ hybridization. *Genomics* 18:160–161, 1993.
- [79380] 17710.Nguyen Van Cong; Weil, D.; Finaz, C.; Cohen–Haguenauer, O.; Gross,M.–S.; Jegou–Foubert, C.; de Tand, M.–F.; Cochet, C.; de Grouchy,J.; Frezal, J.: Panel of twenty–five independent man–rodent hybridsfor human genetic marker mapping. *Ann. Genet.* 29: 20–26, 1986.
- [79381] 17711.Pervaiz, S.; Brew, K.: Homology and structure–function correlationsbetween alpha–1–acid glycoprotein and serum retinol–binding proteinand its relatives. *FASEB J.* 1: 209–214, 1987.
- [79382] 17712.Redl, B.; Holzfeind, P.; Lottspeich, F.: cDNA cloning



and sequencing reveals human tear prealbumin to be a member of the lipophilic–ligand carrier protein superfamily. *J. Biol. Chem.* 28: 20282–20287, 1992.

- [79383] 17713. Redl, B.; Wojnar, P.; Ellemunter, H.; Feichtinger, H.: Identification of a lipocalin in mucosal glands of the human tracheobronchial tree and its enhanced secretion in cystic fibrosis. *Lab. Invest.* 78:1121–1129, 1998.
- [79384] 17714. van't Hof, W.; Blankenvoorde, M. F. J.; Veerman, E. C. I.; Nieuw Amerongen, A. V.: The salivary lipocalin Von Ebner's Gland protein is a cysteine proteinase inhibitor. *J. Biol. Chem.* 272: 1837–1841, 1997.
- [79385] 17715. Huebner, K.; Cannizzaro, L. A.; Croce, C. M.; Frey, A. Z.; Wallner, B. P.; Hecht, B. K.; Hecht, F.: Chromosome localization of the human genes for lipocortin I and the lipocortin II family. (Abstract) *Cytogenet. Cell Genet.* 46: 631 only, 1987.
- [79386] 17716. Huebner, K.; Cannizzaro, L. A.; Frey, A. Z.; Hecht, B. K.; Hecht, F.; Croce, C. M.; Wallner, B. P.: Chromosomal localization of the human genes for lipocortin I and lipocortin II. *Oncogene Res.* 2:299–310, 1988.
- [79387] 17717. Spano, F.; Raugei, G.; Palla, E.; Colella, C.; Melli, M.: Characterization of the human lipocortin–2–encoding multigene family: its structure suggests the existence of a

short amino acid unit undergoing duplication. Gene 95:243–251, 1990.

- [79388] 17718.Takahashi, S.; Reddy, S. V.; Chirgwin, J. M.; Devlin, R.; Haipek,C.; Anderson, J.; Roodman, G. D.: Cloning and identification of annexinII as an autocrine/paracrine factor that increases osteoclast formationand bone resorption. J. Biol. Chem. 269: 28696–28701, 1994.
- [79389] 17719.Harman, S. M.; Metter, E. J.; Blackman, M. R.; Landis, P. K.;Carter, H. B.: Serum levels of insulin–like growth factor I (IGF–I),IGF–II, IGF–binding protein–3, and prostate–specific antigen as predictorsof clinical prostate cancer. J. Clin. Endocr. Metab. 85: 4258–4265,2000.
- [79390] 17720.Holly, J.: Insulin–like growth factor–I and new opportunitiesfor cancer prevention. Lancet 351: 1373–1375, 1998.
- [79391] 17721.Hoppener, J. W. M.; de Pagter–Holthuizen, P.; Geurts van Kessel,A. H. M.; Jansen, M.; Kittur, S. D.; Antonarakis, S. E.; Lips, C.J. M.; Sussenbach, J. S.: The human gene encoding insulin–like growthfactor I is located on chromosome 12. Hum. Genet. 69: 157–160, 1985.
- [79392] 17722.Le Bouc, Y.; Dreyer, D.; Jaeger, F.; Binoux, M.; Sondermeyer,P.: Complete characterization of the human IGF–I nucleotide sequenceisolated from a newly constructed

adult liver cDNA library. FEBSLett. 196: 108–112, 1986.

[79393] 17723. Hellstrom, A.; Perruzzi, C.; Ju, M.; Engstrom, E.; Hard, A.-L.; Liu, J.-L.; Albertsson-Wikland, K.; Carlsson, B.; Niklasson, A.; Sjodell, L.; LeRoith, D.; Senger, D. R.; Smith, L. E. H.: Low IGF-I suppresses VEGF-survival signaling in retinal endothelial cells: direct correlation with clinical retinopathy of prematurity. Proc. Nat. Acad. Sci. 98:5804–5808, 2001.

[79394] 17724. Lembo, G.; Rockman, H. A.; Hunter, J. J.; Steinmetz, H.; Koch, W. J.; Ma, L.; Printz, M. P.; Ross, J., Jr.; Chien, K. R.; Powell-Braxton, L.: Elevated blood pressure and enhanced myocardial contractility in mice with severe IGF-1 deficiency. J. Clin. Invest. 98: 2648–2655, 1996.

[79395] 17725. Li, C. H.; Yamashiro, D.; Gospodarowicz, D.; Kaplan, S. L.; VanVliet, G.: Total synthesis of insulin-like growth factor I (somatomedin C). Proc. Nat. Acad. Sci. 80: 2216–2220, 1983.

[79396] 17726. Liu, J.-P.; Baker, J.; Perkins, A. S.; Robertson, E. J.; Efstratiadis, A.: Mice carrying null mutations of the genes encoding insulin-like growth factor I (Igf-1) and type 1 IGF receptor (Igf1r). Cell 75:59–72, 1993.

[79397] 17727. Mathews, L. S.; Norstedt, G.; Palmiter, R. D.: Regulation of insulin-like growth factor I gene expression by

growth hormone. Proc.Nat. Acad. Sci. 83: 9343–9347, 1986.

[79398] 17728.Morton, C.; Rall, L.; Bell, G.; Shows, T.: Human insulin-like growth factor-1 (IGF1) is encoded at 12q22–q24.1, and insulin-like growth factor-2 (IGF2) is at 11p15. (Abstract) Cytogenet. Cell Genet. 40:703 only, 1985.

[79399] 17729.Musaro, A.; McCullagh, K.; Paul, A.; Houghton, L.; Dobrowolny, G.; Molinaro, M.; Barton, E. R.; Sweeney, H. L.; Rosenthal, N.: Localized IGF-1 transgene expression sustains hypertrophy and regeneration in senescent skeletal muscle. Nature Genet. 27: 195–200, 2001.

[79400] 17730.Musaro, A.; McCullagh, K. J. A.; Naya, F. J.; Olson, E. N.; Rosenthal, N.: IGF-1 induces skeletal myocyte hypertrophy through calcineurin association with GATA-2 and NF-ATc1. Nature 400: 581–585, 1999.

[79401] 17731.Playford, M. P.; Bicknell, D.; Bodmer, W. F.; Macaulay, V. M.: Insulin-like growth factor 1 regulates the location, stability, and transcriptional activity of beta-catenin. Proc. Nat. Acad. Sci. 97:12103–12108, 2000.

[79402] 17732.Powell-Braxton, L.; Hollingshead, P.; Warburton, C.; Dowd, M.; Pitts-Meek, S.; Dalton, D.; Gillett, N.; Stewart, T. A.: IGF-I is required for normal embryonic growth in mice.

Genes Dev. 7: 2609–2617,1993.

- [79403] 17733.Rapp, R.; Deger, A.; Blum, W.; Koch, R.; Weber, U.: Characterization of the protein which binds insulin-like growth factor in human serum. *Europ.J. Biochem.* 172: 421–425, 1988.
- [79404] 17734.Rotwein, P.: Two insulin-like growth factor I messenger RNAs are expressed in human liver. *Proc. Nat. Acad. Sci.* 83: 77–81, 1986.
- [79405] 17735.Schoenle, E. J.; Zenobi, P. D.; Torresani, T.; Werder, E. A.; Zachmann, M.; Froesch, E. R.: Recombinant human insulin-like growth factor I (rhIGF I) reduces hyperglycaemia in patients with extreme insulin resistance. *Diabetologia* 34: 675–679, 1991.
- [79406] 17736.Semsarian, C.; Wu, M.-J.; Ju, Y.-K.; Marciniak, T.; Yeoh, T.; Allen, D. G.; Harvey, R. P.; Graham, R. M.: Skeletal muscle hypertrophy is mediated by a  $\text{Ca}^{2+}$ -dependent calcineurin signalling pathway. *Nature* 400:576–581, 1999.
- [79407] 17737.Svoboda, M. E.; Van Wyk, J. J.; Klapper, D. G.; Fellows, R. E.; Grissom, F. E.; Schleuter, R. J.: Purification of somatomedin-C from human plasma: chemical and biological properties, partial sequence analysis, and relationship to other somatomedins. *Biochemistry* 19:790–797,

1980.

- [79408] 17738. Taylor, B. A.; Grieco, D.: Localization of the gene encoding insulin-like growth factor I on mouse chromosome 10. *Cytogenet. Cell Genet.* 56: 57–58, 1991.
- [79409] 17739. Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T. B.: Insulin-like growth factor genes: chromosome organization and association with disease. (Abstract) *Am. J. Hum. Genet.* 36: 121S only, 1984.
- [79410] 17740. Tricoli, J. V.; Rall, L. B.; Scott, J.; Bell, G. I.; Shows, T. B.: Localization of insulin-like growth factor genes to human chromosomes 11 and 12. *Nature* 310: 784–786, 1984.
- [79411] 17741. Ullrich, A.; Berman, C. H.; Dull, T. J.; Gray, A.; Lee, J. M.: Isolation of the human insulin-like growth factor I gene using a single synthetic DNA probe. *EMBO J.* 3: 361–364, 1984.
- [79412] 17742. Gough, N. M.; Rakar, S.; Harpur, A.; Wilks, A. F.: Localization of genes for two members of the JAK family of protein tyrosine kinases to murine chromosomes 4 and 19. *Mammalian Genome* 6: 247–248, 1995.
- [79413] 17743. Ihle, J. N.: Cytokine receptor signalling. *Nature* 377: 591–594, 1995.
- [79414] 17744. Pritchard, M. A.; Baker, E.; Callen, D. F.; Sutherland,

G. R.;Wilks, A. F.: Two members of the JAK family of protein tyrosine kinases map to chromosomes 1p31.3 and 9p24. *Mammalian Genome* 3: 36–38, 1992.

[79415] 17745.Campbell, G. S.; Argetsinger, L. S.; Ihle, J. N.; Kelly, P. A.;Rillema, J. A.; Carter–Su, C.: Activation of JAK2 tyrosine kinase by prolactin receptors in Nb2 cells and mouse mammary gland explants. *Proc.Nat. Acad. Sci.* 91: 5232–5236, 1994.

[79416] 17746.Huang, L. J.; Constantinescu, S. N.; Lodish, H. F.: The N-terminal domain of Janus kinase 2 is required for Golgi processing and cell surface expression of erythropoietin receptor. *Molec. Cell* 8: 1327–1338, 2001.

[79417] 17747.Neubauer, H.; Cumano, A.; Muller, M.; Wu, H.; Huffstadt, U.; Pfeffer, K.: Jak2 deficiency defines an essential developmental checkpoint in definitive hematopoiesis. *Cell* 93: 397–409, 1998.

[79418] 17748.Parganas, E.; Wang, D.; Stravopodis, D.; Topham, D. J.; Marine, J.–C.; Teglund, S.; Vanin, E. F.; Bodner, S.; Colamonici, O. R.; vanDeursen, J. M.; Grosveld, G.; Ihle, J. N.: Jak2 is essential for signaling through a variety of cytokine receptors. *Cell* 93: 385–395, 1998.

[79419] 17749.Saltzman, A.; Stone, M.; Franks, C.; Searfoss, G.; Munro, R.; Jaye, M.; Ivashchenko, Y.: Cloning and charac–

terization of human Jak-2kinase: high mRNA expression in immune cells and muscle tissue. *Biochem.Biophys. Res. Commun.* 246: 627-633, 1998.

- [79420] 17750.Schwaller, J.; Parganas, E.; Wang, D.; Cain, D.; Aster, J. C.;Williams, I. R.; Lee, C.-K.; Gerthner, R.; Kitamura, T.; Frantsve,J.; Anastasiadou, E.; Loh, M. L.; Levy, D. E.; Ihle, J. N.; Gilliland,D. G.: Stat5 is essential for the myelo- and lymphoproliferativedisease induced by TEL/ JAK2. *Molec. Cell* 6: 693-704, 2000.
- [79421] 17751.Watling, D.; Guschin, D.; Muller, M.; Silvennoinen, O.; Witthuhn,B. A.; Quelle, F. W.; Rogers, N. C.; Schindler, C.; Stark, G. R.;Ihle, J. N.; Kerr, I. M.: Complementation by the protein tyrosinekinase JAK2 of a mutant cell line defective in the interferon-gammasignal transduction pathway. *Nature* 366: 166-170, 1993.
- [79422] 17752.Cihakova, D.; Trebusak, K.; Heino, M.; Fadeyev, V.; Tiulpakov,A.; Battelino, T.; Tar, A.; Halasz, Z.; Blumel, P.; Tawfik, S.; Krohn,K.; Lebl, J.; Peterson, P.; MEWPE-APECED Study Group: Novel AIREmutations and P450 cytochrome autoantibodies in Central and EasternEuropean patients with APECED. *Hum. Mutat.* 18: 225-232, 2001.
- [79423] 17753.Clemente, M. G.; Obermayer-Straub, P.; Meloni, A.; Strassburg,C. P.; Arangino, V.; Tukey, R. H.; de Virgiliis, S.;



Manns, M. P.: Cytochrome P450 1A2 is a hepatic autoantigen in autoimmune polyglandular syndrome type 1. *J. Clin. Endocr. Metab.* 82: 1353–1361, 1997.

[79424] 17754. Craig, J. M.; Schiff, L. H.; Boone, J. E.: Chronic moniliasis associated with Addison's disease. *Am. J. Dis. Child.* 89: 669–684, 1955.

[79425] 17755. Ekwall, O.; Hedstrand, H.; Haavik, J.; Perheentupa, J.; Betterle, C.; Gustafsson, J.; Husebye, E.; Rorsman, F.; Kampe, O.: Pteridin-dependent hydroxylases as autoantigens in autoimmune polyendocrine syndrome type I. *J. Clin. Endocr. Metab.* 85: 2944–2950, 2000.

[79426] 17756. Cetani, F.; Barbesino, G.; Borsari, S.; Pardi, E.; Cianferotti, L.; Pinchera, A.; Marcocci, C.: A novel mutation of the autoimmune regulator gene in an Italian kindred with autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy, acting in a dominant fashion and strongly cosegregating with hypothyroid autoimmune thyroiditis. *J. Clin. Endocr. Metab.* 86: 4747–4752, 2001.

[79427] 17757. Finnish–German APECED Consortium: An autoimmune disease, APECED, caused by mutations in a novel gene featuring two PHD-type zinc-finger domains. *Nature Genet.* 17: 399–403, 1997.

[79428] 17758. Foz, M.; Mirada, A.; Guardia, J.: Endocrine disorder–

ders in a family.(Letter) Lancet II: 269 only, 1970.

- [79429] 17759.Gass, J. D. M.: The syndrome of keratoconjunctivitis, superficialmoniliasis, idiopathic hypoparathyroidism and Addison's disease. Am.J. Ophthal. 54: 660–674, 1962.
- [79430] 17760.Gylling, M.; Tuomi, T.; Bjorses, P.; Kontiainen, S.; Partanen,J.; Christie, M. R.; Knip, M.; Perheentupa, J.; Miettinen, A.: Beta-cellautoantibodies, human leukocyte antigen II alleles, and type 1 diabetesin autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy. J.Clin. Endocr. Metab. 85: 4434–4440, 2000.
- [79431] 17761.Hedstrand, H.; Ekwall, O.; Haavik, J.; Landgren, E.; Betterle,C.; Perheentupa, J.; Gustafsson, J.; Husebye, E.; Rorsman, F.; Kampe,O.: Identification of tyrosine hydroxylase as an autoantigen in autoimmune polyendocrine syndrome type I. Biochem. Biophys. Res. Commun. 267:456–461, 2000.
- [79432] 17762.Heino, M.; Peterson, P.; Kudoh, J.; Shimizu, N.; Antonarakis,S. E.; Scott, H. S.; Krohn, K.: APECED mutations in the autoimmuneregulator (AIRE) gene. Hum. Mutat. 18: 205–211, 2001.
- [79433] 17763.Heino, M.; Scott, H. S.; Chen, Q.; Peterson, P.; Maenpaa, U.;Papasavvas, M.–P.; Mittaz, L.; Barras, C.; Rossier, C.; Chrousos,G. P.; Stratakis, C. A.; Nagamine, K.;

Kudoh, J.; Shimizu, N.; Maclaren, N.; Antonarakis, S. E.; Krohn, K.: Mutation analyses of North American APS-1 patients. *Hum. Mutat.* 13: 69–74, 1999.

[79434] 17764. Hendrix, T. R.: Personal Communication. Baltimore, Md. 1/14/1985.

[79435] 17765. Hiekkala, H.: Idiopathic hypoparathyroidism, adrenal insufficiency and moniliasis in children. *Ann. Paediat. Fenn.* 10: 213–222, 1964.

[79436] 17766. Hung, W.; Migeon, C. J.; Parrott, R. H.: A possible autoimmune basis for Addison's disease in three siblings, one with idiopathic hypoparathyroidism, pernicious anemia and superficial moniliasis. *New Eng. J. Med.* 269: 658–663, 1963.

[79437] 17767. Husebye, E. S.; Gebre-Medhin, G.; Tuomi, T.; Perheentupa, J.; Landin-Olsson, M.; Gustafsson, J.; Rorsman, F.; Kampe, O.: Autoantibodies against aromatic L-amino acid decarboxylase in autoimmune polyendocrine syndrome type I. *J. Clin. Endocr. Metab.* 82: 147–150, 1997.

[79438] 17768. Ishii, T.; Suzuki, Y.; Ando, N.; Matsuo, N.; Ogata, T.: Novel mutations of the autoimmune regulator gene in two siblings with autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy. *J. Clin. Endocr. Metab.* 85: 2922–2926, 2000.

- [79439] 17769.Kenny, F. M.; Holliday, M. D.: Hypoparathyroidism, moniliasis, Addison's and Hashimoto's disease. Hypercalcemia treated with intravenously administered sodium sulfate. *New Eng. J. Med.* 271: 708–713, 1964.
- [79440] 17770.Krohn, K.; Uibo, R.; Aavik, E.; Peterson, P.; Savilahti, K.: Identification by molecular cloning of an autoantigen associated with Addison's disease as steroid 17-alpha-hydroxylase. *Lancet* 339: 770–773, 1992.
- [79441] 17771.Kunin, A. S.; MacKay, B. R.; Burns, S. L.; Halberstam, M. J.: The syndrome of hypoparathyroidism and adrenocortical insufficiency, a possible sequel of hepatitis: case report and review of the literature. *Am. J. Med.* 34: 856–866, 1963.
- [79442] 17772.Louria, D. B.; Shannon, D. C.; Johnson, G.; Caroline, L.; Okas, A.; Taschdjian, C.: The susceptibility to moniliasis in children with endocrine hypofunction. *Trans. Assoc. Am. Phys.* 80: 236–249, 1967.
- [79443] 17773.Maghnie, M.; Cosi, G.; Genovese, E.; Manca-Bitti, M. L.; Cohen, A.; Zecca, S.; Tinelli, C.; Gallucci, M.; Bernasconi, S.; Boscherini, B.; Severi, F.; Arico, M.: Central diabetes insipidus in children and young adults. *New Eng. J. Med.* 343: 998–1007, 2000.
- [79444] 17774.Marieb, N. J.; Melby, J. C.; Lyall, S. S.: Isolated hy-

poaldosteronism associated with idiopathic hypoparathyroidism. Arch. Intern. Med. 134:424–429, 1974.

- [79445] 17775. McKusick, V. A.: Personal Communication. Baltimore, Md. 1985.
- [79446] 17776. Meloni, A.; Perniola, R.; Faa, V.; Corvaglia, E.; Cao, A.; Rosatelli, M. C.: Delineation of the molecular defects in the AIRE gene in autoimmune polyendocrinopathy–candidiasis–ectodermal dystrophy patients from Southern Italy. J. Clin. Endocr. Metab. 87: 841–846, 2002.
- [79447] 17777. Nagamine, K.; Peterson, P.; Scott, H. S.; Kudoh, J.; Minoshima, S.; Heino, M.; Krohn, K. J. E.; Lalioti, M. D.; Mullis, P. E.; Antonarakis, S. E.; Kawasaki, K.; Asakawa, S.; Ito, F.; Shimizu, N.: Positional cloning of the APECED gene. Nature Genet. 17: 393–398, 1997.
- [79448] 17778. Neufeld, M.; Maclaren, N.; Blizzard, R.: Autoimmune polyglandular syndrome. Pediat. Ann. 9: 154–162, 1980.
- [79449] 17779. Neufeld, M.; Maclaren, N. K.; Blizzard, R. M.: Two types of autoimmune Addison's disease associated with different polyglandular autoimmune (PGA) syndromes. Medicine 60: 355–362, 1981.
- [79450] 17780. Nithiyananthan, R.; Heward, J. M.; Allahabadia, A.; Barnett, A. H.; Franklyn, J. A.; Gough, S. C. L.: A heterozy-

gous deletion of the autoimmune regulator (AIRE1) gene, autoimmune thyroid disease, and type 1 diabetes: no evidence for association. *J. Clin. Endocr. Metab.* 85: 1320–1322, 2000.

[79451] 17781. Okano, M.; Bell, D. W.; Haber, D. A.; Li, E.: DNA methyltransferases Dnmt3a and Dnmt3b are essential for de novo methylation and mammalian development. *Cell* 99: 247–257, 1999.

[79452] 17782. Wijmenga, C.; van den Heuvel, L. P. W. J.; Strengman, E.; Luyten, J. A. F. M.; van der Burgt, I. J. A. M.; de Groot, R.; Smeets, D. F. C. M.; Draaisma, J. M. T.; van Dongen, J. J.; De Abreu, R. A.; Pearson, P. L.; Sandkuijl, L. A.; Weemaes, C. M. R.: Localization of the ICF syndrome to chromosome 20 by homozygosity mapping. *Am. J. Hum. Genet.* 63: 803–809, 1998.

[79453] 17783. Xu, G.-L.; Bestor, T. H.; Bourc'his, D.; Hsieh, C.-L.; Tommerup, N.; Bugge, M.; Hulten, M.; Qu, X.; Russo, J. J.; Viegas-Pequignot, E.: Chromosome instability and immunodeficiency syndrome caused by mutations in a DNA methyltransferase gene. *Nature* 402: 187–191, 1999.

[79454] 17784. Chuang, D. T.; Shih, V. E.: Disorders of branched chain amino acid and keto acid metabolism. In: Scriver, C. R.; Beaudet, A. L.; Sly, W. S.; Valle, D.: *Metabolic and*

Molecular Bases of Inherited Disease. New York: McGraw-Hill (pub.) (7th ed.): 1995. Pp. 1239–1277.

[79455] 17785. Smeitink, J.; van den Heuvel, L.: Human mitochondrial complex I in health and disease. *Am. J. Hum. Genet.* 64: 1505–1510, 1999.

[79456] 17786. Pizarro, T. T.; Michie, M. H.; Bentz, M.; Woraratanadharm, J.; Smith, M. F., Jr.; Foley, E.; Moskaluk, C. A.; Bickston, S. J.; Cominelli, F.: IL-18, a novel immunoregulatory cytokine, is up-regulated in Crohn's disease: expression and localization in intestinal mucosal cells. *J. Immun.* 162: 6829–6835, 1999.

[79457] 17787. Le Coniat, M.; Kinet, J. P.; Berger, R.: The human genes for the  $\alpha$  and  $\gamma$  subunits of the mast cell receptor for immunoglobulin E are located on human chromosome band 1q23. *Immunogenetics* 32: 183–186, 1990.

[79458] 17788. 't Hart, L. M.; Stolk, R. P.; Heine, R. J.; Grobbee, D. E.; vander Does, F. E. E.; Maassen, J. A.: Association of the insulin-receptor variant met-985 with hyperglycemia and non-insulin-dependent diabetes mellitus in the Netherlands: a population-based study. *Am. J. Hum. Genet.* 59: 1119–1125, 1996.

[79459] 17789. Accili, D.; Drago, J.; Lee, E. J.; Johnson, M. D.; Cool,

M. H.;Salvatore, P.; Asico, L. D.; Jose, P. A.; Taylor, S. I.; Westphal,H.: Early neonatal death in mice homozygous for a null allele ofthe insulin receptor gene. *Nature Genet.* 12: 106–109, 1996.

[79460] 17790.Accili, D.; Frapier, C.; Mosthaf, L.; McKeon, C.; El-bein, S. C.;Permutt, M. A.; Ramos, E.; Lander, E.; Ullrich, A.; Taylor, S. I.: A mutation in the insulin receptor gene that impairs transport ofthe receptor to the plasma membrane and causes insulin-resistant diabetes. *EMBOJ.* 8: 2509–2517, 1989.

[79461] 17791.Al-Gazali, L. I.; Khalil, M.; Devadas, K.: A syndrome of insulinresistance resembling leprechaunism in five sibs of consanguineousparents. *J. Med. Genet.* 30: 470–475, 1993.

[79462] 17792.Bar, R. S.; Muggeo, M.; Roth, J.; Kahn, C. R.; Havrankova, J.;Imperato-McGinley, J.: Insulin resistance, acanthosis nigricans,and normal insulin receptors in a young woman: evidence for a postreceptordefect. *J. Clin. Endocr. Metab.* 47: 620–625, 1978.

[79463] 17793.Barbetti, F.; Gejman, P. V.; Taylor, S. I.; Raben, N.; Cama, A.;Bonora, E.; Pizzo, P.; Moghetti, P.; Muggeo, M.; Roth, J.: Detectionof mutations in insulin receptor gene by denaturing gradient gel electrophoresis. *Diabetes*



41:408–415, 1992.

- [79464] 17794. Barnes, N. D.; Palumbo, P. J.; Hayles, A. B.; Folgar, H.: Insulin resistance, skin changes, and virilization: a recessively inherited syndrome possibly due to pineal gland dysfunction. *Diabetologia* 10:285–289, 1974.
- [79465] 17795. Belke, D. D.; Betuing, S.; Tuttle, M. J.; Gravelleau, C.; Young, M. E.; Pham, M.; Zhang, D.; Cooksey, R. C.; McClain, D. A.; Litwin, S. E.; Taegtmeyer, H.; Severson, D.; Kahn, C. R.; Abel, E. D.: Insulin signaling coordinately regulates cardiac size, metabolism, and contractile protein isoform expression. *J. Clin. Invest.* 109: 629–639, 2002.
- [79466] 17796. Benecke, H.; Flier, J. S.; Moller, D. E.: Alternatively spliced variants of the insulin receptor protein: expression in normal and diabetic human tissues. *J. Clin. Invest.* 89: 2066–2070, 1992.
- [79467] 17797. Bruning, J. C.; Gautam, D.; Burks, D. J.; Gillette, J.; Schubert, M.; Orban, P. C.; Klein, R.; Krone, W.; Muller-Wieland, D.; Kahn, C. R.: Role of brain insulin receptor in control of body weight and reproduction. *Science* 289: 2122–2125, 2000.
- [79468] 17798. Bruning, J. C.; Michael, M. D.; Winnay, J. N.; Hayashi, T.; Horsch, D.; Accili, D.; Goodyear, L. J.; Kahn, C. R.: A muscle-specific insulin receptor knockout exhibits

features of the metabolic syndrome of NIDDM without altering glucose tolerance. *Molec. Cell* 2: 559–569, 1998.

[79469] 17799. Cama, A.; de la Luz Sierra, M.; Ottini, L.; Kadowaki, T.; Gorden, P.; Imperato-McGinley, J.; Taylor, S. I.: A mutation in the tyrosine kinase domain of the insulin receptor associated with insulin resistance in an obese woman. *J. Clin. Endocr. Metab.* 73: 894–901, 1991.

[79470] 17800. Henry, I.; Humphries, S. E.; Tata, F.; Barichard, F.; Holm, M.; Williamson, R.; Junien, C.: The gene for HMG CoA reductase (HMGCR) is on human chromosome 5. (Abstract) *Cytogenet. Cell Genet.* 40:649–650, 1985.

[79471] 17801. Humphries, S. E.; Tata, F.; Henry, I.; Barichard, F.; Holm, M.; Junien, C.; Williamson, R.: The isolation, characterisation, and chromosomal assignment of the gene for human 3-hydroxy-3-methylglutarylcoenzyme A reductase, (HMG-CoA reductase). *Hum. Genet.* 71: 254–258, 1985.

[79472] 17802. Julier, C.; Delepine, M.; Keavney, B.; Terwilliger, J.; Davis, S.; Weeks, D. E.; Bui, T.; Jeunemaitre, X.; Velho, G.; Froguel, P.; Ratcliffe, P.; Corvol, P.; Soubrier, F.; Lathrop, G. M.: Genetic susceptibility for human familial essential hypertension in a region of homology with blood pressure linkage on rat chromosome 10. *Hum. Molec. Genet.* 6:

2077–2085, 1997.

- [79473] 17803.Rutherford, S.; Johnson, M. P.; Curtain, R. P.; Griffiths, L.R.: Chromosome 17 and the inducible nitric oxide synthase gene in human essential hypertension. *Hum. Genet.* 109: 408–415, 2001.
- [79474] 17804.Aprelikova, O.; Pajusola, K.; Partanen, J.; Armstrong, E.; Alitalo, R.; Bailey, S. K.; McMahon, J.; Wasmuth, J.; Huebner, K.; Alitalo, K.: FLT4, a novel class III receptor tyrosine kinase in chromosome 5q33–qter. *Cancer Res.* 52: 746–748, 1992.
- [79475] 17805.Dumont, D. J.; Jussila, L.; Taipale, J.; Lymboussaki, A.; Mustonen, T.; Pajusola, K.; Breitman, M.; Alitalo, K.: Cardiovascular failure in mouse embryos deficient in VEGF receptor-3. *Science* 282: 946–949, 1998.
- [79476] 17806.Evans, A. L.; Brice, G.; Sotirova, V.; Mortimer, P.; Beninson, J.; Burnand, K.; Rosbotham, J.; Child, A.; Sarfarazi, M.: Mapping of primary congenital lymphedema to the 5q35.3 region. *Am. J. Hum. Genet.* 64: 547–555, 1999.
- [79477] 17807.Ferrell, R. E.; Levinson, K. L.; Esman, J. H.; Kimak, M. A.; Lawrence, E. C.; Barmada, M. M.; Finegold, D. N.: Hereditary lymphedema: evidence for linkage and genetic heterogeneity. *Hum. Molec. Genet.* 7: 2073–2078, 1998.
- [79478] 17808.Galland, F.; Karamysheva, A.; Mattei, M.-G.; Ros-

net, O.; Marchetto,S.; Birnbaum, D.: Chromosomal localization of FLT4, a novel receptor–typetyrosine kinase gene. *Genomics* 13: 475–478, 1992.

[79479] 17809.Irrthum, A.; Karkkainen, M. J.; Devriendt, K.; Alitalo, K.; Vikkula,M.: Congenital hereditary lymphedema caused by a mutation that inactivatesVEGFR3 tyrosine kinase. *Am. J. Hum. Genet.* 67: 295–301, 2000.

[79480] 17810.Kaipainen, A.; Korhonen, J.; Mustonen, T.; van Hinsbergh, V. W.M.; Fang, G.–H.; Dumont, D.; Breitman, M.; Alitalo, K.: Expressionof the fms–like tyrosine kinase 4 gene becomes restricted to lymphaticendothelium during development. *Proc. Nat. Acad. Sci.* 92: 3566–3570,1995.

[79481] 17811.Karkkainen, M. J.; Ferrell, R. E.; Lawrence, E. C.; Kimak, M. A.;Levinson, K. L.; McTigue, M. A.; Alitalo, K.; Finegold, D. N.: Missensemutations interfere with VEGFR–3 signalling in primary lymphoedema. *NatureGenet.* 25: 153–159, 2000.

[79482] 17812.Karkkainen, M. J.; Saaristo, A.; Jussila, L.; Karila, K. A.; Lawrence,E. C.; Pajusola, K.; Bueler, H.; Eichmann, A.; Kauppinen, R.; Kettunen,M. I.; Yla–Herttuala, S.; Finegold, D. N.; Ferrell, R. E.; Alitalo,K.: A model for gene therapy of human hereditary lymphedema. *Proc.Nat. Acad. Sci.* 98: 12677–12682, 2001.

- [79483] 17813.Lee, J.; Gray, A.; Yuan, J.; Luoh, S.-M.; Avraham, H.; Wood, W.I.: Vascular endothelial growth factor-related protein: a ligand and specific activator of the tyrosine kinase receptor Flt4. *Proc.Nat. Acad. Sci.* 93: 1988–1992, 1996.
- [79484] 17814.Lyon, M. F.; Glenister, P. H.: New Mutants. *Mouse Newsletter* 71:26 only, 1984.
- [79485] 17815.Lyon, M. F.; Glenister, P. H.: Gene order of Chyt-Re on chromosome 11. *Mouse Newsletter* 74: 96 only, 1986.
- [79486] 17816.Milroy, W. F.: An undescribed variety of hereditary oedema. *New York Med. J.* 56: 505–508, 1892.
- [79487] 17817.Offori, T. W.; Platt, C. C.; Stephens, M.; Hopkinson, G. B.: Angiosarcoma in congenital hereditary lymphoedema (Milroy's disease): diagnostic beacons and a review of the literature. *Clin. Exp. Derm.* 18:174–177, 1993.
- [79488] 17818.Pajusola, K.; Aprelikova, O.; Korhonen, J.; Kaipainen, A.; Pertovaara, L.; Alitalo, R.; Alitalo, K.: FLT4 receptor tyrosine kinase contains seven immunoglobulin-like loops and is expressed in multiple human tissues and cell lines. *Cancer Res.* 52: 5738–5743, 1992.
- [79489] 17819.Pajusola, K.; Aprelikova, O.; Pelicci, G.; Weich, H.; Claesson-Welsh, L.; Alitalo, K.: Signalling properties of

FLT4, a proteolytically processed receptor tyrosine kinase related to two VEGF receptors. *Oncogene* 9:3545–3555, 1994.

[79490] 17820. Walter, J. W.; North, P. E.; Waner, M.; Mizeracki, A.; Blei, F.; Walker, J. W. T.; Reinisch, J. F.; Marchuk, D. A.: Somatic mutation of vascular endothelial growth factor receptors in juvenile hemangioma. *Genes Chromosomes Cancer* 33: 295–303, 2002.

[79491] 17821. Ragoussis, J.; Senger, G.; Trowsdale, J.; Campbell, I. G.: Genomic organization of the human folate receptor genes on chromosome 11q13. *Genomics* 14:423–430, 1992.

[79492] 17822. Barber, R. C.; Shaw, G. M.; Lammer, E. J.; Greer, K. A.; Biela, T. A.; Lacey, S. W.; Wasserman, C. R.; Finnell, R. H.: Lack of association between mutations in the folate receptor- $\alpha$  gene and spina bifida. *Am. J. Med. Genet.* 76: 310–317, 1998.

[79493] 17823. Bowcock, A. M.; Lacey, S.; Saltman, D.; Mohandas, T. K.; Kamen, B. A.; Taggart, R. T.: Localization of the folate receptor gene to chromosome 11q13. (Abstract) *Cytogenet. Cell Genet.* 58: 1955 only, 1991.

[79494] 17824. Campbell, I. G.; Jones, T. A.; Foulkes, W. D.; Trowsdale, J.: Folate-binding protein is a marker for ovarian can-

cer. Cancer Res. 51:5329–5338, 1991.

[79495] 17825.Kranes, A.; Balogh, K., Jr.: Liver disease in a patient with von Hippel–Lindau disease. New Eng. J. Med. 275: 950–959, 1966.

[79496] 17826.Lamiell, J. M.: Personal Communication. San Francisco, Calif. 9/1987.

[79497] 17827.Lamiell, J. M.; Salazar, F. G.; Hsia, Y. E.: Von Hippel–Lindau disease affecting 43 members of a single kindred. Medicine 68: 1–29, 1989.

[79498] 17828.Latif, F.; Tory, K.; Gnarr, J.; Yao, M.; Duh, F.–M.; Orcutt, M. L.; Stackhouse, T.; Kuzmin, I.; Modi, W.; Geil, L.; Schmidt, L.; Zhou, F.; Li, H.; Wei, M. H.; Chen, F.; Glenn, G.; Choyke, P.; Walther, M. M.; Weng, Y.; Duan, D.–S. R.; Dean, M.; Glavac, D.; Richards, F.M.; Crossey, P. A.; Ferguson–Smith, M. A.; Le Paslier, D.; Chumakov, I.; Cohen, D.; Chinnault, A. C.; Maher, E. R.; Linehan, W. M.; Zbar, B.; Lerman, M. I.: Identification of the von Hippel–Lindau disease tumor suppressor gene. Science 260: 1317–1320, 1993.

[79499] 17829.Lee, S.; Chen, D. Y. T.; Humphrey, J. S.; Gnarr, J. R.; Linehan, W. M.; Klausner, R. D.: Nuclear/cytoplasmic localization of the von Hippel–Lindau tumor suppressor gene product is determined by cell density. Proc. Nat. Acad. Sci. 93: 1770–1775, 1996.

[79500] 17830.Lenz, T.; Thiede, H. M.; Nussberger, J.; Atlas, S. A.; Distler,A.; Schulte, K. L.: Hyperreninemia and secondary hyperaldosteronism in a patient with pheochromocytoma and von Hippel–Lindau disease. Nephron 62:345–350, 1992.



- [79501] 17831.Lindau, A.: Zur Frage der Angiomatosis Retinae und Ihrer Hirncomplication. ActaOphthal. 4: 193–226, 1927.
- [79502] 17832.Loeb, D. B.; Pericak–Vance, M. A.; Stajich, J. M.; Vance, J. M.: A novel mutation in the von Hippel–Lindau gene. Hum. Molec. Genet. 3:1423–1424, 1994.
- [79503] 17833.Lui, W. O.; Chen, J.; Glasker, S.; Bender, B. U.; Madura, C.;Khoo, S. K.; Kort, E.; Larsson, C.; Neumann, H. P. H.; Teh, B. T.: Selective loss of chromosome 11 in pheochromocytomas associatedwith the VHL syndrome. Oncogene 21: 1117–1122, 2002.
- [79504] 17834.Maddock, I. R.; Moran, A.; Maher, E. R.; Teare, M. D.; Norman,A.; Payne, S. J.; Whitehouse, R.; Dodd, C.; Lavin, M.; Hartley, N.;Super, M.; Evans, D. G. R.: A genetic register for von Hippel–Lindaudisease. J. Med. Genet. 33: 120–127, 1996.
- [79505] 17835.Maher, E. R.; Bentley, E.; Yates, J. R. W.; Barton, D.; Jennings,A.; Fellows, I. W.; Ponder, M. A.; Ponder, B. A. J.; Benjamin, C.;Harris, R.; Ferguson–Smith, M. A.: Mapping of von Hippel–Lindau diseaseto chromosome 3p confirmed by genetic linkage analysis. J. Neurol.Sci. 100: 27–30, 1990.
- [79506] 17836.Maher, E. R.; Bentley, E.; Yates, J. R. W.; Latif, F.;

Lerman,M.; Zbar, B.; Affara, N. A.; Ferguson-Smith, M. A.: Mapping of the von Hippel-Lindau disease locus to a small region of chromosome 3p by genetic linkage analysis. *Genomics* 10: 957-960, 1991.

[79507] 17837. Maher, E. R.; Iselius, L.; Yates, J. R. W.; Littler, M.; Benjamin, C.; Harris, R.; Sampson, J.; Williams, A.; Ferguson-Smith, M. A.; Morton, N.: Von Hippel-Lindau disease: a genetic study. *J. Med. Genet.* 28: 443-447, 1991.

[79508] 17838. Maher, E. R.; Yates, J. R. W.; Ferguson-Smith, M. A.: Statistical analysis of the two stage mutation model in von Hippel-Lindau disease, and in sporadic cerebellar haemangioblastoma and renal cell carcinoma. *J. Med. Genet.* 27: 311-314, 1990.

[79509] 17839. Mahon, P. C.; Hirota, K.; Semenza, G. L.: FIH-1: a novel protein that interacts with HIF-1- $\alpha$  and VHL to mediate repression of HIF-1 transcriptional activity. *Genes Dev.* 15: 2675-2686, 2001. 100. Manski, T. J.; Heffner, D. K.; Glenn, G. M.; Patronas, N. J.; Pikus, A. T.; Katz, D.; Lebovics, R.; Sledjeski, K.; Choyke, P. L.; Zbar, B.; Linehan, W. M.; Oldfield, E. H.: Endolymphatic sac tumors: a source of morbid hearing loss in von Hippel-Lindau disease. *J. A. M. A.* 277: 1461-1466, 1997. 101. Maxwell, P. H.; Wiesener, M. S.; Chang, G.-W.; Clifford, S. C.; Vaux, E. C.;

Cockman, M. E.; Wykoff, C. C.; Pugh, C. W.; Maher, E.R.; Ratcliffe, P. J.: The tumour suppressor protein VHL targets hypoxia-inducible factors for oxygen-dependent proteolysis. *Nature* 399: 271–275, 1999.102. McCabe, C. M.; Flynn, H. W., Jr.; Shields, C. L.; Shields, J.A.; Regillo, C. D.; McDonald, H. R.; Berrocal, M. H.; Gass, J. D.M.; Mieler, W. F.: Juxtapapillary capillary hemangiomas: clinical features and visual acuity outcomes. *Ophthalmology* 107: 2240–2249, 2000.103. McKusick, V. A.; Abbey, H.; Bettersby, E. J.; Borhani, N. O.; Boyer, S. H., IV; Cohen, B. H.; Ferguson-Smith, M. A.; Franke, F.R.; Gordon, H.; Handmaker, S. D.; Harris, W. S.; Hawkins, M. R.; and 13 others: Medical genetics 1960. *J. Chronic Dis.* 14: 1–198, 1961. Fig. 71.104. Melmon, K. L.; Rosen, S. W.: Lindau's disease: review of the literature and study of a large kindred. *Am. J. Med.* 36: 595–617, 1964.105. Min, J.-H.; Yang, H.; Ivan, M.; Gertler, F.; Kaelin, W. G., Jr.; Pavletich, N. P.: Structure of an HIF-1- $\alpha$ -pVHL complex: hydroxyproline recognition in signaling. *Science* 296: 1886–1889, 2002.106. Mukhopadhyay, D.; Knebelmann, B.; Cohen, H. T.; Ananth, S.; Sukhatme, V. P.: The von Hippel-Lindau tumor suppressor gene product interacts with Sp1 to repress vascular endothelial growth factor promoter activity. *Molec.*

Cell. Biol. 17: 5629–5639, 1997.107. Neumann, H. P. H.; Berger, D. P.; Sigmund, G.; Blum, U.; Schmidt, D.; Parmer, R. J.; Volk, B.; Kirste, G.: Pheochromocytomas, multiple endocrine neoplasia type 2, and von Hippel–Lindau disease. New Eng. J. Med. 329: 1531–1538, 1993.108. Neumann, H. P. H.; Eng, C.; Mulligan, L. M.; Glavac, D.; Zauner, I.; Ponder, B. A. J.; Crossey, P. A.; Maher, E. R.; Brauch, H.: Consequences of direct genetic testing for germline mutations in the clinical management of families with multiple endocrine neoplasia, type II. J. A. M. A. 274: 1149–1151, 1995.109. Neumann, H. P. H.; Wiestler, O. D.: Clustering of features of von Hippel–Lindau syndrome: evidence for a complex genetic locus. Lancet 337: 1052–1054, 1991.110. Nibbelink, D. W.; Peters, B. H.; McCormick, W. F.: On the association of pheochromocytoma and cerebellar hemangioblastoma. Neurology 19: 455–460, 1969.111. Oberstrass, J.; Reifenberger, G.; Reifenberger, J.; Wechsler, W.; Collins, V. P.: Mutation of the von Hippel–Lindau tumour suppressor gene in capillary haemangioblastomas of the central nervous system. J. Path. 179: 151–156, 1996.112. Ohh, M.; Yauch, R. L.; Lonergan, K. M.; Whaley, J. M.; Stemmer-Rachamimov, A. O.; Louis, D. N.; Gavin, B. J.; Kley, N.; Kaelin, W. G., Jr.; Iliopoulos, O.: The von Hippel–

Lindau tumor suppressor protein is required for proper assembly of an extracellular fibronectin matrix. *Molec. Cell* 1:959–968, 1998.113. Olschwang, S.; Richard, S.; Boisson, C.; Giraud, S.; Laurent–Puig, P.; Resche, F.; Thomas, G.: Germline mutation profile of the VHL gene in von Hippel–Lindau disease and in sporadic hemangioblastoma. *Hum. Mutat.* 12: 424–430, 1998.114. Otenasek, F. J.; Silver, M. L.: Spinal hemangioma (hemangioblastoma) in Lindau's disease: report of six cases in a single family. *J. Neurosurg.* 18:295–300, 1961.115. Pack, S. D.; Zbar, B.; Pak, E.; Ault, D. O.; Humphrey, J. S.; Pham, T.; Hurley, K.; Weil, R. J.; Park, W.–S.; Kuzmin, I.; Stolle, C.; Glenn, G.; Liotta, L. A.; Lerman, M. I.; Klausner, R. D.; Linehan, W. M.; Zhuang, Z.: Constitutional von Hippel–Lindau (VHL) gene deletions detected in VHL families by fluorescence in situ hybridization. *Cancer Res.* 59: 5560–5564, 1999.116. Pause, A.; Lee, S.; Lonergan, K. M.; Klausner, R. D.: The von Hippel–Lindau tumor suppressor gene is required for cell cycle exit upon serum withdrawal. *Proc. Nat. Acad. Sci.* 95: 993–998, 1998.117. Price, E. B., Jr.: Papillary cystadenoma of the epididymis: a clinicopathologic analysis of 20 cases. *Arch. Path.* 91: 456–470, 1971.118. Probst, A.; Lotz, M.; Heitz, P.: Von Hippel–Lindau's dis–

ease, syringomyelia and multiple endocrine tumors: a complex neuroendocrinopathy. *Virchows Arch. Path. Anat. Histol.* 378: 265–272, 1978.119. Prowse, A. H.; Webster, A. R.; Richards, F. M.; Richard, S.; Olschwang, S.; Resche, F.; Affara, N. A.; Maher, E. R.: Somatic inactivation of the VHL gene in Von Hippel–Lindau disease tumors. *Am. J. Hum. Genet.* 60: 765–771, 1997.120. Rho, Y. M.: Von Hippel–Lindau's disease: a report of 5 cases. *Canad. Med. Assoc. J.* 101: 135–142, 1969.121. Richard, S.; Croisille, L.; Yvart, J.; Casadeval, N.; Eschwege, P.; Aghakhani, N.; David, P.; Gaudric, A.; Scigalla, P.; Hermine, O.: Paradoxical secondary polycythemia in von Hippel–Lindau patient treated with anti-vascular endothelial growth factor receptor therapy. *Blood* 99:3851–3853, 2002.122. Richards, F. M.; Crossey, P. A.; Phipps, M. E.; Foster, K.; Latif, F.; Evans, G.; Sampson, J.; Lerman, M. I.; Zbar, B.; Affara, N. A.; Ferguson–Smith, M. A.; Maher, E. R.: Detailed mapping of germline deletions of the von Hippel–Lindau disease tumour suppressor gene. *Hum. Molec. Genet.* 3: 595–598, 1994.123. Richards, F. M.; Maher, E. R.; Latif, F.; Phipps, M. E.; Tory, K.; Lush, M.; Crossey, P. A.; Oostra, B.; Gustavson, K. H.; Green, J.; Turner, G.; Yates, J. R. W.; Linehan, W. M.; Affara, N. A.; Lerman, M.; Zbar, B.; Ferguson–

Smith, M. A.: Detailed genetic mapping of the von Hippel-Lindau disease tumour suppressor gene. *J. Med. Genet.* 30:104-107, 1993.124. Richards, F. M.; Phipps, M. E.; Latif, F.; Yao, M.; Crossey, P. A.; Foster, K.; Linehan, W. M.; Affara, N. A.; Lerman, M. I.; Zbar, B.; Ferguson-Smith, M. A.; Maher, E. R.: Mapping the von Hippel-Lindau disease tumour suppressor gene: identification of germline deletions by pulsed field gel electrophoresis. *Hum. Molec. Genet.* 2: 879-882, 1993.125. Richards, F. M.; Schofield, P. N.; Fleming, S.; Maher, E. R.: Expression of the von Hippel-Lindau disease tumour suppressor gene during human embryogenesis. *Hum. Molec. Genet.* 5: 639-644, 1996.126. Rubenstein, J. L.; Yaari, H.: von Hippel-Lindau and the genetics of astrocytoma. *J. Nat. Cancer Inst.* 86: 142-143, 1994.127. Sander, S.; Normann, T.; Mathisen, W.: Pheochromocytoma associated with von-Hippel-Lindau's disease in a family. *Scand. J. Urol. Nephrol.* 4:259-263, 1970.128. Schechterman, L.: Lindau's disease: report of an unusual case and two additional cases in a Negro family. *Med. Ann. D.C.* 30: 64-76, 1961.129. Schimke, R. N.; Collins, D. L.; Rothberg, P. G.: Functioning carotid paraganglioma in the von Hippel-Lindau syndrome. (Letter) *Am. J. Med. Genet.* 80: 533-534, 1998.130.

Schoenfeld, A.; Davidowitz, E. J.; Burk, R. D.: A second major native von Hippel–Lindau gene product, initiated from an internal translation start site, functions as a tumor suppressor. *Proc. Nat. Acad. Sci.* 95: 8817–8822, 1998.131. Seizinger, B. R.; Farmer, G.; Haines, J.; Anderson, K.; Whaley, J.; Hettlich, C.; Decker, J.; Rouleau, G.; Smith, D.; Drabkin, H.; Filling–Katz, M.; Neumann, H.; Collins, D.; Hsia, E.; Green, J.; Waziri, M.; Gusella, J.; Li, F.: Isolating the gene(s) for von Hippel–Lindau disease and renal cell carcinoma. (Abstract) *Am. J. Hum. Genet.* 45(suppl.): A32, 1989.132. Seizinger, B. R.; Rouleau, G. A.; Ozelius, L. J.; Lane, A. H.; Farmer, G. E.; Lamiell, J. M.; Haines, J.; Yuen, J. W.; Collins, D.; Majoor–Krakauer, D.; et al.: Von Hippel–Lindau disease maps to the region of chromosome 3 associated with renal cell carcinoma. *Nature* 332: 268–269, 1988.133. Seizinger, B. R.; Smith, D. I.; Filling–Katz, M. R.; Neumann, H.; Green, J. S.; Choyke, P. L.; Anderson, K. M.; Freiman, R. N.; Klauck, S. M.; Whaley, J.; Decker, H.–J. H.; Hsia, Y. E.; Collins, D.; Halperin, J.; Lamiell, J. M.; Oostra, B.; Waziri, M. H.; Gorin, M. B.; Scherer, G.; Drabkin, H. A.; Aronin, N.; Schinzel, A.; Martuza, R. L.; Gusella, J. F.; Haines, J. L.: Genetic flanking markers refined diagnostic criteria and provide insights into



the genetics of VonHippel Lindau disease. Proc. Nat. Acad. Sci. 88: 2864–2868, 1991.134. Sgambati, M. T.; Stolle, C.; Choyke, P. L.; Walther, M. M.; Zbar, B.; Linehan, W. M.; Glenn, G. M.: Mosaicism in von Hippel–Lindau disease: lessons from kindreds with germline mutations identified in offspring with mosaic parents. Am. J. Hum. Genet. 66: 84–91, 2000.135. Sharp, W. V.; Platt, R. L.: Familial pheochromocytoma: association with von–Hippel–Lindau's disease. Angiology 22: 141–146, 1971.136. Shokeir, M. H. K.: Von Hippel–Lindau syndrome: a report on three kindreds. J. Med. Genet. 7: 155–157, 1970.137. Silver, M. L.: Hereditary vascular tumors of the nervous system. J.A.M.A. 156: 1053–1056, 1954.138. Stolle, C.; Glenn, G.; Zbar, B.; Humphrey, J. S.; Choyke, P.; Walther, M.; Pack, S.; Hurley, K.; Audrey, C.; Klausner, R.; Linehan, W. M.: Improved detection of germline mutations in the von Hippel–Lindau disease tumor suppressor gene. Hum. Mutat. 12: 417–423, 1998.139. Thomas, M.; Burnside, R. M.: Von Hippel–Lindau disease. Am. J. Ophthal. 51: 140–146, 1961.140. Tisherman, S. E.; Gregg, F. J.; Danowski, T. S.: Familial pheochromocytoma. J.A.M.A. 182: 152–156, 1962.141. Tisherman, S. E.; Tisherman, B. G.; Tisherman, S. A.; Dunmire, S.; Levey, G. S.; Mulvihill, J. J.: Three–

decade investigation of familial pheochromocytoma: an allele of von Hippel-Lindau disease? *Arch. Int. Med.* 153: 2550-2556, 1993.142. Tory, K.; Brauch, H.; Linehan, M.; Barba, D.; Oldfield, E.; Filling-Katz, M.; Seizinger, B.; Nakamura, Y.; White, R.; Marshall, F. F.; Lerman, M. I.; Zbar, B.: Specific genetic change in tumors associated with von Hippel-Lindau disease. *J. Nat. Cancer Inst.* 81: 1097-1101, 1989.143. Tsuda, H.; Fukushima, S.; Takahashi, M.; Hikosaka, Y.; Hayashi, K.: Familial bilateral papillary cystadenoma of the epididymis: report of three cases in siblings. *Cancer* 37: 1831-1839, 1976.144. Tyers, M.; Willems, A. R.: One ring to rule a superfamily of E3 ubiquitin ligases. *Science* 284: 602-604, 1999.145. Vance, J. M.; Small, K. W.; Jones, M. A.; Stajich, J. M.; Yamaoka, L. H.; Roses, A. D.; Hung, W.-Y.; Pericak-Vance, M. A.: Confirmation of linkage in von Hippel-Lindau disease. *Genomics* 6: 565-567, 1990.146. van der Harst, E.; de Krijger, R. R.; Dinjens, W. N. M.; Weeks, L. E.; Bonjer, H. J.; Bruining, H. A.; Lamberts, S. W. J.; Koper, J. W.: Germline mutations in the *vhl* gene in patients presenting with pheochromocytomas. *Int. J. Cancer.* 77: 337-340, 1998.147. Versteeg, R.: Aberrant methylation in cancer. (Editorial) *Am. J. Hum. Genet.* 60: 751-754, 1997.148. Vogelstein, B.: Personal

Communication. Baltimore, Md. 1/6/1995.149. von Hippel, E.: Ueber eine sehr seltene Erkrankung der Netzhaut. Albrecht von Graefes Arch. Ophthalm. 59: 83–106, 1904.150. Webster, A. R.; Maher, E. R.; Bird, A. C.; Moore, A. T.: Risk of multisystem disease in isolated ocular angioma (haemangioblastoma). J. Med. Genet. 37: 62–63, 2000.151. Webster, A. R.; Richards, F. M.; MacDonald, F. E.; Moore, A. T.; Maher, E. R.: An analysis of phenotypic variation in the familial cancer syndrome von Hippel–Lindau disease: evidence for modifier effects. Am. J. Hum. Genet. 63: 1025–1035, 1998.152. Wells, R. A.; Reeders, S. T.; Green, J.; Johnson, G. J.; Robson, K. J. H.: Linkage studies of von Hippel–Lindau syndrome using multiple-locus hypervariable DNA probes. (Abstract) Cytogenet. Cell Genet. 46:714, 1987.153. Wesolowski, D. P.; Ellwood, R. A.; Schwab, R. E.; Farah, J.: Hippel–Lindau syndrome in identical twins. Brit. J. Radiol. 54:982–986, 1981.154. Wiesener, M. S.; Seyfarth, M.; Warnecke, C.; Jurgensen, J. S.; Rosenberger, C.; Morgan, N. V.; Maher, E. R.; Frei, U.; Eckardt, K.-U.: Paraneoplastic erythrocytosis associated with an inactivating point mutation of the von Hippel–Lindau gene in a renal cell carcinoma. Blood 99:3562–3565, 2002.155. Wise, K. S.; Gibson, J. A.: Von Hippel–Lindau's

disease andphaeochromocytoma. Brit. Med. J. 1: 441, 1971.156. Wyburn–Mason, R.: Arteriovenous aneurysm of mid–brain and retina, facial naevi and mental changes. Brain 66: 163–203, 1943.157. Zatyka, M.; da Silva, N. F.; Clifford, S. C.; Morris, M. R.; Wiesener, M. S.; Eckardt, K.–U.; Houlston, R. S.; Richards, F. M.; Latif, F.; Maher, E. R.: Identification of cyclin D1 and other novel targets for the von Hippel–Lindau tumor suppressor gene by expression array analysis and investigation of cyclin D1 genotype as a modifier in von Hippel–Lindau disease. Cancer Res. 62: 3803–3811, 2002.158. Zbar, B.; Kishida, T.; Chen, F.; Schmidt, L.; Maher, E. R.; Richards, F. M.; Crossey, P. A.; Webster, A. R.; Affara, N. A.; Ferguson–Smith, M. A.; Brauch, H.; Glavac, D.; and 14 others: Germline mutation–sin the Von Hippel–Lindau disease (VHL) gene in families from North America, Europe, and Japan. Hum. Mutat. 8: 348–357, 1996.159. Zhuang, Z.; Emmert–Buck, M. R.; Roth, M. J.; Gnarr, J.; Linehan, W. M.; Liotta, L. A.; Luben–sky, I. A.: Von Hippel–Lindau disease gene deletion detected in microdissected sporadic human colon carcinomaspecimens. Hum. Path. 27: 152–156, 1996.

[79510] 17840. Thomas, J. T.; Kilpatrick, M. W.; Lin, K.; Erlacher, L.; Lembessis, P.; Costa, T.; Tsipouras, P.; Luyten, F. P.: Dis–

ruption of human limb morphogenesis by a dominant negative mutation in CDMP1. *Nature Genet.* 58–64, 1997.

- [79511] 17841.Devlin, R. H.; Deeb, S.; Brunzell, J.; Hayden, M. R.: Partial gene duplication involving exon–Alu interchange results in lipoprotein lipase deficiency. *Am. J. Hum. Genet.* 46: 112–119, 1990.
- [79512] 17842.Dichek, H. L.; Fojo, S. S.; Beg, O. U.; Skarlatos, S. I.; Brunzell, J. D.; Cutler, G. B., Jr.; Brewer, H. B., Jr.: Identification of 2 separate allelic mutations in the lipoprotein lipase gene of a patient with the familial hyperchylomicronemia syndrome. *J. Biol. Chem.* 266:473–477, 1991.
- [79513] 17843.Eckel, R. H.: Lipoprotein lipase: a multifunctional enzyme relevant to common metabolic diseases. *New Eng. J. Med.* 320: 1060–1068, 1989.
- [79514] 17844.Emi, M.; Hata, A.; Robertson, M.; Iverius, P.–H.; Hegele, R.; Lalouel, J.–M.: Lipoprotein lipase deficiency resulting from a nonsense mutation in exon 3 of the lipoprotein lipase gene. *Am. J. Hum. Genet.* 47:107–111, 1990.
- [79515] 17845.Emi, M.; Wilson, D. E.; Iverius, P.–H.; Wu, L.; Hata, A.; Hegele, R.; Williams, R. R.; Lalouel, J.–M.: Missense mutation (gly–to–glu188) of human lipoprotein lipase imparting functional deficiency. *J. Biol. Chem.* 265: 5910–5916,

1990.

- [79516] 17846. Faustinella, F.; Chang, A.; Van Biervliet, J. P.; Rosseneu, M.; Vinaimont, N.; Smith, L. C.; Chen, S.-H.; Chan, L.: Catalytic triad residue mutation (asp156-to-gly) causing familial lipoprotein lipase deficiency: co-inheritance with a nonsense mutation (ser447-to-ter) in a Turkish family. *J. Biol. Chem.* 266: 14418–14424, 1991.
- [79517] 17847. Feoli-Fonseca, J. C.; Levy, E.; Godard, M.; Lambert, M.: Familial lipoprotein lipase deficiency in infancy: clinical, biochemical, and molecular study. *J. Pediatr.* 133: 417–423, 1998.
- [79518] 17848. Fisher, R. M.; Humphries, S. E.; Talmud, P. J.: Common variation in the lipoprotein lipase gene: effects on plasma lipids and risk of atherosclerosis. *Atherosclerosis* 135: 145–159, 1997.
- [79519] 17849. Franklin, S. M.: Splenomegaly with lipaemia. *Proc. Roy. Soc. Med.* 30: 711, 1937.
- [79520] 17850. Funke, H.; Klug, J.; Assmann, G.: Hind III RFLP in the lipoprotein lipase gene, (LPL). *Nucleic Acids Res.* 15: 9102, 1987.
- [79521] 17851. Funke, H.; Reckwerth, A.; Stapenhorst, D.; Schulze Beiering, M.; Jansen, M.; Assmann, G.: Bst NI (Eco RII) RFLP in the lipoprotein lipase gene (LPL). *Nucleic Acids Res.* 16:

2741, 1988.

- [79522] 17852. Gagne, C.; Brun, L. D.; Julien, P.; Moorjani, S.; Lupien, P. J.: Primary lipoprotein lipase activity deficiency: clinical investigation of a French Canadian population. *Canad. Med. Assoc. J.* 140: 405–411, 1989.
- [79523] 17853. Gilbert, B.; Rouis, M.; Griglio, S.; de Lumley, L.; Laplaud, P.–M.: Lipoprotein lipase (LPL) deficiency: a new patient homozygote for the preponderant mutation gly88-to-glu in the human LPL gene and review of reported mutations: 75% are clustered in exons 5 and 6. *Ann. Genet.* 44:25–32, 2001.
- [79524] 17854. Ginzinger, D. G.; Lewis, M. E. S.; Ma, Y.; Jones, B. R.; Liu, G.; Jones, S. D.; Hayden, M. R.: A mutation in the lipoprotein lipase gene is the molecular basis of chylomicronemia in a colony of domestic cats. *J. Clin. Invest.* 97: 1257–1266, 1996.
- [79525] 17855. Gotoda, T.; Yamada, N.; Kawamura, M.; Kozaki, K.; Mori, N.; Ishibashi, S.; Shimano, H.; Takaku, F.; Yazaki, Y.; Furuichi, Y.; Murase, T.: Heterogeneous mutations in the human lipoprotein lipase gene in patients with familial lipoprotein lipase deficiency. *J. Clin. Invest.* 88:1856–1864, 1991.
- [79526] 17856. Gotoda, T.; Yamada, N.; Murase, T.; Miyake, S.;

Murakami, R.; Kawamura, M.; Kozaki, K.; Mori, N.; Shimano, H.; Shimada, M.; Yazaki, Y.: A newly identified null allelic mutation in the human lipoprotein lipase (LPL) gene of a compound heterozygote with familial LPL deficiency. *Biochim. Biophys. Acta* 1138: 353–356, 1992.

- [79527] 17857. Hata, A.; Emi, M.; Luc, G.; Basdevant, A.; Gambert, P.; Iverius, P.-H.; Lalouel, J.-M.: Compound heterozygote for lipoprotein lipase deficiency: ser-to-thr(244) and transition in 3-prime splice site of intron 2 (AG-to-AA) in the lipoprotein lipase gene. *Am. J. Hum. Genet.* 47: 721–726, 1990.
- [79528] 17858. Hata, A.; Ridinger, D. N.; Sutherland, S. D.; Emi, M.; Kwong, L. K.; Shuhua, J.; Lubbers, A.; Guy-Grand, B.; Basdevant, A.; Iverius, P. H.; Wilson, D. E.; Lalouel, J.-M.: Missense mutations in exon 5 of the human lipoprotein lipase gene: inactivation correlates with loss of dimerization. *J. Biol. Chem.* 267: 20132–20139, 1992.
- [79529] 17859. Haubenwallner, S.; Horl, G.; Shachter, N. S.; Presta, E.; Fried, S. K.; Hofler, G.; Kostner, G. M.; Breslow, J. L.; Zechner, R.: A novel missense mutation in the gene for lipoprotein lipase resulting in a highly conservative amino acid substitution (asp180-to-glu) causes familial chylomicronemia (type I hyperlipoproteinemia). *Genomics*



18:392–396, 1993.

- [79530] 17860.Havel, R. J.; Gordon, R. S.: Idiopathic hyperlipemia: metabolic studies in an affected family. *J. Clin. Invest.* 39: 1777–1790, 1960.
- [79531] 17861.Heaney, A. P.; Sharer, N.; Rameh, B.; Braganza, J. M.; Durrington, P. N.: Prevention of recurrent pancreatitis in familial lipoprotein lipase deficiency with high-dose antioxidant therapy. *J. Clin. Endocr. Metab.* 84: 1203–1205, 1999.
- [79532] 17862.Heinzmann, C.; Ladas, J.; Antonarakis, S.; Kirchgessner, T.; Schotz, M.; Lusis, A. J.: RFLP for the human lipoprotein lipase (LPL) gene: HindIII. *Nucleic Acids Res.* 15: 6763, 1987.
- [79533] 17863.Heizmann, C.; Kirchgessner, T.; Kwiterovich, P. O.; Ladas, J.A.; Derby, C.; Antonarakis, S. E.; Lusis, A. J.: DNA polymorphism haplotypes of the human lipoprotein lipase gene: possible association with high density lipoprotein levels. *Hum. Genet.* 86: 578–584, 1991.
- [79534] 17864.Henderson, H.; Ma, Y.; Kastelein, J.; Roederer, G.; Julien, P.; Brunzell, J.; Hayden, M. R.: Identification of the molecular defects underlying chylomicronemia in the majority of 75 separate probands with LPL deficiency. (Abstract) *Clin. Res.* 39: 336A, 1991.

- [79535] 17865.Henderson, H. E.; Bijvoet, S. M.; Mannens, M. A. M. M.; Bruin,T.; Erkelens, D. W.; Hayden, M. R.; Kastelein, J. J. P.: Ile225-to-thrloop mutation in the lipoprotein lipase (LPL) gene is a de novo event. *Am.J. Med. Genet.* 78: 313–316, 1998.
- [79536] 17866.Sjalander, A.; Birgander, R.; Rannug, A.; Alexandrie, A.-K.; Tornling,G.; Beckman, G.: Association between the p21 codon 31A1 (arg) alleleand lung cancer. *Hum. Hered.* 46: 221–225, 1996.
- [79537] 17867.Fryns, J. P.; van den Berghe, K.; van Assche, A.; van den Berghe,H.: Prenatal diagnosis of campomelic dwarfism. *Clin. Genet.* 19:199–201, 1981.
- [79538] 17868.Mellows, H. J.; Pryse-Davies, J.; Bennett, M. J.; Carter, C. O.: The camptomelic syndrome in two female siblings. *Clin. Genet.* 18:137–141, 1980.
- [79539] 17869.Millauer, B.; Shawver, L. K.; Plate, K. H.; Risau, W.; Ullrich,A.: Glioblastoma growth inhibited in vivo by a dominant-negativeFlk-1 mutant. *Nature* 367: 576–579, 1994.
- [79540] 17870.Cho, K. R.; Oliner, J. D.; Simons, J. W.; Hedrick, L.; Fearon,E. R.; Preisinger, A. C.; Hedge, P.; Silverman, G. A.; Vogelstein,B.: The DCC gene: structural analysis and mutations in colorectalcarcinomas. *Genomics* 19: 525–531,

1994.

- [79541] 17871.Fazeli, A.; Dickinson, S. L.; Hermiston, M. L.; Tighe, R. V.; Steen, R. G.; Small, C. G.; Stoeckli, E. T.; Keino-Masu, K.; Masu, M.; Rayburn, H.; Simons, J.; Bronson, R. T.; Gordon, J. I.; Tessier-Lavigne, M.; Weinberg, R. A.: Phenotype of mice lacking functional Deleted in colorectal cancer (Dcc) gene. *Nature* 386: 796–804, 1997.
- [79542] 17872.Fearon, E. R.; Cho, K. R.; Nigro, J. M.; Kern, S. E.; Simons, J.W.; Ruppert, J. M.; Hamilton, S. R.; Preisinger, A. C.; Thomas, G.; Kinzler, K. W.; Vogelstein, B.: Identification of a chromosome 18q gene that is altered in colorectal cancers. *Science* 247: 49–56, 1990.
- [79543] 17873.Gotley, D. C.; Reeder, J. A.; Fawcett, J.; Walsh, M. D.; Bates, P.; Simmons, D. L.; Antalis, T. M.: The deleted in colon cancer (DCC) gene is consistently expressed in colorectal cancers and metastases. *Oncogene* 13:787–795, 1996.
- [79544] 17874.Hohne, M. W.; Halatsch, M.-E.; Kahl, G. F.; Weinel, R. J.: Frequent loss of expression of the potential tumor suppressor gene DCC in ductal pancreatic adenocarcinoma. *Cancer Res.* 52: 2616–2619, 1992.
- [79545] 17875.Jen, J.; Kim, H.; Piantadosi, S.; Liu, Z.-F.; Levitt, R. C.; Sistonen, P.; Kinzler, K. W.; Vogelstein, B.; Hamilton, S.

R.: Allelic loss of chromosome 18q and prognosis in colorectal cancer. *New Eng. J. Med.* 331: 213–221, 1994.

- [79546] 17876. Keino-Masu, K.; Masu, M.; Hinck, L.; Leonardo, E. D.; Chan, S.S.-Y.; Culotti, J. G.; Tessier-Lavigne, M.: Deleted in colorectal cancer (DCC) encodes a netrin receptor. *Cell* 87: 175–185, 1996.
- [79547] 17877. Keynes, R.; Cook, G. M. W.: Axon guidance molecules. *Cell* 83: 161–169, 1995.
- [79548] 17878. Kolodziej, P. A.; Timpe, L. C.; Mitchell, K. J.; Fried, S. R.; Goodman, C. S.; Jan, L. Y.; Jan, Y. N.: Frazzled encodes a Drosophila member of the DCC immunoglobulin subfamily and is required for CNS and motor axon guidance. *Cell* 87: 197–204, 1996.
- [79549] 17879. Maesawa, C.; Tamura, G.; Ogasawara, S.; Suzuki, Y.; Sakata, K.; Sugimura, J.; Nishizuka, S.; Sato, N.; Ishida, K.; Saito, K.; Satodate, R.: Loss of heterozygosity at the DCC gene locus is not crucial for the acquisition of metastatic potential in oesophageal squamous cell carcinoma. *Europ. J. Cancer* 32A: 896–898, 1996.
- [79550] 17880. Mehlen, P.; Rabizadeh, S.; Snipas, S. J.; Assa-Munt, N.; Salvesen, G. S.; Bredesen, D. E.: The DCC gene product induces apoptosis by a mechanism requiring receptor proteolysis. *Nature* 395: 801–804, 1998.

- [79551] 17881.Miyake, S.; Nagai, K.; Yoshino, K.; Oto, M.; Endo, M.; Yuasa,Y.: Point mutations and allelic deletion of tumor suppressor geneDCC in human esophageal squamous cell carcinomas and their relationto metastasis. Cancer Res. 54: 3007–3010, 1994.
- [79552] 17882.Nigro, J. M.; Cho, K. R.; Fearon, E. R.; Kern, S. E.; Ruppert,J. M.; Oliner, J. D.; Kinzler, K. W.; Vogelstein, B.: Scrambled exons. Cell 64:607–613, 1991.
- [79553] 17883.Shibata, D.; Reale, M. A.; Lavin, P.; Silverman, M.; Fearon, E.R.; Steele, G., Jr.; Jessup, J. M.; Loda, M.; Summerhayes, I. C.:The DCC protein and prognosis in colorectal cancer. New Eng. J. Med. 335:1727–1732, 1996.
- [79554] 17884.Stein, E.; Tessier–Lavigne, M.: Hierarchical organization ofguidance receptors: silencing of netrin attraction by Slit througha Robo/DCC receptor complex. Science 291: 1928–1938, 2001.
- [79555] 17885.Tanaka, K.; Oshimura, M.; Kikuchi, R.; Seki, M.; Hayashi, T.;Miyaki, M.: Suppression of tumorigenicity in human colon carcinomacells by introduction of normal chromosome 5 or 18. Nature 349:340–342, 1991.
- [79556] 17886.Uchino, S.; Tsuda, H.; Noguchi, M.; Yokota, J.; Terada, M.; Saito,T.; Kobayashi, M.; Sugimura, T.; Hirohashi, S.: Frequent loss ofheterozygosity at the DCC locus in

gastric cancer. *Cancer Res.* 52:3099–3102, 1992.

[79557] 17887.Vogelstein, B.: Personal Communication. Baltimore, Md. 11/30/1995.

[79558] 17888.Zetter, B. R.: Adhesion molecules in tumor metastasis. *Semin.Cancer Biol.* 4: 219–229, 1993.

[79559] 17889.Devor, E. J.; Grandy, D. K.; Civelli, O.; Litt, M.; Burgess, A.K.; Isenberg, K. E.; van de Wetering, B. J. M.; Oostra, B.: Genetic linkage is excluded for the D2-dopamine receptor lambda-HD2G1 and flanking loci on chromosome 11q22–q23 in Tourette syndrome. *Hum.Hered.* 40: 105–108, 1990.

[79560] 17890.Freemantle, S. J.; Taylor, S. M.; Krystal, G.; Moran, R. G.: Upstream organization of and multiple transcripts from the human folylpoly-gamma-glutamatesynthetase gene. *J. Biol. Chem.* 270: 9579–9584, 1995.

[79561] 17891.Garrow, T. A.; Admon, A.; Shane, B.: Expression cloning of a human cDNA encoding folylpoly(gamma-glutamate) synthetase and determination of its primary structure. *Proc. Nat. Acad. Sci.* 89: 9151–9155, 1992.

[79562] 17892.Jones, C.; Kao, F.-T.: Regional mapping of the folylpolyglutamatesynthetase gene (FPGS) to 9cen–q34. (Abstract) *Cytogenet. Cell Genet.* 37:499–500, 1984.

- [79563] 17893.Jones, C.; Kao, F.-T.; Taylor, R. T.: Chromosomal assignment of the gene for folylpolyglutamate synthetase to human chromosome 9. *Cytogenet.Cell Genet.* 28: 181-194, 1980.
- [79564] 17894.Jones, C.; Kao, F. T.: Assignment of the human gene complementing the auxotrophic marker GAT-minus to chromosome 9. (Abstract) *Cytogenet.Cell Genet.* 25: 168, 1979.
- [79565] 17895.Kao, F. T.; Puck, T. T.: Genetics of somatic mammalian cells.VII. Induction and isolation of nutritional mutants in Chinese hamster cells. *Proc. Nat. Acad. Sci.* 60: 1275-1281, 1968.
- [79566] 17896.Sussman, D. J.; Milman, G.; Shane, B.: Characterization of human folylpolyglutamate synthetase expressed in Chinese hamster ovary cells. *Somat.Cell Molec. Genet.* 12: 531-540, 1986.
- [79567] 17897.Taylor, R. T.; Hanna, M. L.: Folate-dependent enzymes in cultured Chinese hamster cells: folylpolyglutamate synthetase and its absence in mutants auxotrophic for glycine, adenosine and thymidine. *Arch.Biochem. Biophys.* 181: 331-344, 1977.
- [79568] 17898.Taylor, S. M.; Freemantle, S. J.; Moran, R. G.: Structural organization of the human folylpoly-

gamma-glutamate synthetase gene: evidence for a single genomic locus. *Cancer Res.* 55: 6030–6034, 1995.

- [79569] 17899. Matsuo, K.; Owens, J. M.; Tonko, M.; Elliott, C.; Chambers, T.J.; Wagner, E. F.: Fos1 is a transcriptional target of c-Fos during osteoclast differentiation. *Nature Genet.* 24: 184–187, 2000.
- [79570] 17900. Buters, J. T. M.; Tang, B.-K.; Pineau, T.; Gelboin, H. V.; Kimura, S.; Gonzalez, F. J.: Role of CYP1A2 in caffeine pharmacokinetics and metabolism: studies using mice deficient in CYP1A2. *Pharmacogenetics* 6:291–296, 1996.
- [79571] 17901. Butler, M. A.; Iwasaki, M.; Guengerich, F. P.; Kadlubar, F. F.: Human cytochrome P-450(PA) (P-450IA2), the phenacetin O-deethylase, is primarily responsible for the hepatic 3-demethylation of caffeine and N-oxidation of carcinogenic arylamines. *Proc. Nat. Acad. Sci.* 86:7696–7700, 1989.
- [79572] 17902. Christiansen, L.; Bygum, A.; Jensen, A.; Thomsen, K.; Brandrup, F.; Horder, M.; Petersen, N. E.: Association between CYP1A2 polymorphism and susceptibility to porphyria cutanea tarda. *Hum. Genet.* 107:612–614, 2000.
- [79573] 17903. Devonshire, H. W.; Kong, I.; Cooper, M.; Sloan, T. P.; Idle, J.R.; Smith, R. L.: The contribution of genetically determined oxidation status to inter-individual variation in



phenacetin disposition. *Brit.J. Clin. Pharm.* 16: 157–166, 1983.

- [79574] 17904.Ikeya, K.; Jaiswal, A. K.; Owens, R. A.; Jones, J. E.; Nebert,D. W.; Kimura, S.: Human CYP1A2: sequence, gene structure, comparisonwith the mouse and rat orthologous gene, and differences in liver1A2 mRNA expression. *Molec. Endocr.* 3: 1399–1408, 1989.
- [79575] 17905.Jaiswal, A. K.; Nebert, D. W.; McBride, O. W.; Gonzalez, F. J.: Human P(3)450: cDNA and complete protein sequence, repetitive Alusequences in the 3–prime non–translated region, and localization ofgene to chromosome 15. *J. Exp. Path.* 3: 1–17, 1987.
- [79576] 17906.Liang, H.–C. L.; Li, H.; McKinnon, R. A.; Duffy, J. J.; Potter,S. S.; Puga, A.; Nebert, D. W.: Cyp1a2(–/–) null mutant mice developnormally but show deficient drug metabolism. *Proc. Nat. Acad. Sci.* 93:1671–1676, 1996.
- [79577] 17907.Nebert, D. W.: Personal Communication. Bethesda, Md. 2/3/1988.
- [79578] 17908.Sesardic, D.; Boobis, A. R.; Edwards, R. J.; Davies, D. S.: Aform of cytochrome P450 in man, orthologous to form d in the rat,catalyses the O–deethylation of phenacetin and is inducible by cigarettesmoking. *Brit. J. Clin. Pharm.* 26: 363–372, 1988.

- [79579] 17909.Tantcheva-Poor, I.; Zaigler, M.; Rietbrock, S.; Fuhr, U.: Estimation of cytochrome P-450 CYP1A2 activity in 863 healthy Caucasians using a saliva-based caffeine test. *Pharmacogenetics* 9: 131-144, 1999.
- [79580] 17910.Wooding, S. P.; Watkins, W. S.; Bamshad, M. J.; Dunn, D. M.; Weiss, R. B.; Jorde, L. B.: DNA sequence variation in a 3.7-kb noncoding sequence 5-prime of the CYP1A2 gene: implications for human population history and natural selection. *Am. J. Hum. Genet.* 71: 528-542, 2002.
- [79581] 17911.Bale, A. E.; Mitchell, A. L.; Gonzalez, F. J.; McBride, O. W.: Localization of CYP2F1 by multipoint linkage analysis and pulsed-field gel electrophoresis. *Genomics* 10: 284-286, 1991.
- [79582] 17912.Mitchell, A. L.; Bale, A. E.; Gonzalez, F.; McBride, O. W.: Mapping of cytochrome p450 IIA, IIB, and IIF subfamilies on chromosome 19 by linkage analysis. (Abstract) *Cytogenet. Cell Genet.* 51: 1045, 1989.
- [79583] 17913.Nhamburo, P. T.; Kimura, S.; McBride, O. W.; Kozak, C. A.; Gelboin, H. V.; Gonzalez, F. J.: The human CYP2F gene subfamily: identification of a cDNA encoding a new cytochrome P450, cDNA-directed expression, and chromosome mapping. *Biochemistry* 29: 5491-5499,

1990.

- [79584] 17914.Imaoka, S.; Yoneda, Y.; Sugimoto, T.; Hiroi, T.; Yamamoto, K.;Nakatani, T.; Funae, Y.: CYP4B1 is a possible risk factor for bladder cancer in humans. *Biochem. Biophys. Res. Commun.* 277: 776–780, 2000.
- [79585] 17915.Nhamburo, P. T.; Gonzalez, F. J.; McBride, O. W.; Gelboin, H. V.;Kimura, S.: Identification of a new P450 expressed in human lung:complete cDNA sequence, cDNA-directed expression, and chromosome mapping. *Biochemistry* 28:8060–8066, 1989.
- [79586] 17916.Thum, T.; Borlak, J.: Gene expression in distinct regions of the heart. *Lancet* 355: 979–983, 2000.
- [79587] 17917.Xu, L.; Xia, J.; Jiang, H.; Zhou, J.; Li, H.; Wang, D.; Pan, Q.;Long, Z.; Fan, C.; Deng, H.–X.: Mutation analysis of hereditary multiple exostoses in the Chinese. *Hum. Genet.* 105: 45–50, 1999.
- [79588] 17918.Joutel, A.; Favrole, P.; Labauge, P.; Chabriat, H.; Lescoat, C.;Andreux, F.; Domenga, V.; Cecillon, M.; Vahedi, K.; Ducros, A.; Cave-Riant, F.; Bousser, M. G.; Tournier-Lasserre, E.: Skin biopsy immunostaining with a Notch3 monoclonal antibody for CADASIL diagnosis. *Lancet* 358:2049–2051, 2001.
- [79589] 17919.Barinaga, M.: An intriguing new lead on Hunting–

ton's disease. *Science* 271:1233–1234, 1996.

[79590] 17920. Burke, J. R.; Enghild, J. J.; Martin, M. E.; Jou, Y.-S.; Myers, R. M.; Roses, A. D.; Vance, J. M.; Strittmatter, W. J.: Huntingtin and DRPLA proteins selectively interact with the enzyme GAPDH. *Nature Med.* 2: 347–350, 1996.

[79591] 17921. Eubanks, J. H.; Djabali, M.; Selleri, L.; Grandy, D. K.; Civelli, O.; McElligott, D. L.; Evans, G. A.: Structure and linkage of the D2 dopamine receptor and neural cell adhesion molecule genes on human chromosome 11q23. *Genomics* 14: 1010–1018, 1992.

[79592] 17922. Gejman, P. V.; Ram, A.; Gelernter, J.; Friedman, E.; Cao, Q.; Pickar, D.; Blum, K.; Noble, E. P.; Kranzler, H. R.; O'Malley, S.; Hamer, D. H.; Whitsitt, F.; Rao, P.; DeLisi, L. E.; Virkkunen, M.; Linnoila, M.; Goldman, D.; Gershon, E. S.: No structural mutation in the dopamine D2 receptor gene in alcoholism or schizophrenia: analysis using denaturing gradient gel electrophoresis. *J.A.M.A.* 271: 204–208, 1994.

[79593] 17923. Gelernter, J.; Grandy, D. K.; Bunzow, J.; Pakstis, A. J.; Civelli, O.; Retief, A. E.; Litt, M.; Kidd, K. K.: D(2) dopamine receptor locus (probe hD2G1) maps close to D11S29 (probe L7) and is also linked to PBGD (probe PBGD0.0) and D11S84 (probe p-2-7-ID6) on 11q. (Abstract) *Cytogenet. Cell Genet.* 51: 1002 only, 1989.

- [79594] 17924.Gelernter, J.; Pakstis, A. J.; Grandy, D.; Litt, M.; Retief, A.E.; Kennedy, J. L.; Hing-Loh, A.; Schoolfield, G.; Civelli, O.; Kidd, K. K.: Linkage map of eight human chromosome 11q markers, including DRD2, spanning 60 cM. *Cytogenet. Cell Genet.* 60: 26–28, 1992.
- [79595] 17925.Gelernter, J.; Pakstis, A. J.; Pauls, D. L.; Kurlan, R.; Gancher, S. T.; Civelli, O.; Grandy, D.; Kidd, K. K.: Gilles de la Tourette syndrome is not linked to D2-dopamine receptor. *Arch. Gen. Psychiat.* 47:1073–1077, 1990.
- [79596] 17926.Giros, B.; Sokoloff, P.; Martres, M.-P.; Riou, J.-F.; Emorine, L. J.; Schwartz, J.-C.: Alternative splicing directs the expression of two D(2) dopamine receptor isoforms. *Nature* 342: 923–926, 1989.
- [79597] 17927.Grandy, D. K.; Litt, M.; Allen, L.; Bunzow, J. R.; Marchionni, M.; Makam, H.; Reed, L.; Magenis, R. E.; Civelli, O.: The human dopamine D(2) receptor gene is located on chromosome 11 at q22–q23 and identifies a TaqI RFLP. *Am. J. Hum. Genet.* 45: 778–785, 1989.
- [79598] 17928.Grandy, D. K.; Marchionni, M. A.; Makam, H.; Stofko, R. E.; Alfano, M.; Frothingham, L.; Fischer, J. B.; Burke-Howie, K. J.; Bunzow, J.R.; Server, A. C.; Civelli, O.: Cloning of the cDNA and gene for a human D2 dopamine receptor. *Proc. Nat. Acad. Sci.* 86: 9762–9766, 1989.

- [79599] 17929.Klein, C.; Brin, M. F.; Kramer, P.; Sena-Esteves, M.; de Leon,D.; Doheny, D.; Bressman, S.; Fahn, S.; Breakefield, X. O.; Ozelius,L. J.: Association of a missense change in the D2 dopamine receptorwith myoclonus dystonia. Proc. Nat. Acad. Sci. 96: 5173–5176, 1999.
- [79600] 17930.Maldonado, R.; Salardi, A.; Valverde, O.; Samad, T. A.; Roques,B. P.; Borrelli, E.: Absence of opiate rewarding effects in micelacking dopamine D2 receptors. Nature 388: 586–589, 1997.
- [79601] 17931.Milligan, G.: Receptors as kissing cousins. Science 288: 65–67,2000.
- [79602] 17932.Monsma, F. J., Jr.; McVittie, L. D.; Gerfen, C. R.; Mahan, L.C.; Sibley, D. R.: Multiple D(2) dopamine receptors produced by alternativeRNA splicing. Nature 342: 926–929, 1989.
- [79603] 17933.Oliveri, R. L.; Annesi, G.; Zappia, M.; Civitelli, D.; Montesanti,R.; Branca, D.; Nicoletti, G.; Spadafora, P.; Pasqua, A. A.; Cittadella,R.; Andreoli, V.; Gambardella, A.; Aguglia, U.; Quattrone, A.: DopamineD2 receptor gene polymorphism and the risk of levodopa-induced dyskinesiasin PD. Neurology 53: 1425–1430, 1999.
- [79604] 17934.Rocheville, M.; Lange, D. C.; Kumar, U.; Patel, S. C.; Patel,R. C.; Patel, Y. C.: Receptors for dopamine and so–

matostatin: formation of hetero-oligomers with enhanced functional activity. *Science* 288:154–157, 2000.

- [79605] 17935. Smith, M.; Wasmuth, J.; McPherson, J. D.; Wagner, C.; Grandy, D.; Civelli, O.; Potkin, S.; Litt, M.: Cosegregation of an 11q22.3–9p22 translocation with affective disorder: proximity of the dopamine D2 receptor gene relative to the translocation breakpoint. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A220 only, 1989.
- [79606] 17936. St Clair, D.; Blackwood, D.; Muir, W.; Carothers, A.; Walker, M.; Spowart, G.; Gosden, C.; Evans, H. J.: Association within a family of a balanced autosomal translocation with major mental illness. *Lancet* 336:13–16, 1990.
- [79607] 17937. Suarez, B. K.; Parsian, A.; Hampe, C. L.; Todd, R. D.; Reich, T.; Cloninger, C. R.: Linkage disequilibria at the D2 dopamine receptor locus (DRD2) in alcoholics and controls. *Genomics* 19: 12–20, 1994.
- [79608] 17938. Usiello, A.; Baik, J.-H.; Rouge-Pont, F.; Picetti, R.; Dierich, A.; LeMeur, M.; Piazza, P. V.; Borrelli, E.: Distinct functions of the two isoforms of dopamine D2 receptors. *Nature* 408: 199–203, 2000.
- [79609] 17939. Wang, J.; Liu, Z.-L.; Chen, B.: Association study of dopamine D2, D3 receptor gene polymorphisms with motor fluctuations in PD. *Neurology* 56:1757–1759, 2001.

- [79610] 17940. Accili, D.; Fishburn, C. S.; Drago, J.; Steiner, H.; Lachowicz, J. E.; Park, B.-H.; Gauda, E. B.; Lee, E. J.; Cool, M. H.; Sibley, D. R.; Gerfen, C. R.; Westphal, H.; Fuchs, S.: A targeted mutation of the D3 dopamine receptor gene is associated with hyperactivity in mice. *Proc. Nat. Acad. Sci.* 93: 1945–1949, 1996.
- [79611] 17941. Asico, L. D.; Ladines, C.; Fuchs, S.; Accili, D.; Carey, R. M.; Semeraro, C.; Pocchiari, F.; Felder, R. A.; Eisner, G. M.; Jose, P. A.: Disruption of the dopamine D3 receptor gene produces renin-dependent hypertension. *J. Clin. Invest.* 102: 493–498, 1998.
- [79612] 17942. Crocq, M.-A.; Mant, R.; Asherson, P.; Williams, J.; Hode, Y.; Mayerova, A.; Collier, D.; Lannfelt, L.; Sokoloff, P.; Schwartz, J.-C.; Gill, M.; Macher, J.-P.; McGuffin, P.; Owen, M. J.: Association between schizophrenia and homozygosity at the dopamine D3 receptor gene. *J. Med. Genet.* 29: 858–860, 1992.
- [79613] 17943. Guillin, O.; Diaz, J.; Carroll, P.; Griffon, N.; Schwartz, J.-C.; Sokoloff, P.: BDNF controls dopamine D3 receptor expression and triggers behavioural sensitization. *Nature* 411: 86–89, 2001.
- [79614] 17944. Ilani, T.; Ben-Shachar, D.; Strous, R. D.; Mazor, M.; Sheinkman, A.; Kotler, M.; Fuchs, S.: A peripheral marker



for schizophrenia:increased levels of D3 dopamine receptor mRNA in blood lymphocytes. Proc.Nat. Acad. Sci. 98: 625–628, 2001.

- [79615] 17945.Le Coniat, M.; Sokoloff, P.; Hillion, J.; Martres, M.-P.; Giros,B.; Pilon, C.; Schwartz, J.-C.; Berger, R.: Chromosomal localizationof the human D-3 dopamine receptor gene. Hum. Genet. 87: 618–620,1991.
- [79616] 17946.Nanko, S.; Sasaki, T.; Fukuda, R.; Hattori, M.; Dai, X. Y.; Kazamatsuri,H.; Kuwata, S.; Juji, T.; Gill, M.: A study of the association between schizophrenia and the dopamine D3 receptor gene. Hum. Genet. 92:336–338, 1993.
- [79617] 17947.Nothen, M. M.; Cichon, S.; Propping, P.; Fimmers, R.; Schwab, S.G.; Wildenauer, D. B.: Excess of homozygosity at the dopamine D3receptor gene in schizophrenia not confirmed. (Letter) J. Med. Genet. 30:708–712, 1993.
- [79618] 17948.Bevilacqua, M.; Butcher, E.; Furie, B.; Furie, B.; Galatin, M.;Gimbrone, M.; Harlan, J.; Kishimoto, K.; Lasky, L.; McEver, R.; Paulson,J.; Rosen, S.; Seed, B.; Siegelman, M.; Springer, T.; Stoolman, L.;Tedder, T.; Varki, A.; Wagner, D.; Weissman, I.; Zimmerman, G.: Selectins:a family of adhesion receptors. (Letter) Cell 67: 233 only, 1991.
- [79619] 17949.Bevilacqua, M. P.; Stengelin, S.; Gimbrone, M. A.,

Jr.; Seed, B.: Endothelial leukocyte adhesion molecule 1: an inducible receptor for neutrophils related to complement regulatory proteins and lectins. *Science* 243:1160–1165, 1989.

[79620] 17950. Collins, T.; Williams, A.; Johnston, G. I.; Kim, J.; Eddy, R.; Shows, T.; Gimbrone, M. A., Jr.; Bevilacqua, M. P.: Structure and chromosomal location of the gene for endothelial-leukocyte adhesion molecule 1. *J. Biol. Chem.* 266: 2466–2473, 1991.

[79621] 17951. DeLisser, H. M.; Christofidou-Solomidou, M.; Sun, J.; Nakada, M. T.; Sullivan, K. E.: Loss of endothelial surface expression of E-selectin in a patient with recurrent infections. *Blood* 94: 884–894, 1999.

[79622] 17952. Wang, N.; Chintala, S. K.; Fini, M. E.; Schuman, J. S.: Activation of a tissue-specific stress response in the aqueous outflow pathway of the eye defines the glaucoma disease phenotype. *Nature Med.* 7:304–309, 2001.

[79623] 17953. Watson, M. L.; Kingsmore, S. F.; Johnston, G. I.; Siegelman, M. H.; Le Beau, M. M.; Lemons, R. S.; Bora, N. S.; Howard, T. A.; Weissman, I. L.; McEver, R. P.; Seldin, M. F.: Genomic organization of the selectin family of leukocyte adhesion molecules on human and mouse chromosome 1. *J. Exp. Med.* 172: 263–272, 1990.

- [79624] 17954.Wenzel, K.; Felix, S.; Kleber, F. X.; Brachold, R.; Menke, T.;Schattke, S.; Schulte, K. L.; Glaser, C.; Rohde, K.; Baumann, G.;Speer, A.: E-selectin polymorphism and atherosclerosis: an associationstudy. Hum. Molec. Genet. 3: 1935–1937, 1994.
- [79625] 17955.Zheng, F.; Chevalier, J. A.; Zhang, L. Q.; Virgil, D.; Ye, S. Q.;Kwiterovich, P. O.: An HphI polymorphism in the E-selectin gene isassociated with premature coronary artery disease. Clin. Genet. 59:58–64, 2001.
- [79626] 17956.Burgess, W. H.; Mehlman, T.; Marshak, D. R.; Fraser, B. A.; Maciag,T.: Structural evidence that endothelial cell growth factor betais the precursor of both endothelial cell growth factor alpha andacidic fibroblast growth factor. Proc. Nat. Acad. Sci. 83: 7216–7220,1986.
- [79627] 17957.Eckenstein, F. P.: Fibroblast growth factors in the nervous system. J.Neurobiol. 25: 1467–1480, 1994.
- [79628] 17958.Gautschi–Sova, P.; Muller, T.; Bohlen, P.: Amino acid sequenceof human acidic fibroblast growth factor. Biochem. Biophys. Res.Communic. 140: 874–880, 1986.
- [79629] 17959.Jaye, M.; Howk, R.; Burgess, W.; Ricca, G. A.; Chiu, I.–M.; Ravera,M. W.; O'Brien, S. J.; Modi, W. S.; Maciag, T.; Drohan, W. N.: Humanendothelial cell growth factor: cloning, nucleotide sequence, andchromosome localiza–

tion. Science 233: 541–545, 1986.

- [79630] 17960.Jung, J.; Zheng, M.; Goldfarb, M.; Zaret, K. S.: Initiation of mammalian liver development from endoderm by fibroblast growth factors. Science 284:1998–2003, 1999.
- [79631] 17961.Mergia, A.; Eddy, R.; Abraham, J. A.; Fiddes, J. C.; Shows, T.B.: The genes for basic and acidic fibroblast growth factors are on different human chromosomes. Biochem. Biophys. Res. Commun. 138:644–651, 1986.
- [79632] 17962.Pellegrini, L.; Burke, D. F.; von Delft, F.; Mulloy, B.; Blundell, T. L.: Crystal structure of fibroblast growth factor receptor ectodomain bound to ligand and heparin. Nature 407: 1029–1034, 2000.
- [79633] 17963.Plotnikov, A. N.; Hubbard, S. R.; Schlessinger, J.; Mohammadi, M.: Crystal structures of two FGF–FGFR complexes reveal the determinants of ligand–receptor specificity. Cell 101: 413–424, 2000.
- [79634] 17964.Wang, W.–P.; Lehtonen, K.; Varban, M. L.; Krishnan, I.; Chiu, I.–M.: Cloning of the gene coding for human class 1 heparin-binding growth factor and its expression in fetal tissues. Molec. Cell. Biol. 9:2387–2395, 1989.
- [79635] 17965.Gamez, J.; Ferreira, C.; Accarino, M. L.; Guarner, L.; Tadesse, S.; Marti, R. A.; Andreu, A. L.; Raguer, N.; Cervera, C.; Hirano, M.: Phenotypic variability in a Spanish family

with MNGIE. *Neurology* 59:455–457, 2002.

- [79636] 17966.Hagiwara, K.; Stenman, G.; Honda, H.; Sahlin, P.; Andersson, A.;Miyazono, K.; Heldin, C. H.; Ishikawa, F.; Takaku, F.: Organizationand chromosomal localization of the human platelet–derived endothelialcell growth factor gene. *Molec. Cell. Biol.* 11: 2125–2132, 1991.
- [79637] 17967.Nishino, I.; Spinazzola, A.; Hirano, M.: Thymidine phosphorylasegene mutations in MNGIE, a human mitochondrial disorder. *Science* 283:689–692, 1999.
- [79638] 17968.Stenman, G.; Sahlin, P.; Dumanski, J. P.; Hagiwara, K.; Ishikawa,F.; Miyazono, K.; Collins, V. P.; Heldin, C.–H.: Regional localizationof the human platelet–derived endothelial cell growth factor (ECGF1)gene to chromosome 22q13. *Cytogenet. Cell Genet.* 59: 22–23, 1992.
- [79639] 17969.Stenman, G.; Sahlin, P.; Hagiwara, K.; Dumanski, J.; Collins, V.;Heldin, C.–H.: Mapping of the human platelet–derived endothelialcell growth factor (PD–ECGF) gene to chromosome 22q13. (Abstract) *Cytogenet.Cell Genet.* 58: 2051 only, 1991.
- [79640] 17970.Kayes–Wandover, K.; White, P. C.: Steroidogenic enzyme gene expressionin the human heart. *J. Clin. Endocr. Metab.* 85: 2519–2525, 2000.
- [79641] 17971.Sinke, R. J.; Tanigami, A.; Nakamura, Y.; Geurts van

Kessel, A.: Reverse mapping of the gene encoding the human fos-related antigen-1(fra-1) within chromosome band 11q13. Genomics 18: 165 only, 1993.

[79642] 17972.Cox, D. R.; Epstein, L. B.; Epstein, C. J.: Genes coding for sensitivity to interferon (IfRec) and soluble superoxide dismutase (SOD-1) are linked in mouse and man and map to mouse chromosome 16. Proc. Nat. Acad. Sci. 77: 2168-2172, 1980.

[79643] 17973.De Clercq, E.; Edy, V. G.; Cassiman, J.-J.: Chromosome 21 does not code for an interferon receptor. Nature 264: 249-251, 1976.

[79644] 17974.Epstein, L. B.; Epstein, C. J.: Localization of the gene AVG for the antiviral expression of immune and classical interferon to the distal portion of the long arm of chromosome 21. J. Infect. Dis. 133(suppl.): A56-A62, 1976.

[79645] 17975.Faltynek, C. R.; Branca, A. A.; McCandless, S.; Baglioni, C.: Characterization of an interferon receptor on human lymphoblastoid cells. Proc. Nat. Acad. Sci. 80: 3269-3273, 1983.

[79646] 17976.Fournier, A.; Zhang, Z. Q.; Tan, Y. H.: Human beta:alpha but not gamma interferon binding site is a product of the chromosome 21 interferon action gene. So-

mat. Cell Molec. Genet. 11: 291–295, 1985.

- [79647] 17977.Lutfalla, G.; Gardiner, K.; Proudhon, D.; Vielh, E.; Uze, G.:The structure of the human interferon alpha/beta receptor gene. J.Biol. Chem. 267: 2802–2809, 1992.
- [79648] 17978.Lutfalla, G.; Roeckel, N.; Mogensen, K. E.; Mattei, M. G.; Uze,G.: Assignment of the human interferon–alpha receptor gene to chromosome21q22.1 by in situ hybridization. J. Interferon Res. 10: 515–517,1990.
- [79649] 17979.Maroun, L. E.: Interferon action and chromosome 21 trisomy. (Letter) J.Theor. Biol. 86: 603–606, 1980.
- [79650] 17980.Novick, D.; Cohen, B.; Rubinstein, M.: The human interferon alpha/betareceptor: characterization and molecular cloning. Cell 77: 391–400,1994.
- [79651] 17981.Raziuddin, A.; Sarkar, F. H.; Dutkowski, R.; Shulman, L.; Ruddle,F. H.; Gupta, S. L.: Receptors for human alpha and beta interferonbut not for gamma interferon are specified by human chromosome 21. Proc.Nat. Acad. Sci. 81: 5504–5508, 1984.
- [79652] 17982.Revel, M.; Bash, D.; Ruddle, F. H.: Antibodies to a cell–surfacecomponent coded by human chromosome 21 inhibit action of interferon. Nature 260:139–141, 1976.
- [79653] 17983.Topham, M. K.; Prescott, S. M.: Mammalian diacylglycerol kinases,a family of lipid kinases with signaling

functions. J. Biol. Chem. 274:11447–11450, 1999.

- [79654] 17984.Calabrese, G.; Crescenzi, C.; Morizio, E.; Palka, G.; Guerra, E.;Alberti, S.: Assignment of TACSTD1 (alias TROP1, M4S1) to human chromosome2p21 and refinement of mapping of TACSTD2 (alias TROP2, M1S1) to human chromosome 1p32 by in situ hybridization. Cytogenet. Cell Genet. 92:164–165, 2001.
- [79655] 17985.Fornaro, M.; Dell'Arciprete, R.; Stella, M.; Bucci, C.; Nutini,M.; Capri, M. G.; Alberti, S.: Cloning of the gene encoding Trop-2,a cell-surface glycoprotein expressed by human carcinomas. Int. J.Cancer 62: 610–618, 1995.
- [79656] 17986.Linnenbach, A. J.; Seng, B. A.; Wu, S.; Robbins, S.; Scollon, M.;Pyrce, J. J.; Druck, T.; Huebner, K.: Retroposition in a family of carcinoma-associated antigen genes. Molec. Cell Biol. 13: 1507–1515,1993.
- [79657] 17987.Linnenbach, A. J.; Wojciorowski, J.; Wu, S.; Pyrc, J. J.; Ross,A. H.; Dietzschold, B.; Speicher, D.; Koprowski, H.: Sequence investigationof the major gastrointestinal tumor-associated antigen gene family,GA733. Proc. Nat. Acad. Sci. 86: 27–31, 1989.
- [79658] 17988.Ren, Z.; Lin, P.-Y.; Klintworth, G. K.; Iwata, F.; Munier, F. L.;Schorderet, D. F.; El Matri, L.; Theendakara, V.; Basti, S.; Reddy,M.; Hejtmancik, J. F.: Allelic and locus het-



erogeneity in autosomalrecessive gelatinous drop-like corneal dystrophy. Hum. Genet. 110:568–577, 2002.

- [79659] 17989.Tsujikawa, M.; Kurahashi, H.; Tanaka, T.; Nishida, K.; Shimomura,Y.; Tano, Y.; Nakamura, Y.: Identification of the gene responsiblefor gelatinous drop-like corneal dystrophy. Nature Genet. 21: 420–423,1999.
- [79660] 17990.Willecke, K.; Jungbluth, S.; Dahl, E.; Hennemann, H.; Heynkes,R.; Grzeschik, K.–H.: Six genes of the human connexin gene familycoding for gap junctional proteins are assigned to four differenthuman chromosomes. Europ. J. Cell Biol. 53: 275–280, 1990.
- [79661] 17991.Reed, K. E.; Westphale, E. M.; Larson, D. M.; Wang, H.–Z.; Veenstra,R. D.; Beyer, E. C.: Molecular cloning and functional expressionof human connexin37, an endothelial cell gap junction protein. J.Clin. Invest. 91: 997–1004, 1993.
- [79662] 17992.Van Camp, G.; Coucke, P.; Speleman, F.; Van Roy, N.; Beyer, E.C.; Oostra, B. A.; Willems, P. J.: The gene for human gap junctionprotein connexin37 (GJA4) maps to chromosome 1p35.1, in the vicinityof D1S195. Genomics 30: 402–403, 1995.
- [79663] 17993.Gelb, B. D.; Zhang, J.; Cotter, P. D.; Gershin, I. F.; Desnick,R. J.: Physical mapping of the human connexin 40

(GJA5) flavin-containing monooxygenase 5, and natriuretic peptide receptor A genes on 1q21. *Genomics* 39:409–411, 1997.

[79664] 1994. Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Cardiac myocytes express multiple gap junction proteins. *Circ. Res.* 70: 438–444, 1992.

[79665] 1995. Kanter, H. L.; Saffitz, J. E.; Beyer, E. C.: Molecular cloning of two human cardiac gap junction proteins, connexin40 and connexin45. *J. Molec. Cell. Cardiol.* 26: 861–868, 1994.

[79666] 1996. Oviedo-Orta, E.; Hoy, T.; Evans, W. H.: Intercellular communication in the immune system: differential expression of connexin40 and 43, and perturbation of gap junction channel functions in peripheral blood and tonsil human lymphocyte subpopulations. *Immunology* 99: 578–590, 2000.

[79667] 1997. Seul, K. H.; Tadros, P. N.; Beyer, E. C.: Mouse connexin40: gene structure and promoter analysis. *Genomics* 46: 120–126, 1997.

[79668] 1998. Britz-Cunningham, S. H.; Shah, M. M.; Zuppan, C. W.; Fletcher, W. H.: Mutations of the connexin43 gap junction gene in patients with heart malformations and defects of laterality. *New Eng. J. Med.* 332:1323–1329, 1995.

- [79669] 17999.Brueckner, M.; D'Eustachio, P.; Horwich, A. L.: Linkage mapping of a mouse gene, iv, that controls left-right asymmetry of the heart and viscera. *Proc. Nat. Acad. Sci.* 86: 5035–5038, 1989.
- [79670] 18000.Burdine, R. D.; Schier, A. F.: Conserved and divergent mechanisms in left-right axis formation. *Genes Dev.* 14: 763–776, 2000.
- [79671] 18001.Casey, B.; Ballabio, A.: Connexin43 mutations in sporadic and familial defects of laterality. (Letter) *New Eng. J. Med.* 333: 941, 1995.
- [79672] 18002.Corcos, I. A.; Meese, E. U.; Loch-Caruso, R.: Human connexin 43 gene locus, GJA1, sublocalized to band 6q21–q23.2. *Cytogenet. Cell Genet.* 64: 31–32, 1993.
- [79673] 18003.Debrus, S.; Tuffery, S.; Matsuoka, R.; Galal, O.; Sarda, P.; Sauer, U.; Bozio, A.; Tanman, B.; Toutain, A.; Claustres, M.; Le Paslier, D.; Bouvagnet, P.: Lack of evidence for connexin 43 gene mutations in human autosomal recessive lateralization defects. *J. Molec. Cell. Cardiol.* 29: 1423–1431, 1997.
- [79674] 18004.Wijnen, J. T.; Oldenburg, M.; Bloemendal, H.; Meera Khan, P.: GS(gamma-S)–crystallin (CRYGS) assignment to chromosome 3. (Abstract) *Cytogenet. Cell Genet.* 51: 1108 only, 1989.

- [79675] 18005.Hai, T.; Liu, F.; Coukos, W. J.; Green, M. R.: Transcription factor ATF cDNA clones: an extensive family of leucine zipper proteins able to selectively form DNA-binding heterodimers. *Genes Dev.* 3: 2083–2090, 1989.
- [79676] 18006.Zucman, J.; Delattre, O.; Desmaze, C.; Epstein, A. L.; Stenman, G.; Speleman, F.; Fletcher, C. D. M.; Aurias, A.; Thomas, G.: EWS and ATF-1 gene fusion induced by t(12;22) translocation in malignant melanoma of soft parts. *Nature Genet.* 4: 341–345, 1993.
- [79677] 18007.Barco, A.; Alarcon, J. M.; Kandel, E. R.: Expression of constitutively active CREB protein facilitates the late phase of long-term potentiation by enhancing synaptic capture. *Cell* 108: 689–703, 2002.
- [79678] 18008.Barton, C. H.; Ajioka, J. W.; Roach, T. I. A.; Blackwell, J. M.: Mapping Creb-1 to chromosome 1 in the mouse. *Genomics* 14: 790–792, 1992.
- [79679] 18009.Barton, K.; Muthusamy, N.; Chanyangam, M.; Fischer, C.; Clendenin, C.; Leiden, J. M.: Defective thymocyte proliferation and IL-2 production in transgenic mice expressing a dominant-negative form of CREB. *Nature* 379:81–85, 1996.
- [79680] 18010.Carlezon, W. A., Jr.; Thome, J.; Olson, V. G.; Lane-Ladd, S. B.; Brodtkin, E. S.; Hiroi, N.; Duman, R. S.; Neve, R.

L.; Nestler, E.J.: Regulation of cocaine reward by CREB. *Science* 282: 2272–2275,1998.

[79681] 18011.Cole, T. J.; Copeland, N. G.; Gilbert, D. J.; Jenkins, N. A.; Schutz,G.; Ruppert, S.: The mouse CREB (cAMP responsive element bindingprotein) gene: structure, promoter analysis, and chromosomal localization. *Genomics* 13:974–982, 1992.

[79682] 18012.Fentzke, R. C.; Korcarz, C. E.; Lang, R. M.; Lin, H.; Leiden, J.M.: Dilated cardiomyopathy in transgenic mice expressing a dominant–negativeCREB transcription factor in the heart. *J. Clin. Invest.* 101: 2415–2426,1998.

[79683] 18013.Herzig, S.; Long, F.; Jhala, U. S.; Hedrick, S.; Quinn, R.; Bauer,A.; Rudolph, D.; Schutz, G.; Yoon, C.; Puigserver, P.; Spiegelman,B.; Montminy, M.: CREB regulates hepatic gluconeogenesis throughthe coactivator PGC–1. *Nature* 413: 179–183, 2001.

[79684] 18014.Benson, A. M.; Hunkeler, M. J.; Talalay, P.: Increase of NAD(P)H:quinonereductase by dietary antioxidants: possible role in protection againstcarcinogenesis and toxicity. *Proc. Nat. Acad. Sci.* 77: 5216–5220,1980.

[79685] 18015.Chen, L. Z.; Harris, P. C.; Apostolou, S.; Baker, E.; Holman, K.;Lane, S. A.; Nancarrow, J. K.; Whitmore, S. A.; Stallings, R. L.;Hildebrand, C. E.; Richards, R. I.; Suther–

land, G. R.; Callen, D.F.: A refined physical map of the long arm of human chromosome 16. *Genomics* 10:308–312, 1991.

[79686] 18016. Edwards, Y. H.; Hopkinson, D. A.; Carritt, B.: A genetic characterization of the human diaphorase-4 deficiency. *Ann. Hum. Genet.* 47: 97–105, 1983.

[79687] 18017. Grzeschik, K.-H.: Assignment of a structural gene for a fourth human diaphorase (DIA-4) to chromosome 16 in man-mouse somatic cell hybrids. *Hum. Genet.* 53: 189–193, 1980.

[79688] 18018. Jaiswal, A. K.; McBride, O. W.; Adesnik, M.; Nebert, D. W.: Human dioxin-inducible cytosolic NAD(P)H:menadione oxidoreductase: cDNA sequence and localization of gene to chromosome 16. *J. Biol. Chem.* 263:13572–13578, 1988.

[79689] 18019. Kelsey, K. T.; Ross, D.; Traver, R. D.; Christiani, D. C.; Zuo, Z. F.; Spitz, M. R.; Wang, M.; Xu, X.; Lee, B. K.; Schwartz, B. S.; Wiencke, J. K.: Ethnic variation in the prevalence of a common NAD(P)H quinone oxidoreductase polymorphism and its implications for anti-cancer chemotherapy. *Brit. J. Cancer* 76: 852–854, 1997.

[79690] 18020. Lavinha, J.; Morrison, N.; Glasgow, L.; Ferguson-Smith, M. A.: Further evidence for the regional localization

of human APRT and DIA4 on chromosome 16. (Abstract)  
Cytogenet. Cell Genet. 37: 517 only, 1984.

- [79691] 18021. Moran, J. L.; Siegel, D.; Ross, D.: A potential mechanism underlying the increased susceptibility of individuals with a polymorphism in NAD(P)H:quinone oxidoreductase 1 (NQO1) to benzene toxicity. Proc. Nat. Acad. Sci. 96: 8150–8155, 1999.
- [79692] 18022. Povey, S.; Wilson, D.; Edwards, Y. H.: Assignment of a human diaphorase (DIA-4) to chromosome 16. Ann. Hum. Genet. 43: 349–353, 1980.
- [79693] 18023. Radjendirane, V.; Joseph, P.; Lee, Y.-H.; Kimura, S.; Klein-Szanto, A. J. P.; Gonzalez, F. J.; Jaiswal, A. K.: Disruption of the DT diaphorase (NQO1) gene in mice leads to increased menadione toxicity. J. Biol. Chem. 273: 7382–7389, 1998.
- [79694] 18024. Rothman, N.; Smith, M. T.; Hayes, R. B.; Traver, R. D.; Hoener, B.; Campleman, S.; Li, G. L.; Dosemeci, M.; Linet, M.; Zhang, L.; Xi, L.; Wacholder, S.; Lu, W.; Meyer, K. B.; Titenko-Holland, N.; Stewart, J. T.; Yin, S.; Ross, D.: Benzene poisoning, a risk factor for hematological malignancy, is associated with the NQO1 609C–T mutation and rapid fractional excretion of chlorzoxazone. Cancer Res. 57: 2839–2842, 1997.

- [79695] 18025.Smith, M. T.: Benzene, NQO1, and genetic susceptibility to cancer.(Commentary) Proc. Nat. Acad. Sci. 96: 7624–7626, 1999.
- [79696] 18026.Traver, R. D.; Horikoshi, T.; Danenberg, K. D.; Stadlbauer, T.H.; Danenberg, P. V.; Ross, D.; Gibson, N. W.: NAD(P)H:quinone oxidoreductase gene expression in human colon carcinoma cells: characterization of a mutation which modulates DT-diaphorase activity and mitomycin sensitivity. Cancer Res. 52: 797–802, 1992.
- [79697] 18027.Traver, R. D.; Siegel, D.; Beall, H. D.; Phillips, R. M.; Gibson, N. W.; Franklin, W. A.; Ross, D.: Characterization of a polymorphism in NAD(P)H: quinone oxidoreductase (DT-diaphorase). Brit. J. Cancer 75:69–75, 1997.
- [79698] 18028.Zhang, L.; Rothman, N.; Wang, Y.; Hayes, R. B.; Li, G.; Dosemeci, M.; Yin, S.; Kolachana, P.; Titenko-Holland, N.; Smith, M. T.: Increased aneuploidy and long arm deletion of chromosomes 5 and 7 in the lymphocytes of Chinese workers exposed to benzene. Carcinogenesis 19: 1955–1961, 1998.
- [79699] 18029.Zhang, L.; Wang, Y.; Shang, N.; Smith, M. T.: Benzene metabolites induce the loss of long arm deletion of chromosomes 5 and 7 in human lymphocytes. Leukemia Res. 22: 105–113, 1998.



- [79700] 18030.Roychoudhury, A. K.; Nei, M.: Human Polymorphic Genes: World Distribution. New York: Oxford Univ. Press (pub.) 1988.
- [79701] 18031.Anholt, R. R. H.; Pederson, P. L.; Desouza, E. B.; Synder, S. H.: The peripheral-type benzodiazepine receptor: localization to the mitochondrial outer membrane. J. Biol. Chem. 261: 576–583, 1986.
- [79702] 18032.DeBernardi, M. A.; Crowe, R. R.; Mocchetti, I.; Shows, T. B.; Eddy, R. L.; Costa, E.: Chromosomal localization of the human diazepam binding inhibitor gene. Proc. Nat. Acad. Sci. 85: 6561–6565, 1988.
- [79703] 18033.Campanelli, J. T.; Hoch, W.; Rupp, F.; Kreiner, T.; Scheller, R.H.: Agrin mediates cell contact-induced acetylcholine receptor clustering. Cell 67:909–916, 1991.
- [79704] 18034.DeChiara, T. M.; Bowen, D. C.; Valenzuela, D. M.; Simmons, M. V.; Poueymirou, W. T.; Thomas, S.; Kinetz, E.; Compton, D. L.; Rojas, E.; Park, J. S.; Smith, C.; DiStefano, P. S.; Glass, D. J.; Burden, S. J.; Yancopoulos, G. D.: The receptor tyrosine kinase MuSK is required for neuromuscular junction formation in vivo. Cell 85: 501–512, 1996.
- [79705] 18035.Anonymous: Freely associating. (Editorial) Nature Genet. 22:1–2, 1999.
- [79706] 18036.Bell, G. I.; Rall, L. B.; Sanchez-Pescador, R.; Merry–

weather, J.P.; Scott, J.; Eddy, R. L.; Shows, T. B.: Human alpha-2-macroglobulin gene is located on chromosome 12. *Somat. Cell Molec. Genet.* 11:285–289, 1985.

[79707] 18037.Bergqvist, D.; Nilsson, I. M.: Hereditary alpha-2-macroglobulin deficiency. *Scand. J. Haemat.* 23: 433–436, 1979.

[79708] 18038.Blacker, D.; Crystal, A. S.; Wilcox, M. A.; Laird, N. M.; Tanzi, R. E.: Reply to Rudrasingham et al. *Nature Genet.* 22: 21–22, 1999.

[79709] 18039.Blacker, D.; Wilcox, M. A.; Laird, N. M.; Rodes, L.; Horvath, S.M.; Go, R. C. P.; Perry, R.; Watson, B., Jr.; Bassett, S. S.; McInnis, M. G.; Albert, M. S.; Hyman, B. T.; Tanzi, R. E.: Alpha-2 macroglobulin is genetically associated with Alzheimer disease. *Nature Genet.* 19:357–360, 1998.

[79710] 18040.David, F.; Kan, C. C.; Lucotte, G.: Two Taq I RFLPs for human alpha-2 macroglobulin (alpha-2M) using a full length cDNA probe. *Nucleic Acids Res.* 15: 374 only, 1987.

[79711] 18041.Chan, S. Y.; Empig, C. J.; Welte, F. J.; Speck, R. F.; Schmaljohn, A.; Kreisberg, J. F.; Goldsmith, M. A.: Folate receptor-alpha is a cofactor for cellular entry by Marburg and Ebola viruses. *Cell* 106:117–126, 2001.

[79712] 18042.Czeizel, A. E.; Dudas, I.: Prevention of the first oc-

currence of neural-tube defects by periconceptional vitamin supplementation. *New Eng. J. Med.* 327: 1832–1835, 1992.

- [79713] 18043. De Marco, P.; Moroni, A.; Merello, E.; de Franchis, R.; Andreussi, L.; Finnell, R. H.; Barber, R. C.; Cama, A.; Capra, V.: Folate pathway gene alterations in patients with neural tube defects. *Am. J. Med. Genet.* 95: 216–223, 2000.
- [79714] 18044. Elwood, P. C.: Molecular cloning and characterization of the human folate-binding protein cDNA from placenta and malignant tissue culture (KB) cells. *J. Biol. Chem.* 264: 14893–14901, 1989.
- [79715] 18045. Elwood, P. C.; Nachmanoff, K.; Saikawa, Y.; Page, S. T.; Pacheco, P.; Roberts, S.; Chung, K.-N.: The divergent 5-prime termini of the alpha human folate receptor (hFR) mRNAs originate from two tissue-specific promoters and alternative splicing: characterization of the alpha hFR gene structure. *Biochemistry* 36: 1467–1478, 1997.
- [79716] 18046. Boman, B. M.; Wildrick, D. M.; Alfaro, S. R.: Chromosome 18 allele loss at the D18S6 locus in human colorectal carcinomas. *Biochem. Biophys. Res. Commun.* 155: 463–469, 1988.
- [79717] 18047. Chan, S. S.-Y.; Zheng, H.; Su, M.-W.; Wilk, R.;

Killeen, M. T.; Hedgecock, E. M.; Culotti, J. G.: UNC-40, a *C. elegans* homolog of the DCC (deleted in colorectal cancer), is required in motile cells responding to UNC-6 netrin cues. *Cell* 87: 187–195, 1996.

[79718] 18048. Sarmanova, J.; Benesova, K.; Gut, I.; Nedelcheva-Kristensen, V.; Tynkova, L.; Soucek, P.: Genetic polymorphisms of biotransformation enzymes in patients with Hodgkin's and non-Hodgkin's lymphomas. *Hum. Molec. Genet.* 10: 1265–1273, 2001.

[79719] 18049. Nadeau, J. H.; Eicher, E. M.: Conserved linkage of soluble aconitase and galactose-1-phosphate uridyl transferase in mouse and man: assignment of these genes to mouse chromosome 4. *Cytogenet. Cell Genet.* 34: 271–281, 1982.

[79720] 18050. Albrecht, F. E.; Drago, J.; Felder, R. A.; Printz, M. P.; Eisner, G. M.; Robillard, J. E.; Sibley, D. R.; Westphal, H. J.; Jose, P. A.: Role of the D-1A dopamine receptor in the pathogenesis of genetic hypertension. *J. Clin. Invest.* 97: 2283–2288, 1996.

[79721] 18051. Bermak, J. C.; Li, M.; Bullock, C.; Zhou, Q.-Y.: Regulation of transport of the dopamine D1 receptor by a new membrane-associated ER protein. *Nature Cell Biol.* 3: 492–498, 2001.

- [79722] 18052. Boultwood, J.; Lewis, M. S.; Wainscoat, J. S.: Physical linkage of glucocorticoid receptor (GRL) and dopamine D1 receptor (DRD1) on the long arm of chromosome 5. (Abstract) Cytogenet. Cell Genet. 58:1894 only, 1991.
- [79723] 18053. Castner, S. A.; Williams, G. V.; Goldman-Rakic, P. S.: Reversal of antipsychotic-induced working memory deficits by short-term dopamine D1 receptor stimulation. Science 287: 2020–2022, 2000.
- [79724] 18054. Dearth, A.; Gingrich, J. A.; Falardeau, P.; Fremeau, R. T., Jr.; Bates, M. D.; Caron, M. G.: Molecular cloning and expression of the gene for a human D(1) dopamine receptor. Nature 347: 72–76, 1990.
- [79725] 18055. Grandy, D. K.; Zhou, Q.-Y.; Allen, L.; Litt, R.; Magenis, R. E.; Civelli, O.; Litt, M.: A human D(1) dopamine receptor gene is located on chromosome 5 at q35.1 and identifies an EcoRI RFLP. Am. J. Hum. Genet. 47: 828–834, 1990.
- [79726] 18056. Krushkal, J.; Xiong, M.; Ferrell, R.; Sing, C. F.; Turner, S. T.; Boerwinkle, E.: Linkage and association of adrenergic and dopamine receptor genes in the distal portion of the long arm of chromosome 5 with systolic blood pressure variation. Hum. Molec. Genet. 7:1379–1383, 1998.

- [79727] 18057.Mayerhofer, A.; Hemmings, H. C., Jr.; Snyder, G. L.; Greengard,P.; Boddien, S.; Berg, U.; Brucker, C.: Functional dopamine-1 receptorsand DARPP-32 are expressed in human ovary and granulosa luteal cellsin vitro. J. Clin. Endocr. Metab. 84: 257-264, 1999.
- [79728] 18058.Sunahara, R. K.; Niznik, H. B.; Weiner, D. M.; Stormann, T. M.;Brann, M. R.; Kennedy, J. L.; Gelernter, J. E.; Rozmahel, R.; Yang,Y.; Israel, Y.; Seeman, P.; O'Dowd, B. F.: Human dopamine D1 receptorencoded by an intronless gene on chromosome 5. Nature 347: 80-83,1990.
- [79729] 18059.Wilkie, T. M.; Chen, Y.; Gilbert, D. J.; Moore, K. J.; Yu, L.;Simon, M. I.; Copeland, N. G.; Jenkins, N. A.: Identification, chromosomallocation, and genome organization of mammalian G-protein-coupled receptors. Genomics 18:175-184, 1993.
- [79730] 18060.Xu, M.; Moratalla, R.; Gold, L. H.; Hiroi, N.; Koob, G. F.; Graybiel,A. M.; Tonegawa, S.: Dopamine D1 receptor mutant mice are deficientin striatal expression of dynorphin and in dopamine-mediated behavioralresponses. Cell 79: 729-742, 1994.
- [79731] 18061.Zhou, Q.-Y.; Grandy, D. K.; Thambi, L.; Kushner, J. A.; Van Tol,H. H. M.; Cone, R.; Pribnow, D.; Salon, J.; Bunzow, J. R.; Civelli,O.: Cloning and expression of human

and rat D(1) dopamine receptors. *Nature* 347:76–80, 1990.

- [79732] 18062. Balk, J.-H.; Picetti, R.; Salardi, A.; Thirlet, G.; Dierich, A.; Depaulis, A.; Le Meur, M.; Borrelli, E.: Parkinsonian-like locomotor impairment in mice lacking dopamine D2 receptors. *Nature* 377: 424–428, 1995.
- [79733] 18063. Basu, S.; Nagy, J. A.; Pal, S.; Vasile, E.; Eckelhoefer, I. A.; Bliss, V. S.; Manseau, E. J.; Dasgupta, P. S.; Dvorak, H. F.; Mukhopadhyay, D.: The neurotransmitter dopamine inhibits angiogenesis induced by vascular permeability factor/vascular endothelial growth factor. *Nature Med.* 7: 569–574, 2001.
- [79734] 18064. Blum, K.; Noble, E. P.; Sheridan, P. J.; Montgomery, A.; Ritchie, T.; Jagadeeswaran, P.; Nogami, H.; Briggs, A. H.; Cohn, J. B.: Allelic association of human dopamine D(2) receptor gene in alcoholism. *J.A.M.A.* 263:2055–2060, 1990.
- [79735] 18065. Bolos, A. M.; Dean, M.; Lucas-Derse, S.; Ramsburg, M.; Brown, G.L.; Goldman, D.: Population and pedigree studies reveal a lack of association between the dopamine D(2) receptor gene and alcoholism. *J.A.M.A.* 264:3156–3160, 1990.
- [79736] 18066. Bunzow, J. R.; Van Tol, H. H.; Grandy, D. K.; Albert,

P.; Salon, J.; Christie, M.; Machida, C. A.; Neve, K. A.; Civelli, O.: Cloning and expression of a rat D2 dopamine receptor cDNA. *Nature* 336: 783–787, 1988.

- [79737] 18067. Dal Toso, R.; Sommer, B.; Ewert, M.; Herb, A.; Pritchett, D. B.; Bach, A.; Shivers, B. D.; Seeburg, P. H.: The dopamine D(2) receptor: two molecular forms generated by alternative splicing. *EMBO J.* 8:4025–4034, 1989.
- [79738] 18068. Seeman, P.; Guan, H.-C.; Van Tol, H. H. M.: Dopamine D4 receptors elevated in schizophrenia. *Nature* 365: 441–445, 1993.
- [79739] 18069. Seeman, P.; Ulpian, C.; Chouinard, G.; Van Tol, H. H. M.; Dwosh, H.; Lieberman, J. A.; Siminovich, K.; Liu, I. S. C.; Wayne, J.; Voruganti, P.; Hudson, C.; Serjeant, G. R.; Masisbay, A. S.; Seeman, M. V.: Dopamine D4 receptor variant, D4GLYCINE194, in Africans, but not in Caucasians: no association with schizophrenia. *Am. J. Med. Genet.* 54: 384–390, 1994.
- [79740] 18070. Smalley, S. L.; Bailey, J. N.; Palmer, C. G.; Cantwell, D. P.; McGough, J. J.; Del'Homme, M. A.; Asarnow, J. R.; Woodward, J. A.; Ramsey, C.; Nelson, S. F.: Evidence that the dopamine D4 receptor is a susceptibility gene in attention deficit hyperactivity disorder. *Molec. Psychiat.* 3: 427–430, 1998.



- [79741] 18071.Meera Khan, P.: Personal Communication. Leiden, The Netherlands 1977.
- [79742] 18072.Takai, T.; Li, M.; Sylvestre, D.; Clynes, R.; Ravetch, J. V.:FcR gamma chain deletion results in pleiotrophic effector cell defects. Cell 76:519–529, 1994.
- [79743] 18073.Garman, S. C.; Kinet, J.–P.; Jardetzky, T. S.: Crystal structure of the human high–affinity IgE receptor. Cell 95: 951–961, 1998.
- [79744] 18074.Garman, S. C.; Wurzburg, B. A.; Tarchevskaya, S. S.; Kinet, J.–P.;Jardetzky, T. S.: Structure of the Fc fragment of human IgE bound to its high–affinity receptor Fc–epsilon–RI–alpha. Nature 406: 259–266,2000.
- [79745] 18075.Liu, F.–T.; Albrandt, K.; Robertson, M. W.: cDNA heterogeneity suggests structural variants related to the high–affinity IgE receptor. Proc.Nat. Acad. Sci. 85: 5639–5643, 1988.
- [79746] 18076.Shimizu, A.; Tepler, I.; Benfey, P. N.; Berenstein, E. H.; Siraganian,R. P.; Leder, P.: Human and rat mast cell high–affinity immunoglobulinE receptors: characterization of putative alpha–chain gene products. Proc.Nat. Acad. Sci. 85: 1907–1911, 1988.
- [79747] 18077.Miyajima, I.; Dombrowicz, D.; Martin, T. R.; Ravetch, J. V.; Kinet,J.–P.; Galli, S. J.: Systemic anaphylaxis

in the mouse can be mediated largely through IgG1 and Fc-gamma-RIII: assessment of the cardiopulmonary changes, mast cell degranulation, and death associated with active or IgE- or IgG1-dependent passive anaphylaxis. *J. Clin. Invest.* 99:901-914, 1997.

[79748] 18078. Tepler, I.; Morton, C. C.; Shimizu, A.; Holcombe, R. F.; Eddy, R.; Shows, T. B.; Leder, P.: The gene for the human mast cell high-affinity IgE receptor alpha chain: chromosomal localization to 1q21-q23 and RFLP analysis. *Am. J. Hum. Genet.* 45: 761-765, 1989.

[79749] 18079. Zhu, D.; Kepley, C. L.; Zhang, M.; Zhang, K.; Saxon, A.: A novel human immunoglobulin Fc-gamma-Fc-epsilon bifunctional fusion protein inhibits Fc-epsilon-RI-mediated degranulation. *Nature Med.* 8: 518-521, 2002.

[79750] 18080. Bain, G.; Robanus Maandag, E. C.; Izon, D. J.; Amesen, D.; Kruisbeek, A. M.; Weintraub, B. C.; Krop, I.; Schlissel, M. S.; Feeney, A. J.; van Roon, M.; van der Valk, M.; te Riele, H. P. J.; Berns, A.; Murre, C.: E2A proteins are required for proper B cell development and initiation of immunoglobulin gene rearrangements. *Cell* 79: 885-892, 1994.

[79751] 18081. El Ghouzzi, V.; Legeai-Mallet, L.; Aresta, S.;

Benoist, C.; Munnich,A.; de Gunzburg, J.; Bonaventure, J.: Saethre–Chotzen mutations causeTWIST protein degradation or impaired nuclear location. Hum. Molec.Genet. 9: 813–819, 2000.

[79752] 18082.Henthorn, P.; Kiledjian, M.; Kadesch, T.: Two distinct transcriptionfactors that bind the immunoglobulin enhancer mu–E5/kappa–E2 motif. Science 247:467–470, 1990.

[79753] 18083.Kamps, M. P.; Murre, C.; Sun, X.; Baltimore, D.: A new homeoboxgene contributes the DNA binding domain of the t(1;19) translocationprotein in pre–B ALL. Cell 60: 547–555, 1990.

[79754] 18084.Mellentin, J. D.; Murre, C.; Donlon, T. A.; McCaw, P. S.; Smith,S. D.; Carroll, A. J.; McDonald, M. E.; Baltimore, D.; Cleary, M.L.: The gene for enhancer binding proteins E12/E47 lies at the t(1;19)breakpoint in acute leukemias. Science 246: 379–382, 1989.

[79755] 18085.Murre, C.; McCaw, P. S.; Baltimore, D. :Cell 56: 777–783, 1989.

[79756] 18086.Nourse, J.; Mellentin, J. D.; Galili, N.; Wilkinson, J.; Stanbridge,E.; Smith, S. D.; Cleary, M. L.: Chromosomal translocation t(1;19)results in synthesis of a homeobox fusion mRNA that codes for a potentialchimeric transcrip–

tion factor. Cell 60: 535–545, 1990.

- [79757] 18087.Fan, Y.–S.; Eddy, R. L.; Byers, M. G.; Haley, L. L.; Henry, W.M.; Kayano, T.; Shows, T. B.; Bell, G. I.: Assignment of genes encoding three human glucose transporter/transporter–like proteins (GLUT4, GLUT5 and GLUT6) to chromosomes 17, 1 and 5, respectively. (Abstract) Cytogenet. Cell Genet. 51: 997 only, 1989.
- [79758] 18088.Gould, G. W.; Holman, G. D.: The glucose transporter family: structure, function and tissue–specific expression. Biochem. J. 295: 329–341, 1993.
- [79759] 18089.Hauguel–de Mouzon, S.; Challier, J. C.; Kacemi, A.; Cauzac, M.; Malek, A.; Girard, J.: The GLUT3 glucose transporter isoform is differentially expressed within human placental cell types. J. Clin. Endocr. Metab. 82:2689–2694, 1997.
- [79760] 18090.Kayano, T.; Burant, C. F.; Fukumoto, H.; Gould, G. W.; Fan, Y.S.; Eddy, R. L.; Byers, M. G.; Shows, T. B.; Seino, S.; Bell, G. I.: Human facilitative glucose transporters: isolation, functional characterization, and gene localization of cDNAs encoding an isoform (GLUT5) expressed in small intestine, kidney, muscle, and adipose tissue and an unusual glucose transporter pseudogene–like sequence (GLUT6). J. Biol. Chem. 265:13276–13282, 1990.

- [79761] 18091.Kayano, T.; Fukumoto, H.; Eddy, R. L.; Fan, Y.-S.; Byers, M. G.;Shows, T. B.; Bell, G. I.: Evidence for a family of human glucosetransporter-like proteins: sequence and gene localization of a proteinexpressed in fetal skeletal muscle and other tissues. J. Biol. Chem. 263:15245-15248, 1988.
- [79762] 18092.Aitken, D. A.; Ferguson-Smith, M. A.: Gene dosage evidence forthe regional assignment of the GOT-S structural gene locus to 10q24-10q25. Cytogenet.Cell Genet. 22: 468-471, 1978.
- [79763] 18093.Creagan, R.; Tischfield, J.; McMorris, F. A.; Chen, S.-H.; Hirschi,M.; Chen, T.-T.; Ricciuti, F.; Ruddle, F. H.: Assignment of the genesfor human peptidase A to chromosome 18 and cytoplasmic glutamic oxaloacetate-transaminase to chromosome 10 using somatic-cell hybrids. Cytogenet.Cell Genet. 12: 187-198, 1973.
- [79764] 18094.Ford, G. C.; Eichele, G.; Jansonius, J. N.: Three-dimensionalstructure of a pyridoxal-phosphate-dependent enzyme, mitochondrialaspartate amino-transferase. Proc. Nat. Acad. Sci. 77: 2559-2563,1980.
- [79765] 18095.Gitelman, B. J.; Tomkins, D. J.; Partington, M. W.; Roberts, M.H.; Simpson, N. E.: Gene dosage studies of glutamic oxaloacetic transaminase(GOT) and hexokinase

(HK) in two patients with possible partial trisomy10q.

(Abstract) Am. J. Hum. Genet. 32: 41A only, 1980.

[79766] 18096.Junien, C.; Despoisse, S.; Turleau, C.; de Grouchy, J.; Bucher,T.; Fundele, R.: Assignment of phosphoglycerate mutase (PGAMA) to human chromosome 10: regional mapping of GOT1 and PGAMA to subbands10q26.1 (or q25.3). Ann. Genet. 25: 25–27, 1982.

[79767] 18097.Overhauser, J.; Mewar, R.; Rojas, K.; Lia, K.; Kline, A. D.; Silverman,G. A.: STS map of genes and anonymous DNA fragments on human chromosome18 using a panel of somatic cell hybrids. Genomics 15: 387–391,1993.

[79768] 18098.Abdalla, S. A.; Pece–Barbara, N.; Vera, S.; Tapia, E.; Paez, E.;Bernabeu, C.; Letarte, M.: Analysis of ALK–1 and endoglin in newborns from families with hereditary hemorrhagic telangiectasia type 2. Hum.Molec. Genet. 9: 1227–1237, 2000.

[79769] 18099.Bode–Lesniewska, B.; Dours–Zimmermann, M. T.; Odermatt, B. F.;Briner, J.; Heitz, P. U.; Zimmermann, D. R.: Distribution of the large aggregating proteoglycan versican in adult human tissues. J.Histochem. Cytochem. 44: 303–312, 1996.

[79770] 18100.Dours–Zimmermann, M. T.; Zimmermann, D. R.: A novel glycosaminoglycan attachment domain identified in

two alternative splice variants of human versican. J. Biol. Chem. 269: 32992–32998, 1994.

- [79771] 18101. Iozzo, R. V.; Naso, M. F.; Cannizzaro, L. A.; Wasmuth, J. J.; McPherson, J. D.: Mapping of the versican proteoglycan gene (CSPG2) to the long arm of human chromosome 5 (5q12–5q14). Genomics 14: 845–851, 1992.
- [79772] 18102. Kjellen, L.; Lindahl, U.: Proteoglycans: structures and interactions. Annu. Rev. Biochem. 60: 443–475, 1991.
- [79773] 18103. Naso, M. F.; Morgan, J. L.; Buchberg, A. M.; Siracusa, L. D.; Iozzo, R. V.: Expression pattern and mapping of the murine versican gene (Cspg2) to chromosome 13. Genomics 29: 297–300, 1995.
- [79774] 18104. Naso, M. F.; Zimmermann, D. R.; Iozzo, R. V.: Characterization of the complete genomic structure of the human versican gene and functional analysis of its promoter. J. Biol. Chem. 269: 32999–33008, 1994.
- [79775] 18105. Zako, M.; Shinomura, T.; Ujita, M.; Ito, K.; Kimata, K.: Expression of PG-M(V3), an alternatively spliced form of PG-M without a chondroitin sulfate attachment region in mouse and human tissues. J. Biol. Chem. 270: 3914–3918, 1995.
- [79776] 18106. Zimmermann, D. R.; Ruoslahti, E.: Multiple domains of the large fibroblast proteoglycan, versican. EMBO J. 8:

2975–2981, 1989.

- [79777] 18107.Byrne, E.; White, O.; Cook, M.: Familial dystonic choreoathetosis with myokymia; a sleep responsive disorder. *J. Neurol. Neurosurg.Psychiat.* 54: 1090–1092, 1991.
- [79778] 18108.Demirkiran, M.; Jankovic, J.: Paroxysmal dyskinesias: clinical features and classification. *Ann. Neurol.* 38: 571–579, 1995.
- [79779] 18109.Fink, J. K.; Rainier, S.; Wilkowski, J.; Jones, S. M.; Kume, A.; Hedera, P.; Albin, R.; Mathay, J.; Girbach, L.; Varvil, T.; Otterud, B.; Leppert, M.: Paroxysmal dystonic choreoathetosis: tight linkage to chromosome 2q. *Am. J. Hum. Genet.* 59: 140–145, 1996.
- [79780] 18110.Fouad, G. T.; Servidei, S.; Durcan, S.; Bertini, E.; Ptacek, L.J.: A gene for familial paroxysmal dyskinesia (FPD1) maps to chromosome 2q. *Am. J. Hum. Genet.* 59: 135–139, 1996.
- [79781] 18111.Grunder, S.; Geisler, H.-S.; Rainer, S.; Fink, J. K.: Acid-sensing ion channel (ASIC) 4 gene: physical mapping, genomic organisation, and evaluation as a candidate for paroxysmal dystonia. *Europ. J. Hum. Genet.* 9: 672–676, 2001.
- [79782] 18112.Hudgins, R. L.; Corbin, K. B.: An uncommon seizure disorder: familial paroxysmal choreoathetosis. *Brain* 89:



199–204, 1966.

- [79783] 18113.Koch, G.; Lalley, P. A.; McAvoy, M.; Shows, T. B.: Assignment of LIPA, associated with human acid lipase deficiency, to human chromosome 10 and comparative assignment to mouse chromosome 19. *Somat. Cell Genet.* 7: 345–358, 1981.
- [79784] 18114.Kadotani, H.; Hirano, T.; Masugi, M.; Nakamura, K.; Nakao, K.; Katsuki, M.; Nakanishi, S.: Motor discoordination results from combined gene disruption of the NMDA receptor NR2A and NR2C subunits, but not from single disruption of the NR2A or NR2C subunit. *J. Neurosci.* 16:7859–7867, 1996.
- [79785] 18115.Lin, Y. J.; Bovetto, S.; Carver, J. M.; Giordano, T.: Cloning of the cDNA for the human NMDA receptor NR2C subunit and its expression in the central nervous system and periphery. *Molec. Brain Res.* 43:57–64, 1996.
- [79786] 18116.Islam, M. Q.; Platz, A.; Szpirer, J.; Szpirer, C.; Levan, G.; Mannervik, B.: Chromosomal localization of human glutathione transferase genes of classes alpha, mu and pi. *Hum. Genet.* 82: 338–342, 1989.
- [79787] 18117.Albert, L. J.; Inman, R. D.: Molecular mimicry and autoimmunity. *New Eng. J. Med.* 341: 2068–2074, 1999.
- [79788] 18118.Asada, H.; Kawamura, Y.; Maruyama, K.; Kume, H.;

Ding, R.-G.; Ji, F. Y.; Kanbara, N.; Kuzume, H.; Sanbo, M.; Yagi, T.; Obata, K.: Micelacking the 65 kDa isoform of glutamic acid decarboxylase (GAD65) maintain normal levels of GAD67 and GABA in their brains but are susceptible to seizures. *Biochem. Biophys. Res. Commun.* 229: 891–895, 1996.

[79789] 18119. Baekkeskov, S.; Aanstoot, H.-J.; Christgau, S.; Reetz, A.; Solimena, M.; Cascalho, M.; Folli, F.; Richter-Olesen, H.; Camilli, P.-D.: Identification of the 64K autoantigen in insulin-dependent diabetes as the GABA-synthesizing enzyme glutamic acid decarboxylase. *Nature* 347:151–156, 1990.

[79790] 18120. Bu, D.-F.; Erlander, M. G.; Hitz, B. C.; Tillakaratne, N. J. K.; Kaufman, D. L.; Wagner-McPherson, C. B.; Evans, G. A.; Tobin, A. J.: Two human glutamate decarboxylases, 65-kDa GAD and 67-kDa GAD, are each encoded by a single gene. *Proc. Nat. Acad. Sci.* 89: 2115–2119, 1992.

[79791] 18121. Cram, D. S.; Barnett, L. D.; Joseph, J. L.; Harrison, L. C.: Cloning and partial nucleotide sequence of human glutamic acid decarboxylase cDNA from brain and pancreatic islets. *Biochem. Biophys. Res. Commun.* 176:1239–1244, 1991.

[79792] 18122. De Aizpurua, H. J.; Wilson, Y. M.; Harrison, L. C.:

Glutamic aciddecarboxylase autoantibodies in preclinical insulin-dependent diabetes. Proc.Nat. Acad. Sci. 89: 9841–9845, 1992.

[79793] 18123.Edelhoff, S.; Grubin, C. E.; Karlsen, A. E.; Adler, D. A.; Foster,D.; Disteché, C. M.; Lernmark, A.: Mapping of glutamic acid decarboxylase(GAD) genes. Genomics 17: 93–97, 1993.

[79794] 18124.Jones, D. B.; Armstrong, N. W.: Coxsackie virus and diabetes revisited.(Letter) Nature Med. 1: 284 only, 1995.

[79795] 18125.Karlsen, A. E.; Hagopian, W. A.; Grubin, C. E.; Dube, S.; Disteché,C. M.; Adler, D. A.; Barmeier, H.; Mathewes, S.; Grant, F. J.; Foster,D.; Lernmark, A.: Cloning and primary structure of a human isletisoform of glutamic acid decarboxylase from chromosome 10. Proc.Nat. Acad. Sci. 88: 8337–8341, 1991.

[79796] 18126.Kash, S. F.; Johnson, R. S.; Tecott, L. H.; Noebels, J. L.; Mayfield,R. D.; Hanahan, D.; Baekkeskov, S.: Epilepsy in mice deficient inthe 65–kDa isoform of glutamic acid decarboxylase. Proc. Nat. Acad.Sci. 94: 14060–14065, 1997.

[79797] 18127.Kaufman, D. L.; Erlander, M. G.; Clare–Salzler, M.; Atkinson,M. A.; Maclaren, N. K.; Tobin, A. J.: Autoimmunity to two forms ofglutamate decarboxylase in insulin-dependent diabetes mellitus. J.Clin. Invest. 89: 283–292,

1992.

- [79798] 18128.Lohmann, T.; Hawa, M.; Leslie, R. D. G.; Lane, R.; Picard, J.;Londei, M.: Immune reactivity to glutamic acid decarboxylase 65 instiff-man syndrome and type 1 diabetes mellitus. *Lancet* 356: 31–35,2000.
- [79799] 18129.Richter, W.; Endl, J.; Eiermann, T. H.; Brandt, M.; Kientsch-Engel,R.; Thivolet, C.; Jungfer, H.; Scherbaum, W. A.: Human monoclonalislet cell antibodies from a patient with insulin-dependent diabetesmellitus reveal glutamate decarboxylase as the target antigen. *Proc.Nat. Acad. Sci.* 89: 8467–8471, 1992.
- [79800] 18130.Tian, J.; Lehmann, P. V.; Kaufman, D. L.: T cell cross-reactivitybetween coxsackievirus and glutamate decarboxylase is associated witha murine diabetes susceptibility allele. *J. Exp. Med.* 180: 1979–1984,1994.
- [79801] 18131.McKusick, V. A.: The morbid anatomy of the human genome: a reviewof gene mapping in clinical medicine (part 1). *Medicine* 65: 1–33,1986.
- [79802] 18132.Panteghini, M.: Aspartate aminotransferase isoenzymes. *Clin.Biochem.* 23: 311–319, 1990.
- [79803] 18133.Pol, S.; Bousquet-Lemercier, B.; Pave-Preux, M.; Bulle, F.; Passage,E.; Hanoune, J.; Mattei, M. G.; Barouki, R.: Chromosomal localizationof human aspartate amino-

transferase genes by in situ hybridization. Hum.Genet. 83: 159–164, 1989.

[79804] 18134.Scott, E. M.; Wright, R. C.: An alternate method for demonstration of erythrocytic aminotransferases on starch gels. Am. J. Hum. Genet. 33:561–563, 1981.

[79805] 18135.Spritz, R. A.; Emanuel, B. S.; Chern, C. J.; Mellman, W. J.:Gene dosage effect: intraband mapping of human soluble glutamic oxaloacetic transaminase. Cytogenet. Cell Genet. 23: 149–156, 1979.

[79806] 18136.Tomkins, D. J.; Gitelman, B. J.; Roberts, M. H.: Confirmation of a de novo duplication, dup(10)(q24–q26), by GOT1 gene dosage studies. Hum.Genet. 63: 369–373, 1983.

[79807] 18137.Wang, C.–Y.; Huang, Y.–Q.; Shi, J.–O.; Marron, M. P.; Ruan, Q.–G.; Hawkins–Lee, B.; Ochoa, B.; She, J.–X.: Genetic homogeneity, high–resolution mapping, and mutation analysis of the urofacial (Ochoa) syndrome and exclusion of the glutamate oxaloacetate transaminase gene (GOT1) in the critical region as the disease gene. Am. J. Med. Genet. 84:454–459, 1999.

[79808] 18138.Wurzinger, K. H.; Mohrenweiser, H. W.: Studies on the genetic and non–genetic (physiological) variation of human erythrocyte glutamic oxaloacetic transaminase.

Ann. Hum. Genet. 46: 191–201, 1982.

- [79809] 18139. Abel, E. D.; Kaulbach, H. C.; Tian, R.; Hopkins, J. C. A.; Duffy, J.; Doetschman, T.; Minnemann, T.; Boers, M.-E.; Hadro, E.; Oberste-Berghaus, C.; Quist, W.; Lowell, B. B.; Ingwall, J. S.; Kahn, B. B.: Cardiachypertrophy with preserved contractile function after selective deletion of GLUT4 from the heart. *J. Clin. Invest.* 104: 1703–1714, 1999.
- [79810] 18140. Abel, E. D.; Peroni, O.; Kim, J. K.; Kim, Y.-B.; Boss, O.; Hadro, E.; Minnemann, T.; Shulman, G. I.; Kahn, B. B.: Adipose-selective targeting of the GLUT4 gene impairs insulin action in muscle and liver. *Nature* 409: 729–733, 2001.
- [79811] 18141. Bell, G. I.; Kayano, T.; Buse, J. B.; Burant, C. F.; Takeda, J.; Lin, D.; Fukumoto, H.; Seino, S.: Molecular biology of mammalian glucose transporters. *Diabetes Care* 13: 198–208, 1990.
- [79812] 18142. Bell, G. I.; Murray, J. C.; Nakamura, Y.; Kayano, T.; Eddy, R. L.; Fan, Y.-S.; Byers, M. G.; Shows, T. B.: Polymorphic human insulin-responsive glucose-transporter gene on chromosome 17p13. *Diabetes* 38: 1072–1075, 1989.
- [79813] 18143. Birnbaum, M. J.: Identification of a novel gene encoding an insulin-responsive glucose transporter protein.

Cell 57: 305–315, 1989.

- [79814] 18144. Chiang, S.-H.; Baumann, C. A.; Kanzaki, M.; Thirumond, D. C.; Watson, R. T.; Neudauer, C. L.; Macara, I. G.; Pessin, J. E.; Saltiel, A. R.: Insulin-stimulated GLUT4 translocation requires the CAP-dependent activation of TC10. *Nature* 410: 944–948, 2001.
- [79815] 18145. Garvey, W. T.; Maianu, L.; Zhu, J.-H.; Brechtel-Hook, G.; Wallace, P.; Baron, A. D.: Evidence for defects in the trafficking and translocation of GLUT4 glucose transporters in skeletal muscle as a cause of human insulin resistance. *J. Clin. Invest.* 101: 2377–2386, 1998.
- [79816] 18146. Ikemoto, S.; Thompson, K. S.; Itakura, H.; Lane, M. D.; Ezaki, O.: Expression of an insulin-responsive glucose transporter (GLUT4) minigene in transgenic mice: effect of exercise and role in glucose homeostasis. *Proc. Nat. Acad. Sci.* 92: 865–869, 1995.
- [79817] 18147. Katz, E. B.; Stenbit, A. E.; Hatton, K.; DePinho, R.; Charron, M. J.: Cardiac and adipose tissue abnormalities but not diabetes in mice deficient in GLUT4. *Nature* 377: 151–155, 1995.
- [79818] 18148. Kusari, J.; Verma, U. S.; Buse, J. B.; Henry, R. R.; Olefsky, J. M.: Analysis of the gene sequences of the insulin receptor and the insulin-sensitive glucose trans-

porter (GLUT-4) in patients with common-type non-insulin-dependent diabetes mellitus. *J. Clin. Invest.* 88:1323–1330, 1991.

- [79819] 18149. Muraoka, A.; Sakura, H.; Kim, K.; Kishimoto, M.; Akanuma, Y.; Buse, J. B.; Yasuda, K.; Seino, S.; Bell, G. I.; Yazaki, Y.; Kasuga, M.; Kadowaki, T.: Polymorphism in exon 4a of the human GLUT4/muscle-fat facilitative glucose transporter gene detected by SSCP. *Nucleic Acids Res.* 19: 4313 only, 1991.
- [79820] 18150. Ribon, V.; Printen, J. A.; Hoffman, N. G.; Kay, B. K.; Saltiel, A. R.: A novel, multifunctional c-Cbl binding protein in insulin receptor signaling in 3T3-L1 adipocytes. *Molec. Cell. Biol.* 18:872–879, 1998.
- [79821] 18151. Zisman, A.; Peroni, O. D.; Abel, E. D.; Michael, M. D.; Mauvais-Jarvis, F.; Lowell, B. B.; Wojtaszewski, J. F. P.; Hirshman, M. F.; Virkamaki, A.; Goodyear, L. J.; Kahn, C. R.; Kahn, B. B.: Targeted disruption of the glucose transporter 4 selectively in muscle causes insulin resistance and glucose intolerance. *Nature Med.* 6: 924–928, 2000.
- [79822] 18152. Astrin, K. H.; Arredondo-Vega, F. X.; Desnick, R. J.; Smith, M.: Assignment of the gene for cytosolic alanine aminotransferase (AAT1) to human chromosome 8. *Ann. Hum. Genet.* 46: 125–133, 1982.



- [79823] 18153.Chen, S.-H.; Giblett, E. R.: Polymorphism of soluble glutamic-pyruvictransaminase: a new genetic marker in man. *Science* 173: 148–149,1971.
- [79824] 18154.Cook, P. J. L.; Jeremiah, S. J.; Buckton, K. E.: Exclusion mapping of GPT. (Abstract) *Cytogenet. Cell Genet.* 32: 258 only, 1982.
- [79825] 18155.Mombaerts, P.; Wang, F.; Dulac, C.; Vassar, R.; Chao, S. K.; Nemes,A.; Mendelsohn, M.; Edmondson, J.; Axel, R.: The molecular biology of olfactory perception. *Cold Spring Harbor Symp. Quant. Biol.* 61:135–145, 1996.
- [79826] 18156.Reed, R. R.: Genetic approaches to mammalian olfaction. *ColdSpring Harbor Symp. Quant. Biol.* 61: 165–172, 1996.
- [79827] 18157.Ronnett, G. V.; Snyder, S. H.: Molecular messengers of olfaction. *TrendsNeurosci.* 15: 508–513, 1992.
- [79828] 18158.Schwab, S. G.; Hallmayer, J.; Lerer, B.; Albus, M.; Borrmann, M.;Honig, S.; Strauss, M.; Segman, R.; Lichter-mann, D.; Knapp, M.; Trixler,M.; Maier, W.; Wildenauer, D. B.: Support for a chromosome 18p locus conferring susceptibility to functional psychoses in families with-schizophrenia, by association and linkage analysis. *Am. J. Hum. Genet.* 63:1139–1152, 1998.
- [79829] 18159.Davignon, I.; Barnard, M.; Gavrilova, O.; Sweet, K.;

Wilkie, T.M.: Gene structure of murine Gna11 and Gna15: tandemly duplicated Gq class G protein alpha subunit genes. *Genomics* 31: 359–366, 1996.

[79830] 18160. Jiang, M.; Pandey, S.; Tran, V. T.; Fong, H. K.: Guanine nucleotide-binding regulatory proteins in retinal pigment epithelial cells. *Proc. Nat. Acad. Sci.* 88: 3907–3911, 1991.

[79831] 18161. Offermanns, S.; Zhao, L.-P.; Gohla, A.; Sarosi, I.; Simon, M. I.; Wilkie, T. M.: Embryonic cardiomyocyte hypoplasia and craniofacial defects in G-alpha-q/G-alpha-11-mutant mice. *EMBO J.* 17: 4304–4312, 1998.

[79832] 18162. Strathmann, M. P.; Simon, M. I.: G-alpha-12 and G-alpha-13 subunits define a fourth class of G protein alpha subunits. *Proc. Nat. Acad. Sci.* 88: 5582–5586, 1991.

[79833] 18163. Ahmed, S. F.; Barr, D. G. D.; Bonthron, D. T.: GNAS1 mutations and progressive osseous heteroplasia. (Letter) *New Eng. J. Med.* 346:1669–1670, 2002.

[79834] 18164. Ahmed, S. F.; Dixon, P. H.; Bonthron, D. T.; Stirling, H. F.; Barr, D. G. D.; Kelnar, C. J. H.; Thakker, R. V.: GNAS1 mutational analysis in pseudohypoparathyroidism. *Clin. Endocr.* 49: 525–531, 1998.

[79835] 18165. Ahrens, W.; Hiort, O.; Staedt, P.; Kirschner, T.;

Marschke, C.; Kruse, K.: Analysis of the GNAS1 gene in Albright's hereditary osteodystrophy. *J.Clin. Endocr. Metab.* 86: 4630–4634, 2001.

[79836] 18166. Aldred, M. A.; Trembath, R. C.: Activating and inactivating mutations in the human GNAS1 gene. *Hum. Mutat.* 16: 183–189, 2000.

[79837] 18167. Ashley, P. L.; Ellison, J.; Sullivan, K. A.; Bourne, H. R.; Cox, D. R.: Chromosomal assignment of the murine Gi and Gs genes. (Abstract) *Am.J. Hum. Genet.* 41: A155 only, 1987.

[79838] 18168. Ballare, E.; Mantovani, S.; Lania, A.; Di Blasio, A. M.; Vallar, L.; Spada, A.: Activating mutations of the GS- $\alpha$  gene are associated with low levels of GS- $\alpha$  protein in growth hormone-secreting tumors. *J.Clin. Endocr. Metab.* 83: 4386–4390, 1998.

[79839] 18169. Bastepe, M.; Pincus, J. E.; Sugimoto, T.; Tojo, K.; Kanatani, M.; Azuma, Y.; Kruse, K.; Rosenbloom, A. L.; Koshiyama, H.; Juppner, H.: Positional dissociation between the genetic mutation responsible for pseudohypoparathyroidism type Ib and the associated methylation defect at exon A/B: evidence for a long-range regulatory element within the imprinted GNAS1 locus. *Hum. Molec. Genet.* 10: 1231–1241, 2001.

- [79840] 18170. Bastepe, M.; Juppner, H.: GNAS1 mutations and progressive osseous heteroplasia. (Letter) *New Eng. J. Med.* 346: 1671 only, 2002.
- [79841] 18171. Bastepe, M.; Lane, A. H.; Juppner, H.: Parental uniparental isodisomy of chromosome 20q--and the resulting changes in GNAS1 methylation--as a plausible cause of pseudohypoparathyroidism. *Am. J. Hum. Genet.* 68:1283-1289, 2001.
- [79842] 18172. Bianco, P.; Riminucci, M.; Majolagbe, A.; Kuznetsov, S. A.; Collins, M. T.; Mankani, M. H.; Corsi, A.; Bone, H. G.; Wientroub, S.; Spiegel, A. M.; Fisher, L. W.; Robey, P. G.: Mutations of the GNAS1 gene, stromal cell dysfunction, and osteomalacic changes in non-McCune-Albright fibrous dysplasia of bone. *J. Bone Miner. Res.* 15: 120-128, 2000.
- [79843] 18173. Billestrup, N.; Swanson, L. W.; Vale, W.: Growth hormone-releasing factor stimulates proliferation of somatotrophs in vitro. *Proc. Nat. Acad. Sci.* 83: 6854-6857, 1986.
- [79844] 18174. Candelieri, G. A.; Glorieux, F. H.; Prud'Homme, J.; St.-Arnaud, R.: Increased expression of the c-fos proto-oncogene in bone from patients with fibrous dysplasia. *New Eng. J. Med.* 332: 1546-1551, 1995.
- [79845] 18175. Candelieri, G. A.; Roughley, P. J.; Glorieux, F. H.:

Polymerase chain reaction-based technique for the selective enrichment and analysis of mosaic *arg201* mutations in *Gαs* from patients with fibrous dysplasia of bone. *Bone* 21: 201–206, 1997.

[79846] 18176. Carel, J. C.; Le Stunff, C.; Condamine, L.; Mallet, E.; Chaussain, J. L.; Adnot, P.; Garabedian, M.; Bougneres, P.: Resistance to the lipolytic action of epinephrine: a new feature of protein GS deficiency. *J. Clin. Endocr. Metab.* 84: 4127–4131, 1999.

[79847] 18177. Carter, A.; Bardin, C.; Collins, R.; Simons, C.; Bray, P.; Spiegel, A.: Reduced expression of multiple forms of the  $\alpha$  subunit of the stimulatory GTP-binding protein in pseudohypoparathyroidism type Ia. *Proc. Nat. Acad. Sci.* 84: 7266–7269, 1987.

[79848] 18178. Cattanaach, B. M.; Kirk, M.: Differential activity of maternally and paternally derived chromosome regions in mice. *Nature* 315: 496–498, 1985.

[79849] 18179. Coutant, R.; Lumbroso, S.; Rey, R.; Lahlou, N.; Venara, M.; Rouleau, S.; Sultan, C.; Limal, J.-M.: Macroorchidism due to autonomous hyperfunction of Sertoli cells and *GS-α* gene mutation: an unusual expression of McCune–Albright syndrome in a prepubertal boy. *J. Clin. Endocr. Metab.* 86: 1778–1781, 2001.

- [79850] 18180.Kim, H.-S.; Nagalla, S. R.; Oh, Y.; Wilson, E.; Roberts, C. T., Jr.; Rosenfeld, R. G.: Identification of a family of low-affinity insulin-like growth factor binding proteins (IGFBPs): characterization of connective tissue growth factor as a member of the IGFBP superfamily. *Proc. Nat. Acad. Sci.* 94: 12981–12986, 1997.
- [79851] 18181.Martinerie, C.; Viegas-Pequignot, E.; Guenard, I.; Dutrillaux, B.; Nguyen, V. C.; Bernheim, A.; Perbal, B.: Physical mapping of human loci homologous to the chicken *nov* proto-oncogene. *Oncogene* 7:2529–2534, 1992.
- [79852] 18182.Nakanishi, T.; Yamaai, T.; Asano, M.; Nawachi, K.; Suzuki, M.; Sugimoto, T.; Takigawa, M.: Overexpression of connective tissue growth factor/hypertrophic chondrocyte-specific gene product 24 decreases bone density in adult mice and induces dwarfism. *Biochem. Biophys. Res. Commun.* 281: 678–681, 2001.
- [79853] 18183.O'Donovan, N.; Galvin, M.; Morgan, J. G.: Physical mapping of the CXC chemokine locus on human chromosome 4. *Cytogenet. Cell Genet.* 84:39–42, 1999.
- [79854] 18184.McGill, J. R.; Boyd, D.; Barrett, K. J.; Drysdale, J. W.; Moore, C. M.: Localization of human ferritin H (heavy) and L (light) subunits by *in situ* hybridization. (Abstract) *Am. J.*

Hum. Genet. 36: 146Sonly, 1984.

- [79855] 18185.Murray, M. T.; White, K.; Munro, H. N.: Conservation of ferritinheavy subunit gene structure: implications for the regulation of ferritingene expression. Proc. Nat. Acad. Sci. 84: 7438–7442, 1987.
- [79856] 18186.Papadopoulos, P.; Bhavsar, D.; Zappone, E.; David, V.; Jones,C.; Worwood, M.; Drysdale, J.: A second human ferritin H locus onchromosome 11. Cytogenet. Cell Genet. 61: 107–108, 1992.
- [79857] 18187.Richard, C. W.; Withers, D. A.; Meeker, T. C.; Myers, R. M.:A radiation hybrid map of the proximal long arm of human chromosome11 containing the MEN–1 and bcl–1 disease locus. (Abstract) Cytogenet.Cell Genet. 58: 1970 only, 1991.
- [79858] 18188.Worwood, M.; Brook, J. D.; Cragg, S. J.; Hellkuhl, B.; Jones,B. M.; Perera, P.; Roberts, S. H.; Shaw, D. J.: Assignment of humanferritin genes to chromosomes 11 and 19q13.3–19qter. Hum. Genet. 69:371–374, 1985.
- [79859] 18189.Wu, K.–J.; Polack, A.; Dalla–Favera, R.: Coordinated regulationof iron–controlling genes, H–ferritin and IRP2, by c–MYC. Science 283:676–679, 1999.
- [79860] 18190.Yachou, A.; Mattei, M. G.; Roeckel, N.; Grand–champ, B.; Beaumont,C.: Mouse ferritin H sequences map

to chromosomes 3, 6, and 19. *Genomics* 9:204–206, 1991.

- [79861] 18191.Yachou, A.–K.; Renaudie, F.; Guenet, J.–L.; Simon–Chazottes, D.; Jones, R.; Grandchamp, B.; Beaumont, C.: Mouse ferritin H multigenefamily is polymorphic and contains a single multiallelic functional gene located on chromosome 19. *Genomics* 10: 531–538, 1991.
- [79862] 18192.Youssoufian, H.; Chance, P.; Tuck–Muller, C. M.; Jabs, E. W.: Association of a new chromosomal deletion [del(1)(q32q42)] with diaphragmatic hernia: assignment of a human ferritin gene. *Hum. Genet.* 78: 267–270, 1988.
- [79863] 18193.Aguilar–Martinez, P.; Biron, C.; Masmejean, C.; Jeanjean, P.; Schved, J.–F.: A novel mutation in the iron responsive element of ferritin L–subunit gene as a cause for hereditary hyperferritinemia–cataract syndrome. (Letter) *Blood* 88: 1895–1903, 1996.
- [79864] 18194.Beaumont, C.; Leneuve, P.; Devaux, I.; Scoazec, J.–Y.; Berthier, M.; Loiseau, M.–N.; Grandchamp, B.; Bonneau, D.: Mutation in the iron responsive element of the L ferritin mRNA in a family with dominant hyperferritinaemia and cataract. *Nature Genet.* 11: 444–446, 1995.
- [79865] 18195.Brown, A. J. P.; Leibold, E. A.; Munro, H. N.: Isolation of cDNA clones for the light subunit of rat liver fer–



ritin: evidence that the light subunit is encoded by a multi-gene family. *Proc. Nat. Acad. Sci.* 80: 1265–1269, 1983.

[79866] 18196. Camaschella, C.; Zecchina, G.; Lockitch, G.; Roetto, A.; Campanella, A.; Arosio, P.; Levi, S.: A new mutation (G51C) in the iron-responsive element (IRE) of L-ferritin associated with hyperferritinaemia-cataracts syndrome decreases the binding affinity of the mutated IRE for iron-regulatory proteins. *Brit. J. Haemat.* 108: 480–482, 2000.

[79867] 18197. Cazzola, M.; Bergamaschi, G.; Tonon, L.; Arbustini, E.; Grasso, M.; Vercesi, E.; Barosi, G.; Bianchi, P. E.; Cairo, G.; Arosio, P.: Hereditary hyperferritinemia-cataract syndrome: relationship between phenotypes and specific mutations in the iron-responsive element of ferritin light-chain mRNA. *Blood* 90: 814–821, 1997.

[79868] 18198. Cazzola, M.; Foglieni, B.; Bergamaschi, G.; Levi, S.; Lazzarino, M.; Arosio, P.: A novel deletion of the L-ferritin iron-responsive element responsible for severe hereditary hyperferritinaemia-cataracts syndrome. *Brit. J. Haemat.* 116: 667–670, 2002.

[79869] 18199. Xu, W.; Gorman, P. A.; Rider, S. H.; Hedge, P. J.; Moore, G.; Prichard, C.; Sheer, D.; Solomon, E.: Construction of a genetic map of human chromosome 17 by use of chromosome-mediated gene transfer. *Proc. Nat. Acad. Sci.*

85: 8563–8567, 1988.

- [79870] 18200.Fishman, G. I.; Eddy, R. L.; Shows, T. B.; Rosenthal, L.; Leinwand, L. A.: The human connexin gene family of gap junction proteins: distinct chromosomal locations but similar structures. *Genomics* 10: 250–256, 1991.
- [79871] 18201.Campbell, H. D.; Webb, G. C.; Kono, T.; Taniguchi, T.; Ford, J.H.; Young, I. G.: Assignment of the interleukin-2 receptor beta chain gene (IL-2rb) to band E on mouse chromosome 15. *Genomics* 12: 179–180, 1992.
- [79872] 18202.Bonneau, D.; Huret, J. L.; Godeau, G.; Couet, D.; Putterman, M.; Tanzer, J.; Babin, P.; Larregue, M.: Recurrent ctb(7)(q31.3) and possible laminin involvement in a neonatal cutis laxa with a Marfan phenotype. *Hum. Genet.* 87: 317–319, 1991.
- [79873] 18203.Buchanan, R.; Wyatt, G. P.: Marfan's syndrome presenting as an intrapartum death. *Arch. Dis. Child.* 60: 1074–1076, 1985.
- [79874] 18204.Burgeson, R. E.; Chiquet, M.; Deutzmann, R.; Ekblom, P.; Engel, J.; Kleinman, H.; Martin, G. R.; Meneguzzi, G.; Paulsson, M.; Sanes, J.; Timpl, R.; Tryggvason, K.; Yamada, Y.; Yurchenco, P. D.: A new nomenclature for the laminins. *Matrix Biol.* 14: 209–211, 1994.
- [79875] 18205.Day, D. L.; Burke, B. A.: Pulmonary emphysema in a

neonate with Marfan syndrome. *Pediatr. Radiol.* 16:  
518–521, 1986.

- [79876] 18206. Elliott, R. W.; Barlow, D.; Hogan, B. L. M.: Linkage of genes for laminin B1 and B2 subunits on chromosome 1 in mouse. *In Vitro Cell Dev. Biol.* 21: 477–484, 1985.
- [79877] 18207. Gross, D. M.; Robinson, L. K.; Smith, L. T.; Glass, N.; Rosenberg, H.; Duvic, M.: Severe perinatal Marfan syndrome. *Pediatrics* 84:83–89, 1989.
- [79878] 18208. Hohn, A. R.; Webb, H. M.: Cardiac studies of infant twins with Marfan's syndrome. *Am. J. Dis. Child.* 122: 526–528, 1971.
- [79879] 18209. Hopker, V. H.; Shewan, D.; Tessier-Lavigne, M.; Poo, M.; Holt, C.: Growth-cone attraction to netrin-1 is converted to repulsion by laminin-1. *Nature* 401: 69–73, 1999.
- [79880] 18210. Huret, J.; Bonneau, D.; Godeau, G.; Leheup, B.; Laroche, M.: Neonatal cutis laxa with Marfan habitus: this syndrome maps with B1 laminin at the junction 7q31.3–q32. (Abstract) *Cytogenet. Cell Genet.* 58:1922 only, 1991.
- [79881] 18211. Iwamoto, Y.; Robey, F. A.; Graf, J.; Sasaki, M.; Kleinman, H. K.; Yamada, Y.; Martin, G. R.: YIGSR, a synthetic laminin pentapeptide, inhibits experimental metas-

tasis formation. Science 238: 1132–1134,1987.

- [79882] 18212.Jaye, M.; Modi, W. S.; Ricca, G. A.; Mudd, R.; Chiu, I.-M.; O'Brien,S. J.; Drohan, W. N.: Isolation of a cDNA clone for the human laminin-B1chain and its gene localization. Am. J. Hum. Genet. 41: 605–615,1987.
- [79883] 18213.Kleinman, H. K.: Personal Communication. Bethesda, Md. 1/7/1982.
- [79884] 18214.Lababidi, Z.; Monzon, C.: Early cardiac manifestations of Marfan'ssyndrome in the newborn. Am. Heart J. 102: 943–945, 1981.
- [79885] 18215.Modi, W. S.; Jaye, M.; O'Brien, S. J.: Chromosomal localizationof a cDNA clone for the human B1 laminin chain. (Abstract) Cytogenet.Cell Genet. 46: 663 only, 1987.
- [79886] 18216.Neimann, N.; Rauber, G.; Marchal, C.; Vidailhet, M.; Fall, M.: Maladie de Marfan chez un nouveau-ne avec atteintes polyviscerales:etude anatomo-clinique. Ann. Pae-diat. 15: 619–624, 1968.
- [79887] 18217.Pikkarainen, T.; Eddy, R.; Fukushima, Y.; Byers, M.; Shows, T.;Pihlajaniemi, T.; Saraste, M.; Tryggvason, K.: Human laminin B1 chain:a multidomain protein with gene (LAMB1) locus in the q22 region ofchromosome 7. J. Biol. Chem. 262: 10454–10462, 1987.

- [79888] 18218. Pikkarainen, T.; Savolainen, E.-R.; Tryggvason, K.: Nhe I and Hinc II polymorphisms in the human laminin B1 chain gene on 7q22. *Nucleic Acids Res.* 17: 4424 only, 1989.
- [79889] 18219. Sasaki, M.; Kato, S.; Kohno, K.; Martin, G. R.; Yamada, Y.: Sequence of the cDNA encoding the laminin B1 chain reveals a multidomain protein containing cysteine-rich repeats. *Proc. Nat. Acad. Sci.* 84: 935–939, 1987.
- [79890] 18220. Vuolteenaho, R.; Chow, L. T.; Tryggvason, K.: Structure of the human laminin B1 chain gene. *J. Biol. Chem.* 265: 15611–15616, 1990.
- [79891] 18221. Gremlich, S.; Porret, A.; Hani, E. H.; Cherif, D.; Vionnet, N.; Froguel, P.; Thorens, B.: Cloning, functional expression, and chromosomal localization of the human pancreatic islet glucose-dependent insulinotropic polypeptide receptor. *Diabetes* 44: 1202–1208, 1995.
- [79892] 18222. Miyawaki, K.; Yamada, Y.; Ban, N.; Ihara, Y.; Tsukiyama, K.; Zhou, H.; Fujimoto, S.; Oku, A.; Tsuda, K.; Toyokuni, S.; Hiai, H.; Mizunoya, W.; and 9 others: Inhibition of gastric inhibitory polypeptide signaling prevents obesity. *Nature Med.* 8: 738–742, 2002.
- [79893] 18223. Stoffel, M.; Fernald, A. A.; Le Beau, M. M.; Bell, G. I.: Assignment of the gastric inhibitory polypeptide receptor

gene (GIPR) to chromosome bands 19q13.2–q13.3 by fluorescence in situ hybridization. *Genomics* 28:607–609, 1995.

- [79894] 18224. Usdin, T. B.; Mezey, E.; Button, D. C.; Brownstein, M. J.; Bonner, T. I.: Gastric inhibitory polypeptide receptor, a member of the secretin–vasoactive intestinal peptide receptor family, is widely distributed in peripheral organs and the brain. *Endocrinology* 133: 2861–2870, 1993.
- [79895] 18225. Volz, A.; Groke, R.; Lankat–Buttgereit, B.; Fehmann, H.–C.; Bode, H. P.; Goke, B.: Molecular cloning, functional expression, and signal transduction of the GIP–receptor cloned from a human insulinoma. *FEBS Lett.* 373: 23–29, 1995.
- [79896] 18226. Yamada, Y.; Hayami, T.; Nakamura, K.; Kaisaki, P. J.; Someya, Y.; Wang, C.–Z.; Seino, S.; Seino, Y.: Human gastric inhibitory polypeptide receptor: cloning of the gene (GIPR) and cDNA. *Genomics* 29: 773–776, 1995.
- [79897] 18227. Miyawaki, K.; Yamada, Y.; Yano, H.; Niwa, H.; Ban, N.; Ihara, Y.; Kubota, A.; Fujimoto, S.; Kajikawa, M.; Kuroe, A.; Tsuda, K.; Hashimoto, H.; Yamashita, T.; Jomori, T.; Tashiro, F.; Miyazaki, J.; Seino, Y.: Glucose intolerance caused by a defect in the entero–insular axis: a study in gastric inhibitory polypeptide receptor knockout mice.

Proc.Nat. Acad. Sci. 96: 14843–14847, 1999.

- [79898] 18228.Flejter, W. L.; Barcroft, C. L.; Guo, S.–W.; Lynch, E. D.; Boehnke,M.; Chandrasekharappa, S.; Hayes, S.; Collins, F. S.; Weber, B. L.;Glover, T. W.: Multicolor FISH mapping with Alu–PCR–amplified YACclone DNA determines the order of markers in the BRCA1 region on chromosome17q12–q21. Genomics 17: 624–631, 1993.
- [79899] 18229.Fukushige, S.; Murotsu, T.; Matsubara, K.: Chromosomal assignmentof human genes for gastrin, thyrotropin (TSH)–beta subunit and C–erb–2by chromosome sorting combined with velocity sedimentation and southernhybridization. Biochem. Biophys. Res. Commun. 134: 477–483, 1986.
- [79900] 18230.Justice, M. J.; Gilbert, D. J.; Kinzler, K. W.; Vogelstein, B.;Buchberg, A. M.; Ceci, J. D.; Matsuda, Y.; Chapman, V. M.; Patriotis,C.; Makris, A.; Tsiichlis, P. N.; Jenkins, N. A.; Copeland, N. G.:A molecular genetic linkage map of mouse chromosome 18 reveals extensivelinkage conservation with human chromosomes 5 and 18. Genomics 13:1281–1288, 1992.
- [79901] 18231.Lebacq–Verheyden, A.–M.; Bertness, V.; Kirsch, I.; Hollis, G. F.;McBride, O. W.; Battey, J.: Human gastrin–releasing peptide genemaps to chromosome band 18q21.

Somat. Cell Molec. Genet. 13: 81–86,1987.

[79902] 18232.Naylor, S. L.; Sakaguchi, A. Y.; Spindel, E.; Chin, W. W.: Humangastrin–releasing peptide gene is located on chromosome 18. Somat.Cell Molec. Genet. 13: 87–91, 1987.

[79903] 18233.Naylor, S. L.; Spindel, E.; Chin, W. W.; Sakaguchi, A. Y.: Gastrinreleasing peptide gene is located on human chromosome 18. (Abstract) Cytogenet.Cell Genet. 40: 711 only, 1985.

[79904] 18234.Sausville, E. A.; Lebacq–Verheyden, A.–M.; Spindel, E. R.; Cuttitta,F.; Gazdar, A. F.; Battey, J. F.: Expression of the gastrin–releasingpeptide gene in human small cell lung cancer: evidence for alternativeprocessing resulting in three distinct mRNAs. J. Biol. Chem. 261:2451–2457, 1986.

[79905] 18235.Spindel, E. R.; Chin, W. W.; Price, J.; Rees, L. H.; Besser, G.M.; Habener, J. F.: Cloning and characterization of cDNAs encodinghuman gastrin–releasing peptide. Proc. Nat. Acad. Sci. 81: 5699–5703,1984.

[79906] 18236.Spindel, E. R.; Zilberberg, M. D.; Habener, J. F.; Chin, W. W.: Two prohormones for gastrin–releasing peptide are encoded by twomRNAs differing by 19 nucleotides. Proc. Nat. Acad. Sci. 83: 19–23,1986.



- [79907] 18237.Kudo, S.; Chagnovich, D.; Rearden, A.; Mattei, M. G.; Fukuda, M.: Molecular analysis of a hybrid gene encoding human glycophorin variantMiltenberger V-like molecule. J. Biol. Chem. 265: 13825–13829, 1990.
- [79908] 18238.Kudo, S.; Fukuda, M.: Identification of a novel human glycophorin, glycophorin E, by isolation of genomic clones and complementary DNA clones utilizing polymerase chain reaction. J. Biol. Chem. 265:1102–1110, 1990.
- [79909] 18239.Onda, M.; Fukuda, M.: Detailed physical mapping of the genes encoding glycophorins A, B, and E, as revealed by P1 plasmids containing human genomic DNA. Gene 159: 225–230, 1995.
- [79910] 18240.Onda, M.; Kudo, S.; Rearden, A.; Mattei, M.-G.; Fukuda, M.: Identification of a precursor genomic segment that provided a sequence unique to glycophorin B and E genes. Proc. Nat. Acad. Sci. 90: 7220–7224, 1993.
- [79911] 18241.Vignal, A.; Rahuel, C.; London, J.; Cherif Zahar, B.; Schaff, S.; Hattab, C.; Okubo, Y.; Cartron, J.-P.: A novel gene member of the human glycophorin A and B gene family: molecular cloning and expression. Europ. J. Biochem. 191: 619–625, 1990.
- [79912] 18242.Board, P. G.; Jones, I. M.; Bentley, A. K.: Molecular

cloning and nucleotide sequence of human alpha-1 acid glycoprotein cDNA. *Gene* 44:127-131, 1986.

- [79913] 18243. Cox, D. W.; Francke, U.: Direct assignment of orosomucoid to human chromosome 9 and alpha-2-HS-glycoprotein to chromosome 3 using human fetal liver x rat hepatoma hybrids. *Hum. Genet.* 70: 109-115, 1985.
- [79914] 18244. Dayhoff, M. O.: *Atlas of Protein Sequence and Structure. Orosomucoid.* Washington: National Biomedical Research Foundation (pub.) 5: 1972. Pp. D310-D316.
- [79915] 18245. Dente, L.; Ciliberto, G.; Cortese, R.: Structure of the human alpha-1-acid glycoprotein gene: sequence homology with other human acute phase protein genes. *Nucleic Acids Res.* 13: 3941-3952, 1985.
- [79916] 18246. Dente, L.; Pizza, M. G.; Metspalu, A.; Cortese, R.: Structure and expression of the genes coding for human alpha-1-acid glycoprotein. *EMBO J.* 6: 2289-2296, 1987.
- [79917] 18247. Dente, L.; Ruther, U.; Tripodi, M.; Wagner, E. F.; Cortese, R.: Expression of human alpha-1-acid glycoprotein genes in cultured cells and in transgenic mice. *Genes Dev.* 2: 259-266, 1988.
- [79918] 18248. Eap, C. B.; Cuendet, C.; Baumann, P.: Orosomucoid (alpha-1 acid glycoprotein) phenotyping by use of immo-

bilized pH gradients with 8 M urea and immunoblotting: a new variant encountered in a population study. Hum. Genet. 80: 183–185, 1988.

[79919] 18249. Eiberg, H.; Mohr, J.; Nielsen, L. S.: Linkage of orosomucoid (ORM) to ABO and AK1. (Abstract) Cytogenet. Cell Genet. 32: 272 only, 1982.

[79920] 18250. Johnson, A. M.; Schmid, K.; Alper, C. A.: Inheritance of human  $\alpha$ (1)-acid glycoprotein (orosomucoid) variants. J. Clin. Invest. 48: 2293–2299, 1969.

[79921] 18251. Luckenbach, C.; Kompf, J.; Ritter, H.: Orosomucoid (ORM1) subtyping and formal genetics. Hum. Genet. 87: 429–432, 1991.

[79922] 18252. Rocchi, M.; Roncuzzi, L.; Santamaria, R.; Archidiacono, N.; Dente, L.; Romeo, G.: Mapping through somatic cell hybrids and cDNA probes of protein C to chromosome 2, factor X to chromosome 13, and  $\alpha$ -1-acid glycoprotein to chromosome 9. Hum. Genet. 74: 30–33, 1986.

[79923] 18253. Schmid, K.; Tokita, K.; Yoshizaki, H.: The  $\alpha$ -1-acid glycoprotein variants of normal Caucasian and Japanese individuals. J. Clin. Invest. 44: 1394–1401, 1965.

[79924] 18254. Tokita, K.; Schmid, K.: Variants of  $\alpha$ -1-acid glycoprotein. Nature 200: 266 only, 1963.

- [79925] 18255.Tomei, L.; Eap, C. B.; Baumann, P.; Dente, L.: Use of transgenic mice for the characterization of human alpha-1-acid glycoprotein (orosomucoid) variants. Hum. Genet. 84: 89-91, 1989.
- [79926] 18256.Umetsu, K.; Ikeda, N.; Kashimura, S.; Suzuki, T.: Orosomucoid(ORM) typing by print lectinofixation: a new technique for isoelectric focusing--two common alleles in Japan. Hum. Genet. 71: 223-224, 1985.
- [79927] 18257.Umetsu, K.; Yuasa, I.; Nishimura, H.; Sasaki, H.; Suzuki, T.: Genetic polymorphisms of orosomucoid and alpha-2-HS-glycoprotein in a Philippine population. Hum. Hered. 38: 287-290, 1988.
- [79928] 18258.Webb, G. C.; Earle, M. E.; Merritt, C.; Board, P. G.: Localization of human alpha-1 glycoprotein genes to 9q31-q34.1. Cytogenet. Cell Genet. 47: 18-21, 1988.
- [79929] 18259.Weidinger, S.; Muller, T.; Schwarzfischer, F.; Cleve, H.: Three new orosomucoid (ORM) variants revealed by isoelectric focusing and print immunofixation. Hum. Genet. 77: 286-288, 1987.
- [79930] 18260.Yuasa, I.; Umetsu, K.; Vogt, U.; Nakamura, H.; Nanba, E.; Tamaki, N.; Irizawa, Y.: Human orosomucoid polymorphism: molecular basis of the three common ORM1 alleles, ORM1\*F1, ORM1\*F2, and ORM1\*S.

Hum.Genet. 99: 393–398, 1997.

- [79931] 18261.Yuasa, I.; Weidinger, S.; Umetsu, K.; Suenaga, K.; Ishimoto, G.;Eap, B. C.; Duche, J.–C.; Baumann, P.: Orosomucoid system: 17 additionalorosomucoid variants and proposal for a new nomenclature. Vox Sang. 64:47–55, 1993.
- [79932] 18262.Triantafilou, K.; Triantafilou, M.; Dedrick, R. L.: A CD14–independentLPS receptor cluster. Nature Immun. 2: 338–345, 2001.
- [79933] 18263.Saito, S.; Okui, K.; Tokino, T.; Oshimura, M.; Nakamura, Y.: Isolationand mapping of 68 RFLP markers on human chromosome 6. Am. J. Hum.Genet. 50: 65–70, 1992.
- [79934] 18264.Takahashi, I.; Tanuma, R.; Hirata, M.; Hashimoto, K.: A cosmidclone at the D6S182 locus on human chromosome 6p12 contains the 90–kDaheat shock protein beta–gene (HSP90–beta). Mammalian Genome 5: 121–122,1994.
- [79935] 18265.Durkin, A. S.; Maglott, D. R.; Vamvakopoulos, N. C.; Zoghbi, H.Y.; Nierman, W. C.: Assignment of an intron–containing human heat–shockprotein gene (hsp90–beta, HSPCB) to chromosome 6 near TCTE1 (6p21)and two intronless pseudogenes to chromosomes 4 and 15 by poly–

merase chain reaction amplification from a panel of hybrid cell lines. *Genomics* 18:452–454, 1993.

- [79936] 18266.Klinger, H. P.: Suppression of tumorigenicity in somatic cell hybrids. I. Suppression and reexpression of tumorigenicity in diploid human x D98/AH2 hybrids and independent segregation of tumorigenicity from other cell phenotypes. *Cytogenet. Cell Genet.* 27: 254–266, 1980.
- [79937] 18267.Lichy, J. H.; Modi, W. S.; Seuanez, H. N.; Howley, P. M.: Identification of a human chromosome 11 gene which is differentially regulated in tumorigenic and nontumorigenic somatic cell hybrids of HeLa cells. *Cell Growth Differ.* 3: 541–548, 1992.
- [79938] 18268.Stanbridge, E. J.: Suppression of malignancy in human cells. *Nature* 260:17–20, 1976.
- [79939] 18269.Corrigan, V. M.; Arastu, M.; Khan, S.; Shah, C.; Fife, M.; Smeets, T.; Tak, P.-P.; Panayi, G. S.: Functional IL-2 receptor beta (CD122) and gamma (CD132) chains are expressed by fibroblast-like synoviocytes: activation by IL-2 stimulates monocyte chemoattractant protein-1 production. *J. Immun.* 166: 4141–4147, 2001.
- [79940] 18270.Gnarra, J. R.; Otani, H.; Wang, M. G.; McBride, O. W.; Sharon, M.; Leonard, W. J.: Human interleukin 2 receptor beta-chain gene: chromosomal localization and identi-

fication of 5-prime regulatory sequences. Proc. Nat. Acad. Sci. 87: 3440–3444, 1990.

- [79941] 18271. Hatakeyama, M.; Tsudo, M.; Minamoto, S.; Kono, T.; Doi, T.; Miyata, T.; Miyasaka, M.; Taniguchi, T.: Interleukin-2 receptor beta chain gene: generation of three receptor forms by cloned human alpha and beta chain cDNA's. Science 244: 551–556, 1989.
- [79942] 18272. Lamaze, C.; Dujancourt, A.; Baba, T.; Lo, C. G.; Benmerah, A.; Dautry-Varsat, A.: Interleukin 2 receptors and detergent-resistant membrane domains define a clathrin-independent endocytic pathway. Molec. Cell 7: 661–671, 2001.
- [79943] 18273. Shibuya, H.; Yoneyama, M.; Nakamura, Y.; Harada, H.; Hatakeyama, M.; Minamoto, S.; Kono, T.; Doi, T.; White, R.; Taniguchi, T.: The human interleukin-2 receptor beta-chain gene: genomic organization, promoter analysis and chromosomal assignment. Nucleic Acids Res. 18: 3697–3703, 1990.
- [79944] 18274. Suzuki, H.; Kundig, T. M.; Furlonger, C.; Wakeham, A.; Timms, E.; Matsuyama, T.; Schmits, R.; Simard, J. J. L.; Ohashi, P. S.; Griesser, H.; Taniguchi, T.; Paige, C. J.; Mak, T. W.: Deregulated T cell activation and autoimmunity in mice lacking interleukin-2 receptor beta. Science

268:1472–1476, 1995.

- [79945] 18275. Tsudo, M.; Kitamura, F.; Miyasaka, M.: Characterization of the interleukin 2 receptor beta chain using three distinct monoclonal antibodies. *Proc. Nat. Acad. Sci.* 86: 1982–1986, 1989.
- [79946] 18276. Alitalo, T.; Koistinen, R.; Aalto-Setälä, K.; Kontula, K.; Julkunen, M.; Janne, O.; de la Chapelle, A.: Gene for the human insulin-like growth factor-binding protein/placental protein 12: mapping to 7p12–p13 and demonstration of a BglII RFLP. (Abstract) *Cytogenet. Cell Genet.* 51:950 only, 1989.
- [79947] 18277. Alitalo, T.; Kontula, K.; Koistinen, R.; Aalto-Setälä, K.; Julkunen, M.; Janne, O. A.; Seppälä, M.; de la Chapelle, A.: The gene encoding human low-molecular weight insulin-like growth-factor binding protein (IGF-BP25): regional localization to 7p12–p13 and description of a DNA polymorphism. *Hum. Genet.* 83: 335–338, 1989.
- [79948] 18278. Ballard, F. J.; Baxter, R. C.; Binoux, M.; Clemmons, D. R.; Drop, S. L. S.; Hall, K.; Hintz, R. L.; Rechler, M. M.; Rutanen, E. M.; Schwander, J. C.: Report on the nomenclature of the IGF binding proteins. *J. Clin. Endocr. Metab.* 70: 817–818, 1990.
- [79949] 18279. Brewer, M. T.; Stetler, G. L.; Squires, C. H.; Thomp-



son, R. C.; Busby, W. H.; Clemmons, D. R.: Cloning, characterization, and expression of a human insulin-like growth factor binding protein. *Biochem. Biophys. Res. Commun.* 152: 1289–1297, 1988.

[79950] 18280. Brinkman, A.; Groffen, C.; Kortleve, D. J.; Geurts van Kessel, A.; Drop, S. L. S.: Isolation and characterization of a cDNA encoding the low molecular weight insulin-like growth factor binding protein (IBP-1). *EMBO J.* 7: 2417–2423, 1988.

[79951] 18281. Brinkman, A.; Groffen, C. A. H.; Kortleve, D. J.; Drop, S. L. S.: Organization of the gene encoding the insulin-like growth factor binding protein IBP-1. *Biochem. Biophys. Res. Commun.* 157: 898–907, 1988.

[79952] 18282. Ekstrand, J.; Ehrenborg, E.; Stern, I.; Stellan, B.; Zech, L.; Luthman, H.: The gene for insulin-like growth factor binding protein-1 is localized to human chromosomal region 7p14–p12. *Genomics* 6: 413–418, 1990.

[79953] 18283. Popovici, R. M.; Lu, M.; Bhatia, S.; Faessen, G. H.; Giaccia, A. J.; Giudice, L. C.: Hypoxia regulates insulin-like growth factor-binding protein 1 in human fetal hepatocytes in primary culture: suggestive molecular mechanisms for in utero fetal growth restriction caused by uteroplacental insufficiency. *J. Clin. Endocr. Metab.* 86:

2653–2659,2001.

- [79954] 18284. Agarwal, N.; Hsieh, C. L.; Sills, D.; Swaroop, M.; Desai, B.; Francke, U.; Swaroop, A.: Sequence analysis, expression and chromosomal localization of a gene, isolated from a subtracted human retina cDNA library, that encodes an insulin-like growth factor binding protein (IGFBP2). *Exp. Eye Res.* 52: 549–561, 1991.
- [79955] 18285. Allander, S. V.; Larsson, C.; Ehrenborg, E.; Suwanichkul, A.; Weber, G.; Morris, S. L.; Bajalica, S.; Kiefer, M. C.; Luthman, H.; Powell, D. R.: Characterization of the chromosomal gene and promoter for human insulin-like growth factor binding protein-5. *J. Biol. Chem.* 269:10891–10898, 1994.
- [79956] 18286. Ehrenborg, E.; Vilhelmsdotter, S.; Bajalica, S.; Larsson, C.; Stern, I.; Koch, J.; Brondum-Nielsen, K.; Luthman, H.: Structure and localization of the human insulin-like growth factor-binding protein 2 gene. *Biochem. Biophys. Res. Commun.* 176: 1250–1255, 1991.
- [79957] 18287. Wood, W. I.; Cachianes, G.; Henzel, W. J.; Winslow, G. A.; Spencer, S. A.; Hellmiss, R.; Martin, J. L.; Baxter, R. C.: Cloning and expression of the growth hormone-dependent insulin-like growth factor-binding protein. *Molec. Endocr.* 2: 1176–1185, 1988.

- [79958] 18288.Eklund, L.; Piuhola, J.; Komulainen, J.; Sormunen, R.; Ongvarrasopone,C.; Fassler, R.; Muona, A.; Ilves, M.; Ruskoaho, H.; Takala, T. E.S.; Pihlajaniemi, T.: Lack of type XV collagen causes a skeletalmyopathy and cardiovascular defects in mice. *Proc. Nat. Acad. Sci.* 98:1194–1199, 2001.
- [79959] 18289.Hagg, P. M.; Hagg, P. O.; Peltonen, S.; Autio-Harmainen, H.; Pihlajaniemi,T.: Location of type XV collagen in human tissues and its accumulationin the interstitial matrix of the fibrotic kidney. *Am. J. Path.* 150:2075–2086, 1997.
- [79960] 18290.Hagg, P. M.; Horelli-Kuitunen, N.; Eklund, L.; Palotie, A.; Pihlajaniemi,T.: Cloning of mouse type XV collagen sequences and mapping of thecorresponding gene to 4B1–3: comparison of mouse and human alpha-1(XV)collagen sequences indicates divergence in the number of small collagenousdomains. *Genomics* 45: 31–41, 1997.
- [79961] 18291.Hagg, P. M.; Muona, A.; Lietard, J.; Kivirikko, S.; Pihlajaniemi,T.: Complete exon–intron organization of the human gene for the alpha-1chain of type XV collagen (COL15A1) and comparison with the homologousCol18a1 gene. *J. Biol. Chem.* 273: 17824–17831, 1998.

- [79962] 18292.Huebner, K.; Cannizzaro, L. A.; Jabs, E. W.; Kivirikko, S.; Manzone,H.; Pihlajaniemi, T.; Myers, J. C.: Chromosomal assignment of a geneencoding a new collagen type (COL15A1) to 9q21–q22. *Genomics* 14:220–224, 1992.
- [79963] 18293.Kivirikko, S.; Heinamaki, P.; Rehn, M.; Honkanen, N.; Myers, J.C.; Pihlajaniemi, T.: Primary structure of the alpha–1 chain of humantype XV collagen and exon–intron organization in the 3–prime regionof the corresponding gene. *J. Biol. Chem.* 269: 4773–4779, 1994.
- [79964] 18294.Muragaki, Y.; Abe, N.; Ninomiya, Y.; Olsen, B. R.; Ooshima, A.: The human alpha–1(XV) collagen chain contains a large amino–terminalnon–triple helical domain with a tandem repeat structure and homologyto alpha–1(XVIII) collagen. *J. Biol. Chem.* 269: 4042–4046, 1994.
- [79965] 18295.Myers, J. C.; Dion, A. S.; Abraham, V.; Amenta, P. S.: Type XVcollagen exhibits a widespread distribution in human tissues but adistinct localization in basement membrane zones. *Cell Tissue Res.* 286:493–505, 1996.
- [79966] 18296.Myers, J. C.; Kivirikko, S.; Gordon, M. K.; Dion, A. S.; Pihlajaniemi,T.: Identification of a previously unknown human collagen chain,alpha–1(XV), characterized by extensive interruptions in the triple–helicalregion. *Proc. Nat.*

Acad. Sci. 89: 10144–10148, 1992.

[79967] 18297. Ramchandran, R.; Dhanabal, M.; Volk, R.; Waterman, M. J. F.; Segal, M.; Lu, H.; Knebelmann, B.; Sukhatme, V. P.: Antiangiogenic activity of restin, NC10 domain of human collagen XV: comparison to endostatin.

Biochem. Biophys. Res. Commun. 255: 735–739, 1999.

[79968] 18298. Rehn, M.; Hintikka, E.; Pihlajaniemi, T.: Primary structure of the alpha 1 chain of mouse type XVIII collagen, partial structure of the corresponding gene, and comparison of the alpha 1(XVIII) chain with its homologue, the alpha 1(XV) collagen chain. J. Biol. Chem.

269:13929–13935, 1994.

[79969] 18299. Sasaki, T.; Larsson, H.; Tisi, D.; Claesson-Welsh, L.; Hohenester, E.; Timpl, R.: Endostatins derived from collagens XV and XVIII differ in structural and binding properties, tissue distribution and anti-angiogenic activity. J.

Molec. Biol. 301: 1179–1190, 2000.

[79970] 18300. Pan, T.-C.; Zhang, R.-Z.; Mattei, M.-G.; Timpl, R.; Chu, M.-L.: Cloning and chromosomal location of human alpha-1(XVI) collagen. Proc. Nat. Acad. Sci. 89:

6565–6569, 1992.

[79971] 18301. Yamaguchi, N.; Kimura, S.; McBride, O. W.; Hori, H.; Yamada, Y.; Kanamori, T.; Yamakoshi, H.; Nagai, Y.: Molec-

ular cloning and partial characterization of a novel collagen chain, alpha-1(XVI), consisting of repetitive collagenous domains and cysteine-containing non-collagenous segments. J. Biochem. 112: 856-863, 1992.

[79972] 18302. Curtis, A. R. J.; Fey, C.; Morris, C. M.; Bindoff, L. A.; Ince, P. G.; Chinnery, P. F.; Coulthard, A.; Jackson, M. J.; Jackson, A. P.; McHale, D. P.; Hay, D.; Barker, W. A.; Markham, A. F.; Bates, D.; Curtis, A.; Burn, J.: Mutation in the gene encoding ferritin light polypeptide causes dominant adult-onset basal ganglia disease. Nature Genet. 28: 350-354, 2001.

[79973] 18303. Eiberg, H.; Bisgaard, M. L.; Mohr, J.: Linkage between alpha-1-B-glycoprotein (A1BG) and Lutheran (LU) red blood group system: assignment to chromosome 19: new genetic variants of A1BG. Clin. Genet. 36: 415-418, 1989.

[79974] 18304. Eiberg, H.; Nielsen, L. S.; Gahne, B.; Juneja, R. K.; Mohr, J.: Exclusion data for the alpha-1-B glycoprotein (A1BG) polymorphism. (Abstract) Cytogenet. Cell Genet. 51: 994 only, 1989.

[79975] 18305. Gahne, B.; Juneja, R. K.; Stratil, A.: Genetic polymorphism of human plasma alpha-1-B-glycoprotein: phenotyping by immunoblotting or by a simple method of 2-D

electrophoresis. Hum. Genet. 76: 111–115, 1987.

[79976] 18306. Ishioka, N.; Takahashi, N.; Putnam, F. W.: Amino acid sequence of human plasma alpha-1B-glycoprotein: homology to the immunoglobulin supergene family. Proc. Nat. Acad. Sci. 83: 2363–2367, 1986.

[79977] 18307. Juneja, R. K.; Weitkamp, L. R.; Stratil, A.; Gahne, B.; Guttormsen, S. A.: Further studies of the plasma alpha-1-B-glycoprotein polymorphism: two new alleles and allele frequencies in Caucasians and in American blacks. Hum. Hered. 38: 267–272, 1988.

[79978] 18308. Anderson, L.; Anderson, N. G.: High resolution two-dimensional electrophoresis of human plasma proteins. Proc. Nat. Acad. Sci. 74: 5421–5425, 1977.

[79979] 18309. Boutin, B.; Feng, S. H.; Arnaud, P.: The genetic polymorphism of alpha(2)-HS glycoprotein: study by ultra-thin-layer isoelectric focusing and immunoblot. Am. J. Hum. Genet. 37: 1098–1105, 1985.

[79980] 18310. Cox, D. W.; Andrews, B. J.: Silver stain immunofixation for alpha-2-HS-glycoprotein: a new method for detection of protein heterogeneity. In: Stathakos, D.: Electrophoresis '82. Berlin: Walter de Gruyter (pub.) 1983. Pp. 243–247.

[79981] 18311. Cox, D. W.; Andrews, B. J.; Wills, D. E.: Genetic

polymorphism of alpha-2-HS-glycoprotein. Am. J. Hum. Genet. 38: 699–706, 1986.

[79982] 18312.Cox, D. W.; Francke, U.; Allderdice, P. W.; McAlpine, P. J.: Genemapping of human serum proteins using hepatoma hybrids and human chromosomal deletions and duplications. Genetics 107 (suppl.): s22–s23, 1984.

[79983] 18313.Eiberg, H.; Mohr, J.; Nielsen, L. S.: A2HS: new methods of phenotyping and analysis of linkage relations: assignment to chromosome 3. (Abstract) Cytogenet. Cell Genet. 37: 461 only, 1984.

[79984] 18314.Gejyo, F.; Chang, J.-L.; Burgi, W.; Schmid, K.; Offner, G. D.; Troxler, R. F.; Van Halbeek, H.; Dorland, L.; Gerwig, G. J.; Vliegenthart, J. F. G.: Characterization of the B-chain of human plasma alpha(2)HS-glycoprotein: the complete amino acid sequence and primary structure of its heteroglycan. J. Biol. Chem. 258: 4966–4971, 1983.

[79985] 18315.Jahnen-Dechent, W.: Personal Communication. Mainz, Germany 7/22/1998.

[79986] 18316.Jahnen-Dechent, W.; Schinke, T.; Trindl, A.; Muller-Esterl, W.; Sablitzky, F.; Kaiser, S.; Blessing, M.: Cloning and targeted deletion of the mouse fetuin gene. J. Biol. Chem. 272: 31496–31503, 1997.

[79987] 18317.Lee, C.-C.; Bowman, B. H.; Yang, F.: Human alpha-



2-HS-glycoprotein:the A and B chains with a connecting sequence are encoded by a single mRNA transcript. Proc. Nat. Acad. Sci. 84: 4403–4407, 1987.

- [79988] 18318.Hulsebos, T. J. M.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N.G.: The beta crystallin genes on human chromosome 22 define a new region of homology with mouse chromosome 5. Genomics 25: 574–576, 1995.
- [79989] 18319.Lampi, K. J.; Ma, Z.; Shih, M.; Shearer, T. R.; Smith, J. B.; Smith, D. L.; David, L. L.: Sequence analysis of beta-A3, beta-B3, and beta-A4 crystallins completes the identification of the major proteins in young human lens. J. Biol. Chem. 272: 2268–2275, 1997.
- [79990] 18320.van Rens, G.; Geurts van Kessel, A.; Bloemendal, H.: Localization of the beta-A4 crystallin (CRYBA4) gene on human chromosome 22 in the region q11.2–q13.1. (Abstract) Cytogenet. Cell Genet. 58: 2052 only, 1991.
- [79991] 18321.van Rens, G. L. M.; Geurts van Kessel, A. H. M.; Bloemendal, H.: Localization of the beta-A4-crystallin gene (CRYBA4) on human chromosome 22 in the region q11.2–q13.1. Cytogenet. Cell Genet. 61: 180–183, 1992.
- [79992] 18322.Brakenhoff, R. H.; Henskens, H. A. M.; van Rossum, M. W. P. C.; Lubsen, N. H.; Schoenmakers, J. G. G.: Activation of the gamma-E-crystallin pseudogene in the human

hereditary Coppock-like cataract. Hum. Molec.Genet. 3: 279–283, 1994.

[79993] 18323.den Dunnen, J. T.; Jongbloed, R. J. E.; Geurts van Kessel, A. H.M.; Schoenmakers, J. G. G.: Human lens gamma-crystallin sequences are located in the p12-qter region of chromosome 2. Hum. Genet. 70:217–221, 1985.

[79994] 18324.Hejtmancik, J. F.: The genetics of cataract: our vision becomes clearer. (Editorial) Am. J. Hum. Genet. 62: 520–525, 1998.

[79995] 18325.Heon, E.; Priston, M.; Schorderet, D. F.; Billingsley, G. D.; Girard, P. O.; Lubsen, N.; Munier, F. L.: The gamma-crystallins and human cataracts: a puzzle made clearer. Am. J. Hum. Genet. 65: 1261–1267, 1999. Note: Erratum: Am. J. Hum. Genet. 66: 753 only, 2000.

[79996] 18326.Klopp, N.; Favor, J.; Loster, J.; Lutz, R. B.; Neuhauser-Klaus, A.; Prescott, A.; Pretsch, W.; Quinlan, R. A.; Sandilands, A.; Vrensen, G. F. J. M.; Graw, J.: Three murine cataract mutants (Cat2) are defective in different gamma-crystallin genes. Genomics 52: 152–158, 1998.

[79997] 18327.Piatigorsky, J.: Lens crystallins and their gene families. Cell 38:620–621, 1984.

[79998] 18328.Shiloh, Y.; Donlon, T.; Bruns, G.; Breitman, M. L.; Tsui, L.-C.: Assignment of the human gamma-crystallin

gene cluster (CRYG) to the long arm of chromosome 2, region q33–36. *Hum. Genet.* 73: 17–19, 1986.

- [79999] 18329. Siezen, R. J.; Thomson, J. A.; Kaplan, E. D.; Benedek, G. B.: Human lens gamma-crystallins: isolation, identification, and characterization of the expressed gene products. *Proc. Nat. Acad. Sci.* 84: 6088–6092, 1987.
- [80000] 18330. Skow, L. C.; Donner, M. E.; Huang, S.-M.; Gardner, J. M.; Taylor, B. A.; Beamer, W. G.; Lalley, P. A.: Mapping of mouse gamma crystallin genes on chromosome 1. *Biochem. Genet.* 26: 557–570, 1988.
- [80001] 18331. Tsui, L.-C.; Breitman, M. L.; Meakin, S. O.; Willard, H. F.; Shiloh, Y.; Donlon, T.; Bruns, G.: Localization of the human gamma-crystallin gene cluster (CRYG) to the long arm of chromosome 2, region q33–q35. (Abstract) *Cytogenet. Cell Genet.* 40: 763–764, 1985.
- [80002] 18332. Vidal, S. M.; Epstein, D. J.; Malo, D.; Weith, A.; Vekemans, M.; Gros, P.: Identification and mapping of six microdissected genomic DNA probes to the proximal region of mouse chromosome 1. *Genomics* 14: 32–37, 1992.
- [80003] 18333. Fajans, S. S.; Bell, G. I.; Polonsky, K. S.: Molecular mechanisms and clinical pathophysiology of maturity-onset diabetes of the young. *New Eng. J. Med.* 345: 971–980, 2001.

- [80004] 18334.Lindner, T.; Gragnoli, C.; Furuta, H.; Cockburn, B. N.; Petzold,C.; Rietzsch, H.; Weiss, U.; Schulze, J.; Bell, G. I.: Hepatic functionin a family with a nonsense mutation (R154X) in the hepatocyte nuclearfactor-4-alpha/MODY1 gene. J. Clin. Invest. 100: 1400-1405, 1997.
- [80005] 18335.Moller, A. M.; Dalgaard, L. T.; Ambye, L.; Hansen, L.; Schmitz,O.; Hansen, T.; Pedersen, O.: A novel Phe75fsdelT mutation in thehepatocyte nuclear factor-4-alpha gene in a Danish pedigree with maturity-onsetdiabetes of the young. J. Clin. Endocr. Metab. 84: 367-369, 1999.
- [80006] 18336.Thomas, H.; Jaschkowitz, K.; Bulman, M.; Frayling, T. M.; Mitchell,S. M. S.; Roosen, S.; Lingott-Frieg, A.; Tack, C. J.; Ellard, S.;Ryffel, G. U.; Hattersley, A. T.: A distant up-stream promoter ofthe HNF-4-alpha gene connects the transcription factors involved inmaturity-onset diabetes of the young. Hum. Molec. Genet. 10: 2089-2097,2001.
- [80007] 18337.Joseph, L. J.; Le Beau, M. M.; Jamieson, G. A., Jr.; Acharya, S.;Shows, T. B.; Rowley, J. D.; Sukhatme, V. P.: Molecular cloning,sequencing, and mapping of EGR2, a human early growth response geneencoding a protein with 'zinc-binding finger' structure. Proc. Nat.Acad. Sci. 85: 7164-7168, 1988.

- [80008] 18338.Nagarajan, R.; Svaren, J.; Le, N.; Araki, T.; Watson, M.; Milbrandt,J.: EGR2 mutations in inherited neuropathies dominant–negativelyinhibit myelin gene expression. *Neuron* 30: 355–368, 2001.
- [80009] 18339.Timmerman, V.; De Jonghe, P.; Ceuterick, C.; De Vriendt, E.; Lofgren,A.; Nelis, E.; Warner, L. E.; Lupski, J. R.; Martin, J.–J.; Van Broeckhoven,C.: Novel missense mutation in the early growth response 2 gene associatedwith Dejerine–Sottas syndrome phenotype. *Neurology* 52: 1827–1832,1999.
- [80010] 18340.Warner, L. E.; Mancias, P.; Butler, I.; Lupski, J. R.: Mutationin the early growth response 2 (EGR2) transcription factor associatedwith recessive congenital hypomyelinating neuropathy (CHN). (Abstract) *Am.J. Hum. Genet.* 61 (suppl.): A350 only, 1997.
- [80011] 18341.Warner, L. E.; Mancias, P.; Butler, I. J.; McDonald, C. M.; Keppen,L.; Koob, K. G.; Lupski, J. R.: Mutations in the early growth response2 (EGR2) gene are associated with hereditary myelinopathies. *NatureGenet.* 18: 382–384, 1998.
- [80012] 18342.Warner, L. E.; Svaren, J.; Milbrandt, J.; Lupski, J. R.: Functionalconsequences of mutations in the early growth response 2 gene (EGR2)correlate with severity of human

myelinopathies. Hum. Molec. Genet. 8:1245–1251, 1999.

[80013] 18343.Wu, J.; Joseph, L.; Sukhatme, V. P.; Kidd, K. K.: A HindIII polymorphism identified by the human early growth response gene 2 (EGR2) on chromosome

[80014] 18344.Nucleic Acids Res. 16: 11855 only, 1988.

[80015] 18345.Kaneda, Y.; Hayes, H.; Uchida, T.; Yoshida, M. C.; Okada, Y.:Regional assignment of five genes on human chromosome 19. Chromosoma 95:8–12, 1987.

[80016] 18346.Hoeffler, J. P.; Meyer, T. E.; Yun, Y.; Jameson, J. L.; Habener, J. F.: Cyclic AMP–responsive DNA–binding protein: structure based on a cloned placental cDNA. Science 242: 1430–1433, 1988.

[80017] 18347.Chan, E. F.; Gat, U.; McNiff, J. M.; Fuchs, E.: A common human skin tumour is caused by activating mutations in beta–catenin. Nature Genet. 21: 410–413, 1999.

[80018] 18348.Horelli–Kuitunen, N.; Kvist, A.–P.; Helaakoski, T.; Kivirikko, K.; Pihlajaniemi, T.; Palotie, A.: The order and transcriptional orientation of the human COL13A1 and P4HA genes on chromosome 10 long arm determined by high–resolution FISH. Genomics 46: 299–302, 1997.

[80019] 18349.Pajunen, L.; Tamminen, M.; Solomon, E.; Pihlajaniemi, T.: Assignment of the gene coding for the alpha 1 chain of collagen type XIII (COL13A1) to human chromo–

some region 10q11–qter. Cytogenet. Cell Genet.  
52:190–193, 1989.

- [80020] 18350.Shows, T. B.; Tikka, L.; Byers, M. G.; Eddy, R. L.; Haley, L. L.; Henry, W. M.; Prockop, D. J.; Tryggvason, K.: Assignment of the human collagen alpha-1(XIII) chain gene (COL13A1) to the q22 region of chromosome 10. Genomics 5: 128–133, 1989.
- [80021] 18351.Tikka, L.; Pihlajaniemi, T.; Henttu, P.; Prockop, D. J.; Tryggvason, K.: Gene structure for the alpha-1 chain of a human short-chain collagen (type XIII) with alternatively spliced transcripts and translation termination codon at the 5-prime end of the last exon. Proc. Nat. Acad. Sci. 85: 7491–7495, 1988.
- [80022] 18352.Cunningham, B. A.; Hemperly, J. J.; Murray, B. A.; Prediger, E.A.; Brackenbury, R.; Edelman, G. M.: Neural cell adhesion molecule: structure, immunoglobulin-like domains, cell surface modulation, and alternative RNA splicing. Science 236: 799–806, 1987.
- [80023] 18353.D'Eustachio, P.; Davisson, M. T.: Resolution of the staggerer (sg) mutation from the neural cell adhesion molecule locus (Ncam) on mouse chromosome 9. Mammalian Genome 4: 278–280, 1993.
- [80024] 18354.D'Eustachio, P.; Owens, G. C.; Edelman, G. M.; Cun-

ningham, B. A.: Chromosomal location of the gene encoding the neural cell adhesion molecule (N-CAM) in the mouse. *Proc. Nat. Acad. Sci.* 82: 7631–7635, 1985.

[80025] 18355. Lin, D. M.; Fetter, R. D.; Kopczynski, C.; Grenningloh, G.; Goodman, C. S.: Genetic analysis of fasciclin II in *Drosophila*: defasciculation, refasciculation, and altered fasciculation. *Neuron* 13: 1055–1069, 1994.

[80026] 18356. Altare, F.; Jouanguy, E.; Lamhamedi-Cherradi, S.; Fondaneche, M.-C.; Fizame, C.; Ribierre, F.; Merlin, G.; Dembic, Z.; Schreiber, R.; Lisowska-Grospierre, B.; Fischer, A.; Seboun, E.; Casanova, J.-L.: A causative relationship between mutant IFN $\gamma$ R1 alleles and impaired cellular response to IFN- $\gamma$  in a compound heterozygous child. (Letter) *Am. J. Hum. Genet.* 62:723–726, 1998.

[80027] 18357. Dalton, D. K.; Pitts-Meek, S.; Keshav, S.; Figari, I. S.; Bradley, A.; Stewart, T. A.: Multiple defects of immune cell function in mice with disrupted interferon- $\gamma$  genes. *Science* 259: 1739–1742, 1993.

[80028] 18358. Dessein, A. J.; Hillaire, D.; Elwali, N. E. M. A.; Marquet, S.; Mohamed-Ali, Q.; Mirghani, A.; Henri, S.; Abdelhameed, A. A.; Saeed, O. K.; Magzoub, M. M. A.; Abel, L.: Severe hepatic fibrosis in *Schistosoma mansoni* infection is controlled by a major locus that is closely linked to the in-



terferon-gamma receptor gene. Am. J. Hum. Genet. 65: 709-721,1999.

[80029] 18359.Fellous, M.: Personal Communication. Paris, France 10/24/1986.

[80030] 18360.Fellous, M.; Couillin, P.; Rosa, F.; Metezeau, P.; Foubert, C.;Gross, M. S.; Frezal, J.; Van Cong, N.: Receptor for human gammainterferon is specified by human chromosome 18. (Abstract) Cytogenet.Cell Genet. 40: 627-628, 1985.

[80031] 18361.Jouanguy, E.; Altare, F.; Lamhamedi, S.; Revy, P.; Emile, J.-F.;Newport, M.; Levin, M.; Blanche, S.; Seboun, E.; Fischer, A.; Casanova,J.-L.: Interferon-gamma-receptor deficiency in an infant with fatalbacille Calmette-Guerin infection. New Eng. J. Med. 335: 1956-1961,1996.

[80032] 18362.Jouanguy, E.; Dupuis, S.; Pallier, A.; Doffinger, R.; Fondaneche,M.-C.; Fieschi, C.; Lamhamedi-Cherradi, S.; Altare, F.; Emile, J.-F.;Lutz, P.; Bordigoni, P.; Cokugras, H.; Akcakaya, N.; Landman-Parker,J.; Donnadieu, J.; Camcioglu, Y.; Casanova, J.-L.: In a novel formof IFN-gamma receptor 1 deficiency, cell surface receptors fail tobind IFN-gamma. J. Clin. Invest. 105: 1429-1436, 2000.

[80033] 18363.Jouanguy, E.; Lamhamedi-Cherradi, S.; Lammas, D.; Dorman, S. E.;Fondaneche, M.-C.; Dupuis, S.; Doffin-

ger, R.; Altare, F.; Girdlestone, J.; Emile, J.-F.; Ducoulombier, H.; Edgar, D.; and 10 others: A human IFNGR1 small deletion hotspot associated with dominant susceptibility to mycobacterial infection. *Nature Genet.* 21: 370–378, 1999.

[80034] 18364. Kaplan, D. H.; Shankaran, V.; Dighe, A. S.; Stockert, E.; Aguet, M.; Old, L. J.; Schreiber, R. D.: Demonstration of an interferon gamma-dependent tumor surveillance system in immunocompetent mice. *Proc. Nat. Acad. Sci.* 95: 7556–7561, 1998.

[80035] 18365. Le Coniat, M.; Alcaide-Loridan, C.; Fellous, M.; Berger, R.: Human interferon gamma receptor 1 (IFNGR1) gene maps to chromosome region 6q23–6q24. *Hum. Genet.* 84: 92–94, 1989.

[80036] 18366. Lekstrom-Himes, J. A.; Gallin, J. I.: Immunodeficiency diseases caused by defects in phagocytes. *New Eng. J. Med.* 343: 1703–1714, 2000.

[80037] 18367. Levin, M.; Newport, M. J.; D'Souza, S.; Kalabalikis, P.; Brown, I. N.; Lenicker, H. M.; Agius, P. V.; Davies, E. G.; Thrasher, A.; Klein, N.; Blackwell, J. M.: Familial disseminated atypical mycobacterial infection in childhood: a human mycobacterial susceptibility gene? *Lancet* 345: 79–83, 1995.

- [80038] 18368.Mariano, T. M.; Kozak, C. A.; Langer, J. A.; Pestka, S.: Themouse immune interferon receptor gene is located on chromosome 10. *J.Biol. Chem.* 262: 5812–5814, 1987.
- [80039] 18369.Deloukas, P.; Schuler, G. D.; Gyapay, G.; Beasley, E. M.; Soderlund,C.; Rodriguez–Tome, P.; Hui, L.; Matisse, T. C.; McKusick, K. B.; Beckmann,J. S.; Bentolila, S.; Bihoreau, M.–T.; and 53 others: A physicalmap of 30,000 human genes. *Science* 282: 744–746, 1998.
- [80040] 18370.Wu, W. J.; Erickson, J. W.; Lin, R.; Cerione, R. A.: The gamma–subunitof the coatomer complex binds Cdc42 to mediate transformation. *Nature* 405:800–804, 2000.
- [80041] 18371.Davis, S. T.; Benson, B. G.; Bramson, H. N.; Chapman, D. E.; Dickerson,S. H.; Dold, K. M.; Eberwein, D. J.; Edelstein, M.; Frye, S. V.; Gampe,R. T., Jr.; Griffin, R. J.; Harris, P. A.; and 14 others: Preventionof chemotherapy–induced alopecia in rats by CDK inhibitors. *Science* 291:134–137, 2001.
- [80042] 18372.De Bondt, H. L.; Rosenblatt, J.; Jancarik, J.; Jones, H. D.; Morgan,D. O.; Kim, S.–H.: Crystal structure of cyclin–dependent kinase 2. *Nature* 363:595–602, 1993.
- [80043] 18373.Basu, S.; Binder, R. J.; Ramalingam, T.; Srivastava, P. K.: CD91is a common receptor for heat shock proteins gp96, hsp90, hsp70, andcalreticulin. *Immunity* 14:

303–313, 2001.

- [80044] 18374.Beisiegel, U.; Weber, W.; Ihrke, G.; Herz, J.; Stanley, K. K.:The LDL–receptor–related protein, LRP, is an apolipoprotein E–bindingprotein. *Nature* 341: 162–164, 1989.
- [80045] 18375.Binder, R. J.; Han, D. K.; Srivastava, P. K.: CD91: a receptorfor heat shock protein gp96. *Nature Immun.* 1: 151–155, 2000.
- [80046] 18376.Forus, A.; Maelandsmo, G. M.; Fodstad, Y.; Myklebost, O.: Thegenes for the alpha–2–macroglobulin receptor/LDL receptor–relatedprotein and GLI are located within a chromosomal segment of about300 kilobases and are coamplified in a rhabdomyosarcoma cell line.(Abstract) *Cytogenet. Cell Genet.* 58: 1977 only, 1991.
- [80047] 18377.Forus, A.; Myklebost, O.: A physical map of a 1.3–Mb region onthe long arm of chromosome 12, spanning the GLI and LRP loci. *Genomics* 14:117–120, 1992.
- [80048] 18378.Hilliker, C.; Van Leuven, F.; Van Den Berghe, H.: Assignment ofthe gene coding for the alpha(2)–macroglobulin receptor to mouse chromosome15 and to human chromosome 12q13–q14 by isotopic and nonisotopic insitu hybridization. *Genomics* 13: 472–474, 1992.

- [80049] 18379.Kristensen, T.; Moestrup, S. K.; Gliemann, J.; Bendtsen, L.; Sand,O.; Sottrup-Jensen, L.: Evidence that the newly cloned low-density-lipoproteinreceptor related protein (LRP) is the alpha-2-macroglobulin receptor. FEB-SLett. 276: 151-155, 1990.
- [80050] 18380.Lendon, C. L.; Talbot, C. J.; Craddock, N. J.; Han, S. W.; Wragg,M.; Morris, J. C.; Goate, A. M.: Genetic association studies between dementia of the Alzheimer's type and three receptors for apolipoproteinE in a Caucasian population. Neurosci. Lett. 222: 187-190, 1997.
- [80051] 18381.Myklebost, O.; Arheden, K.; Rogne, S.; Geurts van Kessel, A.;Mandahl, N.; Herz, J.; Stanley, K.; Heim, S.; Mitelman, F.: The gene for the human putative apoE receptor is on chromosome 12 in the segmentq13-14. Genomics 5: 65-69, 1989.
- [80052] 18382.Pericak-Vance, M. A.; Bass, M. P.; Yamaoka, L. H.; Gaskell, P.C.; Scott, W. K.; Terwedow, H. A.; Menold, M. M.; Conneally, P. M.;Small, G. W.; Vance, J. M.; Saunders, A. M.; Roses, A. D.; Haines,J. L.: Complete genomic screen in late-onset familial Alzheimer disease:evidence for a new locus on chromosome 12. J.A.M.A. 278: 1237-1241,1997.
- [80053] 18383.Scott, W. K.; Yamaoka, L. H.; Bass, M. P.; Gaskell, P.

C.; Conneally, P. M.; Small, G. W.; Farrer, L. A.; Auerbach, S. A.; Saunders, A. M.; Roses, A. D.; Haines, J. L.; Pericak-Vance, M. A.: No genetic association between the LRP receptor and sporadic or late-onset familial Alzheimer disease. *Neurogenetics* 1: 179–183, 1998.

[80054] 18384. Strickland, D. K.; Ashcom, J. D.; Williams, S.; Burgess, W. H.; Migliorini, M.; Argraves, W. S.: Sequence identity between the alpha-2-macroglobulin receptor and low density lipoprotein receptor-related protein suggest that this molecule is a multifunctional receptor. *J. Biol. Chem.* 265:17401–17404, 1990.

[80055] 18385. Yochem, J.; Greenwald, I.: A gene for a low density lipoprotein receptor-related protein in the nematode *Caenorhabditis elegans*. *Proc. Nat. Acad. Sci.* 90: 4572–4576, 1993.

[80056] 18386. Ladias, J. A. A.; Karathanasis, S. K.: Regulation of the apolipoprotein AI gene by ARP-1, a novel member of the steroid receptor superfamily. *Science* 251:561–565, 1991.

[80057] 18387. Modi, W. S.; Seuanez, H.; Mietus-Snyder, M.; O'Brien, S. J.; Karathanasis, S. K.: Chromosomal localization of the ARP-1 gene to 15q26. (Abstract) *Cytogenet. Cell Genet.* 58: 1995 only, 1991.

- [80058] 18388.Pereira, F. A.; Qiu, Y.; Zhou, G.; Tsai, M.-J.; Tsai, S. Y.: Theorphan nuclear receptor COUP-TFII is required for angiogenesis and heart development. *Genes Dev.* 13: 1037–1049, 1999.
- [80059] 18389.Widom, R. L.; Ladias, J. A. A.; Kouidou, S.; Karathanasis, S. K.: Synergistic interactions between transcription factors control expression of the apolipoprotein AI gene in liver cells. *Molec. Cell. Biol.* 11:677–687, 1991.
- [80060] 18390.Ikeda, H.; Yamaguchi, M.; Sugai, S.; Aze, Y.; Narumiya, S.; Kakizuka, A.: Expanded polyglutamine in the Machado–Joseph disease protein induces cell death in vitro and in vivo. *Nature Genet.* 13: 196–202, 1996.
- [80061] 18391.Kang, H.; Sun, L. D.; Atkins, C. M.; Soderling, T. R.; Wilson, M. A.; Tonegawa, S.: An important role of neural activity–dependent CaMKIV signaling in the consolidation of long–term memory. *Cell* 106:771–783, 2001.
- [80062] 18392.Raman, V.; Blaeser, F.; Ho, N.; Engle, D. L.; Williams, C. B.; Chatila, T. A.: Requirement for Ca(2+)/calmodulin–dependent kinase type IV/Gr in setting the thymocyte selection threshold. *J. Immun.* 167:6270–6278, 2001.
- [80063] 18393.Santisteban, I.; Arredondo–Vega, F. X.; Kelly, S.; Mary, A.; Fischer, A.; Hummell, D. S.; Lawton, A.; Sorensen,

R. U.; Stiehm, E. R.; Uribe, L.; Weinberg, K.; Hershfield, M. S.: Novel splicing, missense, and deletion mutations in seven adenosine deaminase-deficient patients with late/delayed onset of combined immunodeficiency disease: contribution of genotype to phenotype. *J. Clin. Invest.* 92: 2291–2302, 1993. 100. Schmalstieg, F. C.; Mills, G. C.; Tsuda, H.; Goldman, A. S.: Severe combined immunodeficiency in a child with a healthy adenosine deaminase deficient mother. *Pediat. Res.* 17: 935–940, 1983. 101.

Schrader, W. P.; Pollara, B.; Meuwissen, H. J.: Characterization of the residual adenosine deaminating activity in the spleen of a patient with combined immunodeficiency disease and adenosine deaminase deficiency. *Proc. Nat. Acad. Sci.* 75: 446–450, 1978. 102. Scott, C. R.; Chen, S.-H.; Giblett, E. R.: Deletion of the carrier state in combined immunodeficiency disease associated with deaminase deficiency. *J. Clin. Invest.* 53: 1194–1196, 1974. 103. Shovlin, C. L.; Hughes, J. M. B.; Simmonds, H. A.; Fairbanks, L.; Deacock, S.; Lechler, R.; Roberts, I.; Webster, A. D. B.: Adult presentation of adenosine deaminase deficiency. (Letter) *Lancet* 341:1471, 1993. 104. Shovlin, C. L.; Simmonds, H. A.; Fairbanks, L. D.; Deacock, S. J.; Hughes, J. M. B.; Lechler, R. I.; Webster, A. D. B.; Sun, X.-M.; Webb, J. C.;



Soutar, A. K.: Adult onset immunodeficiency caused by inherited adenosine deaminase deficiency. *J. Immun.* 153: 2331–2339, 1994.105. Spencer, N.; Hopkinson, D. A.; Harris, H.: Adenosine deaminase polymorphism in man. *Ann. Hum. Genet.* 32: 9–14, 1968.106. Stephan, V.; Wahn, V.; Le Deist, F.; Dirksen, U.; Broker, B.; Muller-Fleckenstein, I.; Horneff, G.; Schroten, H.; Fischer, A.; deSaint Basile, G.: Atypical X-linked severe combined immunodeficiency due to possible spontaneous reversion of the genetic defect in T cells. *New Eng. J. Med.* 335: 1563–1567, 1996.107. Tariverdian, G.; Ritter, H.: Adenosine deaminase polymorphism (EC 3.5.4.4): formal genetics and linkage relations. *Humangenetik* 7:176–178, 1969.108. Tischfield, J. A.; Creagan, R. P.; Nichols, E. A.; Ruddle, F.H.: Assignment of a gene for adenosine deaminase to human chromosome 20. *Hum. Hered.* 24: 1–11, 1974.109. Tzall, S.; Ellenbogen, A.; Eng, F.; Hirschhorn, R.: Identification and characterization of nine RFLPs at the adenosine deaminase (ADA) locus. *Am. J. Hum. Genet.* 44: 864–875, 1989.110. Umetsu, D. T.; Schlossman, C. M.; Ochs, H. D.; Hershfield, M.S.: Heterogeneity of phenotype in two siblings with adenosine deaminase deficiency. *J. Allergy Clin. Immun.* 93: 543–550, 1994.111. Valerio, D.; Dekker, B. M. M.;

Duyvesteyn, M. G. C.; van derVoorn, L.; Berkvens, T. M.; van Ormondt, H.; van der Eb, A. J.: Oneadenosine deaminase allele in a patient with severe combined immunodeficiencycontains a point mutation abolishing enzyme activity. EMBO J. 5:113–119, 1986.112. Valerio, D.; Duyvesteyn, M. G. C.; Dekker, B. M. M.; Weeda, G.;Berkvens, T. M.; van der Voorn, L.; van Ormondt, H.; van der Eb, A.J.: Adenosine deaminase: characterization and expression of a genewith a remarkable promoter. EMBO J. 4: 437–443, 1985.113. Valerio, D.; Duyvesteyn, M. G. C.; Meera Khan, P.; Pearson, P.L.; Geurts van Kessel, A.; van Ormondt, H.: Direct assignment ofADA gene to chromosome 20. (Abstract) Cytogenet. Cell Genet. 37:599, 1984.114. Valerio, D.; Duyvesteyn, M. G. C.; van Ormondt, H.; Meera Khan,P.; van der Eb, A. J.: Adenosine deaminase (ADA) deficiency in cellsderived from humans with severe combined immunodeficiency is due toan aberration of the ADA protein. Nucleic Acids Res. 12: 1015–1024,1984.115. Valerio, D.; McIvor, R. S.; Williams, S. R.; Duyvesteyn, M. G.C.; van Ormondt, H.; van der Eb, A. J.; Martin, D. W., Jr.: Cloningof human adenosine deaminase cDNA and expression in mouse cells. Gene 31:147–153, 1984.116. Van der Weyden, M. B.; Kelley, W.

N.: Adenosine deaminase deficiency in severe combined immunodeficiency: evidence for a posttranslational defect. (Abstract) J. Clin. Invest. 53: 81A–82A, 1974.117.

Wakamiya, M.; Blackburn, M. R.; Jurecic, R.; McArthur, M. J.; Geske, R. S.; Cartwright, J., Jr.; Mitani, K.; Vaishnav, S.; Belmont, J. W.; Kellems, R. E.; Finegold, M. J.; Montgomery, Jr., C. A.; Bradley, A.; Caskey, C. T.: Disruption of the adenosine deaminase gene causes hepatocellular impairment and perinatal lethality in mice. Proc. Nat. Acad. Sci. 92: 3673–3677, 1995.118. Weitkamp, L. R.: Further data on the genetic linkage relationships of the adenosine deaminase locus. Hum. Hered. 21: 351–356, 1971.119.

Weitkamp, L. R.: Genetic linkage relationships of the ADA and 6-PGD loci in 'Humangenetik.' (Letter) Humangenetik 15: 359–360, 1972.120. Wiginton, D. A.; Adrian, G. S.; Friedman, R. L.; Suttle, D. P.; Hutton, J. J.: Cloning of cDNA sequences of human adenosine deaminase. Proc. Nat. Acad. Sci. 80: 7481–7485, 1983.121. Wiginton, D. A.; Adrian, G. S.; Hutton, J. J.: Sequence of human adenosine deaminase cDNA including the coding region and a small intron. Nucleic Acids Res. 12: 2439–2446, 1984.122. Wiginton, D. A.; Hutton, J. J.: Immunoreactive protein in adenosine deaminase deficient human lymphoblast cell

lines. J. Biol. Chem. 257:3211–3217, 1982.123. Wiginton, D. A.; Kaplan, D. J.; States, J. C.; Akeson, A. L.; Perme, C. M.; Bilyk, I. J.; Vaughn, A. J.; Lattier, D. L.; Hutton, J. J.: Complete sequence and structure of the gene for human adenosinedeaminase. Biochemistry 25: 8234–8244, 1986.124. Yokoyama, S.; Hayashi, T.; Yoshimura, Y.; Irimada, K.; Saito, T.; Akiba, T.; Tsuchiya, S.: Severe combined immunodeficiency disease with adenosine deaminase deficiency. Tohoku J. Exp. Med. 129: 197–202, 1979.125. Yount, J.; Nichols, P.; Ochs, H. D.; Hammar, S. P.; Scott, C. R.; Chen, S.-H.; Giblett, E. R.; Wedgwood, R. J.: Absence of erythrocyte adenosine deaminase associated with severe combined immunodeficiency. J. Pediatr. 84: 173–177, 1974.126. Ziegler, J. B.; Lee, C. H.; Van Der Weyden, M. B.; Bagnara, A. S.; Beveridge, J.: Severe combined immunodeficiency and adenosinedeaminase deficiency: failure of enzyme replacement therapy. Arch. Dis. Child. 55: 452–457, 1980.127. Ziegler, J. B.; Van Der Weyden, M. B.; Lee, C. H.; Daniel, A.: Prenatal diagnosis for adenosine deaminase deficiency. J. Med. Genet. 18: 154–156, 1981.

[80064] 18394. Sikela, J. M.; Adamson, M. C.; Wilson-Shaw, D.; Kozak, C. A.: Genetic mapping of the gene for

Ca(2+)/calmodulin-dependent protein kinase IV (Camk-4) to mouse chromosome 18. *Genomics* 8: 579–582, 1990.

[80065] 18395. Sikela, J. M.; Law, M. L.; Kao, F.-T.; Hartz, J. A.; Wei, Q.; Hahn, W. E.: Chromosomal localization of the human gene for brain Ca(2+)/calmodulin-dependent protein kinase type IV. *Genomics* 4: 21–27, 1989.

[80066] 18396. Wei, F.; Qiu, C.-S.; Liao, J.; Robinson, D. A.; Ho, N.; Chatila, T.; Zhuo, M.: Calcium-calmodulin-dependent protein kinase IV is required for fear memory. *Nature Neurosci.* 5: 573–579, 2002.

[80067] 18397. Wu, H.; Kanatous, S. B.; Thurmond, F. A.; Gallardo, T.; Isotani, E.; Bassel-Duby, R.; Williams, R. S.: Regulation of mitochondrial biogenesis in skeletal muscle by CaMK. *Science* 296: 349–352, 2002.

[80068] 18398. Wu, J. Y.; Ribar, T. J.; Cummings, D. E.; Burton, K. A.; McKnight, G. S.; Means, A. R.: Spermiogenesis and exchange of basic nuclear proteins are impaired in male germ cells lacking Camk4. *Nature Genet.* 25:448–452, 2000.

[80069] 18399. Dooley, T. P.: Personal Communication. San Antonio, Tex. 7/21/1992.

[80070] 18400. Dooley, T. P.; Weiland, K. L.; Simon, M.: cDNA sequence of human p11 calpactin I light chain. *Genomics* 13:

866–868, 1992.

- [80071] 18401.Harder, T.; Kube, E.; Gerke, V.: Cloning and characterization of the human gene encoding p11: structural similarity to other members of the S-100 gene family. *Gene* 113: 269–274, 1992.
- [80072] 18402.Kube, E.; Weber, K.; Gerke, V.: Primary structure of human, chicken, and *Xenopus laevis* p11, a cellular ligand of the Src-kinase substrate, annexin II. *Gene* 102: 255–259, 1991.
- [80073] 18403.Okuse, K.; Malik-Hall, M.; Baker, M. D.; Poon, W.-Y. L.; Kong, H.; Chao, M. V.; Wood, J. N.: Annexin II light chain regulates sensory neuron-specific sodium channel expression. *Nature* 417: 653–656, 2002.
- [80074] 18404.Saris, C. J. M.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.; Noonan, D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and tissue distribution of the mRNA for bovine and murine p11, the S100-related light chain of the protein-tyrosine kinase substrate p36 (calpactin I). *J. Biol. Chem.* 262: 10663–10671, 1987.
- [80075] 18405.Inazawa, J.; Nakagawa, H.; Misawa, S.; Abe, T.; Minoshima, S.; Fukuyama, R.; Maki, M.; Murachi, T.; Hatanaka, M.; Shimizu, N.: Assignment of the human calpastatin gene (CAST) to chromosome 5 at region

q14-q22.Cytogenet. Cell Genet. 54: 156-158, 1990.

[80076] 18406.Inazawa, J.; Nakagawa, H.; Misawa, S.; Abe, T.; Minoshima, S.;Fukuyama, R.; Maki, M.; Murachi, T.; Hatanaka, M.; Shimizu, N.: Assignmentof the human calpastatin gene (CAST) to chromosome 5 at region q15-q21.(Abstract) Cytogenet. Cell Genet. 58: 1898 only, 1991.

[80077] 18407.Mimori, T.; Suganuma, K.; Tanami, Y.; Nojima, T.; Matsumura, M.;Fujii, T.; Yoshizawa, T.; Suzuki, K.; Akizuki, M.: Autoantibodiesto calpastatin (an endogenous inhibitor for calcium-dependent neutralprotease, calpain) in systemic rheumatic diseases. Proc. Nat. Acad.Sci. 92: 7267-7271, 1995.

[80078] 18408.Pontremoli, S.; Salamino, F.; Sparatore, B.; De Tullio, R.; Pontremoli,R.; Melloni, E.: Characterization of the calpastatin defect in erythrocytesfrom patients with essential hypertension. Biochem. Biophys. Res.Commun. 157: 867-874, 1988.

[80079] 18409.Carritt, B.: Somatic cell genetic evidence for the presence ofa gene for citrullinemia on human chromosome 9. Cytogenet. Cell Genet. 19:44-48, 1977.

[80080] 18410.Carritt, B.; Goldfarb, P. S. G.; Hooper, M. L.; Slack, C.: Chromosomeassignment of a human gene for argini-

nosuccinate synthetase expression in Chinese hamster-human somatic cell hybrids. *Exp. Cell Res.* 106:71-78, 1977.

[80081] 18411. Daiger, S. P.; Hoffman, N. S.; Wildin, R. S.; Su, T.-S.: Multiple, independent restriction site polymorphisms in human DNA detected with a cDNA probe to argininosuccinate synthetase (AS). *Am. J. Hum. Genet.* 36:736-749, 1984.

[80082] 18412. Daiger, S. P.; Wildin, R. S.; Su, T.-S.: Polymorphic variants of restriction fragments of human DNA detected with a probe to argininosuccinate synthetase. (Abstract) *Am. J. Hum. Genet.* 33: 136A only, 1981.

[80083] 18413. Elshourbagy, N. A.; Walker, D. W.; Boguski, M. S.; Gordon, J. I.; Taylor, J. M.: The nucleotide and derived amino acid sequence of human apolipoprotein A-IV mRNA and the close linkage of its gene to the genes of apolipoproteins A-I and C-III. *J. Biol. Chem.* 261:1998-2002, 1986.

[80084] 18414. Elshourbagy, N. A.; Walker, D. W.; Paik, Y.-K.; Boguski, M. S.; Freeman, M.; Gordon, J. I.; Taylor, J. M.: Structure and expression of the human apolipoprotein A-IV gene. *J. Biol. Chem.* 262: 7973-7981, 1987.

[80085] 18415. Green, P. H. R.; Glickman, R. M.; Riley, J. W.; Quinet, E.: Human apolipoprotein A-IV: intestinal origin



and distribution in plasma. J.Clin. Invest. 65: 911–919, 1980.

[80086] 18416.Kamboh, M. I.; Ferrell, R. E.: Genetic studies of human apolipoproteins.I. Polymorphism of apolipoprotein A-IV. Am. J. Hum. Genet. 41: 119–127,1987.

[80087] 18417.Kamboh, M. I.; Williams, E. R.; Law, J. C.; Aston, C. E.; Bunker,C. H.; Ferrell. R. E.; Pollitzer, W. S.: Molecular basis of a uniqueAfrican variant (A-IV 5) of human apolipoprotein A-IV and its significancein lipid metabolism. Genet. Epidemiol. 9: 379–388, 1992.

[80088] 18418.Karathanasis, S. K.: Apolipoprotein multigene family: tandem organizationof human apolipoprotein AI, CIII, and AIV genes. Proc. Nat. Acad.Sci. 82: 6374–6378, 1985.

[80089] 18419.Lohse, P.; Kindt, M. R.; Rader, D. J.; Brewer, H. B., Jr.: Geneticpolymorphism of human plasma apolipoprotein A-IV is due to nucleotidesubstitutions in the apolipoprotein A-IV gene. J. Biol. Chem. 265:10061–10064, 1990.

[80090] 18420.Lohse, P.; Kindt, M. R.; Rader, D. J.; Brewer, H. B., Jr.: Humanplasma apolipoproteins A-IV-0 and A-IV-3: molecular basis for two rare variants of apolipoprotein A-IV-1. J. Biol. Chem. 265: 12734–12739,1990.

[80091] 18421.Menzel, H.-J.; Boerwinkle, E.; Schrangl-Will, S.; Utermann, G.: Human apolipoprotein A-IV polymorphism:

frequency and effect on lipid and lipoprotein levels. Hum. Genet. 79: 368–372, 1988.

- [80092] 18422.Menzel, H.-J.; Kovary, P. M.; Assmann, G.: Apolipoprotein A-IV polymorphism in man. Hum. Genet. 62: 349–352, 1982.
- [80093] 18423.Menzel, H.-J.; Sigurdsson, G.; Boerwinkle, E.; Schrangl-Will, S.; Dieplinger, H.; Utermann, G.: Frequency and effect of human apolipoprotein A-IV polymorphism on lipid and lipoprotein levels in an Icelandic population. Hum. Genet. 84: 344–346, 1990.
- [80094] 18424.Rogne, S.; Myklebost, O.; Olaisen, B.; Gedde-Dahl, T., Jr.; Prydz, H.: Confirmation of the close linkage between the loci for human apolipoproteins AI and AIV by the use of a cloned cDNA probe and two restriction site polymorphisms. Hum. Genet. 72: 68–71, 1986.
- [80095] 18425.Schamaun, O.; Olaisen, B.; Mevag, B.; Gedde-Dahl, T., Jr.; Ehnholm, C.; Teisberg, P.: The two apolipoprotein loci apoA-I and apoA-IV are closely linked in man. Hum. Genet. 68: 181–184, 1984.
- [80096] 18426.Tracy, R. P.; Currie, R. M.; Young, D. S.: Two-dimensional gelelectrophoresis of serum specimens from a normal population. Clin.Chem. 28: 890–899, 1982.
- [80097] 18427.von Eckardstein, A.; Funke, H.; Schulte, M.; Erren,

M.; Schulte,H.; Assmann, G.: Nonsynonymous polymorphic sites in the apolipoprotein(apo) A-IV gene are associated with changes in the concentration of apo B- and apo A-I-containing lipoproteins in a normal population. *Am.J. Hum. Genet.* 50: 1115-1128, 1992.

[80098] 18428.Gautier, T.; Masson, D.; Jong, M. C.; Duverneuil, L.; Le Guern,N.; Deckert, V.; Pais de Barros, J.-P.; Dumont, L.; Bataille, A.;Zak, Z.; Jiang, X.-C.; Tall, A. R.; Havekes, L. M.; Lagrost, L.:Apolipoprotein CI deficiency markedly augments plasma lipoproteinchanges mediated by human cholesteryl ester transfer protein (CETP)in CETP transgenic/ApoCI-knocked out mice. *J. Biol. Chem.* 277: 31354-31363,2002.

[80099] 18429.Brown, S.; Wiebel, F. J.; Gelboin, H. V.; Minna, J. D.: Assignmentof a locus required for flavoprotein-linked monooxygenase expressionto human chromosome 2. *Proc. Nat. Acad. Sci.* 73: 4628-4632, 1976.

[80100] 18430.Chen, Y. T.; Tukey, R. H.; Swan, D. C.; Negishi, N.; Nebert, D.W.: Characterization of the human P1-450 genomic gene. (Abstract) *Clin.Res.* 31: 456A, 1983.

[80101] 18431.Corcherro, J.; Pimprale, S.; Kimura, S.; Gonzalez, F. J.: Organizationof the CYP1A cluster on human chromosome 15: implications for generegulation. *Pharmacoge-*

netics 11: 1–6, 2001.

- [80102] 18432.Gorman, C.; Padmanabhan, R.; Howard, B. H.: High efficiency DNA-mediated transformation of primate cells. Science 221: 551–553, 1983.
- [80103] 18433.Gorman, C. M.: CAT: an easy assay for gene expression (citation classic). Current Contents (Life Sciences) 36(22): 8, 1993.
- [80104] 18434.Hildebrand, C. E.; Gonzalez, F. J.; Kozak, C. A.; Nebert, D. W.: Regional linkage analysis of the dioxin-inducible P-450 gene family on mouse chromosome 9. Biochem. Biophys. Res. Commun. 130: 396–406, 1985.
- [80105] 18435.Lee, S.; Wu, X.; Reid, M. E.; Zelinski, T.; Redman, C. M.: Molecular basis of the Kell (K1) phenotype. Blood 85: 912–916, 1995.
- [80106] 18436.Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.: The human Kell blood group gene maps to chromosome 7q33 and its expression is restricted to erythroid cells. Blood 81: 2804–2809, 1993.
- [80107] 18437.Lee, S.; Zambas, E. D.; Marsh, W. L.; Redman, C. M.: Molecular cloning and primary structure of Kell blood group protein. Proc. Nat. Acad. Sci. 88: 6353–6357, 1991.
- [80108] 18438.Marsh, W. L.: Molecular biology of blood groups: cloning of the Kell gene. Transfusion 32: 98–101, 1992.

- [80109] 18439.Morton, N. E.; Krieger, H.; Steinberg, A. G.; Rosenfield, R. E.: Genetic evidence confirming the localization of Sutter in the Kellblood-group system. *Vox Sang.* 10: 608–613, 1965.
- [80110] 18440.Murphy, M. T.; Morrison, N.; Miles, J. S.; Fraser, R. H.; Spurr, N. K.; Boyd, E.: Regional chromosomal assignment of the Kell bloodgroup locus (KEL) to chromosome 7q33–q35 by fluorescence in situ hybridization:evidence for the polypeptide nature of antigenic variation. *Hum.Genet.* 91: 585–588, 1993.
- [80111] 18441.Parke, J. T.; Riccardi, V. M.; Lewis, R. A.; Ferrell, R. E.:A syndrome of microcephaly and retinal pigmentary abnormalities without mental retardation in a family with coincidental autosomal dominant hyperreflexia. *Am. J. Med. Genet.* 17: 585–594, 1984.
- [80112] 18442.Purohit, K. R.; Weber, J. L.; Ward, L. J.; Keats, B. J. B.: The Kell blood group locus is close to the cystic fibrosis locus on chromosome 7. *Hum. Genet.* 89: 457–458, 1992.
- [80113] 18443.Russo, D.; Wu, X.; Redman, C. M.; Lee, S.: Expression of Kellblood group protein in nonerythroid tissues *Blood* 96: 340–346, 2000.
- [80114] 18444.Stroup, M.; MacIlroy, M.; Walker, R.; Aydelotte, J. V.: Evidence that Sutter belongs to the Kell blood group sys–

tem. Transfusion 5:309–314, 1965.

- [80115] 18445.Wagner, T.; Berer, A.; Lanzer, G.; Geissler, K.: Kell is not restricted to the erythropoietic lineage but is also expressed on myeloid progenitor cells. Brit. J. Haemat. 110: 409–411, 2000.
- [80116] 18446.Weitkamp, L. R.; Townes, P. L.; Johnston, E.: Linkage data on urinary pepsinogen and the Kell blood group. Birth Defects Orig.Art. Ser. 11(3): 281–282, 1975. Note: Alternate: Cytogenet. Cell Genet. 14: 451–452, 1975.
- [80117] 18447.Wimer, B. M.; Marsh, W. L.; Taswell, H. F.: Clinical characteristics of the McLeod blood group phenotype. (Abstract) 19th Annual Meeting of the Am. Soc. Hemat., Boston , 1976.
- [80118] 18448.Yu, L.-C.; Twu, Y.-C.; Chang, C.-Y.; Lin, M.: Molecular basis of the Kell-null phenotype: a mutation at the splice site of human KEL gene abolishes the expression of Kell blood group antigens. J.Biol. Chem. 276: 10247–10252, 2001.
- [80119] 18449.Zelinski, T. A.; Coghlan, G. E.; Myal, Y.; White, L. J.; Philipps, S. E.: Assignment of the Kell blood group locus to chromosome 7q.(Abstract) Cytogenet. Cell Genet. 58: 1927 only, 1991.
- [80120] 18450.Field, L. L.; Marazita, M. L.; Spence, M. A.; Crandall,

B. F.;Sparkes, R. S.: Is JK linked to IGK on chromosome 2?  
(Abstract) Cytogenet.Cell Genet. 40: 628–629, 1985.

[80121] 18451.Gedde–Dahl, T.: Personal Communication. Oslo,  
Norway 9/26/1986.

[80122] 18452.Geitvik, G. A.; Hoyheim, B.; Gedde–Dahl, T.;  
Grzeschik, K. H.;Lothe, R.; Tomter, H.; Olaisen, B.: The  
Kidd (JK) blood group locusassigned to chromosome 18  
by close linkage to a DNA–RFLP. Hum. Genet.  
77:205–209, 1987.

[80123] 18453.Hulten, M.; Lindsten, J.; Pen–Ming, L. M.; Fraccaro,  
M.; Mannini,A.; Trepolo, L.; Robson, E. B.; Heiken, A.;  
Tellingén, K. G.: Possiblelocalization of the genes for the  
Kidd blood group on an autosomeinvolved in a reciprocal  
translocation. Nature 211: 1067–1068, 1968.

[80124] 18454.Leppert, M.; Ferrell, R.; Kamboh, M. I.; Beasley, J.;  
O'Connell,P.; Lathrop, M.; Lalouel, J.–M.; White, R.: Linkage  
of the polymorphicprotein markers F13B, C1S, C1R, and  
blood group antigen Kidd in CEPHreference families.  
(Abstract) Cytogenet. Cell Genet. 46: 647, 1987.

[80125] 18455.Lucien, N.; Chiaroni, J.; Cartron, J.–P.; Bailly, P.:  
Partialdeletion in the JK locus causing a Jk(null) phenotype.  
Blood 99:1079–1081, 2002.

[80126] 18456.Lucien, N.; Sidoux–Walter, F.; Olives, B.; Moulds, J.;

Le Pennec, P.-Y.; Cartron, J.-P.; Bailly, P.: Characterization of the gene encoding the human Kidd blood group/urea transporter protein: evidence for splice site mutations in Jk-null individuals. *J. Biol. Chem.* 273:12973–12980, 1998.

[80127] 18457. Olives, B.; Mattei, M.-G.; Huet, M.; Neau, P.; Martial, S.; Cartron, J.-P.; Bailly, P.: Kidd blood group and urea transport function of human erythrocytes are carried by the same protein. *J. Biol. Chem.* 270:15607–15610, 1995.

[80128] 18458. Womack, J. E.: Personal Communication. College Station, Texas 2/26/1990.

[80129] 18459. Hiraoka, L. R.; Hsu, L.; Hsieh, C.-L.: Assignment of ALDH3 to human chromosome 17p11.2 and ALDH5 to human chromosome 9p13. *Genomics* 25:323–325, 1995.

[80130] 18460. Hsu, L. C.; Chang, W.-C.; Shibuya, A.; Yoshida, A.: Human stomach aldehyde dehydrogenase cDNA and genomic cloning, primary structure, and expression in *Escherichia coli*. *J. Biol. Chem.* 267: 3030–3037, 1992.

[80131] 18461. Kays, W. T.; Piatigorsky, J.: Aldehyde dehydrogenase class 3 expression: identification of a cornea-preferred gene promoter in transgenic mice. *Proc. Nat. Acad. Sci.* 94: 13594–13599, 1997.

[80132] 18462. Rogers, G. R.; Markova, N. G.; De Laurenzi, V.;



Rizzo, W. B.; Compton, J. G.: Genomic organization and expression of the human fatty aldehyde dehydrogenase gene (FALDH). *Genomics* 39: 127–135, 1997.

- [80133] 18463. Santisteban, I.; Povey, S.; West, L. F.; Parrington, J. M.; Hopkinson, D. A.: Chromosome assignment, biochemical and immunological studies on a human aldehyde dehydrogenase, ALDH3. *Ann. Hum. Genet.* 49: 87–100, 1985.
- [80134] 18464. Teng, Y.-S.: Stomach aldehyde dehydrogenase: report of a new locus. *Hum. Hered.* 31: 74–77, 1981.
- [80135] 18465. Vasiliou, V.; Bairoch, A.; Tipton, K. F.; Nebert, D. W.: Eukaryotic aldehyde dehydrogenase (ALDH) genes: human polymorphisms, and recommended nomenclature based on divergent evolution and chromosomal mapping. *Pharmacogenetics* 9: 421–434, 1999.
- [80136] 18466. Wunderle, V. M.; Critcher, R.; Hastie, N.; Goodfellow, P. N.; Schedl, A.: Deletion of long-range regulatory elements upstream of SOX9 causes campomelic dysplasia. *Proc. Nat. Acad. Sci.* 95: 10649–10654, 1998.
- [80137] 18467. Young, I. D.; Zuccollo, J. M.; Maltby, E. L.; Broderick, N. J.: Campomelic dysplasia associated with a de novo 2q;17q reciprocal translocation. *J. Med. Genet.* 29: 251–252, 1992.
- [80138] 18468. von Lindern, M.; Poustka, A.; Lehrach, H.; Grosveld,

G.: The (6;9)chromosome translocation, associated with a specific subtype of acutenonlymphocytic leukemia, leads to aberrant transcription of a targetgene on 9q34. *Molec. Cell. Biol.* 10: 4016–4026, 1990.

[80139] 18469.Lynch, H. T.; Schuelke, G. S.; Kimberling, W. J.; Albano, W. A.; Lynch, J. F.; Biscione, K. A.; Lipkin, M. L.; Deschner, E. E.; Mikol, Y. B.; Sandberg, A. A.; Elston, R. C.; Bailey–Wilson, J. E.; Danes, B. S.: Hereditary nonpolyposis colorectal cancer (Lynch syndromes I and II). II. Biomarker studies. *Cancer* 56: 939–951, 1985.

[80140] 18470.Hanauer, A.; Heilig, R.; Levin, M.; Moisan, J. P.; Grzeschik, K.H.; Mandel, J. L.: The actin gene family in man: assignment of the gene for skeletal muscle alpha-actin to chromosome 1, and presence of actin sequences on autosomes 2 and 3, and on the X and Y chromosomes.(Abstract) *Cytogenet. Cell Genet.* 37: 487–488, 1984.

[80141] 18471.Ilkovski, B.; Cooper, S. T.; Nowak, K.; Ryan, M. M.; Yang, N.; Schnell, C.; Durling, H. J.; Roddick, L. G.; Wilkinson, I.; Kornberg, A. J.; Collins, K. J.; Wallace, G.; Gunning, P.; Hardeman, E. C.; Laing, N. G.; North, K. N.: Nemaline myopathy caused by mutations in the muscle alpha-skeletal-actin gene. *Am. J. Hum. Genet.* 68:1333–1343,

2001.

- [80142] 18472. Nowak, K. J.; Wattanasirichaigoon, D.; Goebel, H. H.; Wilce, M.; Pelin, K.; Donner, K.; Jacob, R. L.; Hubner, C.; Oexle, K.; Anderson, J. R.; Verity, C. M.; North, K. N.; and 13 others: Mutations in the skeletal muscle alpha-actin gene in patients with actin myopathy and nemaline myopathy. *Nature Genet.* 23: 208–212, 1999.
- [80143] 18473. Kedes, L.; Ng, S.-Y.; Lin, C.-S.; Gunning, P.; Eddy, R.; Shows, T.; Leavitt, J.: The human beta-actin multigene family. *Trans. Assoc. Am. Phys.* 98: 42–46, 1985.
- [80144] 18474. Leavitt, J.; Bushar, G.; Kakunaga, T.; Hamada, H.; Hirakawa, T.; Goldman, D.; Merrill, C.: Variations in expression of mutant beta-actin accompanying incremental increases in human fibroblast tumorigenicity. *Cell* 28: 259–268, 1982.
- [80145] 18475. Nakajima-Iijima, S.; Hamada, H.; Reddy, P.; Kakunaga, T.: Molecular structure of the human cytoplasmic beta-actin gene; interspecies homology of sequences in the introns. *Proc. Nat. Acad. Sci.* 82: 6133–6137, 1985.
- [80146] 18476. Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.; Kedes, L.; Shows, T.: Chromosome 7 assignment of the human beta-actin functional gene (ACTB) and the chromosomal dispersion of pseudogenes. (Abstract) *Cyto-*

genet.Cell Genet. 40: 712 only, 1985.

- [80147] 18477.Ng, S.-Y.; Gunning, P.; Eddy, R.; Ponte, P.; Leavitt, J.; Shows,T.; Kedes, L.: Evolution of the functional human beta-actin geneand its multi-pseudogene family: conser-  
vation of the noncoding regionsand chromosomal disper-  
sion of pseudogenes. Molec. Cell. Biol. 5:2720-2732,  
1985.
- [80148] 18478.Toyama, S.; Toyama, S.: A variant form of beta-  
actin in a mutantof KB cells resistant to cytochalasin B.  
Cell 37: 609-614, 1984.
- [80149] 18479.Vandekerckhove, J.; Weber, K.: Mammalian cyto-  
plasmic actins arethe products of at least two genes and  
differ in primary structurein at least 25 identified posi-  
tions from skeletal muscle actins. Proc.Nat. Acad. Sci. 75:  
1106-1110, 1978.
- [80150] 18480.Allderdice, P. W.; Kaita, H.; Lewis, M.; McAlpine, P.  
J.; Wong,P.; Anderson, J.; Giblett, E. R.: Segregation of  
marker loci in familieswith an inherited paracentric inser-  
tion of chromosome 9. Am. J. Hum.Genet. 39: 612-617,  
1986.
- [80151] 18481.Dayhoff, M. O.: Atlas of Protein Sequence and  
Structure. Dehydrogenases. Washington: National  
Biomedical Research Foundation (pub.) 5: 1972.Pp.

D141–D144.

- [80152] 18482.Osier, M. V.; Pakstis, A. J.; Soodyall, H.; Comas, D.; Goldman,D.; Odunsi, A.; Okonofua, F.; Parnas, J.; Schulz, L. O.; Bertranpetit,J.; Bonne–Tamir, B.; Lu, R.–B.; Kidd, J. R.; Kidd, K. K.: A globalperspective on genetic variation at the ADH genes reveals unusualpatterns of linkage dise–quilibrium and diversity. *Am. J. Hum. Genet.* 71:84–99, 2002.
- [80153] 18483.Smith, M.; Hopkinson, D. A.; Harris, H.: Studies on the subunitstructure and molecular size of the human de–hydrogenase isozymes determinedby the different loci, ADH(1), ADH(2), and ADH(3). *Ann. Hum. Genet.* 36:401–414, 1973.
- [80154] 18484.Comings, D. E.; Gade–Andavolu, R.; Gonzalez, N.; Blake, H.; Wu,S.; MacMurray, J. P.: Additive effect of three noradrenergic genes(ADRA2A, ADRA2C, DBH) on atten–tion–deficit hyperactivity disorderand learning disabilities in Tourette syndrome subjects. *Clin. Genet.* 55:160–172, 1999.
- [80155] 18485.Halperin, J. M.; Newcorn, J. H.; Koda, V. H.; Pick, L.; McKay,K. E.; Knott, P.: Noradrenergic mechanisms in ADHD children withand without reading disabilities: a replication and extension. *J.Am. Acad. Child Adolesc. Psy–*

chiat. 36: 1688–1697, 1997.

- [80156] 18486.Hein, L.; Altman, J. D.; Kobilka, B. K.: Two functionally distinct  $\alpha$ -2-adrenergic receptors regulate sympathetic neurotransmission. *Nature* 402:181–184, 1999.
- [80157] 18487.Hoehe, M.; Berrettini, W.; Leppert, M.; Lalouel, J.-M.; Byerley, W.; Gershon, E.; White, R.: Genetic mapping of adrenergic receptor genes. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A143 only, 1989.
- [80158] 18488.Hoehe, M. R.; Berrettini, W. H.; Lentes, K.-U.: Dra I identifies a two allele DNA polymorphism in the human  $\alpha$ -2-adrenergic receptor gene (ADRAR), using a 5.5 kb probe (p ADRAR). *Nucleic Acids Res.* 16:9070 only, 1988.
- [80159] 18489.Kobilka, B. K.; Matsui, H.; Kobilka, T. S.; Yang-Feng, T. L.; Francke, U.; Caron, M. G.; Lefkowitz, R. J.; Regan, J. W.: Cloning, sequencing, and expression of the gene coding for the human platelet  $\alpha$ -2-adrenergic receptor. *Science* 238: 650–656, 1987.
- [80160] 18490.Oakey, R. J.; Caron, M. G.; Lefkowitz, R. J.; Seldin, M. F.: Genomic organization of adrenergic and serotonin receptors in the mouse: linkage mapping of sequence-related genes provides a method for examining mammalian chromosome evolution. *Genomics* 10: 338–344, 1991.
- [80161] 18491.Philipp, M.; Brede, M. E.; Hadamek, K.; Gessler, M.;

Lohse, M.J.; Hein, L.: Placental alpha-2-adrenoceptors control vascular development at the interface between mother and embryo. *Nature Genet.* 31: 311–315, 2002.

[80162] 18492. Surprenant, A.; Horstman, D. A.; Akbarali, H.; Limbird, L. E.: A point mutation of the alpha-2-adrenoceptor that blocks coupling to potassium but not calcium currents. *Science* 257: 977–980, 1992.

[80163] 18493. Yang-Feng, T. L.; Kobilka, B. K.; Caron, M. G.; Lefkowitz, R.J.; Francke, U.: Chromosomal assignment of genes for an alpha-adrenergic receptor (ADRAR) and for another member of this receptor family coupled to guanine nucleotide regulatory proteins (RG21). (Abstract) *Cytogenet. Cell Genet.* 46: 722–723, 1987.

[80164] 18494. Bruno, J. F.; Whittaker, J.; Song, J.; Berelowitz, M.: Molecular cloning and sequencing of a cDNA encoding a human alpha-1A adrenergic receptor. *Biochem. Biophys. Res. Commun.* 179: 1485–1490, 1991.

[80165] 18495. Esbenshade, T. A.; Hirasawa, A.; Tsujimoto, G.; Tanaka, T.; Yano, J.; Minneman, K. P.; Murphy, T. J.: Cloning of the human alpha-1D-adrenergic receptor and inducible expression of three subtypes in SK-N-MC cells. *Molec. Pharm.* 47: 977–985, 1995.

[80166] 18496. Forray, C.; Bard, J. A.; Wetzel, J. M.; Chiu, G.;

Shapiro, E.; Tang, R.; Lepor, H.; Hartig, P. R.; Weinshank, R. L.; Branchek, T.A.; Gluchowski, C.: The alpha-1-adrenergic receptor that mediates smooth muscle contraction in human prostate has the pharmacological properties of the cloned human alpha-1C subtype. *Molec. Pharm.* 45:703–708, 1994.

[80167] 18497. Loftus, S. K.; Shiang, R.; Warrington, J. A.; Bengtsson, U.; McPherson, J. D.; Wasmuth, J. J.: Genes encoding adrenergic receptors are not clustered on the long arm of human chromosome 5. *Cytogenet. Cell Genet.* 67: 69–74, 1994.

[80168] 18498. Lomasney, J. W.; Cotecchia, S.; Lorenz, W.; Leung, W.-Y.; Schwinn, D. A.; Yang-Feng, T. L.; Brownstein, M.; Lefkowitz, R. J.; Caron, M. G.: Molecular cloning and expression of the cDNA for the alpha-1A-adrenergic receptor: the gene for which is located on human chromosome 5. *J. Biol. Chem.* 266: 6365–6369, 1991.

[80169] 18499. Schwinn, D. A.; Johnston, G. I.; Page, S. O.; Mosley, M. J.; Wilson, K. H.; Worman, N. P.; Campbell, S.; Fidock, M. D.; Furness, L. M.; Parry-Smith, D. J.; Peter, B.; Bailey, D. S.: Cloning and pharmacological characterization of human alpha-1 adrenergic receptors: sequence corrections and



direct comparison with other species homologues. J.

Pharm. Exp. Ther. 272: 134–142, 1995.

[80170] 18500. Schwinn, D. A.; Lomasney, J. W.: Pharmacologic characterization of cloned alpha-1-adrenoceptor subtypes: selective antagonists suggest the existence of a fourth subtype. Europ. J. Pharm. 227: 433–436, 1992.

[80171] 18501. Weinberg, D. H.; Trivedi, P.; Tan, C. P.; Mitra, S.; Perkins-Barrow, A.; Borkowski, D.; Strader, C. D.; Bayne, M.: Cloning, expression and characterization of human alpha adrenergic receptors alpha-1A, alpha-1B, and alpha-1C. Biochem. Biophys. Res. Commun. 201: 1296–1304, 1994.

[80172] 18502. Yang-Feng, T. L.; Han, H.; Lomasney, J. W.; Caron, M. G.: Localization of the cDNA for an alpha-1-adrenergic receptor subtype (ADRA1D) to chromosome band 20p13. Cytogenet. Cell Genet. 66: 170–171, 1994.

[80173] 18503. Allen, L. F.; Lefkowitz, R. J.; Caron, M. G.; Cotecchia, S.: G-protein-coupled receptor genes as protooncogenes: constitutively activating mutation of the alpha-1B-adrenergic receptor enhances mitogenesis and tumorigenicity. Proc. Nat. Acad. Sci. 88: 11354–11358, 1991.

[80174] 18504. Markert, M. L.; Hershfield, M. S.; Wiginton, D. A.;

States, J.C.; Ward, F. E.; Bigner, S. H.; Buckley, R. H.; Kaufman, R. E.; Hutton, J. J.: Identification of a deletion in the adenosine deaminase gene in a child with severe combined immunodeficiency. *J. Immun.* 138:3203–3206, 1987.

[80175] 18505. Markert, M. L.; Hutton, J. J.; Wiginton, D. A.; States, J. C.; Kaufman, R. E.: Adenosine deaminase (ADA) deficiency due to deletion of the ADA gene promoter and first exon by homologous recombination between two Alu elements. *J. Clin. Invest.* 81: 1323–1327, 1988.

[80176] 18506. Markert, M. L.; Norby-Slycord, C.; Ward, F. E.: A high proportion of ADA point mutations associated with a specific alanine-to-valine substitution. *Am. J. Hum. Genet.* 45: 354–361, 1989.

[80177] 18507. Meuwissen, H. J.; Pollara, B.; Pickering, R. J.: Combined immunodeficiency disease associated with adenosine deaminase deficiency (report on a workshop held in Albany, New York, October 1, 1973). *J. Pediat.* 86:169–181, 1975.

[80178] 18508. Migchielsen, A. A. J.; Breuer, M. L.; van Roon, M. A.; te Riele, H.; Zurcher, C.; Ossendorp, F.; Toutain, S.; Hersfield, M. S.; Berns, A.; Valerio, D.: Adenosine-deaminase-deficient mice die perinatally and exhibit liver-cell degeneration, atelectasis and small intestinal cell death.

Nature Genet. 10: 279–287, 1995.

[80179] 18509.Mitchell, B. S.; Mejias, E.; Daddona, P. E.; Kelley, W. N.: Purinogenicimmunodeficiency disease: selective toxicity of deoxyribonucleosidesfor T–cells. Proc. Nat. Acad. Sci. 75: 5011–5014, 1978.

[80180] 18510.Moen, R. C.; Horowitz, S. D.; Sondel, P. M.; Borcharding, W. R.;Trigg, M. E.; Billing, R.; Hong, R.: Immunologic reconstitution afterhaploidentical bone marrow transplantation for immune deficiency disorders:treatment of bone marrow cells with monoclonal antibody CT–2 and complement. Blood 70:664–669, 1987.

[80181] 18511.Mohandas, T.; Sparkes, R. S.; Suh, E. J.; Hershfield, M. S.:Regional localization of the human genes for S–adenosylhomocysteinehydrolase (cen–q131) and adenosine deaminase (q131–qter) on chromosome20. Hum. Genet. 66: 292–295, 1984.

[80182] 18512.Nielsen, K. B.; Tommerup, N.; Jespersen, B.; Nygaard, P.; Kleif,L.: Segregation of a t(3;20) translocation through three generationsresulting in unbalanced karyotypes in six persons. J. Med. Genet. 23:446–451, 1986.

[80183] 18513.Orkin, S. H.; Daddona, P. E.; Shewach, D. S.; Markham, A. F.;Bruns, G. A.; Goff, S. C.; Kelley, W. N.: Molecular cloning of humanadenosine deaminase gene

sequences. J. Biol. Chem. 258: 12753–12756,1983.

[80184] 18514.Ozsahin, H.; Arredondo–Vega, F. X.; Santisteban, I.; Fuhrer, H.;Tuchschmid, P.; Jochum, W.; Aguzzi, A.; Leder–man, H. M.; Fleischman,A.; Winkelstein, J. A.; Seger, R. A.; Hershfield, M. S.: Adenosinedeaminase deficiency in adults. Blood 89: 2849–2855, 1997.

[80185] 18515.Onodera, M.; Ariga, T.; Kawamura, N.; Kobayashi, I.; Ohtsu, M.;Yamada, M.; Tame, A.; Furuta, H.; Okano, M.; Matsumoto, S.; Kotani,H.; McGarrity, G. J.; Blaese, R. M.; Sakiyama, Y.: Successful peripheralT–lymphocyte–directed gene transfer for a patient with severe combinedimmune deficiency caused by adenosine deaminase deficiency. Blood 91:30–36, 1998.

[80186] 18516.Palmer, T. D.; Hock, R. A.; Osborne, W. R. A.; Miller, A. D.:Efficient retrovirus–mediated transfer and expres–sion of a human adenosinedeaminase gene in diploid skin fibroblasts from an adenosine deaminase–deficienthuman. Proc. Nat. Acad. Sci. 84: 1055–1059, 1987.

[80187] 18517.Parkman, R.; Gelfand, E. W.; Rosen, F. S.; Sander–son, A.; Hirschhorn,R.: Severe combined immunodeficiency and adenosine deaminase deficiency. NewEng. J. Med. 292: 714–719, 1975.

[80188] 18518.Petersen, M. B.; Tranebjaerg, L.; Tommerup, N.;

Nygaard, P.; Edwards, H.: New assignment of the adenosine deaminase gene locus to chromosome 20q13.11 by study of a patient with interstitial deletion 20q. *J. Med. Genet.* 24: 93–96, 1987.

[80189] 18519. Polmar, S. H.; Stern, R. C.; Schwartz, A. L.; Wetzler, E. M.; Chase, P. A.; Hirschhorn, R.: Enzyme replacement therapy for adenosine deaminase deficiency and severe combined immunodeficiency. *New Eng. J. Med.* 295: 1337–1343, 1976.

[80190] 18520. Ratech, H.; Greco, M. A.; Gallo, G.; Rimoin, D. L.; Kamino, H.; Hirschhorn, R.: Pathologic findings in adenosine–deaminase–deficient severe combined immunodeficiency. I. Kidney, adrenal, and chondro–osseous tissue alterations. *Am. J. Path.* 120: 157–169, 1985.

[80191] 18521. Ritter, H.; Wendt, G. G.; Tariverdian, G.; Zelch, J.; Rube, M.; Kirchberg, G.: Genetics and linkage analysis of adenosine deaminase. *Humangenetik* 14:69–71, 1971.

[80192] 18522. Rogers, M. H.; Lwin, R.; Fairbanks, L.; Gerritsen, B.; Gaspar, H. B.: Cognitive and behavioral abnormalities in adenosine deaminase deficient severe combined immunodeficiency. *J. Pediat.* 139: 44–50, 2001.

[80193] 18523. Rothschild, C. B.; Akots, G.; Hayworth, R.; Pettenati, M. J.; Rao, P. N.; Wood, P.; Stolz, F.–M.; Hansmann, I.;

Serino, K.; Keith, T. P.; Fajans, S. S.; Bowden, D. W.: A genetic map of chromosome 20q12–q13.1: multiple highly polymorphic microsatellite and RFLP markers linked to the maturity-onset diabetes of the young (MODY) locus. *Am. J. Hum. Genet.* 52: 110–123, 1993.

[80194] 18524. Rubinstein, A.; Hirschhorn, R.; Sicklick, M.; Murphy, R. A.: In vivo and in vitro effects of thymosin and adenosine deaminase on adenosine-deaminase-deficient lymphocytes. *New Eng. J. Med.* 300:387–392, 1979.

[80195] 18525. Rudd, N. L.; Bain, H. W.; Giblett, E.; Chen, S.-H.; Worton, R. G.: Partial trisomy 20 confirmed by gene dosage studies. *Am. J. Med. Genet.* 4: 357–364, 1979.

[80196] 18526. Koch, G.; Shows, T. B.: Somatic cell genetics of adenosine deaminase expression and severe combined immune deficiency disease in man. *Proc. Nat. Acad. Sci.* 77: 4211–4215, 1980.

[80197] 18527. Gautam, M.; Noakes, P. G.; Moscoso, L.; Rupp, F.; Scheller, R. H.; Merlie, J. P.; Sanes, J. R.: Defective neuromuscular synaptogenesis in agrin-deficient mutant mice. *Cell* 85: 525–535, 1996.

[80198] 18528. Glass, D. J.; Bowen, D. C.; Stitt, T. N.; Radziejewski, C.; Bruno, J.; Ryan, T. E.; Gies, D. R.; Shah, S.; Mattsson, K.; Burden, S. J.; DiStefano, P. S.; Valenzuela, D. M.; DeChiara,

T. M.; Yancopoulos, G. D.: Agrin acts via a MuSK receptor complex. *Cell* 85: 513–523, 1996.

- [80199] 18529. Khan, A. A.; Bose, C.; Yam, L. S.; Soloski, M. J.; Rupp, F.: Physiological regulation of the immunological synapse by agrin. *Science* 292: 1681–1686, 2001.
- [80200] 18530. Lin, W.; Burgess, R. W.; Dominguez, B.; Pfaff, S. L.; Sanes, J. R.; Lee, K.-F.: Distinct roles of nerve and muscle in postsynaptic differentiation of the neuromuscular synapse. *Nature* 410: 1057–1064, 2001.
- [80201] 18531. McMahan, U. J.: The agrin hypothesis *Cold Spring Harb. Symp. Quant. Biol.* 50: 407–418, 1990.
- [80202] 18532. Rupp, F.; Ozcelik, T.; Linial, M.; Peterson, K.; Francke, U.; Scheller, R.: Structure and chromosomal localization of the mammalian agrin gene. *J. Neurosci.* 12: 3535–3544, 1992.
- [80203] 18533. Rupp, F.; Payan, D. G.; Magill-Solc, C.; Cowan, D. M.; Scheller, R. H.: Structure and expression of a rat agrin. *Neuron* 6: 811–823, 1991.
- [80204] 18534. Umans, L.; Serneels, L.; Overbergh, L.; Lorent, K.; Van Leuven, F.; Van den Berghe, H.: Targeted inactivation of the mouse alpha-2-macroglobulin gene. *J. Biol. Chem.* 270: 19778–19785, 1995.
- [80205] 18535. D'Alfonso, S.; Richiardi, P. M.: A polymorphic varia-

tion in a putative regulation box of the TNFA promoter region. Immunogenetics 39:150–154, 1994.

- [80206] 18536. Mietus-Snyder, M.; Charmley, P.; Korf, B.; Ladas, J. A. A.; Gatti, R. A. and Karathanasis, S. K.: Genetic linkage of the human apolipoprotein A1–CIII–AIV gene cluster and the neural cell adhesion molecule (NCAM) gene. Genomics 7: 633–637, 1990.
- [80207] 18537. Mietus-Snyder, M.; Korf, B.; Ladas, J. A.; Karathanasis, S. K.: Linkage of the human apolipoproteins A1, C3, A4 and the neural cell adhesion molecule (NCAM) genes. (Abstract) Cytogenet. Cell Genet. 51:1044 only, 1989.
- [80208] 18538. Nguyen, C.; Mattei, M. G.; Goridis, C.; Mattei, J. F.; Jordan, B. R.: Localization of the human NCAM gene to chromosome 11 by insitu hybridization with a murine NCAM cDNA probe. (Abstract) Cytogenet. Cell Genet. 40: 713 only, 1985.
- [80209] 18539. Nguyen, C.; Mattei, M. G.; Mattei, J.-F.; Santoni, M.-J.; Goridis, C.; Jordan, B. R.: Localization of the human NCAM gene to band q23 of chromosome 11: the third gene coding for a cell interaction molecule mapped to the distal portion of the long arm of chromosome 11. J. Cell Biol. 102: 711–715, 1986.



- [80210] 18540.Rabinowitz, J. E.; Rutishauser, U.; Magnuson, T.: Targeted mutation of Ncam to produce a secreted molecule results in a dominant embryonic lethality. *Proc. Nat. Acad. Sci.* 93: 6421–6424, 1996.
- [80211] 18541.Rutishauser, U.; Acheson, A.; Hall, A. K.; Mann, D. M.; Sunshine, J.: The neural cell adhesion molecule (NCAM) as a regulator of cell–cell interactions. *Science* 240: 53–57, 1988.
- [80212] 18542.Rutishauser, U.; Goridis, C.: NCAM: the molecule and its genetics. *Trends Genet.* 2: 72–76, 1986.
- [80213] 18543.Telatar, M.; Lange, E.; Uhrhammer, N.; Gatti, R. A.: New localization of NCAM, proximal to DRD2 at chromosome 11q23. *Mammalian Genome* 6:59–60, 1995.
- [80214] 18544.Agellon, L. B.; Drover, V. A. B.; Cheema, S. K.; Gbaguidi, G. F.; Walsh, A.: Dietary cholesterol fails to stimulate the human cholesterol 7- $\alpha$ -hydroxylase gene (CYP7A1) in transgenic mice. *J. Biol. Chem.* 277:20131–20134, 2002.
- [80215] 18545.Angelin, B.; Einarsson, K.; Hellstrom, K.; Leijdt, B.: Bile acid kinetics in relation to endogenous triglyceride metabolism in various types of hyperlipoproteinemia. *J. Lipid Res.* 19: 1004–1016, 1978.
- [80216] 18546.Angelin, B.; Hershon, K. S.; Brunzell, J. D.: Bile acid

metabolism in hereditary forms of hypertriglyceridemia: evidence for an increased synthesis rate in monogenic familial hypertriglyceridemia. *Proc. Nat. Acad. Sci.* 84: 5434–5438, 1987.

[80217] 18547. Cohen, J. C.; Cali, J. J.; Jelinek, D. F.; Mehrabian, M.; Sparkes, R. S.; Lusis, A. J.; Russell, D. W.; Hobbs, H. H.: Cloning of the human cholesterol 7- $\alpha$ -hydroxylase gene (CYP7) and localization to chromosome 8q11–q12. *Genomics* 14: 153–161, 1992.

[80218] 18548. Drover, V. A. B.; Wong, N. C. W.; Agellon, L. B.: A distinct thyroid hormone response element mediates repression of the human cholesterol 7- $\alpha$ -hydroxylase (CYP7A1) gene promoter. *Molec. Endocr.* 16: 14–23, 2002.

[80219] 18549. Goodwin, B.; Jones, S. A.; Price, R. R.; Watson, M. A.; McKee, D. D.; Moore, L. B.; Galardi, C.; Wilson, J. G.; Lewis, M. C.; Roth, M. E.; Maloney, P. R.; Willson, T. M.; Kliewer, S. A.: A regulatory cascade of the nuclear receptors FXR, SHP-1, and LRH-1 represses bile acid biosynthesis. *Molec. Cell* 6: 517–526, 2000.

[80220] 18550. Molowa, D. T.; Chen, W. S.; Cimis, G. M.; Tan, C. P.: Transcriptional regulation of the human cholesterol 7  $\alpha$ -hydroxylase gene. *Biochemistry* 31: 2539–2544, 1992.

- [80221] 18551.Nitta, M.; Ku, S.; Brown, C.; Okamoto, A. Y.; Shan, B.: CPF: an orphan nuclear receptor that regulates liver-specific expression of the human cholesterol 7- $\alpha$ -hydroxylase gene. *Proc. Nat. Acad.Sci.* 96: 6660–6665, 1999.
- [80222] 18552.Noshiro, M.; Okuda, K.: Molecular cloning and sequence analysis of cDNA encoding human cholesterol 7  $\alpha$ -hydroxylase. *FEBS Lett.* 268:137–140, 1990.
- [80223] 18553.Paumgartner, G.; Sauerbruch, T.: Gallstones: pathogenesis. *Lancet* 338:1117–1121, 1991.
- [80224] 18554.Wang, J.; Freeman, D. J.; Grundy, S. M.; Levine, D. M.; Guerra, R.; Cohen, J. C.: Linkage between cholesterol 7- $\alpha$ -hydroxylase and high plasma low-density lipoprotein cholesterol concentrations. *J.Clin. Invest.* 101: 1283–1291, 1998.
- [80225] 18555.Akita, H.; Chiba, H.; Tsuchihashi, K.; Tsuji, M.; Kumagai, M.; Matsuno, K.; Kobayashi, K.: Cholesteryl ester transfer protein gene: two common mutations and their effect on plasma high-density lipoprotein cholesterol content. *J. Clin. Endocr. Metab.* 79: 1615–1618, 1994.
- [80226] 18556.Altshuler, D.; Kruglyak, L.; Lander, E.: Genetic polymorphisms and disease. (Letter) *New Eng. J. Med.* 338: 1626 only, 1998.

- [80227] 18557. Brown, M. L.; Inazu, A.; Hesler, C. B.; Agellon, L. B.; Mann, C.; Whitlock, M. E.; Marcel, Y. L.; Milne, R. W.; Koizumi, J.; Mabuchi, H.; Takeda, R.; Tall, A. R.: Molecular basis of lipid transfer protein deficiency in a family with increased high-density lipoproteins. *Nature* 342:448–451, 1989.
- [80228] 18558. Drayna, D.; Jarnagin, A. S.; McLean, J.; Henzel, W.; Kohr, W.; Fielding, C.; Lawn, R.: Cloning and sequencing of human cholesteryl ester transfer protein cDNA. *Nature* 327: 632–634, 1987.
- [80229] 18559. Drayna, D.; Lawn, R. M.: Multiple RFLPs at the human cholesteryl ester transfer protein (CETP) locus. *Nucleic Acids Res.* 15: 4698 only, 1987.
- [80230] 18560. Haefliger, J.-A.; Bruzzone, R.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Paul, D. L.: Four novel members of the connexin family of gap junction proteins: molecular cloning, expression, and chromosome mapping. *J. Biol. Chem.* 267: 2057–2064, 1992.
- [80231] 18561. Hsieh, C.-L.; Kumar, N. M.; Gilula, N. B.; Francke, U.: Distribution of genes for gap junction membrane channel proteins on human and mouse chromosomes. *Somat. Cell Molec. Genet.* 17: 191–200, 1991.
- [80232] 18562. Spielman, R. S.; McGinnis, R. E.; Ewens, W. J.:

Transmission test for linkage disequilibrium: the insulin gene region and insulin-dependent diabetes mellitus (IDDM). *Am. J. Hum. Genet.* 52: 506–516, 1993.

- [80233] 18563. Altshuler, D.; Hirschhorn, J. N.; Klannemark, M.; Lindgren, C. M.; Vohl, M.-C.; Nemesh, J.; Lane, C. R.; Schaffner, S. F.; Bolk, S.; Brewer, C.; Tuomi, T.; Gaudet, D.; Hudson, T. J.; Daly, M.; Groop, L.; Lander, E. S.: The common PPAR- $\gamma$  pro12ala polymorphism is associated with decreased risk of type 2 diabetes. *Nature Genet.* 76–80, 2000.
- [80234] 18564. Fakhrai-Rad, H.; Nikoshkov, A.; Kamel, A.; Fernstrom, M.; Zierath, J. R.; Norgren, S.; Luthman, H.; Galli, J.: Insulin-degrading enzyme identified as a candidate diabetes susceptibility gene in GK rats. *Hum. Molec. Genet.* 9: 2149–2158, 2000.
- [80235] 18565. Gersuk, V. H.; Rose, T. M.; Todaro, G. J.: Molecular cloning and chromosomal localization of a pseudogene related to the human acyl-CoA binding protein/diazepam binding inhibitor. *Genomics* 25: 469–476, 1995.
- [80236] 18566. DeStefano, A. L.; Baldwin, C. T.; Burzstyn, M.; Gavras, I.; Handy, D. E.; Joost, O.; Martel, T.; Nicolaou, M.; Schwartz, F.; Streeten, D. H. P.; Farrer, L. A.; Gavras, H.: Autosomal dominant orthostatic hypotensive disorder

maps to chromosome 18q. Am. J. Hum. Genet.  
63:1425–1430, 1998.

- [80237] 18567. Barnett, T.; Zimmermann, W.: Workshop report: proposed nomenclature for the carcinoembryonic antigen (CEA) gene family. Tumor Biol. 11:59–63, 1990.
- [80238] 18568. Boulton, I. C.; Gray–Owen, S. D.: Neisserial binding to CEACAM1 arrests the activation and proliferation of CD4+ T lymphocytes. Nature Immun. 3: 229–236, 2002.
- [80239] 18569. Ergun, S.; Kilic, N.; Ziegeler, G.; Hansen, A.; Nollau, P.; Gotze, J.; Wurmbach, J.–H.; Horst, A.; Weil, J.; Fernando, M.; Wagener, C.: CEA–related cell adhesion molecule 1: a potent angiogenic factor and a major effector of vascular endothelial growth factor. Molec. Cell 5: 311–320, 2000.
- [80240] 18570. Hinoda, Y.; Neumaier, M.; Hefta, S. A.; Drzeniek, Z.; Wagener, C.; Shively, L.; Hefta, L. J. F.; Shively, J. E.; Paxton, R. J.: Molecular cloning of a cDNA coding biliary glycoprotein I: primary structure of a glycoprotein immunologically crossreactive with carcinoembryonic antigen. Proc. Nat. Acad. Sci. 85: 6959–6963, 1988.
- [80241] 18571. Neumaier, M.; Paululat, S.; Chan, A.; Matthaes, P.; Wagener, C.: Biliary glycoprotein, a potential human cell adhesion molecule, is down–regulated in colorectal carcinomas. Proc. Nat. Acad. Sci. 90:10744–10748, 1993.

- [80242] 18572.Poy, M. N.; Yang, Y.; Rezaei, K.; Fernstrom, M. A.; Lee, A. D.;Kido, Y.; Erickson, S. K.; Najjar, S. M.: CEACAM1 regulates insulin clearance in liver. *Nature Genet.* 30: 270–276, 2002.
- [80243] 18573.Robbins, J.; Robbins, P. F.; Kozak, C. A.; Callahan, R.: Themouse biliary glycoprotein gene (Bgp): partial nucleotide sequence,expression, and chromosomal assignment. *Genomics* 10: 583–587, 1991.
- [80244] 18574.Thompson, J. A.; Grunert, F.; Zimmermann, W.: Carcinoembryonicantigen gene family: molecular biology and clinical perspectives. *J.Clin. Lab. Anal.* 5: 344–366, 1991.
- [80245] 18575.Willard, H. F.; Meakin, S. O.; Tsui, L.–C.; Breitman, M. L.:Assignment of human gamma crystallin multigene family to chromosome2. *Somat. Cell Molec. Genet.* 11: 511–516, 1985.
- [80246] 18576.Wistow, G.; Piatigorsky, J.: Recruitment of enzymes as lens structuralproteins. *Science* 236: 1554–1556, 1987.
- [80247] 18577.Zneimer, S. M.; Womack, J. E.: Regional localization of the fibronectinand gamma crystallin genes to mouse chromosome 1 by in situ hybridization. *Cytogenet.Cell Genet.* 48: 238–241, 1988.
- [80248] 18578.Daniels, G.; King, M.–J.; Avent, N. D.; Khalid, G.;

Reid, M.;Mallinson, G.; Symthe, J.; Cedergren, B.: A point mutation in theGYPC gene results in the expression of the blood group An(a) antigenon glycophorin D but not on glycophorin C: further evidence that glycophorinD is a product of the GYPC gene. Blood 82: 3198–3203, 1993.

[80249] 18579.El-Maliki, B.; Blanchard, D.; Dahr, W.; Beyreuther, K.; Cartron,J.-P.: Structural homology between gly-cophorins C and D of humanerythrocytes. Europ. J. Biochem. 183: 639–643, 1989.

[80250] 18580.Jorgensen, J.; Drachmann, O.; Gavin, J.: Duch, Dh(a), a low frequencyred cell antigen. Hum. Hered. 32: 73–75, 1982.

[80251] 18581.King, M. J.; Avent, N. D.; Mallinson, G.; Reid, M. E.: Pointmutation in the glycophorin C gene results in the ex-pression of theblood group antigen Dh(a). Vox Sang. 63: 56–58, 1992.

[80252] 18582.Le Van Kim, C.; Colin, Y.; Blanchard, D.; Dahr, W.; London, J.;Cartron, J.-P.: Gerbich blood group deficiency of the Ge:–1,–2,–3and Ge:–1,–2,3 types: immunochemical study and genomic analysis withcDNA probes. Europ. J. Biochem. 165: 571–579, 1987.

[80253] 18583.Mattei, M. G.; Colin, Y.; Le Van Kim, C.; Mattei, J. F.; Cartron,J. P.: Localization of the gene for human erythro–



cyte glycoprotein C to chromosome 2, q14-q21. *Hum. Genet.* 74: 420-422, 1986.

- [80254] 18584. Pasvol, G.; Anstee, D. J.; Tanner, M. J. A.: Glycophorin C and the invasion of red cells by *Plasmodium falciparum*. *Lancet* I: 907-908, 1984.
- [80255] 18585. Race, R. R.; Sanger, R.: *Blood Groups in Man*. Oxford: Blackwell Sci. Publ. (pub.) (6th ed.): 1975. Pp. 416-421.
- [80256] 18586. Reid, M. E.: The Gerbich blood group antigens: a review. *Med. Lab. Sci.* 43: 177-182, 1972.
- [80257] 18587. Reid, M. E.; Sullivan, C.; Taylor, M.; Anstee, D. J.: Inheritance of human-erythrocyte Gerbich blood group antigens. *Am. J. Hum. Genet.* 41: 1117-1123, 1987.
- [80258] 18588. Simmons, R. T.; Albrey, J. A.: A 'new' blood group antigen Webb (Wb) of low frequency found in two Australian families. *Med. J. Aust.* I: 8-10, 1963.
- [80259] 18589. Sondag, D.; Alloisio, N.; Blanchard, D.; Ducluzeau, M.-T.; Colonna, P.; Bachir, D.; Bloy, C.; Cartron, J.-P.; Delaunay, J.: Gerbich reactivity in 4.1(-) hereditary elliptocytosis and protein 4.1 level in blood group Gerbich deficiency. *Brit. J. Haemat.* 65: 43-50, 1987.
- [80260] 18590. Spring, F. A.: Immunochemical characterisation of the low-incidence antigen, Dh(a). *Vox Sang.* 61: 65-68,

1991.

- [80261] 18591.Telen, M. J.; Le Van Kim, C.; Guizzo, M. L.; Cartron, J.-P.; Colin,Y.: Erythrocyte Webb-type glycophorin C variant lacks N-glycosylation due to an asparagine to serine substitution. *Am. J. Hemat.* 37: 51-52,1991.
- [80262] 18592.Winardi, R.; Reid, M.; Conboy, J.; Mohandas, N.: Molecular analysis of glycophorin C deficiency in human erythrocytes. *Blood* 81: 2799-2803,1993.
- [80263] 18593.Bierhuizen, M. F. A.; Mattei, M.-G.; Fukuda, M.: Expression of the developmental I antigen by a cloned human cDNA encoding a member of a beta-1,6-N-acetylglucosaminyltransferase gene family. *Genes-Dev.* 7: 468-478, 1993.
- [80264] 18594.Lin-Chu, M.; Broadberry, R. E.; Okubo, Y.; Tanaka, M.: The i phenotype and congenital cataracts among Chinese in Taiwan (Letter) *Transfusion* 31:676-677, 1991.
- [80265] 18595.Ogata, H.; Okubo, Y.; Akabane, T.: Phenotype i associated with congenital cataract in Japanese. *Transfusion* 19: 166-168, 1979.
- [80266] 18596.Yeh, J.-C.; Ong, E.; Fukuda, M.: Molecular cloning and expression of a novel beta-1,6-N-acetylglucosaminyltransferase that forms core2, core 4, and I branches. *J. Biol. Chem.* 274: 3215-3221,

1999.

- [80267] 18597.Yu, L.-C.; Twu, Y.-C.; Chang, C.-Y.; Lin, M.: Molecular basis of the adult i phenotype and the gene responsible for the expression of the human blood group I antigen. *Blood* 98: 3840–3845, 2001.
- [80268] 18598.Allen, F. H., Jr.; Krabbe, S. M.; Corcoran, P. A.: A new phenotype (McLeod) in the Kell blood-group system. *Vox Sang.* 6: 555–560, 1961.
- [80269] 18599.Avent, N. D.; Martin, P. G.: Kell typing by allele-specific PCR (ASP). *Brit. J. Haemat.* 93: 728–730, 1996.
- [80270] 18600.Chown, B.; Lewis, M.; Kaita, K.: A 'new' Kell blood-group phenotype. *Nature* 180:711 only, 1957.
- [80271] 18601.Conneally, P. M.; Nance, W. E.; Huntzinger, R. S.: Linkage analysis of Kell–Sutter and PTC loci. (Abstract) *Am. J. Hum. Genet.* 26: 22A only, 1974.
- [80272] 18602.Abdelhak, S.; Kalatzis, V.; Heilig, R.; Compain, S.; Samson, D.; Vincent, C.; Weil, D.; Cruaud, C.; Sahly, I.; Leibovici, M.; Bitner-Glindzicz, M.; Francis, M.; Lacombe, D.; Vigneron, J.; Charachon, R.; Boven, K.; Bedbeder, P.; Van Regemorter, N.; Weissenbach, J.; Petit, C.: A human homologue of the *Drosophila* eyes absent gene underlies branchio-oto-renal (BOR) syndrome and identifies a novel gene family. *Nature Genet.* 15:157–164, 1997.

- [80273] 18603.Tommerup, N.; Schempp, W.; Meinecke, P.; Peder-  
sen, S.; Bolund,L.; Brandt, C.; Goodpasture, C.; Guldberg,  
P.; Held, K.; Reinwein,H.; Saugstad, O. D.; Scherer, G.;  
Skjeldal, O.; Toder, R.; Westvik,J.; van der Hagen, C. B.;  
Wolf, U.: Assignment of an autosomal sexreversal locus  
(SRA1) and campomelic dysplasia (CMPD1) to  
17q24.3–q25.1. NatureGenet. 4: 170–174, 1993.
- [80274] 18604.Azuma, N.; Nishina, S.; Yanagisawa, H.; Okuyama,  
T.; Yamada, M.: PAX6 missense mutation in isolated foveal  
hypoplasia. (Letter) NatureGenet. 13: 141–142, 1996.
- [80275] 18605.Curran, R. E.; Robb, R. M.: Isolated foveal hypopla-  
sia. Arch.Ophthal. 94: 48–50, 1976.
- [80276] 18606.O'Donnell, F. E., Jr.; Pappas, H. R.: Autosomal dom-  
inant fovealhypoplasia and presenile cataracts: a new syn-  
drome. Arch. Ophthal. 100:279–281, 1982.
- [80277] 18607.Anderson, M. J.; Shelton, G. D.; Cavenee, W. K.; Ar-  
den, K. C.:Embryonic expression of the tumor-associated  
PAX3–FKHR fusion proteininterferes with the developmen-  
tal functions of Pax3. Proc. Nat. Acad.Sci. 98: 1589–1594,  
2001.
- [80278] 18608.Anderson, M. J.; Viars, C. S.; Czekay, S.; Cavenee,  
W. K.; Arden,K. C.: Cloning and characterization of three  
human forkhead genesthat comprise an FKHR–like gene

subfamily. Genomics 47: 187–199,1998.

[80279] 18609.Barr, F. G.; Galili, N.; Holick, J.; Biegel, J. A.; Rovera, G.; Emanuel, B. S.: Rearrangement of the PAX3 paired box gene in the paediatric solid tumor alveolar rhabdomyosarcoma. Nature Genet. 3:113–117, 1993.

[80280] 18610.Davis, R. J.; Barr, F. G.: Fusion genes resulting from alternative chromosomal translocations are overexpressed by gene-specific mechanisms in alveolar rhabdomyosarcoma. Proc. Nat. Acad. Sci. 94: 8047–8051,1997.

[80281] 18611.Fredericks, W. J.; Galili, N.; Mukhopadhyay, S.; Rovera, G.; Bennicelli, J.; Barr, F. G.; Rauscher, F. J., III: The PAX3–FKHR fusion protein created by the t(2;13) translocation in alveolar rhabdomyosarcoma is a more potent transcriptional activator than PAX3. Molec. Cell.Biol. 15: 1522–1535, 1995.

[80282] 18612.Galili, N.; Davis, R. J.; Fredericks, W. J.; Mukhopadhyay, S.; Rauscher, F. J., III; Emanuel, B. S.; Rovera, G.; Barr, F. G.: Fusion of a fork head domain gene to PAX3 in the solid tumour alveolar rhabdomyosarcoma. Nature Genet. 5: 230–235, 1993.

[80283] 18613.Khan, J.; Bittner, M. L.; Saal, L. H.; Teichmann, U.; Azorsa, D.O.; Gooden, G. C.; Pavan, W. J.; Trent, J. M.; Meltzer, P. S.: cDNA microarrays detect activation of a

myogenic transcription program by the PAX3–FKHR fusion oncogene. *Proc. Nat. Acad. Sci.* 96: 13264–13269, 1999.

[80284] 18614. Angle, C. R.: Congenital bowing and angulation of the long bones. *Pediatrics* 13:257–268, 1954.

[80285] 18615. Bain, A. D.; Barrett, H. S.: Congenital bowing of the long bones: report of a case. *Arch. Dis. Child.* 34: 516–524, 1959.

[80286] 18616. Bell, D. M.; Leung, K. K. H.; Wheatley, S. C.; Ng, L. J.; Zhou, S.; Ling, K. W.; Sham, M. H.; Koopman, P.; Tam, P. P. L.; Cheah, K. S. E.: SOX9 directly regulates the type-II collagen gene. *Nature Genet.* 16: 174–178, 1997.

[80287] 18617. Bi, W.; Huang, W.; Whitworth, D. J.; Deng, J. M.; Zhang, Z.; Behringer, R. R.; de Crombrughe, B.: Haploinsufficiency of Sox9 results in defective cartilage primordia and premature skeletal mineralization. *Proc. Nat. Acad. Sci.* 98: 6698–6703, 2001.

[80288] 18618. Bishop, C. E.; Whitworth, D. J.; Qin, Y.; Agoulnik, A. I.; Agoulnik, I. U.; Harrison, W. R.; Behringer, R. R.; Overbeek, P. A.: A transgenic insertion upstream of Sox9 is associated with dominant XX sex reversal in the mouse. *Nature Genet.* 26: 490–494, 2000.

[80289] 18619. Caffey, J. P.: Prenatal bowing and thickening of tubular bones, with multiple cutaneous dimples in arms

and legs: a congenital syndrome of mechanical origin. Am. J. Dis. Child. 74: 543–562, 1947.

- [80290] 18620. Cameron, F. J.; Hageman, R. M.; Cooke–Yarborough, C.; Kwok, C.; Goodwin, L. L.; Sillence, D. O.; Sinclair, A. H.: A novel germ line mutation in SOX9 causes familial campomelic dysplasia and sex reversal. Hum. Molec. Genet. 5: 1625–1630, 1996.
- [80291] 18621. Cameron, F. J.; Sinclair, A. H.: Mutations in SRY and SOX9: testis–determining genes. Hum. Mutat. 9: 388–395, 1997.
- [80292] 18622. Cooke, C. T.; Mulcahy, M. T.; Cullity, G. J.; Watson, M.; Sprague, P.: Campomelic dysplasia with sex reversal: morphological and cytogenetic studies of a case. Pathology 17: 526–529, 1985.
- [80293] 18623. Cremin, B. J.; Orsmond, G.; Beighton, P.: Autosomal recessive inheritance in campomelic dwarfism. (Letter) Lancet I: 488–489, 1973.
- [80294] 18624. Dagna Bricarelli, F.; Fraccaro, M.; Lindsten, J.; Muller, U.; Baggio, P.; Carbone, L. D. L.; Hjerpe, A.; Lindgren, F.; Mayerova, A.; Ringertz, H.; Ritzen, E. M.; Rovetta, D. C.; Siccherò, C.; Wolf, U.: Sex–reversed XY females with campomelic dysplasia are H–Y negative. Hum. Genet. 57: 15–22, 1981.

- [80295] 18625.Ebensperger, C.; Jager, R. J.; Lattermann, U.; Dagna Bricarelli,F.; Keutel, J.; Lindsten, J.; Rehder, H.; Muller, U.; Wolf, U.: Noevidence of mutations in four candidate genes for male sex determination/differentiationin sex-reversed XY females with campomelic dysplasia. *Ann. Genet.* 34:233–238, 1991.
- [80296] 18626.Fontaine, G.; Walbaum, R.; Farriaux, J. P.; Tilmont, P.; Peuzin,F.; Delecour, M.: Le conseil genetique dans la dysplasie campomelique(a propos de deux observations). *J. Genet. Hum.* 28: 267–279, 1980.
- [80297] 18627.Foster, J. W.; Dominguez–Steglich, M. A.; Guioli, S.; Kwok, C.;Weller, P. A.; Stevanovic, M.; Weissenbach, J.; Mansour, S.; Young,I. D.; Goodfellow, P. N.; Brook, J. D.; Schafer, A. J.: Campomelicdysplasia and autosomal sex reversal caused by mutations in an SRY–relatedgene. *Nature* 372: 525–530, 1994.
- [80298] 18628.Friedrich, U.; Schaefer, E.; Meinecke, P.: Cam–pomelic dysplasiawithout overt campomelia. *Clin. Dys–morph.* 1: 172–178, 1992.
- [80299] 18629.Gasca, S.; Canizares, J.; de Santa Barbara, P.; Mejean, C.; Poulat,F.; Berta, P.; Boizet–Bonhoure, B.: A nuclear export signal withinthe high mobility group domain regulates the nucleocytoplasmic translocationof SOX9



during sexual determination. Proc. Nat. Acad. Sci. 99: 11199–11204, 2002.

- [80300] 18630. Glass, R. B. J.; Rosenbaum, K. N.: Acampomelic campomelic dysplasia: further radiographic variations. Am. J. Med. Genet. 69: 29–32, 1997.
- [80301] 18631. Hall, B.; Spranger, J. W.: Campomelic dysplasia: further elucidation of a distinct entity. Am. J. Dis. Child. 134: 285–289, 1980.
- [80302] 18632. Hoefnagel, D.; Wurster-Hill, D. H.; Dupree, W. B.; Benirschke, K.; Fuld, G. L.: Campomelic dwarfism associated with XY-gonadal dysgenesis and chromosome anomalies. Clin. Genet. 13: 489–499, 1978.
- [80303] 18633. Houston, C. S.; Opitz, J. M.; Spranger, J. W.; Macpherson, R. I.; Reed, M. H.; Gilbert, E. F.; Herrmann, J.; Schinzel, A.: The campomelic syndrome: review, report of 17 cases, and follow-up on the currently 17-year-old boy first reported by Maroteaux et al in 1971. Am. J. Med. Genet. 15: 3–28, 1983.
- [80304] 18634. Hovmoller, M. L.; Osuna, A.; Eklof, O.; Fredga, K.; Hjerpe, A.; Lindsten, J.; Ritzen, M.; Stanescu, V.; Svenningsen, N.: Campomelic dwarfism. A genetically determined mesenchymal disorder combined with sex reversal. Hereditas 86: 51–62, 1977.

- [80305] 18635.Huang, W.; Chung, U.; Kronenberg, H. M.; de Crombrughe, B.:The chondrogenic transcription factor Sox9 is a target of signalingby the parathyroid hormone–related peptide in the growth plate ofendochondral bones. Proc. Nat. Acad. Sci. 98: 160–165, 2001.
- [80306] 18636.Kanai, Y.; Koopman, P.: Structural and functional characterizationof the mouse Sox9 promoter: implications for campomelic dysplasia. Hum.Molec. Genet. 8: 691–696, 1999.
- [80307] 18637.Medema, R. H.; Kops, G. J. P. L.; Bos, J. L.; Burger–ing, B. M.T.: AFX–like forkhead transcription factors medi–ate cell–cycle regulationby Ras and PKB through p27(kip1). Nature 404: 782–787, 2000.
- [80308] 18638.Eicher, E. M.; Womack, J. E.: Chromosomal location of solubleglutamic–pyruvic transaminase–1 (Gpt–1) in the mouse. Biochem. Genet. 15:1–8, 1977.
- [80309] 18639.Falk, C. T.; Huss, J.: Linkage data on the chromo–some 16 markersHP and PGP: no additional support for or against the mapping of GPT.(Abstract) Cytogenet. Cell Genet. 40: 626 only, 1985.
- [80310] 18640.Ferrell, R. E.: Personal Communication. Pittsburgh, Pa. 1/22/1988.
- [80311] 18641.Jeremiah, S. J.; Kielty, C.; Povey, S.; McLellan, T.:

Immunological characterization of human GPT in hybrids  
supports assignment to chromosome

- [80312] 18642.(Abstract) Cytogenet. Cell Genet. 37: 498 only,  
1984.8. Kalimanovska, V.; Majkic-Singh, N.; Stojanov, M.;  
Grozdanic, V.; Vucetic, G.; Andelic, M.; Gligorovic, V.;  
Tomasevic, R.: Human redcell glutamic-pyruvic transami-  
nase polymorphism in Serbia, Yugoslavia. Hum.Hered. 33:  
319-321, 1983.
- [80313] 18643.Kielty, C. M.; Povey, S.; Hopkinson, D. A.: Regula-  
tion of expression of liver-specific enzymes: II. Activation  
and chromosomal localization of soluble glutamate-pyru-  
vate transaminase. Ann. Hum. Genet. 46:135-143, 1982.
- [80314] 18644.King, M.-C.; Go, R. C. P.; Elston, R. C.; Lynch, H. T.;  
Petrakis, N. L.: Allele increasing susceptibility to human  
breast cancer maybe linked to the glutamate-pyruvate  
transaminase locus. Science 208:406-408, 1980.
- [80315] 18645.Kompf, J.: Population genetics of soluble glutamic-  
pyruvic transaminase (EC:2.6.1.2): gene frequencies in  
southwestern Germany. Humangenetik 14:76-77, 1971.
- [80316] 18646.Lahav, M.; Szeinberg, A.: A red-cell glutamic-pyru-  
vic transaminase polymorphism in several population  
groups in Israel. Hum. Hered. 22:533-538, 1972.
- [80317] 18647.Marazita, M. L.; Spence, M. A.; Sparkes, R. S.; Field,

L. L.;Crandall, B. F.; Sparkes, M. C.; Crist, M.: Linkage relations ofGPT (glutamic-pyruvate transaminase). (Abstract) Cytogenet. CellGenet. 40: 690 only, 1985.

[80318] 18648.McLellan, T.: Two previously undetected variants of glutamic-pyruvictransaminase found by acidic polyacrylamide gel electrophoresis. Am.J. Hum. Genet. 34: 623-628, 1982.

[80319] 18649.McLellan, T.; Cannon, L. A.; Bishop, D. T.; Skolnick, M. H.:The cumulative lod score between a breast cancer susceptibility locusand GPT is -3.86. (Abstract) Cytogenet. Cell Genet. 37: 536-537,1984.

[80320] 18650.Mithal, Y.; Lane, A. B.; Jenkins, T.: Absence of red cell glutamic-pyruvatetransaminase: discovery of a 'silent' allele homozygote. Am. J. Hum.Genet. 32: 42-46, 1980.

[80321] 18651.O'Connell, P.; Nakamura, Y.; Lathrop, G. M.; Lepert, M.; Cartwright,P.; Lalouel, J.-M.; White, R.: Three genetic linkage groups on chromosome8. (Abstract) Cyto-genet. Cell Genet. 46: 672 only, 1987.

[80322] 18652.Olaisen, B.: Genetics of the GPT system: family, mother-childand association studies. Clin. Genet. 7: 245-254, 1975.

[80323] 18653.Pelzer, C. F.; Norum, R. A.: Identification of human red cellglutamate-pyruvate transaminase (GPT) pheno-

types by isoelectric focusing. Am.J. Hum. Genet. 37: 147–152, 1985.

- [80324] 18654.Rocha, J.; Amorim, A.; Almeida, V. M.; Oliveira, J. P.; Leao,M.; Tavares, M. C.; Pereira, M. S.; Vidal–Pinheiro, L.: Gene dosage evidence for the regional assignment of GPT (glutamate–pyruvate transaminase;E.C. 2.6.1.2) locus to 8q24.2–8qter. Hum. Genet. 80: 299–300, 1988.
- [80325] 18655.Sanders, M. F.; King, M. C.; Lattanzio, D.; Crandall, J.; Leung,R.: Absence of linkage between HP and GPT. (Abstract) Cytogenet.Cell Genet. 37: 536–537, 1984.
- [80326] 18656.Santachiara Benerecetti, A. S.; Beretta, M.; Pampiglione, S.:Red cell glutamic–pyruvic transaminase polymorphism in a sample ofthe Italian population: a new variant allele: GPT(8). Hum. Hered. 25:276–278, 1975.
- [80327] 18657.Sohocki, M. M.; Sullivan, L. S.; Harrison, W. R.; Sodergren, E.J.; Elder, F. F. B.; Weinstock, G.; Tanase, S.; Daiger, S. P.: Humanglutamate pyruvate transaminase (GPT): localization to 8q24.3, cDNAand genomic sequences, and polymorphic sites. Genomics 40: 247–252,1997.
- [80328] 18658.Sparkes, M. C.; Crist, M.; Sparkes, R. S.: Glutamate pyruvatetransaminase null allele in seven new families. Hum. Genet. 65:147–148, 1983.

- [80329] 18659.Wijnen, L. M. M.; Meera Khan, P.: Assignment of GPT to human chromosome 16. (Abstract) Cytogenet. Cell Genet. 32: 327 only, 1982.
- [80330] 18660.Burant, C. F.; Takeda, J.; Brot-Laroche, E.; Bell, G. I.; Davidson, N. O.: Fructose transporter in human spermatozoa and small intestine is GLUT5. J. Biol. Chem. 267: 14523–14526, 1992.
- [80331] 18661.Davidson, N. O.; Hausman, A. M. L.; Ifkovits, C. A.; Buse, J. B.; Gould, G. W.; Burant, C. F.; Bell, G. I.: Human intestinal glucose transporter expression and localization of GLUT5. Am. J. Physiol. 262:C795–C800, 1992.
- [80332] 18662.Cremonini, N.; Graziano, E.; Chiarini, V.; Sforza, A.; Zampa, G. A.: Atypical McCune–Albright syndrome associated with growth hormone–prolactin pituitary adenoma: natural history, long-term follow-up, and SMS 201–995–bromocriptine combined treatment results. J. Clin. Endocr. Metab. 75: 1166–1169, 1992.
- [80333] 18663.DeChiara, T.; Robertson, E. J.; Efstratiadis, A.: Parental imprinting of the mouse insulin-like growth factor II gene. Cell 64: 849–859, 1991.
- [80334] 18664.Eddy, M. C.; Jan de Beur, S. M.; Yandow, S. M.; McAlister, W. H.; Shore, E. M.; Kaplan, F. S.; Whyte, M. P.; Levine, M. A.: Deficiency of the alpha-subunit of the stim-

ulatory G protein and severe extraskeletal ossification. *J. Bone Miner. Res.* 15: 2074–2083, 2000.

[80335] 18665. Falconer, M. A.; Cope, C. L.; Robb-Smith, A. H. T.: Fibrous dysplasia of bone with endocrine disorders and cutaneous pigmentation (Albright's disease). *Quart. J. Med.* 11: 121–154, 1942.

[80336] 18666. Albertsen, H. M.; Smith, S. A.; Mazoyer, S.; Fujimoto, E.; Stevens, J.; Williams, B.; Rodriguez, P.; Cropp, C. S.; Slijepcevic, P.; Carlson, M.; Robertson, M.; Bradley, P.; Lawrence, E.; Harrington, T.; Mei Sheng, Z.; Hoopes, R.; Sternberg, N.; Brothman, A.; Callahan, R.; Ponder, B. A. J.; White, R.: A physical map and candidate genes in the BRCA1 region on chromosome 17q12–21. *Nature Genet.* 7: 472–479, 1994.

[80337] 18667. Kato, M.; Araki, S.: Paroxysmal kinesigenic choreoathetosis. *Arch. Neurol.* 20: 508–513, 1969.

[80338] 18668. Pilla, M.; Perachon, S.; Sautel, F.; Garrido, F.; Mann, A.; Wermuth, C. G.; Schwartz, J.-C.; Everitt, B. J.; Sokoloff, P.: Selective inhibition of cocaine-seeking behaviour by a partial dopamine D3 receptor agonist. *Nature* 400: 371–375, 1999.

[80339] 18669. Demange, P.; Voges, D.; Benz, J.; Liemann, S.; Gotting, P.; Berendes, R.; Burger, A.; Huber, R.: Annexin V: the

key to understanding ionselectivity and voltage regulation? Trends Biochem. Sci. 19: 272–276,1994.

- [80340] 18670.Funakoshi, T.; Heimark, R. L.; Hendrickson, L. E.; McMullen, B.A.; Fujikawa, K.: Human placental anticoagulant protein: isolationand characterization. Biochemistry 26: 5572–5578, 1987.
- [80341] 18671.Grundmann, U.; Abel, K.–J.; Bohn, H.; Lobermann, H.; Lottspeich,F.; Kupper, H.: Characterization of cDNA encoding human placentalanticoagulant protein (PP4): homology with the lipocortin family. Proc.Nat. Acad. Sci. 85: 3708–3712, 1988.
- [80342] 18672.Kaplan, R.; Jaye, M.; Burgess, W. H.; Schlaepfer, D. D.; Haigler,H. T.: Cloning and expression of cDNA for human endonexin II, a  $\text{Ca}(2+)$ and phospholipid binding protein. J. Biol. Chem. 263: 8037–8043,1988.
- [80343] 18673.Modi, W. S.; Seuanez, H.; Jaye, M.; Kaplan, R.; Haigler, H.; O'Brien,S. J.: Chromosomal mapping of the endonexin II gene. (Abstract) Cytogenet.Cell Genet. 51: 1046, 1989.
- [80344] 18674.Modi, W. S.; Seuanez, H. N.; Jaye, M.; Haigler, H. J.; Kaplan,R.; O'Brien, S. J.: The human endonexin II (ENX2) gene is locatedat 4q28–q32. Cytogenet. Cell Genet. 52: 167–169, 1989.



- [80345] 18675.Rodriguez-Garcia, M. I.; Kozak, C. A.; Morgan, R. O.; Fernandez,M. P.: Mouse annexin V chromosomal localization, cDNA sequence conservation,and molecular evolution. *Genomics* 31: 151-157, 1996.
- [80346] 18676.Tait, J. F.; Frankenberry, D. A.; Shiang, R.; Murray, J. C.; Adler,D. A.; Disteché, C. M.: Chromosomal localization of the human gene for annexin V (placental anticoagulant protein I) to 4q26-q28. *Cytogenet.Cell Genet.* 57: 187-192, 1991.
- [80347] 18677.Arinami, T.; Ishikawa, M.; Inoue, A.; Yanagisawa, M.; Masaki, T.;Yoshida, M. C.; Hamaguchi, H.: Chromosomal assignments of the human endothelin family genes: the endothelin-1 gene (EDN1) to 6p23-p24,the endothelin-2 gene (EDN2) to 1p34, and the endothelin-3 gene (EDN3)to 20q13.2-q13.3. *Am. J. Hum. Genet.* 48: 990-996, 1991.
- [80348] 18678.Benatti, L.; Bonecchi, L.; Cozzi, L.; Sarmientos, P.: Two preproendothelin1 mRNAs transcribed by alternative promoters. *J. Clin. Invest.* 91:1149-1156, 1993.
- [80349] 18679.Berge, K. E.; Berg, K.: No effect of a TaqI polymorphism in DNA at the endothelin I (EDN1) locus on normal blood pressure level or variability. *Clin. Genet.* 41: 90-95, 1992.

- [80350] 18680.Bloch, K. D.; Friedrich, S. P.; Lee, M.-E.; Eddy, R. L.; Shows,T. B.; Quertermous, T.: Structural organization and chromosomal assignmentof the gene encoding endothe-  
lin. J. Biol. Chem. 264: 10851–10857,1989.
- [80351] 18681.Bourgeois, C.; Robert, B.; Rebourcet, R.; Mondon, F.; Mignot, T.-M.;Duc-Goiran, P.; Ferre, F.: Endothelin–1 and ET(A) receptor expressionin vascular smooth muscle cells from human placenta: a new ET(A) receptormessen-  
ger ribonucleic acid is generated by alternative splicing ofexon 3. J. Clin. Endocr. Metab. 82: 3116–3123, 1997.
- [80352] 18682.Clouthier, D. E.; Hosoda, K.; Richardson, J. A.; Williams, S. C.;Yanagisawa, H.; Kuwaki, T.; Kumada, M.; Hammer, R. E.; Yanagisawa,M.: Cranial and cardiac neural crest defects in endothelin–A receptor–deficientmice. De-  
velopment 125: 813–824, 1998.
- [80353] 18683.Giaid, A.; Gibson, S. J.; Ibrahim, N. B. N.; Legon, S.; Bloom,S. R.; Yanagisawa, M.; Masaki, T.; Varndell, I. M.; Polak, J. M.:Endothelin 1, an endothelium–derived peptide, is expressed in neuronsof the human spinal cord and dor-  
sal root ganglia. Proc. Nat. Acad.Sci. 86: 7634–7638, 1989.
- [80354] 18684.Inoue, A.; Yanagisawa, M.; Kimura, S.; Kasuya, Y.; Miyauchi, T.;Goto, K.; Masaki, T.: The human endothelin

family: three structurally and pharmacologically distinct isopeptides predicted by three separate genes. *Proc. Nat. Acad. Sci.* 86: 2863–2867, 1989.

- [80355] 18685. Inoue, A.; Yanagisawa, M.; Takuwa, Y.; Mitsui, Y.; Kobayashi, M.; Masaki, T.: The human preproendothelin-1 gene: complete nucleotide sequence and regulation of expression. *J. Biol. Chem.* 264: 14954–14959, 1989.
- [80356] 18686. Shaikh, S.; Ball, D.; Craddock, N.; Castle, D.; Hunt, N.; Mant, R.; Owen, M.; Collier, D.; Gill, M.: The dopamine D3 receptor gene: no association with bipolar affective disorder. *J. Med. Genet.* 30: 308–309, 1993.
- [80357] 18687. Sokoloff, P.; Giros, B.; Martres, M.-P.; Bouthenet, M.-L.; Schwartz, J.-C.: Molecular cloning and characterization of a novel dopamine receptor (D-3) as a target for neuroleptics. *Nature* 347: 146–151, 1990.
- [80358] 18688. Spurlock, G.; Williams, J.; McGuffin, P.; Aschauer, H. N.; Lenzinger, E.; Fuchs, K.; Sieghart, W. C.; Meszaros, K.; Fathi, N.; Laurent, C.; Mallet, J.; Macciardi, F.; Pedrini, S.; Gill, M.; Hawi, Z.; Gibson, S.; Jazin, E. E.; Yang, H.-T.; Adolfsson, R.; Pato, C. N.; Dourado, A. M.; Owen, M. J.: European Multicentre Association Study of Schizophrenia: a study of the DRD2 ser311-to-cys and DRD3 ser9-to-gly polymorphisms. *Am. J. Med. Genet.* 81: 24–28, 1998.

- [80359] 18689.Asghari, V.; Sanyal, S.; Buchwaldt, S.; Paterson, A.; Jovanovic,V.; Van Tol, H. H.: Modulation of intracellular cyclic AMP levelsby different human dopamine D4 receptor variants. *J. Neurochem.* 65:1157–1165, 1995.
- [80360] 18690.Benjamin, J.; Li, L.; Patterson, C.; Greenberg, B. D.; Murphy,D. L.; Hamer, D. H.: Population and familial association between the D4 dopamine receptor gene and measures of novelty seeking. *NatureGenet.* 12: 81–84, 1996.
- [80361] 18691.Castellanos, F. X.; Lau, E.; Tayebi, N.; Lee, P.; Long, R. E.;Giedd, J. N.; Sharp, W.; Marsh, W. L.; Walter, J. M.; Hamburger, S.D.; Ginns, E. I.; Rapoport, J. L.; Sidransky, E.: Lack of an associationbetween a dopamine–4 receptor polymorphism and attention–deficit/hyperactivitydisorder: genetic and brain morphometric analyses. *Molec. Psychiat.* 3:431–434, 1998.
- [80362] 18692.Chang, F.–M.; Kidd, J. R.; Livak, K. J.; Pakstis, A. J.; Kidd,K. K.: The world–wide distribution of allele frequencies at the humandopamine D4 receptor locus. *Hum. Genet.* 98: 91–101, 1996.
- [80363] 18693.Cloninger, C. R.; Adolfsson, R.; Svrakic, N. M.: Mapping genesfor human personality. *Nature Genet.* 12: 3–4, 1996.
- [80364] 18694.Cloninger, C. R.; Svrakic, D. M.; Przybeck, T. R.: A

psychobiological model of temperament and character.

Arch. Gen. Psychiat. 50: 975–990, 1993.

[80365] 18695. Cook, E. H.; Stein, M. A.; Krasowski, M. D.; Cox, N. J.; Olkon, D. M.; Kieffer, J. E.; Leventhal, B. L.: Association of attention-deficit disorder and the dopamine transporter gene. Am. J. Hum. Genet. 56:993–998, 1995.

[80366] 18696. De Luca, A.; Rizzardi, M.; Torrente, I.; Alessandroni, R.; Salvioli, G. P.; Filograsso, N.; Dallapiccola, B.; Novelli, G.: Dopamine D4 receptor (DRD4) polymorphism and adaptability trait during infancy: a longitudinal study in 1- to 5-month-old neonates. Neurogenetics 3:79–82, 2001.

[80367] 18697. Ding, Y.-C.; Chi, H.-C.; Grady, D. L.; Morishima, A.; Kidd, J. R.; Kidd, K. K.; Flodman, P.; Spence, M. A.; Schuck, S.; Swanson, J. M.; Zhang, Y.-P.; Moyzis, R. K.: Evidence of positive selection acting at the human dopamine receptor D4 gene locus. Proc. Nat. Acad. Sci. 99: 309–314, 2002.

[80368] 18698. Ebstein, R. P.; Novick, O.; Umansky, R.; Pirelli, B.; Osher, Y.; Blaine, D.; Bennett, E. R.; Nemanov, L.; Katz, M.; Belmaker, R. H.: Dopamine D4 receptor exon III polymorphism associated with the human personality trait of novelty seeking. Nature Genet. 12: 78–80, 1996.

[80369] 18699. Eisenberg, J.; Zohar, A.; Mei-Tal, G.; Steinberg, A.;

Tartakovsky, E.; Gritsenko, I.; Nemanov, L.; Ebstein, R. P.: A haplotype relative risk study of the dopamine D4 receptor (DRD4) exon III repeat polymorphism and attention deficit hyperactivity disorder (ADHD). *Am. J. Med. Genet. (Neuropsychiatr. Genet.)* 96: 258–261, 2000.

[80370] 18700. Gelernter, J.; Kennedy, J. L.; Van Tol, H. H. M.; Civelli, O.; Kidd, K. K.: The D4 dopamine receptor (DRD4) maps to distal 11p close to HRAS. *Genomics* 13: 208–210, 1992.

[80371] 18701. Gelernter, J.; Kennedy, J. L.; Van Tol, H. H. M.; Niznik, H. B.; Civelli, O.; Kidd, K. K.: The D4 dopamine receptor (DRD4) maps to distal 11p close to HRAS. (Abstract) *Cytogenet. Cell Genet.* 58:1960, 1991.

[80372] 18702. Gelernter, J.; Kranzler, H.; Coccaro, E.; Siever, L.; New, A.; Mulgrew, C. L.: D4 dopamine-receptor (DRD4) alleles and novelty seeking in substance-dependent, personality-disorder, and control subjects. *Am. J. Hum. Genet.* 61: 1144–1152, 1997.

[80373] 18703. Harpending, H.; Cochran, G.: In our genes. *Proc. Nat. Acad. Sci.* 99: 10–12, 2002.

[80374] 18704. Jovanovic, V.; Guan, H.-C.; Van Tol, H. H. M.: Comparative pharmacological and functional analysis of the human dopamine D4.2 and D4.10 receptor variants. *Phar-*

macogenetics 9: 561–568, 1999.

- [80375] 18705.LaHoste, G. J.; Swanson, J. M.; Wigal, S. B.; Glabe, C.; Wigal,T.; King, N.; Kennedy, J. L.: Dopamine D4 receptor gene polymorphismis associated with attention deficit hyperactivity disorder. *Molec.Psychiat.* 1: 121–124, 1996.
- [80376] 18706.Liu, I. S. C.; Seeman, P.; Sanyal, S.; Ulpian, C.; Rodgers–Johnson,P. E. B.; Serjeant, G. R.; Van Tol, H. H. M.: Dopamine D4 receptorvariant in Africans, D4(valine194glycine), is insensitive to dopamineand clozapine: report of a homozygous individual. *Am. J. Med. Genet.* 61:277–282, 1996.
- [80377] 18707.Malhotra, A. K.; Virkkunen, M.; Rooney, W.; Eggert, M.; Linnoila,M.; Goldman, D.: The association between the dopamine D(4) receptor(D4DR) 16 amino acid repeat polymorphism and novelty seeking. *Molec.Psychiat.* 1: 388–391, 1996.
- [80378] 18708.Muramatsu, T.; Higuchi, S.; Murayama, M.; Matsushita, S.; Hayashida,M.: Association between alcoholism and the dopamine D4 receptor gene. *J.Med. Genet.* 33: 113–115, 1996.
- [80379] 18709.Nothen, M. M.; Cichon, S.; Hemmer, S.; Hebebrand, J.; Remschmidt,H.; Lehmkuhl, G.; Poustka, F.; Schmidt, M.; Catalano, M.; Fimmers,R.; Korner, J.; Rietschel, M.; Prop–

ping, P.: Human dopamine D4 receptorgene: frequent occurrence of a null allele and observation of homozygosity. Hum.Molec. Genet. 3: 2207–2212, 1994.

[80380] 18710.Petronis, A.; Van Tol, H. H. M.; Lichter, J. B.; Livak, K. J.;Kennedy, J. L.: The D4 dopamine receptor gene maps on 11p proximalto HRAS. Genomics 18: 161–163, 1993.

[80381] 18711.Rowe, D. C.; Stever, C.; Giedinghagen, L. N.; Gard, J. M. C.;Cleveland, H. H.; Terris, S. T.; Mohr, J. H.; Sherman, S.; Abramowitz,A.; Waldman, I. D.: Dopamine DRD4 receptor polymorphism and attentiondeficit hyperactivity disorder. Molec. Psychiat. 3: 419–426, 1998.

[80382] 18712.Rubinstein, M.; Phillips, T. J.; Bunzow, J. R.; Falzone, T. L.;Dziewczapolski, G.; Zhang, G.; Fang, Y.; Larson, J. L.; McDougall,J. A.; Chester, J. A.; Saez, C.; Pugsley, T. A.; Gershanik, O.; Low,M. J.; Grandy, D. K.: Mice lacking dopamine D4 receptors are supersensitiveto ethanol, cocaine, and methamphetamine. Cell 90: 991–1001, 1997.

[80383] 18713.Raskind, W. H.; Bolin, T.; Wolff, J.; Fink, J.; Matsushita, M.;Litt, M.; Lipe, H.; Bird, T. D.: Further localization of a gene forparoxysmal dystonic choreoathetosis to a 5–cM region on chromosome2q34. Hum. Genet. 102: 93–97, 1998.

[80384] 18714.Richards, R. N.; Barnett, H. J.: Paroxysmal dystonic



choreoathetosis:a family study and review of the literature. *Neurology* 18: 461–469,1968.

- [80385] 18715.Stevens, H. F.: Paroxysmal choreo–athetosis: a form of reflexepilepsy. *Arch. Neurol.* 14: 415–420, 1966.
- [80386] 18716.Wagner, G. S.; McLees, B. D.; Hatcher, M. A., Jr.: Familial paroxysmalchoreo–athetosis. (Abstract) *Neurology* 16: 307, 1966.
- [80387] 18717.Williams, J.; Stevens, H.: Familial paroxysmal choreo–athetosis. *Pediatrics* 31:656–659, 1963.
- [80388] 18718.Barsh, G. S.; Seeburg, P. H.; Gelinas, R. E.: The human growthhormone gene family:structure and evolution of the chromosomal locus. *NucleicAcids Res.* 11: 3939–3958, 1983.
- [80389] 18719.Cremers, F. P. M.; Molloy, C. M.; van de Pol, D. J. R.; van denHurk, J. A. J. M.; Bach, I.; Geurts van Kessel, A. H. M.; Ropers,H.–H.: An autosomal homologue of the choroideremia gene colocalizeswith the Usher syndrome type II locus on the distal part of chromosome1q. *Hum. Molec. Genet.* 1: 71–75, 1992.
- [80390] 18720.Halford, S.; Freedman, M. S.; Bellingham, J.; Inglis, S. L.; Poopalasundaram,S.; Soni, B. G.; Foster, R. G.; Hunt, D. M.: Characterization ofa novel human opsin gene with wide tissue expression and identificationof embedded and

flanking genes on chromosome 1q43. *Genomics* 72:203–208, 2001.

- [80391] 18721.van Bokhoven, H.; van Genderen, C.; Molloy, C. M.; van de Pol, D. J. R.; Cremers, C. W. R. J.; van Aarem, A.; Schwartz, M.; Rosenberg, T.; Geurts van Kessel, A. H. M.; Ropers, H.-H.; Cremers, F. P. M.: Mapping of the chorioideremia-like (CHML) gene at 1q42-qter and mutation-analysis in patients with Usher syndrome type II. *Genomics* 19: 385–387, 1994.
- [80392] 18722.Amato, F.; Warnes, G. M.; Kirby, C. A.; Norman, R. J.: Infertility caused by hCG autoantibody. *J. Clin. Endocr. Metab.* 87: 993–997, 2002.
- [80393] 18723.Naylor, S. L.; Chin, W. W.; Goodman, H. M.; Lalley, P. A.; Grzeschik, K.-H.; Sakaguchi, A. Y.: Chromosome assignment of the genes encoding the alpha and beta subunits of the glycoprotein hormones in man and mouse. *Somat. Cell Genet.* 9: 757–770, 1983.
- [80394] 18724.Hart, A. W.; Baeza, N.; Apelqvist, A.; Edlund, H.: Attenuation of FGF signalling in mouse beta-cells leads to diabetes. *Nature* 408:864–868, 2000.
- [80395] 18725.Mammarella, S.; Romano, F.; Di Valerio, A.; Creati, B.; Esposito, D. L.; Palmirotta, R.; Capani, F.; Vitullo, P.; Volpe, G.; Battista, P.; Della Loggia, F.; Mariani-Costantini,

R.; Cama, A.: Interaction between the G1057D variant of IRS-2 and overweight in the pathogenesis of type 2 diabetes. *Hum. Molec. Genet.* 9: 2517–2521, 2000.

[80396] 18726. Triggs-Raine, B. L.; Kirkpatrick, R. D.; Kelly, S. L.; Norquay, L. D.; Cattini, P. A.; Yamagata, K.; Hanley, A. J. G.; Zinman, B.; Harris, S. B.; Barrett, P. H.; Hegele, R. A.: HNF1- $\alpha$  G319S, a transactivation-deficient mutant, is associated with altered dynamics of diabetes onset in an Oji-Cree community. *Proc. Nat. Acad. Sci.* 99:4614–4619, 2002.

[80397] 18727. Tuomi, T.; Carlsson, A.; Li, H.; Isomaa, B.; Miettinen, A.; Nilsson, A.; Nissen, M.; Ehrnstrom, B.-O.; Forsen, B.; Snickars, B.; Lahti, K.; Forsblom, C.; Saloranta, C.; Taskinen, M.-R.; Groop, L. C.: Clinical and genetic characteristics of type 2 diabetes with and without GAD antibodies. *Diabetes* 48: 150–157, 1999.

[80398] 18728. Yuan, M.; Konstantopoulos, N.; Lee, J.; Hansen, L.; Li, Z.-W.; Karin, M.; Shoelson, S. E.: Reversal of obesity- and diet-induced insulin resistance with salicylates or targeted disruption of I $\kappa$ B- $\beta$ . *Science* 293:1673–1677, 2001.

[80399] 18729. Hart, T. C.; Champagne, C.; Zhou, J.; Van Dyke, T. E.: Assignment of the gene for diacylglycerol kinase

(DAGK) to human chromosome 12. *MammalianGenome* 5: 123–124, 1994.

- [80400] 18730.Hart, T. C.; Zhou, J.; Champagne, C.; Van Dyke, T. E.; Rao, P.N.; Pettenati, M. J.: Assignment of the human diacylglycerol kinase gene (DAGK) to 12q13.3 using fluorescence in situ hybridization analysis. *Genomics* 22:246–247, 1994.
- [80401] 18731.Pilz, A.; Schaap, D.; Hunt, D.; Fitzgibbon, J.: Chromosomal localization of three mouse diacylglycerol kinase (DAGK) genes: genes sharing sequence homology to the *Drosophila* retinal degeneration A (rdgA) gene. *Genomics* 26:599–601, 1995.
- [80402] 18732.Schaap, D.; de Widt, J.; van der Wal, J.; Vandekerckhove, J.; van Damme, J.; Gussow, D.; Ploegh, H. L.; van Blitterswijk, W. J.; vander Bend, R. L.: Purification, cDNA-cloning and expression of human diacylglycerol kinase. *FEBS Lett.* 275: 151–158, 1990.
- [80403] 18733.Barker, P. E.; Shipp, M. A.; D'Adamio, L.; Masteller, E. L.; Reinherz, E. L.: The common acute lymphoblastic leukemia antigen gene maps to chromosomal region 3(q21–q27). *J. Immun.* 142: 283–287, 1989.
- [80404] 18734.D'Adamio, L.; Shipp, M. A.; Masteller, E. L.; Reinherz, E. L.: Organization of the gene encoding common

acute lymphoblastic leukemia antigen (neutral endopeptidase 24.11): multiple miniexons and separate 5'-prime untranslated regions. Proc. Nat. Acad. Sci. 86: 7103-7107, 1989.

[80405] 18735. Debiec, H.; Guignon, V.; Mougnot, B.; Decobert, F.; Haymann, J.-P.; Bensman, A.; Deschenes, G.; Ronco, P. M.: Antenatal membranous glomerulonephritis due to anti-neutral endopeptidase antibodies. New Eng. J. Med. 346: 2053-2060, 2002.

[80406] 18736. Letarte, M.; Vera, S.; Tran, R.; Addis, J. B. L.; Onizuka, R. J.; Quackenbush, E. J.; Jongeneel, C. V.; McInnes, R. R.: Common acute lymphocytic leukemia antigen is identical to neutral endopeptidase. J. Exp. Med. 168: 1247-1253, 1988.

[80407] 18737. Shipp, M. A.; Vijayaraghavan, J.; Schmidt, E. V.; Masteller, E. L.; D'Adamio, L.; Hersch, L. B.; Reinherz, E. L.: Common acute lymphoblastic leukemia antigen (CALLA) is active neutral endopeptidase 24.11 ('enkephalinase'): direct evidence by cDNA transfection analysis. Proc. Nat. Acad. Sci. 86: 297-301, 1989.

[80408] 18738. Tran-Paterson, R.; Willard, H. F.; Letarte, M.: The common acute lymphoblastic leukemia antigen (neutral endopeptidase--3.4.24.11) gene is located on human

chromosome 3. *Cancer Genet. Cytogenet.* 42:129–134, 1989.

[80409] 18739.Lahn, B. T.; Page, D. C.: Four evolutionary strata on the humanX chromosome. *Science* 286: 964–967, 1999.

[80410] 18740.Lee, F. A.; Issacs, H.; Strauss, J.: The 'campomelic' syndrome.Short life–span dwarfism with respiratory distress, hypotonia, peculiarfacies, and multiple skeletal and cartilaginous deformities. *Am.J. Dis. Child.* 124: 485–496, 1972.

[80411] 18741.Lynch, S. A.; Gaunt, M. L.; Minford, A. M. B.: Campomelic dysplasia:evidence of autosomal dominant inheritance. *J. Med. Genet.* 30: 683–686,1993.

[80412] 18742.Macpherson, R. I.; Skinner, S. A.; Donnenfeld, A. E.: Acampomeliccampomelic dysplasia. *Pediat. Radiol.* 20: 90–93, 1989.

[80413] 18743.Mansour, S.; Hall, C. M.; Pembrey, M. E.; Young, I. D.: A clinicaland genetic study of campomelic dysplasia. *J. Med. Genet.* 32: 415–420,1995.

[80414] 18744.Maraia, R.; Saal, H. M.; Wangsa, D.: A chromosome 17q de novoparacentric inversion in a patient with campomelic dysplasia; casereport and etiologic hypothesis. *Clin. Genet.* 39: 401–408, 1991.

[80415] 18745.Maroteaux, P.; Spranger, J. W.; Opitz, J. M.; Kucera,

J.; Lowry, R. B.; Schimke, R. N.; Kagan, S. M.: Le syndrome campomelique. *PresseMed.* 22: 1157–1162, 1971.

[80416] 18746. Meyer, J.; Sudbeck, P.; Held, M.; Wagner, T.; Schmitz, M. L.; Bricarelli, F. D.; Eggermont, E.; Friedrich, U.; Haas, O. A.; Kobelt, A.; Leroy, J. G.; van Maldergem, L.; Michel, E.; Mitulla, B.; Pfeiffer, R. A.; Schinzel, A.; Schmidt, H.; Scherer, G.: Mutational analysis of the SOX9 gene in campomelic dysplasia and autosomal sex reversal: lack of genotype/phenotype correlations. *Hum. Molec. Genet.* 6: 91–98, 1997.

[80417] 18747. Moedjono, S. J.; Crandall, B. F.; Sparkes, R. S.; Feldman, G. M.; Austin, G. E.; Perry, S.: The campomelic syndrome in a singleton and monozygotic twins. *Clin. Genet.* 18: 397–401, 1980.

[80418] 18748. Moog, U.; Jansen, N. J. G.; Scherer, G.; Schrandt-Stumpel, C. T. R. M.: A campomelic campomelic syndrome. *Am. J. Med. Genet.* 104: 239–245, 2001.

[80419] 18749. Morais da Silva, S.; Hacker, A.; Harley, V.; Goodfellow, P.; Swain, A.; Lovell-Badge, R.: Sox9 expression during gonadal development implies a conserved role for the gene in testis differentiation in mammals and birds. *Nature Genet.* 14: 62–68, 1996.

[80420] 18750. Murakami, S.; Kan, M.; McKeenan, W. L.; de Crom-

brugghe, B.: Up-regulation of the chondrogenic Sox9 gene by fibroblast growth factors is mediated by the mitogen-activated protein kinase pathway. *Proc. Nat. Acad. Sci.* 97: 1113–1118, 2000.

- [80421] 18751. Ninomiya, S.; Isomura, M.; Narahara, K.; Seino, Y.; Nakamura, Y.: Isolation of a testis-specific cDNA on chromosome 17q from a region adjacent to the breakpoint of t(12;17) observed in a patient with acampomelic campomelic dysplasia and sex reversal. *Hum. Molec. Genet.* 5: 69–72, 1996.
- [80422] 18752. Ninomiya, S.; Yokoyama, Y.; Teraoka, M.; Mori, R.; Inoue, C.; Yamashita, S.; Tamai, H.; Funato, M.; Seino, Y.: A novel mutation (296 del G) of the SOX9 gene in a patient with campomelic syndrome and sex reversal. *Clin. Genet.* 58: 224–227, 2000.
- [80423] 18753. Olney, P. N.; Kean, L. S.; Graham, D.; Elsas, L. J.; May, K. M.: Campomelic syndrome and deletion of SOX9. *Am. J. Med. Genet.* 84:20–24, 1999.
- [80424] 18754. Ozkilic, A.; Seven, M.; Yuksel, A.: A case of acampomelic campomelic dysplasia. *Genet. Counsel.* 13: 23–28, 2002.
- [80425] 18755. Patel, M.; Dorman, K. S.; Zhang, Y.-H.; Huang, B.-L.; Arnold, A. P.; Sinsheimer, J. S.; Vilain, E.; McCabe, E.



R. B.: Primate DAX1, SRY, and SOX9: evolutionary stratification of sex-determination pathway. *Am. J. Hum. Genet.* 68: 275–280, 2001.

[80426] 18756. Pfeifer, D.; Kist, R.; Dewar, K.; Devon, K.; Lander, E. S.; Birren, B.; Korniszewski, L.; Back, E.; Scherer, G.: Campomelic dysplasia translocation breakpoints are scattered over 1 Mb proximal to SOX9: evidence for an extended control region. *Am. J. Hum. Genet.* 65: 111–124, 1999.

[80427] 18757. Puck, S. M.; Haseltine, F. P.; Francke, U.: Absence of H-Y antigen in an XY female with campomelic dysplasia. *Hum. Genet.* 57: 23–27, 1981.

[80428] 18758. Rimoin, D. L.: Personal Communication. Torrance, Calif. 8/12/1976.

[80429] 18759. Rodriguez, J. I.: Vascular anomalies in campomelic syndrome. *Am. J. Med. Genet.* 46: 185–192, 1993.

[80430] 18760. Savarirayan, R.; Bankier, A.: A campomelic campomelic dysplasia with de novo 5q;17q reciprocal translocation and severe phenotype. *J. Med. Genet.* 35: 597–599, 1998.

[80431] 18761. Shimke, R. N.: XY sex-reversed campomelia – possibly an X-linked disorder? (Letter) *Clin. Genet.* 16: 62–63, 1979.

[80432] 18762. Shafai, T.; Schwartz, L.: Campomelic syndrome in

siblings. J.Pediat. 89: 512–513, 1976.

[80433] 18763.Spranger, J.: Advances in bone dysplasias.

(Abstract) SixthInt. Cong. Hum. Genet. Jerusalem , 1981.

[80434] 18764.Stuve, A.; Wiedemann, H.–R.: Congenital bowing of the long bones in two sisters. (Letter) Lancet I: 495, 1971.

[80435] 18765.Sudbeck, P.; Schmitz, M. L.; Baeuerle, P. A.; Scherer, G.: Sexreversal by loss of the C-terminal transactivation domain of humanSOX9. Nature Genet. 13: 230–232, 1996.

[80436] 18766.Thong, M.–K.; Scherer, G.; Kozlowski, K.; Haan, E.; Morris, L.: Acampomelic campomelic dysplasia with SOX9 mutation. Am. J. Med.Genet. 93: 421–425, 2000.

[80437] 18767.Thurmon, T. F.; De Fraites, E. B.; Anderson, E. E.: Familialcampomelic dwarfism. J. Pediat. 83: 841–843, 1973.

[80438] 18768.Vidal, V. P. I.; Chaboissier, M.–C.; de Rooij, D. G.; Schedl,A.: Sox9 induces testis development in XX transgenic mice. NatureGenet. 28: 216–217, 2001.

[80439] 18769.Wagner, T.; Wirth, J.; Meyer, J.; Zabel, B.; Held, M.; Zimmer,J.; Pasantes, J.; Dagna Bricarelli, F.; Keutel, J.; Huser, E.; Wolf,U.; Tommerup, N.; Schempp, W.; Scherer, G.: Autosomal sex reversal and campomelic dysplasia are caused by mutations in and around theSRY-related gene

SOX9. Cell 79: 1111–1120, 1994.

[80440] 18770.Weller, S. D. V.: Hypophosphatasia with congenital dimples. Proc.Roy. Soc. Med. 52: 637, 1959.

[80441] 18771.Wirth, J.; Wagner, T.; Meyer, J.; Pfeiffer, R. A.; Tietze, H.–U.;Schempp, W.; Scherer, G.: Translocation breakpoints in three patientswith campomelic dysplasia and autosomal sex reversal map more than130 kb from SOX9. Hum. Genet. 97: 186–193, 1996.

[80442] 18772.Wright, E.; Hargrave, M. R.; Christiansen, J.; Cooper, L.; Kun,J.; Evans, T.; Gangadharan, U.; Greenfield, A.; Koopman, P.: TheSry–related gene Sox9 is expressed during chondrogenesis in mouseembryos. Nature Genet. 9: 15–20, 1995.

[80443] 18773.Albritton, L. M.; Bowcock, A. M.; Eddy, R. L.; Morton, C. C.; Tseng,L.; Farrer, L. A.; Cavalli–Sforza, L. L.; Shows, T. B.; Cunningham,J. M.: The human cationic amino acid transporter (ATRC1): physicaland genetic mapping to 13q12–q14. Genomics 12: 430–434, 1992.

[80444] 18774.Bowcock, A. M.; Gerken, S. C.; Barnes, R. I.; Shiang, R.; Jabs,E. W.; Warren, A. C.; Antonarakis, S.; Retief, A. E.; Vergnaud, G.;Leppert, M.; Lalouel, J.–M.; White, R. L.; Cavalli–Sforza, L. L.:The CEPH consortium linkage map of human chromosome 13. Genomics 16:486–496, 1993.

- [80445] 18775.Kim, J. W.; Closs, E. I.; Albritton, L. M.; Cunningham, J. M.:Transport of cationic amino acids by the mouse ecotropic retrovirusreceptor. *Nature* 352: 725–728, 1991.
- [80446] 18776.Kozak, C. A.; Albritton, L. M.; Cunningham, J. M.: Genetic mapping of a cloned sequence responsible for susceptibility to ecotropic murine leukemia viruses. *J. Virol.* 64: 3119–3121, 1990.
- [80447] 18777.Oie, H. K.; Gazdar, A. F.; Lalley, P. A.; Russell, E. K.; Minna, J. D.; DeLarco, J.; Todaro, G. J.; Francke, U.: Mouse chromosome 5 codes for ecotropic murine leukaemia virus cell–surface receptor. *Nature* 274:60–62, 1978.
- [80448] 18778.Ruddle, N. H.; Conta, B. S.; Leinwand, L.; Kozak, C.; Ruddle, F.; Besmer, P.; Baltimore, D.: Assignment of the receptor for ecotropic murine leukemia virus to mouse chromosome 5. *J. Exp. Med.* 148: 451–465, 1978.
- [80449] 18779.Boldog, F.; Erlandsson, R.; Klein, G.; Sumegi, J.: Long–range restriction enzyme maps of DNF15S2, D3S2 and c–raf1 loci on the short arm of human chromosome 3. *Cancer Genet. Cytogenet.* 42: 295–306, 1989.
- [80450] 18780.Cook, R. M.; Burke, B. J.; Buchhagen, D. L.; Minna, J. D.; Miller, Y. E.: Human aminoacylase–1: cloning, sequence, and expression analysis of a chromosome 3p21 gene inactivated in small cell lung cancer. *J. Biol. Chem.*

268: 17010–17017, 1993.

- [80451] 18781. Gemmill, R. M.; Varella-Garcia, M.; Smith, D. I.; Erickson, P.; Golembieski, W.; Miller, Y.; Coyle-Morris, J.; Tommerup, N.; Drabkin, H. A.: A 2.5 Mb physical map within 3p21.1 spans the breakpoint associated with Greig cephalopolysyndactyly syndrome. *Genomics* 11: 93–102, 1991.
- [80452] 18782. Ginzinger, D. G.; Shridhar, V.; Baldini, A.; Taggart, R. T.; Miller, O. J.; Smith, D. I.: The human loci DNF15S2 and D3S94 have a high degree of sequence similarity to acyl-peptide hydrolase and are located at 3p21.3. *Am. J. Hum. Genet.* 50: 826–833, 1992.
- [80453] 18783. Jones, W. M.; Scaloni, A.; Bossa, F.; Popowicz, A. M.; Schneewind, O.; Manning, J. M.: Genetic relationship between acylpeptide hydrolase and acylase, two hydrolytic enzymes with similar binding but different catalytic specificities. *Proc. Nat. Acad. Sci.* 88: 2194–2198, 1991.
- [80454] 18784. Miller, Y. E.; Drabkin, H.; Jones, C.; Fisher, J. H.: Aminoacylase-1: cDNA isolation, regional assignment to chromosome 3p21.1 and identification of a cross-hybridizing sequence on chromosome 18. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A28 only, 1989.
- [80455] 18785. Miller, Y. E.; Drabkin, H.; Jones, C.; Fisher, J. H.:

Human aminoacylase-1:cloning, regional assignment to distal chromosome 3p21.1, and identification of a cross-hybridizing sequence on chromosome 18. *Genomics* 8: 149-154,1990.

[80456] 18786.Miller, Y. E.; Minna, J. D.; Gazdar, A. F.: Lack of expression of aminoacylase-1 in small cell lung cancer: evidence for inactivation of genes encoded by chromosome 3p. *J. Clin. Invest.* 83: 2120-2124,1989.

[80457] 18787.Nadeau, J. H.: A chromosomal segment conserved since divergence of lineages leading to man and mouse: the gene order of aminoacylase-1, transferrin, and beta-galactosidase on mouse chromosome 9. *Genet.Res.* 48: 175-178, 1986.

[80458] 18788.Naylor, S. L.; Elliott, R. W.; Brown, J. A.; Shows, T. B.: Mapping of aminoacylase-1 and beta-galactosidase-A to homologous regions of human chromosome 3 and mouse chromosome 9 suggests location of additional genes. *Am. J. Hum. Genet.* 34: 235-244, 1982.

[80459] 18789.Naylor, S. L.; Shows, T. B.; Klebe, R. J.: Bioautographic visualization of aminoacylase-1: assignment of the structural gene ACY-1 to chromosome 3 in man. *Somat. Cell Genet.* 5: 11-21, 1979.

[80460] 18790.Voss, R.; Lerer, I.; Povey, S.; Solomon, E.; Bobrow,

M.: Confirmation and further regional assignment of aminoacylase 1 (ACY-1) on human chromosome 3 using a simplified detection method. *Ann. Hum. Genet.* 44:1-10, 1980.

[80461] 18791. Cook, P. W.; Piepkorn, M.; Clegg, C. H.; Plowman, G. D.; DeMay, J. M.; Brown, J. R.; Pittelkow, M. R.: Transgenic expression of the human amphiregulin gene induces a psoriasis-like phenotype. *J. Clin. Invest.* 100: 2286-2294, 1997.

[80462] 18792. Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly, J.; Bowen-Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of the amphiregulin and the platelet-growth factor receptor alpha genes to the proximal long arm of chromosome 4. (Abstract) *Cytogenet. Cell Genet.* 51:990 only, 1989.

[80463] 18793. Kimura, H.; Fischer, W. H.; Schubert, D.: Structure, expression and function of a schwannoma-derived growth factor. *Nature* 348:257-260, 1990.

[80464] 18794. Gillessen-Kaesbach, G.; Demuth, S.; Thiele, H.; Theile, U.; Lich, C.; Horsthemke, B.: A previously unrecognised phenotype characterised by obesity, muscular hypotonia, and ability to speak in patients with Angelman syndrome caused by an imprinting defect. *Europ. J.*

Hum.Genet. 7: 638–644, 1999.

- [80465] 18795.Cockayne, D. A.; Muchamuel, T.; Grimaldi, J. C.; Muller–Steffner,H.; Randall, T. D.; Lund, F. E.; Murray, R.; Schuber, F.; Howard,M. C.: Mice deficient for the ecto–nicotinamide adenine dinucleotideglycohydrolase CD38 exhibit altered humoral immune responses. Blood 92:1324–1333, 1998.
- [80466] 18796.Ferrero, E.; Malavasi, F.: Human CD38, a leukocyte receptor andectoenzyme, is a member of a novel eukary–otic gene family of nicotinamideadenine dinucleotide +/–converting enzymes: extensive structure homology–with the genes for murine bone marrow stromal cell anti–gen 1 and aplysianADP–ribosyl cyclase. J. Immun. 159: 3858–3865, 1997.
- [80467] 18797.Ferrero, E.; Saccucci, F.; Malavasi, F.: The human CD38 gene:polymorphism, CpG island, and linkage to the CD157 (BST–1) gene. Immunogenetics 49:597–604, 1999.
- [80468] 18798.Fukushi, Y.; Kato, I.; Takasawa, S.; Sasaki, T.; Ong, B. H.; Sato,M.; Ohsaga, A.; Sato, K.; Shirato, K.; Okamoto, H.; Maruyama, Y.:Identification of cyclic ADP–ri–bose–dependent mechanisms in pancreaticmuscarinic Ca(2+) signaling using CD38 knockout mice. J. Biol. Chem. 276:649–655, 2001.



- [80469] 18799.Jackson, D. G.; Bell, J. I.: Isolation of a cDNA encoding the human CD38 (T10) molecule, a cell surface glycoprotein with an unusual discontinuous pattern of expression during lymphocyte differentiation. *J.Immun.* 144: 2811–2815, 1990.
- [80470] 18800.Kaisho, T.; Ishikawa, J.; Oritani, K.; Inazawa, J.; Tomizawa, H.; Muraoka, O.; Ochi, T.; Hirano, T.: BST–1, a surface molecule of bone marrow stromal cell lines that facilitates pre–B–cell growth. *Proc.Nat. Acad. Sci.* 91: 5325–5329, 1994.
- [80471] 18801.Nakagawara, K.; Mori, M.; Takasawa, S.; Nata, K.; Takamura, T.; Berlova, A.; Tohgo, A.; Karasawa, T.; Yonekura, H.; Takeuchi, T.; Okamoto, H.: Assignment of CD38, the gene encoding human leukocyte antigen CD38 (ADP–ribosyl cyclase/cyclic ADP–ribose hydrolase), to chromosome 4p15. *Cytogenet. Cell Genet.* 69: 38–39, 1995.
- [80472] 18802.Takasawa, S.; Nata, K.; Yonekura, H.; Okamoto, H.: Cyclic ADP–ribose in insulin secretion from pancreatic beta cells. *Science* 259: 370–373, 1993.
- [80473] 18803.Takasawa, S.; Tohgo, A.; Noguchi, N.; Koguma, T.; Nata, K.; Sugimoto, T.; Yonekura, H.; Okamoto, H.: Synthesis and hydrolysis of cyclic ADP–ribose by human leuko–

cyte antigen CD38 and inhibition of the hydrolysis by ATP.  
J. Biol. Chem. 268: 26052–26054, 1993.

- [80474] 18804. Acosta, J.; Hettinga, J.; Fluckiger, R.; Krumrei, N.; Goldfine, A.; Angarita, L.; Halperin, J.: Molecular basis for a link between complement and the vascular complications of diabetes. Proc. Nat. Acad. Sci. 97: 5450–5455, 2000.
- [80475] 18805. Bickmore, W. A.; Longbottom, D.; Oghene, K.; Fletcher, J. M.; van Heyningen, V.: Colocalization of the human CD59 gene to 11p13 with the MIC11 cell surface antigen. Genomics 17: 129–135, 1993.
- [80476] 18806. Davies, A.; Simmons, D. L.; Hale, G.; Harrison, R. A.; Tighe, H.; Lachmann, P. J.; Waldmann, H.: CD59, an LY-6-like protein expressed in human lymphoid cells, regulates the action of the complement membrane attack complex on homologous cells. J. Exp. Med. 170: 637–654, 1989.
- [80477] 18807. Forsberg, U. H.; Bazil, V.; Stefanova, I.; Schroder, J.: Gene for human CD59 (likely Ly-6 homologue) is located on the short arm of chromosome 11. Immunogenetics 30: 188–193, 1989.
- [80478] 18808. Harada, R.; Okada, N.; Fujita, T.; Okada, H.: Purification of 1F5 antigen that prevents complement attack on homologous cell membranes. J. Immun. 144: 1823–1828,

1990.

- [80479] 18809.Heckl–Ostreicher, B.; Ragg, S.; Drechsler, M.; Scherthan, H.; Royer–Pokora,B.: Localization of the human CD59 gene by fluorescence in situ hybridizationand pulsed–field gel electrophoresis. *Cytogenet. Cell Genet.* 63:144–146, 1993.
- [80480] 18810.Holt, D. S.; Botto, M.; Bygrave, A. E.; Hanna, S. M.; Walport,M. J.; Morgan, B. P.: Targeted deletion of the CD59 gene causes spontaneousintravascular hemolysis and hemoglobinuria. *Blood* 98: 442–449, 2001.
- [80481] 18811.Holt, D. S.; Powell, M. B.; Rushmere, N. K.; Morgan, B. P.: Genomicstructure and chromosome location of the gene encoding mouse CD59. *Cytogenet.Cell Genet.* 89: 264–267, 2000.
- [80482] 18812.Huppi, K.; Duncan, R.; Potter, M.: Myc–1 is centromeric to thelinkage group Ly–6–Sis–Gdc–1 on mouse chromosome 15. *Immunogenetics* 27:215–219, 1988.
- [80483] 18813.Kamiura, S.; Nolan, C. M.; Meruelo, D.: Long–range physical mapof the Ly–6 complex: mapping the Ly–6 multigene family by field–inversionand two–dimensional gel electrophoresis. *Genomics* 12: 89–105, 1992.
- [80484] 18814.Low, M. G.; Saltiel, A. R.: Structural and functional roles ofglycosyl–phosphatidylinositol in membranes. *Sci–*

ence 239: 268–275,1988.

- [80485] 18815.Mahoney, J. F.; Urakaze, M.; Hall, S.; DeGasperi, R.; Chang, H.–M.;Sugiyama, E.; Warren, C. D.; Borowitz, M.; Nicholson–Weller, A.; Rosse,W. F.; Yeh, E. T. H.: Defective glycosylphosphatidylinositol anchorsynthesis in paroxysmal nocturnal hemoglobinuria granulocytes. *Blood* 79:1400–1403, 1992.
- [80486] 18816.Mao, M.; Yu, M.; Tong, J.–H.; Ye, J.; Zhu, J.; Huang, Q.–H.; Fu,G.; Yu, L.; Zhao, S.–Y.; Waxman, S.; Lanotte, M.; Wang, Z.–Y.; Tan,J.–Z.; Chan, S.–J.; Chen, Z.: RIG–E, a human homolog of the murineLy–6 family, is induced by retinoic acid during the differentiationof acute promyelocytic leukemia cell. *Proc. Nat. Acad. Sci.* 93:5910–5914, 1996.
- [80487] 18817.Meri, S.; Morgan, B. P.; Davies, A.; Daniels, R. H.; Olavesen,M. G.; Waldmann, H.; Lachmann, P. J.: Human protectin (CD59), an18,000–20,000 MW complement lysis restricting factor, inhibits C5b–8catalysed insertion of C9 into lipid bilayers. *Immunology* 71: 1–9,1990.
- [80488] 18818.Meri, S.; Morgan, B. P.; Wing, M.; Jones, J.; Davies, A.; Podack,E.; Lachmann, P. J.: Human protectin (CD59), an 18–20–kD homologouscomplement restriction factor, does not restrict perforin–mediatedlysis. *J. Exp. Med.* 172:

367–370, 1990.

- [80489] 18819.Motoyama, N.; Okada, N.; Yamashina, M.; Okada, H.: Paroxysmalnocturnal hemoglobinuria due to hereditary nucleotide deletion in the HRF20 (CD59) gene. *Europ. J. Immun.* 22: 2669–2673, 1992.
- [80490] 18820.Okada, N.; Harada, R.; Fujiita, T.; Okada, H.: A novel membraneglycoprotein capable of inhibiting membrane attack by homologous complement. *Int.Immun.* 1: 205–208, 1989.
- [80491] 18821.Anderson, L.; Anderson, N. G.: High resolution two–dimensionalelectrophoresis of human plasma proteins. *Proc. Nat. Acad. Sci.* 12:5421–5425, 1977.
- [80492] 18822.Cohen, R. D.; Castellani, L. W.; Qiao, J.–H.; Van Lenten, B. J.;Lusis, A. J.; Reue, K.: Reduced aortic lesions and elevated highdensity lipoprotein levels in transgenic mice overexpressing mouseapolipoprotein A–IV. *J. Clin. Invest.* 99: 1906–1916, 1997.
- [80493] 18823.Duverger, N.; Tremp, G.; Caillaud, J.–M.; Emmanuel, F.; Castro,G.; Fruchart, J.–C.; Steinmetz, A.; Deneffe, P.: Protection againstatherogenesis in mice mediated by human apolipoprotein A–IV. *Science* 273:966–968, 1996.
- [80494] 18824.Szpirer, C.; Riviere, M.; Cortese, R.; Nakamura, T.;

Islam, M.Q.; Levan, G.; Szpirer, J.: Chromosomal localization in man and rat of the genes encoding the liver-enriched transcription factors C/EBP, DBP, and HNF1/LFB-1 (CEBP, DBP, and transcription factor 1, TCF1, respectively) and of the hepatocyte growth factor/scatter factor gene (HGF). *Genomics* 13: 293–300, 1992.

[80495] 18825. Boorstein, W. R.; Vamvakopoulos, N. C.; Fiddes, J. C.: Human chorionic gonadotropin beta-subunit is encoded by at least eight genes arranged in tandem and inverted pairs. *Nature* 300: 419–422, 1982.

[80496] 18826. Fiddes, J. C.; Goodman, H. M.: The cDNA for the beta-subunit of human chorionic gonadotropin suggests evolution of a gene by readthrough into the 3-prime-untranslated region. *Nature* 286: 684–687, 1980.

[80497] 18827. Graham, M. Y.; Otani, T.; Boime, I.; Olson, M. V.; Carle, G. F.; Chaplin, D. D.: Cosmid mapping of the human chorionic gonadotropin beta subunit genes by field-inversion gel electrophoresis. *Nucleic Acids Res.* 15: 4437–4448, 1987.

[80498] 18828. Julier, C.; Weil, D.; Couillin, P.; Cote, J. C.; Boue, A.; Thririon, J. P.; Kaplan, J. C.; Junien, C.: Confirmation of the assignment of the genes coding for human chorionic go-

nadotropin beta subunit to chromosome 19. (Abstract) Cytogenet. Cell Genet. 37: 501–502, 1984.

- [80499] 18829. Gersten, K. M.; Natsuka, S.; Trinchera, M.; Petryniak, B.; Kelly, R. J.; Hiraiwa, N.; Jenkins, N. A.; Gilbert, D. J.; Copeland, N. G.; Lowe, J. B.: Molecular cloning, expression, chromosomal assignment, and tissue-specific expression of a murine alpha-(1,3)-fucosyltransferase locus corresponding to the human ELAM-1 ligand fucosyl transferase. J. Biol. Chem. 270: 25047–25056, 1995.
- [80500] 18830. Julier, C.; Weil, D.; Couillin, P.; Cote, J. C.; Van Cong, N.; Foubert, C.; Boue, A.; Thirion, J. P.; Kaplan, J. C.; Junien, C.: The beta chorionic gonadotropin-beta luteinizing gene cluster maps to human chromosome 19. Hum. Genet. 67: 174–177, 1984.
- [80501] 18831. Lunardi-Iskander, Y.; Bryant, J. L.; Zeman, R. A.; Lam, V. H.; Samaniego, F.; Besnier, J. M.; Hermans, P.; Thierry, A. R.; Gill, P.; Gallo, R. C.: Tumorigenesis and metastasis of neoplastic Kaposi's sarcoma cell line in immunodeficient mice blocked by a human pregnancy hormone. Nature 375: 64–68, 1995.
- [80502] 18832. Policastro, P.; Ovitt, C. E.; Hoshina, M.; Fukuoka, H.; Boothby, M. R.; Bieme, I.: The beta-subunit of human chorionic gonadotropin is encoded by multiple genes. J.

Biol. Chem. 258: 11492–11499, 1983.

- [80503] 18833.Policastro, P. F.; Daniels–McQueen, S.; Carle, G.; Boime, I.:A map of the hCG–beta–LH–beta gene cluster. J. Biol. Chem. 261:5907–5916, 1986.
- [80504] 18834.Talmadge, K.; Vamvakopoulos, N. C.; Fiddes, J. C.: Evolutionof the genes for the beta subunits of human chorionic gonadotropinand luteinizing hormone. Nature 307: 37–40, 1984.
- [80505] 18835.Warburton, D.; Gersen, S.; Yu, M.–T.; Jackson, C.; Handelin, B.;Housman, D.: Monochromosomal rodent–human hybrids from microcellfusion of human lymphoblastoid cells containing an inserted dominantselectable marker. Genomics 6: 358–366, 1990.
- [80506] 18836.Reseland, J. E.; Larsen, F.; Solheim, J.; Eriksen, J. A.; Hanssen,L. E.; Prydz, H.: A novel human chymotrypsin–like digestive enzyme. J.Biol. Chem. 272: 8099–8104, 1997.
- [80507] 18837.Angeletti, R. H.: Chromogranins and neuroendocrine secretion.(Editorial) Lab. Invest. 55: 387–390, 1986.
- [80508] 18838.Bhargava, G.; Russell, J.; Sherwood, L. M.: Phosphorylation ofparathyroid secretory protein. Proc. Nat. Acad. Sci. 80: 878–881,1983.
- [80509] 18839.Cetin, Y.; Aunis, D.; Bader, M.–F.; Galindo, E.; Jorns,



A.; Bargsten,G.; Grube, D.: Chromostatin, a chromogranin A-derived bioactive peptide,is present in human pancreatic insulin (beta) cells. Proc. Nat. Acad.Sci. 90: 2360-2364, 1993.

[80510] 18840.Lu, Q.; Lemke, G.: Homeostatic regulation of the immune systemby receptor tyrosine kinases of the Tyro 3 family. Science 293:306-311, 2001.

[80511] 18841.Abrahamson, M.: Human cysteine proteinase inhibitors: isolation,physiological importance, inhibitory mechanism, gene structure andrelation to hereditary cerebral hemorrhage. Scand. J. Clin. Lab.Invest. 48 (suppl. 191): 21-31, 1988.

[80512] 18842.Abrahamson, M.; Grubb, A.: Increased body temperature acceleratesaggregation of the leu68-to-gln mutant cystatin C, the amyloid-formingprotein in hereditary cystatin C amyloid angiopathy. Proc. Nat. Acad.Sci. 91: 1416-1420, 1994.

[80513] 18843.Abrahamson, M.; Grubb, A.; Olafsson, I.; Lundwall, A.: Molecularcloning and sequence analysis of cDNA coding for the precursor ofthe human cysteine proteinase inhibitor cystatin C. FEBS Lett. 216:229-233, 1987.

[80514] 18844.Abrahamson, M.; Islam, M. Q.; Szpirer, J.; Szpirer, C.; Levan,G.: The human cystatin C gene (CST3), mutated

in hereditary cystatinC amyloid angiopathy, is located on chromosome 20. Hum. Genet. 82:223–226, 1989.

- [80515] 18845.Abrahamson, M.; Jonsdottir, S.; Olafsson, I.; Jensson, O.; Grubb,A.: Hereditary cystatin C amyloid angiopathy: identification of the disease-causing mutation and specific diagnosis by polymerase chain reaction based analysis. Hum. Genet. 89: 377–380, 1992.
- [80516] 18846.Cohen, D. H.; Feiner, H.; Jensson, O.; Frangione, B.: Amyloid fibril in hereditary cerebral hemorrhage with amyloidosis (HCHWA) is related to the gastroentero-pancreatic neuroendocrine protein, gamma trace. J. Exp. Med. 158: 623–628, 1983.
- [80517] 18847.Ghiso, J.; Jensson, O.; Frangione, B.: Amyloid fibrils in hereditary cerebral hemorrhage with amyloidosis of Icelandic type is a variant of gamma-trace basic protein (cystatin C). Proc. Nat. Acad. Sci. 83:2974–2978, 1986.
- [80518] 18848.Ghiso, J.; Pons-Estel, B.; Frangione, B.: Hereditary cerebral amyloid angiopathy: the amyloid fibrils contain a protein which is a variant of cystatin C, an inhibitor of lysosomal cysteine proteases. Biochem.Biophys. Res. Commun. 136: 548–554, 1986.
- [80519] 18849.Grubb, A.; Jensson, O.; Gudmundsson, G.; Arnason, A.; Lofberg,H.; Malm, J.: Abnormal metabolism of gamma-

trace alkaline microprotein:the basic defect in hereditary cerebral hemorrhage with amyloidosis. NewEng. J. Med. 311: 1547–1549, 1984.

- [80520] 18850. Galbraith, G. M. P.; Pandey, J. P.: Tumor necrosis factor alpha (TNF-alpha) gene polymorphism in alopecia areata. *Hum. Genet.* 96:433-436, 1995.
- [80521] 18851. Tazi-Ahnini, R.; di Giovine, F. S.; McDonagh, A. J. G.; Messenger, A. G.; Amadou, C.; Cox, A.; Duff, G. W.; Cork, M. J.: Structure and polymorphism of the human gene for the interferon-induced p78 protein (MX1): evidence of association with alopecia areata in the Down syndrome region. *Hum. Genet.* 106: 639-645, 2000.
- [80522] 18852. Webb, D. J.; Wen, J.; Lysiak, J. J.; Umans, L.; Van Leuven, F.; Gonias, S. L.: Murine alpha-macroglobulins demonstrate divergent activities as neutralizers of transforming growth factor-beta and as inducers of nitric oxide synthesis: a possible mechanism for the endotoxin insensitivity of the alpha-2-macroglobulin gene knockout mouse. *J. Biol. Chem.* 271: 24982-24988, 1996.
- [80523] 18853. Wilson, A. G.; di Giovine, F. S.; Blakemore, A. I. F.; Duff, G. W.: Single base polymorphism in the human tumour necrosis factor alpha (TNF-alpha) gene detectable by *Nco*I restriction of PCR product. *Hum. Molec. Genet.* 1: 353 only, 1992.
- [80524] 18854. Belanger, L.; Roy, S.; Allard, D.: New albumin gene 3-prime adjacent to the alpha-1-fetoprotein locus. *J. Biol.*

Chem. 269: 5481–5484,1994.

- [80525] 18855.Lichenstein, H. S.; Lyons, D. E.; Wurfel, M. M.; Johnson, D. A.;McGinley, M. D.; Leidli, J. C.; Trollinger, D. B.; Mayer, J. P.; Wright,S. D.; Zukowski, M. M.: Afamin is a new member of the albumin, alpha-fetoprotein,and vitamin D-binding protein gene family. J. Biol. Chem. 269: 18149–18154,1994.
- [80526] 18856.Nishio, H.; Dugaiczky, A.: Complete structure of the human alpha-albumingene, a new member of the serum albumin multigene family. Proc. Nat.Acad. Sci. 93: 7557–7561, 1996.
- [80527] 18857.Song, Y.–H.; Naumova, A. K.; Liebhaver, S. A.; Cooke, N. E.: Physicaland meiotic mapping of the region of human chromosome 4q11–q13 encompassingthe vitamin D binding protein DBP/Gc-globulin and albumin multigenecluster. Genome Res. 9: 581–587, 1999.
- [80528] 18858.Beattie, W. G.; Dugaiczky, A.: Structure and evolution of humanalpha-fetoprotein deduced from partial sequence of cloned cDNA. Gene 20:415–422, 1982.
- [80529] 18859.Belayew, A.; Tilghman, S. M.: Genetic analysis of alpha-fetoproteinsynthesis in mice. Molec. Cell. Biol. 2: 1427–1435, 1982.
- [80530] 18860.D'Eustachio, P.; Ingram, R. S.; Tilghman, S. M.;

Ruddle, F. H.: Murine alpha-fetoprotein and albumin: two evolutionarily linked proteins encoded on the same mouse chromosome. *Somat. Cell Genet.* 7:289–294, 1981.

[80531] 18861. Eiferman, F. A.; Young, P. R.; Scott, R. W.; Tilghman, S. M.: Intragenic amplification and divergence in the mouse alpha-fetoprotein gene. *Nature* 294: 713–718, 1981.

[80532] 18862. Faucett, W. A.; Greenberg, F.; Rose, E.; Alpert, E.; Bancalari, L.; Kardon, N. B.; Mizjewski, G.; Knight, G.; Haddow, J. E.: Congenital deficiency of alpha-fetoprotein. (Abstract) *Am. J. Hum. Genet.* 45(suppl.): A259, 1989.

[80533] 18863. Ferguson-Smith, M. A.; May, H. M.; Aitken, D. A.; O'Hare, E.; Yates, J. R. W.; Gallagher, J.; Krumlauf, R.; Tilghman, S. M.: Hereditary persistence of alpha-fetoprotein (HPAFP); linkage studies with chromosome 4 markers. (Abstract) *Cytogenet. Cell Genet.* 37: 469, 1984.

[80534] 18864. Ferguson-Smith, M. A.; Yates, J. R. W.; Kelly, D.; Aitken, D. A.; May, H. M.; Krumlauf, R.; Tilghman, S. M.: Hereditary persistence of alpha-fetoprotein maps to the long arm of chromosome 4. (Abstract) *Cytogenet. Cell Genet.* 40: 628, 1985.

[80535] 18865. Gabant, P.; Forrester, L.; Nichols, J.; Van Reeth, T.; De Mees, C.; Pajack, B.; Watt, A.; Smits, J.; Alexandre, H.;

Szpirer, C.; Szpirer, J.: Alpha-fetoprotein, the major fetal serum protein, is not essential for embryonic development but is required for female fertility. *Proc. Nat. Acad. Sci.* 99: 12865–12870, 2002.

[80536] 18866. Gibbs, P. E. M.; Zielinski, R.; Boyd, C.; Dugaiczyk, A.: Structure, polymorphism, and novel repeated DNA elements revealed by a complete sequence of the human alpha-fetoprotein gene. *Biochemistry* 26: 1332–1343, 1987.

[80537] 18867. Seri, M.; Celli, I.; Betsos, N.; Claudiani, F.; Camera, G.; Romeo, G.: A cys634gly substitution of the RET proto-oncogene in a family with recurrence of multiple endocrine neoplasia type 2A and cutaneous lichen amyloidosis. *Clin. Genet.* 51: 86–90, 1997.

[80538] 18868. Kamal, A.; Almenar-Queralt, A.; LeBlanc, J. F.; Roberts, E. A.; Goldstein, L. S. B.: Kinesin-mediated axonal transport of a membrane compartment containing beta-secretase and presenilin-1 requires APP. *Nature* 414: 643–648, 2001.

[80539] 18869. Kang, D. E.; Soriano, S.; Xia, X.; Eberhart, C. G.; De Strooper, B.; Zheng, H.; Koo, E. H.: Presenilin couples the paired phosphorylation of beta-catenin independent of Axin: implications for beta-catenin inactivation in tumorigenesis. *Cell* 110: 751–762, 2002.

- [80540] 18870.Hofstra, R. M. W.; Sijmons, R. H.; Stelwagen, T.; Stulp, R. P.;Kousseff, B. G.; Lips, C. J. M.; Steijlen, P. M.; Van Voorst Vader,P. C.; Buys, C. H. C. M.: RET mutation screening in familial cutaneouslichen amyloidosis and in skin amyloidosis associated with multipleendocrine neoplasia. J. Invest. Derm. 107: 215–218, 1996.
- [80541] 18871.Bordet, T.; Lesbordes, J.–C.; Rouhani, S.; Castelnau–Ptakhine,L.; Schmalbruch, H.; Haase, G.; Kahn, A.: Protective effects of cardiotrophin–1adenoviral gene transfer on neuromuscular degeneration in transgenicALS mice. Hum. Molec. Genet. 10: 1925–1933, 2001.
- [80542] 18872.Fantes, J. A.; Bickmore, W. A.; Fletcher, J. M.; Ballesta, F.;Hanson, I. M.; van Heyningen, V.: Submicroscopic deletions at theWAGR locus, revealed by nonradioactive in situ hybridization. Am.J. Hum. Genet. 51: 1286–1294, 1992.
- [80543] 18873.Glaser, T.; Jepeal, L.; Edwards, J. G.; Young, S. R.; Favor, J.;Maas, R. L.: PAX6 gene dosage effect in a family with congenitalcataracts, aniridia, anophthalmia and central nervous system defects. NatureGenet. 7: 463–471, 1994.
- [80544] 18874.Glaser, T.; Lane, J.; Housman, D.: A mouse model of the aniridia–Wilmstumor deletion syndrome. Science



250: 823–827, 1990.

- [80545] 18875.Gronskov, K.; Olsen, J. H.; Sand, A.; Pedersen, W.; Carlsen, N.;Jylling, A. M. B.; Lyngbye, T.; Brondum-Nielsen, K.; Rosenberg, T.: Population-based risk estimates of Wilms tumor in sporadic aniridia:a comprehensive mutation screening procedure of PAX6 identifies 80%of mutations in aniridia. Hum. Genet. 109: 11–18, 2001.
- [80546] 18876.Hanson, I. M.; Seawright, A.; Hardman, K.; Hodgson, S.; Zaletayev,D.; Fekete, G.; van Heyningen, V.: PAX6 mutations in aniridia. Hum.Molec. Genet. 2: 915–920, 1993.
- [80547] 18877.Hill, R. E.; Favor, J.; Hogan, B. L. M.; Ton, C. C. T.; Saunders,G. F.; Hanson, I. M.; Prosser, J.; Jordan, T.; Hastie, N. D.; vanHeyningen, V.: Mouse small eye results from mutations in a paired-likehomeobox-containing gene. Nature 354: 522–525, 1991.
- [80548] 18878.Jordan, T.; Hanson, I.; Zaletayev, D.; Hodgson, S.; Prosser, J.;Seawright, A.; Hastie, N.; van Heyningen, V.: The human PAX6 geneis mutated in two patients with aniridia. Nature Genet. 1: 328–332,1992.
- [80549] 18879.Karpen, G. H.: Position effect variegation and the new biologyof heterochromatin. Curr. Opin. Genet. Dev. 4:

281–291, 1994.

- [80550] 18880.Lyon, M. F.: Personal Communication. Harwell, England 6/9/1988.
- [80551] 18881.Martha, A.; Strong, L. C.; Ferrell, R. E.; Saunders, G. F.: Threenovel aniridia mutations in the human PAX6 gene. Hum. Mutat. 6:44–49, 1995.
- [80552] 18882.Matsuo, T.; Osumi–Yamashita, N.; Noji, S.; Ohuchi, H.; Koyama,E.; Myokai, F.; Matsuo, N.; Taniguchi, S.; Doi, H.; Iseki, S.; Ninomiya,Y.; Fujiwara, M.; Watanabe, T.; Eto, K.: A mutation in the Pax–6gene in rat small eye is associated with impaired migration of midbraincrest cells. Nature Genet. 3: 299–304, 1993.
- [80553] 18883.Oliver, M. D.; Dotan, S. A.; Chemke, J.; Abraham, F. A.: Isolatedfoveal hypoplasia. Brit. J. Ophthal. 71: 926–930, 1987.
- [80554] 18884.Prosser, J.; van Heyningen, V.: PAX6 mutations reviewed. Hum.Mutat. 11: 93–108, 1998.
- [80555] 18885.Quiring, R.; Walldorf, U.; Kloter, U.; Gehring, W. J.: Homologyof the eyeless gene of Drosophila to the small eye gene in mice andaniridia in humans. Science 265: 785–789, 1994.
- [80556] 18886.Salvini–Plawen, L.; Mayr, E.: On the evolution of photoreceptorsand eyes.In: Hecht, M. K.; Steere, W.; Wal–

lace, B.: Evolutionary Biology. New York: Plenum Pub.

(pub.) 10: 1977. Pp. 207–263.

[80557] 18887.Schedl, A.; Ross, A.; Lee, M.; Engelkamp, D.; Rashbass, P.; vanHeyningen, V.; Hastie, N. D.: Influence of PAX6 gene dosage on development: overexpression causes severe eye abnormalities. Cell 86: 71–82, 1996.

[80558] 18888.Stone, D. L.; Kenyon, K. R.; Green, W. R.; Ryan, S. J.: Congenital central corneal leukoma (Peters' anomaly). Am. J. Ophthalm. 81: 173–193, 1976.

[80559] 18889.van der Meer–de Jong, R.; Dickinson, M. E.; Woychik, R. P.; Stubbs, L.; Hetherington, C.; Hogan, B. L. M.: Location of the gene involving the small eye mutation on mouse chromosome 2 suggests homology with human aniridia 2 (AN2). Genomics 7: 270–275, 1990.

[80560] 18890.Newport, M. J.; Huxley, C. M.; Huston, S.; Hawrylowicz, C. M.; Oostra, B. A.; Williamson, R.; Levin, M.: A mutation in the interferon–gamma–receptor gene and susceptibility to mycobacterial infection. New Eng. J. Med. 335: 1941–1949, 1996.

[80561] 18891.Cowan, C. A.; Yokoyama, N.; Bianchi, L. M.; Henkemeyer, M.; Fritzsche, B.: EphB2 guides axons at the midline and is necessary for normal vestibular function. Neuron 26: 417–430, 2000.

- [80562] 18892.Ton, C. C. T.; Hirvonen, H.; Miwa, H.; Weil, M. M.; Monaghan,P.; Jordan, T.; van Heyningen, V.; Hastie, N. D.; Meijers–Heijboer,H.; Drechsler, M.; Royer–Pokora, B.; Collins, F.; Swaroop, A.; Strong,L. C.; Saunders, G. F.: Positional cloning and characterization of a paired box– and homeobox–containing gene from the aniridia region. *Cell* 67:1059–1074, 1991.
- [80563] 18893.Novick, D.; Orchansky, P.; Revel, M.; Rubinstein, M.: The humaninterferon–gamma receptor: purification, characterization, and preparationof antibodies. *J. Biol. Chem.* 262: 8483–8487, 1987.
- [80564] 18894.Orchansky, P.; Rubinstein, M.; Fischer, D. G.: The interferon–gammareceptor in human monocytes is different from the one in nonhematopoieticcells. *J. Immun.* 136: 169–173, 1986.
- [80565] 18895.Papanicolaou, G. J.; Parsa, N. Z.; Meltzer, P. S.; Trent, J. M.: Assignment of interferon gamma receptor (IFNGR1) to human chromosomebands 6q24.1–q24.2 by in situ hybridization. *Cytogenet. Cell Genet.* 76:181–182, 1997. Note: Erratum: *Cytogenet. Cell Genet.* 78: 132 only,1997.
- [80566] 18896.Pierre–Audigier, C.; Jouanguy, E.; Lamhamedi, S.; Altare, F.;Rauzier, J.; Vincent, V.; Canioni, D.; Emile, J. F.;

Fischer, A.;Blanche, S.; Gaillard, J. L.; Casanova, J. L.: Fatal disseminatedMycobacterium smegmatis infection in a child with inherited interferongamma receptor deficiency. Clin. Infect. Dis. 24: 982–984, 1997.

- [80567] 18897.Rashidbaigi, A.; Langer, J. A.; Jung, V.; Jones, C.; Morse, H.G.; Tischfield, J. A.; Trill, J. J.; Kung, H.–F.; Pestka, S.: Thegene for the human immune interferon receptor is located on chromosome6. Proc. Nat. Acad. Sci. 83: 384–388, 1986.
- [80568] 18898.Rettig, W. J.; Grzeschik, K.–H.; Yenamandra, A. K.; Garcia, E.;Old, L. J.: Definition of selectable cell surface markers for humanchromosomes and chromosome segments in rodent–human hybrids. Somat.Cell Molec. Genet. 14: 223–231, 1988.
- [80569] 18899.Shankaran, V.; Ikeda, H.; Bruce, A. T.; White, J. M.; Swanson,P. E.; Old, L. J.; Schreiber, R. D.: IFN–gamma and lymphocytes preventprimary tumour development and shape tumour immunogenicity. Nature 410:1107–1111, 2001.
- [80570] 18900.Bauer, R.; Imhof, A.; Pscherer, A.; Kopp, H.; Moser, M.; Seegers,S.; Kerscher, M.; Tainsky, M. A.; Hofstaedter, F.; Buettner, R.:The genomic structure of the human AP–2 transcription factor. NucleicAcids Res. 22: 1413–1420,

1994.

- [80571] 18901. Buettner, R.; Kannan, P.; Imhof, A.; Bauer, R.; Yim, S. O.; Glockshuber, R.; Van Dyke, M. W.; Tainsky, M. A.: An alternatively spliced mRNA from the AP-2 gene encodes a negative regulator of transcriptional activation by AP-2. *Molec. Cell. Biol.* 13: 4174–4185, 1993.
- [80572] 18902. Ahmad, I.: Mash-1 is expressed during ROD photoreceptor differentiation and binds an E-box, E(opsin-1), in the rat opsin gene. *Brain Res. Dev. Brain Res.* 90: 184–189, 1995.
- [80573] 18903. Ball, D. W.; Azzoli, C. G.; Baylin, S. B.; Chi, D.; Dou, S.; Donis-Keller, H.; Cumaraswamy, A.; Borges, M.; Nelkin, B. D.: Identification of a human achaete-scute homolog highly expressed in neuroendocrine tumors. *Proc. Nat. Acad. Sci.* 90: 5648–5652, 1993.
- [80574] 18904. Guillemot, F.; Lo, L.-C.; Johnson, J. E.; Auerbach, A.; Anderson, D. J.; Joyner, A. L.: Mammalian achaete-scute homolog 1 is required for the early development of olfactory and autonomic neurons. *Cell* 75:463–476, 1993.
- [80575] 18905. Renault, B.; Lieman, J.; Ward, D.; Krauter, K.; Kucherlapati, R.: Localization of the human achaete-scute homolog gene (ASCL1) distal to phenylalanine hydroxylase (PAH) and proximal to tumor rejection antigen (TRA1) on

chromosome 12q22–q23. *Genomics* 30: 81–83, 1995.

[80576] 18906. Yang, B.; Ma, T.; Verkman, A. S.: Erythrocyte water permeability and renal function in double knockout mice lacking aquaporin–1 and aquaporin–3. *J. Biol. Chem.* 276: 624–628, 2001.

[80577] 18907. Bunz, F.; Kobayashi, R.; Stillman, B.: cDNAs encoding the large subunit of human replication factor C. *Proc. Nat. Acad. Sci.* 90: 11014–11018, 1993.

[80578] 18908. Lossie, A. C.; Haugen, B. R.; Wood, W. M.; Camper, S. A.; Gordon, D. F.: Chromosomal localization of the large subunit of mouse replication factor C in the mouse and human. *Mammalian Genome* 6: 58–59, 1995.

[80579] 18909. Luckow, B.; Bunz, F.; Stillman, B.; Lichter, P.; Schutz, G.: Cloning, expression, and chromosomal localization of the 140–kilodalton subunit of replication factor C from mice and humans. *Molec. Cell. Biol.* 14: 1626–1634, 1994.

[80580] 18910. Pennaneach, V.; Salles–Passador, I.; Munshi, A.; Brickner, H.; Regazzoni, K.; Dick, F.; Dyson, N.; Chen, T.–T.; Wang, J. Y. J.; Fotedar, R.; Fotedar, A.: The large subunit of replication factor C promotes cell survival after DNA damage in an LxCxE motif– and Rb–dependent manner. *Molec. Cell* 7: 715–727, 2001.

- [80581] 18911.Uchiumi, F.; Ohta, T.; Tanuma, S.: Replication factor C recognizes 5-prime-phosphate ends of telomeres. *Biochem. Biophys. Res. Commun.* 229:310–315, 1996.
- [80582] 18912.Schmickel, R. D.: Contiguous gene syndromes: a component of recognizable syndromes. *J. Pediat.* 109: 231–241, 1986.
- [80583] 18913.Olson, E.; Srivastava, D.: Molecular pathways controlling heart development. *Science* 272: 671–676, 1996.
- [80584] 18914.Lamballe, F.; Klein, R.; Barbacid, M.: TRKC, a new member of the TRK family of tyrosine protein kinases, is a receptor for neurotrophin-3. *Cell* 66:967–979, 1991.
- [80585] 18915.McGregor, L. M.; Baylin, S. B.; Griffin, C. A.; Hawkins, A. L.; Nelkin, B. D.: Molecular cloning of the cDNA for human TrkC (NTRK3), chromosomal assignment, and evidence for a splice variant. *Genomics* 22:267–272, 1994.
- [80586] 18916.Segal, R. A.; Goumnerova, L. C.; Kwon, Y. K.; Stiles, C. D.; Pomeroy, S. L.: Expression of the neurotrophin receptor TrkC is linked to a favorable outcome in medulloblastoma. *Proc. Nat. Acad. Sci.* 91:12867–12871, 1994.
- [80587] 18917.Valent, A.; Danglot, G.; Bernheim, A.: Mapping of the tyrosine kinase receptors trkA (NTRK1), trkB (NTRK2) and trkC (NTRK3) to human chromosomes 1q22, 9q22 and



15q25 by fluorescence in situ hybridization. *Europ.J. Hum. Genet.* 5: 102–104, 1997.

[80588] 18918.Kielkopf, C. L.; Rodionova, N. A.; Green, M. R.; Burley, S. K.: A novel peptide recognition mode revealed by the X-ray structure of a core U2AF35/U2AF65 heterodimer. *Cell* 106: 595–605, 2001.

[80589] 18919.Lalioti, M. D.; Gos, A.; Green, M. R.; Rossier, C.; Morris, M.A.; Antonarakis, S. E.: The gene for human U2 snRNP auxiliary factor small 35-kDa subunit (U2AF1) maps to the progressive myoclonus epilepsy (EPM1) critical region on chromosome 21q22.3. *Genomics* 33: 298–300, 1996.

[80590] 18920.Hayashizaki, Y.; Shibata, H.; Hirotsune, S.; Sugino, H.; Okazaki, Y.; Sasaki, N.; Hirose, K.; Imoto, H.; Okuizumi, H.; Muramatsu, M.; Komatsubara, H.; Shiroishi, T.; Moriwaki, K.; Katsuki, M.; Hatano, N.; Sasaki, H.; Ueda, T.; Mise, N.; Takagi, N.; Plass, C.; Chapman, V. M.: Identification of an imprinted U2af binding protein related sequence on mouse chromosome 11 using the RLGS method. *Nature Genet.* 6:33–40, 1994.

[80591] 18921.Zhang, M.; Zamore, P. D.; Carmo-Fonseca, M.; Lamond, A. I.; Green, M. R.: Cloning and intracellular localization of the U2 small nuclear ribonucleoprotein auxiliary

factor small subunit. Proc. Nat. Acad.Sci. 89: 8769–8773, 1992.

[80592] 18922.Duncan, A. M. V.; Anderson, L.; Duff, C.; Ozawa, T.; Suzuki, H.;Worton, R.; Rozen, R.: Assignment of the gene (UQCRFS1) for the Rieskeiron–sulfur protein subunit of the mitochondrial cytochrome bc–1 complex to the 22q13 and 19q12–q13.1 regions of the human genome. Genomics 21:281–283, 1994.

[80593] 18923.Pennacchio, L. A.; Bergmann, A.; Fukushima, A.; Okubo, K.; Salemi,A.; Lennon, G. G.: Structure, sequence and location of the UQCRFS1gene for the human Rieske Fe–S protein. Gene 155: 207–211, 1995.

[80594] 18924.Cunningham, J. M.; Vanin, E. F.; Tran, N.; Valentine, M.; Jane,S. M.: The human transcription factor CP2 (TFCP2), a component ofthe human gamma–globin stage selector protein, maps to chromosomeregion 12q13 and is within 250 kb of the NF–E2 gene. Genomics 30:398–399, 1995.

[80595] 18925.Jane, S. M.; Nienhuis, A. W.; Cunningham, J. M.: Hemoglobin switchingin man and chicken is mediated by a heteromeric complex between theubiquitous transcrip–tion factor CP2 and a developmentally specificprotein. EMBO J. 13: 197–105, 1995.

- [80596] 18926.Lambert, J.-C.; Goumidi, L.; Wavrant-De Vrieze, F.; Frigard, B.;Harris, J. M.; Cummings, A.; Coates, J.; Pasquier, F.; Cotel, D.;Gaillac, M.; St. Clair, D.; Mann, D. M. A.; Hardy, J.; Lendon, C.L.; Amouyel, P.; Chartier-Harlin, M.-C.: The transcriptional factorLBP-1c/CP2/LSF gene on chromosome 12 is a genetic determinant of Alzheimer'sdisease. Hum. Molec. Genet. 9: 2275-2280, 2000.
- [80597] 18927.Swendeman, S. L.; Spielholz, C.; Jenkins, N. A.; Gilbert, D. J.;Copeland, N. G.; Sheffery, M.: Characterization of the genomic structure,chromosomal location, promoter, and developmental expression of thealpha-globin transcription factor CP2. J. Biol. Chem. 269: 11663-11671,1994.
- [80598] 18928.Taylor, A. E.; Yip, A.; Brayne, C.; Easton, D.; Evans, J. G.; Xuereb,J.; Cairns, N.; Esiri, M. M.; Rubinsztein, D. C.: Genetic associationof an LBP-1c/CP2/LSF gene polymorphism with late onset Alzheimer'sdisease. J. Med. Genet. 38: 232-233, 2001.
- [80599] 18929.Chen, C.-R.; Kang, Y.; Siegel, P. M.; Massague, J.: E2F4/5 andp107 as Smad cofactors linking the TGF-beta receptor to c-myc repression. Cell 110:19-32, 2002.
- [80600] 18930.Zhang, Y.; Venkatraj, V. S.; Fischer, S. G.; Warbur-

ton, D.; Chellappan, S. P.: Genomic cloning and chromosomal assignment of the E2F dimerization partner TFDP gene family. *Genomics* 39: 95–98, 1997.

- [80601] 18931. Moloney, D. J.; Panin, V. M.; Johnston, S. H.; Chen, J.; Shao, L.; Wilson, R.; Wang, Y.; Stanley, P.; Irvine, K. D.; Haltiwanger, R. S.; Vogt, T. F.: Fringe is a glycosyltransferase that modifies Notch. *Nature* 406: 369–375, 2000.
- [80602] 18932. Sestan, N.; Artavanis-Tsakonas, S.; Rakic, P.: Contact-dependent inhibition of cortical neurite growth mediated by Notch signaling. *Science* 286: 741–746, 1999.
- [80603] 18933. Tanigaki, K.; Nogaki, F.; Takahashi, J.; Tashiro, K.; Kurooka, H.; Honjo, T.: Notch1 and Notch3 instructively restrict bFGF-responsive multipotent neural progenitor cells to an astroglial fate. *Neuron* 29: 45–55, 2001.
- [80604] 18934. Barton, D. E.; Foellmer, B. E.; Du, J.; Tamm, J.; Derynck, R.; Francke, U.: Chromosomal mapping of genes for transforming growth factors beta-2 and beta-3 in man and mouse: dispersion of TGF-beta gene family. *Oncogene Res.* 3: 323–331, 1988.
- [80605] 18935. Dean, M.; Park, M.; Vande Woude, G. F.: Characterization of the rearranged TPR-MET oncogene breakpoint. *Molec. Cell. Biol.* 7: 921–924, 1987.
- [80606] 18936. Robinson, B. H.; Taylor, J.; Sherwood, W. G.: The

genetic heterogeneity of lactic acidosis: occurrence of recognisable inborn errors of metabolism in a pediatric population with lactic acidosis. *Pediat. Res.* 14:950–962, 1980.

[80607] 18937. Hosokawa, Y.; Suzuki, H.; Nishikimi, M.; Matsukage, A.; Yoshida, M. C.; Ozawa, T.: Chromosomal assignment of the gene for the ubiquinone-binding protein of human mitochondrial cytochrome bc<sub>1</sub> complex. *Biochem. Int.* 21:41–44, 1990.

[80608] 18938. Suzuki, H.; Hosokawa, Y.; Toda, H.; Nishikimi, M.; Ozawa, T.: Cloning and sequencing of a cDNA for human mitochondrial ubiquinone-binding protein of complex III. *Biochem. Biophys. Res. Commun.* 156: 987–994, 1988.

[80609] 18939. van Leeuwen, F. W.; de Kleijn, D. P. V.; van den Hurk, H. H.; Neubauer, A.; Sonnemans, M. A. F.; Sluijs, J. A.; Koycu, S.; Ramdjielal, R. D. J.; Salehi, A.; Martens, G. J. M.; Grosveld, F. G.; Burbach, J. P. H.; Hol, E. M.: Frameshift mutants of beta-amyloid precursor protein and ubiquitin-B in Alzheimer's and Down patients. *Science* 279: 242–247, 1998.

[80610] 18940. Webb, G. C.; Baker, R. T.; Fagan, K.; Board, P. G.: Localization of the human UbB polyubiquitin gene to chromosome band 17p11.1–17p12. *Am. J. Hum. Genet.* 46:

308–315, 1990.

- [80611] 18941.Vidal, R.; Frangione, B.; Rostagno, A.; Mead, S.; Revesz, T.; Plant,G.; Ghiso, J.: A stop-codon mutation in the BRI gene associated withfamilial British dementia. *Nature* 399: 776–781, 1999.
- [80612] 18942.Adler, D. A.; Tseng, B. Y.; Wang, T. S.–F.; Disteché, C. M.: Physicalmapping of the genes for three components of the mouse DNA replicationcomplex: polymerase alpha to the X chromosome, primase p49 subunitto chromosome 10, and primase p58 subunit to chromosome 1. *Genomics* 9:642–646, 1991.
- [80613] 18943.Ogilvie, A. D.; Battersby, S.; Bubb, V. J.; Fink, G.; Harmar,A. J.; Goodwin, G. M.; Smith, C. A. D.: Polymorphism in serotonintransporter gene associated with susceptibility to major depression. *Lancet* 347:731–733, 1996.
- [80614] 18944.Doniger, J.; DiPaolo, J. A.: Coordinate N–RAS mRNA up-regulationwith mutational activation in tumorigenic guinea pig cells. *NucleicAcids Res.* 16: 969–980, 1988.
- [80615] 18945.Jeffers, M.; Paciucci, R.; Pellicer, A.: Characterization of UNR:a gene closely linked to N–RAS. *Nucleic Acids Res.* 18: 4891–4899,1990.
- [80616] 18946.Mirnics, K.; Middleton, F. A.; Marquez, A.; Lewis, D.

A.; Levitt,P.: Molecular characterization of schizophrenia viewed by microarrayanalysis of gene expression in pre-frontal cortex. *Neuron* 28: 53–67,2000.

[80617] 18947.Engelkamp, D.; Schafer, B. W.; Mattei, M. G.; Erne, P.; Heizmann,C. W.: Six S100 genes are clustered on human chromosome 1q21: identificationof two genes coding for the two previously unreported calcium-bindingproteins S100D and S100E. *Proc. Nat. Acad. Sci.* 90: 6547–6551, 1993.

[80618] 18948.Herrera, G. A.; Turbat-Herrera, E. A.; Lott, R. L.: S-100 proteinexpression by primary and metastatic adenocarcinomas. *Am. J. Clin.Path.* 89: 168–176, 1988.

[80619] 18949.Isobe, T.; Tsugira, A.; Okuyama, T.: Amino acid sequence of thesubunit structure of bovine brain S-100 protein (PAP 1-b). *J. Neurochem.* 30:921–923, 1978.

[80620] 18950.Moore, B. W.: A soluble protein characteristic of the nervoussystem. *Biochem. Biophys. Res. Commun.* 19: 739–744, 1965.

[80621] 18951.Morii, K.; Tanaka, R.; Takahashi, Y.; Minoshima, S.; Fukuyama,R.; Shimizu, N.; Kuwano, R.: Structure and chromosome assignmentof human S100 alpha and beta subunit genes. *Biochem. Biophys. Res.Comm.* 175: 185–191, 1991.

- [80622] 18952. Most, P.; Bernotat, J.; Ehlermann, P.; Pleger, S. T.; Reppel, M.; Borries, M.; Niroomand, F.; Pieske, B.; Janssen, P. M. L.; Eschenhagen, T.; Karczewski, P.; Smith, G. L.; Koch, W. J.; Katus, H. A.; Remppis, A.: S100A1: a regulator of myocardial contractility. *Proc. Nat. Acad. Sci.* 98: 13889–13894, 2001.
- [80623] 18953. Asada, Y.; Nadeau, J. H.: Fert is on mouse chromosome 11, not chromosome 17. *Mammalian Genome* 5: 830 only, 1994.
- [80624] 18954. Hao, Q.-L.; Heisterkamp, N.; Groffen, J.: Isolation and sequence analysis of a novel human tyrosine kinase gene. *Molec. Cell. Biol.* 9: 1587–1593, 1989.
- [80625] 18955. Dryja, T. P.; Grondin, V. J.; Ringens, P.; Cotran, P.; Berson, E. L.; Travis, G.: Isolation of human retinal cDNA fragments homologous to the murine rds gene transcript. (Abstract) *Invest. Ophthalmol. Vis. Sci.* 30 (suppl.): 43 only, 1989.
- [80626] 18956. Farrar, G. J.; Jordan, S. A.; Kenna, P.; Humphries, M. M.; Kumar-Singh, R.; McWilliam, P.; Allamand, V.; Sharp, E.; Humphries, P.: Autosomal dominant retinitis pigmentosa: localization of a disease gene (RP6) to the short arm of chromosome 6. *Genomics* 11: 870–874, 1991.
- [80627] 18957. Farrar, G. J.; Kenna, P.; Jordan, S. A.; Kumar-Singh,



R.; Humphries, M. M.; Sharp, E. M.; Sheils, D. M.;  
Humphries, P.: A three-base-pair deletion in the peripherin-RDS gene in one form of retinitis pigmentosa. *Nature* 354:478–480, 1991.

[80628] 18958. Feist, R. M.; White, M. F., Jr.; Skalka, H.; Stone, E. M.: Choroidal neovascularization in a patient with adult foveomacular dystrophy and a mutation in the retinal degeneration slow gene (pro210-to-arg). *Am. J. Ophthalmol.* 118: 259–260, 1994.

[80629] 18959. Felber, U.; Schilling, H.; Weber, B. H. F.: Adult vitelliform macular dystrophy is frequently associated with mutations in the peripherin/RDS gene. *Hum. Mutat.* 10: 301–309, 1997.

[80630] 18960. Gass, J. D. M.: A clinicopathologic study of a peculiar foveomacular dystrophy. *Trans. Am. Ophthalmol. Soc.* 72: 139–155, 1974.

[80631] 18961. Jackson, K. E.; Mitchell, E. B.; Stone, E. M.; Ferrell, R. E.; Gorin, M. B.: The identification of an exon-2 peripherin mutation in a family with heterogeneous manifestations of a butterfly pattern macular dystrophy. (Abstract) *Am. J. Hum. Genet.* 53 (suppl.): 1177 only, 1993.

[80632] 18962. Jordan, S. A.; Farrar, G. J.; Kumar-Singh, R.; Kenna, P.; Humphries, M. M.; Allamand, V.; Sharp, E. M.;

Humphries, P.: Autosomal dominant retinitis pigmentosa (adRP; RP6): cosegregation of RP6 and the peripherin-RD-S locus in a late-onset family of Irish origin. *Am. J. Hum. Genet.* 50:634–639, 1992.

[80633] 18963. Okamoto, H.; Yonemori, F.; Wakitani, K.; Minowa, T.; Maeda, K.; Shinkai, H.: A cholesteryl ester transfer protein inhibitor attenuates atherosclerosis in rabbits. *Nature* 406: 203–207, 2000.

[80634] 18964. Oliveira, H. C. F.; Chouinard, R. A.; Agellon, L. B.; Bruce, C.; Ma, L.; Walsh, A.; Breslow, J. L.; Tall, A. R.: Human cholesteryl ester transfer protein gene proximal promoter contains dietary cholesterol positive responsive elements and mediates expression in small intestine and periphery while predominant liver and spleen expression is controlled by 5-prime-distal sequences. *J. Biol. Chem.* 271: 31831–31838, 1996.

[80635] 18965. Sakai, N.; Yamashita, S.; Hirano, K.; Menju, M.; Arai, T.; Kobayashi, K.; Ishigami, M.; Yoshida, Y.; Hoshino, T.; Nakajima, N.; Kameda-Takemura, K.; Matsuzawa, Y.: Frequency of exon 15 missense mutation (442D:G) in cholesteryl ester transfer protein gene in hyperlipoproteinemic Japanese subjects. *Atherosclerosis* 114: 139–145, 1995.

- [80636] 18966. Sparkes, R. S.; Drayna, D.; Mohandas, T.; Klisak, I.; Heinzmann, C.; Lawn, R.; Lusis, A. J.: Assignment of cholesterol ester transfer protein (CETP) gene to human 16q21. (Abstract) Cytogenet. Cell Genet. 46:696 only, 1987.
- [80637] 18967. Takahashi, K.; Jiang, X.-C.; Sakai, N.; Yamashita, S.; Hirano, K.; Bujo, H.; Yamazaki, H.; Kusunoki, J.; Miura, T.; Kussie, P.; Matsuzawa, Y.; Saito, Y.; Tall, A.: A missense mutation in the cholesteryl ester transfer protein gene with possible dominant effects on plasma high density lipoproteins. J. Clin. Invest. 92: 2060–2064, 1993.
- [80638] 18968. Yamashita, S.; Hui, D. Y.; Sprecher, D. L.; Matsuzawa, Y.; Sakai, N.; Tarui, S.; Kaplan, D.; Wetterau, J. R.; Harmony, J. A.: Total deficiency of plasma cholesteryl ester transfer protein in subjects homozygous and heterozygous for the intron 14 splicing defect. Biochem. Biophys. Res. Commun. 170: 1346–1351, 1990.
- [80639] 18969. Zhong, S.; Sharp, D. S.; Grove, J. S.; Bruce, C.; Yano, K.; Curb, J. D.; Tall, A. R.: Increased coronary heart disease in Japanese–American men with mutation in the cholesteryl ester transfer protein gene despite increased HDL levels. J. Clin. Invest. 97: 2917–2923, 1996.
- [80640] 18970. Chung, B.-C.; Matteson, K. J.; Voutilainen, R.; Mo-

handas, T. K.; Miller, W. L.: Human cholesterol side-chain cleavage enzyme, P450scc:cDNA cloning, assignment of the gene to chromosome 15, and expression in the placenta. *Proc. Nat. Acad. Sci.* 83: 8962–8966, 1986.

[80641] 18971. Durocher, F.; Morissette, J.; Simard, J.: Genetic linkage mapping of the CYP11A1 gene encoding the cholesterol side-chain cleavage P450(scc) close to the CYP1A1 gene and D15S204 in the chromosome 15q22.33–q23 region. *Pharmacogenetics* 8: 49–53, 1998.

[80642] 18972. Billingsley, G. D.; Walter, M. A.; Hammond, G. L.; Cox, D. W.: Physical mapping of four serpin genes: alpha-1-antitrypsin, alpha-1-antichymotrypsin, corticosteroid-binding globulin, and protein C inhibitor, within a 280-kb region on chromosome 14q32.1. *Am. J. Hum. Genet.* 52: 343–353, 1993.

[80643] 18973. Wang, J.; Hannon, G. J.; Beach, D. H.: Risky immortalization by telomerase. (Letter) *Nature* 405: 755–756, 2000.

[80644] 18974. Wu, K.-J.; Grandori, C.; Amacker, M.; Simon-Vermot, N.; Polack, A.; Lingner, J.; Dalla-Favera, R.: Direct activation of TERT transcription by c-MYC. *Nature Genet.* 21: 220–224, 1999.

[80645] 18975. Greco, A.; Ittmann, M.; Barletta, C.; Basilico, C.;

Croce, C. M.; Cannizzaro, L. A.; Huebner, K.: Chromosomal localization of humangen es required for G(1) progression in mammalian cells. *Genomics* 4:240–245, 1989.

[80646] 18976. Ittmann, M.; Greco, A.; Basilico, C.: Isolation of the human gene that complements a temperature-sensitive cell cycle mutation in BHK cells. *Molec. Cell. Biol.* 7: 3386–3393, 1987.

[80647] 18977. Ip, N. Y.; Stitt, T. N.; Tapley, P.; Klein, R.; Glass, D. J.; Fandl, J.; Greene, L. A.; Barbacid, M.; Yancopoulos, G. D.: Similarities and differences in the way neurotrophins interact with the Trk receptors in neuronal and nonneuronal cells. *Neuron* 10: 137–149, 1993.

[80648] 18978. Knezevich, S. R.; McFadden, D. E.; Tao, W.; Lim, J. F.; Sorensen, P. H. B.: A novel ETV6–NTRK3 gene fusion in congenital fibrosarcoma. *Nature Genet.* 18: 184–187, 1998.

[80649] 18979. Bonnefont, J.–P.; Chretien, D.; Rustin, P.; Robinson, B.; Vassault, A.; Aupetit, J.; Charpentier, C.; Rabier, D.; Saudubray, J.–M.; Munnich, A.: Alpha-ketoglutarate dehydrogenase deficiency presenting as congenital lactic acidosis. *J. Pediat.* 121: 255–258, 1992.

[80650] 18980. Guffon, N.; Lopez–Mediavilla, C.; Dumoulin, R.; Mousson, B.; Godinot, C.; Carrier, H.; Collombet, J. M.; Di–

vry, P.; Mathieu, M.; Guibaud, P.: 2-Ketoglutarate dehydrogenase deficiency, a rare cause of primary hyperlactataemia: report of a new case. *J. Inherit. Metab. Dis.* 16:821–830, 1993.

[80651] 18981. Kohlschütter, A.; Behbehani, A.; Langenbeck, U.; Albani, M.; Heidemann, P.; Hoffmann, G.; Kleineke, J.; Lehnert, W.; Wendel, U.: A familial progressive neurodegenerative disease with 2-oxoglutaric aciduria. *Europ. J. Pediatr.* 138: 32–37, 1982.

[80652] 18982. Koike, K.: The gene encoding human 2-oxoglutarate dehydrogenase: structural organization and mapping to chromosome 7p13–p14. *Gene* 159:261–266, 1995.

[80653] 18983. Koike, K.; Urata, Y.; Goto, S.: Cloning and nucleotide sequence of the cDNA encoding human 2-oxoglutarate dehydrogenase (lipoamide). *Proc. Nat. Acad. Sci.* 89: 1963–1967, 1992.

[80654] 18984. Szabo, P.; Cai, X.; Ali, G.; Blass, J. P.: Localization of the gene (OGDH) coding for the E1 $\alpha$  component of the  $\alpha$ -ketoglutarate dehydrogenase complex to chromosome 7p13–p11.2. *Genomics* 20: 324–326, 1994.

[80655] 18985. Kury, S.; Dreno, B.; Bezieau, S.; Giraudet, S.; Kharfi, M.; Kamoun, R.; Moisan, J.-P.: Identification of SLC39A4, a

gene involved inacrodermatitis enteropathica. Nature Genet. 31: 239–240, 2002.

- [80656] 18986.Rogers, E. E.; Eide, D. J.; Guerinot, M. L.: Altered selectivityin an Arabidopsis metal transporter. Proc. Nat. Acad. Sci. 97: 12356–12360,2000.
- [80657] 18987.Wang, K.; Pugh, E. W.; Griffen, S.; Doheny, K. F.; Mostafa, W.Z.; al-Aboosi, M. M.; el-Shanti, H.; Gitschier, J.: Homozygositymapping places the acrodermatitis entero-pathica gene on chromosomalregion 8q24.3. Am. J. Hum. Genet. 68: 1055–1060, 2001.
- [80658] 18988.Haberle, J.; Pauli, S.; Linnebank, M.; Kleijer, W. J.; Bakker,H. D.; Wanders, R. J. A.; Harms, E.; Koch, H. G.: Structure of thehuman argininosuccinate synthetase gene and an improved system formolecular diagnostics in patients with classical and mild citrullinemia. Hum.Genet. 110: 327–333, 2002.
- [80659] 18989.Barbosa, J.; Rich, S.; Dunsworth, T.; Swanson, J.: Linkage disequilibriumbetween insulin-dependent diabetes and the Kidd blood group Jk(b)allele. J. Clin. Endocr. Metab. 55: 193–195, 1982.
- [80660] 18990.Harper, P. A.; Healy, P. J.; Dennis, J. A.; O'Brien, J. J.; Rayward,D. H.: Citrullinaemia as a cause of neurological disease in neonatalFriesian calves. Aust. Vet. J. 63:

378–379, 1986.

- [80661] 18991. Harper, P. A. W.; Healy, P. J.; Dennis, J. A.: Animal model of human disease: citrullinemia (argininosuccinate synthetase deficiency). *Am. J. Path.* 135: 1213–1215, 1989.
- [80662] 18992. Jackson, M. J.; Surh, L. C.; O'Brien, W. E.; Beaudet, A. L.: Assignment of the structural gene for argininosuccinate synthetase to proximal mouse chromosome 2. *Genomics* 6: 545–547, 1990.
- [80663] 18993. Kobayashi, K.; Ichiki, H.; Saheki, T.; Tatsuno, M.; Uchiyama, C.; Nukada, O.; Yoda, T.: Structure of an abnormal messenger RNA for argininosuccinate synthetase in citrullinemia. *Hum. Genet.* 76:27–32, 1987.
- [80664] 18994. Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.; Beaudet, A. L.: Characterization of nine mutant alleles causing citrullinemia. (Abstract) *Am. J. Hum. Genet.* 45 (suppl.): A201 only, 1989.
- [80665] 18995. Kobayashi, K.; Jackson, M. J.; Tick, D. B.; O'Brien, W. E.; Beaudet, A. L.: Heterogeneity of mutations in argininosuccinate synthetase causing human citrullinemia. *J. Biol. Chem.* 265: 11361–11367, 1990.
- [80666] 18996. Kobayashi, K.; Kakinoki, H.; Fukushige, T.; Shaheen, N.; Terazono, H.; Saheki, T.: Nature and frequency of



mutations in the argininosuccinatesynthetase gene that cause classical citrullinemia. Hum. Genet. 96:454–463, 1995.

- [80667] 18997.Kobayashi, K.; Rosenbloom, C.; Beaudet, A. L.; O'Brien, W. E.: Additional mutations in argininosuccinate synthetase causing citrullinemia. Molec.Biol. Med. 8: 95–100, 1991.
- [80668] 18998.Kobayashi, K.; Saheki, T.; Imamura, Y.; Noda, T.; Inoue, I.; Matuo,S.; Hagihara, S.; Nomiya, H.; Jinno, Y.; Shimada, K.: MessengerRNA coding for argininosuccinate synthetase in citrullinemia. Am.J. Hum. Genet. 38: 667–680, 1986.
- [80669] 18999.Kobayashi, K.; Shaheen, N.; Terazono, H.; Saheki, T.: Mutationsin argininosuccinate synthetase mRNA of Japanese patients, causingclassical citrullinemia. Am. J. Hum. Genet. 55: 1103–1112, 1994.
- [80670] 19000.Li, C.–M.; Chao, H.–K.; Liu, Y.–F.; Su, T.–S.: A non-sense mutationis responsible for the RNA-negative phenotype in human citrullinaemia. Europ.J. Hum. Genet. 9: 685–689, 2001.
- [80671] 19001.Northrup, H.; Lathrop, M.; Lu, S.–Y.; Daiger, S. P.; Beaudet,A. L.; O'Brien, W. E.: Multilocus linkage analysis with the humanargininosuccinate synthetase gene. Ge–

nomics 5: 442–444, 1989.

- [80672] 19002.Sase, M.; Kobayashi, K.; Imamura, Y.; Saheki, T.; Nakano, K.; Miura, S.; Mori, M.: Level of translatable messenger RNA coding for argininosuccinate synthetase in the liver of the patients with quantitative-type citrullinemia. Hum. Genet. 69: 130–134, 1985.
- [80673] 19003.Todd, S.; Naylor, S. L.: New chromosomal mapping assignments for argininosuccinate synthetase pseudogene 1, interferon- $\beta$ -3 gene, and the diazepam binding inhibitor gene. Somat. Cell Molec. Genet. 18:381–385, 1992.
- [80674] 19004.Rousseau-Merck, M. F.; Pizon, V.; Tavitian, A.; Berger, R.: Chromosome mapping of the human RAS related RAP1A, RAP1B and RAP2 genes to chromosomes 1p13–12, 12q14 and 13q34, respectively. (Abstract) Cytogenet. Cell Genet. 51: 1070 only, 1989.
- [80675] 19005.Rousseau-Merck, M. F.; Pizon, V.; Tavitian, A.; Berger, R.: Chromosome mapping of the human RAS-related RAP1A, RAP1B, and RAP2 genes to chromosomes 1p12–p13, 12q14, and 13q34, respectively. Cytogenet. Cell Genet. 53:2–4, 1990.
- [80676] 19006.Sebzda, E.; Bracke, M.; Tugal, T.; Hogg, N.; Cantrell, D. A.: Rap1a positively regulates T cells via inte-

grin activation rather than inhibiting lymphocyte signaling.  
Nature Immun. 3: 251–258, 2002.

- [80677] 19007. Takai, S.; Nishino, N.; Kitayama, H.; Ikawa, Y.; Noda, M.: Mapping of the KREV1 transformation suppressor gene and its pseudogene (KREV1P) to human chromosome 1p13.3 and 14q24.3, respectively, by fluorescence in situ hybridization. Cytogenet. Cell Genet. 63: 59–61, 1993.
- [80678] 19008. Zhu, J. J.; Qin, Y.; Zhao, M.; Van Aelst, L.; Malinow, R.: Ras and Rap control AMPA receptor trafficking during synaptic plasticity. Cell 110: 443–455, 2002.
- [80679] 19009. Ahmad, F.; Goldstein, B. J.: Functional association between the insulin receptor and the transmembrane protein-tyrosine phosphatase LAR in intact cells. J. Biol. Chem. 272: 448–457, 1997.
- [80680] 19010. Disteche, C. M.; Adler, D. A.; Tedder, T. F.; Saito, H.: Mapping of the genes for LYAM1, a new lymphocyte adhesion molecule, and for LAR, a new receptor-linked protein tyrosine phosphatase, to human chromosome 1 (Abstract) Cytogenet. Cell Genet. 51: 990 only, 1989.
- [80681] 19011. Harder, K. W.; Saw, J.; Miki, N.; Jirik, F.: Coexisting amplification of the chromosome 1p32 genes (PTPRF and MYCL1) encoding protein tyrosine phosphatase LAR and L-

myc in a small cell lung cancer line. Genomics  
27:552–553, 1995.

[80682] 19012.Jirik, F. R.; Harder, K. W.; Melhado, I. G.; Anderson, L. L.; Duncan, A. M. V.: The gene for leukocyte antigen-related tyrosine phosphatase(LAR) is localized to human chromosome 1p32, a region frequently deleted in tumors of neuroectodermal origin. Cytogenet. Cell Genet. 61:266–268, 1992.

[80683] 19013.Nam, H.-J.; Poy, F.; Krueger, N. X.; Saito, H.; Frederick, C. A.: Crystal structure of the tandem phosphatase domains of RPTP LAR. Cell 97:449–457, 1999.

[80684] 19014.O'Grady, P.; Krueger, N. X.; Streuli, M.; Saito, H.: Genomic organization of the human LAR protein tyrosine phosphatase gene and alternative splicing in the extracellular fibronectin type-III domains. J. Biol.Chem. 269: 25193–25199, 1994.

[80685] 19015.Schaapveld, R. Q.; Schepens, J. T.; Robinson, G. W.; Attema, J.; Oerlemans, F. T.; Fransen, J. A.; Streuli, M.; Wieringa, B.; Hennighausen, L.; Hendriks, W. J.: Impaired mammary gland development and function in mice lacking LAR receptor-like tyrosine phosphatase activity. Dev.Biol. 188: 134–146, 1997.

[80686] 19016.Schaapveld, R. Q. J.; van den Maagdenberg, A. M. J.

M.; Schepens, J. T. G.; Olde Weghuis, D.; Geurts van Kessel, A.; Wieringa, B.; Hendriks, W. J. A. J.: The mouse gene *Ptprf* encoding the leukocyte common antigen-related molecule LAR: cloning, characterization, and chromosomal localization. *Genomics* 27:124–130, 1995.

[80687] 19017. Tsujikawa, K.; Kawakami, N.; Uchino, Y.; Ichijo, T.; Furukawa, T.; Saito, H.; Yamamoto, H.: Distinct functions of the two protein tyrosine phosphatase domains of LAR (leukocyte common antigen-related) on tyrosine dephosphorylation of insulin receptor. *Molec. Endocr.* 15:271–280, 2001.

[80688] 19018. Ali, R. R.; Sarra, G.-M.; Stephens, C.; de Alwis, M.; Bainbridge, J. W. B.; Munro, P. M.; Fauser, S.; Reichell, M. B.; Kinnon, C.; Hunt, D. M.; Bhattacharya, S. S.; Thrasher, A. J.: Restoration of photoreceptor ultrastructure and function in retinal degeneration slow mice by gene therapy. *Nature Genet.* 25: 306–310, 2000.

[80689] 19019. Bascom, R. A.; Connell, G.; Garcia-Heras, J.; Collins, L.; Ledbetter, D.; Molday, R. S.; Kalnins, V.; McInnes, R. R.: Molecular and ultrastructural characterization of the products of the human retinopathy candidate genes ROM1 and RDS. (Abstract) *Am. J. Hum. Genet.* 47 (suppl.): A101 only, 1990.

- [80690] 19020.Connell, G.; Bascom, R.; Molday, L.; Reid, D.; McInnes, R. R.;Molday, R. S.: Photoreceptor peripherin is the normal product ofthe gene responsible for retinal de-generation in the rds mouse. *Proc.Nat. Acad. Sci.* 88: 723–726, 1991.
- [80691] 19021.Demant, P.; Ivanyi, D.; van Nie, R.: The map posi-tion of the rdsgene on the 17th chromosome of the mouse. *Tissue Antigens* 13: 53–55,1979.
- [80692] 19022.Makela, T. P.; Kere, J.; Winqvist, R.; Alitalo, K.: In-trachromosomalrearrangements fusing L-myc and rlf in small-cell lung cancer. *Molec.Cell. Biol.* 11: 4015–4021, 1991.
- [80693] 19023.Makela, T. P.; Saksela, K.; Evan, G.; Alitalo, K.: A fu-sion proteinformed by L-myc and a novel gene in SCLC. *EMBO J.* 10: 1331–1335,1991.
- [80694] 19024.Brody, L. C.; Abel, K. J.; Castilla, L. H.; Couch, F. J.; McKinley,D. R.; Yin, G.–Y.; Ho, P. P.; Merajver, S.; Chan-drasekharappa, S.C.; Xu, J.; Cole, J. L.; Struewing, J. P.; Valdes, J. M.; Collins,F. S.; Weber, B. L.: Construction of a transcription map surroundingthe BRCA1 locus of human chromosome 17. *Genomics* 25: 238–247, 1995.
- [80695] 19025.Hloch, P.; Schiedner, G.; Stahl, H.: Complete cDNA sequence ofthe human p68 protein. *Nucleic Acids Res.* 18:

3045, 1990.

- [80696] 19026.Iggo, R.; Gough, A.; Xu, W.; Lane, D. P.; Spurr, N. K.: Chromosomemapping of the human gene encoding the 68-kDa nuclear antigen (p68)by using the polymerase chain reaction. *Proc. Nat. Acad. Sci.* 86:6211–6214, 1989.
- [80697] 19027.Iggo, R. D.; Xu, W.; Lane, D. P.; Gough, A.; Spurr, N. K.: Assignmentof P68 to chromosome 17 using PCR. (Abstract) *Cytogenet. Cell Genet.* 51:1017, 1989.
- [80698] 19028.Acker, J.; Mattei, M.–G.; Wintzerith, M.; Roeckel, N.; Depetris,D.; Vigneron, M.; Kedingen, C.: Chromosomal localization of humanRNA polymerase II subunit genes. *Genomics* 20: 496–499, 1994.
- [80699] 19029.Bourquin, J.–P.; Stagljär, I.; Meier, P.; Moosmann, P.; Silke,J.; Baechi, T.; Georgiev, O.; Schaffner, W.: A serine/arginine-richnuclear matrix cyclophilin interacts with the C-terminal domain ofRNA polymerase II. *Nucleic Acids Res.* 25: 2055–2061, 1997.
- [80700] 19030.Schulze, A.; Hansen, C.; Skakkebaek, N. E.; Brøndum–Nielsen, K.;Ledbetter, D. H.; Tommerup, N.: Exclusion of SNRPN as a major determinantof Prader–Willi syndrome by a translocation breakpoint. *Nature Genet.* 12:452–454, 1996.
- [80701] 19031.Schweizer, J.; Zynger, D.; Francke, U.: In vivo nu–

clease hypersensitivity studies reveal multiple sites of parental origin-dependent differential chromatin conformation in the 150 kb SNRPN transcription unit.

Hum.Molec. Genet. 8: 555–566, 1999.

[80702] 19032.Shemer, R.; Hershko, A. Y.; Perk, J.; Mostoslavsky, R.; Tsuberi, B.; Cedar, H.; Buiting, K.; Razin, A.: The imprinting box of the Prader–Willi/Angelman syndrome domain. Nature Genet. 26: 440–443, 2000.

[80703] 19033.Sun, Y.; Nicholls, R. D.; Butler, M. G.; Saitoh, S.; Hainline, B. E.; Palmer, C. G.: Breakage in the SNRPN locus in a balanced 46,XY,t(15;19) Prader–Willi syndrome patient. Hum. Molec. Genet. 5: 517–524, 1996.

[80704] 19034.Sutcliffe, J. S.; Nakao, M.; Christian, S.; Orstavik, K. H.; Tommerup, N.; Ledbetter, D. H.; Beaudet, A. L.: Deletions of a differentially methylated CpG island at the SNRPN gene define a putative imprinting control region. Nature Genet. 8: 52–58, 1994.

[80705] 19035.Wirth, J.; Back, E.; Huttenhofer, A.; Nothwang, H.–G.; Lich, C.; Gross, S.; Menzel, C.; Schinzel, A.; Kioschis, P.; Tommerup, N.; Ropers, H.–H.; Horsthemke, B.; Buiting, K.: A translocation breakpoint cluster disrupts the newly defined 3–prime end of the SNURF–SNRPN transcription unit on chromosome 15. Hum. Molec. Genet. 10:



201–210, 2001.

- [80706] 19036.Stevanovic, M.; Zuffardi, O.; Collignon, J.; Lovell-Badge, R.;Goodfellow, P.: The cDNA sequence and chromosomal location of thehuman SOX2 gene. *Mammalian Genome* 5: 640–642, 1994.
- [80707] 19037.Critcher, R.; Stitson, R. N. M.; Wade-Martins, R.; Easty, D. J.;Farr, C. J.: Assignment of Sox4 to mouse chromosome 13 bands A3–A5by fluorescence in situ hybridization; refinement of the human SOX4location to 6p22.3 and of SOX20 to chromosome 17p12.3. *Cytogenet.Cell Genet.* 81: 294–295, 1998.
- [80708] 19038.Denny, P.; Swift, S.; Connor, F.; Ashworth, A.: An SRY-relatedgene expressed during spermatogenesis in the mouse encodes a sequence-specificDNA-binding protein. *EMBO J.* 11: 3705–3712, 1992.
- [80709] 19039.Farr, C. J.; Easty, D. J.; Ragoussis, J.; Collignon, J.; Lovell-Badge,R.; Goodfellow, P. N.: Characterization and mapping of the humanSOX4 gene. *Mammalian Genome* 4: 577–584, 1993.
- [80710] 19040.Hansen, G. M.; Skapura, D.; Justice, M. J.: Genetic profile ofinsertion mutations in mouse leukemias and lymphomas. *Genome Res.* 10:237–243, 2000.
- [80711] 19041.Li, J.; et al; et al: Leukaemia disease genes: large–

scale cloning and pathway predictions. *Nature Genet.* 23: 348–353, 1999.

[80712] 19042. Lund, A. H.; Turner, G.; Trubetskoy, A.; Verhoeven, E.; Wientjens, E.; Hulsman, D.; Russell, R.; DePinho, R. A.; Lenz, J.; van Lohuizen, M.: Genome-wide retroviral insertional tagging of genes involved in cancer in *Cdkn2a*-deficient mice. *Nature Genet.* 32: 160–165, 2002.

[80713] 19043. Suzuki, T.; Shen, H.; Akagi, K.; Morse, H. C., III; Malley, J. D.; Naiman, D. Q.; Jenkins, N. A.; Copeland, N. G.: New genes involved in cancer identified by retroviral tagging. *Nature Genet.* 32: 166–174, 2002. Note: Erratum: *Nature Genet.* 32: 331 only, 2002.

[80714] 19044. van de Wetering, M.; Oosterwegel, M.; van Norren, K.; Clevers, H.: Sox-4, an Sry-like HMG box protein, is a transcriptional activator in lymphocytes. *EMBO J.* 12: 3847–3854, 1993.

[80715] 19045. Shoyab, M.; McDonald, V. L.; Bradley, J. G.; Todaro, G. J.: Amphiregulin: a bifunctional growth-modulating glycoprotein produced by the phorbol 12-myristate 13-acetate-treated human breast adenocarcinoma cell line MCF-7. *Proc. Nat. Acad. Sci.* 85: 6528–6532, 1988.

[80716] 19046. Shoyab, M.; Plowman, G. D.; McDonald, V. L.; Bradley, J. G.; Todaro, G. J.: Structure and function of hu-

man amphiregulin: a member of the epidermal growth factor family. *Science* 243: 1074–1076, 1989.

[80717] 19047. Donahue, R. P.; Bias, W. B.; Renwick, J. H.; McKusick, V. A.: Probable assignment of the Duffy blood group locus to chromosome 1 in man. *Proc. Nat. Acad. Sci.* 61: 949–955, 1968.

[80718] 19048. Dracopoli, N. C.; Meisler, M. H.: Mapping the human amylase gene cluster on the proximal short arm of chromosome 1 using a highly informative (CA)<sub>n</sub> repeat. *Genomics* 7: 97–102, 1990.

[80719] 19049. Knoll, J. H. M.; Nicholls, R. D.; Magenis, R. E.; Glatt, K.; Graham, J. M., Jr.; Kaplan, L.; Lalande, M.: Angelman syndrome: three molecular classes identified with chromosome 15q11q13-specific DNA markers. *Am. J. Hum. Genet.* 47: 149–155, 1990.

[80720] 19050. Ono, H.; Kuno, Y.; Tanaka, H.; Yamashina, M.; Tsuyoshi, T.; Kondo, N.; Orii, T.: A case of paroxysmal nocturnal hemoglobinuria without deficiency of decay-accelerating factor on erythrocytes. *Blood* 75: 1746–1747, 1990.

[80721] 19051. Petranka, J. G.; Fleenor, D. E.; Sykes, K.; Kaufman, R. E.; Rosse, W. F.: Structure of the CD59-encoding gene: further evidence of a relationship to murine lymphocyte

antigen Ly-6 protein. Proc. Nat.Acad. Sci. 89: 7876–7879, 1992.

[80722] 19052.Rosse, W. F.: Personal Communication. Durham, N. C. 6/3/1993.

[80723] 19053.Rosse, W. F.; Parker, C. J.: Paroxysmal nocturnal hemoglobinuria. Clin.Haemat. 14: 105–125, 1985.

[80724] 19054.Rother, R. P.; Rollins, S. A.; Mennone, J.; Chodera, A.; Fidel,S. A.; Bessler, M.; Hillmen, P.; Squinto, S. P.: Expression of recombinanttransmembrane CD59 in paroxysmal nocturnal hemoglobinuria B cellsconfers resistance to human complement. Blood 84: 2604–2611, 1994.

[80725] 19055.Tone, M.; Walsh, L. A.; Waldmann, H.: Gene structure of humanCD59 and demonstration that discrete mRNAs are generated by alternativepolyadenylation. J. Molec. Biol. 227: 971–976, 1992.

[80726] 19056.Walsh, L. A.; Tone, M.; Thiru, S.; Waldmann, H.: The CD59 antigen--amultifunctional molecule. Tissue Antigens 40: 213–220, 1992.

[80727] 19057.Yamashina, M.; Ueda, E.; Kinoshita, T.; Takami, T.; Ojima, A.;Ono, H.; Tanaka, H.; Kondo, N.; Orii, T.; Okada, N.; Okada, H.; Inoue,K.; Kitani, T.: Inherited complete deficiency of 20-kilodalton homologousrestriction factor (CD59) as a cause of paroxysmal nocturnal hemoglobin-

uria. NewEng. J. Med. 323: 1184–1189, 1990.

[80728] 19058.Kumanogoh, A.; Watanabe, C.; Lee, I.; Wang, X.; Shi, W.; Araki,H.; Hirata, H.; Iwahori, K.; Uchida, J.; Yasui, T.; Matsumoto, M.;Yoshida, K.; Yakura, H.; Pan, C.; Parnes, J. R.; Kikutani, H.: Identification of CD72 as a lymphocyte receptor for the class IV semaphorin CD100:a novel mechanism for regulating B cell signaling. Immunity 13:621–631, 2000.

[80729] 19059.Von Hoegen, I.; Hsieh, C.–L.; Scharting, R.; Francke, U.; Parnes,J. R.: Identity of human Lyb–2 and CD72 and localization of the gene to chromosome 9. Europ. J. Immun. 21: 1425–1431, 1991.

[80730] 19060.Cambiaggi, C.; Scupoli, M. T.; Cestari, T.; Gerosa, F.; Carra,G.; Tridente, G.; Accolla, R. S.: Constitutive expression of CD69in interspecies T–cell hybrids and locus assignment to human chromosome12. Immunogenetics 36: 117–120, 1992.

[80731] 19061.Lopez–Cabrera, M.; Santis, A. G.; Fernandez–Ruiz, E.; Blacher,R.; Esch, F.; Sanchez–Mateos, P.; Sanchez–Madrid, F.: Molecular cloning,expression, and chromosomal localization of the human earliest lymphocyteactivation antigen AIM/CD69, a new member of the C–type animal lectinsuperfamily of signal–transmitting receptors. J.

Exp. Med. 178:537–547, 1993.

- [80732] 19062.Chandra, T.; Stackhouse, R.; Kidd, V. J.; Robson, K. J. H.; Woo, S. L. C.: Sequence homology between human alpha-1-antichymotrypsin, alpha-1-antitrypsin, and antithrombin III. *Biochemistry* 22: 5055–5061, 1983.
- [80733] 19063.Eriksson, S.; Lindmark, B.; Lilia, H.: Familial alpha-1-antichymotrypsin deficiency. *Acta Med. Scand.* 220: 447–453, 1986.
- [80734] 19064.Gilfix, B. M.; Briones, L.: Absence of the A1252G mutation in alpha 1-antichymotrypsin in a North American population suffering from dementia. *J. Cereb. Blood Flow Metab.* 17: 233–235, 1997.
- [80735] 19065.Haines, J. L.; Pritchard, M. L.; Saunders, A. M.; Schildkraut, J. M.; Growdon, J. H.; Gaskell, P. C.; Farrer, L. A.; Auerbach, S. A.; Gusella, J. F.; Locke, P. A.; Rosi, B. L.; Yamaoka, L.; Small, G. W.; Conneally, P. M.; Roses, A. D.; Pericak-Vance, M. A.: No genetic effect of alpha-1-antichymotrypsin in Alzheimer disease. *Genomics* 33:53–56, 1996.
- [80736] 19066.Haines, J. L.; Scott, W. K.; Pericak-Vance, M. A.: Reply to 'Genetic effect of alpha-1-antichymotrypsin on the risk of Alzheimer disease.' (Letter) *Genomics* 40: 384–385, 1997.

- [80737] 19067.Kamboh, M. I.; Aston, C. E.; Ferrell, R. E.; Dekosky, S. T.: Genetic effect of alpha-1-antichymotrypsin on the risk of Alzheimer disease.(Letter) *Genomics* 41: 382–385, 1997.
- [80738] 19068.Kamboh, M. I.; Sanghera, D. K.; Ferrell, R. E.; DeKosky, S. T.: APOE\*4-associated Alzheimer's disease risk is modified by alpha-1-antichymotrypsin polymorphism. *Nature Genet.* 10: 486–488, 1995.
- [80739] 19069.Kelsey, G. D.; Abeliovich, D.; McMahon, C. J.; Whitehouse, D.; Corney, G.; Povey, S.; Hopkinson, D. A.; Wolfe, J.; Mieli-Vergani, G.; Mowat, A. P.: Cloning of the human alpha-1 antichymotrypsin gene and genetic analysis of the gene in relation to alpha-1 antitrypsin deficiency. *J. Med. Genet.* 25: 361–368, 1988.
- [80740] 19070.Morgan, K.; Licastro, F.; Tilley, L.; Ritchie, A.; Morgan, L.; Pedrini, S.; Kalsheker, N.: Polymorphism in the alpha-1-antichymotrypsin(ACT) gene promoter: effect on expression in transfected glial and liver cell lines and plasma ACT concentrations. *Hum. Genet.* 109:303–310, 2001.
- [80741] 19071.Morgan, K.; Morgan, L.; Carpenter, K.; Lowe, J.; Lam, L.; Cave, S.; Xuereb, J.; Wischik, C.; Harrington, C.;

Kalsheker, N. A.: Microsatellite polymorphism of the alpha-1-antichymotrypsin gene locus associated with sporadic Alzheimer's disease. Hum. Genet. 99: 27-31, 1997.

[80742] 19072. Munoz, E.; Obach, V.; Oliva, R.; Marti, M. J.; Ezquerro, M.; Pastor, P.; Ballesta, F.; Tolosa, E.: Alpha-1-antichymotrypsin gene polymorphism and susceptibility to Parkinson's disease. Neurology 52: 297-301, 1999.

[80743] 19073. Poller, W.; Faber, J.-P.; Scholz, S.; Weidinger, S.; Bartholome, K.; Olek, K.; Eriksson, S.: Mis-sense mutation of alpha-1-antichymotrypsin gene associated with chronic lung disease. (Letter) Lancet 339: 1538, 1992.

[80744] 19074. Poller, W.; Faber, J.-P.; Weidinger, S.; Tief, K.; Scholz, S.; Fischer, M.; Olek, K.; Kirchgesser, M.; Heidtmann, H.-H.: A leucine-to-proline substitution causes a defective alpha-1-antichymotrypsin allele associated with familial obstructive lung disease. Genomics 17: 740-743, 1993.

[80745] 19075. Rabin, M.; Watson, M.; Breg, W. R.; Kidd, V.; Woo, S. L. C.; Ruddle, F. H.: Human alpha-1-antichymotrypsin and alpha-1-antitrypsin (PI) genes map to the same region on chromosome 14. (Abstract) Cytogenet. Cell Genet. 40: 728, 1985.

[80746] 19076. Rabin, M.; Watson, M.; Kidd, V.; Woo, S. L. C.; Breg,



W. R.; Ruddle, F. H.: Regional location of alpha-1-antichymotrypsin and alpha-1-antitrypsin genes on human chromosome 14. *Somat. Cell Molec. Genet.* 12: 209–214, 1986.

[80747] 19077. Han, P.; Fletcher, C. F.; Copeland, N. G.; Jenkins, N. A.; Yaremko, L. M.; Michaeli, T.: Assignment of the mouse Pde7A gene to the proximal region of chromosome 3 and of the human PDE7A gene to chromosome 8q13. *Genomics* 48:275–276, 1998.

[80748] 19078. Michaeli, T.; Bloom, T. J.; Martins, T.; Loughney, K.; Ferguson, K.; Riggs, M.; Rodgers, L.; Beavo, J. A.; Wigler, M.: Isolation and characterization of a previously undetected human cAMP phosphodiesterase by complementation of cAMP phosphodiesterase-deficient *Saccharomyces cerevisiae*. *J. Biol. Chem.* 268: 12925–12932, 1993.

[80749] 19079. Milatovich, A.; Bolger, G.; Michaeli, T.; Francke, U.: Chromosomal localizations of genes for five cAMP-specific phosphodiesterases in man and mouse. *Somat. Cell Molec. Genet.* 20: 75–86, 1994.

[80750] 19080. Loughney, K.; Martins, T. J.; Harris, E. A. S.; Sadhu, K.; Hicks, J. B.; Sonnenburg, W. K.; Beavo, J. A.; Ferguson, K.: Isolation and characterization of cDNAs corresponding

to two human calcium, calmodulin-regulated, 3'-prime, 5'-prime-cyclic nucleotide phosphodiesterases. *J. Biol. Chem.* 271:796-806, 1996.

[80751] 19081. Wilson, D. E.; McKenna, L.: Assignment of the human gene for phosphodiesterase 1A to chromosome 4. (Abstract) *Am. J. Hum. Genet.* 43: A162 only, 1988.

[80752] 19082. Ritter, J. K.; Chen, F.; Sheen, Y. Y.; Tran, H. M.; Kimura, S.; Yeatman, M. T.; Owens, I. S.: A novel complex locus UGT1 encodes human bilirubin, phenol, and other UDP-glucuronosyltransferase isozymes with identical carboxyl termini. *J. Biol. Chem.* 267: 3257-3261, 1992.

[80753] 19083. Tukey, R. H.; Strassburg, C. P.: Human UDP-glucuronosyltransferases: metabolism, expression, and disease. *Annu. Rev. Pharm. Toxicol.* 40:581-616, 2000.

[80754] 19084. James, P. F.; Grupp, I. L.; Grupp, G.; Woo, A. L.; Askew, G. R.; Croyle, M. L.; Walsh, R. A.; Lingrel, J. B.: Identification of a specific role for the Na,K-ATPase alpha-2 isoform as a regulator of calcium in the heart. *Molec. Cell* 3: 555-563, 1999.

[80755] 19085. Kawakami, K.; Ohta, T.; Nojima, H.; Nagano, K.: Primary structure of the alpha-subunit of human Na,K-ATPase deduced from cDNA sequence. *J. Biochem.* 100: 389-397, 1986.

- [80756] 19086.Kent, R. B.; Fallows, D. A.; Geissler, E.; Glaser, T.; Emanuel, J. R.; Lalley, P. A.; Levenson, R.; Housman, D. E.: Genes encoding alpha and beta subunits of Na,K-ATPase are located on three different chromosomes in the mouse. Proc. Nat. Acad. Sci. 84: 5369–5373, 1987.
- [80757] 19087.Ruiz-Opazo, N.; Barany, F.; Hirayama, K.; Herrera, V. L. M.: Confirmation of mutant alpha-1 Na,K-ATPase gene and transcript in Dahl salt-sensitive/JR rats. Hypertension 24: 260–270, 1994.
- [80758] 19088.Ovchinnikov, Y. A.; Monastyrskaya, G. S.; Broude, N. E.; Allikmets, R. L.; Ushkaryov, Y. A.; Melkov, A. M.; Smirnov, Y. V.; Malyshev, I. V.; Dulubova, I. E.; Petrukhin, K. E.; Gryshin, A. V.; Sverdlov, V. E.; Kiyatkin, N. I.; Kostina, M. B.; Modyanov, N. N.; Sverdlov, E. D.: The family of human Na<sup>+</sup>,K<sup>+</sup>-ATPase genes: a partial nucleotide sequence related to the alpha-subunit. FEBS Lett. 213: 73–80, 1987.
- [80759] 19089.Shull, M. M.; Lingrel, J. B.: Multiple genes encode the human Na<sup>+</sup>,K<sup>+</sup>-ATPase catalytic subunit. Proc. Nat. Acad. Sci. 84: 4039–4043, 1987.
- [80760] 19090.Sverdlov, E. D.; Broude, N. E.; Sverdlov, V. E.; Monastyrskaya, G. S.; Grishin, A. V.; Petrukhin, K. E.; Akopyanz, N. S.; Modyanov, N. N.; Ovchinnikov, Y. A.: Family of Na<sup>+</sup>,K<sup>+</sup>-ATPase genes: intra-individ-

ual tissue-specific restriction fragment length polymorphism. FEBS Lett. 221:129–133, 1987.

- [80761] 19091. Yang-Feng, T. L.; Schneider, J. W.; Lindgren, V.; Shull, M. M.; Benz, E. J., Jr.; Lingrel, J. B.; Francke, U.: Chromosomal localization of human Na<sup>+</sup>,K<sup>+</sup>-ATPase alpha- and beta-subunit genes. Genomics 2:128–138, 1988.
- [80762] 19092. Gloor, S.; Antonicek, H.; Sweadner, K. J.; Pagliusi, S.; Frank, R.; Moos, M.; Schachner, M.: The adhesion molecule on glia (AMOG) is a homologue of the beta subunit of the Na,K-ATPase. J. Cell Biol. 110:165–174, 1990.
- [80763] 19093. Hsieh, C.-L.; Cheng-Deutsch, A.; Gloor, S.; Schachner, M.; Francke, U.: Assignment of Amog (adhesion molecule on glia) gene to mouse chromosome 11 near Zfp-3 and Asgr-1,2 and to human chromosome 17. Somat. Cell Molec. Genet. 16: 401–405, 1990.
- [80764] 19094. Malo, D.; Schurr, E.; Levenson, R.; Gros, P.: Assignment of Na,K-ATPase beta-(2)-subunit gene (Atpb-2) to mouse chromosome 11. Genomics 6:697–699, 1990.
- [80765] 19095. Martin-Vasallo, P.; Dackowski, P.; Emanuel, J. R.; Levenson, R.: Identification of a putative isoform of the Na,K-ATPase beta subunit: primary structure and tissue-specific expression. J. Biol. Chem. 164:4613–4618, 1989.
- [80766] 19096. Pagliusi, S.; Antonicek, H.; Gloor, S.; Frank, R.;

Moos, M.; Schachner, M.: Identification of a cDNA clone specific for the neural adhesion molecule AMOG. *J. Neurosci. Res.* 22: 113–119, 1989.

[80767] 19097. Gallango, M. L.; Muller, A.; Suinaga, R.: Biochemical characterization of a red cell UMP kinase variant found in the Warao Indians of Venezuela. *Biochem. Genet.* 16: 1085–1093, 1978.

[80768] 19098. Gallango, M. L.; Suinaga, R.: Uridine monophosphate kinase polymorphism in two Venezuelan populations. *Am. J. Hum. Genet.* 30: 215–218, 1978.

[80769] 19099. Giblett, E. R.; Anderson, J. E.; Chen, S.-H.; Teng, Y.-S.; Cohen, F.: Uridine monophosphate kinase: a new genetic polymorphism with possible clinical implications. *Am. J. Hum. Genet.* 26: 627–635, 1974.

[80770] 19100. Giblett, E. R.; Anderson, J. E.; Lewis, M.; Kaita, H.: A new polymorphic enzyme, uridine monophosphate kinase: gene frequencies and a linkage analysis. *Birth Defects Orig. Art. Ser.* 11(3): 159–161, 1975. Note: Alternate: *Cytogenet. Cell Genet.* 14: 329–331, 1975.

[80771] 19101. Medrano, L.; Green, H.: A uridine kinase deficient mutant of 3T3 and a selective method for cells containing the enzyme. *Cell* 1: 23–26, 1974.

[80772] 19102. Petersen, G. M.; Silimperi, D. R.; Scott, E. M.; Hall,

D. B.; Rotter, J. I.; Ward, J. I.: Uridine monophosphate kinase 3: a genetic marker for susceptibility to Haemophilus influenzae type B disease. Lancet II:417–418, 1985.

[80773] 19103. Ranzani, G.; Bertolotti, E.; Santachiara-Benerecetti, A. S.: The polymorphism of the red cell uridine monophosphate kinase in two samples of the Italian population.

Hum. Hered. 27: 332–335, 1977.

[80774] 19104. Ruddle, F. H.; Creagan, R. P.: Parasexual approaches to the genetics of man. Ann. Rev. Genet. 9: 407–486, 1975.

[80775] 19105. Satlin, A.; Kucherlapati, R. S.; Ruddle, F. H.: Assignment of the gene for human uridine monophosphate kinase to chromosome 1 using somatic cell hybrid clone panels. Cytogenet. Cell Genet. 15: 146–152, 1975.

[80776] 19106. Anderson, J. A.; Teng, Y.-S.; Giblett, E. R.: Stains for six enzymes potentially applicable to chromosomal assignment by cell hybridization. Cytogenet. Cell Genet. 14: 295–299, 1975.

[80777] 19107. Hoglund, L.; Reichard, P.: J. Biol. Chem. 265: 6589–6595, 1990.

[80778] 19108. Paglia, D. E.; Valentine, W. N.; Brockway, R. A.: Identification of thymidine nucleotidase and deoxyribonucleotidase activities among normal isozymes of

5-prime-nucleotidase in human erythrocytes. *Proc.Nat. Acad. Sci.* 81: 588–592, 1984.

[80779] 19109.Rampazzo, C.; Gallinaro, L.; Milanesi, E.; Frigimelica, E.; Reichard,P.; Bianchi, V.: A deoxyribonucleotidase in mitochondria: involvement in regulation of dNTP pools and possible link to genetic disease. *Proc.Nat. Acad. Sci.* 97: 8239–8244, 2000.

[80780] 19110.Rampazzo, C.; Johansson, M.; Gallinaro, L.; Ferraro, P.; Hellman,U.; Karlsson, A.; Reichard, P.; Bianchi, V.: Mammalian 5-prime(3-prime)-deoxyribonucleotidase, cDNA cloning, and overexpression of the enzyme in *Escherichia coli* and mammalian cells. *J. Biol. Chem.* 275: 5409–5415,2000.

[80781] 19111.Swallow, D. M.; Turner, V. S.; Hopkinson, D. A.: Isozymes of rodent 5-prime-nucleotidase: evidence for two independent structural loci UMPH-1 and UMPH-2. *Ann. Hum. Genet.* 47: 9–17, 1983.

[80782] 19112.Tjernshaugen, H.; Fritzson, P. :*Biochem. J.* 154: 77–80, 1976.

[80783] 19113.Wilson, D. E.; Swallow, D. M.; Povey, S.: Assignment of the human gene for uridine 5-prime-monophosphate phosphohydrolase (UMPH2) to the long arm of chromosome 17. *Ann. Hum. Genet.* 50: 223–227, 1986.

- [80784] 19114.Wilson, D. E.; Woodard, D.; Sandler, A.; Erickson, J.; Gurney,A.: Provisional assignment of the gene for uridine monophosphatase-2(Umph-2) to mouse chromosome 11. *Biochem. Genet.* 25: 1-6, 1987.
- [80785] 19115.Xiong, J.-P.; Stehle, T.; Diefenbach, B.; Zhang, R.; Dunker, R.;Scott, D. L.; Joachimiak, A.; Goodman, S. L.; Arnaout, M. A.: Crystalstructure of the extracellular segment of integrin alpha-V-beta-3. *Science* 294:339-345, 2001.
- [80786] 19116.Xiong, J.-P.; Stehle, T.; Zhang, R.; Joachimiak, A.; Frech, M.;Goodman, S. L.; Arnaout, M. A.: Crystal structure of the extracellularsegment of integrin alpha-V-beta-3 in complex with an Arg-Gly-Aspligand. *Science* 296: 151-155, 2002.
- [80787] 19117.Disteche, C. M.; Plowman, G. D.; Gronwald, R. G. K.; Kelly, J.;Bowen-Pope, D.; Adler, D. A.; Murray, J. C.: Mapping of the amphiregulinand the platelet-growth factor receptor alpha genes to the proximallong arm of chromosome 4. (Abstract) *Cytogenet. Cell Genet.* 51:990, 1989.
- [80788] 19118.Gronwald, R. G. K.; Adler, D. A.; Kelly, J. D.; Distech, C. M.;Bowen-Pope, D. F.: The human PDGF receptor alpha-subunit gene mapsto chromosome 4 in close proximity to c-kit. *Hum. Genet.* 85: 383-385,1990.



- [80789] 19119.Hol, F. A.; Geurds, M. P. A.; Chatkupt, S.; Shugart, Y. Y.; Balling,R.; Schrande–Stumpel, C. T. R. M.; Johnson, W. G.; Hamel, B. C. J.;Mariman, E. C. M.: PAX genes and human neural tube defects: an aminoacid substitution in PAX1 in a patient with spina bifida. J. Med.Genet. 8: 655–660, 1996.
- [80790] 19120.Hsieh, C.–L.; Navankasattusas, S.; Escobedo, J. A.; Williams, L.T.; Francke, U.: Chromosomal localization of the gene for AA–typeplatelet–derived growth factor receptor (PDGFRA) in humans and mice. Cytogenet.Cell Genet. 56: 160–163, 1991.
- [80791] 19121.Ikuno, Y.; Kazlauskas, A.: TGF–beta–1–dependent contraction offibroblasts is mediated by th PDGF–alpha receptor. Invest. Ophthal.Vis. Sci. 43: 41–46, 2002.
- [80792] 19122.Joosten, P. H. L. J.; Hol, F. A.; van Beersum, S. E. C.; Peters,H.; Hamel, B. C. J.; Afink, G. B.; van Zoelen, E. J. J.; Mariman,E. C. M.: Altered regulation of platelet–derived growth factor receptor–alphagene–transcription in vitro by spina bifida–associated mutant Pax1proteins. Proc. Nat. Acad. Sci. 95: 14459–14463, 1998.
- [80793] 19123.Sanke, T.; Bell, G. I.; Sample, C.; Rubenstein, A. H.; Steiner,D. F.: An islet amyloid peptide is derived from an 89–amino acidprecursor by proteolytic processing. J. Biol.

Chem. 263: 17243–17246,1988.

- [80794] 19124.Verchere, C. B.; D'Alessio, D. A.; Palmiter, R. D.; Weir, G. C.;Bonner–Weir, S.; Baskin, D. G.; Kahn, S. E.: Islet amyloid formationassociated with hyperglycemia in trans–genic mice with pancreatic betacell expression of human islet amyloid polypeptide. Proc. Nat. Acad.Sci. 93: 3492–3496, 1996.
- [80795] 19125.Westermarck, P.; Engstrom, U.; Johnson, K. H.; Westermarck, G. T.;Betsholtz, C.: Islet amyloid polypeptide: pinpointing amino acidresidues linked to amyloid fibril formation. Proc. Nat. Acad. Sci. 87:5036–5040, 1990.
- [80796] 19126.Lin, F.; Worman, H. J.: Structural organization of the human gene(LMNB1) encoding nuclear lamin B1. Genomics 27: 230–236, 1995.
- [80797] 19127.Maeno, H.; Sugimoto, K.; Nakajima, N.: Genomic structure of themouse gene (Lmnb1) encoding nuclear lamin B1. Genomics 30: 342–346,1995.
- [80798] 19128.Oklu, R.; Hesketh, R.: The latent transforming growth factor betabinding protein (LTBP) family. Biochem. J. 352: 601–610, 2000.
- [80799] 19129.Brodsky, G. L.; Muntoni, F.; Miocic, S.; Sinagra, G.; Sewry, C.;Mestroni, L.: Lamin A/C gene mutation associated with dilated cardiomyopathywith variable skeletal

muscle involvement. *Circulation* 101: 473–476,2000.

- [80800] 19130.Cao, H.; Hegele, R. A.: Nuclear lamin A/C R482Q mutation in Canadian kindreds with Dunnigan-type familial partial lipodystrophy. *Hum.Molec. Genet.* 9: 109–112, 2000.
- [80801] 19131.De Sandre-Giovannoli, A.; Chaouch, M.; Kozlov, S.; Vallat, J.–M.;Tazir, M.; Kassouri, N.; Szepietowski, P.; Hammadouche, T.; Vandenberghe,A.; Stewart, C. L.; Grid, D.; Levy, N.: Homozygous defects in LMNA,encoding lamin A/C nuclear-envelope proteins, cause autosomal recessive axonal neuropathy in human (Charcot–Marie–Tooth disorder type 2) and mouse. *Am. J. Hum. Genet.* 70: 726–736, 2002. Note: Erratum: *Am.J. Hum. Genet.* 70: 1075 only, 2002.
- [80802] 19132.Fatkin, D.; MacRae, C.; Sasaki, T.; Wolff, M. R.; Porcu, M.; Frenneaux,M.; Atherton, J.; Vidaillet, H. J., Jr.; Spudich, S.; De Girolami,U.; Seidman, J. G.; Seidman, C. E.: Missense mutations in the rod domain of the lamin A/C gene as causes of dilated cardiomyopathy and conduction-system disease. *New Eng. J. Med.* 341: 1715–1724, 1999.
- [80803] 19133.Fisher, D. Z.; Chaudhary, N.; Blobel, G.: cDNA sequencing of nuclear lamins A and C reveals primary and secondary structural homology to intermediate filament

proteins. Proc. Nat. Acad. Sci. 83: 6450–6454, 1986.

[80804] 19134.Flier, J. S.: Pushing the envelope on lipodystrophy. Nature Genet. 24:103–104, 2000.

[80805] 19135.Garg, A.; Vinaitheerthan, M.; Weatherall, P. T.; Bowcock, A. M.: Phenotypic heterogeneity in patients with familial partial lipodystrophy(Dunnigan variety) related to the site of missense mutations in laminA/C gene. J. Clin. Endocr. Metab. 86: 59–65, 2001.

[80806] 19136.Genschel, J.; Schmidt, H. H.–J.: Mutations in the LMNA gene encoding lamin A/C. Hum. Mutat. 16: 451–459, 2000.

[80807] 19137.Guilly, M. N.; Bensussan, A.; Bourge, J. F.; Bornens, M.; Courvalin, J. C.: A human T lymphoblastic cell line lacks lamins A and C. EMBOJ. 6: 3795–3799, 1987.

[80808] 19138.Hegele, R. A.; Cao, H.; Harris, S. B.; Zinman, B.; Hanley, A.J.; Anderson, C. M.: Genetic variation in LMNA modulates plasma leptin and indices of obesity in aboriginal Canadians. Physiol. Genomics 3:39–44, 2000.

[80809] 19139.Hegele, R. A.; Cao, H.; Huff, M. W.; Anderson, C. M.: LMNA R482Q mutation in partial lipodystrophy associated with reduced plasma leptin concentration. J. Clin. Endocr. Metab. 85: 3089–3093, 2000.

[80810] 19140.Hegele, R. A.; Huff, M. W.; Young, T. K.: Common

genomic variation in LMNA modulates indexes of obesity in Inuit. *J. Clin. Endocr. Metab.* 86:2747–2751, 2001.

- [80811] 19141. Krohne, G.; Benavente, R.: The nuclear lamins: a multigene family of proteins in evolution and differentiation. *Exp. Cell Res.* 162:1–10, 1986.
- [80812] 19142. Lebel, S.; Raymond, Y.: Lamin A is not synthesized as a larger precursor polypeptide. *Biochem. Biophys. Res. Commun.* 149: 417–423, 1987.
- [80813] 19143. Lin, F.; Worman, H. J.: Structural organization of the human gene encoding nuclear lamin A and nuclear lamin C. *J. Biol. Chem.* 268:16321–16326, 1993.
- [80814] 19144. Lloyd, D. J.; Trembath, R. C.; Shackleton, S.: A novel interaction between lamin A and SREBP1: implications for partial lipodystrophy and other laminopathies. *Hum. Molec. Genet.* 11: 769–777, 2002.
- [80815] 19145. McKeon, F. D.; Kirschner, M. W.; Caput, D.: Homologies in both primary and secondary structure between nuclear envelope and intermediate filament proteins. *Nature* 319: 463–468, 1986.
- [80816] 19146. Muchir, A.; Bonne, G.; van der Kooi, A. J.; van Meegen, M.; Baas, F.; Bolhuis, P. A.; de Visser, M.; Schwartz, K.: Identification of mutations in the gene encoding lamins A/C in autosomal dominant limb girdle muscular dystrophy

with atrioventricular conduction disturbances(LGMD1B).

Hum. Molec. Genet. 9: 1453–1459, 2000.

[80817] 19147.Novelli, G.; Muchir, A.; Sangiuolo, F.; Helbling-Leclerc, A.;D'Apice, M. R.; Massart, C.; Capon, F.; Sbraccia, P.; Federici, M.;Lauro, R.; Tudisco, C.; Pallotta, R.; Scarano, G.; Dallapiccola, B.;Merlini, L.; Bonne, G.: Mandibuloacral dysplasia is caused by a mutationin LMNA–encoding lamin A/C. Am. J. Hum. Genet. 71: 426–431, 2002.

[80818] 19148.Raffaele di Barletta, M.; Ricci, E.; Galluzzi, G.; Tonali, P.;Mora, M.; Morandi, L.; Romorini, A.; Voit, T.; Orstavik, K. H.; Merlini,L.; Trevisan, C.; Biancalana, V.; Housmanowa–Petrusewicz, I.; Bione,S.; Ricotti, R.; Schwartz, K.; Bonne, G.; Toniolo, D.: Differentmutations in the LMNA gene cause autosomal dominant and autosomalrecessive Emery–Dreifuss muscular dystrophy. Am. J. Hum. Genet. 66:1407–1412, 2000.

[80819] 19149.Schmidt, H. H.–J.; Genschel, J.; Baier, P.; Schmidt, M.; Ockenga,J.; Tietge, U. J. F.; Propsting, M.; Buttner, C.; Manns, M. P.; Lochs,H.; Brabant, G.: Dyslipemia in familial partial lipodystrophy causedby and R482W mutation in the LMNA gene. J. Clin. Endocr. Metab. 86:2289–2295, 2001.

[80820] 19150.Shackleton, S.; Lloyd, D. J.; Jackson, S. N. J.; Evans, R.; Niermeijer,M. F.; Singh, B. M.; Schmidt, H.; Brabant, G.;

Kumar, S.; Durrington, P. N.; Gregory, S.; O'Rahilly, S.; Trembath, R. C.: LMNA, encoding lamin A/C, is mutated in partial lipodystrophy. *Nature Genet.* 24:153–156, 2000.

[80821] 19151. Wydner, K. L.; McNeil, J. A.; Lin, F.; Worman, H. J.; Lawrence, J. B.: Chromosomal assignment of human nuclear envelope protein genes LMNA, LMNB1, and LBR by fluorescence in situ hybridization. *Genomics* 32:474–478, 1996.

[80822] 19152. Furukawa, K.; Hotta, Y.: cDNA cloning of a germ cell specific lamin B3 from mouse spermatocytes and analysis of its function by ectopic expression in somatic cells. *EMBO J.* 12: 97–106, 1993.

[80823] 19153. Furukawa, K.; Inagaki, H.; Hotta, Y.: Identification and cloning of an mRNA coding for a germ cell-specific A-type lamin in mice. *Exp. Cell Res.* 212: 426–430, 1994.

[80824] 19154. Garbarz, M.; Devaux, I.; Bournier, O.; Grandchamp, B.; Dhermy, D.: Protein 4.1 Lille, a novel mutation in the downstream initiation codon of protein 4.1 gene associated with heterozygous 4,1(–) hereditary elliptocytosis. *Hum. Mutat.* 5: 339–340, 1995.

[80825] 19155. Garbarz, M.; Dhermy, D.; Lecomte, M. C.; Feo, C.; Chaverot, I.; Galand, C.; Bournier, O.; Bertrand, O.; Boivin, P.: A variant of erythrocyte membrane skeletal pro-

tein band 4.1 associated with hereditary elliptocytosis.

Blood 64: 1006–1015, 1984.

- [80826] 19156. Geerdink, R. A.; Nijenhuis, L. E.; Huizinga, J.: Hereditary elliptocytosis: linkage data in man. Ann. Hum. Genet. 30: 363–378, 1967.
- [80827] 19157. Jensson, O.; Jonasson, T.; Olafsson, O.: Hereditary elliptocytosis in Iceland. Brit. J. Haemat. 13: 844–854, 1967.
- [80828] 19158. Kan, Y.-W.: Personal Communication. San Francisco, Calif. 2/28/1986.
- [80829] 19159. Kuroda, S.; Takeuchi, T.; Nagamori, H.: Data on the linkage between elliptocytosis and Rh blood type. Jpn. J. Hum. Genet. 5: 112–118, 1960.
- [80830] 19160. Lambert, S.; Conboy, J.; Zail, S.: A molecular study of heterozygous protein 4.1 deficiency in hereditary elliptocytosis. Blood 72: 1926–1929, 1988.
- [80831] 19161. Lambert, S.; Zail, S.: Partial deficiency of protein 4.1 in hereditary elliptocytosis. Am. J. Hemat. 26: 263–272, 1987.
- [80832] 19162. Lipton, E. L.: Elliptocytosis with hemolytic anemia: the effect of splenectomy. Pediatrics 15: 67–82, 1955.
- [80833] 19163. Lux, S. E.; Wolfe, L. C.: Inherited disorders of the red cell membrane skeleton. Pediat. Clin. N. Am. 27:



463–486, 1980.

- [80834] 19164. Marchesi, S. L.; Conboy, J.; Agre, P.; Letsinger, J. T.; Marchesi, V. T.; Speicher, D. W.; Mohandas, N.: Molecular analysis of insertion/deletion mutations in protein 4.1 in elliptocytosis. I. Biochemical identification of rearrangements in the spectrin/actin binding domain and functional characterizations. *J. Clin. Invest.* 86: 516–523, 1990.
- [80835] 19165. McGuire, M.; Agre, P.: Three distinct variants of protein 4.1 in Caucasian hereditary elliptocytosis. (Abstract) *Clin. Res.* 35:428A, 1987.
- [80836] 19166. McGuire, M.; Smith, B. L.; Agre, P.: Distinct variants of erythrocyte protein 4.1 inherited in linkage with elliptocytosis and Rh type in three white families. *Blood* 72: 287–293, 1988.
- [80837] 19167. Morle, L.; Garbarz, M.; Alloisio, N.; Girot, R.; Chaveroche, I.; Boivin, P.; Delaunay, J.: The characterization of protein 4.1 Presles, a shortened variant of RBC membrane protein 4.1. *Blood* 65: 1511–1517, 1985.
- [80838] 19168. Morle, L.; Pothier, B.; Alloisio, N.; Ducluzeau, M.-T.; Marques, S.; Olim, G.; Martins e Silva, J.; Feo, C.; Garbarz, M.; Chaveroche, I.; Boivin, P.; Delaunay, J.: Red cell membrane alteration involving protein 4.1 and protein 3 in a case of recessively inherited haemolytic anemia. *Eu-*

rop. J. Haemat. 38: 447–455, 1987.

- [80839] 19169.Morton, N. E.: The detection and estimation of linkage between the genes for elliptocytosis and the Rh blood type. Am. J. Hum. Genet. 8:80–96, 1956.
- [80840] 19170.Nielsen, J. A.; Strunk, K. W.: Homozygous hereditary elliptocytosis as the cause of haemolytic anemia in infancy. Scand. J. Haemat. 5:486–496, 1968.
- [80841] 19171.Parra, M.; Gascard, P.; Walensky, L. D.; Snyder, S. H.; Mohandas, N.; Conboy, J. G.: Cloning and characterization of 4.1G (EPB41L2), a new member of the skeletal protein 4.1 (EPB41) gene family. Genomics 49:298–306, 1998.
- [80842] 19172.Peters, J. C.; Rowland, M.; Israels, L. G.; Zipursky, A.: Erythrocyte sodium transport in hereditary elliptocytosis. Canad. J. Physiol. Pharm. 44: 817–827, 1966.
- [80843] 19173.Roberts, J. A. F.: Genetic linkage in man, with particular reference to the usefulness of very small bodies of data. Quart. J. Med. 14:27–33, 1945.
- [80844] 19174.Shi, Z.-T.; Afzal, V.; Coller, B.; Patel, D.; Chasis, J. A.; Parra, M.; Lee, G.; Paszty, C.; Stevens, M.; Walensky, L.; Peters, L. L.; Mohandas, N.; Rubin, E.; Conboy, J. G.: Protein 4.1R-deficient mice are viable but have erythroid membrane skeleton abnormalities. J. Clin. Invest. 103:

331–340, 1999.

- [80845] 19175.Takakuwa, Y.; Tchernia, G.; Rossi, M.; Benabadji, M.; Mohandas,N.: Restoration of normal membrane stability to unstable protein4.1-deficient erythrocyte membranes by incorporation of purified protein4.1. J. Clin. Invest. 78: 80–85, 1986.
- [80846] 19176.Tang, C.–J. C.; Tang, T. K.: Rapid localization of membrane skeletalprotein 4.1 (EL1) to human chromosome 1p33–p34.2 by nonradioactivein situ hybridization. Cytogenet. Cell Genet. 57: 119, 1991.
- [80847] 19177.Tang, T. K.; Leto, T. L.; Correas, I.; Alonso, M. A.; Marchesi,V. T.; Benz, E. J., Jr.: Selective expression of an erythroid-specificisoform of protein 4.1. Proc. Nat. Acad. Sci. 85: 3713–3717, 1988.
- [80848] 19178.Tchernia, G.; Mohandas, N.; Shohet, S. B.: Deficiency of skeletalmembrane protein band 4.1 in homozygous hereditary elliptocytosis:implications for erythrocyte membrane stability. J. Clin. Invest. 68:454–460, 1981.
- [80849] 19179.Tomaselli, M. B.; John, K. M.; Lux, S. E.: Elliptical erythrocytemembrane skeletons and heat-sensitive spectrin in hereditary elliptocytosis. Proc.Nat. Acad. Sci. 78: 1911–1915, 1981.
- [80850] 19180.Brands, J. H. G. M.; Maassen, J. A.; Van Hemert, F.

J.; Amons, R.; Moller, W.: The primary structure of the alpha subunit of human elongation factor 1: structural aspects of guanine-nucleotide-binding sites. *Europ. J. Biochem.* 155: 167-171, 1986.

[80851] 19181. Ditzel, H. J.; Masaki, Y.; Nielsen, H.; Farnes, L.; Burton, D. R.: Cloning and expression of a novel human antibody-antigen pair associated with Felty's syndrome. *Proc. Nat. Acad. Sci.* 97: 9234-9239, 2000.

[80852] 19182. Lund, A.; Knudsen, S. M.; Vissing, H.; Clark, B.; Tommerup, N.: Assignment of human elongation factor 1-alpha genes: EEF1A maps to chromosome 6q14 and EEF1A2 to 20q13.3. *Genomics* 36: 359-361, 1996.

[80853] 19183. Opdenakker, G.; Cabeza-Arvelaiz, Y.; Fiten, P.; Dijkmans, R.; Van Damme, J.; Volckaert, G.; Billiau, A.; Van Elsen, A.; Van der Schueren, B.; Van den Berghe, H.; Cassiman, J.-J.: Human elongation factor 1-alpha: a polymorphic and conserved multigene family with multiple chromosomal localizations. *Hum. Genet.* 75: 339-344, 1987.

[80854] 19184. Cohen, A. J.; Li, F. P.; Berg, S.; Marchetto, D. J.; Tsai, S.; Jacobs, S. C.; Brown, R. S.: Hereditary renal-cell carcinoma associated with chromosomal translocation. *New Eng. J. Med.* 301: 592-595, 1979.

[80855] 19185. Gemmill, R. M.; West, J. D.; Boldog, F.; Tanaka, N.;

Robinson, L. J.; Smith, D. I.; Li, F.; Drabkin, H. A.: The hereditary renal cell carcinoma 3;8 translocation fuses FHIT to a patched-related gene, TRC8. *Proc. Nat. Acad. Sci.* 95: 9572–9577, 1998.

[80856] 19186. Nakano, A.; Pulkkinen, L.; Murrell, D.; Rico, J.; Lucky, A. W.; Garzon, M.; Stevens, C. A.; Robertson, S.; Pfendner, E.; Uitto, J.: Epidermolysis bullosa with congenital pyloric atresia: novel mutations in the beta-4 integrin gene (ITGB4) and genotype/phenotype correlations. *Pediatr. Res.* 49: 618–626, 2001.

[80857] 19187. Pulkkinen, L.; Bruckner-Tuderman, L.; August, C.; Uitto, J.: Compound heterozygosity for missense mutation (L156P) and nonsense (R554X) mutations in the beta-4 integrin gene (ITGB4) underlies mild, nonlethal phenotype of epidermolysis bullosa with pyloric atresia. *Am. J. Path.* 152: 935–941, 1998.

[80858] 19188. Pulkkinen, L.; Rouan, F.; Bruckner-Tuderman, L.; Wallerstein, R.; Garzon, M.; Brown, T.; Smith, L.; Carter, W.; Uitto, J.: Novel ITGB4 mutations in lethal and nonlethal variants of epidermolysis bullosa with pyloric atresia: missense versus nonsense. *Am. J. Hum. Genet.* 63: 1376–1387, 1998.

[80859] 19189. Shaw, L. M.; Rabinovitz, I.; Wang, H. H.-F.; Toker,

A., Mercurio, A. M.: Activation of phosphoinositide 3-OH kinase by the alpha-6/beta-4 integrin promotes carcinoma invasion. *Cell* 91: 949–960, 1997.

[80860] 19190. Suzuki, S.; Naitoh, Y.: Amino acid sequence of a novel integrin beta-4 subunit and primary expression of the mRNA in epithelial cells. *EMBO J.* 9: 757–763, 1990.

[80861] 19191. Vidal, F.; Aberdam, D.; Miquel, C.; Christiano, A. M.; Pulkkinen, L.; Uitto, J.; Ortonne, J.-P.; Meneguzzi, G.: Integrin beta-4 mutations associated with junctional epidermolysis bullosa with pyloric atresia. *Nature Genet.* 10: 229–234, 1995.

[80862] 19192. Krissansen, G. W.; Yuan, Q.; Jenkins, D.; Jiang, W.-M.; Rooke, L.; Spurr, N. K.; Eccles, M.; Leung, E.; Watson, J. D.: Chromosomal locations of the genes coding for the integrin beta-6 and beta-7 subunits. *Immunogenetics* 35: 58–61, 1992.

[80863] 19193. Baker, E.; Sutherland, G. R.; Jiang, W.-M.; Yuan, Q.; Leung, E.; Watson, J. D.; Krissansen, G. W.: Mapping of the human integrin beta-7 gene (ITG-beta-7) to 12q13.13 by non-isotopic in situ hybridization. *Mammalian Genome* 2: 272–273, 1992.

[80864] 19194. Erle, D. J.; Ruegg, C.; Sheppard, D.; Pytela, R.: Complete amino acid sequence of an integrin beta subunit

(beta-7) identified in leukocytes. *J. Biol. Chem.* 266: 11009–11016, 1991.

[80865] 19195. Yuan, Q.; Kozak, C. A.; Jiang, W.; Hollander, D.; Watson, J. D.; Krissansen, G. W.: Genetic mapping of the gene coding for the integrin beta-7 subunit to the distal part of mouse chromosome 15. *Immunogenetics* 35:403–407, 1992.

[80866] 19196. Chany, C.; Vignal, M.; Couillin, P.; Van Cong, N.; Boue, J.; Boue, A.: Chromosomal localization of human genes governing the interferon-induced antiviral state. *Proc. Nat. Acad. Sci.* 72: 3129–3133, 1975.

[80867] 19197. Dithmar, S.; Rusciano, D.; Lynn, M. J.; Lawson, D. H.; Armstrong, C. A.; Grossniklaus, H. E.: Neoadjuvant interferon alfa-2b treatment in a murine model for metastatic ocular melanoma: a preliminary study. *Arch. Ophthalmol.* 118: 1085–1089, 2000.

[80868] 19198. Ezekowitz, R. A. B.; Mulliken, J. B.; Folkman, J.: Interferon alfa-2a therapy for life-threatening hemangiomas of infancy. *New Eng. J. Med.* 326: 1456–1463, 1992.

[80869] 19199. Hejny, C.; Sternberg, P., Jr.; Lawson, D. H.; Greiner, K.; Aaberg, T. M., Jr.: Retinopathy associated with high-dose interferon alfa-2b therapy. *Am. J. Ophthalmol.* 131: 782–787, 2001.

- [80870] 19200.Aplin, H. M.; Hirst, K. L.; Crosby, A. H.; Dixon, M. J.: Mapping of the human dentin matrix acidic phosphoprotein gene (DMP1) to the dentinogenesis imperfecta type II critical region at chromosome 4q21. *Genomics* 30:347–349, 1995.
- [80871] 19201.Ho, A. S. Y.; Liu, Y.; Khan, T. A.; Hsu, D.-H.; Bazan, J. F.; Moore, K. W.: A receptor for interleukin 10 is related to interferon receptors. *Proc. Nat. Acad. Sci.* 90: 11267–11271, 1993.
- [80872] 19202.Liu, Y.; Wei, S. H.-Y.; Ho, A. S.-Y.; de Waal Malefyt, R.; Moore, K. W.: Expression cloning and characterization of a human IL-10 receptor. *J. Immun.* 152: 1821–1829, 1994.
- [80873] 19203.Blixt, A.; Mahlapuu, M.; Aitola, M.; Peltö-Huikko, M.; Enerbäck, S.; Carlsson, P.: A forkhead gene, FoxE3, is essential for lens epithelial proliferation and closure of the lens vesicle. *Genes Dev.* 14: 245–254, 2000.
- [80874] 19204.Semina, E. V.; Brownell, I.; Mintz-Hittner, H. A.; Murray, J. C.; Jamrich, M.: Mutations in the human forkhead transcription factor FOXE3 associated with anterior segment ocular dysgenesis and cataracts. *Hum. Molec. Genet.* 10: 231–236, 2001.
- [80875] 19205.Siegelmann-Danieli, N.; Buetow, K. H.: Constitu-



tional genetic variation at the human aromatase gene (Cyp19) and breast cancer risk. *Brit. J. Cancer* 79: 456–463, 1999.

- [80876] 19206. Simpson, E. R.; Michael, M. D.; Agarwal, V. R.; Hinchelwood, M.M.; Bulun, S. E.; Zhao, Y.: Expression of the CYP19 (aromatase) gene: an unusual case of alternative promoter usage. *FASEB J.* 11: 29–36, 1997.
- [80877] 19207. Sparkes, R. S.; Mohandas, T.; Chen, S.; Besman, M. J.; Zollman, S.; Shively, J. E.: Assignment of the aromatase gene to human chromosome 15q21. (Abstract) *Cytogenet. Cell Genet.* 46: 696–697, 1987.
- [80878] 19208. Toda, K.; Merashima, M.; Kawamoto, T.; Sumimoto, H.; Yokoyama, Y.; Kuribayashi, I.; Mitsuuchi, Y.; Maeda, T.; Yamamoto, Y.; Sagara, Y.; Ikeda, H.; Shizuta, Y.: Structural and functional characterization of human aromatase P-450 gene. *Europ. J. Biochem.* 193: 559–565, 1990.
- [80879] 19209. Wang, Z. J.; Jeffs, B.; Ito, M.; Achermann, J. C.; Yu, R. N.; Hales, D. B.; Jameson, J. L.: Aromatase (Cyp19) expression is up-regulated by targeted disruption of Dax1. *Proc. Nat. Acad. Sci.* 98: 7988–7993, 2001.
- [80880] 19210. Whitlock, J. P., Jr.: The regulation of cytochrome P-450 gene expression. *Annu. Rev. Pharm. Toxicol.* 26: 333–369, 1986.

- [80881] 19211.Zhou, D.; Pompon, D.; Chen, S.: Structure–function studies of human aromatase by site–directed mutagenesis: kinetic properties of mutants pro308–to–phe, tyr361–to–phe, tyr361–to–leu, and phe406–to–arg. Proc.Nat. Acad. Sci. 88: 410–414, 1991.
- [80882] 19212.Bruneau, G.; Gross, M.–S.; Krieger, M.; Bernheim, A.; Thibault,J.; Nguyen, V. C.: Preparation of a human DOPA decarboxylase cDNA probe by PCR and its assignment to chromosome 7. Ann. Genet. 33:208–213, 1990.
- [80883] 19213.Christenson, J. G.; Dairman, W.; Udenfriend, S.: On the identity of DOPA decarboxylase and 5–hydroxytryptophan decarboxylase (immunological titration–aromatic L–amino acid decarboxylase–serotonin–dopamine–norepinephrine).Proc. Nat. Acad. Sci. 69: 343–347, 1972.
- [80884] 19214.Craig, S. P.; Le Van Thai, A.; Weber, M.; Craig, I. W.: Localisation of the gene for human aromatic L–amino acid decarboxylase (DDC) to chromosome 7p13–p11 by in situ hybridisation. Cytogenet. Cell Genet. 61:114–116, 1992.
- [80885] 19215.Ichinose, H.; Kurosawa, Y.; Titani, K.; Fujita, K.; Nagatsu, T.: Isolation and characterization of a cDNA clone encoding human aromatic L–amino acid decarboxylase. Biochem. Biophys. Res. Commun. 164:1024–1030, 1989.

- [80886] 19216.Scherer, L. J.; McPherson, J. D.; Wasmuth, J. J.; Marsh, J. L.: Human dopa decarboxylase: localization to human chromosome 7p11 and characterization of hepatic cDNAs. *Genomics* 13: 469–471, 1992.
- [80887] 19217.Sumi-Ichinose, C.; Ichinose, H.; Takahashi, E.; Hori, T.; Nagatsu, T.: Molecular cloning of genomic DNA and chromosomal assignment of the gene for human aromatic L-amino acid decarboxylase, the enzyme for catecholamine and serotonin biosynthesis. *Biochemistry* 31:2229–2238, 1992.
- [80888] 19218.Luttrell, L. M.; Ferguson, S. S. G.; Daaka, Y.; Miller, W. E.; Maudsley, S.; Della Rocca, G. J.; Lin, F.-T.; Kawakatsu, H.; Owada, K.; Luttrell, D. K.; Caron, M. G.; Lefkowitz, R. J.: Beta-arrestin-dependent formation of beta-2 adrenergic receptor-Src protein kinase complexes. *Science* 283:655–661, 1999.
- [80889] 19219.Attramadal, H.; Arriza, J. L.; Aoki, C.; Dawson, T. M.; Codina, J.; Kwatra, M. M.; Snyder, S. H.; Caron, M. G.; Lefkowitz, R. J.: Beta-arrestin-2, a novel member of the arrestin/beta-arrestin gene family. *J. Biol. Chem.* 267: 17882–17890, 1992.
- [80890] 19220.Bohn, L. M.; Gainetdinov, R. R.; Lin, F.-T.; Lefkowitz, R. J.; Caron, M. G.: Mu-opioid receptor desensi-

tization by beta-arrestin-2 determines morphine tolerance but not dependence. *Nature* 408: 720–723, 2000.

- [80891] 19221. Bohn, L. M.; Lefkowitz, R. J.; Gainetdinov, R. R.; Peppel, K.; Caron, M. G.; Lin, F.-T.: Enhanced morphine analgesia in mice lacking beta-arrestin 2. *Science* 286: 2495–2498, 1999.
- [80892] 19222. Fong, F. M.; Premont, R. T.; Richardson, R. M.; Yu, Y.-R. A.; Lefkowitz, R. J.; Patel, D. D.: Defective lymphocyte chemotaxis in beta-arrestin2- and GRK6-deficient mice. *Proc. Nat. Acad. Sci.* 99: 7478–7483, 2002.
- [80893] 19223. Kiselev, A.; Socolich, M.; Vinos, J.; Hardy, R. W.; Zuker, C. S.; Ranganathan, R.: A molecular pathway for light-dependent photoreceptor apoptosis in *Drosophila*. *Neuron* 28: 139–152, 2000.
- [80894] 19224. McDonald, P. H.; Chow, C.-W.; Miller, W. E.; Laporte, S. A.; Field, M. E.; Lin, F.-T.; Davis, R. J.; Lefkowitz, R. J.: Beta-arrestin2: a receptor-regulated MAPK scaffold for the activation of JNK3. *Science* 290: 1574–1577, 2000.
- [80895] 19225. Karathanasis, S. K.; Oettgen, P.; Haddad, I. A.; Antonarakis, S. E.: Structure, evolution, and polymorphisms of the human apolipoprotein A4 gene (APOA4). *Proc. Nat. Acad. Sci.* 83: 8457–8461, 1986.
- [80896] 19226. Carrington, M.; Kissner, T.; Gerrard, B.; Ivanov, S.;

O'Brien, S. J.; Dean, M.: Novel alleles of the chemokine-receptor gene CCR5. *Am. J. Hum. Genet.* 61: 1261–1267, 1997.

[80897] 19227. Chaganti, R. S. K.: Personal Communication. New York, N. Y. 10/22/1993.

[80898] 19228. Chaudhuri, A.; Polyakova, J.; Zbrzezna, V.; Pogo, A. O.: The coding sequence of Duffy blood group gene in humans and simians: restriction fragment length polymorphism, antibody and malarial parasite specificities, and expression in nonerythroid tissues in Duffy-negative individuals. *Blood* 85:615–621, 1995.

[80899] 19229. Chaudhuri, A.; Polyakova, J.; Zbrzezna, V.; Williams, K.; Gulati, S.; Pogo, A. O.: Cloning of glycoprotein D cDNA, which encodes the major subunit of the Duffy blood group system and the receptor for the *Plasmodium vivax* malaria parasite. *Proc. Nat. Acad. Sci.* 90:10793–10797, 1993.

[80900] 19230. Cook, P. J. L.; Page, B. M.; Johnston, A. W.; Stanford, W. K.; Gavin, J.: Four further families informative for 1q and the Duffy blood group. *Cytogenet. Cell Genet.* 22: 378–380, 1978.

[80901] 19231. Cook, P. J. L.; Robson, E. B.; Buckton, K. E.; Jacobs, P. A.; Polani, P. E.: Segregation of genetic markers in fami-

lies with chromosome polymorphisms and structural rearrangements involving chromosome no.1. Ann. Hum. Genet. 37: 261–274, 1974.

- [80902] 19232.Crawford, M. N.; Punnett, H. H.; Carpenter, G. G.: Deletion of the long arm of chromosome 16 and an unexpected Duffy blood group phenotype reveal a possible autosomal linkage. Nature 215: 1075–1076, 1967.
- [80903] 19233.Gelpi, A. P.; King, M. C.: Association of Duffy blood group with the sickle cell trait. Hum. Genet. 32: 65–68, 1976.
- [80904] 19234.Hadley, T. J.; David, P. H.; McGinniss, M. H.; Miller, L. H.: Identification of an erythrocyte component carrying the Duffy blood group Fy-a antigen. Science 223: 597–599, 1984.
- [80905] 19235.Hadley, T. J.; Peiper, S. C.: From malaria to chemokine receptor: the emerging physiologic role of the Duffy blood group antigen. Blood 89:3077–3091, 1997.
- [80906] 19236.Hamblin, M. T.; Di Rienzo, A.: Detection of the signature of natural selection in humans: evidence from the Duffy blood group locus. Am.J. Hum. Genet. 66: 1669–1679, 2000.
- [80907] 19237.Hamblin, M. T.; Thompson, E. E.; Di Rienzo, A.: Complex signatures of natural selection at the Duffy blood

group locus. Am. J. Hum. Genet. 70: 369–383, 2002.

[80908] 19238. Hessner, M. J.; Pircon, R. A.; Johnson, S. T.; Luhm, R. A.: Prenatal genotyping of the Duffy blood group system by allele-specific polymerase chain reaction. Prenatal Diag. 19: 41–45, 1999.

[80909] 19239. Horuk, R.; Chitnis, C. E.; Darbonne, W. C.; Colby, T. J.; Rybicki, A.; Hadley, T. J.; Miller, L. H.: A receptor for the malarial parasite *Plasmodium vivax*: the erythrocyte chemokine receptor. Science 261: 1182–1184, 1993.

[80910] 19240. Howard, P. N.; Stoddard, G. R.; Goddard, M. W.; Seely, J. R.: Giemsa banding of chromosome 1qh+ and linkage analysis. J. Med. Genet. 12: 44–48, 1975.

[80911] 19241. Iwamoto, S.; Li, J.; Sugimoto, N.; Okuda, H.; Kajii, E.: Characterization of the Duffy gene promoter: evidence for tissue-specific abolishment of expression in Fy(a–b–) of black individuals. Biochem. Biophys. Res. Commun. 222: 852–859, 1996.

[80912] 19242. Lautenberger, J. A.; Stephens, J. C.; O'Brien, S. J.; Smith, M. W.: Significant admixture linkage disequilibrium across 30 cM around the FY locus in African Americans. Am. J. Hum. Genet. 66: 969–978, 2000.

[80913] 19243. Lee, C. S. N.; Ying, K. L.; Bowen, P.: Position of the Duffy locus on chromosome 1 in relation to breakpoints

for structural rearrangements. Am.J. Hum. Genet. 26: 93–102, 1974.

[80914] 19244.Li, J.; Iwamoto, S.; Sugimoto, N.; Okuda, H.; Kajii, E.: Dinucleotiderepeat in the 3–prime flanking region provides a clue to the molecular evolution of the Duffy gene. Hum. Genet. 99: 573–577, 1997.

[80915] 19245.Braun, A.; Kammerer, S.; Bohme, E.; Muller, B.; Roscher, A. A.: Identification of polymorphic sites of the human bradykinin B(2)receptor gene. Biochem. Biophys. Res. Commun. 211: 234–240, 1995.

[80916] 19246.Erdmann, J.; Hegemann, N.; Weidemann, A.; Kallisch, H.; Hummel,M.; Hetzer, R.; Fleck, E.; Regitz–Zagrosek, V.: Screening the humanbradykinin B2 receptor gene in patients with cardiovascular diseases:identification of a functional mutation in the promoter and a newcoding variant (T21M). Am. J. Med. Genet. 80: 521–525, 1998.

[80917] 19247.Hess, J. F.; Borkowski, J. A.; Young, G. S.; Strader, C. D.; Ransom,R. W.: Cloning and pharmacological characterization of a human bradykinin(BK–2) receptor. Biochem. Biophys. Res. Commun. 184: 260–268, 1992.

[80918] 19248.Kammerer, S.; Braun, A.; Arnold, N.; Roscher, A. A.: The humanbradykinin B(2) receptor gene: full length



cDNA, genomic organization and identification of the regulatory region. *Biochem. Biophys. Res. Commun.* 211: 226–233, 1995.

[80919] 19249. Ma, J.; Wang, D.; Ward, D. C.; Chen, L.; Dessai, T.; Chao, J.; Chao, L.: Structure and chromosomal localization of the gene (BDKRB2) encoding human bradykinin B2 receptor. *Genomics* 23: 362–369, 1994.

[80920] 19250. Powell, S. J.; Slynn, G.; Thomas, C.; Hopkins, B.; Briggs, I.; Graham, A.: Human bradykinin B2 receptor: nucleotide sequence analysis and assignment to chromosome 14. *Genomics* 15: 435–438, 1993.

[80921] 19251. Taketo, M.; Yokoyama, S.; Rochelle, J.; Kimura, S.; Higashida, H.; Taketo, M.; Seldin, M. F.: Mouse B2 bradykinin receptor gene maps to distal chromosome 12. *Genomics* 27: 222–223, 1995.

[80922] 19252. Chen, H.; Weber, A. J.: BDNF enhances retinal ganglion cell survival in cats with optic nerve damage. *Invest. Ophthalm. Vis. Sci.* 42: 966–974, 2001.

[80923] 19253. Conover, J. C.; Erickson, J. T.; Katz, D. M.; Bianchi, L. M.; Poueymirou, W. T.; McClain, J.; Pan, L.; Helgren, M.; Ip, N. Y.; Boland, P.; Friedman, B.; Wiegand, S.; Vejsada, R.; Kato, A. C.; DeChiara, T. M.; Yancopoulos, G. D.: Neuronal deficits, not involving motor neurons, in mice lacking BDNF

and/or NT4. Nature 375: 235–238, 1995.

- [80924] 19254.Hanson, I. M.; Seawright, A.; van Heyningen, V.: The human BDNF gene maps between FSHB and HVBS1 at the boundary of 11p13–p14. Genomics 13:1331–1333, 1992.
- [80925] 19255.Hofer, M. M.; Barde, Y.–A.: Brain–derived neurotrophic factor prevents neuronal death in vivo. Nature 331: 261–262, 1988.
- [80926] 19256.Huang, Z. J.; Kirkwood, A.; Pizzorusso, T.; Porciatti, V.; Morales, B.; Bear, M. F.; Maffei, L.; Tonegawa, S.: BDNF regulates the maturation of inhibition and the critical period of plasticity in mouse visual cortex. Cell 98: 739–755, 1999.
- [80927] 19257.Kovalchuk, Y.; Hanse, E.; Kafitz, K. W.; Konnerth, A.: Postsynaptic induction of BDNF–mediated long–term potentiation. Science 295:1729–1734, 2002.
- [80928] 19258.Liu, X.; Ernfors, P.; Wu, H.; Jaenisch, R.: Sensory but not motor neuron deficits in mice lacking NT4 and BDNF. Nature 375: 238–241, 1995.
- [80929] 19259.Lu, Q.; Sun, E. E.; Klein, R. S.; Flanagan, J. G.: Ephrin–B reverse signaling is mediated by a novel PDZ–RGS protein and selectively inhibits G protein–coupled chemoattraction. Cell 105: 69–79, 2001.

- [80930] 19260.Masi, L.; Becherini, L.; Colli, E.; Gennari, L.; Mansani, R.; Falchetti,A.; Becorpi, A. M.; Cepollaro, C.; Gonnelli, S.; Tanini, A.; Brandi,M. L.: Polymorphisms of the calcitonin receptor gene are associatedwith bone mineral density in postmenopausal Italian women. *Biochem.Biophys. Res. Commun.* 248: 190–195, 1998.
- [80931] 19261.Masi, L.; Becherini, L.; Gennari, L.; Colli, E.; Mansani, R.; Falchetti,A.; Cepollaro, C.; Gonnelli, S.; Tanini, A.; Brandi, M. L.: Allelicvariants of human calci-  
tonin receptor: distribution and associationwith bone mass in postmenopausal Italian women. *Biochem. Bio-  
phys.Res. Commun.* 245: 622–626, 1998.
- [80932] 19262.Nakamura, M.; Zhang, Z. Q.; Shan, L.; Hisa, T.; Sasaki, M.; Tsukino,R.; Yokoi, T.; Kaname, A.; Kakudo, K.: Allelic variants of humancalcitonin receptor in the Japanese population. *Hum. Genet.* 99:38–41, 1997.
- [80933] 19263.Perez Jurado, L. A.; Li, X.; Francke, U.: The human calcitoninreceptor gene (CALCR) at 7q21.3 is outside the deletion associatedwith the Williams syndrome. *Cyto-  
genet. Cell Genet.* 70: 246–249,1995.
- [80934] 19264.Taboulet, J.; Frendo, J. L.; Delage–Murroux, R.; Pichaud, F.; deVernejoul, M. C.; Jullienne, A.: Evidence for 2 allelic forms ofcalcitonin receptor gene: distribution in

normal and osteoporotic women. (Abstract) J. Bone Miner. Res. 11 (suppl. 1): S204, 1996.

- [80935] 19265.Taboulet, J.; Frenkian, M.; Frendo, J. L.; Feingold, N.; Jullienne,A.; de Vernejoul, M. C.: Calcitonin receptor polymorphism is associated with a decreased fracture risk in post-menopausal women. Hum. Molec.Genet. 7: 2129–2133, 1998.
- [80936] 19266.Amara, S. G.; Arriza, J. L.; Leff, S. E.; Swanson, L. W.; Evans,R. M.; Rosenfeld, M. G.: Expression in brain of a messenger RNA encoding a novel neuropeptide homologous to calcitonin gene-related peptide.Science 229: 1094–1097, 1985.
- [80937] 19267.Hoppener, J. W. M.; Steenbergh, P. H.; Zandberg, J.; Geurts vanKessel, A. H. M.; Baylin, S. B.; Nelkin, B. D.; Jansz, H. S.; Lips,C. J. M.: The second human calcitonin/ CGRP gene is located on chromosome 11. Hum. Genet. 70: 259–263, 1985.
- [80938] 19268.Steenbergh, P. H.; Hoppener, J. W. M.; Zandberg, J.; Lips, C. J.M.; Jansz, H. S.: A second human calcitonin/ CGRP gene. FEBS Lett. 183:403–407, 1985.
- [80939] 19269.Arthur, J. S. C.; Elce, J. S.; Hegadorn, C.; Williams, K.; Greer,P. A.: Disruption of the murine calpain small subunit gene, Capn4:calpain is essential for embryonic

development but not for cell growth and division. *Molec. Cell. Biol.* 20: 4474–4481, 2000.

- [80940] 19270. Ohno, S.; Emori, Y.; Suzuki, K.: Nucleotide sequence of a cDNA coding for the small subunit of human calcium-dependent protease. *Nucleic Acids Res.* 14: 5559 only, 1986.
- [80941] 19271. Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Ohmi-Imajoh, S.; Suzuki, K.; Shimizu, Y.; Shimizu, N.: Four genes for the calpain family locate on four distinct human chromosomes. (Abstract) *Cytogenet. Cell Genet.* 51: 1054–1055, 1989.
- [80942] 19272. Ohno, S.; Minoshima, S.; Kudoh, J.; Fukuyama, R.; Shimizu, Y.; Ohmi-Imajoh, S.; Shimizu, N.; Suzuki, K.: Four genes for the calpain family locate on four distinct human chromosomes. *Cytogenet. Cell Genet.* 53: 225–229, 1990.
- [80943] 19273. Sakihama, T.; Kakidani, H.; Zenita, K.; Yumoto, N.; Kikuchi, T.; Sasaki, T.; Kannagi, R.; Nakanishi, S.; Ohmori, M.; Takio, K.; Titani, K.; Murachi, T.: A putative  $\text{Ca}^{2+}$ -binding protein: structure of the light subunit of porcine calpain elucidated by molecular cloning and protein sequence analysis. *Proc. Nat. Acad. Sci.* 82: 6075–6079, 1985.
- [80944] 19274. Berchtold, M. W.; Egli, R.; Rhyner, J. A.; Hameister,

H.; Strehler, E. E.: Localization of the human bona fide calmodulin genes CALM1, CALM2, and CALM3 to chromosomes 14q24-q31, 2p21.1-p21.3, and 19q13.2-q13.3. *Genomics* 16:461-465, 1993.

[80945] 19275. Chin, D.; Winkler, K. E.; Means, A. R.: Characterization of substrate phosphorylation and use of calmodulin mutants to address implications from the enzyme crystal structure of calmodulin-dependent protein kinase I. *J. Biol. Chem.* 272: 31235-31240, 1997.

[80946] 19276. Drum, C. L.; Yan, S.-Z.; Bard, J.; Shen, Y.-Q.; Lu, D.; Soelaiman, S.; Grabarek, Z.; Bohm, A.; Tang, W.-J.: Structural basis for the activation of anthrax adenyl cyclase exotoxin by calmodulin. *Nature* 415:396-402, 2002.

[80947] 19277. Koller, M.; Schnyder, B.; Strehler, E. E.: Structural organization of the human CaMIII calmodulin gene. *Biochim. Biophys. Acta* 1087:180-189, 1990.

[80948] 19278. Kretsinger, R. H.; Rudnick, S. E.; Weissman, L. J.: Crystal structure of calmodulin. *J. Inorganic Biochem.* 28: 289-302, 1986.

[80949] 19279. McPherson, J. D.; Hickie, R. A.; Wasmuth, J. J.; Meyskens, F. L.; Perham, R. N.; Strehler, E. E.; Graham, M. T.: Chromosomal localization of multiple genes encoding calmodulin. (Abstract) *Cytogenet. Cell Genet.* 58: 1951

only, 1991.

- [80950] 19280. Pegues, J. C.; Friedberg, F.: Multiple mRNAs encoding human calmodulin. *Biochem. Biophys. Res. Commun.* 172: 1145–1149, 1990.
- [80951] 19281. Rhyner, J. A.; Ottiger, M.; Wicki, R.; Greenwood, T. M.; Strehler, E. E.: Structure of the human CALM1 calmodulin gene and identification of two CALM1-related pseudogenes CALM1P1 and CALM1P2. *Europ. J. Biochem.* 225: 71–82, 1994.
- [80952] 19282. Scambler, P. J.; McPherson, M. A.; Bates, G.; Bradbury, N. A.; Dormer, R. L.; Williamson, R.: Biochemical and genetic exclusion of calmodulin as the site of the basic defect in cystic fibrosis. *Hum. Genet.* 76: 278–282, 1987.
- [80953] 19283. Schumacher, M. A.; Rivard, A. F.; Bachinger, H. P.; Adelman, J. P.: Structure of the gating domain of a  $\text{Ca}(2+)$ -activated  $\text{K}^+$  channel complexed with  $\text{Ca}(2+)/$ calmodulin. *Nature* 410: 1120–1124, 2001.
- [80954] 19284. Sen Gupta, B.; Detera-Wadleigh, S. D.; McBride, O. W.; Friedberg, F.: A calmodulin pseudogene on human chromosome 17. *Nucleic Acids Res.* 17: 2868 only, 1989.
- [80955] 19285. Powers, P. A.; Liu, S.; Hogan, K.; Gregg, R. G.: Molecular characterization of the gene encoding the gamma subunit of the human skeletal mus–

cle1,4-dihydropyridine-sensitive  $\text{Ca}^{2+}$  channel (CACNLG), cDNA sequence, gene structure, and chromosomal location. J. Biol. Chem. 268: 9275–9279, 1993.

[80956] 19286. Ambartsumian, N.; Tarabykina, S.; Grigorian, M.; Tulchinsky, E.; Hulgaard, E.; Georgiev, G.; Lukanidin, E.: Characterization of two splice variants of metastasis-associated human mts1 gene. Gene 159:125–130, 1995.

[80957] 19287. Jackson-Grusby, L. L.; Swiergiel, J.; Linzer, D. I.: A growth-related mRNA in cultured mouse cells encodes a placental calcium binding protein. Nucleic Acids Res. 15: 6677–6690, 1987.

[80958] 19288. Iles, D. E.; Segers, B.; Weghuis, D. O.; Suikerbuijk, R.; Wieringa, B.: Localization of the gamma-subunit of the skeletal muscle L-type voltage-dependent calcium channel gene (CACNLG) to human chromosome band 17q24 by in situ hybridization and identification of a polymorphic repetitive DNA sequence at the gene locus. Cytogenet. Cell Genet. 64:227–230, 1993.

[80959] 19289. Saris, C. J.; Kristensen, T.; D'Eustachio, P.; Hicks, L. J.; Noonan, D. J.; Hunter, T.; Tack, B. F.: cDNA sequence and tissue distribution of the mRNA for bovine and murine p11, the S100-related light chain of the protein-tyrosine kinase substrate p36 (calpactin I). J. Biol. Chem. 262:



10663–10671, 1987.

- [80960] 19290.Lefort, A.; Lecocq, R.; Libert, F.; Lamy, F.; Swillens, S.; Vassart,G.; Dumont, J. E.: Cloning and sequencing of a calcium-binding proteinregulated by cyclic AMP in the thyroid. EMBO J. 8: 111–116, 1989.
- [80961] 19291.Loughlin, J.; Irven, C.; Sykes, B.: Exclusion of the cartilagelink protein and the cartilage matrix protein genes as the mutantloci in several heritable chondrodysplasias. Hum. Genet. 94: 698–700,1994.
- [80962] 19292.Osborne–Lawrence, S. L.; Sinclair, A. K.; Hicks, R. C.; Lacey,S. W.; Eddy, R. L., Jr.; Byers, M. G.; Shows, T. B.; Duby, A. D.:Complete amino acid sequence of human cartilage link protein (CRTL1)deduced from cDNA clones and chromosomal assignment of the gene. Genomics 8:562–567, 1990.
- [80963] 19293.Watanabe, H.; Yamada, Y.: Mice lacking link protein develop dwarfismand craniofacial abnormalities. Nature Genet. 21: 225–229, 1999.
- [80964] 19294.Deak, F.; Piecha, D.; Bachrati, C.; Paulsson, M.; Kiss, I.: Primarystructure and expression of matrilin–2, the closest relative of cartilagematrix protein within the von Willebrand factor type A–like modulesuperfamily. J. Biol. Chem. 272: 9268–9274, 1997.

- [80965] 19295.Deak, F.; Wagener, R.; Kiss, I.; Paulsson, M.: The matrilins:a novel family of oligomeric extracellular matrix proteins. *MatrixBiol.* 18: 55–64, 1999.
- [80966] 19296.Hansson, A.–S.; Heinegard, D.; Holmdahl, R.: A new animal model for relapsing polychondritis, induced by cartilage matrix protein(matrilin–1). *J. Clin. Invest.* 104: 589–598, 1999.
- [80967] 19297.Jenkins, R. N.; Osborne–Lawrence, S. L.; Sinclair, A. K.; Eddy, R. L., Jr.; Byers, M. G.; Shows, T. B.; Duby, A. D.: Structure and chromosomal location of the human gene encoding cartilage matrix protein. *J. Biol. Chem.* 265: 19624–19631, 1990.
- [80968] 19298.Lang, B.; Rothenfusser, A.; Lanchbury, J. S.; Rauh, G.; Breedveld, F. C.; Urlacher, A.; Albert, E. D.; Peter, H.–H.; Melchers, I.: Susceptibility to relapsing polychondritis is associated with HLA–DR4. *Arthritis Rheum.* 36: 660–664, 1993.
- [80969] 19299.Muratoglu, S.; Krysan, K.; Balazs, M.; Sheng, H.; Zakany, R.; Modis, L.; Kiss, I.; Deak, F.: Primary structure of human matrilin–2, chromosomal location of the MATN2 gene and conservation of an AT–AC intron in matrilin genes. *Cytogenet. Cell Genet.* 90: 323–327, 2000.
- [80970] 19300.Wagener, R.; Kobbe, B.; Paulsson, M.: Primary

structure of matrilin-3, a new member of a family of extracellular matrix proteins related to cartilage matrix protein (matrilin-1) and von Willebrand factor. FEBS Lett. 413: 129-134, 1997.

- [80971] 19301. Zeuner, M.; Straub, R. H.; Rauh, G.; Albert, E. D.; Scholmerich, J.; Lang, B.: Relapsing polychondritis: clinical and immunogenetic analysis of 62 patients. J. Rheum. 24: 96-101, 1997.
- [80972] 19302. Boldyreff, B.; Klett, C.; Gottert, E.; Geurts van Kessel, A.; Hameister, H.; Issinger, O.-G.: Assignment of casein kinase 2 alpha sequence to two different human chromosomes. Hum. Genet. 89: 79-82, 1992.
- [80973] 19303. Doray, B.; Ghosh, P.; Griffith, J.; Geuze, H. J.; Kornfeld, S.: Cooperation of GGAs and AP-1 in packaging MPRs at the trans-Golgi network. Science 297: 1700-1703, 2002.
- [80974] 19304. Meisner, H.; Heller-Harrison, R.; Buxton, J.; Czech, M. P.: Molecular cloning of the human casein kinase II alpha subunit. Biochemistry 28: 4072-4076, 1989.
- [80975] 19305. Wirkner, U.; Voss, H.; Lichter, P.; Ansorge, W.; Pyerin, W.: The human gene (CSNK2A1) coding for the casein kinase II subunit alpha is located on chromosome 20 and contains tandemly arranged Alu repeats. Genomics

19:257–265, 1994.

- [80976] 19306.Wirkner, U.; Voss, H.; Lichter, P.; Weitz, S.; An-  
sorge, W.; Pyerin,W.: Human casein kinase II subunit al-  
pha: sequence of a processed(pseudo)gene and its local-  
ization on chromosome 11. *Biochim. Biophys.Acta* 1131:  
220–222, 1992.
- [80977] 19307.Yang–Feng, T. L.; Zheng, K.; Kopatz, I.; Naiman, T.;  
Canaani, D.: Mapping of the human casein kinase II cat-  
alytic subunit genes: twoloci carrying the homologous se-  
quences for the alpha subunit. *NucleicAcids Res.* 19:  
7125–7129, 1991.
- [80978] 19308.Lozman, F. J.; Litchfield, D. W.; Piening, C.; Takio,  
K.; Walsh,K. A.; Krebs, E. G.: Isolation and characterization  
of human cDNAclones encoding the alpha and the alpha-  
prime subunits of casein kinasell. *Biochemistry* 29:  
8436–8447, 1990.
- [80979] 19309.Xu, X.; Toselli, P. A.; Russell, L. D.; Seldin, D. C.:  
Globozoospermia in mice lacking the casein kinase II al-  
pha–prime catalytic subunit. *NatureGenet.* 23: 118–121,  
1999.
- [80980] 19310.Abrieu, A.; Kahana, J. A.; Wood, K. W.; Cleveland,  
D. W.: CENP–Eas an essential component of the mitotic  
checkpoint in vitro. *Cell* 102:817–826, 2000.

- [80981] 19311.Wood, K. W.; Sakowicz, R.; Goldstein, L. S. B.; Cleveland, D. W.: CENP-E is a plus end-directed kinetochore motor required for metaphase chromosome alignment. *Cell* 97: 357–366, 1997.
- [80982] 19312.Yao, X.; Abrieu, A.; Zheng, Y.; Sullivan, K. F.; Cleveland, D.W.: CENP-E forms a link between attachment of spindle microtubulesto kinetochores and the mitotic checkpoint. *Nature Cell Biol.* 2:484–491, 2000.
- [80983] 19313.Yen, T. J.; Compton, D. A.; Wise, D.; Zinkowski, R. P.; Brinkley, B. R.; Earnshaw, W. C.; Cleveland, D. W.: CENP-E, a novel human centromere-associated protein required for progression from metaphase to anaphase. *EMBOJ.* 10: 1245–1254, 1991.
- [80984] 19314.Yen, T. J.; Li, G.; Schaar, B. T.; Szilak, I.; Cleveland, D. W.: CENP-E is a putative kinetochore motor that accumulates just before mitosis. *Nature* 359: 536–539, 1992.
- [80985] 19315.Geissler, E. N.; Liao, M.; Brook, J. D.; Martin, F. H.; Zsebo, K. M.; Housman, D. E.; Galli, S. J.: Stem cell factor (SCF), a novel hematopoietic growth factor and ligand for c-kit tyrosine kinase receptor, maps on human chromosome 12 between 12q14.3 and 12qter. *Somat. Cell Molec. Genet.* 17: 207–214, 1991.
- [80986] 19316.Heissig, B.; Hattori, K.; Dias, S.; Friedrich, M.; Fer-

ris, B.;Hackett, N. R.; Crystal, R. G.; Besmer, P.; Lyden, D.; Moore, M. A.S.; Werb, Z.; Rafii, S.: Recruitment of stem and progenitor cellsfrom the bone marrow niche requires MMP-9 mediated release of Kit-ligand. Cell 109:625-637, 2002.

[80987] 19317.Martin, F. H.; Suggs, S. V.; Langley, K. E.; Lu, H. S.; Ting, J.;Okino, K. H.; Morris, C. F.; McNiece, I. K.; Jacobsen, F. W.; Mendiaz,E. A.; Birkett, N. C.; Smith, K. A.; and 15 others: Primary structureand functional expression of rat and human stem cell factor DNAs. Cell 63:203-211, 1990.

[80988] 19318.Mathew, S.; Murty, V. V. V. S.; Hunziker, W.; Chaganti, R. S. K.: Subregional mapping of 13 single-copy genes on the long arm of chromosome12 by fluorescence in situ hybridization. Genomics 14: 775-779,1992.

[80989] 19319.Li, X.-Y.; Mattei, M. G.; Zaleska-Rutczynska, Z.; Hooft van Huijsduijnen,R.; Figueroa, F.; Nadeau, J.; Benoist, C.; Mathis, D.: One subunitof the transcription factor NF-Y maps close to the major histocompatibility-complex in murine and human chromosomes. Genomics 11: 630-634, 1991.

[80990] 19320.Johnston, J.; Bollekens, J.; Allen, R. H.; Berliner, N.: Structureof the cDNA encoding transcobalamin I, a neu-

trophil granule protein. J.Biol. Chem. 264: 15754–15757, 1989.

- [80991] 19321. Johnston, J.; Yang–Feng, T.; Berliner, N.: Genomic structure and mapping of the chromosomal gene for transcobalamin I (TCN1): comparison to human intrinsic factor. Genomics 12: 459–464, 1992.
- [80992] 19322. Yang–Feng, T. L.; Berliner, N.; Deverajan, P.; Johnston, J.: Assignment of two human neutrophil secondary granule protein genes, transcobalamin I and neutrophil collagenase to chromosome 11. (Abstract) Cytogenet. Cell Genet. 58: 1974 only, 1991.
- [80993] 19323. Courey, A. J.; Holtzman, D. A.; Jackson, S. P.; Tjian, R.: Synergistic activation by the glutamine–rich domains of human transcription factor Sp1. Cell 59: 827–836, 1989.
- [80994] 19324. Gaynor, R. B.; Shieh, B.–H.; Klisak, I.; Sparkes, R. S.; Lusk, A. J.: Localization of the transcription factor SP1 gene to human chromosome 12q12–q13.2. Cytogenet. Cell Genet. 64: 210–212, 1993.
- [80995] 19325. Kadonaga, J. T.; Carner, K. R.; Masiarz, F. R.; Tjian, R.: Isolation of a cDNA encoding transcription factor Sp1 and functional analysis of the DNA binding domain. Cell 51: 1079–1090, 1987.
- [80996] 19326. Matera, A. G.; Ward, D. C.: Localization of the hu–

- man Sp1 transcriptionfactor gene to 12q13 by fluorescence in situ hybridization. *Genomics* 17:793–794, 1993.
- [80997] 19327.Szpirer, J.; Szpirer, C.; Riviere, M.; Levan, G.; Marynen, P.;Cassiman, J.–J.; Wiese, R.; DeLuca, H. F.: The Sp1 transcriptionfactor gene (SP1) and the 1,25–dihydroxyvitamin D(3) receptor gene(VDR) are colocalized on human chromosome arm 12q and rat chromosome
- [80998] 19328.*Genomics* 11: 168–173, 1991.7. Wimmer, E. A.; Jackle, H.; Pfeifle, C.; Cohen, S. M.: A Drosophilahomologue of human Sp1 is a head–specific segmentation gene. *Nature* 366:690–694, 1993.
- [80999] 19329.Abbott, C.; Piaggio, G.; Ammendola, R.; Solomon, E.; Povey, S.;Gounari, F.; De Simone, V.; Cortese, R.: Mapping of the gene TCF2for the transcription factor LFB3 to human chromosome 17 by polymerasechain reaction. *Genomics* 8: 165–167, 1990.
- [81000] 19330.Bach, I.; Mattei, M.–G.; Cereghini, S.; Yaniv, M.: Two membersof an HNF1 homeoprotein family are expressed in human liver. *NucleicAcids Res.* 19: 3553–3559, 1991.
- [81001] 19331.Bingham, C.; Ellard, S.; Allen, L.; Bulman, M.; Shepherd, M.; Frayling,T.; Berry, P. J.; Clark, P. M.; Lindner, T.; Bell, G. I.; Ryffel,G. U.; Nicholls, A. J.; Hattersley, A. T.: Ab–



normal nephron development associated with a frameshift mutation in the transcription factor hepatocyte nuclear factor-1-beta. *Kidney Int.* 57: 898-907, 2000.

[81002] 19332. Horikawa, Y.; Iwasaki, N.; Hara, M.; Furuta, H.; Hinokio, Y.; Cockburn, B. N.; Lindner, T.; Yamagata, K.; Ogata, M.; Tomonaga, O.; Kuroki, H.; Kasahara, T.; Iwamoto, Y.; Bell, G. I.: Mutation in hepatocyte nuclear factor-1-beta gene (TCF2) associated with MODY. (Letter) *Nature Genet.* 17: 384-385, 1997.

[81003] 19333. Menzel, R.; Kaisaki, P. J.; Rjasanowski, I.; Heinke, P.; Kerner, W.; Menzel, S.: A low renal threshold for glucose in diabetic patients with a mutation in the hepatocyte nuclear factor-1-alpha (HNF-1-alpha) gene. *Diabet. Med.* 15: 816-820, 1998.

[81004] 19334. Nishigori, H.; Yamada, S.; Kohama, T.; Tomura, H.; Sho, K.; Horikawa, Y.; Bell, G. I.; Takeuchi, T.; Takeda, J.: Frameshift mutation, A263fsinsGG, in the hepatocyte nuclear factor-1-beta gene associated with diabetes and renal dysfunction. *Diabetes* 47: 1354-1355, 1998.

[81005] 19335. Wild, W.; Pogge von Strandmann, E.; Nastos, A.; Senkel, S.; Lingott-Frieg, A.; Bulman, M.; Bingham, C.; El-lard, S.; Hattersley, A. T.; Ryffel, G. U.: The mutated human gene encoding hepatocyte nuclear factor 1-beta inhibits

kidney formation in developing *Xenopus* embryos.

Proc.Nat. Acad. Sci. 97: 4695–4700, 2000.

- [81006] 19336.Aplan, P. D.; Jones, C. A.; Chervinsky, D. S.; Zhao, X.; Ellsworth,M.; Wu, C.; McGuire, E. A.; Gross, K. W.: An scl gene product lackingthe transactivation domain induces bony abnormalities and cooperateswith LMO1 to generate T-cell malignancies in transgenic mice. EMBOJ. 16: 2408–2419, 1997.
- [81007] 19337.Aplan, P. D.; Lombardi, D. P.; Ginsberg, A. M.; Cossman, J.; Bertness,V. L.; Kirsch, I. L.: Disruption of the human SCL locus by 'illegitimate'V–(D)–J recombinase activity. Science 250: 1426–1429, 1990.
- [81008] 19338.Aplan, P. D.; Lombardi, D. P.; Reaman, G. H.; Sather, H. N.; Hammond,G. D.; Kirsch, I. R.: Involvement of the putative hematopoietic transcriptionfactor SCL in T-cell acute lymphoblastic leukemia. Blood 79: 1327–1333,1992.
- [81009] 19339.Brown, L.; Cheng, J.–T.; Chen, Q.; Siciliano, M. J.; Crist, W.;Buchanan, G.; Baer, R.: Site–specific recombination of the tal–1gene is a common occurrence in human T-cell leukemia. EMBO J. 9:3343–3351, 1990.
- [81010] 19340.Collazo–Garcia, N.; Scherer, P.; Aplan, P. D.: Cloning and characterizationof a murine SIL gene. Ge–

nomics 30: 506–513, 1995.

- [81011] 19341.Izraeli, S.; Lowe, L. A.; Bertness, V. L.; Good, D. J.; Dorward, D. W.; Kirsch, I. R.; Kuehn, M. R.: The SIL gene is required for mouse embryonic axial development and left-right specification. *Nature* 399:691–694, 1999.
- [81012] 19342.Roy, A. L.; Du, H.; Gregor, P. D.; Novina, C. D.; Martinez, E.; Roeder, R. G.: Cloning of an Inr- and E-box binding protein, TFIID-1, that interacts physically and functionally with USF1. *EMBO J.* 16:7091–7104, 1997.
- [81013] 19343.Steingrimsson, E.; Sawadogo, M.; Gilbert, D. J.; Zervos, A. S.; Brent, R.; Blumberg, M. A.; Fisher, D. E.; Copeland, N. G.; Jenkins, N. A.: Murine chromosomal location of five bHLH-Zip transcription factor genes. *Genomics* 28: 179–183, 1995.
- [81014] 19344.Zhong, G.; Fan, P.; Ji, H.; Dong, F.; Huang, Y.: Identification of a chlamydial protease-like activity factor responsible for the degradation of host transcription factors. *J. Exp. Med.* 193: 935–942, 2001.
- [81015] 19345.Aasland, R.; Olsen, L. C.; Spurr, N. K.; Krokan, H. E.; Helland, D. E.: Chromosomal assignment of human uracil-DNA glycosylase to chromosome 12. *Genomics* 7: 139–141, 1990.
- [81016] 19346.Dinner, A. R.; Blackburn, G. M.; Karplus, M.: Uracil-

DNA glycolase acts by substrate autocatalysis. *Nature* 413: 752–755, 2001.

- [81017] 19347. Haug, T.; Skorpen, F.; Kvaloy, K.; Eftedal, I.; Lund, H.; Krokan, H. E.: Human uracil–DNA glycosylase gene: sequence organization, methylation pattern, and mapping to chromosome 12q23–q24.1. *Genomics* 36:408–416, 1996.
- [81018] 19348. Haug, T.; Skorpen, F.; Lund, H.; Krokan, H. E.: Structure of the gene for human uracil–DNA glycosylase and analysis of the promoter function. *FEBS Lett.* 353: 180–184, 1994.
- [81019] 19349. Meyer–Siegler, K.; Mauro, D. J.; Seal, G.; Wurzer, J.; deRiel, J. K.; Sirover, M. A.: A human nuclear uracil DNA glycosylase is the 37–kDa subunit of glyceraldehyde–3–phosphate dehydrogenase. *Proc. Nat. Acad. Sci.* 88: 8460–8464, 1991.
- [81020] 19350. Nilsen, H.; Rosewell, I.; Robins, P.; Skjelbred, C. F.; Andersen, S.; Slupphaug, G.; Daly, G.; Krokan, H. E.; Lindahl, T.; Barnes, D. E.: Uracil–DNA glycosylase (UNG)–deficient mice reveal a primary role of the enzyme during DNA replication. *Molec. Cell* 5: 1059–1065, 2000.
- [81021] 19351. Olsen, L. C.; Aasland, R.; Wittwer, C. U.; Krokan, H. E.; Helland, D. E.: Molecular cloning of human uracil–DNA glycosylase, a highly conserved DNA repair enzyme. *EMBO*

J. 8: 3121–3125, 1989.

- [81022] 19352.Vollberg, T. M.; Siegler, K. M.; Cool, B. L.; Sirover, M. A.: Isolation and characterization of the human uracil DNA glycosylase gene. Proc. Nat. Acad. Sci. 86: 8693–8697, 1989.
- [81023] 19353.Momeni, P.; Glockner, G.; Schmidt, O.; von Holtum, D.; Albrecht, B.; Gillessen-Kaesbach, G.; Hennekam, R.; Meinecke, P.; Zabel, B.; Rosenthal, A.; Horsthemke, B.; Ludecke, H.-J.: Mutations in a new gene, encoding a zinc-finger protein, cause tricho-rhino-phalangeal syndrome type I. Nature Genet. 24: 71–74, 2000.
- [81024] 19354.Komuro, I.; Wenninger, K. E.; Philipson, K. D.; Izumo, S.: Molecular cloning and characterization of the human cardiac Na(+)/Ca(2+) exchanger cDNA. Proc. Nat. Acad. Sci. 89: 4769–4773, 1992.
- [81025] 19355.Kraev, A.; Chumakov, I.; Carafoli, E.: The organization of the human gene NCX1 encoding the sodium-calcium exchanger. Genomics 37:105–112, 1996.
- [81026] 19356.McDaniel, L. D.; Lederer, W. J.; Kofuji, P.; Schulze, D. H.; Kieval, R.; Schultz, R. A.: Mapping of the human cardiac Na<sup>+</sup>/Ca<sup>2+</sup> exchanger gene (NCX1) by fluorescent in situ hybridization to chromosome region 2p22–p23. Cytogenet. Cell Genet. 63: 192–193, 1993.

- [81027] 19357.Shieh, B.-H.; Xia, Y.; Sparkes, R. S.; Klisak, I.; Lusk, A. J.; Nicoll, D. A.; Philipson, K. D.: Mapping of the gene for the cardiac sarcolemmal Na(+)-Ca(2+) exchanger to human chromosome 2p21-p23. *Genomics* 12:616-617, 1992.
- [81028] 19358.Wakimoto, K.; Kobayashi, K.; Kuro-o, M.; Yao, A.; Iwamoto, T.; Yanaka, N.; Kita, S.; Nishida, A.; Azuma, S.; Toyoda, Y.; Omori, K.; Imahie, H.; Oka, T.; Kudoh, S.; Kohmoto, O.; Yazaki, Y.; Shigekawa, M.; Imai, Y.; Nabeshima, Y.; Komura, I.: Targeted disruption of Na(+)/Ca(2+) exchanger gene leads to cardiomyocyte apoptosis and defects in heartbeat. *J. Biol. Chem.* 275: 36991-36998, 2000.
- [81029] 19359.Chong, S. S.; Kristjansson, K.; Zoghbi, H. Y.; Hughes, M. R.: Molecular cloning of the cDNA encoding a human renal sodium phosphate transport protein and its assignment to chromosome 6p21.3-p23. *Genomics* 18:355-359, 1993.
- [81030] 19360.Kos, C. H.; Tihy, F.; Murer, H.; Lemieux, N.; Tenenhouse, H. S.: Comparative mapping of Na(+)-phosphate cotransporter genes, NPT1 and NPT2, in human and rabbit. *Cytogenet. Cell Genet.* 75: 22-24, 1996.
- [81031] 19361.Zhang, X.-X.; Tenenhouse, H. S.; Hewson, A. S.; Ey-

doux, P.: Assignment of renal specific Na(+)-phosphate cotransporter gene Slc17a1 to mouse chromosome bands 13A3–A4 by in situ hybridization. *Cytogenet. Cell Genet.* 76: 180 only, 1997.

- [81032] 19362. Ghishan, F. K.; Knobel, S.; Dasuki, M.; Butler, M.; Phillips, J.: Chromosomal localization of the human renal sodium phosphate transporter to chromosome 5: implications for X-linked hypophosphatemia. *Pediatr. Res.* 35: 510–513, 1994.
- [81033] 19363. Hartmann, C. M.; Hewson, A. S.; Kos, C. H.; Hilfiker, H.; Soumounou, Y.; Murer, H.; Tenenhouse, H. S.: Structure of murine and human renal type II Na(+)-phosphate cotransporter genes (Npt2 and NPT2). *Proc. Nat. Acad. Sci.* 93: 7409–7414, 1996.
- [81034] 19364. Kos, C. H.; Tihy, F.; Econs, M. J.; Murer, H.; Lemieux, N.; Tenenhouse, H. S.: Localization of a renal sodium-phosphate cotransporter gene to human chromosome 5q35. *Genomics* 19: 176–177, 1994.
- [81035] 19365. Magagnin, S.; Werner, A.; Markovich, D.; Sorribas, V.; Stange, G.; Biber, J.; Murer, H.: Expression cloning of human and rat renal cortex Na/Pi cotransport. *Proc. Nat. Acad. Sci.* 90: 5979–5983, 1993.
- [81036] 19366. McPherson, J. D.; Krane, M. C.; Wagner-McPherson,

C. B.; Kos, C.H.; Tenenhouse, H. S.: High resolution mapping of the renal sodium–phosphate cotransporter gene (NPT2) confirms its localization to human chromosome 5q35. *Pediat. Res.* 41: 632–634, 1997.

- [81037] 19367. Tenenhouse, H. S.; Werner, A.; Biber, J.; Ma, S.; Martel, J.; Roy, S.; Murer, H.: Renal Na(+)-phosphate cotransport in murine X-linked hypophosphatemic rickets: molecular characterization. *J. Clin. Invest.* 93: 671–676, 1994.
- [81038] 19368. Zhang, X.-X.; Tenenhouse, H. S.; Hewson, A. S.; Murer, H.; Eydoux, P.: Assignment of renal-specific Na(+)-phosphate cotransporter gene *Slc17a2* to mouse chromosome band 13B by in situ hybridization. *Cytogenet. Cell Genet.* 77: 304–305, 1997.
- [81039] 19369. Barnard, R.; Kelly, G.; Manzetti, S. O.; Harris, E. L.: Neither the New Zealand genetically hypertensive strain nor Dahl salt-sensitive strain has an A1079T transversion in the alpha-1 isoform of the Na(+),K(+)-ATPase gene. *Hypertension* 38: 786–792, 2001.
- [81040] 19370. Chehab, F. F.; Kan, Y. W.; Law, M. L.; Hartz, J.; Kao, F.-T.; Blostein, R.: Human placental Na<sup>+</sup>,K<sup>+</sup>-ATPase alpha subunit: cDNA cloning, tissue expression, DNA polymorphism, and chromosomal localization. *Proc. Nat. Acad. Sci.*



84: 7901–7905, 1987.

- [81041] 19371.Dahl, L. K.; Heine, M.; Tassinari, L.: Role of genetic factors in susceptibility to experimental hypertension due to chronic excess salt ingestion. *Nature* 194: 480–482, 1972.
- [81042] 19372.Dahl, L. K.; Heine, M.; Thompson, K.: Genetic influence of the kidneys on blood pressure: evidence from chronic renal homografts in rats with opposite predispositions to hypertension. *Circ. Res.* 40:94–101, 1974.
- [81043] 19373.Glorioso, N.; Filigheddu, F.; Troffa, C.; Soro, A.; Parpaglia, P. P.; Tsikoudakis, A.; Myers, R. H.; Herrera, V. L. M.; Ruiz-Opazo, N.: Interaction of alpha-1-Na,K-ATPase and Na,K,2Cl-cotransporter genes in human essential hypertension. *Hypertension* 38: 204–209, 2001.
- [81044] 19374.Herrera, V. L. M.; Ruiz-Opazo, N.: Alteration of alpha-1 Na<sup>+</sup>,K<sup>(+)</sup>-ATPase 86R-beta(+) influx by a single amino acid substitution. *Science* 249:1023–1026, 1990.
- [81045] 19375.Andersson, S.; Russell, D. W.: Structural and biochemical properties of cloned and expressed human and rat steroid 5-alpha-reductases. *Proc.Nat. Acad. Sci.* 87: 3640–3644, 1990.
- [81046] 19376.Ellis, J. A.; Stebbing, M.; Harrap, S. B.: Genetic analysis of male pattern baldness and the 5-alpha-reductase

genes. J. Invest.Derm. 110: 849–853, 1998.

- [81047] 19377.Harris, G.; Azzolina, B.; Baginsky, W.; Cimis, G.; Rasmusson, G.H.; Tolman, R. L.; Raetz, C. R. H.; Ellsworth, K.: Identification and selective inhibition of an isozyme of steroid 5–alpha–reductase in human scalp. Proc. Nat. Acad. Sci. 89: 10787–10791, 1992.
- [81048] 19378.Hsieh, C.–L.; Milatovich, A.; Russell, D.; Francke, U.: Chromosomal mapping of human steroid 5 alpha–reductase gene (SRD5A1) and pseudogene (SRD5AP1) in human and mouse. (Abstract) Cytogenet. Cell Genet. 58:1897 only, 1991.
- [81049] 19379.Jenkins, E. P.; Andersson, S.; Imperato–McGinley, J.; Wilson, J.D.; Russell, D. W.: Genetic and pharmacological evidence for more than one human steroid 5–alpha–reductase. J. Clin. Invest. 89: 293–300, 1992.
- [81050] 19380.Jenkins, E. P.; Hsieh, C.–L.; Milatovich, A.; Normington, K.; Berman, D. M.; Francke, U.; Russell, D. W.: Characterization and chromosomal mapping of a human steroid 5–alpha–reductase gene and pseudogene and mapping of the mouse homologue. Genomics 11: 1102–1112, 1991.
- [81051] 19381.Thigpen, A. E.; Silver, R. I.; Guileyardo, J. M.; Casey, M. L.; McConnell, J. D.; Russell, D. W.: Tissue distribution

and ontogeny of steroid 5- $\alpha$ -reductase isozyme expression. *J. Clin. Invest.* 92:903–910, 1993.

- [81052] 19382. He, Z.; Yamamoto, R.; Furth, E. E.; Schantz, L. J.; Naylor, S. L.; George, H.; Billheimer, J. T.; Strauss, J. F., III: cDNAs encoding members of a family of proteins related to human sterol carrier protein 2 and assignment of the gene to human chromosome 1p21-pter. *DNA Cell Biol.* 10: 559–569, 1991.
- [81053] 19383. Vesa, J.; Hellsten, E.; Barnoski, B. L.; Emanuel, B. S.; Billheimer, J. T.; Mead, S.; Cowell, J. K.; Strauss, J. F., III; Peltonen, L.: Assignment of sterol carrier protein X/sterol carrier protein 2 to 1p32 and its exclusion as the causative gene for infantile neuronal ceroid lipofuscinosis. *Hum. Molec. Genet.* 3: 341–346, 1994.
- [81054] 19384. Welch, C. L.; Xia, Y.-R.; Billheimer, J. T.; Strauss, J. F., III; Lusi, A. J.: Assignment of the mouse sterol carrier protein gene (Scp2) to chromosome 4. *Mammalian Genome* 7: 624–625, 1996.
- [81055] 19385. Yamamoto, R.; Kallen, C. B.; Babalola, G. O.; Renert, H.; Billheimer, J. T.; Strauss, J. F., III: Cloning and expression of a cDNA encoding human sterol carrier protein 2. *Proc. Nat. Acad. Sci.* 88: 463–467, 1991.
- [81056] 19386. Yamamoto, R.; Naylor, S. L.; George, H.; Billheimer,

J. T.; Strauss, J. F., III: Assignment of the gene encoding sterol carrier protein<sup>2</sup> to human chromosome 1pter-p21. (Abstract) Cytogenet. Cell Genet. 58:1866-1867, 1991.

[81057] 19387. DeBose-Boyd, R. A.; Brown, M. S.; Li, W.-P.; Nothurfft, A.; Goldstein, J. L.; Espenshade, P. J.: Transport-dependent proteolysis of SREBP: relocation of Site-1 protease from Golgi to ER obviates the need for SREBP transport to Golgi. Cell 99: 703-712, 1999.

[81058] 19388. Hua, X.; Wu, J.; Goldstein, J. L.; Brown, M. S.; Hobbs, H. H.: Structure of the human gene encoding sterol regulatory element binding protein-1 (SREBF1) and localization of SREBF1 and SREBF2 to chromosomes 17p11.2 and 22q13. Genomics 25: 667-673, 1995.

[81059] 19389. Osborne, T. F.: CREating a SCAP-less liver keeps SREBPs pinned in the ER membrane and prevents increased lipid synthesis in response to low cholesterol and high insulin. Genes Dev. 15: 1873-1878, 2001.

[81060] 19390. Shimano, H.; Shimomura, I.; Hammer, R. E.; Herz, J.; Goldstein, J. L.; Brown, M. S.; Horton, J. D.: Elevated levels of SREBP-2 and cholesterol synthesis in livers of mice homozygous for a targeted disruption of the SREBP-1 gene. J. Clin. Invest. 100: 2115-2124, 1997.

[81061] 19391. Shimomura, I.; Hammer, R. E.; Richardson, J. A.;

Ikemoto, S.; Bashmakov, Y.; Goldstein, J. L.; Brown, M. S.:  
Insulin resistance and diabetes mellitus in transgenic mice  
expressing nuclear SREBP-1c in adipose tissue: model for  
congenital generalized lipodystrophy. *Genes Dev.*  
12:3182–3194, 1998.

[81062] 19392. Tobe, K.; Suzuki, R.; Aoyama, M.; Yamauchi, T.;  
Kamon, J.; Kubota, N.; Terauchi, Y.; Matsui, J.; Akanuma,  
Y.; Kimura, S.; Tanaka, J.; Abe, M.; Ohsumi, J.; Nagai, R.;  
Kadowaki, T.: Increased expression of the sterol regulatory  
element-binding protein-1 gene in insulin receptor sub-  
strate-2  $-/-$  mouse liver. *J. Biol. Chem.* 276:  
38337–38340, 2001.

[81063] 19393. Wang, X.; Sato, R.; Brown, M. S.; Hua, X.; Goldstein,  
J. L.: SREBP-1, a membrane-bound transcription factor re-  
leased by sterol-regulated proteolysis. *Cell* 77: 53–62,  
1994.

[81064] 19394. Yokoyama, C.; Wang, X.; Briggs, M. R.; Admon, A.;  
Wu, J.; Hua, X.; Goldstein, J. L.; Brown, M. S.: SREBP-1, a  
basic-helix-loop-helix-leucine zipper protein that controls  
transcription of the low density lipoprotein receptor gene.  
*Cell* 75: 187–197, 1993.

[81065] 19395. Joosten, P. H. L. J.; Toepoel, M.; Mariman, E. C. M.;  
Van Zoelen, E. J. J.: Promoter haplotype combinations of

the platelet-derived growth factor alpha-receptor gene predispose to human neural tube defects. *Nature Genet.* 27: 215–217, 2001.

[81066] 19396. Travis, G. H.; Sutcliffe, J. G.; Bok, D.: The retinal degeneration slow (rds) gene product is a photoreceptor disc membrane-associated glycoprotein. *Neuron* 6: 61–70, 1991.

[81067] 19397. Morris, C.; Heisterkamp, N.; Hao, Q. L.; Testa, J. R.; Groffen, J.: The human tyrosine kinase gene (FER) maps to chromosome 5 and is deleted in myeloid leukemias with a del(5q). *Cytogenet. Cell Genet.* 53:196–200, 1990.

[81068] 19398. Herrera, V. L. M.; Xie, H. X.; Lopez, L. V.; Schork, N. J.; Ruiz-Opazo, N.: The alpha-1 Na,K-ATPase gene is a susceptibility hypertension gene in the Dahl salt-sensitive-HSD rat. *J. Clin. Invest.* 102: 1102–1111, 1998.

[81069] 19399. Williams, T. M.; Montoya, G.; Wu, Y.; Eddy, R. L.; Byers, M. G.; Shows, T. B.: The TCF8 gene encoding a zinc finger protein (Nil-2-a) resides on human chromosome 10p11.2. *Genomics* 14: 194–196, 1992.

[81070] 19400. van Nie, R.; Ivanyi, D.; Demant, P.: A new H-2-linked mutation, rds, causing retinal degeneration in the mouse. *Tissue Antigens* 12:106–108, 1978.

[81071] 19401. Vine, A. K.; Schatz, H.: Adult-onset foveomacular

pigment epithelial dystrophy. *Am. J. Ophthalmol.* 89: 680–691, 1980.

[81072] 19402. Weleber, R. G.; Carr, R. E.; Murphey, W. H.; Sheffield, V. C.; Stone, E. M.: Phenotypic variation including retinitis pigmentosa, pattern dystrophy, and fundus flavimaculatus in a single family with a deletion of codon 153 or 154 of the peripherin/RDS gene. *Arch. Ophthalmol.* 111: 1531–1542, 1993.

[81073] 19403. Wells, J.; Wroblewski, J.; Keen, J.; Inglehearn, C.; Jubb, C.; Eckstein, A.; Jay, M.; Arden, G.; Bhattacharya, S.; Fitzke, F.; Bird, A.: Mutations in the human retinal degeneration slow (RDS) gene can cause either retinitis pigmentosa or macular dystrophy. *Nature Genet.* 3:213–218, 1993.

[81074] 19404. Wroblewski, J. J.; Wells, J. A., III; Eckstein, A.; Fitzke, F.; Jubb, C.; Keen, J.; Inglehearn, C.; Bhattacharya, S.; Arden, G. B.; Jay, M.; Bird, A. C.: Macular dystrophy associated with mutations at codon 172 in the human retinal degeneration slow gene. *Ophthalmology* 101:12–22, 1994.

[81075] 19405. Eph Nomenclature Committee: Unified nomenclature for the Eph family receptors and their ligands, the ephrins. *Cell* 90: 403–404, 1997.

[81076] 19406. Hattori, M.; Osterfield, M.; Flanagan, J. G.: Regu-

lated cleavage of a contact-mediated axon repellent. *Science* 289: 1360–1365, 2000.

- [81077] 19407. Maru, Y.; Hirai, H.; Yoshida, M. C.; Takaku, F.: Evolution, expression, and chromosomal location of a novel receptor tyrosine kinase gene, *eph*. *Molec. Cell. Biol.* 8: 3770–3776, 1988.
- [81078] 19408. Yoshida, M. C.; Maru, H.; Hirai, H.; Takau, F.: Chromosomal location of a novel receptor tyrosine kinase gene, *EPH*, on chromosome 7. (Abstract) *Cytogenet. Cell Genet.* 51: 1113 only, 1989.
- [81079] 19409. Boyd, A. W.; Ward, L. D.; Wicks, I. P.; Simpson, R. J.; Salvaris, E.; Wilks, A.; Welch, K.; Loudovaris, M.; Rockman, S.; Busmanis, I.: Isolation and characterization of a novel receptor-type protein tyrosine kinase (*hek*) from a human pre-B cell line. *J. Biol. Chem.* 267:3262–3267, 1992.
- [81080] 19410. Fox, G. M.; Holst, P. L.; Chute, H. T.; Lindberg, R. A.; Janssen, A. M.; Basu, R.; Welcher, A. A.: cDNA cloning and tissue distribution of five human *EPH*-like receptor protein-tyrosine kinases. *Oncogene* 10:897–905, 1995.
- [81081] 19411. Wicks, I. P.; Lapsys, N. M.; Baker, E.; Campbell, L. J.; Boyd, A. W.; Sutherland, G. R.: Localization of a human receptor tyrosine kinase (*ETK1*) to chromosome region 3p11.2. *Genomics* 19: 38–41, 1994.



- [81082] 19412.Wicks, I. P.; Wilkinson, D.; Salvaris, E.; Boyd, A. W.: Molecularcloning of HEK, the gene encoding a receptor tyrosine kinase expressedby human lymphoid tumor cell lines. Proc. Nat. Acad. Sci. 89: 1611–1615,1992.
- [81083] 19413.Elsea, S. H.; Mykytyn, K.; Ferrell, K.; Coulter, K. L.; Das, P.;Dubiel, W.; Patel, P. I.; Metherall, J. E.: Hemizygosity for theCOP9 signalosome subunit gene, SGN3, in the Smith–Magenis syndrome. Am.J. Med. Genet. 87: 342–348, 1999.
- [81084] 19414.Koyama, K.; Fukushima, Y.; Inazawa, J.; Tomot-sune, D.; Takahashi,N.; Nakamura, Y.: The human homologue of the murine Llglh gene (LLGL)maps within the Smith–Magenis syndrome region in 17p11.2. Cyto-genet.Cell Genet. 72: 78–82, 1996.
- [81085] 19415.Forsberg, E.; Hirsch, E.; Frohlich, L.; Meyer, M.; Ek-blom, P.;Aszodi, A.; Werner, S.; Fassler, R.: Skin wounds and severed nervesheal normally in mice lacking tenascin–C. Proc. Nat. Acad. Sci. 93:6594–6599, 1996.
- [81086] 19416.Gulcher, J. R.; Alexakos, M. J.; Le Beau, M. M.; Lemons, R. S.;Stefansson, K.: Chromosomal localization of the human hexabrachion(tenascin) gene and evidence for recent reduplication within the gene. Genomics 6:616–622, 1990.

- [81087] 19417. Gulcher, J. R.; Nies, D. E.; Alexakos, M. J.; Ravikant, N. A.; Sturgill, M. E.; Marton, L. S.; Stefansson, K.: Structure of the human hexabrachion (tenascin) gene. *Proc. Nat. Acad. Sci.* 88: 9438–9442, 1991.
- [81088] 19418. Mitrovic, N.; Schachner, M.: Detection of tenascin-C in the nervous system of the tenascin-C mutant mouse. *J. Neurosci. Res.* 42: 710–717, 1995.
- [81089] 19419. Nies, D. E.; Hemesath, T. J.; Kim, J.-H.; Gulcher, J. R.; Stefansson, K.: The complete cDNA sequence of human hexabrachion (tenascin): a multidomain protein containing unique epidermal growth factor repeats. *J. Biol. Chem.* 266: 2818–2823, 1991.
- [81090] 19420. Williams, T. M.; Moolten, D.; Burlein, J.; Romano, J.; Bhaerman, R.; Godillot, A.; Mellon, M.; Rauscher, F. J., III; Kant, J. A.: Identification of a zinc finger protein that inhibits IL-2 gene expression. *Science* 254: 1791–1794, 1991.
- [81091] 19421. Ludecke, H.-J.; Schaper, J.; Meinecke, P.; Momeni, P.; Gross, S.; von Holtum, D.; Hirche, H.; Abramowicz, M. J.; Albrecht, B.; Apacik, C.; Christen, H.-J.; Claussen, U.; and 28 others: Genotypic and phenotypic spectrum in trichorhino-phalangeal syndrome types I and III. *Am. J. Hum. Genet.* 68: 81–91, 2001.

- [81092] 19422. Chiang, L.; Contreras, L.; Chiang, J.; Ward, P. H.: Human prostatic gastricsinogen: the precursor of seminal fluid acid proteinase. *Arch. Biochem. Biophys.* 210: 14–20, 1981.
- [81093] 19423. Hayano, T.; Sogawa, K.; Ichihara, Y.; Fujii-Kuriyama, Y.; Takahashi, K.: Primary structure of human pepsinogen C gene. *J. Biol. Chem.* 263:1382–1385, 1988.
- [81094] 19424. Pals, G.; Azuma, T.; Mohandas, T. K.; Bell, G. I.; Bacon, J.; Samloff, I. M.; Walz, D. A.; Barr, P. J.; Taggart, R. T.: Human pepsinogen C (progastricsin) polymorphism: evidence for a single locus located at 6p21.1–pter. *Genomics* 4: 137–145, 1989.
- [81095] 19425. Randolph, L. M.; Azuma, T.; Petersen, G. M.; Sparkes, R. S.; Toyoda, M.; Wang, S. J.; Taggart, R. T.: Assignment of pepsinogen C (PGC) to 6p21.3–p21.1. (Abstract) *Cytogenet. Cell Genet.* 51: 1063–1064, 1989.
- [81096] 19426. Szymura, J. M.; Klein, J.: Linkage of a gene controlling urinary pepsinogen with the major histocompatibility complex of the mouse. *Immunogenetics* 13: 267–271, 1981.
- [81097] 19427. Taggart, R. T.; Cass, L. G.; Mohandas, T. K.; Derby, P.; Barr, P. J.; Pals, G.; Bell, G. I.: Human pepsinogen C (progastricsin): isolation of cDNA clones, localization to

chromosome 6, and sequence homology with pepsinogen  
A. J. Biol. Chem. 264: 375–379, 1989.

- [81098] 19428. Taggart, R. T.; Mohandas, T. K.; Bell, G. I.: Assignment of human preprogastricsin (PGC) to chromosome 6 and regional localization of PGC (6pter–p21.1), prolactin PRL (6pter–p21.1). (Abstract) Cytogenet. Cell Genet. 46: 701–702, 1987.
- [81099] 19429. Takahara, K.; Fukushige, S.; Murotsu, T.; Ichihara, Y.; Hayano, T.; Ishihara, T.; Takahashi, K.: Assignment of human pepsinogen C (PGC) gene to chromosome 6. Cytogenet. Cell Genet. 52: 100–101, 1989.
- [81100] 19430. Miwa, S.; Sato, T.; Murao, H.; Kozuru, M.; Ibayashi, H.: A new type of phosphofructokinase deficiency: hereditary nonspherocytic hemolytic anemia. Acta Haemat. Jpn. 35: 113–118, 1972.
- [81101] 19431. Bernardi, G.: Personal Communication. Paris, France 5/13/1992.
- [81102] 19432. Boulard, M. R.; Bois, M.; Reviron, M.; Najean, Y.: Red-cell phosphofructokinase deficiency. New Eng. J. Med. 291: 978–979, 1974.
- [81103] 19433. Chadeaux, B.; Rethore, M. O.; Allard, D.: Regional mapping of liver type 6-phosphofructokinase isoenzyme on chromosome 21. Hum. Genet. 68: 136–137, 1984.

- [81104] 19434.Cox, D. R.; Kawashima, H.; Vora, S.; Epstein, C. J.: Regional mapping of SOD-1, PRGS, and PFK-L on human chromosome 21. (Abstract) Cytogenet.Cell Genet. 37: 441-442, 1984.
- [81105] 19435.Elson, A.; Levanon, D.; Brandeis, M.; Dafni, N.; Bernstein, Y.;Danciger, E.; Groner, Y.: The structure of the human liver-type phosphofructokinase gene. Genomics 7: 47-56, 1990.
- [81106] 19436.Etiemble, J.; Kahn, A.; Boivin, P.; Bernard, J. F.; Goudemand,M.: Hereditary hemolytic anemia with erythrocyte phosphofructokinase deficiency. Hum. Genet. 31: 83-91, 1976.
- [81107] 19437.Etiemble, J.; Picat, C.; Simeon, J.; Blatrix, C.; Boivin, P.:Inherited erythrocyte phosphofructokinase deficiency: molecular mechanism. Hum.Genet. 55: 383-390, 1980.
- [81108] 19438.Gehnrich, S. C.; Gekakis, N.; Sul, H. S.: Liver (B-type) phosphofructokinase mRNA: cloning, structure, and expression. J. Biol. Chem. 263: 11755-11759,1988.
- [81109] 19439.Kahn, A.; Etiemble, J.; Meienhofer, M. C.; Boivin, P.: Erythrocyte phosphofructokinase deficiency associated with an unstable variant of muscle phosphofructokinase. Clin. Chim. Acta 61: 415-419, 1975.
- [81110] 19440.Levanon, D.; Danciger, E.; Dafni, N.; Groner, Y.:

Genomic clones of the human liver-type phosphofructokinase. *Biochem. Biophys. Res. Commun.* 141: 374–380, 1986.

- [81111] 19441. Pantelakis, S. N.; Karaklis, A. G.; Alexiou, D.; Vardas, E.; Valaes, T.: Red cell enzymes in trisomy 21. *Am. J. Hum. Genet.* 22: 184–193, 1970.
- [81112] 19442. Van Keuren, M.; Drabkin, H.; Hart, I.; Harker, D.; Patterson, D.; Vora, S.: Regional assignment of human liver-type 6-phosphofructokinase to chromosome 21q22.3 by using somatic cell hybrids and a monoclonal anti-L antibody. *Hum. Genet.* 74: 34–40, 1986.
- [81113] 19443. Vora, S.; Davidson, M.; Seaman, C.; Miranda, A. F.; Noble, N. A.; Tanaka, K. R.; Frenkel, E. P.; DiMauro, S.: Heterogeneity of the molecular lesions in inherited phosphofructokinase deficiency. *J. Clin. Invest.* 72: 1995–2006, 1983.
- [81114] 19444. Vora, S.; Durham, S.; de Martinville, B.; Francke, U.: Assignment of the genes for liver type phosphofructokinase (PFKL) to chromosome 21 and for muscle type (PFKM) to region p32–q32 of chromosome 1. (Abstract) *Cytogenet. Cell Genet.* 32: 324 only, 1982.
- [81115] 19445. Vora, S.; Francke, U.: Assignment of the human gene for liver type phosphofructokinase isozyme to chro-

mosome 21 using somatic cellhybrids. (Abstract) Pediat. Res. 15: 570 only, 1981.

[81116] 19446.Vora, S.; Francke, U.: Assignment of the human gene for liver-type6-phosphofructokinase isozyme (PFKL) to chromosome 21 by using somaticcell hybrids and monoclonal anti-L antibody. Proc. Nat. Acad. Sci. 78:3738–3742, 1981.

[81117] 19447.Wang, D.; Fang, H.; Cantor, C. R.; Smith, C. L.: A contiguousNotI restriction map of band q22.3 of human chromosome 21. Proc.Nat. Acad. Sci. 89: 3222–3226, 1992.

[81118] 19448.Vidal, R.; Revesz, T.; Rostagno, A.; Kim, E.; Holton, J. L.; Bek,T.; Bojsen-Moller, M.; Braendgaard, H.; Plant, G.; Ghiso, J.; Frangione,B.: A decamer duplication in the 3-prime region of the BRI gene originatesan amyloid peptide that is associated with dementia in a Danish kindred. Proc.Nat. Acad. Sci. 97: 4920–4925, 2000.

[81119] 19449.Fathallah-Shaykh, H.; Wolf, S.; Wong, E.; Posner, J. B.: Cloningof a leucine-zipper protein recognized by the sera of patients withantibody-associated paraneoplastic cerebellar degeneration. Proc.Nat. Acad. Sci. 88: 3451–3454, 1991.

[81120] 19450.Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.;

Posner, J. B.;Siniscalco, M.: In situ mapping of the gene coding for a leucinezipper DNA binding protein (CDR 2) to the region between two rarefragile sites of autosome 16 (16p12–p13.1). (Abstract) Cytogenet.Cell Genet. 58: 1999–2000, 1991.

- [81121] 19451.Gress, T.; Baldini, A.; Rocchi, M.; Furneaux, H.; Posner, J. B.;Siniscalco, M.: In situ mapping of the gene coding for a leucinezipper DNA binding protein (CDR62) to 16p12–16p13.1. Genomics 13:1340–1342, 1992.
- [81122] 19452.Raimondi, E.; Rubboli, F.; Moralli, D.; Chini, B.; Fornasari, D.;Tarroni, P.; De Carli, L.; Clementi, F.: Chromosomal localizationand physical linkage of the genes encoding the human alpha–3, alpha–5,and beta–4 neuronal nicotinic receptor subunits. Genomics 12: 849–850,1992.
- [81123] 19453.Anand, R.; Lindstrom, J.: Nucleotide sequence of the human nicotinicacetylcholine receptor beta–2 subunit gene. Nucleic Acids Res. 18:4272, 1990.
- [81124] 19454.De Fusco, M.; Becchetti, A.; Patrignani, A.; Annesi, G.; Gambardella,A.; Quattrone, A.; Ballabio, A.; Wanke, E.; Casari, G.: The nicotinicreceptor beta–2 subunit is mutant in nocturnal frontal lobe epilepsy. NatureGenet. 26: 275–276, 2000.
- [81125] 19455.Lueders, K. K.; Elliott, R. W.; Marenholz, I.; Mischke,



D.; DuPree, M.; Hamer, D.: Genomic organization and mapping of the human and mouse neuronal beta-1-nicotinic acetylcholine receptor genes. *Mammalian Genome* 10: 900–905, 1999.

- [81126] 19456. Phillips, H. A.; Favre, I.; Kirkpatrick, M.; Zuberi, S. M.; Goudie, D.; Heron, S. E.; Scheffer, I. E.; Sutherland, G. R.; Berkovic, S. F.; Bertrand, D.; Mulley, J. C.: CHRNA2 is the second acetylcholine receptor subunit associated with autosomal dominant nocturnal frontal lobe epilepsy. *Am. J. Hum. Genet.* 68: 225–231, 2001.
- [81127] 19457. Picciotto, M. R.; Zoli, M.; Lena, C.; Bessis, A.; Lallemand, Y.; LeNovere, N.; Vincent, P.; Pich, E. M.; Brulet, P.; Changeux, J.-P.: Abnormal avoidance learning in mice lacking functional high-affinity nicotine receptor in the brain. *Nature* 374: 65–67, 1995.
- [81128] 19458. Zoli, M.; Picciotto, M. R.; Ferrari, R.; Cocchi, D.; Changeux, J.-P.: Increased neurodegeneration during ageing in mice lacking high-affinity nicotine receptors. *EMBO J.* 18: 1235–1244, 1999.
- [81129] 19459. Tarroni, P.; Rubboli, F.; Chini, B.; Zwart, R.; Oortgiesen, M.; Sher, E.; Clementi, F.: Neuronal-type nicotinic receptors in human neuroblastoma and small-cell lung carcinoma cell lines. *FEBS Lett.* 312:66–70, 1992.

- [81130] 19460. Bonner, T. I.; Buckley, N. J.; Young, A. C.; Brann, M. R.: Identification of a family of muscarinic acetylcholine receptor genes. *Science* 237:527–532, 1987.
- [81131] 19461. Schwenk, W. F.; Rizza, R. A.; Mandarino, L. J.; Gerich, J. E.; Hayles, A. B.; Haymond, M. W.: Familial insulin resistance and acanthosis nigricans: presence of a post-binding defect. *Diabetes* 35: 33–37, 1986.
- [81132] 19462. Speckman, R. A.; Garg, A.; Du, F.; Bennett, L.; Veile, R.; Arioglu, E.; Taylor, S. I.; Lovett, M.; Bowcock, A. M.: Mutational and haplotype analyses of families with familial partial lipodystrophy (Dunnigan variety) reveal recurrent missense mutations in the globular C-terminal domain of lamin A/C. *Am. J. Hum. Genet.* 66: 1192–1198, 2000.  
Note: Erratum: *Am. J. Hum. Genet.* 67: 775 only, 2000.
- [81133] 19463. Chan, J.; Watt, V. M.: Eek and erk, new members of the eph subclass of receptor protein-tyrosine kinases. *Oncogene* 6: 1057–1061, 1991.
- [81134] 19464. Park, S.; Frisen, J.; Barbacid, M.: Aberrant axonal projections in mice lacking EphA8 (Eek) tyrosine protein kinase receptors. *EMBO J.* 16: 3106–3114, 1997.
- [81135] 19465. Ganju, P.; Shigemoto, K.; Brennan, J.; Entwistle, A.; Reith, A. D.: The Eck receptor tyrosine kinase is implicated in pattern formation during gastrulation, hindbrain seg-

mentation and limb development. *Oncogene* 9:1613–1624, 1994.

- [81136] 19466.Lindberg, R. A.; Hunter, T.: cDNA cloning and characterization of eck, an epithelial cell receptor–tyrosine kinase in the eph/elk family of protein kinases. *Molec. Cell. Biol.* 10: 6316–6324, 1990.
- [81137] 19467.Sulman, E. P.; Tang, X. X.; Allen, C.; Biegel, J. A.; Pleasure, D. E.; Brodeur, G. M.; Ikegaki, N.: ECK, a human EPH–related gene, maps to 1p36.1, a common region of alteration in human cancers. *Genomics* 40:371–374, 1997.
- [81138] 19468.Colucci, F.; Schweighoffer, E.; Tomasello, E.; Turner, M.; Ortaldo, J. R.; Vivier, E.; Tybulewicz, V. L. J.; Di Santo, J. P.: Natural cytotoxicity uncoupled from the Syk and ZAP–70 intracellular kinases. *Nature Immun.* 3: 288–294, 2002.
- [81139] 19469.Ku, G.; Malissen, B.; Mattei, M.–G.: Chromosomal location of the Syk and ZAP–70 tyrosine kinase genes in mice and humans. *Immunogenetics* 40:300–302, 1994.
- [81140] 19470.Toyabe, S.–I.; Watanabe, A.; Harada, W.; Karasawa, T.; Uchiyama, M.: Specific immunoglobulin E responses in ZAP–70–deficient patients are mediated by Syk–dependent T–cell receptor signalling. *Immunology* 103:164–171,

2001.

- [81141] 19471.Sudbrak, R.; Golla, A.; Hogan, K.; Powers, P.; Gregg, R.; Du Chesne,I.; Lehmann–Horn, F.; Deufel, T.: Exclusion of malignant hyperthermiasusceptibility (MHS) from a putative MHS2 locus on chromosome 17qand of the alpha–1, beta–1, and gamma subunits of the dihydropyridinereceptor calcium channel as candidates for the molecular defect. *Hum.Molec. Genet.* 2: 857–862, 1993.
- [81142] 19472.Naim, H. Y.; Sterchi, E. E.; Lentze, M. J.: Structure, biosynthesis,and glycosylation of human small intestinal maltase–glucoamylase. *J.Biol. Chem.* 263: 19709–19717, 1988.
- [81143] 19473.Nichols, B. L.; Eldering, J.; Avery, S.; Hahn, D.; Quaroni, A.;Sterchi, E.: Human small intestinal maltase–glucoamylase cDNA cloning:homology to sucrase–isomaltase. *J. Biol. Chem.* 273: 3076–3081, 1998.
- [81144] 19474.Schuler, G. D.; Boguski, M. S.; Stewart, E. A.; Stein, L. D.; Gyapay,G.; Rice, K.; White, R. E.; Rodriguez–Tome, P.; Aggarwal, A.; Bajorek,E.; Bentolila, S.; Birren, B. B.; Butler, A.; Castle, A. B.; Chiannilkulchai,N.; Chu, A.; Clee, C.; Cowles, S.; Day, P. J. R.; Dibling, T.; and84 others: A gene map of the human genome. *Science* 274: 540–546,1996.
- [81145] 19475.Tanahashi, N.; Suzuki, M.; Fujiwara, T.; Takahashi,

E.; Shimbara, N.; Chung, C. H.; Tanaka, K.: Chromosomal localization and immunological analysis of a family of human 26S proteasomal ATPases. *Biochem. Biophys. Res. Commun.* 243: 229–232, 1998.

[81146] 19476. Levine, A.; Cantoni, G. L.; Razin, A.: Inhibition of promoter activity by methylation: possible involvement of protein mediators. *Proc. Nat. Acad. Sci.* 88: 6515–6518, 1991.

[81147] 19477. Nan, X.; Meehan, R. R.; Bird, A.: Dissection of the methyl-CpG binding domain from the chromosomal protein MeCP2. *Nucleic Acids Res.* 21: 4886–4892, 1993.

[81148] 19478. Ohki, I.; Shimotake, N.; Fujita, N.; Jee, J.-G.; Ikegami, T.; Nakao, M.; Shirakawa, M.: Solution structure of the methyl-CpG binding domain of human MBD1 in complex with methylated DNA. *Cell* 105: 487–497, 2001.

[81149] 19479. Carrera, C. J.; Eddy, R. L.; Shows, T. B.; Carson, D. A.: Assignment of the gene for methylthioadenosine phosphorylase to human chromosome 9 by mouse-human somatic cell hybridization. *Proc. Nat. Acad. Sci.* 81: 2665–2668, 1984.

[81150] 19480. Chilcote, R. R.; Brown, E.; Rowley, J. D.: Lymphoblastic leukemia with lymphomatous features associated with abnormalities of the short arm of chromosome 9.

New Eng. J. Med. 313: 286–291, 1985.

- [81151] 19481.Nobori, T.; Takabayashi, K.; Tran, P.; Orvis, L.; Batova, A.; Yu, A. L.; Carson, D. A.: Genomic cloning of methylthioadenosine phosphorylase: a purine metabolic enzyme deficient in multiple different cancers. Proc. Nat. Acad. Sci. 93: 6203–6208, 1996.
- [81152] 19482.Olopade, O. I.; Jenkins, R. B.; Ransom, D. T.; Malik, K.; Pomykala, H.; Nobori, T.; Cowan, J. M.; Rowley, J. D.; Diaz, M. O.: Molecular analysis of deletions of the short arm of chromosome 9 in human gliomas. Cancer Res. 52: 2523–2529, 1992.
- [81153] 19483.Olopade, O. I.; Pomykala, H. M.; Hagos, F.; Sveen, L. W.; Espinosa, R., III; Dreyling, M. H.; Gursky, S.; Stadler, W. M.; Le Beau, M. M.; Bohlander, S. K.: Construction of a 2.8-megabase yeast artificial chromosome contig and cloning of the human methylthioadenosine phosphorylase gene from the tumor suppressor region on 9p21. Proc. Nat. Acad. Sci. 92: 6489–6493, 1995.
- [81154] 19484.Ragione, F. D.; Takabayashi, K.; Mastropietro, S.; Mercurio, C.; Oliva, A.; Russo, G. L.; Pietra, V. D.; Borriello, A.; Nobori, T.; Carson, D. A.; Zappia, V.: Purification and characterization of recombinant human 5-prime-methylthioadenosine phosphorylase: definite

identification of coding cDNA. *Biochem. Biophys. Res. Commun.* 223: 514–519, 1996.

- [81155] 19485. Williams–Ashman, H. G.; Seidenfeld, J.; Galletti, P.: Trends in the biochemical pharmacology of 5–prime–deoxy–5–prime–methylthioadenosine. *Biochem. Pharm.* 31: 277–288, 1982.
- [81156] 19486. Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human gene encoding methionyl–tRNA synthetase to chromosome 12. (Abstract) *Cytogenet. Cell Genet.* 37: 437 only, 1984.
- [81157] 19487. Cirullo, R. E.; Wasmuth, J. J.: Assignment of the human MARS gene, encoding methioninyl–tRNA synthetase, to chromosome 12 using human x Chinese hamster cell hybrids. *Somat. Cell Molec. Genet.* 10: 225–234, 1984.
- [81158] 19488. Grewal, P. K.; Holzfeind, P. J.; Bittner, R. E.; Hewitt, J. E.: Mutant glycosyltransferase and altered glycosylation of alpha–dystroglycan in the myodystrophy mouse. *Nature Genet.* 28: 151–154, 2001.
- [81159] 19489. Zabel, B. U.; Eddy, R. L.; Scott, J.; Shows, T. B.: The human nerve growth factor gene (NGF) is located on the short arm of chromosome 1. (Abstract) *Cytogenet. Cell Genet.* 37: 614, 1984.
- [81160] 19490. Adams, D. S.; Kiyokawa, M.; Getman, M. E.;

Shashoua, V. E.: Genes encoding giant danio and golden shiner ependymin. *Neurochem. Res.* 21:377–384, 1996.

[81161] 19491. de Groen, P. C.; Eggen, B. J. L.; Gispen, W. H.; Schotman, P.; Schrama, L. H.: Cloning and promoter analysis of the human B-50/GAP-43 gene. *J. Molec. Neurosci.* 6: 109–119, 1995.

[81162] 19492. Kosik, K. S.; Orecchio, L. D.; Bruns, G. A. P.; Benowitz, L. I.; MacDonald, G. P.; Cox, D. R.; Neve, R. L.: Human GAP-43: its deduced amino acid sequence and chromosomal localization in mouse and human. *Neuron* 1:127–132, 1988.

[81163] 19493. Reeves, R. H.; O'Hara, B. F.; Gearhart, J. D.: Localization on mouse chromosome 16 of Smst and Gap43: a conserved synteny with human chromosome 3. (Abstract) *Cytogenet. Cell Genet.* 51: 1064, 1989.

[81164] 19494. Strittmatter, S. M.; Fankhauser, C.; Huang, P. L.; Mashimo, H.; Fishman, M. C.: Neuronal pathfinding is abnormal in mice lacking the neuronal growth cone protein GAP-43. *Cell* 80: 445–452, 1995.

[81165] 19495. Bernalova, I. N.; Farjo, Q.; Mortlock, D. P.; Jackson, A. U.; Meisler, M. H.; Swaroop, A.; Burmeister, M.: Mapping of the neural retinal leucine zipper gene, *Nrl*, to mouse chromosome 14. *Mammalian Genome* 4:618–620, 1993.



- [81166] 19496. Bessant, D. A. R.; Payne, A. M.; Mitton, K. P.; Wang, Q.-L.; Swain, P. K.; Plant, C.; Bird, A. C.; Zack, D. J.; Swaroop, A.; Bhattacharya, S. S.: A mutation in NRL is associated with autosomal dominant retinitis pigmentosa. *Nature Genet.* 21: 355–356, 1999.
- [81167] 19497. Bessant, D. A. R.; Payne, A. M.; Plant, C.; Bird, A. C.; Swaroop, A.; Bhattacharya, S. S.: NRL S50T mutation and the importance of 'founder effects' in inherited retinal dystrophies. *Europ. J. Hum. Genet.* 8: 783–787, 2000.
- [81168] 19498. Dahl, S. P.; Jackson, A.; Kimberling, W. J.; Blackwood, D.; Swaroop, A.: Genetic mapping of NRL, a human retina-specific gene located on chromosome 14. (Abstract) *Am. J. Hum. Genet.* 51 (suppl.): A185 only, 1992.
- [81169] 19499. Farjo, Q.; Jackson, A.; Pieke-Dahl, S.; Scott, K.; Kimberling, W. J.; Sieving, P. A.; Richards, J. E.; Swaroop, A.: Human bZIP transcription factor gene NRL: structure, genomic sequence, and fine linkage mapping at 14q11.2 and negative mutation analysis in patients with retinal degeneration. *Genomics* 45: 395–401, 1997.
- [81170] 19500. Farjo, Q.; Jackson, A. U.; Xu, J.; Gryzenia, M.; Skolnick, C.; Agarwal, N.; Swaroop, A.: Molecular characterization of the murine neural retina leucine zipper gene, *Nrl*. *Genomics* 18: 216–222, 1993.

- [81171] 19501.Mears, A. J.; Kondo, M.; Swain, P. K.; Takada, Y.; Bush, R. A.;Saunders, T. L.; Sieving, P. A.; Swaroop, A.: Nrl is required forrod photoreceptor development. *Nature Genet.* 29: 447–452, 2001.
- [81172] 19502.Swaroop, A.; Xu, J.; Agarwal, N.; Weissman, S. M.: A simple andefficient cDNA library subtraction procedure: isolation of human retina–specificcDNA clones. *Nucleic Acids Res.* 19: 1954 only, 1991.
- [81173] 19503.Swaroop, A.; Xu, J.; Pawar, H.; Jackson, A.; Skolnick, C.; Agarwal,N.: A conserved retina–specific gene encodes a basic motif/leucinezipper domain. *Proc. Nat. Acad. Sci.* 89: 266–270, 1992.
- [81174] 19504.Wright, A. F.; Mansfield, D. C.; Bruford, E. A. Teague, P. W.;Thomson, K. L.; Riise, R.; Jay, M.; Patton, M. A.; Jeffery, S.; Schinzel,A.; Tommerup, N.: Fossarello, M.: Genetic studies in autosomal recessiveforms of retinitis pigmentosa.In: Anderson, R. E.; LaVail, M. M.;Hollyfield, J. G. (eds.): *Degenerative diseases of the retina*. New York: Plenum Press (pub.) 1995. Pp. 293–302.
- [81175] 19505.Yang–Feng, T. L.; Swaroop, A.: Neural retina–specific leucinezipper gene NRL (D14S46E) maps to human chromosome 14q11.1–q11.2. *Genomics* 14:491–492, 1992.

- [81176] 19506.Eddy, R. L.; Kretschmer, P. J.; Fairhurst, J. L.; Shows, T. B.; Bohlen, P.; O'Hara, B.; Kovesdi, I.: A human gene family of neuriteoutgrowth-promoting proteins: the gene for a heparin binding neuriteoutgrowth-promoting factor maps to 7q22-qter. (Abstract) Cytogenet.Cell Genet. 58: 1920 only, 1991.
- [81177] 19507.Lai, S.; Czubayko, F.; Riegel, A. T.; Wellstein, A.: Structureof the human heparin-binding growth factor gene pleiotrophin. Biochem.Biophys. Res. Commun. 187: 1113-1122, 1992.
- [81178] 19508.Li, Y.-S.; Hoffman, R. M.; Le Beau, M. M.; Espinosa, R., III; Jenkins,N. A.; Gilbert, D. J.; Copeland, N. G.; Deuel, T. F.: Characterizationof the human pleiotrophin gene: promoter region and chromosomal localization. J.Biol. Chem. 267: 26011-26016, 1992.
- [81179] 19509.Li, Y.-S.; Milner, P. G.; Chauhan, A. K.; Watson, M. A.; Hoffman,R. M.; Kodner, C. M.; Milbrandt, J.; Deuel, T. F.: Cloning and expressionof a developmentally regulated protein that induces mitogenic andneurite outgrowth activity. Science 250: 1690-1694, 1990.
- [81180] 19510.Milner, P. G.; Shah, D.; Veile, R.; Donis-Keller, H.; Kumar, B.V.: Cloning, nucleotide sequence, and chromosome localization ofthe human pleiotrophin gene. Bio-

chemistry 31: 12023–12028, 1992.

- [81181] 19511. Brightbill, H. D.; Libraty, D. H.; Krutzik, S. R.; Yang, R.-B.; Bellsie, J. T.; Bieharski, J. R.; Maitland, M.; Norgard, M. V.; Plevy, S. E.; Smale, S. T.; Brennan, P. J.; Bloom, B. R.; Godowski, P. J.; Modlin, R. L.: Host defense mechanisms triggered by microbial lipoproteins through toll-like receptors. *Science* 285: 732–736, 1999.
- [81182] 19512. Jankovic, D.; Kullberg, M. C.; Hieny, S.; Caspar, P.; Collazo, C. M.; Sher, A.: In the absence of IL-12, CD4<sup>+</sup> T cell responses to intracellular pathogens fail to default to a Th2 pattern and are host protective in an IL-10<sup>-/-</sup> setting. *Immunity* 16: 429–439, 2002.
- [81183] 19513. Jouanguy, E.; Lamhamedi-Cherradi, S.; Altare, F.; Fondaneche, M.-C.; Tuerlinckx, D.; Blanche, S.; Emile, J.-F.; Gaillard, J.-L.; Schreiber, R.; Levin, M.; Fischer, A.; Hivroz, C.; Casanova, J.-L.: Partial interferon- $\gamma$  receptor 1 deficiency in a child with tuberculoid bacillus Calmette-Guerin infection and a sibling with clinical tuberculosis. *J. Clin. Invest.* 100: 2658–2664, 1997.
- [81184] 19514. Warrington, J. A.; Bengtsson, U.: High-resolution physical mapping of human 5q31–q33 using three methods: radiation hybrid mapping, interphase fluorescence in situ hybridization, and pulsed-field gel electrophoresis.

Genomics 24:395–398, 1994.

- [81185] 19515. Anderson, S. K.; Gallinger, S.; Roder, J.; Frey, J.; Young, H.A.; Ortaldo, J. R.: A cyclophilin-related protein involved in the function of natural killer cells. Proc. Nat. Acad. Sci. 90: 542–546, 1993.
- [81186] 19516. Frey, J. L.; Bino, T.; Kantor, R. R. S.; Segal, D. M.; Giardina, S. L.; Roder, J.; Anderson, S.; Ortaldo, J. R.: Mechanism of target cell recognition by natural killer cells: characterization of a novel triggering molecule restricted to CD3– large granular lymphocytes. J. Exp. Med. 174: 1527–1536, 1991.
- [81187] 19517. Rinfret, A.; Anderson, S. K.: IL-2 regulates the expression of the NK-TR gene via an alternate RNA splicing mechanism. Molec. Immun. 30:1307–1313, 1993.
- [81188] 19518. Simons-Evelyn, M.; Young, H. A.; Anderson, S. K.: Characterization of the mouse Nktr gene and promoter. Genomics 40: 94–100, 1997.
- [81189] 19519. Young, H. A.; Jenkins, N. A.; Copeland, N. G.; Simek, S.; Lerman, M. I.; Zbar, B.; Glenn, G.; Ortaldo, J. R.; Anderson, S. K.: Localization of a novel natural killer triggering receptor locus to human chromosome 3p23–p21 and mouse chromosome 9. Genomics 16: 548–549, 1993.
- [81190] 19520. Scott, I. C.; Halila, R.; Jenkins, J. M.; Mehan, S.;

Apostolou,S.; Winqvist, R.; Callen, D. F.; Prockop, D. J.; Peltonen, L.; Kadler,K. E.: Molecular cloning, expression and chromosomal localization of a human gene encoding a 33 kDa putative metallopeptidase (PRSM1). *Gene* 174:135–143, 1996.

[81191] 19521.Aliprantis, A. O.; Yang, R.-B.; Mark, M. R.; Suggett, S.; Devaux,B.; Radolf, J. D.; Klimpel, G. R.; Godowski, P.; Zychlinsky, A.:Cell activation and apoptosis by bacterial lipoproteins through toll-like receptor-2. *Science* 285: 736–739, 1999.

[81192] 19522.Baldwin, A. S., Jr.: The NF-kappa-B and I-kappa-B proteins: new discoveries and insights. *Annu. Rev. Immun.* 14: 649–683, 1996.

[81193] 19523.Barnes, P. J.; Karin, M.: Nuclear factor-kappa-B -- A pivotal transcription factor in chronic inflammatory diseases. *New Eng. J.Med.* 336: 1066–1071, 1997.

[81194] 19524.Chen, F.; Castranova, V.; Shi, X.; Demers, L. M.: New insights into the role of nuclear factor-kappa-B, a ubiquitous transcription factor in the initiation of diseases. *Clin. Chem.* 45: 7–17, 1999.

[81195] 19525.Heron, E.; Deloukas, P.; van Loon, A. P. G. M.: The complete exon-intron structure of the 156-kb human gene NFKB1, which encodes the p105 and p50 proteins of

transcription factors NF-kappa-B and I-kappa-B-gamma: implications for NF-kappa-B-mediated signal transduction. *Genomics* 30:493-505, 1995.

- [81196] 19526. Huxford, T.; Huang, D.-B.; Malek, S.; Ghosh, G.: The crystal structure of the I-kappa-B-alpha/NF-kappa-B complex reveals mechanisms of NF-kappa-B inactivation. *Cell* 95: 759-770, 1998.
- [81197] 19527. Lin, L.; DeMartino, G. N.; Greene, W. C.: Cotranslational biogenesis of NF-kappa-B p50 by the 26S proteasome. *Cell* 92: 819-828, 1998.
- [81198] 19528. Liptay, S.; Schmid, R. M.; Perkins, N. D.; Meltzer, P.; Altherr, M. R.; McPherson, J. D.; Wasmuth, J. J.; Nabel, G. J.: Related subunits of NF-kappa-B map to two distinct loci associated with translocations in leukemia, NFKB1 and NFKB2. *Genomics* 13: 287-292, 1992.
- [81199] 19529. Mathew, S.; Murty, V. V. V. S.; Dalla-Favera, R.; Chaganti, R. S. K.: Chromosomal localization of genes encoding the transcription factors, c-rel, NF-kappa-Bp50, NF-kappa-Bp65, and Iy-10 by fluorescence in situ hybridization. *Oncogene* 8: 191-193, 1993.
- [81200] 19530. Meyer, R.; Hatada, E. N.; Hohmann, H. P.; Haiker, M.; Bartsch, C.; Rothlisberger, U.; Lahm, H. W.; Schlaeger, E. J.; van Loon, A. P.; Scheidereit, C.: Cloning of the DNA-

binding subunit of human nuclear factor kappa B: the level of its mRNA is strongly regulated by phorbol ester or tumor necrosis factor alpha. Proc. Nat. Acad. Sci. 88: 966–970, 1991.

[81201] 19531. Ozes, O. N.; Mayo, L. D.; Gustin, J. A.; Pfeffer, S. R.; Pfeffer, L. M.; Donner, D. B.: NF-kappa-B activation by tumor necrosis factor requires the Akt serine-threonine kinase. Nature 401: 82–85, 1999.

[81202] 19532. Romashkova, J. A.; Makarov, S. S.: NF-kappa-B is a target of AKT in anti-apoptotic PDGF signalling. Nature 401: 86–90, 1999.

[81203] 19533. Ryan, K. M.; Ernst, M. K.; Rice, N. R.; Voudsen, K. H.: Role of NF-kappa-B in p53-mediated programmed cell death. Nature 404: 892–897, 2000.

[81204] 19534. Sen, R.; Baltimore, D.: Multiple nuclear factors interact with the immunoglobulin enhancer sequences. Cell 46: 705–716, 1986.

[81205] 19535. Sha, W. C.; Liou, H.-C.; Tuomanen, E. I.; Baltimore, D.: Targeted disruption of the p50 subunit of NF-kappa-B leads to multifocal defects in immune responses. Cell 80: 321–330, 1995.

[81206] 19536. Zhong, H.; May, M. J.; Jimi, E.; Ghosh, S.: The phosphorylation status of nuclear NF-kappa-B determines its



association with CBP/p300 or HDAC-1. *Molec. Cell* 9: 625–636, 2002.

- [81207] 19537. Claudio, E.; Brown, K.; Park, S.; Wang, H.; Siebenlist, U.: BAFF-induced NEMO-independent processing of NF- $\kappa$ B2 in maturing B cells. *Nature Immun.* 3: 958–965, 2002.
- [81208] 19538. Neri, A.; Chang, C.-C.; Lombardi, L.; Salina, M.; Corradini, P.; Maiolo, A. T.; Chaganti, R. S. K.; Dalla-Favera, R.: B cell lymphoma-associated chromosomal translocation involves candidate oncogene *lyt-10*, homologous to NF- $\kappa$ B p50. *Cell* 67: 1075–1087, 1991.
- [81209] 19539. Taparowsky, E.; Shimizu, K.; Goldfarb, M.; Wigler, M.: Structure and activation of the human N-ras gene. *Cell* 34: 581–586, 1983.
- [81210] 19540. Yuasa, Y.; Gol, R. A.; Chang, A.; Chiu, I.-M.; Reddy, E. P.; Tronick, S. R.; Aaronson, S. A.: Mechanism of activation of an N-ras oncogene of SW-1271 human lung carcinoma cells. *Proc. Nat. Acad. Sci.* 81: 3670–3674, 1984.
- [81211] 19541. Bakin, A. V.; Curran, T.: Role of DNA 5-methylcytosine transferase in cell transformation by fos. *Science* 283: 387–390, 1999.
- [81212] 19542. Barker, P. E.; Rabin, M.; Watson, M.; Breg, W. R.; Ruddle, F. H.; Verma, I. M.: Human c-fos oncogene

mapped within chromosomal region 14q21–q31. Proc. Nat. Acad. Sci. 81: 5826–5830, 1984.

[81213] 19543. Dony, C.; Gruss, P.: Proto-oncogene c-fos expression in growth regions of fetal bone and mesodermal web tissue. Nature 328: 711–714, 1987.

[81214] 19544. Ekstrand, A. J.; Zech, L.: Human c-fos proto-oncogene mapped to chromosome 14, band q24.3–q31: possibilities for oncogene activation by chromosomal rearrangements in human neoplasms. Exp. Cell Res. 169:262–266, 1987.

[81215] 19545. Glover, J. N. M.; Harrison, S. C.: Crystal structure of the heterodimeric bZIP transcription factor c-Fos–c-Jun bound to DNA. Nature 373:257–261, 1995.

[81216] 19546. Grigoriadis, A. E.; Wang, Z.-Q.; Cecchini, M. G.; Hofstetter, W.; Felix, R.; Fleisch, H. A.; Wagner, E. F.: c-Fos: a key regulator of osteoclast–macrophage lineage determination and bone remodeling. Science 266:443–448, 1994.

[81217] 19547. Grigoriadis, A. E.; Wang, Z.-Q.; Wagner, E. F.: Fos and bone cell development: lessons from a nuclear oncogene. Trends Genet. 11:436–441, 1995.

[81218] 19548. Hayflick, L.: The limited in vitro lifetime of human diploid cell strains. Exp. Cell Res. 37: 614–636, 1965.

[81219] 19549. Johnson, R. S.; Spiegelman, B. M.; Papaioannou, V.:

Pleiotropic effects of a null mutation in the c-fos proto-oncogene. *Cell* 71:577–586, 1992.

[81220] 19550. Muller, R.; Tremblay, J. M.; Adamson, E. D.; Verma, I. M.: Tissue and cell type-specific expression of two human c-onc genes. *Nature* 304:454–456, 1983.

[81221] 19551. Rogaev, E. I.; Lukiw, W. J.; Vaula, G.; Haines, J. L.; Rogaeva, E. A.; Tsuda, T.; Alexandrova, N.; Liang, Y.; Mortilla, M.; Amaducci, L.; Bergamini, L.; Bruni, A. C.; Foncin, J.-F.; Macciardi, F.; Montesi, M. P.; Sorbi, S.; Rainero, I.; Pinessi, L.; Polinsky, R. J.; Frommelt, P.; Duara, R.; Lopez, R.; Pollen, D.; Gusella, J. F.; Tanzi, R.; Crapper MacLachlan, D.; St. George-Hyslop, P. H.: Analysis of the c-FOS gene on chromosome 14 and the promoter of the amyloid precursor protein gene in familial Alzheimer's disease. *Neurology* 43: 2275–2279, 1993.

[81222] 19552. Ruther, U.; Garber, C.; Komitowski, D.; Muller, R.; Wagner, E. F.: Deregulated c-fos expression interferes with normal bone development in transgenic mice. *Nature* 325: 412–416, 1987.

[81223] 19553. Saez, E.; Rutberg, S. E.; Mueller, E.; Oppenheim, H.; Smoluk, J.; Yuspa, S. H.; Spiegelman, B. M.: c-fos is required for malignant progression of skin tumors. *Cell* 82: 721–732, 1995.

- [81224] 19554.Sanyal, S.; Sandstrom, D. J.; Hoeffler, C. A.; Ramaswami, M.:AP-1 function upstream of CREB to control synaptic plasticity in *Drosophila*. *Nature* 416:870–874, 2002.
- [81225] 19555.Seshadri, T.; Campisi, J.: Repression of c-fos transcription and an altered genetic program in senescent human fibroblasts. *Science* 247:205–209, 1990.
- [81226] 19556.van Straaten, F.; Muller, R.; Curran, T.; Van Beveren, C.; Verma, I. M.: Complete nucleotide sequence of a human c-onc gene: deduced amino acid sequence of the human c-fos protein. *Proc. Nat. Acad.Sci.* 80: 3183–3187, 1983.
- [81227] 19557.Visvader, J.; Sassone-Corsi, P.; Verma, I. M.: Two adjacent promoter elements mediate nerve growth factor activation of the c-fos gene and bind distinct nuclear complexes. *Proc. Nat. Acad. Sci.* 85: 9474–9478, 1988.
- [81228] 19558.Wang, Z.-Q.; Grigoriadis, A. E.; Mohle-Steinlein, U.; Wagner, E. F.: A novel target cell for c-fos-induced oncogenesis: development of chondrogenic tumours in embryonic stem cell chimeras. *EMBO J.* 10:2437–2450, 1991.
- [81229] 19559.Wang, Z.-Q.; Ovitt, C.; Grigoriadis, A. E.; Mohle-Steinlein, U.; Ruther, U.; Wagner, E. F.: Bone and haematopoietic defects in mice lacking c-fos. *Nature* 360:

741–745, 1992.

- [81230] 19560.Zhang, J.; Zhang, D.; McQuade, J. S.; Behbehani, M.; Tsien, J.Z.; Xu, M.: c-fos regulates neuronal excitability and survival. *NatureGenet.* 30: 416–420, 2002.
- [81231] 19561.Arheden, K.; Mandahl, N.; Strombeck, B.; Isaksson, M.; Mitelman, F.: Chromosome localization of the human oncogene INT1 to 12q13 by in situ hybridization. *Cytogenet. Cell Genet.* 47: 86–87, 1988.
- [81232] 19562.Gavin, B. J.; McMahon, J. A.; McMahon, A. P.: Expression of multiple novel Wnt-1/int-1-related genes during fetal and adult mouse development. *GenesDev.* 4: 2319–2332, 1990.
- [81233] 19563.Kirikoshi, H.; Sekihara, H.; Katoh, M.: WNT10A and WNT6, clustered in human chromosome 2q35 region with head-to-tail manner, are strongly coexpressed in SW480 cells. *Biochem. Biophys. Res. Commun.* 283:798–805, 2001.
- [81234] 19564.McMahon, A. P.: The Wnt family of developmental regulators. *TrendsGenet.* 8: 236–242, 1992.
- [81235] 19565.Nusse, R.; Brown, A.; Papkoff, J.; Scambler, P.; Shackleford, G.; McMahon, A.; Moon, R.; Varmus, H.: A new nomenclature for int-1 and related genes: the Wnt gene family. (Letter) *Cell* 64: 231–232, 1991.

- [81236] 19566.Nusse, R.; van't Veer, L.; Geurts van Kessel, A.; van Agthoven,A.; Bootsma, D.; Varmus, H.: Chromosomal localization of a humanhomologue of a putative mammary tumor oncogene. (Abstract) Cytogenet.Cell Genet. 37: 556–557, 1984.
- [81237] 19567.Pellegrino, J. E.; Lensch, M. W.; Muenke, M.; Chance, P. F.: Clinicaland molecular analysis in Joubert syndrome. Am. J. Med. Genet. 72:59–62, 1997.
- [81238] 19568.Thomas, K. R.; Capecchi, M. R.: Targeted disruption of the murineint–1 proto–oncogene resulting in severe abnormalities in midbrainand cerebellar development. Nature 346: 847–850, 1990.
- [81239] 19569.Braaten, D.; Wellington, S.; Warburton, D.; Luban, J.: Assignmentof cyclophilin A (PPIA) to human chromosome band 7p13 by in situ hybridization. Cytogenet.Cell Genet. 74: 262 only, 1996.
- [81240] 19570.Liu, J.; Albers, M. W.; Chen, C.–M.; Schreiber, S. L.; Walsh, C.T.: Cloning, expression, and purification of human cyclophilin inEscherichia coli and assessment of the catalytic role of cysteinesby site–directed mutagenesis. Proc. Nat. Acad. Sci. 87: 2304–2308,1990.
- [81241] 19571.Luban, J.; Bossolt, K. L.; Franke, E. K.; Kalpana, G. V.; Goff,S. P.: Human immunodeficiency virus type 1 gag

protein binds to cyclophilins A and B. *Cell* 73: 1067–1078, 1993.

[81242] 19572. Takahashi, N.; Hayano, T.; Suzuki, M.: Peptidyl-prolyl cis-trans isomerase is the cyclosporin A-binding protein cyclophilin. *Nature* 337:473–475, 1989.

[81243] 19573. Willenbrink, W.; Halaschek, J.; Schuffenhauer, S.; Kunz, J.; Steinkasserer, A.: Cyclophilin A, the major intracellular receptor for the immunosuppressant cyclosporin A, maps to chromosome 7p11.2–p13: four pseudogenes map to chromosomes 3, 10, 14, and 18. *Genomics* 28: 101–104, 1995.

[81244] 19574. Dickinson, D. P.; Zhao, Y.; Thiesse, M.; Siciliano, M. J.: Direct mapping of seven genes encoding human type 2 cystatins to a single site located at 20p11.2. *Genomics* 24: 172–175, 1994.

[81245] 19575. Saitoh, E.; Isemura, S.; Sanada, K.; Ohnishi, K.: The human cysteine gene family: cloning of three members and evolutionary relationship between cystatins and Bowman-Birk type proteinase inhibitors. *Biomed. Biochim. Acta* 50: 599–605, 1991.

[81246] 19576. Takahashi, E.; Hori, T.; O'Connell, P.; Leppert, M.; White, R.: R-banding and nonisotopic in situ hybridization: precise localization of the human type II collagen

gene (COL2A1). Hum. Genet. 86: 14–16,1990.

[81247] 19577.Takahashi, E.; Yamauchi, M.; Tsuji, H.; Hitomi, A.; Meuth, M.;Hori, T.: Chromosome mapping of the human cytidine–5–prime–triphosphatesynthetase (CTPS) gene to band 1p34.1–p34.3 by fluorescence in situhybridization. Hum. Genet. 88: 119–121, 1991.

[81248] 19578.Takahashi, E.–I.; Yamauchi, M.; Ayusawa, D.; Kaneda, S.; Seno,T.; Meuth, M.; Hori, T.–A.: Chromosome mappings of the human cytidine–5–prime–triphosphatesynthetase (CTPS) gene and the human ubiquitin–activating enzyme UBE1gene by fluorescence in situ hybridization. (Abstract) Cytogenet.Cell Genet. 58: 1864 only, 1991.

[81249] 19579.Thomas, P. E.; Sen, S.; Lamb, B. J.; Chu, E. H. Y.: Cloning andexpression of mammalian CTP synthetase genes. (Abstract) Am. J. Hum.Genet. 45 (suppl.): A11 only, 1989.

[81250] 19580.Whelan, J.; Phear, G.; Yamauchi, M.; Meuth, M.: Clustered basesubstitutions in CTP synthetase conferring drug resistance in Chinesehamster ovary cells. Nature Genet. 3: 317–322, 1993.

[81251] 19581.Yamauchi, M.; Takahashi, E.; Whelan, J.; Phear, G.; Meuth, M.:Mapping and functional analysis of the cytidine



triphosphate synthetase(CTPS) gene. (Abstract) Human Genome Mapping Workshop 93 1 only,1993.

- [81252] 19582.Nebert, D. W.; Gonzalez, F. J.: P450 genes: structure, evolution,and regulation. Annu. Rev. Biochem. 56: 945–993, 1987.
- [81253] 19583.Paolini, M.; Cantelli–Forti, G.; Perocco, P.; Pedulli, G. F.;Abdel–Rahman, S. Z.; Legator, M. S.: Co–carcinogenic effect of beta–carotene.(Letter) Nature 398: 760–761, 1999.
- [81254] 19584.Espinoza, H.; Cox, C. J.; Semina, E. V.; Amendt, B. A.: A molecularbasis for differential developmental anomalies in Axenfeld–Riegersyndrome. Hum. Molec. Genet. 11: 743–753, 2002.
- [81255] 19585.Leticic, K.; Zoncu, R.; Rakic, P.: Origin of GABAergic neuronsin the human neocortex. Nature 417: 645–649, 2002.
- [81256] 19586.McGuinness, T.; Porteus, M. H.; Smiga, S.; Bulfone, A.; Kingsley,C.; Qiu, M.; Liu, J. K.; Long, J. E.; Xu, D.; Rubenstein, J. L. R.: Sequence, organization, and transcription of the Dlx–1 and the Dlx–2locus. Genomics 35: 473–485, 1996.
- [81257] 19587.Ozcelik, T.; Porteus, M. H.; Rubenstein, J. L. R.; Francke, U.: DLX2 (TES1), a homeobox gene of the Distal–

- less family, assigned to conserved regions on human and mouse chromosomes 2. *Genomics* 13:1157–1161, 1992.
- [81258] 19588. Porteus, M. H.; Brice, A. E. J.; Bulfone, A.; Usdin, T. B.; Ciaranello, R. D.; Rubenstein, J. L. R.: Isolation and characterization of a library of cDNA clones that are preferentially expressed in the embryonic telencephalon. *Molec. Brain Res.* 12: 7–22, 1992.
- [81259] 19589. Qiu, M.; Bulfone, A.; Martinez, S.; Meneses, J. J.; Shimamura, K.; Pedersen, R. A.; Rubenstein, J. L.: Null mutation of *Dlx-2* results in abnormal morphogenesis of proximal first and second branchial arch derivatives and abnormal differentiation in the forebrain. *Genes Dev.* 9: 2523–2538, 1995.
- [81260] 19590. Salvador, J. M.; Hollander, M. C.; Nguyen, A. T.; Kopp, J. B.; Barisoni, L.; Moore, J. K.; Ashwell, J. D.; Fornace, A. J., Jr.: Mice lacking the p53-effector gene *Gadd45a* develop a lupus-like syndrome. *Immunity* 16:499–508, 2002.
- [81261] 19591. Takekawa, M.; Saito, H.: A family of stress-inducible GADD45-like proteins mediate activation of the stress-responsive MTK1/MEKK4 MAPKKK. *Cell* 95:521–530, 1998.
- [81262] 19592. Dhermy, D.; Garbarz, M.; Lecomte, M. C.; Feo, C.;

Bournier, O.;Chaveroche, I.; Gautero, H.; Galand, C.;  
Boivin, P.: Hereditary elliptocytosis:clinical, morphological  
and biochemical studies of 38 cases. *Nouv.Rev. Franc.  
Hemat.* 28: 129–140, 1986.

[81263] 19593.Fairbanks, G.; Steck, T. L.; Wallach, D. F. H.: Elec-  
trophoreticanalysis of the major polypeptides of the hu-  
man erythrocyte membrane. *Biochemistry* 10:2606–2617,  
1971.

[81264] 19594.Feo, C. J.; Fischer, S.; Piau, J. P.; Grange, M. J.; Tch-  
ernia,G.: Premiere observation de l'absence d'une proteine  
de la membraneerythrocytaire (bande 4–1) dans un cas  
d'anemie elliptocytaire familiale. *Nouv.Rev. Franc. Hemat.*  
22: 315–325, 1980.

[81265] 19595.Eddy, R. L.; Mahnke–Zizelman, D. K.; Bausch–Ju-  
rken, M. T.; Sabina,R. L.; Shows, T. B.: Distribution of the  
AMP deaminase multigenefamily within the human  
genome: assignment of the AMPD2 to chromo-  
some1p21–p34 and AMPD3 to chromosome 11p13–pter.  
(Abstract) *Human GenomeMapping Workshop* 93 24 only,  
1993.

[81266] 19596.Ogasawara, N.; Goto, H.; Yamada, Y.; Nishigaki, I.;  
Itoh, T.; Hasegawa,I.; Park, K. S.: Deficiency of AMP deam-  
inase in erythrocytes. *Hum.Genet.* 75: 15–18, 1987.

- [81267] 19597.Sermsuvitayawong, K.; Wang, X.; Nagabukuro, A.; Matsuda, Y.; Morisaki,H.; Toyama, K.; Mukai, T.; Morisaki, T.: Genomic organization ofAmpd3, heart-type AMPD gene, located in mouse chromosome 7. Mam-malianGenome 8: 767-769, 1997.
- [81268] 19598.Yamada, Y.; Goto, H.; Ogasawara, N.: Cloning and nucleotide sequenceof the cDNA encoding human erythrocyte-specific AMP deaminase. Biochim.Biophys. Acta 1171: 125-128, 1992.
- [81269] 19599.Yamada, Y.; Goto, H.; Ogasawara, N.: A point mutation responsiblefor human erythrocyte AMP deaminase deficiency. Hum. Molec. Genet. 3:331-334, 1994.
- [81270] 19600.Allegrucci, C.; Liguori, L.; Mezzasoma, I.; Minelli, A.: A1 adenosinereceptor in human spermatozoa: its role in the fertilization process. Molec.Genet. Metab. 71: 381-386, 2000.
- [81271] 19601.Boman, H.; Hermodson, M.; Hammond, C. A.; Motulsky, A. G.: Analbuminemiain an American Indian girl. Clin. Genet. 9: 513-526, 1976.